Moduli Axions, Stabilizing Moduli and the Large Field Swampland Conjecture in Heterotic M-Theory

Cédric Deffayet,^a Burt A. Ovrut^b and Paul J. Steinhardt^c

 a Laboratoire de Physique de l'Ecole Normale Supérieure, ENS, Université PSL, CNRS Sorbonne Université, Université Paris Cité, F-75005 Paris, France b Department of Physics, University of Pennsylvania Philadelphia, PA 19104, USA c Department of Physics, Princeton University

Princeton, New Jersey 08544, USA

 d Jefferson Physical Laboratory, Harvard University Cambridge MA 02138 USA

Abstract

We compute the F- and D-term potential energy for the dilaton, complex structure and K¨ahler moduli of realistic vacua of heterotic Mtheory compactified on Calabi-Yau threefolds where, for simplicity, we choose $h^{1,1} = h^{2,1} = 1$. However, the formalism is immediately applicable to the "universal" moduli of Calabi-Yau threefolds with $h^{1,1} = h^{1,2} > 1$ as well. The F-term potential is computed using the non-perturbative complex structure, gaugino condensate and "worldsheet instanton" superpotentials in theories in which the hidden sector contains an anomalous $U(1)$ structure group. The Green-Schwarz anomaly cancellation induces inhomogeneous "axion" transformations for the imaginary components of the dilaton and Kähler modulus $-$

which then produce a D-term potential. V_D is a function of the real components of the dilaton and Kähler modulus (s and t) that is minimized and precisely vanishes along a unique line in the s-t plane. Excitations transverse to this line have a mass m_{anom} which is an explicit function of t. The F-term potential energy is then evaluated along the $V_D = 0$ line. For values of t small enough that $m_{\text{anom}} \gtrsim M_U$ – where M_U is the compactification scale – we plot V_F for a realistic choice of coefficients as a function of the Pfaffian parameter p . We find values of p for which V_F has a global minimum at negative or zero vacuum density or a metastable minimum with positive vacuum density. In all three cases, the s, t and associated "axion" moduli are completely stabilized. Finally, we show that, for any of these vacua, the large t behavior of the potential energy satisfies the "large scalar field" Swampland conjecture.

cedric.deffayet@phys.ens.fr, ovrut@elcapitan.hep.upenn.edu, steinh@princeton.edu

1 Introduction

In this paper, we calculate and discuss the potential energy for the complex structure moduli, the K¨ahler moduli and the dilaton in heterotic Mtheory vacua [1, 2, 3, 4] whose hidden sector contains a line bundle with an anomalous $U(1)$ structure group appropriately embedded into E_8 [5]. The Green-Schwarz cancellation $[6]$ of this $U(1)$ anomaly induces an inhomogeneous transformation of the imaginary components of the dilaton and each of the complex Kähler moduli. Thus, these imaginary components act as moduli "axions". The formalism introduced here is valid for any Calabi-Yau threefold with $h^{1,1} = h^{2,1} \geq 1$. However, for simplicity, we present our specific results for Calabi-Yau threefolds X with $h^{1,1} = h^{2,1} = 1$. These results are immediately applicable to the "universal" Kähler and complex structure moduli of Calabi-Yau compactifications with $h^{1,1} = h^{2,1} > 1$ as well. We note that moduli stabilization has been discussed in a number of papers whose contexts differ from the present work – see $[7, 8, 9, 10]$ for example.

The Kähler potential, K , and superpotential, W_{flux} , for the complex structure moduli of a Calabi-Yau threefold with any $h^{2,1} \geq 1$ are well-known for "large" values of these moduli [11, 12]. However, these quantities are not known analytically for "small" values. Hence, for specificity, we will conduct our analysis using the large modulus formalism. We do not expect this to change the basic results of the potential energy calculation. In Section 2, assuming $h^{2,1} = 1$, we compute the F-term potential energy V_{flux} for the complex structure modulus alone and show that there exists an infinite number of local minima that do *not* spontaneously break $N = 1$ supersymmetry. Our complete extended calculation including both the dilaton and Kähler modulus will be carried out under the assumption that the complex structure modulus has been *fixed* to any one of these supersymmetric minima.

Having done this, in Section 3 we extend the calculation of V_F to include the dilaton. This is accomplished under the assumption that the commutant of the anomalous $U(1)$ structure group contains a non-Abelian group that becomes strongly coupled at a high scale, thus leading to gaugino condensation [13, 14, 15, 16]. This produces a non-perturbative superpotential W_G which, to lowest order, depends only on the complex dilaton field S. Using this, and the associated dilaton Kahler potential K_S , we extend our calculation of the V_F potential energy using $W = W_{flux} + W_G$.

The cohomology $H^{1,1}$ is associated with the homology group H_2 . For $h^{1,1} = 1$, H_2 contains a single homology class. We denote this by [C], where C is an isolated, genus-zero holomorphic curve. It is well-known that the string can wrap itself around such a curve, producing a non-perturbative "instanton" superpotential $[17, 18, 19]$ given by an exponential of the Kähler modulus T multiplied by the "Pfaffian" associated with the Dirac operator. The Pfaffian is a holomorphic function of a subset of the vector bundle moduli evaluated at the curve C [20, 21]. Generically, C is not unique – with the number of such isolated, genus-zero curves in $[C]$ given by the Gromov-Witten invariant. As shown by Beasley and Witten [22], under a range of circumstances, the instanton superpotentials generated by all such curves can cancel exactly. However, for a wide set of vacua [23, 24, 25, 26], such as those whose Calabi-Yau threefold has a finite isotropy group [27], this cancellation does not occur. Such theories then have an additional nonperturbative superpotential which is the sum of the instanton potentials over all isolated curves. We denote this superpotential, which depends on T , as W_T . Using this, and the associated Kähler potential K_T , in Section 4 we extend our calculation of the potential energy V_F further by using $W =$ $W_{flux} + W_G + W_T$.

Having computed the F -term potential, we recognize that the inhomogeneous transformation of the S and T moduli "axions" under the anomalous $U(1)$ structure group [28, 29] leads to a second contribution to the potential energy - a D-term potential V_D – that must be added to V_F to obtain the total potential energy V . The exact form of V_D was derived in previous work [28, 30]. In Section 5 we present this potential and show that it is minimized, and $N = 1$ supersymmetry is left unbroken, in a specific direction in field space with $s = \text{const.} \times t$ – where $s = \text{Re } S, t = \text{Re } T$ and the constant is a fixed function of various parameters of the chosen vacuum. Along this direction, described as setting the Fayet-Iliopoulos term $FI = 0$, the potential V_D is minimized and exactly vanishes. For any given pair of $\langle s \rangle$ and $\langle t \rangle$ satisfying $FI = 0$, we can expand both S and T around the associated local minimum. As discussed in detail in [28, 29], the complex fluctuations around this minimum can be unitarily rotated to two new complex fields that have canonical kinetic energy and are mass eigenstates. We show that one of these new complex fields has a mass m_{anom} , that is, the mass of the anomalous $U(1)$ gauge field [5]. For sufficiently small values of $\langle t \rangle$, this mass satisfies $m_{\text{anom}} \gtrsim M_U$, where M_U is the compactification scale. Hence, both the gauge field and this diagonalized complex scalar can be integrated out of the low energy theory. However, the second diagonalized complex scalar, with real and imaginary components η and ϕ respectively, has canonical kinetic energy and, with respect to V_D only, has vanishing mass. When V_F is included, each of η and ϕ get a non-vanishing mass substantially smaller than M_U . Hence, they remain in the low energy effective theory. In Section 5 we present the expressions for these lower mass scalars.

In Section 6, we now combine the results from both V_F and V_D and search for minima. We first impose the relation $s = const. \times t$ along which V_D is minimized and equal to zero; then we search for local extrema of V_F along that specific direction. For sufficiently small values of t, where $m_{\text{anom}} \gtrsim M_U$, we insert the results for the single light complex scalar field derived in the previous Section. Even with this reduced number of scalar fields, V_F is a complicated function which we plot numerically using Mathematica. We find that the V_F potential can have a unique minimum along $s = const. \times t$ for a wide range of input parameters. Specifically, we find that the potential energy can have a global minimum at $\langle t \rangle$ and, hence, $\langle s \rangle$ with a vanishing or negative cosmological constant or a metastable minimum with positive vacuum density – depending on the explicit values of the input parameters. In this paper, for specificity, we choose an explicit set of physically motivated parameters and present the potential energy function for various choices of the instanton Pfaffian parameter p . Within this context, we present a plot of V_F for five different choices of p which exhibit these characteristics. In addition to stabilizing the real components s, t respectively, we show that the imaginary, axionic component of the light scalar is simultaneously stabilized.

Finally, in Section 7, using the complete expression for V_F valid at large values of t where $m_{\text{anom}} \ll M_U$, we show that V_F easily satisfies the conjectured Swampland lower bound on the potential gradient [31, 32, 33, 34, 35, 36. This is the case for a wide range of values of Pfaffian parameter p , including cases where V_F has a metastable minimum with positive vacuum energy.

In conclusion, we have shown that in heterotic M-theory vacua compactified on Calabi-Yau threefolds with $h^{1,1} = h^{2,1} = 1$ and, more generically, for the universal Kähler and complex structure moduli of compactifications on any Calabi-Yau threefold with $h^{1,1} = h^{2,1}$, the dilaton and geometric moduli can be completely stabilized. These vacua can have positive, zero or negative vacuum density depending on the precise parameters chosen for the theory. It is clear that these hidden sector vacua, in combination with the vacuum of the observable sector, will lead to a range of values for the cosmological constant. For hidden sector vacua that have positive, but small, vacuum density, one can expect a cosmological constant that can be adjusted

to give the experimentally observed value. This will be discussed in detail in [37]. When redefined so as to have canonical kinetic energy, we find that the fluctuations around these vacua have positive definite masses with values sufficiently below the unification scale so as to allow these moduli fields to be in the low energy theory. Finally, we show that at large values of t , the Swampland "large moduli field" conjecture is satisfied – even when the V has a metastable minimum at small t with positive vacuum density.

2 Complex Structure Moduli

We begin our analysis by considering the Kähler potential, superpotential and potential energy function for the complex structure moduli of a Calabi-Yau threefold X within the context of heterotic M-theory. As discussed in the Introduction, in this paper we will restrict our analysis to Calabi-Yau threefolds with cohomology $h^{2,1} = 1$; that is, to a single complex structure modulus. However, it is instructive to begin our discussion by presenting the Kähler potential and superpotential for an arbitrary number of complex structure moduli, which we then restrict to the $h^{2,1} = 1$ case.

2.1 Kähler Potential

The Kähler potential for an arbitrary number of complex structure moduli z^a , $a = 1, \ldots, h^{2,1} \ge 1$ was presented in [11] and given by

$$
\kappa_4^2 \mathcal{K}(z) = -\ln \left[2i(\mathcal{G} - \bar{\mathcal{G}}) - i(z^a - \bar{z}^a)(\frac{\partial \mathcal{G}}{\partial z^a} + \frac{\partial \bar{\mathcal{G}}}{\partial \bar{z}^a}) \right]
$$
(1)

where

$$
\mathcal{G} = -\frac{1}{6}\tilde{d}_{abc}z^a z^b z^c \,,\tag{2}
$$

 d_{abc} are the intersection numbers for the threefold X ,

$$
\kappa_4^2 = \frac{8\pi}{M_P^2} \tag{3}
$$

and

$$
M_P = 1.22 \times 10^{19} \text{ GeV}
$$
 (4)

is the unreduced Planck mass.

Limiting the Calabi-Yau threefolds to those with $h^{2,1} = 1$, that is, with a single complex modulus z only, one can then write

$$
\tilde{d}_{111} = 6\tilde{d} \tag{5}
$$

where \tilde{d} is a positive integer. The prepotential $\mathcal G$ then becomes

$$
\mathcal{G} = -\tilde{d}z^3 \tag{6}
$$

It follows that the expression for the Kähler potential in (1) simplifies to

$$
\kappa_4^2 \mathcal{K}(z) = -\ln\left[i\tilde{d}(z-\bar{z})^3\right] \tag{7}
$$

2.2 Superpotential

The flux generated superpotential in heterotic M -theory is given, at the classical level, by the Gukov-Vafa-Witten expression

$$
W_{flux} = \frac{\sqrt{2}}{\kappa_4^2} \frac{1}{\pi \rho v^{1/2}} \int_{X \times S^1/\mathbb{Z}_2} \Omega \wedge G , \qquad (8)
$$

where Ω is the holomorphic three-form on the Calabi-Yau threefold X and G is the four-form G-flux. Generically, W_{flux} can depend on both the dilaton S and the complex structure moduli z^a , $a = 1, \ldots, h^{2,1} \geq 1$. However, W_{flux} was explicitly calculated within the context of heterotic M -theory in [12] and found to be independent of S and given by

$$
W_{flux} = \mathcal{C}(\frac{1}{6}\tilde{d}_{abc}z^a z^b z^c n^0 - \frac{1}{2}\tilde{d}_{abc}z^a z^b n^c - z^a n_a - n_0)
$$
 (9)

where n^0, n^a, n_0, n_a are units of flux and arbitrary independent integers in \mathbb{Z} ,

$$
\mathcal{C} = \frac{\sqrt{2}v^{1/6}\epsilon_0}{\kappa_4^2(\pi\rho)^2} \tag{10}
$$

and

$$
\epsilon_0 = (2\sqrt{2}\pi)^{2/3} \frac{(\pi \rho)^{4/3}}{v^{1/3} M_P^{2/3}} \,. \tag{11}
$$

The parameters v and $\pi \rho$ set the scale for the volume of the Calabi-Yau threefold X and the length of the 5-th dimension respectively. We will choose these parameters as follows. First, let

$$
v^{1/6} = \frac{1}{M_U} \,,\tag{12}
$$

where M_U is the unification scale of the effective low energy theory in the observable sector. As discussed in [38, 39], a canonical value for M_U , which we will use in this paper, is given by

$$
M_U = 3.15 \times 10^{16} \text{ GeV} \ . \tag{13}
$$

Second, we set

$$
\pi \rho = 5Fv^{1/6} \tag{14}
$$

where F is, at the moment, an arbitrary real number. To conform to standard values for $\pi \rho$ in the literature, see for example [40, 41], we will generically restrict

$$
0.6 \lesssim F \lesssim 2 \tag{15}
$$

However, as discussed in detail at the beginning of Section 6.4, both larger and smaller values for F are acceptable for the specific vacua analyzed in this paper. Note that the "physical" volume of the Calabi-Yau threefold and the "physical" length of the 5-th dimension are given by

$$
\mathbb{V} = vV \quad , \quad \mathbb{L} = \pi \rho \hat{R} \tag{16}
$$

where V and \hat{R} are the moduli for the Calabi-Yau volume and fifth dimensional length respectively. Using (12) and (14) , as well as the values of M_P and M_U given in (4) and (13), one can calculate the dimensionless coefficient ϵ_0 in (11). We find that

$$
\epsilon_0 = 0.690 \ F^{4/3} \ . \tag{17}
$$

Again using (4) , (12) , (13) , (14) as well as (17) , we find that coefficient C in (10) is given by

$$
\mathcal{C} = \frac{232}{F^{2/3}} M_U^3 \tag{18}
$$

We have written C as proportional to M_U^3 to emphasize that C and, hence, W_{flux} has mass dimension three set by the compactification scale. Finally, let us restrict the above superpotential to Calabi-Yau threefolds with $h^{2,1} = 1$. Expression (9) then reduces to

$$
W_{flux} = \mathcal{C}(\tilde{d}z^3 n^0 - 3\tilde{d}z^2 n^1 - zn_1 - n_0)
$$
 (19)

where all coefficients remain unchanged and n^0, n^1, n_0, n_1 are independent, arbitrary elements of Z.

Before continuing on to the potential energy function for the complex structure modulus, we first compute the covariant derivative of W_{flux} in (19) with respect to z defined by

$$
D_z W_{flux} = \frac{\partial W_{flux}}{\partial z} + \kappa_4^2 \frac{\partial \mathcal{K}}{\partial z} W_{flux}.
$$
 (20)

Using (7) and (19), we find that

$$
D_z W_{flux} = \frac{\mathcal{C}}{z - \bar{z}} \left(3\tilde{d}z^2 n^1 + 2zn_1 - 3\tilde{d}z^2 \bar{z}n^0 + 6\tilde{d}z \bar{z}n^1 + n_1 \bar{z} + 3n_0 \right). \tag{21}
$$

In order for a complex structure moduli vacuum to preserve $N = 1$ supersymmetry, it must satisfy

$$
D_z W_{flux} = 0 \tag{22}
$$

and, hence, from (21) that

$$
3\tilde{d}z^2n^1 + 2zn_1 - 3\tilde{d}z^2\bar{z}n^0 + 6\tilde{d}z\bar{z}n^1 + n_1\bar{z} + 3n_0 = 0.
$$
 (23)

Defining

$$
z = r + ic \t{24}
$$

we find that this will be the case if and only if

$$
r = -\frac{3n_0n^0 + n_1n^1}{2(3\tilde{d}(n^1)^2 + n^0n_1)}
$$
\n(25)

and

$$
c = \pm \left(-r^2 + \frac{2n^1r}{n^0} + \frac{n_1}{3\tilde{d}n^0} \right)^{1/2} . \tag{26}
$$

There are, of course, an infinite number of solution to r and c of this form, indexed by the choices of integers n^0, n^1, n_0, n_1 . As a simple example, let us consider solutions where

$$
r = 0 \Rightarrow 3n_0 n^0 = -n_1 n^1
$$
 and $c = \pm \left(\frac{n_1}{3\tilde{d}n^0}\right)^{1/2}$. (27)

Furthermore, by taking

$$
n_1 = n^0 = -n^1 \Rightarrow c = \pm \frac{1}{\sqrt{3}\tilde{d}}
$$
 and $3n_0 = n_1 = n^0 = -n^1$. (28)

Therefore, any choice of this one parameter set of integers gives a unique complex structure modulus, with $r = 0$ and a simple value of c, that satisfies $D_z W_{flux} = 0.$

As we proceed, it will become essential to know the value of W_{flux} at a point where $D_zW_{flux} = 0$. It follows from $(19),(20),(22)$ and (7) that at a supersymmetric point

$$
W_{flux} = \mathcal{C}\tilde{d}c\big(-4(n^0r - n^1)c + i[2n^0(r^2 - c^2) - 4n^1r - \frac{2n_1}{3\tilde{d}}]\big). \tag{29}
$$

Using (26), this simplifies to

$$
W_{flux} = -4\mathcal{C}\tilde{d}c^2\big((n^0r - n^1) + in^0c\big) \ . \tag{30}
$$

We find it convenient to re-express this as

$$
W_{flux} = -4\mathcal{C}\tilde{d}(A+iB) , \qquad (31)
$$

where

$$
A = c2(n0r - n1), B = c3n0, C = \frac{232}{F2/3} M_U3
$$
 (32)

and r and c are given in (25) and (26) respectively. As a concrete example, consider the choice of integer parameters given in (28). Then

$$
r = 0
$$
, $c = \pm \frac{1}{\sqrt{3}\tilde{d}}$ \Rightarrow $A = \frac{n^0}{3\tilde{d}}$, $B = \pm \frac{n^0}{3\tilde{d}\sqrt{3}\tilde{d}}$ for $n^0 \in \mathbb{Z}$. (33)

2.3 Complex Structure F-Term Potential Energy

In this subsection, we want to discuss the part of the F-term potential energy function arising from the flux superpotential W_{flux} only. To do this, however, it is essential to include contributions that arise from terms associated with the Kähler potentials of the dilaton S and the $h^{1,1} = 1$ Kähler modulus T, as well as the complex structure z. That is, the relevant terms in the potential energy function are

$$
V_{flux} = e^{\kappa_4^2 K} \left[K^{z\bar{z}} D_z W_{flux} D_{\bar{z}} \bar{W}_{flux} + K^{S\bar{S}} D_S W_{flux} D_{\bar{S}} \bar{W}_{flux} \right. \left. + K^{T\bar{T}} D_T W_{flux} D_{\bar{T}} \bar{W}_{flux} - 3\kappa_4^2 |W_{flux}|^2 \right] \,, \tag{34}
$$

where

$$
\kappa_4^2 K = \kappa_4^2 (K_S + K_T + \mathcal{K})
$$

$$
\kappa_4^2 K_S = -\ln(S + \bar{S}) , \ \kappa_4^2 K_T = -3\ln(T + \bar{T}) , \ \kappa_4^2 \mathcal{K} = -\ln(i\tilde{d}(z - \bar{z})^3)
$$
 (35)

and

$$
D_i W_{flux} = \partial_i W_{flux} + \kappa_4^2 (\partial_i K) W_{flux} \text{ for } i = S, T, z. \qquad (36)
$$

We emphasize that here, and *throughout this paper*, we always use the *tree* level expressions–given in (35) –for the Kähler potentials. This is motivated by the fact that, in the absence of five-branes–which we are assuming–the leading order string perturbative corrections vanish. See, for example, [29]. Evaluating $K^{S\bar{S}}$ and $K^{T\bar{T}}$ and using the fact that W_{flux} is independent of both S and T , expression (34) becomes

$$
V_{flux} = e^{\kappa_4^2 K} \left[K^{z\bar{z}} D_z W_{flux} D_{\bar{z}} \bar{W}_{flux} + \kappa_4^2 |W_{flux}|^2 + 3\kappa_4^2 |W_{flux}|^2 - 3\kappa_4^2 |W_{flux}|^2 \right],
$$
\n(37)

which, noting the cancellation of the last two $|W_{flux}|^2$ terms, simplifies to

$$
V_{flux} = e^{\kappa_4^2 K} \left[K^{z\bar{z}} D_z W_{flux} D_{\bar{z}} \bar{W}_{flux} + \kappa_4^2 |W_{flux}|^2 \right] \,. \tag{38}
$$

Finally, using

$$
e^{\kappa_4^2 K} = \frac{-i}{(S+\bar{S})(T+\bar{T})^3 \tilde{d}(z-\bar{z})^3}, \quad K^{z\bar{z}} = -\frac{\kappa_4^2 (z-\bar{z})^2}{3}, \quad (39)
$$

it follows that

$$
V_{flux} = \frac{1}{(S+\bar{S})(T+\bar{T})^3} \left[\frac{-i}{\tilde{d}(z-\bar{z})^3} \left(-\frac{\kappa_4^2 (z-\bar{z})^2}{3} D_z W_{flux} D_{\bar{z}} \bar{W}_{flux} + \kappa_4^2 |W_{flux}|^2 \right) \right]
$$
(40)

Note that this is the potential energy function for arbitrary values of complex structure modulus z. Hence, $D_{\bar{z}} \overline{W}_{flux}$ does not generically vanish.

Although complex structure moduli satisfying constraints (25) and (26) have vanishing covariant derivative D_zW_{flux} , it is essential to determine whether or not they are local minima of V_{flux} . We begin by considering the first derivative of the potential function (40). We find that

$$
\frac{\partial V_{flux}}{\partial z} = \frac{-i\kappa_4^2}{(S+\bar{S})(T+\bar{T})^3\tilde{d}(z-\bar{z})^3} \left[\frac{\partial W_{flux}}{\partial z} - \frac{3}{(z-\bar{z})}W_{flux}\right]\bar{W}_{flux} \quad (41)
$$

Using (7) and (20) , it follows that

$$
\frac{\partial W_{flux}}{\partial z} - \frac{3}{(z - \bar{z})} W_{flux} = D_z W_{flux} \tag{42}
$$

and, hence,

$$
\frac{\partial V_{flux}}{\partial z} = \frac{-i\kappa_4^2}{(S+\bar{S})(T+\bar{T})^3\tilde{d}(z-\bar{z})^3} \left[D_z W_{flux}\right]\bar{W}_{flux} \ . \tag{43}
$$

Therefore, all complex structure moduli with vanishing covariant derivative $D_zW_{flux} = 0$ satisfy

$$
\frac{\partial V_{flux}}{\partial z} = 0 \tag{44}
$$

Since V_{flux} is real, this implies that

$$
\frac{\partial V_{flux}}{\partial r} = \frac{\partial V_{flux}}{\partial c} = 0.
$$
 (45)

Are these extrema with $D_zW_{flux} = 0$ also local minima of V_{flux} ? The answer is yes, *provided* $c > 0$. This can be proven by a tedious but straightforward evaluation of the second derivatives of V_{flux} at the extrema which shows that

$$
\frac{\partial^2 V_{flux}}{\partial^2 r} > 0, \frac{\partial^2 V_{flux}}{\partial^2 c} > 0, \text{ and } \frac{\partial^2 V_{flux}}{\partial r \partial c} = 0 \text{ for } \langle c \rangle > 0. \tag{46}
$$

This suffices to prove that the determinant of the Hessian and the second derivative V_{flux} with respect to r are both positive at the extrema and, therefore, the extrema must be minima.

Henceforth, we will chose the vacuum expectation value $\langle z \rangle$ of the complex structure modulus to be at one of the local minima of V_{flux} . It follows from $(31),(32)$ and (40) that, at any such minimum,

$$
V_{flux} = \frac{M_U^4}{st^3 \langle c \rangle^3} \left(\frac{1.14}{F^{4/3}}\right) \tilde{d}(A^2 + B^2) \tag{47}
$$

2.4 Non-Renormalization of W_{flux}

The Gukov-Vafa-Witten formula in (8) gives the tree level expression for W_{flux} . Can W_{flux} receive string and α' perturbative corrections? Within the context of heterotic M -theory the answer is no. As shown in [43], low energy Peccei-Quinn symmetry forbids W_{flux} from containing the dilaton S both at the tree level–thus explaining why S does not appear in (9) –and at any perturbative level–thus ensuring that W_{flux} receives no corrections in string perturbation theory. Similarly, axion-like symmetries of the Kähler moduli

preclude W_{flux} from acquiring a T-dependence. Hence, W_{flux} can receive no α' corrections. That is, the expression for W_{flux} in (9) can receive no string and α' perturbative corrections.

Be that as it may, these arguments no longer apply to non-perturbative corrections to W_{flux} , to which we now turn.

3 Gaugino Condensation and $N = 1$ Supersymmetry Breaking

To spontaneously break $N = 1$ supersymmetry, following, for example, the approach in the heterotic M-theory $B-L$ MSSM vacuum [5, 42], we consider theories containing a line bundle with an anomalous $U(1)$ structure group in the hidden sector. Furthermore, we assume that the commutant subgroup of $U(1)$ contains a non-Abelian group which becomes strongly coupled at a mass scale Λ near M_U .

3.1 Gaugino Condensate Superpotential

As is well-known [13, 14, 15, 16], the condensation of the associated gauginos at scale Λ produces a non-perturbative superpotential which, to lowest order, is given by

$$
W_G = M_U^3 e^{-bS} \t\t(48)
$$

where

$$
b = \frac{6\pi}{b_L \hat{\alpha}_{GUT}} \quad , \quad \hat{\alpha}_{GUT} = \frac{\left(8\pi (\pi \rho) \kappa_4^2\right)^{2/3}}{v^{1/3}} \tag{49}
$$

 $\frac{1}{2}$

and b_L is the renomalization group coefficient associated with the specific non-Abelian commutant. Using $(3),(12),(13)$ and (14) , we find that

$$
\hat{\alpha}_{GUT} = 0.0762 \ F^{2/3} \tag{50}
$$

where F can run over the range given in (15). The value of b_L depends on the specific commutant non-Abelian group in the hidden sector. For example, in [5, 39], the non-Abelian group is E_7 with the renormalization group coefficient

$$
b_L = 6 \tag{51}
$$

The non-perturbative superpotential W_G leads to spontaneous breaking of $N = 1$ supersymmetry in the S and T moduli, which is then gravitationally

mediated to the observable matter sector. The scale of SUSY breaking in the low-energy observable sector is of order

$$
m_{\text{susy}} \sim \kappa_4^2 \Lambda^3 = 8\pi \frac{\Lambda^3}{M_P^2} \ . \tag{52}
$$

Let us now extend the superpotential to include both the complex structure and gaugino condensate superpotentials. That is, take

$$
W = W_{flux} + W_G . \t\t(53)
$$

3.2 F-Term Potential Including Gaugino Condensation

Generically, the F-term potential energy function is given by

$$
V_F = e^{\kappa_4^2 K} \left[K^{i\bar{j}} D_i W D_{\bar{j}} \bar{W} - 3\kappa_4^2 |W|^2 \right] \tag{54}
$$

where

$$
D_i W = \partial_i W + \kappa_4^2 (\partial_i K) W , \qquad (55)
$$

the Kähler potential is given in (35) and the indices i, j run over S, T and z. In this Section, we choose the superpotential to be that given in (53).

In our evaluation of V_F we will, henceforth, assume that the complex structure modulus z is always fixed to be a local minimum $\langle z \rangle$ of V_{flux} satisfying $D_zW_{flux} = 0$. For example, the evaluation of the z covariant derivative in (54) is given by

$$
D_z W = \partial_z (W_{flux} + W_G) + \kappa_4^2 \partial_z K (W_{flux} + W_G)
$$

=
$$
D_z W_{flux} + \left(\frac{-3}{(z - \bar{z})}\right) W_G
$$

=
$$
\frac{-3}{\langle z - \bar{z} \rangle} W_G
$$
 (56)

where we have used the fact that the gaugino condensate superpotential (48) is independent of complex structure. Then, using W_{flux} given in (31), (32), the gaugino condensate superpotential W_G in (48), the Kähler potentials in (35) and defining

$$
S = s + i\sigma \quad , \quad T = t + i\chi \tag{57}
$$

we find, after a moderate calculation, that

$$
V_F = \frac{M_U^4}{st^3 \langle c \rangle^3} \left[\left(\frac{1.1376}{F^{4/3}} \right) \tilde{d} (A^2 + B^2) + 1.32 \times 10^{-6} \tilde{d}^{-1} \left((1 + 2bs)^2 + 3 \right) e^{-2bs} - \left(\frac{2.43 \times 10^{-3}}{F^{2/3}} \right) \left(1 + 2bs \right) e^{-bs} \left(A \cos(b\sigma) - B \sin(b\sigma) \right) \right].
$$
\n(58)

Once again, note that we are using the tree level expressions for the Kähler potentials given in (35).

We conclude that adding the non-perturbative superpotential W_G substantially alters the F-term effective potential V_F presented in (47) based on W_{flux} alone and evaluated at $D_zW_{flux} = 0$. It is important, therefore, to search for and include any other non-perturbative superpotentials in the calculation of the F-term potential energy.

4 Worldsheet Instanton Superpotential

It is well-known [17, 18, 19] that a non-perturbative contribution to the superpotential can also be generated by the superstring wrapping around isolated, genus zero, holomorphic curves in the Calabi-Yau compactification threefold. The leading order contribution to this superpotential arises from the superstring wrapping once around each such curve. In this Section, we introduce this leading order, non-perturbative "worldsheet instanton" contribution.

4.1 Single Isolated, Genus-Zero Curve

Let C be an holomorphic, isolated, genus zero curve in the Calabi-Yau threefold X with $h^{1,1} = 1$. Then, as discussed in [44], the general form of the instanton superpotential induced by a string wrapping C is given by

$$
W_I = \mathcal{P}e^{-\tau T} \tag{59}
$$

where

$$
\tau = \frac{1}{2} T_M(\pi \rho) v_C , \qquad (60)
$$

 v_C is the volume of the holomorphic curve C and T_M is the string membrane tension

$$
T_M = \frac{1}{(v\rho \kappa_4^2)^{1/3}} \ . \tag{61}
$$

Using $(3),(12),(13),(14)$ one finds

$$
T_M = \frac{15.5}{F^{1/3}} M_U^3 \tag{62}
$$

and, hence, that

$$
\tau = 38.8 \ F^{2/3} \frac{v_C}{v^{1/3}}
$$
 (63)

The factor P in (59) is given by

$$
\mathcal{P} \propto \text{Pfaff}(\mathcal{D}_-),\tag{64}
$$

where Pfaff(\mathcal{D}_-) is the Pfaffian of the chiral Dirac operator constructed using the hermitian Yang-Mills connection associated with the holomorphic vector bundle on both the observable and hidden sectors evaluated at the curve C. For a generic bundle, one expects the Pfaffian to be proportional to a holomorphic, polynomial function of a subset of the vector bundle moduli, as shown in several contexts in [20, 21, 27]. In the previous Section, we specified that the vector bundle in the hidden sector must contain a line bundle with an anomalous $U(1)$ structure group. If the hidden sector bundle is strictly a single line bundle L (more exactly, its extension to $L \oplus L^{-1}$ whose structure group is embedded in E_8), we note that it will have no vector bundle moduli and, hence, vanishing Pfaffian. However, it is important to note that the "string instanton" arises from all bundle gauge connections in both the hidden and the observable sectors. That is, P is the Pfaffian associated with the gauge connections of the vector bundles of both the observable and hidden sectors, which, as a rule, will have multiple vector bundle moduli and, potentially, a non-vanishing Pfaffian. Finally, we note that the proportionality factor in (64) was defined in detail, for example in [27, 45], and shown to be an explicit function of the complex structure moduli. In this paper, since the value of the complex structure modulus has been fixed to be at a supersymmetry preserving minimum of V_{flux} , this function becomes a constant that could, in principle, be explicitly computed. However, for simplicity, we will simply absorb it into the expression for the Pfaffian and write

$$
\mathcal{P} = \text{Pfaff}(\mathcal{D}_-) \tag{65}
$$

4.2 Multiple Isolated, Genus-Zero Curves

When the Calabi-Yau threefold has $h^{1,1} = 1$, there is a single homology class in H_2 which, given the above discussion, we denote by $[C]$. It is well known that, in general, this class can contain a finite number of holomorphic, isolated, genus zero curves in addition to C. The total number of such curves is specified by the Gromov-Witten invariant, which can be computed given a specific Calabi-Yau threefold – see, for example $[27, 46, 47]$. Denote the number of such curves by $n_{[C]}$ and write them as C_i , $i = 1, \ldots n_{[C]}$. All such curves in the same homology class have the same area, the same classical action and the same exponential factor as in (59). However, their Pfaffians are, in general, different. Hence, the contribution to the superpotential from all curves C_i in the homology class $[C]$ is given by

$$
W_I = \left(\sum_{i=1}^{n_{[C]}} \mathcal{P}_i\right) e^{-\tau T} \tag{66}
$$

An important theorem of Beasley and Witten [22] proved that, under a range of circumstances, the sum over the Pfaffians in any given homology class must vanish. It then follows that the associated instanton superpotential in (66) would be zero. However, as shown in a number of papers – see [23, 24, 25, 26] for example – there are a substantial number of phenomenologically acceptable vacua that violate one or more of the assumptions in the Beasley-Witten theorem. For example, in the heterotic M-theory $B - L$ MSSM vacuum in [27], the existence of $\mathbb{Z}_3 \times \mathbb{Z}_3$ discrete torsion violates the Beasley-Witten theorem and has non-vanishing instanton superpotential W_I . Henceforth, we will assume that the vacua we are considering violate the Beasley-Witten theorem; that is, that $\sum_{i=1}^{n_{[C]}} \mathcal{P}_i \neq 0$.

The exact functional form of $\sum_{i=1}^{n_{[C]}} \mathcal{P}_i$ has only been calculated in a few specific examples, see [20, 21, 27], and is unknown for most physically acceptable heterotic vacua. This is also the case for the vector bundle moduli Kähler potential. Hence, the precise formalism for computing the supersymmetry preserving vector bundle moduli vacua is unknown. Be that as it may, we will now assert–similarly to the complex structure modulus–that the values of the vector bundle moduli are independently fixed to be constants in a *vacuum state that does not break* $N = 1$ *supersymmetry.* It follows that the Pfaffian factor is simply a complex number, which we will express as

$$
\sum_{i=1}^{n_{[C]}} \mathcal{P}_i = p e^{i\theta_p} \tag{67}
$$

where, since it is unknown how to compute the Pfaffian for the theories discussed in this paper, the values of p and θ_p are, a priori, unrestricted.

That is, in this paper, we simply parameterize the Pfaffian factor in terms of the real coefficients p and θ_p and deduce what their values must be in order to stabilize the dilaton, real Kähler modulus and the T axion respectively within our context.

In summary, we will henceforth assume that $W_I \neq 0$, and simply denote (66) as

$$
W_I = M_U^3 p e^{i\theta_p} e^{-\tau T} \tag{68}
$$

where, using the fact that the Calabi-Yau threefold in this paper has a mass scale of order M_U , we have restored natural units.

4.3 F-Term Potential Function With String Instantons

In this subsection, we extend the F-term potential energy function given in (58) to include contributions from the instanton superpotential. That is, we evaluate the potential energy function in $(54),(55)$ but now taking the superpotential to be

$$
W = W_{flux} + W_G + W_I , \qquad (69)
$$

where W_{flux} , W_C and W_I are given in (31), (48) and (68) respectively. As above, the Kähler potential is presented in (35) and the indices i, j run over $S, T \text{ and } z.$ We emphasize once again that throughout this paper we always use the tree level expressions for the Kähler potentials. Importantly, as we did in the previous Section, we will assume in our evaluation of V_F that the complex structure z is always fixed to be a local minimum $\langle z \rangle$ of V_{flux} satisfying $D_zW_{flux} = 0$. Again defining

$$
S = s + i\sigma \quad , \quad T = t + i\chi \tag{70}
$$

we find, after a detailed calculation, that

$$
V_F = \frac{M_U^4}{st^3 \langle c \rangle^3} \left[\left(\frac{1.14}{F^{4/3}} \right) \tilde{d}(A^2 + B^2) + 1.32 \times 10^{-6} \tilde{d}^{-1} \left((1 + 2bs)^2 + 3 \right) e^{-2bs} - \left(\frac{2.43 \times 10^{-3}}{F^{2/3}} \right) \left(1 + 2bs \right) e^{-bs} sgn(A) \sqrt{A^2 + B^2} \cos(b\sigma + \arctan(\frac{B}{A})) + 2.62 \times 10^{-6} p \tilde{d}^{-1} \left(1 + 2bs + 3(\tau t + \frac{3}{2}) \right) e^{-bs - \tau t} \cos(b\sigma - \tau \chi + \theta_p) + 4.36 \times 10^{-7} p^2 \tilde{d}^{-1} \left(3 + (2\tau t + 3)^2 \right) e^{-2\tau t} - \left(\frac{2.43 \times 10^{-3}}{F^{2/3}} \right) p (1 + 2\tau t) e^{-\tau t} sgn(A) \sqrt{A^2 + B^2} \cos(\tau \chi - \theta_p + \arctan(\frac{B}{A})) \right],
$$
\n(71)

where A, B, $\langle c \rangle$ are defined in (32), F is defined in (14),(15), b and τ are given in (49) and (63), respectively, and p and θ_p arise in (68). In addition, we have used the trigonometric relation

$$
A\cos(x) - B\sin(x) = sgn(A)\sqrt{A^2 + B^2}\cos(x + \arctan(\frac{B}{A}))
$$
 (72)

We conclude that adding the "string instanton" superpotential W_I greatly alters the F-term potential energy function – which, for $W_{flux} + W_G$ was presented in (58). The potential V_F in (71) is – to lowest order – the most general expression for the F-term potential energy for the dilaton and Kähler modulus when the complex structure z is evaluated at a local minimum of V_{flux} for which $D_zW_{flux} = 0$.

5 Anomalous $U(1)$ and σ, χ Axions

In the previous Sections, we have considered only the superpotentials and the F-term potential energy associated with the dilaton, complex structure and Kähler moduli. This was motivated by the fact that none of these moduli fields transform homogeneously under any low-energy gauge group. However, as we discuss in this Section, the anomalous $U(1)$ gauge group does produce an inhomogeneous transformation of the imaginary component of both the dilation and the Kähler modulus – thus inducing a D-term potential energy V_D . In this Section, we define and discuss this D-term potential only. In Section 6, we will reintroduce V_F in combination with V_D and discuss the associated moduli vacua.

5.1 Inhomogeneous $U(1)$ Transformations

As discussed in Section 2, in this paper we will assume that the hidden sector vector bundle contains a line bundle L with an anomalous $U(1)$ structure group. Within the context of compactification on a Calabi-Yau threefold with $h^{1,1} = 1$, it was shown in [29] that the Green-Schwarz mechanism [6] cancels this anomaly by producing an inhomogeneous transformation of the dilaton S and the Kähler modulus T. It was shown in [28, 29] that for a $U(1)$ parameter θ , these transformations are given by

$$
\delta_{\theta} S = 2i\pi a \epsilon_S^2 \epsilon_R^2 \beta l \theta \equiv k_S \theta \quad , \quad \delta_{\theta} T = -2i a \epsilon_S \epsilon_R^2 l \theta \equiv k_T \theta \tag{73}
$$

where

$$
\epsilon_S = \frac{2\pi \rho^{4/3}}{v^{1/3} M_P^{2/3}} \quad , \quad \epsilon_R = \frac{v^{1/6}}{\pi \rho} \tag{74}
$$

are strong coupling expansion parameters. The integer l defines the line bundle as $\mathcal{O}_X(l) = L$, β is the gauge charge on the hidden sector and

$$
a = \frac{1}{4}tr_{E_8}Q^2\tag{75}
$$

is determined given the embedding matrix Q of $U(1)$ into E_8 in the hidden sector.

Using (3) , (12) , (13) and (14) one can determine the expansion parameters in (74). We find that

$$
\epsilon_S = 0.220 \ F^{4/3} \ , \ \epsilon_R = \frac{0.2}{F} \ .
$$
\n(76)

Note that the ϵ_S often occurs multiplied by π , so we introduce

$$
\epsilon_S' = \epsilon_S \pi = 0.690 \ F^{4/3} \ . \tag{77}
$$

For simplicity, we will assume henceforth that the line bundle structure group embeds into the $SU(2)$ subgroup of E_8 in the hidden sector. It follows that $a = 1$. Finally, we will defer a discussion of the values of l and β until later in the paper.

We see from (73) that these $U(1)$ transformations are purely imaginary and, hence, represent inhomogeneous transformations of the σ and χ imaginary components of S and T respectively. That is, the σ and χ fields behave as "axions" under the anomalous $U(1)$ transformation.

5.2 Anomalous $U(1)$ Induced Potential Energy V_D

In addition to the canonical D-term potential energy involving all hidden sector matter fields carrying non-vanishing $U(1)$ charge, the inhomogeneous $U(1)$ transformations of S and T presented in (73) induce a potential energy for the dilaton and modulus fields even though they are neutral under linear $U(1)$ transformations. As discussed in detail in [29], assuming the vacuum expectation values for all hidden matter scalars are zero, then these matter scalars "decouple" from the S and T moduli D-term potential energy. We will assume that this is the case in this paper. Then, as shown in [29, 30], ignoring the matter scalars, the D-term potential energy is given by

$$
V_D = \frac{1}{2s} \mathbb{P}^2 \tag{78}
$$

where

$$
\mathbb{P} = ik_S \partial_S K + ik_T \partial_T K . \tag{79}
$$

It follows from (35) that

$$
\partial_S K = -\kappa_4^{-2} \frac{1}{S+\bar{S}} = -\kappa_4^{-2} \frac{1}{2s} \ , \ \ \partial_T K = -3\kappa_4^{-2} \frac{1}{T+\bar{T}} = -3\kappa_4^{-2} \frac{1}{2t} \quad (80)
$$

and from (73) that

$$
k_S = i2\pi \epsilon_S^2 \epsilon_R^2 \beta l \quad , \quad k_T = -i2\epsilon_S \epsilon_R^2 l \; . \tag{81}
$$

Inserting these results into (79), we find that

$$
\mathbb{P} = -\frac{\epsilon_S \epsilon_R^2}{\kappa_4^2} \left(-\frac{1}{s} \epsilon_S' \beta l + \frac{3l}{t} \right) . \tag{82}
$$

Using (76), (77) as well as (3) and (13), we can re-express $\mathbb P$ as

$$
\mathbb{P} = -\left(\frac{53.4}{F^{2/3}}\right) M_U^2 \left(-\frac{1}{s}(.690F^{4/3})\beta l + \frac{3l}{t}\right) \,. \tag{83}
$$

Hence, it follows from (78) that

$$
V_D = \left(\frac{1.42 \times 10^3}{F^{4/3}}\right) \frac{M_U^4}{s} \left(-\frac{1}{s}(.690F^{4/3})\beta l + \frac{3l}{t}\right)^2. \tag{84}
$$

Note that V_D is a function of s and t (the real parts of S and T respectively) and is independent of the "axions" σ and χ .

Finally, we note that the part of $\mathbb P$ which is independent of the charged matter scalars, that is, $\mathbb P$ given in (82), when evaluated for some fixed values of $\langle s \rangle$ and $\langle t \rangle$ is customarily referred to as the Fayet-Iliopoulos (FI) term. We will do so, henceforth, in this paper.

5.3 Supersymmetry Preserving Vacua of V_D

Minimizing the D-term potential (84) defines D-flat, $N = 1$ supersymmetry preserving vacuum states $\langle s \rangle, \langle t \rangle$ for which

$$
\langle \mathbb{P} \rangle = FI = 0. \tag{85}
$$

It follows from (82) that this will be the case for

$$
\langle s \rangle = \frac{\epsilon_S' \beta}{3} \langle t \rangle \tag{86}
$$

or, using (77), for

$$
\langle s \rangle = .230F^{4/3}\beta \langle t \rangle . \tag{87}
$$

Integer l which defines the line bundle L has canceled out of this expression. Importantly, note that the values of $\langle s \rangle \langle t \rangle$ are not completely determined by requiring that $FI = 0$. Rather, they form a straight one-dimensional line in s and t space where $V_D = 0$. That is, a priori, $\langle t \rangle$ can take any value whereas $\langle s \rangle$ is constrained to satisfy (87). The value for $\langle t \rangle$ will be fixed to be a minimum of the F-term potential V_F in Section 6. Finally, note that since V_D is independent of the axions σ and χ , there is, as yet, no constraints on their vacuum expectation values.

Let us choose *any* point along the D -flat direction and expand the complex fields S and T around the associated vacuum expectation values. Expanding

$$
S = \langle s \rangle + \delta S \quad , \quad T = \langle t \rangle + \delta T \tag{88}
$$

one finds that the Lagrangian for δS and δT has off-diagonal kinetic energy and mass terms. However, as shown in [29], one can define two new complex fields ξ^1 and ξ^2 which have canonically normalized kinetic energy and are mass eigenstates.

Following [29], we define

$$
\begin{pmatrix} \xi^1 \\ \xi^2 \end{pmatrix} = U \begin{pmatrix} \delta S \\ \delta T \end{pmatrix}
$$
 (89)

where

$$
U = \frac{1}{\langle \Sigma \rangle} \begin{pmatrix} \langle g_{S\bar{S}} \bar{k}_{S} \rangle & \langle g_{T\bar{T}} \bar{k}_{T} \rangle \\ \sqrt{\langle g_{S\bar{S}} g_{T\bar{T}} \rangle} \langle \bar{k}_{T} \rangle & -\sqrt{\langle g_{S\bar{S}} g_{T\bar{T}} \rangle} \langle \bar{k}_{S} \rangle \end{pmatrix}
$$
(90)

and

$$
\Sigma^2 = g_{S\bar{S}}k_S\bar{k}_S + g_{T\bar{T}}k_T\bar{k}_T.
$$
\n(91)

It then follows that the Lagrangian for ξ^1 and ξ^2 is given by

$$
\mathcal{L} = -\partial^{\mu}\bar{\xi}^{1}\partial_{\mu}\xi^{1} - \partial^{\mu}\bar{\xi}^{2}\partial_{\mu}\xi^{2} - m_{\text{anom}}^{2}\bar{\xi}^{1}\xi^{1} , \qquad (92)
$$

where

$$
m_{\text{anom}} = \sqrt{2 \langle g_2^2 \Sigma^2 \rangle} \tag{93}
$$

and ξ^2 is massless. As discussed below, m_{anom} is the mass of the gauge connection of the anomalous $U(1)$ structure group.

Let us evaluate m_{anom} . The gauge coupling is defined by

$$
g_2^2 = \frac{\pi \hat{\alpha}_{GUT}}{s} \tag{94}
$$

Using(50) and (87), we find that in the chosen $FI = 0$ vacuum

$$
\langle g_2 \rangle = \frac{.489 F^{1/3}}{\langle s \rangle^{1/2}} = \left(\frac{1.02}{F^{1/3} \beta^{1/2}} \right) \frac{1}{\langle t \rangle^{1/2}} \,. \tag{95}
$$

Computing Σ^2 in (91) using (35),(76) and (81) we find that

$$
\Sigma^2 = \frac{M_P^2}{8\pi} l^2 \left(\frac{3.68 \times 10^{-5} F^{4/3} \beta^2}{s^2} + \frac{2.32 \times 10^{-4}}{t^2 F^{4/3}} \right) \,. \tag{96}
$$

Now evaluate this for the chosen $FI = 0$ vacua. Using (87), we find that the coefficient β exactly cancels out of the expression. Finally, using (3) and (13) , we find

$$
\langle \Sigma \rangle = \left(\frac{2.35 \, l}{F^{2/3}} \right) \frac{M_U}{\langle t \rangle} \,. \tag{97}
$$

It then follows from (93), (95) and (97) that

$$
m_{\text{anom}} = \left(\frac{3.39 \, l}{F \beta^{1/2}}\right) \frac{M_U}{\langle t \rangle^{3/2}} \,. \tag{98}
$$

Note that m_{anom} is a function of the expectation value $\langle t \rangle$.

Since complex fields ξ^1 and ξ^2 are mass eigenstates with canonical kinetic energy, it is useful to express both S and T in terms of ξ^1 and ξ^2 . This can be done by inverting expression (89). Then

$$
\begin{pmatrix} \delta S \\ \delta T \end{pmatrix} = U^{-1} \begin{pmatrix} \xi^1 \\ \xi^2 \end{pmatrix} \tag{99}
$$

where

$$
U^{-1} = \frac{1}{\langle \Sigma \rangle} \begin{pmatrix} \langle k_S \rangle & \sqrt{\langle \frac{g_{T\bar{T}}}{g_{S\bar{S}}}} \rangle \langle k_T \rangle \\ \langle k_T \rangle & -\sqrt{\langle \frac{g_{S\bar{S}}}{g_{T\bar{T}}}} \rangle \langle k_S \rangle \end{pmatrix}
$$
(100)

Using $(35),(76),(77),(81),(97)$, as well as (3) and (13) , we find that

$$
U^{-1} = \frac{i\langle t \rangle}{M_P} \begin{pmatrix} 2.00F^{4/3}\beta & -1.15F^{4/3}\beta \\ -2.90 & -5.00 \end{pmatrix}
$$
 (101)

Writing

$$
\delta S = \delta s + i\sigma \quad , \quad \delta T = \delta t + i\chi \quad , \tag{102}
$$

it follows from (99) and (101) that

$$
\delta s + i\sigma = \frac{i\langle t \rangle}{M_P} \left(2.00 F^{4/3} \beta \xi^1 - 1.15 F^{4/3} \beta \xi^2 \right),
$$

$$
\delta t + i\chi = \frac{i\langle t \rangle}{M_P} \left(-2.90 \xi^1 - 5.00 \xi^2 \right).
$$
 (103)

To proceed, we recall that ξ^1 is a massive complex field with mass m_{anom} given in (98), while ξ^2 is massless. Defining

$$
\xi^1 = \eta^1 + i\phi^1 \,, \tag{104}
$$

it was shown in detail in [5, 28, 29] that the anomalous $U(1)$ hidden sector gauge group is spontaneously broken such that: (1) the $U(1)$ gauge field $A_{U(1)}$ attains a mass m_{anom} ; (2) the real scalar ϕ^1 obtains the same mass m_{anom} based on (92); and, (3) real scalar $\eta¹$ acts as the zero mass Goldstone boson associated with this spontaneous breaking. In combination with the fermionic superpartner ψ^1 which also, by supersymmetry, must obtain mass m_{anom} , the combination $(\phi^1, A_{U(1)}, \psi^1)$ forms a vector supermultiplet with mass m_{anom} . As we will demonstrate in a concrete example presented below, physically realistic values of F, β , l and $\langle t \rangle$ are typically such that

$$
m_{\text{anom}} \gtrsim M_U \tag{105}
$$

It follows from (12) that M_U sets the Calabi-Yau threefold mass scale. In this paper, we are interested only in the effective low-energy theory composed of fields with mass substantially less than this scale, Hence, we will integrate out all fields with mass greater than or approximately equal to M_U . Thus, it follows from (105), that we will integrate the anomalous $U(1)$ massive vector superfield, which includes the scalar field component ϕ^1 , out of the theory. Recalling that the Goldstone boson η^1 can be gauged away, it follows that the entire complex scalar field ξ^1 can be integrated out of the low energy theory.

Therefore, in the low energy effective theory, the expressions in (103) simplify to \cdot / \cdot

$$
\delta s + i\sigma = -\frac{i\langle t \rangle}{M_P} \left(1.15 F^{4/3} \beta \xi^2 \right),
$$

$$
\delta t + i\chi = -\frac{i\langle t \rangle}{M_P} \left(5.00 \xi^2 \right).
$$
 (106)

Writing

$$
\xi^2 = \eta + i\phi \tag{107}
$$

it follows that

$$
\delta s = \frac{\langle t \rangle}{M_P} 1.15 F^{4/3} \beta \phi , \quad \sigma = -\frac{\langle t \rangle}{M_P} 1.15 F^{4/3} \beta \eta
$$

$$
\delta t = \frac{\langle t \rangle}{M_P} 5.00 \phi , \quad \chi = -\frac{\langle t \rangle}{M_P} 5.00 \eta .
$$
 (108)

It then follows from (88), (102) and (108) that

$$
s = \langle s \rangle + \frac{\langle t \rangle}{M_P} 1.12 F^{4/3} \beta \phi , \quad t = \langle t \rangle + \frac{\langle t \rangle}{M_P} 5.00 \phi
$$

$$
\sigma = -\frac{\langle t \rangle}{M_P} 1.15 F^{4/3} \beta \eta , \quad \chi = -\frac{\langle t \rangle}{M_P} 5.00 \eta .
$$
 (109)

Now recall from (87) that

$$
\langle s \rangle = .230F^{4/3}\beta \langle t \rangle \tag{110}
$$

Inserting this into (109) we obtain finally that

$$
s = \langle t \rangle F^{4/3} \beta (0.230 + 1.15 \frac{\phi}{M_P}) \quad , \quad t = \langle t \rangle (1 + 5.00 \frac{\phi}{M_P})
$$

$$
\sigma = -\langle t \rangle 1.15 F^{4/3} \beta \frac{\eta}{M_P} \quad , \quad \chi = -\langle t \rangle 5.00 \frac{\eta}{M_P} \quad . \tag{111}
$$

It is *important* to note that s and t in this expression satisfy

$$
s = .230F^{4/3}\beta t , \qquad (112)
$$

that is, the same relationship as in (89) . Hence, the expansion of s and t around vacuum expectation values $\langle s \rangle$ and $\langle t \rangle$ satisfying $FI = 0$, and integrating out the heavy scalar ξ^1 while keeping the light scalar ξ^2 in the effective theory, restricts all values of s and t to lie along the D -flat direction

$$
s = \frac{\epsilon_S' \beta}{3} t = .230 F^{4/3} \beta t \tag{113}
$$

in which $V_D = 0$.

6 Determining the Moduli Vacua

In the previous Section, we constructed the D-term potential energy V_D for the dilaton and Kähler modulus induced by the their inhomogeneous transformations under the anomalous $U(1)$ gauge group. The minima of this D-term potential energy was shown to lie along a one-dimensional line in s and t space for which $V_D = 0$. Specifically, $FI = 0$ and, hence, $N = 1$ supersymmetry is preserved for s and t satisfying relation (113).

In this Section, we will explicitly assume that parameters F , β , l and the values of $\langle t \rangle$ are restricted so that $m_{\text{anom}} \gtrsim M_U$. It then follows from (98) that, for fixed values of F, β and l, there is a maximum value of $\langle t \rangle$, which we denote $\langle t \rangle_{bound}$, given by

$$
\langle t \rangle_{bound} = \left(\frac{3.39l}{F\beta^{1/2}}\right)^{2/3} \,. \tag{114}
$$

Below this bound, one can integrate complex scalar ξ^1 out of the low energy theory, leaving s, t, σ and χ to be expressed in terms of $\langle t \rangle$, ϕ and η as in (111). In this Section, we analyze the F-term potential given in (71) along the $V_D = 0$ line for $\langle t \rangle \lesssim \langle t \rangle_{bound}$ using (111). We will discuss the case where $\langle t \rangle \gg \langle t \rangle_{bound}$ in the next Section.

Finally, for concreteness, we henceforth assume that the commutant subgroup to the anomalous $U(1)$ is E_7 and, therefore, that $b_L = 6$ – as stated in (51). Using the expression for α_{GUT} in (50), it follows that the parameter b in (49) is given by

$$
b = \frac{41.2}{F^{2/3}} \tag{115}
$$

6.1 The F-term Potential Along the $V_D = 0$ Line

Inserting expressions (111) for s, t, σ and χ into the F-term potential energy V_F given in (71), and using b given in (115), we find

$$
V_F(\langle t \rangle, \tilde{\eta}, \tilde{\phi}) = \frac{M_0^4}{F^{4/3} \beta \langle t \rangle^4 \langle c \rangle^3 (0.230 + 1.15\tilde{\phi})(1 + 5.00\tilde{\phi})^3}
$$

\n
$$
\times \left[1.138 F^{-4/3} \tilde{d}(A^2 + B^2) + 1.32 \times 10^{-6} \tilde{d}^{-1} \left((1 + 19.0F^{2/3} \beta \langle t \rangle (1 + 5.01\tilde{\phi})^2 + 3\right) + 1.32 \times 10^{-8} \tilde{d}^{-1} \left((1 + 19.0F^{2/3} \beta \langle t \rangle (1 + 5.01\tilde{\phi})\right) - (2.43 \times 10^{-3} F^{-2/3}) \left(1 + 19.0 F^{2/3} \beta \langle t \rangle (1 + 5.01\tilde{\phi})\right) + 2.62 \times 10^{-8} \tilde{d}^{-2/3} \beta \langle t \rangle (1 + 5.00\tilde{\phi})
$$

\n
$$
\times sgn(A) \sqrt{A^2 + B^2} \cos[47.5 F^{2/3} \beta \langle t \rangle \tilde{\eta} - arctan(\frac{B}{A})]
$$

\n
$$
+ 2.62 \times 10^{-6} \tilde{d}^{-1} p \left(5.50 + \langle t \rangle (19.0 F^{2/3} \beta (1 + 5.01\tilde{\phi}) + 3\tau (1 + 5.00\tilde{\phi})\right)) + 2.62 \times 10^{-6} \tilde{d}^{-1} p \left(5.50 + \langle t \rangle (19.0 F^{2/3} \beta (1 + 5.01\tilde{\phi}) + 3\tau (1 + 5.00\tilde{\phi})\right) - 2.43 F^{2/3} \beta \langle t + 5.00 \tau \rangle \langle t \rangle \tilde{\eta} + \theta_p]
$$

\n
$$
+ 4.36 \times 10^{-7} \tilde{d}^{-1} p^2 (3 + (3 + 2 \tau \langle t \rangle (1 + 5.005\tilde{\phi}))^2) + 2.43 \times 10^{-3} F^{-2/3} p (1 + 2\tau \langle t \rangle (
$$

where $\tilde{\eta} = \eta/M_P$ and $\tilde{\phi} = \phi/M_P$ are dimensionless. As in (71), coefficients A, B, $\langle c \rangle$ are defined in (32), F is defined in (14),(15), τ is given in (63) and p, θ_p arise in (68).

This is a rather complicated expression which is difficult to analyze analytically. We find it most illuminating to simply choose some fixed values for the flux parameters A, B, $\langle c \rangle$, and the coefficients F, β subject to the constraints on them discussed above. We will also choose a realistic fixed value for τ and set $\tilde{d} = 1$. The potential V_F will then be evaluated at these fixed parameters, initially allowing parameters p and θ_p to take arbitrary values. The choices of F, β and l will also, using (114), determine a fixed value for

 $\langle t \rangle_{bound}$. We then plot V_F in the range $0 \le \langle t \rangle \lesssim \langle t \rangle_{bound}$ for various values of p and θ_p .

6.2 Stabilizing the Axion

It is important to note that the "light" axion $\tilde{\eta}$ enters V_F in (116) via three cosine terms, specifically in the third, fourth and sixth terms, each with a different coefficient that depends on the parameters listed above as well as on $\langle t \rangle$. Therefore, before discussing the stabilization of $\langle t \rangle$, it is essential to determine the vacuum expectation value of $\tilde{\eta}$. To do this, we first note that the coefficient of each of the three cosines has an exponentially suppressed multiplicative factor. For the third, fourth and sixth terms in (116) – that is, the terms containing the cosines – these factors are

$$
\exp[-9.49 F^{2/3} \beta \langle t \rangle], \exp[-(9.49 F^{2/3} \beta + \tau) \langle t \rangle], \exp[-\tau \langle t \rangle] \quad (117)
$$

respectively, where we have ignored the ϕ field contribution which simply enhances the suppression of each term equally. For the physically acceptable range of parameters F, β and τ discussed above, we find that the value of $\langle t \rangle$ is such that the first two exponentials in (117) are greatly suppressed relative to the third entry – typically by a factor of 10^{-6} or smaller. Hence, a very good approximation for determining $\langle \tilde{\eta} \rangle$ is to drop the third and fourth terms in (116) and to evaluate $\langle \tilde{\eta} \rangle$ using the sixth term only. Then, using the fact that there is a minus sign in front of the sixth term, we find

$$
\frac{\partial V_F}{\partial \tilde{\eta}} = 0, \quad \frac{\partial^2 V_F}{\partial^2 \tilde{\eta}} > 0 \quad \Rightarrow \quad \langle \tilde{\eta} \rangle = \frac{2\pi n + \arctan(\frac{B}{A}) - \theta_p}{5.005\tau \langle t \rangle} \tag{118}
$$

where $n \in \mathbb{Z}$. We conclude that once we have found the local minimum for $\langle t \rangle$ in the next subsection, substituting its value into (118) gives one a very good approximation for the local minima of $\tilde{\eta}$ as a function of the Pfaffian parameter θ_p . As a check, we calculated the minima for $\tilde{\eta}$ including all three cosine terms using Mathematica. We find that expression (118) is indeed the correct expression for $\langle \tilde{\eta} \rangle$ to a very high degree of accuracy. That is, for any *value of* θ_p we have stabilized the "light" axion. Having done this, we now proceed to computing the vacuum expectation value for $\langle t \rangle$ which minimizes the V_F potential energy.

6.3 Stabilizing $\langle t \rangle$: An Explicit Example

In this paper, we will present only a single, but physically representative, set of parameters as an example. A much more comprehensive discussion of V_F will be presented elsewhere.

First, recall that V_F is a function of $\langle t \rangle$, $\tilde{\phi}$ and $\tilde{\eta}$. It is clear from the (111) that ϕ is simply the fluctuation around $\langle t \rangle$ in the t direction. Hence, to evaluate $\langle t \rangle$ one can simply set $\phi = 0$. Furthermore, as stated above, to a very high degree of accuracy one can approximate $\tilde{\eta}$ by the vacuum expectation value given in (118)–which sets the associated cosine factor to unity for any value of θ_p . Having done this, V_F becomes a function of $\langle t \rangle$ only. We emphasize that, although we have used (118) as an approximation to $\langle \tilde{\eta} \rangle$, the following calculation uses all terms for V_F given in (116). In Figure 1, we show a sequence of potentials V_F corresponding to fixed flux coefficients we show a sequence or poten
 $A = 2/3$, $B = A/\sqrt{3}$, $c = \frac{1}{\sqrt{3}}$ $\frac{1}{3}$, as well as choosing $\beta = 2$, $l = 2$, $F = 2$ and $\tau = 2$. However, we allow the Pfaffian coefficient p to vary over a range of values. It follows from (114) that, for this choice of parameters,

$$
\langle t \rangle_{bound} = 1.79 \tag{119}
$$

Therefore, we plot these curves in the region where $0 \leq \langle t \rangle \leq 1.79$. From bottom to top, the shapes range from having a global minimum with $V_F < 0$; a global minimum with $V_F = 0$; a local minimum with $V_F > 0$; an inflection point; and no extremum at all. We find that a similar range of potential shapes satisfying these same approximation conditions can be obtained for $1 < \tau < 3.75$ (keeping all other parameters fixed).

Whether there is a V_F extremum at a given $\langle t \rangle$ depends on the value of p. Choosing an explicit value of p, one can attempt to solve $dV_F/d\langle t\rangle = 0$ over the range $0 \le \langle t \rangle \le \langle t \rangle_{bound}$. For some choices of p, there will be no solution and, hence, there are no extrema of the potential V_F over the allowed range of t. This is the case for the green curve in Figure 1. For a special choice of p, there will be a solution of $dV_F/d(t) = 0$ at a single value of $\langle t \rangle$ in the given range. This leads to an inflection point in the potential energy. This is the case for the dashed purple curve in Figure 1. For a finite range of values of p , we find solutions with two extrema, one corresponding to a minimum and the other to a local maximum of the potential energy. This is the case for the blue, red and solid purple curves in Figure 1. The values of p leading to each of these results are given in the caption for Figure 1. Note that the value of p is progressively decreasing from the bottom curve

Figure 1: Range of potential shapes for fixed $A = 2/3$, $B = A/\sqrt{3}$, $c = \frac{1}{\sqrt{3}}$ $\frac{1}{3}$ as well as $l = 2$, $\beta = 2$, $F = 2$, $\tau = 2$ and an arbitrary parameter θ_p , but varying the Pfaffian coefficient p . The values of p for the curves from the bottom to the top are $p = (370, 347, 333, 316, 307)$ respectively. $\langle t \rangle_{bound}$ represents the value of $\langle t \rangle$ above which $m_{\text{anom}} < sM_U$ and so the potential shape may not be precise.

(blue, with a negative energy global minimum) through the red curve (zero energy global minimum), solid purple curve (positive energy local minimum), dashed purple curve (positive energy inflection point) and green curve (no extremum). Experimenting over a substantial range of parameters p , we find that a large number give a potential energy with a negative global minimum while a smaller, but substantial, number lead to a zero or positive potential energy minimum. Finally, as a check on these results, we performed the above calculations using all three cosines to determine $\langle \tilde{\eta} \rangle$, rather than simply inserting (118). The results are identical to the above to at least decimal places.

Given potential energy (116), one can determine the masses of the dimen-

	$m_{\rm anom}$	$m_{\scriptscriptstyle \phi}$	m_{n}
blue $(V_{\min} < 0)$	1.5×10^{17} GeV	GeV 1.6×10^{15}	1.2×10^{15} GeV
red $(V_{\min} = 0)$	1.4×10^{17} GeV	1.3×10^{15} GeV	1.7×10^{15} GeV
solid purple $(V_{\min} > 0)$	GeV 1.0×10^{17}	1.1×10^{15} GeV	1.4×10^{15} GeV

Table 1: The values for m_{anom} and m_{ϕ} , m_{η} at the minima of the blue, red and solid purple curves in Figure 1.

sionful fluctuation fields ϕ (= $\tilde{\phi}M_P$) and η (= $\tilde{\eta}M_P$) by computing $\partial^2V_F/\partial^2\phi$ and $\partial^2 V_F/\partial^2 \eta$ evaluated at the minima at $\langle t \rangle \langle t \rangle_{bound}$ for the blue, red and solid purple curves, respectively. These masses, along with the associated value of m_{anom} computed at each such $\langle t \rangle$ using (98), are presented in Table 1. Note that all values of m_{anom} exceed M_U , as they must. Importantly, in all three cases the values of the m_{ϕ} and m_{η} are each over an order of magnitude smaller than $M_U = 3.15 \times 10^{16} \text{GeV}$. Hence, the ϕ and η moduli remain in the low energy effective field theory.

6.4 Finding the Range of the Pfaffian Coefficient p

Note that the coefficient $F = 2$ chosen for V_F associated with Figure 1, is the largest value of F in the "standard" range given in (15). However, as alluded to in Subsection 2.2, it is possible in specific vacua for the value of the coefficient F to either exceed or be less than the upper and lower bounds respectively presented in (15). The vacua discussed in this paper are exactly of this type. The reason is the following. It has been shown [4, 48] that in generic heterotic M-theory vacua, the "effective" expansion parameter is given by

$$
\epsilon_S^{\text{eff}} = 2\epsilon_S' \frac{t}{s} \,,\tag{120}
$$

where we have used the fact that $s = V$ and that, in the $h^{1,1} = 1$ case, $\hat{R} = 2t$. It then follows from (113) that

$$
\epsilon_S^{\text{eff}} = \frac{6}{\beta} \,. \tag{121}
$$

We conclude that in the vacua we are considering – that is, the $h^{1,1} = 1$ case where s and t satisfy $FI = 0$, thus setting $V_D = 0$ – the size of the effective strong coupling parameter is set by parameter β , but is *independent* of the coefficient F . Hence, F need not be bounded by the constraints in (15) and can, in principle, take any value – including values larger than 2 and smaller than 0.6.

However, in considering possible shapes of the potential energy curve V_F , the value of F cannot be made arbitrarily large. This is because the condition $m_{anom} \geq M_U$ has been assumed in deriving V_F , and this can only be satisfied for $0 \le \langle t \rangle \lesssim \langle t \rangle_{bound}$ where, from (114), $\langle t \rangle_{bound} \propto 1/F^{2/3}$. To be sure of potentials with stable or metastable minima and local maxima, like the bottom three curves in Figure 1, it must be that $\langle t \rangle_{bound}$ exceeds $\langle t \rangle$ at the local maximum for each curve, which sets an upper bound for F . As a concrete example, let us consider the blue curve in Figure 1. This curve was determined using $F = 2$, leading to $\langle t \rangle_{bound} = 1.79$ given in (119). It will be shown in a subsequent publication [37] that the values of $\langle t \rangle$ at both the minima and maxima of any V_F curve will remain unchanged under a change in parameter F , as long as the Pfaffian parameter is appropriately adjusted.

Let us denote the value of $\langle t \rangle$ at the local maximum of the blue curve by $\langle t \rangle_{max}$. We see from Figure 1 that $\langle t \rangle_{max} = 1.1$. Let us now gradually raise the value of F to lower the value of $\langle t \rangle_{bound}$ – but appropriately adjusting p at each stage so that the extrema of the blue curve remain at the same values of $\langle t \rangle$. We can continue until we reach the point where

$$
\langle t \rangle_{bound} = \langle t \rangle_{max} = 1.1 \tag{122}
$$

This occurs when F reaches $F = 4$, in accordance with (114) (and using the values of l and β cited in the caption of Figure 1). Note that this is significantly larger than the conventional upper bound $F = 2$ given in (15) and used in constructing Figure 1. As will be shown in [37], the appropriate adjustment of p is to rescale, $p \propto F^{-2/3}$. In other words, in changing from $F = 2$ to $F = 4$, the values of $\langle t \rangle$ at the minima and maxima do not change if one adjusts the corresponding value of Pfaffian parameter substantially from $p = 370$ to $p = 233$. A similar adjustment is required for the two higher curves–that is, the red and solid purple curves–in Figure 1. The values of $\langle t \rangle$ at their maxima, the values of F_{max} and the associated values of p for each of these three curves are given in Table 2. Having done this, it is important to check that the masses of the fields ϕ and η at the local minima of each of these three curves continue to be considerably smaller than the value of m_{anom} evaluated at each such local minimum. This turns out to to be the case, as is shown explicitly in Table 3.

	$\langle t \rangle$ at max	F_{max}	
blue $(V_{\text{min}} < 0)$			233
red $(V_{\text{min}} = 0)$	10		205
solid $(V_{\min} > 0)$	1.97	4.X	185.

Table 2: The values of $\langle t \rangle$ at the local maximum and the associated values for F_{max} and p when $\langle t \rangle_{bound}$ is set equal to $\langle t \rangle$ maximum, for the blue, red and solid purple curves in Figure 1 respectively.

	m_{anom}	m_{ϕ}	$m_{\scriptscriptstyle\cal n}$
blue $(V_{\min} < 0)$	7.1×10^{16} GeV	6.5×10^{14} GeV	4.7×10^{14} GeV
red $(V_{\min} = 0)$	6.5×10^{16} GeV	5.8×10^{14} GeV	4.2×10^{14} GeV
solid purple $(V_{\min} > 0)$	5.9×10^{16} GeV	5.1×10^{14} GeV	3.7×10^{14} GeV

Table 3: The values of m_{anom} and m_{ϕ} , m_{η} for each of the three curves defined in Table 2. Note that, in all cases, m_{ϕ} and m_{η} are each $\ll m_{anom}$.

We conclude that by lowering $\langle t \rangle_{bound}$ to the values of the local maxima for each of the blue, red and solid purple curves in Figure 1, the values of the Pfaffian coefficient p vary through a substantial range. As a second example, a red curve with vanishing potential energy at the same value of $\langle t \rangle$ is possible for the Pfaffian coefficient range $205 < p < 347$. The ability to stabilize the expectation values of dilaton and geometric moduli at the same value for a wide range of the Pfaffian coefficient p has an important implication. As mentioned in Section 4, when fixed at a supersymmetry preserving minimum of the vector bundle moduli, the Pfaffian becomes a complex number specified in (67) by an amplitude p and a phase θ_p . As discussed above, it is not known how to explicitly calculate their values. However, it is trivial to show that the values of $\langle s \rangle$ and $\langle t \rangle$ at the potential minimum do not depend on θ_p at all; and now, as we have demonstrated, their values can also be obtained for a wide range of p . The fact that the dilaton and geometric moduli can have the same expectation values for a wide range of Pfaffian parameter p means there is more likely to be some vector bundle moduli vacuum that produces a p in that range and stabilizes those values. The explicit method for determining the Pfaffian parameter p for each type of potential in Figure 1 will be presented in detail in a subsequent publication [37].

7 Swampland Bound on the Potential at Large Values of t

Our explicit construction of potentials for heterotic M-theory compactified on Calabi-Yau threefolds with $h^{1,1} = h^{2,1} = 1$ provides interesting test cases for the Swampland conjectures [31, 32, 33, 34, 35, 36]. Among the different conjectures, the Transplanckian Censorship Conjecture [35] and the Strong de Sitter Conjecture [36] both postulate that, for large values of the moduli fields (with canonically normalized kinetic energy), there is a positive lower bound on the gradient of the potential when $V > 0$, namely

$$
\frac{|\nabla V|}{V} \ge \frac{2}{\sqrt{d-2}},\tag{123}
$$

where d is the spacetime dimension. Since $d = 4$ in our case, the Swampland where *a* is the spacetime din
lower bound is equal to $\sqrt{2}$.

To evaluate whether our potential satisfies this Swampland conjecture we first recall that the total potential energy density is $V = V_D + V_F$, where V_D is the function of s and t given in (84) . Since we are interested in the large t limit where $t \gg t_{bound}$, V_F is given in (71) (and not by (116), which assumes $t \lesssim t_{bound}$). Now, as discussed in Subsection 5.3, $V_D = 0$ for $s = .2301F^{4/3}\beta t$, which we assume henceforth. Inserting this into (71) , it follows that V_F is a function of the modulus field t and the axions σ and χ . Although this V_F has numerous terms, the task of evaluating its large field limit is straightforward. Except for the first term, all other terms in V_F are suppressed by a factor of $\exp(-ct)$ for some positive coefficient c (which differs for each of the terms). This includes all three terms containing the axions σ and χ , so it is not necessary to consider their large field limits. Hence, the first term dominates all other terms in the large t limit. Keeping only the first term in V_F , we have

$$
V_F \propto \frac{1}{st^3} \propto \frac{1}{t^4} \tag{124}
$$

where have imposed the condition that $s \propto t$ along the D-flat direction. We do not need to use the exact proportionality constant because the Swampland condition uses the logarithmic derivative of V_F , so any constant factors drop out.

To proceed, we note that the Swampland condition (123) requires the moduli fields to have canonically normalized kinetic energy. The kinetic energy for $t = \text{Re } T$ is given by

$$
\kappa_4^2 \frac{\partial^2 K}{\partial T \partial \bar{T}} (\partial T \partial \bar{T})|_{\text{Im} T=0} = \frac{3}{4} \frac{(\partial t)^2}{t^2}
$$
 (125)

where $\kappa_4^2 K = -3 \ln (T + \bar{T})$ is the Kähler potential given in (35). To rewrite this kinetic energy in terms of a field Φ with canonical kinetic energy, we use the ansatz $\Phi \equiv q \ln t$, where q is a constant to be determined by the condition that Φ has canonical kinetic energy. That is, we set

$$
\frac{1}{2}(\partial \Phi)^2 = \frac{q^2}{2} \frac{(\partial t)^2}{t^2}.
$$
\n(126)

equal to Eqs. (125) and obtain:

$$
\frac{q^2}{2} = \frac{3}{4} \Rightarrow q = \sqrt{3/2}.
$$
 (127)

Hence,

$$
\Phi = \sqrt{3/2} \ln t \tag{128}
$$

What remains, then, is to rewrite V_F in terms of canonical field Φ in the limit of large t and to check if the Swampland constraint (123) is satisfied. Using (128), we can rewrite the potential in (124) as

$$
V_F \propto e^{-4\sqrt{2/3}\Phi},\tag{129}
$$

from which we obtain

$$
\frac{|\nabla V|}{V} = \frac{|dV_F/d\Phi|}{V_F} = 4\sqrt{2/3} > \sqrt{2} .
$$
 (130)

That is, our theory exceeds the Swampland lower bound of $\sqrt{2}$ in the large field limit [35]. A discussion of how our results relate to Swampland conjectures concerning conditions at small values of the $\langle t \rangle$ field near the center of moduli space will be presented elsewhere [37].

Acknowledgements

We thank Alek Bedroya and Cumrun Vafa for useful comments and for reviewing the manuscript. Burt Ovrut is supported in part by both the research

grant DOE No. DESC0007901 and SAS Account 020-0188-2-010202-6603- 0338. Ovrut would like to acknowledge the hospitality of the CCPP at New York University where much of this work was carried out. Paul Steinhardt supported in part by the DOE grant number DEFG02-91ER40671 and by the Simons Foundation grant number 654561. Steinhardt thanks Cumrun Vafa and the High Energy Physics group in the Department of Physics at Harvard University for graciously hosting him during his sabbatical leave.

References

- [1] P. Horava and E. Witten, "Eleven-dimensional supergravity on a manifold with boundary," Nucl. Phys. B 475, 94-114 (1996) doi:10.1016/0550-3213(96)00308-2 [arXiv:hep-th/9603142 [hep-th]]
- [2] P. Horava and E. Witten, "Heterotic and type I string dynamics from eleven-dimensions," Nucl. Phys. B 460, 506-524 (1996) doi:10.1016/0550-3213(95)00621-4 [arXiv:hep-th/9510209 [hep-th]].
- [3] A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, "The Universe as a domain wall," Phys. Rev. D 59, 086001 (1999) doi:10.1103/PhysRevD.59.086001 [arXiv:hep-th/9803235 [hep-th]]
- [4] A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, "Heterotic M theory in five-dimensions," Nucl. Phys. B 552, 246-290 (1999) doi:10.1016/S0550-3213(99)00196-0 [arXiv:hep-th/9806051 [hep-th]].
- [5] A. Ashmore, S. Dumitru and B. A. Ovrut, "Line Bundle Hidden Sectors for Strongly Coupled Heterotic Standard Models," Fortsch. Phys. 69, no.7, 2100052 (2021) doi:10.1002/prop.202100052 [arXiv:2003.05455 [hep-th]].
- [6] M. B. Green and J. H. Schwarz, "Anomaly Cancellation in Supersymmetric D=10 Gauge Theory and Superstring Theory," Phys. Lett. B 149, 117-122 (1984) doi:10.1016/0370-2693(84)91565-X
- [7] S. Gukov, S. Kachru, X. Liu and L. McAllister, "Heterotic moduli stabilization with fractional Chern-Simons invariants," Phys. Rev. D 69, 086008 (2004) doi:10.1103/PhysRevD.69.086008 [arXiv:hep-th/0310159 [hep-th]].
- [8] V. Balasubramanian, P. Berglund, J. P. Conlon and F. Quevedo, "Systematics of moduli stabilisation in Calabi-Yau flux compactifications," JHEP 03, 007 (2005) doi:10.1088/1126-6708/2005/03/007 [arXiv:hepth/0502058 [hep-th]].
- [9] L. B. Anderson, J. Gray, A. Lukas and B. Ovrut, "Stabilizing All Geometric Moduli in Heterotic Calabi-Yau Vacua," Phys. Rev. D 83, 106011 (2011) doi:10.1103/PhysRevD.83.106011 [arXiv:1102.0011 [hep-th]].
- [10] M. Cicoli, S. de Alwis and A. Westphal, "Heterotic Moduli Stabilisation," JHEP 10, 199 (2013) doi:10.1007/JHEP10(2013)199 [arXiv:1304.1809 [hep-th]].
- [11] P. Candelas and X. de la Ossa, "Moduli Space of Calabi-Yau Manifolds," Nucl. Phys. B 355, 455-481 (1991) doi:10.1016/0550-3213(91)90122-E
- [12] J. Gray, A. Lukas and B. Ovrut, "Flux, gaugino condensation and anti-branes in heterotic M-theory," Phys. Rev. D 76, 126012 (2007) doi:10.1103/PhysRevD.76.126012 [arXiv:0709.2914 [hep-th]].
- [13] M. Dine, R. Rohm, N. Seiberg and E. Witten, "Gluino Condensation in Superstring Models," Phys. Lett. B 156, 55-60 (1985) doi:10.1016/0370- 2693(85)91354-1
- [14] H. P. Nilles, "Gaugino Condensation and Supersymmetry Breakdown," Int. J. Mod. Phys. A 5, 4199-4224 (1990) doi:10.1142/S0217751X90001744
- [15] P. Horava, "Gluino condensation in strongly coupled heterotic string theory," Phys. Rev. D 54, 7561-7569 (1996) doi:10.1103/PhysRevD.54.7561 [arXiv:hep-th/9608019 [hep-th]]
- [16] A. Lukas, B. A. Ovrut and D. Waldram, "Gaugino condensation in M theory on $s^{**}1 / Z(2)$," Phys. Rev. D 57, 7529-7538 (1998) doi:10.1103/PhysRevD.57.7529 [arXiv:hep-th/9711197 [hep-th]].
- [17] M. Dine, N. Seiberg, X. G. Wen and E. Witten, "Nonperturbative Effects on the String World Sheet," Nucl. Phys. B 278, 769-789 (1986) doi:10.1016/0550-3213(86)90418-9
- [18] M. Dine, N. Seiberg, X. G. Wen and E. Witten, "Nonperturbative Effects on the String World Sheet. 2.," Nucl. Phys. B 289, 319-363 (1987) doi:10.1016/0550-3213(87)90383-X
- [19] E. Witten, "Nonperturbative superpotentials in string theory," Nucl. Phys. B 474, 343-360 (1996) doi:10.1016/0550-3213(96)00283-0 $[\text{arXiv:hep-th}/9604030$ $[\text{hep-th}]]$.
- [20] E. I. Buchbinder, R. Donagi and B. A. Ovrut, "Superpotentials for vector bundle moduli," Nucl. Phys. B 653, 400-420 (2003) doi:10.1016/S0550-3213(02)01093-3 [arXiv:hep-th/0205190 [hep-th]].
- [21] G. Curio, "Perspectives on Pfaffians of Heterotic World-sheet Instantons," JHEP 09, 131 (2009) doi:10.1088/1126-6708/2009/09/131 [arXiv:0904.2738 [hep-th]].
- [22] C. Beasley and E. Witten, "Residues and world sheet instantons," JHEP 10, 065 (2003) doi:10.1088/1126-6708/2003/10/065 [arXiv:hepth/0304115 [hep-th]].
- [23] E. Buchbinder, A. Lukas, B. Ovrut and F. Ruehle, "Heterotic Instanton Superpotentials from Complete Intersection Calabi-Yau Manifolds," JHEP 10, 032 (2017) doi:10.1007/JHEP10(2017)032 [arXiv:1707.07214 [hep-th]].
- [24] E. I. Buchbinder, L. Lin and B. A. Ovrut, "Non-vanishing Heterotic Superpotentials on Elliptic Fibrations," JHEP 09, 111 (2018) doi:10.1007/JHEP09(2018)111 [arXiv:1806.04669 [hep-th]].
- [25] E. I. Buchbinder, A. Lukas, B. A. Ovrut and F. Ruehle, "Heterotic Instantons for Monad and Extension Bundles," JHEP 02, 081 (2020) doi:10.1007/JHEP02(2020)081 [arXiv:1912.07222 [hep-th]].
- [26] E. I. Buchbinder, A. Lukas, B. A. Ovrut and F. Ruehle, "Instantons and Hilbert Functions," Phys. Rev. D 102, no.2, 026019 (2020) doi:10.1103/PhysRevD.102.026019 [arXiv:1912.08358 [hep-th]].
- [27] E. I. Buchbinder and B. A. Ovrut, "Non-vanishing Superpotentials in Heterotic String Theory and Discrete Torsion," JHEP 01, 038 (2017) doi:10.1007/JHEP01(2017)038 [arXiv:1611.01922 [hep-th]].
- [28] S. Dumitru and B. A. Ovrut, "Heterotic M-Theory Hidden Sectors with an Anomalous $U(1)$ Gauge Symmetry," [arXiv:2109.13781 [hep-th]].
- [29] S. Dumitru and B. A. Ovrut, "Moduli and Hidden Matter in Heterotic M-Theory with an Anomalous $U(1)$ Hidden Sector," $\arcsin 2201.01624$ [hep-th]].
- [30] D. Z. Freedman and A. Van Proeyen, "Supergravity," Cambridge Univ. Press, 2012, ISBN 978-1-139-36806-3, 978-0-521-19401-3
- [31] H. Ooguri and C. Vafa, "On the Geometry of the String Landscape and the Swampland," Nucl. Phys. B 766, 21-33 (2007) doi:10.1016/j.nuclphysb.2006.10.033 [arXiv:hep-th/0605264 [hep-th]].
- [32] H. Ooguri, E. Palti, G. Shiu and C. Vafa, "Distance and de Sitter Conjectures on the Swampland," Phys. Lett. B 788, 180-184 (2019) doi:10.1016/j.physletb.2018.11.018 [arXiv:1810.05506 [hep-th]].
- [33] D. Lüst, E. Palti and C. Vafa, "AdS and the Swampland," Phys. Lett. B 797, 134867 (2019) doi:10.1016/j.physletb.2019.134867 [arXiv:1906.05225 [hep-th]].
- [34] E. Palti, "The Swampland: Introduction and Review," Fortsch.Phys. 67, 1900037 (2019) doi:10.10.1002/prop.201900037 [arXiv:1903.06239 $\vert \text{hep-th} \vert$.
- [35] A. Bedroya and C. Vafa, "Trans-Planckian Censorship and the Swampland," JHEP 09, 123 (2020) doi:10.1007/JHEP09(2020)123 [arXiv:1909.11063 [hep-th]].
- [36] T. Rudelius, "Asymptotic observables and the swampland," PRD 104, 126023 (2021) doi:10.1103/PhysRevD.104.126023 [arXiv:/2106.09026 [hep-th]].
- [37] C. Deffayet, B.A. Ovrut and P.J. Steinhardt "Stable and Metastable Vacua in Heterotic M-theory Models that Incorporate Standard Model Physics and ΛCDM Cosmology," in preparation.
- [38] R. Deen, B. A. Ovrut and A. Purves, "The minimal SUSY B L model: simultaneous Wilson lines and string thresholds," JHEP 07, 043 (2016) doi:10.1007/JHEP07(2016)043 [arXiv:1604.08588 [hep-ph]].
- [39] A. Ashmore, S. Dumitru and B. A. Ovrut, "Explicit soft supersymmetry breaking in the heterotic M-theory $B - L$ MSSM," JHEP 08, 033 (2021) doi:10.1007/JHEP08(2021)033 [arXiv:2012.11029 [hep-th]].
- [40] T. Banks and M. Dine, "Couplings and scales in strongly coupled heterotic string theory," Nucl. Phys. B 479, 173-196 (1996) doi:10.1016/0550-3213(96)00457-9 [arXiv:hep-th/9710208 [hep-th]]
- [41] A. Lukas, B. A. Ovrut and D. Waldram, "On the four-dimensional effective action of strongly coupled heterotic string theory," Nucl. Phys. B 532, 43-82 (1998) doi:10.1016/S0550-3213(98)00463-5 [arXiv:hepth/9710208 [hep-th]].
- [42] V. Braun, Y. H. He, B. A. Ovrut and T. Pantev, "The Exact MSSM spectrum from string theory," JHEP 05, 043 (2006) doi:10.1088/1126- $6708/2006/05/043$ [arXiv:hep-th/0512177 [hep-th]].
- [43] C. P. Burgess, C. Escoda and F. Quevedo, "Nonrenormalization of flux superpotentials in string theory," JHEP 06, 044 (2006) doi:10.1088/1126-6708/2006/06/044 [arXiv:hep-th/0510213 [hep-th]].
- [44] E. I. Buchbinder and B. A. Ovrut, "Vacuum stability in heterotic M theory," Phys. Rev. D 69, 086010 (2004) doi:10.1103/PhysRevD.69.086010 $[\text{arXiv:hep-th}/0310112 \text{ [hep-th]}.$
- [45] G. Curio, "On the Heterotic World-sheet Instanton Superpotential and its individual Contributions," JHEP 08, 092 (2010) doi:10.1007/JHEP08(2010)092 [arXiv:1006.5568 [hep-th]].
- [46] V. Braun, M. Kreuzer, B. A. Ovrut and E. Scheidegger, "Worldsheet instantons and torsion curves, part A: Direct computation," JHEP 10, 022 (2007) doi:10.1088/1126-6708/2007/10/022 [arXiv:hep-th/0703182 $\vert \text{hep-th} \vert$.
- [47] V. Braun, M. Kreuzer, B. A. Ovrut and E. Scheidegger, "Worldsheet instantons, torsion curves, and non-perturbative superpotentials," Phys. Lett. B 649, 334-341 (2007) doi:10.1016/j.physletb.2007.03.066 $[\text{arXiv:hep-th}/0703134$ $[\text{hep-th}].$

[48] Burt A. Ovrut, "Vacuum Constraints for Realistic Strongly Coupled Heterotic M-Theories," Symmetry 10, 723 (2018) doi:10.3390/sym10120723 [arXiv:hep-th/1811.08892[hep-th]].