Manifestation of heavy 3/2-spin lepton in large-angle $e^+e^-\to \gamma\gamma$ reaction at high energies

G. I. Gakh |, M.I. Konchatnij,^{*} and N.P. Merenkov^{[†](#page-0-1)}

National Science Centre, Kharkov Institute of Physics and Technology, Akademicheskaya 1, and V. N. Karazin Kharkov National University,

Dept. of Physics and Technology, 31 Kurchatov, 61108 Kharkov, Ukraine

A. G. Gakh [‡](#page-0-2)

V. N. Karazin Kharkov National University, Dept. of Physics and Technology, 31 Kurchatov, 61108 Kharkov, Ukraine

Abstract

Manifestation of the heavy 3/2-spin lepton (h^{\pm}) , as possible virtual intermediate state in Feynmann diagrams, have been searched in the $e^+e^- \rightarrow \gamma\gamma$ reaction at high energies and large photon angles. The spin-vector field Ψ_{α} of the 3/2-lepton is described by the Rarita-Schwinger one and phenomenological Lagrangian of $h^{\pm} e^{\pm} \gamma$ -interaction is chosen similarly to interaction of Δ isobar with nucleon and γ quant. It is described by two constants with dimension $[M^{-1}]$ and $[M^{-2}]$. The differential cross section and polarization asymmetries have been calculated for the case when both beams are polarized longitudinally along their directions, as well transversally, in the reaction plane, and normally, perpendicularly to it. Numerical estimations are performed in wide diapason of the collision energy and parameters entering phenomenological Lagrangian.

[∗] konchatnij@kipt.kharkov.ua

[†] merenkov@@kipt.kharkov.ua

[‡] agakh@karazin.ua

I. INTRODUCTION

Recent analysis of experimental data collected from 1989 up to 2003, to search the nonpointness of the electron in e^+e^- annihilation reactions (see [\[1\]](#page-15-0) and references therein) showed a significant signal at a confidence level 5 standard deviations. To explain this finding authors suggest the existence of an exited electron with mass about 310 GeV and some contact interaction with a cutoff scale about 1250 GeV. Although the more recent experimental data obtained in electron-proton [\[2](#page-15-1)] and proton-proton [\[3](#page-15-2)[–5\]](#page-15-3) collisions do not see an evidence for the exited lepton, this result is awaiting confirmation at future circular and linear electron-positron colliders such as FCC [\[6\]](#page-15-4), CEPC [\[7\]](#page-15-5), ILC [\[8](#page-15-6)] and ClIC [\[9\]](#page-15-7).

In present paper we consider one more mechanism that can influence such deviation (it may be more exotic one): effects due to existence of a heavy lepton h^{\pm} with spin 3/2 which, like a heavy exited electron, appears as a virtual intermediate state in Feynman diarams. We assume that 3/2-spin heavy lepton is described by the Rarita-Schwinger spin-vector field [\[10](#page-15-8)] and it interacts with at least the QED sector of the Standard Model and can decay into electron and γ quant just similarly to decay of the Δ isobar into nucleon and photon (see, for example, [\[11](#page-15-9), [12\]](#page-15-10)). Therefore it is quite natural to look for the manifestation of this lepton in the large-angle high energy annihilation reaction

$$
e^-(p_1) + e^+(p_2) \to \gamma(k_1) + \gamma(k_2). \tag{1}
$$

The possibility of the existence 3/2-spin heavy lepton within composite models [\[13\]](#page-15-11) and searching for a signal associated with it have been discussed early in a number works [\[14](#page-15-12)– [18\]](#page-15-13). Here we mainly focus on the effects due to arbitrary beam polarizations and consider the influence of the virtual intermediate heavy lepton on various polarization asymmetries in the angular distribution of photons.

II. FORMALISM

To estimate effects due to $3/2$ -spin heavy lepton in the reaction (1) we restrict ourselves with standard QED diagrams (Fig. 1 a)) and diagrams with intermediate heavy lepton h^{\pm} (Fig. 1 b)) instead of electron (HL diagrams). We consider the case when both, electron and positron, can be arbitrary polarized, namely: longitudinally (along their 3-momentum direction) with 4-vector $S^l_{1\mu}$, transversally (in the reaction plane) with $S^t_{1\mu}$ and normal (perpendicularly to the reaction plane) with $S_{1\mu}^n$. Our aim is to search manifestation of the HL diagrams on the

level differential cross section and different polarization asymmetries at high beam energies achievable at future linear colliders ILC and CLIC.

A. Normalization and polarization 4-vectors

We choose a normalization such that, being averaged over the electron and positron polarizations, the differential angular cross section reads

$$
\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{16 s} |M^2|, \ s = (p_1 + p_2)^2, \ t = (p_1 - k_1)^2, \ u = (p_1 - k_2)^2,
$$
\n(2)

where $\alpha = e^2/(4\pi) = 1/137$ fine structure constant and the matrix element squared is defined by a sum of the QED, QED-HL interference and HL contributions. The same expression is valid when electron and positron are polarized. When calculating traces in $|M^2|$ at chosen normalization we have to use the electron and positron projections as (we omit the spin indecies)

$$
u(p_1)\,\bar{u}(p_1) = (\hat{p}_1 + m)(1 + \gamma_5\,\hat{S}_1), \quad v(p_2)\,\bar{v}(p_2) = (\hat{p}_2 - m)(1 + \gamma_5\,\hat{S}_2),\tag{3}
$$

where $u(p_1)$ and $v(p_2)$ are the electron and positron bispinors, respectively.

At high energies, neglecting the electron mass, we can express 4-vectors $S_{1,2\mu}^{l,t,n}$ in terms of particle 4-momenta, namely

$$
S_{1\mu}^{l} = \lambda_{1} \frac{p_{1\mu}}{m}, \ S_{2\mu}^{l} = \lambda_{2} \frac{p_{2\mu}}{m}, \ S_{1\mu}^{t} = \eta_{1} \sqrt{\frac{s}{ut}} \left(\frac{u}{s} p_{1\mu} + \frac{t}{s} p_{2\mu} + k_{1\mu}\right), \ S_{2\mu}^{t} = \frac{\eta_{2}}{\eta_{1}} S_{1\mu}^{t},
$$

$$
S_{1\mu}^{n} = \xi_{1} \frac{\varepsilon_{\mu\nu\lambda\rho} p_{1}^{\nu} p_{2}^{\lambda} k_{1}^{\rho}}{K}, \ K = \frac{1}{2} \sqrt{stu}, \ S_{2\mu}^{n} = \frac{\xi_{2}}{\xi_{1}} S_{1\mu},
$$
 (4)

where $\lambda_{1,2}, \eta_{1,2}$ and $\xi_{1,2}$ are corresponding polarization degrees. In the case of longitudinal polarization one must first perform multiplication in [\(3\)](#page-2-0) and then go to the limit $m \to 0$. Take attention that part of the differential cross section which does not depend on polarizations is the same for unpolarized and polarized beams.

Let $(d\sigma/d\Omega)_{un}$ is the cross section in unpolarized case and $(d\sigma/d\Omega)_{ij}$, $i, j = l, t, n$ is the cross section in the case of positron polarization S_2^i and electron one S_1^j . Then we define the polarization asymmetries A_{ll} , A_{tt} , A_{nn} , A_{lt} and A_{tl} as follows

$$
\frac{d\sigma_{ll}}{d\Omega} = \frac{d\sigma_{un}}{d\Omega} \left(1 \pm \lambda_1 \lambda_2 A_{ll}\right), \quad \frac{d\sigma_{tt}}{d\Omega} = \frac{d\sigma_{un}}{d\Omega} \left(1 \pm \eta_1 \eta_2 A_{tt}\right), \quad \frac{d\sigma_{nn}}{d\Omega} = \frac{d\sigma_{un}}{d\Omega} \left(1 \pm \xi_1 \xi_2 A_{nn}\right), \quad (5)
$$
\n
$$
\frac{d\sigma_{lt}}{d\Omega} = \frac{d\sigma_{un}}{d\Omega} \left(1 \pm \eta_1 \lambda_2 A_{lt}\right), \quad \frac{d\sigma_{tl}}{d\Omega} = \frac{d\sigma_{un}}{d\Omega} \left(1 \pm \lambda_1 \eta_2 A_{tl}\right).
$$

FIG. 1. Feynman diagrams for the process $e^+e^- \rightarrow \gamma \gamma$. a) QED diagrams, b) diagrams with the heavy 3/2-spin lepton h in $t-$ and $u-$ channels.

Signs in front A_{ij} can be changed by changing the sign of one of the 4-vector polarization to the opposite; sign $" +"$ corresponds to choice given by definitions [\(4\)](#page-2-1).

Note that $A_{nt} = A_{nl} = A_{tn} = A_{ln} = 0$ for obvious reason. In the limit $m \to 0$ the Born approximation gives $A_{lt}^{B} = A_{tl}^{B} = 0$ and

$$
\left(\frac{d\sigma}{d\Omega}\right)_{un}^{B} = \frac{\alpha^2}{s} \left(\frac{1+\cos^2\theta}{1-\cos^2\theta}\right), \quad A_{ll}^{B} = -1, \quad A_{tt}^{B} = A_{nn}^{B} = \frac{1-\cos^2\theta}{1+\cos^2\theta}.
$$
\n
$$
(6)
$$

B. About 3/2-spin heavy lepton

We suggest that even-parity $3/2$ -spin lepton h^{\pm} is described by the Rarita-Schwinger spinvector field $\Psi_p^{\alpha}(x) = u^{\alpha}(p) \exp(-ip \cdot x)$ and Lagrangian its interaction with electron and photon is very similar to the Lagrangian describing electromagnetic decay of ∆ isobar into nucleon and γ quant: $\Delta \to N + \gamma$. There are a lot possibilities to choose this Lagrangian, see for example, [\[19](#page-15-14)[–21\]](#page-16-0) but we will use well known one from [\[11\]](#page-15-9)

$$
L_{\Psi^{\alpha} e \gamma} = e \bar{\psi} \gamma_5 \left(\frac{ig_1}{M} \gamma^{\mu} - \frac{g_2}{M^2} \partial^{\mu} \right) \Psi^{\alpha} F_{\alpha \mu} + h.c., \ F_{\alpha \mu} = \partial_{\mu} A_{\alpha} - \partial_{\alpha} A_{\mu}, \tag{7}
$$

where dimensionless constants g_1 and g_2 are real (CP parity is conserved), parameter M has the dimension of mass, and $F_{\alpha\mu}$ is tensor of an electromagnetic field. Below we take mass of the h^{\pm} as the parameter M and assume $M^2 \approx s$.

Lagrangian [\(7\)](#page-3-0) defines an amplitude A_D of the decay $h^-(p_1) \to e^-(p_2) + \gamma(k)$ in terms of constants g_1 and g_2 as follows

$$
A_D = e \bar{u}(p_2) \gamma_5 \Gamma^{\alpha \mu} u_{\alpha}(p_1) \varepsilon_{\mu}^*(k), \qquad (8)
$$

where

$$
\Gamma^{\alpha\mu} = \frac{g_1}{M} (k^{\alpha} \gamma^{\mu} - \hat{k} g^{\alpha \mu}) + \frac{g_2}{M^2} [k^{\alpha} p_1^{\mu} - (k \cdot p_1) g^{\alpha \mu}].
$$

The square of this amplitude reads

$$
|A_D|^2 = e^2 Sp\{(\hat{p}_2 - m) \Gamma^{\alpha \mu} (\hat{p}_1 + M) P_{\alpha \beta}(p_1) \Gamma^{\beta}_{\mu}\},
$$
\n
$$
P_{\alpha \beta}(p_1) = g_{\alpha \beta} - \frac{1}{3} \gamma_{\alpha} \gamma_{\beta} - \frac{1}{3 p_1^2} (\hat{p}_1 \gamma_{\alpha} p_{1\beta} + p_{1\alpha} \gamma_{\beta} \hat{p}_1), \quad \Gamma^{\beta}_{\mu} = \Gamma^{\beta \lambda} g_{\lambda \mu},
$$
\n(9)

where we taken into account that $\sum_{pol} \varepsilon^*_{\mu} \varepsilon_{\rho} = -g_{\mu\rho}$. Using conservation low one obtains the partial decay width neglecting the electron mass

$$
\Gamma(h^{\pm} \to e^{\pm} \gamma) = \frac{\alpha M}{24} (3g_1^2 + g_2^2 + 3g_1 g_2).
$$
 (10)

Note that in this limit the analytic form of the corresponding negative-parity HL decay width coincides with [\(10\)](#page-4-0).

C. Unpolarized cross section

In accordance with Feynmann rules, contribution HL diagrams to the reaction amplitude can be written as follows (we extracted factor α^2 in accordance with the definition [\(2\)](#page-2-2) of the differential cross section)

$$
M_h = \bar{v}(p_2)\gamma_5 \left[\Gamma_{22}^{\beta\nu}\Delta_{\beta\alpha}(p)\Gamma_{11}^{\alpha\mu} + \Gamma_{21}^{\beta\mu}\Delta_{\beta\alpha}(q)\Gamma_{12}^{\alpha\nu}\right]\gamma_5 u(p_1)\epsilon^*_{\mu}(k_1)\,\epsilon^*_{\nu}(k_2),\tag{11}
$$

where $\epsilon_{\mu}(k_1)$ and $\epsilon_{\nu}(k_2)$ are 4-vectors of photon polarizations and

$$
p = p_1 - k_1 = k_2 - p_2, \ q = p_1 - k_2 = k_1 - p_2.
$$

The sort notation for all Γ 's in [\(11\)](#page-4-1) is

$$
\Gamma_{ij}^{\beta\nu} = \frac{g_1}{M} (k_j^{\beta} \gamma^{\nu} - \hat{k}_j g^{\beta \nu}) + \frac{g_2}{M^2} [k_j^{\beta} p_i^{\nu} - (p_i \cdot k_j) g^{\beta \nu}], (i, j = 1, 2)
$$
\n(12)

Propagator $\Delta_{\alpha\beta}$ of the particle with spin 3/2 reads (see Eqs. (13), (14) in Ref. [?])

$$
\Delta_{\alpha\beta}(p) = -\frac{\hat{p} + M}{p^2 - M^2} P_{\alpha\beta}(p), \ P_{\alpha\beta}(p) p^{\beta} = p^{\alpha} P_{\alpha\beta}(p) = 0, \ P_{\alpha\beta}(p) \gamma^{\beta} = \gamma^{\alpha} P_{\alpha\beta}(p) = 0. \tag{13}
$$

To obtain $|M_h|^2$ one has to compute rather long trace

$$
|M_h|^2 = Spur\left\{(\hat{p}_2 - m)\gamma_5 \left[\Gamma_{22}^{\beta\nu}\Delta_{\beta\alpha}(p)\Gamma_{11}^{\alpha\mu} + \Gamma_{21}^{\beta\mu}\Delta_{\beta\alpha}(q)\Gamma_{12}^{\alpha\nu}\right]\gamma_5 \times \tag{14}
$$

$$
(\hat{p}_1 + m)\gamma_5 \left[\Gamma_{11\,\mu}^{\delta} \widetilde{\Delta}_{\delta\sigma}(p) \Gamma_{22\,\nu}^{\sigma} + \Gamma_{12\,\nu}^{\delta} \widetilde{\Delta}_{\delta\sigma}(q) \Gamma_{21\,\mu}^{\sigma} \right] \gamma_5 \}, \ \widetilde{\Delta}_{\delta\sigma}(p) = -P_{\delta\sigma}(p)\frac{\hat{p} + M}{p^2 - M^2}.
$$

As concerns the interference of the pure QED and HL diagrams, it can be written as

$$
2 M_h M_e^* = 2 Spur\left\{ (\hat{p}_2 - m)\gamma_5 \left[\Gamma_{22}^{\beta \nu} \Delta_{\beta \alpha}(p) \Gamma_{11}^{\alpha \mu} + \Gamma_{21}^{\beta \mu} \Delta_{\beta \alpha}(q) \Gamma_{12}^{\alpha \nu} \right] \gamma_5 \times \tag{15}
$$

$$
(\hat{p}_1 + m) \left[\gamma_\mu \frac{\hat{p} + m}{p^2 - m^2} \gamma_\nu + \gamma_\nu \frac{\hat{q} + m}{q^2 - m^2} \gamma_\mu \right] \}.
$$

In what follows we neglect the electron mass and introduce the dimensionless invariant variables

$$
s = \frac{2(p_1 \cdot p_2)}{M^2}, \ t = -\frac{2(p_1 \cdot k_1)}{M^2}, \ u = -\frac{2(p_1 \cdot k_2)}{M^2}, \ s + t + u = 0, \ \cos \theta = 1 + \frac{2t}{s}.\tag{16}
$$

These variables are very convenient especially in the limit $m \to 0$. In terms of these variables we obtain for HL diagrams contribution into matrix element squared

$$
|M_h|^2 = 2g_1^4 A_{11}(s,t) + \frac{1}{3}g_1^2 g_2^2 A_{12}(s,t) + \frac{1}{36}g_2^4 A_{22}(s,t),
$$
\n(17)

where

$$
A_{11}(s,t) = -2[3+2s^2+t^2+s(5+t)] - (1+3s+3s^2)\left[\frac{1}{(1-t)^2} + \frac{1}{(1+s+t)^2}\right] +
$$

$$
\frac{8+22s+21s^2+10s^3+s^4}{2+s}\left(\frac{1}{1-t} + \frac{1}{1+s+t}\right),
$$

$$
A_{12}(s,t) = 4s^3+3s^2(11+6t)+12(2+t)^2+6s(11+2t+3t^2)+3(1+2s)^2\left[\frac{1}{(1-t)^2} + \frac{1}{(1+s+t)^2}\right]
$$

$$
-\frac{30+114s+126s^2+46s^3+s^4}{2+s}\left(\frac{1}{1-t} + \frac{1}{1+s+t}\right),
$$

$$
A_{22}(s,t) = s^3(23-5t) - s^2(74+44t+21t^2) - 4s(21+9t+11t^2+8t^3) - 4(14+9t^2+4t^4)
$$

$$
-(5+6s+6s^2)\left[\frac{1}{(1-t)^2} + \frac{1}{(1+s+t)^2}\right]
$$

$$
+\frac{66+152s+175s^2+52s^3-18s^4}{2+s}\left(\frac{1}{1-t} + \frac{1}{1+s+t}\right).
$$

The QED-HL interference contributes as follows

$$
2\operatorname{Re}[M_h M_e^*] = 8\,g_1^2 \left[\frac{(1+s)^2}{1-t} + \frac{(1+s)^2}{1+s+t} - 2 - 3s \right] + \tag{18}
$$

$$
\frac{2 g_2^2}{3} \left[8 + 14s + s^2 + 8st + 8t^2 - (4 + 9s + 3s^2) \left(\frac{1}{1 - t} + \frac{1}{1 + s + t} \right) \right].
$$

To calculate the angular distribution in high energy limit one can use

$$
A_{11}(s,\theta) = \frac{4(8+22s+21s^2+10s^3+s^4)}{(2+s)^2-s^2\cos^2\theta} - 6 - 10s - \frac{s^2(7+\cos^2\theta)}{2}
$$

$$
-4(1+3s+3s^2)\left[\frac{1}{(2+s-s\cos\theta)^2} + \frac{1}{(2+s+s\cos\theta)^2}\right],
$$

$$
A_{12}(s,\theta) = 24 + 66s + 30s^2 - \frac{s^3}{2} + 3s^2\cos^2\theta\left(1+\frac{3s}{2}\right)
$$

$$
+12(1+2s)^2 \Big[\frac{1}{(2+s-s\cos\theta)^2} + \frac{1}{(2+s+s\cos\theta)^2} \Big] - \frac{4[30+114s+126s^2+46s^3+s^4]}{(2+s)^2-s^2\cos^2\theta},
$$

$$
A_{22}(s,\theta) = -56-84s-65s^2+34s^3+\frac{s^4}{4}-\left(9+11s-\frac{3s^2}{4}\right)s^2\cos^2\theta - s^4\cos^4\theta +
$$

$$
\frac{4(66+152s+175s^2+52s^3-18s^4)}{(2+s)^2-s^2\cos^2\theta} - (5+6s+6s^2) \Big[\frac{4}{(2+s-s\cos\theta)^2} + \frac{4}{(2+s+s\cos\theta)^2} \Big].
$$

Regards the QED-HL interference we have

3

$$
2\operatorname{Re}[M_{3/2}^+M^{e*}] = 8e^2 g_1^2 \Big[-2 - 3s + \frac{4(1+s)^2(2+s))}{(2+s)^2 - s^2 \cos^2 \theta} \Big] +
$$
\n
$$
\frac{2e^2 g_2^2}{3} \Big[8 + 14s + s^2 - 2s^2 \sin^2 \theta - \frac{4(4+9s+3s^2)(2+s)}{(2+s)^2 - s^2 \cos^2 \theta} \Big].
$$
\n(19)

 $(2 + s)^2 - s^2 \cos^2 \theta$

Note that in unpolarized case neither $|M_h|^2$ nor $Re[M_h M_e^*]$ contains product of the odd powers g_1 and g_2 , so they do not depend on relative sign of these constants. With accounting for all the contribution, we can write down the the differential over the photon solid angle cross section in unpolarized case as

$$
\frac{d\sigma_{un}}{d\,\Omega} = \frac{\alpha^2}{16M^2} \left[\frac{16(1+\cos^2\theta)}{s(1-\cos^2\theta)} + \frac{2\Re\left[M_h M_e^*\right]}{s} + \frac{|M_h|^2}{s} \right].\tag{20}
$$

In Fig. 2 we plot the differential cross section [\(20\)](#page-6-0) (in units M^{-2}) at different values of constant g_1 : 1, 0.35, 0.1 and two angles $\theta = 45^{\circ}$, 90° as a function of the ratio $g = g_2/g_1$ and dimensionless invariant variable s.

III. POLARIZATION ASYMMETRIES

To estimate different polarization asymmetries one has to compute corresponding polarization dependent parts of the cross section, using the electron and positron projections as given by Eq.[\(3\)](#page-2-0) and expressions for different polarization 4-vectors defined by relations [\(4\)](#page-2-1).

A. Case of longitudinal polarization

The direct calculation in the case of longitudinal polarization of the electron and positron along beam directions results

$$
\left|M_h^{+ll}\right|^2 = 2g_1^4 L_{11}(s,t) + \frac{1}{3}L_{12}(s,t) + \frac{1}{36}L_{22}(s,t),
$$
\n
$$
L_{11}(s,t) = 2[3 + 2s^2 + t^2 + s(5+t)] + (1 + 3s + 3s^2 + 2s^3) \left[\frac{1}{(1-t)^2} + \frac{1}{(1+s+t)^2}\right]
$$
\n(21)

 $(1 + s + t)^2$

FIG. 2. Unpolarized differential cross section defined by Eq. [\(20\)](#page-6-0) (in units M^{-2}) at two values of the photon angle: $\theta = 45^{\circ}$, the upper row; $\theta = 90^{\circ}$, the lower row; and three values of constants g1: 1−left panel; 0.35−middle panel; 0.1− right panel. g is the ratio g_2/g_1 .

$$
-\frac{8+22s+21s^2+6s^3+s^4}{2+s}\left(\frac{1}{1-t}+\frac{1}{1+s+t}\right),
$$

\n
$$
L_{12}(s,t) = 2s^3+3s^2(5+2t)-12(2+t^2)-6s(3+2t-t^2)-3(1+2s)\left[\frac{1}{(1-t)^2}+\frac{1}{(1+s+t)^2}\right]+
$$

\n
$$
\frac{30+54s+6s^2-14s^3+s^4}{2+s}\left(\frac{1}{1-t}+\frac{1}{1+s+t}\right),
$$

\n
$$
L_{22}(s,t) = s^3(47+5t)+s^2(178+44t+21t^2)+4s(43+9t+11t^2+8t^3)+4(14+9t^2+4t^4)+
$$

\n
$$
(5+28s+54s^2+36s^3)\left[\frac{1}{(1-t)^2}+\frac{1}{(1+s+t)^2}\right]
$$

\n
$$
-\frac{66+284s+463s^2+320s^3+78s^4}{2+s}\left(\frac{1}{1-t}+\frac{1}{1+s+t}\right).
$$

For the corresponding contribution into QED HL interference we have

$$
2\operatorname{Re}\left[M_h M_e^*\right]^{ll} = -8\,g_1^2 \frac{\left[s^3 + 2t^2 + s^2(1+3t) + st(2+3t)\right]}{(1-t)(1+s+t)} +
$$
\n
$$
2\,g_2^2 \frac{8t^4 + s^3(2+t) + 3s^2t(2+3t) + 2st^2(3+8t)}{3(1-t)(1+s+t)}.
$$
\n(22)

As one can see, the odd powers of constant g_1 and g_2 are also absent as in the unpolarized case.

To study the (s, θ) distribution one can use for contribution into $\left|M_h^{+ll}\right|$ 2

$$
L_{11}(s,\theta) = 6 + 10s + 4s^{2} - \frac{s^{2}}{2}\sin^{2}\theta - \frac{4(8 + 22s + 21s^{2} + 6s^{3} + s^{4})}{(2 + s)^{2} - s^{2}\cos^{2}\theta} +
$$

$$
4(1 + 3s + 3s^{2} + 2s^{3})\Big[\frac{1}{(2 + s - s\cos\theta)^{2}} + \frac{1}{(2 + s - s\cos\theta)^{2}}\Big],
$$

$$
L_{12}(s,\theta) = -24 - 18s + 15s^{2} + 23s^{3} + 3s^{2}\Big(1 - \frac{s}{2}\Big)\sin^{2}\theta + \frac{4(30 + 54s + 6s^{2} - 14s^{3} + s^{4})}{(2 + s)^{2} - s^{2}\cos^{2}\theta} -
$$

$$
-12(1 + 2s)\Big[\frac{1}{(2 + s - s\cos\theta)^{2}} + \frac{1}{(2 + s - s\cos\theta)^{2}}\Big],
$$

$$
L_{22}(s,\theta) = 56 + 172s + 178s^{2} + 47s^{3} + \Big(s^{2}\sin^{2}\theta - 9 - 11s - \frac{5s^{2}}{4}\Big)s^{2}\sin^{2}\theta +
$$

$$
4(5 + 28s + 54s^{2} + 36s^{3})\Big[\frac{1}{(2 + s - s\cos\theta)^{2}} + \frac{1}{(2 + s - s\cos\theta)^{2}}\Big]
$$

$$
-\frac{4(66 + 284s + 463s^{2} + 320s^{3} + 78s^{4})}{(2 + s)^{2} - s^{2}\cos^{2}\theta}.
$$

The corresponding result for the QED-HL interference reads

$$
2\operatorname{Re}[M_h M_e^*]^{ll} = -8\,g_1^2 s^2 \frac{4(1+s) - (2+3s)\sin^2\theta}{(2+s)^2 - s^2\cos^2\theta} + \frac{2}{3}\,g_2^2 s^3 \frac{8 + (2s\sin^2\theta - 6 - s)\sin^2\theta}{(2+s)^2 - s^2\cos^2\theta}.\tag{23}
$$

Asymmetry A_{ll} , in the case of longitudinally polarized electron and positron along their beam directions, is plotted In Fig. 3 at the same values of the angle θ and constant g_1 as in Fig. 2.

B. Transverse polarization of beams in the reaction plane

As one can see from [\(4\)](#page-2-1), polarization 4-vector S_1^t in the rest frame of the electron has only space component which (for $\lambda_1 = 1$) is $S_1 = (0, \mathbf{n}_{1t})$ and in this system $\mathbf{n}_{1t} \cdot \mathbf{p}_2 = 0$. Transition to c.m.s. does not change S_1^t , and because the same is valid for S_2 (with the change $\mathbf{n}_{1t} \rightleftarrows \mathbf{n}_{2t}, \ \mathbf{p}_2 \rightleftarrows \mathbf{p}_1$, it is possible $\mathbf{n}_{1t} = \pm \mathbf{n}_{2t}$ only. The same is valid for the normal 4-vectors S_1^n and S_2^n .

The $|M_h^{tt}|^2$, when both, the electron and positron are transversely polarized in the reaction plane along the same direction can be written as

$$
\left|M_{3/2}^{+tt}\right|^2 = e^4 \left[2 g_1^4 T_{11}(s,t) + \frac{1}{3} g_1^2 g_2^2 T_{12}(s,t) + \frac{1}{36} g_2^4 T_{22}(s,t)\right]
$$
\n
$$
T_{11}(s,t) = \frac{2t^2(s+t)^2}{(1-t)(1+s+t)},
$$
\n(24)

FIG. 3. Asymmetry A_{ll} as defined by Eq. [\(5\)](#page-2-3) at the same conditions as in Fig. 2 for unpolarized cross section.

$$
T_{12}(s,t) = \frac{(s+t)[s^3 - 4st^3 + t^4 - 5s^2t(1+t)]}{(1+s+t)^2} - \frac{t[8st^3 + t^4 + 6s^3(1+t) + s^2t(5+13t)]}{(1-t)^2} + \frac{s[s^3 + 9t^4 + 2st^2(1+9t) + s^2t(2+9t)]}{(1-t)(1+s+t)},
$$

$$
T_{22}(s,t) = \frac{3(s+t)^2[s^2t(t-2) + st^2(t-3) + t^4 + s^3(1+3t)]}{(1+s+t)^2} + \frac{3t^2[t^4 + 6s^3(1+t) + st^2(3+5t) + 2s^2t(5t+4)]}{(1-t)^2} - \frac{2t(s+t)[s^3(6-12t) + 5t^4 - s^2t(9+7t) + st^2(10t-9)]}{(1-t)(1+s+t)}.
$$

The QED-HL interference in this case reads

$$
2\operatorname{Re}\left[M_h M_e^*\right] = \frac{2t(s+t)}{3(1-t)(1+s+t)}[-12g_1^2(2+s)+g_2^2[3s^2t+8t^2+s(2+8t)].\tag{25}
$$

The corresponding expressions for angular distribution read

$$
T_{11}(s,\theta) = \frac{s^4 \sin^4 \theta}{2[(2+s)^2 - s^2 \cos^2 \theta]},
$$

\n
$$
T_{12}(s,\theta) = \frac{s^4 (1 + \cos \theta)[56 - 11s + 4(3s - 10) \cos \theta + s(10 \cos^2 \theta - 12 \cos^3 \theta + \cos^4 \theta)]}{8(2+s+s \cos \theta)^2}
$$

\n
$$
+ \frac{s^4 [8 + 9s + (8 - 18s) \cos^2 \theta + 9s \cos^4 \theta]}{4[(2+s)^2 - s^2 \cos^2 \theta]}
$$

FIG. 4. The same as in Fig. 3 but for asymmetry A_{tt} .

$$
+\frac{s^{4}(1-\cos\theta)[56-11s+(40-12s)\cos\theta+s(10\cos^{2}\theta+12\cos^{3}\theta+\cos^{4}\theta)]}{8(2+s-s\cos\theta)^{2}},
$$

\n
$$
T_{22}(s,\theta)=\frac{s^{5}\sin^{2}\theta[132+53s-36\cos^{2}\theta-58\cos^{2}\theta+5s\cos^{4}\theta]}{8[(2+s)^{2}-s^{2}\cos^{2}\theta]}
$$

\n
$$
+\frac{3s^{5}(1+\cos\theta)^{2}[44-17s+(6s-40)\cos\theta+4(3+4s)\cos^{2}\theta-6s\cos^{3}\theta+s\cos^{4}\theta]}{16(2+s+s\cos\theta)^{2}}
$$

\n
$$
+\frac{3s^{5}(1-\cos\theta)^{2}[44-17s+40\cos\theta+12\cos^{2}\theta+s(-6\cos\theta+16\cos^{2}\theta+6\cos^{3}\theta+\cos^{6})]}{16(2+s-s\cos\theta)^{2}}
$$

The interference with the QED amplitude is

$$
2\operatorname{Re}\left[M_h M_e^*\right]^{tt} = \frac{2s^2 \sin^2 \theta}{3[(2+s)^2 - s^2 \cos^2 \theta]} \left[12\,g_1^2(2+s) - g_2^2 s(2+s+2s\cos^2 \theta)\right].\tag{26}
$$

.

Asymmetry caused by the transverse polarizations of beams along the same direction is shown in Fig. 4. It changes sign if directions of polarizations are opposite.

C. longitudinal-transverse correlation

If the positron beam is polarized longitudinally and the electron beam − transversally (or vise versa), the corresponding spin-dependent contribution is absent in the Born approximation. But contribution caused by the intermediate $3/2$ -spin heavy lepton is nonzero in this case and

has a form

$$
|M_h^{+lt}|^2 = (-st)\sqrt{\frac{s+t}{-st}} \Big(2g_1^3 g_2 P_{31}^{lt}(s,t) + g_1^2 g_2^2 P_{22}^{lt}(s,t) + \frac{1}{3}g_1 g_2^3 P_{13}^{lt}(s,t) + \frac{1}{3}g_2^4 P_{04}^{lt}(s,t)\Big), (27)
$$

\n
$$
P_{31}^{lt}(s,t) = (1+2s+2s^2) \Big[\frac{1}{(1+s+t)^2} - \frac{1}{(1-t)^2}\Big] - (4+5s+s^2) \Big[\frac{1}{1+s+t} - \frac{1}{1-t}\Big] - 2(s+2t),
$$

\n
$$
P_{22}^{lt}(s,t) = -(1+2s+2s^2) \Big[\frac{1}{(1+s+t)^2} - \frac{1}{(1-t)^2}\Big] + (2+3s+s^2) \Big[\frac{1}{1+s+t} - \frac{1}{1-t}\Big],
$$

\n
$$
P_{13}^{lt}(s,t) = (2-s)(s+2t) - \Big[\frac{1}{(1+s+t)^2} - \frac{1}{(1-t)^2}\Big] + 4\Big[\frac{1}{1+s+t} - \frac{1}{1-t}\Big],
$$

\n
$$
P_{04}^{lt}(s,t) = -(2+s)(s+2t) + (1+3s+3s^2) \Big[\frac{1}{(1+s+t)^2} - \frac{1}{(1-t)^2}\Big] + (4+8s+5s^2) \Big[\frac{1}{1+s+t} - \frac{1}{1-t}\Big].
$$

\nAs one can see term with q_1^4 is absent and terms appear with the odd powers of q_1 and q_2 . If

As one can see, term with g_1^4 is absent and terms appear with the odd powers of g_1 and g_2 . If contrary, the electron beam is polarized longitudinally and the positron beam transversally, we have

$$
P_{31}^{tl}(s,t) = P_{31}^{lt}(s,t), \ P_{13}^{tl}(s,t) = P_{13}^{lt}(s,t), \ P_{22}^{tl}(s,t) = -P_{22}^{lt}(s,t), \ P_{04}^{tl}(s,t) = -P_{04}^{lt}(s,t)
$$

(terms with even powers g_1 and $_2$ change sign).

The QED-HL interference is also nonzero and reads

$$
2Re\left[M_h M_e^*\right]^{lt} = 2e^4s^2\sqrt{\frac{-t(s+t)}{s}}\left[2g_1g_2\left(\frac{1}{1+s+t} - \frac{1}{1-t}\right) + g_2^2\left(\frac{s+t}{1+s+t} + \frac{s+t}{1-t}\right)\right].\tag{28}
$$

In the case when the positron beam is polarized transversally and the electron one longitudinally, term with $g_1 g_2$ change sign.

Taking into account that

$$
(-st)\sqrt{\frac{s+t}{-st}} = \frac{s^{3/2}}{2}\sin\theta,
$$

one can rewrite corresponding the (s, θ) −distribution

$$
P_{31}^{lt}(s, \theta) = 2s \cos \theta \Big[-1 + \frac{2(4+5s+s^2)}{(2+s)^2 - s^2 \cos^2 \theta} - \frac{8(2+s)(1+2s+2s^2)}{[(2+s)^2 - s^2 \cos^2 \theta]^2} \Big],
$$

\n
$$
P_{22}^{lt}(s, \theta) = 4s \cos \theta \Big[-\frac{2+3s+s^2}{(2+s)^2 - s^2 \cos^2 \theta} + \frac{4(2+s)(1+2s+2s^2)}{[(2+s)^2 - s^2 \cos^2 \theta]^2} \Big],
$$

\n
$$
P_{13}^{lt}(s, \theta) = s \cos \theta \Big[2 - s - \frac{16}{(2+s)^2 - s^2 \cos^2 \theta} + \frac{16(2+s)}{[(2+s)^2 - s^2 \cos^2 \theta]^2} \Big],
$$

\n
$$
P_{04}^{lt}(s, \theta) = s \cos \theta \Big[-2 - s - \frac{4(4+8s+5s^2)}{(2+s)^2 - s^2 \cos^2 \theta} - \frac{16(2+s)(1+3s+3s^2)}{[(2+s)^2 - s^2 \cos^2 \theta]^2} \Big].
$$

The QED-HL interference for this distribution is

$$
2\operatorname{Re}\left[M_{3/2}^{+}M^{e*}\right]^{lt} = 4\operatorname{s}^{5/2}\sin\theta\cos\theta\frac{-2g_{1}g_{2}s + g_{2}^{2}(2+s)}{(2+s)^{2} - s^{2}\cos^{2}\theta}.
$$
\n(29)

It is clear that corresponding asymmetries go to zero if θ go to 90^o. They are shown in Figs. 5, 6.

FIG. 5. Asymmetry A_{lt} at photon angle $\theta = 45^{\circ}$ (the upper row); $\theta = 60^{\circ}$ (lower row) and the values of constants g_1 as in Fig. 2. It arises due to intermediate $3/2$ -spin heavy lepton and goes to zero at $\theta \to \pi/2.$

FIG. 6. The same as in Fig. 5 but for asymmetry A_{tl} .

D. normal polarizations

As it noted above, at $\eta_1 = \eta_2 = 1$, the 4-vectors of normal polarization are $S_1^n = (0, n_{1n})$ and $S_2^n = (0, n_{2n})$ with $n_{1n} = \pm n_{2n}$. If both beams are polarized perpendicularly to the reaction plane in the same direction we have

$$
|M_h|^2 = 2g_1^4 N_{11}(s,t) + \frac{1}{3}g_1^2 g_2^2 N_{12}(s,t) + \frac{1}{36}N_{22}(s,t),
$$
\n(30)
\n
$$
N_{11}(s,t) = \frac{2t^2(s+t)^2}{(1-t)(1+s+t)},
$$
\n
$$
N_{12}(s,t) = \frac{[s^3 - 4st^3 + t^4 - 5s^2t(1+t)][s+t)}{(1+s+t)^2} - \frac{t[8st^3 + t^4 + 6s^3(1+t) + s^2t(5+13t)]}{(1-t)^2} + \frac{s[s^3 + 9t^4 + 2st^2(1+9t) + s^2t(2+9t)]}{(1-t)(1+s+t)},
$$
\n
$$
N_{22}(s,t) = \frac{3(s+t)^2[s^2t(t-2) + st^2(t-3) + t^4 + s^3(1+3t)]}{(1+s+t)^2} + \frac{3t^2[t^4 + 6s^3(1+t) + st^2(3+5t) + 2s^2t(4+5t)]}{(1-t)^2} - \frac{2t[s^3(6-12t) + 5t^4 - s^2t(9+7t) + st^2(10t-9)]}{(1-t)(1+s+t)}.
$$
\n(4.1)

The QED-HL interference in this case coincides with [\(25\)](#page-9-0).

$$
N_{11}(s,\theta) = -2(1+s) + \frac{s^2 \sin^2 \theta}{2} + \frac{8(1+s)}{(2+s)^2 - s^2 \cos^2 \theta},
$$

\n
$$
N_{12}(s,\theta) = -8 - 6s - 10s^2 - \frac{s^2}{2} + \frac{s^2(s-2)\cos^2 \theta}{2} + \frac{4(10+18s+22s^2+12s^3-s^4)}{(2+s)^2 - s^2 \cos^2 \theta} - \frac{8(1+2s+2s^2)[(2+s)^2 + s^2 \cos^2 \theta]}{[(2+s)^2 - s^2 \cos^2 \theta]^2},
$$

\n
$$
N_{22}(s,\theta) = 40 + 92s + 63s^2 - 34s^3 + \frac{s^4}{4} + \frac{s^2(28+52s-5s^2)\cos^2 \theta}{4} + s^4 \cos^4 \theta - \frac{4(46+144s+177s^2+52s^3-18s^4)}{(2+s)^2 - s^2 \cos^2 \theta} + \frac{24(1+2s+2s^2)((2+s)^2+s^2\cos^2 \theta)}{[(2+s)^2 - s^2 \cos^2 \theta]^2}.
$$

As well as longitudinal and transverse asymmetries the normal one does not contain the odd powers of g_1 and g_2 . It plotted in Fig. 7.

FIG. 7. the same asin Fig. 3 but for asymmetry A_{nn} .

IV. DISCUSSION AND CONCLUSION

Manifestation of the 3/2-spin heavy lepton h^{\pm} has been considered in large-angle $e^+e^-\rightarrow \gamma\,\gamma$ reaction at high energies. We suggest that lepton h^{\pm} can interact with QED sector of the SM and appears as a virtual intermediate state in Feynmann diagrams, changing the intermediate electrons in t and u channels of this reaction. Phenomenological Lagrangian of the $h^{\pm} e^{\pm} \gamma$ interaction chosen similarly to one describing decay of the Δ isobar into electron and photon. It contains two unknown real constants which are dimensionless by dividing by the parameter with the dimension of mass, and we take this parameter as the heavy lepton mass M.

At energies $\sqrt{s} \approx 100 \,\text{GeV}$ the born differential section [\(6\)](#page-3-1) is about 2 pb and decreases with increasing energy as s^{-1} . Electroweak radiative corrections (EWRC) to the total cross section at such energies consist of about few percent and the positive QED share partially compensated by a negative weak correction [\[22,](#page-16-1) [23](#page-16-2)]. Contribution due to 3/2-spin heavy lepton increases very quickly with the energy and therefore can, in principle, be competitive with EWRC at ILC and CLIC colliders.

Our finding is that at $g_1 = 1$ and $M = 100 \,\text{GeV}$ contribution due to HL and QED-HL interference dominates already at \sqrt{s} = 50 GeV and such situation seems implausible. At $g_1 = 0.35$ it is of the order born cross section what is unrealistic too, and if $g_1 = 0.1$ it can be on the level of EWRC. Very interesting is the possibility to probe effects due to heavy lepton by

measuring the longitudinal-transverse asymmetries A_{lt} and A_{tl} Figs. 5, 6 where the pure QED mechanism does not contribute.

Acknowledgment

The authors acknowledge support by the National Academy of Sciences of Ukraine via the program "Participation in the international projects in high energy and nuclear physics" (project no. 0121U111693).

- [1] Y. Chen, C.-H. Lin, M. Liu, A. S. Sakharov, J. Ulbricht, and J. Zhao, Physics 5, 752 (2023), 2210.11149.
- [2] F. D. Aaron et al. (H1), Phys. Lett. B 666, 131 (2008), 0805.4530.
- [3] V. M. Abazov et al. (D0), Phys. Rev. D 77, 091102 (2008), 0801.0877.
- [4] G. Aad et al. (ATLAS), New J. Phys. 15, 093011 (2013), 1308.1364.
- [5] V. Khachatryan et al. (CMS), JHEP 03, 125 (2016), 1511.01407.
- [6] A. Abada et al. (FCC), Eur. Phys. J. C 79, 474 (2019).
- [7] M. Dong et al. (CEPC Study Group) (2018), 1811.10545.
- [8] The International Linear Collider Technical Design Report - Volume 1: Executive Summary (2013), 1306.6327.
- [9] A Multi-TeV Linear Collider Based on CLIC Technology: CLIC Conceptual Design Report (2012).
- [10] W. Rarita and J. Schwinger, Phys. Rev. 60, 61 (1941), URL <https://link.aps.org/doi/10.1103/PhysRev.60.61>.
- [11] H. F. Jones and M. D. Scadron, Annals Phys. 81, 1 (1973).
- [12] V. Pascalutsa, M. Vanderhaeghen, and S. N. Yang, Phys. Rept. 437, 125 (2007), hep-ph/0609004.
- [13] H. Harari, Phys. Rept. **104**, 159 (1984).
- [14] J. Leite Lopes, J. A. Martins Simoes, and D. Spehler, Phys. Lett. B 94, 367 (1980).
- [15] N. Fleury, J. Leite Lopes, and D. Spehler, Lett. Nuovo Cim. 36, 401 (1983).
- [16] S. R. Choudhury, R. G. Ellis, and G. C. Joshi, Phys. Rev. D 31, 2390 (1985).
- [17] F. M. L. Almeida, Jr., J. H. Lopes, J. A. Martins Simoes, and A. J. Ramalho, Phys. Rev. D 53, 3555 (1996), hep-ph/9509364.
- [18] O. Cakir and A. Ozansoy, Phys. Rev. D 77, 035002 (2008), 0709.2134.
- [19] V. Pascalutsa, Phys. Rev. D 58, 096002 (1998), hep-ph/9802288.
- [20] V. Pascalutsa and R. Timmermans, Phys. Rev. C 60, 042201 (1999), nucl-th/9905065.
- [21] J. Kristiano, S. Clymton, and T. Mart, Phys. Rev. C 96, 052201 (2017), 1710.07930.
- [22] C. M. Carloni Calame, M. Chiesa, G. Montagna, O. Nicrosini, and F. Piccinini, Phys. Lett. B 798, 134976 (2019), 1906.08056.
- [23] S. Bondarenko, Y. Dydyshka, L. Kalinovskaya, A. Kampf, L. Rumyantsev, R. Sadykov, and V. Yermolchyk, Phys. Rev. D 107, 073003 (2023), 2211.11467.