JOINT MOMENTS OF DERIVATIVES OF CHARACTERISTIC POLYNOMIALS OF RANDOM SYMPLECTIC AND ORTHOGONAL MATRICES

JULIO C. ANDRADE AND CHRISTOPHER G. BEST

ABSTRACT. We investigate the joint moments of derivatives of characteristic polynomials over the unitary symplectic group Sp(2N) and the orthogonal ensembles SO(2N) and $O^{-}(2N)$. We prove asymptotic formulae for the joint moments of the n_1 -th and n_2 -th derivatives of the characteristic polynomials for all three matrix ensembles. Our results give two explicit formulae for each of the leading order coefficients, one in terms of determinants of hypergeometric functions and the other as combinatorial sums over partitions. We use our results to put forward conjectures on the joint moments of derivatives of L-functions with symplectic and orthogonal symmetry.

1. INTRODUCTION

Let $G(2N) \in \{Sp(2N), SO(2N), O^{-}(2N)\}$, where Sp(2N) is the the group of $2N \times 2N$ unitary symplectic matrices and SO(2N) and $O^{-}(2N)$ are the subsets of $2N \times 2N$ orthogonal matrices with determinant +1 and -1, respectively. Also, denote the characteristic polynomial of a matrix $A \in G(2N)$ by

$$\Lambda_A(s) = \det\left(I - As\right)$$

In this paper we consider the joint moments

$$\int_{G(2N)} \left(\Lambda_A^{(n_1)}(1) \right)^{k_1} \left(\Lambda_A^{(n_2)}(1) \right)^{k_2} dA \tag{1.1}$$

of the n_1 -th and n_2 -th derivatives of the characteristic polynomials, where n_1, n_2 are non-negative integers and dA denotes the Haar measure on the relevant matrix ensemble. Using techniques developed in [1, 9, 21], we obtain asymptotic formulae for (1.1) for each G(2N) and for all non-negative integers k_1, k_2 . Our main results give two explicit expressions for the leading order coefficients for each of the matrix ensembles under consideration and are detailed in section 2.

The problem we study here is part of a general problem to obtain exact formulae for the complex moments of the derivatives of characteristic polynomials. A key motivation is the link between random matrix theory and the study of families of *L*-functions and their value distribution in analytic number theory. Specifically, one can use formulae obtained for characteristic polynomials of the various matrix ensembles to predict formulae for the corresponding quantities for *L*-functions with the same symmetry type. The complex moments of the derivatives of characteristic polynomials and *L*-functions can then be used to infer information on the zeros of the derivative of characteristic polynomials and on the horizontal distribution of the zeros of the derivative of characteristic function, see, for example, [11, 23] and [25, 26], respectively.

Additionally on the number theory side, the order of vanishing of an L-function at the central point, which is controlled by the derivatives of the L-function, is widely believed to contain deep arithmetic and geometric information. The Birch and Swinnerton-Dyer Conjecture for example,

Date: December 11, 2023.

²⁰¹⁰ Mathematics Subject Classification. Primary 60B20; Secondary 11M06, 11M50.

Key words and phrases. Random matrix theory, joint moments, characteristic polynomials, random symplectic matrices, random orthogonal matrices, Riemann zeta function, L-functions.

famously states that the order of vanishing of an L-function attached to an elliptic curve over \mathbb{Q} is equal to the rank of the curve.

For the ensemble of random unitary matrices U(N), Conrey, Rubinstein and Snaith [9] proved that for integer $k \ge 1$,

$$\int_{U(N)} |\Lambda'_A(1)|^{2k} dA \sim c_k N^{k^2 + 2k}$$

where

$$c_{k} = (-1)^{k(k+1)/2} \sum_{h=0}^{k} {\binom{k}{h}} \left(\frac{d}{dx}\right)^{k+h} \left(e^{-x} x^{-k^{2}/2} \det_{k \times k} \left(I_{i+j-1}(2\sqrt{x})\right)\right),$$

with $I_n(x)$ denoting the modified Bessel function of the first kind. Here, and throughout the paper, the indices *i* and *j* of the matrix in the determinant range from 1 to *k*. Also proven in [9] is a similar asymptotic formula for the 2*k*-th moment of the derivative of an analogue of Hardy's Z-function. As an application, the authors use their results to make a conjecture for the moments of the derivative of the Riemann zeta function and of the Z-function. Forrester and Witte [13] have given alternate expressions for the leading order coefficients obtained in [9] in terms of solutions to Painlevé III differential equations.

Concerning the joint moments over the unitary ensemble, one is interested in the quantity

$$\int_{U(N)} |\Lambda_A^{(n_1)}(1)|^{2M} |\Lambda_A^{(n_2)}(1)|^{2M-2k} dA.$$
(1.2)

When $n_1 = 1$ and $n_2 = 0$, Hughes [17] was able to show that the limit of $(1.2)/N^{k^2+2M}$ as $N \to \infty$ exists when k and M are integers. This result was extended to all suitable, real k and M by Assiotis, Keating and Warren in [4]. In [5], Bailey et al. obtain an asymptotic formula for (1.2) when $k \ge M$ are both non-negative integers.

In the case of general n_1, n_2 , Barhoumi-Andréani [6] gave an asymptotic formula for (1.2) for integer k and M with $k \ge M$ and $k \ge 2$ where the leading order coefficient is given in the form of a certain (k-1)-fold real integral. Recently, Keating and Wei [21] have obtained asymptotic formulae for (1.2) and for the joint moments of the n_1 -th and n_2 -th of the analogue of Hardy's Z-function for all integers $k \ge M \ge 0$. They give two explicit expressions for the leading order coefficients, one in terms of derivatives of determinants involving the modified Bessel function and the other as combinatorial sums involving partitions. They also use their results to motivate conjectures for the joint moments of the n_1 -th derivatives of the Riemann zeta function and of the Zfunction. The conjectures made in [21] are shown to agree with the known results of [15, 16, 18]. In [22], Keating and Wei further explore the structure and properties of their leading order coefficients. They establish recursive relations that the coefficients satisfy and also build a connection to a solution of the σ -Painlevé III' equation.

Turning to the symplectic and orthogonal matrix ensembles, Altuğ et al. [1] considered the moments of the *m*-th derivative

$$M_k(G(2N),m) := \int_{G(2N)} \left(\Lambda_A^{(m)}(1)\right)^k dA.$$

Extending the results of [9] to these ensembles, they prove asymptotic formulae for $M_k(G(2N), m)$ as $N \to \infty$ for integer $k \ge 1$ when G(2N) = Sp(2N) or G(N) = SO(2N) and m = 2, and when $G(2N) = O^-(2N)$ with m = 3. One considers the second derivative rather than the first in the case of Sp(2N) and SO(2N) since $\Lambda'_A(1)$ can be expressed simply in terms of $\Lambda_A(1)$. Thus, the moments of the first derivative can be computed using the result of Keating and Snaith [20] on the moments of $\Lambda_A(1)$. If $A \in O^-(2N)$, then $\Lambda_A(1) = 0$ and $\Lambda''_A(1)$ has as simple expression in terms of $\Lambda'_A(1)$ Hence, in this case, it is the moments of $\Lambda''_A(1)$ that are of interest. The leading order coefficients obtained in [1] are given in terms of derivatives of determinants involving hypergeometric functions. These determinants are shown to satisfy a differential recurrence relation similar to a Toda lattice equation connected to τ -function theory in the study of Painlevé differential equations. An interesting question put forward in [1] is whether there is a differential equation in the symplectic and orthogonal cases which plays a part analogous to Painlevé III in the unitary setting. Gharakhloo and Witte [14] have made promising progress in this direction in their study of 2j - k and j - 2k bi-orthogonal polynomial systems on the unit circle.

The authors of [1] also use their results to make conjectures for the asymptotic behaviour of the moments of derivatives at the central point of L-functions with symplectic or orthogonal symmetry. After stating our results in section 2, we will extend these to give general conjectures for the joint moments of the derivatives of L-functions with these symmetry types.

1.1. Notation. Recall that an $N \times N$ matrix A is said to be unitary if $AA^* = I$, where A^* is the conjugate transpose of A. The unitary symplectic group Sp(2N) is the subgroup of $2N \times 2N$ unitary matrices A which satisfy $A^T \Omega A = \Omega$, where

$$\Omega = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix},$$

with I the $N \times N$ identity matrix. The special orthogonal group SO(2N) and $O^{-}(2N)$ are the subsets of orthogonal $2N \times 2N$ matrices with determinant +1 and -1, respectively. Each of these matrix ensembles is endowed with the normalised Haar measure dA.

We will set $i^2 = -1$ and the variable *i* will only be used as an index. For real numbers *x*, we use [x] to denote the greatest integer less than or equal to *x*. We write S_k for the set of permutations on $\{1, 2, \ldots, k\}$. The multinomial coefficient is defined as

$$\binom{n}{l_1,\ldots,l_k} = \frac{n!}{l_1!\cdots l_k!}$$

for integers n and l_1, \ldots, l_k with $l_1 + \cdots + l_k = n$. Also, for integer n, whenever we write $l_1 + \cdots + l_k = n$ or $l_1 + \cdots + l_k \leq n$ this means that $l_i \geq 0$ are taken to be integers.

For any $w = (w_1, \ldots, w_k) \in \mathbb{C}^k$, the Vandermonde determinant is denoted by

$$\Delta(w) := \det_{k \times k} (w_i^{j-1}) = \prod_{1 \le i < j \le k} (w_i - w_j),$$

and we write $w^2 = (w_i^2)_{1 \le i \le k}$. We will also make use of Vandermonde determinants of differential operators, written as

$$\Delta\left(\frac{d}{dx}\right) := \det_{k \times k} \left(\frac{d^{j-1}}{dx_i^{j-1}}\right) = \prod_{1 \le i < j \le k} \left(\frac{d}{dx_i} - \frac{d}{dx_j}\right).$$

Lastly, for $u \in \mathbb{C}$ and $m \in \mathbb{Z}$, we let

$$g_m(u) := \frac{1}{2\pi 1} \oint_{|w|=1} \frac{e^{w+u/w^2}}{w^{m+1}} dw$$

= $\frac{1}{\Gamma(m+1)} {}_0F_2\left(\frac{m}{2}+1, \frac{m+1}{2}; \frac{u}{4}\right).$ (1.3)

These hypergeometric functions will play the role that the modified Bessel function plays in the unitary case. For negative m, one should interpret the above expression as the limit.

2. Main results

We now state our main results. Our first two theorems give an asymptotic formula for the joint moments of derivatives of characteristic polynomials of matrices over Sp(2N).

Theorem 2.1. Let $0 \le n_1 \le n_2$ be integers and let k_1, k_2 be non-negative integers, not both 0. Set $k = k_1 + k_2$. Then, we have

$$\int_{Sp(2N)} \left(\Lambda_A^{(n_1)}(1) \right)^{k_1} \left(\Lambda_A^{(n_2)}(1) \right)^{k_2} dA = b_{k_1,k_2}^{Sp}(n_1,n_2) \cdot (2N)^{k(k+1)/2+k_1n_1+k_2n_2} \left(1 + O(N^{-1}) \right),$$

where

$$b_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) = \frac{(-1)^{k_{1}n_{1}+k_{2}n_{2}}}{2^{k(k+1)/2+k_{1}n_{1}+k_{2}n_{2}}} \sum_{u_{1}+\dots+u_{P}=k_{1}} \binom{k_{1}}{u_{1}\dots,u_{P}} \frac{(n_{1}!)^{u_{1}}}{\prod_{i=1}^{P}(\boldsymbol{a}_{i}!)^{u_{i}}(\prod_{j=1}^{[n_{1}/2]}j\Sigma_{i=1}^{P}u_{i}a_{i,j})} \\ \times \sum_{v_{1}+\dots+v_{Q}=k_{2}} \binom{k_{2}}{v_{1}\dots,v_{Q}} \frac{(n_{2}!)^{k_{2}}}{\prod_{i=1}^{Q}(\boldsymbol{b}_{i}!)^{v_{i}}(\prod_{j=1}^{[n_{2}/2]}j\Sigma_{i=1}^{Q}v_{i}b_{i,j})} \\ \times \sum_{\substack{\sum_{i=1}^{k}m_{s,i}=m_{s}\\s=2,\dots,n}} \left(\prod_{s=2}^{n}\binom{m_{s}}{m_{s,1},\dots,m_{s,k}}\right) \binom{d}{du}^{m_{1}} \det_{k\times k} \left(g_{2i-j+2\sum_{s=2}^{n}sm_{s,i}}(u)\right)\Big|_{u=0}.$$

Here, for i = 1, ..., P, we define $\mathbf{a}_i = (a_{i,0}, a_{i,1}, ..., a_{i,[n_1/2]})$ to be the P distinct tuples of integers satisfying

$$a_{i,j} \ge 0$$
 and $a_{i,0} + 2\sum_{j=1}^{[n_1/2]} ja_{i,j} = n_1.$

Similarly, the $\mathbf{b}_i = (b_{i,0}, b_{i,1}, \dots, b_{i,\lfloor n_2/2 \rfloor})$ for $i = 1, \dots, Q$ are defined to be the Q tuples of integers satisfying

$$b_{i,j} \ge 0$$
 and $b_{i,0} + 2\sum_{j=1}^{[n_2/2]} jb_{i,j} = n_2.$

Also, we define $\mathbf{a}_i! := \prod_{j=1}^{[n_1/2]} a_{i,j}!$ and $\mathbf{b}_i! := \prod_{j=1}^{[n_2/2]} b_{i,j}!$. Finally, $m_j := \sum_{i=1}^{P} u_i a_{i,j} + \sum_{i=1}^{Q} v_i b_{i,j}$ for $j = 1, \ldots, [n_1/2]$ and $m_j := \sum_{i=1}^{Q} v_i b_{i,j}$ for $j = [n_1/2] + 1, \ldots, [n_2/2]$.

Theorem 2.2. Let $0 \le n_1 \le n_2$ be integers and let k_1, k_2 be non-negative integers, not both 0. Set $k = k_1 + k_2$. Then, we have

$$\int_{Sp(2N)} \left(\Lambda_A^{(n_1)}(1) \right)^{k_1} \left(\Lambda_A^{(n_2)}(1) \right)^{k_2} dA = b_{k_1,k_2}^{Sp}(n_1,n_2) \cdot (2N)^{k(k+1)/2+k_1n_1+k_2n_2} \left(1 + O(N^{-1}) \right),$$

where

$$b_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) = \frac{(-1)^{k(k-1)/2+k_{1}n_{1}+k_{2}n_{2}}}{2^{k(k+1)/2+k_{1}n_{1}+k_{2}n_{2}}} (n_{1}!)^{k_{1}} (n_{2}!)^{k_{2}}$$

$$\times \sum_{\substack{2\sum_{j=1}^{k} l_{i,j} \le n_{1}}{2\sum_{j=1}^{k} m_{i,j} \le n_{2}}} \sum_{\substack{i=1,\dots,k_{2}}} \left(\prod_{i=1}^{k_{1}} \frac{1}{(n_{1}-2\sum_{j=1}^{k} l_{i,j})!} \right) \left(\prod_{i=1}^{k_{2}} \frac{1}{(n_{2}-2\sum_{j=1}^{k} m_{i,j})!} \right)$$

$$\times \prod_{j=1}^{k} \frac{1}{(2k+V_{j}-2j+1)!} \prod_{1 \le i < j \le k} (V_{j}-V_{i}-2j+2i).$$

Here $V_j := 2 \sum_{i=1}^{k_1} l_{i,j} + 2 \sum_{i=1}^{k_2} m_{i,j}$ for $j = 1, \dots, k$.

Our next two theorems give an asymptotic formula for the joint moments over SO(2N).

Theorem 2.3. With notation as in Theorem 2.1, we have

$$\int_{SO(2N)} \left(\Lambda_A^{(n_1)}(1) \right)^{k_1} \left(\Lambda_A^{(n_2)}(1) \right)^{k_2} dA = b_{k_1,k_2}^{SO}(n_1,n_2) \cdot (2N)^{k(k-1)/2 + k_1 n_1 + k_2 n_2} \left(1 + O(N^{-1}) \right),$$

where

$$b_{k_{1},k_{2}}^{SO}(n_{1},n_{2}) = \frac{1}{2^{k(k-3)/2+k_{1}n_{1}+k_{2}n_{2}}} \sum_{u_{1}+\dots+u_{P}=k_{1}} \binom{k_{1}}{u_{1}\dots,u_{P}} \frac{(n_{1}!)^{u_{1}}}{\prod_{i=1}^{P}(a_{i}!)^{u_{i}}(\prod_{j=1}^{[n_{1}/2]}j\sum_{i=1}^{P}u_{i}a_{i,j})} \\ \times \sum_{v_{1}+\dots+v_{Q}=k_{2}} \binom{k_{2}}{v_{1}\dots,v_{Q}} \frac{(n_{2}!)^{k_{2}}}{\prod_{i=1}^{Q}(b_{i}!)^{v_{i}}(\prod_{j=1}^{[n_{2}/2]}j\sum_{i=1}^{Q}v_{i}b_{i,j})} \\ \times \sum_{\substack{\sum_{i=1}^{k}m_{s,i}=m_{s}\\s=2,\dots,n}} \left(\prod_{s=2}^{n}\binom{m_{s}}{m_{s,1},\dots,m_{s,k}}\right) \binom{d}{du}^{m_{1}} \det_{k\times k} \left(g_{2i-j-1+2\sum_{s=2}^{n}sm_{s,i}}(u)\right)\Big|_{u=0}.$$

Theorem 2.4. With notation as in Theorem 2.2, we have

$$\int_{SO(2N)} \left(\Lambda_A^{(n_1)}(1) \right)^{k_1} \left(\Lambda_A^{(n_2)}(1) \right)^{k_2} dA = b_{k_1,k_2}^{SO}(n_1,n_2) \cdot (2N)^{k(k-1)/2 + k_1 n_1 + k_2 n_2} \left(1 + O(N^{-1}) \right),$$

where

$$b_{k_{1},k_{2}}^{SO}(n_{1},n_{2}) = \frac{(-1)^{k(k-1)/2}}{2^{k(k-3)/2+k_{1}n_{1}+k_{2}n_{2}}} (n_{1}!)^{k_{1}} (n_{2}!)^{k_{2}}$$

$$\times \sum_{\substack{2\sum_{j=1}^{k} l_{i,j} \le n_{1}}{2\sum_{j=1}^{k} m_{i,j} \le n_{2}}} \sum_{\substack{j=1\\i=1,\dots,k_{1}}} \left(\prod_{i=1}^{k} \frac{1}{(n_{1}-2\sum_{j=1}^{k} l_{i,j})!} \right) \left(\prod_{i=1}^{k_{2}} \frac{1}{(n_{2}-2\sum_{j=1}^{k} m_{i,j})!} \right)$$

$$\times \prod_{j=1}^{k} \frac{1}{(2k+V_{j}-2j)!} \prod_{1 \le i < j \le k} (V_{j}-V_{i}-2j+2i).$$

Our final theorem gives an asymptotic formula for the joint moments over $O^-(2N)$ with the leading order coefficient expressed in terms of $b_{k_1,k_2}^{Sp}(n_1,n_2)$.

Theorem 2.5. Let $1 \le n_1 \le n_2$ be integers and let k_1, k_2 be non-negative integers, not both 0. Set $k = k_1 + k_2$. Then, we have

$$\int_{O^{-}(2N)} \left(\Lambda_{A}^{(n_{1})}(1) \right)^{k_{1}} \left(\Lambda_{A}^{(n_{2})}(1) \right)^{k_{2}} dA = b_{k_{1},k_{2}}^{O^{-}}(n_{1},n_{2}) \cdot (2N)^{k(k+1)/2+k_{1}(n_{1}-1)+k_{2}(n_{2}-1)} \left(1 + O(N^{-1}) \right),$$
where

where

$$b_{k_1,k_2}^{O^-}(n_1,n_2) = (-1)^{k_1(n_1-1)+k_2(n_2-1)} 2^k n_1^{k_1} n_2^{k_2} b_{k_1,k_2}^{Sp}(n_1-1,n_2-1)$$

with $b_{k_1,k_2}^{Sp}(n_1,n_2)$ as defined in Theorems 2.1 and 2.2.

Our Theorems 2.1, 2.3 exhibit the same structure as the asymptotic formulae obtained in [1]. Namely, the leading order coefficients are expressed in terms of derivatives of determinants of the hypergeometric functions $g_m(u)$. As mentioned in the introduction, these determinants were shown to satisfy a differential recurrence relation [1, Theorem 1.5] which allows the leading order coefficients to be computed much more quickly as k_1, k_2 get large. However, similarly to the unitary case considered in [21], the formulae given for the leading order coefficients in Theorems 2.1 and 2.3 may not be computationally efficient when n_1, n_2 are large since one has to compute the tuples \mathbf{a}_i and \mathbf{b}_i . Aside from giving an alternate expression for the leading order coefficients, the advantage of Theorems 2.2 and 2.4 is that the formulae are more computationally effective when n_1, n_2 are large and k_1, k_2 are small.

Using the standard random matrix philosophy allows us to make conjectures based on our results for the joint moments of derivatives of L-functions with symmetry type Sp, SO or O^- in the sense of [19]. We give an example conjecture for the family of quadratic Dirichlet L-functions at s = 1/2below. This is an example of a family with symplectic symmetry so we use our results for Sp(2N)as a model.

Conjecture 2.6. Let $\mathcal{D}(X) = \{d \text{ a fundamental discriminant } : |d| < X\}$, and let $L(s, \chi_d)$ be the Dirichlet L-function attached to the quadratic character χ_d . Then, for $0 \le n_1 \le n_2$ and $k_1, k_2 \ge 0$ integers with k_1, k_2 not both 0, we have that as $X \to \infty$,

$$\frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_1} L^{(n_2)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_1} L^{(n_2)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_1} L^{(n_2)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_1} L^{(n_2)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_1} L^{(n_2)} (1/2, \chi_d)^{k_2} \sim a_k \cdot b_{k_1, k_2}^{Sp} (n_1, n_2) \cdot (\log X)^{k(k+1)/2 + k_1 n_1 + k_2 n_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} = \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} = \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} = \frac{1}{|\mathcal{D}(X)|} \sum_{d \in \mathcal{D}(X)} L^{(n_1)} (1/2, \chi_d)^{k_2} + \frac{1}{|\mathcal{D}(X)|} \sum_{d$$

where $k = k_1 + k_2$ and $b_{k_1,k_2}^{Sp}(n_1, n_2)$ is the random matrix theory coefficient defined in Theorems 2.1 and 2.2. Also,

$$a_k = \prod_{p \text{ prime}} \frac{(1 - 1/p)^{k(k+1)/2}}{1 + 1/p} \left(\frac{(1 - 1/\sqrt{p})^{-k} + (1 + 1/\sqrt{p})^{-k}}{2} + \frac{1}{p} \right),$$

is an arithmetic constant depending on the family of L-functions. We note that a_k is the same coefficient appearing in the conjectures for the moments of $L(1/2, \chi_d)$, see [8, 20].

To the best of our knowledge, there are no results on the moments of derivatives of these quadratic Dirichlet L-functions over number fields. However, the moments of derivatives of L-functions over function fields have been investigated. For this discussion, we let \mathbb{F}_q denote a finite field with qelements and let \mathcal{H}_{2g+1} be the subset of square-free, monic polynomials of degree 2g + 1 in the polynomial ring $\mathbb{F}_q[x]$. For each $D \in \mathcal{H}_{2g+1}$, we have a Dirichlet L-function $L(s, \chi_D)$ attached to the quadratic character χ_D with conductor $|D| := q^{2g+1}$. This family of L-functions also exhibits symplectic symmetry. Andrade and Rajagopal [3] and subsequently Andrade and Jung [2] studied the mean values of $L^{(n)}(1/2, \chi_D)$ and an asymptotic formula for the first moment of $L^{(n)}(1/2, \chi_D)$ is given in [2] which implies that for any positive integer n as $g \to \infty$,

$$\frac{1}{|\mathcal{H}_{2g+1}|} \sum_{D \in \mathcal{H}_{2g+1}} L^{(n)}(1/2, \chi_D) \sim \frac{(-1)^n}{2(n+1)} \mathcal{A}(1) \cdot (2g+1)^{n+1}.$$
(2.1)

Here, $\mathcal{A}(1)$ is given as an Euler product over the monic, irreducible polynomials in $\mathbb{F}_q[x]$ and is the same arithmetic factor that appears in the asymptotic formula for the first moment of $L(1/2, \chi_D)$. The following proposition shows that (2.1) is indeed the asymptotic formula predicted by our conjecture in this case.

Proposition 2.7. For $n \ge 1$ an integer, we have that

$$b_{0,1}^{Sp}(0,n) = \frac{(-1)^n}{2(n+1)}.$$

Also, for $n \ge 1$, we have

$$b_{0,1}^{SO}(0,n) = 1.$$

A mixed second moment involving a second derivative of these quadratic Dirichlet L-functions over function fields was considered by Djanković and Đokić [10]. In particular, Theorem 1.1 and Remark 1.2 in [10], along with Florea's asymptotic formula for the second moment of $L(1/2, \chi_D)$ in [12], imply that

$$\frac{1}{|\mathcal{H}_{2g+1}|} \sum_{D \in \mathcal{H}_{2g+1}} \frac{L(1/2, \chi_D) L''(1/2, \chi_D)}{\log^2 q} \sim \frac{(1-q^{-1})}{80} \mathcal{B}(1/q) \cdot (2g+1)^5.$$

Similarly to the previous asymptotic formula, $\mathcal{B}(1/q)$ is given as an Euler product over monic, irreducible polynomials and it agrees with the arithmetic factor appearing in the asymptotic formula for the second moment of $L(1/2, \chi_D)$. This result is therefore consistent with the prediction of our conjecture as we compute that $b_{1,1}^{Sp}(0,2) = 1/80$.

One can naturally use our Theorems 2.1-2.5 to make analogous conjectures for the joint moments of derivatives at the central point for any family of L-functions with symplectic or orthogonal symmetry.

3. Preliminaries

We begin with the following two lemmas concerning Vandermonde determinants of differential operators. The first is quoted from [1, Lemma 2.8] and follows from the definition.

Lemma 3.1. Let $f_1(x), \ldots, f_k(x)$ be k-1 times differentiable. Then

$$\Delta\left(\frac{d}{dx}\right)\prod_{i=1}^{k}f_i(x_i) = \det_{k \times k}\left(f_i^{(j-1)}(x_i)\right).$$

Lemma 3.2. Let $f_1(x, y), \ldots, f_k(x, y)$ be k - 1 times differentiable in x and y. Then

$$\Delta\left(\frac{d}{dx}\right)\Delta\left(\frac{d}{dy}\right)\prod_{i=1}^{k}f_{i}(x_{i},y_{i})\bigg|_{\substack{x_{1}=\cdots=x_{k}=X,\\y_{1}=\cdots=y_{k}=Y}}=\sum_{\mu\in S_{k}}\det_{k\times k}\left(\frac{d^{i+j-2}}{dX^{i-1}dY^{j-1}}f_{\mu(i)}(X,Y)\right).$$

In particular, when $f_1 = \cdots = f_k = f$, we have

$$\Delta\left(\frac{d}{dx}\right)\Delta\left(\frac{d}{dy}\right)\prod_{i=1}^{k}f(x_{i},y_{i})\bigg|_{\substack{x_{1}=\cdots=x_{k}=X,\\y_{1}=\cdots=y_{k}=Y}}=k!\det_{k\times k}\left(\frac{d^{i+j-2}}{dX^{i-1}dY^{j-1}}f(X,Y)\right).$$

Proof. The case when $f_1 = \cdots = f_k = f$ is the result of [1, Lemma 2.9] and the proof of the general case follows the same lines. By Lemma 3.1, we have

$$\Delta\left(\frac{d}{dx}\right)\prod_{i=1}^{k}f_{i}(x_{i}, y_{i}) = \det_{k \times k}\left(\frac{d^{j-1}}{dx_{i}^{j-1}}f_{i}(x_{i}, y_{i})\right) = \sum_{\mu \in S_{k}}\operatorname{sign}(\mu)\prod_{i=1}^{k}\frac{d^{\mu(i)-1}}{dx_{i}^{\mu(i)-1}}f_{i}(x_{i}, y_{i})$$

Then, by Lemma 3.1 again, we have that

$$\begin{split} \Delta\left(\frac{d}{dx}\right)\Delta\left(\frac{d}{dy}\right)\prod_{i=1}^{k}f_{i}(x_{i},y_{i})\bigg|_{\substack{x_{i}=X,\\y_{i}=Y}} &=\sum_{\mu\in S_{k}}\operatorname{sign}(\mu)\Delta\left(\frac{d}{dy}\right)\prod_{i=1}^{k}\frac{d^{\mu(i)-1}}{dx_{i}^{\mu(i)-1}}f_{i}(x_{i},y_{i})\bigg|_{\substack{x_{i}=X,\\y_{i}=Y}} \\ &=\sum_{\mu\in S_{k}}\operatorname{sign}(\mu)\det_{k\times k}\left(\frac{d^{\mu(i)+j-2}}{dx_{i}^{\mu(i)-1}dy_{i}^{j-1}}f_{i}(x_{i},y_{i})\right)\bigg|_{\substack{x_{i}=X,\\y_{i}=Y}} \\ &=\sum_{\mu\in S_{k}}\operatorname{sign}(\mu)\det_{k\times k}\left(\frac{d^{\mu(i)+j-2}}{dX^{\mu(i)-1}dY^{j-1}}f_{i}(X,Y)\right) \\ &=\sum_{\mu\in S_{k}}\det_{k\times k}\left(\frac{d^{i+j-2}}{dX^{i-1}dY^{j-1}}f_{\mu(i)}(X,Y)\right), \end{split}$$

where we have interchanged the rows of the matrix to obtain the final line.

We next express a certain contour integral in terms of the hypergeometric functions $g_m(u)$.

Lemma 3.3. Let $k \in \mathbb{Z}$ and let $n \geq 1$ be an integer. Then, for complex numbers u_1, \ldots, u_n , we have

$$\frac{1}{2\pi i} \oint_{|w|=1} \exp\left(w + \sum_{j=1}^{n} \frac{u_j}{w^{2j}}\right) \frac{dw}{w^{k+1}} = \sum_{m_2,\dots,m_n=0}^{\infty} \left(\prod_{j=2}^{n} \frac{u_j^{m_j}}{m_j!}\right) g_{k+2\sum_{j=2}^{n} jm_j}(u_1),$$

where $g_m(u)$ is the hypergeometric function defined in (1.3).

Proof. We compute the integral by determining the coefficient of w^k in the exponential factor of the integrand. So, let $a_n(k)$ be the coefficient of w^k in $\exp(w + \sum_{j=1}^n \frac{u_j}{w^{2j}})$. Then,

$$\exp\left(w + \sum_{j=1}^{n} \frac{u_j}{w^{2j}}\right) = \exp\left(\frac{u_n}{w^{2n}}\right) \exp\left(w + \sum_{j=1}^{n-1} \frac{u_j}{w^{2j}}\right)$$
$$= \left(\sum_{m=0}^{\infty} \frac{u_n^m}{m!} w^{-2nm}\right) \left(\sum_{m=-\infty}^{\infty} a_{n-1}(m) w^m\right).$$

From this it follows that

$$a_n(k) = \sum_{m_n=0}^{\infty} \frac{u_n^{m_n}}{m_n!} a_{n-1}(k+2nm_n)$$
$$= \sum_{m_2,\dots,m_n=0}^{\infty} \left(\prod_{j=2}^n \frac{u_j^{m_j}}{m_j!}\right) a_1(k+2\sum_{j=2}^n jm_j).$$

We then see that by definition, $a_1(k+2\sum_{j=2}^n jm_j) = g_{k+2\sum_{j=2}^n jm_j}(u_1)$ and hence

$$a_n(k) = \sum_{m_2,\dots,m_n=0}^{\infty} \left(\prod_{j=2}^n \frac{u_j^{m_j}}{m_j!} \right) g_{k+2\sum_{j=2}^n j m_j}(u_1),$$

as required.

The next lemma allows us to take higher order derivatives of determinants of functions.

Lemma 3.4 (Lemma 13 in [21]). Let $s \ge 0$, $k \ge 1$ be integers and $a_{i,j}(x)$ be s-th differentiable functions of x. Then

$$\left(\frac{d}{dx}\right)^{s} \det_{k \times k} \left(a_{i,j}(x)\right) = \sum_{l_1 + \dots + l_k = s} \left(s \atop l_1, \dots, l_k\right) \det_{k \times k} \left(a_{i,j}^{(l_i)}(x)\right),$$

where $a_{i,j}^{(l_i)}(x)$ means that we take the l_i -th derivative of $a_{i,j}(x)$.

Now, the shifted moments of the characteristic polynomials are defined as

$$I(G(2N); z_1, \dots, z_k) := \int_{G(2N)} \Lambda_A(z_1) \cdots \Lambda_A(z_k) \, dA.$$

These shifted moments have been computed by Conrey et al. [7] and can be expressed in the form of a multiple contour integral. We will use the following approximate versions of their formulae which follow easily from the results of [7] and the fact that $(1 - e^{-x})^{-1} = x^{-1} + O(1)$ for small x.

Lemma 3.5 (Corollary 2.4 in [1]). Let $\alpha_1, \ldots, \alpha_k$ be complex numbers such that $|\alpha_j| \ll 1/N$ for $j = 1, 2, \ldots, k$. Then

$$I(Sp(2N); e^{-\alpha_1}, \dots, e^{-\alpha_k}) = \frac{(-1)^{k(k-1)/2}}{(2\pi 1)^k k!} \oint \cdots \oint_{|w_i|=1} \frac{\Delta(w) \Delta(w^2) e^{N \sum_{i=1}^k (w_i - \alpha_i)}}{\prod_{1 \le i, j \le k} (w_i^2 - \alpha_j^2)} \prod_{i=1}^k dw_i \left(1 + O(N^{-1})\right).$$

Lemma 3.6 (Corollary 2.5 in [1]). Let $\alpha_1, \ldots, \alpha_k$ be complex numbers such that $|\alpha_j| \ll 1/N$ for $j = 1, 2, \ldots, k$. Then

$$I(SO(2N); e^{-\alpha_1}, \dots, e^{-\alpha_k}) = \frac{(-1)^{k(k-1)/2} 2^k}{(2\pi 1)^k k!} \oint \cdots \oint_{|w_i|=1} \frac{\Delta(w) \Delta(w^2) (\prod_{i=1}^k w_i) e^{N \sum_{i=1}^k (w_i + \alpha_i)}}{\prod_{1 \le i, j \le k} (w_i^2 - \alpha_j^2)} \prod_{i=1}^k dw_i \left(1 + O(N^{-1})\right).$$

Lemma 3.7 (Corollary 2.6 in [1]). Let $\alpha_1, \ldots, \alpha_k$ be complex numbers such that $|\alpha_j| \ll 1/N$ for $j = 1, 2, \ldots, k$. Then

$$I(O^{-}(2N); e^{-\alpha_{1}}, \dots, e^{-\alpha_{k}}) = \frac{(-1)^{k(k-1)/2} 2^{k}}{(2\pi 1)^{k} k!} \oint \cdots \oint_{|w_{i}|=1} \frac{\Delta(w) \Delta(w^{2}) (\prod_{i=1}^{k} \alpha_{i}) e^{N \sum_{i=1}^{k} (w_{i} + \alpha_{i})}}{\prod_{1 \le i, j \le k} (w_{i}^{2} - \alpha_{j}^{2})} \prod_{i=1}^{k} dw_{i} \left(1 + O(N^{-1})\right).$$

Below we give two expressions for the derivatives of these contour integral expressions for shifted moments with respect to the shifts α_j .

Lemma 3.8. Let $n \ge 0$ and $k \ge 1$ be integers. Then

$$\frac{d^n}{d\alpha^n} \left. \frac{e^{-N\alpha}}{\prod_{i=1}^k (w_i^2 - \alpha^2)} \right|_{\alpha=0} = \left(\prod_{i=1}^k \frac{1}{w_i^2} \right) \sum_{m=0}^n \binom{n}{m} (-N)^{n-m} m! \sum_{\substack{l_1 + \dots + l_k = m \\ l_j \text{ even}}} \prod_{i=1}^k \frac{1}{w_i^{l_i}} \sum_{m=0}^{k-1} \frac{1}{w_i^{l_i}}} \sum_{m=0}^{k-1} \frac{1}{w_i^{l_i}} \sum_{m=0}^{k-1} \frac{1}{w_i^{$$

Proof. This follows from the proof of [1, Lemma 2.7] where we have corrected a typo. Lemma 3.9. Let $n \ge 0$ and $k \ge 1$ be integers. Then

$$\frac{d^n}{d\alpha^n} \left. \frac{e^{-N\alpha}}{\prod_{i=1}^k (w_i^2 - \alpha^2)} \right|_{\alpha=0} = \left(\prod_{i=1}^k \frac{1}{w_i^2} \right) \sum_{\substack{m_1+2m_2+\dots+nm_n=n\\m_3=m_5=\dots=0}} \frac{n!}{m_1!\dots m_n!} (-N)^{m_1} \prod_{j=1}^{\lfloor n/2 \rfloor} \left(\frac{1}{j} \sum_{i=1}^k \frac{1}{w_i^{2j}} \right)^{m_{2j}}.$$

Proof. The proof is similar to that of [21, Lemma 9]. First, we have

$$\frac{d}{d\alpha} \frac{e^{-N\alpha}}{\prod_{i=1}^{k} (w_i^2 - \alpha^2)} = \frac{e^{-N\alpha}}{\prod_{i=1}^{k} (w_i^2 - \alpha^2)} f_1(\alpha),$$

where

$$f_1(\alpha) = -N + 2\alpha \sum_{i=1}^k \frac{1}{w_i^2 - \alpha^2}.$$

We can then write

$$\frac{d^{n}}{d\alpha^{n}} \frac{e^{-N\alpha}}{\prod_{i=1}^{k} (w_{i}^{2} - \alpha^{2})} = \frac{e^{-N\alpha}}{\prod_{i=1}^{k} (w_{i}^{2} - \alpha^{2})} f_{n}(\alpha),$$
(3.1)

where $f_n(\alpha)$ is defined recursively by

$$f_{n+1}(\alpha) = f_n(\alpha)f_1(\alpha) + f'_n(\alpha).$$

Now, let $g(\alpha)$ be a function such that $g'(\alpha) = f_1(\alpha)$. Then, we have that

$$\frac{d^n}{d\alpha^n}e^{g(\alpha)} = e^{g(\alpha)}f_n(\alpha).$$

But, by Faà di Bruno's formula, we also have that

$$\frac{d^{n}}{d\alpha^{n}}e^{g(\alpha)} = e^{g(\alpha)} \sum_{m_{1}+2m_{2}+\dots+nm_{n}=n} \frac{n!}{m_{1}!\dots m_{n}!} \prod_{j=1}^{n} \left(\frac{g^{(j)}(\alpha)}{j!}\right)^{m_{j}}$$
$$= e^{g(\alpha)} \sum_{m_{1}+2m_{2}+\dots+nm_{n}=n} \frac{n!}{m_{1}!\dots m_{n}!} \prod_{j=1}^{n} \left(\frac{f_{1}^{(j-1)}(\alpha)}{j!}\right)^{m_{j}}$$

Comparing the above two expressions for $(d/d\alpha)^n e^{g(\alpha)}$, we see that

$$f_n(\alpha) = \sum_{\substack{m_1+2m_2+\dots+nm_n=n\\ \text{for } i > 1}} \frac{n!}{m_1! \cdots m_n!} \prod_{j=1}^n \left(\frac{f_1^{(j-1)}(\alpha)}{j!}\right)^{m_j}.$$

One can check that for $j \ge 1$, we have

$$f_1^{(j)}(0) = \begin{cases} 0 & \text{if } j \text{ even,} \\ 2j! \sum_{i=1}^k w_i^{-(1+j)} & \text{if } j \text{ odd.} \end{cases}$$

Hence, we have that

$$f_n(0) = \sum_{\substack{m_1+2m_2+\dots+nm_n=n\\m_3=m_5=\dots=0}} \frac{n!}{m_1!\cdots m_n!} (-N)^{m_1} \prod_{j=1}^{[n/2]} \left(\frac{1}{j} \sum_{i=1}^k \frac{1}{w_i^{2j}}\right)^{m_{2j}}.$$

Evaluating (3.1) at $\alpha = 0$ using this expression for $f_n(0)$ yields the desired result.

In the next two propositions we will compute the main contour integrals that we need to evaluate. **Proposition 3.10.** Let $k \ge 1$ and $n \ge 1$ be integers. Also, let (m_1, \ldots, m_n) be a tuple of nonnegative integers. Then, we have

$$\frac{1}{(2\pi i)^{k}} \oint \cdots \oint_{|w_{i}|=1} \Delta(w) \Delta(w^{2}) e^{N \sum_{i=1}^{k} w_{i}} \prod_{j=1}^{n} \left(\sum_{i=1}^{k} \frac{1}{w_{i}^{2j}} \right)^{m_{j}} \prod_{i=1}^{k} \frac{dw_{i}}{w_{i}^{2k}} \\
= (-1)^{k(k-1)/2} k! N^{k(k+1)/2+2 \sum_{j=1}^{n} jm_{j}} \sum_{\substack{\sum_{i=1}^{k} m_{s,i} = m_{s} \\ s = 2, \dots, n}} \left(\prod_{s=2}^{n} \binom{m_{s}}{m_{s,1}, \dots, m_{s,k}} \right) \\
\times \left(\frac{d}{du} \right)^{m_{1}} \det_{k \times k} \left(g_{2i-j+2 \sum_{s=2}^{n} sm_{s,i}}(u) \right) \Big|_{u=0}.$$
(3.2)

Also, we have

$$\frac{1}{(2\pi 1)^{k}} \oint \cdots \oint_{|w_{i}|=1} \Delta(w) \Delta(w^{2}) e^{N \sum_{i=1}^{k} w_{i}} \prod_{j=1}^{n} \left(\sum_{i=1}^{k} \frac{1}{w_{i}^{2j}} \right)^{m_{j}} \prod_{i=1}^{k} \frac{dw_{i}}{w_{i}^{2k-1}} \\
= (-1)^{k(k-1)/2} k! N^{k(k-1)/2+2 \sum_{j=1}^{n} jm_{j}} \sum_{\substack{\sum_{i=1}^{k} m_{s,i} = m_{s} \\ s = 2, \dots, n}} \left(\prod_{s=2}^{n} \binom{m_{s}}{m_{s,1}, \dots, m_{s,k}} \right) \right) \\
\times \left(\frac{d}{du} \right)^{m_{1}} \det_{k \times k} \left(g_{2i-j-1+2 \sum_{s=2}^{n} sm_{s,i}}(u) \right) \Big|_{u=0}.$$
(3.3)

Proof. First, note that

$$\Delta(w^2) = \Delta\left(\frac{d}{dX}\right) \exp\left(\sum_{i=1}^k w_i^2 X_i\right) \bigg|_{X_i=0},$$

and

$$\Delta(w) e^{N \sum_{i=1}^{k} w_i} = \Delta\left(\frac{d}{dY}\right) \exp\left(\sum_{i=1}^{k} w_i Y_i\right) \bigg|_{Y_i = N}.$$

We may also write

$$\prod_{j=1}^{n} \left(\sum_{i=1}^{k} \frac{1}{w_i^{2j}} \right)^{m_j} = \prod_{j=1}^{n} \left(\frac{d}{dt_j} \right)^{m_j} \exp\left(\sum_{j=1}^{n} t_j \sum_{i=1}^{k} \frac{1}{w_i^{2j}} \right) \bigg|_{t_j=0}.$$

Then, we have that

$$\begin{split} \frac{1}{(2\pi i)^k} \oint \cdots \oint \Delta(w) \Delta(w^2) \, e^{N\sum_{i=1}^k w_i} \prod_{j=1}^n \left(\sum_{i=1}^k \frac{1}{w_i^{2j}}\right)^{m_j} \prod_{i=1}^k \frac{dw_i}{w_i^{2k}} \\ &= \Delta\left(\frac{d}{dX}\right) \Delta\left(\frac{d}{dY}\right) \prod_{j=1}^n \left(\frac{d}{dt_j}\right)^{m_j} \\ &\times \frac{1}{(2\pi i)^k} \oint \cdots \oint \exp\left(\sum_{i=1}^k \left(w_i^2 X_i + w_i Y_i + \sum_{j=1}^n \frac{t_j}{w_i^{2j}}\right)\right) \prod_{i=1}^k \frac{dw_i}{w_i^{2k}} \bigg|_{\substack{X_i = 0, \\ Y_i = N, \\ t_j = 0}} \\ &= \prod_{j=1}^n \left(\frac{d}{dt_j}\right)^{m_j} \Delta\left(\frac{d}{dX}\right) \Delta\left(\frac{d}{dY}\right) \prod_{i=1}^k \left(\frac{1}{2\pi i} \oint_{|w|=1} \exp\left(w^2 X_i + wY_i + \sum_{j=1}^n \frac{t_j}{w^{2j}}\right) \frac{dw}{w^{2k}}\right) \bigg|_{\substack{X_i = 0, \\ Y_i = N, \\ t_j = 0}} \\ &= \prod_{j=1}^n \left(\frac{d}{dt_j}\right)^{m_j} \, k! \det\left(\frac{d^{i+j-2}}{dX^{i-1}dY^{j-1}} \frac{1}{2\pi i} \oint_{|w|=1} \exp\left(w^2 X + wY + \sum_{l=1}^n \frac{t_l}{w^{2l}}\right) \frac{dw}{w^{2k}}\right) \bigg|_{\substack{X_i = 0, \\ Y_i = N, \\ t_j = 0}} , \end{split}$$

where the last line is by Lemma 3.2. Now,

$$\frac{d^{i+j-2}}{dX^{i-1}dY^{j-1}} \frac{1}{2\pi_{1}} \oint \exp\left(w^{2}X + wY + \sum_{l=1}^{n} \frac{t_{l}}{w^{2l}}\right) \frac{dw}{w^{2k}}\Big|_{\substack{X=0, \\ Y=N}}$$
$$= \frac{1}{2\pi_{1}} \oint \exp\left(wN + \sum_{l=1}^{n} \frac{t_{l}}{w^{2l}}\right) \frac{dw}{w^{2k-2i-j+3}}$$
$$= \frac{N^{2k-2i-j+2}}{2\pi_{1}} \oint \exp\left(w + \sum_{l=1}^{n} \frac{N^{2l}t_{l}}{w^{2l}}\right) \frac{dw}{w^{2k-2i-j+3}}.$$

Therefore, we have

$$\begin{split} & \frac{1}{(2\pi i)^k} \oint \cdots \oint \Delta(w) \Delta(w^2) \, e^{N\sum_{i=1}^k w_i} \prod_{j=1}^n \left(\sum_{i=1}^k \frac{1}{w_i^{2j}}\right)^{m_j} \prod_{i=1}^k \frac{dw_i}{w_i^{2k}} \\ &= k! \prod_{j=1}^n \left(\frac{d}{dt_j}\right)^{m_j} \det\left(\frac{N^{2k-2i-j+2}}{2\pi i} \oint \exp\left(w + \sum_{l=1}^n \frac{N^{2l}t_l}{w^{2l}}\right) \frac{dw}{w^{2k-2i-j+3}}\right) \Big|_{t_j=0} \\ &= k! N^{k(k+1)/2} \prod_{j=1}^n \left(\frac{d}{dt_j}\right)^{m_j} \det\left(\frac{1}{2\pi i} \oint \exp\left(w + \sum_{l=1}^n \frac{N^{2l}t_l}{w^{2l}}\right) \frac{dw}{w^{2k-2i-j+3}}\right) \Big|_{t_j=0} \\ &= k! N^{k(k+1)/2} + \sum_{j=1}^n 2^{jm_j} \prod_{j=1}^n \left(\frac{d}{du_j}\right)^{m_j} \det\left(\frac{1}{2\pi i} \oint \exp\left(w + \sum_{l=1}^n \frac{u_l}{w^{2l}}\right) \frac{dw}{w^{2k-2i-j+3}}\right) \Big|_{u_j=0} \\ &= (-1)^{k(k-1)/2} k! N^{k(k+1)/2 + \sum_{j=1}^n 2^{jm_j}} \prod_{j=1}^n \left(\frac{d}{du_j}\right)^{m_j} \det\left(\frac{1}{du_j}\right)^{m_j} \det\left(\frac{1}{2\pi i} \oint \exp\left(w + \sum_{l=1}^n \frac{u_l}{w^{2l}}\right) \frac{dw}{w^{2k-2i-j+3}}\right) \Big|_{u_j=0} \\ &= (-1)^{k(k-1)/2} k! N^{k(k+1)/2 + \sum_{j=1}^n 2^{jm_j}} \prod_{j=1}^n \left(\frac{d}{du_j}\right)^{m_j} \det\left(\frac{1}{2\pi i} \oint \exp\left(w + \sum_{l=1}^n \frac{u_l}{w^{2l}}\right) \frac{dw}{w^{2i-j+1}}\right) \Big|_{u_j=0} , \end{split}$$

where we have used the fact that $\det_{k\times k}(N^{-2i-j}a_{i,j}) = N^{-3k(k+1)/2} \det_{k\times k}(a_{i,j})$. Also, the fourth line follows from the change of variables $u_j = N^{2j}t_j$ and in the last line we have interchanged the rows of the matrix.

Next, by Lemma 3.3, the contour integral appearing in the determinant is

$$\frac{1}{2\pi i} \oint \exp\left(w + \sum_{l=1}^{n} \frac{u_l}{w^{2l}}\right) \frac{dw}{w^{2i-j+1}} = \sum_{l_2,\dots,l_n=0}^{\infty} \left(\prod_{s=2}^{n} \frac{u_s^{l_s}}{l_s!}\right) g_{2i-j+2\sum_{s=2}^{n} sl_s}(u_1).$$

We use Lemma 3.4 to carry out the differentiation of the determinant with respect to u_2, \ldots, u_n . This gives us

$$\begin{split} \prod_{j=1}^{n} \left(\frac{d}{du_{j}}\right)^{m_{j}} \det_{k \times k} \left(\frac{1}{2\pi i} \oint \exp\left(w + \sum_{l=1}^{n} \frac{u_{l}}{w^{2l}}\right) \frac{dw}{w^{2i-j+1}}\right) \bigg|_{u_{j}=0} \\ &= \sum_{\substack{\sum_{i=1}^{k} m_{s,i}=m_{s} \\ s=2,...,n}} \left(\prod_{s=2}^{n} \binom{m_{s}}{m_{s,1},...,m_{s,k}}\right) \right) \\ &\times \left(\frac{d}{du_{1}}\right)^{m_{1}} \det_{k \times k} \left(\prod_{s=2}^{n} \left(\frac{d}{du_{s}}\right)^{m_{s,i}} \sum_{l_{2},...,l_{n}=0}^{\infty} \left(\prod_{s=2}^{n} \frac{u_{s}^{l_{s}}}{l_{s}!}\right) g_{2i-j+2\sum_{s=2}^{n} sl_{s}}(u_{1}) \right) \bigg|_{u_{j}=0} \\ &= \sum_{\substack{\sum_{i=1}^{k} m_{s,i}=m_{s} \\ s=2,...,n}} \left(\prod_{s=2}^{n} \binom{m_{s}}{m_{s,1},...,m_{s,k}}\right) \left(\frac{d}{du}\right)^{m_{1}} \det_{k \times k} \left(g_{2i-j+2\sum_{s=2}^{n} sm_{s,i}}(u)\right) \bigg|_{u=0}. \end{split}$$

Putting it all together yields (3.2). The proof of (3.3) is similar.

Proposition 3.11. Let $k \ge 1$ and m_j be integers for $j = 1, \ldots, k$. Then we have

$$\frac{1}{(2\pi i)^k} \oint \cdots \oint_{|w_i|=1} \frac{\Delta(w)\Delta(w^2) e^{N\sum_{i=1}^k w_i}}{\prod_{j=1}^k w_j^{2k+m_j}} \prod_{i=1}^k dw_i = \sum_{\mu \in S_k} \det_{k \times k} \left(\frac{N^{2k+m_{\mu(i)}-2i-j+2}}{\Gamma(2k+m_{\mu(i)}-2i-j+3)} \right).$$

Proof. As in the proof of Proposition 3.10, we write

$$\Delta(w)\Delta(w^2) e^{N\sum_{i=1}^k w_i} = \Delta\left(\frac{d}{dX}\right)\Delta\left(\frac{d}{dY}\right) \exp\left(\sum_{i=1}^k w_i^2 X_i + w_i Y_i\right) \bigg|_{\substack{X_i=0, \\ Y_i=N}}$$

Then, we have that

$$\frac{1}{(2\pi 1)^k} \oint \cdots \oint \frac{\Delta(w)\Delta(w^2) e^{N\sum_{i=1}^k w_i}}{\prod_{j=1}^k w_j^{2k+m_j}} \prod_{i=1}^k dw_i = \Delta\left(\frac{d}{dX}\right)\Delta\left(\frac{d}{dY}\right) \prod_{i=1}^k f_i(X_i, Y_i) \bigg|_{\substack{X_i=0, \\ Y_i=N}},$$

where

$$f_i(X_i, Y_i) = \frac{1}{2\pi i} \oint_{|w|=1} \frac{e^{(w^2 X_i + wY_i)}}{w^{2k+m_i}} dw$$

So, by Lemma 3.2, we have

$$\frac{1}{(2\pi i)^k} \oint \cdots \oint \frac{\Delta(w)\Delta(w^2) e^{N\sum_{i=1}^k w_i}}{\prod_{j=1}^k w_j^{2k+m_j}} \prod_{i=1}^k dw_i = \sum_{\mu \in S_k} \det_{k \times k} \left(\frac{d^{i+j-2}}{dX^{i-1}dY^{j-1}} f_{\mu(i)}(X,Y) \right) \Big|_{\substack{X=0, \\ Y=N}}$$

Now,

$$\frac{d^{i+j-2}}{dX^{i-1}dY^{j-1}} f_{\mu(i)}(X,Y) \Big|_{\substack{X=0,\\Y=N}} = \frac{1}{2\pi i} \oint_{|w|=1} \frac{e^{Nw}}{w^{2k+m_{\mu(i)}-2i-j+3}} dw = \frac{N^{2k+m_{\mu(i)}-2i-j+2}}{\Gamma(2k+m_{\mu(i)}-2i-j+3)},$$

and the proposition follows.

Lastly, we include a lemma that allows us to explicitly evaluate certain determinants whose entries are reciprocals of the Gamma function.

Lemma 3.12. Let $k \ge 1$ and $m_j \ge 0$ be integers for $j = 1, \ldots, k$. Then, we have

$$\det_{k \times k} \left(\frac{1}{\Gamma(2k+m_i-2i-j+2)} \right) = \prod_{j=1}^k \frac{1}{(2k+m_j-2j)!} \prod_{1 \le i,j \le k} (m_j-m_i-2j+2i).$$

Proof. With our notation, equation (4.13) in [24] can be written as

$$\det_{k \times k} \left(\frac{1}{\Gamma(z_i - j + 1)} \right) = \frac{\Delta(z_1, \dots, z_k)}{\prod_{j=1}^k \Gamma(z_j)}.$$

We take $z_i = 2k + m_i - 2i + 1$ for $i = 1, \ldots, k$. Then, we have

$$\det_{k \times k} \left(\frac{1}{\Gamma(2k + m_i - 2i - j + 2)} \right) = \prod_{i=1}^k \Gamma(2k + m_i - 2i + 1)^{-1} \prod_{1 \le i < j \le k} (m_j - 2j - m_i + 2i).$$

Since $2k + m_i - 2i + 1 \ge 1$ for $1 \le i \le k$, we have that $\Gamma(2k + m_i - 2i + 1) = (2k + m_i - 2i)!$ which completes the proof.

4. PROOFS OF THE MAIN RESULTS

In this section we will present the proofs of our main results. Our strategy is to obtain the joint moments by differentiating the corresponding shifted moments with respect to the shifts. Indeed, one may check by induction that for $G(2N) \in \{Sp(2N), SO(2N), O^{-}(2N)\}$, we have that

$$\int_{G(2N)} \left(\Lambda_A^{(n_1)}(1) \right)^{k_1} \left(\Lambda_A^{(n_2)}(1) \right)^{k_2} dA$$

= $\prod_{j=1}^{k_1} \left(\frac{d}{d\alpha_j} \right)^{n_1} \prod_{j=k_1+1}^k \left(\frac{d}{d\alpha_j} \right)^{n_2} I(G(2N); e^{-\alpha_1}, \dots, e^{-\alpha_k}) \big|_{\alpha_j=0} \left(1 + O(N^{-1}) \right),$

where $k = k_1 + k_2$. Also, the error terms in Lemmas 3.5-3.7 are uniform in α so we obtain an asymptotic formula after performing the differentiation.

4.1. The unitary symplectic group Sp(2N).

Proof of Theorem 2.1. By the above argument and Lemma 3.5, we have that

$$\int_{Sp(2N)} \left(\Lambda_A^{(n_1)}(1)\right)^{k_1} \left(\Lambda_A^{(n_2)}(1)\right)^{k_2} dA = \frac{(-1)^{k(k-1)/2}}{k!} J_{k_1,k_2}^{Sp}(n_1,n_2) \left(1+O(N^{-1})\right), \tag{4.1}$$

where

$$J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) = \prod_{j=1}^{k_{1}} \left(\frac{d}{d\alpha_{j}}\right)^{n_{1}} \prod_{j=k_{1}+1}^{k} \left(\frac{d}{d\alpha_{j}}\right)^{n_{2}} \frac{1}{(2\pi)^{k}} \oint \cdots \oint_{|w_{i}|=1} \frac{\Delta(w)\Delta(w^{2}) e^{N\sum_{i=1}^{k}(w_{i}-\alpha_{i})}}{\prod_{1\leq i,j\leq k}(w_{i}^{2}-\alpha_{j}^{2})} \prod_{i=1}^{k} dw_{i} \bigg|_{\alpha_{j}=0}.$$
(4.2)

We use Lemma 3.9 to carry out the differentiation and obtain

$$\begin{split} J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) \\ &= \frac{1}{(2\pi 1)^{k}} \oint \cdots \oint \left(\sum_{\substack{a_{1}+2a_{2}+\dots+n_{1}a_{n_{1}}=n_{1}\\a_{3}=a_{5}=\dots=0}} \frac{n_{1}!}{a_{1}!\cdots a_{n_{1}}!} (-N)^{a_{1}} \prod_{j=1}^{[n_{1}/2]} \left(\frac{1}{j} \sum_{i=1}^{k} \frac{1}{w_{i}^{2j}}\right)^{a_{2j}} \right)^{k_{1}} \\ &\times \left(\sum_{\substack{b_{1}+2b_{2}+\dots+n_{2}b_{n_{2}}=n_{2}\\b_{3}=b_{5}=\dots=0}} \frac{n_{2}!}{b_{1}!\cdots b_{n_{2}}!} (-N)^{b_{1}} \prod_{j=1}^{[n_{2}/2]} \left(\frac{1}{j} \sum_{i=1}^{k} \frac{1}{w_{i}^{2j}}\right)^{b_{2j}} \right)^{k_{2}} \Delta(w) \Delta(w^{2}) e^{N \sum_{i=1}^{k} w_{i}} \prod_{i=1}^{k} \frac{dw_{i}}{w_{i}^{2k}} dw_{i}^{2k} dw_{i}^$$

Recall the definition of the tuples \mathbf{a}_i and \mathbf{b}_i defined in the statement of the theorem. Then, we can expand the brackets in the integrand of $J_{k_1,k_2}^{Sp}(n_1,n_2)$ as

$$\left(\sum_{\substack{a_1+2a_2+\dots+n_1a_{n_1}=n_1\\a_3=a_5=\dots=0}}\frac{n_1!}{a_1!\dots a_{n_1}!}(-N)^{a_1}\prod_{j=1}^{[n_1/2]}\left(\frac{1}{j}\sum_{i=1}^k\frac{1}{w_i^{2j}}\right)^{a_{2j}}\right)^{k_1}$$
$$=\sum_{u_1+\dots+u_P=k_1}\binom{k_1}{u_1,\dots,u_P}\frac{(n_1)!^{k_1}}{\prod_{i=1}^P(\mathbf{a}_i!)^{u_i}}(-N)^{\sum_{i=1}^Pu_ia_{i,0}}\prod_{j=1}^{[n_1/2]}\left(\frac{1}{j}\sum_{l=1}^k\frac{1}{w_l^{2j}}\right)^{\sum_{i=1}^Pu_ia_{i,j}},$$

with a similar expression for the bracket to the power of k_2 in the integrand. Using these expansions, our expression for $J_{k_1,k_2}^{Sp}(n_1,n_2)$ becomes

$$J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) = \sum_{u_{1}+\dots+u_{P}=k_{1}} \binom{k_{1}}{u_{1},\dots,u_{P}} \frac{(n_{1})!^{k_{1}}}{\prod_{i=1}^{P}(\mathbf{a}_{i}!)^{u_{i}}(\prod_{j=1}^{[n_{1}/2]}j\Sigma_{i=1}^{P}u_{i}a_{i,j})} (-N)^{\sum_{i=1}^{P}u_{i}a_{i,0}}$$

$$\times \sum_{v_{1}+\dots+v_{Q}=k_{2}} \binom{k_{2}}{v_{1},\dots,v_{Q}} \frac{(n_{2})!^{k_{2}}}{\prod_{i=1}^{Q}(\mathbf{b}_{i}!)^{v_{i}}(\prod_{j=1}^{[n_{2}/2]}j\Sigma_{i=1}^{Q}v_{i}b_{i,j})} (-N)^{\sum_{i=1}^{Q}v_{i}b_{i,0}}$$

$$\times \frac{1}{(2\pi 1)^{k}} \oint \cdots \oint_{|w_{i}|=1} \Delta(w)\Delta(w^{2}) e^{N\sum_{i=1}^{k}w_{i}} \prod_{j=1}^{[n_{2}/2]} \left(\sum_{l=1}^{k}\frac{1}{w_{l}^{2j}}\right)^{m_{j}} \prod_{i=1}^{k}\frac{dw_{i}}{w_{i}^{2k}},$$

where $m_j = \sum_{i=1}^{P} u_i a_{i,j} + \sum_{i=1}^{Q} v_i b_{i,j}$ for $j = 1, \ldots, [n_1/2]$ and $m_j = \sum_{i=1}^{Q} v_i b_{i,j}$ for $j = [n_1/2] + 1, \ldots, [n_2/2]$. We now apply Proposition 3.10 to the contour integral above with these m_j which gives us that

$$\begin{aligned} J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) &= (-1)^{k(k-1)/2} k! N^{k(k+1)/2} \sum_{u_{1}+\dots+u_{P}=k_{1}} \binom{k_{1}}{u_{1},\dots,u_{P}} \frac{(n_{1})!^{k_{1}}(-N)^{\sum_{i=1}^{P}u_{i}a_{i,0}}}{\prod_{j=1}^{P}(\mathbf{a}_{i}!)^{u_{i}}(\prod_{j=1}^{[n_{1}/2]}j^{\sum_{i=1}^{P}u_{i}a_{i,j}})} \\ &\times \sum_{v_{1}+\dots+v_{Q}=k_{2}} \binom{k_{2}}{v_{1},\dots,v_{Q}} \frac{(n_{2})!^{k_{2}}(-N)^{\sum_{i=1}^{Q}v_{i}b_{i,0}}}{\prod_{i=1}^{Q}(\mathbf{b}_{i}!)^{v_{i}}(\prod_{j=1}^{[n_{2}/2]}j^{\sum_{i=1}^{Q}v_{i}b_{i,j}})} \cdot N^{2\sum_{j=1}^{[n_{2}/2]}jm_{j}} \\ &\times \sum_{\substack{\sum_{i=1}^{k}m_{s,i}=m_{s}\\s=2,\dots,n}} \binom{n}{k_{s}} \binom{m_{s}}{m_{s,1},\dots,m_{s,k}} \binom{d}{du}^{m_{1}} \det_{k\times k} \left(g_{2i-j+2\sum_{s=2}^{n}sm_{s,i}}(u)\right)\Big|_{u=0}. \end{aligned}$$

Using the definition of m_i , we compute the power of N in the summand as

$$\sum_{i=1}^{P} u_i a_{i,0} + \sum_{i=1}^{Q} v_i b_{i,0} + 2 \sum_{j=1}^{[n_2/2]} jm_j = \sum_{i=1}^{P} u_i \left(a_{i,0} + 2 \sum_{j=1}^{[n_1/2]} ja_{i,j} \right) + \sum_{i=1}^{Q} v_i \left(b_{i,0} + 2 \sum_{j=1}^{[n_2/2]} jb_{i,j} \right)$$
$$= n_1 \sum_{i=1}^{P} u_i + n_2 \sum_{i=1}^{Q} v_i$$
$$= k_1 n_1 + k_2 n_2.$$

Also, since $a_{i,0} \equiv n_1 \pmod{2}$ and $b_{i,0} \equiv n_2 \pmod{2}$ for all *i*, the factor of (-1) in the summand is

$$(-1)^{\sum_{i=1}^{P} u_{i}a_{i,0} + \sum_{i=1}^{Q} v_{i}b_{i,0}} = (-1)^{n_{1}\sum_{i=1}^{P} u_{i} + n_{2}\sum_{i=1}^{Q} v_{i}} = (-1)^{k_{1}n_{1} + k_{2}n_{2}}$$

Combining these two observations with our final expression for $J_{k_1,k_2}^{Sp}(n_1,n_2)$ completes the proof.

Proof of Theorem 2.2. We begin as in the proof of Theorem 2.1 with (4.1) and (4.2). We use Lemma 3.8 for the derivatives in this case which gives us that

$$J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) = \frac{1}{(2\pi)^{k}} \oint \cdots \oint_{|w_{i}|=1} \left(\sum_{m=0}^{n_{1}} \binom{n_{1}}{m} (-N)^{n_{1}-m} m! \sum_{\substack{l_{1}+\dots+l_{k}=m\\l_{j} \text{ even}}} \prod_{i=1}^{k} \frac{1}{w_{i}^{l_{i}}} \right)^{k_{1}} \\ \times \left(\sum_{m=0}^{n_{2}} \binom{n_{2}}{m} (-N)^{n_{2}-m} m! \sum_{\substack{l_{1}+\dots+l_{k}=m\\l_{j} \text{ even}}} \prod_{i=1}^{k} \frac{1}{w_{i}^{l_{i}}} \right)^{k_{2}} \Delta(w) \Delta(w^{2}) e^{N \sum_{i=1}^{k} w_{i}} \prod_{i=1}^{k} \frac{dw_{i}}{w_{i}^{2k}}$$

Rather than expand the brackets in the integrand, we write them as

$$\left(\sum_{m=0}^{n_1} \binom{n_1}{m} (-N)^{n_1-m} m! \sum_{\substack{l_1+\dots+l_k=m \ i=1}} \prod_{i=1}^k \frac{1}{w_i^{l_i}} \right)^{k_1}$$

$$= \left(\sum_{\substack{\sum_{j=1}^k l_j \le n_1 \\ l_j \text{ even}}} (-N)^{n_1-\sum_{j=1}^k l_j} \binom{n_1}{\sum_{j=1}^k l_j} \left(\sum_{j=1}^k l_j \right) \left(\sum_{j=1}^k l_j \right)! \prod_{j=1}^k \frac{1}{w_j^{l_j}} \right)^{k_1}$$

$$= \sum_{\substack{2\sum_{j=1}^k l_{i,j} \le n_1 \\ i=1,\dots,k_1}} \prod_{i=1}^{k_1} \left((-N)^{n_1-2\sum_{j=1}^k l_{i,j}} \binom{n_1}{2\sum_{j=1}^k l_{i,j}} \left(2\sum_{j=1}^k l_{i,j} \right)! \right) \prod_{j=1}^k \frac{1}{w_j^{2\sum_{i=1}^{k_1} l_{i,j}}},$$

with a similar expression for the second bracket to the k_2 . We then have that

$$\begin{split} J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) &= \sum_{\substack{2\sum_{j=1}^{k} l_{i,j} \leq n_{1}} \prod_{i=1}^{k_{1}} \left((-N)^{n_{1}-2\sum_{j=1}^{k} l_{i,j}} \binom{n_{1}}{2\sum_{j=1}^{k} l_{i,j}} \right) \left(2\sum_{j=1}^{k} l_{i,j} \right)! \right) \\ &\times \sum_{\substack{2\sum_{j=1}^{k} m_{i,j} \leq n_{2}} \prod_{i=1}^{k_{2}} \left((-N)^{n_{2}-2\sum_{j=1}^{k} m_{i,j}} \binom{n_{2}}{2\sum_{j=1}^{k} m_{i,j}} \right) \left(2\sum_{j=1}^{k} m_{i,j} \right)! \right) \\ &\times \frac{1}{(2\pi 1)^{k}} \oint \cdots \oint_{|w_{i}|=1} \frac{\Delta(w)\Delta(w^{2}) e^{N\sum_{i=1}^{k} w_{i}}}{\prod_{j=1}^{k} w_{j}^{2k+2\sum_{i=1}^{k-1} l_{i,j}+2\sum_{i=1}^{k-2} m_{i,j}} \prod_{i=1}^{k} dw_{i} \\ &= \sum_{\substack{2\sum_{j=1}^{k} l_{i,j} \leq n_{1}} 2\sum_{j=1}^{k} m_{i,j} \leq n_{2}} (n_{1}!)^{k_{1}} (n_{2}!)^{k_{2}} (-N)^{k_{1}n_{1}+k_{2}n_{2}-2\sum_{j=1}^{k} \sum_{l_{i,j}} l_{i,j}+\sum_{i=1}^{k_{2}} m_{i,j})} \\ &\times \left(\prod_{i=1}^{k} \frac{1}{(n_{1}-2\sum_{j=1}^{k} l_{i,j})!}\right) \left(\prod_{i=1}^{k_{2}} \frac{1}{(n_{2}-2\sum_{j=1}^{k} m_{i,j})!}\right) \\ &\times \frac{1}{(2\pi 1)^{k}} \oint \cdots \oint_{|w_{i}|=1} \frac{\Delta(w)\Delta(w^{2}) e^{N\sum_{i=1}^{k} w_{i}}}{\prod_{j=1}^{k} w_{j}^{2k+2\sum_{i=1}^{k-1} l_{i,j}+2\sum_{i=1}^{k-2} m_{i,j}}} \prod_{i=1}^{k} dw_{i}. \end{split}$$

We set $V_j = 2 \sum_{i=1}^{k_1} l_{i,j} + 2 \sum_{i=1}^{k_2} m_{i,j}$ for j = 1, ..., k. Then, by Proposition 3.11, the contour integral in the last line above is equal to

$$\sum_{\mu \in S_k} \det_{k \times k} \left(\frac{N^{2k+V_{\mu(i)}-2i-j+2}}{\Gamma(2k+V_{\mu(i)}-2i-j+3)} \right) = N^{k(k+1)/2+\sum_{j=1}^k V_j} \sum_{\mu \in S_k} \det_{k \times k} \left(\frac{1}{\Gamma(2k+V_{\mu(i)}-2i-j+3)} \right).$$

Hence, our expression for $J^{Sp}_{k_1,k_2}(n_1,n_2)$ becomes

$$\begin{split} J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) &= (-1)^{k_{1}n_{1}+k_{2}n_{2}}(n_{1}!)^{k_{1}}(n_{2}!)^{k_{2}}N^{k(k+1)/2+k_{1}n_{1}+k_{2}n_{2}} \\ &\times \sum_{\mu \in S_{k}} \sum_{\substack{2\sum_{j=1}^{k} l_{i,j} \leq n_{1}}} \sum_{\substack{2\sum_{j=1}^{k} m_{i,j} \leq n_{2} \\ i=1,\dots,k_{1}}} \left(\prod_{i=1}^{k_{1}} \frac{1}{(n_{1}-2\sum_{j=1}^{k} l_{i,j})!} \right) \left(\prod_{i=1}^{k_{2}} \frac{1}{(n_{2}-2\sum_{j=1}^{k} m_{i,j})!} \right) \\ &\times \det_{k \times k} \left(\frac{1}{\Gamma(2k+V_{\mu(i)}-2i-j+3)} \right). \end{split}$$

Now, by an argument similar to that given at the end of the proof of [21, Theorem 25], we have that the sums over $l_{i,j}$ and $m_{i,j}$ do not depend on the choice of permutation μ . Thus, we may take μ to be the identity and replace the sum over $\mu \in S_k$ by k!. To finish, we apply Lemma 3.12 with $m_j = V_j + 1$ to the last determinant which gives us

$$J_{k_{1},k_{2}}^{Sp}(n_{1},n_{2}) = (-1)^{k_{1}n_{1}+k_{2}n_{2}}(n_{1}!)^{k_{1}}(n_{2}!)^{k_{2}}N^{k(k+1)/2+k_{1}n_{1}+k_{2}n_{2}}$$

$$\times \sum_{\mu \in S_{k}} \sum_{\substack{2\sum_{j=1}^{k} l_{i,j} \leq n_{1}}} \sum_{\substack{2\sum_{j=1}^{k} m_{i,j} \leq n_{2}}} \left(\prod_{i=1}^{k_{1}} \frac{1}{(n_{1}-2\sum_{j=1}^{k} l_{i,j})!} \right) \left(\prod_{i=1}^{k_{2}} \frac{1}{(n_{2}-2\sum_{j=1}^{k} m_{i,j})!} \right)$$

$$\times \prod_{j=1}^{k} \frac{1}{(2k+V_{j}-2j+1)!} \prod_{1 \leq i < j \leq k} (V_{j}-V_{i}-2j+2i).$$

The theorem follows.

4.2. The special orthogonal group SO(2N) and $O^{-}(2N)$.

Proof of Theorem 2.3. The proof is similar to that of Theorem 2.1 but we now use Lemma 3.6 for the shifted moments. We again use Lemma 3.9 for the derivatives and apply (3.3) in Proposition 3.10 to the resulting contour integral.

Proof of Theorem 2.4. We follow the proof of Theorem 2.2 using Lemma 3.6 for the shifted moments and Lemma 3.8 for the derivatives. We then use Proposition 3.11 with $m_j = V_j - 1$ for the contour integral and conclude the proof similarly using Lemma 3.12.

Proof of Theorem 2.5. Using the argument at the beginning of the section and Lemma 3.7, we have that

$$\int_{O^{-}(2N)} \left(\Lambda_A^{(n_1)}(1) \right)^{k_1} \left(\Lambda_A^{(n_2)}(1) \right)^{k_2} dA = \frac{(-1)^{k(k-1)/2} 2^k}{k!} J_{k_1,k_2}^{O^{-}(2N)}(n_1,n_2) \left(1 + O(N^{-1}) \right),$$

where

For the derivatives, we use the fact that for $n \ge 1$,

$$\frac{d^n}{d\alpha^n} \left. \frac{\alpha e^{N\alpha}}{\prod_{i=1}^k (w_i^2 - \alpha^2)} \right|_{\alpha=0} = n \frac{d^{n-1}}{d\alpha^{n-1}} \left. \frac{e^{N\alpha}}{\prod_{i=1}^k (w_i^2 - \alpha^2)} \right|_{\alpha=0}$$

Hence, we have that

$$J_{k_{1},k_{2}}^{O^{-}}(n_{1},n_{2}) = n_{1}^{k_{1}}n_{2}^{k_{2}} \times \prod_{j=1}^{k_{1}} \left(\frac{d}{d\alpha_{j}}\right)^{n_{1}-1} \prod_{j=k_{1}+1}^{k} \left(\frac{d}{d\alpha_{j}}\right)^{n_{2}-1} \frac{1}{(2\pi i)^{k}} \oint \cdots \oint \frac{\Delta(w)\Delta(w^{2}) e^{N\sum_{i=1}^{k}(w_{i}+\alpha_{i})}}{\prod_{1\leq i,j\leq k}(w_{i}^{2}-\alpha_{j}^{2})} \prod_{i=1}^{k} dw_{i} \Big|_{\alpha_{j}=0}.$$

This integral expression is very similar to the expression for $J_{k_1,k_2}^{Sp(2N)}(n_1-1,n_2-1)$ given in (4.2) and so we can proceed as in the proof of Theorem 2.1 or 2.2, simply replacing (-N) by N when we use Lemma 3.8 or 3.9. In either case, we obtain the statement of the theorem.

4.3. **Proof of Proposition 2.7.** We conclude this setion by proving Proposition 2.7. Let $n \ge 1$ be an integer. Then, by Theorem 2.2, we have that

$$\begin{split} b_{0,1}^{Sp}(0,n) &= \frac{(-1)^n n!}{2^{n+1}} \sum_{2l \le n} \frac{1}{(n-2l)!(2l+1)!} \\ &= \frac{(-1)^n n!}{2^{n+1}(n+1)!} \sum_{2l \le n} \binom{n+1}{2l+1} \\ &= \frac{(-1)^n}{2^{n+1}(n+1)} \left(\sum_{2l \le n} \binom{n}{2l} + \sum_{2l \le n-1} \binom{n}{2l+1} \right) \\ &= \frac{(-1)^n}{2^{n+1}(n+1)} \sum_{l=0}^n \binom{n}{l} \\ &= \frac{(-1)^n}{2(n+1)}, \end{split}$$

where we have used standard properties of the binomial coefficient. In the same manner, by Theorem 2.4, we have

$$\begin{split} b_{0,1}^{SO}(0,n) &= 2^{1-n} n! \sum_{2l \le n} \frac{1}{(n-2l)!(2l)!} \\ &= 2^{1-n} \sum_{2l \le n} \binom{n}{2l} \\ &= 2^{1-n} \left(\sum_{2l \le n-1} \binom{n-1}{2l} + \sum_{2l \le n} \binom{n-1}{2l-1} \right) \\ &= 2^{1-n} \sum_{l=0}^{n-1} \binom{n-1}{l} \\ &= 1. \end{split}$$

5. Numerical results

Below we give some numerical values for $b_{k_1,k_2}^{Sp}(n_1,n_2)$ and $b_{k_1,k_2}^{SO}(n_1,n_2)$. Values of $b_{k_1,k_2}^{O^-}(n_1,n_2)$ follow from Theorem 2.5 so are omitted. Numerical values for $b_{0,k}^{Sp}(0,2)$ and $b_{0,k}^{SO}(0,2)$ for $k \leq 10$ are given in [1, Section 4]. The following are $b_{0,k}^{Sp}(0,3)$ for $k = 1, \dots, 4$:

$$-\frac{1}{2^{3}}$$

$$-\frac{23}{2^{7} \cdot 3 \cdot 5 \cdot 7}$$

$$-\frac{1}{2^{8} \cdot 5^{2} \cdot 7 \cdot 11}$$

$$-\frac{233}{2^{18} \cdot 3^{4} \cdot 5^{3} \cdot 7^{2} \cdot 11}$$

 $b_{0,k}^{Sp}(0,4)$ for $k = 1, \dots, 4$:

$$\frac{\frac{1}{2 \cdot 5}}{\frac{251}{2^4 \cdot 3^2 \cdot 5^2 \cdot 7 \cdot 11}}$$

$$\frac{89 \cdot 13103}{2^9 \cdot 3^5 \cdot 5^3 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17}$$

$$\frac{1627 \cdot 693731}{2^{10} \cdot 3^5 \cdot 5^5 \cdot 7^3 \cdot 11^2 \cdot 13^2 \cdot 17 \cdot 19 \cdot 23}.$$
We also have $b_{1,1}^{Sp}(n_1, 1)$ for $n_1 = 0, 1$:

$$-\frac{1}{48}, \frac{1}{96}$$

 $b_{1,1}^{Sp}(n_1, 2)$ for $n_1 = 0, 1, 2$:

$$\frac{1}{80}, -\frac{1}{160}, \frac{19}{5040}.$$

 $b_{1,1}^{Sp}(n_1,3)$ for $n_1 = 0, 1, 2, 3$:

$$-\frac{1}{120}, \frac{1}{240}, -\frac{17}{6720}, \frac{23}{13440}.$$

 $b_{1,2}^{Sp}(n_1, 1)$ for $n_1 = 0, 1$:

$$\frac{1}{11520}, -\frac{1}{23040}$$

 $b_{1,2}^{Sp}(n_1, 2)$ for $n_1 = 0, 1, 2$:

$$\frac{103}{3628800}, \ -\frac{103}{7257600}, \ \frac{487}{59875200}$$

 $b_{1,2}^{Sp}(n_1,3)$ for $n_1 = 0, 1, 2, 3$:

$$\frac{1}{89600}, \ -\frac{1}{179200}, \ \frac{19}{5913600}, \ -\frac{1}{492800}$$

The following are $b_{0,k}^{SO}(0,3)$ for $k = 1, 2, 3, 4$:

	1
	3
	$2^2 \cdot 5$
	$\frac{1}{2^4 \cdot 3 \cdot 7}$
	$\frac{1613}{2^9 \cdot 3 \cdot 5^2 \cdot 7^2 \cdot 11 \cdot 13}.$
$b_{0,k}^{SO}(0,4)$ for $k = 1, 2, 3, 4$:	
	1
	71
	$2\cdot 3^2\cdot 5\cdot 7$
	$23 \cdot 2657$

$$\frac{2 \cdot 3^3 \cdot 5^3 \cdot 7^2 \cdot 11 \cdot 13}{7159 \cdot 316201}
 \frac{76 \cdot 3^5 \cdot 5^4 \cdot 7^3 \cdot 11^2 \cdot 13 \cdot 17 \cdot 19}{76 \cdot 35 \cdot 5^4 \cdot 7^3 \cdot 11^2 \cdot 13 \cdot 17 \cdot 19}$$

We also have $b_{1,1}^{SO}(n_1, 1)$ for $n_1 = 0, 1$:

	$1, \frac{1}{2}.$
$b_{1,1}^{SO}(n_1, 2)$ for $n_1 = 0, 1, 2$:	0 1 7
$b_{1,1}^{SO}(n_1,3)$ for $n_1 = 0, 1, 2, 3$:	$\frac{2}{3}, \frac{1}{3}, \frac{7}{30}.$
$b_{1,2}^{SO}(n_1, 1)$ for $n_1 = 0, 1$:	$\frac{1}{2}, \frac{1}{4}, \frac{11}{60}, \frac{3}{20}.$
	$\frac{1}{12}, \ \frac{1}{24}.$
$b_{1,2}^{5O}(n_1,2)$ for $n_1 = 0, 1, 2$:	$\frac{19}{620}, \frac{19}{1260}, \frac{26}{2825}.$
$b_{1,2}^{SO}(n_1,3)$ for $n_1 = 0, 1, 2, 3$:	030 1200 2835
	$\frac{23}{1680}, \ \frac{23}{3360}, \ \frac{43}{10080}, \ \frac{1}{336}.$

Acknowledgments. The first author is grateful to the Leverhulme Trust (RPG-2017-320) for the support through the research project grant "Moments of L-functions in Function Fields and Random Matrix Theory". The research of the second author is supported by an EPSRC Standard Research Studentship (DTP) at the University of Exeter.

References

- S. A. Altuğ, S. Bettin, I. Petrow, Rishikesh and I. Whitehead, A Recursion Formula for Moments of Derivatives of Random Matrix Polynomials, Quart. J. Math. 65 (2014), 1111–1125.
- [2] J. C. Andrade and H. Jung, Mean values of derivatives of L-functions in function fields: IV, J. Korean Math. Soc. 58 (2021), no. 6, 1529–1547.
- [3] J. Andrade and S. Rajagopal, Mean values of derivatives of L-functions in function fields: I, J. Math. Anal. Appl. 443 (2016), 526–541.
- [4] T. Assiotis, J. P. Keating and J. Warren On the joint moments of the characteristic polynomials of random unitary matrices, Int. Math. Res. Notices 2022 (2022), no. 18, 14564–14603.
- [5] E. C. Bailey, S. Bettin, G. Blower, J. B. Conrey, A. Prokhorov, M. O. Rubinstein and N. C. Snaith, *Mixed moments of characteristic polynomials of random unitary matrices*, J. Math. Phys. **60** (2019), no. 8, 083509.
- [6] Y. Barhoumi-Andréani, A new approach to the characteristic polynomial of a random unitary matrix, arXiv:2011.02465, (2020).
- [7] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein and N. C. Snaith, Autocorrelation of Random Matrix Polynomials, Commun. Math. Phys. 237 (2003), no. 3, 365–395.
- [8] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein and N. C. Snaith, Integral moments of L-functions, Proc. London Math. Soc. 91 (2005), no. 1, 33–104.
- [9] J. B. Conrey, M. O. Rubinstein and N. C. Snaith, Moments of the derivative of characteristic polynomials with an application to the Riemann zeta function, Commun. Math. Phys. 267 (2006), no. 3, 611–629. Theory 2 (2008), no. 1, 31–68.
- [10] G. Djanković and D. Đokić, The mixed second moment of quadratic Dirichlet L-functions over function fields, Rocky Mountain J. Math. **51** (2021), no. 6, 2003–2017.
- [11] E. Dueñez, D. W. Farmer, S. Froehlich, C. P. Hughes, F. Mezzadri and T. Phan, Roots of the derivative of the Riemann-zeta function and of characteristic polynomials, Nonlinearity 23 (2010), no. 10, 2599–2621.
- [12] A. Florea, The second and third moment of $L(1/2, \chi)$ in the hyperelliptic ensemble, Forum Math. **29** (2017), no. 4, 873–892.
- [13] P. J. Forrester and N. S. Witte, Boundary conditions associated with the Painlevé III' and V evaluations of some random matrix averages, J. Phys. A: Math. Gen. **39** (2006), no. 28, 8983–8995.
- [14] R. Gharakhloo and N. S. Witte, Modulated Bi-orthogonal Polynomials on the Unit Circle: The 2j k and j 2k Systems, Constr. Approx. 58 (2023), 1–74.
- [15] R. R. Hall, The behaviour of the Riemann zeta-function on the critical line, Mathematika **46** (1999), no. 2, 281–313.
- [16] R. R. Hall, On the stationary points of Hardy's function Z(t), Acta Arith. 111 (2004), 125–140.
- [17] C. P. Hughes, On the characteristic polynomial of a random unitary matrix and the Riemann zeta function, PhD thesis, University of Bristol, 2001.
- [18] A. E. Ingham Mean-value theorems in the theory of the Riemann zeta-function, Proc. London Math. Soc. 2 (1928), no. 1, 273–300.
- [19] N. M. Katz and P. Sarnak, *Random matrices, Frobenius eigenvalues, and monodromy*, American Mathematical Society Colloquium Publications **45** (1999).
- [20] J. P. Keating and N. C. Snaith, Random Matrix Theory and L-Functions at s = 1/2, Commun. Math. Phys. 214 (2000), 91–110.
- [21] J. P. Keating and F. Wei, Joint moments of higher order derivatives of CUE characteristic polynomials I: asymptotic formulae, arXiv:2307.01625, (2023).
- [22] J. P. Keating and F. Wei, Joint moments of higher order derivatives of CUE characteristic polynomials II: Structures, recursive relations, and applications, arXiv:2307.02831, (2023).
- [23] F. Mezzadri, Random matrix theory and the zeros of $\zeta'(s)$, J. Phys. A: Math. Gen. **36** (2003), no. 12, 2945–2962.
- [24] J.-M. Normand, Calculation of some determinants using the s-shifted factorial, J. Phys. A: Math. Gen. **37** (2004), no. 22, 5737–5762.
- [25] K. Soundararajan, The horizontal distribution of zeros of $\zeta'(s)$, Duke Math. J. 91 (1998), no. 1, 33–59.
- [26] Y. Zhang, On the zeros of $\zeta'(s)$ near the critical line, Duke Math. J. **110** (2001), no. 3, 555–572.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EXETER, EXETER, EX4 4QF, UNITED KINGDOM *Email address*: j.c.andrade@exeter.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EXETER, EXETER, EX4 4QF, UNITED KINGDOM *Email address*: cgb212@exeter.ac.uk