A LEAVE-ONE-OUT APPROACH TO APPROXIMATE MESSAGE PASSING

ZHIGANG BAO, QIYANG HAN, AND XIAOCONG XU

Abstract. Approximate message passing (AMP) has emerged both as a popular class of iterative algorithms and as a powerful analytic tool in a wide range of statistical estimation problems and statistical physics models. A well established line of AMP theory proves Gaussian approximations for the empirical distributions of the AMP iterate in the high dimensional limit, under the GOE random matrix model and its variants.

This paper provides a non-asymptotic, leave-one-out representation for the AMP iterate that holds under a broad class of Gaussian random matrix models with general variance profiles. In contrast to the typical AMP theory that describes the empirical distributions of the AMP iterate via a low dimensional state evolution, our leave-one-out representation yields an intrinsically high dimensional state evolution formula which provides non-asymptotic characterizations for the possibly heterogeneous, entrywise behavior of the AMP iterate under the prescribed random matrix models.

To exemplify some distinct features of our AMP theory in applications, we analyze, in the context of regularized linear estimation, the precise stochastic behavior of the Ridge estimator for independent and non-identically distributed observations whose covariates exhibit general variance profiles. We find that its finite-sample distribution is characterized via a weighted Ridge estimator in a heterogeneous Gaussian sequence model. Notably, in contrast to the i.i.d. sampling scenario, the effective noise and regularization are now full dimensional vectors determined via a high dimensional system of equations.

Our leave-one-out method of proof differs significantly from the widely adopted conditioning approach for rotational invariant ensembles, and relies instead on an inductive method that utilizes almost solely integration-by-parts and concentration techniques.

CONTENTS

Date: December 27, 2023.

²⁰⁰⁰ *Mathematics Subject Classification.* 60E15, 60G15.

Key words and phrases. approximate message passing, first-order iterative algorithm, leave-oneout, random matrix theory, regularized least squares, ridge regression, state evolution.

1. Introduction

1.1. Overview. Approximate message passing (AMP), originated from statistical physics and graphical ideas [\[TAP77,](#page-53-0) [KF09,](#page-52-0) [Mon12\]](#page-52-1), has emerged as a popular class of first-order iterative algorithms that find diverse applications in both statistical estimation problems and probabilistic analyses of statistical physics models. Some non-exhaustive examples include vector/matrix estimation problems [\[DMM09,](#page-51-0) [Ran11,](#page-53-1) [BM12,](#page-50-2) [KMS](#page-52-2)+12, [DM14,](#page-51-1) [SR15,](#page-53-2) [DM16,](#page-51-2) [KKM](#page-52-3)+16, [MR16,](#page-53-3) [DAM17,](#page-51-3) [LKZ17,](#page-52-4) [FR18,](#page-51-4) [BKM](#page-50-3)+19, [SC19,](#page-53-4) [MV21,](#page-53-5) [CM22a,](#page-50-4) [MV22\]](#page-53-6), and the analyses of spin glass and perceptron models [\[Bol14,](#page-50-5) [Bol19,](#page-50-6) [DS19,](#page-51-5) [FW21,](#page-51-6) [BNSX22,](#page-50-7) [FLS22\]](#page-51-7). We refer the readers to [\[FVRS22\]](#page-51-8) for a recent survey on the AMP and more references can be found therein.

In its simplest form, with a symmetric input matrix $A \in \mathbb{R}^{n \times n}$ distributed according to GOE(*n*), an initialization $z^{(0)} \in \mathbb{R}^n$ independent of *A*, and a sequence of possibly non-linear functions { $F_t : \mathbb{R} \to \mathbb{R}$ }, the AMP algorithm iteratively generates $z^{(1)}, z^{(2)}, \ldots \in \mathbb{R}^n$, according to

$$
z^{(t+1)} \equiv AF_t(z^{(t)}) - b_t F_{t-1}(z^{(t-1)}), \quad t = 0, 1, 2, \dots
$$
 (1.1)

Here for a vector $z \in \mathbb{R}^n$, $F_t(z) \in \mathbb{R}^n$ is understood as applied component-wise. The most crucial part in the AMP iterate [\(1.1\)](#page-1-1) rests in the choice of the scalar factor $b_t \approx n^{-1} \sum_{\ell \in [n]} \mathbb{E} F'_t(z_{\ell}^{(t)})$ $\ell^{(t)}$) $\in \mathbb{R}$. For this choice of *b_t*, a common heuristic in the literature (cf. [\[BM11,](#page-50-8) [BLM15\]](#page-50-9)) postulates a similar distributional behavior of the AMP iterate $\{z^{(t)}\}$ to the iterate $\{\bar{z}^{(t)}\}$ defined by

$$
\tilde{z}^{(t+1)} = A_t \mathsf{F}_t(\tilde{z}^{(t)}), \quad t = 0, 1, 2, \dots,
$$
\n(1.2)

where $\{A_t\}$ are i.i.d. copies of *A*. A well established line of AMP theory [\[BM11,](#page-50-8) [JM13,](#page-52-5) [BLM15,](#page-50-9) [BMN20\]](#page-50-10) makes this heuristic precise, by proving that in the high dimensional limit $n \to \infty$, the empirical distribution of the AMP iterate $\{z^{(t)}\}$ matches that of $\{\overline{z}^{(t)}\}$, which is easily seen to be Gaussian with variance recursively characterized by the so-called *state evolution*. More precisely, [\[BM11\]](#page-50-8) proves that for any fixed $t = 0, 1, 2, \ldots$, and any good enough test function $\psi : \mathbb{R} \to \mathbb{R}$, it holds almost surely that

$$
\lim_{n \to \infty} \frac{1}{n} \sum_{k \in [n]} \psi(z_k^{(t+1)}) = \mathbb{E} \psi(N(0, \sigma_{t+1}^2)), \tag{1.3}
$$

where for $t \geq 1$, $\{\sigma_t\} \subset \mathbb{R}_{\geq 0}$ is defined via the one-dimensional state evolution

$$
\sigma_{t+1}^2 \equiv \mathbb{E} \, \mathsf{F}_t^2(\mathcal{N}(0, \sigma_t^2)), \quad \sigma_1^2 \equiv \lim_{n \to \infty} \frac{\| \mathsf{F}_0(z^{(0)}) \|^2}{n}.
$$

The method of proof for $(1.3)-(1.4)$ $(1.3)-(1.4)$ in [\[BM11\]](#page-50-8) is based on a conditioning technique initiated in [\[Bol14\]](#page-50-5), which, in an extreme synthesis, iteratively computes the

conditional distribution of $A = GOE(n)$ conditioned on random linear constraints. This approach allows for generalizations to more complicated AMP iterates under rotationally invariant Gaussian matrices/tensors, with initialization $z^{(0)}$ that can be possibly correlated with *A*, cf. [\[Ran11,](#page-53-1) [JM13,](#page-52-5) [BMN20,](#page-50-10) [MV21,](#page-53-5) [EAMS21,](#page-51-9) [GB23\]](#page-52-6). This intimate connection to the rotational invariance property of $GOE(n)$ facilitates adaptation of this technique to other random matrix ensembles with such invariance properties, cf. [\[MP17,](#page-53-7) [RSF19,](#page-53-8) [Tak21,](#page-53-9) [ZWF21,](#page-53-10) [Fan22\]](#page-51-10). Moreover, the conditioning technique is flexible enough to provide non-asymptotic versions of [\(1.3\)](#page-1-2) and [\(1.4\)](#page-1-3), cf. [\[RV18,](#page-53-11) [LW22,](#page-52-7) [CR23\]](#page-51-11).

On the other hand, the requisite rotational invariance property poses non-trivial technical challenges in generalizing [\(1.3\)](#page-1-2)-[\(1.4\)](#page-1-3) to other random matrix ensembles without such properties. A simple test example in this regard is to provide a direct proof to [\(1.3\)](#page-1-2)-[\(1.4\)](#page-1-3) for the classical mean 0 variance 1/*n* Wigner matrices. Even for this basic universality problem, existing direct proofs resort to the completely different method of moments [\[BLM15\]](#page-50-9), which inevitably involves combinatorial calculations for the highly complicated matrix polynomials appearing in the unfolded AMP iterates. Although conceptually simple and amenable to extensions to other random matrix models (cf. [\[WZF22,](#page-53-12) [DLS23\]](#page-51-12)), the moment method is confined to provide weak convergence in distribution in a suitable limiting sense, and therefore falls short of capturing the non-asymptotic, algorithmic behavior of the AMP that are particularly important in its applications to statistical and signal processing problems.

The goal of this paper is to introduce a new, leave-one-out method to understand the finer-scale, non-asymptotic behavior of the AMP iterate. In particular, our method provides a direct representation of the AMP iterate $\{z^{(t)}\}$ in [\(1.1\)](#page-1-1) in terms of a suitable version of [\(1.2\)](#page-1-4) via an associated leave-one-out AMP. As will be clear below, our representation holds for a large class of Gaussian random matrix models with general variance profiles, down to the level of each individual coordinates. Crucially, our leave-one-out method requires neither the rotational invariance property as in the conditioning technique, nor combinatorially complex calculations as in the method of moments. In view of the ubiquity of the leave-one-out idea in the broader context of random matrix theory, and its well-known technical robustness and flexibility compared to, e.g., the moment method, in various random matrix models [\[BS10,](#page-50-11) [BGK16,](#page-50-12) [BES17\]](#page-50-13), we anticipate further developments of our method in analyzing the behavior of more complicated AMP algorithms beyond the random matrix model worked out in this paper.

1.2. A leave-one-out representation of the AMP iterate. For a Gaussian symmetric matrix *A* with independent upper triangular entries, we show that each coordinate of the AMP iterate [\(1.1\)](#page-1-1), where the correction scalar term $b_t F_{t-1}(z^{(t-1)})$ is now replaced by $b_t \circ F_{t-1}(z^{(t-1)})$ with a vector $(b_{t,k} =$ $\sum_{\ell \in [n]} \text{Var}(A_{k\ell}) \mathbb{E}\left[F'_{t,\ell}(z^{(t)}_{\ell})\right]$ $\binom{n}{\ell} |z^{(0)}| \big)_{k \in [n]}$, delivers an efficient representation via an associated leave-one-out AMP iterate. Specifically, for any $k \in [n]$, let $\{z_{\mathfrak{l}-}^{(t)}\}$ $\binom{[t]}{[-k]}$ be the leave-one-out AMP iterate, starting from the same initialization and then subsequently updated by replacing *A* in [\(1.1\)](#page-1-1) via its leave-one-out version *A*[−*k*] that sets

all elements in the *k*-th row and column to be 0. We prove in Theorem [2.1](#page-7-1) that for $t = 0, 1, 2, \ldots$

$$
z_k^{(t+1)} = \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle + O_\mathbf{P}\big((\log n)^{c(1 \vee t)^3} \cdot n^{-1/2} \big). \tag{1.5}
$$

While we have used O_P above for simplicity of presentation, the leave-one-out representation in [\(1.5\)](#page-3-0) is intrinsically non-asymptotic, and holds for the prescribed random, symmetric Gaussian matrices *A* with very general variance profiles, along with general separable nonlinear functions $F_t : \mathbb{R}^n \to \mathbb{R}^n$, $F_t(z) = (F_{t,\ell}(z_{\ell}))_{\ell \in [n]},$ whose components $F_{t,\ell} : \mathbb{R} \to \mathbb{R}$ may differ significantly from each other.

An immediate consequence of [\(1.5\)](#page-3-0) is an efficient method to compute the nonasymptotic distribution of an individual coordinate $z_k^{(t+1)}$ $(k+1)$ ∈ ℝ at iterate $t + 1$ via *z*^(*t*) ∈ ℝ^{*n*} at iterate *t*. To proceed, as *A_k* is independent of $F_t(z_{[-k]}^{(t)})$, by first conditioning on $A_{[-k]}$ and then using concentration of the average along with the approximation $z^{(t)} \approx z_{[-t]}^{(t)}$ $\binom{U}{[-k]}$ (in an ℓ_2 sense), we are then naturally led to $z_k^{(t+1)}$ *k* $\stackrel{d}{\approx}$ **N**(0, ∑_{$\ell \in [n]}$} Var($A_{k\ell}$) \mathbb{E} $\mathsf{F}^2_{t,\ell}(z^{(t)}_{\ell})$ $\binom{r}{\ell}$). Formalizing these heuristics, we prove in Theorem [2.3](#page-8-0) that for $t = 0, 1, 2, ...$,

$$
z_k^{(t+1)} \stackrel{d}{\approx} \mathcal{N}(0, \sigma_{t+1,k}^2), \tag{1.6}
$$

where for $t \geq 1$, $\{\sigma_t\} \subset \mathbb{R}^n$ are now vectors determined via the following, possibly high dimensional state evolution

$$
\sigma_{t+1,k}^2 \equiv \sum_{\ell \in [n]} \text{Var}(A_{k\ell}) \cdot \mathbb{E}\left[F_{t,\ell}^2(\mathcal{N}(0, \sigma_{t,\ell}^2))|z^{(0)}\right], \quad k \in [n],\tag{1.7}
$$

with the initial condition $\sigma_{1,k}^2 \equiv \sum_{\ell \in [n]} \text{Var}(A_{k\ell}) \cdot \mathsf{F}_{0,\ell}^2(z_{\ell}^{(0)})$ $\binom{0}{k}$ for $k \in [n]$.

The behavior of the AMP iterate—primarily in the formalism [\(1.3\)](#page-1-2)—under our random matrix models with general variance profiles, has been previously examined in [\[BLM15\]](#page-50-9) via the method of moments, and a further adaptation of this moment method allowing for sparse matrices was performed in [\[Hac23\]](#page-52-8). As mentioned above, the method of moments is designed for proving asymptotic weak convergence, and therefore does not capture the actual, non-asymptotic behavior of the AMP iterate, which is expected to exhibit different behavior across its coordinates as well as with respect to the problem dimension in our setting. From this perspective, our result in $(1.6)-(1.7)$ $(1.6)-(1.7)$ can be naturally viewed as a non-asymptotic, second-order formula to [\(1.3\)](#page-1-2)-[\(1.4\)](#page-1-3) that provides entrywise distributional characterizations for the AMP iterates in these heterogeneous situations.

1.3. Proof techniques. We now provide a technical glimpse into the core of our new leave-one-out method. Upon using the leave-one-out AMP, we show in Propo-sition [4.1](#page-18-0) that $z_k^{(t+1)}$ $\chi_k^{(t+1)} - \langle A_k, \mathsf{F}_t(z_{[-}^{(t)} \rangle)$ $\binom{t}{[-k]}$ admits an exact error decomposition. This decomposition then essentially boils the problem down to controlling the normalized trace of a sequence of random matrices $M_1^{(t)}$ $\mathbf{M}_t^{(t)}$, \ldots , $M_t^{(t)} \in \mathbb{R}^{n \times n}$, defined via the following matrix recursion: initialized with $M_{-1}^{(t)}$ $\binom{t}{-1} = 0_{n \times n}, M_0^{(t)}$ $\mathbf{D}^{(t)}$ = $\mathbf{D}^{(t)}$, for $s \in [t]$,

$$
M_s^{(t)} \equiv (M_{s-1}^{(t)}A - M_{s-2}^{(t)}B_{t-s+1})D^{(t-s)},
$$
\n(1.8)

where $D^{(s)}$ is a random diagonal matrix that collects all elements ${F'_{s,\ell}}(z_{\ell}^{(s)})$ $\begin{pmatrix} (s) \\ \ell \end{pmatrix}$ $\{e[n]\}$ (or some of its variants), and B_s is a diagonal matrix that collects those of $\{b_{s,\ell}\}_{{\ell \in [n]}}$.

The underlying mechanism for the normalized trace n^{-1} tr $(M_s^{(t)})$ to be small for 1 ≤ *s* ≤ *t* is non-trivial. In the simplest case where $A = GOE(n)$, and F_t 's are identity maps, the matrix couple $(D^{(s)}, B_s)$ becomes $D^{(\cdot)} = B = I_n$, and therefore the cancellations in the trace calculations for $M_s^{(t)}$ can be directly attributed to the orthogonality of Chebyshev polynomials of the second kind with respect to the (scaled) semicircle density on $[-1, 1]$ (see the discussion after the statement of Proposition [4.4](#page-23-0) for details).

In the general case where the variance profile of A is arbitrary and F_t 's are general non-linear functions, there appears no obvious analogue to the above global reduction to special orthogonal polynomials. We instead prove the trace control $n^{-1} \mathbb{E}$ tr($M_s^{(t)}$) = $O((\log n)^{ct^3} \cdot n^{-1/2})$ via an inductive argument (cf. Proposition [4.4\)](#page-23-0). At a high level, our inductive argument recursively reduces the degree of the complicated matrix polynomial in $n^{-1} \mathbb{E}$ tr $(M_s^{(t)})$ via repeated applications of Gaussian integration by parts to the recursion [\(1.8\)](#page-3-3). The feasibility of this strategy relies on the fact that applying Gaussian integration by parts once to [\(1.8\)](#page-3-3) results in a reduced matrix polynomial that shares a similar form with $\{M^{(\cdot)}_k\}$, due to the special coupling of $(D^{(s)}, B_s)$. Therefore, by repeating the procedure, the large complexity of the matrix polynomial in $n^{-1} \mathbb{E} \text{tr}(M_s^{(t)})$ for a general pair $1 \le s \le t$ can be recursively reduced all the way down to the base cases $1 \leq s \leq t \leq 2$. The latter problem can then be handled via straightforward Gaussian concentration of linear and quadratic forms.

The key technical advantage of our leave-one-out method, as briefly described above, lies in its soft analytic nature that explores the Gaussianity of *A* only via Gaussian integration by parts, along with some Gaussian concentration. It is now well understood that a large array of Gaussian random matrix results proved via such integration-by-parts techniques can be generalized to non-Gaussian settings including the sparse matrices and even the non-Gaussian matrices with correlated entries, using the method of cumulant expansion, cf. [\[KKP96,](#page-52-9) [LP09,](#page-52-10) [HK17,](#page-52-11) [LS18,](#page-52-12) [EKS19\]](#page-51-13). We expect similar analytic techniques to be applicable in our setting as well. To avoid distraction in presenting the key proof ingredients of our new leaveone-out method, we focus here on the Gaussian setting and refrain from pursuing these extensions unrelated to our new techniques.

We also expect that the analytic advantages of our leave-one-method will be widely useful in AMP applications to accommodate more general class of random matrix models without rotational invariance. A notable example in this direction is the problem of estimating a low rank matrix corrupted by heterogeneous noise via the AMP with spectral initialization; see, e.g., [\[BR22,](#page-50-14) [GKKZ22,](#page-52-13) [PKK23\]](#page-53-13) for some recent developments. By analyzing a suitable leave-one-out version of this AMP, we expect that our approach can be used to bypass the crucial rotation invariance property of $GOE(n)$ required in the existing analysis for the homogeneous noise case, cf. [\[MV21\]](#page-53-5). We will explore this direction in a future work.

1.4. An application to regularized estimation. Apart from the aforementioned technical advantages, our new leave-one-out method also holds promise for a diverse range of direct applications through the leave-one-out AMP representation [\(1.5\)](#page-3-0) and its consequential, non-asymptotic second-order theory [\(1.6\)](#page-3-1)-[\(1.7\)](#page-3-2).

Here, as an illustration of some new features resulting from the high dimensionality of the state evolution [\(1.7\)](#page-3-2), we focus on one specific example of Ridge regression in the context of regularized estimation. Suppose we observe the pairs of data $\{(A_i, Y_i)\}_{i \in [m]}$ from the linear model

$$
Y_i = A_i^{\top} \mu_0 + \xi_i, \quad i \in [m].
$$
 (1.9)

Here $A_i \in \mathbb{R}^n$, $\xi_i \in \mathbb{R}$ and $Y_i \in \mathbb{R}$ are the covariate vector, the measurement error, and the response associated with the sample $i \in [m]$, respectively. We are interested in estimation/recovery of the unknown signal $\mu_0 \in \mathbb{R}^n$ via the following Ridge estimator [\[HK70\]](#page-52-14): for a given tuning level $\lambda > 0$,

$$
\widehat{\mu} \equiv \widehat{\mu}(\lambda) \equiv \underset{\mu \in \mathbb{R}^n}{\arg \min} \left\{ \frac{1}{2} \sum_{i \in [m]} \left(Y_i - A_i^\top \mu \right)^2 + \frac{\lambda}{2} ||\mu||^2 \right\}.
$$
 (1.10)

The Ridge estimator [\(1.10\)](#page-5-0) has a long history in statistical applications, and its precise asymptotic properties have been investigated under the i.i.d. sampling setting from various aspects, cf. [\[Dic16,](#page-51-14) [DW18,](#page-51-15) [WX20,](#page-53-14) [HMRT22,](#page-52-15) [CM22b,](#page-51-16) [HX23\]](#page-52-16).

Using our AMP theory [\(1.6\)](#page-3-1)-[\(1.7\)](#page-3-2), we may describe the finite-sample distribution of $\widehat{\mu}$ for independent, non-identically distributed (i.n.i.d.) observations with general variance profiles on the covariate A_i 's. Let $y^{\text{seq}}(\gamma^*) \equiv \mu_0 + \gamma^* \circ Z_n$, $Z_n \sim N(0, I_n)$, be the (heteroscedastic) Gaussian sequence model with 'effective noise' $\gamma^* \in \mathbb{R}^n_{\geq 0}$. With the observation $y^{\text{seq}}(\gamma^*)$, let the sequence Ridge estimator with 'effective regularization' $\lambda \tau_{b^*} \in \mathbb{R}^n_{>0}$ be defined via

$$
\widehat{\mu}^{\text{seq}}(\gamma^*; \tau_{b^*}) \equiv \underset{\mu \in \mathbb{R}^n}{\arg \min} \left\{ \frac{1}{2} ||y^{\text{seq}}(\gamma^*) - \mu||^2 + \frac{\lambda}{2} \sum_{j \in [n]} \tau_{b^*, j} \mu_j^2 \right\}.
$$
 (1.11)

Here the crucial, vector-valued parameters $\gamma^* \in \mathbb{R}^n_{\geq 0}$ and $\tau_{b^*} \in \mathbb{R}^n_{\geq 0}$ are determined implicitly via a high dimensional system of *m* + *n* equations in Eqn. [\(3.1\)](#page-13-1) ahead. Interestingly, under a certain transformation, we may recast the system of equations for τ_{b^*} into a specific quadratic vector equation that arises from the random matrix theory literature [\[AEK17,](#page-50-15) [AEK19\]](#page-50-16). With the above notation, we prove in Theorem [3.2](#page-14-0) the distributional approximation $\hat{\mu} \stackrel{d}{\approx} \hat{\mu}^{\text{seq}}(\gamma^*; \tau_{b^*})$, in the sense that for any good enough test function $\psi : \mathbb{R}^2 \to \mathbb{R}$,

$$
\frac{1}{n}\sum_{j\in[n]}\psi(\widehat{\mu}_j,\mu_{0,j})\approx \frac{1}{n}\sum_{j\in[n]}\mathbb{E}\,\psi(\widehat{\mu}_j^{\text{seq}}(\gamma^*;\tau_{b^*}),\mu_{0,j})\tag{1.12}
$$

holds with high probability.

The essential new feature of [\(1.12\)](#page-5-1) lies in the inherent high dimensionality of the effective noise γ^* and regularization $\lambda \tau_{b^*}$ for the equivalent, sequence Ridge estimator $\widehat{\mu}^{\text{seq}}(\gamma^*; \tau_{b^*})$, when the sampling scheme of [\(1.9\)](#page-5-2) is i.n.i.d. with a general variance profile on the covariate A_i 's. In contrast, in the i.i.d. sampling scheme with either isotropic or anisotropic Gaussian designs, it is now well understood that variants of the formula [\(1.12\)](#page-5-1) can also be proved for a large class of regularized least squares estimators via Gaussian comparison methods, with a pair of *scalar* effective noise and regularization determined via a system of two equations in two unknowns, cf. [\[TAH18,](#page-53-15) [MM21,](#page-52-17) [CMW23,](#page-51-17) [HS23,](#page-52-18) [HX23\]](#page-52-16). Unfortunately, these random process methods require i.i.d. sampling scheme in an essential way, and therefore do not generalize, at least in its current form, to general i.n.i.d sampling schemes. As such, while we have only verified the formula [\(1.12\)](#page-5-1) for the Ridge estimator (1.10) to avoid excessive technicalities unrelated to our main AMP theory, we anticipate our theory to be useful in proving similar formulae as in [\(1.12\)](#page-5-1) for a broad class of regularized least squares estimators under general i.n.i.d. sampling schemes that are otherwise beyond the reach of existing methods.

1.5. Organization. The rest of the paper is organized as follows. We provide formal statements in Section [2](#page-7-0) for the leave-one-out representation [\(1.5\)](#page-3-0) and its consequences $(1.6)-(1.7)$ $(1.6)-(1.7)$ $(1.6)-(1.7)$ (in a more general form) that hold for both the symmetric and asymmetric AMP algorithms. The application into regularized estimation for the Ridge estimator is detailed in Section [3.](#page-13-0) We provide proofs for the leave-oneout representation [\(1.5\)](#page-3-0) in Section [4,](#page-17-0) whereas the proofs of all other results are contained in Sections [5](#page-33-0)[-6](#page-39-0) and the Appendix.

1.6. **Notation.** For any two integers $m, n \in \mathbb{Z}$, let $[m : n] \equiv \{m, m+1, \ldots, n\}$ when *m* ≤ *n* and ∅ otherwise. Let $[m : n] \equiv [m : n-1]$, $(m : n] \equiv [m+1 : n]$, and we write $[n] \equiv [1 : n]$. For $a, b \in \mathbb{R}$, $a \vee b \equiv \max\{a, b\}$ and $a \wedge b \equiv \min\{a, b\}$. For $a \in \mathbb{R}$, let $a_{\pm} \equiv (\pm a) \vee 0$.

For a vector $x \in \mathbb{R}^n$, let $||x||_p$ denote its *p*-norm $(0 \le p \le \infty)$, and we simply write $||x|| \equiv ||x||_2$. For vectors $x, y \in \mathbb{R}^n$, let $\langle x, y \rangle \equiv x^\top y = \sum_{i \in [n]} x_i y_i$. We often write x^{-1} ≡ $(x_i^{-1})_{i \in [n]}$ ∈ \mathbb{R}^n , x^2 ≡ *x* ◦ *x* = $(x_i^2)_{i \in [n]}$ ∈ \mathbb{R}^n and \mathfrak{D}_x ≡ diag(*x*) ≡ $(x_i 1_{i=j})_{i,j \in [n]} \in \mathbb{R}^{n \times n}$. We use $\{e_k\}$ to denote the canonical basis of the Euclidean space, whose dimension should be self-clear from the context.

For a matrix $M \in \mathbb{R}^{m \times n}$, let $||M||_{op}$, $||M||_F$ denote the spectral and Frobenius norm of *M*, respectively. I_n is reserved for an $n \times n$ identity matrix, written simply as *I* (in the proofs) if no confusion arises. For two matrices $M, N \in \mathbb{R}^{m \times n}$ of the same size, let $\hat{M} \circ N \equiv (M_{ij}N_{ij}) \in \mathbb{R}^{m \times n}$ be their Hadamard product. For squared matrices $M, N \in \mathbb{R}^{n \times n}$, let $\langle M, N \rangle \equiv \text{tr}(M^{\top}N) = \sum_{i,j \in [n]} M_{ij} N_{ij}$.

We use C_x to denote a generic constant that depends only on x , whose numeric value may change from line to line unless otherwise specified. $a \leq x$ *b* and $a \geq x$ *b* mean $a \leq C_x b$ and $a \geq C_x b$, abbreviated as $a = O_x(b)$, $a = \Omega_x(b)$ respectively; $a \times_{x} b$ means $a \leq_{x} b$ and $a \geq_{x} b$, abbreviated as $a = \Theta_{x}(b)$. O and ϕ (resp. O_P) and Φ) denote the usual big and small O notation (resp. in probability). For a random variable *X*, we use \mathbb{P}_X , \mathbb{E}_X (resp. \mathbb{P}^X , \mathbb{E}^X) to indicate that the probability and expectation are taken with respect to *X* (resp. conditional on *X*).

For $\Lambda > 0$ and $\mathfrak{p} \in \mathbb{N}$, a measurable map $f : \mathbb{R}^n \to \mathbb{R}$ is called Λ -pseudo-*Lipschitz of order* p iff

$$
|f(x) - f(y)| \le \Lambda \cdot (1 + ||x|| + ||y||)^{p-1} \cdot ||x - y||, \quad \forall x, y \in \mathbb{R}^n.
$$
 (1.13)

Moreover, *f* is called Λ*-Lipschitz* iff *f* is Λ-pseudo-Lipschitz of order 1, and in this case we often write $||f||_{\text{Lip}} \le L$, where $||f||_{\text{Lip}} \equiv \sup_{x \ne y} |f(x) - f(y)| / ||x - y||$.

For a proper, closed convex function f defined on \mathbb{R}^n , its *proximal operator* prox_{*f*}(\cdot ; τ) for any $\tau > 0$ is defined by prox_{*f*}(x ; τ) = arg min $\tau \in \mathbb{R}^n$ { $\frac{1}{2\pi}$ $\frac{1}{2\tau}||x-z||^2 + f(z)].$

2. Main results

2.1. Symmetric AMP. Consider the general non-linear AMP

$$
z^{(t+1)} = AF_t(z^{(t)}) - b_t \circ F_{t-1}(z^{(t-1)}), \quad t = 0, 1, 2, ..., \tag{2.1}
$$

where $F_t: \mathbb{R}^n \to \mathbb{R}^n$ is 'separable' in the sense that $F_t(z) = (F_{t,\ell}(z_{\ell}))_{\ell \in [n]}$. Here $z^{(-1)} = 0_n$, $F_{-1} \equiv 0$ and the initialization $z^{(0)} \in \mathbb{R}^n$ is either fixed or independent of *A*. We consider the following random matrix model for *A*: for some deterministic, symmetric matrix $V \in \mathbb{R}_{\geq 0}^{n \times n}$ with non-negative entries,

$$
A \equiv V \circ G_n, \quad \{G_{n,ij}\}_{i \leq j} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1/n).
$$

For $t = 1, 2, \ldots$, the correction vector $b_t = (b_{t,k})_{k \in [n]}$ is defined via

$$
b_{t,k} \equiv \frac{1}{n} \sum_{\ell \in [n]} V_{k\ell}^2 \mathbb{E} \left[F'_{t,\ell}(z_{\ell}^{(t)}) | z^{(0)} \right], \quad k \in [n].
$$

Consider the following leave-one-out AMP: for $k \in [n]$,

$$
z_{[-k]}^{(t+1)} = A_{[-k]} \mathsf{F}_t(z_{[-k]}^{(t)}) - b_t \circ \mathsf{F}_{t-1}(z_{[-k]}^{(t-1)}), \quad t = 0, 1, 2, \dots,
$$
 (2.2)

with the same $z_{[-k]}^{(-1)}$ $\begin{bmatrix} (-1) \\ [-k] \end{bmatrix} = 0_n$ and initialization $z_{[-k]}^{(0)}$ $[(-k)] = z^{(0)}$. Here $A_{[-k]} \equiv (A_{ij} \mathbf{1}_{i,j \neq k})_{i,j \in [n]}$ is obtained by setting all elements in the *k*-th row and column of *A* to be 0, and keeping all other elements of *A* unchanged.

As the leave-one-out AMP iterate $\{z_{[-]}^{(t)}\}$ $\begin{bmatrix} (t) \\ [-k] \end{bmatrix}$ still makes use of the randomness of almost the whole matrix *A* except for its *k*-th row and column, we expect that $\{z_{\mathfrak{l}-}^{(t)}\}$ [−*k*] } is close to the original AMP iterate $\{z^{(t)}\}$ except for their *k*-th coordinates. Actually, the following theorem shows that after one more step of iteration, the information on the $z_k^{(t+1)}$ $\binom{(t+1)}{k}$ can be extracted from $z_{[-]}^{(t)}$ $\binom{t}{[-k]}$ instead of using the original $z^{(t)}$.

Theorem 2.1. *Suppose the following hold.*

 $(S1)$ max_{*i*, *j*∈[*n*_{*j*}]} $|V_{ij}|$ ≤ *K holds for some K* ≥ 2*.* $(S2)$ $\mathsf{F}_{s,\ell} \in C^2(\mathbb{R})$ *for all* $s \in [0 : t]$, $\ell \in [n]$, and there exists some $\Lambda \geq 2$ *such that*

$$
\max_{s\in[0:t]}\max_{\ell\in[n]}\left\{\left|\mathsf{F}_{s,\ell}(0)\right|+\max_{q=1,2}\left|\left|\mathsf{F}_{s,\ell}^{(q)}\right|\right|_{\infty}\right\}\leq\Lambda.
$$

Then for any D > 0, there exists some universal constant c_0 > 0 *and another constant* $c_1 = c_1(D)$ *such that with* $\mathbb{P}(\cdot|z^{(0)})$ *-probability at least* $1 - c_1 n^{-D}$ *, uniformly in* k ∈ $[n]$,

$$
\left|z_k^{(t+1)} - \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle\right| \le (c_1 K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0(1 \vee t)^3} \cdot n^{-1/2}.
$$

The above theorem is proved in Section [4.](#page-17-0)

Remark 1*.* A few remarks are in order.

- (1) Theorem [2.1](#page-7-1) covers the matrix-variate AMP iterate studied in [\[JM13\]](#page-52-5), where $\{z^{(t)}\}\subset \mathbb{R}^{n \times r}$, and $F_t: \mathbb{R}^{n \times r} \to \mathbb{R}^{n \times r}$ is defined by $F_t(z) \equiv$ $(F_{t,(.,1)}(z_{.,1}), \ldots, F_{t,(.,r)}(z_{.,r}))$ with 'separable functions' $\{F_{t,(.,u)}(z_{.,u}) : \mathbb{R}^n \to \mathbb{R}^n \}$ \mathbb{R}^n _{*u*∈[*r*]} as those in the vector iterate [\(2.1\)](#page-7-2). Theorem [2.1](#page-7-1) may then be applied to each column of the matrix-variate AMP iterates $\{z^{(t)}\} \subset \mathbb{R}^{n \times r}$. These matrix-variate AMP iterates find applications in generalized linear models [\[Ran11,](#page-53-1) [SCC19,](#page-53-16) [SC19,](#page-53-4) [MV22\]](#page-53-6), and matrix estimation problems [\[MV21,](#page-53-5) [ZWF21,](#page-53-10) [ZSF22,](#page-53-17) [PKK23\]](#page-53-13). For simplicity of presentation, below we shall only focus on the vector case $r = 1$.
- (2) Theorem [2.1](#page-7-1) can be formulated in a slightly more general leave-*k*-out form as follows. Let $\emptyset \neq \mathcal{P} \subset [n]$, and consider the following leave- \mathcal{P} -out AMP: starting from $z_{1-\phi}^{(-1)}$ $\begin{bmatrix} (-1) \\ [-\mathcal{P}] \end{bmatrix} = 0_n$ and the same initialization $z_{[-\mathcal{P}]}^{(0)}$ $\binom{0}{[-P]}$ = $z^{(0)}$, let

$$
z_{[-\mathcal{P}]}^{(t+1)} = A_{[-\mathcal{P}]} \mathsf{F}_t(z_{[-\mathcal{P}]}^{(t)}) - b_t \circ \mathsf{F}_{t-1}(z_{[-\mathcal{P}]}^{(t-1)}), \quad t = 0, 1, 2, \dots,
$$

where $A_{[-P]} \equiv (A_{ij} \mathbf{1}_{i,j \notin P})_{i,j \in [n]}$. Then under the same assumptions as in The-orem [2.1,](#page-7-1) with $\mathbb{P}(\cdot|z^{(0)})$ -probability at least $1 - c_1 n^{-D}$, uniformly in $k \in \mathcal{P}$,

$$
\left|z_k^{(t+1)} - \langle A_k, F_t(z_{[-P]}^{(t)}) \rangle\right| \le (c_1 K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0(1 \vee t)^3} \cdot n^{-1/2}.
$$

The constants c_0 , c_1 may depend further on $|\mathcal{P}|$. The above leave- \mathcal{P} -out representation may be used to obtain joint normal approximations for $(z_k^{(t+1)})$ *k*)*k*∈P ∈ $\mathbb{R}^{|\mathcal{P}|}$. To keep notation simple, below we only work with the case $|\mathcal{P}| = 1$.

As mentioned in the introduction, the leave-one-out representation in Theorem [2.1](#page-7-1) provides a principled method to approximate the distribution for each coordinate of the AMP iterate $\{z^{(t)}\}$, in that $\overline{z_k^{(t+1)}}$ $\binom{n+1}{k}$ is approximately a centered Gaussian random variable whose variance is completely determined by that of $\{z_{\ell}^{(t)}\}$ $\binom{n}{\ell}$ _{$\ell \in [n]$}. The following high dimensional state evolution describes the entire covariance structure of $(z_k^{(t+1)}$ $\zeta_k^{(t+1)}, \zeta_k^{(t)}$ $\frac{1}{k}, \ldots, z_k^{(1)}$ $\binom{1}{k}$ for any $k \in [n]$.

Definition 2.2 (*State evolution for symmetric AMP*). Let $Z^{(1)}, Z^{(2)}, \ldots \in \mathbb{R}^n$ be a sequence of correlated *n*-dimensional centered Gaussian vectors with independent entries. The correlation structure along the iteration path is determined with the initial condition $Var(Z_k^{(1)})$ $h_k^{(1)}$) $\equiv n^{-1} \sum_{\ell \in [n]} V_{k\ell}^2 \mathsf{F}_{0,\ell}^2(z_{\ell}^{(0)})$ $\binom{00}{\ell}$ for $k \in [n]$, and the subsequent recursive updates: for $s_1, s_2 \geq 1$,

$$
Cov(Z_k^{(s_1+1)}, Z_k^{(s_2+1)}) \equiv \frac{1}{n} \sum_{\ell \in [n]} V_{k\ell}^2 \mathbb{E} \left[\mathsf{F}_{s_1,\ell}(Z_{\ell}^{(s_1)}) \mathsf{F}_{s_2,\ell}(Z_{\ell}^{(s_2)}) | z^{(0)} \right], \quad k \in [n].
$$

The following theorem provides a formal distributional characterization for $(z_k^{(t+1)}$ $\zeta_k^{(t+1)}, \zeta_k^{(t)}$ $\binom{t}{k}, \ldots, z_k^{(1)}$ $\binom{1}{k}$ $\stackrel{d}{\approx}$ $(Z_k^{(t+1)}$ $Z_k^{(t+1)}, Z_k^{(t)}$ $z_k^{(t)}, \ldots, Z_k^{(1)}$ $\binom{1}{k}$ for each $k \in [n]$; its proof can be found in Section [5.1.](#page-33-1)

Theorem 2.3. *Suppose* (S1)-(S2) hold for some $K, \Lambda \geq 2$. Fix any Λ -pseudo-*Lipschitz function* $\psi : \mathbb{R}^{t+1} \to \mathbb{R}$ *of order* $\mathfrak{p} \in \mathbb{N}$ *. Then there exist some universal constant c*₀ > 1*, and another constant c*_p > 1 *depending on* p *only, such that*

$$
\max_{k \in [n]} \left| \mathbb{E} \left[\psi(z_k^{(t+1)}, z_k^{(t)}, \dots, z_k^{(1)}) | z^{(0)} \right] - \mathbb{E} \left[\psi(Z_k^{(t+1)}, Z_k^{(t)}, \dots, Z_k^{(1)}) | z^{(0)} \right] \right|
$$

$$
\leq (K\Lambda \log n \cdot (1+||z^{(0)}||_{\infty}))^{c_{\mathfrak{p}}(1\vee t)^3} \cdot n^{-1/c_0^{1\vee t}}.
$$

An important feature of the above theorem is that the distributional characterization holds for *individual* coordinates, as opposed to the typical AMP theory that holds for the average of all coordinates, cf. [\[BM11,](#page-50-8) [JM13,](#page-52-5) [BMN20\]](#page-50-10). This nonasymptotic, entrywise distributional characterization is particularly relevant with the presence of a general variance profile matrix *V*, as the behavior of the AMP iterate $\{z^{(t)}\}$ may differ significantly across different coordinates and with respect to the problem dimension *n*.

We note that the worsened error bound in the above Theorem [2.3](#page-8-0) (compared to that in Theorem [2.1\)](#page-7-1) is fundamentally due to the possible singular nature of the covariance of $(Z_k^{(t+1)}$ $Z_k^{(t+1)}, Z_k^{(t)}$ $z_k^{(t)}, \ldots, Z_k^{(1)}$ $\binom{1}{k}$. In fact, in typical applications where the AMP iterate $\{z^{(t)}\}$ is expected to converge as $t \to \infty$, as $z^{(t+1)}$ becomes increasingly indistinguishable from $z^{(t)}$, the smallest eigenvalue of Cov (($Z_t^{(s+1)}$) $\binom{(s+1)}{k}$ _{s∈[0:*t*]}) is expected to decay as *t* grows.

The C^2 smoothness condition on $\{F_t\}$ in (S2) can be substantially relaxed to a Lipschitz condition, when the main interest lies in the empirical distribution of the AMP iterate $\{z^{(t)}\}$. For technical reasons, in the following theo-rem (proved in Section [5.2\)](#page-35-0), we work with a modified AMP iterate $\{\overline{z}^{(t)}\}$ whose initialization remains the same $\overline{z}^{(0)} = z^{(0)}$, and the correction vectors $\{\overline{b}_{t,k}\}$ *n*^{−1} ∑_{t€[n]} $V_{k\ell}^2 \mathbb{E} [F'_{t,\ell}(Z^{(t)}_{\ell})]$ $\mathcal{L}^{(t)}(\mathcal{L}^{(0)})|z^{(0)}\rangle_{k\in[n]}$ are now defined via the Gaussian random vectors $\{Z^{(\cdot)}\}$ in the state evolution in Definition [2.2,](#page-8-1) with $\{F'_{t,\cdot}\}$'s taken as the weak derivatives of {F*t*,·}.

Theorem 2.4. *Suppose (S1) holds and (S2) is replaced by the following: (SA2)* There exists some $\Lambda \geq 2$ *such that*

$$
\max_{s\in[0:t]}\max_{\ell\in[n]}\left\{|\mathsf{F}_{s,\ell}(0)|+||\mathsf{F}_{s,\ell}||_{\mathrm{Lip}}\right\}\leq\Lambda.
$$

Fix any Λ -pseudo-Lipschitz function $\psi : \mathbb{R}^{t+1} \to \mathbb{R}$ of order 2. Then for any $D > 0$, *there exist a universal constant* $c_0 > 1$ *and another constant* $c_1 = c_1(D) > 0$ *, such that with* $\mathbb{P}(\cdot|z^{(0)})$ *-probability at least* $1 - c_1 n^{-D}$ *,*

$$
\left| \frac{1}{n} \sum_{k \in [n]} \left(\psi(\overline{z}_{k}^{(t+1)}, \overline{z}_{k}^{(t)}, \dots, \overline{z}_{k}^{(1)}) - \mathbb{E} \left[\psi(Z_{k}^{(t+1)}, Z_{k}^{(t)}, \dots, Z_{k}^{(1)}) | z^{(0)} \right] \right) \right|
$$

$$
\leq (c_1 K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0(1 \vee t)^3} \cdot n^{-1/c_0^{1 \vee t}}
$$

holds whenever the right hand side of the above display is smaller or equal to $(1 \wedge \sigma_{*,\psi})^{c_0}$. Here $\sigma_{*,\psi}^2 \equiv \min_{s \in S_{\psi} \cap \mathbb{Z}_{\geq 2}} \min_{k \in [n]} \text{Var}(Z_k^{(s-1)} | z^{(0)})$, where S_{ψ} is the *minimum-sized subset of* $[t + 1]$ *so that* $\psi(z)$ *depends on z only via* $z_{\mathcal{S}_{\psi}}$ *.*

Remark 2*.* A few remarks are in order:

(1) The range of *t* for both Theorems [2.3](#page-8-0) and [2.4](#page-9-0) to be effective is probably sub-optimal. In the GOE case, the averaged distributional characterization for the AMP iterate $\{z^{(t)}\}$ (as in Theorem [2.4](#page-9-0) above) is known to hold up to $t \ll \log n / \log \log n$, cf. [\[RV18\]](#page-53-11), and up to $t = O(n / \text{polylog}(n))$ in the

rank one spiked Wigner model, cf. [\[LW22,](#page-52-7) [LFW23\]](#page-52-19). A different method of analysis in [\[CFM23\]](#page-50-17) proves the convergence of an AMP algorithm with spectral initialization as $t \to \infty$ in the \mathbb{Z}_2 synchronization model. It remains open to examine the validity of our theory, in particular in the regime *t* ≫ log *n*, under a general variance profile matrix *V* and nonlinear ${F_t}$'s.

- (2) We compare Theorem [2.4](#page-9-0) above to related results in the literature.
	- (a) For *V* satisfying $\max_{i \in [n]} |n^{-1} \sum_{j \in [n]} V_{ij}^2 1| \approx 0$, [\[WZF22,](#page-53-12) Theorem 2.4] obtained an asymptotic characterization for the AMP iterate that matches the GOE case with the scalar state evolution [\(1.4\)](#page-1-3). For such *V*'s, the limiting spectrum of $A = V \circ G_n$ is the same semicircle law as in the GOE case, cf. [\[EYY12,](#page-51-18) Theorem 2.1].
	- (b) When both ${F_t}$'s and ψ are polynomials, by applying the moment method to the matrix polynomials appearing in the unfolded AMP iterate, [\[BLM15,](#page-50-9) Theorem 3] proved that the limiting behavior of $n^{-1} \sum_{k \in [n]} \psi((\overline{z}_k^{(s)})$ (k) _{s∈[*t*+1]}) is universal across different random matrix models whose upper triangular entries are independent, mean 0, with matching second moments and admit subgaussian tails. Using the same method of proof as in [\[BLM15\]](#page-50-9) and further polynomial approximation techniques, [\[Hac23\]](#page-52-8) refined these results to accommodate general ${F_t}$'s and ψ , as well as possibly sparse *V*'s, along with an explicit asymptotic state evolution characterization.

Our Theorem [2.4](#page-9-0) allows for complicated patterns of *V* whose corresponding spectrum of *A* can be significantly different from the semicircle law as in (a) (cf. [\[AEK19\]](#page-50-16)). Moreover, as mentioned in the introduction, our Theorem [2.4](#page-9-0) is non-asymptotic in nature, whereas the moment method used in [\[BLM15,](#page-50-9) [Hac23\]](#page-52-8) as in (b) is intrinsically confined to obtain asymptotic results.

We stress again that while we state the results in Theorems [2.1,](#page-7-1) [2.3](#page-8-0) and [2.4](#page-9-0) only for the Gaussian case, as the proof only involves Gaussian integration-by-parts and some Gaussian concentration, universality can be obtained by the well-developed cumulant expansion method [\[KKP96,](#page-52-9) [LP09,](#page-52-10) [HK17,](#page-52-11) [LS18,](#page-52-12) [EKS19\]](#page-51-13). We refrain from these detours here to focus on the new leave-one-out analysis.

2.2. Asymmetric AMP. We consider an asymmetric version of the AMP, which is initialized with $(u^{(0)}, v^{(0)}) \in \mathbb{R}^m \times \mathbb{R}^n$ (here $u^{(0)} \in \mathbb{R}^m$ is a dummy variable with $G_0 \equiv 0$ introduced merely for notational consistency; the algorithm is actually initialized at $v^{(0)}$), and subsequently updated for $t = 0, 1, 2, \ldots$, according to

$$
\begin{cases} u^{(t+1)} = A F_t(v^{(t)}) - b_t^{\mathsf{F}} \circ \mathsf{G}_t(u^{(t)}) \in \mathbb{R}^m, \\ v^{(t+1)} = A^{\mathsf{T}} \mathsf{G}_{t+1}(u^{(t+1)}) - b_{t+1}^{\mathsf{G}} \circ \mathsf{F}_t(v^{(t)}) \in \mathbb{R}^n. \end{cases} \tag{2.3}
$$

Here we consider the following random matrix model for $A \in \mathbb{R}^{m \times n}$: for some deterministic matrix $V \in \mathbb{R}_{\geq 0}^{m \times n}$ with non-negative entries,

$$
A \equiv V \circ G_{m \times n}, \quad \{ (G_{m \times n})_{ij} \}_{i \in [m], j \in [n]} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1/m).
$$

The variance scaling $1/m$ is chosen in accordance to the AMP literature [\[BM11,](#page-50-8) [BMN20\]](#page-50-10). We also follow the same convention that the functions $F_t = (F_{t,\ell})_{\ell \in [n]}$:

 $\mathbb{R}^n \to \mathbb{R}^n$, $G_t = (G_{t,k})_{k \in [m]} : \mathbb{R}^m \to \mathbb{R}^m$ are understood as applied component-wise. For $t = 1, 2, \ldots$, the correction vectors $b_t^{\mathsf{F}} \in \mathbb{R}^m$, $b_t^{\mathsf{G}} \in \mathbb{R}^n$ are defined via

$$
\begin{cases} b_{t,k}^{\mathsf{F}} \equiv \frac{1}{m} \sum_{\ell \in [n]} V_{k\ell}^2 \mathbb{E} \left[\mathsf{F}_{t,\ell}'(v_{\ell}^{(t)}) | v^{(0)} \right], \qquad \ \ k \in [m], \\ b_{t,\ell}^{\mathsf{G}} \equiv \frac{1}{m} \sum_{k \in [m]} V_{k\ell}^2 \mathbb{E} \left[\mathsf{G}_{t,k}'(u_k^{(t)}) | v^{(0)} \right], \qquad \ \ \ell \in [n]. \end{cases}
$$

In the canonical applications of the asymmetric $AMP(2.3)$ into high dimensional linear estimation problems, *m*, *n* are understood as the sample size and the signal dimension, respectively. Moreover, $u^{(t+1)} \in \mathbb{R}^m$ is the residual vector, and $v^{(t+1)} \in$ \mathbb{R}^n is the AMP iterate for the statistical estimator of interest.

We shall first consider leave-one-out representation of $(u^{(t+1)}, v^{(t+1)}) \in \mathbb{R}^m \times \mathbb{R}^n$ in a similar vein to Theorem [2.1.](#page-7-1) There are two leave-one-out versions associated with the asymmetric AMP (2.3) :

• (*Leave-one-row-out version*): For $k \in [m]$, with the same initialization $(u_{L}^{(0)}$ [−*k*] , *v* (0) $\binom{0}{[-k]}$ = $(u^{(0)}, v^{(0)})$ ∈ ℝ^{*m*} × ℝ^{*n*}, let for *t* = 0, 1, 2, ...

$$
\begin{cases}\n u_{[-k]}^{(t+1)} = A_{[-k]}, F_t(v_{[-k]}^{(t)}) - b_t^{\mathsf{F}} \circ \mathsf{G}_t(u_{[-k]}^{(t)}) \in \mathbb{R}^m, \\
 v_{[-k]}^{(t+1)} = A_{[-k], \cdot}^{\top} \mathsf{G}_{t+1}(u_{[-k]}^{(t+1)}) - b_{t+1}^{\mathsf{G}} \circ \mathsf{F}_t(v_{[-k]}^{(t)}) \in \mathbb{R}^n.\n\end{cases} \tag{2.4}
$$

• (*Leave-one-column-out version*): For all $\ell \in [n]$, with the same initialization $(u_{(l)}^{(0)}$ $\overset{(0)}{{_{(\ell)}}}, \overset{(0)}{{\nu_{(\ell)}}}$ $(v^{(0)}_{(\ell)}) = (u^{(0)}, v^{(0)}) \in \mathbb{R}^m \times \mathbb{R}^n$, let for $t = 0, 1, 2, ...$

$$
\begin{cases}\n u_{(-\ell)}^{(t+1)} = A_{\cdot, (-\ell)} \mathsf{F}_t(v_{(-\ell)}^{(t)}) - b_t^{\mathsf{F}} \circ \mathsf{G}_t(u_{(-\ell)}^{(t)}) \in \mathbb{R}^m, \\
 v_{(-\ell)}^{(t+1)} = A_{\cdot, (-\ell)}^{\top} \mathsf{G}_{t+1}(u_{(-\ell)}^{(t+1)}) - b_{t+1}^{\mathsf{G}} \circ \mathsf{F}_t(v_{(-\ell)}^{(t)}) \in \mathbb{R}^n.\n\end{cases} \tag{2.5}
$$

Here for any $k \in [m]$ (resp. $\ell \in [n]$), $A_{[-k], \cdot} \equiv (A_{ij} \mathbf{1}_{i \neq k})_{i \in [m], j \in [n]}$ (resp. $A_{\cdot, (-\ell)} \equiv$ $(A_{ij}1_{i\neq\ell})_{i\in[m],i\in[n]}$) is obtained by setting all elements in the *k*-th row (resp. ℓ -th column) of *A* to be 0 and keeping all other elements of *A* unchanged.

In statistical languages, [\(2.4\)](#page-11-0) and [\(2.5\)](#page-11-1) are obtained via the original AMP iterate (2.3) by discarding the *k*-th sample and the ℓ -th predictor, respectively.

Theorem 2.5. *Suppose the following hold for* $\phi \equiv m/n$.

 (S^*1) (ϕ^{-1} + 1)^{1/2} max_{*k*∈[*m*], ℓ ∈[*n*]| $V_{k\ell}$ | ≤ *K* holds for some *K* ≥ 2*.*}

 (S^*2) $\mathsf{F}_{s,\ell}, \mathsf{G}_{s,k} \in C^2(\mathbb{R})$ *for all* $s \in [0 : t]$, $k \in [m]$, $\ell \in [n]$, and there exists some Λ ≥ 2 *such that*

$$
\max_{s\in [0:t]}\max_{k\in [m], \ell\in [n]}\left\{| {\sf F}_{s,\ell}(0)|+|{\sf G}_{s,k}(0)|+\max_{q=1,2}\left(\|{\sf F}_{s,\ell}^{(q)}\|_\infty+\|{\sf G}_{s,k}^{(q)}\|_\infty\right)\right\}\leq \Lambda.
$$

Then for any D > 0, there exists some universal constant c_0 *> 0 and another constant* $c_1 = c_1(D)$ *such that with* $\mathbb{P}(\cdot|v^{(0)})$ *-probability at least* $1 - c_1 n_{\phi}^{-D}$ *where* $n_{\phi} \equiv n(1 + \phi) = m + n$, uniformly in $k \in [m], \ell \in [n]$,

$$
\begin{aligned} \left| u_k^{(t+1)} - \langle A_k, \mathsf{F}_t (v_{[-k]}^{(t)}) \rangle \right| &\vee \left| v_\ell^{(t+1)} - \langle A_{\cdot \ell}, \mathsf{G}_{t+1} (u_{(-\ell)}^{(t+1)}) \rangle \right| \\ &\leq \left(c_1 K \Lambda \log n_\phi \cdot (1 + \| z^{(0)} \|_\infty) \right)^{c_0 (1 \vee t)^3} \cdot n_\phi^{-1/2}. \end{aligned}
$$

Note that in the above theorem, we have not assumed apriori $m \times n$, although this will be usually required in concrete applications.

Similar to Theorems [2.1](#page-7-1) and [2.3,](#page-8-0) the leave-one-out representation above also leads to distributional characterizations for $\{u_k^{(t+1)}\}$ $\{v_{\ell}^{(t+1)}\}, \{v_{\ell}^{(t+1)}\}$ $\binom{(i+1)}{l}$ via a high dimensional state evolution, defined as follows.

Definition 2.6 (*State evolution for asymmetric AMP*). Let $U^{(1)}, U^{(2)}, \ldots \in \mathbb{R}^m$ and $V^{(1)}$, $V^{(2)}$, ... $\in \mathbb{R}^n$ be two sequences of correlated, centered Gaussian vectors with independent entries. The correlation structure for each sequence along the iteration is determined with the initial condition

$$
\begin{cases} \text{Var}(U_{k}^{(1)}) \equiv \frac{1}{m} \sum_{\ell \in [n]} V_{k\ell}^2 \mathsf{F}^2_{0,\ell}(v_{\ell}^{(0)}), & k \in [m]; \\ \text{Var}(V_{\ell}^{(1)}) \equiv \frac{1}{m} \sum_{k \in [m]} V_{k\ell}^2 \mathbb{E} \big[\mathsf{G}^2_{1,k}(U_{k}^{(1)}) | v^{(0)} \big], & \ell \in [n], \end{cases}
$$

and subsequently updated recursively as follows: for s_1 , $s_2 \geq 1$ and $k \in [m], \ell \in [n]$,

$$
\begin{cases} \text{Cov}(U_{k}^{(s_{1}+1)},U_{k}^{(s_{2}+1)})\equiv\frac{1}{m}\sum_{\ell\in[n]}V_{k\ell}^{2}\,\mathbb{E}\left[\mathsf{F}_{s_{1},\ell}\big(V_{\ell}^{(s_{1})}\big)\mathsf{F}_{s_{2},\ell}\big(V_{\ell}^{(s_{2})}\big)|\nu^{(0)}\right],\\ \text{Cov}(V_{\ell}^{(s_{1}+1)},V_{\ell}^{(s_{2}+1)})\equiv\frac{1}{m}\sum_{k\in[m]}V_{k\ell}^{2}\,\mathbb{E}\left[\mathsf{G}_{s_{1}+1,k}\big(U_{k}^{(s_{1}+1)}\big)\mathsf{G}_{s_{2}+1,k}\big(U_{k}^{(s_{2}+1)}\big)|\nu^{(0)}\right]. \end{cases}
$$

We note that the above recursion does not specify the correlation between the two sequences $(U^{(1)}, U^{(2)}, \ldots)$ and $(V^{(1)}, V^{(2)}, \ldots)$.

The following theorem provides an analogue of Theorem [2.3](#page-8-0) for the asymmetric AMP iterate $\{u_k^{(\bar{t}+1)}\}$ $\{v_{\ell}^{(t+1)}\}, \{v_{\ell}^{(t+1)}\}$ $_{\ell}^{(\ell+1)}$.

Theorem 2.7. *Suppose* (S*1)-(S*2) hold for some $K, \Lambda \ge 2$. Fix any Λ -pseudo-*Lipschitz function* $\psi : \mathbb{R}^{t+1} \to \mathbb{R}$ *of order* $\mathfrak{p} \in \mathbb{N}$ *. Then there exist some universal constant c*₀ > 1*, and another constant c*_p > 1 *depending on* \mathfrak{p} *only, such that*

$$
\max_{k \in [m]} \left| \mathbb{E} \left[\psi(u_k^{(t+1)}, u_k^{(t)}, \dots, u_k^{(1)}) | v^{(0)} \right] - \mathbb{E} \left[\psi(U_k^{(t+1)}, U_k^{(t)}, \dots, U_k^{(1)}) | v^{(0)} \right] \right|
$$

+
$$
\max_{\ell \in [n]} \left| \mathbb{E} \left[\psi(v_{\ell}^{(t+1)}, v_{\ell}^{(t)}, \dots, v_{\ell}^{(1)}) | v^{(0)} \right] - \mathbb{E} \left[\psi(V_{\ell}^{(t+1)}, V_{\ell}^{(t)}, \dots, V_{\ell}^{(1)}) | v^{(0)} \right] \right|
$$

$$
\leq (K \Lambda \log n_\phi \cdot (1 + ||v^{(0)}||_\infty))^{c_p(1 \vee t)^3} \cdot n_\phi^{-1/c_0^{1 \vee t}}.
$$

Similar to Theorem [2.4,](#page-9-0) we may relax the C^2 smoothness condition in (S^*2) when looking at the behavior of the empirical distributions of the modified AMP iterate $\{\overline{u}^{(t)}\}$, $\{\overline{v}^{(t)}\}$ (whose correction vectors $\{\overline{b}_t^{\mathsf{F}}\}$ $\overline{b}_t^{\mathsf{B}}$, $\{\overline{b}_t^{\mathsf{G}}\}$ \tilde{t} are defined by the Gaussian random vectors $\{U^{(\cdot)}\}, \{V^{(\cdot)}\}$ in the state evolution in Definition [2.6\)](#page-12-0).

Theorem 2.8. *Suppose (S*1) holds and (S*2) is replaced by the following: (SA*2)* There exists some $\Lambda \geq 2$ *such that*

$$
\max_{s \in [0:t]} \max_{k \in [m], \ell \in [n]} \left\{ | \mathsf{F}_{s,\ell}(0) | + | \mathsf{G}_{s,k}(0) | + \| \mathsf{F}_{s,\ell} \|_{\mathrm{Lip}} + \| \mathsf{G}_{s,k} \|_{\mathrm{Lip}} \right\} \leq \Lambda.
$$

Fix any Λ -*pseudo-Lipschitz function* $\psi : \mathbb{R}^{t+1} \to \mathbb{R}$ *of order* 2*. Then for any* $D > 0$ *, there exists some universal constant* $c_0 > 0$ *and another constant* $c_1 = c_1(D)$ *such that with* $\mathbb{P}(\cdot | v^{(0)})$ *-probability at least* $1 - c_1 n_{\phi}^{-D}$ *,*

$$
\left| \frac{1}{n_{\phi}} \sum_{k \in [m]} \left(\psi(\overline{u}_k^{(t+1)}, \overline{u}_k^{(t)}, \dots, \overline{u}_k^{(1)}) - \mathbb{E} \left[\psi(U_k^{(t+1)}, U_k^{(t)}, \dots, U_k^{(1)}) | v^{(0)} \right] \right) \right|
$$

$$
\vee \left| \frac{1}{n_{\phi}} \sum_{\ell \in [n]} \left(\psi(\overline{v}_{\ell}^{(t+1)}, \overline{v}_{\ell}^{(t)}, \dots, \overline{v}_{\ell}^{(1)}) - \mathbb{E} \left[\psi(V_{\ell}^{(t+1)}, V_{\ell}^{(t)}, \dots, V_{\ell}^{(1)}) | v^{(0)} \right] \right) \right|
$$

$$
\leq (c_1 K \Lambda \log n_{\phi} \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0(1 \vee t)^3} \cdot n_{\phi}^{-1/c_0^{1 \vee t}}
$$

holds whenever the right hand side of the above display is smaller than or $equal to (1 \wedge \sigma_{*,\psi})^{c_0}.$ *Here* $\sigma_{*,\psi}^2 \equiv \min_{s \in S_{\psi} \cap \mathbb{Z}_{\geq 2}} \min_{k \in [m], \ell \in [n]} \text{Var}(U_k^{(s-1)} | v^{(0)}) \wedge$ $Var(V_{\ell}^{(s-1)} | v^{(0)})$, with S_{ψ} defined in Theorem [2.4.](#page-9-0)

The proofs of Theorems [2.5,](#page-11-2) [2.7](#page-12-1) and [2.8](#page-12-2) for the asymmetric AMP [\(2.3\)](#page-10-0) can be reduced to the symmetric case via a reduction scheme (cf. [\[JM13,](#page-52-5) [BMN20\]](#page-50-10)). For sake of completeness, we provide some details for this reduction in Section [5.3.](#page-39-1)

3. Applications to regularized estimation

Recall the linear model [\(1.9\)](#page-5-2) and the Ridge estimator $\hat{\mu}$ in [\(1.10\)](#page-5-0). We write $A \in \mathbb{R}^{m \times n}$ as the design matrix that collects all A_i 's as its rows, and $\xi, Y \in \mathbb{R}^m$ as the error and response vector that collects all elements $\{\xi_i\}$ and $\{Y_i\}$. We will usually omit notational dependence on $\lambda > 0$ for simplicity.

3.1. The fixed point equation. To describe our fixed point equation, let

$$
\mathscr{V}_m \equiv m^{-1}V \circ V = \mathbb{E} A \circ A \in \mathbb{R}^{m \times n}
$$

be the variance profile matrix of *A*. Consider the following system of (non-linear) equations in $(\gamma, b) \in \mathbb{R}^n_{\geq 0} \times [0, 1)^m$:

$$
\begin{cases}\n\gamma^2 = \mathfrak{D}_{\tau_b}^2 \mathscr{V}_m^{\top} \mathfrak{D}_{1_m-b}^2 (\xi^2 + \mathscr{V}_m \mathfrak{D}_{\lambda \tau_b \circ (1_n + \lambda \tau_b)^{-1}}^2 \mu_0^2) \\
+ \mathfrak{D}_{\tau_b}^2 \mathscr{V}_m^{\top} \mathfrak{D}_{1_m-b}^2 \mathscr{V}_m \mathfrak{D}_{1_n + \lambda \tau_b}^{-2} \gamma^2 \in \mathbb{R}^n, \\
b \circ (1_m - b)^{-1} = \mathscr{V}_m(\tau_b \circ (1_n + \lambda \tau_b)^{-1}) \in \mathbb{R}^m.\n\end{cases} \tag{3.1}
$$

Here $\tau_b \equiv (\mathcal{V}_m^\top (1_m - b))^{-1} \in \mathbb{R}^n$, and recall for a generic vector $x \in \mathbb{R}^n$, we write $x^{-1} = (x_i^{-1})_{i \in [n]} \in \mathbb{R}^n$, $x^2 = (x_i^2)_{i \in [n]} \in \mathbb{R}^n$ and $\mathfrak{D}_x = (x_i \mathbf{1}_{i=j})_{i,j \in [n]} \in \mathbb{R}^{n \times n}$.

The following proposition proves the existence and uniqueness of the solution pair (γ^*, b^*) to the high dimensional system of equations [\(3.1\)](#page-13-1).

Proposition 3.1. *Suppose that* $||V_k||, ||V_{\ell}|| > 0$ *for all* $k \in [m], \ell \in [n]$ *. Then there exists a unique pair* $(\gamma^*, b^*) \in C(\mathbb{R}_{>0} \to \mathbb{R}^n_{\geq 0} \times [0, 1)^m)$ *that solves the system of equations [\(3.1\)](#page-13-1).*

In fact, the above proposition proves a stronger statement in that the solution $\lambda \mapsto (\gamma^*(\lambda), b^*(\lambda))$ exists uniquely as a continuous function of λ . As a consequence, the 'effective regularization' $\tau_{b^*(\cdot)} \equiv (\mathcal{V}_m^{\top}(1_m - b^*(\cdot)))^{-1} : \mathbb{R}_{>0} \to \mathbb{R}_{>0}^n$ (the precise meaning of this terminology will be clear in the next subsection) is locally bounded away from both 0 and ∞ .

It is worth noting that for the Gaussian i.i.d. design, i.e., $V = 1_{m \times n}$ (or equivalently, $\mathcal{V}_m = 1_{m \times n}/m$, any solution (γ^*, b^*) to [\(3.1\)](#page-13-1) must take the form $(\gamma^*, b^*) = (\gamma_0^* 1_n, b_0^* 1_n)$ for some scalars $\gamma_0^* \ge 0, b_0^* \in [0, 1)$. In this case, [\(3.1\)](#page-13-1) reduces to a system of two equations in two unknowns, and can be solved with a closed form by direct calculations (see, e.g., Eqn. [\(3.5\)](#page-17-1) below).

The proof of Proposition [3.1](#page-13-2) is fairly non-trivial due to the high dimensionality of the system of equations (3.1) . The key to the proof is to identify appropriate metric-like structures, under which certain transformations of the solutions (γ^*, b^*) for [\(3.1\)](#page-13-1) are fixed points of a strict contraction mapping. More concretely:

- For the first equation of [\(3.1\)](#page-13-1), with a suitable linear transformation of γ , we will prove strict contraction with respect to the ℓ_{∞} metric;
- For the second equation of (3.1) , inspired by random matrix techniques that deal with vector-valued Stieljes transforms (see, e.g., [\[AEK17,](#page-50-15) [AEK19\]](#page-50-16) and references therein), under a suitable non-linear transformation of *b*, we will first prove strict contraction with respect to a weighted- ℓ_{∞} , symmetric function that does not strictly define a metric. We then use a hyperbolic transformation of this symmetric function that leads to a well-defined metric to induce the existence of a fixed point.

The details of the arguments can be found in Section [6.1.](#page-39-2)

3.2. Distribution of the Ridge estimator. Recall $\widehat{\mu}^{\text{seq}}(\gamma^*; \tau_{b^*})$ defined in [\(1.11\)](#page-5-3). The following theorem provides a formal description of [\(1.12\)](#page-5-1).

Theorem 3.2. *Suppose* $1/K \le m/n, \lambda \le K$ *for some* $K \ge 2$ *. Further suppose* $m^{-1/2}(\|V_k\| \wedge \|V_{\cdot \ell}\|) \geq 1/K$, and $V_{k\ell} \leq K$ holds for all $k \in [m], \ell \in [n]$. Fix any Λ -pseudo-Lipschitz function $\psi : \mathbb{R}^2 \to \mathbb{R}$ of order 2 with $\Lambda \geq 2$. Then for any $D > 1$, there exists some constant $c_1 = c_1(D, K, \Lambda) > 0$, with \mathbb{P}^{ξ} -probability at $least 1 - c_1 n^{-D}$,

$$
\left|\frac{1}{n}\sum_{j\in [n]}\psi(\widehat{\mu}_j,\mu_{0,j}) - \frac{1}{n}\sum_{j\in [n]} \mathbb{E}^{\xi}\,\psi(\widehat{\mu}_j^{\text{seq}}(\gamma^*;\tau_{b^*}),\mu_{0,j})\right| \leq c_1 L_{\mu_0,\xi}^2\cdot e^{-(\log\log n)^{0.99}/c_1}.
$$

Here $L_{\mu_0,\xi} \equiv 1 + n^{-1/2} (\|\mu_0\| \vee \|\xi\|).$

We note that here the error ξ is treated as fixed in the above theorem. It is easy to generalize to the 'usual' statistical setting with random ξ 's; we omit these details.

For the Gaussian i.i.d. design case $V = 1_{m \times n}$, distributional results as in Theorem [3.2](#page-14-0) can be proven via Gaussian comparison methods [\[TAH18,](#page-53-15) [HS23,](#page-52-18) [HX23\]](#page-52-16). In this case, for more specific choices of ψ , for instance $\psi(\mu, \mu_0) = ||\mu - \mu_0||^2$ (which corresponds to the ℓ_2 error of $\widehat{\mu}$), analysis of $n^{-1} \sum_{j \in [n]} \psi(\widehat{\mu}_j, \mu_{0,j})$ can also be appealed directly to existing random matrix methods (and therefore Gaussianity is typically not required); see e.g., [\[Dic16,](#page-51-14) [DW18,](#page-51-15) [HMRT22\]](#page-52-15).

Below we present some simulation for Theorem [3.2.](#page-14-0) In the simulation, we use $m = 100, n = 200, \mu_0 = 1_n$, and fix a variance profile matrix $V \in \mathbb{R}^{m \times n}$ and an error vector $\xi \in \mathbb{R}^m$, whose entries form a random draw from i.i.d. $|N(1, 1)|$ and $N(0, 1)$ respectively. In the left panel of Figure [1,](#page-15-0) we verify that the ℓ_2 estimation error for the Ridge estimator $\hat{\mu}$ matches the prediction in Theorem [3.2,](#page-14-0) both under Gaussian and non-Gaussian designs with a general variance profile. Moreover, while not covered exactly by Theorem [3.2,](#page-14-0) the bias and variance for the single coordinate $\hat{\mu}_1$ in the middle and right panels of Figure [1](#page-15-0) are seen to match the theoretical counterpart $\widehat{\mu}_1^{\text{seq}}$ seq, in that $\mathbb{E}\widehat{\mu}_1 \approx \mathbb{E}\widehat{\mu}_1^{\text{seq}}$ seq and Var $(\widehat{\mu}_1) \approx \text{Var}(\widehat{\mu}_1^{\text{seq}})$ $_1^{\text{eq}}$).

FIGURE 1. *Left panel*: Empirical vs. theoretical ℓ_2 estimation error of the Ridge estimator $\widehat{\mu}$. *Middle and right panels*: Empirical vs. theoretical values of $\mathbb{E} \widehat{\mu}_1$ and Var $(\widehat{\mu}_1)$. *Simulation parameters*: $m = 100$, $n = 200$ with $B = 5000$ Monte Carlo repetitions.

3.3. Main steps for the proof of Theorem [3.2](#page-14-0) via AMP. We will prove Theorem [3.2](#page-14-0) by approximating the Ridge estimator $\hat{\mu}$ via an AMP iterate, whose distributional properties can be analyzed via the AMP theory developed in Section [2.](#page-7-0) This proof strategy is developed in [\[BM12\]](#page-50-2) for the Lasso estimator, and in [\[DM16\]](#page-51-2) for robust estimators, both under Gaussian i.i.d. designs (i.e. $V = 1_{m \times n}$). The key technical difference in our setting is that we need to work with a suitable transformation of the random matrix *A* in the constructed AMP iterate, due to the general variance profile matrix *V*. To this end, let

$$
A_{b^*} \equiv \mathfrak{D}_{1_m-b^*}^{1/2} A \mathfrak{D}_{7_{b^*}}^{1/2}, \quad \xi_{b^*} \equiv \mathfrak{D}_{1_m-b^*}^{1/2} \xi, \quad \theta_{0;b^*} \equiv \mathfrak{D}_{7_{b^*}}^{-1/2} \mu_0,
$$

and let $f_{\eta}: \mathbb{R}^n \to \mathbb{R}$ be defined by $f_{\eta}(\mu) \equiv 2^{-1} \lambda \sum_{\ell \in [n]} \eta_{\ell} \mu_{\ell}^2$. Consider the following iterative algorithm, initialized with $r^{(0)} = 0_m$, $\theta^{(0)} = \theta_{0;b^*}$, and updated according to

$$
\begin{cases} r^{(t+1)} \equiv A_{b^*}(\theta_{0;b^*} - \theta^{(t)}) + \xi_{b^*} + b^* \circ r^{(t)} \in \mathbb{R}^m, \\ \theta^{(t+1)} \equiv \text{prox}_{\mathfrak{f}_{\tau_{b^*}}}(\theta^{(t)} + A_{b^*}^\top r^{(t+1)}) \in \mathbb{R}^n. \end{cases} \tag{3.2}
$$

Let $(\widehat{r}, \widehat{\theta})$ be a stationary point of [\(3.2\)](#page-15-1), and $\mu^{(t+1)} \equiv \mathfrak{D}_{\tau_{b^*}}^{1/2} \theta^{(t+1)}$.

The following proposition provides the precise connection between both (i) the stationary point of [\(3.2\)](#page-15-1) to the original Ridge estimator $\hat{\mu}$, and (ii) (3.2) and the standard form of an asymmetric AMP iterate.

Proposition 3.3. *The following hold.*

- *(1) The stationary point* $\widehat{\theta}$ *of* [\(3.2\)](#page-15-1) *satisfies* $\widehat{\mu} = \mathfrak{D}_{\tau_{b^*}}^{1/2} \widehat{\theta}$.
- *(2) [\(3.2\)](#page-15-1) can be recast into the standard asymmetric AMP iterate [\(2.3\)](#page-10-0) via the following identification: with initialization* $v^{(0)} = 0$ *, for* $t = 0, 1, 2, \ldots$ *, the* $sequence (u^{(t+1)}, v^{(t+1)}) \in \mathbb{R}^m \times \mathbb{R}^n$ *is updated according to*

$$
\begin{cases}\nu^{(t+1)} = \xi_{b^*} - r^{(t+1)}, & \nu^{(t+1)} = \theta_{0;b^*} - \theta^{(t)} - A_{b^*}^T r^{(t+1)}, \\
G_t(u) = (u - \xi_{b^*}) \mathbf{1}_{t \ge 1}, & F_t(v) = (\text{prox}_{\mathfrak{f}_{\tau_{b^*}}}(\theta_{0;b^*} - v) - \theta_{0;b^*}) \mathbf{1}_{t \ge 1}.\n\end{cases} \tag{3.3}
$$

Moreover, we have $\mu^{(t+1)} = \text{prox}_{\lambda \|\cdot\|^2/2}(\mu_0 - \mathfrak{D}_{\tau_{b^*}}^{1/2} v^{(t+1)}; \tau_{b^*}).$

Proof. (1). By the first-order optimality condition for $\widehat{\mu}$, for any $\eta \in \mathbb{R}^n_{>0}$,

$$
\widehat{\mu} = \text{prox}_{f_{\eta}} \left(\widehat{\mu} + \mathfrak{D}_{\eta} A^{\top} (A(\mu_0 - \widehat{\mu}) + \xi) \right).
$$

On the other hand, a stationary point $\widehat{\theta}$ of [\(3.2\)](#page-15-1) satisfies

$$
\widehat{\theta} = \text{prox}_{\mathbf{f}_{\tau_{b^*}}} \left[\widehat{\theta} + A_{b^*}^{\top} \mathfrak{D}_{1_m - b^*}^{-1} \Big(A_{b^*} (\theta_{0; b^*} - \widehat{\theta}) + \xi_{b^*} \Big) \right],
$$

\n
$$
\Leftrightarrow \quad \mathfrak{D}_{\tau_{b^*}}^{1/2} \widehat{\theta} = \text{prox}_{\mathbf{f}_{\tau_{b^*}}} \left[\mathfrak{D}_{\tau_{b^*}}^{1/2} \widehat{\theta} + \mathfrak{D}_{\tau_{b^*}}^{1/2} A_{b^*}^{\top} \mathfrak{D}_{1_m - b^*}^{-1} \Big(A_{b^*} (\theta_{0; b^*} - \widehat{\theta}) + \xi_{b^*} \Big) \right],
$$

\n
$$
\Leftrightarrow \quad \mathfrak{D}_{\tau_{b^*}}^{1/2} \widehat{\theta} = \text{prox}_{\mathbf{f}_{\tau_{b^*}}} \left[\mathfrak{D}_{\tau_{b^*}}^{1/2} \widehat{\theta} + \mathfrak{D}_{\tau_{b^*}} A^{\top} \Big(A (\mu_0 - \mathfrak{D}_{\tau_{b^*}}^{1/2} \widehat{\theta}) + \xi \Big) \right].
$$

Comparing the above two equations leads to (1).

(2). From [\(3.3\)](#page-15-2), it is easily verified that for $t \ge 1$, $F_t(v^{(t)}) = \theta^{(t)} - \theta_{0;b^*}$ and $G_t(u^{(t)}) = -r^{(t)}$. Furthermore, the correction vector $b_t^G = 1_n$, as for all $\ell \in [n]$, $b_{t,\ell}^G = \sum_{k \in [m]} \mathbb{E}(A_{b^*})_{k\ell}^2 = 1$ due to the normalization τ_{b^*} . Meanwhile, the other correction vector $b_t^F = b^*$, as for $k \in [m]$,

$$
b_{t,k}^{\mathsf{F}} = \sum_{\ell \in [n]} \mathbb{E}(A_{b^*})_{k\ell}^2 (1 + \lambda \tau_{b^*,\ell})^{-1} = \frac{1 - b_k^*}{m} \sum_{\ell \in [n]} V_{k\ell}^2 \frac{\tau_{b^*,\ell}}{1 + \lambda \tau_{b^*,\ell}} = b_k^*,
$$

where the last identity uses the second equation of [\(3.1\)](#page-13-1). Due to the linearity of F_t , G_t , it makes no difference to use either ${b^F}$. $\mathsf{F}, b^{\mathsf{G}}$ $\left\{ \begin{matrix} G \\ G \end{matrix} \right\}$ or $\{ \overline{b}^{\mathsf{F}}_G \}$ $\mathsf{F}, \overline{b}^{\mathsf{G}}$ · }.

Next we shall quantify the ℓ_2 error between the iterate $\theta^{(t)}$ and the stationary point $\widehat{\theta} = \mathfrak{D}_{\tau_{b^*}}^{-1/2} \widehat{\mu}$ in [\(3.2\)](#page-15-1).

Proposition 3.4. *Suppose* $1/K \leq m/n, \lambda \leq K$ *for some* $K \geq 2$ *. Further suppose* $m^{-1/2}(\|V_k\| \wedge \|V_{\cdot \ell}\|) \geq 1/K$, and $V_{k\ell} \leq K$ holds for all $k \in [m], \ell \in [n]$. Then *for any D > 1, there exist some universal constant* c_0 > 0 *and another constant* $c_1 = c_1(D) > 0$ *, such that for all* $1 \le t \le \log n/c_0$ *, with* $\mathbb{P}^{\xi}(\cdot|\theta^{(0)})$ *-probability at* $least 1 - c_1 n^{-D}$,

$$
n^{-1} \max \{ ||\theta^{(t)} - \widehat{\theta}||^2, ||r^{(t)} - \widehat{r}||^2 \}
$$

$$
\leq K^{c_0} \cdot n^{-1} (||\mu_0||^2 \vee ||\xi||^2) \cdot (1 - K^{-c_0})^t + (K \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot n^{-1/c_0^t}.
$$

An important step in the proof of Proposition [3.4](#page-16-0) is to relate the first equation of [\(3.1\)](#page-13-1) to the high dimensional state evolution in Definition [2.6,](#page-12-0) which then allows applications of the joint distributional characterizations in Theorem [2.8.](#page-12-2)

Under Gaussian i.i.d. designs, similar ℓ_2 error controls between a suitably designed AMP iterate and a given regularized regression estimator of interest are obtained in several cases, typically in the high dimensional limit $m/n \to \phi \in (0, \infty)$ followed by $t \to \infty$. Examples along this line include the Lasso estimator [\[BM12\]](#page-50-2), robust estimators [\[DM16\]](#page-51-2), the SLOPE estimator [\[BKRS21\]](#page-50-18), ℓ_1 -minimum norm interpolator [\[LW21\]](#page-52-20) in the linear model, and the maximum likelihood estimator in logistic regression [\[SCC19,](#page-53-16) [SC19\]](#page-53-4). Here we provide non-asymptotic ℓ_2 error controls in the Ridge case under a general variance profile on the design matrix.

With Proposition [3.4](#page-16-0) in hand, we may then use our AMP theory in Theorem [2.8](#page-12-2) to derive the distribution of $\hat{\mu}$. The details of the proofs for both Proposition [3.4](#page-16-0) and Theorem [3.2](#page-14-0) can be found in Section [6.2.](#page-44-0)

3.4. Conjectured fixed point equations for general regularized least squares. The above method of analysis for the Ridge estimator $\hat{\mu}$ in [\(1.10\)](#page-5-0) via our AMP theory suggests a broader paradigm for a general class of regularized least squares estimators.

Let $f : \mathbb{R} \to \mathbb{R}_{\geq 0}$ be a non-negative, proper and closed convex function, and let $f_n(\mu) \equiv \sum_{j \in [n]} f(\mu_j)$. We are interested in the behavior of the regularized least squares estimator $\widehat{\mu}_{f} \in \arg \min_{\mu \in \mathbb{R}^{n}} \{2^{-1} || Y - A\mu||^{2} + f_{n}(\mu)\}\)$. We conjecture that $\widehat{\mu}_{f} \approx$ ≈ prox_{f_n}($\mu_0 + \gamma_f^* \circ Z_n$; $\tau_{b_f^*}$) in a similar sense to Theorem [3.2,](#page-14-0) where $(\gamma_f^*, b_f^*) \in \mathbb{R}_{\geq 0}^n \times$ $[0, 1)^m$ is now the solution to the following high dimensional system of equations (whenever exists uniquely and is suitably bounded):

$$
\begin{cases}\n\gamma^2 = \mathfrak{D}_{\tau_b}^2 \mathscr{V}_m^{\top} \mathfrak{D}_{1_m-b}^2 \Big[\xi^2 + \mathscr{V}_m \mathbb{E}^{\xi} \big(\text{prox}_{\mathfrak{f}_n} (\mu_0 + \gamma \circ Z_n; \tau_b) - \mu_0 \big)^2 \Big] \in \mathbb{R}^n, \\
b \circ (1_m - b)^{-1} = \mathscr{V}_m \mathfrak{D}_{\tau_b} \mathbb{E}^{\xi} \text{prox}_{\mathfrak{f}_n} (\mu_0 + \gamma \circ Z_n; \tau_b) \in \mathbb{R}^m.\n\end{cases} (3.4)
$$

In the Gaussian i.i.d. design case $V = 1_{m \times n}$ (i.e., $\mathcal{V}_m = 1_{m \times n}/m$), [\(3.4\)](#page-17-2) reduces to a system of two equations in two unknowns $(\gamma, \tau) \in \mathbb{R}_{\geq 0} \times [1, \infty)$: with $\sigma_{\xi}^2 \equiv ||\xi||^2/m$,

$$
\begin{cases}\n\gamma^2 = \sigma_{\xi}^2 + \frac{1}{m} \mathbb{E}^{\xi} ||\text{prox}_{\mathfrak{f}_n}(\mu_0 + \gamma Z_n; \tau) - \mu_0||^2, \\
1 - 1/\tau = \frac{1}{m} \mathbb{E}^{\xi} \operatorname{div}_{\text{prox}_{\mathfrak{f}_n}(\cdot; \tau)}(\mu_0 + \gamma Z_n).\n\end{cases} (3.5)
$$

The above system of equations integrate its limiting version in [\[EK18,](#page-51-19) [TAH18\]](#page-53-15), the Lasso case in [\[MM21\]](#page-52-17) with $f_n(\mu) = \lambda ||\mu||_1$, and a class of general non-separable, non-smooth projection-type regularizers in [\[Han23\]](#page-52-21).

Due to the apparent complications in the analysis of the fixed point equations [\(3.4\)](#page-17-2), it remains an outstanding open question to formally establish the prescribed distributional properties of $\widehat{\mu}_{f}$ via [\(3.4\)](#page-17-2) for general regularizers f_n .

4. Proof of Theorem [2.1](#page-7-1)

In the proofs, we write $\mathbb{E}[\cdot] = \mathbb{E}[\cdot|z^{(0)}]$ for simplicity. For notational simplicity, *t*'s appearing in the estimates are understood as 1 ∨ *t*.

4.1. Decomposition via leave-one-out AMP. Recall the leave-one-out AMP $\{z_{[-}^{(t)}\}$ $\begin{bmatrix} (b) \\ [-k] \end{bmatrix}$ defined in [\(2.2\)](#page-7-3). Let us define some further notation. Let

$$
\Delta z_{[-k]}^{(t)} \equiv z^{(t)} - z_{[-k]}^{(t)},
$$

$$
z_{[-k]}^{(t)}(u) \equiv (1 - u)z_{[-k]}^{(t)} + uz^{(t)}, \quad u \in [0, 1].
$$

We also write

$$
\Delta A_{[-k]} \equiv A - A_{[-k]} = e_k (A_k - A_{kk} e_k)^{\top} + A_k e_k^{\top}.
$$

Recall for any vector $x \in \mathbb{R}^n$, $\mathfrak{D}_x = (x_i \mathbf{1}_{i=j})_{i,j \in [n]}$. Let

$$
(\mathsf{F}')^{(t)} \equiv \mathsf{F}'_t(z^{(t)}) \in \mathbb{R}^n, \quad \mathsf{D}^{(t)} \equiv \mathfrak{D}_{(\mathsf{F}')^{(t)}} \in \mathbb{R}^{n \times n},
$$

$$
(\mathsf{F}')_{[-k]}^{(t)} \equiv \mathsf{F}'_t(z_{[-k]}^{(t)}) \in \mathbb{R}^n, \quad \mathsf{D}_{[-k]}^{(t)} \equiv \mathfrak{D}_{(\mathsf{F}')}^{(t)} \in \mathbb{R}^{n \times n}, \tag{4.1}
$$

and with \mathbb{E}_U being the expectation taken over $U \sim \text{Unif}[0, 1]$, where *U* is independent of *A*, let

$$
\overline{\mathsf{F}'}_{[-k]}^{(t)} \equiv \mathbb{E}_U \,\mathsf{F}'_t(z_{[-k]}^{(t)}(U)) \in \mathbb{R}^n, \quad \overline{\mathsf{D}}_{[-k]}^{(t)} \equiv \mathfrak{D}_{\overline{\mathsf{F}'}_{[-k]}^{(t)}} \in \mathbb{R}^{n \times n}.
$$

For notational convenience, we also write $B_t = D_{b_t} \in \mathbb{R}^{n \times n}$. Using these notation, a first-order Taylor expansion yields that for $r = 1, 2, \ldots$,

$$
\Delta z_{[-k]}^{(r)} = A_{[-k]} \overline{D}_{[-k]}^{(r-1)} \Delta z_{[-k]}^{(r-1)} - B_{r-1} \overline{D}_{[-k]}^{(r-2)} \Delta z_{[-k]}^{(r-2)} + \Delta A_{[-k]} F_{r-1}(z^{(r-1)}).
$$
(4.2)

For any given *t*, we define a sequence of matrices $\overline{M}_{0,1}^{(t)}$ $\overline{M}_{0,[-k]}^{(t)}, \overline{M}_{1,[k]}^{(t)}$ $\overline{M}_{1,[-k]}^{(t)}, \ldots, \overline{M}_{t,[k]}^{(t)}$ *^t*,[−*k*] ∈ $\mathbb{R}^{n \times n}$ recursively as follows: let $\overline{M}_{-1,[-k]}^{(t)} = 0_{n \times n}, \overline{M}_{0,[-k]}^{(t)} \equiv \overline{D}_{[-k]}^{(t)}$ $\begin{aligned} \binom{y}{[-k]} \text{, and for } s \in [t], \end{aligned}$

$$
\overline{M}_{s,[-k]}^{(t)} \equiv (\overline{M}_{s-1,[-k]}^{(t)} A_{[-k]} - \overline{M}_{s-2,[-k]}^{(t)} B_{t-s+1}) \overline{D}_{[-k]}^{(t-s)}.
$$
\n(4.3)

The next proposition provides an exact error decomposition for the leave-one-out representation in Theorem [2.1,](#page-7-1) via the matrix sequence defined above.

Proposition 4.1. *Suppose* $\mathsf{F}_{s,\ell} \in C^1(\mathbb{R})$ *for all* $s \in [0 : t]$ *and* $\ell \in [n]$ *. Then for any k* ∈ [*n*]*,*

$$
z_k^{(t+1)} - \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle = \sum_{s \in [t]} \text{Rem}_k^{(s)},
$$

where for $s \in [t]$ *,*

$$
\text{Rem}_k^{(s)} \equiv \langle A_k, \overline{M}_{t-s, [-k]}^{(t)} e_k \rangle \cdot \left(z_k^{(s)} + b_{s-1,k} \mathsf{F}_{s-2,k} (z_k^{(s-2)}) - A_{kk} \mathsf{F}_{s-1,k} (z_k^{(s-1)}) \right) + \left(\langle A_k, \overline{M}_{t-s, [-k]}^{(t)} A_k \rangle - b_{t,k} \mathbf{1}_{s=t} \right) \cdot \mathsf{F}_{s-1,k} (z_k^{(s-1)}).
$$

Suppose further the following hold for some K, $\Lambda \geq 2$ *.*

- *(1)* max_{*i*, *j*∈[*n*]} $|V_{ij}|$ ≤ *K*.
- *(2)* **F**_{*s*, ℓ} ∈ $C^1(\mathbb{R})$ *for all s* ∈ [0 : *t*], ℓ ∈ [*n*]*, and*

$$
\max_{s\in[0:t]}\max_{\ell\in[n]}\{|F_{s,\ell}(0)|+||F'_{s,\ell}||_{\infty}\}\leq\Lambda.
$$

Then there exists some universal constant $c_0 > 0$ *such that*

$$
\left|z_k^{(t+1)} - \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle \right| \le (K\Lambda)^{c_0 t} \cdot (1 + |A_{kk}|) \cdot \max_{s \in [0:t]} (1 + |z_k^{(s)}|)
$$

$$
\times \Big(\max_{s \in [t]} \Big| \langle \langle A_k, \overline{M}_{t-s, [-k]}^{(t)} A_k \rangle - b_{t,k} \mathbf{1}_{s=t} \rangle \Big| + |A_{kk}| \Big).
$$

Proof. First, using [\(4.2\)](#page-18-1) for $r = t$,

$$
z_k^{(t+1)} = \langle A \mathsf{F}_t(z^{(t)}) - \mathsf{B}_t \mathsf{F}_{t-1}(z^{(t-1)}), e_k \rangle
$$

\n
$$
= \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle + \langle A_k, \overline{M}_{0,[-k]}^{(t)} \Delta z_{[-k]}^{(t)} \rangle - b_{t,k} \mathsf{F}_{t-1,k}(z_k^{(t-1)})
$$

\n
$$
= \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle + \langle A_k, \overline{M}_{1,[-k]}^{(t)} \Delta z_{[-k]}^{(t-1)} - \overline{M}_{0,[-k]}^{(t)} \mathsf{B}_{t-1} \overline{\mathsf{D}}_{[-k]}^{(t-2)} \Delta z_{[-k]}^{(t-2)} \rangle + \text{Rem}_k^{(t)}
$$

.

Next, using [\(4.2\)](#page-18-1) for $r = t - 1$,

$$
\langle A_k, \overline{M}_{1, [-k]}^{(t)} \Delta z_{[-k]}^{(t-1)} - \overline{M}_{0, [-k]}^{(t)} \mathsf{B}_{t-1} \overline{\mathsf{D}}_{[-k]}^{(t-2)} \Delta z_{[-k]}^{(t-2)} \rangle
$$

=
$$
\langle A_k, \overline{M}_{2, [-k]}^{(t)} \Delta z_{[-k]}^{(t-2)} - \overline{M}_{1, [-k]}^{(t)} \mathsf{B}_{t-2} \overline{\mathsf{D}}_{[-k]}^{(t-3)} \Delta z_{[-k]}^{(t-3)} \rangle + \text{Rem}_k^{(t-1)}.
$$

The recursion terminates when [\(4.2\)](#page-18-1) is used until $r = 1$, where by using the terminating condition $\Delta z_{L}^{(0)}$ $\binom{(0)}{[-k]} = 0$, we arrive at

$$
z_k^{(t+1)} = \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle + \text{Rem}_k^{(t)} + \cdots + \text{Rem}_k^{(1)}.
$$

The claimed identity follows. The claimed estimate follows by noting that $|\langle A_k, \overline{M}_{t-}^{(t)}\rangle$ $\left|\frac{u}{t-s,[-k]}e_k\right|$ ≤ $(K\Lambda)^{2t}$ $|A_{kk}|$.

In view of the error bound in Proposition [4.1,](#page-18-0) in the next two subsections we will then (i) provide coordinate-wise controls for the AMP iterate $\{z^{(t)}\}$, as well as (ii) compute the normalized trace of the matrix $\overline{M}_{s}^{(t)}$ s _{*s*}[−*k*] defined via the recursion [\(4.3\)](#page-18-2), as $\overline{M}_{s}^{(t)}$ s _{*s*}[−*k*] and A_k are approximately independent and thus the estimate of the quadratic forms such as $\langle A_k, \overline{M}^{(t)}_{t-s, [-k]} A_k \rangle$ boils down to the estimate of tracial quantities of $\overline{M}_{t-s,[-k]}^{(t)}$ via Hanson-Wright inequality.

4.2. Some apriori estimates. We shall first prove delocalization estimates for the AMP iterate $\{z^{(t)}\}.$

Proposition 4.2. *Suppose the following hold for some K,* $\Lambda \geq 2$ *.*

(1) max_{*i*, *j*∈[*n*]} $|V_{ij}|$ ≤ *K*. *(2)* **F**_{*s*, ℓ} ∈ $C^1(\mathbb{R})$ *for all s* ∈ $[0 : t]$, ℓ ∈ $[n]$, *and*

$$
\max_{s\in[0:t]}\max_{\ell\in[n]}\left\{|\mathsf{F}_{s,\ell}(0)|+||\mathsf{F}_{s,\ell}'||_\infty\right\}\leq\Lambda.
$$

Then there exists some universal constant $c_0 > 0$ *such that for* $x \ge 1$ *,*

$$
\max_{k\in[n]}\mathbb{P}\left(\|z^{(t)}\|_{\infty}\vee\|z^{(t)}-z^{(t)}_{[-k]}\|\geq (K\Lambda)^{c_0t}x\cdot(1+\|z^{(0)}\|_{\infty})\big|z^{(0)}\right)\leq c_0\cdot tne^{-(x^2\wedge n)/c_0}.
$$

Moreover, for any p \geq 1*, there exists some* C_p $>$ 0 *such that for* $t \leq n/(C_p \log n)$ *,*

$$
\mathbb{E}^{1/p}\left[\|z^{(t)}\|_{\infty}^p |z^{(0)}| \vee \max_{k \in [n]} \mathbb{E}^{1/p}\left[\|z^{(t)} - z^{(t)}_{[-k]}\|^p |z^{(0)}\right] \le (K\Lambda)^{C_p t} \sqrt{\log n} \cdot (1 + \|z^{(0)}\|_{\infty}).
$$

Proof. Using the AMP recursion [\(2.1\)](#page-7-2), we have

$$
\begin{aligned} ||\Delta z^{(t+1)}_{[-k]}|| &\leq ||A\mathsf{F}_t(z^{(t)}) - A_{[-k]}\mathsf{F}_t(z^{(t)}_{[-k]})|| + ||b_t||_{\infty}||\Delta z^{(t-1)}_{[-k]}|| \\ &\leq (K\Lambda)^2 (1 + ||A||_{\text{op}})(||\Delta z^{(t)}_{[-k]}|| \vee ||\Delta z^{(t-1)}_{[-k]}||) + ||\Delta A_{[-k]}\mathsf{F}_t(z^{(t)}_{[-k]})||, \end{aligned}
$$

where

$$
\|\Delta A_{[-k]} \mathsf{F}_t(z_{[-k]}^{(t)})\| \leq |A_k^{\top} \mathsf{F}_t(z_{[-k]}^{(t)})| + 2\Lambda \|A_k\| (1 + |z_{k, [-k]}^{(t)}|).
$$

On the other hand, using [\(2.2\)](#page-7-3),

 $\bullet \ \ |z_{k,1}^{(t)}$ $|k(k-1)|$ ≤ $(K\Lambda)^2(1+|z_{k-1-k}^{(t-2)})$ $(k, [-k])$ ≤ ··· ≤ $(K\Lambda)^{ct}(1 + |z_k^{(0)}|)$ $\binom{(0)}{k}$.

•
$$
||z_{[-k]}^{(t)}|| \le (K\Lambda)^2 (1 + ||A_{[-k]}||_{op}) (\sqrt{n} + ||z_{[-k]}^{(t-1)}|| \vee ||z_{[-k]}^{(t-2)}||) \le \cdots \le (K\Lambda(1 + ||A_{[-k]}||_{op}))^{ct} (\sqrt{n} + ||z_{\perp}^{(0)}||).
$$

Using the independence of A_k and $F_t(z_{t-1}^{(t)})$ $\begin{bmatrix} \begin{bmatrix} L \\ -k \end{bmatrix}$ and the standard subgaussian tail estimate for $||A||_{op}$, $||A_{[-k]}||_{op}$ (cf. Lemma [A.1\)](#page-47-1), for $x \ge 1$, with probability at least 1 − $ce^{-(x^2 \land n)/c}$, both $\sqrt{n} |A_k^{\top} \mathsf{F}_t(z_{[-}^{(t)})|)$ $\left| \begin{array}{c} (t) \\ [-k] \end{array} \right|$ ≤ $cKx \cdot ||F_t(z_{[}^{(t)})||$ $\binom{L}{[-k]}$ || and $||A||_{op}$ ∨ $||A_{[-k]}||_{op}$ ≤ *cK* hold. So with the same probability,

$$
\begin{split} \|\Delta z_{[-k]}^{(t+1)}\| &\leq c(K\Lambda)^2 \big(\|\Delta z_{[-k]}^{(t)}\| \vee \|\Delta z_{[-k]}^{(t-1)}\|\big) \\ &+ (K\Lambda)^{ct}(1+|z_k^{(0)}|) + (K\Lambda)^{ct}x \cdot \big(1+n^{-1/2}\|z^{(0)}\|\big) \\ &\leq c(K\Lambda)^2 \big(\|\Delta z_{[-k]}^{(t)}\| \vee \|\Delta z_{[-k]}^{(t-1)}\|\big) + (K\Lambda)^{ct}x \cdot \big(1+\|z^{(0)}\|_{\infty}\big). \end{split}
$$

Iterating the bound, with probability at least $1 - cte^{-(x^2 \wedge n)/c}$,

$$
\|\Delta z_{[-k]}^{(t+1)}\| \le (K\Lambda)^{ct} x \cdot (1 + \|z^{(0)}\|_{\infty}),
$$

and therefore with the same probability,

$$
|z_k^{(t+1)}| \le ||\Delta z_{[-k]}^{(t+1)}|| + |z_{k,[-k]}^{(t+1)}| \le (K\Lambda)^{ct} x \cdot (1 + ||z^{(0)}||_{\infty}).
$$

Consequently, with probability at least $1 - c \cdot tne^{-(x^2 \wedge n)/c}$,

$$
||z^{(t+1)}||_{\infty} \le (K\Lambda)^{ct} x \cdot (1 + ||z^{(0)}||_{\infty}).
$$

The moment estimate for $||z^{(t+1)}||_{\infty}$ follows by combining the above display and the simple apriori estimate $||z^{(t+1)}||$ ≤ $||z^{(t+1)}||$ ≤ $(\Lambda(1 + ||A||_{op}))^{ct}(\sqrt{n} + ||z^{(0)}||)$.

For *i*, *j* ∈ [*n*], let $\partial_{ij} = \partial/\partial A_{ij}$, and

$$
\Delta_{ij} \equiv 2^{-1} (1 + \mathbf{1}_{i \neq j}) (e_i e_j^\top + e_j e_i^\top) = \partial_{ij} A. \tag{4.4}
$$

The following proposition provides delocalization estimates for $\{\partial_{ij}z^{(t)}\}$.

Proposition 4.3. *Suppose the following hold for some K,* $\Lambda \geq 2$ *.*

(1) max_{*i*, *j*∈[*n*_]| V *i*_j| ≤ *K*.} *(2)* **F**_{*s*, ℓ} ∈ $\overline{C^2}(\mathbb{R})$ *for all s* ∈ [0 : *t*], ℓ ∈ [*n*]*, and*

$$
\max_{s\in[0:t]}\max_{\ell\in[n]}\left\{\left|\mathsf{F}_{s,\ell}(0)\right|+\max_{q=1,2}\left|\mathsf{F}_{s,\ell}^{(q)}\right|\right|_{\infty}\right\}\leq\Lambda.
$$

For any p \geq 1*, there exists some* C_p $>$ 0 *such that for* $t \leq n/(C_p \log n)$ *,*

$$
\mathbb{E}^{1/p}\left[||\partial_{ij}z^{(t)}||^p|z^{(0)}\right]\leq (K\Lambda)^{C_pt}\sqrt{\log n}\cdot (1+||z^{(0)}||_{\infty}),
$$

and for $t \leq \log n/C_p$ *,*

$$
\mathbb{E}^{1/p} \left[\max_{i,j,\ell \in [n], \ell \neq i,j} \left(\sqrt{n} |\partial_{ij} z_{\ell}^{(s)}| \right)^p \Big| z^{(0)} \right] \le \left((K\Lambda)^t \log n \cdot (1 + \| z^{(0)} \|_{\infty}) \right)^{C_p t}.
$$

Proof. (1). By the definition of the AMP recursion [\(2.1\)](#page-7-2), we have

$$
\partial_{ij} z^{(t+1)} = \Delta_{ij} \mathsf{F}_t(z^{(t)}) + A(\mathsf{F}'_t(z^{(t)}) \circ \partial_{ij} z^{(t)}) - \mathsf{B}_t(\mathsf{F}'_{t-1}(z^{(t-1)}) \circ \partial_{ij} z^{(t-1)}).
$$

Consequently,

$$
\|\partial_{ij} z^{(t+1)}\| \leq |\mathsf{F}_{t,i}(z_i^{(t)})| + |\mathsf{F}_{t,j}(z_j^{(t)})| + \Lambda \|A\|_{\text{op}} \|\partial_{ij} z^{(t)}\| + (K\Lambda)^2 \|\partial_{ij} z^{(t-1)}\|
$$

$$
\leq (K\Lambda)^2 (1 + ||A||_{op}) (||\partial_{ij} z^{(t)}|| \vee ||\partial_{ij} z^{(t-1)}||) + 2\Lambda (1 + ||z^{(t)}||_{\infty}).
$$

Iterating the bound, for any $p \ge 1$, there exists some $C_p > 0$ such that

$$
\mathbb{E}^{1/p} \|\partial_{ij} z^{(t+1)}\|^p \le C_p \, \mathbb{E}^{1/2p} \left(K\Lambda(1 + \|A\|_{\text{op}}) \right)^{C_p t} \max_{s \in [0:t]} \mathbb{E}^{1/2p} (1 + \|z^{(s)}\|_{\infty})^{2p}.
$$

The claim follows from Proposition [4.2.](#page-19-0)

(2). For any $u \in [t]$, we let $\mathfrak{A}_{1}^{(u)}$ $y_1^{(u)} \equiv A, \mathfrak{A}_2^{(u)}$ $\mathbf{B}_{2}^{(u)} \equiv \mathbf{B}_{u+1}$. Note that for $\ell \neq i, j, \partial_{ij} z_{\ell}^{(t+1)}$ consists of at most $(K\Lambda)^{ct}$ terms of the form

$$
\left\langle e_i, \prod_{k \in [s]} \left(\mathfrak{A}^{(t_k)}_{\tau_{t_k}} \mathsf{D}^{(t_k)} \right) A e_j \right\rangle, \tag{4.5}
$$

where $s \in [0 : t]$, $1 \le t_s < t_{s-1} < \cdots < t_1 \le t$, $\tau_{t_k} \in \{1, 2\}$ with $\tau_{t_1} = 1$, and the product $\prod_{k \in [s]}$ is taken in the order from 1 to *s*. Consequently, it suffices to derive the moment estimate for the above terms.

Suppose $s \ge 1$ (the case $s = 0$ is trivial). For $r \in [s]$, let

$$
\mathcal{A}_{r} \equiv \max_{i \neq j \in [n]} \max_{w \in [r]} \max_{\substack{1 \leq t_w < t_{w-1} < \dots < t_1 \leq t, \\ \tau_{t_k} \in \{1, 2\}, \tau_{t_1} = 1}} \left| \left\langle e_i, \prod_{k \in [w]} (\mathfrak{A}_{\tau_{t_k}}^{(t_k)} \mathsf{D}^{(t_k)}) A e_j \right\rangle \right|.
$$

Let us fix $i \neq j \in [n]$, and let $\mathcal{P} \equiv \{i, j\}$. We first replace $D^{(\cdot)}$ by $D^{(\cdot)}_{[-]}$ $\binom{(\cdot)}{[-\mathcal{P}]}$, where $\mathsf{D}_{\mathsf{r_-}}^{(\cdot)}$ $\sum_{[\text{-}P]}$ \equiv $\mathfrak{D}_{F'(\text{z}_{[\text{-}P]}^{(\cdot)})}$. Let $\Delta D_{[\text{-}]}^{(\cdot)}$ $\frac{(\cdot)}{[-\mathcal{P}]} = D^{(\cdot)} - D^{(\cdot)}_{[-]}$ $\binom{(\cdot)}{(-\mathcal{P})}$. Then for any sequence $1 \leq t_s$ $t_{s-1} < \cdots < t_1 \leq t$ and $\tau_{t_k} \in \{1, 2\}$, $\langle e_i, A D^{(t_1)} \mathfrak{A}^{(t_2)}_{\tau_2} D^{(t_2)} \cdots \mathfrak{A}^{(t_s)}_{\tau_{t_s}} D^{(t_s)} A e_j \rangle - \langle e_i, A D^{(t_1)}_{\{-\mathcal{F}\}}$ $\left\{ \frac{(t_1)}{[-P]} \mathfrak{A}_{\tau_{t_2}}^{(t_2)} \mathsf{D}^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}}^{(t_s)} \mathsf{D}^{(t_s)} A e_j \right\}$ = $\overline{\nabla}$ *k*∈[*n*] A_{ik} ∆D $_{\mathsf{I} - 4}^{(t_1)}$ $\left[\begin{matrix} (t_1) \\ (-\mathcal{P}], k k \end{matrix}\right]$ $\left(\mathfrak{A}^{(t_2)}_{{\tau}_{t_2}} \mathsf{D}^{(t_2)} \cdots \mathfrak{A}^{(t_s)}_{{\tau}_{t_s}} \mathsf{D}^{(t_s)} A \right)_{kj}$ \leq $||A_i|| \cdot \Lambda ||z^{(t_1)} - z^{(t_1)}_{[-\mathcal{I}]}$ $\int_{[-\mathcal{P}]}^{(t_1)} || \cdot (K\Lambda)^{ct} \cdot \mathcal{A}_{s-1} + |A_{ij}| (K\Lambda ||A||_{op})^{ct}.$

Repeating the procedure of replacing one $D^{(\cdot)}$ factor by a corresponding $D^{(\cdot)}_{I-}$ $\int_{[-\mathcal{P}]}^{(+)}$ factor till the end, with $||A||_{\infty} \equiv \max_{i,j \in [n]} |A_{ij}|$, we have

$$
\left| \langle e_i, A D^{(t_1)} \mathfrak{A}^{(t_2)}_{\tau_{t_2}} D^{(t_2)} \cdots \mathfrak{A}^{(t_s)}_{\tau_{ts}} D^{(t_s)} A e_j \rangle - \langle e_i, A D^{(t_1)}_{[-\mathcal{P}]} \mathfrak{A}^{(t_2)}_{\tau_{t_2}} D^{(t_2)}_{[-\mathcal{P}]} \cdots \mathfrak{A}^{(t_s)}_{\tau_{ts}} D^{(t_s)}_{[-\mathcal{P}]} A e_j \rangle \right|
$$

\n
$$
\leq (K\Lambda (1 + ||A||_{op}))^{ct} \cdot \left(\max_{s \in [t]} \|z^{(s)} - z^{(s)}_{[-\mathcal{P}]} \| \cdot \mathcal{A}_{s-1} + ||A||_{\infty} \right). \tag{4.6}
$$

Next we replace $\mathfrak{A}_{\tau_u}^{(u)}$ with $\mathfrak{A}_{\tau_u}^{(u)}$ τ_{u} ,[- φ], where $A_{[-\varphi]} \equiv (A_{k\ell} \mathbf{1}_{k,\ell \notin \varphi})$ and $B_{\cdot,[-\varphi]} \equiv B$. Then it is easy to see

$$
\Delta \mathfrak{A}^{(u)}_{\tau_u,[-\mathcal{P}]} \equiv \mathfrak{A}^{(u)}_{\tau_u} - \mathfrak{A}^{(u)}_{\tau_u,[-\mathcal{P}]} = \mathbf{1}_{\tau_u=1} \sum_{(a,b)} \left(e_a e_a^\top A + Ae_a e_a^\top - A_{ab} e_a e_b^\top - A_{aa} e_a e_a^\top \right),
$$

where the summation is running over $(a, b) = \{(i, j), (j, i)\}$. Then

$$
\left| \langle e_i, A D_{[-\mathcal{P}]}^{(t_1)} \mathfrak{A}_{\tau_{t_2}}^{(t_2)} D_{[-\mathcal{P}]}^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}}^{(t_s)} D_{[-\mathcal{P}]}^{(t_s)} A e_j \rangle - \langle e_i, A D_{[-\mathcal{P}]}^{(t_1)} \mathfrak{A}_{\tau_{t_2}, [-\mathcal{P}]} D_{[-\mathcal{P}]}^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}}^{(t_s)} D_{[-\mathcal{P}]}^{(t_s)} A e_j \rangle \right|
$$

\n
$$
\leq (K\Lambda (1 + ||A||_{op}))^{ct} \cdot \left(||A||_{\infty} + \max_{w \in [s-1]} \max_{\substack{1 \leq t_w < t_w - 1 < \cdots < t_1 \leq t, \\ \tau_{t_k} \in \{1, 2\}, \tau_{t_1} = 1}} \left| \langle e_i, \prod_{k \in [w]} (\mathfrak{A}_{\tau_{t_k}}^{(t_k)} D_{[-\mathcal{P}]}^{(t_k)}) A e_j \rangle \right| \right)
$$

$$
\leq (K\Lambda(1+\|A\|_{\text{op}}))^{ct}\cdot\Big[(1+\max_{s\in[t]}|z^{(s)}-z^{(s)}_{[-\mathcal{P}]}||)\cdot\mathcal{A}_{s-1}+||A||_{\infty}\Big].
$$

Here in the last inequality we used [\(4.6\)](#page-21-0) to reduce the estimate to \mathcal{A}_{s-1} . Repeating the procedure of replacing one $\mathfrak{A}^{(\cdot)}$ factor by a corresponding $\mathfrak{A}^{(\cdot)}_{\cdot, \cdot}$ \cdot ,[– φ] factor till the end, we have

$$
\left| \langle e_i, A D_{[-\mathcal{P}]}^{(t_1)} \mathfrak{A}_{\tau_{t_2}}^{(t_2)} D_{[-\mathcal{P}]}^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}}^{(t_s)} D_{[-\mathcal{P}]}^{(t_s)} A e_j \rangle \right|
$$

\n
$$
- \langle e_i, A D_{[-\mathcal{P}]}^{(t_1)} \mathfrak{A}_{\tau_{t_2}, [-\mathcal{P}]}^{(t_2)} D_{[-\mathcal{P}]}^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}, [-\mathcal{P}]}^{(t_s)} D_{[-\mathcal{P}]}^{(t_s)} A e_j \rangle \right|
$$

\n
$$
\leq (K\Lambda (1 + ||A||_{op}))^{ct} \cdot \left[(1 + \max_{s \in [t]} ||z^{(s)} - z_{[-\mathcal{P}]}^{(s)} ||) \cdot \mathcal{A}_{s-1} + ||A||_{\infty} \right].
$$
 (4.7)

Combining [\(4.6\)](#page-21-0)-[\(4.7\)](#page-22-0), Proposition [4.2,](#page-19-0) and standard concentration estimates, for *t* ≤ *n*, uniformly in $i \neq j \in [n]$, with probability at least $1 - cn^{-D}$,

$$
\left| \langle e_i, A \mathsf{D}^{(t_1)} \mathfrak{A}^{(t_2)}_{\tau_{t_2}} \mathsf{D}^{(t_2)} \cdots \mathfrak{A}^{(t_s)}_{\tau_{t_s}} \mathsf{D}^{(t_s)} A e_j \rangle \right|
$$

\n
$$
- \langle e_i, A \mathsf{D}^{(t_1)}_{[-\mathcal{P}]} \mathfrak{A}^{(t_2)}_{\tau_{t_2}, [-\mathcal{P}]} \mathsf{D}^{(t_2)}_{[-\mathcal{P}]} \cdots \mathfrak{A}^{(t_s)}_{\tau_{t_s}, [-\mathcal{P}]} \mathsf{D}^{(t_s)}_{[-\mathcal{P}]} A e_j \rangle \right|
$$

\n
$$
\leq (K\Lambda)^{ct} \sqrt{\log n} \cdot (1 + ||z^{(0)}||_{\infty}) \cdot (\mathcal{A}_{s-1} + n^{-1/2}). \tag{4.8}
$$

On the other hand, as A_i , A_j are independent of $\mathfrak{A}^{(\cdot)}$ $\sum_{j=1}^{(\cdot)}$ and D $\sum_{j=1}^{(\cdot)}$ $\binom{(\cdot)}{[-\mathcal{P}]}$, using Hanson-Wright inequality, with probability at least $1 - cn^{-D}$,

$$
\left| \langle e_i, AD_{[-\mathcal{P}]}^{(t_1)} \mathfrak{A}_{\tau_{t_2}, [-\mathcal{P}]}^{(t_2)} D_{[-\mathcal{P}]}^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}, [-\mathcal{P}]}^{(t_s)} D_{[-\mathcal{P}]}^{(t_s)} A e_j \rangle \right|
$$

=
$$
\left| A_i^\top (D_{[-\mathcal{P}]}^{(t_1)} \mathfrak{A}_{\tau_{t_2}, [-\mathcal{P}]}^{(t_2)} D_{[-\mathcal{P}]}^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}, [-\mathcal{P}]}^{(t_s)} D_{[-\mathcal{P}]}^{(t_s)} A_j \right| \leq (K\Lambda)^{ct} \log n \cdot n^{-1/2}.
$$
 (4.9)

Combining [\(4.8\)](#page-22-1) and [\(4.9\)](#page-22-2), for $t \le n$, uniformly in $i \ne j \in [n]$, with probability at least $1 - cn^{-D}$,

$$
\left| \langle e_i, A D^{(t_1)} \mathfrak{A}_{\tau_{t_2}}^{(t_2)} D^{(t_2)} \cdots \mathfrak{A}_{\tau_{t_s}}^{(t_s)} D^{(t_s)} A e_j \rangle \right| \leq (K\Lambda)^{ct} \log n \cdot (1 + ||z^{(0)}||_{\infty}) \cdot (\mathcal{A}_{s-1} + n^{-1/2}).
$$

Now by a union bound via cardinality calculations, we conclude that for $t \leq$ $\log n/c$, with probability at least $1 - cn^{-D}$,

$$
\mathcal{A}_s \le (K\Lambda)^{ct} \log n \cdot (1 + ||z^{(0)}||_{\infty}) \cdot (\mathcal{A}_{s-1} + n^{-1/2}).
$$

Iterating the above bound shows that for $t \leq \log n/c$, with probability at least $1 - cn^{-D}$,

$$
\mathcal{A}_s \le ((K\Lambda)^t \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct} \cdot n^{-1/2}.
$$

The moment estimate for $\mathbb{E}^{1/p}$ [$|(4.5)|^p$ $|(4.5)|^p$ $|(4.5)|^p$] follows from the above high probability control, and the simple apriori estimate $|(4.5)| \le (K\Lambda ||A||_{op})^{ct}$ $|(4.5)| \le (K\Lambda ||A||_{op})^{ct}$ $|(4.5)| \le (K\Lambda ||A||_{op})^{ct}$. — Первой Станингии II <mark>п</mark>

4.3. The trace control. Recall the $D^{(.)}$ defined in [\(4.1\)](#page-18-3). Consider a sequence of matrices $M_0^{(t)}$ $\stackrel{(t)}{0},$ $M_1^{(t)}$ $\binom{t}{1}, \ldots, M_t^{(t)}$ $A_t^{(t)}$ ∈ ℝ^{*n*×*n*} defined recursively as follows: $M_{-1}^{(t)}$ $\binom{(t)}{-1} = 0_{n \times n},$ $M_0^{(t)}$ $b_0^{(t)} = D^{(t)}$, and for $s \in [t]$,

$$
M_s^{(t)} \equiv (M_{s-1}^{(t)}A - M_{s-2}^{(t)}B_{t-s+1})D^{(t-s)}.
$$
\n(4.10)

The following proposition provides the key trace control for $M_s^{(t)}$.

Proposition 4.4. *Suppose the following hold for some K,* $\Lambda \geq 2$ *.*

(1) max_{*i*, *j*∈[*n*_]| V *i*_j| ≤ *K*.} *(2)* **F**_{*s*, ℓ} ∈ $\overline{C^2}(\mathbb{R})$ *for all s* ∈ [0 : *t*], ℓ ∈ [*n*]*, and*

$$
\max_{s\in[0:t]}\max_{\ell\in[n]}\left\{\left|\mathsf{F}_{s,\ell}(0)\right|+\max_{q=1,2}\left|\mathsf{F}_{s,\ell}^{(q)}\right|\right|_{\infty}\right\}\leq\Lambda.
$$

Then there exists some universal constant $c_0 > 0$ *such that for* $t \leq \log n/c_0$,

$$
\max_{s\in[t]} \frac{1}{n} \Big| \mathbb{E} \big[\operatorname{tr}(M_s^{(t)}) |z^{(0)} \big] \Big| \le (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot n^{-1/2}.
$$

4.3.1. *The GOE and* $F = id$ *case.* Before presenting the proof for the above proposition, let us examine the simplest possible case, where $V = 1_{n \times n}$, and $F_t = id$ is the coordinate-wise identity map. In this case, $D^{(\cdot)} = B = I_n$, and the matrix recursion [\(4.10\)](#page-22-3) reduces to

$$
M_s = 2M_{s-1}A_0 - M_{s-2}, \quad M_0 = I_n, \quad M_{-1} = 0_{n \times n}, \tag{4.11}
$$

where $A_0 \equiv A/2$. Let $U_s(\cdot)$ be the Chebyshev polynomials of the second kind, defined recursively via

$$
U_s(x) = 2xU_{s-1}(x) - U_{s-2}(x), \quad U_0(x) \equiv 1, \quad U_{-1}(x) \equiv 0. \tag{4.12}
$$

The structural similarity between [\(4.11\)](#page-23-1) and [\(4.12\)](#page-23-2) leads to the following observation. Let $\mu_n \equiv n^{-1} \sum_{i \in [n]} \delta_{\lambda_i(A_0)}$ be the empirical spectral distribution of A_0 . As A_0 is approximately (scaled) GOE, μ_n is close to the (scaled) semicircle law μ_∞ on $[-1, 1]$: $\mu_n(dx) \approx \mu_\infty(dx) = (2/\pi) \sqrt{1 - x^2} dx$. Now using the fact that $\{U_s\}_{s \in \mathbb{Z}_{\geq 0}}$ are orthogonal polynomials in $L_2(\mu_\infty)$, we have for any $s \ge 1$,

$$
\frac{1}{n}\operatorname{tr}(M_s) = \frac{1}{n}\sum_{i\in[n]}U_s(\lambda_i(A_0)) = \int_{-1}^1 U_s(x)\mu_n(\mathrm{d}x) \approx \int_{-1}^1 U_s(x)U_0(x)\mu_\infty(\mathrm{d}x) = 0.
$$

Unfortunately, the essential use of the linearity of F_t and the GOE property of *A* in the above (rough) argument makes it less amenable to further generalizations to cover the general cases in Proposition [4.4.](#page-23-0) Note that even without the cancellation mechanism for orthogonal polynomial, we can still write $M_s^{(t)}$ as a polynomial of *A*, B 's and $D^{(\cdot)}$'s, by solving the recursion [\(4.10\)](#page-22-3). For instance, $M_s^{(t)}$ can be written as the sum of matrix polynomials that take the form of $\pm D^{(t)}$ followed by a product of $AD^{(1)}$ factors and $B.D^{(1)}$ factors. However, it appears difficult to explore the cancellation mechanism systematically among these terms in the polynomial due to the appearance of a large number of the random A factors and $D^{(*)}$ factors, and also the iteration-dependence of B and $D^{(*)}$ factors.

In the proof below, instead of estimating the trace of this complicated matrix polynomial at once, we turn to a recursive estimate based on the recursion [\(4.10\)](#page-22-3). Namely, we eliminate the high complexity randomness in $M_s^{(t)}$ step by step via Gaussian integration by parts, and eventually arrive at simple linear and quadratic forms which can be estimated via Gaussian concentration directly.

4.3.2. *Some further notation and preliminary facts.* For generic matrices $M_0, M_1, M_2, M_3 \in \mathbb{R}^{n \times n}$, we write

$$
\langle \mathsf{M}_1 \partial .. \mathsf{M}_0 \mathsf{M}_2 , \mathsf{M}_3 \rangle \equiv \sum_{i,j \in [n]} \big(\mathsf{M}_1 \partial_{ij} \mathsf{M}_0 \mathsf{M}_2 \big)_{ij} (\mathsf{M}_3)_{ij}.
$$

Recall $\Delta_{ij} = \partial_{ij}A$ defined in [\(4.4\)](#page-20-0). We have the derivative formula: for *i*, *j* ∈ [*n*], and $r \in [t]$,

$$
\partial_{ij}M_r^{(t)} = (\partial_{ij}M_{r-1}^{(t)}A - \partial_{ij}M_{r-2}^{(t)}B_{t-r+1})D^{(t-r)} + M_{r-1}^{(t)}\Delta_{ij}D^{(t-r)} + (M_{r-1}^{(t)}A - M_{r-2}^{(t)}B_{t-r+1})\mathfrak{D}_{\mathsf{F}_{i-r}^{(2)}(z^{(t-r)})}\mathfrak{D}_{\partial_{ij}z^{(t-r)}}.
$$
\n(4.13)

For a fixed $s \in [t]$, we consider another sequence of matrices $N_0^{(s)}$ $N_s^{(s)}, \ldots, N_s^{(s)} \in \mathbb{R}^{n \times n},$ defined recursively as follows: $N_{-1}^{(s)}$ $\binom{s}{-1} \equiv 0_{n \times n}, N_0^{(s)}$ $\binom{S}{0}$ $\equiv I_n$, and for $u \in [s]$,

$$
N_u^{(s)} \equiv A D^{(t-s+u)} N_{u-1}^{(s)} - B_{t-s+u} D^{(t-s+u-1)} N_{u-2}^{(s)}.
$$
 (4.14)

Then following derivative formula holds:

$$
\partial_{ij} N_u^{(s)} = (AD^{(t-s+u)} \partial_{ij} N_{u-1}^{(s)} - B_{t-s+u} D^{(t-s+u-1)} \partial_{ij} N_{u-2}^{(s)}) + \Delta_{ij} D^{(t-s+u)} N_{u-1}^{(s)}
$$

+ $A \mathfrak{D}_{\mathsf{F}_{t-s+u}^{(2)}(z^{(t-s+u)})} \mathfrak{D}_{\partial_{ij} z^{(t-s+u)}} N_{u-1}^{(s)}$
- $B_{t-s+u} \mathfrak{D}_{\mathsf{F}_{t-s+u-1}^{(2)}(z^{(t-s+u-1)})} \mathfrak{D}_{\partial_{ij} z^{(t-s+u-1)}} N_{u-2}^{(s)}$. (4.15)

The sequence of matrices $\{N^{(\cdot)}\}$ can be viewed, in a certain sense, a reversed recursion to $\{M^{(*)}\}$, the role of which will be clear in the proof in the next subsection.

We need the following two preliminary results on $\{M_{\cdot}^{(1)}\}$ and $\{N_{\cdot}^{(1)}\}$.

Lemma 4.5. *Assume the same conditions as in Proposition [4.4.](#page-23-0) There exists some universal* $c_0 > 0$ *such that for* $t \leq n/(c_0 \log n)$ *and any* $q \geq 1$ *,*

$$
n^{-1/2} \left\{ \mathbb{E}^{1/q} ||M_{\cdot}^{(t)}||_F^q \vee \mathbb{E}^{1/q} ||N_{\cdot}^{(s)}||_F^q \right\} + (\mathbb{E}^{1/q} ||M_{\cdot}^{(t)}||_{op}^q \vee \mathbb{E}^{1/q} ||N_{\cdot}^{(s)}||_{op}^q) \le C_q (K\Lambda)^{c_0 t}.
$$
 (4.16)

Here the constant $C_q > 0$ *may depend on q.*

Proof. In view of the recursions in [\(4.10\)](#page-22-3) and [\(4.14\)](#page-24-0), we have an easy estimate

$$
n^{-1/2}\{||M_{\cdot}^{(t)}||_F \vee ||N_{\cdot}^{(s)}||_F\} + (||M_{\cdot}^{(t)}||_{op} \vee ||N_{\cdot}^{(s)}||_{op}) \leq (K\Lambda(1 + ||A||_{op}))^{ct}.
$$

The claim follows by using the subgaussian tail of $||A||_{op}$, cf. Lemma [A.1.](#page-47-1) \Box

Lemma 4.6. *Assume the same conditions as in Proposition [4.4.](#page-23-0) There exists some universal* $c_0 > 0$ *such that for any two symmetric matrices* $A_1, A_2 \in \mathbb{R}^{n \times n}$,

$$
||M^{(t)}(A_1) - M^{(t)}(A_2)||_F \vee ||N^{(s)}(A_1) - N^{(s)}(A_2)||_F
$$

$$
\leq (K\Lambda(1 + ||z^{(0)}||_{\infty})(1 + ||A_1||_{op} + ||A_2||_{op}))^{c_0t^2} \cdot \sqrt{n}||A_1 - A_2||_{op}. \tag{4.17}
$$

Proof. We shall only prove the case for $M_s^{(t)}$; the other cases follow from minor modifications. Using the recursion [\(4.10\)](#page-22-3), we have

$$
||M_{s}^{(t)}(A_{1}) - M_{s}^{(t)}(A_{2})||_{F} \leq ||M_{s-1}^{(t)}(A_{1})A_{1}D^{(t-s)}(A_{1}) - M_{s-1}^{(t)}(A_{2})A_{2}D^{(t-s)}(A_{2})||_{F}
$$

+
$$
||M_{s-2}^{(t)}(A_1)B_{t-s+1}D^{(t-s)}(A_1) - M_{s-2}^{(t)}(A_2)B_{t-s+1}D^{(t-s)}(A_2)||_F \equiv (I) + (II).
$$

\nLet $\Gamma(A_1, A_2; c) \equiv (K\Lambda(1 + ||z^{(0)}||_{\infty})(1 + ||A_1||_{op} + ||A_2||_{op}))^c$. For (I) , we have
\n
$$
(I) \leq ||M_{s-1}^{(t)}(A_1) - M_{s-1}^{(t)}(A_2)||_F \cdot ||A_1D^{(t-s)}(A_1)||_{op}
$$
\n
$$
+ ||M_{s-1}^{(t)}(A_2)||_{op} \cdot ||A_1D^{(t-s)}(A_1) - A_2D^{(t-s)}(A_2)||_F
$$
\n
$$
\leq \Gamma(A_1, A_2; ct) \cdot (||M_{s-1}^{(t)}(A_1) - M_{s-1}^{(t)}(A_2)||_F
$$
\n
$$
+ ||A_1 - A_2||_F + ||D^{(t-s)}(A_1) - D^{(t-s)}(A_2)||_F
$$
\n
$$
\leq \Gamma(A_1, A_2; c_1 t) \cdot (||M_{s-1}^{(t)}(A_1) - M_{s-1}^{(t)}(A_2)||_F + \sqrt{n}||A_1 - A_2||_{op}).
$$

Here in the last inequality we used the estimate $\|D^{(t-s)}(A_1) - D^{(t-s)}(A_2)\|_F \leq \Lambda$. $||z^{(t-s)}(A_1) - z^{(t-s)}(A_2)||$ ≤ Γ(*A*₁, *A*₂; *ct*) · $\sqrt{n}||A_1 - A_2||_{op}$. Using the same argument, we may estimate (*II*):

$$
(II) \leq \Gamma(A_1, A_2; c_2 t) \cdot (||M_{s-2}^{(t)}(A_1) - M_{s-2}^{(t)}(A_2)||_F + \sqrt{n}||A_1 - A_2||_{op}).
$$

Combining the estimates, there exists some universal $c_0 > 0$ such that

$$
\max_{u \in [0:s]} ||M_u^{(t)}(A_1) - M_u^{(t)}(A_2)||_F
$$
\n
$$
\leq \Gamma(A_1, A_2; c_0 t) \cdot \Big(\max_{u \in [0:s-1]} ||M_u^{(t)}(A_1) - M_u^{(t)}(A_2)||_F + \sqrt{n} ||A_1 - A_2||_{\text{op}} \Big).
$$

Iterating the bound and using the initial condition $||M_0^{(t)}||$ $M_0^{(t)}(A_1) - M_0^{(t)}$ $0^{(t)}(A_2)||_F =$ $||D^{(t)}(A_1) - D^{(t)}(A_2)||_F \le \Gamma(A_1, A_2; c't) \cdot \sqrt{n} ||A_1 - A_2||_{op}$ to conclude. □

4.3.3. *Proof of Proposition [4.4.](#page-23-0)* First, for $3 \leq s \leq t$, we show that the normalized trace of $M_s^{(t)}$ can be reduced to that of $\{N_s^{(t)}\}$, the most complicated one being $N_{s-1}^{(s)}$ The following formulation also allows for the base case $s = 1, 2$. *s*−2 .

Lemma 4.7. *Assume the same conditions as in Proposition [4.4.](#page-23-0) There exists some universal constant* $c_0 > 0$ *such that for any fixed* $d_0 \in \mathbb{R}^n$ *and an index set* $\mathcal{T} \subset$ $[-1 : t]$ *allowing for multiplicity, and any* $1 \leq s \leq t \leq \log n/c_0$ *,*

$$
\frac{1}{n} \left| \mathbb{E} \operatorname{tr} \left(\mathfrak{D}_{d_0} M_s^{(t)} \mathsf{D}^{(\mathcal{T})} \right) \right| \le (K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^2} \cdot n^{-1/2} \tag{4.18}
$$
\n
$$
+ (K \Lambda)^{c_0 t^2} \sum_{m \in [s-2]} \max_{d_{[c_0, m]} \in \{V^2_{[1, m]}, b_{[0, t]}, 1_n\}, \atop \mathcal{S} \in [-1: t]^{c_0 m}} \frac{1}{n} \left| \mathbb{E} \operatorname{tr} \left(\mathfrak{D}_{d_0 \circ \otimes_{q \in [c_0, m]} d_q} N_{s-m-1}^{(s-m+1)} \mathsf{D}^{(\mathcal{T})} \mathsf{D}^{(\mathcal{S})} \right) \right|.
$$

Here we write $D^{(\mathcal{T})} \equiv \prod_{s \in \mathcal{T}} D^{(s)}$ *and* $D^{(-1)} \equiv I_n$ *for notational convenience.*

Proof. (Step 1). First consider the case $s = 1, 2$. Using the recursion [\(4.10\)](#page-22-3),

$$
M_1^{(t)} = D^{(t)}AD^{(t-1)}, \quad M_2^{(t)} = D^{(t)}(AD^{(t-1)}A - B_{t-1})D^{(t-2)}.
$$

For $s = 1$, a trivial bound suffices:

$$
\frac{1}{n} \left| \mathbb{E} tr(M_1^{(t)}) \right| \leq \frac{1}{n} \sum_{i \in [n]} \mathbb{E} \left| F'_{t,i}(z_i^{(t)}) F'_{t-1,i}(z_i^{(t-1)}) A_{ii} \right| \leq \frac{(K\Lambda)^c}{\sqrt{n}}.
$$

For $s = 2$, note that

$$
\frac{1}{n} \left| \mathbb{E} \, \text{tr}(M_2^{(t)}) \right| \le \frac{\Lambda^2}{n} \sum_{k \in [n]} \mathbb{E} |(A \mathsf{D}^{(t-1)} A)_{kk} - b_{t-1,k}|. \tag{4.19}
$$

For each $k \in [n]$,

$$
\mathbb{E}|(AD^{(t-1)}A)_{kk} - b_{t-1,k}|
$$
\n
$$
\leq \mathbb{E}\left|\sum_{j\in[n]} (A_{kj}^2 - V_{kj}^2/n)\mathsf{F}'_{t-1,j}(z_j^{(t-1)})\right| + \mathbb{E}\left|\frac{1}{n}\sum_{j\in[n]} V_{kj}^2(\text{id} - \mathbb{E})\mathsf{F}'_{t-1,j}(z_j^{(t-1)})\right|
$$
\n
$$
\equiv \mathfrak{Q}_1^{(t)} + \mathfrak{Q}_2^{(t)}.
$$
\n(4.20)

For $\mathfrak{Q}_1^{(t)}$ $_1^{(t)}$, we have for $t \leq \log n/c$,

$$
\mathfrak{Q}_{1}^{(t)} \leq \mathbb{E} \left| \sum_{j \in [n]} (A_{kj}^{2} - V_{kj}^{2}/n)(F'_{t-1,j}(z_{j}^{(t-1)}) - F'_{t-1,j}(z_{[-k],j}^{(t-1)})) \right|
$$

+
$$
\mathbb{E} \left| \sum_{j \in [n]} (A_{kj}^{2} - V_{kj}^{2}/n)F'_{t-1,j}(z_{[-k],j}^{(t-1)}) \right|
$$

$$
\leq (K\Lambda)^{c} \cdot n^{-1/2} (\mathbb{E}^{1/2} || z^{(t-1)} - z_{[-k]}^{(t-1)} ||^{2} + 1)
$$

$$
\leq (K\Lambda)^{ct} (1 + ||z^{(0)}||_{\infty}) \sqrt{\log n} \cdot n^{-1/2}.
$$
 (4.21)

Here in the last line we used Proposition [4.2.](#page-19-0)

For $\mathfrak{Q}_2^{(t)}$ (*t*)</sup>, let *H*₀(*A*) ≡ *n*⁻¹ ∑_{*j*∈[*n*]} V_{kj}^2 **F**_{*t*−1, *j*}($z_j^{(t-1)}$ (*A*)). Then for any two symmetric matrices $A_1, A_2 \in \mathbb{R}^{n \times n}$,

$$
|H_0(A_1) - H_0(A_2)| \le (K\Lambda)^c \cdot n^{-1/2} ||z^{(t-1)}(A_1) - z^{(t-1)}(A_2)||
$$

$$
\le (K\Lambda(1 + ||A_1||_{op} + ||A_2||_{op}))^{ct} \cdot ||A_1 - A_2||_{op}.
$$

Moreover,

$$
|H_0(A)| \le (K\Lambda)^c \cdot (1 + n^{-1/2} ||z^{(t-1)}||) \le (K\Lambda(1 + ||A||_{op}))^{ct} (1 + ||z^{(0)}||_{\infty}).
$$

By applying the Gaussian concentration Lemma [A.2,](#page-48-0) we have for $t \leq \log n/c$,

$$
\mathfrak{Q}_2^{(t)} \le \text{Var}^{1/2}(H_0(A)) \le (K\Lambda)^{ct}(1 + ||z^{(0)}||_{\infty})\log n \cdot n^{-1/2}.
$$
 (4.22)

Combining $(4.19)-(4.22)$ $(4.19)-(4.22)$, we have

1

$$
\frac{1}{n} \left| \mathbb{E} \, \text{tr}(M_2^{(t)}) \right| \le (K\Lambda)^{ct} (1 + ||z^{(0)}||_{\infty}) \log n \cdot n^{-1/2}.
$$

The claimed, slightly more general form follows from minor modifications. (Step 2). Next assume $s \geq 3$. For $r \in [s-1]$, let

$$
R_r^{(s)} \equiv \frac{1}{n^2} \mathbb{E} \left\langle D^{(t-s)} M_{r-1}^{(t)} \Delta_{\cdot} D^{(t-r)} N_{s-1-r}^{(s)}, V \circ V \right\rangle
$$

+
$$
\frac{1}{n^2} \mathbb{E} \left\langle D^{(t-s)} (M_{r-1}^{(t)} A - M_{r-2}^{(t)} B_{t-r+1}) \mathfrak{D}_{F_{t-r}^{(2)}(z^{(t-r)})} \mathfrak{D}_{\partial_{\cdot} z^{(t-r)}} N_{s-1-r}^{(s)}, V \circ V \right\rangle.
$$

By Gaussian integration by parts,

$$
\frac{1}{n} \mathbb{E} tr(M_s^{(t)}) = \frac{1}{n} \sum_{i,j \in [n]} \mathbb{E} D_{ii}^{(t-s)} M_{s-1,ij}^{(t)} A_{ij} - \frac{1}{n} \mathbb{E} tr(D^{(t-s)} M_{s-2}^{(t)} B_{t-s+1})
$$
\n
$$
= \frac{1}{n^2} \langle \mathbb{E} D^{(t-s)} \partial ... M_{s-1}^{(t)}, V \circ V \rangle - \frac{1}{n} \mathbb{E} tr(D^{(t-s)} M_{s-2}^{(t)} B_{t-s+1})
$$
\n
$$
+ \frac{1}{n^2} \sum_{i,j \in [n]} \mathbb{E} F_{t-s,i}^{(2)} (z_i^{(t-s)}) \partial_{ij} z_i^{(t-s)} M_{s-1,ij}^{(t)} V_{ij}^2.
$$

Using [\(4.13\)](#page-24-1) with $r = s - 1$, the first term above equals

$$
\frac{1}{n^2} \mathbb{E} \left\langle D^{(t-s)} \Big(\partial \ldots M_{s-2}^{(t)} N_1^{(s)} - \partial \ldots M_{s-3}^{(t)} B_{t-s+2} D^{(t-s+1)} N_0^{(s)} \Big), V \circ V \right\rangle + R_{s-1}^{(s)}.
$$

Using [\(4.13\)](#page-24-1) with $r = s - 2$, the first term of the above display equals

$$
\frac{1}{n^2} \mathbb{E} \left\langle D^{(t-s)} \Big(\partial_{\cdot \cdot} M^{(t)}_{s-3} N^{(s)}_2 - \partial_{\cdot \cdot} M^{(t)}_{s-4} \mathsf{B}_{t-s+3} D^{(t-s+2)} N^{(s)}_1 \Big), V \circ V \right\rangle + R^{(s)}_{s-2}.
$$

Consequently, iterating this procedure until (4.13) is applied with $r = 1$, we obtain

$$
\frac{1}{n} \mathbb{E} \operatorname{tr}(M_s^{(t)}) = R_{s-1}^{(s)} + \dots + R_1^{(s)} - \frac{1}{n} \mathbb{E} \operatorname{tr} (\mathsf{D}^{(t-s)} M_{s-2}^{(t)} \mathsf{B}_{t-s+1}) \n+ \frac{1}{n^2} \sum_{i,j \in [n]} \mathbb{E} \mathsf{F}'_{t-s,i} (z_i^{(t-s)}) \mathsf{F}^{(2)}_{t,i} (z_i^{(t)}) \partial_{ij} z_i^{(t)} N_{s-1,ij}^{(s)} V_{ij}^2 \n+ \frac{1}{n^2} \sum_{i,j \in [n]} \mathbb{E} \mathsf{F}^{(2)}_{t-s,i} (z_i^{(t-s)}) \partial_{ij} z_i^{(t-s)} M_{s-1,ij}^{(t)} V_{ij}^2.
$$
\n(4.23)

(Step 2-(a)). We provide a bound for the last two terms in (4.23) : using the moment estimates in [\(4.16\)](#page-24-2) and Proposition [4.3,](#page-20-1) for $t \le \log n/c$,

Last two terms of [\(4.23\)](#page-27-0)

$$
\leq (K\Lambda)^c \cdot \max_{u \in [0:t]} \mathbb{E}^{1/2} \max_{i,j \in [n]} ||\partial_{ij} z^{(u)}||_{\infty}^2 \cdot n^{-1} \Big(\mathbb{E}^{1/2} ||M_{s-1}^{(t)}||_F^2 + \mathbb{E}^{1/2} ||N_{s-1}^{(s)}||_F^2 \Big)
$$

$$
\leq \frac{(K\Lambda)^{ct}}{\sqrt{n}} \log^c n \cdot (1 + ||z^{(0)}||_{\infty}).
$$
 (4.24)

(**Step 2-(b**)). We provide a bound for $R_r^{(s)}$ for $r \in [s-1]$. Note that

$$
|R_r^{(s)}| \leq \frac{1}{n^2} \left| \mathbb{E} \sum_{i,j \in [n]} \mathsf{D}_{ii}^{(t-s)} M_{r-1,ii}^{(t)} \mathsf{D}_{jj}^{(t-r)} N_{s-1-r,jj}^{(s)} V_{ij}^2 \right| + \frac{1}{n^2} \left| \mathbb{E} \sum_{i \neq j \in [n]} \mathsf{D}_{ii}^{(t-s)} M_{r-1,ij}^{(t)} \mathsf{D}_{ii}^{(t-r)} N_{s-1-r,j}^{(s)} V_{ij}^2 \right| + \left| \frac{1}{n^2} \mathbb{E} \sum_{i,j,\ell} \mathsf{D}_{ii}^{(t-s)} (M_{r-1}^{(t)} A - M_{r-2}^{(t)} \mathsf{B}_{t-r+1})_{i\ell} \mathsf{F}_{t-r,\ell}^{(2)} (z_{\ell}^{(t-r)}) \partial_{ij} z_{\ell}^{(t-r)} N_{s-1-r,\ell j}^{(s)} V_{ij}^2 \right| \equiv \mathfrak{R}_{r,1}^{(s)} + \mathfrak{R}_{r,2}^{(s)} + \mathfrak{R}_{r,3}^{(s)}.
$$
(4.25)

We shall handle the three terms $\mathfrak{R}_{r;1}^{(s)}$, $\mathfrak{R}_{r;2}^{(s)}$, $\mathfrak{R}_{r;3}^{(s)}$ separately.

Term $\mathfrak{R}_{r;3}^{(s)}$: By writing $P_r^{(t)} \equiv M_{r-1}^{(t)}$ $f_{r-1}^{(t)}A - M_{r-1}^{(t)}$ *r*−2 B*t*−*r*+¹ for notational simplicity,

$$
\begin{split} \mathfrak{R}^{(s)}_{r;3}&\leq \frac{(K\Lambda)^c}{n^2}\,\mathbb{E}\sum_{i,j,\ell}|P^{(t)}_{r,i\ell}||\partial_{ij}z^{(t-r)}_{\ell}||N^{(s)}_{s-1-r,\ell j}|\\ &\leq \frac{(K\Lambda)^c}{n^2}\,\mathbb{E}\Big[\max_{i,j\in[n],\ell\neq i,j}|\partial_{ij}z^{(t-r)}_{\ell}|\cdot n||P^{(t)}_r||_F||N^{(s)}_{s-1-r}||_F\\ &\qquad\qquad+\max_{i,j\in[n]}\|\partial_{ij}z^{(t-r)}\|_\infty\cdot n\bigg(||P^{(t)}_r||_{{\rm op}}||N^{(s)}_{s-1-r}||_F+||P^{(t)}_r||_F||N^{(s)}_{s-1-r}||_{{\rm op}}\bigg)\Big]. \end{split}
$$

Using the moment estimates in [\(4.16\)](#page-24-2) and Proposition [4.3,](#page-20-1) for $t \le \log n/c$,

$$
\mathfrak{R}_{r;3}^{(s)} \le (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct^2} \cdot n^{-1/2}.
$$
 (4.26)

Term $\mathfrak{R}_{r;2}^{(s)}$: By a simple Cauchy-Schwarz inequality and the moment estimates in [\(4.16\)](#page-24-2), for *t* ≤ log *n*/*c*,

$$
\mathfrak{R}_{r;2}^{(s)} \le (K\Lambda)^2 n^{-2} \cdot \mathbb{E}^{1/2} ||M_{r-1}^{(t)}||_F^2 \cdot \mathbb{E}^{1/2} ||N_{s-1-r}^{(s)}||_F^2 \le \frac{(K\Lambda)^{ct}}{n}.\tag{4.27}
$$

Term $\mathbb{R}_{r,1}^{(s)}$ with $r = 1$: Note that

$$
\mathfrak{R}_{1;1}^{(s)} \leq \Lambda \, \mathbb{E}^{1/2} \|M_0^{(t)}\|_{\text{op}}^2 \cdot \frac{1}{n} \max_{\ell \in [n]} \mathbb{E}^{1/2} \, \text{tr}^2 \big(\mathfrak{D}_{V_\ell^2} N_{s-2}^{(s)} \mathsf{D}^{(t-1)} \big).
$$

Clearly $\mathbb{E}^{1/2} || M_0^{(t)}$ $||_{0}^{(t)}||_{0}^{2} \leq \Lambda$. Let $H(A) \equiv n^{-1}$ tr $(\mathfrak{D}_{V_{\ell}^{2}}N_{s-1}^{(s)})$ ^{(*s*})</sup> D ^{(*t*−1)</sub>). Then using [\(4.17\)](#page-24-3),} for any two symmetric $A_1, A_2 \in \mathbb{R}^{n \times n}$,

$$
|H(A_1) - H(A_2)| \le n^{-1} \left| \operatorname{tr} \left(\mathfrak{D}_{V_\ell^2}(N_{s-2}^{(s)}(A_1) - N_{s-2}^{(s)}(A_2)) \mathsf{D}^{(t-1)}(A_1) \right) \right|
$$

+ $n^{-1} \left| \operatorname{tr} \left(\mathfrak{D}_{V_\ell^2} N_{s-2}^{(s)}(A_2) (\mathsf{D}^{(t-1)}(A_1) - \mathsf{D}^{(t-1)}(A_2)) \right) \right|$
 $\le (K\Lambda)^c \cdot n^{-1/2} ||N_{s-2}^{(s)}(A_1) - N_{s-2}^{(s)}(A_2)||_F$
+ $(K\Lambda)^c \cdot (1 + ||N_{s-2}^{(s)}(A_1)||_{op} + ||N_{s-2}^{(s)}(A_2)||_{op}) \cdot n^{-1/2} ||z^{(t-1)}(A_1) - z^{(t-1)}(A_2)||_F$
 $\le (K\Lambda(1 + ||z^{(0)}||_{\infty})(1 + ||A_1||_{op} + ||A_2||_{op}))^{c t^2} \cdot ||A_1 - A_2||_{op}.$

Moreover,

$$
|H(A)| \le (K\Lambda)^2 ||N_{s-2}^{(s)}||_{op} \le (K\Lambda(1 + ||A||_{op}))^{ct^2}.
$$

Now applying the Gaussian concentration Lemma [A.2,](#page-48-0) we have

$$
\text{Var}(H(A)) \le n^{-1} (K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty})^{ct^2}.
$$

Combining the above two displays, we have for $3 \leq s \leq t \leq \log n/c$,

$$
\mathfrak{R}_{1;1}^{(s)} \le \frac{(K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct^2}}{\sqrt{n}} + (K\Lambda)^{ct} \max_{\ell \in [n]} \frac{1}{n} \Big| \mathbb{E} \operatorname{tr} \big(\mathfrak{D}_{V_{\ell}^2} N_{s-2}^{(s)} \mathsf{D}^{(t-1)} \big) \Big| . \tag{4.28}
$$

Term $\mathfrak{R}_{r;1}^{(s)}$ with $r \in [2 : s - 1]$: Note that

$$
\mathfrak{R}_{r;1}^{(s)} \le \frac{1}{n^2} \bigg| \mathbb{E} \sum_{j \in [n]} \mathsf{D}_{jj}^{(t-r)} N_{s-1-r,jj}^{(s)} \cdot \text{tr} \left(\mathfrak{D}_{V_j^2} M_{r-1}^{(t)} \mathsf{D}^{(t-s)} \right) \bigg|
$$

$$
\leq \Lambda \, \mathbb{E}^{1/2} \| N_{s-1-r}^{(s)} \|_{\text{op}}^2 \cdot \frac{1}{n} \max_{\ell \in [n]} \mathbb{E}^{1/2} \, \text{tr}^2 \, (\mathfrak{D}_{V_\ell^2} M_{r-1}^{(t)} \mathsf{D}^{(t-s)}).
$$

From here, we may use arguments similarly as above to conclude that, for $3 \leq s \leq$ $t \leq \log n/c$,

$$
\mathfrak{R}_{r;1}^{(s)} \le \frac{(K\Lambda \log n(1+||z^{(0)}||_{\infty}))^{ct^2}}{\sqrt{n}} + (K\Lambda)^{ct} \max_{\ell \in [n]} \frac{1}{n} \Big| \mathbb{E} \operatorname{tr} \big(\mathfrak{D}_{V_{\ell}^2} M_{r-1}^{(t)} \mathsf{D}^{(t-s)} \big) \Big|. \tag{4.29}
$$

Combining [\(4.25\)](#page-27-1)-[\(4.29\)](#page-29-0), for $3 \le s \le t \le \log n/c$,

$$
|R_r^{(s)}| \le (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct^2} \cdot n^{-1/2}
$$

+ $(K\Lambda)^{ct} \max_{\ell \in [n]} \frac{1}{n} \mathbb{E} \text{tr} (\mathfrak{D}_{V_{\ell}^2} N_{s-2}^{(s)} \mathsf{D}^{(t-1)}) \Big| \mathbf{1}_{r=1}$
+ $(K\Lambda)^{ct} \max_{\ell \in [n]} \frac{1}{n} \mathbb{E} \text{tr} (\mathfrak{D}_{V_{\ell}^2} M_{r-1}^{(t)} \mathsf{D}^{(t-s)}) \Big| \mathbf{1}_{r \in [2:s-1]}.$

(Step 2-(c)). In view of [\(4.23\)](#page-27-0), we then conclude that for $3 \le s \le t \le \log n/c$,

$$
\frac{1}{n} \left| \mathbb{E} \operatorname{tr}(M_s^{(t)}) \right| \le (K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct^2} \cdot n^{-1/2} \n+ (K \Lambda)^{ct} \max_{d \in \{V_{[1:n]}^2, b_{[0:t]}\}} \frac{1}{n} \left| \mathbb{E} \operatorname{tr} (\mathfrak{D}_d N_{s-2}^{(s)} \mathsf{D}^{(t-1)}) \right| \n+ (K \Lambda)^{ct} \sum_{r \in [s-2]} \max_{d \in \{V_{[1:n]}^2, b_{[0:t]}\}} \frac{1}{n} \left| \mathbb{E} \operatorname{tr} (\mathfrak{D}_d M_r^{(t)} \mathsf{D}^{(t-s)}) \right|.
$$

Now repeating the above arguments for the term $r = s - 2$, $s - 3$, ..., $r \ge 3$ in the first summation, it follows that for $3 \leq s \leq t \leq \log n/c$,

$$
\begin{split} & \frac{1}{n} \Big| \mathbb{E} \, \text{tr}(M_s^{(t)}) \Big| \leq \big(K \Lambda \log n \cdot (1 + \|z^{(0)}\|_\infty)\big)^{ct^2} \cdot n^{-1/2} \\ & + (K \Lambda)^{ct} \sum_{r=1,2} \max_{d \in \{V_{[1:n]}^2, b_{[0:t]}\}} \frac{1}{n} \Big| \mathbb{E} \, \text{tr} \, \big(\mathfrak{D}_d M_r^{(t)} \mathsf{D}^{(t-s)}\big) \Big| \\ & + (K \Lambda)^{ct^2} \sum_{m \in [s-2]} \max_{d_{[cm]} \in \{V_{[1:n]}^2, b_{[0:t]}, 1_n\}} \frac{1}{n} \Big| \mathbb{E} \, \text{tr} \, \big(\mathfrak{D}_{\otimes_{q \in [cm]} d_q} N_{s-m-1}^{(s-m+1)} \mathsf{D}^{(S)}\big) \Big|. \end{split}
$$

The first summation term can be assimilated into the first term by similar arguments in Step 1. This proves the desired estimate [\(4.18\)](#page-25-0) (the slightly more general form claimed there follows from minor modifications).

Next, for $3 \le u \le s \le t$, we show that the normalized trace of $N_u^{(s)}$ can be reduced back to those of $\{M^{(s)}_s\}$, the most complicated one being $M^{(t-s+u)}_{u-2}$ ^(*u*−*s*+*u*). Again the formulation below allows for $u = 1, 2$.

Lemma 4.8. *Assume the same conditions as in Proposition [4.4.](#page-23-0) There exists some universal constant* $c_0 > 0$ *such that for any fixed* $d_0 \in \mathbb{R}^n$ *and an index set* $\mathcal{T} \subset$ $[-1 : t]$ *allowing for multiplicity, and any* $1 \le u \le s \le t \le \log n/c_0$ *,*

$$
\frac{1}{n} \left| \mathbb{E} \, \text{tr} \left(\mathfrak{D}_{d_0} N_u^{(s)} \mathsf{D}^{(\mathcal{T})} \right) \right| \le (K \Lambda \, \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^2} \cdot n^{-1/2} \tag{4.30}
$$

$$
+ \ (K\Lambda)^{c_0t^2} \sum_{m \in [u-2]} \max_{\substack{d_{[c_0m]} \in \{V_{[1:n]}^2, b_{[0:t]}, 1_n\}, \\ \mathcal{S} \in [-1:t]^{c_0m}}} \frac{1}{n} \Bigg| \mathbb{E} \, \text{tr} \left(\mathfrak{D}_{d_0 \circ \otimes_{q \in [c_0m]} d_q} M_{u-m-1}^{(t-s+u-m+1)} \mathsf{D}^{(\mathcal{T})} \mathsf{D}^{(\mathcal{S})} \right) \Bigg|.
$$

Here recall $D^{(T)} \equiv \prod_{s \in T} D^{(s)}$ *and* $D^{(-1)} \equiv I_n$ *.*

Proof. The proof of [\(4.30\)](#page-29-1) is largely similar to that in [\(4.18\)](#page-25-0) in the previous Lemma [4.7.](#page-25-1) We only sketch the key reduction. Let for $v \in [u-1]$,

$$
Q_{\nu}^{(u)} = \frac{1}{n^2} \Big\langle \mathbb{E} M_{u-1-\nu}^{(t-s+u)} \Delta \cdot D^{(t-s+\nu)} N_{\nu-1}^{(s)}, V \circ V \Big\rangle + \frac{1}{n^2} \mathbb{E} \Big\langle A \mathfrak{D}_{\mathsf{F}_{t-s+\nu}^{(2)}(z^{(t-s+\nu)})} \mathfrak{D}_{\partial \cdot z^{(t-s+\nu)}} N_{\nu-1}^{(s)}, V \circ V \Big\rangle - \frac{1}{n^2} \mathbb{E} \Big\langle \mathsf{B}_{t-s+\nu} \mathfrak{D}_{\mathsf{F}_{t-s+\nu-1}^{(2)}(z^{(t-s+\nu-1)})} \mathfrak{D}_{\partial \cdot z^{(t-s+\nu-1)}} N_{\nu-2}^{(s)}, V \circ V \Big\rangle.
$$

By Gaussian integration by parts,

$$
\frac{1}{n} \mathbb{E} tr(N_u^{(s)}) = \frac{1}{n} \sum_{i,j \in [n]} \mathbb{E} D_{ii}^{(t-s+u)} N_{u-1,ij}^{(s)} A_{ij} - \frac{1}{n} \mathbb{E} tr (B_{t-s+u} D^{(t-s+u-1)} N_{u-2}^{(s)})
$$
\n
$$
= \frac{1}{n^2} \mathbb{E} \langle D^{(t-s+u)} \partial ... N_{u-1}^{(s)}, V \circ V \rangle - \frac{1}{n} \mathbb{E} tr (B_{t-s+u} D^{(t-s+u-1)} N_{u-2}^{(s)})
$$
\n
$$
+ \frac{1}{n^2} \sum_{i,j \in [n]} \mathbb{E} F_{t-s+u,i}^{(2)} (z_i^{(t-s+u)}) \partial_{ij} z_i^{(t-s+u)} N_{u-1,ij}^{(s)} V_{ij}^2.
$$

Using [\(4.15\)](#page-24-4) with *u* replaced by $u - 1$, the first term above equals

$$
\frac{1}{n^2}\Big\langle \mathbb{E} M_1^{(t-s+u)}\partial_{u} N_{u-2}^{(s)} - \mathbb{E} M_0^{(t-s+u)} \mathsf{B}_{t-s+u-1} \mathsf{D}^{(t-s+u-2)}\partial_{u} N_{u-3}^{(s)}, V \circ V \Big\rangle + Q_{u-1}^{(u)}.
$$

Iterating this procedure until [\(4.15\)](#page-24-4) is applied with *u* replaced by 1, we have

$$
\frac{1}{n} \mathbb{E} \operatorname{tr}(N_u^{(s)}) = Q_{u-1}^{(u)} + \dots + Q_1^{(u)} - \frac{1}{n} \mathbb{E} \operatorname{tr} \left(\mathsf{B}_{t-s+u} \mathsf{D}^{(t-s+u-1)} N_{u-2}^{(s)} \right) + \frac{1}{n^2} \sum_{i,j \in [n]} \mathbb{E} \mathsf{F}_{t-s+u,i}^{(2)} (z_i^{(t-s+u)}) \partial_{ij} z_i^{(t-s+u)} N_{u-1,ij}^{(s)} V_{ij}^2.
$$

Note that compared to [\(4.23\)](#page-27-0) that has two remainder terms in the second and third lines therein, due to ∂ . $N_0^{(s)}$ $S_0^{(s)} = 0$ we only have one remainder term in the second line of the above display. From here we may use almost the same arguments as in the proof of Lemma [4.7](#page-25-1) to prove the claim [\(4.30\)](#page-29-1). We omit these repetitive details. \square

Lemmas [4.7](#page-25-1) and [4.8](#page-29-2) taken together, the normalized traces of $\{M_{\cdot}^{(t)}\}$ may be reduced to those of ${M^{(s)}}$ for $s \le t - 2$, and therefore eventually to those of ${M⁽¹⁾, M⁽²⁾}$. We make this precise below.

Proof of Proposition [4.4.](#page-23-0) For any $t \in \mathbb{N}$ and $k_{\ell}, k_u \in \mathbb{Z}_{\geq 0}$, let

$$
\mathcal{M}_t(k_\ell, k_u) \equiv \max_{1 \leq r \leq s \leq t} \max_{d_{[k_\ell]} \in \{V^2_{[1:n]}, b_{[0:n]}, 1_n\}} \max_{w_{[k_u]} \in [-1:n]^{k_u}} \frac{1}{n} \Big| \mathbb{E} \operatorname{tr} \big(\mathfrak{D}_{\otimes_{q \in [k_\ell]} d_q} M_r^{(s)} D^{(w_{[k_u]})} \big) \Big|.
$$

Here $D^{(-1)} = I_n$, and we write $D^{(w_{k}(\mu))} \equiv \prod_{q \in [k_u]} D^{(w_q)}$ for notational simplicity. Using [\(4.18\)](#page-25-0) in Lemma [4.7](#page-25-1) and [\(4.30\)](#page-29-1) in Lemma [4.8,](#page-29-2) with some calculations, there exists some universal constant $c_0 > 0$ such that

$$
\mathcal{M}_t(k_\ell, k_u) \le (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0t^2} \cdot n^{-1/2} + (K\Lambda)^{c_0t^2} \cdot \mathcal{M}_{t-2}(k_\ell + c_0t, k_u + c_0t).
$$

Iterating the bound with the following initial condition (which can be proved in a similar way as in Step 1 in Lemma [4.7\)](#page-25-1),

$$
\mathcal{M}_1(k_\ell, k_u) \vee \mathcal{M}_2(k_\ell, k_u) \leq (K\Lambda)^{c_1(k_\ell + k_u)} (\log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_1} \cdot n^{-1/2},
$$

we conclude that

$$
\mathcal{M}_t(0,0) \le (K\Lambda \log n \cdot (1+||z^{(0)}||_{\infty}))^{c_2 t^3} \cdot n^{-1/2},
$$

as desired. \Box

4.4. Proof of Theorem [2.1.](#page-7-1) We shall now relate the estimate for the trace calcu-lations in Proposition [4.4](#page-23-0) to the original sequence $\{M_{\cdot,\cdot}^{(t)}\}$ $\binom{1}{r}$,[-k] defined in [\(4.3\)](#page-18-2). To this end, consider an auxiliary sequence of matrices $M_{0}^{(t)}$ 0,[−*k*] , *M* (*t*) $\sum_{1,\lbrack -k \rbrack}^{(t)}, \ldots, M_{t,\lbrack k \rbrack}^{(t)}$ *t*,[−*k*] ∈ R *n*×*n* , defined recursively as follows: $M_{-1}^{(t)}$ $\binom{t}{-1, [-k]} = 0_{n \times n}, M_{0, [0]}^{(t)}$ $\binom{0}{0,[-k]} = \mathsf{D}_{[-k]}^{(t)}$ $\binom{r}{[-k]}$, and for *s* ∈ [*t*],

$$
M_{s,[-k]}^{(t)} \equiv (M_{s-1,[-k]}^{(t)}A_{[-k]} - M_{s-2,[-k]}^{(t)}B_{t-s+1})D_{[-k]}^{(t-s)}.
$$
\n(4.31)

The sequence $\{M_{\cdot,\cdot}^{(t)}\}$ $\{h^{(t)}(V_{[-k]})\}$ is different from $\{M^{(t)}(V_{[-k]})\}$ in [\(4.10\)](#page-22-3) due to the use of B. rather than $B_{\cdot, [-k]} = \text{diag}([n^{-1} \sum_{\ell \in [n]} V_{[-k], \ell}^2 \mathbb{E} F'_{\cdot, \ell}(z_{[-k], \ell}^{(\cdot)})]_{j \in [n]}$). However, their trace differences are negligible, as shown in the proof of the following result.

Corollary 4.9. *Assume the same conditions as in Proposition [4.4.](#page-23-0) Then the trace estimate in Proposition [4.4](#page-23-0) also holds for* {*M* (*t*) ·,[−*k*] } *defined in [\(4.31\)](#page-31-0), i.e., there exists some universal constant* $c_0 > 0$ *such that for* $t \le \log n/c_0$,

$$
\max_{s\in[t]} \frac{1}{n} \left| \mathbb{E} \left[\text{tr}(M_{s,[-k]}^{(t)}) | z^{(0)} \right] \right| \leq (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot n^{-1/2}.
$$

Proof. We write $\{M_{\cdot}^{(t)}(V_{[-k]})\} \equiv \{M_{\cdot,k}^{(t)}(V_{\cdot,k})\}$ $\{f(t), f(t)$,*V*_[−*k*]} in the proof. Then with $\Delta M_{s, [t]}^{(t)}$ $s, [-k]$ \equiv $M_{\rm eff}^{(t)}$ $S_{s,[-k]}^{(t)} - M_{s,1}^{(t)}$ $\sum_{s, V_{[-k]}}^{(t)}$, we have

$$
\|\Delta M_{s,[-k]}^{(t)}\|_F \le \Lambda \|A_{[-k]}\|_{\text{op}} \|\Delta M_{s-1,[-k]}^{(t)}\|_F + (K\Lambda)^2 \|\Delta M_{s-2,[-k]}^{(t)}\|_F + \Lambda \|B_{t-s+1} - B_{t-s+1,[-k]} \|_F \|M_{s-2,[-k]}^{(t)}\|_{\text{op}}.
$$

Using $||M_{s,[}^{(t)}||$ *(t)*</sup>_{*s*},[−*k*][|]lop ≤ $(KΛ(1 + ||A_{[-k]}||_{op}))^{ct}$ and

$$
\begin{split} \|\mathbf{B}_{\cdot} - \mathbf{B}_{\cdot, [-k]} \|_{F}^{2} &\leq 2 \sum_{j \in [n]} \left| \frac{1}{n} \sum_{\ell \in [n]} V_{[-k], j\ell}^{2} \left(\mathbb{E} \, \mathsf{F}'_{\cdot,\ell}(z_{\ell}^{(\cdot)}) - \mathbb{E} \, \mathsf{F}'_{\cdot,\ell}(z_{[-k], \ell}^{(\cdot)}) \right) \right|^{2} \\ &\quad + 2 \sum_{j \in [n]} \left| \frac{1}{n} \sum_{\ell \in [n]} \left(V_{j\ell}^{2} - V_{[-k], j\ell}^{2} \right) \mathbb{E} \, \mathsf{F}'_{\cdot,\ell}(z_{\ell}^{(\cdot)}) \right|^{2} \\ &\leq (K\Lambda)^{c} \cdot \left(1 + \| z^{(\cdot)} - z_{[-k]}^{(\cdot)} \|^{2} \right), \end{split}
$$

we obtain via the above recursion and Proposition [4.2](#page-19-0) that

$$
\max_{s \in [t]} \mathbb{E} ||\Delta M_{s,[-k]}^{(t)}||_F \le (K\Lambda)^{ct} (1 + ||z^{(0)}||_{\infty}) \sqrt{\log n}.
$$

The claim follows by using n^{-1} |E[tr($M_{s,[}^{(t)}$ ^(*t*)_{*s*,[−*k*]})]–E[tr(*M*^(*t*)_{*s*,*l*} $\left| \sum_{s, V_{[-k]}}^{(t)} \right|$]| ≤ $n^{-1/2}$ E||∆ $M_{s, [t]}^{(t)}$ \int *s*,[−*k*]^{$||F$} and noting that Proposition [4.4](#page-23-0) applies to $n^{-1} \mathbb{E}[\text{tr}(\mathbf{M}_{s}^{(t)})]$ $\cup_{s, V_{[-k]}}(t)$]. □

More importantly, the sequence $\{M_{\cdot,\cdot}^{(t)}\}$ $\sum_{k=1}^{(t)}$ can be related to $\{\overline{M}_{k}^{(t)}\}$ ·,[−*k*] } in [\(4.3\)](#page-18-2) (defined via a different multiplicative matrix factor $D_{t-k_1}^{(t-s)}$ $\begin{bmatrix} (-k) \\ (-k) \end{bmatrix}$ in the quadratic form of interest in the leave-one-out decomposition in Proposition [4.1.](#page-18-0)

Lemma 4.10. *Suppose the following hold for some K,* $\Lambda \geq 2$ *.*

(1) max_{*i*, *j*∈[*n*_]| V *i*_{*j*}| ≤ *K*.} *(2)* **F**_{*s*, ℓ} ∈ $C^2(\mathbb{R})$ *for all s* ∈ [0 : *t*], ℓ ∈ [*n*]*, and* max *s*∈[0:*t*] max ℓ∈[*n*] $\left\{ |F_{s,\ell}(0)| + \max_{q=1,2}^{\infty} ||F_{s,\ell}^{(q)}||_{\infty} \right\}$ $\big\} \leq \Lambda.$

Then for any D > 0, there exist some universal constant c_0 *> 0 and another constant* $c_1 = c_1(D) > 0$ *such that for* $0 \le s \le t \le \log n/c_0$ *, with* $\mathbb{P}(\cdot | z^{(0)})$ *probability at least* $1 - c_1 n^{-D}$ *,*

$$
\left| \langle A_k, \overline{M}_{s, [-k]}^{(t)} A_k \rangle - \langle A_k, M_{s, [-k]}^{(t)} A_k \rangle \right| \leq c_1 (K \Lambda)^{c_0 t} \log^{c_0} n \cdot (1 + ||z^{(0)}||_{\infty}) \cdot n^{-1/2}.
$$

Proof. Recall the definition of $\mathfrak{A}_{\tau_u}^{(u)}$ in the proof of Proposition [4.3,](#page-20-1) where $\tau_u \in \{1, 2\}$ and $\mathfrak{A}_1^{(u)}$ $\mathfrak{A}_{1}^{(u)} \equiv A, \mathfrak{A}_{2}^{(u)}$ $\mathcal{L}_2^{(u)} \equiv \mathsf{B}_{u+1}$. Recall also $\mathfrak{A}_{\tau_u}^{(u)}$ τ_{u} ,[-*k*] = $A_{[-k]} \mathbf{1}_{\tau_u=1} + \mathbf{B}_{u+1} \mathbf{1}_{\tau_u=2}$.

Fix $s \in [t]$. Note that $\langle A_k, \overline{M}_{s,[-k]}^{(t)} A_k \rangle$ consists of at most c^t terms of the form $\langle A_k, \prod_{w \in [1:r-1]} (\overline{\mathsf{D}}_{[-k]}^{(t_w)} \mathfrak{A}_{\tau_{t_w}}^{(t_w)})$ $(\tau_{t_w}, \tau_{t_w}, [-k]) \overline{D}_{[-k]}^{(t_r)} A_k$, where $r \in [s]$. Let $\Delta \overline{D}_{[-k]}^{(\cdot)} \equiv D_{[-k]}^{(\cdot)}$ $\overline{\mathsf{D}}_{[-k]}^{(\cdot)} - \overline{\mathsf{D}}_{[-k]}^{(\cdot)}$ [−*k*] . Then we have $\|\Delta \overline{D}_{[-]}^{(\cdot)}\|$ $\sum_{[-k]}^{(·)}$ ||*F* ≤ Λ||*z*^(·) − *z*_{[−} $\binom{(\cdot)}{[-k]}$ Consequently,

$$
\left| \langle A_{k}, D_{[-k]}^{(t_{1})} \mathfrak{A}_{\tau_{1},[-k]}^{(t_{1})} D_{[-k]}^{(t_{2})} \mathfrak{A}_{\tau_{2},[-k]}^{(t_{2})} \cdots D_{[-k]}^{(t_{r})} A_{k} \rangle \right|
$$

\n
$$
- \langle A_{k}, \overline{D}_{[-k]}^{(t_{1})} \mathfrak{A}_{\tau_{1},[-k]}^{(t_{1})} D_{[-k]}^{(t_{2})} \mathfrak{A}_{\tau_{2},[-k]}^{(t_{2})} \cdots D_{[-k]}^{(t_{r})} A_{k} \rangle \right|
$$

\n
$$
= \left| \sum_{\ell \in [n] \setminus \{k\}} A_{k\ell} \Delta \overline{D}_{[-k],\ell\ell}^{(t_{1})} \cdot \left\langle (\mathfrak{A}_{\tau_{t_{1}},[-k]}^{(t_{1})})_{\ell}, D_{[-k]}^{(t_{2})} \mathfrak{A}_{\tau_{t_{2}},[-k]}^{(t_{2})} \cdots D_{[-k]}^{(t_{r})} A_{k} \right\rangle \right|
$$

\n
$$
+ \left| A_{kk} \Delta \overline{D}_{[-k],kk}^{(t_{1})} \cdot \left\langle (\mathfrak{A}_{\tau_{t_{1}},[-k]}^{(t_{1})})_{k}, D_{[-k]}^{(t_{2})} \mathfrak{A}_{\tau_{t_{2}},[-k]}^{(t_{2})} \cdots D_{[-k]}^{(t_{r})} A_{k} \right\rangle \right|
$$

\n
$$
\leq ||A_{k}|| \cdot \Lambda ||z^{(t_{1})} - z_{[-k]}^{(t_{1})}|| \cdot \max_{\ell \in [n] \setminus \{k\}} \left| \left\langle (\mathfrak{A}_{\tau_{t_{1}},[-k]}^{(t_{1})})_{\ell}, D_{[-k]}^{(t_{2})} \mathfrak{A}_{\tau_{t_{2}},[-k]}^{(t_{2})} \cdots D_{[-k]}^{(t_{r})} A_{k} \right\rangle \right|
$$

\n
$$
+ \left| A_{kk} \right| \cdot \left(K \Lambda (1 + ||A||_{\text{op}}) \right)^{ct}.
$$

\n(4.32)

Note that for $\ell \in [n] \setminus \{k\}, (\mathfrak{A}_{\tau_{t_1}}^{(t_1)})$ (τ_1) _{τ_{t_1}},[−*k*])_{*ℓ*}. and D_{[−*k*}] (t_2) $\mathfrak{A}^{(t_2)}$
 $[-k]$ ^{τ} τ_2 $\tau_{t_2,[-k]}^{(t_2)} \cdots D_{[-k]}^{(t_r)}$ $\begin{bmatrix} (l_r) \\ [-k] \end{bmatrix}$ are independent of A_k , so by subgaussian inequality first conditional on $A_{[-k]}$ and then unconditionally, with probability at least $1 - cn^{-D}$,

$$
\max_{\ell \in [n] \setminus \{k\}} \left| \langle (\mathfrak{A}_{\tau_{t_1}, [-k]}^{(t_1)})_{\ell}, D_{[-k]}^{(t_2)} \mathfrak{A}_{\tau_{t_2}, [-k]}^{(t_2)} \cdots D_{[-k]}^{(t_r)} A_k \rangle \right| \leq c_D (K\Lambda)^{ct} \sqrt{\log n} \cdot n^{-1/2}.
$$

Combined with Proposition [4.2,](#page-19-0) for $t \leq \log n/c$, with probability at least $1 - cn^{-D}$,

LHS of (4.32)
$$
\leq c_D(K\Lambda)^{ct} \log^c n \cdot (1 + ||z^{(0)}||_{\infty}) \cdot n^{-1/2}
$$

Iterating the bound proves the claim.

We are now finally in a position to prove Theorem [2.1.](#page-7-1)

Proof of Theorem [2.1.](#page-7-1) By Propositions [4.1,](#page-18-0) [4.2](#page-19-0) and Lemma [4.10,](#page-32-1) for $t \le \log n/c$, with probability at least $1 - cn^{-D}$,

$$
\left| z_k^{(t+1)} - \langle A_k, F_t(z_{[-k]}^{(t)}) \rangle \right| \le (K\Lambda)^{ct} \log n \cdot (1 + ||z^{(0)}||_{\infty})
$$

$$
\times \Big(\max_{s \in [t]} \Big| \langle \langle A_k, M_{t-s, [-k]}^{(t)} A_k \rangle - b_{t,k} \mathbf{1}_{s=t} \rangle \Big| + n^{-1/2} \Big).
$$

For $s \le t - 1$, by Hanson-Wright inequality and Corollary [4.9,](#page-31-1) for $t \le \log n/c$, with probability at least $1 - cn^{-D}$,

$$
\max_{s\in[t-1]} \left| \left(\langle A_k, M_{t-s, [-k]}^{(t)} A_k \rangle \right) \right| \le \left(K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}) \right)^{ct^3} \cdot n^{-1/2}.
$$

For $s = t$, by Hanson-Wright inequality and using Proposition [4.2](#page-19-0) again, for $t \leq$ log *n*/*c*, with probability at least $1 - cn^{-D}$,

$$
\left| (\langle A_k, M_{0, [-k]}^{(t)} A_k \rangle - b_{t,k}) \right| \leq \frac{1}{n} \sum_{\ell \in [n]} V_{k\ell}^2 \left| \mathbb{E} \left[F'_{t,\ell}(z_{[-k],\ell}^{(t)}) - \mathbb{E} \left[F'_{t,\ell}(z_{\ell}^{(t)}) \right] + \frac{(K\Lambda)^c \log n}{\sqrt{n}} \right] \right|
$$

$$
\leq (K\Lambda)^{ct} \log^c n \cdot (1 + ||z^{(0)}||_{\infty}) \cdot n^{-1/2}.
$$

Combining the above three displays proves that for $t \leq \log n/c$, with probability at least $1 - cn^{-D}$,

$$
\left| z_k^{(t+1)} - \langle A_k, \mathsf{F}_t(z_{[-k]}^{(t)}) \rangle \right| \le (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct^3} \cdot n^{-1/2}.
$$
 (4.33)

The proof is complete.

5. Proofs of remaining results in Section [2](#page-7-0)

5.1. **Proof of Theorem [2.3.](#page-8-0)** By apriori estimates for $(z_k^{(s+1)})$ $\binom{(S+1)}{k}$ _{s∈[0:*t*]} (cf. Proposition [4.2\)](#page-19-0) and $(A_k, F_s(z_{[-k]}^{(s)})$ $\binom{s}{[-k]}$))_{s∈[0:*t*]}, [\(4.33\)](#page-33-2) also holds for moment estimates. Consequently, for $t \leq \log n/c$,

$$
\begin{split} \left| \mathbb{E} \psi \big[(z_k^{(s+1)})_{s \in [0:r]} \big] - \mathbb{E} \psi \big[(\langle A_k, \mathsf{F}_s(z_{[-k]}^{(s)}) \rangle)_{s \in [0:r]} \big] \right| \\ &\le \Lambda \cdot \mathbb{E}^{1/2} \left(1 + \left\| (z_k^{(s+1)})_{s \in [0:r]} \right\| \right) + \left\| (\langle A_k, \mathsf{F}_s(z_{[-k]}^{(s)}) \rangle)_{s \in [0:r]} \right\| \right)^{2p} \\ &\times \mathbb{E}^{1/2} \left\| (z_k^{(s+1)} - \langle A_k, \mathsf{F}_s(z_{[-k]}^{(s)}) \rangle)_{s \in [0:r]} \right\|^2 \\ &\le (K\Lambda \log n \cdot (1 + \|z^{(0)}\|_{\infty}))^{c_p r^3} \cdot n^{-1/2} . \end{split} \tag{5.1}
$$

Below we shall replace $\mathbb{E}\,\psi[(\langle A_k, \mathsf{F}_s(z_{t-1}^{(s)})\rangle)$ $\binom{s}{[-k]}$))_{s∈[0:*t*]} in the left hand side of the above display by the quantity $\mathbb{E}\psi[(Z_k^{(s+1)})]$ $\binom{(s+1)}{k}$ _{*s*∈[0:*t*]} and its variant.

$$
\Box
$$

.

(**Step 1**). For $s_1, s_2 \in [0 : t]$, let for $k \in [n]$

$$
\begin{cases}\n\alpha_k^{(s_1+1,s_2+1)} \equiv \frac{1}{n} \sum_{\ell \in [n]} V_{k\ell}^2 \mathbb{E} \mathsf{F}_{s_1,\ell}(z_{\ell}^{(s_1)}) \mathsf{F}_{s_2,\ell}(z_{\ell}^{(s_2)}), \\
\alpha_{[-k],k}^{(s_1+1,s_2+1)} \equiv \frac{1}{n} \sum_{\ell \in [n]} V_{k\ell}^2 \mathsf{F}_{s_1,\ell}(z_{[-k],\ell}^{(s_1)}) \mathsf{F}_{s_2,\ell}(z_{[-k],\ell}^{(s_2)}). \n\end{cases} \tag{5.2}
$$

Note that $\alpha_{[-k]k}^{(s_1+1,s_2+1)}$ $[-k]$, k is random (without expectation). Let $Z'_{k;\alpha} \sim \mathcal{N}(0, I_{t+1})$ be a standard Gaussian vector, independent of all other random variables. Let

$$
\Sigma_{k;\alpha} \equiv (\alpha_k^{(s_1+1,s_2+1)})_{s_1,s_2 \in [0:t]}, \quad \Sigma_{[-k],k;\alpha} \equiv (\alpha_{[-k],k}^{(s_1+1,s_2+1)})_{s_1,s_2 \in [0:t]}
$$

It is easy to see that $\Sigma_{k;\alpha}, \Sigma_{[-k],k;\alpha} \in \mathbb{R}^{(t+1)\times(t+1)}$ are covariance matrices, so we may further define

$$
(Z_{k;\alpha}^{(s+1)})_{s\in[0:t]} \equiv \Sigma_{k;\alpha}^{1/2} Z_{k;\alpha}', \quad (Z_{[-k],k;\alpha}^{(s+1)})_{s\in[0:t]} \equiv \Sigma_{[-k],k;\alpha}^{1/2} Z_{k;\alpha}' \in \mathbb{R}^{t+1}.
$$

Using $\mathbb{E}\,\psi[(\langle A_k, \mathsf{F}_s(z_{[})}^{(s)}\rangle$ $\langle S \rangle_{s \in [0:t]}$ $\big] = \mathbb{E} \psi[(Z_{[-k],k}^{(s+1)}]$ $\binom{(s+1)}{[-k],k;\alpha}$ _{*s*∈[0:*t*]}, we then have

$$
\begin{aligned} & \left| \mathbb{E} \, \psi \big[(\langle A_k, \mathsf{F}_s(z_{[-k]}^{(s)}) \rangle)_{s \in [0:t]} \big] - \mathbb{E} \, \psi \big[(Z_{k;\alpha}^{(s+1)})_{s \in [0:t]} \big] \right| \\ &= \left| \mathbb{E} \, \psi \big[(Z_{k;\alpha}^{(s+1)})_{s \in [0:t]} \big] - \mathbb{E} \, \psi \big[(Z_{[-k],k;\alpha}^{(s+1)})_{s \in [0:t]} \big] \right| \\ &\le (K\Lambda)^{c_p t} \cdot \mathbb{E}^{1/2} \left\| \sum_{k;\alpha}^{1/2} - \sum_{[-k],k;\alpha}^{1/2} \right\|_F^2 \\ &\stackrel{(*)}{\le} (K\Lambda)^{c_p t} \cdot \left(\log n \cdot (1 + \|z^{(0)}\|_{\infty}) \right)^c \cdot n^{-1/4} . \end{aligned} \tag{5.3}
$$

In (∗) above, we used the following estimate: By Lemma [A.3,](#page-49-0)

$$
\mathbb{E} \left\| \sum_{k;\alpha}^{1/2} - \sum_{[-k],k;\alpha}^{1/2} \right\|_F^2 \le (t+1)^{1/2} \cdot \mathbb{E}^{1/2} \left\| \sum_{k;\alpha} - \sum_{[-k],k;\alpha} \right\|_F^2
$$

\n
$$
\le (K\Lambda)^{ct} \cdot \max_{s_1, s_2 \in [0:t]} \left\{ \max_{u,v \in \{s_1, s_2\}} \mathbb{E}^{1/2} \left(\frac{1}{n} \sum_{\ell \in [n]} (1 + |z_{\ell}^{(u)}| + |z_{[-k],\ell}^{(u)})| z_{\ell}^{(v)} - z_{[-k],\ell}^{(v)} \right) \right\}^2
$$

\n
$$
+ \mathbb{E}^{1/2} \left(\frac{1}{n} \sum_{\ell \in [n]} V_{k\ell}^2 \cdot (\text{id} - \mathbb{E}) \mathsf{F}_{s_1,\ell} (z_{\ell}^{(s_1)}) \mathsf{F}_{s_2,\ell} (z_{\ell}^{(s_2)}) \right)^2 \right\}
$$

\n
$$
\le (K\Lambda)^{ct} \cdot (\log n \cdot (1 + ||z^{(0)}||_{\infty}))^c \cdot n^{-1/2}.
$$

Here in the last inequality of the above display, we bound the second term in the bracket by applying the Gaussian concentration Lemma [A.2](#page-48-0) to $H_0(A) \equiv$ *n*⁻¹ ∑_{$\ell \in [n]$} $V_{k\ell}^2$ **F**_{*s*₁, ℓ ($z_{\ell}^{(s_1)}$} $(\frac{s_1}{\ell})\mathsf{F}_{s_2,\ell}(z_{\ell}^{(s_2)})$ $\binom{s_2}{\ell}$, upon computing, for any symmetric matrices *A*, *A*₁, *A*₂, the following estimates: (i) $|H_0(A_1) - H_0(A_2)| \le (K\Lambda(1 + ||A_1||_{op} + ||A_2||_{op} + ||A_1||_{op} + ||A_2||_{op} + ||A_2||_{op} + ||A_1||_{op} + ||A_2||_{op} + ||A_2||_{op} + ||A_1||_{op} + ||A_2||_{op}$ $||A_2||_{op})$ ^{ct}(1+ $||z^{(0)}||_{\infty}$)^c⋅ $||A_1 - A_2||_{op}$, and (ii) $|H_0(A)| \le (K\Lambda(1+||A||_{op}))^{ct}(1+||z^{(0)}||_{\infty})^c$.

Combining (5.1) and (5.3) , we conclude

$$
\max_{k \in [n]} \left| \mathbb{E} \psi[(z_k^{(s+1)})_{s \in [0:t]}] - \mathbb{E} \psi[(Z_{k;\alpha}^{(s+1)})_{s \in [0:t]}] \right|
$$

$$
\leq (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_p t^3} \cdot n^{-1/4} \equiv \delta_n^{(t+1)}.
$$
 (5.4)

(**Step 2**). Recall the definition of $Z^{(1)}, Z^{(2)}, \ldots$ with correlation structures specified in Definition [2.2.](#page-8-1) For $k \in [n]$, let $\Sigma_k \equiv (\text{Cov}(Z_k^{(s_1+1)})$ $Z_k^{(s_1+1)}, Z_k^{(s_2+1)}$ ${k^{(s_2+1)} \choose k}$ _{s_{1,*s*2∈[0:*t*]}. Then using} Lemma [A.3](#page-49-0) again,

$$
\left| \mathbb{E} \, \psi \big[(Z_{k;\alpha}^{(s+1)})_{s \in [0:t]} \big] - \mathbb{E} \, \psi \big[(Z_k^{(s+1)})_{s \in [0:t]} \big] \right|
$$

.

$$
\leq (K\Lambda)^{c_{\mathfrak{p}}t} \cdot \|\Sigma_{k;\alpha}^{1/2} - \Sigma_{k}^{1/2}\|_{F} \leq (K\Lambda)^{c_{\mathfrak{p}}t} \cdot \|\Sigma_{k;\alpha} - \Sigma_{k}\|_{F}^{1/2}
$$

$$
\leq (K\Lambda)^{c_{\mathfrak{p}}t} \cdot \max_{s_1, s_2 \in [0:t]} \left\{\frac{1}{n} \sum_{\ell \in [n]} \left| \mathbb{E} \mathsf{F}_{s_1,\ell}(z_{\ell}^{(s_1)}) \mathsf{F}_{s_2,\ell}(z_{\ell}^{(s_2)}) - \mathbb{E} \mathsf{F}_{s_1,\ell}(Z_{\ell}^{(s_1)}) \mathsf{F}_{s_2,\ell}(Z_{\ell}^{(s_2)}) \right| \right\}^{1/2}.
$$

Here we adopt the convention that $Z_{\cdot}^{(0)} = z_{\cdot}^{(0)}$. Now with $\mathscr{F}_{t+1}(L, \mathfrak{p}) \equiv {\psi : \mathbb{R}^{t+1} \rightarrow \mathbb{R}^{t}}$ $\mathbb{R}, |\psi(x) - \psi(y)| \leq L \cdot (1 + ||x|| + ||y||)^p \cdot ||x - y||$ for all $x, y \in \mathbb{R}^{t+1}$ and

$$
\gamma_{t+1}(L,\mathfrak{p}) \equiv \max_{k \in [n]} \sup_{\psi \in \mathscr{F}_{t+1}(L,\mathfrak{p})} \left| \mathbb{E} \psi \big[\big(Z_{k;\alpha}^{(s+1)} \big)_{s \in [0:r]} \big] - \mathbb{E} \psi \big[\big(Z_{k}^{(s+1)} \big)_{s \in [0:r]} \big] \right|,
$$

using [\(5.4\)](#page-34-1) at iteration *t*, we obtain the recursion:

$$
\gamma_{t+1}(\Lambda, \mathfrak{p}) \le (K\Lambda)^{c_{\mathfrak{p}}t} \cdot [(\delta_n^{(t)})^{1/2} + \gamma_t^{1/2}(2\Lambda, \mathfrak{p})].
$$

Now iterating the bound and using the initial condition $\gamma_1 \equiv 0$ (equivalently, $\mathbb{E}\,\psi[Z^{(1)}_{k\cdot\alpha}$ $\chi_{k;\alpha}^{(1)}$] = $\mathbb{E}\psi[Z_k^{(1)}]$ k ⁽¹⁾) for all $k \in [n]$), we have

$$
\gamma_{t+1}(\Lambda, \mathfrak{p}) \le (K\Lambda \log n \cdot (1+||z^{(0)}||_{\infty}))^{c_{\mathfrak{p}}t^3} \cdot n^{-1/c_0'},\tag{5.5}
$$

where $c_0 > 0$ is a universal constant. The claimed error estimate now follows from [\(5.4\)](#page-34-1) and [\(5.5\)](#page-35-1), and noting that $t \leq \log n/c$ can be removed for free as otherwise we may use the (trivial) estimate in Proposition [4.2](#page-19-0) instead.

5.2. **Proof of Theorem [2.4.](#page-9-0)** Recall the state evolution and $Z^{(1)}, Z^{(2)}, \ldots$ in Definition [2.2.](#page-8-1) For $k \in [n]$, let $\sigma_{t,k}^2 \equiv \text{Var}(Z_k^{(t)})$ $b_{t,k}$ and recall $b_{t,k}$ = *n*^{−1} ∑ $\ell \in [n]$ $V_{k\ell}^2 \mathbb{E} [F'_{t,\ell}(Z^{(t)}_{\ell})]$ $(\bar{v}_\ell^{(t)})|z^{(0)}|$. The modified AMP iterate associated with $\{\bar{b}_t\}$ is denoted $\{\overline{z}^{(t)}\}$.

Lemma 5.1. *Assume the same conditions as in Theorem [2.4.](#page-9-0) Fix any* Λ*-pseudo-Lipschitz function* $\psi : \mathbb{R}^{t+1} \to \mathbb{R}$ *of order 2. Then there exists some universal constant* $c_0 > 0$ *, such that*

$$
\begin{aligned} & \left| \frac{1}{n} \sum_{k \in [n]} \left(\mathbb{E} \left[\psi(\overline{z}_{k}^{(t+1)}, \overline{z}_{k}^{(t)}, \dots, \overline{z}_{k}^{(1)}) | z^{(0)} \right] - \mathbb{E} \left[\psi(Z_{k}^{(t+1)}, Z_{k}^{(t)}, \dots, Z_{k}^{(1)}) | z^{(0)} \right] \right) \right| \\ &\leq (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot n^{-1/c_0^t}. \end{aligned}
$$

Proof. We assume for notational simplicity that ψ is full dimensional (i.e., ψ depends on all of its arguments), and write $\sigma_{*,\psi}$ as σ_* . Let $\varphi \in C^{\infty}(\mathbb{R})$ be a mollifier supported in [-1, 1] such that $\varphi \ge 0$ and $\int \varphi = 1$, and $\varphi_{\delta}(\cdot) = \delta^{-1} \varphi(\cdot/\delta)$. Let $\mathsf{F}_{\delta;s,\ell} \equiv \mathsf{F}_{s,\ell} * \varphi_{\delta}$. Then:

- For $q = 1$, $|F'_{\delta;s,\ell}(x)| = |$ $\int \mathsf{F}'_{s,\ell}(x-z)\varphi_{\delta}(z) dz \leq \Lambda.$
- For $q \ge 2, q \in \mathbb{N}$ and $\delta \in (0, 1]$, $\left|\mathsf{F}^{(q)}_{\delta;s,\ell}(x)\right| = \frac{1}{\delta^q}$ δ *q* $\begin{array}{c} \n \downarrow \\ \n \downarrow \\ \n \downarrow \n \end{array}$ \overline{a} [*x*±δ] $\mathsf{F}_{s,\ell}'(z)\varphi^{(q-1)}((x-z)/\delta) dz$ $\left| \leq 2\Lambda ||\varphi^{(q-1)}||_{\infty} \delta^{-(q-1)}.$
- $|F_{\delta;s,\ell}(x) F_{s,\ell}(x)| = |$ \int_1^1 -1 $\left(\mathsf{F}_{s,\ell}(x-\delta z)-\mathsf{F}_{s,\ell}(x)\right)\varphi(z)\,\mathrm{d}z\big|\leq\Lambda\|\varphi\|_{\infty}\delta.$

(Step 1). Let { $z_{\delta}^{(\cdot)}$ } be the AMP iterate associated with the non-linearities { $F_{\delta,s}$ } $_{s\in[0:t]}$ and the same initialization $z^{(0)}$. We define the state evolution associated with $z_{\delta}^{(\cdot)}$ as follows. Let $Z_{\delta}^{(1)}$ $z_\delta^{(1)}, Z_\delta^{(2)}$ $\delta^{(2)}$, ... $\in \mathbb{R}^n$ be a sequence of correlated *n*-dimensional centered Gaussian vectors with independent entries, whose correlation structure along the iteration path is determined with the initial condition $Var(Z_{\delta \cdot k}^{(1)})$ $\binom{11}{\delta;k}$ ≡ *n*⁻¹ ∑_{$\ell \in [n]$} $V_{k\ell}^2$ **F**_{δ}, $0, \ell$ ($z_{\ell}^{(0)}$ ⁽⁰⁾) for $k \in [n]$, and the subsequent recursive updates:

$$
Cov(Z_{\delta;k}^{(s_1+1)}, Z_{\delta;k}^{(s_2+1)}) \equiv \frac{1}{n} \sum_{\ell \in [n]} V_{k\ell}^2 \mathbb{E} \left[\mathsf{F}_{\delta;s_1,\ell}(Z_{\delta;\ell}^{(s_1)}) \mathsf{F}_{\delta;s_2,\ell}(Z_{\delta;\ell}^{(s_2)}) \right], \quad k \in [n]. \quad (5.6)
$$

For notational convenience, we shall identify $Z_{0}^{(\cdot)}$ $Q_{0}^{(1)} = Z_{0}^{(1)}$ as defined in Definition [2.2](#page-8-1) for the (unsmoothed) AMP iterate. Using almost the same proof as in Step 2 of Theorem [2.3,](#page-8-0) there exists some universal constant $c_0 > 0$ such that

$$
\max_{k \in [n]} \left| \mathbb{E} \, \psi(Z_k^{(t+1)}, \dots, Z_k^{(1)}) - \mathbb{E} \, \psi(Z_{\delta; k}^{(t+1)}, \dots, Z_{\delta; k}^{(1)}) \right| \leq (K \Lambda \log n)^{c_0 t^2} \delta^{1/c_0^t}.
$$
 (5.7)

In particular, with $\sigma_{\delta;t,k}^2 \equiv \text{Var}(Z_{\delta;k}^{(t)})$ $\delta_{k}^{(l)}$, we have

$$
\max_{k \in [n]} \left| \sigma_{\delta; t, k}^2 - \sigma_{t, k}^2 \right| \le (K \Lambda \log n)^{c_0 t^2} \cdot \delta^{1/c_0^t} \equiv \text{err}_{n, t}(\delta). \tag{5.8}
$$

(Step 2). Let $\{\overline{z}_{\delta}^{(\cdot)}\}$ be the modified AMP iterate using the correction vector $\{\overline{b}_{\delta;t}\}$ with the above state evolution, i.e., $\overline{b}_{\delta;t,k} = n^{-1} \sum_{\ell \in [n]} V_{k\ell}^2 \mathbb{E}[\mathbf{F}'_{\delta;t,\ell}(Z_{\delta;\ell}^{(t)})]$ $\frac{d^{(t)}}{d^{(t)}}$ = $|z^{(0)}|$. In this step, we prove that for some universal constant $c_0 > 0$, if $t \le \log n/c_0$,

$$
\left| \frac{1}{n} \sum_{k \in [n]} \left(\mathbb{E} \psi(z_{\delta;k}^{(t+1)}, z_{\delta;k}^{(t)}, \dots, z_{\delta;k}^{(1)}) - \mathbb{E} \psi(\overline{z}_{\delta;k}^{(t+1)}, \overline{z}_{\delta;k}^{(t)}, \dots, \overline{z}_{\delta;k}^{(1)}) \right) \right|
$$
\n
$$
\leq (K \Lambda \delta^{-1} \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot n^{-1/c_0^t}.
$$
\n(5.9)

To this end, note that by applying Theorem [2.3](#page-8-0) to the smoothed AMP iterate $\{z_{\delta}^{(\cdot)}\},$

$$
\begin{split} ||z_{\delta}^{(t+1)} - \overline{z}_{\delta}^{(t+1)}|| \\ &\leq ||A\mathsf{F}_{\delta;t}(z_{\delta}^{(t)}) - A\mathsf{F}_{\delta;t}(\overline{z}_{\delta}^{(t)})|| + K^{2}\Lambda \cdot ||\mathsf{F}_{\delta;t-1}(z_{\delta}^{(t-1)}) - \mathsf{F}_{\delta;t-1}(\overline{z}_{\delta}^{(t-1)})|| \\ &+ K^{2} \cdot \frac{1}{n} \sum_{\ell \in [n]} \left| \mathbb{E} \mathsf{F}_{\delta;t,\ell}'(z_{\delta;\ell}^{(t)}) - \mathbb{E} \mathsf{F}_{\delta;t,\ell}'(Z_{\delta;\ell}^{(t)}) \right| \cdot ||\mathsf{F}_{\delta;t-1}(\overline{z}_{\delta}^{(t-1)})|| \\ &\leq (||A||_{\text{op}} + K^{2}\Lambda) \cdot \left(\max_{s=t,t-1} ||\mathsf{F}_{\delta;s}(z_{\delta}^{(s)}) - \mathsf{F}_{\delta;s}(\overline{z}_{\delta}^{(s)})|| \right. \\ &\left. + (K\Lambda \delta^{-1} \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c t^{3}} \cdot n^{-1/c^{t}} \cdot (\sqrt{n} + ||\overline{z}_{\delta}^{(t-1)}||) \right). \end{split}
$$

Consequently,

$$
\max_{s \in [0:t+1]} ||z_{\delta}^{(s)} - \overline{z}_{\delta}^{(s)}|| \le (K\Lambda)^2 (1 + ||A||_{\text{op}}) \cdot \max_{s \in [0:t]} ||z_{\delta}^{(s)} - \overline{z}_{\delta}^{(s)}||
$$

+ $(1 + ||A||_{\text{op}})(K\Lambda\delta^{-1} \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c t^3} \cdot n^{-1/c^t} \cdot (\sqrt{n} + \max_{s \in [0:t]} ||\overline{z}_{\delta}^{(s)}||).$

Iterating the bound, we arrive at

$$
\max_{s \in [0:t+1]} n^{-1/2} \|z_{\delta}^{(s)} - \overline{z}_{\delta}^{(s)}\| \le (1 + \|A\|_{\text{op}})^{ct} \cdot (K\Lambda \delta^{-1} \log n \cdot (1 + \|z^{(0)}\|_{\infty}))^{ct^3}
$$

$$
\times n^{-1/c'} \cdot \left(1 + \max_{s \in [0:t]} |\overline{z}_{\delta}^{(s)}\|_{\infty}\right). \tag{5.10}
$$

On the other hand, using the same leave-one-out argument as in the proof of Propo-sition [4.2](#page-19-0) (where for the current modified AMP iterate, we only need to require $F_{\cdot,\cdot}$ to be Lipschitz), with probability at least $1 - (nt)^{c_0} e^{-(x^2 \wedge n)/c_0}$,

$$
\max_{s \in [t+1]} (||\overline{z}_{\delta}^{(s)}||_{\infty} \vee ||\overline{z}^{(s)}||_{\infty}) \le (K\Lambda)^{c_0 t} x \cdot (1 + ||z^{(0)}||_{\infty}). \tag{5.11}
$$

Finally note that with $\psi_0(z_\delta^{(\cdot)}) \equiv n^{-1} \sum_{k \in [n]} \psi(z_{\delta;k}^{(t+1)})$ $\zeta_{\delta;k}^{(t+1)}, \ldots, z_{\delta;k}^{(1)}$ $\phi_{\delta;k}^{(1)}$ and similarly $\psi_0(\overline{z}_{\delta}^{(\cdot)}) \equiv$ n^{-1} $\sum_{k \in [n]} \psi(\overline{z}_{\delta;k}^{(t+1)})$ $\overline{z}_{\delta;k}^{(t+1)}, \ldots, \overline{z}_{\delta;k}^{(1)}$ $\binom{1}{\delta;k}$, we have

$$
\begin{split} \left| \psi_0(z_{\delta}^{(\cdot)}) - \psi_0(\overline{z}_{\delta}^{(\cdot)}) \right| \\ &\leq \Lambda t^{c_0} \cdot \max_{s \in [0:t+1]} \left(1 + \|z_{\delta}^{(s)}\|_{\infty} + \|\overline{z}_{\delta}^{(s)}\|_{\infty} \right)^{c_0} \cdot n^{-1/2} \|z_{\delta}^{(s)} - \overline{z}_{\delta}^{(s)}\|. \end{split} \tag{5.12}
$$

Combining [\(5.10\)](#page-37-0)-[\(5.12\)](#page-37-1), for any $D > 0$, there exists some $c_1 = c_1(D) > 0$, such that for $t \leq \log n/c_0$, on an event with probability at least $1 - c_1 n^{-D}$,

$$
\left|\psi_0(z_{\delta}^{(\cdot)}) - \psi_0(\overline{z}_{\delta}^{(\cdot)})\right| \le (c_1 K \Lambda \delta^{-1} \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct^3} \cdot n^{-1/c^t}.
$$

The moment estimate in [\(5.9\)](#page-36-0) follows by the following simple apriori estimates: by assuming $\psi(0) = 0$ without loss of generality, using Proposition [4.2](#page-19-0) we have

$$
\mathbb{E}^{1/2} |\psi_0(z_{\delta}^{(\cdot)})|^2 \le \Lambda \mathbb{E}^{1/2} \max_{s \in [t+1]} (1 \vee ||z_{\delta}^{(s)}||_{\infty})^c \cdot (n^{-1/2} ||z_{\delta}^{(s)}||)^2
$$

$$
\le (K\Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{ct}.
$$

A completely similar estimate holds for $\mathbb{E}^{1/2} |\psi_0(\overline{z}_\delta^{(\cdot)})|^2$. (Step 3). In this step, we prove that for $t \le \log n/c_0$ and $err_{n,t}(\delta) \le \min\{1, \sigma_*^2/2\}$,

$$
\left| \frac{1}{n} \sum_{k \in [n]} \left(\mathbb{E} \psi(\overline{z}_{k}^{(t+1)}, \overline{z}_{k}^{(t)}, \dots, \overline{z}_{k}^{(1)}) - \mathbb{E} \psi(\overline{z}_{\delta;k}^{(t+1)}, \overline{z}_{\delta;k}^{(t)}, \dots, \overline{z}_{\delta;k}^{(1)}) \right) \right|
$$

$$
\leq (K \Lambda \sigma_*^{-1} \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^2} \cdot (\text{err}_{n,t}(\delta) + \delta + n^{-100}). \tag{5.13}
$$

The proof has certain similarities to Step 2. For $err_{n,t}(\delta) \le \min\{1, \sigma_*^2/2\}$,

$$
\begin{split} ||\overline{z}_{\delta}^{(t+1)} - \overline{z}^{(t+1)}|| \\ &\leq ||A\mathsf{F}_{\delta;t}(\overline{z}_{\delta}^{(t)}) - A\mathsf{F}_{t}(\overline{z}^{(t)})|| + K^{2}\Lambda \cdot ||\mathsf{F}_{\delta;t-1}(\overline{z}_{\delta}^{(t-1)}) - \mathsf{F}_{t-1}(\overline{z}^{(t-1)})|| \\ &+ K^{2} \cdot \frac{1}{n} \sum_{\ell \in [n]} \left| \mathbb{E} \mathsf{F}_{\delta;t,\ell}'(Z_{\delta;\ell}^{(t)}) - \mathbb{E} \mathsf{F}_{t,\ell}'(Z_{\ell}^{(t)}) \right| \cdot ||\mathsf{F}_{t-1}(\overline{z}^{(t-1)})|| \\ &\stackrel{(*)}{\leq} (||A||_{\text{op}} + K^{2}\Lambda) \cdot \left(\max_{s=t,t-1} ||\mathsf{F}_{\delta;s}(\overline{z}_{\delta}^{(s)}) - \mathsf{F}_{\delta;s}(\overline{z}^{(s)})|| + \max_{s=t,t-1} ||\mathsf{F}_{\delta;s}(\overline{z}^{(s)}) - \mathsf{F}_{s}(\overline{z}^{(s)})|| \\ &+ (K\Lambda)^{ct} \sigma_{*}^{-c} \cdot (\text{err}_{n,t}(\delta) + \delta) \cdot (\sqrt{n} + ||\overline{z}^{(t-1)}||) \right) \end{split}
$$

$$
\leq (||A||_{op} + 1)(K\Lambda \sigma_*^{-1})^{ct} \cdot \Big(\max_{s=t,t-1}||\overline{z}_{\delta}^{(s)} - \overline{z}^{(s)}|| + (\text{err}_{n,t}(\delta) + \delta) \cdot (\sqrt{n} + ||\overline{z}^{(t-1)}||)\Big).
$$

Here in (*) we used the following estimate: for $err_{n,t}(\delta) \le \min\{1, \sigma_*^2/2\}$, by [\(5.8\)](#page-36-1),

$$
\frac{1}{n} \sum_{\ell \in [n]} \left| \mathbb{E} \mathsf{F}'_{\delta;t,\ell}(Z_{\delta;\ell}^{(t)}) - \mathbb{E} \mathsf{F}'_{t,\ell}(Z_{\ell}^{(t)}) \right|
$$
\n
$$
= \frac{1}{n} \sum_{\ell \in [n]} \left| \sigma_{\delta;t,k}^{-2} \mathbb{E} Z_{\delta;\ell}^{(t)} \mathsf{F}_{\delta;t,\ell}(Z_{\delta;\ell}^{(t)}) - \sigma_{t,k}^{-2} \mathbb{E} Z_{\ell}^{(t)} \mathsf{F}_{t,\ell}(Z_{\ell}^{(t)}) \right|
$$
\n
$$
\leq \frac{(K\Lambda)^{ct}}{\sigma_*^2(\sigma_*^2 - \text{err}_{n,t}(\delta))_+} \cdot \text{err}_{n,t}(\delta) + \frac{1}{\sigma_*^2} \cdot \frac{1}{n} \sum_{\ell \in [n]} \left| \mathbb{E} Z_{\delta;\ell}^{(t)} \mathsf{F}_{\delta;t,\ell}(Z_{\delta;\ell}^{(t)}) - \mathbb{E} Z_{\ell}^{(t)} \mathsf{F}_{t,\ell}(Z_{\ell}^{(t)}) \right|
$$
\n
$$
\leq (K\Lambda)^{ct} \sigma_*^{-c} \cdot (\text{err}_{n,t}(\delta) + \delta).
$$

Consequently,

$$
\max_{s \in [0:t+1]} ||\overline{z}_{\delta}^{(s)} - \overline{z}^{(s)}|| \le (||A||_{\text{op}} + 1)(K\Lambda \sigma_*^{-1})^{ct}
$$

$$
\times \Big(\max_{s \in [0:t]} ||\overline{z}_{\delta}^{(s)} - \overline{z}^{(s)}|| + (\text{err}_{n,t}(\delta) + \delta) \cdot (\sqrt{n} + \max_{s \in [0:t]} ||\overline{z}^{(s)}||) \Big).
$$

Iterating the bound, we obtain

$$
\max_{s \in [0:t+1]} n^{-1/2} ||\overline{z}_{\delta}^{(s)} - \overline{z}^{(s)}||
$$

$$
\leq (K\Lambda \sigma_*^{-1} \cdot (1 \vee ||A||_{\text{op}}))^{c_0 t^2} \cdot (\text{err}_{n,t}(\delta) + \delta) \cdot (1 + \max_{s \in [0:t]} ||\overline{z}^{(s)}||_{\infty}).
$$

From here the proof proceeds as in Step 2.

(**Step 4**). Applying Theorem [2.3](#page-8-0) to the smoothed AMP iterate $\{z_{\delta}^{(\cdot)}\}$, we obtain

$$
\left| \frac{1}{n} \sum_{k \in [n]} \left(\mathbb{E} \psi(z_{\delta;k}^{(t+1)}, z_{\delta;k}^{(t)}, \dots, z_{\delta;k}^{(1)}) - \mathbb{E} \psi(Z_{\delta;k}^{(t+1)}, Z_{\delta;k}^{(t)}, \dots, Z_{\delta;k}^{(1)}) \right) \right|
$$

$$
\leq (K \Lambda \delta^{-1} \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot n^{-1/c_0'}.
$$

Combined with (the averaged version of) [\(5.7\)](#page-36-2), [\(5.9\)](#page-36-0) and [\(5.13\)](#page-37-2), we have for $t \le$ $\log n/c_0$ and $err_{n,t}(\delta) \le \min\{1, \sigma_*^2/2\},\$

$$
\left| \frac{1}{n} \sum_{k \in [n]} \left(\mathbb{E} \psi(\overline{z}_{k}^{(t+1)}, \overline{z}_{k}^{(t)}, \dots, \overline{z}_{k}^{(1)}) - \mathbb{E} \psi(Z_{k}^{(t+1)}, Z_{k}^{(t)}, \dots, Z_{k}^{(1)}) \right) \right|
$$

$$
\leq (K \Lambda \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot (\delta^{-c_0 t^3} n^{-1/c_0^t} + \delta^{1/c_0^t}).
$$

Choosing $\delta \equiv n^{-1/c_*^t}$ for sufficiently large c_* to conclude.

Proof of Theorem [2.4.](#page-9-0) Without loss of generality we assume $\psi(0) = 0$ (otherwise consider $\psi - \psi(0)$). By Lemma [5.1,](#page-35-2) we only need to prove concentration of $H(A) \equiv$ $n^{-1} \sum_{k \in [n]} \psi(\overline{z}_{k}^{(t+1)})$ $\bar{z}_k^{(t+1)}(A), \bar{z}_k^{(t)}$ $\chi_k^{(t)}(A), \ldots, \overline{z}_k^{(1)}$ $\binom{1}{k}(A)$). It is easy to verify that

$$
|H(A)| \leq ct\Lambda \cdot \left(1 + \max_{s \in [t+1]} n^{-1} \|\overline{z}^{(s)}\|^2 \right) \leq \left(K\Lambda(1 + \|A\|_{\mathrm{op}})\right)^{ct} \left(1 + \|z^{(0)}\|_{\infty}\right)^c,
$$

and that for any two symmetric matrices $A_1, A_2 \in \mathbb{R}^{n \times n}$,

$$
|H(A_1) - H(A_2)| \le (K\Lambda(1 + ||A_1||_{op} + ||A_2||_{op}))^{ct}(1 + ||z^{(0)}||_{\infty})^c \cdot ||A_1 - A_2||_{op}.
$$

Now we may apply the Gaussian concentration Lemma [A.2](#page-48-0) to conclude. \Box

5.3. Proofs of Theorems [2.5,](#page-11-2) [2.7](#page-12-1) and [2.8.](#page-12-2) We will reduce the asymmetric AMP iterate [\(2.3\)](#page-10-0) into the symmetric AMP iterate [\(2.1\)](#page-7-2). The reduction scheme is well understood, cf. [\[BMN20,](#page-50-10) Section 6]; we spell out some details for the convenience of the reader. Let $z^{(-1)} \equiv \begin{pmatrix} u^{(0)} \\ 0_n \end{pmatrix}$ 0*n* $(x, z^{(0)} \equiv \begin{pmatrix} 0_m \\ v^{(0)} \end{pmatrix}$ $\left(\begin{array}{c} 0_m \\ v^{(0)} \end{array} \right) \in \mathbb{R}^{m+n}$ (recall $z^{(-1)}$ is a dummy variable; the algorithm is initialized at $z^{(0)}$, and for $t = 1, 2, \dots$, let

$$
z^{(2t-1)} \equiv \begin{pmatrix} u^{(t)} \\ 0_n \end{pmatrix}, \quad z^{(2t)} \equiv \begin{pmatrix} 0_m \\ v^{(t)} \end{pmatrix} \in \mathbb{R}^{m+n}.
$$

Let $\overline{F}_{2t-1}, \overline{F}_{2t} : \mathbb{R}^{m+n} \to \mathbb{R}^{m+n}$ be defined via

$$
\overline{\mathsf{F}}_{2t-1}(z^{(2t-1)})\equiv\binom{\mathsf{G}_t(u^{(t)})}{0_n},\quad \overline{\mathsf{F}}_{2t}(z^{(2t)})\equiv\binom{0_m}{\mathsf{F}_t(v^{(t)})}.
$$

Furthermore, recall with $\phi = m/n$, we let

$$
\overline{V} \equiv (\phi^{-1} + 1)^{1/2} \begin{pmatrix} 0_{m \times m} & V \\ V^\top & 0_{n \times n} \end{pmatrix} \in \mathbb{R}^{(m+n) \times (m+n)}.
$$

Using these notation, we may rewrite the asymmetric AMP iterate [\(2.3\)](#page-10-0) into the standard symmetric AMP iterate [\(2.1\)](#page-7-2):

$$
\begin{aligned} z^{(2t+1)} &= (\overline{V} \circ G_{m+n}) \overline{\mathsf{F}}_{2t}(z^{(2t)}) - b_{2t}(\overline{V}) \circ \overline{\mathsf{F}}_{2t-1}(z^{(2t-1)}), \\ z^{(2t+2)} &= (\overline{V} \circ G_{m+n}) \overline{\mathsf{F}}_{2t+1}(z^{(2t+1)}) - b_{2t+1}(\overline{V}) \circ \overline{\mathsf{F}}_{2t}(z^{(2t)}). \end{aligned}
$$

Here for $k \in [m+n]$ and $s \in \mathbb{N}$, $b_{s,k}(\overline{V}) = \frac{1}{m+n} \sum_{\ell \in [m+n]} \overline{V}_{k\ell}^2 \mathbb{E} \overline{F}_{s,\ell}'(z_{\ell}^{(s)})$ $\binom{s}{l}$. It is easy to check that, when $s = 2t$, $b_s(\overline{V}) = \begin{pmatrix} b_t^F \\ 0_n \end{pmatrix}$), and when $s = 2t + 1$, $b_s(\overline{V}) = \begin{pmatrix} 0_m \\ b^{\text{G}} \end{pmatrix}$ $\binom{0_m}{b_{t+1}^G}$. The state evolution in Definition [2.6](#page-12-0) follows from Definition [2.2](#page-8-1) and the above identification. We may now apply Theorems [2.1,](#page-7-1) [2.3](#page-8-0) and [2.4](#page-9-0) to conclude. \square

6. Proofs for Section [3](#page-13-0)

6.1. Proof of Proposition [3.1.](#page-13-2) The following lemma is needed.

Lemma 6.1. *Let* $\mathfrak{d}_{\mathbb{R}} : \mathbb{R}_{>0} \times \mathbb{R}_{>0} \to \mathbb{R}_{>0}$ *defined via* $\mathfrak{d}_{\mathbb{R}}(x, y) = |x - y|^2 / (xy)$ *. The following properties hold for* $\mathfrak{d}_\mathbb{R}$ *:*

- *(1) For any x, y* > 0, $\delta_{\mathbb{R}}(1/x, 1/y) = \delta_{\mathbb{R}}(x, y)$.
- *(2) For any x, y,* $\beta > 0$ *,* $\delta_{\mathbb{R}}(x + \beta, y + \beta) = (1 + \beta/x)^{-1}(1 + \beta/y)^{-1}\delta_{\mathbb{R}}(x, y)$ *.*
- (3) For any $x, y, \alpha \in \mathbb{R}^m_{>0}$, $\delta_{\mathbb{R}}(\sum_{k \in [m]} \alpha_k x_k, \sum_{k \in [m]} \alpha_k y_k) \leq \max_{k \in [m]} \delta_{\mathbb{R}}(x_k, y_k)$.

Proof. (1) and (2) follow by direct calculations; we provide the proof of (3). Let us fix $\alpha, x, y \in \mathbb{R}_{>0}^m$. First note that by AM-GM inequality, we have $(x_iy_ix_jy_j \cdot \delta_{\mathbb{R}}(x_i, y_i)\delta_{\mathbb{R}}(x_j, y_j))^{1/2} \leq 2^{-1}(x_iy_j\delta_{\mathbb{R}}(x_i, y_i) + x_jy_i\delta_{\mathbb{R}}(x_j, y_j)).$ This means

$$
\bigg|\sum_{i\in[m]} \alpha_i x_i - \sum_{j\in[m]} \alpha_j y_j\bigg|^2 \le \sum_{i,j\in[m]} \alpha_i \alpha_j \cdot |x_i - y_i| \cdot |x_j - y_j|
$$

$$
= \sum_{i,j\in[m]} \alpha_i \alpha_j \cdot (x_i y_i x_j y_j \cdot b_{\mathbb{R}}(x_i, y_i) b_{\mathbb{R}}(x_j, y_j))^{1/2}
$$

\n
$$
\leq \frac{1}{2} \Biggl(\sum_{i,j\in[m]} \alpha_i \alpha_j x_i y_j b_{\mathbb{R}}(x_i, y_i) + \sum_{i,j\in[m]} \alpha_i \alpha_j x_j y_i b_{\mathbb{R}}(x_j, y_j) \Biggr)
$$

\n
$$
\leq \max_{k\in[m]} b_{\mathbb{R}}(x_k, y_k) \cdot \Biggl(\sum_{i\in[m]} \alpha_i x_i \Biggr) \cdot \Biggl(\sum_{j\in[m]} \alpha_j y_j \Biggr).
$$

The claim follows by the definition of $\delta_{\mathbb{R}}(\sum_{k \in [m]} \alpha_k x_k, \sum_{k \in [m]} \alpha_k y_k)$

*Proof of Proposition 3.1: Existence and uniqueness of b*_∗*.* We shall first define several notations used in the proof. Let $u = \lambda^{-1}(1_m - b)$. Then viewed as a function of $\lambda > 0$, $u(\lambda) \in \mathbb{R}^m$ satisfies the system of equations

$$
\frac{1}{\mathfrak{u}_k(\lambda)} = \lambda + \sum_{\ell \in [n]} \frac{V_{k\ell}^2}{m} \left(1 + \sum_{k' \in [m]} \frac{V_{k'\ell}^2}{m} \cdot \mathfrak{u}_{k'}(\lambda) \right)^{-1}, \quad k \in [m]. \tag{6.1}
$$

The above system of equations can be rewritten in a vector form as a quadratic vector equation of $u(\lambda)$. The existence and uniqueness of the solution to this type of equation with λ in the upper half complex plane has been discussed in [\[AEK17,](#page-50-15) [AEK19\]](#page-50-16). Below we prove analogous results when λ ranges over the half real line for the above system of equations.

Let us fix some $\eta \in (0, 1)$, and let $\mathbb{R}_{\eta} \equiv [\eta, 1/\eta]$. Fix any pair $0 < \underline{L}_{\eta} \le \overline{L}_{\eta} < \infty$ such that $\underline{L}_{\eta} \leq \eta/(1 + m^{-1}\eta \max_{k \in [m]} ||V_k||^2)$ and $\overline{L}_{\eta} \geq 1/\eta$. Let the function space $\mathfrak{B}_{\eta}(\underline{L}_{\eta}, \overline{L}_{\eta})$ be defined by

$$
\mathfrak{B}_{\eta}(\underline{L}_{\eta},\overline{L}_{\eta})\equiv\Big\{ \mathfrak{u}:\mathbb{R}_{\eta}\rightarrow\mathbb{R}_{\geq0}^m\text{ s.t. }\inf_{\lambda\in\mathbb{R}_{\eta},k\in[m]}\mathfrak{u}_{k}(\lambda)\geq\underline{L}_{\eta},\sup_{\lambda\in\mathbb{R}_{\eta}}\| \mathfrak{u}(\lambda)\|_{\infty}\leq\overline{L}_{\eta}\Big\}.
$$

We will write $\mathfrak{B}_{\eta} \equiv \mathfrak{B}_{\eta}(\underline{L}_{\eta}, \overline{L}_{\eta})$ below for notational simplicity.

Recall $\delta_{\mathbb{R}}$ defined in Lemma [6.1.](#page-39-3) Let $\delta_{\mathfrak{B}_\eta}: \mathfrak{B}_\eta \times \mathfrak{B}_\eta \to \mathbb{R}_{\geq 0}$ be a symmetric, non-negative map defined by

$$
\mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u},\mathfrak{w})\equiv \sup_{\lambda\in\mathbb{R}_{\eta}}\max_{k\in[m]}\mathfrak{d}_{\mathbb{R}}(\mathfrak{u}_{k}(\lambda),\mathfrak{w}_{k}(\lambda))\equiv \sup_{\lambda\in\mathbb{R}_{\eta}}\max_{k\in[m]}\frac{|\mathfrak{u}_{k}(\lambda)-\mathfrak{w}_{k}(\lambda)|^{2}}{\mathfrak{u}_{k}(\lambda)\cdot\mathfrak{w}_{k}(\lambda)}.
$$

The idea of the above construction essentially dates back to [\[EH70\]](#page-51-20). Furthermore, let the map $\Phi : \mathfrak{B}_{\eta} \to (\mathbb{R}_{\eta} \to \mathbb{R}^m_{\geq 0})$ be defined via

$$
\Phi(\mathfrak{u})(\lambda) \equiv \frac{1}{\lambda 1_m + \mathscr{V}_m(1_n + \mathscr{V}_m^{\top} \mathfrak{u}(\lambda))^{-1}}, \quad \mathfrak{u} \in \mathfrak{B}_\eta, \ \lambda \in \mathbb{R}_\eta.
$$

Then the system of nonlinear equations [\(6.1\)](#page-40-0) takes the form $u = \Phi(u)$. (Step 1). We first verify that $\Phi(\mathfrak{B}_\eta) \subset \mathfrak{B}_\eta$, so is well-defined as a map from \mathfrak{B}_η to itself. Note that for any $k \in [m]$, $\lambda \in \mathbb{R}_\eta$ and $u \in \mathfrak{B}_\eta$, we have $\eta \leq \lambda + e_k^\top \mathcal{V}_m(1_n +$ $\mathcal{W}_m^{\top} \mathfrak{u}(\lambda)$ ⁻¹ ≤ $\lambda + e_k^{\top} \mathcal{V}_m \leq \eta^{-1} + ||V_k||^2/m$, so

$$
\frac{1}{\eta} \geq \Phi(\mathfrak{u})_k(\lambda) = \frac{1}{\lambda + e_k^{\top} \mathcal{V}_m(\mathbf{1}_n + \mathcal{V}_m^{\top} \mathfrak{u}(\lambda))^{-1}} \geq \frac{\eta}{1 + m^{-1} \eta \max_{k \in [m]} ||V_k||^2}.
$$

. — П

This verifies $\Phi(\mathfrak{B}_\eta) \subset \mathfrak{B}_\eta$.

(Step 2). Next we prove that for any $u, w \in \mathcal{B}_\eta$, the map $\Phi : \mathcal{B}_\eta \to \mathcal{B}_\eta$ has the following contraction property with respect to $\delta_{\mathfrak{B}_\eta}$: for any $\mathfrak{u}, \mathfrak{w} \in \mathfrak{B}_\eta$,

$$
\mathfrak{d}_{\mathfrak{B}_{\eta}}(\Phi(\mathfrak{u}),\Phi(\mathfrak{w})) \le \left(1 + \frac{1/\overline{L}_{\eta}}{m^{-1}\max_{k\in[m],\ell\in[n]}\|V_{k\cdot}\|^2 \vee \|V_{\cdot\ell}\|^2}\right)^{-4} \cdot \mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u},\mathfrak{w}).\tag{6.2}
$$

To this end, for any $k \in [m], \ell \in [n], \lambda \in \mathbb{R}_\eta$ and $\mathfrak{u} \in \mathfrak{B}_\eta$, let

$$
c_{k;u(\lambda)} \equiv \left(1 + \frac{\lambda}{e_k^{\top} \mathcal{V}_m(1_n + \mathcal{V}_m^{\top}u(\lambda))^{-1}}\right)^{-1} \le \left(1 + \frac{\eta}{\max_{k \in [m]} ||V_k||^2/m}\right)^{-1},
$$

$$
d_{\ell;u(\lambda)} \equiv \left(1 + \frac{1}{e_\ell^{\top} \mathcal{V}_m^{\top}u(\lambda)}\right)^{-1} \le \left(1 + \frac{1/\overline{L}_\eta}{\max_{\ell \in [n]} ||V_{\ell}||^2/m}\right)^{-1}.
$$

Then, for u, $w \in \mathcal{B}_\eta$, by applying Lemma [6.1](#page-39-3) (where (i)-(iii) correspond to (1)-(3) therein respectively), we have for any $k \in [m]$,

$$
\begin{split}\n&\Phi_{\mathbb{R}}(\Phi(\mathfrak{u})_k(\lambda),\Phi(\mathfrak{w})_k(\lambda)) \\
&\stackrel{\text{(i)}}{=} \mathfrak{d}_{\mathbb{R}}\Big(\lambda + e_k^\top \mathscr{V}_m(1_n + \mathscr{V}_m^\top \mathfrak{u}(\lambda))^{-1}, \lambda + e_k^\top \mathscr{V}_m(1_n + \mathscr{V}_m^\top \mathfrak{w}(\lambda))^{-1}\Big) \\
&\stackrel{\text{(ii)}}{=} c_{k;\mathfrak{u}(\lambda)} c_{k;\mathfrak{w}(\lambda)} \cdot \mathfrak{d}_{\mathbb{R}}\Big(e_k^\top \mathscr{V}_m(1_n + \mathscr{V}_m^\top \mathfrak{u}(\lambda))^{-1}, e_k^\top \mathscr{V}_m(1_n + \mathscr{V}_m^\top \mathfrak{w}(\lambda))^{-1}\Big) \\
&\stackrel{\text{(iii)}}{=} c_{k;\mathfrak{u}(\lambda)} c_{k;\mathfrak{w}(\lambda)} \cdot \max_{\ell \in [n]} \mathfrak{d}_{\mathbb{R}}\Big(e_\ell^\top (1_n + \mathscr{V}_m^\top \mathfrak{u}(\lambda))^{-1}, e_\ell^\top (1_n + \mathscr{V}_m^\top \mathfrak{w}(\lambda))^{-1}\Big) \\
&\stackrel{\text{(i)}}{=} c_{k;\mathfrak{u}(\lambda)} c_{k;\mathfrak{w}(\lambda)} \cdot \max_{\ell \in [n]} \mathfrak{d}_{\mathbb{R}}\Big(e_\ell^\top (1_n + \mathscr{V}_m^\top \mathfrak{u}(\lambda)), e_\ell^\top (1_n + \mathscr{V}_m^\top \mathfrak{w}(\lambda))\Big) \\
&\stackrel{\text{(ii)}}{=} c_{k;\mathfrak{u}(\lambda)} c_{k;\mathfrak{w}(\lambda)} \cdot \max_{\ell \in [n]} d_{\ell;\mathfrak{u}(\lambda)} d_{\ell;\mathfrak{w}(\lambda)} \cdot \mathfrak{d}_{\mathbb{R}}\Big(e_\ell^\top \mathscr{V}_m^\top \mathfrak{u}(\lambda), e_\ell^\top \mathscr{V}_m^\top \mathfrak{w}(\lambda)\Big) \\
&\stackrel{\text{(iii)}}{=} c_{k;\mathfrak{u}(\lambda)} c_{k;\mathfrak{w}(\lambda)} \cdot \max_{\ell \in [n]} d_{\ell;\mathfrak{u}(\lambda)} d_{\ell;\mathfrak{w}(\lambda)}
$$

Taking maxima over $k \in [m]$ and $\lambda \in \mathbb{R}_n$ completes the proof of [\(6.2\)](#page-41-0). (Step 3). We now wish to use the contraction property proved in Step 2 to conclude the existence and uniqueness.

We start with the existence proof. For any continuous $u^{(0)} \in \mathcal{B}_{\eta}$, we iteratively define $u^{(t+1)} \equiv \Phi(u^{(t)})$. Then the contraction property in [\(6.2\)](#page-41-0) yields that, for some constants $\varepsilon_0 \equiv \varepsilon_0(V) \in (0, 1)$ and $C_0 = C_0(V, \eta) > 1$,

$$
\mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u}^{(t+1)}, \mathfrak{u}^{(t)}) \le (1 - \varepsilon_0)^t \cdot \mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u}^{(1)}, \mathfrak{u}^{(0)}) \le C_0 (1 - \varepsilon_0)^t. \tag{6.3}
$$

Note that $\delta_{\mathfrak{B}_\eta}$ does *not* define a metric due to the failure of the triangle inequality, so the contraction property with respect to $\delta_{\mathfrak{B}_n}$ does not immediately lead to the existence of the solution. To this end, with the inverse hyperbolic function $\arccosh(x) = \log(x + \sqrt{x^2 - 1})$ defined for $x \ge 1$, let $\exp(x, \sin x) = \exp(x)$ be defined by

$$
e_{\mathfrak{B}_{\eta}}(u, w) \equiv \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{k \in [m]} \operatorname{arcosh}\left(\frac{1}{2} \mathfrak{d}_{\mathbb{R}}(u_k(\lambda), w_k(\lambda)) + 1\right) \equiv \sup_{\lambda \in \mathbb{R}_{\eta}} \left\| \log u(\lambda) - \log w(\lambda) \right\|_{\infty}.
$$

Then $e_{\mathfrak{B}_{\eta}}(\cdot, \cdot)$ is a metric on $\mathfrak{B}_{\eta} \times \mathfrak{B}_{\eta}$ that in particular satisfies the triangle inequality. Moreover, $(\mathfrak{B}_{\eta}, \mathfrak{e}_{\mathfrak{B}_{\eta}})$ is a complete metric space, as $\mathfrak{e}_{\mathfrak{B}_{\eta}}$ induces the same topology as that of L_{∞} on \mathfrak{B}_η . On the other hand, using the simple inequality that for any $z > 0$, arcosh $(1 + z) = \log(1 + z + \sqrt{z^2 + 2z}) \le z + \sqrt{z^2 + 2z} \le 2z + \sqrt{2z}$, we have for u, $w \in \mathfrak{B}_n$,

$$
\mathfrak{e}_{\mathfrak{B}_{\eta}}(\mathfrak{u},\mathfrak{w}) \leq \mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u},\mathfrak{w}) + \mathfrak{d}_{\mathfrak{B}_{\eta}}^{1/2}(\mathfrak{u},\mathfrak{w}).\tag{6.4}
$$

Consequently, for any $t < s$, using triangle inequality for $e_{\mathfrak{B}_\eta}(\cdot, \cdot)$ and [\(6.3\)](#page-41-1)-[\(6.4\)](#page-42-0),

$$
\begin{split} &\mathfrak{e}_{\mathfrak{B}_{\eta}}(\mathfrak{u}^{(s)}, \mathfrak{u}^{(t)}) \leq \sum_{r \in [t:s)} \mathfrak{e}_{\mathfrak{B}_{\eta}}(\mathfrak{u}^{(r+1)}, \mathfrak{u}^{(r)}) \\ & \leq \sum_{r \in [t:s)} \mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u}^{(r+1)}, \mathfrak{u}^{(r)}) + \sum_{r \in [t:s)} \mathfrak{d}_{\mathfrak{B}_{\eta}}^{1/2}(\mathfrak{u}^{(r+1)}, \mathfrak{u}^{(r)}) \leq C_1 (1 - \varepsilon_0)^{t/2}. \end{split}
$$

Here $C_1 = C_1(V, \eta) > 1$. This means $\{u^{(t)}\}$ is a Cauchy sequence in the complete metric space $(\mathfrak{B}_{\eta}, \mathfrak{e}_{\mathfrak{B}_{\eta}})$, and therefore the limit $\mathfrak{u}^{(\infty)} \equiv \lim_{t \to \infty} \mathfrak{u}^{(t)}$ is well defined in $(\mathfrak{B}_{\eta}, \mathfrak{e}_{\mathfrak{B}_{\eta}})$. By continuity of Φ with respect to $\mathfrak{e}_{\mathfrak{B}_{\eta}}$ on \mathfrak{B}_{η} , we may take limit $t \to \infty$ in the equation $u^{(t+1)} = \Phi(u^{(t)})$ to obtain $u^{(\infty)} = \Phi(u^{(\infty)})$. As $u^{(0)}$ is continuous, and Φ preserves continuity, the limit $u^{(\infty)}$ induced by L_{∞} -topology must also be continuous. From [\(6.1\)](#page-40-0), it is easy to see that $\max_{k \in [m]} |u_k^{(\infty)}(\lambda)| < 1/\lambda$, and therefore $b^{(\infty)} \equiv 1_m - \lambda u^{(\infty)}(\lambda) \in [0, 1)^m$ is well-defined. This completes the existence proof.

Next we provide the uniqueness proof. The uniqueness proof does not require the triangle inequality property for $b_{\mathfrak{B}_\eta}(\cdot, \cdot)$, so is much easier. Let us take $\mathfrak{u}, \mathfrak{u}' \in \mathfrak{B}_\eta$ with $u = \Phi(u)$ and $u' = \Phi(u')$. Then the contraction property [\(6.2\)](#page-41-0) shows that there exists some $\varepsilon_0 = \varepsilon_0(V) \in (0, 1)$ such that

$$
0 \leq \mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u},\mathfrak{u}') = \mathfrak{d}_{\mathfrak{B}_{\eta}}(\Phi(\mathfrak{u}),\Phi(\mathfrak{u}')) \leq (1-\varepsilon_0) \cdot \mathfrak{d}_{\mathfrak{B}_{\eta}}(\mathfrak{u},\mathfrak{u}').
$$

This means $\delta_{\mathfrak{B}_\eta}(u, u') = 0$. Using the definition of $\delta_{\mathfrak{B}_\eta}$ and the fact that $u, u' \in \mathfrak{B}_\eta$,

$$
\underline{L}_{\eta}^{-2} \sup_{\lambda \in \mathbb{R}_{\eta}} ||u(\lambda) - u'(\lambda)||_{\infty} \leq b_{\mathfrak{B}_{\eta}}(u, u') = 0.
$$

This implies the uniqueness claim $u \equiv u'$ as elements in \mathfrak{B}_{η} .

Combining both claims, we have proven that for any $\eta \in (0, 1)$, there exists a unique solution $u^* \in \mathfrak{B}_{\eta} \cap C(\mathbb{R}_{\eta} \to \mathbb{R}^m_{\geq 0})$. The claim now follows as $\eta \in (0, 1)$ is $arbitrary.$

Proof of Proposition [3.1:](#page-13-2) Existence and uniqueness of γ_* . Let $b^* = b^*(\lambda)$ be the unique solution of the second equation of (3.1) as proved above. It is easy to obtain the estimate

$$
\max_{k \in [m]} b_k^*(\lambda) \le \left(1 + m\lambda / \max_{k \in [m]} ||V_{k\cdot}||^2\right)^{-1},
$$

$$
\max_{\ell \in [n]} \tau_{b^*(\lambda), \ell} \le \left(1 - \max_{k \in [m]} b_k^*(\lambda)\right)^{-1} \cdot \left(\min_{\ell \in [n]} ||V_{\cdot \ell}||^2 / m\right)^{-1},
$$

$$
\min_{\ell \in [n]} \tau_{b^*(\lambda), \ell} \ge \left(\max_{k \in [m]} ||V_{k\cdot}||^2 / m\right)^{-1}.
$$

In particular, for any $\eta \in (0, 1)$, with $\mathbb{R}_{\eta} = [\eta, 1/\eta]$ as before, we may find some $\varepsilon_0 = \varepsilon_0(V, \eta) \in (0, 1)$ such that

$$
\sup_{\lambda \in \mathbb{R}_{\eta}} \max_{k \in [m]} b_k^*(\lambda) \le 1 - \varepsilon_0, \quad \varepsilon_0 \le \inf_{\lambda \in \mathbb{R}_{\eta}} \min_{\ell \in [n]} \tau_{b^*(\lambda), \ell} \le \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{\ell \in [n]} \tau_{b^*(\lambda), \ell} \le 1/\varepsilon_0. \tag{6.5}
$$

The first equation of (3.1) of interest can be written as

$$
\gamma^{2}(\lambda) = q_{b^{*}(\lambda)} + \mathfrak{D}_{\tau_{b^{*}}(\lambda)}^{2} \mathcal{V}_{m}^{\top} \mathfrak{D}_{1_{m} - b^{*}(\lambda)}^{2} \mathcal{V}_{m} \mathfrak{D}_{1_{n} + \lambda \tau_{b^{*}(\lambda)}}^{-2} \gamma^{2}(\lambda),
$$
(6.6)

where $q_{b^*}: \mathbb{R}_{>0} \to \mathbb{R}_{\geq 0}^n$ is locally bounded. In particular, for any $\eta \in (0, 1)$, there exists $\varepsilon_1 = \varepsilon_1(V, \eta) \in (0, 1)$,

$$
\sup_{\lambda \in \mathbb{R}_\eta} \|q_{b^*}(\lambda)\|_{\infty} \le 1/\varepsilon_1. \tag{6.7}
$$

 $\overline{}$ $\overline{}$ $\overline{}$ $\overline{}$ \overline{a}

We remark here that simply inverting [\(6.6\)](#page-43-0) to solve $\gamma^2(\lambda)$ will not guarantee that the solution is in $\mathbb{R}^n_{\geq 0}$. Equivalently, we may write [\(6.6\)](#page-43-0) as

$$
\zeta(\lambda) = \mathfrak{D}_{\tau_{b^*(\lambda)}}^{-1} q_{b^*(\lambda)} + \mathfrak{D}_{\tau_{b^*(\lambda)}} \mathscr{V}_m^{\top} \mathfrak{D}_{1_m - b^*(\lambda)}^2 \mathscr{V}_m \mathfrak{D}_{\tau_{b^*(\lambda)} \circ (1_n + \lambda \tau_{b^*(\lambda)})^{-2}} \zeta(\lambda)
$$

\n
$$
\equiv \Psi(\zeta)(\lambda), \tag{6.8}
$$

where $\zeta \equiv \mathfrak{D}_{\tau_b}^{-1} \gamma^2 : \mathbb{R}_{>0} \to \mathbb{R}_{\geq 0}^n$. The essential merit of working with ζ instead of γ is that Ψ will be a strict contraction with respect to a natural L_{∞} metric. In particular, consider the function class

$$
\mathfrak{Z}_{\eta} \equiv \Big\{ \zeta : \mathbb{R}_{\eta} \to \mathbb{R}_{\geq 0}^{n} \text{ s.t. } \sup_{\lambda \in \mathbb{R}_{\eta}} ||\zeta(\lambda)||_{\infty} \leq \frac{1}{\varepsilon_{0}^{2} \varepsilon_{1}} \Big\},\,
$$

and the metric (with slight abuse of notation) \mathfrak{d}_{3n} : $\mathfrak{Z}_\eta \times \mathfrak{Z}_\eta \to \mathbb{R}_{\geq 0}$ defined via

$$
\mathfrak{d}_{\mathfrak{Z}_{\eta}}(\zeta_{1},\zeta_{2}) \equiv \sup_{\lambda \in \mathbb{R}_{\eta}} ||\zeta_{1}(\lambda) - \zeta_{2}(\lambda)||_{\infty}.
$$

In the sequel, we shall adopt a similar 3-step strategy as in the proof of existence and uniqueness for *b* ∗ .

(Step 1). First, we shall verify that $\Psi(3_\eta) \subset 3_\eta$. Using the definition of Ψ in [\(6.8\)](#page-43-1), and the estimates in [\(6.5\)](#page-43-2) and [\(6.7\)](#page-43-3), for any $\zeta \in \mathfrak{Z}_n$,

$$
\sup_{\lambda \in \mathbb{R}_{\eta}} \|\Psi(\zeta)(\lambda)\|_{\infty} \n\leq \frac{1}{\varepsilon_{0}\varepsilon_{1}} + \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{\ell \in [n]} \left| \tau_{b^{*},\ell} \sum_{k \in [m]} \frac{V_{k\ell}^{2}}{m} (1 - b_{k}^{*})^{2} \sum_{\ell' \in [n]} \frac{V_{k\ell'}^{2}}{m} \cdot \frac{\tau_{b^{*},\ell'}}{(1 + \lambda \tau_{b^{*},\ell'})^{2}} \zeta_{\ell'}(\lambda) \right| \n\leq \frac{1}{\varepsilon_{0}\varepsilon_{1}} + \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{\ell \in [n]} \tau_{b^{*},\ell} \sum_{k \in [m]} \frac{V_{k\ell}^{2}}{m} (1 - b_{k}^{*})^{2} \cdot \frac{b_{k}^{*}}{1 - b_{k}^{*}} \cdot \sup_{\lambda \in \mathbb{R}_{\eta}} |\zeta(\lambda)| \big|_{\infty} \n\leq \frac{1}{\varepsilon_{0}\varepsilon_{1}} + \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{k \in [m]} k_{k}^{*}(\lambda) \cdot \max_{\ell \in [n]} \left(\tau_{b^{*},\ell} \sum_{k \in [m]} \frac{V_{k\ell}^{2}}{m} (1 - b_{k}^{*}) \right) \cdot \sup_{\lambda \in \mathbb{R}_{\eta}} |\zeta(\lambda)| \big|_{\infty} \n\stackrel{(**)}{=} \frac{1}{\varepsilon_{0}\varepsilon_{1}} + \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{k \in [m]} b_{k}^{*}(\lambda) \cdot \sup_{\lambda \in \mathbb{R}_{\eta}} |\zeta(\lambda)| \big|_{\infty} \n\leq \frac{1}{\varepsilon_{0}\varepsilon_{1}} + \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{k \in [m]} b_{k}^{*}(\lambda) \cdot \sup_{\lambda \in \mathbb{R}_{\eta}} |\zeta(\lambda)| \big|_{\infty}
$$

$$
\leq \frac{1}{\varepsilon_0 \varepsilon_1} + (1 - \varepsilon_0) \cdot \sup_{\lambda \in \mathbb{R}_\eta} ||\zeta(\lambda)||_{\infty} \leq \frac{1}{\varepsilon_0^2 \varepsilon_1}.
$$

Here in (*) we used the characterizing equation for b^* , and in (**) we used the definition of τ_{b^*} . This verifies $\Psi(\mathfrak{Z}_\eta) \subset \mathfrak{Z}_\eta$.

(Step 2). Next, we shall verify the following contraction property for $\Psi : \mathfrak{Z}_\eta \to \mathfrak{Z}_\eta$ with respect to b_{3n} : for any $\zeta_1, \zeta_2 \in \mathcal{S}_n$,

$$
\mathfrak{d}_{\mathfrak{Z}_{\eta}}(\Psi(\zeta_1),\Psi(\zeta_2)) \leq (1-\varepsilon_0) \cdot \mathfrak{d}_{\mathfrak{Z}_{\eta}}(\zeta_1,\zeta_2).
$$

The above display follows by essentially the same calculations as in the Step 1, but without the additive term $1/(\varepsilon_0 \varepsilon_1)$ and replacing ζ therein with $\zeta_1 - \zeta_2$:

$$
\sup_{\lambda \in \mathbb{R}_{\eta}} \left\| \Psi(\zeta_1)(\lambda) - \Psi(\zeta_2)(\lambda) \right\|_{\infty} \n= \sup_{\lambda \in \mathbb{R}_{\eta}} \max_{\ell \in [n]} \left| \tau_{b^*,\ell} \sum_{k \in [m]} \frac{V_{k\ell}^2}{m} (1 - b_k^*)^2 \sum_{\ell' \in [n]} \frac{V_{k\ell'}^2}{m} \cdot \frac{\tau_{b^*,\ell'}}{(1 + \lambda \tau_{b^*,\ell'})^2} (\zeta_{1,\ell'}(\lambda) - \zeta_{2,\ell'}(\lambda)) \right| \n\le (1 - \varepsilon_0) \cdot \sup_{\lambda \in \mathbb{R}_{\eta}} ||\zeta_1(\lambda) - \zeta_2(\lambda)||_{\infty}.
$$

(Step 3). As $(3_{\eta}, \delta_{3_{\eta}})$ is a complete metric space, $\Psi(\zeta) = \zeta$ admits a unique solution $\zeta^* \in \mathcal{S}_\eta$ by the contraction mapping theorem. Clearly ζ^* can be taken as continuous. The claim follows as $\eta \in (0, 1)$ is arbitrary.

6.2. Proof of Theorem [3.2.](#page-14-0) We shall first prove Proposition [3.4.](#page-16-0) To this end, we need two lemmas.

Lemma 6.2. *Suppose for some K* ≥ 2, $m^{-1/2}(\|V_k\| \wedge \|V_{\ell}\|)$ ≥ 1/*K*, $V_{k\ell}$ ≤ *K and hold for all* $k \in [m], \ell \in [n]$ *, and* $1/K \leq \lambda \leq K$ *holds. Then there exists some universal constant c*⁰ > 1 *such that*

$$
\|\theta^{(t)} - \widehat{\theta}\| \le K^{c_0} \cdot (1 \vee \|A\|_{\text{op}}) \cdot (\|u^{(t+1)} - u^{(t)}\| + \|v^{(t+1)} - v^{(t)}\|),
$$

and

$$
||r^{(t)} - \widehat{r}|| \leq K^{c_0} (1 \vee ||A||_{op})^2 \sum_{s \in [t-1]} (1 - K^{-c_0})^{t-1-s} (||u^{(s+1)} - u^{(s)}|| + ||v^{(s+1)} - v^{(s)}||).
$$

Proof. By the first equation of [\(3.2\)](#page-15-1), with $\Xi_r^{(t)} \equiv \mathfrak{D}_{b^* \circ (1_m - b^*)^{-1}}(r^{(t)} - r^{(t+1)}),$

$$
r^{(t+1)} = \mathfrak{D}_{1_m-b^*}^{-1}(A_{b^*}(\theta_{0;b^*}-\theta^{(t)})+\xi_{b^*})+\Xi_r^{(t)}.
$$

Plugging into the second equation of [\(3.2\)](#page-15-1), we have

$$
\theta^{(t+1)} = \text{prox}_{\mathsf{f}_{\tau_{b^*}}} \left[\theta^{(t)} + A_{b^*}^\top \mathfrak{D}_{1_m-b^*}^{-1} \Big(A_{b^*} (\theta_{0;b^*} - \theta^{(t)}) + \xi_{b^*} \Big) + A_{b^*}^\top \Xi_r^{(t)} \right].
$$

Consequently, with

$$
\delta^{(t)} \equiv (1_n + \tau_{b^*} \lambda)^{-1} A_{b^*}^\top \mathfrak{D}_{b^* \circ (1_m - b^*)^{-1}}(r^{(t)} - r^{(t+1)}) + (\theta^{(t)} - \theta^{(t+1)}),
$$

we have

$$
\theta^{(t)} = \text{prox}_{\mathfrak{f}_{r_{b^*}}} \left[\theta^{(t)} + A_{b^*}^{\top} \mathfrak{D}_{1_m - b^*}^{-1} \left(A_{b^*} (\theta_{0; b^*} - \theta^{(t)}) + \xi_{b^*} \right) \right] + \delta^{(t)},
$$

\n
$$
\Leftrightarrow \quad \mathfrak{D}_{\tau_{b^*}}^{1/2} \theta^{(t)} = \text{prox}_{\mathfrak{f}_{r_{b^*}}} \left[\mathfrak{D}_{\tau_{b^*}}^{1/2} \theta^{(t)} + \mathfrak{D}_{\tau_{b^*}} A^{\top} \left(A(\mu_0 - \mathfrak{D}_{\tau_{b^*}}^{1/2} \theta^{(t)}) + \xi \right) \right] + \mathfrak{D}_{\tau_{b^*}}^{1/2} \delta^{(t)},
$$

$$
\Leftrightarrow \quad \lambda(\mathfrak{D}_{\tau_{b^*}}^{1/2}\theta^{(t)}) + A^{\top}(A(\mathfrak{D}_{\tau_{b^*}}^{1/2}\theta^{(t)}) - Y) = \mathfrak{D}_{1_n + \lambda \tau_{b^*}} \mathfrak{D}_{\tau_{b^*}}^{-1/2} \delta^{(t)}.
$$

On the other hand, with $L(\mu) = 2^{-1} (||Y - A\mu||^2 + \lambda ||\mu||^2)$, $\nabla L(\mu) = A^T (A\mu - Y) +$ $λμ$, and $∇² L ≥ λI_n$. Moreover, the above display is equivalent to $∇L(Ω_{τ_{β[*]}}^{1/2}θ^(t))$ = $\mathfrak{D}_{1_n + \lambda \tau_{b^*}} \mathfrak{D}_{\tau_{b^*}}^{-1/2} \delta^{(t)}$. So by the cost optimality,

$$
\mathsf{L}(\mathfrak{D}_{\tau_{b^*}}^{1/2}\theta^{(t)})\geq\mathsf{L}(\widehat{\mu})\geq\mathsf{L}(\mathfrak{D}_{\tau_{b^*}}^{1/2}\theta^{(t)})+\nabla\mathsf{L}(\mathfrak{D}_{\tau_{b^*}}^{1/2}\theta^{(t)})\cdot(\widehat{\mu}-\mathfrak{D}_{\tau_{b^*}}^{1/2}\theta^{(t)})+\frac{\lambda}{2}\big\Vert\widehat{\mu}-\mathfrak{D}_{\tau_{b^*}}^{1/2}\theta^{(t)}\big\Vert^2
$$

.

Under the assumed conditions on *V*, $\|\tau_{b^*}^{-1}\|_{\infty} \vee \|\tau_{b^*}\|_{\infty} \vee \|(1_m - b^*)^{-1}\|_{\infty} \leq K^c$, so

$$
\|\widehat{\mu} - \mathfrak{D}_{\tau_{b^*}}^{1/2} \theta^{(t)}\| \le 2\lambda^{-1} \|\nabla L(\mathfrak{D}_{\tau_{b^*}}^{1/2} \theta^{(t)})\| \le K^c \cdot \|\delta^{(t)}\|
$$

$$
\le K^c \cdot (\|A\|_{\text{op}} \cdot \|r^{(t+1)} - r^{(t)}\| + \|\theta^{(t+1)} - \theta^{(t)}\|).
$$

Now using that $||r^{(t+1)} - r^{(t)}|| = ||u^{(t+1)} - u^{(t)}||$ and that $||\theta^{(t+1)} - \theta^{(t)}|| \le ||v^{(t+1)} - v^{(t)}|| +$ $||A_{b^*}||_{op}||r^{(t+1)} - r^{(t)}||$, we have

$$
\|\theta^{(t)} - \widehat{\theta}\| \le K^c \cdot (1 \vee \|A\|_{\text{op}}) \cdot (\|u^{(t+1)} - u^{(t)}\| + \|v^{(t+1)} - v^{(t)}\|).
$$

For the bound for $||r^{(t)} - \hat{r}||$, using the first equation of [\(3.2\)](#page-15-1),

$$
||r^{(t)} - \widehat{r}|| \le ||A_{b^*}||_{op} ||\theta^{(t-1)} - \widehat{\theta}|| + ||b^*||_{\infty} ||r^{(t-1)} - \widehat{r}||.
$$

Iterating the bound, we arrive at

$$
||r^{(t)} - \widehat{r}|| \le ||A_{b^*}||_{op} \sum_{s \in [t-1]} ||b^*||_{\infty}^{t-1-s} ||\theta^{(s)} - \widehat{\theta}||
$$

$$
\le K^c (1 \vee ||A||_{op})^2 \sum_{s \in [t-1]} (1 - K^{-c})^{t-1-s} (||u^{(s+1)} - u^{(s)}|| + ||v^{(s+1)} - v^{(s)}||),
$$

as desired. \Box

Lemma 6.3. *Suppose* $1/K \le m/n, \lambda \le K$ *for some* $K \ge 2$ *. Further suppose* $m^{-1/2}(\|V_k\| \wedge \|V_{\cdot \ell}\|) \geq 1/K$, and $V_{k\ell} \leq K$ holds for all $k \in [m], \ell \in [n]$. Then *for any D > 1, there exist some universal constant* c_0 > 0 *and another constant* $c_1 = c_1(D) > 0$ *, such that for all* $1 \le t \le \log n/c_0$ *, with* $\mathbb{P}^{\xi}(\cdot|\theta^{(0)})$ *-probability at* $least 1 - c_1 n^{-D}$,

$$
n^{-1} \max \{ ||u^{(t+1)} - u^{(t)}||^2, ||v^{(t+1)} - v^{(t)}||^2 \}
$$

$$
\leq K^{c_0} \cdot n^{-1} (||\mu_0||^2 \vee ||\xi||^2) \cdot (1 - K^{-c_0})^t + (K \log n \cdot (1 + ||z^{(0)}||_{\infty}))^{c_0 t^3} \cdot n^{-1/c_0^t}.
$$

Proof. In this proof the expectation is all taken conditional on ξ . First, with $(\sigma_u^{(s)})^2 \equiv (\text{Var}(U_k^{(s)})$ (k) *k*∈[*m*] ∈ ℝ^{*m*} and $(\sigma_v^{(s)})^2$ </sup> ≡ (Var($V_{\ell}^{(s)}$) $(\epsilon^{(s)})_{\ell \in [n]} \in \mathbb{R}^n$, using the state evolution in Definition [2.6,](#page-12-0) the first equation therein reduces to

$$
\begin{aligned} (\sigma_u^{(t+1)})^2 &= \big(\mathfrak{D}_{1_m-b^*}\mathscr{V}_m\mathfrak{D}_{\tau_{b^*}}\big) \,\mathbb{E}\, \mathsf{F}_t^2(\sigma_v^{(t)} Z_n) \\ &= \big(\mathfrak{D}_{1_m-b^*}\mathscr{V}_m\mathfrak{D}_{\tau_{b^*}}\big)\mathfrak{D}_{1+\lambda\tau_{b^*}}^{-2}\big(\mathfrak{D}_{\lambda\tau_{b^*}}^2\theta_{0;b^*}^2 + (\sigma_v^{(t)})^2\big), \end{aligned}
$$

and the second equation reduces to

$$
(\sigma_v^{(t+1)})^2 = (\mathfrak{D}_{1_m-b^*} \mathscr{V}_m \mathfrak{D}_{\tau_{b^*}})^\top \mathbb{E} \mathsf{G}_{t+1}^2(\sigma_u^{(t+1)} Z_m)
$$

= $\mathfrak{D}_{\tau_{b^*}} \mathscr{V}_m^\top \mathfrak{D}_{1_m-b^*} [(\sigma_u^{(t+1)})^2 + \xi_{b^*}^2]$

A LEAVE-ONE-OUT APPROACH TO AMP 47

$$
= \mathfrak{D}_{\tau_{b^*}} \mathscr{V}_m^{\top} \mathfrak{D}_{1_m-b^*}^2 (\mathscr{V}_m \mathfrak{D}_{\lambda \tau_{b^*} \circ (1+\lambda \tau_{b^*})^{-1}}^2 \mu_0^2 + \xi^2) + \mathfrak{D}_{\tau_{b^*}} \mathscr{V}_m^{\top} \mathfrak{D}_{1_m-b^*}^2 \mathscr{V}_m \mathfrak{D}_{1+\lambda \tau_{b^*}}^{-2} \mathfrak{D}_{\tau_{b^*}} (\sigma_v^{(t)})^2.
$$

Comparing the above display to [\(6.8\)](#page-43-1) and the proofs therein,

$$
\|(\sigma_{\nu}^{(t+1)})^2 - \zeta_{\nu}^*\|_{\infty} \leq K^c (1 - K^{-c})^t \cdot \|(\sigma_{\nu}^{(1)})^2 - (\sigma_{\nu}^{(0)})^2\|_{\infty}.
$$

Here (with slight abuse of notation) $\zeta_v^* = \zeta_v^*(\lambda) \in \mathbb{R}_{\geq 0}^n$ is the unique fixed point for the equation (6.8) . Under the assumed conditions on V ,

$$
\| (\sigma_{\nu}^{(t+1)})^2 - \zeta_{\nu}^* \|_{\infty} \le K^c \cdot m^{-1} (\|\mu_0\|^2 \vee \|\xi\|^2) \cdot (1 - K^{-c})^t. \tag{6.9}
$$

Next, let us define the correlation vectors $\mathbf{r}_{u}^{(s+1)} \equiv (\text{Cov}(U_k^{(s+1)})$ $\binom{(s+1)}{k}$, $U_k^{(s)}$ $\binom{n}{k}$)_{$k \in [m]$} ∈ \mathbb{R}^m and $r_v^{(s+1)} = (Cov(V_\ell^{(s+1)})$ $v_{\ell}^{(s+1)}, V_{\ell}^{(s)}$ $(\epsilon^{(s)})_{\ell \in [n]} \in \mathbb{R}^n$. Using the state evolution in Definition [2.6](#page-12-0) again, with some calculations along the above lines we obtain the exact same recursion:

$$
\mathbf{r}_{v}^{(t+1)} = \mathfrak{D}_{\tau_{b^*}} \mathscr{V}_m^{\top} \mathfrak{D}_{1_m-b^*}^2 (\mathscr{V}_m \mathfrak{D}_{\lambda \tau_{b^*} \circ (1 + \lambda \tau_{b^*})^{-1}}^2 \mu_0^2 + \xi^2) + \mathfrak{D}_{\tau_{b^*}} \mathscr{V}_m^{\top} \mathfrak{D}_{1_m-b^*}^2 \mathscr{V}_m \mathfrak{D}_{1 + \lambda \tau_{b^*}}^{-2} \mathfrak{D}_{\tau_{b^*}} \mathfrak{r}_{v}^{(t)}.
$$

While the difference lies in the initialization, the following bound remains valid:

$$
\|\mathbf{r}_{\nu}^{(t+1)} - \zeta_{\nu}^*\|_{\infty} \le K^c \cdot m^{-1} (\|\mu_0\|^2 \vee \|\xi\|^2) \cdot (1 - K^{-c})^t. \tag{6.10}
$$

Similarly, with $\zeta_u^* \equiv (\mathfrak{D}_{1_m-b^*} \mathcal{V}_m \mathfrak{D}_{\tau_{b^*}}) \mathfrak{D}_{1+\lambda \tau_{b^*}}^{-2} (\mathfrak{D}_{\lambda \tau_{b^*}}^2 \theta_{0;b^*}^2 + \zeta_v^*),$ we have the estimate

$$
\| (\sigma_u^{(t+1)})^2 - \zeta_u^* \|_{\infty} \vee \| \mathfrak{r}_u^{(t+1)} - \zeta_u^* \|_{\infty} \leq K^c \cdot m^{-1} (\|\mu_0\|^2 \vee \|\xi\|^2) \cdot (1 - K^{-c})^t. \tag{6.11}
$$

Now using the above display [\(6.11\)](#page-46-0) and Theorem [2.8,](#page-12-2) with $c_0 > 0$ a proper universal constant, $c_1 = c_1(D) > 0$ depending on $D > 1$ only, and $err_n^{(t)} \equiv$ $(c_1 K \log n_\phi \cdot (1 + ||z^{(0)}||_\infty))^{c_0 t^3} \cdot n^{-1/c_0^t}$, we have with probability $1 - c_1 n^{-D}$,

$$
n_{\phi}^{-1} ||u^{(t+1)} - u^{(t)}||^2 \le n_{\phi}^{-1} \mathbb{E} ||U^{(t+1)} - U^{(t)}||^2 + \text{err}_n^{(t)}
$$

\n
$$
\le K^c \cdot \max_{k \in [m]} |(\sigma_{u,k}^{(t+1)})^2 + (\sigma_{u,k}^{(t)})^2 - 2\mathfrak{r}_{u,k}^{(t+1)}| + \text{err}_n^{(t)}
$$

\n
$$
\le K^c \cdot m^{-1} (||\mu_0||^2 \vee ||\xi||^2) \cdot (1 - K^{-c})^t + \text{err}_n^{(t)}.
$$

A completely similar argument via the help of [\(6.9\)](#page-46-1)-[\(6.10\)](#page-46-2) establishes the same estimate for $n_{\phi}^{-1} ||v^{(t+1)} - v^{(t)}||^2$. The contract of the contract

Proof of Proposition [3.4.](#page-16-0) The claim follows from Lemmas [6.2](#page-44-1) and [6.3](#page-45-0) and the standard subgaussian tail estimate for $||A||_{op}$, cf. Lemma [A.1.](#page-47-1)

Now we may prove Theorem [3.2.](#page-14-0)

Proof of Theorem [3.2.](#page-14-0) In this proof expectation is all taken conditional on ξ. Constants unspecified in the proof may depend on *K*, Λ . Recall $\widehat{\mu} = (A^{\top}A + \lambda I)^{-1}A^{\top}Y$, so $\|\widehat{\mu}\| \leq \lambda^{-1} (1 + \|A\|_{op})^2 (\|\mu_0\| \vee \|\xi\|)$. By subgaussian tail of $\|A\|_{op}$ (cf. Lemma [A.1\)](#page-47-1), with probability at least $1 - e^{-n/c}$,

$$
n^{-1/2}|\widehat{\mu}|| \leq c \cdot n^{-1/2}(\|\mu_0\| \vee \|\xi\|).
$$

So combined with Proposition [3.4,](#page-16-0) there exists some $\varepsilon_0 = \varepsilon_0(K) \in (0, 1)$ such that with probability $1 - e^{-n/c}$,

$$
\left| \frac{1}{n} \sum_{j \in [n]} \psi(\widehat{\mu}_j, \mu_{0,j}) - \frac{1}{n} \sum_{j \in [n]} \psi(\mu_j^{(t)}, \mu_{0,j}) \right|
$$
\n
$$
\leq c \cdot (1 + n^{-1/2} (||\widehat{\mu}|| \vee ||\mu^{(t)}|| \vee ||\mu_{0}||)) \cdot n^{-1/2} ||\mu^{(t)} - \widehat{\mu}||
$$
\n
$$
\leq c \cdot \left(1 + n^{-1/2} (||\widehat{\mu}|| \vee ||\mu_{0}||) + n^{-1/2} ||\mu^{(t)} - \widehat{\mu}||\right) \cdot n^{-1/2} ||\mu^{(t)} - \widehat{\mu}||
$$
\n
$$
\leq c \cdot (L_{\mu_0, \xi} + \text{err}_n^{(t)}) \cdot (L_{\mu_0, \xi} (1 - \varepsilon_0)^t + \text{err}_n^{(t)}).
$$
\n(6.12)

Here $\text{err}_n^{(t)}(c_1) \equiv (c_1 \log n)^{c_0 t^3} \cdot n^{-1/c_0^t}, \text{err}_n^{(t)} \equiv \text{err}_n^{(t)}(1).$

On the other hand, by Proposition [3.3,](#page-15-3) $\mu^{(t)} = \text{prox}_{\lambda \| \cdot \|^{2}/2}(\mu_0 - \mathfrak{D}_{\tau_{b^*}}^{1/2} v^{(t)}; \tau_{b^*}) =$ $(1_n + \lambda \tau_{b^*})^{-1} \circ (\mu_0 - \mathfrak{D}_{\tau_{b^*}}^{1/2} \mathcal{V}^{(t)})$. Now applying Theorem [2.8](#page-12-2) to $\{\mathcal{V}^{(t)}\}$ and recalling [\(6.9\)](#page-46-1) with $(\zeta_v^*)^{1/2} = \mathfrak{D}_{\tau_{b^*}}^{-1/2} \gamma^*$ therein, for $t \leq \log n/c$, with probability at least $1 - c_1 n^{-D}$ where $c_1 = c_1(D) > 0$,

$$
\left| \frac{1}{n} \sum_{j \in [n]} \psi(\mu_j^{(t)}, \mu_{0,j}) - \frac{1}{n} \sum_{j \in [n]} \mathbb{E} \psi(\text{prox}_{\lambda | \cdot |^2 / 2} (\mu_{0,j} + \gamma_j^* Z; \tau_{b^*,j}), \mu_{0,j}) \right|
$$

 $\leq c \cdot (\text{err}_n^{(t)}(c_1) + L_{\mu_0, \xi} \cdot (1 - \varepsilon_0)^t).$ (6.13)

By noting $prox_{\lambda|\cdot|^2/2} (\mu_{0,j} + \gamma_j^* Z; \tau_{b^*,j}) = \widehat{\mu}_j^{\text{seq}}$ $j_j^{\text{seq}}(\gamma^*; \tau_{b^*})$, combining [\(6.12\)](#page-47-2) and [\(6.13\)](#page-47-3), for $t \le \log n/c$, with probability at least $1 - c_1 n^{-D}$ where $c_1 = c_1(D) > 0$,

$$
\left| \frac{1}{n} \sum_{j \in [n]} \psi(\widehat{\mu}_j, \mu_{0,j}) - \frac{1}{n} \sum_{j \in [n]} \mathbb{E} \psi(\widehat{\mu}_j^{\text{seq}}(\gamma^*; \tau_{b^*}), \mu_{0,j}) \right|
$$

 $\leq c \cdot (L_{\mu_0, \xi} + \text{err}_n^{(t)}) \cdot (L_{\mu_0, \xi}(1 - \varepsilon_0)^t + \text{err}_n^{(t)}(c_1)).$

The claim follows by choosing, e.g., $t = (\log \log n)^{0.99}$

. В последните последните село в последните село в сел
Село в село в село

Appendix A. Auxiliary results

Let $\mathcal{M}_n(\mathbb{R})$ be the space of $n \times n$ real symmetric matrices. The following result is frequently referred to 'subgaussian tail of $||A||_{op}$ ' in the proof.

Lemma A.1. *Let the upper triangular elements of* $G_n \in \mathcal{M}_n(\mathbb{R})$ *be i.i.d.* $N(0, 1/n)$ *, and* $V \in \mathcal{M}_n(\mathbb{R})$ *be a deterministic matrix such that* $0 \leq V_{ij} \leq K$ *for some* $K > 1$ *. Let* $A = V \circ G_n$ *. Then there exists some universal constant* $c_0 > 0$ *such that the following hold:*

(1) For $x \ge 0$, $\mathbb{P}(|A||_{op}/K \ge c_0 + \sqrt{x/n}) \le c_0 e^{-x/c_0}$. *(2) For* $t \le n/(c_0 \log n)$ *,* $\mathbb{E}(\|A\|_{op}/K)^t \le c_0^t$.

Proof. (1) follows from, e.g., [\[Ver18,](#page-53-18) Theorem 4.4.5]. For (2), let $E_0 = {\frac{||A||_{op}}{K \le \epsilon}}$ *c*₁}. Then $\mathbb{P}(E_0^c) \le c_1 e^{-n/c_1}$, and therefore for the prescribed range of *t*,

$$
\mathbb{E}(\|A\|_{\text{op}}/K)^{t} \le c_{1}^{t} + \mathbb{E}(\|A\|_{\text{op}}/K)^{t} \mathbf{1}_{E_{0}^{c}} \le c_{1}^{t} + (c_{2}n^{1/2})^{t} e^{-n/c_{2}} \le c_{3}^{t}
$$

as desired. \Box

,

The following version of the Gaussian concentration inequality, allowing for 'high probability bounded Lipschitz constant', is useful.

Lemma A.2. Let $H : \mathcal{M}_n(\mathbb{R}) \to \mathbb{R}$ be a measurable map. Suppose there exist $t, L \geq 2$ *such that the following hold:*

- *(1) For all* $A \in \mathcal{M}_n(\mathbb{R})$, $|H(A)| \le L(1 + ||A||_{op})^t$.
- *(2) For all* $A_1, A_2 \in \mathcal{M}_n(\mathbb{R})$,

$$
|H(A_1) - H(A_2)| \le L(1 + ||A_1||_{op} + ||A_2||_{op})^t \cdot ||A_1 - A_2||_{op}.
$$

Let the upper triangular elements of $G_n \in \mathcal{M}_n(\mathbb{R})$ *be i.i.d.* $N(0, 1/n)$ *, and* $V \in$ $\mathcal{M}_n(\mathbb{R})$ *be a deterministic matrix such that* $0 \leq V_{ij} \leq K$ *for some* $K \geq 2$ *. Then there exists some universal constant* $c_0 > 0$ *such that if* $t \leq n/(c_0 \log n)$ *and* $x \geq$ $c_0 L K^{c_0 t} e^{-n/c_0},$

$$
\mathbb{P}\left(|H(V\circ G_n)-\mathbb{E}\,H(V\circ G_n)|\geq x\right)\leq c_0e^{-n\cdot\min\{x^2/L^2K^{c_0t},1/c_0\}}.
$$

Moreover, there exists some universal $c_1 > 0$ *such that for* $t \leq n/(c_1 \log n)$,

$$
\text{Var}\left(H(V\circ G_n)\right) \leq L^2 K^{c_1 t} \log n \cdot n^{-1}.
$$

Proof. The proof below is inspired by that of [\[CL21,](#page-50-19) Lemma 3.5]. Fix $M > 0$ to be chosen later. Let

$$
\mathscr{H}(A) \equiv \inf_{A' \in \mathscr{M}_n(\mathbb{R}), ||A'||_{\text{op}} \le M} \Big\{ H(A') + L \cdot \Gamma(A, A') (||A - A'||_{\text{op}} \wedge (2M)) \Big\},\
$$

where

$$
\Gamma(A, A') \equiv \left(\left(\left\| A \right\|_{\text{op}} \wedge M \right) + \left\| A' \right\|_{\text{op}} + 1 \right)^t.
$$

Note that for any $A \in \mathcal{M}_n(\mathbb{R})$ with $||A||_{op} \leq M$, the definition of \mathcal{H} entails that $\mathcal{H}(A) \leq H(A)$. On the other hand, the condition (2) implies that $H(A) \leq \mathcal{H}(A)$. In summary,

$$
\mathcal{H}(A) = H(A) \text{ for all } A \in \mathcal{M}_n(\mathbb{R}) \text{ such that } ||A||_{op} \le M. \tag{A.1}
$$

Next, for $A_1, A_2 \in \mathcal{M}_n(\mathbb{R}),$

$$
|\mathcal{H}(A_1) - \mathcal{H}(A_2)|/L
$$

\n
$$
\leq \sup_{A' \in \mathcal{M}_n(\mathbb{R}), \|A'\|_{op} \leq M} \left| \Gamma(A_1, A')(\|A_1 - A'\|_{op} \wedge 2M) - \Gamma(A_2, A')(\|A_2 - A'\|_{op} \wedge 2M) \right|
$$

\n
$$
\leq 2M \cdot \sup_{A' \in \mathcal{M}_n(\mathbb{R}), \|A'\|_{op} \leq M} \left| \Gamma(A_1, A') - \Gamma(A_2, A') \right| + (2M + 1)^t \cdot \|A_1 - A_2\|_{op}
$$

\n
$$
\leq 2t(2M + 1)^t \cdot \|A_1 - A_2\|_{op} \leq 2t(2M + 1)^t \cdot \|A_1 - A_2\|_{F}.
$$

For the random matrix model $A = V \circ G_n$, we then have

$$
|\mathscr{H}(V\circ G_{n,1})-\mathscr{H}(V\circ G_{n,2})|\leq KLM^{c_0t}\cdot ||G_{n,1}-G_{n,2}||_F.
$$

Using Gaussian concentration inequality, for any $x > 0$, $M > c'K$,

$$
2 \exp \left(-\frac{nx^2}{(KL)^2 M^{ct}}\right) \ge \mathbb{P}\left(|\mathcal{H}(A) - \mathbb{E}|\mathcal{H}(A)| \ge x\right)
$$

$$
\ge \mathbb{P}\left(|\mathcal{H}(A) - \mathbb{E}|\mathcal{H}(A)| \ge x, \|A\|_{\text{op}} \le M\right)
$$

$$
\geq \mathbb{P}\left(|H(A) - \mathbb{E}\,\mathcal{H}(A)| \geq x\right) - ce^{-nM^2/cK^2}.\tag{A.2}
$$

Here in the last inequality we used [\(A.1\)](#page-48-1) and the subgaussian tail estimate for $||A||_{op}$, cf. Lemma [A.1.](#page-47-1) On the other hand, as $E \mathcal{H}(A) = E H(A) 1_{||A||_{op} \leq M}$ + $\mathbb{E}\mathcal{H}(A)\mathbf{1}_{\|A\|_0}$, using the condition (1), we have for $t \leq n/(c \log n)$,

$$
\left| \mathbb{E} \mathcal{H}(A) - \mathbb{E} H(A) \right| \leq \mathbb{E} \left(|\mathcal{H}(A)| + |H(A)| \right) \mathbf{1}_{\|A\|_{op} \geq M}
$$

\n
$$
\leq \mathbb{E}^{1/2} \left(|H(0)| + L(1 + \|A\|_{op})^{t+1} + |H(A)| \right)^{2} \cdot \mathbb{P}^{1/2} (\|A\|_{op} \geq M)
$$

\n
$$
\leq cL \mathbb{E} (1 + \|A\|_{op})^{ct} \cdot e^{-nM^{2}/cK^{2}} \leq L K^{c_{1}t} e^{-nM^{2}/c_{1}K^{2}}.
$$

So for any $x > 0$, $M > c'_1 K$ such that $x \ge L K^{c_1 t} e^{-nM^2/c_1 K^2}$, and $t \le n/(c_2 \log n)$,

$$
\mathbb{P}\left(|H(A)-\mathbb{E} H(A)|\geq 2x\right)\leq c_2\exp\left(-n\cdot\min\left\{\frac{x^2}{c_2(KL)^2M^{c_2t}}\wedge\frac{M^2}{c_2K^2}\right\}\right),
$$

proving the subgaussian estimate by choosing $M = c'_2 K$ for a large universal c'_2 1, and adjusting the constant.

For the variance bound, note that for a large enough, universal $c_3 > 0$, the above inequality yields that the event $E_0 = \left\{ |H(A) - \mathbb{E}[H(A)] \right\} \le L K^{c_3 t} \sqrt{\log n} \cdot n^{-1/2}$ occurs with $\mathbb{P}(E_0) \ge 1 - c_3 n^{-100}$. Consequently, for $t \le n/(c_3 \log n)$,

$$
\begin{aligned} &\text{Var}(H(A)) \le L^2 K^{ct} \log n \cdot n^{-1} + \mathbb{E}|H(A) - \mathbb{E}|H(A)|^2 \mathbf{1}_{E_0^c} \\ &\le L^2 K^{ct} \log n \cdot n^{-1} + L^2 \mathbb{E}^{1/2} (1 + ||A||_{op})^{ct} \cdot \mathbb{P}^{1/2}(E_0^c) \le L^2 K^{c_4 t} \log n \cdot n^{-1}, \end{aligned}
$$

as desired.

We also need the following elementary lemma on the matrix square root.

Lemma A.3. Let $\Sigma_1, \Sigma_2 \in \mathbb{R}^{t \times t}$ be two positive semi-definite matrices. Then it *holds that* $\|\Sigma_1^{1/2}\|$ $\frac{1}{2}$ – $\Sigma_2^{1/2}$ $\frac{1}{2}$ ^{1/2} $\Big| \frac{2}{F} \leq$ $\sqrt{t} \cdot ||\Sigma_1 - \Sigma_2||_F.$

Proof. The result is well-known, see e.g., [\[Wih09,](#page-53-19) Eqn. (3.2)] for a more general formulation. For the convenience of the reader, we provide a direct and short proof below. Let $\Sigma_{\ell} = U_{\ell} \Lambda_{\ell} U_{\ell}^{\top}$ ($\ell = 1, 2$) be the spectral decomposition of Σ_{ℓ} . By writing $\Lambda_{\ell} = \text{diag}(\{\lambda_{\ell,i}\}_{i \in [t]})$ where $\lambda_{\ell,i} \geq 0$, we have

$$
\begin{split} \left\| \Sigma_{1}^{1/2} - \Sigma_{2}^{1/2} \right\|_{F}^{2} &= \left\| U_{1} \Lambda_{1}^{1/2} U_{1}^{\top} - U_{2} \Lambda_{2}^{1/2} U_{2}^{\top} \right\|_{F}^{2} = \left\| \Lambda_{1}^{1/2} U_{1}^{\top} U_{2} - U_{1}^{\top} U_{2} \Lambda_{2}^{1/2} \right\|_{F}^{2} \\ &= \sum_{i,j \in [t]} \left(\lambda_{1,i}^{1/2} - \lambda_{2,j}^{1/2} \right)^{2} (U_{1}^{\top} U_{2})_{ij}^{2} \stackrel{(*)}{\leq} \sum_{i,j \in [t]} \left| \lambda_{1,i} - \lambda_{2,j} \right| (U_{1}^{\top} U_{2})_{ij}^{2} \\ &\leq \left(\sum_{i,j \in [t]} \left(\lambda_{1,i} - \lambda_{2,j} \right)^{2} (U_{1}^{\top} U_{2})_{ij}^{2} \right)^{1/2} \cdot \left\| U_{1}^{\top} U_{2} \right\|_{F} = \sqrt{t} \cdot \left\| \Sigma_{1} - \Sigma_{2} \right\|_{F}. \end{split}
$$

Here in (*) we used the fact that for any $a, b \ge 0$, $(\sqrt{a} - \sqrt{b})^2 \le |a - b|$.

ACKNOWLEDGMENTS

The research of Z. Bao is partially supported by Hong Kong RGC Grant 16303922, NSFC12222121 and NSFC12271475. The research of Q. Han is partially supported by NSF Grant DMS-2143468. The research of X. Xu is partially supported by Hong Kong RGC Grant 16305421.

REFERENCES

- [AEK17] Johannes Alt, László Erdős, and Torben Krüger, *Local law for random Gram matrices*, Electron. J. Probab. 22 (2017), Paper No. 25, 41.
- [AEK19] Oskari Ajanki, László Erdős, and Torben Krüger, *Quadratic vector equations on complex upper half-plane*, vol. 261, American Mathematical Society, 2019.
- [BES17] Zhigang Bao, László Erdős, and Kevin Schnelli, *Local law of addition of random matrices on optimal scale*, Comm. Math. Phys. 349 (2017), no. 3, 947–990.
- [BGK16] Florent Benaych-Georges and Antti Knowles, *Lectures on the local semicircle law for wigner matrices*, arXiv preprint arXiv:1601.04055 (2016).
- [BKM⁺19] Jean Barbier, Florent Krzakala, Nicolas Macris, Léo Miolane, and Lenka Zdeborová, *Optimal errors and phase transitions in high-dimensional generalized linear models*, Proc. Natl. Acad. Sci. 116 (2019), no. 12, 5451–5460.
- [BKRS21] Zhiqi Bu, Jason M. Klusowski, Cynthia Rush, and Weijie J. Su, *Algorithmic analysis and statistical estimation of SLOPE via approximate message passing*, IEEE Trans. Inform. Theory 67 (2021), no. 1, 506–537.
- [BLM15] Mohsen Bayati, Marc Lelarge, and Andrea Montanari, *Universality in polytope phase transitions and message passing algorithms*, Ann. Appl. Probab. 25 (2015), no. 2, 753– 822.
- [BM11] Mohsen Bayati and Andrea Montanari, *The dynamics of message passing on dense graphs, with applications to compressed sensing*, IEEE Trans. Inform. Theory 57 (2011), no. 2, 764–785.
- [BM12] , *The LASSO risk for Gaussian matrices*, IEEE Trans. Inform. Theory 58 (2012), no. 4, 1997–2017.
- [BMN20] Raphaël Berthier, Andrea Montanari, and Phan-Minh Nguyen, *State evolution for approximate message passing with non-separable functions*, Inf. Inference 9 (2020), no. 1, 33–79.
- [BNSX22] Erwin Bolthausen, Shuta Nakajima, Nike Sun, and Changji Xu, *Gardner formula for Ising perceptron models at small densities*, Conference on Learning Theory, PMLR, 2022, pp. 1787–1911.
- [Bol14] Erwin Bolthausen, *An iterative construction of solutions of the TAP equations for the Sherrington-Kirkpatrick model*, Comm. Math. Phys. 325 (2014), no. 1, 333–366.
- [Bol19] , *A Morita type proof of the replica-symmetric formula for SK*, Statistical mechanics of classical and disordered systems, Springer Proc. Math. Stat., vol. 293, Springer, Cham, 2019, pp. 63–93.
- [BR22] Joshua K Behne and Galen Reeves, *Fundamental limits for rank-one matrix estimation with groupwise heteroskedasticity*, International Conference on Artificial Intelligence and Statistics, PMLR, 2022, pp. 8650–8672.
- [BS10] Zhidong Bai and Jack W. Silverstein, *Spectral analysis of large dimensional random matrices*, second ed., Springer Series in Statistics, Springer, New York, 2010.
- [CFM23] Michael Celentano, Zhou Fan, and Song Mei, *Local convexity of the TAP free energy and AMP convergence for* \mathbb{Z}_2 -synchronization, Ann. Statist. **51** (2023), no. 2, 519–546.
- [CL21] Wei-Kuo Chen and Wai-Kit Lam, *Universality of approximate message passing algorithms*, Electron. J. Probab. 26 (2021), Paper No. 36, 44.
- [CM22a] Michael Celentano and Andrea Montanari, *Fundamental barriers to high-dimensional regression with convex penalties*, Ann. Statist. 50 (2022), no. 1, 170–196.

52 Z. BAO, Q. HAN, AND X. XU

- [CM22b] Chen Cheng and Andrea Montanari, *Dimension free ridge regression*, arXiv preprint arXiv:2210.08571 (2022).
- [CMW23] Michael Celentano, Andrea Montanari, and Yuting Wei, *The Lasso with general Gaussian designs with applications to hypothesis testing*, Ann. Statist., to appear. Available at arXiv:2007.13716v2 (2023+).
- [CR23] Collin Cademartori and Cynthia Rush, *A non-asymptotic analysis of generalized approximate message passing algorithms with right rotationally invariant designs*, arXiv preprint arXiv:2302.00088 (2023).
- [DAM17] Yash Deshpande, Emmanuel Abbe, and Andrea Montanari, *Asymptotic mutual information for the balanced binary stochastic block model*, Inf. Inference 6 (2017), no. 2, 125–170.
- [Dic16] Lee H. Dicker, *Ridge regression and asymptotic minimax estimation over spheres of growing dimension*, Bernoulli 22 (2016), no. 1, 1–37.
- [DLS23] Rishabh Dudeja, Yue M. Lu, and Subhabrata Sen, *Universality of approximate message passing with semirandom matrices*, Ann. Probab. 51 (2023), no. 5, 1616–1683.
- [DM14] Yash Deshpande and Andrea Montanari, *Information-theoretically optimal sparse PCA*, 2014 IEEE International Symposium on Information Theory, IEEE, 2014, pp. 2197– 2201.
- [DM16] David Donoho and Andrea Montanari, *High dimensional robust M-estimation: asymptotic variance via approximate message passing*, Probab. Theory Related Fields 166 (2016), no. 3-4, 935–969.
- [DMM09] David L Donoho, Arian Maleki, and Andrea Montanari, *Message-passing algorithms for compressed sensing*, Proceedings of the National Academy of Sciences 106 (2009), no. 45, 18914–18919.
- [DS19] Jian Ding and Nike Sun, *Capacity lower bound for the Ising perceptron*, STOC'19— Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing, ACM, New York, 2019, pp. 816–827.
- [DW18] Edgar Dobriban and Stefan Wager, *High-dimensional asymptotics of prediction: ridge regression and classification*, Ann. Statist. 46 (2018), no. 1, 247–279.
- [EAMS21] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke, *Optimization of mean-field spin glasses*, Ann. Probab. 49 (2021), no. 6, 2922–2960.
- [EH70] Clifford J. Earle and Richard S. Hamilton, *A fixed point theorem for holomorphic mappings*, Global Analysis (Proc. Sympos. Pure Math., Vols. XIV, XV, XVI, Berkeley, Calif., 1968), Proc. Sympos. Pure Math., vol. XIV-XVI, Amer. Math. Soc., Providence, RI, 1970, pp. 61–65.
- [EK18] Noureddine El Karoui, *On the impact of predictor geometry on the performance on highdimensional ridge-regularized generalized robust regression estimators*, Probab. Theory Related Fields 170 (2018), no. 1-2, 95–175.
- [EKS19] László Erdős, Torben Krüger, and Dominik Schröder, *Random matrices with slow correlation decay*, Forum Math. Sigma 7 (2019), Paper No. e8, 89.
- [EYY12] László Erdős, Horng-Tzer Yau, and Jun Yin, *Bulk universality for generalized Wigner matrices*, Probab. Theory Related Fields 154 (2012), no. 1-2, 341–407.
- [Fan22] Zhou Fan, *Approximate message passing algorithms for rotationally invariant matrices*, Ann. Statist. 50 (2022), no. 1, 197–224.
- [FLS22] Zhou Fan, Yufan Li, and Subhabrata Sen, *TAP equations for orthogonally invariant spin glasses at high temperature*, arXiv preprint arXiv:2202.09325 (2022).
- [FR18] Alyson K. Fletcher and Sundeep Rangan, *Iterative reconstruction of rank-one matrices in noise*, Inf. Inference 7 (2018), no. 3, 531–562.
- [FVRS22] Oliver Y. Feng, Ramji Venkataramanan, Cynthia Rush, and Richard J. Samworth, *A unifying tutorial on approximate message passing*, Foundations and Trends® in Machine Learning 15 (2022), no. 4, 335–536.
- [FW21] Zhou Fan and Yihong Wu, *The replica-symmetric free energy for Ising spin glasses with orthogonally invariant couplings*, arXiv preprint arXiv:2105.02797 (2021).
- [GB23] Cédric Gerbelot and Raphaël Berthier, *Graph-based approximate message passing iterations*, Inf. Inference 12 (2023), no. 4, Paper No. iaad020, 67.
- [GKKZ22] Alice Guionnet, Justin Ko, Florent Krzakala, and Lenka Zdeborov´a, *Low-rank matrix estimation with inhomogeneous noise*, arXiv preprint arXiv:2208.05918 (2022).
- [Hac23] Walid Hachem, *Approximate message passing for sparse matrices with application to the equilibria of large ecological lotka-volterra systems*, arXiv preprint arXiv:2302.09847 (2023).
- [Han23] Qiyang Han, *Noisy linear inverse problems under convex constraints: Exact risk asymptotics in high dimensions*, Ann. Statist. 51 (2023), no. 4, 1611–1638.
- [HK70] Arthur E. Hoerl and Robert W. Kennard, *Ridge regression: Biased estimation for nonorthogonal problems*, Technometrics 12 (1970), no. 1, 55–67.
- [HK17] Yukun He and Antti Knowles, *Mesoscopic eigenvalue statistics of Wigner matrices*, Ann. Appl. Probab. 27 (2017), no. 3, 1510–1550.
- [HMRT22] Trevor Hastie, Andrea Montanari, Saharon Rosset, and Ryan J. Tibshirani, *Surprises in high-dimensional ridgeless least squares interpolation*, Ann. Statist. 50 (2022), no. 2, 949–986.
- [HS23] Qiyang Han and Yandi Shen, *Universality of regularized regression estimators in high dimensions*, Ann. Statist. 51 (2023), no. 4, 1799–1823.
- [HX23] Qiyang Han and Xiaocong Xu, *The distribution of ridgeless least squares interpolators*, arXiv preprint arXiv:2307.02044 (2023).
- [JM13] Adel Javanmard and Andrea Montanari, *State evolution for general approximate message passing algorithms, with applications to spatial coupling*, Inf. Inference 2 (2013), no. 2, 115–144.
- [KF09] Daphne Koller and Nir Friedman, *Probabilistic graphical models: principles and techniques*, MIT press, 2009.
- [KKM+16] Yoshiyuki Kabashima, Florent Krzakala, Marc Mézard, Ayaka Sakata, and Lenka Zdeborov´a, *Phase transitions and sample complexity in Bayes-optimal matrix factorization*, IEEE Trans. Inform. Theory 62 (2016), no. 7, 4228–4265.
- [KKP96] Alexei M. Khorunzhy, Boris A. Khoruzhenko, and Leonid A. Pastur, *Asymptotic properties of large random matrices with independent entries*, J. Math. Phys. 37 (1996), no. 10, 5033–5060.
- [KMS⁺12] Florent Krzakala, Marc Mézard, Francois Sausset, Yifan Sun, and Lenka Zdeborová, *Probabilistic reconstruction in compressed sensing: algorithms, phase diagrams, and threshold achieving matrices*, Journal of Statistical Mechanics: Theory and Experiment 2012 (2012), no. 08, P08009.
- [LFW23] Gen Li, Wei Fan, and Yuting Wei, *Approximate message passing from random initialization with applications to* \mathbb{Z}_2 *synchronization*, Proc. Natl. Acad. Sci. USA 120 (2023), no. 31, Paper No. e2302930120, 7.
- [LKZ17] Thibault Lesieur, Florent Krzakala, and Lenka Zdeborov´a, *Constrained low-rank matrix estimation: phase transitions, approximate message passing and applications*, J. Stat. Mech. Theory Exp. (2017), no. 7, 073403, 86.
- [LP09] A. Lytova and L. Pastur, *Central limit theorem for linear eigenvalue statistics of random matrices with independent entries*, Ann. Probab. 37 (2009), no. 5, 1778–1840.
- [LS18] Ji Oon Lee and Kevin Schnelli, *Local law and Tracy-Widom limit for sparse random matrices*, Probab. Theory Related Fields 171 (2018), no. 1-2, 543–616.
- [LW21] Yue Li and Yuting Wei, *Minimum* ℓ_1 *-norm interpolators: Precise asymptotics and multiple descent*, arXiv preprint arXiv:2110.09502 (2021).
- [LW22] Gen Li and Yuting Wei, *A non-asymptotic framework for approximate message passing in spiked models*, arXiv preprint arXiv:2208.03313 (2022).
- [MM21] Léo Miolane and Andrea Montanari, *The distribution of the Lasso: uniform control over sparse balls and adaptive parameter tuning*, Ann. Statist. 49 (2021), no. 4, 2313–2335.
- [Mon12] Andrea Montanari, *Graphical models concepts in compressed sensing*, Compressed sensing, Cambridge Univ. Press, Cambridge, 2012, pp. 394–438.

54 Z. BAO, Q. HAN, AND X. XU

- [MP17] Junjie Ma and Li Ping, *Orthogonal AMP*, IEEE Access **5** (2017), 2020–2033.
- [MR16] Andrea Montanari and Emile Richard, *Non-negative principal component analysis: message passing algorithms and sharp asymptotics*, IEEE Trans. Inform. Theory 62 (2016), no. 3, 1458–1484.
- [MV21] Andrea Montanari and Ramji Venkataramanan, *Estimation of low-rank matrices via approximate message passing*, Ann. Statist. 49 (2021), no. 1, 321–345.
- [MV22] Marco Mondelli and Ramji Venkataramanan, *Approximate message passing with spectral initialization for generalized linear models*, J. Stat. Mech. Theory Exp. (2022), no. 11, Paper No. 114003, 41.
- [PKK23] Aleksandr Pak, Justin Ko, and Florent Krzakala, *Optimal algorithms for the inhomogeneous spiked wigner model*, arXiv preprint arXiv:2302.06665 (2023).
- [Ran11] Sundeep Rangan, *Generalized approximate message passing for estimation with random linear mixing*, 2011 IEEE International Symposium on Information Theory Proceedings, IEEE, 2011, pp. 2168–2172.
- [RSF19] Sundeep Rangan, Philip Schniter, and Alyson K. Fletcher, *Vector approximate message passing*, IEEE Trans. Inform. Theory 65 (2019), no. 10, 6664–6684.
- [RV18] Cynthia Rush and Ramji Venkataramanan, *Finite sample analysis of approximate message passing algorithms*, IEEE Trans. Inform. Theory 64 (2018), no. 11, 7264–7286.
- [SC19] Pragya Sur and Emmanuel J. Candès, *A modern maximum-likelihood theory for highdimensional logistic regression*, Proc. Natl. Acad. Sci. 116 (2019), no. 29, 14516– 14525.
- [SCC19] Pragya Sur, Yuxin Chen, and Emmanuel J. Candès, *The likelihood ratio test in highdimensional logistic regression is asymptotically a rescaled chi-square*, Probab. Theory Related Fields 175 (2019), no. 1-2, 487–558.
- [SR15] Philip Schniter and Sundeep Rangan, *Compressive phase retrieval via generalized approximate message passing*, IEEE Trans. Signal Process. 63 (2015), no. 4, 1043–1055.
- [TAH18] Christos Thrampoulidis, Ehsan Abbasi, and Babak Hassibi, *Precise error analysis of regularized M-estimators in high dimensions*, IEEE Trans. Inform. Theory 64 (2018), no. 8, 5592–5628.
- [Tak21] Keigo Takeuchi, *Bayes-optimal convolutional AMP*, IEEE Trans. Inform. Theory 67 (2021), no. 7, 4405–4428.
- [TAP77] David J. Thouless, Philip W. Anderson, and Robert G. Palmer, *Solution of 'solvable model of a spin glass'*, Philosophical Magazine 35 (1977), no. 3, 593–601.
- [Ver18] Roman Vershynin, *High-dimensional probability: An introduction with applications in data science*, Cambridge Series in Statistical and Probabilistic Mathematics, vol. 47, Cambridge University Press, Cambridge, 2018.
- [Wih09] Thomas P. Wihler, *On the Hölder continuity of matrix functions for normal matrices*, JIPAM. J. Inequal. Pure Appl. Math. 10 (2009), no. 4, Article 91, 5.
- [WX20] Denny Wu and Ji Xu, *On the optimal weighted* ℓ_2 *regularization in overparameterized linear regression*, Advances in Neural Information Processing Systems 33 (2020), 10112–10123.
- [WZF22] Tianhao Wang, Xinyi Zhong, and Zhou Fan, *Universality of approximate message passing algorithms and tensor networks*, arXiv preprint arXiv:2206.13037 (2022).
- [ZSF22] Xinyi Zhong, Chang Su, and Zhou Fan, *Empirical Bayes PCA in high dimensions*, J. R. Stat. Soc. Ser. B. Stat. Methodol. 84 (2022), no. 3, 853–878.
- [ZWF21] Xinyi Zhong, Tianhao Wang, and Zhou Fan, *Approximate message passing for orthogonally invariant ensembles: Multivariate non-linearities and spectral initialization*, arXiv preprint arXiv:2110.02318 (2021).

(Z. Bao) Department of Mathematics, Hong Kong University of Science and Technology, Hong Kong.

Email address: mazgbao@ust.hk

(Q. Han) Department of Statistics, Rutgers University, Piscataway, NJ 08854, USA. *Email address*: qh85@stat.rutgers.edu

(X. Xu) Department of Mathematics, Hong Kong University of Science and Technology, Hong Kong.

Email address: xxuay@connect.ust.hk