CATEGORICAL EQUIVALENCES FOR HOPF TRUSSES AND THEIR MODULES

Ramón González Rodríguez¹ and Ana Belén Rodríguez Raposo².

¹ [https://orcid.org/0000-0003-3061-6685].

CITMAga, 15782 Santiago de Compostela, Spain.

Universidade de Vigo, Departamento de Matemática Aplicada II, E. E. Telecomunicación, E-36310 Vigo, Spain.

Spam

email: rgon@dma.uvigo.es

² [https://orcid.org/0000-0002-8719-5159]

Universidade de Santiago de Compostela, Departamento de Didácticas Aplicadas, Facultade C. C. Educación, E-15782 Santiago de Compostela, Spain.

email: anabelen.rodriguez.raposo@usc.es

ABSTRACT. In this paper we introduce the notion of generalized invertible 1-cocycle in a strict braided monoidal category C, and we prove that the category of Hopf trusses is equivalent to the category of generalized invertible 1-cocycles. On the other hand, we also introduce the notions of module for a Hopf truss and for a generalized invertible 1-cocycle. We prove some functorial results involving these categories of modules and we show that the category of modules associated to a generalized invertible 1-cocycle is equivalent to a category of modules associated to a suitable Hopf truss. Finally, assuming that in C we have equalizers, we introduce the notion of Hopf-module in the Hopf truss setting and we obtain the Fundamental Theorem of Hopf modules associated to a Hopf truss.

KEYWORDS: Braided monoidal category, skew truss, Hopf truss, generalized invertible 1-cocycle, module, Hopf-module.

2020 MATHEMATICS SUBJECT CLASSIFICATION: 18M15, 16T05.

1. INTRODUCTION

The notion of skew brace was introduced recently in [6]. This algebraic object consists of two different group structures (T, \diamond) and (T, \circ) on the same set T and that satisfy $\forall a, b, c \in T$ the compatibility condition

$$a \circ (b \diamond c) = (a \circ b) \diamond a^{\diamond} \diamond (a \circ c)$$

where a^{\diamond} denotes the inverse with respect to \diamond . On the other hand, Hopf braces were introduced in [2] as the linearisation of skew braces and then consists of two structures (H_1, H_2) of Hopf algebras defined on the same object that share a common coalgebra structure. As a consequence of the compatibility condition, the Hopf algebra H_1 can be endowed with a structure of module algebra over H_2 . The relevance of this structure comes through bijective 1-cocycles $\sigma : H_1 \to B$, where B is a Hopf algebra that acts on H_1 (see [2]). In fact, the category of Hopf braces with fixed H_1 is equivalent to the category of invertible 1-cocycles $\sigma : H_1 \to B$ (see [2, Theorem 1.12]). Thus, Hopf braces are nothing more than coalgebra isomorphisms between Hopf algebras that share the underlying coalgebra and related by a module algebra structure. In [3] T. Brzeziński introduces the notion of skew truss as a generalization of the notion of skew brace in the following way: A skew truss consist of a group structure (T, \diamond) and a semigroup structure (T, \circ) defined on the same set T, and a map (called the cocycle of the skew truss) $\omega : T \to T$ that satisfy $\forall a, b, c \in T$:

$$a \circ (b \diamond c) = (a \circ b) \diamond \omega(a)^{\diamond} \diamond (a \circ c)$$

where $\omega(a)^{\diamond}$ is the inverse in (T, \diamond) . This object can be linearised to obtain the so-called Hopf truss (H_1, H_2, σ) that consist of a Hopf algebra H_1 and a non-unital bialgebra H_2 with the same underlying coalgebra structure, and a cocycle $\sigma : H_2 \to H_1$ that must satisfy the corresponding compatibility condition. Due to it, H_1 can be endowed with a non-unital H_2 -module algebra structure. Observe that if H_2 is a Hopf algebra and σ is the identity, the Hopf truss is actually a Hopf brace. In the second section of this paper we prove that there exists a relation between the categories of Hopf trusses and skew trusses in the following way (see Theorem 2.17): The category of skew trusses is equivalent to the full subcategory of Hopf trusses whose objects are pointed cosemisimple Hopf trusses.

Taking into account the equivalences between the categories of Hopf braces an invertible 1-cocycles and with the intention of extending them to the context of Hopf trusses, in the third section of this paper we introduce the notion of generalized invertible 1-cocycle in a strict braided monoidal category C as a generalization of the bijective 1-cocycles proposed by I. Angiono, C. Galindo and L. Vendramín in [2]. Thanks to this, in Theorem 3.2 we prove that the category of Hopf trusses is equivalent to the category of generalized invertible 1-cocycles. Moreover, in Section 4 we also introduce the notions of module for a Hopf truss and for a generalized invertible 1-cocycle. After doing this, in this section we prove some functorial results involving these categories of modules and in Theorem 4.12 we show that the category of modules associated to a generalized invertible 1-cocycle is equivalent to a category of modules associated to a suitable Hopf truss that we can build thanks to the generalized invertible 1-cocycle. This results are a generalization to the Hopf truss setting of the ones proved for modules associates to Hopf braces in [4].

On the other hand, it is well known that, if H is a Hopf algebra in a category \mathbb{F} -Vect of vector spaces over a field \mathbb{F} and X is an object in C, the tensor product $H \otimes X$, with the action and coaction induced by the product and the coproduct of H, is an object in the category H-Hopf-Mod, i.e. the category of left H-Hopf modules. The objects and morphisms of this category are defined in the following way: Let M be a left H-module and a left H-comodule. In this setting, if for all $m \in M$ and $h \in H$, we write h.m for the left action and we use the Heyenman-Sweedler notation $\rho_M(m) = m_{[0]} \otimes m_{[1]}$ for the coaction, we will say that M is a left H-Hopf module if the equality

$$\rho_M(h.m) = h_{(1)}m_{[0]} \otimes h_{(2)}.m_{[1]}$$

holds, where $\delta_H(h) = h_{(1)} \otimes h_{(2)}$ is the coproduct of H and $h_{(1)}m_{[0]}$ is the product in H of $h_{(1)}$ and $m_{[0]}$. A morphism between two left H-Hopf modules is a \mathbb{F} -linear map that is H-linear and H-colinear.

This construction introduced for $H \otimes X$ in the previous paragraph is functorial and, as a consequence, we have a functor, called the induction functor, $F = H \otimes -: \mathbb{C} \to \mathsf{H}\text{-}\mathsf{Hopf}\text{-}\mathsf{Mod}$. Moreover, for all $M \in H\text{-}\mathsf{Hopf}$, the construction of subobject of coinvariants $M^{coH} = \{m \in M \mid m_{[0]} \otimes m_{[1]} = 1_H \otimes m\}$ also is functorial. Thus, there exists a functor of coinvariants $G = (\)^{coH} : \mathsf{H}\text{-}\mathsf{Hopf}\text{-}\mathsf{Mod} \to \mathbb{C}$ such that $F \dashv G$. Moreover, $H \otimes M^{coH}$ and M are isomorphic in $\mathsf{H}\text{-}\mathsf{Hopf}\text{-}\mathsf{Mod}$ (see [7] and [10]) and F and G induces an equivalence between the categories $\mathsf{H}\text{-}\mathsf{Hopf}\text{-}\mathsf{Mod}$ and $\mathbb{F}\text{-}\mathsf{Vect}$. The existence of the isomorphism between $H \otimes M^{coH}$ and M is the main statement of the Fundamental Theorem of Hopf modules.

The Fundamental Theorem of Hopf modules and the categorical equivalence of the previous paragraph remain valid for weak Hopf algebras, Hopf quasigroups, weak Hopf quasigroups and in [5] we can find that which can also be obtained for Hopf braces. In the last section of this paper, assuming that the braided monoidal category C admits equalizers, we extend the previous results to the Hopf truss setting, i.e. we introduce the notion of Hopf-module associated to a Hopf truss, we obtain the Fundamental Theorem of Hopf modules associated to a Hopf truss and we prove that there exists a categorical equivalence as in the case of Hopf algebras, weak Hopf algebras, Hopf quasigroups, weak Hopf quasigroups, Hopf braces, etc.

2. Hopf trusses

Throughout this paper C denotes a strict braided monoidal category with tensor product \otimes , unit object K and braiding c. Recall that a monoidal category is a category C together with a functor $\otimes : C \times C \rightarrow C$, called tensor product, an object K of C, called the unit object, and families of natural isomorphisms

$$a_{M,N,P}: (M \otimes N) \otimes P \to M \otimes (N \otimes P), \quad r_M: M \otimes K \to M, \quad l_M: K \otimes M \to M,$$

in C, called associativity, right unit and left unit constraints, respectively, satisfying the Pentagon Axiom and the Triangle Axiom, i.e.,

$$a_{M,N,P\otimes Q} \circ a_{M\otimes N,P,Q} = (id_M \otimes a_{N,P,Q}) \circ a_{M,N\otimes P,Q} \circ (a_{M,N,P} \otimes id_Q),$$
$$(id_M \otimes l_N) \circ a_{M,K,N} = r_M \otimes id_N,$$

where for each object X in C, id_X denotes the identity morphism of X (see [9]). A monoidal category is called strict if the constraints of the previous paragraph are identities. It is a well-known fact (see for example [8]) that every non-strict monoidal category is monoidal equivalent to a strict one. This lets us to treat monoidal categories as if they were strict and, as a consequence, the results proved in a strict setting hold for every non-strict monoidal category, for example the category \mathbb{F} -Vect of vector spaces over a field \mathbb{F} , the category *R*-Mod of left modules over a commutative ring *R*, or Set the category of sets.

For simplicity of notation, given objects M, N, P in C and a morphism $f : M \to N$, we will write $P \otimes f$ for $id_P \otimes f$ and $f \otimes P$ for $f \otimes id_P$.

A braiding for a strict monoidal category C is a natural family of isomorphisms

$$c_{M,N}: M \otimes N \to N \otimes M$$

subject to the conditions

$$c_{M,N\otimes P} = (N \otimes c_{M,P}) \circ (c_{M,N} \otimes P), \ c_{M\otimes N,P} = (c_{M,P} \otimes N) \circ (M \otimes c_{N,P}).$$

A strict braided monoidal category C is a strict monoidal category with a braiding. Note that, as a consequence of the definition, the equalities $c_{M,K} = c_{K,M} = id_M$ hold, for all object M of C. If the braiding satisfies that $c_{N,M} \circ c_{M,N} = id_{M\otimes N}$, for all M, N in C, we will say that C is symmetric. In this case, we call the braiding c a symmetry for the category C.

Definition 2.1. A monoid in C is a triple $A = (A, \eta_A, \mu_A)$ where A is an object in C and $\eta_A : K \to A$ (unit), $\mu_A : A \otimes A \to A$ (product) are morphisms in C such that $\mu_A \circ (A \otimes \eta_A) = id_A = \mu_A \circ (\eta_A \otimes A)$ (unit property) and $\mu_A \circ (A \otimes \mu_A) = \mu_A \circ (\mu_A \otimes A)$ (associative property) hold.

Given two monoids $A = (A, \eta_A, \mu_A)$ and $B = (B, \eta_B, \mu_B)$, a morphism $f : A \to B$ in C is an monoid morphism if $\mu_B \circ (f \otimes f) = f \circ \mu_A$ (f is multiplicative) and $f \circ \eta_A = \eta_B$ (f preserves the unit).

If A, B are monoids in C, the tensor product $A \otimes B$ is also an algebra in C where $\eta_{A \otimes B} = \eta_A \otimes \eta_B$ and $\mu_{A \otimes B} = (\mu_A \otimes \mu_B) \circ (A \otimes c_{B,A} \otimes B)$.

We will say that a monoid A is commutative if $\mu_A \circ c_{A,A} = \mu_A$.

Definition 2.2. A comonoid in C is a triple $D = (D, \varepsilon_D, \delta_D)$ where D is an object in C and $\varepsilon_D : D \to K$ (counit), $\delta_D : D \to D \otimes D$ (coproduct) are morphisms in C such that $(\varepsilon_D \otimes D) \circ \delta_D = id_D = (D \otimes \varepsilon_D) \circ \delta_D$ (counit property) and $(\delta_D \otimes D) \circ \delta_D = (D \otimes \delta_D) \circ \delta_D$ (coassociative property) hold.

If $D = (D, \varepsilon_D, \delta_D)$ and $E = (E, \varepsilon_E, \delta_E)$ are comonoids, a morphism $f : D \to E$ in C is a comonoid morphism if $(f \otimes f) \circ \delta_D = \delta_E \circ f$ (f is comultiplicative) and $\varepsilon_E \circ f = \varepsilon_D$ (f preserves the counit). Given D, E comonoids in C, the tensor product $D \otimes E$ is a comonoid in C where $\varepsilon_{D \otimes E} = \varepsilon_D \otimes \varepsilon_E$

and $\delta_{D\otimes E} = (D \otimes c_{D,E} \otimes E) \circ (\delta_D \otimes \delta_E).$

We will say that a comonoid D is cocommutative if $\delta_D = c_{D,D} \circ \delta_D$.

Definition 2.3. Let $D = (D, \varepsilon_D, \delta_D)$ be a comonoid in C. We will say that a morphism $g: K \to D$ is a grouplike morphism if satisfy

(1)
$$\delta_D(g) = g \otimes g$$

and

(2)
$$\varepsilon_D \circ g = id_K$$

Remark 2.4. In the category of vector spaces over a field \mathbb{F} we can find interesting examples of comonoids. For example, if S is a set, with $\mathbb{F}[S]$ we will denote the free \mathbb{F} -vector space on S, i.e.,

$$\mathbb{F}[S] = \bigoplus_{s \in S} \mathbb{F}s.$$

This vector space has a comonoid structure determined by

(3)
$$\delta_{\mathbb{F}[S]}(s) = s \otimes s, \quad \varepsilon_{\mathbb{F}[S]}(s) = 1_{\mathbb{F}}.$$

Let $(C, \varepsilon_C, \delta_C)$ be a comonoid in \mathbb{F} -Vect. A grouplike element c of C is a $c \in C$ such that the linear map $g_c : \mathbb{F} \to C$ defined by $g_c(1_{\mathbb{F}}) = c$ is a grouplike morphism in \mathbb{F} -Vect. Therefore, $c \in C$ is a grouplike element if (3) holds for c, ε_C and δ_C . In the following we will denote by $\mathsf{G}(C)$ the set of grouplike elements of C and it is well known that they are linearly independent [1, Theorem 2.1.2]. If S is a set, the comonoid $\mathbb{F}[S]$ is called the grouplike comonoid of S and satisfies $\mathsf{G}(\mathbb{F}[S]) = S$. Moreover the grouplike comonoid of $\mathsf{G}(C)$ is a subcomonoid of C (D is a subcomonoid of C if $\delta_C(D) \subset D \otimes D$ or, in other words, D is a comonoid with the restriction of the coproduct δ_C and the counit ε_C).

A pointed comonoid in \mathbb{F} -Vect is a comonoid C whose simple subcomonoids are one-dimensional. Then, C is pointed if and only if its coradical C_0 (the sum of the simple subcomonoids of C) is the grouplike comonoid of $\mathsf{G}(C)$, i.e., $C_0 = \mathbb{F}[\mathsf{G}(C)]$. We will say that the comonoid C is cosemisimple if $C = C_0$. Therefore, if C is pointed cosemisimple, $C = \mathbb{F}[\mathsf{G}(C)]$. On the other hand, if G is a group and $C = \mathbb{F}[G]$, we have that C is pointed and cosemisimple. Finally, if \mathbb{F} is algebraically closed and C is cocommutative, C is pointed.

Definition 2.5. Let $D = (D, \varepsilon_D, \delta_D)$ be a comonoid and let $A = (A, \eta_A, \mu_A)$ be an monoid. By $\mathcal{H}(D, A)$ we denote the set of morphisms $f : D \to A$ in C. With the convolution operation

$$f * g = \mu_A \circ (f \otimes g) \circ \delta_D,$$

 $\mathcal{H}(D, A)$ is an monoid where the unit element is $\eta_A \circ \varepsilon_D = \varepsilon_D \otimes \eta_A$. We will say that $f: D \to A$ is convolution invertible if there exists $f^{-1}: D \to A$ such that $f * f^{-1} = f^{-1} * f = \varepsilon \otimes \eta$.

Definition 2.6. Let A be an monoid. The pair (M, φ_M) is a left A-module if M is an object in C and $\varphi_M : A \otimes M \to M$ is a morphism in C satisfying $\varphi_M \circ (\eta_A \otimes M) = id_M$ and

$$\varphi_M \circ (A \otimes \varphi_M) = \varphi_M \circ (\mu_A \otimes M).$$

Given two left A-modules (M, φ_M) and (N, φ_N) , $f : M \to N$ is a morphism of left A-modules if $\varphi_N \circ (A \otimes f) = f \circ \varphi_M$ (left A-linearity).

Then left A-modules with morphisms of left A-modules form a category that we will denote by $_{A}Mod$.

Let B an object in C such that there exists an associative product $\mu_B : B \otimes B \to B$. We will say that (M, ϕ_M) is a non-unital left B-module if $\phi_M \circ (B \otimes \phi_M) = \phi_M \circ (\mu_B \otimes M)$. A morphism between non-unital left B-modules is a left B-linear morphism as in the case of morphisms for modules over a monoid. Then non-unital left B-modules form a category that we will denote by Bmod.

Definition 2.7. A non-unital bimonoid in the category C is a comonoid $(B, \varepsilon_B, \delta_B)$ with an associative product $\mu_B : B \otimes B \to B$ such that μ_B is a comonoid morphism. Then the following identities hold:

(4)
$$\varepsilon_B \circ \mu_B = \varepsilon_B \otimes \varepsilon_B,$$

(5)
$$\delta_B \circ \mu_B = (\mu_B \otimes \mu_B) \circ \delta_{B \otimes B}.$$

A bimonoid in C is a monoid (B, η_B, μ_B) and a comonoid $(B, \varepsilon_B, \delta_B)$ such that η_B and μ_B are comonoid morphisms. Then, (4), (5),

(6) $\varepsilon_B \circ \eta_B = id_K,$

and

(7)
$$\delta_B \circ \eta_B = \eta_B \otimes \eta_B$$

hold.

A morphism between non-unital bimonoids H and B is a morphism $f: H \to B$ in C of comonoids and multiplicative. A morphism between bimonoids H and B is a morphism $f: H \to B$ in C of monoids and comonoids.

With the composition of morphisms in C we can define a category whose objects are non-unital bimonoids (bimonoids) and whose morphisms are morphisms of non-unital bimonoids (bimonoids). We denote this category by bimod (BiMod).

Definition 2.8. Let H be a bimonoid in C. If there exists a morphism $\lambda_H : H \to H$ in C , called the antipode of H, satisfying that λ_H is the inverse of id_H in $\mathcal{H}(H, H)$, i.e.,

(8)
$$id_H * \lambda_H = \eta_H \circ \varepsilon_H = \lambda_H * id_H,$$

we say that H is a Hopf monoid.

A morphism of Hopf monoids is an bimonoid morphism. With the composition of morphisms in C we can define a category whose objects are Hopf monoids and whose morphisms are morphisms of Hopf monoids. We denote this category by Hopf.

Remark 2.9. If H is a Hopf monoid, the antipode is antimultiplicative and anticomultiplicative

$$\lambda_H \circ \mu_H = \mu_H \circ (\lambda_H \otimes \lambda_H) \circ c_{H,H}, \quad \delta_H \circ \lambda_H = c_{H,H} \circ (\lambda_H \otimes \lambda_H) \circ \delta_H,$$

and leaves the unit and counit invariant, i.e.,

$$\lambda_H \circ \eta_H = \eta_H, \ \varepsilon_H \circ \lambda_H = \varepsilon_H.$$

A Hopf monoid is commutative if it is commutative as monoid and cocommutative if it is cocommutative as comonoid. It is easy to see that in both cases $\lambda_H \circ \lambda_H = id_H$.

Note that, if $f: H \to D$ is a Hopf monoid morphism the following equality holds:

(9)
$$\lambda_D \circ f = f \circ \lambda_H.$$

Remark 2.10. In the category Set the tensor product is the cartesian product and the unit element is a set $\{\star\}$ with an unique element \star . Taking all this into account, we have that a set T is a non-unital bimonoid in Set if and only if T is a semigroup in Set with product \diamond . If (T, \diamond) is a semigroup the non-unital bimonoid structure is the following: the product is the one induced by \diamond , the coproduct is defined by $\delta_T(a) = (a, a)$ and the counit by $\varepsilon_T(a) = \star$. Moreover, T is a bimonoid in Set if and only if T is a monoid in Set with product \diamond and unit 1_{\diamond} . In this case we define the non-unital bimonoid structure as in the case of semigroups and the unit is defined by $\eta_T(\star) = 1_{\diamond}$. Finally, T is a Hopf monoid in Set if and only if it is a group. The bimonoid structure is defined as for monoids and the antipode by $\lambda_T(a) = a^{\diamond}$ where a^{\diamond} is the inverse of a in T.

On the other hand, if \mathbb{F} is a field and (T, \diamond) is a semigroup in Set, the direct sum

$$\mathbb{F}[T] = \bigoplus_{a \in T} \mathbb{F}a$$

is a non-unital bimonoid in \mathbb{F} -Vect where $\mu_{\mathbb{F}[T]}$ is the unique linear map such that $\mu_{\mathbb{F}[T]}(a \otimes b) = a \diamond b$ and the comonoid structure is the one defined in (3). Also, if (T, \diamond) is a monoid in Set with unit 1_{\diamond} , $\mathbb{F}[T]$ is a bimonoid in \mathbb{F} -Vect, where $\eta_{\mathbb{F}[T]}$ is the unique linear map such that $\eta_{\mathbb{F}[T]}(1_{\mathbb{F}}) = 1_{\diamond}$, and the non-unital bimonoid structure is the one introduced for semigroups. Finally, if (T, \diamond) is a group, $\mathbb{F}[T]$ is a Hopf monoid in \mathbb{F} -Vect with the previous bimonoid structure and antipode the unique linear map satisfying $\lambda_{\mathbb{F}[T]}(a) = a^{\diamond}$, where a^{\diamond} denotes the inverse of a.

Let (T, \diamond) , (S, \diamond) be semigroups in Set and let $f: T \to S$ be a semigroup morphism in Set. Then, if $\mathbb{F}[f]$ denotes the linear extension of f, $\mathbb{F}[f]$ is a non-unital bimonoid morphism between $\mathbb{F}[T]$ and $\mathbb{F}[S]$ in \mathbb{F} -Vect. The same property holds for a morphisms of monoids f in Set, i.e. $\mathbb{F}[f]$ is a bimonoid morphism, and for a morphism f of groups, i.e. $\mathbb{F}[f]$ is a group morphism.

Therefore, if sGpr is the category of semigroups in Set, Mon is the category of monoids in Set and Gpr denotes the category of groups, there exists three functors

$$L_{sa}: sGpr \rightarrow bimon, \quad L_m: Mon \rightarrow BiMon, \quad L_a: Gpr \rightarrow Hopf,$$

where bimon is the category of non-unital bimonoids in \mathbb{F} -Vect, BiMon is the category of bimonoids in \mathbb{F} -Vect and, finally, Hopf denotes the category of Hopf monoids in \mathbb{F} -Vect. In this setting, the functor L_m is the restriction of L_{sg} to the category of monoids in Set and L_g is the restriction of L_m to the category of groups.

In any case non-unital bimonoids, bimonoids and Hopf monoids in \mathbb{F} -Vect are comonoids. Taking this into account, if B is an object in bimon, it is possible to define a semigroup structure on G(B), with product \diamond induced by μ_B $(a \diamond b = \mu_B(a \otimes b))$. Moreover, if $f : B \to B'$ is a morphism of non-unital bimonoids, the image of the restriction of f to G(B) lies into G(B'). Thus we have a functor G_{sg} between bimon and sGpr defined by $G_{sg}(B) = G(B)$ on objects and by $G_{sg}(f) = G(B)$ on morphisms. Also, if A is an object in BiMon it is possible to define a monoid structure on G(A), with product \diamond as in the case of non-unital bimonoids and unit 1_{\diamond} induced by η_A ($1_{\diamond} = \eta_A(1_{\mathbb{F}})$). Moreover, if $f : A \to A'$ is a morphism of bimonoids, the image of the restriction of f to G(A) lies into G(A'). Then, we have a new functor G_m between BiMon and Mon defined by $G_m(A) = G(A)$ on objects and by $G_m(f) = G(f)$ on morphisms. Finally, If H is a Hopf monoid, G(H) is a group where the monoid structure is the one defined for bimonoids and the inverse is defined by the antipode, i.e., the inverse of $h \in G(H)$ is $\lambda_H(h)$. As in the two previous cases, this construction works well with morphisms of Hopf monoids and, as a consequence, there exists a functor G_g between Hopf and Gpr defined by $G_q(H) = G(H)$ on objects and by $G_q(f) = G(f)$ on morphisms.

It is true that $L_{sg} \dashv G_{sg}$, $L_m \dashv G_m$ and $L_g \dashv G_g$. In the three adjunctions the unit of every one of them is the identity. Thus, the first adjoint pair induces an equivalence of categories between sGpr and the full subcategory of bimon of all pointed cosemisimple non-unital bimonoids in \mathbb{F} -Vect, the

second one induces an equivalence of categories between Mon and the full subcategory of BiMon of all pointed cosemisimple bimonoids in \mathbb{F} -Vect, and the third one induces an equivalence of categories between Gpr and the full subcategory of Hopf of all pointed cosemisimple non-unital Hopf monoids in \mathbb{F} -Vect. Also we have this commutative diagrams



where I_m and I_b and I_g denote the corresponding inclusion functors.

Definition 2.11. Let B a bimonoid and let A be a monoid in C. We will say that (A, ϕ_A) is a left *B*-module monoid if it is a left *B*-module with action $\varphi_A : B \otimes A \to A$ such that

(10)
$$\varphi_A \circ (B \otimes \eta_A) = \varepsilon_B \otimes \eta_A$$

and

(11)
$$\varphi_A \circ (B \otimes \mu_A) = \mu_A \circ (\varphi_A \otimes \varphi_A) \circ (B \otimes c_{B,A} \otimes A) \circ (\delta_B \otimes A \otimes A)$$

hold.

If B is a non-unital bimonoid, we will say that (A, ϕ_A) is a non-unital left B-module monoid if (A, ϕ_A) is a non-unital left *B*-module and (10) and (11) hold.

The notion of Hopf truss was introduced by T. Brzeziński in [3] in the category \mathbb{F} -Vect as the linearisation of the notion of skew truss. In the monoidal setting the definition of Hopf truss is the following:

Definition 2.12. Let $(H, \varepsilon_H, \delta_H)$ be a comonoid in C. Assume that there are a monoid structure (H, η_H, μ_H^1) , a product $\mu_H^2 : H \otimes H \to H$ and two endomorphism of H denoted by λ_H and σ_H . We will say that

$$(H, \eta_H, \mu_H^1, \mu_H^2, \varepsilon_H, \delta_H, \lambda_H, \sigma_H)$$

is a Hopf truss if:

- (i) $H_1 = (H, \eta_H, \mu_H^1, \varepsilon_H, \delta_H, \lambda_H)$ is a Hopf monoid in C. (ii) $H_2 = (H, \mu_H^2, \varepsilon_H, \delta_H,)$ is a non-unital bimonoid in C.
- (iii) The morphism σ_H is a comonoid morphism and the following equality holds:

$$\mu_{H}^{2} \circ (H \otimes \mu_{H}^{1}) = \mu_{H}^{1} \circ (\mu_{H}^{2} \otimes \Gamma_{H_{1}}^{\sigma_{H}}) \circ (H \otimes c_{H,H} \otimes H) \circ (\delta_{H} \otimes H \otimes H),$$

where

$$\Gamma_{H_1}^{\sigma_H} = \mu_H^1 \circ ((\lambda_H \circ \sigma_H) \otimes \mu_H^2) \circ (\delta_H \otimes H).$$

We will say that a Hopf truss is cocommutative if the comonoid $(H, \varepsilon_H, \delta_H)$ is cocommutative.

Note that, a Hopf truss is a Hopf brace in the sense of I. Angiono, C. Galindo and L. Vendramin (see [2]) if σ_H is the identity and there exists a morphism $S_H : H \to H$ such that $H_2 = (H, \eta_H, \mu_H^2, \varepsilon_H, \delta_H, S_H)$ is a Hopf monoid.

Notation 2.13. Given a Hopf truss, we will denote it by $\mathbb{H} = (H_1, H_2, \sigma_H)$. The morphism σ_H is called the cocycle of \mathbb{H} .

The proofs that we can find in Section 6 of [3] can be replicated in the braided monoidal setting since they do not depend on the symmetry of the category \mathbb{F} -Vect. Then we have the following properties: By [3, Lemma 6.2] the cocycle σ_H of a Hopf truss \mathbb{H} in C is fully determined by η_H and the product μ_H^2 in the following way:

(12)
$$\sigma_H = \mu_H^2 \circ (H \otimes \eta_H)$$

Then, as a consequence of the associativity for the product μ_H^2 , we have that

(13)
$$\sigma_H \circ \mu_H^2 = \mu_H^2 \circ (H \otimes \sigma_H)$$

holds. Finally, by [3, Theorem 6.5] we know that the monoid H_1 is a non-unital left H_2 -module monoid for the action $\Gamma_{H_1}^{\sigma_H}$.

Definition 2.14. Given two Hopf trusses \mathbb{H} and \mathbb{B} , a morphism f between the two underlying objects is called a morphism of Hopf trusses if $f : H_1 \to B_1$ is a Hopf monoid morphism and $f : H_2 \to B_2$ is a morphism of non-unital bimonoids. Then by [3, Proposition 6.8]

(14)
$$\sigma_B \circ f = f \circ \sigma_H$$

holds.

Hopf trusses together with morphisms of Hopf trusses form a category which we denote by HTr. It is obvious that Hopf braces with morphisms of Hopf braces form a category which we denote by HBr that is a subcategory of HTr.

Remark 2.15. Following [3] a skew truss is a set T with two binary operations \diamond_1 and \diamond_2 and a map $\omega_T : T \to T$ (called the cocycle) such that the pair $T_1 = (T, \diamond_1)$ is a group with unit 1_{\diamond_1} , $T_2 = (T, \diamond_2)$ is a semigroup and the following identity

(15)
$$a\diamond_2(b\diamond_1 c) = (a\diamond_2 b)\diamond_1\omega(a)^{\diamond_1}\diamond_1(a\diamond_2 c)$$

holds for all $a, b, c \in T$. We will denote the previous skew truss by $\mathbb{T} = (T_1, T_2, \omega_T)$. Then, if \mathbb{T} is a skew truss \mathbb{T} is a Hopf truss in the category Set. A morphism f between two skew trusses $\mathbb{T} = (T_1, T_2, \omega_T)$ and $\mathbb{S} = (S_1, S_2, \omega_S)$ is a map f between the two underlying sets such that f is a morphism of groups between T_1 and S_1 and of semigroups between T_2 and S_2 . Then, by [3, Proposition 2.8], the equality $\omega_S \circ f = f \circ \omega_T$ holds. With SkTr we will denote the category of skew trusses.

Let \mathbb{F} be a field. Let \mathbb{T} be a skew truss. Then $\mathbb{F}[T]$ admits a structure of Hopf truss in \mathbb{F} -Vect where the products, coproduct, counit and antipode are defined as in Remark 2.10 and the comonoid morphism $\sigma_{\mathbb{F}[T]}$ is the linear extension of ω_T . Also, if f is a morphism between skew trusses, its linear extension is a morphism of Hopf trusses. As a consequence, there exists a functor

$$P_{st}$$
 : SkTr \rightarrow HTr

given by

$$\mathsf{P}_{st}(\mathbb{T}) = (\mathsf{L}_g(T_1), \mathsf{L}_{sg}(T_2), \sigma_{\mathbb{F}[T]})$$

where L_g , L_{sg} are the functors defined in Remark 2.10 and $\lambda_{\mathbb{F}[T]} = ()^{\diamond_1}$.

Let \mathbb{F} be a field and $\mathbb{H} = (H_1, H_2, \sigma_H)$ a Hopf truss in \mathbb{F} -Vect. Let G(H) be the set of grouplike elements of the comonoid $(H, \varepsilon_H, \delta_H)$. In light of Remark 2.10, on the one hand we have that G(H) is a group. On the other hand, G(H) is a semigroup. As σ_H is a comonoid morphism, $\sigma_H(h) \in G(H) \forall h \in G(H)$. If we denote by ω_{σ_H} the restriction of σ_H to G(H), we have that $(G_g(H_1), G_{sg}(H_2), \omega_{\sigma_H})$ is an object in SkTr because equality 15 holds as a consequence of (iii) of Definition 2.12. By the functoriality of G_g and G_{sg} any morphism $f : \mathbb{H} \to \mathbb{B}$ of Hopf trusses induces a morphism of skew trusses between $(\mathsf{G}_g(H_1), \mathsf{G}_{sg}(H_2), \omega_{\sigma_H})$ and $(\mathsf{G}_g(B_1), \mathsf{G}_{sg}(B_2), \omega_{\sigma_B})$ defined by $\mathsf{G}_q(f)$ or by $\mathsf{G}_{sq}(f)$. Therefore

$$R_{ht}$$
 : HTr \rightarrow SkTr

defined by $\mathsf{R}_{ht}(\mathbb{H}) = (\mathsf{G}_g(H_1), \mathsf{G}_{sg}(H_2), \omega_{\sigma_H})$ on objects and by $\mathsf{R}_{ht}(f) = \mathsf{G}_{sg}(f)$ on morphisms is a functor between HTr and SkTr.

Definition 2.16. Let \mathbb{F} be a field and let \mathbb{H} be a Hopf truss in $\mathsf{Vect}_{\mathbb{F}}$. We will say that \mathbb{H} is pointed cosemisimple if the its subjacent coalgebra $(H, \varepsilon_H, \delta_H)$ is pointed and cosemisimple.

Theorem 2.17. Let P_{st} and R_{ht} be the functors defined in the previous remark. Then, $P_{st} \dashv R_{ht}$ and this adjunction induces an equivalence of categories between SkTr and the full subcategory of HTr of all pointed cosemisimple Hopf trusses.

Proof. Let \mathbb{T} be an object in SkTr and let \mathbb{H} be an object in HTr. We define a map

 $\Gamma_{\mathbb{T},\mathbb{H}}:\mathsf{Hom}_{\mathsf{HTr}}(\mathsf{P}_{st}(\mathbb{T}),\mathbb{H})\to\mathsf{Hom}_{\mathsf{SkTr}}(\mathbb{T},\mathsf{R}_{ht}(\mathbb{H}))$

in the following way: if $q: \mathsf{P}_{st}(\mathbb{T}) \to \mathbb{H}$ is a morphism in $\mathsf{Hom}_{\mathsf{HTr}}(\mathsf{P}_{st}(\mathbb{T}), \mathbb{H})$, $\Gamma_{\mathbb{T},\mathbb{H}}(q): T \to \mathsf{G}(\mathsf{H})$ is the map defined by $\Gamma_{\mathbb{T},\mathbb{H}}(q)(a) = q(a)$. Then, using that q Hopf monoid morphism between $\mathsf{L}_g(T_1)$ and H_1 and a non-unital bimonoid morphism between $\mathsf{L}_{sg}(T_2)$ and H_2 , we obtain that $\Gamma_{\mathbb{T},\mathbb{H}}(q)$ is a group morphism between T_1 and $\mathsf{G}_g(H_1)$ and a semigroup morphism between T_2 and $\mathsf{G}_{sg}(H_2)$. Therefore, $\Gamma_{\mathbb{T},\mathbb{H}}(q)$ is well defined. It is easy to show that it is natural in both components and injective. Moreover, it is surjective because, if $h:\mathbb{T}\to\mathsf{R}_{ht}(\mathbb{H})$ is a morphism in SkTr , $h=\Gamma_{\mathbb{T},\mathbb{H}}(l_h)$, where l_h is the unique morphisms in HTr such that $l_h(a) = h(a)$ for all $a \in T$. Indeed, it is easy to show that l_h is a Hopf monoid morphism between $\mathsf{L}_g(T_1)$ and H_1 and a morphism of non-unital bimonoids between $\mathsf{L}_{sg}(T_2)$ and H_2 . Finally, the identity $h = \Gamma_{\mathbb{T},\mathbb{H}}(l_h)$ follows trivially from the definition of l_h .

Therefore $\mathsf{P}_{st} \dashv \mathsf{R}_{ht}$ and the unit of this adjuction is the identity while the counit is not a natural isomorphism in general. In any case, if \mathbb{H} is a pointed cosemisimple Hopf truss the counit is an isomorphism, as the counits for the adjunctions given in Remark 2.10 are, and thus we have the equivalence.

3. Hopf trusses and generalized invertible 1-cocycles

In [2] the authors proved that there exists a closed relation between Hopf braces and 1-cocycles. In this section we will prove that this connection remains valid for Hopf trusses. First we will introduce the notion of generalized invertible 1-cocycle between a non-unital bimonoid B and a Hopf monoid H in the braided monoidal category C.

Definition 3.1. Let $H = (H, \eta_H, \mu_H, \varepsilon_H, \delta_H, \lambda_H)$ be a Hopf monoid in C and let $B = (B, \mu_B, \varepsilon_B, \delta_B)$ be a non-unital bimonoid in C. Assume that H is a non-unital left B-module monoid with action ϕ_H . Let $\pi : B \to H$ be comonoid morphism. We will say that π is an generalized invertible 1-cocycle if it is an isomorphism and there exist a comonoid endomorphism $\theta_{\pi} : B \to B$ such that

(16)
$$\pi \circ \mu_B = \mu_H \circ ((\pi \circ \theta_\pi) \otimes \phi_H) \circ (\delta_B \otimes \pi)$$

holds.

Let $\pi : B \to H$ and $\pi' : B' \to H'$ be generalized invertible 1-cocycles. A morphism between them is a pair (f,g) where $f : B \to B'$ is a morphism of non-unital bimonoids and $g : H \to H'$ is a morphism of Hopf monoids satisfying the following identities:

(17)
$$f \circ \theta_{\pi} = \theta_{\pi'} \circ f,$$

CATEGORICAL EQUIVALENCES FOR HOPF TRUSSES AND THEIR MODULES

(18)
$$g \circ \pi = \pi' \circ f,$$

(19)
$$g \circ \phi_H = \phi_{H'} \circ (f \otimes g).$$

Then, with these morphisms, generalized invertible 1-cocycles form a category denoted by GIC. In the following lines an object in GIC will also be denoted by the triple $(\pi : B \to H, \theta_{\pi})$.

Note that if $(\pi : B \to H, \theta_{\pi})$ is a generalized invertible 1-cocycle such that B is a Hopf monoid, (H, ϕ_H) is a left B-module monoid and $\theta_{\pi} = id_B$, $(\pi : B \to H, \theta_{\pi})$ is an invertible 1-cocycle in the sense of [2]. If we denote the category of invertible 1-cocycles by GIC, it is obvious that it is a subcategory of GIC.

Theorem 3.2. The categories GIC and HTr are equivalent.

Proof. Let $\mathbb{H} = (H_1, H_2, \sigma_H)$ be an object in HTr. Then, $(id_H : H_2 \to H_1, \theta_{id_H} = \sigma_H)$ is a generalized invertible 1-cocycle. Indeed, trivially id_H is a comonoid isomorphism and H_1 is a non unital left H_2 -module monoid with the action $\Gamma_{H_1}^{\sigma_H}$ defined in (iii) of Definiton (2.12). Finally,

$$\begin{split} & \mu_{H}^{1} \circ \left(\left(id_{H} \circ \theta_{id_{H}} \right) \otimes \Gamma_{H_{1}}^{\sigma_{H}} \right) \circ \left(\delta_{H} \otimes id_{H} \right) \\ &= \mu_{H}^{1} \circ \left(\sigma_{H} \otimes \left(\mu_{H}^{1} \circ \left(\left(\lambda_{H} \circ \sigma_{H} \right) \otimes \mu_{H}^{2} \right) \circ \left(\delta_{H} \otimes H \right) \right) \right) \circ \left(\delta_{H} \otimes H \right) \text{ (by the definition of } \Gamma_{H_{1}}^{\sigma_{H}} \right) \\ &= \mu_{H}^{1} \circ \left(\left(\mu_{H}^{1} \circ \left(\sigma_{H} \otimes \left(\lambda_{H} \circ \sigma_{H} \right) \circ \delta_{H} \right) \right) \otimes \mu_{H}^{2} \right) \circ \delta_{H} \text{ (by the associativity of } \mu_{H}^{1} \text{ and the coassociativity of } \\ & \delta_{H} \right) \\ &= \mu_{H}^{1} \circ \left(\left(id_{H} * \lambda_{H} \right) \otimes \mu_{H}^{2} \right) \circ \delta_{H} \text{ (by the condition of comonoid morphism for } \sigma_{H} \right) \\ &= id_{H} \circ \mu_{H}^{2} \text{ (by (8) and the properties of } \varepsilon_{H} \text{ and } \eta_{H} \text{)}. \end{split}$$

On the other hand, let $\mathbb{H} = (H_1, H_2, \sigma_H)$ and $\mathbb{H}' = (H'_1, H'_2, \sigma_{H'})$ be objects in HTr and let $f : \mathbb{H} \to \mathbb{H}'$ be a morphism between them. The pair (f, f) is a morphism in GIC between $(id_H : H_2 \to H_1, \sigma_H)$ and $(id_{H'} : H'_2 \to H'_1, \sigma_{H'})$ because $f : H_1 \to H'_1$ is a Hopf monoid morphism, $f : H_2 \to H'_2$ is a non-unital bimonoid morphisn, (17) holds trivially, (18) holds by (14), and, finally, (19) follows from

$$\begin{split} & f \circ \Gamma_{H_1}^{\sigma_H} \\ &= \mu_H^1 \circ \left((\lambda_H \circ f \circ \sigma_H) \otimes (\mu_{H'}^2 \circ (f \otimes f)) \right) \circ (\delta_H \otimes H) \text{ (by the condition of multiplicative morphism for } f \\ & \text{and } (9)) \\ &= \mu_H^1 \circ \left((\lambda_H \circ \sigma_{H'} \circ f) \otimes (\mu_{H'}^2 \circ (f \otimes f)) \right) \circ (\delta_H \otimes H) \text{ (by } (14)) \\ &= \Gamma_{H'}^{\sigma_{H'}} \circ (f \otimes f) \text{ (by the condition of comonoid morphism for } f). \end{split}$$

Therefore, there exists a functor

$$E: HTr \rightarrow GIC$$

defined on objects by $\mathsf{E}(\mathbb{H}) = (id_H : H_2 \to H_1, \sigma_H)$ and on morphisms by $\mathsf{E}(f) = (f, f)$.

The next step is to define the functor

$$Q: IC \rightarrow HTr$$

Let $(\pi : B \to H, \theta_{\pi})$ be an object in GIC. Define $\mu_{H}^{\pi} := \pi \circ \mu_{B} \circ (\pi^{-1} \otimes \pi^{-1})$ and $\sigma_{\pi} := \pi \circ \theta_{\pi} \circ \pi^{-1}$. Then, $\mathbb{H}_{\pi} = (H, H_{\pi}, \sigma_{\pi})$, where $H_{\pi} = (H, \mu_{H}^{\pi}, \varepsilon_{H}, \delta_{H})$, is an object in HTr. Indeed, trivially H_{π} is a non-unital bimonoid in C because π is a comonoid isomorphism and B is a non-unital bimonoid. The morphism σ_{π} is a comonoid morphism because is defined as a composition of comonoid morphisms. Finally,

$$\mu_{H} \circ (\mu_{H}^{\pi} \otimes \Gamma_{H}^{\sigma_{\pi}}) \circ (H \otimes c_{H,H} \otimes H) \circ (\delta_{H} \otimes H \otimes H)$$

$$= \mu_{H} \circ ((\mu_{H} \circ ((\pi \circ \theta_{\pi}) \otimes \phi_{H}) \circ (\delta_{B} \otimes H)) \otimes (\mu_{H} \circ ((((\lambda_{H} * id_{H}) \circ \pi \circ \theta_{\pi}) \otimes \phi_{H}) \circ (\delta_{B} \otimes H))))$$

$$\circ (B \otimes c_{B,H} \otimes H) \circ ((\delta_{B} \circ \pi^{-1}) \otimes H \otimes H) \text{ (by (16), associativity of } \mu_{H}, \text{ coassociativity of } \delta_{H} \text{ and naturality}$$
of c)
$$= \mu_{H} \circ ((\pi \circ \theta_{\pi}) \otimes (\mu_{H} \circ (\phi_{H} \otimes \phi_{H}) \circ (B \otimes c_{B,H} \otimes H) \circ (\delta_{B} \otimes H \otimes H))) \circ ((\delta_{B} \circ \pi^{-1}) \otimes H \otimes H)$$

10

(by (8), the condition of comonoid morphism for θ_{π} and π , the properties of the unit η_H and the counit ε_B and the coassociativity of δ_H)

 $= \mu_H^{\pi} \circ (H \otimes \mu_H)$ (by the condition of non-unital left *B*-module monoid for *H* and (16)).

Also, if (f,g) is a morphism in GIC between $(\pi : B \to H, \theta_{\pi})$ and $(\pi' : B' \to H', \theta_{\pi'})$, g is a morphism in HTr between \mathbb{H}_{π} and $\mathbb{H}'_{\pi'}$ because, $g : H \to H'$ is a morphism of Hopf monoids and $g : H_{\pi} \to H'_{\pi'}$ is a morphism in bimod because

$$\begin{split} g \circ \mu_H^{\pi} \\ &= g \circ \pi \circ \mu_B \circ \left(\pi^{-1} \otimes \pi^{-1} \right) \text{ (by definition of } \mu_H^{\pi} \text{)} \\ &= \pi' \circ f \circ \mu_B \circ \left(\pi^{-1} \otimes \pi^{-1} \right) \text{ (by (18))} \\ &= \pi' \circ \mu_{B'} \circ \left(\left(f \circ \pi^{-1} \right) \otimes \left(f \circ \pi^{-1} \right) \right) \text{ (by the condition of multiplicative morphism for } f \text{)} \\ &= \pi' \circ \mu_{B'} \circ \left(\left(\pi'^{-1} \circ g \right) \otimes \left(\pi'^{-1} \circ g \right) \right) \text{ (by (18))} \\ &= \mu_{H'}^{\pi'} \circ \left(g \otimes g \right) \text{ (by definition of } \mu_{H'}^{\pi'} \text{).} \end{split}$$

As a consequence of these facts, we have a functor $Q : IC \to HTr$ defined by $Q((\pi : B \to H, \theta_{\pi})) = \mathbb{H}_{\pi}$ on objects and by Q((f, g)) = g on morphisms.

These functors induce an equivalence between the two categories because, clearly, $QE = id_{HTr}$. On the other hand, we have that

$$(\pi, id_H) : (\pi : B \to H, \theta_\pi) \to (id_H : H_\pi \to H, \sigma_\pi) = \mathsf{EQ}((\pi : B \to H, \theta_\pi))$$

is an isomorphism in GIC because $\pi : B \to H_{\pi}$ is a isomorphism between B and H_{π} in bimod, $\pi \circ \theta_{\pi} = \sigma_{\pi} \circ \pi$, (18) holds trivially, and

 $\Gamma_{H}^{\sigma_{\pi}} \circ (\pi \otimes id_{H})$ $= \mu_{H} \circ ((\lambda_{H} \circ \pi \circ \theta_{\pi} \circ \pi^{-1}) \otimes (\pi \circ \mu_{B} \circ (\pi^{-1} \otimes \pi^{-1}))) \circ ((\delta_{H} \circ \pi) \otimes H)$ (by definition of σ_{π} and μ_{H}^{π}) $= \mu_{H} \circ (((\lambda_{H} * id_{H}) \circ \pi \circ \theta_{\pi}) \otimes \phi_{H}) \circ (\delta_{B} \otimes H)$ (by (16), the condition of comonoid morphism for θ_{π} and π) $= \phi_{\pi} (h_{\pi} \otimes h_{\pi}) \otimes h_{\pi} \otimes h_{$

= ϕ_H (by (8), the condition of comonoid morphism for θ_{π} and π and the properties of the unit η_H and the counit ε_B).

Therefore, $EQ \simeq Id_{GIC}$.

Remark 3.3. As is easy to see, the previous theorem is a generalization of [2, Theorem 1.12] because if we particularize this equivalence to the Hopf brace setting, we obtain the quoted result.

4. Modules for Hopf trusses and generalized invertible 1-cocycles

In this section we introduce the notion of module over a Hopf truss and over a generalized invertible 1-cocycle. Also we will prove that, thanks to the categorical equivalence of Theorem 3.2, there exist some interesting categorical connections between these categories of modules.

Definition 4.1. Let \mathbb{H} be a Hopf truss. A left \mathbb{H} -module is a triple (M, ψ_M^1, ψ_M^2) , where (M, ψ_M^1) is a left H_1 -module, (M, ψ_M^2) is a non-unital left H_2 -module and the following identity

(20)
$$\psi_M^2 \circ (H \otimes \psi_M^1) = \psi_M^1 \circ (\mu_H^2 \otimes \Gamma_M^{\sigma_H}) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes M)$$

holds, where

$$\Gamma_M^{\sigma_H} = \psi_M^1 \circ ((\lambda_H \circ \sigma_H) \otimes \psi_M^2) \circ (\delta_H \otimes M).$$

Given two left \mathbb{H} -modules (M, ψ_M^1, ψ_M^2) and (N, ψ_N^1, ψ_N^2) , a morphism $f : M \to N$ is called a morphism of left \mathbb{H} -modules if f is a morphism of left H_1 -modules and left non-unital H_2 -modules. Left \mathbb{H} -modules with morphisms of left \mathbb{H} -modules form a category which we denote by $\mathbb{H}Mod$.

Example 4.2. Let \mathbb{H} be a Hopf truss. The triple $(H, \psi_H^1 = \mu_H^1, \psi_H^2 = \mu_H^2)$ is an example of left \mathbb{H} -module. Also, if K is the unit object of C, $(K, \psi_K^1 = \varepsilon_H, \psi_K^2 = \varepsilon_H)$ is a left \mathbb{H} -module called the trivial module.

If X is an object in C, the triple

$$\mathbb{H} \otimes X = (H \otimes X, \psi^1_{H \otimes X} = \mu^1_H \otimes X, \psi^2_{H \otimes X} = \mu^2_H \otimes X)$$

is an example of left \mathbb{H} -module. Also, if $f : X \to X'$ is a morphism in $\mathsf{C}, H \otimes f$ is a morphism in $\mathbb{H}\mathsf{M}\mathsf{od}$ between $\mathbb{H} \otimes X$ and $\mathbb{H} \otimes X'$. Therefore, there exist a functor, called the induction functor,

 $\mathbb{H}\otimes -:\mathsf{C}\to\ _{\mathbb{H}}\mathsf{Mod}$

defined on objects by $\mathbb{H} \otimes -(X) = \mathbb{H} \otimes X$ and by $\mathbb{H} \otimes -(f) = \mathbb{H} \otimes f$ on morphisms.

Remark 4.3. If the a Hopf truss \mathbb{H} is a Hopf brace and in Definition 4.1 we assume that a (M, ψ_M^2) is a left H_2 module, we obtain the definition of module over a Hopf brace introduced in [5].

Remark 4.4. Using the naturality of c and the coassociativity of δ_H , it is easy to show that (20) is equivalent to

(21)
$$\psi_M^2 \circ (H \otimes \psi_M^1) = \psi_M^1 \circ (\Lambda_{H_1}^{\sigma_H} \otimes \psi_M^2) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes M)$$

where

$$\Lambda_{H_1}^{\sigma_H} = \mu_H^1 \circ (\mu_H^2 \otimes (\lambda_H \circ \sigma_H)) \circ (H \otimes c_{H,H}) \circ (\delta_H \otimes H).$$

Also, the following equality

(22)
$$\Gamma_M^{\sigma_H} \circ (H \otimes \psi_M^1) = \psi_M^1 \circ (\Gamma_{H_1}^{\sigma_H} \otimes \Gamma_M^{\sigma_H}) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes M).$$

holds. Indeed:

$$\begin{split} &\psi_{M}^{1}\circ\left(\Gamma_{H_{1}}^{\sigma_{H}}\otimes\Gamma_{M}^{\sigma_{H}}\right)\circ\left(H\otimes c_{H,H}\otimes M\right)\circ\left(\delta_{H}\otimes H\otimes M\right)\\ &=\psi_{M}^{1}\circ\left(\left(\lambda_{H}\circ\sigma_{H}\right)\otimes\left(\psi_{M}^{1}\circ\left(\Lambda_{H_{1}}^{\sigma_{H}}\otimes\psi_{M}^{2}\right)\circ\left(H\otimes c_{H,H}\otimes M\right)\circ\left(\delta_{H}\otimes H\otimes M\right)\right)\right)\circ\left(\delta_{H}\otimes H\otimes M\right)\\ &\text{ (by naturality of c, coassociativity of δ_{H}, the condition of left H_{1}-module for (M,ψ_{M}^{1}) and the associativity of μ_{H}^{1})}\\ &=\Gamma_{M}^{\sigma_{H}}\circ\left(H\otimes\psi_{M}^{1}\right) \text{ (by (21))}. \end{split}$$

Definition 4.5. Let $(\pi : B \to H, \theta_{\pi})$ be a generalized invertible 1-cocycle. A left module over $(\pi : B \to H, \theta_{\pi})$ is a 6-tuple $(M, N, \phi_M, \varphi_M, \phi_N, \gamma)$ where

- (i) $\phi_M : B \otimes M \to M$ is a morphism in C.
- (ii) (M, φ_M) is a left *H*-module.
- (iii) (N, ϕ_N) is a non-unitary left *B*-module.
- (iv) The equality

(23)
$$\phi_M \circ (B \otimes \varphi_M) = \varphi_M \circ (\phi_H \otimes \phi_M) \circ (B \otimes c_{A,H} \otimes M) \circ (\delta_B \otimes H \otimes M).$$

holds.

(v) $\gamma: N \to M$ is an isomorphism in C such that

(24)
$$\gamma \circ \phi_N = \varphi_M \circ ((\pi \circ \theta_\pi) \otimes \phi_M) \circ (\delta_B \otimes \gamma).$$

Definition 4.6. Let $(M, N, \phi_M, \varphi_M, \phi_N, \gamma)$ and $(M', N', \phi_{M'}, \varphi_{M'}, \phi_{N'}, \gamma')$ be left modules over a generalized invertible 1-cocycle $(\pi : B \to H, \theta_{\pi})$. A morphism between them is a pair (h, l) of morphisms in C such that:

- (i) The morphism $h: M \to M'$ satisfies $f \circ \phi_M = \phi_{M'} \circ (B \otimes f)$ and is left *H*-linear.
- (ii) The morphism $l: N \to N'$ is left *B*-linear.
- (iii) The following identity holds:

$$h \circ \gamma = \gamma' \circ l.$$

Note that, by (25), the morphism l is determined by h because $l = (\gamma')^{-1} \circ h \circ \gamma$.

With the obvious composition of morphisms, left modules over a generalized invertible 1-cocycle $(\pi : B \to H, \theta_{\pi})$ with action ϕ_H form a category that we will denote by $(\pi, \phi_H, \theta_{\pi})$ Mod.

Remark 4.7. If $(M, N, \phi_M, \varphi_M, \phi_N, \gamma)$ is a left module over the generalized invertible 1-cocycle $(\pi : B \to H, \theta_\pi)$, by (24), we obtain that ϕ_N is determined by ϕ_M and φ_M in the following way:

(26)
$$\phi_N = \gamma^{-1} \circ \varphi_M \circ ((\pi \circ \theta_\pi) \otimes \phi_M) \circ (\delta_B \otimes \gamma).$$

Also, composing in both sides of the equality (24) with $((\lambda_H \circ \pi) \otimes B) \circ \delta_B) \otimes \gamma^{-1}$ on the right and with φ_M on the left we obtain the identity

(27)
$$\phi_M = \varphi_M \circ ((\lambda_H \circ \pi \circ \theta_\pi) \otimes (\gamma \circ \phi_N)) \circ (\delta_B \otimes \gamma^{-1})$$

because:

 $\begin{aligned} \varphi_M \circ \left((\lambda_H \circ \pi \circ \theta_\pi) \otimes (\gamma \circ \phi_N) \right) \otimes (\delta_B \otimes \gamma^{-1}) \\ &= \varphi_M \circ \left((\lambda_H \circ \pi \circ \theta_\pi) \otimes (\varphi_M \circ ((\pi \circ \theta_\pi) \otimes \phi_M) \circ (\delta_B \otimes \gamma)) \otimes (\delta_B \otimes \gamma^{-1}) \text{ (by (24))} \\ &= \varphi_M \circ \left(\left((\lambda_H * id_H) \circ \pi \circ \theta_\pi \right) \otimes \phi_M \right) \otimes (\delta_B \otimes M) \text{ (by the condition of comonoid morphism for } \theta_\pi \text{ and } \pi, \\ \text{the coassociativity of } \delta_B) \\ &= \phi_M \text{ (by (8) and the unit counit properties)).} \end{aligned}$

Example 4.8. It is easy to see that if $(\pi : B \to H, \theta_{\pi})$ is a generalized invertible 1-cocycle, the 6-tuple $(H, B, \phi_H, \mu_H, \mu_B, \pi)$ is an example of left module over $(\pi : B \to H, \theta_{\pi})$.

Theorem 4.9. Assume that (f,g) is a morphism between the generalized invertible 1-cocycles $(\pi : B \to H, \theta_{\pi})$ and $(\pi : B' \to H', \theta_{\pi'})$. Then, there exists a functor

$$\mathsf{M}_{(f,g)}: \ _{(\pi',\phi_{H'},\theta_{\pi'})}\mathsf{Mod} \ o \ _{(\pi,\phi_H,\theta_{\pi})}\mathsf{Mod}$$

defined on objects by

$$\mathsf{M}_{(f,g)}((P,Q,\phi_P,\varphi_P,\phi_Q,\tau)) = (P,Q,\phi_P^{\pi} = \phi_P \circ (f \otimes P), \varphi_P^{\pi} = \varphi_P \circ (g \otimes P), \phi_Q^{\pi} = \phi_Q \circ (f \otimes Q), \tau)$$

and on morphisms by the identity.

Proof. The existence of the functor $\mathsf{M}_{(f,g)}$ is a consequence of the following facts: Trivially (P, φ_P^{π}) is a left *H*-module and (Q, ϕ_Q^{π}) is a non-unitary left *B*-modules and . Also,

$$\begin{split} \varphi_P^{\pi} \circ (\phi_H \otimes \phi_P^{\pi}) \circ (B \otimes c_{B,H} \otimes P) \circ (\delta_B \otimes H \otimes P) \\ &= \varphi_P \circ ((g \circ \phi_H) \otimes (\phi_P \circ (f \otimes P))) \circ (B \otimes c_{B,H} \otimes P) \circ (\delta_B \otimes H \otimes P) \text{ (by definition of } \phi_P^{\pi} \text{ and } \varphi_P^{\pi}) \\ &= \varphi_P \circ (\phi_{H'} \otimes \phi_P) \circ (B' \otimes c_{B',H'} \otimes P) \circ (((f \otimes f) \circ \delta_B) \otimes g \otimes P) \text{ (by (19) and naturality of } c) \\ &= \varphi_P \circ (\phi_{H'} \otimes \phi_P) \circ (B' \otimes c_{B',H'} \otimes P) \circ ((\delta_{B'} \circ f) \otimes g \otimes P) \text{ (by the comonoid morphism condition for } f) \\ &= \phi_P^{\pi} \circ (B \otimes \varphi_P^{\pi}) \text{(by (23))}, \end{split}$$

and

 $\begin{aligned} \varphi_P^{\pi} \circ \left(\left(\pi \circ \theta_{\pi} \right) \otimes \phi_P^{\pi} \right) \circ \left(\delta_B \otimes \tau \right) \\ &= \varphi_P \circ \left(\left(g \circ \pi \circ \theta_{\pi} \right) \otimes \left(\phi_P \circ \left(f \otimes P \right) \right) \right) \circ \left(\delta_B \otimes \tau \right) \text{ (by definition of } \phi_P^{\pi} \text{ and } \varphi_P^{\pi} \right) \\ &= \varphi_P \circ \left(\left(\pi' \circ \theta_{\pi'} \circ f \right) \otimes \left(\phi_P \circ \left(f \otimes P \right) \right) \right) \circ \left(\delta_B \otimes \tau \right) \text{ (by (17) and (18))} \\ &= \varphi_P \circ \left(\left(\pi' \circ \theta_{\pi'} \right) \otimes \phi_P \right) \circ \left(\left(\delta_{B'} \circ f \right) \otimes \tau \right) \text{ (by the comonoid morphism condition for } f) \\ &= \tau \circ \phi_Q^{\pi} \text{ (by (24))} \end{aligned}$

Then $(P, Q, \phi_P^{\pi}, \varphi_P^{\pi}, \phi_Q^{\pi}, \tau)$ is an object in $(\pi, \phi_H, \theta_{\pi})$ Mod. Finally, it is obvious that if (h, l) is a morphism in $(\pi', \phi_{H'}, \theta_{\pi'})$ Mod, (h, l) is a morphism in $(\pi, \phi_H, \theta_{\pi})$ Mod.

Remark 4.10. If (f,g) is an isomorphism defined between the generalized invertible 1-cocycles $(\pi: B \to H, \theta_{\pi})$ and $(\pi: B' \to H', \theta_{\pi'})$ with inverse (f^{-1}, g^{-1}) , the functor $\mathsf{M}_{(f,g)}$ is an isomorphism of categories with inverse $\mathsf{M}_{(f^{-1},g^{-1})}$. For example, in the proof of Theorem 3.2 we proved that, for all generalized invertible 1-cocycle $(\pi: B \to H, \theta_{\pi})$, (π, id_H) is an isomorphism between the generalized invertible 1-cocycles $(\pi: B \to H, \theta_{\pi})$, (π, id_H) and $(id_H: H_{\pi} \to H, \sigma_{\pi})$ where $\sigma_{\pi} = \pi \circ \theta_{\pi} \circ \pi^{-1}$ and the action is $\Gamma_{H}^{\sigma_{\pi}}$, i.e. the action is $\phi_{H} \circ (\pi^{-1} \otimes H)$ where ϕ_{H} is the action associated to $(\pi: B \to H, \theta_{\pi})$. Therefore, the functor

$$\mathsf{M}_{(\pi,id_H)}: {}_{(id_H,\Gamma^{\sigma_{\pi}}_{_{_{\mathcal{H}}}},\sigma_{\pi})}\mathsf{Mod} \rightarrow {}_{(\pi,\phi_H,\theta_{\pi})}\mathsf{Mod}$$

is an isomorphism of categories with inverse

$$\mathsf{M}_{(\pi^{-1}, id_H)} : \ _{(\pi, \phi_H, \theta_\pi)}\mathsf{Mod} \ \rightarrow \ _{(id_H, \Gamma_H^{\sigma_\pi}, \sigma_\pi)}\mathsf{Mod}$$

Theorem 4.11. Let \mathbb{H} be a Hopf truss and let $\mathsf{E}(\mathbb{H})$ be the invertible 1-cocycle induced by the functor E introduced in the proof of Theorem 3.2. There exists a functor

$$\mathsf{G}_{\mathbb{H}}: \mathbb{H}\mathsf{Mod} \to (id_H, \Gamma_{H_1}^{\sigma_H}, \sigma_H)\mathsf{Mod}$$

defined on objects by

$$\mathsf{G}_{\mathbb{H}}((M,\psi_{M}^{1},\psi_{M}^{2})) = (M,M,\widehat{\phi}_{M} = \Gamma_{M}^{\sigma_{H}},\widehat{\varphi}_{M} = \psi_{M}^{1},\overline{\phi}_{M} = \psi_{M}^{2}, id_{M})$$

and on morphisms by $G_{\mathbb{H}}(f) = (f, f)$.

Proof. By assumption, $(M, \widehat{\varphi}_M = \psi_M^1)$ is a left H_1 -module and $(M, \overline{\phi}_M = \psi_M^2)$ is a non-unital left H_2 -module. On the other hand, by (22) we have that

$$\widehat{\phi}_M \circ (H \otimes \widehat{\varphi}_M) = \widehat{\varphi}_M \circ (\Gamma_{H_1} \otimes \widehat{\phi}_M) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes M)$$

and then, (23) holds. Also,

$$\begin{aligned} \widehat{\varphi}_{M} &\circ (\sigma_{H} \otimes \widehat{\phi}_{M}) \circ (\delta_{H} \otimes M) \\ &= \psi_{M}^{1} \circ (\sigma_{H} \otimes \Gamma_{M}^{\sigma_{H}}) \circ (\delta_{H} \otimes M) \text{ (by definition of } \widehat{\varphi}_{M} \text{ and } \widehat{\phi}_{M}) \\ &= \psi_{M}^{1} \circ (\sigma_{H} \otimes (\psi_{M}^{1} \circ ((\lambda_{H} \circ \sigma_{H}) \otimes \psi_{M}^{2}) \circ (\delta_{H} \otimes M))) \circ (\delta_{H} \otimes M) \text{ (by definition of } \Gamma_{M}^{\sigma_{H}}) \\ &= \psi_{M}^{1} \circ (((id_{H} * \lambda_{H}^{1}) \circ \sigma_{H}) \otimes \psi_{M}^{2}) \circ (\delta_{H} \otimes M) \text{ (by the condition of left } H_{1}\text{-module of } (M, \psi_{M}^{1}), \text{ the } \\ &= \overline{\phi}_{M} \text{ (by (8), the counit properties, the condition of left } H_{1}\text{-module of } (M, \psi_{M}^{1}) \text{ and the definition of } \overline{\phi}_{M}). \end{aligned}$$

Finally, it is easy to show that if f is a morphism in \mathbb{H} Mod between the objects (M, ψ_M^1, ψ_M^2) and $(M', \psi_{M'}^1, \psi_{M'}^2)$, the pair (f, f) is a morphism in $_{(id_H, \Gamma_{H_1}^{\sigma_H}, \sigma_H)}$ Mod between $\mathsf{G}_{\mathbb{H}}((M, \psi_M^1, \psi_M^2))$ and $\mathsf{G}_{\mathbb{H}}((M', \psi_{M'}^1, \psi_{M'}^2))$.

Theorem 4.12. Let $(\pi : B \to H, \theta_{\pi})$ be a generalized invertible 1-cocycle. Then the categories $(\pi, \phi_H, \theta_{\pi})$ Mod and \mathbb{H}_{π} Mod are equivalent.

Proof. First of all we will prove that there exists a functor

$$\mathsf{H}^{\pi}_{\mathsf{tr}}: \ _{(\pi,\phi_{H}, heta_{\pi})}\mathsf{Mod} \ o \ _{\mathbb{H}_{\pi}}\mathsf{Mod}$$

defined on objects by

$$\mathsf{H}^{\pi}_{\mathsf{tr}}((M, N, \phi_M, \varphi_M, \phi_N, \gamma)) = (M, \overline{\psi}^1_M = \varphi_M, \overline{\psi}^2_M = \gamma \circ \phi_N \circ (\pi^{-1} \otimes \gamma^{-1}))$$

and on morphisms by $\mathsf{H}^{\pi}_{\mathsf{tr}}((h,l)) = h$. Indeed: By assumption, $(M, \overline{\psi}^1_M = \varphi_M)$ is a left *H*-module and, using the condition of non-unital left *B*-module of *N*, we obtain that

$$(M, \overline{\psi}_M^2 = \gamma \circ \phi_N \circ (\pi^{-1} \otimes \gamma^{-1}))$$

is a non-unital left H_{π} -module. Also, by (27), we have that the identity

(28)
$$\Gamma_M^{\sigma_{\pi}} = \phi_M \circ (\pi^{-1} \otimes M)$$

holds, where
$$\Gamma_M^{0\pi} = \psi_M^- \circ ((\lambda_H \circ \sigma_\pi) \otimes \psi_M^-) \circ (\delta_H \otimes M)$$
. Then, (20) holds because:
 $\overline{\psi}_M^2 \circ (H \otimes \overline{\psi}_M^1)$
 $= \varphi_M \circ ((\pi \circ \theta_\pi) \otimes \phi_M) \circ ((\delta_B \circ \pi^{-1}) \otimes \varphi_M) \text{ (by (26))}$
 $= \varphi_M \circ ((\pi \circ \theta_\pi) \otimes (\varphi_M \circ (\phi_H \otimes \phi_M) \circ (B \otimes c_{B,H} \otimes M) \circ (\delta_B \otimes H \otimes M))) \circ ((\delta_B \circ \pi^{-1}) \otimes H \otimes M)$
(by (23))
 $= \varphi_M \circ ((\mu_H \circ ((\pi \circ \theta_\pi) \otimes \phi_H) \circ (\delta_B \otimes \pi)) \otimes \phi_M) \circ (B \otimes c_{B,B} \otimes M) \circ ((\delta_B \circ \pi^{-1}) \otimes \pi^{-1} \otimes M)$
(by the condition of left *H*-module for *M*, the coassociativity of δ_B , the naturality of *c* and the condition of
isomorphism for π)
 $= \varphi_M \circ ((\pi \circ \mu_B) \otimes \phi_M) \circ (B \otimes c_{B,B} \otimes M) \circ ((\delta_B \circ \pi^{-1}) \otimes \pi^{-1} \otimes M) \text{ (by (16))}$
 $= \varphi_M \circ ((\mu_{H_\pi} \otimes (\phi_M \circ (\pi^{-1} \otimes M)))) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes M) \text{ (by the condition of coalgebra
isomorphism for π and the naturality of *c*)
 $= \overline{\psi}_M^1 \circ (\mu_{H_\pi} \otimes \Gamma_M^{\sigma_\pi}) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes M) \text{ (by (28))}$$

On the other hand, if (h, l) is a morphisms in $(\pi, \phi_H, \theta_\pi)$ Mod between $(M, N, \phi_M, \varphi_M, \phi_N, \gamma)$ and $(M', N', \phi_{M'}, \varphi_{M'}, \phi_{N'}, \gamma')$, we have that h is a morphism in \mathbb{H}_{π} Mod between $(M, \overline{\psi}_M^1, \overline{\psi}_M^2)$ and $(M', \overline{\psi}_{M'}^1, \overline{\psi}_{M'}^2)$ because, using that h is a morphism of left H-modules, we have $h \circ \overline{\psi}_M^1 = \overline{\psi}_{M'}^1 \circ (H \otimes h)$ and, by (25) and the condition of morphism of non-unital left B-modules for l, we have that $h \circ \overline{\psi}_M^2 = \overline{\psi}_{M'}^2 \circ (H \otimes h)$.

Taking into account the functors H_{tr}^{π} , $G_{\mathbb{H}_{\pi}}$ and $M_{(\pi,id_{H})}$, it is easy to show that

$$\mathsf{H}_{\mathsf{br}}^{\pi} \circ (\mathsf{M}_{(\pi, id_{H})} \circ \mathsf{G}_{\mathbb{H}_{\pi}}) = \mathsf{id}_{\mathbb{H}_{\pi}\mathsf{Mod}}$$

and

 $((\mathsf{M}_{(\pi,id_H)} \circ \mathsf{G}_{\mathbb{H}_{\pi}}) \circ \mathsf{H}_{\mathsf{br}}^{\pi})((M, N, \phi_M, \varphi_M, \phi_N, \gamma)) = (M, M, \phi_M, \varphi_M, \overline{\phi}_M^{\pi} = \gamma \circ \phi_N \circ (A \otimes \gamma^{-1}), id_M)$ holds. Then,

$$(\mathsf{M}_{(\pi,id_H)}\circ\mathsf{G}_{\mathbb{H}_\pi})\circ\mathsf{H}^{\pi}_{\mathsf{br}} \simeq \mathsf{id}_{(\pi,\phi_H,\theta_\pi)}\mathsf{Mod}$$

because (id_M, γ) is an isomorphism in $(\pi, \phi_H, \theta_\pi)$ Mod between the objects $(M, N, \phi_M, \varphi_M, \phi_N, \gamma)$ and $(M, M, \phi_M, \varphi_M, \overline{\phi}_M^{\pi}, id_M)$.

Remark 4.13. When we particularize the previous results to Hopf braces we obtain the categorical equivalences obtained in [4]. More concretely, [4, Theorem 2.26] is the Hopf brace version of Theorem 28.

5. The fundamental theorem of Hopf modules for Hopf trusses

In this section we will assume that C admits equalizers. As a consequence every idempotent morphism in C splits, i.e., if $q: M \to M$ is a morphism in C such that $q = q \circ q$, there exists an object I(q), called the image of q, and morphisms $i: I(q) \to M$ and $p: M \to I(q)$ such that $q = i \circ p$ and $p \circ i = id_{I(q)}$. The morphisms p and i will be called a factorization of q. Note that I(q), p and i are unique up to isomorphism.

Under the previous condition, we will introduce the category \mathbb{H} -Hopf of left Hopf modules over a Hopf truss \mathbb{H} and we will to obtain the Fundamental Theorem of Hopf modules for Hopf trusses that is a generalization of the one proved in [5] for Hopf brace. **Definition 5.1.** Let D be a comonoid in C. The pair (M, ρ_M) is a left D-comodule if M is an object in C and $\rho_M: M \to D \otimes M$ is a morphism in C satisfying $(\varepsilon \otimes M) \circ \rho_M = id_M, (D \otimes \rho_M) \circ \rho_M =$ $(\delta \otimes M) \circ \rho_M$. Given two left D-comodules (M, ρ_M) and (N, ρ_N) , a morphism $f: M \to N$ in C is a morphism of left D-comodules if $(D \otimes f) \circ \rho_M = \rho_N \circ f$. Left D-comodules with morphisms of left D-comodules form a category which we denote by ^DComod.

Definition 5.2. Let D be a comonoid such that there exits a comonoid morphism $e: K \to D$. Let (M, ρ_M) be a left *D*-comodule. We define the subobject of coinvariants of *M*, denoted by M_e^{coD} , as the equalizer object of ρ_M and $e \otimes M$. Then, we have an equalizer diagram

$$M_e^{coD} \xrightarrow{j_M^e} M \xrightarrow{\rho_M} D \otimes M$$

where j_M^e denotes the equalizer (inclusion) morphism.

If H is a Hopf monoid, the unit η_H is a comonoid morphism. Then, Let (M, ρ_M) be a left D-comodule, we will denote the equalizer object of ρ_M and $\eta_H \otimes M$ by M^{coD} and the equalizer morphism by j_M .

Definition 5.3. Let B be a non-unital bimonoid. A non-unital left H-Hopf module over B is a triple (M, φ_M, ρ_M) where (M, φ_M) is a non-unital left B-module, (M, ρ_M) is a left B-comodule and

(29)
$$\varphi_M \circ \rho_M = (\mu_B \otimes \varphi_M) \circ (B \otimes c_{B,B} \otimes M) \circ (\delta_B \otimes \rho_M)$$

holds. Non-unital left B-Hopf modules over B with left linear and colinear morphisms form a category which we denote by B-Hopf-mod.

The definition for left H-Hopf modules over a Hopf monoid H is similar changing non-unital left H-modules by left H-modules. Then, in this case we will denote the category of H-Hopf modules over H by H-Hopf-Mod

Remark 5.4. Let H be a Hopf monoid, it easy to show that, if (M, φ_M, ρ_M) is a left H-Hopf module over H, the endomorphism $q_M: M \to M$, defined by

$$q_M = \varphi_M \circ (\lambda_H \otimes M) \circ \rho_M$$

is idempotent and satisfies $\rho_M \circ q_M = \eta_H \otimes q_M$. Therefore, there exists a unique morphism

$$t_M: M \to M^{coH}$$

such that $t_M \circ j_M = q_M$. Let $I(q_M)$ be the image of the idempotent morphism q_M and let i_M : $I(q_M) \to M$ and $p_M: M \to I(q_M)$ be the morphisms such that $p_M = i_M \circ p_M$ and $p_M \circ i_M = i_M \circ p_M$ $id_{I(q_M)}$. The morphism

$$\omega_M = t_M \circ i_M : I(q_M) \to M^{coH}$$

is an isomorphism with inverse $\omega_M^{-1} = p_M \circ j_M$. Moreover, $t_M \circ \varphi_M = \varepsilon_H \otimes t_M$ and, as a consequence,

 (M^{coH}, t_M) is the coequalizer of φ_M and $\varepsilon_H \otimes M$. On the other hand, the object $H \otimes M^{coH}$ is a left *H*-Hopf module with action $\varphi_{H \otimes M^{coH}} =$ $\mu_H \otimes M^{coH}$ and coaction $\rho_{H \otimes M^{coH}} = \delta_H \otimes M^{coH}$. The Fundamental Theorem of Hopf modules asserts that $H \otimes M^{coH}$ is isomorphic to M in the category H-Hopf-Mod. The isomorphism is defined by

$$\theta_M = \varphi_M \circ (H \otimes j_M) : H \otimes M^{coH} \to M$$

where $\theta_M^{-1} = (H \otimes t_M) \circ \rho_M$. In the same way as in the case of M^{coH} , if X is an object in C, the tensor product $H \otimes X$, with the action and coaction induced by the product and the coproduct of H, is a left H-Hopf module. Then, there exists a functor $\mathsf{F} = H \otimes -: \mathsf{C} \to \mathsf{H}\text{-}\mathsf{Hopf}\text{-}\mathsf{Mod}$ called the induction functor. Also, for all $M \in \mathsf{H}\text{-}\mathsf{Hopf}\text{-}\mathsf{Mod}$, the construction of M^{coH} is functorial. Thus,

there exists a new functor $G = ()^{coH} : H-Hopf-Mod \to C$, called the functor of coinvariants, such that $F \dashv G$. Moreover, F and G induce an equivalence between the categories H-Hopf-Mod and C.

Let us see now the definition of left Hopf module over a Hopf truss \mathbb{H} .

Definition 5.5. Let $\mathbb{H} = (H_1, H_2, \sigma_H)$ be a Hopf truss. A left Hopf module over \mathbb{H} (left \mathbb{H} -Hopf module) is a 4-tuple $(M, \psi_M^1, \psi_M^2, \rho_M)$ such that:

- (i) The triple (M, ψ_M^1, ψ_M^2) is a left \mathbb{H} -module.
- (ii) The triple (M, ψ_M^1, ρ_M) is a left H_1 -Hopf module.
- (iii) The triple (M, ψ_M^2, ρ_M) is a non-unital left H_2 -Hopf module.
- (iv) If j_M is the equalizer morphism of ρ_M and $\eta_H \otimes M$, the identity

$$\psi_M^1 \circ (\sigma_H \otimes j_M) = \psi_M^2 \circ (H \otimes j_M)$$

holds.

A morphism of left Hopf modules over \mathbb{H} is a morphism of left \mathbb{H} -modules and left H-comodules.

Left Hopf modules over \mathbb{H} with morphisms of left Hopf modules form a category which we denote by \mathbb{H} -Hopf-Mod. Note that, this definition is a generalization to the Hopf truss setting of the notion of Hopf module over a Hopf brace introduced in [5].

Example 5.6. Let X be an object in C and let $\mathbb{H} = (H_1, H_2, \sigma_H)$ be a Hopf truss. Then, the 4-tuple

$$(H \otimes X, \psi^1_{H \otimes X} = \mu^1_H \otimes X, \psi^2_{H \otimes X} = \mu^2_H \otimes X, \rho_{H \otimes X} = \delta_H \otimes X)$$

is a left \mathbb{H} -Hopf module. Indeed, by Example 4.2, the triple $(H \otimes X, \psi^1_{H \otimes X}, \psi^2_{H \otimes X})$ is a left \mathbb{H} -module. Moreover, $(H \otimes X, \psi^1_{H \otimes X}, \rho_{H \otimes X})$ is a left H_1 -Hopf module and $(H \otimes X, \psi^2_{H \otimes X}, \rho_{H \otimes X})$ is a non-unital left H_2 -Hopf module. Finally,

$$X \xrightarrow{\eta_H \otimes X} H \otimes X \xrightarrow{\delta_H \otimes X} H \otimes H \otimes X$$

is an equalizer diagram and this implies that $(H \otimes X)^{coH_1} = X$ and $j_{H \otimes X} = \eta_H \otimes X$. Then, (iv) of Definition 5.5 holds, because by (12), we have that $\varphi_{H \otimes X} \circ (\sigma_H \otimes j_{H \otimes X}) = \sigma_H \otimes X = \psi_{H \otimes X} \circ (H \otimes j_{H \otimes X})$. Note that, for X = K, we obtain that

$$(H, \mu_H^1, \mu_H^2, \delta_H)$$

is an object in \mathbb{H} -Hopf-Mod where $H^{coH_1} = K$.

On the other hand, $q_{H\otimes X} = \varepsilon_H \otimes \eta_H \otimes X$ and, as a consequence, $I(q_{H\otimes X}) = X$, $p_{H\otimes X} = \varepsilon_H \otimes X$ and $i_{H\otimes X} = \eta_H \otimes X$.

Finally, it is obvious that, if $f: X \to Y$ is a morphism in $\mathsf{C}, H \otimes f$ is a morphism in \mathbb{H} -Hopf-Mod.

Then, as a consequence of the facts presented in the previous example, we have the following results:

Theorem 5.7. Let \mathbb{H} be a Hopf truss. There exists a functor $\mathsf{F} = H \otimes -: \mathsf{C} \to \mathbb{H}$ -Hopf, called the induction functor, defined on objects by $\mathsf{F}(X) = (H \otimes X, \psi^1_{H \otimes X}, \psi^2_{H \otimes X}, \rho_{H \otimes X})$ and on morphisms by $\mathsf{F}(f) = H \otimes f$.

Theorem 5.8. (Fundamental Theorem of Hopf modules) Let $\mathbb{H} = (H_1, H_2, \sigma_H)$ be a Hopf truss and let $(M, \varphi_M, \psi_M, \rho_M)$ be an object in \mathbb{H} -Hopf-Mod. Then $(M, \psi_M^1, \psi_M^2, \rho_M)$ and $F(M^{coH_1})$ are isomorphic in \mathbb{H} -Hopf-Mod. *Proof.* Let $(M, \psi_M^1, \psi_M^2, \rho_M)$ be an object in \mathbb{H} -Hopf. By Theorem 5.7, the 4-tuple

$$(H \otimes M^{coH_1}, \psi^1_{H \otimes M^{coH_1}}, \psi^2_{H \otimes M^{coH_1}}, \rho_{H \otimes M^{coH_1}})$$

is a left \mathbb{H} -Hopf module. Let $\theta_M : H \otimes M^{coH_1} \to M$ be the morphism defined by $\theta_M = \psi_M^1 \circ (H \otimes j_M)$. Then, by the general theory of Hopf modules exposed at the beginning of this section, θ_M is an isomorphism of H_1 -Hopf modules with inverse $\theta_M^{-1} = (H \otimes t_M) \circ \rho_M$. Also, θ_M is an isomorphism of non-unital H_2 -Hopf modules because

$$\begin{split} &\psi_M^2 \circ (H \otimes \theta_M) \\ &= \psi_M^1 \circ (\mu_H^2 \otimes \Gamma_M^{\sigma_H}) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes j_M) \text{ (by 20)} \\ &= \psi_M^1 \circ (\mu_H^2 \otimes (\psi_M^1 \circ (H \otimes \psi_M^1) \circ ((((\lambda_H \circ \sigma_H) \otimes \sigma_H) \circ \delta_H) \otimes M))) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes j_M) \\ &\text{(by (v) of definition 5.5)} \\ &= \psi_M^1 \circ (\mu_H^2 \otimes (\psi_M^1 \circ ((\lambda_H * id_H) \circ \sigma_H) \otimes M)) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes H \otimes j_M) \text{ (by the condition of comonoid morphism for } \sigma_H \text{ and the condition of left } H_1\text{-module for } M) \\ &= \psi_M^1 \circ (\mu_H^2 \otimes j_M) \text{ (by (8), the naturality of } c, \text{ the counit properties and the condition of left } H_1\text{-module for } M) \\ &= \theta_M \circ \psi_{H \otimes M^{coH_1}}^2 \text{ (by definition.)} \end{split}$$

Therefore, θ_M is an isomorphism in \mathbb{H} -Hopf-Mod because, by (ii) of Definition 5.5, the properties of j_M , the naturality of c and the properties of η_H , we have that θ_M is a morphism of left H-comodules.

In Theorem 5.7 we construct the induction functor $\mathsf{F} = H \otimes -: \mathsf{C} \to \mathbb{H}$ -Hopf-Mod for a Hopf truss $\mathbb{H} = (H_1, H_2, \sigma_H)$. As in the Hopf case, for all $M \in \mathbb{H}$ -Hopf-Mod, the construction of the subobject of coinvariants $M^{co\mathbb{H}} := M^{coH_1}$ is functorial. Thus, there exists a functor of coinvariants $\mathsf{W} = ()^{co\mathbb{H}} : \mathbb{H}$ -Hopf-Mod $\to \mathsf{C}$ defined on objects by $= \mathsf{W}((M, \psi_M^1, \psi_M^2, \rho_M)) = M^{co\mathbb{H}}$ where $M^{co\mathbb{H}} = M^{coH_1}$ and on morphisms $f: M \to N$ by $\mathsf{W}(f) = f^{co\mathbb{H}}$, where $f^{co\mathbb{H}} := f^{coH_1}$ is the unique morphism such that $j_N \circ f^{coH_1} = f \circ j_M$.

In the end of this section we prove that there exists a categorical equivalence between C and the category \mathbb{H} -Hopf for a Hopf truss \mathbb{H} .

Theorem 5.9. Let \mathbb{H} a Hopf truss. The induction functor $\mathsf{F} = H \otimes -: \mathsf{C} \to \mathbb{H}$ -Hopf-Mod is left adjoint of the functor of coinvariants $\mathsf{W} = (\)^{co\mathbb{H}} : \mathbb{H}$ -Hopf-Mod $\to \mathsf{C}$ and they induce a categorical equivalence between \mathbb{H} -Hopf and C .

Proof. Let X be an object in C. Then the unit of the adjunction is $\alpha_X = id_X$ because as we proved in Example 5.6, W(F(X)) = X. For all $(M, \psi_M^1, \psi_M^2, \rho_M)$ in \mathbb{H} -Hopf-Mod, the counit is defined by $\beta_M = \theta_M$ where θ_M is the isomorphism introduced in the proof of Theorem 5.8. The triangular identities hold, because they hold for the adjunction between the categories C and H_1 -Hopf-Mod. \Box

If we particularize the previous theorems to the case of Hopf braces we have the Fundamental Theorem of Hopf Modules and the associated categorical equivalence obtained in [5].

Funding

The authors were supported by Ministerio de Ciencia e Innovación of Spain. Agencia Estatal de Investigación. Unión Europea - Fondo Europeo de Desarrollo Regional (FEDER). Grant PID2020-115155GB-I00: Homología, homotopía e invariantes categóricos en grupos y álgebras no asociativas.

References

- [1] Abe E.: Hopf algebras, University Press, Cambridge, 1980.
- [2] Angiono I., Galindo C., Vendramin L.: Hopf braces and Yang-Baxter operators, Proc. Am. Math. Soc. 145, 1981-1995 (2017).

- [3] Brzezńiski T.: Trusses: between braces and rings, Trans. Am. Math. Soc. 372, 4149-4176 (2019).
- [4] Fernández Vilaboa J.M., González Rodríguez R., Ramos Pérez, B., Rodríguez Raposo A.B., Modules for invertible 1-cocycles, arXiv:2311.05233 [math.RA] (2023).
- [5] González Rodríguez R.: The fundamental theorem of Hopf modules for Hopf braces, Linear Multilinear Algebra, 70, 5146-5156 (2022).
- [6] Guarneri L., Vendramin L.: Skew braces and the Yang-Baxter equation, Math. Comput. 86, 2519-2534 (2017).
- [7] Larson R.G., Sweedler M. E., An associative orthogonal bilinear form for Hopf algebras, Amer. J. Math. 91 (1969), 75-93.
- [8] Kassel C.: Quantum Groups, Quantum groups. Graduate Texts in Mathematics, 155, Springer-Verlag, New York, 1995.
- [9] Mac Lane S.: Categories for the working mathematician. Second edition. Graduate Texts in Mathematics, 5, Springer-Verlag, New York, 1998.
- [10] Sweedler M.E., Hopf algebras, Benjamin, New York (1969).