Quantum to Classical Cavity Chemistry Electrodynamics

Leonardo F. Calderón,*^{,†,‡} Humberto Triviño,[†] and Leonardo A. Pachón*^{,†,¶}

† Grupo de Física Teórica y Matemática Aplicada, Instituto de Física, Facultad de Ciencias Exactas y Naturales, Universidad de Antioquia; Calle 70 No. 52-21, Medellín, Colombia.
‡ Grupo de Física Computacional en Materia Condensada, Escuela de Física, Facultad de Ciencias, Universidad Industrial de Santander UIS; Cra 27 Calle 9 Ciudad Universitaria, Bucaramanga, Colombia.


¶ Grupo de Física Atómica y Molecular, Instituto de Física, Facultad de Ciencias Exactas y Naturales, Universidad de Antioquia UdeA; Calle 70 No. 52-21, Medellín, Colombia.

E-mail: leonardo.calderon@utoronto.ca; leonardo.pachon@udea.edu.co

Abstract

Polaritonic chemistry has ushered in new avenues for controlling molecular dynamics. However, two key questions remain: (i) Can classical light sources elicit the same effects as certain quantum light sources on molecular systems? (ii) Can semiclassical treatments of light-matter interaction capture nontrivial quantum effects observed in molecular dynamics? This work presents a quantum-classical approach addressing issues of realizing cavity chemistry effects without actual cavities. It also highlights the limitations of the standard semiclassical light-matter interaction. It is demonstrated that classical light sources can mimic quantum effects up to the second order of lightmatter interaction, provided that the mean-field contribution, symmetrized two-time correlation function, and the linear response function are the same in both situations. Numerical simulations show that the quantum-classical method aligns more closely with exact quantum molecular-only dynamics for quantum light states such as Fock states, superpositions of Fock states, and vacuum squeezed states than the conventional semiclassical approach.

TOC Graphic



Introduction.—Quantum cavity chemistry explores the potential to transform the chemical landscape, paving the way for novel photophysical phenomena.^{1–12} This transformation is achieved by coupling molecular systems to the confined photonic degrees of freedom within microcavities.¹³ As a result, it offers a promising avenue for manipulating room-temperature photophysical processes in organic molecules. Instances of these novel phenomena have been examined in the context of singlet fission processes,¹⁴ triplet harvesting,¹⁵ energy transfer,^{16–20} remote regulation of chemical reactions,²¹ and nonadiabatic effects.^{22–24}

In classical control theory,²⁵ properties of light, such as intensity and phase, are employed to alter the inherent dynamics of molecular reactions using classical laser-control schemes. In contrast, quantum cavity chemistry leverages the quantum nature of light as a vital tool for reshaping the chemical behavior of molecular reactions. However, despite the nonadiabatic effects observed in the adiabatic ground state populations, evidence from the NaI molecule^{23,26} suggests minimal discernible differences between dynamics driven by classical laser radiation and those induced by cavity interactions. This observation has piqued interest in exploring how classical radiation fields might replicate the molecular dynamics typically attributed to quantum light states.²⁷

Traditionally, in the study of light-matter interactions, ^{28,29} semiclassical methods have involved a full quantum treatment of the molecular degrees of freedom, while adopting various classical dynamics schemes for the photonic degrees of freedom. ^{30–38} However, recent advances employing mixed quantum-classical dynamics techniques ^{39,40} to solve the Ehrenfest mean-field dynamics have made strides. These techniques have demonstrated an ability to account, at least partially, for phenomena such as spontaneous emission, interference, strong coupling, and correlated light-matter dynamics. ^{30–38}

Utilizing a mixed representation of molecule operators and radiation phase-space distributions^{41–43}—and by tracing out the radiation degrees of freedom—an effective master equation exclusive to the molecular degrees of freedom was derived. To achieve the closeness condition for this master equation, contributions were considered up to the second order in light-matter interaction. This closed master equation facilitates a direct juxtaposition of the full quantum dynamics, the conventional semiclassical light-matter interaction, and an alternative semiclassical description that is non-local in time. It should be noted that the primary focus of this paper is not on demonstrating the application of trajectory methods.^{31,33,34} Instead, the emphasis is on elucidating the impact of photonic degrees of freedom on molecular-only dynamics, framed in the context of radiation field statistics.

In a nutshell, up to the second order in light-matter interaction, quantum effects can be mimicked by classical light sources if the mean-field contribution, the symmetrized two-time correlation function, and the linear response function (defined through the antisymmetrized two-time correlation function) are consistent across both scenarios. This condition does not pertain to the classical limit of the quantum light source. Instead, it highlights the potential to design classical light sources that mimic the radiation's effect on the molecular degrees of freedom up to the second order.

In this context, the impact of quantum fluctuations from three distinct quantum light field states—a Fock state, a Fock state superposition, and a squeezed vacuum state—is analyzed using system models. These states possess a non-trivial quantum character^{44–46} and facilitate the exploration of the influence of both vanishing and non-vanishing mean-field contributions and the symmetrized/antisymmetrized two-time correlation functions on molecular-only dynamics.

This letter is structured as follows. First, a description of a molecular system coupled to a quantized light field is provided, accompanied by the formulation of a second-order quantum master equation that treats the light degrees of freedom quantum mechanically. Then, a quantum-classical projection operator formalism is introduced, from which a quantumclassical second-order master equation is derived for the molecular system, considering the light degrees of freedom classically. Lastly, numerical results that juxtapose the quantumclassical, standard semiclassical, and precise quantum dynamics of Hamiltonians employed in theoretical molecular polaritonics are presented, considering various light field states with nonclassical characteristics.

Quantum cavity molecular-only dynamics.—Consider a molecular system that interacts with a quantized light field. Assuming the dipole approximation is valid, the interaction is framed in the length gauge.^{3,47} In this configuration (Pauli-Fierz model), the correlated electron-nuclear-photon Hamiltonian, which comprises $n_{\rm el}$ electrons, $n_{\rm nul}$ nuclei, and $n_{\rm ph}$ photon modes, can be expressed as a sum of the electro-nuclear (or bare molecular) Hamiltonian, denoted as $\hat{H}_{\rm bm}$, and the photon Hamiltonian, represented as $\hat{H}_{\rm ph}$. The bare molecular Hamiltonian, $\hat{H}_{\rm bm}$, includes five components: the kinetic energy of the electrons $\hat{T}_{\rm el}$, the kinetic energy of the nucleons $\hat{T}_{\rm nuc}$, the interaction between electrons $\hat{V}_{\rm el-el}$, between nucleons $\hat{V}_{\rm nuc-nuc}$ and the electron-nucleons interaction $\hat{V}_{\rm el-nuc}$, it reads $\hat{H}_{\rm bm} =$ $\hat{T}_{\rm el} + \hat{T}_{\rm nuc} + \hat{V}_{\rm el-el} + \hat{V}_{\rm nuc-nuc} + \hat{V}_{\rm el-nuc}$. The photon Hamiltonian incorporates the coupling to the molecular degrees of freedom and reads

$$\hat{H}_{\rm ph} = \frac{1}{2} \sum_{\alpha=1}^{n_{\rm ph}} \left[\hat{p}_{\alpha}^2 + \omega_{\alpha}^2 (\hat{q}_{\alpha} + e \,\boldsymbol{\lambda}_{\alpha} \cdot \hat{\mathbf{R}} / \omega_{\alpha})^2 \right]$$

$$= \hat{H}_{\rm bph} + \hat{H}_{\rm ph-m} + \hat{H}_{\rm rm},$$
(1)

with $\hat{\mathbf{R}} = \sum_{I=1}^{n_{\text{nucc}}} Z_I \hat{\mathbf{X}}_I - \sum_{i=1}^{n_{\text{el}}} \hat{\mathbf{x}}_i$ being the total dipole operator including both sets of electronic $\{\hat{\mathbf{x}}_i\}$ and nuclear $\{\hat{\mathbf{X}}_I\}$ coordinates and Z_I stand for the nuclear charges. $\hat{H}_{\text{bph}} = \frac{1}{2} \sum_{\alpha=1}^{n_{\text{ph}}} (\hat{\mathbf{p}}_{\alpha}^2 + \omega_{\alpha}^2 \hat{\mathbf{q}}_{\alpha}^2)$ denotes the bare photon Hamiltonian, $\hat{H}_{\text{ph-m}} = \sum_{\alpha=1}^{n_{\text{ph}}} \omega_{\alpha} \hat{\mathbf{q}}_{\alpha} \boldsymbol{\lambda}_{\alpha} \cdot e \hat{\mathbf{R}}$ does so for the interaction between photons and molecular degrees of freedom, with $\boldsymbol{\lambda}$ as the dipole coupling strength. Finally, $\hat{H}_{\text{rm}} = \frac{1}{2} \sum_{\alpha=1}^{n_{\text{ph}}} (\boldsymbol{\lambda}_{\alpha} \cdot e \hat{\mathbf{R}})^2$ accounts for the renormalization of the bare molecular Hamiltonian due to the coupling to light and corresponds to the dipole self-energy.

Another more conventional formulation for studying molecule-cavity physics corresponds to the multipolar macroscopic model of QED under the Power-Zineau-Woolley (PZW) unitary transformation from the minimal coupling.^{48–50} Macroscopic QED allows for quantizing the electromagnetic field in any geometry, considering realistic cavities with lossy materials.^{51–53} Light-matter interactions are addressed directly through the electric **E** and magnetic **B** fields, and the field-emitter interactions are incorporated by multipole moments.^{51,52} In multipolar QED, the dipole self-energy contribution is contained in the molecular Hamiltonian and does not directly depend on the vacuum field.^{11,52,54,55} In addition, for realistic cavities where the interaction is driven mainly by longitudinal fields (e.g., plasmonic nanocavities), the dipole self-energy does not appear.^{52,56}

To provide a general insight into the quantum aspects of light in molecular systems, consider the molecular system-light field Hamiltonian

$$\hat{H} = \hat{H}_{\rm m} + \hat{H}_{\rm ph-m} + \hat{H}_{\rm bph},\tag{2}$$

with the molecular Hamiltonian $\hat{H}_{\rm m} = \hat{H}_{\rm bm} + \hat{H}_{\rm rm}$. The dynamics of the state $\hat{\rho}$ of the entire system described by \hat{H} follows from the von Neumann equation, $\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho} = -\frac{\mathrm{i}}{\hbar}[\hat{H},\hat{\rho}]$. Because interest here is in the influence that coupling to photons induces on the molecular systems, the partial trace over the states of the photons is applied to $\hat{\rho}$, so that the reduced state of the molecular systems follows from $\hat{\rho}_{\rm m} = \mathrm{tr}_{\rm ph}\hat{\rho}$. To focus exclusively on the dynamics induced by the light-matter coupling, it is convenient to calculate the equation of motion of $\hat{\rho}_{\rm m}$ in the interaction picture by introducing the unitary transformation $\hat{\rho}^{(\mathrm{I})}(t) = \hat{U}^{\dagger}(t-t_0)\hat{\rho}(t_0)\hat{U}(t-t_0)$ with $\hat{U}(t-t_0) = \exp\left[\frac{\mathrm{i}}{\hbar}\hat{H}_{\rm m}(t-t_0)\right] \exp\left[\frac{\mathrm{i}}{\hbar}\hat{H}_{\rm bph}(t-t_0)\right]$. Therefore,

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t) = -\frac{\mathrm{i}}{\hbar}\mathrm{tr}_{\mathrm{ph}}\left[\hat{H}_{\mathrm{ph-m}}^{(\mathrm{I})}(t),\hat{\rho}^{(\mathrm{I})}(t)\right].$$
(3)

By continuing the discussion in the interaction picture, the sheer complexity of the intrinsic molecular dynamics is left out, and the light-matter interaction becomes the focus of the description below.

Quantum-classical cavity molecular-only dynamics.—Once the light fields have been traced out, the core proposition of this work emerges: if the reduced equation of motion for $\hat{\rho}_{\rm m}$ remains consistent under both quantum and classical light, then it becomes feasible to *effectively* simulate the molecular dynamics triggered by quantum light fields using classical counterparts. To further explore the capability of classical fields to replicate the dynamics stimulated by quantum fields, the radiation degrees of freedom will be translated into phase-space distributions in the subsequent sections.

To achieve this, an initial step involves executing a partial Wigner-Weyl transform ^{57,58} on the light field's Hilbert space. This process provides a phase-space depiction of the light field's degrees of freedom. The partial Wigner-Weyl representation for the state $\hat{\rho}$ is given as follows: $\hat{\rho}^{W}(\mathbf{q}, \mathbf{p}) = \frac{1}{(2\pi\hbar)^{n_{\mathrm{ph}}}} \int d\mathbf{u} \langle \mathbf{q} + \frac{\mathbf{u}}{2} | \hat{\rho} | \mathbf{q} - \frac{\mathbf{u}}{2} \rangle e^{-\frac{i}{\hbar} \mathbf{p} \cdot \mathbf{u}}$, where the integration is performed over the entire phase-space of light field degrees of freedom $(\mathbf{q}, \mathbf{p}) = (\{\mathbf{q}_{\alpha}\}, \{\mathbf{p}_{\alpha}\})$. The state $\hat{\rho}^{W}$ retains its operator nature since the molecular degrees of freedom still represent quantum operators in the molecule Hilbert space \mathcal{H}_{m} . After performing the Wigner-Weyl transform, the molecule-light field dynamics is governed by the von-Neumann-Moyal equation ^{59,60}

$$\frac{\partial \hat{\rho}^{\mathrm{W}}}{\partial t} = \left\{ \hat{H}^{\mathrm{W}}, \hat{\rho}^{\mathrm{W}} \right\}_{\mathrm{M}} = -\frac{\mathrm{i}}{\hbar} (\hat{H}^{\mathrm{W}} \mathrm{e}^{\frac{\mathrm{i}\hbar}{2}\Lambda} \hat{\rho}^{\mathrm{W}} - \hat{\rho}^{\mathrm{W}} \mathrm{e}^{\frac{\mathrm{i}\hbar}{2}\Lambda} \hat{H}^{\mathrm{W}}), \tag{4}$$

where $\{\cdot, \cdot\}_{\mathrm{M}}$ symbolizes the Moyal bracket, with $\mathrm{e}^{\frac{\mathrm{i}\hbar}{2}\Lambda} = \lim_{N\to\infty} \sum_{l=0}^{N} \frac{1}{l!} (\frac{\mathrm{i}\hbar}{2}\Lambda)^{l}$, and $\Lambda = \sum_{\alpha=1}^{n_{\mathrm{ph}}} \frac{\overleftarrow{\partial}}{\partial \mathbf{q}_{\alpha}} \frac{\overrightarrow{\partial}}{\partial \mathbf{p}_{\alpha}} - \frac{\overleftarrow{\partial}}{\partial \mathbf{p}_{\alpha}} \frac{\overrightarrow{\partial}}{\partial \mathbf{q}_{\alpha}}$. The arrows indicate the direction in which a derivative acts. In this representation there is no approximation invoked.

The classical limit for the light degrees of freedom of Eq. (4) corresponds to the von-Neumann–Poisson equation that is reached by disregarding $\mathcal{O}(\hbar^2)$ terms in the expansion of the phase-space operator $e^{\frac{i\hbar}{2}\Lambda}$. Thus, the quantum-classical dynamics^{41,42} of the state $\hat{\rho}^{W}$ reads

$$\frac{\partial \hat{\rho}^{\mathrm{W}}}{\partial t} = -\mathrm{i}\mathcal{L}^{\mathrm{W}}\hat{\rho}^{\mathrm{W}} = -\frac{\mathrm{i}}{\hbar} \left[\hat{H}^{\mathrm{W}}, \hat{\rho}^{\mathrm{W}} \right] + \frac{1}{2} \left(\left\{ \hat{H}^{\mathrm{W}}, \hat{\rho}^{\mathrm{W}} \right\}_{\mathrm{P}} - \left\{ \hat{\rho}^{\mathrm{W}}, \hat{H}^{\mathrm{W}} \right\}_{\mathrm{P}} \right), \tag{5}$$

where $\{\cdot, \cdot\}_P$ denotes the classical Poisson bracket. It is essential to point out that the dynamics generated by the quantum-classical Liouvillian superoperator \mathcal{L}_W coincides with the Moyal bracket's quantum dynamics as the photon Hamiltonian is quadratic in its quadratures.

In this partial Wigner representation, the Hamiltonian of the Eq. (2) reads

$$\hat{H}^{W} = \hat{H}_{m} + \hat{H}^{W}_{ph-m} + H^{W}_{ph}.$$
 (6)

The interaction between the light field and the molecular degrees of freedom is defined by the Hamiltonian of the general form $\hat{H}_{ph-m}^{W} = \sum_{u} \hat{K}_{u} \Phi_{u}^{W}$, where \hat{K}_{u} represents an operator in the Hilbert space of the molecular system, and Φ_{u}^{W} the phase-space representation of operator $\hat{\Phi}_{u}$ of the light field. Note that the photon Hamiltonian H_{ph}^{W} is not an operator anymore.

The partial trace over the light degrees of freedom is obtained as $\hat{\rho}_{\rm m} = {\rm tr}_{\rm ph} \hat{\rho}^{\rm W} = \int d\mathbf{q} d\mathbf{p} \hat{\rho}^{\rm W}$. After replacing the Hamiltonian of the Eq. (6) in the Eq. (5), and considering that $[H_{\rm ph}^{\rm W}, \hat{\rho}^{\rm W}] = \{\hat{H}_{\rm m}, \hat{\rho}^{\rm W}\}_{\rm P} = 0$, the evolution of the reduced state of the molecular system reads

$$\frac{\partial \hat{\rho}_{\rm m}}{\partial t} = -\frac{\mathrm{i}}{\hbar} \left[\hat{H}_{\rm m}, \hat{\rho}_{\rm m} \right] - \frac{\mathrm{i}}{\hbar} \sum_{u} \left[\hat{K}_{u}, \operatorname{tr}_{\rm ph} \left(\Phi_{u}^{\rm W} \hat{\rho}^{\rm W} \right) \right]
+ \frac{1}{2} \sum_{u} \left[\hat{K}_{u}, \operatorname{tr}_{\rm ph} \left\{ \Phi_{u}^{\rm W}, \hat{\rho}^{\rm W} \right\}_{\rm P} \right]_{+} + \operatorname{tr}_{\rm ph} \left\{ H_{\rm ph}^{\rm W}, \hat{\rho}^{\rm W} \right\}_{\rm P}.$$
(7)

The full influence of the quantum/classical light field into the molecular dynamics is encoded in Φ_u^W provided that no approximation is involved in Eq. (7). Therefore, deviations from the quantum description can be discussed in terms of the quantum/classical correlations of Φ_u^W .

Derivation of the Reduced Master Equation in the Mixed Wigner Representation.—Since the dynamics of Φ_u^W , at first-order, depend on the interaction with the molecule itself, Eq. (7) is not closed from a mathematical viewpoint. This can be overcome after introducing a projection operator technique in the quantum-classical Hilbert–phase-space of the entire molecular-light field system (see section S1 in the Supporting Information for an equivalent treatment in the standard full Hilbert space description). For simplicity, consider the secondorder regime in the molecule-light interaction through the Liouvillian superoperators in the interaction picture of the partial Wigner transform $\hat{\rho}^{W(I)}(t) = e^{i\mathcal{L}_0^W t} \hat{\rho}^W(t)$, with $\mathcal{L}_0^W = \mathcal{L}_m + \mathcal{L}_{ph}^W$ Therefore, the quantum-classical Liouville equation in the interaction picture reads

$$\frac{\partial}{\partial t}\hat{\rho}^{\mathrm{W}(\mathrm{I})}(t) = -\mathrm{i}\mathcal{L}_{\mathrm{ph-m}}^{\mathrm{W}(\mathrm{I})}(t)\hat{\rho}^{\mathrm{W}(\mathrm{I})}(t),\tag{8}$$

where $\mathcal{L}_{\rm ph-m}^{\rm W(I)}(t) = e^{i\mathcal{L}_0^{\rm Wt}}\mathcal{L}_{\rm ph-m}^{\rm W}$. The factorized interaction Hamiltonian in the interaction representation assumes the time-dependent form $\hat{H}_{\rm ph-m}^{\rm W(I)}(t) = \sum_u \hat{K}_u^{(I)}(t) \Phi_{uW}^{(I)}(t)$. Thus, the interaction Liouvillian in the interaction representation reads $\mathcal{L}_{\rm ph-m}^{\rm W(I)}(t) \bullet = \frac{1}{\hbar} [\hat{H}_{\rm ph-m}^{\rm W(I)}(t), \bullet] + \frac{i}{2} (\{\hat{H}_{\rm ph-m}^{\rm W(I)}(t), \bullet\}_{\rm P} - \{\bullet, \hat{H}_{\rm ph-m}^{\rm W(I)}(t)\}_{\rm P})$. Using the concept of partial trace over the phase-space light-field-degrees-of-freedom, and since $\mathrm{tr}_{\rm ph}\rho_{\rm ph}^{\rm W(I)}(t_0) = \int \mathrm{d}\mathbf{q}\mathrm{d}\mathbf{p}\rho_{\rm ph}^{\rm W(I)}(t_0) = 1$, it is then possible to define the quantum-classical projection superoperator $\mathcal{P}\hat{\rho}^{\rm W(I)}(t) = \rho_{\rm ph}^{\rm W(I)}(t_0)\mathrm{tr}_{\rm ph}\hat{\rho}^{\rm W(I)}(t)$, with orthogonal complement $\mathcal{Q} = \hat{1} - \mathcal{P}$. The equations of motion for both subspaces read

$$\operatorname{tr}_{\mathrm{ph}}\left(\mathcal{P}\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}^{\mathrm{W}(\mathrm{I})}(t)\right) = -\operatorname{itr}_{\mathrm{ph}}\left(\mathcal{P}\mathcal{L}_{\mathrm{ph-m}}^{\mathrm{W}(\mathrm{I})}(t)\rho_{\mathrm{ph}}^{\mathrm{W}(\mathrm{I})}(t_{0})\times\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t) + \mathcal{P}\mathcal{L}_{\mathrm{ph-m}}^{\mathrm{W}(\mathrm{I})}(t)\mathcal{Q}\hat{\rho}^{\mathrm{W}(\mathrm{I})}(t)\right), \quad (9)$$

$$\mathcal{Q}\frac{\mathrm{d}\hat{\rho}^{\mathrm{W}(\mathrm{I})}(t)}{\mathrm{d}t} = -\mathrm{i}\Big(\mathcal{Q}\mathcal{L}_{\mathrm{ph-m}}^{\mathrm{W}(\mathrm{I})}(t)\rho_{\mathrm{ph}}^{\mathrm{W}(\mathrm{I})}(t_0)\hat{\rho}_{\mathrm{m}}^{\mathrm{(I)}}(t) + \mathcal{Q}\mathcal{L}_{\mathrm{ph-m}}^{\mathrm{W}(\mathrm{I})}(t)\mathcal{Q}\hat{\rho}^{\mathrm{W}(\mathrm{I})}(t)\Big).$$
(10)

To second-order, the term $\mathcal{QL}_{ph-m}^{W(I)}(t)\mathcal{Q}\hat{\rho}^{W(I)}(t)$ is disregarded. The mixed-Wigner-representation, Eqs. (9) and 10) are analog to Eqs. (S1) and (S2) in the Supporting Information, respectively.

Considering the factorized initial condition $\mathcal{Q}\hat{\rho}^{W(I)}(t_0) = 0$, the solution of Eq. (10) reads $\mathcal{Q}\hat{\rho}^{W(I)}(t) = -i \int_{t_0}^t dt' \mathcal{QL}_{ph-m}^{W(I)}(t') \rho_{ph}^{W(I)}(t_0) \hat{\rho}_m^{(I)}(t')$. After replacing this result in Eq. (9), the first and second-order reduced master equations for the molecular system read

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})(1)}(t) = -\frac{\mathrm{i}}{\hbar}\sum_{u} \left\langle \Phi_{u}^{\mathrm{W}(\mathrm{I})}(t) \right\rangle \left[\hat{K}_{u}^{(\mathrm{I})}(t), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t)\right],\tag{11}$$

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})(2)}(t) = -\frac{\mathrm{i}}{\hbar}\sum_{u} \left\langle \Phi_{u}^{\mathrm{W}(\mathrm{I})}(t) \right\rangle \left[\hat{K}_{u}^{(\mathrm{I})}(t), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t)\right]
- \frac{1}{\hbar^{2}} \int_{t_{0}}^{t} \mathrm{d}t' \sum_{u,v} \left(\mathcal{C}_{u,v}^{\mathrm{W}}(t,t') - \left\langle \Phi_{u}^{\mathrm{W}(\mathrm{I})}(t) \right\rangle \left\langle \Phi_{v}^{\mathrm{W}(\mathrm{I})}(t') \right\rangle \right) \left[\hat{K}_{u}^{(\mathrm{I})}(t), \left[\hat{K}_{v}^{(\mathrm{I})}(t'), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t')\right] \right]
- \frac{\mathrm{i}}{2\hbar} \int_{t_{0}}^{t} \mathrm{d}t' \sum_{u,v} \chi_{u,v}^{\mathrm{W}}(t,t') \left[\hat{K}_{u}^{(\mathrm{I})}(t), \left[\hat{K}_{v}^{(\mathrm{I})}(t'), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t')\right]_{+}\right],$$
(12)

where

$$\mathcal{C}_{u,v}^{\mathrm{W}}(t,t') = \frac{1}{2} \Big(\left\langle \Phi_{u}^{\mathrm{W}(\mathrm{I})}(t) \Phi_{v}^{\mathrm{W}(\mathrm{I})}(t') \right\rangle + \left\langle \Phi_{v}^{\mathrm{W}(\mathrm{I})}(t') \Phi_{u}^{\mathrm{W}(\mathrm{I})}(t) \right\rangle \Big), \tag{13}$$

$$\chi_{u,v}^{\mathrm{W}}(t,t') = \left\langle \left\{ \Phi_{u}^{\mathrm{W}(\mathrm{I})}(t), \Phi_{v}^{\mathrm{W}(\mathrm{I})}(t') \right\}_{\mathrm{P}} \right\rangle, \tag{14}$$

represent the phase-space version of real (symmetrized) part of the two-time correlation function g(t,t') and the linear response function^{61,62} of the light field operator that couples to the molecular system. $\langle \Phi_u^{W(I)}(t) \rangle = tr_{ph} [\Phi_u^{W(I)}(t) \rho_{ph}^{W(I)}(t_0)]$ represents the expected value of the light field observable that couples to the molecular system.

The reduced master equation in Eq. (12) admits analyzing the influence of classical or quantum descriptions of light fields on the same foot-stage. The correlations $C_{u,v}^{W}(t,t')$ and $\chi_{u,v}^{W}(t,t')$, derived in the quantum-classical projection operator approach, correspond to the classical limit of the quantities $C_{u,v}(t,t')$ and $\chi_{u,v}(t,t')$ in the Eq. (S5) of the Supporting Information, respectively. The analogy can be understood in terms of the light field two-time correlation function given by

$$g_{u,v}^{W}(t,t') = \left\langle \Phi_{u}^{W(I)}(t)\Phi_{v}^{W(I)}(t') \right\rangle = \int dqdp \,\Phi_{u}^{W}(q,p;t)\Phi_{v}^{W}(q,p;t')\rho_{ph}^{W(I)}(q,p;t_{0}).$$
(15)

In the classical limit, the symmetrized correlation function coincides with the two-time correlation function $C_{u,v}^{W}(t,t') = g_{u,v}^{W}(t,t')$. The antisymmetrized correlation function is defined through the phase-space Poisson bracket $\mathcal{A}_{u,v}^{W}(t,t') = \frac{1}{2i} \left\{ \Phi_{u}^{W(I)}(t), \Phi_{v}^{W(I)}(t') \right\}_{P}$. It is useful to

express the symmetrized correlation function in the form given in Eq. (13) for the comparison with the quantum case discussed below. The quantum limit of the Poisson bracket in the phasespace linear response function corresponds to $\{\Phi_u^{W(I)}(t), \Phi_v^{W(I)}(t')\}_P \rightarrow \frac{1}{i\hbar} [\hat{\Phi}_u^{(I)}(t), \hat{\Phi}_v^{(I)}(t')].$

Since the state of the photon field is aimed not to be measured, up to second-order in the light-molecule interaction, it is then not possible to distinguish between the dynamics induced by a classical electric field having the same statistics as the quantum electric field Eq. (12) and the actual dynamics generated by the quantum photon field (see Eq. (S5) of the Supporting Information). However, since the light degrees of freedom evolve independently of the molecular dynamics, Eq. (12) and Eq. (S5) of the Supporting Information do not consider the back-action of the molecular dynamics on the light field.²⁷

This formalism can be extended to (i) generalize the effect of classical vibrational baths composed of harmonic oscillators or to (ii) include higher orders in the light-matter interaction; thus, a quantum-classical master equation of the type Nakajima-Zwanzig or time-convolutionless could be deduced through this approach, allowing to treat strong coupling effects that can lead to deviations from canonical statistics,^{63–68} as recently reported in the modification of ground-state chemical reactivity in infrared cavities.⁶⁹ In addition, if interest is in the dynamics of the light state instead of the molecule state, the role of the system could be inverted above (see Section S2 of the Supporting Information). However, these extensions are beyond the scope of this Letter.

Traditional Semiclassical Approach.—To fully appreciate the difference between the present approach to the standard semiclassical approach and the conundrum posed by the results in the NaI molecule,^{23,26} the standard semiclassical approach is reviewed next. The semiclassical Hamiltonian reads

$$\hat{H}_{\rm sc} = \hat{H}_{\rm m} - \hat{\boldsymbol{\mu}} \cdot \mathbf{E}(t) \tag{16}$$

with $\hat{\mu}$ representing the total molecular dipole operator and the classical electric given by $\mathbf{E}(t) = \mathbf{E}_0 \cos(\omega_c t + \phi)$. The classical field amplitude \mathbf{E}_0 comprises the polarization state which it is assumed parallel to the total molecular dipole operator. The molecular dynamics follow the unitary evolution given by the master equation

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t) = -\frac{\mathrm{i}}{\hbar}\mathbf{E}(t)\cdot\left[-\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t),\hat{\rho}^{(\mathrm{I})}(t)\right].$$
(17)

whose formal solution is given by

$$\hat{\rho}_{\rm m}^{\rm (I)}(t) = \hat{\rho}_{\rm m}^{\rm (I)}(0) - \frac{\mathrm{i}}{\hbar} \int_0^t \mathrm{d}t' \mathbf{E}(t') \cdot \left[-\hat{\boldsymbol{\mu}}^{\rm (I)}(t'), \hat{\rho}^{\rm (I)}(t') \right].$$
(18)

For further analysis, It is convenient to replace Eq. (18) in Eq. (17) to have an alternative equation of motion for the molecule density matrix

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t) = -\frac{\mathrm{i}}{\hbar} \left[-\mathbf{E}(t) \cdot \hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(0) \right] - \frac{1}{\hbar^{2}} \left[-\mathbf{E}(t) \cdot \hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t), \int_{0}^{t} \mathrm{d}t' \left[-\mathbf{E}(t') \cdot \hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t'), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t') \right] \right]$$
(19)

To incorporate the classical statistical properties of light (cf. Chap. 4 in Ref. 44), an ensemble average is performed

$$\frac{\mathrm{d}}{\mathrm{d}t} \left\langle \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t) \right\rangle_{\mathrm{light}} = \frac{\mathrm{i}}{\hbar} \left\langle \mathbf{E}(t) \right\rangle_{\mathrm{light}} \left[\cdot \hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(0) \right]
- \frac{1}{\hbar^{2}} \int_{0}^{t} \mathrm{d}t' \left\langle \mathbf{E}(t) \cdot \mathbf{E}(t') \right\rangle_{\mathrm{light}} \left[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t), \left[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t'), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t') \right] \right].$$
(20)

If $\langle \mathbf{E}(t) \rangle_{\text{light}} = 0$, as in the case of incoherent light or monochromatic plane wave, then

$$\frac{\mathrm{d}}{\mathrm{d}t} \left\langle \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t) \right\rangle_{\mathrm{light}} = -\frac{1}{\hbar^2} \int_0^t \mathrm{d}t' \mathcal{C}^{\mathrm{cl}}(t,t') \left[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t), \left[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t'), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t') \right] \right], \tag{21}$$

with $C^{\text{cl}}(t,t') = \langle \mathbf{E}(t) \cdot \mathbf{E}(t') \rangle_{\text{light}}$. For the sake of comparison, consider only one coupling operator in Eq. (12), identify $\langle \Phi_u^{\text{W}(I)}(t) \rangle = \mathbf{E}(t)$ and $\hat{K}^{(I)}(t) = -\hat{\boldsymbol{\mu}}^{(I)}(t)$ and reduce for the

case $\langle \hat{\Phi}_{u}^{(\mathrm{I})} \rangle = \langle \Phi_{u}^{\mathrm{W}(\mathrm{I})}(t) \rangle = 0$, so that,

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})(2)}(t) = -\frac{1}{\hbar^{2}}\int_{0}^{t}\mathrm{d}t'\mathcal{C}^{\mathrm{W}}(t,t') \Big[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t), \left[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t'), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t')\right]\Big]
- \frac{\mathrm{i}}{2\hbar}\int_{0}^{t}\mathrm{d}t'\chi^{\mathrm{W}}(t,t') \Big[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t), \left[\hat{\boldsymbol{\mu}}^{(\mathrm{I})}(t'), \hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t')\right]_{+}\Big].$$
(22)

The standard semiclassical description of the light-matter interaction in Eq. (17) perfectly replicates the molecular dynamics ignited by light, provided that $C^{W}(t,t') = C^{cl}(t,t')$ and $\chi^{W}(t,t') = 0$. This condition is characteristic of light sources without quantum correlations. Even in scenarios where $\langle \hat{\Phi}_{u}^{(I)} \rangle = 0$, it is anticipated that classical light sources can, to a certain degree, emulate molecular dynamics as long as $C^{W}(t,t') = C^{cl}(t,t')$. It should be noted that corrections arising from purely quantum correlations of light primarily enter via $\chi^{W}(t,t')$.

From Eq.(19), it emerges that for quantum states like the Fock state where $\langle \hat{\Phi} u^{(I)} \rangle = \langle \Phi u^{W(I)}(t) \rangle = 0$, a simulation of the reduced dynamics of the molecule using classical light sources becomes feasible if $\mathcal{C}^{W}(t,t') \approx \mathcal{C}^{cl}(t,t')$. The fidelity of this simulation is contingent upon the light's characteristics, the strength of the coupling, and the inherent complexity of the molecular structure. Notably, the compatibility of the standard semiclassical description in Eq. (17) with the quantum results for the NaI molecule, as documented in Ref. 23, infers the potential accuracy of the effective field description for a variety of system models and coupling strengths.

Results for Model Systems.—The subsequent analysis evaluates the influence of light fields on the reduced molecular dynamics in Eq. (12) across three system models: (i) the Rabi model, (ii) the Dicke model, and (iii) a vibronic dimer model. The introduction of an effective electric field \mathbf{E}_{eff} facilitates a discussion on the results from Eq. (21).

Rabi Model.—The general framework analyzed in this study is exemplified by considering a concrete model system, the Rabi model. This model involves a two-level system interacting

with a cavity and includes the counter-rotating terms, as described by the Hamiltonian

$$\hat{H}_{\rm R} = \frac{1}{2}\hbar\omega_0\hat{\sigma}_z + \hbar g\hat{\sigma}_x(\hat{a}^{\dagger} + \hat{a}) + \hbar\omega_c\hat{a}^{\dagger}\hat{a}.$$
(23)

Following Eq. (22), the classical description of the light field yields the following master equation for the two-level system tracing over the light degrees of freedom

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})(2)}(t) = -\frac{1}{\hbar^{2}}\int_{t_{0}}^{t}\mathrm{d}t'\mathcal{C}^{\mathrm{W}}(t,t')\left[\hat{\sigma}_{x}^{(\mathrm{I})}(t),\left[\hat{\sigma}_{x}^{(\mathrm{I})}(t'),\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t')\right]\right]
-\frac{\mathrm{i}}{2\hbar}\int_{t_{0}}^{t}\mathrm{d}t'\chi^{\mathrm{W}}(t,t')\left[\hat{\sigma}_{x}^{(\mathrm{I})}(t),\left[\hat{\sigma}_{x}^{(\mathrm{I})}(t'),\hat{\rho}_{\mathrm{m}}^{(\mathrm{I})}(t')\right]_{+}\right].$$
(24)

In this context, the quantum equivalent of the phase-space light field observable $\Phi^{W}(t)$ is defined as $\hat{\Phi}^{(I)}(t) = \hbar g(\hat{a}^{\dagger} e^{i\omega_{c}t} + \hat{a} e^{-i\omega_{c}t})$, with the positive(negative)-frequency component expressed by $\hat{\Phi}^{(+)}(t) = \hbar g \hat{a} e^{-i\omega_{c}t}$, $(\hat{\Phi}^{(-)}(t) = \hbar g \hat{a}^{\dagger} e^{i\omega_{c}t})$. This analysis excludes many-modecavity-effects, concentrating solely on one frequency mode ω_{c} . Also, any space dependence of the light-matter coupling strength g is not considered.¹³ Additionally, all Hamiltonians discussed in this section do not include the dipole self-energy term.^{55,70}

The choice for the classical radiation field in the semiclassical approximation corresponds to an electric field whose two-time correlation function reproduces the symmetrized correlation function C(t, t') (see Section S3 of the Supporting Information)

$$E_{\rm cl} = 2\hbar g \sqrt{\langle n_{\rm c} \rangle + \frac{1}{2}} \cos \omega_{\rm c} t, \qquad (25)$$

where $\langle n_c = \hat{a}^{\dagger} \hat{a} \rangle$ represents the average photon number. In the few-photon regime, the semiclassical dynamics driven by the classical electric field of Eq. 25 deviate significantly from the exact quantum dynamics. Therefore, this choice for the electric field displays satisfactory results only in the many-photon regime.

Enhancing the results in the few-photon regime from the semiclassical dynamics necessitates the incorporation of the emission $\langle \hat{\Phi}^{(+)}(t) \hat{\Phi}^{(-)}(t') \rangle = \hbar^2 g^2 \langle \hat{a}^{\dagger} \hat{a} + 1 \rangle \exp(-i\omega_c(t-t'))$ and absorption $\langle \hat{\Phi}^{(-)}(t) \hat{\Phi}^{(+)}(t') \rangle = \hbar^2 g^2 \langle \hat{a}^{\dagger} \hat{a} \rangle \exp(i\omega_c(t-t'))$ processes in the molecular system.^{45,62} The expression for the effective electric field is thus

$$E_{\rm eff} = 2\hbar g \sqrt{\langle n_{\rm c} \rangle + n_{\rm a}} \cos(\omega_{\rm c} t), \qquad (26)$$

where $n_{\rm a} = 0$ $(n_{\rm a} = 1)$ corresponds to the ground $(|g\rangle)$ (excited $|e\rangle$) state of the two-level system. This effective electric field is contingent upon both the initial state of the matter and the state of light. The semiclassical Hamiltonian,

$$\hat{H}_{\text{eff}} = \frac{1}{2}\hbar\omega_0\sigma_z + \hat{\sigma}_x E_{\text{eff}},\tag{27}$$

is adept at mimicking the quantum dynamics up to the second order. A comparative analysis is conducted with this approach, the quantum-classical master-equation formalism [Eq. (24)], and the full quantum results.

The forthcoming simulations assume that the phase-space correlations^{71,72} in Eq. (24) are aligned with their quantum-mechanical counterparts. Figures 1 and 2 illustrate the dynamics of the ground and excited electronic state populations of the two-level system. The exact quantum (blue), quantum-classical master equation (magenta), and standard semiclassical (green) dynamics, induced by the corresponding effective electric field, are compared. The analysis includes various states of the light field exhibiting non-classical signatures.^{44–46} The numerical integration of Eq. (24) extends beyond the secular approximation^{73,74} and encompasses non-Markovian effects due to the non-locality in time of the dynamics.⁶⁴

Initially, the case of a Fock state $|\psi_c\rangle = |n_c\rangle$, with n_c photons in the cavity, is examined; the initial condition is $\hat{\rho}_{\rm ph}(t_0) = |n_c\rangle\langle n_c|$. The Wigner representation of this state features negative regions, indicative of a non-classical character.^{44–46} For this state of the light field, the expected value of the light field operator is $\langle \hat{\Phi}^{(I)}(t) \rangle = 0$. The symmetrized correlation function is expressed as $C(t, t') = \hbar^2 g^2 (2n_c + 1) \cos(\omega_c(t - t'))$, and the antisymmetrized correlation function is $\mathcal{A}(t, t') = -\hbar^2 g^2 \sin(\omega_c(t - t'))$. The value of $\mathcal{A}(t, t')$ is independent of the cavity state and is determined by the commutation relation $[\hat{a}, \hat{a}^{\dagger}] = \hat{1}$ (refer to Section S3 of the Supporting Information).




Figure 1: Populations of the excited (ground) state of the Rabi model as functions of time, assuming a Fock state with n_c photons in the cavity. Initially, the entire population is in the ground (excited) state $\langle g | \hat{\rho}_m(t=0) | g \rangle = 1$ ($\langle e | \hat{\rho}_m(t=0) | e \rangle = 1$). (A) $\omega_c = 0.75 \omega_0$, $g = 0.01\omega_0$ and $n_c = 1$. (B) $\omega_c = 0.9 \omega_0$, $g = 0.01\omega_0$ and $n_c = 1$. (C) $\omega_c = 0.75 \omega_0$, $g = 0.01\omega_0$ and $n_c = 0$. (D) $\omega_c = 0.9 \omega_0$, $g = 0.01\omega_0$ and $n_c = 0$. Color coding is shown on top.

Figures 1(A) and 1(B) illustrate the population dynamics of the excited electronic state $\langle e | \hat{\rho}_{\rm m}(t) | e \rangle$ for a one-photon Fock state $| \psi_{\rm c} \rangle = | 1 \rangle$ in the cavity. Two distinct frequencies, $\omega_{\rm c} = 0.75 \,\omega_0$ (Fig. 1(A)) and $\omega_{\rm c} = 0.9\omega_0$ (Fig. 2(B)), and a coupling strength of $g = 0.01\omega_0$ are considered. The assumption here is an initial total population in the ground electronic state, given by $\langle g | \hat{\rho}_{\rm m}(t=0) | g \rangle = 1$. Conversely, Figures 1(C) and 1(D) depict the population dynamics of the ground electronic state $\langle g | \hat{\rho}_{\rm m}(t) | g \rangle$ for a zero-photon Fock state $| \psi_{\rm c} \rangle = | 0 \rangle$

(vacuum state) in the cavity. The same frequencies and coupling strength as the prior case are considered, and the initial total population is assumed to be in the excited electronic state, expressed as $\langle e | \hat{\rho}_{\rm m}(t=0) | e \rangle = 1$.

A comparison with the exact quantum results reveals that the dynamics described by the quantum-classical master equation are more accurate than those obtained from the standard semiclassical approach. The latter exhibits an oscillatory behavior that is overestimated compared to the exact quantum results, a discrepancy attributable to the exclusion of the complex part of the two-time correlation function in the effective electric field. Consider now a Fock state superposition $|\psi_c\rangle = c_n |n\rangle + c_{n+1} |n+1\rangle$. The symmetrized correlation function for this state is given by $C(t, t') = \hbar^2 g^2 (|c_n|^2 (2n_c+1) + |c_{n+1}|^2 (2n_c+3)) \cos \omega_c (t-t')$, while the antisymmetrized correlation function $\mathcal{A}(t, t')$ remains constant at $-\hbar^2 g^2 \sin \omega_c (t-t')$ as it is independent of the light field state. This state also yields a non-zero expected value of the light field operator $\langle \hat{\Phi}^{(I)}(t) \rangle = 2\hbar g \sqrt{n+1} c_n c_{n+1} \cos \omega t$. This light field state exhibits non-trivial quantum characteristics, as indicated by a negative value of the Mandel parameter.⁴⁴⁻⁴⁶

Fig. 2 illustrates the ground state population (Cases A and B) and ground-excited state coherence (Cases C and D) dynamics of the two-level system. In Figures 2(A) and 2(C), a superposition of the vacuum and one-photon Fock states $|\psi_c\rangle = \sqrt{0.2}|0\rangle + \sqrt{0.8}|1\rangle$ is considered, with a frequency $\omega_c = 0.75\omega_0$ and a coupling strength $g = 0.015\omega_0$. In contrast, Figures 2(B) and 2(D) involve a light field state $|\psi_c\rangle = \sqrt{0.5}|4\rangle + \sqrt{0.5}|5\rangle$, frequency $\omega_c = 0.9\omega_0$, and a coupling strength $g = 0.0025\omega_0$. One might expect that the non-classicality of the Fock state's Wigner representation, indicated by negative regions in the phase space, would result in dynamics distinct from those produced by classical sources. However, this expectation does not hold, corroborated by previous works.^{23,26,27} A comparison to the exact quantum results reveals that the quantum-classical master equation offers more accurate dynamics than the standard semiclassical approach, which exhibits an overestimated oscillatory behavior.

Considering a squeezed vacuum state $|\psi_c\rangle = |\xi\rangle$,^{46,75} where $\varphi = 0$ and thus $\xi = r$,




Figure 2: Ground state populations and ground-excited state coherences of the Rabi model as functions of time, considering different Fock state superpositions: (A) and (C) $\omega_{\rm c} = 0.75\omega_0$, $g = 0.015\omega_0$ and $|\psi_{\rm c}\rangle = \sqrt{0.2}|0\rangle + \sqrt{0.8}|1\rangle$. (B) and (D) $\omega_{\rm c} = 0.9\omega_0$, $g = 0.0025\omega_0$ and $|\psi_{\rm c}\rangle = \sqrt{0.5}|4\rangle + \sqrt{0.5}|5\rangle$. Initially, the entire population is in the excited state $\langle e | \hat{\rho}_{\rm m}(t=0) | e \rangle = 1$. Color coding is shown on top.




Figure 3: Ground state populations of the Rabi model as functions of time, considering different squeezed vacuum states: (A) $\omega_{\rm c} = 0.75\omega_0$, $g = 0.005\omega_0$ and $|\psi_{\rm c}\rangle = |\xi; r = 0.2\rangle$. (B) $\omega_{\rm c} = 0.9\omega_0$, $g = 0.005\omega_0$ and $|\psi_{\rm c}\rangle = |\xi; r = 0.2\rangle$. (C) $\omega_{\rm c} = 0.9\omega_0$, $g = 0.005\omega_0$ and $|\psi_{\rm c}\rangle = |\xi; r = 1.2\rangle$. Initially, the entire population is in the excited state $\langle e | \hat{\rho}_{\rm m}(t=0) | e \rangle = 1$. Color coding is shown on top.

and r denotes the squeezing parameter (see section S3 of the Supporting Information), the symmetrized correlation function becomes $C(t, t') = \hbar^2 g^2 ((2 \sinh^2 r + 1) \cos \omega_c (t - t') - 2(\cosh r \sinh r) \cos \omega_c (t + t'))$, with the expected value of the light field operator given by $\langle \hat{\Phi}^{(I)}(t) \rangle = 0$. The non-classical nature of the squeezed vacuum state is highlighted by a negative squeezing parameter $S = -\frac{1}{2}(1 - e^{-2r})$.⁴⁶

Figures 3(A) and 3(B) illustrate the ground state population of the two-level system for a squeeze parameter r = 0.2. The quantum-classical master equation offers superior results compared to the standard semiclassical approach. The latter exhibits an exaggerated oscillatory behavior in contrast with the exact quantum results. However, it's noteworthy that for higher values of r, even under weak light-matter coupling, both the quantum-classical master equation and the standard semiclassical approaches fail to accurately model the molecular-only dynamics (see Figure 3C). This deviation underscores a distinct quantum signature within the molecular-only dynamics, diverging significantly from the dynamics influenced by classical light sources.⁷⁶

Dicke Model.—The Dicke model, describing an ensemble of N two-level atoms collectively coupled to a single quantized cavity mode, is renowned for its ground-state transition to a




Figure 4: Ground state populations of the site one as functions of time in the Dicke model of 4 atoms initially in their excited states It is assumed a Fock state with n_c photons in the cavity. (A) $\omega_c = 0.75 \omega_0$, $g = 0.015 \omega_0$ and $n_c = 0$. (B) $\omega_c = 0.9 \omega_0$, $g = 0.005 \omega_0$ and $n_c = 0$. (C) $\omega_c = 0.75 \omega_0$, $g = 0.008 \omega_0$ and $n_c = 10$. (D) $\omega_c = 0.99 \omega_0$, $g = 0.0005 \omega_0$ and $n_c = 10$. Color coding is shown on top.

superradiant phase.^{74,77} The Hamiltonian for the Dicke model is given by

$$\hat{H}_{\rm D} = \frac{1}{2}\hbar\omega_0 \sum_{i=1}^{N} \hat{\sigma}_{z(i)} + \hbar\omega_{\rm c} \hat{a}^{\dagger} \hat{a} + \hbar g \sum_{i=1}^{N} \hat{\sigma}_{x(i)} (\hat{a}^{\dagger} + \hat{a}).$$
(28)

This model, a cornerstone of many-body dynamics in quantum optics, has recently been explored within the realm of atom-only dynamics utilizing the non-secular Redfield master equation approach.⁷⁴

Fig. 4 illustrates the ground state population of one of the four two-level atoms, all initially in their excited states and considering a Fock state in the cavity. Results for a quantized cavity mode corresponding to a vacuum state $|\psi_c\rangle = |0\rangle$ with frequencies $\omega_c = 0.75 \omega_0$ and $\omega_c = 0.9 \omega_0$, and coupling strengths $g = 0.015\omega_0$ and $g = 0.005\omega_0$ are presented in Fig. 4(A) and 4(B) respectively. Fig. 4(C) depicts a scenario where the quantized cavity mode is in a Fock state of ten photons $|\psi_c\rangle = |10\rangle$, with a frequency of $\omega_c = 0.75 \omega_0$, and a coupling strength of $g = 0.008\omega_0$. Fig. 4(D) considers a cavity mode also in a ten-photon Fock state but in resonance with the two-level atom excitation energy, $\omega_c = 0.99 \omega_0$, and a significantly reduced coupling strength $g = 0.0005\omega_0$.

Across all considered scenarios, the quantum-classical master equation approach outperforms the standard semiclassical dynamics in approximation accuracy to the exact quantum dynamics.

Conclusions.—This study has presented a second-order quantum-classical master equation formalism, laying down the criteria for classical light field states to emulate the effects of quantum light sources on molecular systems. The comparative analysis was anchored in the simulation of molecular-only dynamics, juxtaposing the quantum-classical, quantum-exact, and semiclassical approaches across various systems within the scope of molecular polaritonics.

A discernible alignment is observed with the quantum-classical approach resonating more accurately with the quantum exact molecular-only dynamics at the second-order in lightmatter interaction relative to the semiclassical approach. A pivotal realization surfaces that the induced dynamics between a classical electric field mirroring the statistics of the quantum electric field are indistinguishable.

The implication is profound, underscoring that the intrinsic non-classical character of a light quantum state does not predicate a distinctive molecular dynamic relative to excitations emanating from a classical source, a phenomenon evident in the behavior of Fock states. This observation finds corroboration in other investigative forays into polaritonic chemistry.

A curious capability emerges, where classical light fields exterior to the cavity convincingly replicate effects intrinsic to the cavity's confines. A caveat, however, surfaces in scenarios marked by weak light-matter coupling. Here, a vacuum squeezed state, characterized by a high value of the squeeze parameter r, precipitates a departure where both the quantum-classical master equation and the standard semiclassical approach falter in capturing the nuanced molecular-only dynamics. This underscores the emergence of complex, non-trivial quantum molecular-only dynamics.

An avenue of exploration and expansion beckons, where the quantum-classical open quantum system approach can be extrapolated to embrace multiple cavity modes. This integration promises to weave in the often sidestepped dissipation and decoherence effects, injecting a layer of realism often overlooked in the prevailing light-matter interaction wave function methodologies.

Acknowledgement

Discussions with Profs. Paul Brumer, José Luis Sanz-Vicario, Alessandro Sergi and Johan F. Triana are acknowledged with pleasure. L.F.C. acknowledges financial support from the *Departamento Administrativo de Ciencia, Tecnología e Innovación* –COLCIENCIAS–. This work was supported by the *Vicerrectoría de Investigación* at Universidad de Antioquia, Colombia.

Supporting Information Available

Second Order Master Equation; Light-field dynamics; Light field correlation functions.

References

- Ebbesen, T. W. Hybrid Light-Matter States in a Molecular and Material Science Perspective. Acc. Chem. Res. 2016, 49, 2403–2412.
- (2) Ribeiro, R. F.; Martínez-Martínez, L. A.; Du, M.; Campos-Gonzalez-Angulo, J.; Yuen-Zhou, J. Polariton Chemistry: Controlling Molecular Dynamics with Optical Cavities. *Chem. Sci.* 2018, 9, 6325–6339.
- (3) Ruggenthaler, M.; Tancogne-Dejean, N.; Flick, J.; Appel, H.; Rubio, A. From a Quantum-Electrodynamical Light–Matter Description to Novel Spectroscopies. *Nat. Rev. Chem.* 2018, 2, 1–16.
- (4) Flick, J.; Rivera, N.; Narang, P. Strong Light-Matter Coupling in Quantum Chemistry and Quantum Photonics. *Nanophotonics* 2018, 7, 1479–1501.
- (5) Dovzhenko, D.; Ryabchuk, S.; Rakovich, Y. P.; Nabiev, I. Light–Matter Interaction in the Strong Coupling Regime: Configurations, Conditions, and Applications. *Nanoscale* 2018, 10, 3589–3605.
- (6) Herrera, F.; Spano, F. C. Theory of Nanoscale Organic Cavities: The Essential Role of Vibration-Photon Dressed States. ACS Photonics 2018, 5, 65–79.
- (7) Feist, J.; Galego, J.; Garcia-Vidal, F. J. Polaritonic Chemistry with Organic Molecules. ACS Photonics 2018, 5, 205–216.
- (8) Kockum, A. F.; Miranowicz, A.; De Liberato, S.; Savasta, S.; Nori, F. Ultrastrong Coupling Between Light and Matter. *Nat. Rev. Phys.* **2019**, *1*, 19–40.

- (9) Hertzog, M.; Wang, M.; Mony, J.; Börjesson, K. Strong Light–Matter Interactions: A New Direction Within Chemistry. *Chem. Soc. Rev.* **2019**, *48*, 937–961.
- (10) Herrera, F.; Owrutsky, J. Molecular Polaritons for Controlling Chemistry with Quantum Optics. J. Chem. Phys. 2020, 152, 100902.
- (11) Mandal, A.; Taylor, M. A.; Weight, B. M.; Koessler, E. R.; Li, X.; Huo, P. Theoretical Advances in Polariton Chemistry and Molecular Cavity Quantum Electrodynamics. *Chem. Rev.* 2023, 123, 9786–9879.
- (12) Campos-Gonzalez-Angulo, J.; Poh, Y.; Du, M.; Yuen-Zhou, J. Swinging Between Shine and Shadow: Theoretical Advances on Thermally Activated Vibropolaritonic Chemistry. J. Chem. Phys. 2023, 158, 230901.
- (13) Kavokin, A.; Baumberg, J. J.; Malpuech, G.; Laussy, F. P. *Microcavities*; Oxford University Press, New York, 2007.
- (14) Martínez-Martínez, L. A.; Du, M.; Ribeiro, R. F.; Kéna-Cohen, S.; Yuen-Zhou, J. Polariton-Assisted Singlet Fission in Acene Aggregates. J. Phys. Chem. Lett. 2018, 9, 1951–1957.
- (15) Martínez-Martínez, L. A.; Eizner, E.; Kéna-Cohen, S.; Yuen-Zhou, J. Triplet Harvesting in the Polaritonic Regime: A Variational Polaron Approach. J. Chem. Phys. 2019, 151, 054106.
- (16) Zhong, X.; Chervy, T.; Wang, S.; George, J.; Thomas, A.; Hutchison, J. A.; Devaux, E.; Genet, C.; Ebbesen, T. W. Non-Radiative Energy Transfer Mediated by Hybrid Light-Matter States. Angew. Chem. 2016, 128, 6310–6314.
- (17) Zhong, X.; Chervy, T.; Zhang, L.; Thomas, A.; George, J.; Genet, C.; Hutchison, J. A.;
 Ebbesen, T. W. Energy Transfer Between Spatially Separated Entangled Molecules.
 Angew. Chem. Int. Ed. 2017, 56, 9034–9038.

- (18) Sáez-Blázquez, R.; Feist, J.; Fernández-Domínguez, A.; García-Vidal, F. Organic Polaritons Enable Local Vibrations to Drive Long-Range Energy Transfer. *Phys. Rev. B* 2018, 97, 241407.
- (19) Du, M.; Martínez-Martínez, L. A.; Ribeiro, R. F.; Hu, Z.; Menon, V. M.; Yuen-Zhou, J. Theory for Polariton-Assisted Remote Energy Transfer. *Chem. Sci.* 2018, 9, 6659–6669.
- (20) Sáez-Blázquez, R.; Feist, J.; Romero, E.; Fernández-Domínguez, A. I.; García-Vidal, F. J. Cavity-Modified Exciton Dynamics in Photosynthetic Units. J. Phys. Chem. Lett. 2019, 10, 4252–4258.
- (21) Du, M.; Ribeiro, R. F.; Yuen-Zhou, J. Remote Control of Chemistry in Optical Cavities. Chem 2019, 5, 1167–1181.
- (22) Galego, J.; Garcia-Vidal, F. J.; Feist, J. Cavity-Induced Modifications of Molecular Structure in the Strong-Coupling Regime. *Phys. Rev. X* 2015, 5, 041022.
- (23) Kowalewski, M.; Bennett, K.; Mukamel, S. Cavity Femtochemistry: Manipulating Nonadiabatic Dynamics at Avoided Crossings. J. Phys. Chem. Lett. 2016, 7, 2050–2054.
- (24) Gu, B.; Mukamel, S. Manipulating Nonadiabatic Conical Intersection Dynamics by Optical Cavities. *Chem. Sci.* 2020, 11, 1290–1298.
- (25) Shapiro, M.; Brumer, P. Quantum control of molecular processes; Wiley-VCH, Wertheim, 2012.
- (26) Csehi, A.; Halász, G. J.; Cederbaum, L. S.; Vibók, A. Competition Between Light-Induced and Intrinsic Nonadiabatic Phenomena in Diatomics. J. Phys. Chem. Lett. 2017, 8, 1624–1630.
- (27) Csehi, A.; Vibók, A.; Halász, G. J.; Kowalewski, M. Quantum Control with Quantum Light of Molecular Nonadiabaticity. *Phys. Rev. A* 2019, 100, 053421.

- (28) Calderón, L. F.; Pachón, L. A. Nonadiabatic Sunlight Harvesting. Phys. Chem. Chem. Phys. 2020, 22, 12678–12687.
- (29) Calderón, L. F.; Chuang, C.; Brumer, P. Electronic-Vibrational Resonance Does Not Significantly Alter Steady-State Transport in Natural Light-Harvesting Systems. J. Phys. Chem. Lett. 2023, 14, 1436–1444.
- (30) Hoffmann, N. M.; Schäfer, C.; Rubio, A.; Kelly, A.; Appel, H. Capturing Vacuum Fluctuations and Photon Correlations in Cavity Quantum Electrodynamics with Multitrajectory Ehrenfest Dynamics. *Phys. Rev. A* 2019, *99*, 063819.
- (31) Hoffmann, N. M.; Schäfer, C.; Säkkinen, N.; Rubio, A.; Appel, H.; Kelly, A. Benchmarking Semiclassical and Perturbative Methods for Real-Time Simulations of Cavity-Bound Emission and Interference. J. Chem. Phys. 2019, 151, 244113.
- (32) Chen, H.-T.; Li, T. E.; Sukharev, M.; Nitzan, A.; Subotnik, J. E. Ehrenfest+ R Dynamics.
 I. A Mixed Quantum-classical Electrodynamics Simulation of Spontaneous Emission. J. Chem. Phys. 2019, 150, 044102.
- (33) Li, T. E.; Chen, H.-T.; Nitzan, A.; Subotnik, J. E. Quasiclassical Modeling of Cavity Quantum Electrodynamics. *Phys. Rev. A* 2020, 101, 033831.
- (34) Rosenzweig, B.; Hoffmann, N. M.; Lacombe, L.; Maitra, N. T. Analysis of the Classical Trajectory Treatment of Photon Dynamics for Polaritonic Phenomena. J. Chem. Phys. 2022, 156, 054101.
- (35) He, X.; Wu, B.; Shang, Y.; Li, B.; Cheng, X.; Liu, J. New Phase Space Formulations and Quantum Dynamics Approaches. Wiley Interdiscip. Rev. Comput. Mol. Sci. 2022, 12, e1619.
- (36) Li, T. E.; Tao, Z.; Hammes-Schiffer, S. Semiclassical Real-Time Nuclear-Electronic

Orbital Dynamics for Molecular Polaritons: Unified Theory of Electronic and Vibrational Strong Couplings. J. Chem. Theory Comput. **2022**, 18, 2774–2784.

- (37) Hsieh, M.-H.; Krotz, A.; Tempelaar, R. A Mean-Field Treatment of Vacuum Fluctuations in Strong Light–Matter Coupling. J. Phys. Chem. Lett. 2023, 14, 1253–1258.
- (38) Hsieh, M.-H.; Krotz, A.; Tempelaar, R. Ehrenfest Modeling of Cavity Vacuum Fluctuations and How to Achieve Emission from a Three-Level Atom. arXiv preprint arXiv:2309.01912 2023,
- (39) Gerasimenko, V. I. Dynamical Equations of Quantum-Classical Systems. Theor. Math. Phys. 1982, 50, 49–55.
- (40) Grunwald, A., Robbieand Kelly; Kapral, R. Energy Transfer Dynamics in Biomaterial Systems; Springer Berlin Heidelberg, Berlin, Heidelberg, 2009; pp 383–413.
- (41) Kapral, R.; Ciccotti, G. Mixed Quantum-Classical Dynamics. J. Chem. Phys. 1999, 110, 8919–8929.
- (42) Kapral, R. Quantum Dynamics in Open Quantum-Classical Systems. J. Phys. Condens. Matter 2015, 27, 073201.
- (43) Toutounji, M.; Kapral, R. Subsystem Dynamics in Mixed Quantum–Classical Systems. Chem. Phys. 2001, 268, 79–89.
- (44) Mandel, L.; Wolf, E. Optical Coherence and Quantum Optics; Cambridge University Press, New York, 1995.
- (45) Gerry, C.; Knight, P. L. Introductory Quantum Optics; Cambridge University Press, New York, 2005.
- (46) Agarwal, G. S. *Quantum Optics*; Cambridge University Press, New York, 2012.

- (47) Flick, J.; Ruggenthaler, M.; Appel, H.; Rubio, A. Atoms and Molecules in Cavities, From Weak to Strong Coupling in Quantum-Electrodynamics (QED) Chemistry. Proc. Natl. Acad. Sci. U.S.A. 2017, 114, 3026–3034.
- (48) Power, E. A.; Zienau, S. Coulomb Gauge in Non-Relativistic Quantum Electro-Dynamics and the Shape of Spectral Lines. *Philos. Trans. R. Soc. A* 1959, 251, 427–454.
- (49) Atkins, P.; Woolley, R. The Interaction of Molecular Multipoles with the Electromagnetic Field in the Canonical Formulation of Non-Covariant Quantum Electrodynamics. *Proc. R. Soc. A* 1970, *319*, 549–563.
- (50) Woolley, R. G. Molecular Quantum Electrodynamics. Proc. R. Soc. A 1971, 321, 557–572.
- (51) Scheel, S.; Buhmann, S. Y. Macroscopic Quantum Electrodynamics Concepts and Applications. Acta Phys. Slovaca 2008, 58, 675–809.
- (52) Feist, J.; Fernández-Domínguez, A. I.; García-Vidal, F. J. Macroscopic QED for Quantum Nanophotonics: Emitter-Centered Modes as a Minimal Basis for Multiemitter Problems. Nanophotonics 2020, 10, 477–489.
- (53) Svendsen, M. K.; Thygesen, K. S.; Rubio, A.; Flick, J. Molecules in Real Cavities with Quantum Electroynamical Density Functional Theory. arXiv preprint arXiv:2305.02391 2023,
- (54) Hernández, F. J.; Herrera, F. Multi-level Quantum Rabi Model for Anharmonic Vibrational Polaritons. J. Chem. Phys. 2019, 151, 144116.
- (55) Rokaj, V.; Welakuh, D. M.; Ruggenthaler, M.; Rubio, A. Light–Matter Interaction in the Long-Wavelength Limit: No Ground-State Without Dipole Self-Energy. J. Phys. B: At. Mol. Opt. Phys. 2018, 51, 034005.

- (56) Fregoni, J.; Garcia-Vidal, F. J.; Feist, J. Theoretical Challenges in Polaritonic Chemistry. ACS photonics 2022, 9, 1096–1107.
- (57) Weyl, H. Quantenmechanik und Gruppentheorie. Z. Phys. 1927, 46, 1–46.
- (58) Wigner, E. P. On the Quantum Correction For Thermodynamic Equilibrium. *Phys. Rev.* 1932, 40, 749.
- (59) J. E. Moyal, Quantum Mechanics as a Statistical Theory. Proc. Camb. Phil. Soc. 1949, 45, 99–124.
- (60) Beck, G. M.; Sergi, A. Quantum Dynamics in the Partial Wigner Picture. J. Phys. A 2013, 46, 395305.
- (61) Ingold, G.-L. In Coherent Evolution in Noisy Environments; Buchleitner, A., Hornberger, K., Eds.; Springer-Verlag, Berlin-Heidelberg-New York, 2002; pp 1–53.
- (62) Cohen-Tannoudji, C.; Dupont-Roc, J.; Grynberg, G. Atom-photon Interactions: Basic Processes and Applications; John Wiley & Sons, New York, 1998.
- (63) Pachón, L. A.; Brumer, P. Incoherent Excitation of Thermally Equilibrated Open Quantum Systems. *Phys. Rev. A* 2013, *87*, 022106.
- (64) Pachón, L. A.; Yu, L.; Brumer, P. Coherent One-Photon Phase Control in Closed and Open Quantum Systems: A General Master Equation Approach. *Faraday Discuss.* 2013, 163, 485–495.
- (65) Pachón, L. A.; Brumer, P. Quantum Driven Dissipative Parametric Oscillator in a Blackbody Radiation Field. J. Math. Phys. 2014, 55, 012103.
- (66) Pachón, L. A.; Triana, J. F.; Zueco, D.; Brumer, P. Influence of Non-Markovian Dynamics in Equilibrium Uncertainty-Relations. J. Chem. Phys. 2019, 150, 034105.

- (67) Trushechkin, A.; Merkli, M.; Cresser, J.; Anders, J. Open Quantum System Dynamics and the Mean Force Gibbs State. AVS Quantum Sci. 2022, 4, 012301.
- (68) Anto-Sztrikacs, N.; Nazir, A.; Segal, D. Effective-Hamiltonian Theory of Open Quantum Systems at Strong Coupling. *PRX Quantum* **2023**, *4*, 020307.
- (69) Ahn, W.; Triana, J. F.; Recabal, F.; Herrera, F.; Simpkins, B. S. Modification of Ground-State Chemical Reactivity via Light–Matter Coherence in Infrared Cavities. *Science* 2023, 380, 1165–1168.
- (70) Schäfer, C.; Ruggenthaler, M.; Rokaj, V.; Rubio, A. Relevance of the Quadratic Diamagnetic and Self-Polarization Terms in Cavity Quantum Electrodynamics. ACS Photonics 2020, 7, 975–990.
- (71) Carmichael, H. J. Statistical Methods in Quantum Optics 1: Master Equations and Fokker-Planck Equations; Springer-Verlag, Berlin-Heidelberg-New York, 1999.
- (72) Carmichael, H. J. Statistical Methods in Quantum Optics 2: Non-Classical Fields;
 Springer Verlag, Berlin-Heidelberg-New York, 2009.
- (73) Dodin, A.; Tscherbul, T.; Alicki, R.; Vutha, A.; Brumer, P. Secular Versus Nonsecular Redfield Dynamics and Fano Coherences in Incoherent Excitation: An Experimental Proposal. *Phys. Rev. A* 2018, *97*, 013421.
- (74) Damanet, F.; Daley, A. J.; Keeling, J. Atom-Only Descriptions of the Driven-Dissipative Dicke Model. Phys. Rev. A 2019, 99, 033845.
- (75) Breuer, H.; Petruccione, F. The Theory of Open Quantum Systems; Oxford University Press, New York, 2002.
- (76) Triana, J. F.; Pelaez Ruiz, D.; Sanz-Vicario, J. L. Entangled Photonic-Nuclear Molecular Dynamics of LiF in Quantum Optical Cavities. J. Phys. Chem. A 2018, 122, 2266–2278.

(77) Kirton, P.; Roses, M. M.; Keeling, J.; Dalla Torre, E. G. Introduction to the Dicke Model: From Equilibrium to Nonequilibrium, and Vice Versa. Adv. Quantum Technol. 2019, 2, 1800043.