NOTES ON SEMISIMPLE TENSOR CATEGORIES OF RANK TWO

HUA SUN, HUI-XIANG CHEN, AND YINHUO ZHANG

ABSTRACT. In this paper, we show that there are infinitely many semisimple tensor (or monoidal) categories of rank two over an algebraically closed field \mathbb{F} .

Introduction

In recent years, great breakthroughs have been made in the classification of fusion categories. Ostrik classified the fusion categories of rank 2 [9]. He proved that there exist only 4 fusion categories of rank 2 up to tensor equivalence. Furthermore, he classified the pivotal fusion categories of rank 3 [10]. Larson investigated the Pseudo-unitary non-self-dual fusion categories of rank 4 [6]. Rowell, Strong and Wang proved that there exist 35 unitary modular categories of rank less than 4 up to ribbon tensor equivalence [11]. Recently, Chen and Zhang showed that a finite rank tensor category C is uniquely determined by three structure invariants [2]: the Green ring r(C), the Auslander algebra A(C) and the associator constraints. In particular, if C is semisimple tensor category of finite rank, then C is uniquely determined by the Green ring r(C) and the associator constraints. We ask ourselves the question: given a \mathbb{Z}_+ -ring R, is there a semisimple tensor category such that its Green ring is isomorphic to R? Such a tensor category is called a categorification of the \mathbb{Z}_+ -ring. If it exists, then how many tensor categories are there such that their Green rings are isomorphic to R?

In this paper, we consider the categorization of the \mathbb{Z}_+ -rings R with a \mathbb{Z}_+ -basis $\{r_1 = 1, r_2\}$ such that $r_2^2 \neq 0$. Let $r_2^2 = mr_1 + nr_2$ for some nonnegative integers m, n with m + n > 0. We divide it into three cases: m > 0, n = 0; m = 0, n > 0; m, n > 0. We show that there is no semisimple tensor category C such that the Green ring r(C) is isomorphic to R with m > 1 and n = 0. If m = 1 and n = 0, then there is only two semisimple tensor categories C_+ and C_- (up to tensor equivalent) when char($\mathbb{F} \neq 2$. Moreover, $C_+ = C_-$ if char($\mathbb{F} = 2$. If m = 0 and n = 1, then there is only one semisimple tensor categories C_n and C'_n for any n. If m = n = 1, then there is only two semisimple tensor categories $C_{(0)}$ and $C'_{(0)}$ when char($\mathbb{F} \neq 5$, and there is only one semisimple tensor category $C_{(5)}$ when char($\mathbb{F} = 5$. Finally, we proved that if m > 1 and $2m > n^2$, then there is no semisimple tensor category C such that r(C) is isomorphic to R.

²⁰¹⁰ Mathematics Subject Classification. 18D10, 16T05.

Key words and phrases. Green ring, Auslander algebra, associator, tensor category.

The paper is organized as follows. In Section 1, we introduce some basic notations including a (\mathbf{m}, \mathbf{s}) -type matrix and a \mathbb{Z}_+ -ring. In Section 2, we characterize the structure invariants of the semisimple tensor category of rank two using 6-*j* symbols. We prove that the pentagon commutative diagram of a tensor category is equivalent to the Biedenharn-Elliot identity. In Section 3, we consider the categorizations of R with a \mathbb{Z}_+ -basis $\{r_1 = 1, r_2\}$ such that $r_2^2 \neq 0$.

1. Preliminaries and Notations

Throughout, let \mathbb{F} be an algebraically closed field, \mathbb{Z} the ring of integers, and \mathbb{N} the set of nonnegative integers. Unless otherwise stated, all algebras are defined over \mathbb{F} , dim means dim_{\mathbb{F}}. All rings and algebras are assumed to be associative with identity. The definitions of a \mathbb{Z}_+ -basis and a \mathbb{Z}_+ -ring can be found in [3, 7, 8]. For the theory of (tensor) categories, we refer the reader to [1, 4, 5].

Definition 1.1. (1) Let R be a ring free as a module over \mathbb{Z} . A \mathbb{Z}_+ -basis of R is a \mathbb{Z} -basis $B = \{b_i\}_{i \in I}$ such that for any $i, j \in I$, $b_i b_j = \sum_l c_{ijl} b_l$ with $c_{ijl} \in \mathbb{N}$. (2) A \mathbb{Z}_+ -ring is a \mathbb{Z} -algebra with identity 1 endowed with a fixed \mathbb{Z}_+ -basis. (3) A \mathbb{Z}_+ -ring with a \mathbb{Z}_+ -basis B is a unital \mathbb{Z}_+ -ring if $1 \in B$.

For an \mathbb{F} -algebra A and two positive integers m and n, we denote by $M_{m \times n}(A)$ the \mathbb{F} -space consisting of all $m \times n$ -matrices over A, and by $M_m(A) := M_{m \times m}(A)$ the full matrix algebra of $m \times m$ -matrices over A. If B is a subspace of A, let $M_{m \times n}(B)$ denote the subspace of $M_{m \times n}(A)$ consisting of all $m \times n$ -matrices over A with entries contained in B.

In the following, assume that A is the \mathbb{F} -algebra \mathbb{F}^2 with a given set of orthogonal primitive idempotents $\{e_1, e_2\}$ such that $e_1 + e_2 = 1$. In this case, $Ae_i = \mathbb{F}e_i$, i = 1, 2

For any $\mathbf{m} = (m_1, m_2) \in \mathbb{N}^2$, let $|\mathbf{m}| := m_1 + m_2$. For $\mathbf{m} = (m_1, m_2), \mathbf{s} = (s_1, s_2) \in \mathbb{N}^2$ with $|\mathbf{m}| > 0$ and $|\mathbf{s}| > 0$, an (\mathbf{m}, \mathbf{s}) -type matrix over A is a matrix $X \in M_{|\mathbf{m}| \times |\mathbf{s}|}(A)$ defined by $X = \begin{pmatrix} X_1 & 0 \\ 0 & X_2 \end{pmatrix}$ with $X_i \in M_{m_i \times s_i}(\mathbb{F}e_i), i = 1, 2$.

A $(\mathbf{0}, \mathbf{m})$ -type matrix over A means a $1 \times |\mathbf{m}|$ zero matrix over A, and an $(\mathbf{m}, \mathbf{0})$ -type matrix over A means an $|\mathbf{m}| \times 1$ zero matrix over A. Of course, a $(\mathbf{0}, \mathbf{0})$ -type matrix over A means a 1×1 zero matrix over A. Here $\mathbf{0} = (0, 0)$.

For any positive integer m, let I_m denote the $m \times m$ identity matrix over A (or over \mathbb{F}). Then the $m \times m$ identity matrix over $\mathbb{F}e_i$ is $e_i I_m$ (or $I_m e_i$), where i = 1, 2.

For $\mathbf{m}, \mathbf{s} \in \mathbb{N}^2$ with $\mathbf{m} \neq \mathbf{0}$ and $\mathbf{s} \neq \mathbf{0}$, let $M_{\mathbf{m} \times \mathbf{s}}(A)$ be the \mathbb{F} -space consisting of all (\mathbf{m}, \mathbf{s}) -type matrices over A. Then $M_{\mathbf{m}}(A) := M_{\mathbf{m} \times \mathbf{m}}(A)$ is an associative \mathbb{F} -algebra with the identity $E_{\mathbf{m}}$ given by $E_{\mathbf{m}} = \begin{pmatrix} e_1 I_{m_1} & 0 \\ 0 & e_2 I_{m_2} \end{pmatrix}$, here we regard $E_{\mathbf{m}} = e_2 I_{m_2}$ if $m_1 = 0$ and $E_{\mathbf{m}} = e_1 I_{m_1}$ if $m_2 = 0$. For convenience, we let $E_{\mathbf{0}} := 0$, the 1×1 zero matrix over A. Then $M_{\mathbf{0}}(A) = E_{\mathbf{0}} = 0$.

Note that \mathbb{N}^2 is a monoid with respect to the addition: $(m_1, m_2) + (s_1, s_2) = (m_1 + s_1, m_2 + s_2).$

Let $\mathbf{m}, \mathbf{m}', \mathbf{s} \in \mathbb{N}^2$. For any $X = \begin{pmatrix} X_1 & 0 \\ 0 & X_2 \end{pmatrix} \in M_{\mathbf{m} \times \mathbf{s}}(A)$ with $X_i \in M_{m_i \times s_i}(\mathbb{F}e_i)$ and $Y = \begin{pmatrix} Y_1 & 0 \\ 0 & Y_2 \end{pmatrix} \in M_{\mathbf{m}' \times \mathbf{s}}(A)$ with $Y_i \in M_{m'_i \times s_i}(\mathbb{F}e_i)$, define $X \oplus Y \in M_{(\mathbf{m}+\mathbf{m}') \times \mathbf{s}}(A)$ by

$$X \underline{\oplus} Y := \begin{pmatrix} X_1 & 0 \\ Y_1 & 0 \\ 0 & X_2 \\ 0 & Y_2 \end{pmatrix}$$

It is obvious that $\underline{\oplus}$ is associative, but not commutative in general.

Let $P_{\mathbf{m},\mathbf{m}'}$ be an $(|\mathbf{m} + \mathbf{m}'|) \times (|\mathbf{m} + \mathbf{m}'|)$ permutation matrix defined by

$$P_{\mathbf{m},\mathbf{m}'} = \begin{pmatrix} I_{m_1} & 0 & 0 & 0 \\ 0 & 0 & I_{m'_1} & 0 \\ 0 & I_{m_2} & 0 & 0 \\ 0 & 0 & 0 & I_{m'_2} \end{pmatrix}.$$

It is obvious that $X \oplus Y = P_{\mathbf{m},\mathbf{m}'} \begin{pmatrix} X \\ Y \end{pmatrix}$. In a similar way, we define another sum $X \oplus Y \in M_{\mathbf{m} \times (\mathbf{s} + \mathbf{s}')}(A)$ for two matrices $X \in M_{\mathbf{m} \times \mathbf{s}}(A)$ and $Y \in M_{\mathbf{m} \times \mathbf{s}'}(A)$ as follows:

$$X \overline{\oplus} Y := \left(\begin{array}{ccc} X_1 & Y_1 & 0 & 0\\ 0 & 0 & X_2 & Y_2 \end{array}\right)$$

Like the sum $\underline{\oplus}$, the sum $\overline{\oplus}$ is associative, but not commutative in general. Obviously, $X\overline{\oplus}Y = (X,Y)P_{\mathbf{s},\mathbf{s}'}^T$, where P^T denotes the transposed matrix of P.

Let $\mathbf{m}_1, \mathbf{m}_2, \cdots, \mathbf{m}_r, \mathbf{s}_1, \mathbf{s}_2, \cdots, \mathbf{s}_l \in \mathbb{N}^2$. For a matrix

$$X = \begin{pmatrix} X_{11} & X_{12} & \cdots & X_{1l} \\ X_{21} & X_{22} & \cdots & X_{2l} \\ \cdots & \cdots & \cdots & \cdots \\ X_{r1} & X_{r2} & \cdots & X_{rl} \end{pmatrix}$$

over A with $X_{ij} \in M_{\mathbf{m}_i \times \mathbf{s}_j}(A)$, $1 \leq i \leq r$, $1 \leq j \leq l$, we define an (\mathbf{m}, \mathbf{s}) -type matrix $\Pi(X)$ over A by setting

 $\Pi(X) := (X_{11} \overline{\oplus} X_{12} \overline{\oplus} \cdots \overline{\oplus} X_{1l}) \underline{\oplus} (X_{21} \overline{\oplus} X_{22} \overline{\oplus} \cdots \overline{\oplus} X_{2l}) \underline{\oplus} \cdots \underline{\oplus} (X_{r1} \overline{\oplus} X_{r2} \overline{\oplus} \cdots \overline{\oplus} X_{rl}),$ where $\mathbf{m} = \mathbf{m}_1 + \mathbf{m}_2 + \cdots + \mathbf{m}_r$ and $\mathbf{s} = \mathbf{s}_1 + \mathbf{s}_2 + \cdots + \mathbf{s}_l$. Obviously, we have $\Pi(X) = (X_{11} \underline{\oplus} X_{21} \underline{\oplus} \cdots \underline{\oplus} X_{r1}) \overline{\oplus} (X_{12} \underline{\oplus} X_{22} \underline{\oplus} \cdots \underline{\oplus} X_{r2}) \overline{\oplus} \cdots \overline{\oplus} (X_{1l} \underline{\oplus} X_{2l} \underline{\oplus} \cdots \underline{\oplus} X_{rl}).$

For $\mathbf{m}_1, \mathbf{m}_2, \cdots, \mathbf{m}_r \in \mathbb{N}^2$, define a permutation matrix $P_{\mathbf{m}_1, \cdots, \mathbf{m}_r}$ recursively on r as follows: for r = 1, $P_{\mathbf{m}_1} = I_{|\mathbf{m}_1|}$; for r = 2, $P_{\mathbf{m}_1, \mathbf{m}_2}$ is defined as before; for r > 2,

$$P_{\mathbf{m}_{1},\cdots,\mathbf{m}_{r}} := P_{\mathbf{m}_{1}+\cdots+\mathbf{m}_{r-1},\mathbf{m}_{r}} \begin{pmatrix} P_{\mathbf{m}_{1},\cdots,\mathbf{m}_{r-1}} & 0\\ 0 & I_{|\mathbf{m}_{r}|} \end{pmatrix}.$$
 Then one can see that
$$\Pi(X) = P_{\mathbf{m}_{1},\cdots,\mathbf{m}_{r}} X P_{\mathbf{s}_{1},\cdots,\mathbf{s}_{l}}^{T}.$$

Let A and B be two algebras over \mathbb{F} . Let $X = (x_{ij}) \in M_{m \times s}(A)$ and $Y = (y_{ij}) \in M_{m' \times s'}(B)$. Then one can define a matrix $X \otimes_{\mathbb{F}} Y \in M_{mm' \times ss'}(A \otimes_{\mathbb{F}} B)$ in a natural way:

$$X \otimes_{\mathbb{F}} Y := \begin{pmatrix} X \otimes_{\mathbb{F}} y_{11} & X \otimes_{\mathbb{F}} y_{12} & \cdots & X \otimes_{\mathbb{F}} y_{1s'} \\ X \otimes_{\mathbb{F}} y_{21} & X \otimes_{\mathbb{F}} y_{22} & \cdots & X \otimes_{\mathbb{F}} y_{2s'} \\ \cdots & \cdots & \cdots \\ X \otimes_{\mathbb{F}} y_{m'1} & X \otimes_{\mathbb{F}} y_{m'2} & \cdots & X \otimes_{\mathbb{F}} y_{m's'} \end{pmatrix}$$

where

$$X \otimes_{\mathbb{F}} y_{ij} := \begin{pmatrix} x_{11} \otimes_{\mathbb{F}} y_{ij} & x_{12} \otimes_{\mathbb{F}} y_{ij} & \cdots & x_{1s} \otimes_{\mathbb{F}} y_{ij} \\ x_{21} \otimes_{\mathbb{F}} y_{ij} & x_{22} \otimes_{\mathbb{F}} y_{ij} & \cdots & x_{2s} \otimes_{\mathbb{F}} y_{ij} \\ \cdots & \cdots & \cdots \\ x_{m1} \otimes_{\mathbb{F}} y_{ij} & x_{m2} \otimes_{\mathbb{F}} y_{ij} & \cdots & x_{ms} \otimes_{\mathbb{F}} y_{ij} \end{pmatrix}$$

for any $1 \leq i \leq m'$ and $1 \leq j \leq s'$. It is obvious that the above tensor product is associative. If $A = B = \mathbb{F}$, then we may regard $\mathbb{F} \otimes_{\mathbb{F}} \mathbb{F} = \mathbb{F}$ and $x_{ij} \otimes_{\mathbb{F}} y_{i'j'} = x_{ij}y_{i'j'}$, and so $X \otimes_{\mathbb{F}} Y \in M_{mm' \times ss'}(\mathbb{F})$. The following property of the tensor product is easy to check.

Lemma 1.2. Assume that A and B are two \mathbb{F} -algebras. Let $X \in M_{m \times s}(A)$, $X_1 \in M_{s \times t}(A)$, $Y \in M_{m' \times s'}(B)$ and $Y_1 \in M_{s' \times t'}(B)$. Then $(X \otimes_{\mathbb{F}} Y)(X_1 \otimes_{\mathbb{F}} Y_1) = (XX_1) \otimes_{\mathbb{F}} (YY_1)$.

Let *n* be a positive integer. For any $1 \leq i \leq n$, let $\varepsilon_{n,i} = (\delta_{i1} \ \delta_{i2} \ \cdots \ \delta_{in}) \in M_{1 \times n}(\mathbb{F})$, where $\delta_{ij} = 1$ for j = i and $\delta_{ij} = 0$ for $j \neq i$. For any $1 \leq i, j \leq n$, let $I_{i,j}^{[n]} = \varepsilon_{n,i}^T \varepsilon_{n,j} \in M_n(\mathbb{F})$. Let

$$I_{[n]} = \sum_{i,j=1}^{n} I_{i,j}^{[n]} \otimes_{\mathbb{F}} I_{j,i}^{[n]} = \begin{pmatrix} I_{1,1}^{[n]} & I_{2,1}^{[n]} & \cdots & I_{n,1}^{[n]} \\ I_{1,2}^{[n]} & I_{2,2}^{[n]} & \cdots & I_{n,2}^{[n]} \\ \cdots & \cdots & \cdots & \cdots \\ I_{1,n}^{[n]} & I_{2,n}^{[n]} & \cdots & I_{n,n}^{[n]} \end{pmatrix} \in M_{n^2}(\mathbb{F}).$$

If there is no ambiguity, $I_{i,j}^{[n]}$ is simply written as $I_{i,j}$. For any positive integers m and n, the zero matrix 0 in $M_{m \times n}(\mathbb{F})$ (or in $M_n(\mathbb{F})$) is sometimes denoted $0_{m \times n}$ (or 0_n) in order to indicate its order.

2. Construction of semisimple tensor categories of rank two

Throughout this section, assume that R is a \mathbb{Z}_+ -ring with a unital \mathbb{Z}_+ -basis $\{r_1 = 1, r_2\}_{i \in I}$ such that $r_2^2 \neq 0$, and $A = \mathbb{F}^2$ is a 2-dimensional \mathbb{F} -algebra as given in the last section. Let \mathbb{F}^{\times} be the multiplicative group of nonzero elements of \mathbb{F} .

Now we construct a category C as follows: (1) $\operatorname{Ob}(C) := \mathbb{N}^2$; (2) $\operatorname{Mor}(C)$: for $\mathbf{m} = (m_1, m_2)$ and $\mathbf{s} = (s_1, s_2)$ in $\operatorname{Ob}(C) = \mathbb{N}^2$, define $\operatorname{Hom}_{\mathcal{C}}(\mathbf{m}, \mathbf{s}) := M_{\mathbf{s} \times \mathbf{m}}(A)$; (3) Composition: for s, m, $\mathbf{t} \in \mathbb{N}^2$, define the composition

$$\operatorname{Hom}_{\mathcal{C}}(\mathbf{m}, \mathbf{s}) \times \operatorname{Hom}_{\mathcal{C}}(\mathbf{t}, \mathbf{m}) \to \operatorname{Hom}_{\mathcal{C}}(\mathbf{t}, \mathbf{s}), \ (X, Y) \mapsto X \circ Y$$

by $X \circ Y = XY \in M_{\mathbf{s} \times \mathbf{t}}(A) = \operatorname{Hom}_{\mathcal{C}}(\mathbf{t}, \mathbf{s})$, the usual product of matrices.

Let $\mathbf{e}_1 = (1,0)$, $\mathbf{e}_2 = (0,1) \in \text{Ob}(\mathcal{C})$. Then by the definition of \mathcal{C} , $\text{Hom}_{\mathcal{C}}(\mathbf{e}_1, \mathbf{e}_2) = 0$, $\text{Hom}_{\mathcal{C}}(\mathbf{e}_2, \mathbf{e}_1) = 0$, and $\text{End}_{\mathcal{C}}(\mathbf{e}_i) = e_i A e_i = \mathbb{F} e_i$, i = 1, 2. By [2, Proposition 2.14 and Corollary 2.15], we have the following proposition.

Proposition 2.1. C is a semisimple category over \mathbb{F} . Moreover, \mathbf{e}_1 , \mathbf{e}_2 are the only two non-isomorphic simple objects of C.

Note that R is a \mathbb{Z}_+ -ring with the unital \mathbb{Z}_+ -basis $\{r_1 = 1, r_2\}$. We have $r_2^2 = mr_1 + nr_2$ for some $m, n \in \mathbb{N}$ with m + n > 0. Let $\mathbf{c}_{22} = (m, n)$ and $\mathbf{c}_{1i} = \mathbf{c}_{i1} = \mathbf{e}_i$, i = 1, 2. Then $\mathbf{c}_{ij} = (c_{ij1}, c_{ij2}) \in \mathbb{N}^2$ and $r_i r_j = c_{ij1} r_1 + c_{ij2} r_2$, where $1 \leq i, j \leq 2$.

For any $\mathbf{m} = (m_1, m_2)$ and $\mathbf{s} = (s_1, s_2)$ in $Ob(\mathcal{C})$, define $\mathbf{m} \otimes \mathbf{s} \in Ob(\mathcal{C})$ by

$$\mathbf{m} \otimes \mathbf{s} := \sum_{i,j=1}^{2} m_i s_j \mathbf{c}_{ij}.$$
 (1)

Then $\mathbf{m} \otimes \mathbf{s} = \mathbf{0}$ if and only if $\mathbf{m} = \mathbf{0}$ or $\mathbf{s} = \mathbf{0}$. Moreover, $\mathbf{m} \otimes \mathbf{e}_1 = \mathbf{e}_1 \otimes \mathbf{m} = \mathbf{m}$. By the associativity of the ring R, one can see that $(\mathbf{m} \otimes \mathbf{s}) \otimes \mathbf{t} = \mathbf{m} \otimes (\mathbf{s} \otimes \mathbf{t})$, denoted $\mathbf{m} \otimes \mathbf{s} \otimes \mathbf{t}$, where $\mathbf{m}, \mathbf{s}, \mathbf{t} \in Ob(\mathcal{C})$.

Define a vector space M(R, A) over \mathbb{F} by

$$M(R,A) := \bigoplus_{1 \leq i,i',j,j' \leq 2} M_{\mathbf{c}_{i',i'} \times \mathbf{c}_{ij}}(A)$$

Then M(R, A) is an associative \mathbb{F} -algebra with the multiplication defined as follows: if $X \in M_{\mathbf{c}_{i'j'} \times \mathbf{c}_{ij}}(A)$ and $Y \in M_{\mathbf{c}_{i''j''} \times \mathbf{c}_{ij}}(A)$, then XY is the usual matrix product for (i, j) = (i'', j''), and XY = 0 for $(i, j) \neq (i'', j'')$. The identity of M(R, A) is $(E_{\mathbf{c}_{ij}})_{i,j \in \{1,2\}}$, where $E_{\mathbf{c}_{ij}} \in M_{\mathbf{c}_{ij}}(A)$ is given in the last section.

Note that $A \otimes_{\mathbb{F}} A$ has an \mathbb{F} -basis $\{e_i \otimes e_j | 1 \leq i, j \leq 2\}$. Define an \mathbb{F} -linear map $\phi : A \otimes_{\mathbb{F}} A \to M(R, A)$ by $\phi(e_i \otimes_{\mathbb{F}} e_j) = E_{\mathbf{c}_{ij}} \in M_{\mathbf{c}_{ij}}(A), 1 \leq i, j \leq 2$. Then ϕ is the unique algebra map from $A \otimes_{\mathbb{F}} A$ to M(R, A) such that the two conditions $(\phi 1)$ and $(\phi 2)$ in [2, page 266] are satisfied. Moreover, we have $\phi(x) \in M_{\mathbf{c}_{i'j'} \times \mathbf{c}_{ij}}(A)$ for any $x \in e_{i'}Ae_i \otimes_{\mathbb{F}} e_{j'}Ae_j \subset A \otimes_{\mathbb{F}} A$, where $1 \leq i', i, j', j \leq 2$.

For any homogeneous matrix $X = (x_{ij})$ over $A \otimes_{\mathbb{F}} A$, $\phi(X) = (\phi(x_{ij}))$ is a welldefined homogeneous matrix over A. Let $X \in \text{Hom}_{\mathcal{C}}(\mathbf{m}_1, \mathbf{s}_1)$ and $Y \in \text{Hom}_{\mathcal{C}}(\mathbf{m}_2, \mathbf{s}_2)$. Then one can also define $X \otimes Y \in \text{Hom}_{\mathcal{C}}(\mathbf{m}_1 \otimes \mathbf{m}_2, \mathbf{s}_1 \otimes \mathbf{s}_2)$ as follows:

$$X \otimes Y := \Pi(\phi(X \otimes_{\mathbb{F}} Y)) = P(\mathbf{s}_1, \mathbf{s}_2)\phi(X \otimes_{\mathbb{F}} Y)P(\mathbf{m}_1, \mathbf{m}_2)^T$$
(2)

for $\mathbf{m}_1 \otimes \mathbf{m}_2 \neq \mathbf{0}$ and $\mathbf{s}_1 \otimes \mathbf{s}_2 \neq \mathbf{0}$, and $X \otimes Y := 0$ otherwise.

Let $\mathbf{0} \neq \mathbf{m} = (m_1, m_2) \in \mathbb{N}^2$. For any $1 \leq i \leq 2$ with $m_i > 0$, and any $1 \leq k \leq m_i$, define a matrix $Y_{i,k}^{\mathbf{m}} = (y_1, y_2, \cdots, y_{|\mathbf{m}|}) \in M_{\mathbf{e}_i \times \mathbf{m}}(A)$ by

$$y_j = \begin{cases} e_i, & j = k + \delta_{i2}m_1, \\ 0, & \text{otherwise,} \end{cases}$$

where δ_{i2} is the Kronecker symbol. Let $X_{i,k}^{\mathbf{m}} = (Y_{i,k}^{\mathbf{m}})^T$, the transposed matrix of $Y_{i,k}^{\mathbf{m}}$. Then $X_{i,k}^{\mathbf{m}} \in M_{\mathbf{m} \times \mathbf{e}_i}(A)$. We have

$$Y_{i,k}^{\mathbf{m}} X_{i',k'}^{\mathbf{m}} = \begin{cases} e_i = E_{\mathbf{e}_i}, & (i,k) = (i',k'), \\ 0, & \text{otherwise,} \end{cases}$$

and $\sum_{i=1}^{2} \sum_{1 \leq k \leq m_i} X_{i,k}^{\mathbf{m}} Y_{i,k}^{\mathbf{m}} = \sum_{1 \leq i \leq 2, m_i > 0} \sum_{k=1}^{m_i} X_{i,k}^{\mathbf{m}} Y_{i,k}^{\mathbf{m}} = E_{\mathbf{m}}.$

For any $1 \leq i, j \leq 2$, let $a_{1,i,j} = a_{i,1,j} = a_{i,j,1} = E_{\mathbf{e}_i \otimes \mathbf{e}_j} \in M_{\mathbf{e}_i \otimes \mathbf{e}_j}(A)$. Note that $\mathbf{e}_1 \otimes \mathbf{e}_i \otimes \mathbf{e}_j = \mathbf{e}_i \otimes \mathbf{e}_1 \otimes \mathbf{e}_j = \mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_1 = \mathbf{c}_{ij}$ and $\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2 = (mn, m + n^2)$.

From now on, we assume that $a_{2,2,2}$ is an invertible element in $M_{\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2}(A)$.

Lemma 2.2. Let $1 \leq i, j, l, i', j', l' \leq 2$, $x \in e_{i'}Ae_i$, $y \in e_{j'}Ae_j$ and $z \in e_{l'}Ae_l$. Then $(x \otimes (y \otimes z))a_{i,j,l} = a_{i',j',l'}((x \otimes y) \otimes z)$.

Proof. If $(i', j', l') \neq (i, j, l)$, then $(x \otimes (y \otimes z)) = ((x \otimes y) \otimes z) = 0$ since $e_1Ae_2 = e_2Ae_1 = 0$, so $(x \otimes (y \otimes z))a_{i,j,l} = a_{i',j',l'}((x \otimes y) \otimes z)$. If (i', j', l') = (i, j, l), then $x = \alpha e_i, y = \beta e_j$ and $z = \gamma e_l$ for some $\alpha, \beta, \gamma \in \mathbb{F}$. In this case, $(x \otimes (y \otimes z)) = ((x \otimes y) \otimes z) = \alpha \beta \gamma E_{\mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_l}$ by [2, Corollary 3.6]. Thus $(x \otimes (y \otimes z))a_{i,j,l} = a_{i,j,l}((x \otimes y) \otimes z)$.

Now let us consider the equality Ass(4) in [2]:

$$\sum_{j=1}^{2} \sum_{1 \leqslant k \leqslant c_{i_{2}i_{3}j}} (e_{i_{1}} \otimes a_{i_{2},i_{3},i_{4}} (X_{j,k}^{\mathbf{c}_{i_{2}i_{3}}} \otimes e_{i_{4}})) a_{i_{1},j,i_{4}} ((e_{i_{1}} \otimes Y_{j,k}^{\mathbf{c}_{i_{2}i_{3}}}) a_{i_{1},i_{2},i_{3}} \otimes e_{i_{4}})$$

$$= \sum_{j,j'=1}^{2} \sum_{1 \leqslant k \leqslant c_{i_{3}i_{4}j}} \sum_{1 \leqslant k' \leqslant c_{i_{1}i_{2}j'}} (e_{i_{1}} \otimes (e_{i_{2}} \otimes X_{j,k}^{\mathbf{c}_{i_{3}i_{4}}})) a_{i_{1},i_{2},j} (X_{j',k'}^{\mathbf{c}_{i_{1}i_{2}}} \otimes Y_{j,k}^{\mathbf{c}_{i_{3}i_{4}}})$$

$$\cdot a_{j',i_{3},i_{4}} ((Y_{j',k'}^{\mathbf{c}_{i_{1}i_{2}}} \otimes e_{i_{3}}) \otimes e_{i_{4}}), \qquad (3)$$

where $1 \le i_1, i_2, i_3, i_4 \le 2$.

Lemma 2.3. If one of i_1, i_2, i_3, i_4 is equal to 1, then the equality (3) holds.

Proof. Follows from a straightforward verification.

When $i_1 = i_2 = i_3 = i_4 = 2$, the equality (3) becomes

$$\sum_{j=1}^{2} \sum_{1 \leqslant k \leqslant c_{22j}} (e_2 \otimes a_{2,2,2} (X_{j,k}^{\mathbf{c}_{22}} \otimes e_2)) a_{2,j,2} ((e_2 \otimes Y_{j,k}^{\mathbf{c}_{22}}) a_{2,2,2} \otimes e_2)$$

$$= \sum_{j,j'=1}^{2} \sum_{1 \leqslant k \leqslant c_{22j}} \sum_{1 \leqslant k' \leqslant c_{22j'}} (e_2 \otimes (e_2 \otimes X_{j,k}^{\mathbf{c}_{22}})) a_{2,2,j} (X_{j',k'}^{\mathbf{c}_{22}} \otimes Y_{j,k}^{\mathbf{c}_{22}})$$

$$\cdot a_{j',2,2} ((Y_{j',k'}^{\mathbf{c}_{22}} \otimes e_2) \otimes e_2). \tag{4}$$

To simplify the equality (4), we use the 6-*j* symbols of semisimple tensor categories. Let $H_k^{ij} = \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_k, \mathbf{e}_i \otimes \mathbf{e}_j) = M_{\mathbf{c}_{ij} \times \mathbf{e}_k}(A)$ be the multiplicity spaces. Then $\dim(H_k^{ij}) = c_{ijk}$, and the decomposition of the tensor product of simple objects can be described as follows:

$$\mathbf{e}_i \otimes \mathbf{e}_j \cong \bigoplus_{k=1}^2 \dim(H_k^{ij}) \mathbf{e}_k.$$

Hence we have

$$(\mathbf{e}_i \otimes \mathbf{e}_j) \otimes \mathbf{e}_l \cong \bigoplus_{s=1}^2 \bigoplus_{k=1}^2 \dim(H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij}) \mathbf{e}_s$$

and

$$\mathbf{e}_i \otimes (\mathbf{e}_j \otimes \mathbf{e}_l) \cong \bigoplus_{s=1}^2 \bigoplus_{k=1}^2 \dim(H_s^{ik} \otimes_{\mathbb{F}} H_k^{jl}) \mathbf{e}_s.$$

Lemma 2.4. For any $1 \leq i, j, k, l, s \leq 2$, there is a linear injection

$$\sigma: H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} \to \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_s, (\mathbf{e}_i \otimes \mathbf{e}_j) \otimes \mathbf{e}_l), \ f \otimes_{\mathbb{F}} g \mapsto (g \otimes \operatorname{id}_{\mathbf{e}_l})f,$$

which induces a linear isomorphism:

$$\oplus_{k=1}^{2} H_{s}^{kl} \otimes_{\mathbb{F}} H_{k}^{ij} \cong \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_{s}, (\mathbf{e}_{i} \otimes \mathbf{e}_{j}) \otimes \mathbf{e}_{l}).$$

Proof. If $H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} = 0$, then σ is obviously injective. If $H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} \neq 0$, then H_s^{kl} has an \mathbb{F} -basis $\{X_{s,p}^{c_{kl}} | 1 \leq p \leq c_{kls}\}$, and H_k^{ij} has an \mathbb{F} -basis $\{X_{k,q}^{c_{ij}} | 1 \leq q \leq c_{ijk}\}$.

We first consider the case of l = 1. In this case, $(\mathbf{e}_i \otimes \mathbf{e}_j) \otimes \mathbf{e}_l = \mathbf{e}_i \otimes \mathbf{e}_j = \mathbf{c}_{ij}$. If $s \neq k$, then $H_s^{k_1} = 0$. Hence σ is injective. If s = k, then $c_{k_1k} = 1$ and $(X_{k,q}^{\mathbf{c}_{ij}} \otimes \mathrm{id}_{\mathbf{e}_1}) X_{k,1}^{\mathbf{c}_{k_1}} = (X_{k,q}^{\mathbf{c}_{ij}} \otimes e_1) e_k = X_{k,q}^{\mathbf{c}_{ij}}$ for all $1 \leq q \leq c_{ijk}$. Thus, σ is bijective, and the lemma holds in this case.

Now suppose that l = 2. First let i = j = 1. Then $(\mathbf{e}_i \otimes \mathbf{e}_j) \otimes \mathbf{e}_l = \mathbf{e}_2$. In this case, if k = 2 or s = k = 1, then $H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} = 0$; if k = 1 and s = 2, then $c_{122} = c_{111} = 1$ and $(X_{1,1}^{c_{11}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{2,1}^{c_{12}} = (e_1 \otimes e_2) e_2 = X_{2,1}^{c_{12}}$. Thus, the lemma holds for l = 2 and i = j = 1. Next, let $i \neq j$. Then $\mathbf{e}_i \otimes \mathbf{e}_j = \mathbf{c}_{ij} = \mathbf{e}_2$ and $(\mathbf{e}_i \otimes \mathbf{e}_j) \otimes \mathbf{e}_l = \mathbf{e}_2 \otimes \mathbf{e}_2 = \mathbf{c}_{22}$. Note that $c_{221} = m$ and $c_{222} = n$. If k = 1, then $H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} = 0$. If k = 2 and s = 1, then by [2, Corollary 3.6], $(X_{2,1}^{c_{ij}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{1,p}^{c_{22}} = (e_2 \otimes e_2) X_{1,p}^{c_{22}} = E_{\mathbf{c}_{22}} X_{1,p}^{c_{22}} = X_{2,p}^{c_{22}}$ for all $1 \leq p \leq m$. If s = k = 2, then similarly, $(X_{2,1}^{c_{ij}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{2,p}^{c_{22}} = X_{2,p}^{c_{22}}$ for all $1 \leq p \leq m$. Thus, the lemma holds for l = 2 and $i \neq j$. Finally, let i = j = 2. Then $\mathbf{e}_i \otimes \mathbf{e}_j = \mathbf{c}_{22}$ and $\mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_l = \mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2 = mn\mathbf{e}_1 \oplus (m + n^2)\mathbf{e}_2$. If s = k = 1, then $H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} = 0$. If s = 1 and k = 2, then $(X_{2,q}^{c_{22}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{1,p}^{c_{22}} = X_{1,m(q-1)+p}^{\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2}$ and $\mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_i = \mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2 = mn\mathbf{e}_1 \oplus (m + n^2)\mathbf{e}_2$. If s = k = 1, then $H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} = 0$. If s = 1 and k = 2, then $(X_{2,q}^{c_{22}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{1,p}^{c_{22}} = X_{1,m(q-1)+p}^{\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2}$ and k = 1, then $(X_{1,q}^{c_{22}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{2,1}^{c_{12}} = X_{2,q}^{\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2}$ for all $1 \leq q \leq m$. If s = 2 and k = 1, then $(X_{1,q}^{c_{22}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{2,1}^{c_{12}} = X_{2,q}^{\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2}$ for all $1 \leq q \leq m$. If s = 2 and k = 2, then $(X_{2,q}^{c_{22}} \otimes \mathrm{id}_{\mathbf{e}_2}) X_{2,p}^{c_{22}} = X_{2,m+n(q-1)+p}^{c_{2}}$ for all $1 \leq p, q \leq m$. Thus, the lemma holds for l = i = j = s = 2. This co

Lemma 2.5. For any $1 \leq i, j, k, l, s \leq 2$, there is a linear injection

$$H_s^{ik} \otimes_{\mathbb{F}} H_k^{jl} \to \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_s, \mathbf{e}_i \otimes (\mathbf{e}_j \otimes \mathbf{e}_l)), \ f \otimes_{\mathbb{F}} g \mapsto (\operatorname{id}_{\mathbf{e}_i} \otimes g)f$$

which induces a linear isomorphism

$$\oplus_{k=1}^{2} H_{s}^{ik} \otimes_{\mathbb{F}} H_{k}^{jl} \cong \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_{s}, \mathbf{e}_{i} \otimes (\mathbf{e}_{j} \otimes \mathbf{e}_{l})).$$

Proof. Similar to the proof of Lemma 2.4.

Let $1 \leq i, j, l, s \leq 2$. Then the associator $a_{i,j,l} : (\mathbf{e}_i \otimes \mathbf{e}_j) \otimes \mathbf{e}_l \to \mathbf{e}_i \otimes (\mathbf{e}_j \otimes \mathbf{e}_l)$ induces a k-linear isomorphism

$$\phi_{i,j}^{l,s}$$
: Hom _{\mathcal{C}} ($\mathbf{e}_s, (\mathbf{e}_i \otimes \mathbf{e}_j) \otimes \mathbf{e}_l$)) \rightarrow Hom _{\mathcal{C}} ($\mathbf{e}_s, \mathbf{e}_i \otimes (\mathbf{e}_j \otimes \mathbf{e}_l)$), $f \mapsto a_{i,j,l}f$

We may regard the isomorphisms in Lemmas 2.4 and 2.5 as identity maps. Then the above isomorphism can be regarded as the following isomorphism

$$\phi_{i,j}^{l,s}:\oplus_{k=1}^2 H_s^{kl}\otimes_{\mathbb{F}} H_k^{ij}\to\oplus_{t=1}^2 H_s^{it}\otimes_{\mathbb{F}} H_t^{jl}.$$

In this case, the 6-*j* symbols are defined to be the component maps of $\phi_{i,j}^{l,s}$:

$$\left\{\begin{array}{ccc}i & j & k\\ l & s & t\end{array}\right\}: H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij} \to H_s^{it} \otimes_{\mathbb{F}} H_t^{jl},$$

where $1 \leq k, t \leq 2$. Note that $\{X_{s,p}^{\mathbf{c}_{kl}} | 1 \leq p \leq c_{kls}\}, \{X_{k,q}^{\mathbf{c}_{ij}} | 1 \leq q \leq c_{ijk}\}, \{X_{s,d}^{\mathbf{c}_{ij}} | 1 \leq d \leq c_{its}\}$ and $\{X_{t,r}^{\mathbf{c}_{jl}} | 1 \leq r \leq c_{jlt}\}$ are the bases of $H_s^{kl}, H_k^{ij}, H_s^{it}$ and H_t^{jl} , respectively. Under the identification described above, one can identify $X_{s,p}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,q}^{\mathbf{c}_{ij}}$ with $X_{s,u}^{\mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_l}$, and $X_{s,d}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,r}^{\mathbf{c}_{jl}}$ with $X_{s,v}^{\mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_l}$, respectively, where $u = \sum_{1 \leq w \leq k-1} c_{wls}c_{ijw} + (q-1)c_{kls} + p$ and $v = \sum_{1 \leq w \leq t-1} c_{iws}c_{jlw} + (r-1)c_{its} + d$.

Let $B = \{B^1, B^2\}$ and $C = \{C^1, C^2\}$ be the (ordered) bases of $\bigoplus_{k=1}^2 H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij}$ and $\bigoplus_{t=1}^2 H_s^{it} \otimes_{\mathbb{F}} H_t^{jl}$ given by

$$B^{k} = \left\{ \begin{array}{c} X_{s,1}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,1}^{\mathbf{c}_{ij}}, X_{s,2}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,1}^{\mathbf{c}_{ij}}, \cdots, X_{s,c_{kls}}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,1}^{\mathbf{c}_{ij}}, \\ X_{s,1}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,2}^{\mathbf{c}_{ij}}, X_{s,2}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,2}^{\mathbf{c}_{ij}}, \cdots, X_{s,c_{kls}}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,2}^{\mathbf{c}_{ij}}, \\ & \dots \dots \dots , \\ X_{s,1}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,c_{ijk}}^{\mathbf{c}_{ij}}, X_{s,2}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,c_{ijk}}^{\mathbf{c}_{ij}}, \cdots, X_{s,c_{kls}}^{\mathbf{c}_{kl}} \otimes_{\mathbb{F}} X_{k,c_{ijk}}^{\mathbf{c}_{ij}} \end{array} \right\}, \ k = 1, 2,$$

and

$$C^{t} = \left\{ \begin{array}{c} X_{s,1}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,1}^{\mathbf{c}_{jl}}, X_{s,2}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,1}^{\mathbf{c}_{it}}, \cdots, X_{s,c_{its}}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,1}^{\mathbf{c}_{jl}}, \\ X_{s,1}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,2}^{\mathbf{c}_{jl}}, X_{s,2}^{\mathbf{c}_{jl}} \otimes_{\mathbb{F}} X_{t,2}^{\mathbf{c}_{jl}}, \cdots, X_{s,c_{its}}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,2}^{\mathbf{c}_{jl}}, \\ & \dots \dots \dots , \\ X_{s,1}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,c_{jlt}}^{\mathbf{c}_{jl}}, X_{s,2}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,c_{jlt}}^{\mathbf{c}_{jl}}, \cdots, X_{s,c_{its}}^{\mathbf{c}_{it}} \otimes_{\mathbb{F}} X_{t,c_{jlt}}^{\mathbf{c}_{jl}} \end{array} \right\}, \ t = 1, 2.$$

Under the identification as before, both B and C are the canonical bases of

$$\operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_s, \mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_l) = M_{\mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_l \times \mathbf{e}_s}(A).$$

With respect to two bases B and C, the matrix of the linear isomorphism $\phi_{i,j}^{l,s}$ is A_s , i.e.,

$$\phi_{i,j}^{l,s}(X_{s,1}^{\mathbf{c}_{1l}} \otimes_{\mathbb{F}} X_{1,1}^{\mathbf{c}_{ij}}, \cdots, X_{s,c_{2ls}}^{\mathbf{c}_{2l}} \otimes_{\mathbb{F}} X_{2,c_{ij2}}^{\mathbf{c}_{ij}}) \\
= (X_{s,1}^{\mathbf{c}_{i1}} \otimes_{\mathbb{F}} X_{1,1}^{\mathbf{c}_{jl}}, \cdots, X_{s,c_{i2s}}^{\mathbf{c}_{i2}} \otimes_{\mathbb{F}} X_{2,c_{jl2}}^{\mathbf{c}_{jl}})A_s,$$
(5)

where A_s is a square matrix over $\mathbb{F}e_s$ such that $a_{i,j,l} = \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix}$. In the following, we call B^k and C^t the standard bases of vector space $H_s^{kl} \otimes_{\mathbb{F}} H_k^{ij}$ and $H_s^{it} \otimes_{\mathbb{F}} H_t^{jl}$, respectively.

Proposition 2.6. The commutative pentagon diagram is equivalent to the Biedenharn-Elliot identity. That is, the equality (3) is equivalent to

$$\sum_{j=1}^{2} (\operatorname{id}_{j_{0}}^{j_{1}j_{7}} \otimes_{\mathbb{F}} \left\{ \begin{array}{ccc} j_{2} & j_{3} & j \\ j_{4} & j_{7} & j_{8} \end{array} \right\}) \left(\left\{ \begin{array}{ccc} j_{1} & j & j_{6} \\ j_{4} & j_{0} & j_{7} \end{array} \right\} \otimes_{\mathbb{F}} \operatorname{id}_{j}^{j_{2}j_{3}}) \left(\operatorname{id}_{j_{0}}^{j_{6}j_{4}} \otimes_{\mathbb{F}} \left\{ \begin{array}{ccc} j_{1} & j_{2} & j_{5} \\ j_{3} & j_{6} & j \end{array} \right\} \right) \\ = \left(\left\{ \begin{array}{ccc} j_{1} & j_{2} & j_{5} \\ j_{8} & j_{0} & j_{7} \end{array} \right\} \otimes_{\mathbb{F}} \operatorname{id}_{j_{8}}^{j_{3}j_{4}}) P_{23} \left(\left\{ \begin{array}{ccc} j_{5} & j_{3} & j_{6} \\ j_{4} & j_{0} & j_{8} \end{array} \right\} \otimes_{\mathbb{F}} \operatorname{id}_{j_{5}}^{j_{1}j_{2}} \right) \\ \end{array} \right\}$$

for all $j_i = 1, 2, 0 \leq i \leq 8$, where id_k^{jl} denotes the identity map on H_k^{jl} , P_{23} denotes the flip map of the second and the third tensor factors. Both sides of this equality are linear maps from $H_{j_0}^{j_6j_4} \otimes_{\mathbb{F}} H_{j_6}^{j_5,j_3} \otimes_{\mathbb{F}} H_{j_5}^{j_1,j_2}$ to $H_{j_0}^{j_1,j_7} \otimes_{\mathbb{F}} H_{j_7}^{j_2,j_8} \otimes_{\mathbb{F}} H_{j_8}^{j_3,j_4}$.

Proof. The equation (3) means that the diagram

$$\begin{array}{c} ((\mathbf{e}_{j_1} \otimes \mathbf{e}_{j_2}) \otimes \mathbf{e}_{j_3}) \otimes \mathbf{e}_{j_4} \xrightarrow{a_{\mathbf{c}_{j_1j_2}, \mathbf{e}_{j_3}, \mathbf{e}_{j_4}}} (\mathbf{e}_{j_1} \otimes \mathbf{e}_{j_2}) \otimes (\mathbf{e}_{j_3} \otimes \mathbf{e}_{j_4}) \\ a_{j_1, j_2, j_3} \otimes \mathrm{id}_{j_4} \downarrow & & \\ (\mathbf{e}_{j_1} \otimes (\mathbf{e}_{j_2} \otimes \mathbf{e}_{j_3})) \otimes \mathbf{e}_{j_4} & & a_{\mathbf{e}_{j_1}, \mathbf{e}_{j_2}, \mathbf{c}_{j_3j_4}} \\ a_{\mathbf{e}_{j_1}, \mathbf{c}_{j_2j_3}, \mathbf{e}_{j_4}} \downarrow & & \downarrow \\ \mathbf{e}_{j_1} \otimes ((\mathbf{e}_{j_2} \otimes \mathbf{e}_{j_3}) \otimes \mathbf{e}_{j_4}) \xrightarrow{\mathrm{id}_{j_1} \otimes a_{\mathbf{e}_{j_2}, \mathbf{e}_{j_3}, \mathbf{e}_{j_4}}} \mathbf{e}_{j_1} \otimes (\mathbf{e}_{j_2} \otimes (\mathbf{e}_{j_3} \otimes \mathbf{e}_{j_4})) \end{array}$$

commutes for any $j_1, j_2, j_3, j_4 \in \{1, 2\}$, where $id_j = id_{\mathbf{e}_j} = e_j$. Since C is a semisimple category, the above commutative diagram is equivalent to the following commutative diagram

$$\begin{array}{c|c} \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_{j_{0}}, ((\mathbf{e}_{j_{1}} \otimes \mathbf{e}_{j_{2}}) \otimes \mathbf{e}_{j_{3}}) \otimes \mathbf{e}_{j_{4}}) & \xrightarrow{a_{*}} \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_{j_{0}}, (\mathbf{e}_{j_{1}} \otimes \mathbf{e}_{j_{2}}) \otimes (\mathbf{e}_{j_{3}} \otimes \mathbf{e}_{j_{4}})) \\ & (a \otimes e_{j_{4}})_{*} \downarrow \\ \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_{j_{0}}, (\mathbf{e}_{j_{1}} \otimes (\mathbf{e}_{j_{2}} \otimes \mathbf{e}_{j_{3}})) \otimes \mathbf{e}_{j_{4}}) & a_{*} \downarrow \\ & a_{*} \downarrow \\ \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_{j_{0}}, \mathbf{e}_{j_{1}} \otimes ((\mathbf{e}_{j_{2}} \otimes \mathbf{e}_{j_{3}}) \otimes \mathbf{e}_{j_{4}})) \xrightarrow{(e_{1} \otimes a)_{*}} \operatorname{Hom}_{\mathcal{C}}(\mathbf{e}_{j_{0}}, \mathbf{e}_{j_{1}} \otimes (\mathbf{e}_{j_{2}} \otimes (\mathbf{e}_{j_{3}} \otimes \mathbf{e}_{j_{4}}))) \end{array}$$

where $j_0 \in \{1, 2\}$, and we omit the subscripts of a. Then by Lemmas 2.4-2.5 and the above identification, we conclude that the last commutative diagram is equivalent to the Biedenharn-Elliot identity.

By Proposition 2.6, the equation (4) is equivalent to

$$\sum_{j=1}^{2} (\operatorname{id}_{j_{0}}^{2j_{7}} \otimes_{\mathbb{F}} \left\{ \begin{array}{ccc} 2 & 2 & j \\ 2 & j_{7} & j_{8} \end{array} \right\}) \left(\left\{ \begin{array}{ccc} 2 & j & j_{6} \\ 2 & j_{0} & j_{7} \end{array} \right\} \otimes_{\mathbb{F}} \operatorname{id}_{j}^{22}\right) \left(\operatorname{id}_{j_{0}}^{j_{6}2} \otimes_{\mathbb{F}} \left\{ \begin{array}{ccc} 2 & 2 & j_{5} \\ 2 & j_{6} & j \end{array} \right\} \right) \right) = \left(\left\{ \begin{array}{ccc} 2 & 2 & j_{5} \\ j_{8} & j_{0} & j_{7} \end{array} \right\} \otimes_{\mathbb{F}} \operatorname{id}_{j_{8}}^{22}\right) P_{23} \left(\left\{ \begin{array}{ccc} j_{5} & 2 & j_{6} \\ 2 & j_{0} & j_{8} \end{array} \right\} \otimes_{\mathbb{F}} \operatorname{id}_{j_{5}}^{22}\right),$$
(6)

where $j_0, j_5, j_6, j_7, j_8 \in \{1, 2\}$.

3. Categorization of \mathbb{Z}_+ -ring

In this section, we study the categorization of \mathbb{Z}_+ -ring R as stated in last section.

Let
$$a_{2,2,2} = \begin{pmatrix} \Lambda_1 e_1 & 0 \\ 0 & \Lambda_2 e_2 \end{pmatrix}$$
 for some matrices $\Lambda_1 \in M_{mn}(\mathbb{F}), \Lambda_2 \in M_{m+n^2}(\mathbb{F}),$
and $\Lambda_2 = \begin{pmatrix} \Lambda_{22} & \Lambda_{23} \\ \Lambda_{32} & \Lambda_{33} \end{pmatrix}$ with $\Lambda_{22} \in M_m(\mathbb{F})$ and $\Lambda_{33} \in M_{n^2}(\mathbb{F})$. Since $a_{1,i,j} = a_{i,1,j} = a_{i,j,1} = E_{\mathbf{e}_i \otimes \mathbf{e}_j}$, with respect to the standard bases B and C , one can regard the above 6- j symbols as matrices as follow:

$$\begin{cases} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \\ \end{cases} = \begin{cases} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 2 & 2 & 2 \\ \end{cases} = \begin{cases} 1 & 2 & 2 \\ 1 & 2 & 2 \\ 2 & 1 & 1 \\ \end{cases} = \begin{cases} 2 & 1 & 2 \\ 2 & 1 & 2 \\ 2 & 1 & 2 \\ \end{cases} = \begin{cases} 2 & 1 & 2 \\ 2 & 1 & 2 \\ 2 & 1 & 2 \\ \end{cases} = \begin{cases} 2 & 2 & 1 \\ 1 & 1 & 2 \\ 1 & 2 & 2 \\ 2 & 2 & 2 \\ \end{cases} = \begin{cases} 2 & 1 & 2 \\ 2 & 2 & 2 \\ 2 & 2 & 2 \\ \end{cases} = \begin{cases} 2 & 2 & 2 \\ 1 & 2 & 2 \\ 1 & 2 & 2 \\ \end{cases} = I_m,$$
$$\begin{cases} 1 & 2 & 2 \\ 2 & 2 & 2 \\ 2 & 2 & 2 \\ 2 & 1 & 2 \\ \end{cases} = \Lambda_1, \begin{cases} 2 & 2 & 1 \\ 2 & 2 & 1 \\ 2 & 2 & 1 \\ \end{cases} = \Lambda_{22}, \begin{cases} 2 & 2 & 2 \\ 2 & 2 & 1 \\ 2 & 2 & 1 \\ \end{cases} = \Lambda_{23},$$
$$\begin{cases} 2 & 2 & 1 \\ 2 & 2 & 1 \\ 2 & 2 & 2 \\ \end{cases} = \Lambda_{33}$$

and the rest $\left\{ \begin{array}{cc} i & j & k \\ l & m & n \end{array} \right\} = 0.$

In the following, we regard each component map of Equation (6) as a matrix with respect to the standard bases. Since P_{23} is a flip map from $H_{j_0}^{j_5 j_8} \otimes H_{j_8}^{j_3 j_4} \otimes H_{j_5}^{j_1 j_2} \longrightarrow H_{j_0}^{j_5 j_8} \otimes H_{j_8}^{j_1 j_2} \otimes H_{j_8}^{j_3 j_4}$, $P_{23} = I_a \otimes_{\mathbb{F}} I_{[c,b]}$, where $a = c_{j_5 j_8 j_0}$, $b = c_{j_3 j_4 j_8}$, $c = c_{j_1 j_2 j_5}$, $I_{i,j}^{[c,b]} = \varepsilon_{c,i}^T \varepsilon_{b,j} \in M_{c \times b}(\mathbb{F})$ and

$$I_{[c,b]} = \begin{pmatrix} I_{1,1}^{[c,b]} & I_{2,1}^{[c,b]} & \cdots & I_{c,1}^{[c,b]} \\ I_{1,2}^{[c,b]} & I_{2,2}^{[c,b]} & \cdots & I_{c,2}^{[c,b]} \\ \cdots & \cdots & \cdots & \cdots \\ I_{1,b}^{[c,b]} & I_{2,b}^{[c,b]} & \cdots & I_{c,b}^{[c,b]} \end{pmatrix} \in M_{bc \times bc}(\mathbb{F})$$

In particularly, if b = c = n then $I_{[c,b]} = I_{[n]}$.

In order to make C into a tensor category, we need to investigate when the equation (6) holds. We will consider it in three cases: (1) m > 0 and n = 0, (2) m = 0 and n > 0, (3) m > 0 and n > 0, respectively.

3.1.
$$m > 0$$
 and $n = 0$. In this case, $\mathbf{e}_2 \otimes \mathbf{e}_2 = \mathbf{c}_{22} = (m, 0) = m\mathbf{e}_1$ and $\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2 = (0, m) = m\mathbf{e}_2$. Moreover, $\begin{cases} 1 & 2 & 2 \\ 2 & 2 & 2 \end{cases} = \begin{cases} 2 & 1 & 2 \\ 2 & 2 & 2 \end{cases} = \begin{cases} 2 & 1 & 2 \\ 2 & 2 & 2 \end{cases} = \begin{cases} 2 & 2 & 2 \\ 1 & 2 & 2 \end{cases} = \begin{cases} 2 & 2 & 2 \\ 1 & 2 & 2 \end{cases}$

$$\left\{ \begin{array}{cc} 2 & 2 & 2 \\ 2 & 1 & 2 \end{array} \right\} = \left\{ \begin{array}{cc} 2 & 2 & 2 \\ 2 & 2 & 2 \end{array} \right\} = \left\{ \begin{array}{cc} 2 & 2 & 2 \\ 2 & 2 & 1 \end{array} \right\} = \left\{ \begin{array}{cc} 2 & 2 & 1 \\ 2 & 2 & 1 \end{array} \right\} = \left\{ \begin{array}{cc} 2 & 2 & 1 \\ 2 & 2 & 2 \end{array} \right\} = 0, \ a_{2,2,2} =$$

 $\Lambda_{22}e_2$ and $\Lambda_{22} = \left\{ \begin{array}{ccc} 2 & 2 & 1 \\ 2 & 2 & 1 \end{array} \right\}$. Hence the equation (6) is equivalent to

$$I_m \otimes_{\mathbb{F}} \Lambda_{22}^2 = P_{23}.$$

Let $a_{222} = (\alpha_{ij})e_2$ and $\beta_{ij} = \sum_{k=1}^m \alpha_{ik}\alpha_{kj}$ for all $1 \leq i, j \leq m$. That is, $(\beta_{ij}) = \Lambda_{22}^2 \in M_m(\mathbb{F})$. Thus, the above equation becomes

$$\begin{pmatrix} \beta_{11}I_m & \beta_{12}I_m & \cdots & \beta_{1m}I_m \\ \beta_{21}I_m & \beta_{22}I_m & \cdots & \beta_{2m}I_m \\ \cdots & \cdots & \cdots & \cdots \\ \beta_{m1}I_m & \beta_{m2}I_m & \cdots & \beta_{mm}I_m \end{pmatrix} = \begin{pmatrix} I_{1,1} & I_{2,1} & \cdots & I_{m,1} \\ I_{1,2} & I_{2,2} & \cdots & I_{m,2} \\ \cdots & \cdots & \cdots \\ I_{1,m} & I_{2,m} & \cdots & I_{m,m} \end{pmatrix}$$

where $I_{i,j} = I_{i,j}^{[m]}$. If m > 1, it is easy to see that there is no invertible matrix (β_{ij}) that satisfies the above equation. Now assume m = 1. Then $a_{2,2,2} = \alpha e_2$ for some $\alpha \in \mathbb{F}^{\times}$. In this case, it follows from the above computation that the equation (6) holds if and only if $\alpha^2 = 1$, which is equivalent to that $a_{2,2,2} = e_2$ or $a_{2,2,2} = -e_2$.

Let $a_{2,2,2} = e_2$. Then by Lemmas 2.2, 2.3 and the above discussion, the family $\{a_{i,j,l} \in M_{\mathbf{e}_i \otimes \mathbf{e}_j \otimes \mathbf{e}_l}(A) | 1 \leq i, j, l \leq 2\}$ satisfies the conditions Ass (1)-(4). For any $\mathbf{m} = (m_1, m_2), \mathbf{s} = (s_1, s_2), \mathbf{t} = (t_1, t_2) \in \mathrm{Ob}(\mathcal{C})$, define a matrix $a_{\mathbf{m},\mathbf{s},\mathbf{t}} \in M_{\mathbf{m} \otimes \mathbf{s} \otimes \mathbf{t}}(A)$ by $a_{\mathbf{m},\mathbf{s},\mathbf{t}} = E_{\mathbf{0}} = 0$ when $\mathbf{m} \otimes \mathbf{s} \otimes \mathbf{t} = \mathbf{0}$, and

$$=\sum_{i,j,l=11\leqslant k_1\leqslant m_i 1\leqslant k_2\leqslant s_j}^{2}\sum_{1\leqslant k_3\leqslant t_l}\sum_{(X_{i,k_1}^{\mathbf{m}}\otimes (X_{j,k_2}^{\mathbf{s}}\otimes X_{l,k_3}^{\mathbf{t}}))a_{i,j,l}((Y_{i,k_1}^{\mathbf{m}}\otimes Y_{j,k_2}^{\mathbf{s}})\otimes Y_{l,k_3}^{\mathbf{t}})$$

when $\mathbf{m} \otimes \mathbf{s} \otimes \mathbf{t} \neq \mathbf{0}$. Thus, by [2, Theorem 3.16], we have the following proposition.

Lemma 3.1. Assume m = 1. Then with the tensor products (1) and (2), C is a tensor category over \mathbb{F} , where \mathbf{e}_1 is the unit object, the associativity constraint a is induced by $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ as above, the left and right unit constraints are the identities. Denote the tensor category by C_+ .

Let $a'_{2,2,2} = -e_2$ and $a'_{1,i,j} = a'_{i,1,j} = a'_{i,j,1} = E_{\mathbf{e}_i \otimes \mathbf{e}_j} \in M_{\mathbf{e}_i \otimes \mathbf{e}_j}(A)$ for any $1 \leq i, j \leq 2$. Then similarly, we have the following proposition.

Lemma 3.2. Assume m = 1. Then with the tensor products (1) and (2), C is a tensor category over \mathbb{F} , where \mathbf{e}_1 is the unit object, the associativity constraint a' is induced by $\{a'_{i,j,l}|1 \leq i, j, l \leq 2\}$ similarly as above, the left and right unit constraints are the identities. Denote the tensor category by C_{-} .

Note that $\mathcal{C}_+ = \mathcal{C}_-$ in case char(\mathbb{F}) = 2. In case char(\mathbb{F}) \neq 2, we have the following lemma.

Lemma 3.3. If $char(\mathbb{F}) \neq 2$, then \mathcal{C}_+ and \mathcal{C}_- are not tensor equivalent.

Proof. We first notice that if σ is a permutation of $\{1,2\}$ such that the additive group endmorphism $R \to R$, $r_i \mapsto r_{\sigma(i)}$, is a ring automorphism, then σ must be the

identity permutation of $\{1, 2\}$. Now assume that \mathcal{C}_+ and \mathcal{C}_- are tensor equivalent. Then by [2, Theorem 3.22], there exists an $\alpha \in \mathbb{F}^{\times}$ and a family of invertible elements $\{\varphi_{i,j} \in M_{\mathbf{c}_{ij}}(A) | 1 \leq i, j \leq 2\}$ such that $\varphi_{1,i} = \varphi_{i,1} = \alpha E_{\mathbf{e}_i} = \alpha e_i$ and

$$\sum_{t=1}^{2} \sum_{1 \leqslant k \leqslant c_{ijt}} a_{i,j,l} (X_{t,k}^{\mathbf{c}_{ij}} \otimes e_l) \varphi_{t,l} (Y_{t,k}^{\mathbf{c}_{ij}} \varphi_{i,j} \otimes e_l)$$
$$= \sum_{t=1}^{2} \sum_{1 \leqslant k \leqslant c_{jlt}} (e_i \otimes X_{t,k}^{\mathbf{c}_{jl}}) \varphi_{i,t} (e_i \otimes Y_{t,k}^{\mathbf{c}_{jl}} \varphi_{j,l}) a'_{i,j,l}$$

for all $1 \leq i, j, l \leq 2$. Since $\mathbf{c}_{22} = \mathbf{e}_1$, $\varphi_{2,2} = \beta e_1$ for some $\beta \in \mathbb{F}^{\times}$. Putting i = j = l = 2 in the above equation, one obtains

$$e_2(e_1 \otimes e_2)(\alpha e_2)(e_1(\beta e_1) \otimes e_2) = (e_2 \otimes e_1)(\alpha e_2)(e_2 \otimes e_1(\beta e_1))(-e_2),$$

which implies $\alpha\beta e_2 = -\alpha\beta e_2$. This is impossible since char(\mathbb{F}) $\neq 2$.

Summarizing the above discussion, we have the following proposition.

Proposition 3.4. Let \widehat{C} be a semisimple tensor category of rank two over \mathbb{F} , and V a simple object not isomorphic to the unit object **1**. If $V \otimes V \cong m\mathbf{1}$ with m > 0, then m = 1 and \widehat{C} is tensor equivalent to \mathcal{C}_+ or \mathcal{C}_- .

Proof. Follows from [2, Corollary 3.15 and Theorem 3.25] and the above discussion. \Box

3.2.
$$m = 0$$
 and $n > 0$. In this case, $\mathbf{e}_2 \otimes \mathbf{e}_2 = \mathbf{c}_{22} = (0, n) = n\mathbf{e}_2$ and $\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2 = (0, n^2) = n^2 \mathbf{e}_2$. Moreover, $\begin{cases} 1 & 2 & 2 \\ 2 & 1 & 1 \end{cases} = \begin{cases} 2 & 1 & 2 \\ 2 & 1 & 2 \end{cases} = \begin{cases} 2 & 1 & 2 \\ 2 & 1 & 2 \end{cases} = \begin{cases} 2 & 2 & 1 \\ 1 & 1 & 2 \end{cases} = \begin{cases} 2 & 2 & 1 \\ 1 & 1 & 2 \end{cases} = \begin{cases} 2 & 2 & 1 \\ 2 & 2 & 1 \end{cases} = \begin{cases} 2 & 2 & 1 \\ 2 & 2 & 1 \end{cases} = \begin{cases} 2 & 2 & 1 \\ 2 & 2 & 2 \end{cases} = 0, a_{2,2,2} = \Lambda_{33}e_2 \text{ and } \Lambda_{33} = \begin{cases} 2 & 2 & 2 \\ 2 & 2 & 2 \end{cases}$. Thus the equation (6) is equivalent to

$$(I_n \otimes_{\mathbb{F}} \Lambda_{33})(\Lambda_{33} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{33}) = (\Lambda_{33} \otimes_{\mathbb{F}} I_n)P_{23}(\Lambda_{33} \otimes_{\mathbb{F}} I_n).$$
(7)

where $P_{23} = I_n \otimes_{\mathbb{F}} I_{[n]}$. In case n = 1, one can check that $\Lambda_{33} = I_1 \in M_1(\mathbb{F})$ is the unique matrix satisfying the equation (7). In this case, $a_{2,2,2} = e_2$, \mathcal{C} becomes a tensor category, denoted \mathcal{C}_1 , with the associativity constraint a induced by $\{a_{i,j,l} | 1 \leq i, j, l \leq 2\}$ as before. Thus, we have the following proposition.

Proposition 3.5. Let \widehat{C} be a semisimple tensor category of rank two over \mathbb{F} , and V a simple object not isomorphic to the unit object **1**. If $V \otimes V \cong V$, then \widehat{C} is tensor equivalent to C_1 .

Proof. Follows from [2, Corollary 3.15 and Theorem 3.25] and the above discussion. \Box

For n > 1, put $\Lambda_{33} = I_{[n]}$. Then it is straightforward to verify that the matrix Λ_{33} satisfies the equation (7). In this case, $a_{2,2,2} = I_{[n]}e_2$, and C becomes a tensor

category, denoted by C_n , with the associativity constraint *a* induced by $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ as before. Similarly, putting

$$\Lambda_{33} = \begin{pmatrix} I_{1,1} & I_{2,1} & \cdots & I_{n-1,1} & I_{n,1} \\ I_{2,2} & I_{3,2} & \cdots & I_{n,2} & I_{1,2} \\ \cdots & \cdots & \cdots & \cdots \\ I_{n-1,n-1} & I_{n,n-1} & \cdots & I_{n-3,n-1} & I_{n-2,n-1} \\ I_{n,n} & I_{1,n} & \cdots & I_{n-2,n} & I_{n-1,n} \end{pmatrix}.$$

Let $a'_{2,2,2} = \Lambda_{33}e_2$ and $a'_{1,i,j} = a'_{i,1,j} = a'_{i,j,1} = E_{\mathbf{e}_i \otimes \mathbf{e}_j}$ for $1 \leq i, j \leq 2$. Then \mathcal{C} is also a tensor category, denoted \mathcal{C}'_n , with the associativity constraint a' induced by $\{a'_{i,j,l} | 1 \leq i, j, l \leq 2\}$ as above. Similarly to Lemma 3.3, one can prove the following proposition.

Proposition 3.6. If n > 1, then the tensor categories C_n and C'_n are not tensor equivalent.

Proof. It can be shown by using [2, Theorem 3.22].

3.3. m > 0 and n > 0. In this case, $\mathbf{e}_2 \otimes \mathbf{e}_2 = \mathbf{c}_{22} = (m, n) = m\mathbf{e}_1 + n\mathbf{e}_2$ and $\mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2 = (mn, m + n^2) = mn\mathbf{e}_1 + (m + n^2)\mathbf{e}_2$. Hence the equation (6) holds if and only if the following equations are satisfied:

- (i) $I_m \otimes_{\mathbb{F}} \Lambda_{22}^2 + (I_m \otimes_{\mathbb{F}} \Lambda_{23})(\Lambda_1 \otimes_{\mathbb{F}} I_n)(I_m \otimes_{\mathbb{F}} \Lambda_{32}) = I_{[m]},$
- (ii) $I_m \otimes_{\mathbb{F}} \Lambda_{22} \Lambda_{23} + (I_m \otimes_{\mathbb{F}} \Lambda_{23})(\Lambda_1 \otimes_{\mathbb{F}} I_n)(I_m \otimes_{\mathbb{F}} \Lambda_{33}) = 0,$
- (iii) $I_m \otimes_{\mathbb{F}} \Lambda_{32} \Lambda_{22} + (I_m \otimes_{\mathbb{F}} \Lambda_{33})(\Lambda_1 \otimes_{\mathbb{F}} I_n)(I_m \otimes_{\mathbb{F}} \Lambda_{32}) = 0,$
- (iv) $I_m \otimes_{\mathbb{F}} \Lambda_{32} \Lambda_{23} + (I_m \otimes_{\mathbb{F}} \Lambda_{33}) (\Lambda_1 \otimes_{\mathbb{F}} I_n) (I_m \otimes_{\mathbb{F}} \Lambda_{33})$
 - $= (\Lambda_1 \otimes_{\mathbb{F}} I_n)(I_m \otimes_{\mathbb{F}} I_{[n]})(\Lambda_1 \otimes_{\mathbb{F}} I_n),$
- (v) $\Lambda_1(\Lambda_{22} \otimes_{\mathbb{F}} I_n)\Lambda_1 = (\Lambda_{23} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} I_{[n]})(\Lambda_{32} \otimes_{\mathbb{F}} I_n),$
- (vi) $\Lambda_1(\Lambda_{23} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{32}) = (\Lambda_{22} \otimes_{\mathbb{F}} I_n)I_{[m,n]},$
- (vii) $\Lambda_1(\Lambda_{23} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{33}) = (\Lambda_{23} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} I_{[n]})(\Lambda_{33} \otimes_{\mathbb{F}} I_n),$
- (viii) $(I_n \otimes_{\mathbb{F}} \Lambda_{23})(\Lambda_{32} \otimes_{\mathbb{F}} I_n)\Lambda_1 = I_{[m,n]}^T(\Lambda_{22} \otimes_{\mathbb{F}} I_n),$
- (ix) $I_n \otimes_{\mathbb{F}} \Lambda_{22}^2 + (I_n \otimes_{\mathbb{F}} \Lambda_{23})(\Lambda_{33} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{32}) = 0,$
- (x) $I_n \otimes_{\mathbb{F}} \Lambda_{22} \Lambda_{23} + (I_n \otimes_{\mathbb{F}} \Lambda_{23})(\Lambda_{33} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{33}) = I_{[m,n]}^T (\Lambda_{23} \otimes_{\mathbb{F}} I_n),$
- (xi) $(I_n \otimes_{\mathbb{F}} \Lambda_{33})(\Lambda_{32} \otimes_{\mathbb{F}} I_n)\Lambda_1 = (\Lambda_{33} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} I_{[n]})(\Lambda_{32} \otimes_{\mathbb{F}} I_n),$
- (xii) $I_n \otimes_{\mathbb{F}} \Lambda_{32} \Lambda_{22} + (I_n \otimes_{\mathbb{F}} \Lambda_{33})(\Lambda_{33} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{32}) = (\Lambda_{32} \otimes_{\mathbb{F}} I_n)I_{[m,n]},$
- (xiii) $I_n \otimes_{\mathbb{F}} \Lambda_{32} \Lambda_{23} + (I_n \otimes_{\mathbb{F}} \Lambda_{33})(\Lambda_{33} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{33})$

$$= (\Lambda_{33} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} I_{[n]})(\Lambda_{33} \otimes_{\mathbb{F}} I_n)$$

Assume that m = n = 1. Then $\Lambda_1 = (\alpha)$ and $\Lambda_2 = \begin{pmatrix} \alpha_{22} & \alpha_{23} \\ \alpha_{32} & \alpha_{33} \end{pmatrix}$ for some

 $\alpha, \alpha_{ij} \in \mathbb{F}$ with $\alpha \neq 0$ and $\alpha_{22}\alpha_{33} - \alpha_{23}\alpha_{32} \neq 0$. In this case, a straightforward computation shows that the equations (i)-(xiii) are equivalent to that $\alpha = 1, \alpha_{33} = -\alpha_{22}, \alpha_{23}\alpha_{32} = \alpha_{22}$ and $\alpha_{22}^2 + \alpha_{22} = 1$. Let $f(x) = x^2 + x - 1 \in \mathbb{F}[x]$.

In case char(\mathbb{F}) = 5, the polynomial f(x) has a unique root 2 in \mathbb{F} . Let

$$a_{2,2,2} = \begin{pmatrix} e_1 & 0 & 0\\ 0 & 2e_2 & e_2\\ 0 & 2e_2 & 3e_2 \end{pmatrix}.$$

Then by the above discussion, the equation (4) is satisfied, and hence C becomes a tensor category, denoted by $C_{(5)}$, with the associativity constraint *a* induced by $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ as before. For any $\alpha \in \mathbb{F}^{\times}$, let

$$a_{2,2,2}' = \begin{pmatrix} e_1 & 0 & 0\\ 0 & 2e_2 & \alpha e_2\\ 0 & 2\alpha^{-1}e_2 & 3e_2 \end{pmatrix}$$

and $a'_{1,i,j} = a'_{I,1,j} = a'_{i,j,1} = E_{\mathbf{e}_i \otimes \mathbf{e}_j}$ for $1 \leq i, j \leq 2$. Similarly, \mathcal{C} also becomes a tensor category, denoted $\mathcal{C}_{[\alpha]}$, with the associativity constraint a' induced by $\{a'_{i,j,l} | 1 \leq i, j, l \leq 2\}$ as before.

Proposition 3.7. Assume that $char(\mathbb{F}) = 5$. Let $\widehat{\mathcal{C}}$ be a semisimple tensor category of rank two over \mathbb{F} , and V a simple object not isomorphic to the unit object **1**. If $V \otimes V \cong \mathbf{1} \oplus V$, then $\widehat{\mathcal{C}}$ is tensor equivalent to $\mathcal{C}_{(5)}$.

Proof. By [2, Corollary 3.15 and Theorem 3.25] and the discussion above, there exists a nonzero scale α in \mathbb{F} such that $\widehat{\mathcal{C}}$ is tensor equivalent to $\mathcal{C}_{[\alpha]}$. Let $a_{2,2,2}$ and $a'_{2,2,2}$ be the matrices given above, and let $a'_{1,i,j} = a'_{i,1,j} = a'_{i,j,1} = E_{\mathbf{e}_i \otimes \mathbf{e}_j}$ for $1 \leq i, j \leq 2$. Then the associativity constraints of $\mathcal{C}_{(5)}$ and $\mathcal{C}_{[\alpha]}$ are induced by $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ and $\{a'_{i,j,l}|1 \leq i, j, l \leq 2\}$, respectively. Let $\eta(1,1) = e_1$, $\eta(1,2) = \eta(2,1) = e_2$ and $\eta(2,2) = \begin{pmatrix} \alpha e_1 & 0 \\ 0 & \alpha e_2 \end{pmatrix}$. Then it is straightforward to check that the two families $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ and $\{a'_{i,j,l}|1 \leq i, j, l \leq 2\}$ are equivalent via the family of matrices $\{\eta(i,j) \in M_{\mathbf{c}_{ij}}(A)|1 \leq i, j \leq 2\}$ (see [2, Definition 3.19]). Thus, it follows from [2, Proposition 3.20] that $\mathcal{C}_{(5)}$ is tensor equivalent to $\mathcal{C}_{[\alpha]}$, hence $\widehat{\mathcal{C}}$ is tensor equivalent to $\mathcal{C}_{(5)}$.

In case char(\mathbb{F}) $\neq 5$, the polynomial f(x) has two distinct roots ω_1 and ω_2 in \mathbb{F} . If char(\mathbb{F}) $\neq 2$, then $\omega_1 = \frac{-1+\sqrt{5}}{2}$ and $\omega_2 = \frac{-1-\sqrt{5}}{2}$. If char(\mathbb{F}) = 2, then ω_1 is a 3^{th} primitive root of unity and $\omega_2 = \omega_1^2$. Let

$$a_{2,2,2} = \left(\begin{array}{ccc} e_1 & 0 & 0\\ 0 & \omega_1 e_2 & e_2\\ 0 & \omega_1 e_2 & -\omega_1 e_2 \end{array}\right)$$

Then by the discussion before, C becomes a tensor category, denoted $C_{(0)}$, with the associativity constraint a induced by $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$. Similarly, let

$$a_{2,2,2}' = \begin{pmatrix} e_1 & 0 & 0 \\ 0 & \omega_2 e_2 & e_2 \\ 0 & \omega_2 e_2 & -\omega_2 e_2 \end{pmatrix}$$

and $a'_{1,i,j} = a'_{i,1,j} = a'_{i,j,1} = E_{\mathbf{e}_i \otimes \mathbf{e}_j}$ for $1 \leq i, j \leq 2$. Then \mathcal{C} also becomes a tensor category, denoted $\mathcal{C}'_{(0)}$ with the associativity constraint a' induced by $\{a'_{i,i,l} | 1 \leq i, j, l \leq 2\}$. For any $0 \neq \alpha \in \mathbb{F}$, let

$$a_{2,2,2}'' = \begin{pmatrix} e_1 & 0 & 0\\ 0 & \omega_1 e_2 & \alpha e_2\\ 0 & \alpha^{-1} \omega_1 e_2 & -\omega_1 e_2 \end{pmatrix}$$

and $a_{1,i,j}' = a_{i,1,j}' = a_{i,j,1}' = E_{\mathbf{e}_i \otimes \mathbf{e}_j}$ for $1 \leq i, j \leq 2$. Then by an argument similar to the proof of Proposition 3.7, one can check that $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ and $\{a_{i,j,l}'|1 \leq i, j, l \leq 2\}$ are equivalent. Similarly, if we set

$$a_{2,2,2}^{\prime\prime\prime} = \left(\begin{array}{ccc} e_1 & 0 & 0\\ 0 & \omega_2 e_2 & \alpha e_2\\ 0 & \alpha^{-1} \omega_2 e_2 & -\omega_2 e_2 \end{array} \right)$$

and $a_{1,i,j}^{\prime\prime\prime} = a_{i,1,j}^{\prime\prime\prime} = a_{i,j,1}^{\prime\prime\prime} = E_{\mathbf{e}_i \otimes \mathbf{e}_j}$ for $1 \leq i, j \leq 2$, then $\{a_{i,j,l}^{\prime} | 1 \leq i, j, l \leq 2\}$ and $\{a_{i,j,l}^{\prime\prime\prime} | 1 \leq i, j, l \leq 2\}$ are equivalent. Thus, we have the following proposition.

Proposition 3.8. Assume that $char(\mathbb{F}) \neq 5$. Let \widehat{C} be a semisimple tensor category of rank two over \mathbb{F} , and V a simple object not isomorphic to the unit object **1**. If $V \otimes V \cong \mathbf{1} \oplus V$, then \widehat{C} is tensor equivalent to $\mathcal{C}_{(0)}$ or $\mathcal{C}'_{(0)}$. Moreover, $\mathcal{C}_{(0)}$ and $\mathcal{C}'_{(0)}$ are not tensor equivalent.

Proof. The first statement follows from an argument similar to the proof of Proposition 3.7 and the above discussion. By [2, Theorem 3.22], one can see that $C_{(0)}$ and $C'_{(0)}$ are tensor equivalent if and only if $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ and $\{a'_{i,j,l}|1 \leq i, j, l \leq 2\}$ are equivalent. However, it is easy to check that $\{a_{i,j,l}|1 \leq i, j, l \leq 2\}$ and $\{a'_{i,j,l}|1 \leq i, j, l \leq 2\}$ are not equivalent. Therefore, $C_{(0)}$ and $C'_{(0)}$ are not tensor equivalent.

Now assume that m > 1. Let $G = (\Lambda_1 \otimes_{\mathbb{F}} I_n)(I_m \otimes_{\mathbb{F}} \Lambda_{32}) \in M_{mn^2 \times m^2}(\mathbb{F})$. Then (i) and (iii) are equivalent to

$$(I_m \otimes_{\mathbb{F}} \Lambda_2) \begin{pmatrix} I_m \otimes_{\mathbb{F}} \Lambda_{22} \\ G \end{pmatrix} = \begin{pmatrix} I_{[m]} \\ 0 \end{pmatrix}.$$

Let $\Lambda_2 = (\alpha_{ij})$ with $\alpha_{ij} \in \mathbb{F}$ and $G = \begin{pmatrix} G_{1,1} & G_{1,2} & \cdots & G_{1,m} \\ G_{2,1} & G_{2,2} & \cdots & G_{2,m} \\ \cdots & \cdots & \cdots & \cdots \\ G_{n^2,1} & G_{n^2,2} & \cdots & G_{n^2,m} \end{pmatrix}$ with $G_{i,j} \in M_m(\mathbb{F})$. Then $\Lambda_{22} = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1m} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{2m} \\ \cdots & \cdots & \cdots & \cdots \\ \alpha_{m1} & \alpha_{m2} & \cdots & \alpha_{mm} \end{pmatrix}$. Thus, for any $1 \leq s \leq m + n^2$

and $1 \leq t \leq m$, it follows from the matrix equation above that

$$\sum_{j=1}^{m} \alpha_{sj} \alpha_{jt} I_m + \sum_{j=m+1}^{m+n^2} \alpha_{sj} G_{j-m,t} = \begin{cases} I_{t,s}, & \text{if } 1 \leq s \leq m, \\ 0, & \text{if } m+1 \leq s \leq m+n^2. \end{cases}$$

Hence for any $1 \leq p, q \leq m$, by comparing the (p,q)-entries of the matrices on the both sides of the above equation, one gets

$$\sum_{j=1}^{m} \alpha_{sj} \alpha_{jt} \delta_{pq} + \sum_{j=m+1}^{m+n^2} \alpha_{sj} G_{j-m,t}(p,q) = \begin{cases} \delta_{tp} \delta_{sq}, & \text{if } 1 \leqslant s \leqslant m, \\ 0, & \text{if } m+1 \leqslant s \leqslant m+n^2, \end{cases}$$
(8)

where δ_{pq} 's are Kronecker symbols, $G_{j-m,t}(p,q)$ is the (p,q)-entry of the matrix $G_{j-m,t}$. Let $1 \leq k \leq m$. Putting t = p = q = k in the equation (8), one finds that

 $(\alpha_{1k}, \alpha_{2k}, \cdots, \alpha_{mk}, G_{1,k}(k,k), G_{2,k}(k,k), \cdots, G_{n^2,k}(k,k))^T$

is a solution of the system of linear equations $\Lambda_2 X = \varepsilon_{m+n^2,k}^T$. Now assume that m > 1. Then one can choose an integer l with $1 \leq l \leq m$ such that $l \neq k$. Putting t = p = l and q = k in the equation (8), one finds that

$$(0, 0, \cdots, 0, G_{1,l}(l, k), G_{2,l}(l, k), \cdots, G_{n^2,l}(l, k))^T$$

is also a solution of the system of linear equations $\Lambda_2 X = \varepsilon_{m+n^2,k}^T$. However, the solution of the system of linear equations $\Lambda_2 X = \varepsilon_{m+n^2,k}^T$ is unique since its coefficient matrix Λ_2 is invertible. This implies that $\alpha_{1k} = \alpha_{2k} = \cdots = \alpha_{mk} = 0$ and $\Lambda_{22} = 0$. Again by the invertibility of Λ_2 , the ranks of the matrices Λ_{23} and Λ_{32} are both equal to m. Then by the equation (vi), one gets $(\Lambda_{23} \otimes_{\mathbb{F}} I_n)(I_n \otimes_{\mathbb{F}} \Lambda_{32}) = 0$, which implies $2mn \leq n^3$, and hence $2m \leq n^2$.

Proposition 3.9. Let $\widehat{\mathcal{C}}$ be a semisimple tensor category of rank two over \mathbb{F} , and V a simple object not isomorphic to the unit object 1. Suppose that $V \otimes V \cong m\mathbf{1} \oplus nV$ with m > 0 and n > 0. If m > 1, then $2m \leq n^2$. Furthermore, if m = 2, then n > 2.

Proof. The first statement follows from the above discussion. Now let m = 2. Then $n \ge 2$. Suppose n = 2. Then $\Lambda_{23} \in M_{2 \times 4}(\mathbb{F})$ and $\Lambda_{32} \in M_{4 \times 2}(\mathbb{F})$. By the discussion

above, the ranks of Λ_{23} and Λ_{32} are both equal to 2. Let $\Lambda_{32} = \begin{pmatrix} \beta_{11} & \beta_{12} \\ \beta_{21} & \beta_{22} \\ \beta_{31} & \beta_{32} \\ \beta_{31} & \beta_{32} \\ \beta_{31} & \beta_{32} \end{pmatrix}$.

Then by the equations (v) and (vi), the following vectors are solutions of the system of linear equations $\Lambda_{23}X = 0$:

$$\begin{pmatrix} \beta_{1i} \\ \beta_{2i} \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ \beta_{1i} \\ \beta_{2i} \end{pmatrix}, \begin{pmatrix} \beta_{1i} \\ 0 \\ \beta_{2i} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \beta_{1i} \\ 0 \\ \beta_{2i} \end{pmatrix}, \begin{pmatrix} \beta_{3i} \\ \beta_{4i} \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ \beta_{3i} \\ \beta_{4i} \end{pmatrix}, \begin{pmatrix} \beta_{3i} \\ 0 \\ \beta_{4i} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \beta_{3i} \\ 0 \\ \beta_{4i} \end{pmatrix}$$

where $1 \leq i \leq 2$. Since $(\beta_{1i}, \beta_{2i}, \beta_{3i}, \beta_{4i}) \neq (0, 0, 0, 0)$, the rank of the above vector group is larger than or equal to 3, which contradicts to the fact that the rank of the matrix Λ_{23} is 2. Therefore, n > 2. Ostrik proved that there are only 4 fusion categories of rank 2 [9]. By above discussion, we have following remark.

Remark 3.10. The fusion categories of rank two are C_+ , C_- , $C_{(0)}$ and $C'_{(0)}$.

The following table display our main results.

m = 1, n = 0	$\mathcal{C}_+,\mathcal{C}$	$\begin{array}{c} \mathcal{C}_+ \ncong \mathcal{C} \text{ , if chark} \neq 2\\ \hline\\ \mathcal{C}_+ \cong \mathcal{C} \text{, if chark} = 2 \end{array}$
m > 1, n = 0	No	
m = 0, n = 1	\mathcal{C}_1	
m=0,n>1	$\mathcal{C}_n \ncong \mathcal{C}'_n$	
m = 1, n = 1	$\mathcal{C}_{(5)}$, if chark=5	$\mathcal{C}_{(0)} \cong \mathcal{C}'_{(0)}, \text{ if chark} \neq 5$
$m > 1, 2m > n^2$	No	
m = 2, n = 2	No	

ACKNOWLEDGMENTS

This work is supported by NNSF of China (Nos. 12071412,12201545).

References

- B. Bakalov and A. Kirillov, Jr., Lectures on tensor categories and modular functor, Amer. Math. Soc., 2000.
- [2] H. X. Chen and Y. H. Zhang, Reconstruction of tensor categories from their structure invariants, Bull.Belg. Math. Soc. Simon Stevin 27(2020), no 2,245-279.
- [3] P. Etingof and M. Khovanov, Representations of tensor categories and Dynkin diagrams, Inter. Math. Res. Let. 5 (1995), 235-247.
- [4] P. Etingof, D. Nikschych and V. Ostrik, Tensor categories, Mathematical Survey and Monographs Vol. 205, AMS 2015.
- [5] C. Kassel, Quantum groups, Springer-Verlag, New York, 1995.
- [6] H. K. Larson, Pseudo-unitary non-self-dual fusion categories of rank 4, J. Algebra 415 (2014), 184-213.
- [7] G. Lusztig, Leading coefficients of character values of Hecke algebras, Proc. Symp. in Pure Math. 47 (1987), 235-262.
- [8] V. Ostrik, module categories, weak Hopf algebras and modular invariants, Transform. Groups 8(2) (2003), 177-206.
- [9] V. Ostrik, Fusion categories of rank 2, Math. Res. Let. 10 (2003), 177-183.
- [10] V. Ostrik, Pivotal fusion category of rank 3, arXiv:13094822.
- [11] E. Rowell, R. Stong and Z, Wang, On classification of modular tensor categories, Comm. Math. Phys. 292 (2009), 343-389.

College of Mathematical Science, Yangzhou University, Yangzhou 225002, China

Email address: huasun@yzu.edu.cn

College of Mathematical Science, Yangzhou University, Yangzhou 225002, China

Email address: hxchen@yzu.edu.cn

Department of Mathematics & Statistics, University of Hasselt, Universitaire Campus, 3590 Diepeenbeek, Belgium

Email address: yinhuo.zhang@uhasselt.be