A RANDOM MATRIX MODEL TOWARDS THE QUANTUM CHAOS TRANSITION CONJECTURE

BERTRAND STONE, FAN YANG, AND JUN YIN

ABSTRACT. Consider D random systems that are modeled by independent $N \times N$ complex Hermitian Wigner matrices. Suppose they are lying on a circle and the neighboring systems interact with each other through a deterministic matrix A. We prove that in the asymptotic limit $N \to \infty$, the whole system exhibits a quantum chaos transition when the interaction strength $||A||_{\text{HS}}$ varies. Specifically, when $||A||_{\text{HS}} \ge N^{\varepsilon}$, we prove that the bulk eigenvalue statistics match those of a $DN \times DN$ GUE asymptotically and each bulk eigenvector is approximately equally distributed among the D subsystems with probability 1 - o(1). These phenomena indicate quantum chaos of the whole system. In contrast, when $||A||_{\text{HS}} \le N^{-\varepsilon}$, we show that the system is integrable: the bulk eigenvalue statistics behave like D independent copies of GUE statistics asymptotically and each bulk eigenvector is localized on only one subsystem. In particular, if we take $D \to \infty$ after the $N \to \infty$ limit, the bulk statistics converge to a Poisson point process under the DN scaling.

CONTENTS

1.	Introduction	1
2.	Main results	7
3.	Chaotic regime: eigenvectors	13
4.	Gaussian divisible case	15
5.	Characteristic flow estimates	26
6.	Chaotic regime: eigenvalues	32
7.	Integrable regime: eigenvectors	35
8.	Integrable regime: eigenvalues	38
Ap	opendix A. Some deterministic estimates	46
Ap	opendix B. Green's function comparison	48
Ap	opendix C. Proof of local laws	49
Re	ferences	52

1. INTRODUCTION

Consider a quantum Hamiltonian $H_{\lambda} = H + \lambda H_I$, where H represents the Hamiltonian of some separable (or independent) subsystems in a specific coordinate system, and H_I is a non-separable perturbation describing the interactions between these subsystems. In the study of quantum chaos, researchers aim to answer the following questions. (i) Does the behavior of the system become "chaotic" when the interaction (measured by the parameter $\lambda > 0$) becomes sufficiently strong? If the chaotic phase indeed exists, then (ii) what kind of features of energy levels, eigenstates, or some other physical observables can be used to distinguish between the integrable (or non-chaotic) and chaotic phases, and (iii) when the quantum chaos transition occurs as λ varies.

In the context of a generic integrable model, the well-known Berry-Tabor conjecture [18, 19, 87] states that the local energy level statistics should follow a Poisson process, i.e., neighboring energy levels should be uncorrelated. Conversely, for a generic chaotic model, the renowned Bohigas-Giannoni-Schmit (BGS) conjecture [23,24] suggests that its local energy level statistics should resemble those of Gaussian orthogonal ensemble (GOE). Specifically, it predicts the presence of level repulsion between neighboring energy levels, and the nearest neighbor spacing is expected to align with the "Wigner surmise" predicted by [106] for Wigner matrices. The eigenfunctions of chaotic models are predicted to satisfy a fundamental property called *quantum ergodicity* [51, 101, 112], i.e., almost all eigenfunctions become equidistributed as the energy tends to ∞ . Later, a stronger notion called *quantum unique ergodicity* (QUE) states that the above equipartition principle holds for all eigenfunctions in the high-energy limit [92]. We refer the readers to [21, 22, 31] for a more comprehensive review of the above conjectures and the relation between quantum chaos and random matrix theory. There have been numerous studies on the Berry-Tabor conjecture [38, 39, 88, 89, 104] and quantum unique ergodicity [74, 85], to name just a few. However, to the best of our knowledge, the BGS conjecture remains unproven in the existing literature.

In the past few decades, there has been great progress in understanding the chaotic behaviors of some classical ensembles of random matrices, including the delocalization of eigenvectors, QUE, the GOE/GUE statistics of eigenvalues (or the Wigner surmise), and so on. They are now considered a good paradigm for exploring many ideas related to quantum chaos and the BGS conjecture. The extensive body of literature explores these behaviors across various random matrix ensembles, such as Wigner matrices (see, for example, [8, 27, 33, 41–43, 45, 59–63, 65–67, 81, 82, 103]), random graphs (see, for example, [10, 11, 14–16, 30, 55, 57, 76]), heavy-tailed random matrices (see, for example, [4–6]), and non-mean-field random matrices (see, for example, [28,34,56,107–109,111]), to name just a few. For a historical review of seminal works that have led to the proof of the Wigner surmise and the delocalization of eigenvectors for random matrices, readers can refer to the book [64]. However, most of these models (except for non-mean-field random matrices) are chaotic in nature, so they generally do not exhibit integrable features and hence are not suitable for the investigation of the transition between integrable and chaotic phases. On the other hand, random band matrices are proposed for this purpose. The random band matrix ensemble with band width W and system size N is conjectured to exhibit a quantum chaos transition as W varies (see, for example, [25, 35, 36, 68, 69, 102]): there exists a critical band width $W_c = \sqrt{L}$ such that

- if $W \ll W_c$, H has localized bulk eigenvectors, and the bulk eigenvalues satisfy Poisson statistics;
- if $W \gg W_c$, *H* has delocalized bulk eigenvectors satisfying QUE, and the bulk eigenvalues satisfy GOE/GUE statistics depending on the symmetry of the system.

This conjecture remains unresolved; however, significant progress has been made in the past decade towards establishing its validity (see, for example, [13, 28, 34, 37, 50, 52, 53, 56, 70, 73, 91, 94–100, 111]).

1.1. Overview of the main results. In this paper, we consider a simpler random matrix model (than random band matrices) that also naturally exhibits the quantum chaos transition. More precisely, fix an integer $D \ge 2$, we take D independent random subsystems, assuming that they can be modeled by $N \times N$ Wigner matrices, whose entries have mean zero, variance N^{-1} , and fulfill certain moment conditions. Without introducing interactions, the whole system is modeled by a block diagonal matrix H with diagonal blocks being independent Wigner matrices H_a , $a = 1, \ldots, D$. Next, we assume that the subsystems are aligned along a cycle and the neighboring subsystems interact with each other through an arbitrary deterministic $N \times N$ matrix A (non-nearest-neighbor interactions can also be allowed; see Remark 1.1 below). We denote the interaction Hamiltonian by Λ , which is a tridiagonal block matrix with non-diagonal blocks being A or A^* , and denote the whole system by H_{Λ} . In other words, H_{Λ} is defined as

$$H_{\Lambda} = H + \Lambda, \tag{1.1}$$

where H and A are $D \times D$ block matrices defined as

$$H = \begin{pmatrix} H_1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & H_2 & 0 & \cdots & 0 & 0 \\ 0 & 0 & H_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & H_{D-1} & 0 \\ 0 & 0 & 0 & \cdots & 0 & H_D \end{pmatrix}, \quad \Lambda = \begin{pmatrix} 0 & A & 0 & \cdots & 0 & A^* \\ A^* & 0 & A & \cdots & 0 & 0 \\ 0 & A^* & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & A \\ A & 0 & 0 & \cdots & A^* & 0 \end{pmatrix}.$$
(1.2)

Furthermore, we assume that H_{Λ} is a perturbation of H, i.e., $||A|| \ll \mathbb{E}||H|| \sim 1$.

In this paper, we prove that the bulk eigenvalues and eigenvectors of H_{Λ} exhibit a transition from the integrable phase to the chaotic phase, and this transition occurs at $||A||_{\text{HS}} \sim 1$. For simplicity, we introduce the index sets $\mathcal{I}_a := [\![(a-1)N+1, aN]\!]$, $a \in \{1, \ldots, D\}$, for the subsystems and let $\mathcal{I} := [\![DN]\!]$ be the index set for the whole system. Hereafter, for any $n, m \in \mathbb{R}$, we denote $[\![n, m]\!] := [\![n, m] \cap \mathbb{Z}$ and $[\![n]\!] := [\![1, n]\!]$. Roughly speaking, we prove the following results:

- If $||A||_{\text{HS}} \ll 1$, then every bulk eigenvector **v** is concentrated on only one subsystem, i.e., there exists an \mathcal{I}_a such that $\sum_{k \in \mathcal{I}_a} |\mathbf{v}(k)|^2 = 1 o(1)$ with probability 1 o(1) (Theorem 2.4). The bulk eigenvalues of H_{Λ} are negligible perturbations of those of H (Theorem 2.5); in particular, if D is large, the eigenvalue statistics around a bulk energy is almost a Poisson point process under the DN scaling.
- If $||A||_{\text{HS}} \gg 1$, then every bulk eigenvector **v** extends to the whole system (Theorem 2.2): on each \mathcal{I}_a ,

$$\sum_{k \in \mathcal{I}_a} |\mathbf{v}(k)|^2 = D^{-1} + o(1) \quad \text{with probability } 1 - o(1).$$
(1.3)

The bulk eigenvalue statistics of H_{Λ} match those of a $DN \times DN$ GOE/GUE asymptotically (Theorem 2.3). For ease of presentation, we will, with a slight departure from strict notation, refer to (1.3) as the "QUE estimate" within the context of this paper. It is important to note, however, that QUE typically denotes a stronger property; see the discussion below Theorem 2.2.

We would like to emphasize that these results apply to any deterministic interaction A—no specific structure about A will be used in the proof except for the operator norm bound $||A|| \ll 1$ and the conditions on $||A||_{\text{HS}}$.

As $N \to \infty$, when D is fixed, the typical spectral distances between D independent Wigner matrices are of order N^{-1} . Intuitively, quantum chaos arises when the interaction Λ mixes the spectra of subsystems. In the proof, we will show that Λ induces a shift in the eigenvalues of magnitude $N^{-1} ||A||_{\text{HS}}$, thereby rigorously confirming the aforementioned intuition: when $||A||_{\text{HS}} \ll 1$, the spectra of subsystems remain separate, indicating integrability of the entire system; when $||A||_{\text{HS}} \gg 1$, the subsystems become mixed, resulting in chaotic behavior.



FIGURE 1.1. The entries $|\mathbf{v}(k)|^2$ for $k \in \mathcal{I}$, where \mathbf{v} is the (DN/2)-th eigenvector of H_{Λ} . We choose D = 10, N = 200, and $A = \lambda I$. In (a) and (b), the red lines mark the value N^{-1} ; in (c) and (d), the red lines mark the value $(DN)^{-1}$.

We now perform some simulations to illustrate the theoretical results. We choose the D independent $N \times N$ Wigner matrices to be GOE, and take $A = \lambda I$ (so $||A||_{\text{HS}} = \lambda \sqrt{N}$ and the transition occurs at $\lambda = N^{-1/2}$). In Figure 1.1, we show the squared magnitudes of the entries of a bulk eigenvector. One can observe that in the subcritical cases, the eigenvector entries are localized in only one block, while in the supercritical cases, the eigenvector is delocalized. In Figure 1.2, we depict the simulated distributions of the energy gaps between neighboring bulk eigenvalues under the scaling DN. In the subcritical cases, the simulated gap distribution exhibits a good resemblance to the exponential distribution, providing evidence

that the behavior of bulk eigenvalues locally resembles that of a Poisson process. This finding substantiates the Berry-Tabor conjecture within our framework. Conversely, in the supercritical cases, the simulated gap distribution closely aligns with the Wigner surmise, indicating that the bulk eigenvalues follow the GOE statistics. This result confirms the BGS conjecture within our context.



FIGURE 1.2. The bulk eigenvalue gap distributions under the same setting as Figure 1.1. The normalized histograms plot the rescaled eigenvalue gaps $DN\rho_{sc}(\lambda_k)(\lambda_{k+1} - \lambda_k)$, where λ_k are the eigenvalues of H_{Λ} with $k \in [DN/20, 19DN/20]$ and ρ_{sc} is the semicircle density given by (2.17) below. In (a) and (b), the red curves plot the probability density function for the exponential distribution: $f(x) = e^{-x}$. In (c) and (d), the red curves plot the Wigner surmise: $f(x) = \frac{\pi x}{2} e^{-\pi x^2/4}$. Note that repulsions between eigenvalues already appear to be present in the $\lambda = N^{-0.6}$ case. We attribute this phenomenon to the finite-N effect.

Remark 1.1. There are some direct extensions of our results to various other models. Our method relies primarily on the block diagonal structure of matrix H and the block translation invariance of the interactions. Hence, the proofs in our paper can be readily extended to more general cases encompassing non-nearestneighbor interactions (including interactions within each block), as long as the interaction between the *i*-th and *j*-th blocks depends only on j - i (by utilizing the periodic distance on the cycle). We can further relax the condition and only require that the interactions have the same distribution. For instance, the interactions can consist of independent lower triangular matrices with i.i.d. entries, in which case we get random band matrices. Alternatively, the interactions can be independent Ginibre matrices with i.i.d. entries, leading to the Wegner orbital model [91]. In addition, the random matrix characterizing each subsystem can be more intricate models, such as sparse random matrices, generalized Wigner matrices, or deformed Wigner matrices. Finally, our results can be easily extended to higher-dimensional models, where the subsystems reside on a *d*-dimensional lattice with $d \ge 2$. In this setting, all our arguments remain applicable as long as the interactions between different blocks maintain translation symmetry on the lattice.

Remark 1.2. At the critical point where $||A||_{\text{HS}} \sim 1$, simulations show that each block contains a substantial portion of a bulk eigenvector, although certain blocks contain significantly larger amounts (measured in terms of the ℓ^2 -norm) than others. We conjecture that the ℓ^2 -mass distribution across the *D* blocks would follow a polynomial decay. Additionally, we anticipate that the bulk eigenvalue statistics would exhibit an interpolation between the GOE/GUE statistics and *D* independent copies of GOE/GUE statistics. The investigation of this critical scenario is deferred for future research.

1.2. **Physical background.** One of the main physical motivations behind this model is to gain insights into the Anderson metal-insulator transition of the well-known Anderson model [12]. In the Anderson model, the random potential (denoted by V) consists of i.i.d. diagonal entries. However, in our model (and its higher-dimensional counterparts, as mentioned in Remark 1.1), we have replaced V with a block diagonal random matrix H and generalized the Laplacian matrix to a more general matrix Λ that introduces interactions between adjacent blocks. From a physical perspective, our "block potential" is related to the And erson potential through a scaling transformation. Regarding Λ , choosing A = I yields the block Anderson model [91], while incorporating random independent interactions leads to random band matrices or the Wegner orbital model, as mentioned in Remark 1.1 above. These models represent important extensions of the Anderson model and all exhibit the desired metal-insulator transition as the strength of interactions varies. In particular, the Wegner orbital models draw inspiration from the work of Wegner [105], which was further developed in [90, 93]. They are proposed to model the motion of a quantum particle with many internal degrees of freedom in a disordered medium. We expect that investigating the block model H_{Λ} could provide an alternative approach to the renowned Anderson metal-insulator transition conjecture. We will delve into this aspect in an ongoing paper [110], where we study the localization-delocalization transition of the block Anderson and Wegner orbital models as the interaction strength changes. It is worth highlighting that the framework presented in that paper poses greater challenges, as it involves examining "more local" interactions where $D \to \infty$ as $N \to \infty$.

Next, we explore another physical connection, aiming to gain some *heuristic insights* into the mechanism of quantum chaos transition through the lens of random matrix theory. To accomplish this, we begin by considering a simplified toy model. Consider the two-dimensional (2D) quantum billiard inside the unit disk \mathcal{D} :

$$-\Delta\psi(r,\theta) = E\psi(r,\theta), \quad \psi|_{\partial\mathcal{D}} = 0.$$

This is an integrable model with eigenfunctions $\psi_{n,m}(r,\theta) = J_{|m|}(k_{m,n}r)e^{im\theta}$ and eigenvalues $E = k_{m,n}^2$, where $m \in \mathbb{Z}$, $n \in \mathbb{N}$, $J_{|m|}$ is the |m|-th Bessel function, and $k_{m,n}$ is the *n*-th zero of $J_{|m|}$. We then add a potential $V(r,\theta)$ to the Hamiltonian: $H = -\Delta + V(r,\theta)$. Note when V tends to $\infty \cdot \mathbf{1}_{\mathcal{B}}$ for a region $\mathcal{B} \subset \mathcal{D}$, we get a quantum billiard inside the region between $\partial \mathcal{D}$ and $\partial \mathcal{B}$. Our goal is to understand when quantum chaos arises as the potential V varies. In particular, we are interested in the asymptotic behavior of the eigenvalues and corresponding eigenstates around an energy E in the high-energy limit $E \to \infty$.

In the original quantum billiard in \mathcal{D} , there is a conserved physical quantity, namely the *angular momentum* $-i\partial_{\theta}$. As a result, the operator $-\Delta$ can be decomposed according to its invariant subspaces characterized by the quantum numbers for the angular momentum:

$$V_m := \left\{ \psi(r, \theta) : \psi(r, \theta) = f(r)e^{im\theta} \text{ for some radial function } f(r) \right\}$$

where $-\Delta$ restricted to V_m is given by

$$-\Delta \psi = -\Delta^{(m)}\psi, \quad \text{with} \quad \psi(r,\theta) = f(r)e^{\mathrm{i}m\theta}, \quad \Delta^{(m)} := \frac{\partial^2}{\partial r^2} + \frac{1}{r}\frac{\partial}{\partial r} - \frac{m^2}{r^2}.$$
 (1.4)

This decomposition naturally leads to a block structure of H with blocks labeled by the quantum numbers for the angular momentum. More precisely, let $\{f_n(r)\}$ be an orthonormal basis of the space of radial functions with $f_n(1) = 0$. Then, the (m, m')-block of H can be represented by an ∞ -dimensional matrix with entries

$$H_{n',n}^{(m',m)} = \frac{1}{2\pi} \iint \bar{f}_{n'}(r) \left[-\Delta^{(m)} + V(r,\theta) \right] f_n(r) e^{i(m-m')\theta} dr d\theta.$$
(1.5)

By performing a finite-dimensional reduction, we can express H as a block matrix.

The block matrix obtained above is a deterministic matrix, whose behavior seems to be inherently complex to analyze. Intuitively, one would expect that, for a general V, H exhibits characteristics akin to a "quasi-random" matrix, with entries possessing intricate magnitudes and phases. To facilitate a more accessible study of the quantum chaos transition, we propose employing a block random matrix H_{Λ} as a computationally feasible "toy model". This model allows for easier investigation while still capturing essential qualitative features that establish a connection between the aforementioned quantum billiard model and the model (1.1).

(i) Due to the fast-oscillating nature of the functions f_n when n is large, the entries within the diagonal blocks $H^{(m,m)}$ exhibit complicated phases. Furthermore, neighboring diagonal blocks demonstrate slightly different spectra (due to the term $-m^2/r^2$ in (1.4)). Following Wigner's insight [106], we



FIGURE 1.3. A 2D quantum billiard inside a circle with a potential V supported in a compact region. The left picture depicts an integrable system featuring a radially symmetric potential, which does not mix eigenstates with different angular momentum. In the right picture, the rotational symmetry of V is broken, resulting in mixing of eigenstates with different angular momentum, which can induce chaotic behavior. Note when the potential $V \to \infty$, essentially forming an inner boundary, the system becomes a quantum billiard between two boundaries.

propose employing independent Wigner matrices to model these blocks—random matrices serve as effective phenomenological models for complex "quasi-random" matrices, and independent Wigner matrices have distinct yet closely aligned spectra.

- (ii) When the potential $V(r, \theta)$ is zero or exhibits rotational symmetry, as depicted in the left graph of Figure 1.3, the eigenstates with different angular momentum remain unmixed, resulting in an integrable model. This scenario corresponds to the A = 0 case in our model (1.1).
- (iii) When we modify the potential V as shown in the right graph of Figure 1.3, off-diagonal block entries $H_{n',n}^{(m',m)} = (2\pi)^{-1} \iint \bar{f}_{n'}(r)V(r,\theta)f_n(r)e^{i(m-m')\theta}drd\theta$ with $m' \neq m$ will emerge and they reflect the strength of V and the departure of V from rotational symmetry. From (1.5), we first notice that H exhibits a block translation symmetry, meaning that $H^{(m',m)}$ depends on m' and m only through their difference m' m. This aligns with the block translation invariance of Λ in (1.2). Second, the magnitudes of the off-diagonal block entries $H_{n',n}^{(m',m)}$ generally decrease as the difference |m' m| increases, given fixed n and n'. Consequently, we have chosen Λ in (1.2) to be nearest-neighbor interactions to model such "short-range phenomenon". (However, this simplification is not crucial since, as mentioned in Remark 1.1, we do allow for the inclusion of non-nearest-neighbor interactions as well.)

Physically, item (ii) above means that the introduction of a circular hole at the center still gives a block diagonal matrix, so our results in the integrable phase imply *heuristically* the integrability of the quantum billiard model. On the other hand, in this paper, we provide evidence of a quantum chaos transition for H_{Λ} as the interaction strength increases. In particular, as $V \to \infty$, our results suggest, in a heuristic sense, that adding a "generic" non-rotationally symmetric hole will lead to chaos.

We expect that our paper may serve as an initial step towards heuristically understanding the quantum chaos transition from the perspective of random matrix theory. Significant efforts are still needed to attain a more comprehensive understanding and navigate the vast realm awaiting exploration. One important extension is explained in the following remark.

Remark 1.3. In the model (1.1), the quantities D and N correspond to the number of quantum numbers m for the angular momentum and the number of basis functions f_n retained in the finite-dimensional reduction, respectively. On the other hand, given a target energy E, the matrix dimension DN represents the number of energy levels falling within an energy window around E, say [0, 2E]. When |m| or n is large, utilizing the asymptotics of the zeros of Bessel functions, we see that the eigenvalues of $-\Delta$ scale as $k_{m,n}^2 \sim m^2 + n^2$. Consequently, it becomes apparent that the most relevant regime for investigating the quantum chaos transition should be $D \sim N \sim \sqrt{E}$. Note that this regime corresponds to a random band matrix model with a *critical band width*, which is a highly challenging problem as mentioned earlier. As a simplification, this paper focuses on a scenario where D remains fixed as $N \to \infty$, i.e., we only examine the mixing of a finite number of blocks with similar angular momentum. It is possible to relax the assumption to a certain extent with $1 \ll D \ll N$, but we defer such exploration to [110] and more future studies.

The rest of this paper is organized as follows. In Section 2, we define our model and state the main results, which consist of two parts: in the chaotic regime with $||A|| \ge N^{\varepsilon}$, we state the QUE of bulk eigenvectors (Theorem 2.2) and the GUE statistics for the bulk eigenvalues (Theorem 2.3); in the integrable regime, we state the localization of bulk eigenvectors (Theorem 2.4) and the Poisson statistics for the bulk eigenvalues (Theorem 2.5). These theorems (Theorems 2.2–2.5) are proved in Section 3, Section 6, Section 7, and Section 8, respectively. For the proof of Theorem 2.2, we reduce its proof to that of a Gaussian divisible case with a small Gaussian component in Section 4, and then prove the Gaussian divisible case using a characteristic flow argument in Section 5. Finally, the proofs of several estimates used in the main proofs are included in Appendix A–C, including some basic deterministic estimates, a Green's function comparison estimate, and the local laws for the Green's function of H_{Λ} .

Notations. To facilitate the presentation, we introduce some necessary notations that will be used throughout this paper. In this paper, we are interested in the asymptotic regime with $N \to \infty$. When we refer to a constant, it will not depend on N. Unless otherwise noted, we will use C to denote generic large positive constants, whose values may change from line to line. Similarly, we will use ε , δ , τ , c etc. to denote generic small positive constants. For any two (possibly complex) sequences a_N and b_N depending on N, $a_N = O(b_N)$ or $a_N \leq b_N$ means that $|a_N| \leq C|b_N|$ for a constant C > 0, whereas $a_N = o(b_N)$ or $|a_N| \ll |b_N|$ means that $\lim_{N\to\infty} |a_N|/|b_N| \to 0$. We say that $a_N \sim b_N$ if $a_N = O(b_N)$ and $b_N = O(a_N)$. For any $a, b \in \mathbb{R}$, we denote $a \lor b := \max\{a, b\}$ and $a \land b := \min\{a, b\}$. For an event Ξ , we let $\mathbf{1}_{\Xi}$ or $\mathbf{1}(\Xi)$ denote its indicator function. Given a vector \mathbf{v} , $\|\mathbf{v}\| \equiv \|\mathbf{v}\|_2$ denotes the Euclidean norm and $\|\mathbf{v}\|_p$ denotes the ℓ_p -norm. Throughout this paper, we use "*" to denote the Hermitian conjugate of a matrix. Given a matrix $B = (B_{ij})$, we use $\|B\|$, $\|B\|_{\mathrm{HS}}$, and $\|B\|_{\mathrm{max}} := \max_{i,j} |B_{ij}|$ to denote the operator, Hilbert-Schmidt, and maximum norms, respectively. We also adopt the notion of generalized entries: $B_{\mathbf{uv}} \equiv \mathbf{u}^* B\mathbf{v}$ for vectors \mathbf{u}, \mathbf{v} .

Acknowledgement. Fan Yang is supported in part by the National Key R&D Program of China (No. 2023YFA1010400). Jun Yin is supported in part by the Simons Fellows in Mathematics Award 85515. The authors would like to thank László Erdős for helpful comments on an earlier draft of this paper. We are very grateful to two anonymous referees for the helpful comments, which have resulted in a significant improvement of the paper.

2. Main results

2.1. The model and main results. In this paper, we consider a block random matrix model. Fix any integer $D \ge 2$, let H_1, H_2, \ldots, H_D be D independent copies of $N \times N$ Wigner matrices, i.e., the entries of H_a are independent (up to symmetry $H = H^*$) random variables satisfying that

$$\mathbb{E}(H_a)_{ij} = 0, \quad \mathbb{E}|(H_a)_{ij}|^2 = N^{-1}, \quad a \in [\![D]\!], \quad i, j \in [\![N]\!].$$
(2.1)

For the definiteness of notations, in this paper, we consider the complex Hermitian case, while the real case can be proved in the same way with some minor changes in notations. In the complex case, we assume additionally that

$$\mathbb{E}[(H_a)_{ij}^2] = 0, \quad a \in [\![D]\!], \quad i \neq j \in [\![N]\!].$$
(2.2)

We assume that the diagonal entries are i.i.d. real random variables and the entries above the diagonal are i.i.d. complex random variables. Let A be an arbitrary $N \times N$ (real or complex) deterministic matrix. Then, we consider the block random matrix model H_{Λ} defined in (1.1) with H and Λ given in (1.2). Note that the distribution of the model H_{Λ} exhibits block translation symmetry. As we consider H_{Λ} as a perturbation of H, we always assume in this paper that $||A|| \ll \mathbb{E}||H|| \sim 1$. Under this assumption, the empirical measure of H_{Λ} follows approximately the semicircle law with support [-2, 2]. (We believe that the results in the chaotic regime, Theorems 2.2 and 2.3, remain valid even for $||A|| \gtrsim 1$ provided that A has a reasonably large rank. However, they may fail if, for example, A contains only one nonzero entry.) We now summarize the assumptions for the model (1.1).

Assumption 2.1. Fix any integer $D \ge 2$, we consider the model (1.1), where A is an arbitrary $N \times N$ deterministic matrix with $||A|| \le N^{-\delta_A}$ for a constant $\delta_A > 0$, and H_1, H_2, \ldots, H_D are D i.i.d. $N \times N$ complex Hermitian Wigner matrices satisfying (2.1), (2.2), and the following high moment condition: for any $p \in \mathbb{N}$, there exists a constant $C_p > 0$ such that

$$\mathbb{E} |H_{11}|^p + \mathbb{E} |H_{12}|^p \leqslant C_p N^{-p/2}.$$
(2.3)

Let the eigenvalues of H_{Λ} be $\lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_{DN}$ and denote the corresponding (unit) eigenvectors by $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_{DN}$. Now, we are ready to state the first two main results concerning the bulk eigenvectors and eigenvalues in the quantum chaotic phase when $||A||_{\text{HS}} \gg 1$. Our first theorem states that with probability 1 - o(1), each bulk eigenvector is asymptotically uniformly distributed among all $\mathcal{I}_a, a \in [\![D]\!]$.

Theorem 2.2 (Chaotic regime: eigenvectors). Under Assumption 2.1, let $\kappa \in (0, 1/2)$ be an arbitrary constant and suppose there exists a constant $\varepsilon_A > 0$ such that

$$\|A\|_{\mathrm{HS}} \ge N^{\varepsilon_A}.\tag{2.4}$$

Then, for any $k \in [[\kappa DN, (1-\kappa)DN]]$, there exists a constant c > 0 such that

$$\mathbb{P}\left(\max_{a\in[D]}\left|\mathbf{v}_{k}^{*}E_{a}\mathbf{v}_{k}-D^{-1}\right|\geq N^{-c}\right)\leq N^{-c},$$
(2.5)

where $E_a \in \mathbb{C}^{DN \times DN}$ denotes the block identity matrix restricted to \mathcal{I}_a , i.e., $(E_a)_{ij} = \mathbf{1}(i = j \in \mathcal{I}_a)$.

In the proof, we will prove a slightly stronger result than (2.5):

$$\mathbb{P}\left(\max_{i,j\in[[k-N^c,k+N^c]]}\max_{a\in[[D]]}\left|\mathbf{v}_i^*(E_a-D^{-1})\mathbf{v}_j\right|\geq N^{-c}\right)\leq N^{-c}.$$
(2.6)

As mentioned below (1.3), we will refer to (2.5) and (2.6) as the "QUE estimates" of bulk eigenvectors. However, QUE usually represents a stronger property: with probability 1 - o(1), the entries of \mathbf{v}_k are approximately equally distributed on any subset $I \subset \mathcal{I}$ with $|I| \gg 1$. Such probabilistic QUE was first proved for Wigner matrices [33]; later, a stronger notion called the *eigenstate thermalization hypothesis* (ETH) was also established for Wigner matrices [40, 42, 43, 45]. The QUE (or ETH) of eigenvectors has been extended to many other types of mean-field random matrices and random graphs (such as [1, 5, 15, 17, 30, 41, 86], to name just a few) and non-mean-field random matrices [28, 34, 107]. In this paper, we refrain from proving the stronger QUE for brevity, and the current form is already sufficient for our proof of the following result, Theorem 2.3. We leave the exploration of optimal QUE or ETH to future works.

Our second theorem shows that the local bulk eigenvalue statistics of H_{Λ} match those of GUE asymptotically under the condition (2.4), that is, H_{Λ} satisfies the bulk universality. Let $p_{H_{\Lambda}}(\lambda_1, \ldots, \lambda_{DN})$ denote the joint symmetrized probability density of the eigenvalues of H_{Λ} . For any $1 \leq n \leq DN$, define the *n*-point correlation function by

$$p_{H_{\Lambda}}^{(n)}(\lambda_{1},\cdots,\lambda_{n}) := \int_{\mathbb{R}^{DN-n}} p_{H_{\Lambda}}(\lambda_{1},\cdots,\lambda_{DN}) \,\mathrm{d}\lambda_{n+1}\cdots\mathrm{d}\lambda_{DN}.$$

Denote the corresponding *n*-point correlation function for $DN \times DN$ GUE by $p_{GUE}^{(n)}$.

Theorem 2.3 (Chaotic regime: eigenvalues). In the setting of Theorem 2.2, let $O \in C_c^{\infty}(\mathbb{R}^n)$ be an arbitrary smooth, compactly supported function. Then, for any $|E| \leq 2 - \kappa$ and fixed $n \in \mathbb{N}$, there exists a constant c > 0 so that

$$\left| \int_{\mathbb{R}^n} \mathrm{d}\boldsymbol{\alpha} \ O(\boldsymbol{\alpha}) \left[p_{H_{\Lambda}}^{(n)} - p_{GUE}^{(n)} \right] \left(E + \frac{\alpha_1}{DN}, \dots, E + \frac{\alpha_n}{DN} \right) \right| \leq N^{-c}, \quad \boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_n).$$

The next two main results concern the bulk eigenvectors and eigenvalues of H_{Λ} in the integrable (or non-chaotic) phase when $||A||_{\text{HS}} \ll 1$. First, Theorem 2.4 shows that in contrast to Theorem 2.2, each bulk eigenvector is mostly localized in only one \mathcal{I}_a in the integrable phase.

Theorem 2.4 (Integrable regime: eigenvectors). Under Assumption 2.1, let $\kappa \in (0, 1/2)$ be an arbitrary constant and suppose there exists a constant ε_A such that

$$\|A\|_{\mathrm{HS}} \le N^{-\varepsilon_A}.\tag{2.7}$$

Then, for any $k \in [\kappa DN, (1-\kappa)DN]$, there exists a constant c > 0 such that

$$\mathbb{P}\left(\max_{a=1}^{D} \|E_a \mathbf{v}_k\|^2 \le 1 - N^{-c}\right) \le N^{-c}.$$
(2.8)

Next, we show that under the condition (2.7), the bulk eigenvalues of H_{Λ} are negligible perturbations of those of H under the scaling N. As a result, the bulk eigenvalue statistics of H_{Λ} and H match asymptotically. Denote the eigenvalues of H as $\lambda_1(H) \leq \ldots \leq \lambda_{DN}(H)$, and for any $1 \leq n \leq N$, let $p_H^{(n)}$ represent the *n*-point correlation function of them.

Theorem 2.5 (Integrable regime: eigenvalues). Under the setting of Theorem 2.4, for any $k \in [\kappa DN, (1 - \kappa)DN]$, there exists a constant c > 0 such that

$$\mathbb{P}\left(|\lambda_k - \lambda_k(H)| \ge N^{-1-c}\right) \le N^{-c}.$$
(2.9)

As a consequence, it implies that for any $|E| \leq 2 - \kappa$, fixed $n \in \mathbb{N}$, and smooth, compactly supported test function $O \in C_c^{\infty}(\mathbb{R}^n)$, there exists a constant c > 0 so that

$$\left| \int_{\mathbb{R}^n} \mathrm{d}\boldsymbol{\alpha} \, O(\boldsymbol{\alpha}) \left[p_{H_{\Lambda}}^{(n)} - p_{H}^{(n)} \right] \left(E + \frac{\alpha_1}{DN}, \dots, E + \frac{\alpha_n}{DN} \right) \right| \leqslant N^{-c}, \quad \boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_n).$$

The spectrum of H consists of the eigenvalues of D independent copies of Wigner matrices H_1, \ldots, H_D . Recall that it has been shown in a sequence of works (see e.g., [27, 59, 62, 65-67, 103]) that the eigenvalue statistics of each H_a , $a \in \llbracket D \rrbracket$, around a bulk energy $E \in [-2 + \kappa, 2 - \kappa]$ and under a scaling N converge to a determinantal point process with sine-kernel. Hence, the eigenvalue statistics of H_Λ around E and under a scaling DN will converge to D independent copies of sine processes as $N \to \infty$. In particular, if we consider the limit $N \to \infty$ followed by $D \to \infty$, the bulk eigenvalue statistics $DN(\lambda_k - E)$ is expected to converge to a Poisson point process whose density is determined by the limiting density ρ_N defined in (2.18) below.

Remark 2.6. We expect that for $k \in [\![1, \kappa DN]\!]$, the transition threshold from the integrable phase to the chaotic phase should be

$$\|A\|_{\rm HS} \sim N^{1/3} / k^{1/3}. \tag{2.10}$$

In fact, the proofs for the bulk case in this paper suggest that the interaction Λ induces an eigenvalue perturbation of order $N^{-1}||A||_{\text{HS}}$. We expect chaos to enter when this perturbation can mix neighboring energy levels. On the other hand, the typical energy gap between the k-th and (k + 1)-th eigenvalues of H near the spectral edge is known to be of order $N^{-2/3}k^{-1/3}$. Thus, the transition should occur when

$$N^{-1} \|A\|_{\mathrm{HS}} \sim N^{-2/3} k^{-1/3},$$

which leads to the threshold (2.10). We leave the detailed study of the edge case for future research.

To summarize, Theorems 2.2 and 2.3 together indicate the chaotic behavior of the whole system under (2.4), while Theorems 2.4 and 2.5 indicate the non-chaotic/integrable behavior of the whole system under (2.7). Hence, we have established the quantum chaos transition of the random matrix model (1.1) as $||A||_{\text{HS}}$ crosses the threshold 1.

2.2. Local law. One basic tool for our proof is the local law for the Green's function (or resolvent) of H_{Λ} ,

$$G(z) \equiv G(z, H, \Lambda) := (H_{\Lambda} - z)^{-1}, \quad z \in \mathbb{C}_{+} := \{ z \in \mathbb{C} : \text{Im} \, z > 0 \},$$
(2.11)

as we will state in Lemma 2.10 below. Note the model (1.1) can be regarded as a deformed generalized Wigner matrix. Many methods have been developed in the literature (see e.g., [9, 58, 72, 78, 84]) to show that in the $N \to \infty$ limit, G(z) converges to a deterministic matrix M(z) in the sense of local laws (see Lemma 2.10). Moreover, $M(z) \equiv M(z, \Lambda)$ satisfies the matrix Dyson equation:

$$\left(\mathcal{S}(M) + z - \Lambda\right)M + I = 0, \qquad (2.12)$$

where $\mathcal{S}(\cdot)$ is a linear operator acting on M such that $\mathcal{S}(M)$ is a diagonal matrix with entries

$$\mathcal{S}(M)_{ij} = \mathbf{1}(i=j)\sum_{x} s_{ix}M_{xx} = \mathbf{1}(i=j)D\langle ME_a\rangle, \quad i,j\in\mathcal{I}_a.$$

Hereafter, we denote the variances of the entries of H by

$$s_{ij} = \mathbb{E}|H_{ij}|^2 = N^{-1}\mathbf{1}(i, j \in \mathcal{I}_a \text{ for some } a \in \llbracket D \rrbracket),$$
(2.13)

and let $S = (s_{ij} : i, j \in \mathcal{I})$ be the variance matrix. In addition, we use $\langle B \rangle := (DN)^{-1} \operatorname{Tr} B$ to denote the normalized trace of a $DN \times DN$ matrix B. Due to the block translation symmetry of S and Λ , we see that M is also block translation invariant, which implies that $\mathcal{S}(M)$ should be a scalar matrix $\mathcal{S}(M) = mI$, where m(z) is defined as $m(z) := \langle M(z) \rangle$.

Remark 2.7. When D = 2, the block translation symmetry may not hold. In this case, we denote

$$M = \begin{pmatrix} M_{(11)} & M_{(12)} \\ M_{(21)} & M_{(22)} \end{pmatrix}$$

Then, we can derive directly from equation (2.12) that

$$M_{(11)} = \frac{m+z}{AA^* - (m+z)^2}, \quad M_{(22)} = \frac{m+z}{A^*A - (m+z)^2},$$

$$M_{(12)} = \frac{1}{AA^* - (m+z)^2}A, \quad M_{(21)} = \frac{1}{A^*A - (m+z)^2}A^*,$$
(2.14)

where m(z) satisfies the self-consistent equation $m(z) = N^{-1} \operatorname{Tr} M_{(11)}(z) = N^{-1} \operatorname{Tr} M_{(22)}(z)$.

Definition 2.8 (Matrix limit of G). We define $m(z) \equiv m_N(z)$ as the unique solution to

$$m(z) = \left\langle \left(\Lambda - z - m(z)\right)^{-1} \right\rangle \tag{2.15}$$

such that $\operatorname{Im} m(z) > 0$ whenever $z \in \mathbb{C}_+$. Then, we define the matrix $M(z) \equiv M_N(z, \Lambda)$ as

$$M(z) := (\Lambda - z - m(z))^{-1}.$$
(2.16)

Since Λ is Hermitian, we have that $m(\overline{z}) = \overline{m(z)}$ and $M(\overline{z}) = M(z)^*$.

Under this definition, m(z) is actually the Stieltjes transform of a probability measure μ_N , called the *free convolution* of the empirical measure of Λ and the semicircle law with density

$$\rho_{sc}(x) = \frac{1}{2\pi} \sqrt{4 - x^2} \mathbf{1}_{x \in [-2,2]}.$$
(2.17)

Moreover, the probability density ρ_N of μ_N is determined from m(z) by

$$\rho_N(x) = \pi^{-1} \lim_{\eta \downarrow 0} \operatorname{Im} m(x + \mathrm{i}\eta).$$
(2.18)

Under the assumption ||A|| = o(1), the support of ρ_N is a single interval $[-a_N, b_N]$ with $|a_N - 2| + |b_N - 2| = o(1)$, and m(z) is close to the Stieltjes transform of ρ_{sc} given by $m_{sc}(z) = (-z + \sqrt{z^2 - 4})/2$. In this paper, we focus on the eigenvalues and eigenvectors of H_Λ in the bulk $[-2 + \kappa, 2 - \kappa]$ for some constant $\kappa > 0$, away from the edges $-a_N$ and b_N . We define the quantiles γ_k , $k \in \mathcal{I}$, of ρ_N as

$$\gamma_k := \sup_{x \in \mathbb{R}} \left\{ \int_{-\infty}^x \rho_N(x) \mathrm{d}x < \frac{k}{DN} \right\}.$$
(2.19)

To state the local law and streamline the presentation, in this paper, we adopt the following convenient notion of stochastic domination introduced in [54].

Definition 2.9 (Stochastic domination and high probability event). (i) Let

$$\xi = \left(\xi^{(N)}(u) : N \in \mathbb{N}, u \in U^{(N)}\right), \quad \zeta = \left(\zeta^{(N)}(u) : N \in \mathbb{N}, u \in U^{(N)}\right)$$

be two families of non-negative random variables, where $U^{(N)}$ is a possibly N-dependent parameter set. We say ξ is stochastically dominated by ζ , uniformly in u, if for any fixed (small) $\tau > 0$ and (large) D > 0,

$$\mathbb{P}\bigg(\bigcup_{u\in U^{(N)}}\left\{\xi^{(N)}(u)>N^{\tau}\zeta^{(N)}(u)\right\}\bigg)\leq N^{-D}$$

for large enough $N \ge N_0(\tau, D)$, and we will use the notation $\xi \prec \zeta$. If for some complex family ξ we have $|\xi| \prec \zeta$, then we will also write $\xi \prec \zeta$ or $\xi = O_{\prec}(\zeta)$.

(ii) As a convention, for two deterministic non-negative quantities ξ and ζ , we will write $\xi \prec \zeta$ if and only if $\xi \leq N^{\tau} \zeta$ for any constant $\tau > 0$.

(iii) Let A be a family of random matrices and ζ be a family of non-negative random variables. Then, we use $A = O_{\prec}(\zeta)$ to mean that $||A|| \prec \xi$, where $|| \cdot ||$ denotes the operator norm.

(iv) We say an event Ξ holds with high probability (w.h.p.) if for any constant D > 0, $\mathbb{P}(\Xi) \ge 1 - N^{-D}$ for large enough N. More generally, we say an event Ω holds w.h.p. in Ξ if for any constant D > 0, $\mathbb{P}(\Xi \setminus \Omega) \le N^{-D}$ for large enough N.

Lemma 2.10 (Local laws and rigidity of eigenvalues). Under Assumption 2.1, for any small constant $\tau > 0$, the following local laws hold uniformly in $z = E + i\eta$ with $|z| \le \tau^{-1}$ and $\eta \ge N^{-1+\tau}$.

• Anisotropic local law: For any deterministic unit vectors $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{DN}$, we have

$$(G(z) - M(z))_{\mathbf{uv}} \prec \sqrt{\frac{\operatorname{Im} m(z)}{N\eta} + \frac{1}{N\eta}}.$$
(2.20)

• Averaged local law: For any deterministic matrix $B \in \mathbb{C}^{DN \times DN}$ with $||B|| \leq 1$, we have

$$\langle (G-M)B \rangle \prec \frac{1}{N\eta}.$$
 (2.21)

As a consequence of (2.21) when B = I, we have the rigidity of eigenvalues:

$$\lambda_k - \gamma_k \prec N^{-2/3} \min(k, DN + 1 - k)^{-1/3}, \quad k \in \mathcal{I}.$$
 (2.22)

In addition, all the above estimates remain valid even if we do not assume identical distributions for the diagonal and off-diagonal entries of H.

With the anisotropic local law (2.20), we can derive the following estimate on the products of resolvents. The proofs of Lemma 2.10 and Lemma 2.11 are both presented in Appendix C.

Lemma 2.11. Fix any integer $p \ge 1$. Suppose $(\Lambda_i)_{1 \le i \le p}$ is an arbitrary sequence of $D \times D$ block matrices of the same form as Λ and consisting of $N \times N$ deterministic blocks A_i and A_i^* with $||A_i|| = o(1)$. Let $(B_i)_{1 \leq i \leq p}$ be an arbitrary sequence of deterministic matrices satisfying $||B_i|| \leq 1$. Suppose the anisotropic local law (2.20) holds for all G_i , where $G_i := G(z_i, H, \Lambda_i)$ for a sequence $(z_i)_{1 \leq i \leq p}$ with $z_i \in \{z, \overline{z}\}$. Then, for any deterministic unit vectors $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{DN}$, we have

$$\mathbf{u}^* \left(\prod_{i=1}^p G_i B_i\right) \mathbf{v} \prec \frac{1}{\eta^{p-1}}.$$
(2.23)

2.3. **Proof ideas.** In this subsection, we discuss some key ideas for the proof of the main results.

Integrable regime. We define a sequence of interpolating matrices as

$$H_{\Lambda}(\theta) := H + \theta \Lambda, \quad \theta \in [0, 1], \quad \text{with} \quad H_{\Lambda}(0) = H, \quad H_{\Lambda}(1) = H_{\Lambda}.$$
(2.24)

By the standard perturbation theory for eigenvalues, we have $\lambda'_k(\theta) = \mathbf{v}_k(\theta)^* \Lambda \mathbf{v}_k(\theta)$, where k is a bulk index, $\lambda_k(\theta)$ is the k-th eigenvalue of $H_{\Lambda}(\theta)$, and $\mathbf{v}_k(\theta)$ denotes the corresponding eigenvector. Thus, the shift of the eigenvalue λ_k is of order $\int_0^1 \mathbb{E} |\mathbf{v}_k(\theta)^* \Lambda \mathbf{v}_k(\theta)| d\theta \leq \int_0^1 (\mathbb{E} |\mathbf{v}_k(\theta)^* \Lambda \mathbf{v}_k(\theta)|^2)^{1/2} d\theta$. With the spectral decomposition of $G_{\theta}(z) := (H_{\Lambda}(\theta) - z)^{-1}$, we can show that

$$\mathbb{E}|\mathbf{v}_k(\theta)^* \Lambda \mathbf{v}_k(\theta)|^2 \prec \eta^2 \mathbb{E} \operatorname{Tr}\left[\operatorname{Im} G_{\theta}(z_k(\theta)) \Lambda \operatorname{Im} G_{\theta}(z_k(\theta)) \Lambda\right], \quad z_k(\theta) := \gamma_k(\theta) + \mathrm{i}\eta, \quad (2.25)$$

where Im $G_{\theta} := (G_{\theta} - G_{\theta}^*)/(2i)$, $\eta = N^{-1+c}$ for an arbitrary small constant c > 0, and $\gamma_k(\theta)$ is the classical location of $\lambda_k(\theta)$ defined as in (2.19). Then, we will prove a key technical estimate that

$$\mathbb{E}\operatorname{Tr}\left[\operatorname{Im} G_{\theta}(z_{k}(\theta))\Lambda\operatorname{Im} G_{\theta}(z_{k}(\theta))\Lambda\right] \prec \|A\|_{\mathrm{HS}}^{2}.$$
(2.26)

Together with (2.25), it implies that as long as $||A||_{\text{HS}} \leq N^{-\varepsilon_A}$ and $c < \varepsilon_A/2$, Λ leads to a perturbation of λ_k of order O($N^{-1-\varepsilon_A/2}$) with probability 1-o(1), which is negligible under the scaling DN.

Now, we have shown that $\lambda_k(1)$ is a perturbation of the eigenvalue $\lambda_k(0)$ of H. Suppose $\lambda_k(0)$ is the eigenvalue of a diagonal block H_a , $a \in [D]$. By the bulk universality of Wigner matrices [27], we know that conditioning on $\lambda_k(0)$, the eigenvalue spectra of other blocks H_b , $b \neq a$, are away from λ_k by a distance of order N^{-1} with probability 1 - o(1). Without loss of generality, suppose $a \neq 1$. We write \mathbf{v}_k as $\mathbf{v}_k = (\mathbf{u}_k^\top, \mathbf{w}_k^\top)^\top$ for $\mathbf{u}_k \in \mathbb{C}^N$ and $\mathbf{w}_k \in \mathbb{C}^{(D-1)N}$, and let $\widetilde{A} := (A, \mathbf{0}_{N \times (D-3)N}, A^*)$ denote the upper right $N \times (D-1)N$ block of H_{Λ} . Then, from the eigenvalue equation $H_{\Lambda} \mathbf{v}_k = \lambda_k \mathbf{v}_k$, we get that

$$H_1 \mathbf{u}_k + \widetilde{A} \mathbf{w}_k = \lambda_k \mathbf{u}_k \Rightarrow \mathbf{u}_k = -(H_1 - \lambda_k)^{-1} \widetilde{A} \mathbf{w}_k$$

Our intuition is that H_1 and \mathbf{w}_k are almost independent in the integrable regime, so when dist(λ_k , spec(H_1)) \gtrsim N^{-1} , $\|\mathbf{u}_k\| = \|(H_1 - \lambda_k)^{-1} \widetilde{A} \mathbf{w}_k\|$ should be small. In fact, we will prove that

$$\mathbb{E}\|\mathbf{u}_k\|^2 = \mathbb{E}\|\left(H_1 - \lambda_k\right)^{-1} \widetilde{A} \mathbf{w}_k\|^2 \prec N^{\tau + 2c} \mathbb{E} \operatorname{Tr}\left[\operatorname{Im} G_0(z_k) \Lambda \operatorname{Im} G(z_k) \Lambda\right]_{11}$$

for any small constant $\tau > 0$, where G_0 denotes the resolvent of H and $z_k \equiv z_k(\theta = 1)$ is defined in (2.25). For the right-hand side (RHS), we will prove a similar bound as in (2.26):

$$\mathbb{E}\operatorname{Tr}\left[\operatorname{Im} G_0(z_k)\Lambda\operatorname{Im} G(z_k)\Lambda\right] \prec ||A||_{\mathrm{HS}}^2.$$
(2.27)

Thus, as long as $||A||_{\text{HS}} \leq N^{-\varepsilon_A}$ and $\tau + 2c < \varepsilon_A$, we obtain that $||\mathbf{u}_k|| \leq N^{-\varepsilon_A/2}$ with probability 1 - o(1). Applying this argument to all blocks H_b with $b \neq a$, we can show that the ℓ_2 -norm of \mathbf{v}_k within each \mathcal{I}_b is negligible, implying that \mathbf{v}_k must be concentrated on the subset \mathcal{I}_a .

The proof of the *multi-resolvent estimates* (2.26) and (2.27) constitutes the main technical part of the above argument. We will apply the cumulant expansions formula in [71,77] iteratively to bound the left-hand sides (LHS) of (2.26) and (2.27). A key point of the proof is that we need to find a systematic way to expand the resolvent entries so that after each expansion, the resulting expressions become strictly smaller until they satisfy the desired bound $||A||^2_{\text{HS}}$. For this purpose, we need to keep track of the fine structures of the expressions and find the "correct" resolvent entry to expand; see Section 8.1 for more details.

Chaotic regime. By Markov's inequality, the QUE estimate (2.5) follows immediately from the second moment bound $\mathbb{E}[||E_a \mathbf{v}_k||^2 - D^{-1}]^2 \leq N^{-\delta}$ for a constant $\delta > 0$. To prove this bound, we again need to establish a multi-resolvent estimate for G(z). More precisely, when $||A||_{\text{HS}} \geq N^{\varepsilon_A}$, we aim to show that

$$\mathbb{E}\langle \operatorname{Im} G(z)(E_a - D^{-1}) \operatorname{Im} G(z)(E_a - D^{-1}) \rangle \le N^{-1-\delta} \eta^{-2}, \quad a \in [\![D]\!],$$
(2.28)

for a constant $\delta > 0$ depending on ε_A . With the spectral decomposition of Im G, we obtain from (2.28) that

$$\mathbb{E}\eta^4 \sum_{i,j\in\mathcal{I}} \frac{|\mathbf{v}_i^*(E_a - D^{-1})\mathbf{v}_j|^2}{|\lambda_i - z|^2 |\lambda_j - z|^2} \le N^{-\delta}, \quad z = \gamma_k + \mathrm{i}\eta, \quad a \in \llbracket D \rrbracket.$$

From the above equation, we see that a bound of the form (2.28) at $\eta = N^{-1+c}$ implies that a local sum of $|\mathbf{v}_i^*(E_a - D^{-1})\mathbf{v}_j|^2$ in $|i - k| = O(N\eta)$ and $|j - k| = O(N\eta)$ is bounded by $N^{-\delta}$. In particular, by taking i = j = k, it gives the desired estimate.

Let $z_1, z_2 \in \{z, \overline{z}\}$ and abbreviate $G_i := G(z_i)$. Writing $\operatorname{Im} G(z) = (G(z) - G(\overline{z}))/(2i)$ and $I = \sum_a E_a$, we can expand the LHS of (2.28) into a linear combination of two-resolvent traces of the form $\mathbb{E}L_{ab}, a, b \in [D]$, with $L_{ab} := D\langle G_1 E_a G_2 E_b \rangle$. Then, a key point of our proof is to establish the following two-resolvent local law for $\mathbb{E}L_{ab}$: there exists a deterministic matrix K and a constant $\delta' > 0$ so that

$$|\mathbb{E}L_{ab} - K_{ab}| \le N^{-1-\delta'} \eta^{-2}.$$
(2.29)

This estimate shows that $\mathbb{E}L_{ab}$ is approximated by K_{ab} up to a negligible error. In addition, we will show that the deterministic part K is "flat" when $||A||_{\text{HS}} \gg 1$, that is, $\max_{a,b} |K_{ab} - K_{11}| = O(N/||A||_{\text{HS}}^2)$. It implies that the leading part of the LHS of (2.28) is of order $O(N/||A||_{\text{HS}}^2) \ll N^{-1-\varepsilon_A}\eta^{-2}$ as long as we take $2c < \varepsilon_A$. This observation, combined with (2.26), elucidates the essence of the quantum chaos transition occurring at $||A||_{\text{HS}} \sim 1$.

We will see that, without the expectation, L_{ab} satisfies the following (essentially sharp) local law:

$$|L_{ab} - K_{ab}| \prec N^{-1} \eta^{-2}. \tag{2.30}$$

Multi-resolvent local laws as in (2.30) have been an important development in the random matrix theory literature over the past few years. Specifically, (almost) sharp multi-resolvent local laws have been established for Wigner matrices in [40, 42, 44, 45, 47, 49] and they have played an important role in proving the eigenstate thermalization hypothesis for Wigner matrices [40, 42, 43, 45]. Later, they are extended to deformed Wigner matrices [41] and generalized Wigner matrices with mean-field variance profiles [1]. In these works, the canonical proof is to derive a self-consistent equation for L and then bound the fluctuation error carefully using the cumulant expansion method. However, their proofs depend crucially on the flat (or some special mean-field) variance profiles of the random matrices, which do not apply to our block variance profile setting. Instead, we will adopt a dynamical approach called the method of *characteristic flow*. This idea of estimating resolvents dynamically first appeared in [83], and was later applied to various models [2, 3, 26, 75, 79, 80] to show single-resolvent local laws (or closely related quantities). More recently, its usefulness in establishing multi-resolvent local laws has also been recognized in [29,40,46], as well as the concurrently posted paper [48]. However, in comparison to these works, our specific setting presents two distinct challenges: the non-meanfield nature of the problem and the arbitrary interaction A. Moreover, we need to improve the estimate (2.30) to (2.29) by a small factor $N^{-\delta'}$.

In the proof of (2.30), we will consider a matrix-valued process $H_{\Lambda}(t)$, where each diagonal block of H_{Λ} satisfies an independent matrix-valued Ornstein-Uhlenbeck (OU) process $H_a(t)$ with initial condition H_a , and each off-diagonal block satisfies a matrix-valued linear differential equation. Then, we let the spectral parameter z_t of the resolvent $G_t := (H_{\Lambda}(t) - z_t)^{-1}$ satisfy a first-order differential equation, which is indeed the characteristic flow of the underlying complex Burgers equation. Along this flow, the leading terms in the time evolution of G_t cancel each other, and $\eta_t = \text{Im } z_t$ decreases, which enables us to transfer the estimates (2.29) and (2.30) from larger scales of η to smaller scales at the expense of an added Gaussian component. But, we can easily remove it with a standard Green's function comparison argument. The aforementioned flow approach comprises two components: an initial estimate at t = 0 and large η_0 , followed by a flow estimate for the subsequent time evolution. One interesting observation is that the initial estimate, often believed to be simple at large η , actually becomes a major obstacle when attempting to improve (2.30) to (2.29). More precisely, when we expand $L_{ab} - K_{ab}$ using cumulant expansions, besides some centered random fluctuations, we also find leading deterministic terms in L_{ab} of exact order $N^{-1}\eta^{-2}$. We will need to obtain the exact expressions of these terms and show an intricate cancellation between them through an explicit calculation. However, we remark that this cancellation is not accidental and indeed corresponds to a deep self-energy renormalization mechanism originally identified in the context of random band matrices [108]; see Remark 4.15 below. We believe that this kind of self-energy renormalization is a conceptually universal mechanism behind the chaotic behavior of non-mean-field random models.

Finally, we remark that $L_{ab} = N^{-1} \sum_{x \in \mathcal{I}_b, y \in \mathcal{I}_a} G_{xy}(z_1) G_{yx}(z_2)$ corresponds to the *T*-variable in the study of random band matrices, so the *T*-expansion method developed in [32, 56, 107–109, 111] is potentially useful for the proof of the two-resolvent local laws for *L*. Although it is not as concrete as the flow method in the current setting, the *T*-expansion method can be crucial when we consider extensions to the large *D* case later.

Given the QUE estimates in Theorem 2.2, we can prove the GOE/GUE statistics for the bulk eigenvalues of H_{Λ} adopting an idea developed recently in [107]. More precisely, similar to many previous proofs of bulk universality of random matrices, our starting point is the three-step strategy initiated in [59, 62]. Readers can refer to [64] for an overview of this strategy. The first step is to establish a local law for the resolvent G(z) up to the optimal scale $\eta \gg N^{-1}$, which has been accomplished in Lemma 2.10. For the second step, one considers the matrix Brownian motion $H_{\Lambda}(t) = H_t + \Lambda$, where $H_t = (h_{ij}(t))_{i,j \in \mathcal{I}}$ is a matrix-valued OU process. Note that in contrast to the flow $H_{\Lambda}(t)$ defined above in the characteristic flow approach, H_t here satisfies a $DN \times DN$ matrix-valued OU equation (see (6.1) for the definition) and Λ remains unchanged. With the local law as main input, it was proved in [81,82] that the local bulk statistics of $H_{\Lambda}(t)$ converge to those of GOE/GUE at time $t \gg N^{-1}$. In the third step, one needs to show that the local bulk statistics of the original matrix H_{Λ} are well-approximated by those of $H_{\Lambda}(t)$, which is achieved by comparing the moments of $(\operatorname{Im} G(z))$ with those of $(\operatorname{Im} G_t(z))$, where $G_t(z) := (H_\Lambda(t) - z)^{-1}$ denotes the resolvent of $H_\Lambda(t)$. In the literature (see e.g., [59, 62, 64, 67]), this step is typically accomplished through a standard Green's function comparison approach via moments matching between the entries of H_{Λ} and those of $H_{\Lambda}(t)$. However, this kind of moment-matching argument fails in our specific setting. To address this challenge, we adopt the idea in [107], that is, we utilize the QUE of bulk eigenvectors to perform the Green's function comparison. We will see that the leading terms appearing in the comparison of $\langle \operatorname{Im} G(z) \rangle$ with $\langle \operatorname{Im} G_t(z) \rangle$ will contain factors of the form $\mathbf{v}_i^*(E_a - D^{-1})\mathbf{v}_i$. These factors are small as shown in the QUE estimate (2.6), which will help with bounding the leading terms in the comparison. This allows us to prove that the local bulk statistics of H_{Λ} match those of $H_{\Lambda}(t)$ asymptotically.

3. Chaotic regime: eigenvectors

In this section, we prove Theorem 2.2. As discussed in Section 2.3, the main ingredient for the proof is a precise two-resolvent local law on quantities of the form $\langle G(z_1)E_aG(z_2)E_b\rangle$, $a, b \in [D]$. We first introduce several notations.

Definition 3.1. Define the spectral domain $\mathbf{D}(\kappa, \tau) := \{z = E + i\eta \in \mathbb{C} : |z| \le \tau^{-1}, |E| \le 2-\kappa, |\eta| \ge N^{-1+\tau}\}$ for an arbitrarily small constant $\tau > 0$. For $z_1, z_2 \in \mathbf{D}(\kappa, \tau)$, we define the $D \times D$ matrices \widehat{M} and L as

$$M_{ab}(z_1, z_2, \Lambda) := D\langle M(z_1)E_a M(z_2)E_b \rangle, \quad L_{ab}(z_1, z_2, H, \Lambda) := D\langle G(z_1)E_a G(z_2)E_b \rangle, \tag{3.1}$$

for $a, b \in \llbracket D \rrbracket$, and define the $D \times D$ matrix K by

$$K(z_1, z_2, \Lambda) := \left[1 - \widehat{M}(z_1, z_2, \Lambda)\right]^{-1} \widehat{M}(z_1, z_2, \Lambda).$$
(3.2)

For ease of presentation, we introduce the following simplified notations: given a matrix-valued function (e.g., G, M, \widehat{M}, L , and K) of z, we use subscripts to indicate its dependence on the spectral parameters. For example, we will denote $G_i := G(z_i, H, \Lambda), M_i := M(z_i, \Lambda), \widehat{M}_{(1,2)} := \widehat{M}(z_1, z_2, \Lambda), L_{(1,2)} := L(z_1, z_2, H, \Lambda),$ and $K_{(1,2)} := K(z_1, z_2, \Lambda)$.

Lemma 3.2. Take any $z = E + i\eta \in \mathbf{D}(\kappa, \tau)$ with $\eta = N^{-1+\varepsilon_L}$ for a small constant $\varepsilon_L > 0$. Then, under the assumptions of Theorem 2.2, there exists a constant $c_L > 0$ (depending on ε_L , ε_A , and δ_A) such that

$$\left(\mathbb{E}L_{(1,2)} - K_{(1,2)}\right)_{ab} = \mathcal{O}(N^{-1-c_L}\eta^{-2})$$
(3.3)

for $z_1, z_2 \in \{z, \bar{z}\}$ and $a, b \in [\![D]\!]$.

Before proving this key lemma, we first use it to prove Theorem 2.2.

Proof of Theorem 2.2. For $z = E + i\eta$, using the spectrum decomposition of Im G(z), we get that for any $DN \times DN$ matrix B,

$$\operatorname{Tr}\left[\operatorname{Im} G(z)B\operatorname{Im} G(z)B^*\right] = \eta^2 \sum_{i,j\in\mathcal{I}} \frac{|\mathbf{v}_i^*B\mathbf{v}_j|^2}{|\lambda_i - z|^2|\lambda_j - z|^2}$$

In particular, choosing $B = E_a - D^{-1}I$ and $z_k = \gamma_k + iN^{-1+\varepsilon_L}$ and using the rigidity of eigenvalues in (2.22), we get from this estimate that for any constant $c \in (0, \varepsilon_L)$,

$$\max_{i,j\in[[k-N^c,k+N^c]]} |\mathbf{v}_i^*(E_a - D^{-1})\mathbf{v}_j|^2 \prec N^{-2+2\varepsilon_L} \operatorname{Tr} \left[\operatorname{Im} G(z_k)(E_a - D^{-1}I) \operatorname{Im} G(z_k)(E_a - D^{-1}I)\right].$$
(3.4)

It remains to bound the RHS. By denoting $z_1 = z_k$, $z_2 = \bar{z}_k$ and using (3.3), its expectation is estimated as

$$-\frac{N^{-2+2\varepsilon_L}}{4}\mathbb{E}\operatorname{Tr}\left[(G_1 - G_2)\left(E_a - D^{-1}\sum_b E_b\right)(G_1 - G_2)\left(E_a - D^{-1}\sum_{b'} E_{b'}\right)\right]$$
$$= N^{-1+2\varepsilon_L}\left(\mathbb{E}\mathcal{L}_{aa} - \frac{2}{D}\sum_{b=1}^{D}\mathbb{E}\mathcal{L}_{ab} + \frac{1}{D^2}\sum_{b,b'=1}^{D}\mathbb{E}\mathcal{L}_{bb'}\right)$$
$$= N^{-1+2\varepsilon_L}\left(\mathcal{K}_{aa} - \frac{2}{D}\sum_{b=1}^{D}\mathcal{K}_{ab} + \frac{1}{D^2}\sum_{b,b'=1}^{D}\mathcal{K}_{bb'}\right) + O\left(N^{-c_L}\right), \qquad (3.5)$$

where the $D \times D$ matrices \mathcal{L} and \mathcal{K} are defined as $\mathcal{L} := (L_{(12)} + L_{(21)} - L_{(11)} - L_{(22)})/4$ and $\mathcal{K} := (K_{(12)} + K_{(21)} - K_{(11)} - K_{(22)})/4$. On the other hand, we claim that for $i, j \in \{1, 2\}$,

$$\max_{a,b,a',b' \in [D]} \left| \left(K_{(ij)} \right)_{ab} - \left(K_{(ij)} \right)_{a'b'} \right| = \mathcal{O} \left(N / \|A\|_{\mathrm{HS}}^2 \right).$$
(3.6)

We postpone the proof of this estimate to Lemma A.1 in the appendix. With (3.6), we obtain that

$$N^{-1+2\varepsilon_L} \left(\mathcal{K}_{aa} - \frac{2}{D} \sum_{b=1}^{D} \mathcal{K}_{ab} + \frac{1}{D^2} \sum_{b,b'=1}^{D} \mathcal{K}_{bb'} \right) \lesssim N^{-2\varepsilon_A + 2\varepsilon_L}.$$
(3.7)

Combining (3.4), (3.5), and (3.7), we obtain that for any small constant $\varepsilon > 0$,

$$\mathbb{E} \max_{i,j \in \llbracket k-n^c, k+n^c \rrbracket} |\mathbf{v}_i^* (E_a - D^{-1}) \mathbf{v}_j|^2 \le N^{-c_L + \varepsilon} + N^{-2\varepsilon_A + 2\varepsilon_L + \varepsilon}.$$
(3.8)

If we take $\varepsilon_L < \varepsilon_A/2$ and $\varepsilon < (c_L \wedge \varepsilon_A)/2$, this gives that

$$\mathbb{E} \max_{i,j \in \llbracket k-n^c, k+n^c \rrbracket} |\mathbf{v}_i^*(E_a - D^{-1})\mathbf{v}_j|^2 \le N^{-c_L/2} + N^{-\varepsilon_A/2}.$$

Then, applying Markov's inequality and a simple union bound over $a \in \llbracket D \rrbracket$ concludes (2.6). Taking i = j = k, we obtain (2.5).

For the proof of Lemma 3.2, we first establish it for a special case with Gaussian divisible H. Its proof will be presented in Section 4.

Lemma 3.3. Under the setting of Theorem 2.2, suppose H_a , $a \in [D]$, are of the form

$$H_{a} = \sqrt{1 - N^{-\varepsilon_{g}}} \cdot H_{a}^{(0)} + N^{-\varepsilon_{g}/2} H_{a}^{(g)}, \qquad (3.9)$$

where $H_a^{(0)}$ are independent Wigner matrices satisfying the assumptions for H_a in Assumption 2.1 and $H_a^{(g)}$ are i.i.d. GUE satisfying (2.1) and (2.2). Then, for small enough constant $\varepsilon_g > 0$ (depending on ε_L and ε_A) and $z = E + i\eta$ with $|E| \leq 2 - \kappa$, there exists an absolute constant C > 1 such that

$$\|\mathbb{E}L_{(1,2)} - K_{(1,2)}\|_{\mathrm{HS}} \prec N^{-1 - (\varepsilon_g \wedge \delta_A)} \eta^{-2}, \quad for \quad N^{-1 + C\varepsilon_g} \leq \eta \leq N^{-C\varepsilon_g} \ , \ z_1, z_2 \in \{z, \bar{z}\}.$$
(3.10)

Now, Lemma 3.2 follows from Lemma 3.3 combined with a Green's function comparison result stated in Lemma 3.4. We postpone the proof of Lemma 3.4 to Appendix B.

Lemma 3.4. Let H and H be two matrices satisfying Assumption 2.1. Suppose they satisfy the following moment-matching conditions: for $i, j \in \mathcal{I}$ and integers $k, l \ge 0$,

$$\mathbb{E}(H_{ij})^k (H_{ij}^*)^l - \mathbb{E}(\widetilde{H}_{ij})^k (\widetilde{H}_{ij}^*)^l = 0 \quad for \quad k+l \le 3,$$

$$(3.11)$$

and there exists a constant $\delta \in (0, 1/2)$ such that

$$\left| \mathbb{E}(H_{ij})^k (H_{ij}^*)^l - \mathbb{E}(\widetilde{H}_{ij})^k (\widetilde{H}_{ij}^*)^l \right| \lesssim N^{-2-\delta} \quad for \quad k+l = 4.$$

$$(3.12)$$

Then, for any $z \in \mathbf{D}(\kappa, \tau)$, $z_1, z_2 \in \{z, \overline{z}\}$, and $a, b \in \llbracket D \rrbracket$,

$$\mathbb{E}\langle G_1 E_a G_2 E_b \rangle - \mathbb{E} \langle \widetilde{G}_1 E_a \widetilde{G}_2 E_b \rangle \prec N^{-1-\delta} \eta^{-2}, \tag{3.13}$$

where $\widetilde{G}_i \equiv G(z_i, \widetilde{H}, \Lambda), i \in \{1, 2\}$, denote the Green's functions of \widetilde{H} .

Proof of Lemma 3.2. Given the matrix H considered in Lemma 3.2, we can construct another random matrix \tilde{H} satisfying the setting in Lemma 3.3 and such that the moment-matching conditions (3.11) and (3.12) hold with $\delta = \varepsilon_g$ (see e.g., Lemma 6.5 in [66]). By Lemma 3.3, as long as we choose ε_g small enough such that $C\varepsilon_g \leq \varepsilon_L \leq 1 - C\varepsilon_g$, there is

$$D\mathbb{E}\langle \widetilde{G}_1 E_a \widetilde{G}_2 E_b \rangle - (K_{(1,2)})_{ab} \prec N^{-1 - (\varepsilon_g \wedge \delta_A)} \eta^{-2}$$

for $\eta = N^{-1+\varepsilon_L}$. On the other hand, by Lemma 3.4, we have that

$$\mathbb{E}\langle G_1 E_a G_2 E_b \rangle - \mathbb{E}\langle \widetilde{G}_1 E_a \widetilde{G}_2 E_b \rangle \prec N^{-1-\varepsilon_g} \eta^{-2}$$

Combining the above two estimates, we conclude Lemma 3.2 by choosing $c_L < \varepsilon_q \wedge \delta_A$.

4. Gaussian divisible case

In this section, we prove the key lemma—Lemma 3.3—in the proof of Theorem 2.2. We first introduce some new notations that are similar to those in Definition 3.1 but with three z arguments.

Definition 4.1. Define the $D \times D \times D$ tensors L and K as

$$[L(z_1, z_2, z_3, H, \Lambda)]_{a_1 a_2 a_3} := D \langle G_1 E_{a_1} G_2 E_{a_2} G_3 E_{a_3} \rangle,$$

$$[K(z_1, z_2, z_3, \Lambda)]_{a_1 a_2 a_3} = \sum_{b_1, b_2, b_3} (I - \widehat{M}_{(1,2)})_{a_1 b_1}^{-1} (I - \widehat{M}_{(2,3)})_{a_2 b_2}^{-1} (I - \widehat{M}_{(3,1)})_{a_3 b_3}^{-1} D \langle M_1 E_{b_1} M_2 E_{b_2} M_3 E_{b_3} \rangle,$$

for $a_i \in \llbracket D \rrbracket$, $i \in \{1, 2, 3\}$. Here, we have abused the notations a little bit and still use L and K to denote these tensors. Moreover, we will also abbreviate them by $L_{(1,2,3)}$ and $K_{(1,2,3)}$.

As discussed in Section 2.3, the proof of Lemma 3.3 is based on the characteristic flow method in [83]. We start with several estimates at the initial scale $\eta \sim N^{-\varepsilon_g}$ as stated in Lemmas 4.2 and 4.3. Their proofs will be postponed to Sections 4.1 and 4.2.

Lemma 4.2. Under the assumptions of Theorem 2.2, take any $z = E + i\eta$ with $|E| \le 2 - \kappa$ and $\eta \sim N^{-\varepsilon_g}$. Then, for any constant $\varepsilon_g \in (0, 1/4)$ and $z_i \in \{z, \overline{z}\}$, $i \in \{1, 2, 3\}$, we have that

$$\|L_{(1,2)} - K_{(1,2)}\|_2 \prec N^{-1} \eta^{-3}, \tag{4.1}$$

$$\|L_{(1,2,3)} - K_{(1,2,3)}\|_2 \prec N^{-1}\eta^{-4}, \tag{4.2}$$

where $\|\cdot\|_2$ denotes the ℓ_2 -norm by regarding matrices and tensors as vectors (for matrices, it is the Hilbert-Schmidt norm).

Lemma 4.3. Under the assumptions of Theorem 2.2, take any $z = E + i\eta$ with $|E| \le 2 - \kappa$ and $\eta \sim N^{-\varepsilon_g}$. Then, for any constant $\varepsilon_g \in (0, 1/4)$ and $z_i \in \{z, \overline{z}\}$, $i \in \{1, 2\}$, we have that

$$\max_{a \in \llbracket D \rrbracket} |\mathbb{E} \langle (G(z) - M(z)) E_a \rangle| \prec N^{-1},$$
(4.3)

$$\|\mathbb{E}L_{(1,2)} - K_{(1,2)}\|_2 \prec N^{-1}\eta^{-2} \left(\eta + N^{-\delta_A}\right).$$
(4.4)

Next, we apply the flow method to transfer the bounds at larger scales of η to smaller scales of η .

Definition 4.4 (Characteristic flow). Given a starting time $t_0 \in \mathbb{R}$ and initial values (z_{t_0}, Λ_{t_0}) , we define flows of z and Λ as

$$\frac{\mathrm{d}}{\mathrm{d}t}z_t = -\frac{1}{2}z_t - \langle M_t \rangle, \quad \frac{\mathrm{d}}{\mathrm{d}t}\Lambda_t = -\frac{1}{2}\Lambda_t, \quad t \ge t_0, \tag{4.5}$$

where $M_t := M(z_t, \Lambda_t)$ is the solution to (2.12) with z and Λ replaced by z_t and Λ_t . Let $t_c := \inf\{t \ge t_0 : \lim z_{t_c} = 0\}$ be the first time $\operatorname{Im} z_t$ vanishes. We also introduce the function $Z : \mathbb{C} \times \mathbb{C}^{DN \times DN} \to \mathbb{C}^{DN \times DN}$ as $Z(z, \Lambda) := zI - \Lambda$ and abbreviate that $Z_t := Z(z_t, \Lambda_t)$. Note that Z_t satisfies

$$\frac{\mathrm{d}}{\mathrm{d}t}Z_t = -\frac{1}{2}Z_t - \langle M_t \rangle. \tag{4.6}$$

Given the initial random matrix H_{t_0} satisfying Assumption 2.1 with diagonal blocks $(H_a)_{t_0}$, $a \in [D]$, we define the flow H_t as a $DN \times DN$ random matrix with diagonal blocks $(H_a)_t$ being matrix-valued OU processes

$$d(H_a)_t = -\frac{1}{2}(H_a)_t dt + \frac{1}{\sqrt{N}} d(B_a)_t,$$
(4.7)

where $(B_a)_t$, $a \in \llbracket D \rrbracket$, are independent complex Hermitian matrix Brownian motions (i.e., $\sqrt{2} \operatorname{Re}(B_a)_{ij}$ and $\sqrt{2} \operatorname{Im}(B_a)_{ij}$, i < j, and $(B_a)_{ii}$ are independent standard Brownian motions and $(B_a)_{ji} = (\overline{B}_a)_{ij}$). In particular, for each $t \geq t_0$, $(H_a)_t$ has the same law as

$$e^{-(t-t_0)/2} \cdot H_a^{(0)} + \sqrt{1 - e^{-(t-t_0)}} \cdot H_a^{(g)},$$
(4.8)

where $H_a^{(g)}$, $a \in \llbracket D \rrbracket$, are i.i.d. GUE. Then, we define the Green's function flow $G_t = (H_t + \Lambda_t - z_t)^{-1}$. Finally, with $(z_i)_t$, $i \in \{1, 2, 3\}$, Λ_t , H_t , and M_t , we can define

 $\widehat{M}_{(1,2),t} = \widehat{M}((z_1)_t, (z_2)_t, \Lambda_t), \quad L_{(1,2),t} = L((z_1)_t, (z_2)_t, H_t, \Lambda_t), \quad K_{(1,2),t} = K((z_1)_t, (z_2)_t, \Lambda_t)$

as in Definition 3.1, and define

$$L_{(1,2,3),t} = L((z_1)_t, (z_2)_t, (z_3)_t, H_t, \Lambda_t), \quad K_{(1,2,3),t} = K((z_1)_t, (z_2)_t, (z_3)_t, \Lambda_t)$$

as in Definition 4.1.

We now collect some basic properties of the characteristic flows in (4.5).

Lemma 4.5. Under Definition 4.4, the following properties hold.

• Denote $m_t := \langle M_t \rangle$. Suppose Im $z_t = o(1)$. Then, for any $t \ge t_0$, we have that

$$t_c - t = \frac{\mathrm{Im}\, z_t}{\mathrm{Im}\, m_t} (1 + \mathrm{o}(1)). \tag{4.9}$$

• M_t satisfies the following equation for $t \ge t_0$:

$$\frac{\mathrm{d}}{\mathrm{d}t}M(z_t,\Lambda_t) = \frac{1}{2}M(z_t,\Lambda_t). \tag{4.10}$$

• Conjugate flow: We have $\overline{Z}_t = Z(\overline{z}_t, \Lambda_t)$ and $\overline{M}_t = M(\overline{z}_t, \Lambda_t)$. Moreover, they satisfy the following equations under the conjugate flows $(\overline{z}_t, \Lambda_t)$ for $t \ge t_0$:

$$\frac{\mathrm{d}}{\mathrm{d}t}Z(\bar{z}_t,\Lambda_t) = -\frac{1}{2}Z(\bar{z}_t,\Lambda_t) - \langle M(\bar{z}_t,\Lambda_t)\rangle, \quad \frac{\mathrm{d}}{\mathrm{d}t}M(\bar{z}_t,\Lambda_t) = \frac{1}{2}M(\bar{z}_t,\Lambda_t).$$
(4.11)

• For any $(z_i)_t \in \{z_t, \bar{z}_t\}, i \in \{1, 2, 3\}, \widehat{M}_{(1,2),t} \text{ and } K_{(1,2),t} \text{ satisfy the equations}$

$$\frac{\mathrm{d}}{\mathrm{d}t}\widehat{M}_{(1,2),t} = \widehat{M}_{(1,2),t}, \quad \frac{\mathrm{d}}{\mathrm{d}t}K_{(1,2),t} = \left(K_{(1,2),t}\right)^2 + K_{(1,2),t}, \tag{4.12}$$

and $K_{(1,2,3),t}$ satisfies that for any $a_1, a_2, a_3 \in \llbracket D \rrbracket$,

$$\frac{\mathrm{d}}{\mathrm{d}t}(K_{(1,2,3),t})_{a_1a_2a_3} = \frac{3}{2}K_{(1,2,3),t} + \sum_{a=1}^{D} \left[(K_{(1,2),t})_{a_1a}(K_{(1,2,3),t})_{aa_2a_3} + (K_{(2,3),t})_{a_2a}(K_{(1,2,3),t})_{a_1aa_3} + (K_{(3,1),t})_{a_3a}(K_{(1,2,3),t})_{a_1a_2a_3} \right].$$
(4.13)

Proof. The estimate (4.9) follows by tracking Im m_t solved from the first equation in (4.5) and using the simple facts that $z_s - z_t = o(1)$ and $m_s(z_s) - m_t(z_t) = o(1)$ for all $s \in [t, t_c]$ as long as $|t_c - t| = o(1)$. Now, we prove the identity (4.10). Using $M_t = -(Z_t + m_t)^{-1}$ and equation (4.6), we obtain that

$$\dot{M}_t = M_t (\dot{Z}_t + \dot{m}_t) M_t = M_t (-Z_t/2 - m_t + \dot{m}_t) M_t, \qquad (4.14)$$

where we have used Newton's notation for derivatives with respect to time. Taking trace and using $Z_t M_t = -I - m_t M_t$, we obtain

$$\dot{m}_t = \frac{1}{2}m_t + \frac{1}{2}m_t \langle M_t^2 \rangle - m_t \langle M_t^2 \rangle + \dot{m}_t \langle M_t^2 \rangle \Rightarrow \dot{m}_t = \frac{1}{2}m_t.$$

Plugging it back into (4.14), we conclude (4.10). Taking conjugate transposes of (4.6) and (4.10), we get (4.11). The equations (4.12) and (4.13) follow from direct calculations with (4.6) and (4.10).

For the proof of Lemma 3.3, we need the following several key lemmas, Lemmas 4.6–4.9. These lemmas rely on the previously defined flow and their detailed proofs will be provided in Section 5.

Lemma 4.6. Suppose the assumptions of Theorem 2.2 hold for H_{t_0} and Λ_{t_0} . Under Definition 4.4, take any $z_{t_0} = E + i\eta \in \mathbf{D}(\kappa, \tau)$ such that $t_c - t_0 \sim N^{-\varepsilon_g}$. Let $(z_1)_t, (z_2)_t \in \{z_t, \bar{z}_t\}$ for $t \in [t_0, t_c]$. Then, for any fixed constant C > 4 and $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\|L_{(1,2),t} - K_{(1,2),t}\|_2 \prec \frac{(t_c - t_0)^2}{(t_c - t)^2} \|L_{(1,2),t_0} - K_{(1,2),t_0}\|_2 + \frac{1}{N(t_c - t)^2}.$$
(4.15)

Together with (4.1), it implies that for any $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\left\|L_{(1,2),t} - K_{(1,2),t}\right\|_{2} \prec \frac{N^{\varepsilon_{g}}}{N(t_{c}-t)^{2}}.$$
(4.16)

Lemma 4.7. Under the assumptions of Lemma 4.6, let $(z_1)_t, (z_2)_t, (z_3)_t \in \{z_t, \overline{z}_t\}$ for $t \in [t_0, t_c]$. Then, we have that for any $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\left\|L_{(1,2,3),t} - K_{(1,2,3),t}\right\|_{2} \prec \frac{(t_{c} - t_{0})^{3}}{(t_{c} - t)^{3}} \left\|L_{(1,2,3),t_{0}} - K_{(1,2,3),t_{0}}\right\|_{2} + \frac{N^{\varepsilon_{g}}}{N(t_{c} - t)^{3}}.$$
(4.17)

Together with (4.2), it implies that for any $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\left\|L_{(1,2,3),t} - K_{(1,2,3),t}\right\|_{2} \prec \frac{N^{\varepsilon_{g}}}{N(t_{c}-t)^{3}}.$$
(4.18)

Lemma 4.8. Under the assumptions of Lemma 4.6, we have that for any $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\max_{a \in \llbracket D \rrbracket} \mathbb{E} \langle (G_t - M_t) E_a \rangle \prec \frac{t_c - t_0}{t_c - t} \max_{a \in \llbracket D \rrbracket} |\mathbb{E} \langle (G_{t_0} - M_{t_0}) E_a \rangle| + \frac{N^{\varepsilon_g}}{N^2 (t_c - t)^2}.$$
(4.19)

Together with (4.3), it implies that for any $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\mathbb{E}\langle (G_t - M_t) E_a \rangle \prec \frac{N^{-\varepsilon_g}}{N(t_c - t)}.$$
(4.20)

Armed with the above three lemmas, we can prove the following key result for the proof of Lemma 3.3.

Lemma 4.9. Under the assumptions of Lemma 4.6, we have that for any $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\left\|\mathbb{E}L_{(1,2),t} - K_{(1,2),t}\right\|_{2} \prec \frac{(t_{c} - t_{0})}{(t_{c} - t)^{2}} \left\|\mathbb{E}L_{(1,2),t_{0}} - K_{(1,2),t_{0}}\right\|_{2} + \frac{N^{-1}g}{N(t_{c} - t)^{2}} + \frac{N^{-1}g}{N^{2}(t_{c} - t)^{3}}.$$
(4.21)

Together with (4.4), it implies that for any $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$,

$$\left\|\mathbb{E}L_{(1,2),t} - K_{(1,2),t}\right\|_{2} \prec \frac{N^{-(\varepsilon_{g} \wedge \delta_{A})}}{N(t_{c} - t)^{2}}.$$
(4.22)

Proof of Lemma 3.3. Suppose we want to prove (3.10) for $z = E + i\eta$ and Λ satisfying the assumptions of Lemma 3.3. In particular, we have $|E| \leq 2 - \kappa$ and $N^{-1+C\varepsilon_g} \leq \eta \leq N^{-C\varepsilon_g}$ for some constant C > 4 such that Lemma 4.9 holds. Now, take $t_f = 0$ and let $t_0 = t_f - N^{-\varepsilon_g}/2$. We can find initial values z_{t_0} and Λ_{t_0} such that $z_{t_f} = z$ and $\Lambda_{t_f} = \Lambda$ at $t = t_f$. (In fact, we can first solve the second equation in (4.5) as $\Lambda_t = e^{(t_f - t)/2}\Lambda$ and then plug it into the first equation in (4.5). In the resulting equation, the RHS is a locally Lipschitz function in t and z, so there exists a solution z_{t_0} at $t = t_0$.) Since $m_t(z) = m_{sc}(z) + o(1)$ under $||\Lambda_t|| = o(1)$ (see (A.1) in the appendix), we have $\operatorname{Im} m_t(z_t) \sim 1$. Thus, by (4.9), we know that $t_c - t_f \sim \eta$, which also gives $t_c - t_0 = (t_f - t_0)(1 + o(1)) = N^{-\varepsilon_g}(1/2 + o(1))$.

Under the assumption (3.9) for H, we can find another Gaussian divisible matrix H_{t_0} such that H_{t_f} has the same law as H under the evolution (4.7). With the above choices of H_{t_0} , z_{t_0} and Λ_{t_0} , we have

$$\mathbb{E}L_{(1,2)} = \mathbb{E}L_{(1,2),t_f}, \quad K_{(1,2)} = K_{(1,2),t_f}.$$

Finally, using (4.22) at $t = t_f$, we conclude the desired result (3.10).

4.1. **Proof of Lemma 4.2.** Our proof relies on the following formula derived from the definitions of G and M in (2.12),

$$G - M = -G(H + m)M = -M(H + m)G,$$
(4.23)

and the following complex cumulant expansion formula. We adopt the form stated in [71, Lemma 7.1].

Lemma 4.10. (Complex cumulant expansion) Let h be a complex random variable with all its moments exist. The (p,q)-cumulant of h is defined as

$$\mathcal{C}^{(p,q)}(h) := (-\mathbf{i})^{p+q} \cdot \left(\frac{\partial^{p+q}}{\partial s^p \partial t^q} \log \mathbb{E} e^{\mathbf{i}sh + \mathbf{i}t\bar{h}} \right) \Big|_{s=t=0}.$$

Let $f: \mathbb{C}^2 \to \mathbb{C}$ be a smooth function, and we denote its holomorphic derivatives by

$$f^{(p,q)}(z_1, z_2) := \frac{\partial^{p+q}}{\partial z_1^p \partial z_2^q} f(z_1, z_2)$$

Then, for any fixed $l \in \mathbb{N}$, we have

$$\mathbb{E}f(h,\bar{h})\bar{h} = \sum_{p+q=0}^{l} \frac{1}{p!\,q!} \mathcal{C}^{(p,q+1)}(h) \mathbb{E}f^{(p,q)}(h,\bar{h}) + R_{l+1}\,, \qquad (4.24)$$

given all integrals in (4.24) exist. Here, R_{l+1} is the remainder term depending on f and h, and for any $\tau > 0$, we have the estimate

$$R_{l+1} = \mathcal{O}(1) \cdot \mathbb{E} |h^{l+2} \mathbf{1}_{\{|h| > N^{\tau-1/2}\}} | \cdot \max_{p+q=l+1} ||f^{(p,q)}(z,\bar{z})||_{\infty} + \mathcal{O}(1) \cdot \mathbb{E} |h|^{l+2} \cdot \max_{p+q=l+1} ||f^{(p,q)}(z,\bar{z}) \cdot \mathbf{1}_{\{|z| \le N^{\tau-1/2}\}} ||_{\infty}.$$

With assumptions (2.1), (2.2), and (2.3), we can show that for $i, j \in \mathcal{I}$,

$$\mathcal{C}^{(0,1)}(H_{ij}) = \mathcal{C}^{(1,0)}(H_{ij}) = 0, \quad \mathcal{C}^{(1,1)}(H_{ij}) = s_{ij}, \quad \mathcal{C}^{(0,2)}(H_{ij}) = \mathcal{C}^{(2,0)}(H_{ij}) = s_{ij}\delta_{ij},$$

and that for any fixed $p,q \in \mathbb{N}$ with $p+q \geq 3$, there exists a constant C > 0 such that

$$\max_{i,j\in\mathcal{I}} |\mathcal{C}^{(p,q)}(H_{ij})| \le (CN)^{-(p+q)/2}.$$
(4.25)

We also adopt the following notation from [42, equation (42)].

Definition 4.11. Suppose that f and g are matrix-valued functions. Define

$$\underline{g(H)Hf(H)} := g(H)Hf(H) - \widetilde{\mathbb{E}}g(H)\widetilde{H}(\partial_{\widetilde{H}}f)(H) - \widetilde{\mathbb{E}}(\partial_{\widetilde{H}}g)(H)\widetilde{H}f(H),$$
(4.26)

where \widetilde{H} is an indepdent copy of H, $\widetilde{\mathbb{E}}$ denotes the partial expectation with respect to \widetilde{H} , and $(\partial_{\widetilde{H}}f)(H)$ denotes the directional derivative of the function f in the direction \widetilde{H} at the point H, i.e.,

$$[(\partial_{\widetilde{H}}f)(H)]_{xy} = (\widetilde{H} \cdot \nabla f(H))_{xy} := \sum_{\alpha,\beta \in \mathcal{I}} \widetilde{H}_{\alpha\beta} \frac{\partial f(H)_{xy}}{\partial H_{\alpha\beta}}.$$
(4.27)

The terms subtracted from g(H)Hf(H) are precisely the second-order term in the cumulant expansion. In particular, if all entries of H are Gaussian, we have $\mathbb{E} \underline{g(H)Hf(H)} = 0$. Moreover, if we take g(H) = I and f(H) = G, we have that

$$\underline{HG} = HG + \widetilde{\mathbb{E}}[\widetilde{H}G\widetilde{H}]G, \quad \text{with} \quad \widetilde{\mathbb{E}}[\widetilde{H}G\widetilde{H}] = \sum_{a=1}^{D} D\langle GE_a \rangle E_a.$$
(4.28)

First, we prove the estimate (4.1). Here, we restrict our consideration to the case where $z_1 = z$ and $z_2 = \overline{z}$; the case where $z_1 = z_2 \in \{z, \overline{z}\}$ is simpler and can be treated in a similar manner. For simplicity, we drop the subscripts and abbreviate $\widehat{M} \equiv \widehat{M}_{(1,2)}$, $L \equiv L_{(1,2)}$, and $K = K_{(1,2)}$. Using (4.23) and (4.28), we write that

$$G - M = -M(m+H)G = -M\underline{H}G + M(\widetilde{\mathbb{E}}[\widetilde{H}G\widetilde{H}] - m)G.$$
(4.29)

Applying it to G_2 and multiplying by $G_1 E_a$ from the left gives

$$G_1 E_a G_2 = G_1 E_a M_2 - G_1 E_a M_2 \underline{HG_2} + G_1 E_a M_2 (\tilde{\mathbb{E}}[\tilde{H}G_2\tilde{H}] - m_2) G_2.$$

On the other hand, by definition,

$$\underline{G_1 E_a M_2 H G_2} = G_1 E_a M_2 \underline{H G_2} + G_1 \widetilde{\mathbb{E}} [\widetilde{H} G_1 E_a M_2 \widetilde{H}] G_2, \qquad (4.30)$$

so we have that

$$G_1 E_a G_2 = G_1 E_a M_2 - \underline{G_1 E_a M_2 H G_2} + G_1 \widetilde{\mathbb{E}} [\widetilde{H} G_1 E_a M_2 \widetilde{H}] G_2 + G_1 E_a M_2 (\widetilde{\mathbb{E}} [\widetilde{H} G_2 \widetilde{H}] - m_2) G_2$$

Now, multiplying E_b from the right and taking the trace, we get that

$$L_{ab} = D\langle G_1 E_a M_2 E_b \rangle - D\langle \underline{G_1 E_a M_2 H G_2 E_b} \rangle$$

+ $D \sum_{x=1}^{D} \langle G_1 E_a M_2 E_x \rangle L_{xb} + D^2 \sum_{x=1}^{D} \langle (G_2 - M_2) E_x \rangle \langle G_1 E_a M_2 E_x G_2 E_b \rangle.$ (4.31)

Then, using the averaged local law (2.21) and Lemma 2.11, we can estimate the RHS as

$$L_{ab} = \widehat{M}_{ab} - D\langle \underline{G_1 E_a M_2 H G_2 E_b} \rangle + (\widehat{M}L)_{ab} + \mathcal{O}_{\prec}(\eta^{-1}(N\eta)^{-1}), \quad a, b \in \llbracket D \rrbracket.$$

With the bound on $\|(1-\widehat{M})^{-1}\|$ in (A.7) below, we get from the above equation that

$$L_{ab} = K_{ab} - D \sum_{x=1}^{D} \left(1 - \widehat{M} \right)_{ax}^{-1} \langle \underline{G_1 E_x M_2 H G_2 E_b} \rangle + \mathcal{O}_{\prec} (\eta^{-2} (N\eta)^{-1}).$$
(4.32)

To conclude (4.1), we only need to prove that given any $a, b \in [D]$, there is

$$Z := \langle \underline{G_1 E_a M_2 H G_2 E_b} \rangle \prec (N\eta^2)^{-1}.$$

By Markov's inequality, it suffices to prove the following high-moment bound: for any fixed $p \in 2\mathbb{N}$ with $p \ge 4$,

$$\mathbb{E}|Z|^p \prec (N\eta^2)^{-p}.\tag{4.33}$$

To prove (4.33), we use the cumulant expansion formula in Lemma 4.10:

$$\mathbb{E}|Z|^{p} = \mathbb{E}\langle \underline{G_{1}E_{a}M_{2}HG_{2}E_{b}}\rangle\overline{Z}|Z|^{p-2} = \frac{1}{DN^{2}}\sum_{x=1}^{D}\sum_{\alpha,\beta\in\mathcal{I}_{x}}\mathbb{E}(G_{2}E_{b}G_{1}E_{a}M_{2})_{\beta\alpha}\partial_{\beta\alpha}(\overline{Z}|Z|^{p-2})$$

$$+\sum_{m+n=2}^{l}\frac{1}{DN}\sum_{x=1}^{D}\sum_{\alpha,\beta\in\mathcal{I}_{x}}\frac{1}{m!n!}\mathcal{C}^{(n,m+1)}_{\beta\alpha}\mathbb{E}\partial^{m}_{\alpha\beta}\partial^{n}_{\beta\alpha}((G_{2}E_{b}G_{1}E_{a}M_{2})_{\beta\alpha}\overline{Z}|Z|^{p-2}) + \mathcal{R}_{l+1} \qquad (4.34)$$

$$=:X_{1} + \sum_{m+n=2}^{l}X_{m,n} + \mathcal{R}_{l+1},$$

where $\partial_{\beta\alpha} \equiv \partial_{h_{\beta\alpha}}$, $\mathcal{C}_{\beta\alpha}^{(n,m)} = \mathcal{O}(N^{-(n+m)/2})$ denotes the (n,m)-cumulant of $h_{\beta\alpha}$, and \mathcal{R}_{l+1} denotes the error term involving the (l+2)-th order moment of $h_{\beta\alpha}$. By definitions, we can write Z as

$$Z = \langle G_1 E_a M_2 H G_2 E_b \rangle + D \sum_{x=1}^{D} \langle G_2 E_x \rangle \langle G_1 E_a M_2 E_x G_2 E_b \rangle + D \sum_{x=1}^{D} \langle G_1 E_x G_2 E_b \rangle \langle G_1 E_a M_2 E_x \rangle.$$
(4.35)

With this formula and $HG_2 = I - (\Lambda - z)G_2$, we can check with direct calculations that

$$X_1 = N^{-2} \mathbb{E} \left(\mathcal{X}_{11} |Z|^2 + \mathcal{X}_{12} (\overline{Z})^2 \right) |Z|^{p-4},$$
(4.36)

where \mathcal{X}_{11} and \mathcal{X}_{12} are linear combinations of O(1) many terms taking the following forms with coefficients of order O(1):

- $\langle \prod_{i=1}^{k} G_{j_i} B_i \rangle$ for some $k \leq 5, j_i \in \{1, 2\}$, and deterministic matrices B_i with $||B_i|| = O(1)$;
- $\langle \prod_{i=1}^{k} G_{j_i} B_i \rangle \langle \prod_{i=1}^{k'} G_{j_{i'}} B_{i'} \rangle$ for some $k, k' \ge 1$ with $k + k' \le 6, j_i, j_{i'} \in \{1, 2\}$, and deterministic matrices $B_i, B_{i'}$ with $||B_i|| = O(1), ||B_{i'}|| = O(1)$.

Then, using Lemma 2.11, we can bound (4.36) by

$$X_1 \prec (N\eta^2)^{-2} \mathbb{E}|Z|^{p-2}.$$
 (4.37)

Next, we consider the terms $X_{m,n}$ with $2 \le m + n \le l$. Using (4.35), the anisotropic local law (2.20), and Lemma 2.11, we can check that for any fixed $m, n \in \mathbb{N}$ with $m + n \ge 1$,

$$\partial_{\alpha\beta}^{m}\partial_{\beta\alpha}^{n} \left(G_{2}E_{b}G_{1}E_{a}M_{2}\right)_{\beta\alpha} \prec \frac{1}{\eta}, \qquad \partial_{\alpha\beta}^{m}\partial_{\beta\alpha}^{n}Z \prec \frac{1}{N\eta^{2}}.$$
(4.38)

With these estimates, we obtain that

$$\partial_{\alpha\beta}^{m}\partial_{\beta\alpha}^{n}\left(\left(G_{2}E_{b}G_{1}E_{a}M_{2}\right)_{\beta\alpha}\overline{Z}|Z|^{p-2}\right)\prec\sum_{q=0}^{p-1}\frac{1}{\eta}\left(\frac{1}{N\eta^{2}}\right)^{q}|Z|^{p-1-q},$$

which gives that for any $3 \le m + n \le l$,

$$X_{m,n} \prec \sum_{q=1}^{p} N^{2-(n+m+1)/2} \left(\frac{1}{N\eta^2}\right)^q \mathbb{E}|Z|^{p-q} \le \sum_{q=1}^{p} \left(\frac{1}{N\eta^2}\right)^q \mathbb{E}|Z|^{p-q}.$$
(4.39)

It remains to consider the m + n = 2 case:

(i) If no derivative or both derivatives act on $(G_2 E_b G_1 E_a M_2)_{\beta\alpha}$, using the anisotropic local law (2.20) and Lemma 2.11, we can check directly that

$$\sum_{\alpha,\beta\in\mathcal{I}_x} \left|\partial_{\alpha\beta}^m \partial_{\beta\alpha}^n (G_2 E_b G_1 E_a M_2)_{\beta\alpha}\right|^2 \prec \eta^{-3} N.$$

Together with the Cauchy-Schwarz inequality, it implies that

$$\sum_{\alpha,\beta\in\mathcal{I}_x} \left|\partial^m_{\alpha\beta}\partial^n_{\beta\alpha} \left(G_2 E_b G_1 E_a M_2\right)_{\beta\alpha}\right| \prec \eta^{-3/2} N^{3/2}.$$
(4.40)

(ii) If one derivative acts on $(G_2 E_b G_1 E_a M_2)_{\beta\alpha}$ and the other on Z or \overline{Z} , then we can check that

$$\left|\partial_{\alpha\beta} \left(G_2 E_b G_1 E_a M_2\right)_{\beta\alpha}\right| + \left|\partial_{\beta\alpha} \left(G_2 E_b G_1 E_a M_2\right)_{\beta\alpha}\right| \prec \eta^{-1}, \quad \sum_{\alpha,\beta} \left|\partial_{\alpha\beta} Z\right|^2 \prec N^{-1} \eta^{-5}.$$
(4.41)

To see the second estimate, we express $\partial_{\alpha\beta}Z$ as

$$\partial_{\alpha\beta}Z = \frac{1}{DN} \left(\mathcal{Z}_{\beta\alpha}^{(1)} + \mathcal{Z}_{\beta\alpha}^{(2)} \right),$$

where $\mathcal{Z}_{\beta\alpha}^{(1)}$ comes from the derivative of the first term in (4.35):

$$\mathcal{Z}_{\beta\alpha}^{(1)} := -(G_1 E_a M_2 E_b G_1)_{\beta\alpha} + (G_1 E_a M_2 (\Lambda - z) G_2 E_b G_1)_{\beta\alpha} + (G_2 E_b G_1 E_a M_2 (\Lambda - z) G_2)_{\beta\alpha},$$
²⁰

and $\mathcal{Z}_{\beta\alpha}^{(2)}$ comes from the derivative of the remaining two terms. Moreover, $\mathcal{Z}_{\beta\alpha}^{(2)}$ is a sum of 6 terms with coefficients of order O(1), taking the forms:

$$D\sum_{x=1}^{D} (G_{j_1}B_1G_{j_2}B_2G_{j_1})_{\beta\alpha} \langle G_{j_3}B_3 \rangle, \quad \text{or} \quad D\sum_{x=1}^{D} \langle G_{j_1}B_1G_{j_2}B_2 \rangle (G_{j_3}B_3G_{j_3})_{\beta\alpha},$$

where $j_i \in \{1, 2\}$ and B_i are deterministic matrices such that $||B_i|| = O(1)$ for $i \in \{1, 2, 3\}$. By using Lemma 2.11, we can establish the bounds:

$$\sum_{\alpha,\beta} |\mathcal{Z}_{\beta\alpha}^{(1)}|^2 \prec N\eta^{-5}, \quad \sum_{\alpha,\beta} |(G_{j_1}B_1G_{j_2}B_2G_{j_1})_{\beta\alpha}\langle G_{j_3}B_3\rangle|^2 \prec N\eta^{-5},$$
$$\sum_{\alpha,\beta} |\langle G_{j_1}B_1G_{j_2}B_2\rangle (G_{j_3}B_3G_{j_3})_{\beta\alpha}|^2 \prec \eta^{-2} \sum_{\alpha,\beta} |(G_{j_3}B_3G_{j_3})_{\beta\alpha}|^2 \prec N\eta^{-5}.$$

These conclude the second estimate in (4.41). Now, combining (4.41) with the Cauchy-Schwarz inequality, we obtain that

$$\sum_{\alpha,\beta\in\mathcal{I}_x} \left(\left| \partial_{\alpha\beta} \left(G_2 E_b G_1 E_a M_2 \right)_{\beta\alpha} \right| + \left| \partial_{\beta\alpha} \left(G_2 E_b G_1 E_a M_2 \right)_{\beta\alpha} \right| \right) \left(\left| \partial_{\alpha\beta} Z \right| + \left| \partial_{\beta\alpha} Z \right| \right) \prec \frac{N^{1/2}}{\eta^{7/2}}.$$

$$(4.42)$$

Therefore, combining (4.38), (4.40), and (4.42), we obtain that for m + n = 2,

$$X_{m,n} \prec \sum_{q=1}^{3} \left(\frac{1}{N\eta^2}\right)^q \mathbb{E}|Z|^{p-q}.$$
(4.43)

Finally, for the remainder term \mathcal{R}_{l+1} , with the argument in e.g., [71, Lemma 4.6], it is easy to prove the following estimate: as long as l is sufficiently large depending on τ and p, there is

$$\mathcal{R}_{l+1} \prec N^{-p}.\tag{4.44}$$

Since the argument is standard, we omit the details here.

(

Now, plugging (4.37), (4.39), (4.43), and (4.44) into (4.34) yields that for any small constant $\varepsilon > 0$,

$$\mathbb{E} |Z|^{p} \prec \sum_{q=1}^{p} \left(N^{-1} \eta^{-2} \right)^{q} \mathbb{E} |Z|^{p-q} \le \sum_{q=1}^{p} \left(N^{-1} \eta^{-2} \right)^{q} \left(\mathbb{E} |Z|^{p} \right)^{\frac{p-q}{p}} \prec \left(N^{-1+\varepsilon} \eta^{-2} \right)^{p} + N^{-\varepsilon} \mathbb{E} |Z|^{p},$$

where we have applied Hölder's and Young's inequalities in the second and third steps. This implies (4.33)since ε is arbitrary, which further completes the proof of (4.1).

Remark 4.12. In the above proof, the matrix E_b can be replaced by any deterministic B with ||B|| = O(1). Thus, the above arguments lead to the following estimate that is similar to (4.1):

$$\langle G_1 E_a G_2 B \rangle - \sum_{b \in \llbracket D \rrbracket} \left(1 - \widehat{M}_{(1,2)} \right)_{ab}^{-1} \langle M_1 E_b M_2 B \rangle \prec N^{-1} \eta^{-3}.$$

$$\tag{4.45}$$

Furthermore, in the case where $z_1 = z_2 \in \{z, \overline{z}\}$, since $||(1 - \widehat{M})^{-1}|| \lesssim 1$ due to (A.8), the error term in (4.32) becomes $O_{\prec}(N^{-1}\eta^{-2})$. Thus, we get a better bound

$$||L_{(1,2)} - K_{(1,2)}||_2 \prec N^{-1}\eta^{-2}$$
 when $z_1 = z_2$. (4.46)

Next, we prove the estimate (4.2). Note that two of z_1, z_2, z_3 must take the same value. We assume without loss of generality that $z_1 = z_3 \in \{z, \bar{z}\}$. Then, using (4.23) and Definition 4.11, we write that

$$\begin{split} G_{1}E_{a_{1}}G_{2}E_{a_{2}}G_{3}E_{a_{3}} &= G_{1}E_{a_{1}}G_{2}E_{a_{2}}M_{3}E_{a_{3}} - \underline{G_{1}E_{a_{1}}G_{2}E_{a_{2}}M_{3}HG_{3}E_{a_{3}}} \\ &+ G_{1}E_{a_{1}}G_{2}E_{a_{2}}M_{3}\widetilde{\mathbb{E}}[\widetilde{H}(G_{3}-M_{3})\widetilde{H}]G_{3}E_{a_{3}} \\ &+ G_{1}\widetilde{\mathbb{E}}[\widetilde{H}G_{1}E_{a_{1}}G_{2}E_{a_{2}}M_{3}\widetilde{H}]G_{3}E_{a_{3}} + G_{1}E_{a_{1}}G_{2}\widetilde{\mathbb{E}}[\widetilde{H}G_{2}E_{a_{2}}M_{3}\widetilde{H}]G_{3}E_{a_{3}} \end{split}$$

Taking the trace and applying (2.21) and Lemma 2.11, we derive from this equation that

$$L_{(1,2,3)})_{a_1 a_2 a_3} = D\langle G_1 E_{a_1} G_2 E_{a_2} M_3 E_{a_3} \rangle - D\langle \underline{G_1 E_{a_1} G_2 E_{a_2} M_3 H G_3 E_{a_3}} \rangle \\ + \sum_{x=1}^{D} (\widehat{M}_{(2,3)})_{a_2 x} (L_{(1,2,3)})_{a_1 x a_3} + \sum_{x=1}^{D} D\langle G_1 E_{a_1} G_2 E_{a_2} M_3 E_x \rangle (K_{(3,1)})_{a_3 x} \\ {}_{21}$$

+
$$\sum_{x=1}^{D} D\langle G_1 E_{a_1} G_2 E_{a_2} M_3 E_x \rangle \left(L_{(3,1)} - K_{(3,1)} \right)_{a_3 x} + \mathcal{O}_{\prec} (\eta^{-2} (N\eta)^{-1}).$$
 (4.47)

Applying (4.46) to the last sum on the RHS and using Lemma 2.11, we get that

$$D\sum_{x=1}^{D} \langle G_1 E_{a_1} G_2 E_{a_2} M_3 E_x \rangle \left(L_{(3,1)} - (K_{(3,1)}) \right)_{xa_3} \prec N^{-1} \eta^{-3}.$$

Applying (4.45) to the first term on the RHS of (4.47), we get that

$$D\langle G_1 E_{a_1} G_2 E_{a_2} M_3 E_{a_3} \rangle = \sum_x (I - \widehat{M}_{(1,2)})_{a_1 x}^{-1} D\langle M_1 E_x M_2 E_{a_2} M E_{a_3} \rangle + \mathcal{O}_{\prec} (N^{-1} \eta^{-3}).$$
(4.48)

For simplicity, we will abbreviate the first term on the RHS of (4.48) as $\nu_{a_1a_2a_3}$. The equation (4.48) also applies to the fourth term on the RHS of (4.47) and gives that

$$D\sum_{x=1}^{D} \langle G_1 E_{a_1} G_2 E_{a_2} M_3 E_x \rangle (K_{(3,1)})_{a_3 x} = \sum_{x=1}^{D} \nu_{a_1 a_2 x} (K_{(3,1)})_{a_3 x} + \mathcal{O}_{\prec} (N^{-1} \eta^{-3}),$$

where we used that $\|K_{(3,1)}\| \lesssim 1$ by (A.8) below. Finally, we can prove that

$$\langle \underline{G_1 E_{a_1} G_2 E_{a_2} M_3 H G_3 E_{a_3}} \rangle \prec N^{-1} \eta^{-3}$$

using a similar argument as that for the proof of (4.33) with cumulant expansions. We omit the details here. In sum, we have derived from (4.47) that

$$(L_{(1,2,3)})_{a_1a_2a_3} = \nu_{a_1a_2a_3} + \sum_{x=1}^D \nu_{a_1a_2x}(K_{(3,1)})_{a_3x} + \sum_{x=1}^D (\widehat{M}_{(2,3)})_{a_2x}(L_{(1,2,3)})_{a_1xa_3} + \mathcal{O}_{\prec}(N^{-1}\eta^{-3}).$$

Now, solving for $L_{(1,2,3)}$, using (A.7) and that $I + K_{(3,1)} = (I - \widehat{M}_{(3,1)})^{-1}$, the estimate (4.2) follows.

4.2. **Proof of Lemma 4.3.** For any $z_1, z_2 \in \{z, \overline{z}\}$, we abbreviate that

$$\widehat{M} \equiv \widehat{M}_{(1,2)}, \quad L \equiv L_{(1,2)}, \quad K \equiv K_{(1,2)}, \quad \text{and} \quad \widetilde{M} \equiv \widehat{M}_{(2,1)}, \quad \widetilde{L} \equiv L_{(2,1)}, \quad \widetilde{K} \equiv K_{(2,1)}.$$

Moreover, given any deterministic matrix $B \in \mathbb{C}^{DN \times DN}$, we denote

$$L_{ab}(B) := D\langle G_1 E_a G_2 E_b B \rangle, \quad K_{ab}(B) := \sum_x (1 - \widehat{M})_{ax}^{-1} D\langle M_1 E_x M_2 E_b B \rangle.$$

Similarly, we define $\tilde{L}_{ab}(B)$ and $\tilde{K}_{ab}(B)$ by exchanging 1 and 2. Now, taking the expectation of (4.31), we obtain that

$$\mathbb{E}L_{ab} = \widehat{M}_{ab} + D\mathbb{E}\langle (G_1 - M_1)E_aM_2E_b\rangle - D\mathbb{E}\langle \underline{G_1E_aM_2HG_2E_b}\rangle + \sum_{x=1}^{D}\widehat{M}_{ax}\mathbb{E}L_{xb}$$

$$+ D\sum_{x=1}^{D}\mathbb{E}\langle (G_1 - M_1)E_aM_2E_x\rangle L_{xb} + D^2\sum_{x=1}^{D}\mathbb{E}\langle (G_2 - M_2)E_x\rangle\langle G_1E_aM_2E_xG_2E_b\rangle$$

$$= \widehat{M}_{ab} + D\mathbb{E}\langle (G_1 - M_1)E_aM_2E_b\rangle - D\mathbb{E}\langle \underline{G_1E_aM_2HG_2E_b}\rangle + \sum_{x=1}^{D}\widehat{M}_{ax}\mathbb{E}L_{xb}$$

$$+ D\sum_{x=1}^{D}\mathbb{E}\langle (G_1 - M_1)E_aM_2E_x\rangle K_{xb} + D^2\sum_{x=1}^{D}\mathbb{E}\langle (G_2 - M_2)E_x\rangle \widetilde{K}_{ba}(M_2E_x) + O_{\prec}(N^{-2}\eta^{-4}),$$

$$(4.49)$$

where we used the averaged local law (2.21) and the estimates (4.1) and (4.45) in the second step. The proof of Lemma 4.3 is based on the next two lemmas, Lemmas 4.13 and 4.14, which give precise estimates of the expectations on the RHS of (4.49).

Lemma 4.13. Under the setting of Lemma 4.3, we have that

$$-D\mathbb{E}\langle \underline{G_1 E_a M_2 H G_2 E_b} \rangle = \mathcal{O}_{\prec}(\eta^{-2} N^{-3/2} + \eta^{-3} N^{-2})$$

$$+ \frac{D\kappa^{(2,2)}}{N} \sum_{x=1}^{D} \left[\langle \operatorname{diag}(M_2)^2 E_x \rangle \widetilde{K}_{ba}(M_2 \operatorname{diag}(M_2) E_x) + \langle M_1 \operatorname{diag}(M_2) E_x \rangle \widetilde{K}_{bx}(\operatorname{diag}(M_1 E_a M_2)) \right]$$

$$+ \frac{D\kappa^{(2,2)}}{N} \sum_{x=1}^{D} \left[\langle M_1 E_a M_2 \operatorname{diag}(M_1) E_x \rangle \widetilde{K}_{bx}(\operatorname{diag}(M_1)) + \langle M_1 E_a M_2 \operatorname{diag}(M_2) E_x \rangle \widetilde{K}_{bx}(\operatorname{diag}(M_2)) \right], \quad (4.50)$$

where $\kappa^{(2,2)}$ is the normalized (2,2)-cumulant of h_{12} defined as $\kappa^{(2,2)} := N^2 \mathcal{C}_{12}^{(2,2)}$, and diag(B) is the diagonal matrix consisting of the diagonal entries of the given matrix B.

Proof. We again use the cumulant expansion in Lemma 4.10:

$$-D\mathbb{E}\langle \underline{G_1 E_a M_2 H G_2 E_b} \rangle = \frac{-1}{N} \sum_{m+n=2}^{l} \sum_{\alpha,\beta} \frac{1}{m!} \frac{1}{n!} \mathcal{C}^{(n,m+1)}_{\beta\alpha} \mathbb{E} \partial^m_{\alpha\beta} \partial^n_{\beta\alpha} (G_2 E_b G_1 E_a M_2)_{\beta\alpha} + \mathcal{R}'_{l+1}$$

where we choose l large enough such that the remainder term satisfies $\mathcal{R}'_{l+1} \prec N^{-2}$. Using Lemma 2.11, it is straightforward to check that $\partial^m_{\alpha\beta}\partial^n_{\beta\alpha}(G_2E_bG_1E_aM_2)_{\beta\alpha} \prec \eta^{-1}$. We thus obtain the rough bound

$$Y_{m,n} := \frac{-1}{N} \sum_{\alpha,\beta} \frac{1}{m!} \frac{1}{n!} \mathcal{C}^{(n,m+1)}_{\beta\alpha} \mathbb{E} \partial^m_{\alpha\beta} \partial^n_{\beta\alpha} (G_2 E_b G_1 E_a M_2)_{\beta\alpha} \prec N^{-(m+n-1)/2} \eta^{-1},$$

which is good enough when $m + n \ge 4$. It remains to deal with the cases $m + n \in \{2, 3\}$.

For the m + n = 2 case, we take m = n = 1 as an example:

$$\begin{split} Y_{1,1} &= -\frac{1}{N} \sum_{\alpha,\beta} \mathcal{C}_{\beta\alpha}^{(1,2)} \mathbb{E}\left[(G_2)_{\beta\beta} (G_2)_{\beta\alpha} (G_2 E_b G_1 E_a M_2)_{\alpha\alpha} + (G_2)_{\beta\beta} (G_2)_{\alpha\alpha} (G_2 E_b G_1 E_a M_2)_{\beta\alpha} \right] \\ &- \frac{1}{N} \sum_{\alpha,\beta} \mathcal{C}_{\beta\alpha}^{(1,2)} \mathbb{E}\left[(G_2)_{\beta\beta} (G_2 E_b G_1)_{\alpha\alpha} (G_1 E_a M_2)_{\beta\alpha} + (G_2)_{\beta\alpha} (G_2 E_b G_1)_{\beta\beta} (G_1 E_a M_2)_{\alpha\alpha} \right] \\ &- \frac{1}{N} \sum_{\alpha,\beta} \mathcal{C}_{\beta\alpha}^{(1,2)} \mathbb{E}\left[(G_1)_{\beta\beta} (G_2 E_b G_1)_{\beta\alpha} (G_1 E_a M_2)_{\alpha\alpha} + (G_1)_{\alpha\alpha} (G_2 E_b G_1)_{\beta\beta} (G_1 E_a M_2)_{\beta\alpha} \right]. \end{split}$$

For the first term on the RHS, we write it as

$$\frac{1}{N^{5/2}} \mathbb{E} \sum_{x=1}^{D} \sum_{\alpha,\beta \in \mathcal{I}_x} \left(\kappa^{(1,2)} + u \delta_{\alpha\beta} \right) (G_2)_{\beta\beta} (G_2)_{\beta\alpha} (G_2 E_b G_1 E_a M_2)_{\alpha\alpha}$$
(4.51)

where we used that $C_{\beta\alpha}^{(1,2)} = N^{-3/2}(\kappa^{(1,2)} + u\delta_{\alpha\beta})$ with $\kappa^{(1,2)} := N^{3/2}C_{12}^{(1,2)}$ and u being some fixed number. Then, using $(G_2)_{\beta\beta} \prec 1$ and $(G_2E_bG_1E_aM_2)_{\alpha\alpha} \prec \eta^{-1}$ by Lemma 2.11, applying the trivial bound $||G_2|| \leq \eta^{-1}$ gives that

$$(4.51) \prec \frac{1}{N^{5/2}} \cdot \frac{N}{\eta} + \frac{1}{N^{5/2}} \cdot \frac{1}{\eta} \frac{N}{\eta} \le 2\eta^{-2} N^{-3/2}.$$

The other terms in $Y_{1,1}$ and the terms in the $(m,n) \in \{(0,2), (2,0)\}$ cases can be handled in a similar way.

For the m + n = 3 case, through direct calculations, we see that when $(m, n) \in \{(0, 3), (2, 1), (3, 0)\}$, the corresponding $Y_{m,n}$ consists of terms of the form

$$J_{1} = \frac{1}{N^{3}} \sum_{x=1}^{D} \sum_{\alpha,\beta \in \mathcal{I}_{x}} (G_{s}B)_{ij} \mathcal{F}_{1} \quad \text{or} \quad J_{2} = \frac{1}{N^{3}} \sum_{x=1}^{D} \sum_{\alpha,\beta \in \mathcal{I}_{x}} (G_{2}E_{b}G_{1}B)_{ij} \mathcal{F}_{2}, \tag{4.52}$$

where $s \in \{1, 2\}, (i, j) \in \{(\alpha, \beta), (\beta, \alpha)\}, B$ is some deterministic matrix with ||B|| = O(1), and \mathcal{F}_1 and \mathcal{F}_2 are expressions satisfying $\mathcal{F}_1 = O_{\prec}(\eta^{-1})$ and $\mathcal{F}_2 = O_{\prec}(1)$. Using Cauchy-Schwarz inequality and Lemma 2.11, we get that

$$\sum_{\beta \in \mathcal{I}_x} |(G_s B)_{ij}| \prec \eta^{-1/2} N^{1/2}, \quad \sum_{\beta \in \mathcal{I}_x} |(G_2 E_b G_1 B)_{ij}| \prec \eta^{-3/2} N^{1/2},$$

with which we can bound J_1 and J_2 by $J_1 + J_2 \prec (N\eta)^{-3/2}$.

Finally, we consider the remaining case with m = 1 and n = 2. In this case, some terms in $Y_{1,2}$ will be of the form (4.52), so they can be bounded by $O_{\prec}((N\eta)^{-3/2})$. However, $Y_{1,2}$ also contains some leading terms that contribute to (4.50). We can expand $Y_{1,2}$ as

$$\begin{split} Y_{1,2} &= -\frac{\kappa^{(2,2)}}{2N^3} \sum_{x=1}^D \sum_{\alpha,\beta \in \mathcal{I}_x} \mathbb{E} \partial_{\alpha\beta} \partial_{\beta\alpha}^2 (G_2 E_b G_1 E_a M_2)_{\beta\alpha} \\ &= \frac{\kappa^{(2,2)}}{2N^3} \sum_{x=1}^D \sum_{\alpha,\beta \in \mathcal{I}_x} \mathbb{E} \partial_{\alpha\beta} \partial_{\beta\alpha} \left[(G_2)_{\beta\beta} (G_2 E_b G_1 E_a M_2)_{\alpha\alpha} + (G_2 E_b G_1)_{\beta\beta} (G_1 E_a M_2)_{\alpha\alpha} \right] \\ &= -\frac{\kappa^{(2,2)}}{N^3} \sum_{x=1}^D \sum_{\alpha,\beta \in \mathcal{I}_x} \mathbb{E} \partial_{\alpha\beta} \left[(G_2)_{\beta\beta} (G_2)_{\alpha\beta} (G_2 E_b G_1 E_a M_2)_{\alpha\alpha} + (G_2)_{\beta\beta} (G_2 E_b G_1)_{\alpha\beta} (G_1 E_a M_2)_{\alpha\alpha} \right] \\ &- \frac{\kappa^{(2,2)}}{N^3} \sum_{x=1}^D \sum_{\alpha,\beta \in \mathcal{I}_x} \mathbb{E} \partial_{\alpha\beta} \left[(G_1)_{\alpha\beta} (G_2 E_b G_1)_{\beta\beta} (G_1 E_a M_2)_{\alpha\alpha} \right]. \end{split}$$

When $\partial_{\alpha\beta}$ acts on the off-diagonal terms $(G_2)_{\alpha\beta}$, $(G_2E_bG_1)_{\alpha\beta}$, and $(G_1)_{\alpha\beta}$, it yields 4 leading terms consisting of diagonal terms only. The partial derivative $\partial_{\alpha\beta}$ acting on other terms gives errors, each containing two off-diagonal entries of the form $(G_sB)_{ij}$ and $(G_sE_{\gamma}G_{s'}B)_{ij}$, where $s, s' \in \{1, 2\}$, $(i, j) \in$ $\{(\alpha, \beta), (\beta, \alpha)\}, \gamma \in \{a, b\}$, and B is some deterministic matrix with $||B_i|| = O(1)$. By Lemma 2.11, we have $\sum_{\beta \in \mathcal{I}_x} |(G_sB)_{ij}|^2 \prec \eta^{-1}$ and $\sum_{\beta \in \mathcal{I}_x} |(G_sE_{\gamma}G_{s'}B)_{ij}|^2 \prec \eta^{-3}$. Combining these bounds with Cauchy-Schwarz inequality, we can bound the error terms by $O_{\prec}((N\eta)^{-2})$. In sum, we obtain that

$$Y_{1,2} = \frac{\kappa^{(2,2)}}{N^3} \sum_{x=1}^{D} \sum_{\alpha,\beta \in \mathcal{I}_x} \mathbb{E}(G_2)_{\beta\beta} \left[(G_2)_{\beta\beta} (G_2)_{\alpha\alpha} (G_2 E_b G_1 E_a M_2)_{\alpha\alpha} + (G_2)_{\alpha\alpha} (G_2 E_b G_1)_{\beta\beta} (G_1 E_a M_2)_{\alpha\alpha} \right] \\ + \frac{\kappa^{(2,2)}}{N^3} \sum_{x=1}^{D} \sum_{\alpha,\beta \in \mathcal{I}_x} \mathbb{E}(G_1)_{\beta\beta} \left[(G_2)_{\beta\beta} (G_2 E_b G_1)_{\alpha\alpha} (G_1 E_a M_2)_{\alpha\alpha} + (G_1)_{\alpha\alpha} (G_2 E_b G_1)_{\beta\beta} (G_1 E_a M_2)_{\alpha\alpha} \right] \\ + \mathcal{O}_{\prec} ((N\eta)^{-2}).$$

Then, using the anisotropic local law (2.20), Lemma 2.11, and the estimate (4.45), we get from the above expression that

$$Y_{1,2} = \frac{\kappa^{(2,2)}}{N^2} \sum_{x=1}^{D} \left\{ \sum_{\beta \in \mathcal{I}_x} \left[(M_2)_{\beta\beta} \right]^2 \widetilde{K}_{ba}(M_2 \operatorname{diag}(M_2) E_x) + \sum_{\alpha \in \mathcal{I}_x} (M_2)_{\alpha\alpha} (M_1 E_a M_2)_{\alpha\alpha} \widetilde{K}_{bx}(\operatorname{diag}(M_2)) \right\} \\ + \frac{\kappa^{(2,2)}}{N^2} \sum_{x=1}^{D} \left\{ \sum_{\beta \in \mathcal{I}_x} (M_1)_{\beta\beta} (M_2)_{\beta\beta} \widetilde{K}_{bx}(\operatorname{diag}(M_1 E_a M_2)) + \sum_{\alpha \in \mathcal{I}_x} (M_1)_{\alpha\alpha} (M_1 E_a M_2)_{\alpha\alpha} \widetilde{K}_{bx}(\operatorname{diag}(M_1)) \right\} \\ + \mathcal{O}_{\prec} ((N\eta)^{-3/2} + \eta^{-3} N^{-2}).$$

Rewriting this expression concludes (4.50).

Lemma 4.14. Under the setting of Lemma 4.3, let B be an arbitrary deterministic matrix with $||B|| \leq 1$. Then, we have that

$$\mathbb{E}\langle (G_1 - M_1)B \rangle = \frac{\kappa^{(2,2)} \langle \operatorname{diag}(M_1)^2 \rangle}{N} \left[\langle M_1 B M_1 \operatorname{diag}(M_1) \rangle + \frac{\langle M_1^2 \operatorname{diag}(M_1) \rangle}{1 - \langle M_1^2 \rangle} \langle M_1^2 B \rangle \right] + \mathcal{O}_{\prec} (\eta^{-1} N^{-3/2} + (N\eta)^{-2}).$$
(4.53)

Proof. Using (4.23) and (4.28), we get that

$$\mathbb{E}\langle (G_1 - M_1)B \rangle = -\mathbb{E}\langle M_1 \underline{H} \underline{G}_1 B \rangle + D \sum_{x=1}^{D} \mathbb{E}\langle (G_1 - M_1) E_x \rangle \langle M_1 E_x G_1 B \rangle$$

$$= -\mathbb{E}\langle M_1 \underline{H} \underline{G}_1 B \rangle + D \sum_{x=1}^{D} \mathbb{E}\langle (G_1 - M_1) E_x \rangle \langle M_1 E_x M_1 B \rangle + \mathcal{O}_{\prec}((N\eta)^{-2}),$$
(4.54)

where we used the averaged local law (2.21) in the second step. Taking $B = E_a$ and using (A.8) below, we can solve this equation to get that

$$\mathbb{E}\langle (G_1 - M_1)E_a \rangle = -\sum_{x=1}^{D} (1 - \widehat{M}_{(1,1)})_{ax}^{-1} \mathbb{E}\langle M_1 \underline{HG_1}E_x \rangle + \mathcal{O}_{\prec}((N\eta)^{-2}).$$
(4.55)

With the cumulant expansion in Lemma 4.10, using a similar argument as in the proof of Lemma 4.13, we can derive that

$$-\mathbb{E}\langle M_{1}\underline{HG_{1}}E_{x}\rangle = \frac{\kappa^{(2,2)}}{DN^{3}} \sum_{y=1}^{D} \sum_{\alpha,\beta\in\mathcal{I}_{y}} \mathbb{E}\left[(G_{1})_{\beta\beta}\right]^{2} (G_{1})_{\alpha\alpha} (G_{1}E_{x}M_{1})_{\alpha\alpha} + \mathcal{O}_{\prec}(\eta^{-1}N^{-3/2})$$
$$= \frac{\kappa^{(2,2)}}{N} \langle \operatorname{diag}(M_{1})^{2} \rangle \langle M_{1}E_{x}M_{1}\operatorname{diag}(M_{1}) \rangle + \mathcal{O}_{\prec}(\eta^{-1}N^{-3/2})$$
$$= \frac{\kappa^{(2,2)}}{DN} \langle \operatorname{diag}(M_{1})^{2} \rangle \langle M_{1}^{2}\operatorname{diag}(M_{1}) \rangle + \mathcal{O}_{\prec}(\eta^{-1}N^{-3/2}), \qquad (4.56)$$

where we used the averaged local law (2.21) and the block translation symmetry of M_1 in the above derivation. A similar argument gives that

$$-\mathbb{E}\langle M_1\underline{HG_1}B\rangle = \frac{\kappa^{(2,2)}}{N}\langle \operatorname{diag}(M_1)^2\rangle\langle M_1BM_1\operatorname{diag}(M_1)\rangle + \mathcal{O}_{\prec}(\eta^{-1}N^{-3/2}).$$
(4.57)

Now, plugging (4.56) into (4.55) and using

$$\sum_{x=1}^{D} (1 - \widehat{M}_{(1,1)})_{ax}^{-1} = \frac{1}{1 - \sum_{x=1}^{D} (M_{(1,1)})_{ax}} = \frac{1}{1 - \langle M_1^2 \rangle},$$

we obtain that

$$D\mathbb{E}\langle (G_1 - M_1)E_a \rangle = \frac{\kappa^{(2,2)}}{N} \frac{\langle \operatorname{diag}(M_1)^2 \rangle \langle M_1^2 \operatorname{diag}(M_1) \rangle}{1 - \langle M_1^2 \rangle} + \mathcal{O}_{\prec} (\eta^{-1} N^{-3/2} + (N\eta)^{-2}).$$
(4.58)

Finally, applying (4.57) and (4.58) to (4.54), we conclude (4.53).

Now, we are ready to complete the proof of Lemma 4.3 using Lemmas 4.13 and 4.14.

Proof of Lemma 4.3. Notice that $(1 - \langle M(z)^2 \rangle)^{-1} \leq 1$ by (A.2) and (A.3) below. Hence, the estimate (4.3) follows from Lemma 4.14 immediately for $\eta \sim N^{-\varepsilon_g}$. It remains to show (4.4).

First, when $z_1 = z_2 \in \{z, \overline{z}\}$, by (A.8) below, we have $\max_{a,b \in \llbracket D \rrbracket} \|K_{ab}(B)\| \leq 1$ for any matrix B with $\|B\| = O(1)$. Then, applying Lemma 4.13 and Lemma 4.14 to (4.49), we obtain that

$$\mathbb{E}L_{ab} = \widehat{M}_{ab} + \sum_{x=1}^{D} \widehat{M}_{ax} \mathbb{E}L_{xb} + \mathcal{O}_{\prec} (N^{-1} + \eta^{-2}N^{-3/2} + \eta^{-4}N^{-2}),$$

solving which gives (4.4).

Next, we consider the case $z_1 = \bar{z}_2 \in \{z, \bar{z}\}$. Without loss of generality, we assume that $z_1 = \bar{z}_2 = z$. Again, applying Lemma 4.13 and Lemma 4.14 to (4.49), we obtain that

$$\mathbb{E}L_{ab} = \widehat{M}_{ab} + \sum_{x=1}^{D} \widehat{M}_{ax} \mathbb{E}L_{xb} + \mathcal{O}_{\prec} \left(N^{-\delta_A} (N\eta)^{-1} + N^{-1} + \eta^{-2} N^{-3/2} + \eta^{-4} N^{-2} \right) \\ + \frac{\kappa^{(2,2)}}{N} \left[m_{sc}(z_2)^4 + |m_{sc}(z_1)|^4 + |m_{sc}(z_1)|^2 m_{sc}(z_1)^2 + |m_{sc}(z_1)|^2 m_{sc}(z_2)^2 \right] K_{ab} \\ + \frac{\kappa^{(2,2)}}{N} \left(\frac{m_{sc}(z_1)^4 |m_{sc}(z_1)|^2}{1 - m_{sc}(z_1)^2} + \frac{m_{sc}(z_2)^6}{1 - m_{sc}(z_2)^2} \right) K_{ab},$$
(4.59)

where we have used the estimate (A.2) below to replace M_1 and M_2 by scalar matrices $m_{sc}(z_1)I$ and $m_{sc}(z_2)I$ up to an error of order $O(||A||) = O(N^{-\delta_A})$. Now, using $|m_{sc}(z_1)| = 1 - O(\eta)$ and $m_{sc}(z_2) = \overline{m}_{sc}(z_1)$, we can check that

$$m_{sc}(z_{2})^{4} + |m_{sc}(z_{1})|^{4} + |m_{sc}(z_{1})|^{2}m_{sc}(z_{1})^{2} + |m_{sc}(z_{1})|^{2}m_{sc}(z_{2})^{2} + \frac{m_{sc}(z_{1})^{4}|m_{sc}(z_{1})|^{2}}{1 - m_{sc}(z_{1})^{2}} + \frac{m_{sc}(z_{2})^{6}}{1 - m_{sc}(z_{2})^{2}}$$

$$= m_{sc}(z_2)^4 + m_{sc}(z_1)^2 + m_{sc}(z_2)^2 + 1 + \frac{m_{sc}(z_1)^4}{1 - m_{sc}(z_1)^2} + \frac{m_{sc}(z_2)^6}{1 - m_{sc}(z_2)^2} + \mathcal{O}(\eta) = \mathcal{O}(\eta).$$
(4.60)

Plugging it back to (4.59) and using that $\eta \sim N^{-\varepsilon_g}$ with $\varepsilon_g \in (0, 1/4)$, we get

$$\mathbb{E}L_{ab} = \widehat{M}_{ab} + \sum_{x=1}^{D} \widehat{M}_{ax} \mathbb{E}L_{xb} + \mathcal{O}_{\prec} (N^{-\delta_A} (N\eta)^{-1} + N^{-1}).$$

Solving $\mathbb{E}L$ from this equation and using (A.7) below concludes (4.4).

Remark 4.15. We remark that the cancellation in (4.60) is not a mere coincidence. Instead, there is a robust mechanism behind it that does not depend on the specific properties of m_{sc} . This mechanism has been thoroughly investigated and elucidated in the context of random band matrices in [108], where it is referred to as the sum zero property or self-energy renormalization. To explain the core idea, we recall that by Lemmas 4.13 and 4.14, there is

$$\mathbb{E}L_{ab} = \widehat{M}_{ab} + \sum_{x=1}^{D} \widehat{M}_{ax} \mathbb{E}L_{xb} + \mathcal{D}_{ab} + \mathcal{O}_{\prec}(N^{-1}), \qquad (4.61)$$

for some deterministic matrix \mathcal{D} satisfying the "rough bound" $\|\mathcal{D}\| \prec (N\eta)^{-1}$. Now, summing both sides of this equation over $a \in [D]$, we get that

$$\left(1 - \frac{\operatorname{Im} m}{\operatorname{Im} m + \eta}\right) \frac{\mathbb{E}D\operatorname{Im}\langle GE_b\rangle}{\eta} = \frac{\operatorname{Im} m}{\operatorname{Im} m + \eta} + \sum_{a=1}^{D} \mathcal{D}_{ab} + \mathcal{O}_{\prec}(N^{-1}),$$

where we have used equation (A.4) below and applied Ward's identity (C.11) to $\sum_{x} L_{xb}$. By Lemma 4.14, this equation gives

$$\sum_{a=1}^{D} \mathcal{D}_{ab} = \frac{D \operatorname{Im} \mathbb{E} \langle (G-M) E_b \rangle}{\operatorname{Im} m + \eta} + \mathcal{O}_{\prec}(N^{-1}) = \mathcal{O}_{\prec}(N^{-1}).$$
(4.62)

This already indicates a non-trivial cancellation compared to the rough bound $(N\eta)^{-1}$. In particular, if we can write \mathcal{D}_{ab} as some coefficient c_N times K_{ab} (up to a small error) as shown in the proof of Lemma 4.3 above, then c_N must be of order $O_{\prec}(\eta/N)$. This gives another proof of (4.60).

5. Characteristic flow estimates

In this section, we complete the proofs of Lemmas 4.6–4.9 using the characteristic flow method. Let $\mathbf{B}_t = (b_{ij}(t))_{i,j\in\mathcal{I}}$ be a $D \times D$ block matrix Brownian motion consisting of the diagonal blocks $(B_a)_t$ in (4.7). Then, by (4.7), $H_t = (h_{ij}(t))_{i,j\in\mathcal{I}}$ satisfies the equation

$$\mathrm{d}h_{ij} = -\frac{1}{2}h_{ij}\mathrm{d}t + \frac{1}{\sqrt{N}}\mathrm{d}b_{ij}(t)$$

with initial data $H_{t_0} = H_0$. Let F be any function of t and H with continuous second-order derivatives. Then, by Itô's formula, we have that

$$dF = \partial_t F dt + \sum_{a=1}^D \sum_{k,l \in \mathcal{I}_a} \partial_{h_{kl}} F dh_{kl} + \frac{1}{2N} \sum_{a=1}^D \sum_{k,l \in \mathcal{I}_a} \partial_{h_{kl}} \partial_{h_{lk}} F dt.$$
(5.1)

We will apply this equation to functions of the resolvents $G_{i,t} \equiv (G_i)_t = (H_t - Z_{i,t})^{-1}$ with $Z_{i,t} = (z_i)_t - \Lambda_t$ for $z_i \in \{z, \bar{z}\}$. Using the formula (with the simplified notation $\partial_{kl} \equiv \partial_{h_{kl}}$)

$$\partial_{kl} (G_{i,t})_{k'l'} = - (G_{i,t})_{k'k} (G_{i,t})_{ll'}, \quad k', l' \in \mathcal{I}, \ k, l \in \mathcal{I}_a, \ a \in [\![D]\!],$$
(5.2)

we can easily obtain the following identities (with $M_{i,t} \equiv (M_i)_t$):

$$\partial_t G_{i,t} = G_{i,t} \left(\frac{\mathrm{d}}{\mathrm{d}t} Z_{i,t} \right) G_{i,t}, \quad \text{with} \quad \frac{\mathrm{d}}{\mathrm{d}t} Z_{i,t} = -\frac{1}{2} Z_{i,t} - \langle M_{i,t} \rangle; \tag{5.3}$$

$$\sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_a} h_{kl} \partial_{kl} G_{i,t} = -G_{i,t} H_t G_{i,t} = -G_{i,t} - G_{i,t} Z_{i,t} G_{i,t};$$
(5.4)

$$\sum_{k,l\in\mathcal{I}_a}\partial_{kl} \left(G_{i,t}\right)_{k_1l_1} \cdot \partial_{lk} \left(G_{i',t}\right)_{k_2l_2} = \left(G_{i,t}E_aG_{i',t}\right)_{k_1l_2} \left(G_{i',t}E_aG_{i,t}\right)_{k_2l_1}, \quad k_1, l_1, k_2, l_2\in\mathcal{I}.$$
(5.5)

5.1. **Proof of Lemma 4.6.** In this subsection, we give the proof of Lemma 4.6. For simplicity of notations, we abbreviate $\widehat{M}_{(1,2),t}$, $L_{(1,2),t}$, and $K_{(1,2),t}$ as \widehat{M}_t , L_t , and K_t . Moreover, we denote $z_t = E_t + i\eta_t$ and

$$\widetilde{L}_t \equiv \widetilde{L}_{(1,2),t} := (t_c - t)L_t, \quad \widetilde{K}_t \equiv \widetilde{K}_{(1,2),t} := (t_c - t)K_t.$$
(5.6)

Using Itô's formula (5.1) and the identities (5.2)–(5.5), we can calculate that for $x, y \in [D]$,

$$\begin{split} \mathbf{d}(\widetilde{L}_{t})_{xy} &= -(L_{t})_{xy} \mathbf{d}t + \frac{1}{\sqrt{N}} \sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_{a}} \partial_{kl} (\widetilde{L}_{t})_{xy} \mathbf{d}b_{kl} + D(t_{c} - t) \left\langle G_{1,t} E_{x} G_{2,t} E_{y} \right\rangle \mathbf{d}t \\ &+ D^{2}(t_{c} - t) \sum_{a=1}^{D} \left\langle G_{1,t} E_{x} G_{2,t} E_{a} \right\rangle \left\langle G_{2,t} E_{y} G_{1,t} E_{a} \right\rangle \mathbf{d}t \\ &+ D^{2}(t_{c} - t) \sum_{a=1}^{D} \left\langle (G_{1,t} - M_{1,t}) E_{a} \right\rangle \left\langle G_{1,t} E_{x} G_{2,t} E_{y} G_{1,t} E_{a} \right\rangle \mathbf{d}t \\ &+ D^{2}(t_{c} - t) \sum_{a=1}^{D} \left\langle (G_{2,t} - M_{2,t}) E_{a} \right\rangle \left\langle G_{2,t} E_{y} G_{1,t} E_{x} G_{2,t} E_{a} \right\rangle \mathbf{d}t. \end{split}$$

Using the definitions of \widetilde{L}_t and $L_{(1,2,3),t}$, we can rewrite the above equation as

$$d(\widetilde{L}_{t})_{xy} = \frac{1}{\sqrt{N}} \sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_{a}} \partial_{kl}(\widetilde{L}_{t})_{xy} db_{kl} + \left(1 - \frac{1}{t_{c} - t}\right) (\widetilde{L}_{t})_{xy} dt + \frac{1}{t_{c} - t} \sum_{a=1}^{D} (\widetilde{L}_{t})_{xa} (\widetilde{L}_{t})_{ay} dt + D(t_{c} - t) \sum_{a=1}^{D} \left\{ \left\langle (G_{1,t} - M_{1,t}) E_{a} \right\rangle [L_{(1,2,1),t}]_{xya} + \left\langle (G_{2,t} - M_{2,t}) E_{a} \right\rangle [L_{(2,1,2),t}]_{yxa} \right\} dt.$$
(5.7)

Next, with the averaged local law (2.21) and Lemma 2.11, we can bound the last term by $O_{\prec}((t_c - t) \cdot N^{-1}\eta_t^{-3}) = O_{\prec}(N^{-1}(t_c - t)^{-2})$, where we used $\eta_t \sim t_c - t$ by (4.9). Hence, we can rewrite (5.7) as

$$d\widetilde{L}_t = \frac{1}{\sqrt{N}} \sum_{a=1}^D \sum_{k,l \in \mathcal{I}_a} \partial_{kl} \widetilde{L}_t db_{kl} + \left[\left(1 - \frac{1}{t_c - t} \right) \widetilde{L}_t + \frac{1}{t_c - t} (\widetilde{L}_t)^2 \right] dt + \mathcal{O}_{\prec} \left(\frac{1}{N(t_c - t)^2} \right) dt.$$
(5.8)

On the other hand, by (4.12), we see that K_t satisfies the following equation:

$$\frac{\mathrm{d}}{\mathrm{d}t}\widetilde{K}_t = \left(1 - \frac{1}{t_c - t}\right)\widetilde{K}_t + \frac{1}{t_c - t}(\widetilde{K}_t)^2,\tag{5.9}$$

which matches the drift term in (5.8).

We now study the martingale term in (5.8), which is denoted as \mathcal{L}_t :

$$\mathrm{d}\mathcal{L}_t = \frac{1}{\sqrt{N}} \sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_a} \partial_{kl} \widetilde{L}_t \mathrm{d}b_{kl} \quad \text{with} \quad \mathcal{L}_{t_0} = 0.$$

The quadratic variation of $(\mathcal{L}_t)_{xy}, x, y \in \llbracket D \rrbracket$, is given by

$$[\mathcal{L}_{xy}]_t = \frac{1}{N} \int_{t_0}^t \sum_{a=1}^D \sum_{k,l \in \mathcal{I}_a} |\partial_{kl}(\widetilde{L}_s)_{xy}|^2 \mathrm{d}s.$$
(5.10)

Using (5.2), we can calculate the integrand as

$$\sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_{a}} |\partial_{kl}(\widetilde{L}_{s})_{xy}|^{2} = \frac{(t_{c}-s)^{2}}{N^{2}} \sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_{a}} \left(\left| (G_{1,s}E_{x}G_{2,s}E_{y}G_{1,s})_{lk} \right|^{2} + \left| (G_{2,s}E_{y}G_{1,s}E_{x}G_{2,s})_{lk} \right|^{2} + 2\operatorname{Re} \left[(G_{1,s}E_{x}G_{2,s}E_{y}G_{1,s})_{lk} \overline{(G_{2,s}E_{y}G_{1,s}E_{x}G_{2,s})_{lk}} \right] \right)^{2}$$

$$= \frac{D(t_c - s)^2}{N} \sum_{a=1}^{D} \left(\langle G_{1,s} E_x G_{2,s} E_y G_{1,s} E_a G_{1,s}^* E_y G_{2,s}^* E_x G_{1,s}^* E_a \rangle + \langle G_{2,s} E_y G_{1,s} E_x G_{2,s} E_a G_{2,s}^* E_x G_{1,s}^* E_y G_{2,s}^* E_a \rangle + 2 \operatorname{Re} \langle G_{1,s} E_x G_{2,s} E_y G_{1,s} E_a G_{2,s}^* E_x G_{1,s}^* E_y G_{2,s}^* E_a \rangle \right).$$

Applying Lemma 2.11 and (4.9), we obtain that if $t_0 \leq s \leq t_c - N^{-1+\tau}$ for a constant $\tau > 0$,

$$\sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_a} |\partial_{kl}(\widetilde{L}_s)_{xy}|^2 \prec \frac{|t_c - s|^2}{N} \cdot \frac{1}{\eta_s^5} \lesssim \frac{1}{N(t_c - s)^3}.$$
(5.11)

With a standard continuity argument, we obtain that this estimate holds uniformly in $s \in [t_0, t_c - N^{-1+\tau}]$ (i.e., we first show that (5.11) holds uniformly in t belonging to an N^{-C} -net of $[t_0, t_c - N^{-1+\tau}]$ and then extend it uniformly to the whole interval using the Lipschitz continuity in t). Plugging (5.11) into (5.10), we get the estimate

$$[\mathcal{L}_{xy}]_t \prec \frac{1}{N^2(t_c - t)^2}, \quad \text{if} \quad t_0 \le t \le t_c - N^{-1+\tau}.$$
 (5.12)

On the other hand, we have the trivial bound $|[\mathcal{L}_{xy}]_t| \leq N$ by using $||G_{i,s}|| \leq \eta_s^{-1} \ll N$ for $t \in [t_0, t_c - N^{-1+\tau}]$. Together with (5.12) and Definition 2.9, it implies that for any constant c > 0 and fixed $p \in \mathbb{N}$,

$$\mathbb{E} \left| \left[\mathcal{L}_{xy} \right]_t \right|^p \le \left(\frac{N^c}{N^2 (t_c - t)^2} \right)^p, \quad \text{if} \quad t_0 \le t \le t_c - N^{-1 + \tau}$$

Applying the Burkholder-Davis-Gundy inequality, we obtain a p-th moment bound on $\sup_{s \in [t_0,t]} |(\mathcal{L}_s)_{xy}|$. Then, applying Markov's inequality yields that for any $t \in [t_0, t_c - N^{-1+\tau}]$ and $x, y \in [D]$,

$$\sup_{\in [t_0,t]} |(\mathcal{L}_s)_{xy}| \prec \frac{1}{N(t_c-t)}.$$
(5.13)

Inserting (5.13) back to (5.8), we obtain that for any $t \in [t_0, t_c - N^{-1+\tau}]$ and $x, y \in [D]$,

s

$$\widetilde{L}_t - \widetilde{L}_{t_0} = \int_{t_0}^t \left[\left(1 - \frac{1}{t_c - s} \right) \widetilde{L}_s + \frac{1}{t_c - s} (\widetilde{L}_s)^2 \right] \mathrm{d}s + \mathcal{O}_{\prec} \left(\frac{1}{N(t_c - t)} \right).$$
(5.14)

On the other hand, by (5.9), we have

$$\widetilde{K}_t - \widetilde{K}_{t_0} = \int_{t_0}^t \left[\left(1 - \frac{1}{t_c - s} \right) \widetilde{K}_s + \frac{1}{t_c - s} (\widetilde{K}_s)^2 \right] \mathrm{d}s.$$
(5.15)

For simplicity, we introduce the notation $\widetilde{\Delta}_t := \widetilde{L}_t - \widetilde{K}_t$ and define the linear operator \mathcal{T}_t acting on $D \times D$ matrices as

$$\mathcal{T}_t(V) := \widetilde{K}_t V + V \widetilde{K}_t - [1 - (t_c - t)]V, \quad V \in \mathbb{C}^{D \times D}.$$
(5.16)

Then, subtracting (5.15) from (5.14), we obtain that

$$\begin{split} \widetilde{\Delta}_{t} - \widetilde{\Delta}_{t_{0}} &= \int_{t_{0}}^{t} \left[\left(1 - \frac{1}{t_{c} - s} \right) \widetilde{\Delta}_{s} + \frac{1}{t_{c} - s} \left(\widetilde{K}_{s} \widetilde{\Delta}_{s} + \widetilde{\Delta}_{s} \widetilde{K}_{s} + (\widetilde{\Delta}_{s})^{2} \right) \right] \mathrm{d}s + \mathcal{O}_{\prec} \left(\frac{1}{N(t_{c} - t)} \right) \\ &= \int_{t_{0}}^{t} \left(\left[(t_{c} - s) - 1 \right] \widetilde{\Delta}_{s} + \widetilde{K}_{s} \widetilde{\Delta}_{s} + \widetilde{\Delta}_{s} \widetilde{K}_{s} + (\widetilde{\Delta}_{s})^{2} \right) \frac{\mathrm{d}s}{t_{c} - s} + \mathcal{O}_{\prec} \left(\frac{1}{N(t_{c} - t)} \right) \\ &= \int_{t_{0}}^{t} \left(\mathcal{T}_{s} (\widetilde{\Delta}_{s}) + (\widetilde{\Delta}_{s})^{2} \right) \frac{\mathrm{d}s}{t_{c} - s} + \mathcal{E}_{t}, \end{split}$$
(5.17)

where \mathcal{E}_t is a $D \times D$ random matrix satisfying that $\|\mathcal{E}_t\|_{\mathrm{HS}} \prec [N(t_c - t)]^{-1}$ uniformly in $t \in [t_0, t_c - N^{-1+\tau}]$. Denoting $\widehat{\Delta}_t := \widetilde{\Delta}_t - \mathcal{E}_t$ and noticing that $\mathcal{E}_{t_0} = 0$, we can rewrite (5.17) as

$$\widehat{\Delta}_t - \widehat{\Delta}_{t_0} = \int_{t_0}^t \left(\mathcal{T}_s(\widehat{\Delta}_s) + \mathcal{T}_s(\mathcal{E}_s) + (\widehat{\Delta}_s + \mathcal{E}_s)^2 \right) \frac{\mathrm{d}s}{t_c - s}.$$
(5.18)

Let $\Phi(t; t_0)$ be the standard Peano-Baker series corresponding to the linear operator $\mathcal{T}_t/(t_c - t)$, i.e., it is the unique solution to the following linear integral equation

$$\Phi(t;t_0) = \mathbf{1} + \int_{t_0}^t \frac{\mathcal{T}_s}{t_c - s} \circ \Phi(s;t_0) \,\mathrm{d}s,$$
(5.19)

where 1 denotes the identity operator. Equivalently, $\Phi(t; t_0)$ satisfies the differential equation

$$\frac{\mathrm{d}}{\mathrm{d}t}\Phi\left(t;t_{0}\right)=\frac{\mathcal{T}_{t}}{t_{c}-t}\circ\Phi\left(t;t_{0}\right),\quad\text{with}\quad\Phi\left(t_{0};t_{0}\right)=\mathbf{1}.$$

By Duhamel's principle, the solution $\widehat{\Delta}_t$ to (5.18) can be expressed as

$$\widehat{\Delta}_{t} = \Phi\left(t; t_{0}\right) \widehat{\Delta}_{t_{0}} + \int_{t_{0}}^{t} \Phi\left(t; s\right) \left(\frac{\mathcal{T}_{s}(\mathcal{E}_{s}) + (\widehat{\Delta}_{s} + \mathcal{E}_{s})^{2}}{t_{c} - s}\right) \mathrm{d}s.$$
(5.20)

Suppose the space $\mathbb{C}^{D \times D}$ of $D \times D$ matrices is equipped with the Hilbert-Schmidt norm. Then, we claim that, as a linear operator on $\mathbb{C}^{D \times D}$, \mathcal{T}_t has operator norm at most 1 + o(1):

$$\|\mathcal{T}_t\|_{op} \le 1 + o(1).$$
 (5.21)

Before proving this estimate, we first use it to prove (4.15). With (5.21), we get from (5.19) that

$$\frac{\mathrm{d}}{\mathrm{d}t} \|\Phi(t;s)\|_{op} \le \frac{1+\mathrm{o}(1)}{t_c-t} \|\Phi(t;s)\|_{op}.$$

Using Grönwall's inequality, we conclude that for $t_0 \leq s \leq t \leq t_c - N^{-1+\tau}$,

$$\|\Phi(t;s)\|_{op} \prec \frac{t_c - s}{t_c - t}.$$
 (5.22)

Applying (5.21) and (5.22) to (5.20) and using the bound on $\|\mathcal{E}_t\|_{\text{HS}}$, we obtain that

$$\|\widehat{\Delta}_{t}\|_{2} \prec \frac{t_{c} - t_{0}}{t_{c} - t} \|\widehat{\Delta}_{t_{0}}\|_{2} + \frac{1}{t_{c} - t} \int_{t_{0}}^{t} \|\widehat{\Delta}_{s} + \mathcal{E}_{s}\|_{2}^{2} \mathrm{d}s + \int_{t_{0}}^{t} \frac{\mathrm{d}s}{N(t_{c} - t)(t_{c} - s)}.$$

From this estimate, writing $\widehat{\Delta}_t = \overline{\Delta}_t - \mathcal{E}_t$, we obtain that for $t_c - t_0 \sim N^{-\varepsilon_g}$ and $t_c - t \ge N^{-1+\tau}$,

$$\|\widetilde{\Delta}_{t}\|_{2} \prec \frac{t_{c} - t_{0}}{t_{c} - t} \|\widetilde{\Delta}_{t_{0}}\|_{2} + \frac{1}{t_{c} - t} \int_{t_{0}}^{t} \|\widetilde{\Delta}_{s}\|_{2}^{2} \mathrm{d}s + \frac{1}{N(t_{c} - t)}.$$
(5.23)

By (4.1), we have $\|\widetilde{\Delta}_{t_0}\|_2 \prec N^{-1+2\varepsilon_g}$. Then, from (5.23), we derive the following self-improving estimate for $t \in [t_0, t_c - N^{-1+C\varepsilon_g}]$ when C > 2:

$$\sup_{s \in [t_0,t]} N(t_c - s) \|\widetilde{\Delta}_s\|_2 \le N^{2\varepsilon_g} \implies N(t_c - t) \|\widetilde{\Delta}_t\|_2 \prec N^{\varepsilon_g} + N^{(4-C)\varepsilon_g}.$$
(5.24)

Moreover, defining the stopping time $T = \inf_{t > t_0} \{ N(t_c - t) \| \widetilde{\Delta}_t \|_2 \ge N^{2\varepsilon_g} \}$, we obtain from (5.23) that

$$\|\widetilde{\Delta}_t\|_2 \prec \frac{t_c - t_0}{t_c - t} \|\widetilde{\Delta}_{t_0}\|_2 + \frac{1}{N(t_c - t)}$$

if $t \leq T$ and $t_c - t \geq N^{-1+C\varepsilon_g}$ with C > 4. Now, applying a standard continuity argument with (5.24) gives that $T \ge t_c - N^{-1+C\varepsilon_g}$ with high probability when C > 4 and hence concludes the desired result (4.15).

Finally, we prove the bound (5.21). Notice that

$$\|\widetilde{K}_t\| = (t_c - t)\|K_t\| \le (t_c - t)\|(1 - \widehat{M}_t)^{-1}\|\|\widehat{M}_t\| \lesssim (t_c - t)\|(1 - \widehat{M}_t)^{-1}\|.$$
(5.25)

Hence, when $(z_1)_t = (z_2)_t \in \{z_t, \bar{z}_t\}$, we have $\|\tilde{K}_t\| \lesssim t_c - t$ by (A.8) below, with which we readily derive (5.21). It remains to consider the case $(z_1)_t = (\overline{z}_2)_t \in \{z_t, \overline{z}_t\}$. Since \widehat{M}_t is a circulant matrix, it has an eigendecomposition $\widehat{M}_t = U_t D_t U_t^*$, where D_t is the diagonal matrix of eigenvalues and U_t is a $D \times D$ unitary matrix. Then, \widetilde{K}_t can be written as

$$\widetilde{K}_t = U_t \Xi_t U_t^*, \quad \Xi_t := (t_c - t) \frac{D_t}{1 - D_t}$$

Now, we define the linear operator $\widetilde{\mathcal{T}}_t$ as

$$\widetilde{\mathcal{T}}_t(V) := \Xi_t V + V \Xi_t - [1 - (t_c - t)]V, \quad V \in \mathbb{C}^{D \times D}.$$

It is easy to see $\mathcal{T}_t(V) = U_t[\mathcal{T}_t(U_t^*VU_t)]U_t^*$, which implies that $\|\mathcal{T}_t\|_{op} = \|\mathcal{T}_t\|_{op}$. From the definition of \mathcal{T}_t , we see that

$$\|\widetilde{\mathcal{T}}_{t}\|_{op} \le \max_{k,l \in [D]} |(\Xi_{t})_{kk} + (\Xi_{t})_{ll} - 1| + |t_{c} - t|.$$
(5.26)

It remains to estimate the eigenvalues of \widetilde{K}_t .

Since the entries of \widehat{M}_t are all non-negative when $(z_1)_t = (\overline{z}_2)_t$, it has a Perron–Frobenius eigenvalue

$$d_1 = \frac{\operatorname{Im} m_t(z_t)}{\operatorname{Im} m_t(z_t) + \eta_t}$$

by equation (A.4) below. Moreover, by equation (A.5), the eigenvalues d_k of \widehat{M}_t satisfy $d_k = d_1 - a_k - ib_k$, $k \in \llbracket D \rrbracket$, for some $a_k \ge 0$ and $a_k + |b_k| = o(1)$ (specifically, $a_1 = b_1 = 0$). Thus,

$$\begin{aligned} (\Xi_t)_{kk} + (\Xi_t)_{ll} - 1 &= (t_c - t) \left[\frac{d_1 - a_k - \mathrm{i}b_k}{(1 - d_1) + a_k + \mathrm{i}b_k} + \frac{d_1 - a_l - \mathrm{i}b_l}{(1 - d_1) + a_l + \mathrm{i}b_l} \right] - 1 \\ &= \frac{\eta_t}{\eta_t + a'_k + \mathrm{i}b'_k} + \frac{\eta_t}{\eta_t + a'_l + \mathrm{i}b'_l} - 1 + \mathrm{o}(1), \end{aligned}$$

where we used (4.9) in the second step and abbreviated that $a'_k := (\operatorname{Im} m_t(z_t) + \eta_t)a_k$ and $b'_k := (\operatorname{Im} m_t(z_t) + \eta_t)a_k$ $\eta_t b_k$. Together with the simple fact $|1/(1+z) - 1/2| \le 1/2$ when $\operatorname{Re} z \ge 0$, this equation implies $|(\Xi_t)_{kk} + 1/2| \le 1/2$ $(\Xi_t)_{ll} - 1 \le 1 + o(1)$. Plugging it into (5.26) concludes (5.21).

5.2. **Proof of Lemma 4.7.** The proof of Lemma 4.7 is similar to that of Lemma 4.6 above. Hence, we only describe an outline of the proof without giving all the details. For simplicity, we abuse the notations a little bit and abbreviate $L_{(1,2,3),t}$ and $K_{(1,2,3),t}$ as L_t and K_t . Moreover, we denote $z_t = E_t + i\eta_t$ and

$$\widetilde{L}_t := (t_c - t)^{3/2} L_t, \quad \widetilde{K}_t := (t_c - t)^{3/2} K_t, \quad \widetilde{L}_{(1,2),t} = (t_c - t) L_{(1,2),t}, \quad \widetilde{K}_{(1,2),t} = (t_c - t) K_{(1,2),t}.$$

Similar to (5.8), using Itô's formula (5.1), the identities (5.2)–(5.5), and Lemmas 2.10 and 2.11, we obtain that for $x, y, w \in \llbracket D \rrbracket$,

$$d(\widetilde{L}_{t})_{xyw} = \frac{1}{\sqrt{N}} \sum_{a=1}^{D} \sum_{k,l \in \mathcal{I}_{a}} \partial_{kl} (\widetilde{L}_{t})_{xyw} db_{kl} + \frac{3}{2} \left(1 - \frac{1}{t_{c} - t} \right) (\widetilde{L}_{t})_{xyw} dt + O_{\prec} \left(\frac{1}{N \left(t_{c} - t \right)^{5/2}} \right) dt \\ + \frac{1}{t_{c} - t} \sum_{a=1}^{D} \left[(\widetilde{L}_{(1,2),t})_{xa} (\widetilde{L}_{t})_{ayw} + (\widetilde{L}_{(2,3),t})_{ya} (\widetilde{L}_{t})_{xaw} + (\widetilde{L}_{(3,1),t})_{wa} (\widetilde{L}_{t})_{xya} \right] dt.$$
(5.27)

We again denote the martingale term by

$$\mathrm{d}\mathcal{L}_t := \frac{1}{\sqrt{N}} \sum_{a=1}^D \sum_{k,l \in \mathcal{I}_a} \partial_{kl} (\widetilde{L}_t)_{xyz} \mathrm{d}b_{kl}.$$

Similar to (5.13), we can prove that uniformly in $t \in [t_0, t_c - N^{-1+\tau}]$,

$$\sup_{x,y,w \in [\![D]\!]} \sup_{s \in [t_0,t]} |(\mathcal{L}_s)_{xyw}| \prec \frac{1}{N(t_c - t)^{3/2}}.$$
(5.28)

Plugging it into (5.27), we obtain the following estimate for $x, y, w \in [D]$ and $t \in [t_0, t_c - N^{-1+\tau}]$:

$$\left(\tilde{L}_t - \tilde{L}_{t_0} \right)_{xyw} = \int_{t_0}^t \frac{3}{2} \left(1 - \frac{1}{t_c - s} \right) (\tilde{L}_s)_{xyw} ds + \mathcal{O}_{\prec} \left(\frac{1}{N(t_c - t)^{3/2}} \right)$$

$$+ \int_{t_0}^t \frac{1}{t_c - s} \sum_{a=1}^D \left[(\tilde{L}_{(1,2),s})_{xa} (\tilde{L}_s)_{ayw} + (\tilde{L}_{(2,3),s})_{ya} (\tilde{L}_s)_{xaw} + (\tilde{L}_{(3,1),s})_{wa} (\tilde{L}_s)_{xya} \right] ds.$$

$$(5.29)$$

Using (4.16) and applying Lemma 2.11 to \tilde{L}_s , we obtain from (5.29) that

$$\widetilde{L}_t - \widetilde{L}_{t_0} = \int_{t_0}^t \mathscr{T}_s(\widetilde{L}_s) \frac{\mathrm{d}s}{t_c - s} + \mathcal{O}_{\prec} \left(\frac{N^{\varepsilon_g}}{N(t_c - t)^{3/2}} \right),$$
(5.30)

where $\mathscr{T}_t : \mathbb{C}^{D \times D \times D} \to \mathbb{C}^{D \times D \times D}$ is a linear operator defined as follows: for $V \in \mathbb{C}^{D \times D \times D}$,

$$\mathscr{T}_{t}(V)_{xyw} = \sum_{a=1}^{D} \left[(\widetilde{K}_{(1,2),t})_{xa} V_{ayw} + (\widetilde{K}_{(2,3),t})_{ya} V_{xaw} + (\widetilde{K}_{(3,1),t})_{wa} V_{xya} \right] - \frac{3}{2} [1 - (t_{c} - t)] V_{xyw}.$$

On the other hand, by (4.13), we have

$$\widetilde{K}_t - \widetilde{K}_{t_0} = \int_{t_0}^t \mathscr{T}_s(\widetilde{K}_s) \frac{\mathrm{d}s}{t_c - s}.$$
(5.31)

Now, subtracting (5.31) from (5.30) and denoting $\Delta_t := L_t - K_t$, we obtain that

$$\widetilde{\Delta}_t - \widetilde{\Delta}_{t_0} = \int_{t_0}^t \mathscr{T}_s(\widetilde{\Delta}_s) \frac{\mathrm{d}s}{t_c - s} + \mathcal{E}_t, \qquad (5.32)$$

where \mathcal{E}_t is a $D \times D \times D$ random tensor satisfying that $\|\mathcal{E}_t\|_2 \prec N^{\varepsilon_g}/[N(t_c-t)^{3/2}]$, where $\|\cdot\|_2$ denotes the ℓ^2 -norm by regarding \mathcal{E}_t as a D^3 -dimensional vector. We now claim that

$$\|\mathscr{T}_t\|_{op} \le 3/2 + o(1). \tag{5.33}$$

This bound is trivial when $(z_1)_t = (z_2)_t = (z_3)_t$. Next, we assume that $(z_1)_t = (z_3)_t = (\bar{z}_2)_t \in \{z_t, \bar{z}_t\}$ without loss of generality. Then, $\|\widetilde{K}_{(3,1),t}\| = o(1)$ and $(\widetilde{K}_{(2,3),t})_{ya} = (\widetilde{K}_{(1,2),t})_{ay}$. Hence, we can write \mathscr{T}_t as

$$\mathscr{T}_t(V) = \mathscr{T}_t^{(1)}(V) + \sum_{a=1}^D (\widetilde{K}_{(3,1),t})_{wa} V_{xya} - \frac{1}{2} [1 - (t_c - t)] V_{xyw},$$

where the operator $\mathscr{T}_{t}^{(1)}$ is defined as

$$\mathscr{T}_t^{(1)}(V)_{xyw} := \sum_{a=1}^D \left[(\widetilde{K}_{(1,2),t})_{xa} V_{ayw} + V_{xaw} (\widetilde{K}_{(1,2),t})_{ay} \right] - [1 - (t_c - t)] V_{xyw}$$

Note that when w is fixed, $\mathscr{T}_t^{(1)}$ reduces to the operator (5.16). With (5.21), we can easily conclude (5.33).

Finally, we adopt a similar argument as that below (5.21). Let $\Phi(t; t_0)$ be the standard Peano-Baker series corresponding to the linear operator $\mathscr{T}_t/(t_c-t)$. From (5.33), we can derive that for $t_0 \leq s \leq t \leq t_c - N^{-1+\tau}$,

$$\|\Phi(t;s)\|_{op} \prec \frac{(t_c - s)^{3/2}}{(t_c - t)^{3/2}}.$$
(5.34)

Then, from (5.32) and (5.34), we obtain that

$$\|\widetilde{\Delta}_t\|_2 \prec \frac{(t_c - t_0)^{3/2}}{(t_c - t)^{3/2}} \|\widetilde{\Delta}_{t_0}\|_2 + \frac{N^{\varepsilon_g}}{N(t_c - t)^{3/2}},$$

which implies (4.17) and completes the proof of Lemma 4.7.

5.3. Proof of Lemma 4.8. To mimic our proofs of Lemmas 4.6 and 4.7 above, we abuse the notations again and denote $\widetilde{L}_t, \widetilde{K}_t \in \mathbb{C}^D$ by

$$(\widetilde{L}_t)_x := (t_c - t)^{1/2} \langle GE_x \rangle, \quad (\widetilde{K}_t)_x := (t_c - t)^{1/2} \langle ME_x \rangle, \quad x \in \llbracket D \rrbracket.$$

Again, using Itô's formula (5.1), the identities (5.2)–(5.5), we get that for $x \in [D]$ and $(z_1)_t \equiv z_t = E_t + i\eta_t$,

$$d\mathbb{E}(\widetilde{L}_t)_x = \frac{1}{2} \left(1 - \frac{1}{t_c - t} \right) \mathbb{E}(\widetilde{L}_t)_x dt + \frac{1}{t_c - t} \sum_{a=1}^D \mathbb{E}(\widetilde{L}_{(1,1),t})_{xa} (\widetilde{L}_t - \widetilde{K}_t)_a dt,$$
(5.35)

where recall that $\widetilde{L}_{(1,1),t}$ was defined in (5.6) with $(z_1)_t = z_t$. By (2.21), (4.16), and (4.9), we have that for $t \in [t_0, t_c - N^{-1 + C\varepsilon_g}],$

$$(\widetilde{L}_t - \widetilde{K}_t)_a \prec \frac{1}{N(t_c - t)^{1/2}}, \quad \widetilde{L}_{(1,1),t} - \widetilde{K}_{(1,1),t} \prec \frac{N^{\varepsilon_g}}{N(t_c - t)}$$

Again, these estimates hold uniformly in t due to the N^{-C} -net argument. Therefore, (5.35) now writes

$$\mathrm{d}\,\mathbb{E}\widetilde{L}_t = \frac{1}{2}\left(1 - \frac{1}{t_c - t}\right)\widetilde{K}_t\mathrm{d}t + \frac{1}{t_c - t}T_t\left(\mathbb{E}\widetilde{L}_t - \widetilde{K}_t\right)\mathrm{d}t + \mathcal{O}_{\prec}\left(\frac{N^{\varepsilon_g}}{N^2(t_c - t)^{5/2}}\right)\mathrm{d}t,$$
₃₁

where $T_t : \mathbb{C}^D \to \mathbb{C}^D$ is a linear operator defined as

$$T_t(V) = \widetilde{K}_{(1,1),t}V - \frac{1}{2} [1 - (t_c - t)]V, \quad V \in \mathbb{C}^D.$$

On the other hand, with (4.10), we can derive that

$$\mathrm{d}\widetilde{K}_t = \frac{1}{2} \left(1 - \frac{1}{t_c - t} \right) \widetilde{K}_t \mathrm{d}t$$

Subtracting the above two differential equations and denoting $\widetilde{\Delta}_t := \mathbb{E}\widetilde{L}_t - \widetilde{K}_t$, we get

$$\widetilde{\Delta}_t - \widetilde{\Delta}_{t_0} = \int_{t_0}^t T_s(\widetilde{\Delta}_s) \frac{\mathrm{d}s}{t_c - s} + \mathcal{E}_t, \qquad (5.36)$$

where \mathcal{E}_t is a *D*-dimensional random vector satisfying $\|\mathcal{E}_t\|_2 \prec N^{\varepsilon_g}/[N^2(t_c-t)^{3/2}]$. Since $\|\widetilde{K}_{(1,1),t}\| = o(1)$, we easily see that $\|T_t\|_{op} \leq 1/2 + o(1)$. Again, using the argument with Peano-Baker series, we obtain from (5.36) that

$$\|\widetilde{L}_t - \widetilde{K}_t\|_2 \prec \frac{(t_c - t_0)^{1/2}}{(t_c - t)^{1/2}} \|\widetilde{L}_{t_0} - \widetilde{K}_{t_0}\|_2 + \frac{N^{\varepsilon_g}}{N^2(t_c - t)^{3/2}},$$

which implies (4.19) and completes the proof of Lemma 4.8.

5.4. **Proof of Lemma 4.9.** Finally, in this subsection, we complete the proof of Lemma 4.9 with Lemmas 4.7 and 4.8 and the proof of Lemma 4.6 as main inputs. Taking the expectation of (5.7), we obtain that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}\widetilde{L}_t = \left(1 - \frac{1}{t_c - t}\right)\mathbb{E}\widetilde{L}_t + \frac{1}{t_c - t}\mathbb{E}(\widetilde{L}_t)^2 + \mathcal{O}_{\prec}\left(\frac{N^{-\varepsilon_g}}{N(t_c - t)^2} + \frac{N^{\varepsilon_g}}{N^2(t_c - t)^3}\right).$$
(5.37)

In the derivation of this equation, we have applied (4.18) and (2.21) to get that

$$\mathbb{E} \langle (G_{1,t} - M_{1,t}) E_a \rangle [L_{(1,2,1),t}]_{xya} = \mathbb{E} \langle (G_{1,t} - M_{1,t}) E_a \rangle [K_{(1,2,1),t}]_{xya} + \mathcal{O}_{\prec} \left(\frac{N^{\varepsilon_g}}{N^2 (t_c - t)^4} \right) \\ \prec \frac{N^{-\varepsilon_g}}{N (t_c - t)^3} + \frac{N^{\varepsilon_g}}{N^2 (t_c - t)^4},$$

where in the second step we have used (4.20) to control $\mathbb{E}\langle (G_{1,t} - M_{1,t}) E_a \rangle$ and the bound $|(K_{(1,2,1),t})_{xya}| \lesssim \eta_t^{-2} \lesssim (t_c - t)^{-2}$ due to the estimates (A.7) and (A.8) below. The other term $\mathbb{E} \langle (G_{2,t} - M_{2,t}) E_a \rangle [L_{(2,1,2),t}]_{yxa}$ can be bounded in a similar way.

Next, the estimate (4.16) implies that for any $x, y, a \in \llbracket D \rrbracket$

$$\mathbb{E}(\widetilde{L}_t)_{xa}(\widetilde{L}_t)_{ay} - \mathbb{E}(\widetilde{L}_t)_{xa}\mathbb{E}(\widetilde{L}_t)_{ay} \prec \frac{N^{2\varepsilon_g}}{N^2(t_c - t)^2}$$

Plugging this estimate into (5.37), we can improve the expectation of equation (5.17) to

$$\mathbb{E}\widetilde{\Delta}_t - \mathbb{E}\widetilde{\Delta}_{t_0} = \int_{t_0}^t \mathcal{T}_s(\mathbb{E}\widetilde{\Delta}_s) \frac{\mathrm{d}s}{t_c - s} + \mathcal{E}_t, \quad \mathcal{E}_t = \mathcal{O}_{\prec} \left(\frac{N^{-\varepsilon_g}}{N(t_c - t)} + \frac{N^{2\varepsilon_g}}{N^2(t_c - t)^2} \right).$$

Then, following the proof below (5.18), we can derive that

$$\|\mathbb{E}\widetilde{\Delta}_t\|_2 \prec \frac{t_c - t_0}{t_c - t} \|\mathbb{E}\widetilde{\Delta}_{t_0}\|_2 + \frac{N^{-\varepsilon_g}}{N(t_c - t)} + \frac{N^{2\varepsilon_g}}{N^2(t_c - t)^2},$$

which implies (4.21) and completes the proof of Lemma 4.9.

6. Chaotic regime: eigenvalues

Consider the matrix OU process $H_{\Lambda}(t) = H_t + \Lambda$, where $H_t = (h_{ij}(t))_{i,j \in \mathcal{I}}$ satisfies the OU equation

$$dh_{ij} = -\frac{1}{2}h_{ij}dt + \frac{1}{\sqrt{DN}}db_{ij}(t), \text{ with } H_0 = H,$$
 (6.1)

where $B_t = (b_{ij}(t))_{i,j \in \mathcal{I}}$ denotes a Hermitian matrix whose upper triangular entries are independent complex Brownian motions with variance t. Then, Theorem 2.3 follows immediately from Lemmas 6.1 and 6.2 below. **Lemma 6.1.** Under the assumptions of Theorem 2.3, suppose $\mathfrak{t} = N^{-1+\mathfrak{c}}$ for a constant $\mathfrak{c} \in (0, 1/10)$. Then, for any fixed $n \in \mathbb{N}$, there exist a constant $c_n = c_n(\mathfrak{c}, \delta_A) > 0$ such that

$$\left| \int_{\mathbb{R}^n} \mathrm{d}\boldsymbol{\alpha} \, O(\boldsymbol{\alpha}) \left[p_{H_{\Lambda}(\mathfrak{t})}^{(n)} \left(E + \frac{\boldsymbol{\alpha}}{DN} \right) - p_{GUE}^{(n)} \left(E + \frac{\boldsymbol{\alpha}}{DN} \right) \right] \right| \le N^{-c_n}, \tag{6.2}$$

where $p_{H_{\Lambda}(\mathfrak{t})}^{(n)}$ denotes the n-point correlation function of $H_{\Lambda}(\mathfrak{t})$ and we have abbreviated that $E + \frac{\alpha}{DN} =$ $\left(E+\frac{\alpha_1}{DN},\ldots,E+\frac{\alpha_n}{DN}\right).$

Proof. This lemma is a simple consequence of Theorem 2.2 in [81]. More precisely, H_t in (6.1) has law

$$H_t \stackrel{d}{=} e^{-t/2} \cdot H + \sqrt{1 - e^{-t}} \cdot W, \tag{6.3}$$

where $\stackrel{d}{=}$ means "equal in distribution" and W is a $DN \times DN$ GUE independent of H. Since the matrix $H + e^{t/2}\Lambda$ also satisfies Assumption 2.1, its resolvent satisfies the averaged local law (2.21) with Λ replaced by $e^{t/2}\Lambda$. Then, applying [81, Theorem 2.2] with $V = e^{-t/2}H + \Lambda$ gives that

$$\int_{\mathbb{R}^n} \mathrm{d}\boldsymbol{\alpha} \, O(\boldsymbol{\alpha}) \left[p_{H_{\Lambda}(\mathfrak{t})}^{(n)} \left(E + \frac{\boldsymbol{\alpha}}{DN\widehat{\rho}_{\mathrm{fc},\mathfrak{t}}(E)} \right) - p_{GUE}^{(n)} \left(E + \frac{\boldsymbol{\alpha}}{DN\rho_{sc}(E)} \right) \right] \le N^{-c} \tag{6.4}$$

for a constant c > 0, where $\hat{\rho}_{fc,t}$ denotes the density for the free convolution of the empirical measure of V and the semicircle law with variance $1 - e^{-t}$. More precisely, $\hat{\rho}_{fc,t}$ can be defined through its Stieltjes transform $\widehat{m}_{fc,t}(z)$, which is the unique solution to

$$\widehat{m}_{\mathrm{fc},\mathfrak{t}}(z) = \frac{1}{DN} \operatorname{Tr} \frac{1}{V - z - (1 - e^{-\mathfrak{t}}) \widehat{m}_{\mathrm{fc},\mathfrak{t}}(z)} \quad \text{with} \quad \operatorname{Im} \widehat{m}_{\mathrm{fc},\mathfrak{t}}(z) > 0 \text{ whenever } \operatorname{Im} z > 0.$$

Using the averaged local law for $H + e^{t/2}\Lambda$, we get that

$$\widehat{m}_{\rm fc,t}(z) = e^{t/2} m \left(e^{t/2} z + (e^{t/2} - e^{-t/2}) \widehat{m}_{\rm fc,t}(z), e^{t/2} \Lambda \right) + \mathcal{O}_{\prec} \left((N\mathfrak{t})^{-1} \right) \\ = m_{sc}(z) + \mathcal{O}_{\prec} \left((N\mathfrak{t})^{-1} + \|A\| + \mathfrak{t} \right)$$

for $z = E + i\eta$ with $E \in [-2 + \kappa, 2 - \kappa]$ and $0 \le \eta \le 1$, where in the second step, we used that m is well-approximated by m_{sc} due to the estimate (A.2) below. Taking the imaginary part of the above equation when $\eta = 0$, we obtain that $\hat{\rho}_{fc,t}(E) = \rho_{sc}(E) + O_{\prec}(N^{-c \wedge \delta_A})$. Plugging it into (6.4), applying the change of variables, and using the smoothness of the test function O, we conclude (6.2). \square

Lemma 6.2. Under the assumptions of Theorem 2.3, there exists a constant c > 0 depending on ε_A such that the following holds for $\mathfrak{t} = N^{-1+\mathfrak{c}}$. For any fixed $n \in \mathbb{N}$, there exists a constant $c_n = c_n(\mathfrak{c}, \varepsilon_A)$ such that

$$\int_{\mathbb{R}^n} \mathrm{d}\boldsymbol{\alpha} \, O(\boldsymbol{\alpha}) \left(p_{H_{\Lambda}(\mathfrak{t})}^{(n)} - p_{H_{\Lambda}}^{(n)} \right) \left(E + \frac{\boldsymbol{\alpha}}{N} \right) \leqslant N^{-c_n}.$$
(6.5)

The rest of this section is devoted to the proof Lemma 6.2. We will use the following correlation function comparison theorem stated in [64, Theorem 15.3], which is a slightly modified version of [66, Theorem 6.4].

Lemma 6.3 (Theorem 15.3 of [64]). Under the assumptions of Theorem 2.3, let G and G_t denote the resolvents of H_{Λ} and $H_{\Lambda}(\mathfrak{t})$, respectively. Suppose that for some small constants $\sigma, \delta > 0$, the following two conditions hold.

(i) Fix any $\varepsilon > 0$ and $k \in \mathbb{N}$, there is

$$\mathbb{E}[\mathrm{Im}\langle G(z)\rangle]^k + \mathbb{E}[\mathrm{Im}\langle G_{\mathfrak{t}}(z)\rangle]^k \lesssim 1, \quad for \quad z = E + \mathrm{i}N^{-1+\varepsilon} \quad with \quad |E| \le 2 - \kappa.$$
(6.6)

(ii) For any sequence $z_j = E_j + i\eta_j$, j = 1, ..., n, with $|E_j| \leq 2 - \kappa$ and $\eta_j = N^{-1-\sigma_j}$ for some constants $0 < \sigma_j \leq \sigma$, we have

$$\left| \mathbb{E} \prod_{i=1}^{n} \operatorname{Im} \langle G(z_i) \rangle - \mathbb{E} \prod_{i=1}^{n} \operatorname{Im} \langle G_{\mathfrak{t}}(z_i) \rangle \right| \leqslant N^{-\delta}.$$
(6.7)

Then, for any integer $n \ge 1$, there is a constant $c_n = c_n(\sigma, \delta) > 0$ such that for any $|E| \le 2 - 2\kappa$ and any C^1 -function $O: \mathbb{R}^n \to \mathbb{R}$ with compact support,

$$\int_{\mathbb{R}^n} \, \mathrm{d}\boldsymbol{\alpha} \, O(\boldsymbol{\alpha}) \left(p_{H_{\Lambda}(\mathfrak{t})}^{(n)} - p_{H_{\Lambda}}^{(n)} \right) \left(E + \frac{\boldsymbol{\alpha}}{N} \right) \leqslant N^{-c_n}.$$

Next, we bound the quantities in (6.6) and (6.7) using Lemma 2.10 and Theorem 2.2. Note that we have only proved Lemma 2.10 and Theorem 2.2 for H_{Λ} , but they can be extended to any $H_{\Lambda}(t)$ with $t \in [0, \mathfrak{t}]$. (Heuristically, adding a GUE component will "help" the QUE of eigenvectors, so there is no essential difficulty in making this extension.) Let $M_t(z)$ be the solution to the matrix Dyson equation (2.12) with the operator S replaced by S_t :

$$\mathcal{S}_t(M_t) := e^{-t} \mathcal{S}(M_t) + (1 - e^{-t}) \langle M_t \rangle.$$

However, note that the self-consistent equation (2.15) for $m_t(z) := \langle M_t(z) \rangle$ is unchanged, so we have $m_t(z) = m(z)$ and $M_t(z) = M(z)$ as given by (2.16).

Lemma 6.4. For any $t \in [0, t]$, under the assumptions of Lemma 2.10, the local laws (2.20)–(2.21) holds for G_t and the eigenvalue rigidity estimate (2.22) holds. Under the assumptions of Theorem 2.2, (2.6) holds for the bulk eigenvectors of $H_{\Lambda}(t)$.

Proof. The local laws and eigenvalue rigidity can be proved in the same way as Lemma 2.10 using the methods developed in e.g., [9, 58, 72]; more details will be explained in Appendix C.1 below. The proof of (2.6) is similar to that for Theorem 2.2, and we omit the details.

Proof of Lemma 6.2. The estimate (6.6) follows directly from the local law (2.21) for $G_t(z)$ established in Lemma 6.4. Now, to apply Lemma 6.3 to conclude Lemma 6.2, we only need to prove the Green's function comparison estimate (6.7). Using the local law of $G_t(z)$ and the fact that $\eta \operatorname{Im}(G_t)_{xx}(E+i\eta)$ is an increasing function of η for fixed $x \in \mathcal{I}$ and $E \in \mathbb{R}$, we obtain that for any constant $\varepsilon > 0$,

$$\max_{x \in \mathcal{I}} \sup_{|E| \le 2-\kappa} \sup_{N^{-1-\sigma} \le \eta \le N^{-1+\varepsilon}} \operatorname{Im}(G_t)_{xx}(E+\mathrm{i}\eta) \le \frac{N^{-1+\varepsilon}}{N^{-1-\sigma}} \max_{x \in \mathcal{I}} \sup_{|E| \le 2-\kappa} \operatorname{Im}(G_t)_{xx}(E+\mathrm{i}N^{-1+\varepsilon}) \prec N^{\sigma+\varepsilon}.$$

Then, using [64, Lemma 15.5], we can derive that

$$\max_{x,y\in\mathcal{I}}\sup_{|E|\leq 2-\kappa}\sup_{\eta\geq N^{-1-\sigma}}|(G_t)_{xy}(E+\mathrm{i}\eta)|\prec N^{\sigma+\varepsilon}.$$
(6.8)

Since $\text{Im}\langle G_t(z_i)\rangle = (\langle G_t(z_i)\rangle - \langle G_t(\bar{z}_i)\rangle)/(2i\eta_i)$, to show (6.7), it suffices to prove that for $z_j = E_j \pm i\eta_j$, $j = 1, \ldots, n$, we have

$$\left| \mathbb{E} \prod_{i=1}^{n} \langle G(z_i) \rangle - \mathbb{E} \prod_{i=1}^{n} \langle G_{\mathfrak{t}}(z_i) \rangle \right| \leqslant N^{-\delta}.$$
(6.9)

To prove (6.9), we apply the Itô's formula and get that

$$\partial_t \mathbb{E} \prod_{i=1}^n \langle G_t(z_i) \rangle = \frac{1}{2DN} \mathbb{E} \sum_{x,y \in \mathcal{I}} \partial_{xy} \partial_{yx} \left(\prod_{i=1}^n \langle G_t(z_i) \rangle \right) - \frac{1}{2} \mathbb{E} \sum_{x,y \in \mathcal{I}} h_{xy}(t) \partial_{xy} \left(\prod_{i=1}^n \langle G_t(z_i) \rangle \right),$$

where ∂_{xy} denotes the partial derivative $\partial/\partial_{h_{xy}(t)}$. Then, applying the cumulant expansion in Lemma 4.10 to the second term on the RHS, we get that

$$\partial_t \mathbb{E} \prod_{i=1}^n \langle G_t(z_i) \rangle = \frac{e^{-t}}{2} \mathbb{E} \sum_{x,y \in \mathcal{I}} \left(\frac{1}{DN} - s_{xy} \right) \partial_{xy} \partial_{yx} \left(\prod_{i=1}^n \langle G_t(z_i) \rangle \right) + \sum_{k=3}^l \mathcal{F}_k + \mathcal{E}_{l+1}, \tag{6.10}$$

where we used that $E|h_{xy}(t)|^2 = e^{-t}s_{xy} + (1 - e^{-t})(DN)^{-1}$ by (6.3) (recall that s_{xy} was defined in (2.13)), \mathcal{F}_k is the sum of terms involving the cumulants $\mathcal{C}^{(m,n)}(h_{xy}(t))$ with m + n = k, and \mathcal{E}_{l+1} is the remainder term. We can choose l sufficiently large such that $\mathcal{E}_{l+1} \leq 1$. For the terms \mathcal{F}_k , using the estimates in (6.8) and the fact that ε is arbitrarily small, it is easy to check that

$$\mathcal{F}_k \prec N^{-k/2+2+(n+k)\sigma}, \quad 3 \le k \le l. \tag{6.11}$$

It remains to bound the first term on the RHS of (6.10). To simplify notations, for a fixed $t \in [0, \mathfrak{t}]$, we abbreviate $G_i \equiv G_t(z_i)$. Then, we can write the first term on the RHS of (6.10) as e^{-t} times

$$\mathcal{F}_{2} := \frac{1}{DN^{2}} \mathbb{E} \sum_{x,y \in \mathcal{I}} \left(D^{-1} - Ns_{xy} \right) \left\{ \sum_{i \in \llbracket n \rrbracket} (G_{i}^{2})_{xx} (G_{i})_{yy} \prod_{k \neq i} \langle G_{k} \rangle + \frac{1}{DN} \sum_{i < j \in \llbracket n \rrbracket} (G_{i}^{2})_{xy} (G_{j}^{2})_{yx} \prod_{k \notin \{i,j\}} \langle G_{k} \rangle \right\}$$
$$= D\mathbb{E} \sum_{a \in \llbracket D \rrbracket} \langle G_{i}^{2} (D^{-1} - E_{a}) \rangle \langle G_{i} E_{a} \rangle \prod_{k \neq i} \langle G_{k} \rangle + \frac{1}{DN^{2}} \mathbb{E} \sum_{i < j \in \llbracket n \rrbracket} \langle G_{i}^{2} (D^{-1} - E_{a}) G_{j}^{2} E_{a} \rangle \prod_{k \notin \{i,j\}} \langle G_{k} \rangle.$$

It remains to bound the two terms

$$X(i;a) := \langle G_i^2(D^{-1} - E_a) \rangle \langle G_i E_a \rangle \prod_{k \neq i} \langle G_k \rangle,$$
$$Y(i,j;a) := N^{-2} \langle G_i^2(D^{-1} - E_a) G_j^2 E_a \rangle \prod_{k \notin \{i,j\}} \langle G_k \rangle.$$

With the bound (6.8), we can derive that

$$\langle G_k \rangle \prec N^{\sigma}, \quad \langle G_i E_a \rangle \prec N^{\sigma}, \quad \langle G_i^2 (D^{-1} - E_a) \rangle \prec N^{1+2\sigma}, \quad \langle G_i^2 (D^{-1} - E_a) G_j^2 E_a \rangle \prec N^{3+4\sigma}. \tag{6.12}$$

With these estimates, we get the following rough bounds on X and Y:

$$X(i;a) \prec N^{1+(n+2)\sigma}, \quad Y(i,j;a) \prec N^{1+(n+2)\sigma}.$$
 (6.13)

To improve these estimates, we consider the eigendecompositions

$$\langle G_i^2 \left(D^{-1} - E_a \right) \rangle = \frac{1}{DN} \sum_k \frac{\mathbf{v}_k^* (D^{-1} - E_a) \mathbf{v}_k}{(\lambda_k - z_i)^2},$$
 (6.14)

$$\left\langle G_{i}^{2} \left(D^{-1} - E_{a} \right) G_{j}^{2} E_{a} \right\rangle = \frac{1}{DN} \sum_{k,l} \frac{\mathbf{v}_{k}^{*} (D^{-1} - E_{a}) \mathbf{v}_{l} \cdot \mathbf{v}_{l}^{*} E_{a} \mathbf{v}_{k}}{(\lambda_{k} - z_{i})^{2} (\lambda_{l} - z_{j})^{2}}, \tag{6.15}$$

where $\lambda_k \equiv \lambda_k(t)$ and $\mathbf{v}_k \equiv \mathbf{v}_k(t)$ denote the eigenvalues and eigenvectors of $H_t + \Lambda$, respectively. Now, let $k_0 \in \mathcal{I}$ be the integer such that $|\gamma_{k_0} - E_i| = \min_{k \in \mathcal{I}} |\gamma_k - E_i|$. Using the eigenvalue rigidity (2.22) and the QUE estimate (2.6) for $H_{\Lambda}(t)$ in Lemma 6.4, we can bound (6.14) as follows: with probability $1 - O(N^{-c})$,

$$(6.14) \lesssim \frac{1}{DN} \sum_{\substack{|k-k_0| \le N^{\varepsilon}}} \frac{N^{-c}}{\eta_i^2} + \frac{1}{DN} \sum_{\substack{N^{\varepsilon} < |k-k_0| \le N^{c}}} \frac{N^{-c}}{|k-k_0|^2/N^2} + \frac{1}{DN} \sum_{\substack{|k-k_0| > N^{c}}} \frac{1}{|k-k_0|^2/N^2} \\ \lesssim N^{1-c+2\sigma+\varepsilon},$$

$$(6.16)$$

for any small constant $\varepsilon \in (0, c)$. Similarly, we can bound (6.15) as

$$\mathbb{P}\left(\left|\left\langle G_i^2\left(D^{-1} - E_a\right)G_j^2 E_a\right\rangle\right| \ge N^{3-c+4\sigma+\varepsilon}\right) \lesssim N^{-c}.$$
(6.17)

Combining (6.16) and (6.17) with (6.12), we obtain that for any constant $\varepsilon > 0$,

$$\mathbb{P}\left(|X(i;a)| + |Y(i,j;a)| \ge N^{1-c+(n+2)\sigma+\varepsilon}\right) \le N^{-c}.$$

Together with the rough bound (6.13), it yields that

$$\mathcal{F}_2 \lesssim N^{1-c+(n+2)\sigma+\varepsilon} + N^{1+(n+2)\sigma+\varepsilon} \cdot N^{-c} \leq 2N^{1-c+(n+2)\sigma+\varepsilon}.$$
(6.18)

Finally, plugging (6.11) and (6.18) into (6.10), we conclude that for any small constant $\varepsilon > 0$,

$$\left|\partial_t \mathbb{E}\prod_{i=1}^n \langle G_t(z_i) \rangle \right| \le N^{1-c+(n+2)\sigma+\varepsilon} + N^{1/2+(n+3)\sigma+\varepsilon}, \quad t \in [0,\mathfrak{t}].$$

if we choose $\sigma \in (0, 1/2)$. Integrating over t, we conclude (6.7) with $\delta = (c - (n+2)\sigma - \varepsilon - \mathfrak{c}) \wedge (1/2 - (n+3)\sigma - \varepsilon - \mathfrak{c})$ as long as we choose \mathfrak{c} , σ , and ε sufficiently small depending on c.

7. INTEGRABLE REGIME: EIGENVECTORS

In this section, we prove Theorem 2.4. For notational convenience, we will assume D = 2 in the subsequent proof, although the argument remains the same for the general case of $D \ge 2$.

Take the k-th bulk eigenvalue λ_k with $k \in [[DN, (1 - \kappa)DN]]$. Denote the corresponding eigenvector by $\mathbf{v}_k = \begin{pmatrix} \mathbf{u}_k \\ \mathbf{w}_k \end{pmatrix}$. Then, we have the eigenvalue equation

$$H\begin{pmatrix}\mathbf{u}_k\\\mathbf{w}_k\end{pmatrix} = \begin{pmatrix}H_1 & A\\A^* & H_2\end{pmatrix}\begin{pmatrix}\mathbf{u}_k\\\mathbf{w}_k\end{pmatrix} = \lambda_k\begin{pmatrix}\mathbf{u}_k\\\mathbf{w}_k\end{pmatrix}.$$

From this equation, we derive that

$$\mathbf{w}_{k} = -\mathcal{G}_{2}(\lambda_{k})A^{*}\mathbf{u}_{k}, \quad \mathbf{u}_{k} = -\mathcal{G}_{1}(\lambda_{k})A\mathbf{w}_{k}$$

where we denote the resolvents of H_1 and H_2 by

$$\mathcal{G}_1(z) := (H_1 - z)^{-1}, \quad \mathcal{G}_2(z) := (H_2 - z)^{-1}.$$

Now, given an arbitrarily small constant $\delta > 0$, we define the events

$$\mathscr{E}_1 := \left\{ \operatorname{dist}(\lambda_k, \operatorname{spec}(H_1)) \ge N^{-1-\delta} \right\}, \quad \mathscr{E}_2 := \left\{ \operatorname{dist}(\lambda_k, \operatorname{spec}(H_2)) \ge N^{-1-\delta} \right\}.$$

We claim that

$$\mathbb{P}\left(\mathscr{E}_1 \cup \mathscr{E}_2\right) = 1 - \mathcal{O}(N^{-\delta/2}). \tag{7.1}$$

To prove this claim, notice that

$$\mathbb{P}\left(\left(\mathscr{E}_1 \cup \mathscr{E}_2\right)^c\right) \le \mathbb{P}\left(\exists i, j \in [\![N]\!] \text{ such that } |\lambda_i^{(1)} - \lambda_j^{(2)}| \le 2N^{-1-\delta}\right),$$

where $\lambda_i^{(1)}$ and $\lambda_j^{(2)}$ denote the eigenvalues of H_1 and H_2 , respectively. Using the rigidity of eigenvalues for Wigner matrices [67, Theorem 2.2] (or using (2.22) in the case of D = 1), we get

$$\lambda_i^{(1)} - \gamma_i^{sc} | + |\lambda_i^{(2)} - \gamma_i^{sc}| \prec N^{-2/3} \min(i, N+1-i)^{-1/3}, \quad i \in [\![N]\!],$$
(7.2)

where $\gamma_i^{sc}, i \in [\![N]\!]$, denote the quantiles of the semicircle law:

$$\gamma_i^{sc} := \sup_{x \in \mathbb{R}} \left\{ \int_{-\infty}^x \rho_{sc}(x) \mathrm{d}x < \frac{i}{N} \right\}.$$

By the bulk universality of H_1 , there exists a constant $\delta > 0$ such that for any constant $\tau > 0$ and $j \in [N]$,

$$\mathbb{P}\left(\left|\exists i \in [[N]], \left|\lambda_i^{(1)} - \lambda_j^{(2)}\right| \le 2N^{-1-\delta} \right| H_2\right) \le N^{-\delta+\tau}.$$
(7.3)

In fact, [27, Proposition B.1] shows that (7.3) holds for a Gaussian divisible ensemble; then, applying the Green's function comparison theorem [64, Theorem 15.3 and Theorem 16.1] concludes (7.3). Let $k_0 := \arg \min_{i=1}^{N} |\gamma_i^{sc} - \gamma_k|$. Now, using (7.2) and (7.3), we obtain that for any constants $\tau, C > 0$,

$$\mathbb{P}\left(\left(\mathscr{E}_{1} \cup \mathscr{E}_{2}\right)^{c}\right) \leq \mathbb{P}\left(\exists i, j \in \llbracket k_{0} - N^{\tau}, k_{0} + N^{\tau}\rrbracket \text{ such that } |\lambda_{i}^{(1)} - \lambda_{j}^{(2)}| \leq 2N^{-1-\delta}\right) + \mathcal{O}(N^{-C})$$
$$\leq \sum_{j \in \llbracket k_{0} - N^{\tau}, k_{0} + N^{\tau}\rrbracket} \mathbb{P}\left(\exists i \in \llbracket N\rrbracket, |\lambda_{i}^{(1)} - \lambda_{j}^{(2)}| \leq 2N^{-1-\delta}\right) + \mathcal{O}(N^{-C}) = \mathcal{O}(N^{-\delta+2\tau}).$$

Taking $\tau < \delta/4$ concludes (7.1).

Without loss of generality, suppose \mathscr{E}_1 holds. Then, we claim the following estimate.

Lemma 7.1. Under the assumptions of Theorem 2.4 (with D = 2) and the notations defined above, the following estimate holds for any constants $\varepsilon, C > 0$:

$$\mathbb{P}\left(\|\mathcal{G}_1(\lambda_k)A\mathbf{w}_k\| \ge N^{\delta+\varepsilon}\|A\|_{\mathrm{HS}} + N^{-C}; \mathscr{E}_1\right) \le N^{-\varepsilon}.$$
(7.4)

Remark 7.2. The intuition behind the estimates (7.4) is that we believe \mathbf{w}_k is delocalized in the eigenbasis of H_1 on \mathscr{E}_1 , since H_1 and \mathbf{w}_k should be almost independent in the integrable regime. As a consequence, we expect that on \mathscr{E}_1 ,

$$\begin{aligned} \|\mathbf{u}_k\|^2 &\prec \frac{1}{N} \operatorname{Tr} \left[A^* \mathcal{G}_1(\lambda_k)^* \mathcal{G}_1(\lambda_k) A \right] \prec \|A\|_{\mathrm{HS}}^2 \max_i \frac{(\mathbf{u}_i^a)^* \mathcal{G}_1(\lambda_k)^* \mathcal{G}_1(\lambda_k) \mathbf{u}_i^a}{N} \\ &\prec \frac{\|A\|_{\mathrm{HS}}^2}{N^2} \sum_{i=1}^N \frac{1}{|\lambda_i^{(1)} - \lambda_k|^2} \prec N^{2\delta} \|A\|_{\mathrm{HS}}^2, \end{aligned}$$

where \mathbf{u}_i^a are the eigenvectors of AA^* . Above, in the second step, we used the definition of $||A||_{\text{HS}}^2$ in terms of the singular values of A; in the third step, we used the delocalization of \mathbf{u}_i^a in the eigenbasis of H_1 ; in the last step, we used the definition of \mathscr{E}_1 and the rigidity of eigenvalues (7.2).

Proof of Lemma 7.1. Let $z = E + i\eta$ with $E = \gamma_k$ and $\eta = N^{-1+c}$ for a small constant $c \in (0, 1/2)$. We claim that

$$\mathbb{E}\left(\|\mathcal{G}_{1}(\lambda_{k})A\mathbf{w}_{k}\|^{2};\mathscr{E}_{1}\right) \lesssim N^{2(c+\delta)}\mathbb{E}\operatorname{Tr}\left[\begin{pmatrix}0 & 0\\ 0 & A^{*}\operatorname{Im}\mathcal{G}_{1}(z)A\end{pmatrix}\operatorname{Im}G(z)\right].$$
(7.5)

To see why (7.5) holds, with the spectral decomposition of Im G, we obtain that

$$\mathbb{E}\operatorname{Tr}\left[\begin{pmatrix}0&0\\0&A^*\operatorname{Im}\mathcal{G}_1(z)A\end{pmatrix}\operatorname{Im}G(z)\right] = \mathbb{E}\sum_{j\in\mathcal{I}}\frac{\eta}{(\lambda_j-\gamma_k)^2+\eta^2}\mathbf{w}_j^*A^*\operatorname{Im}\mathcal{G}_1(z)A\mathbf{w}_j$$
$$\geq \mathbb{E}\frac{\eta}{(\lambda_k-\gamma_k)^2+\eta^2}\mathbf{w}_k^*A^*\operatorname{Im}\mathcal{G}_1(z)A\mathbf{w}_k \gtrsim \eta^{-1}\mathbb{E}\mathbf{w}_k^*A^*\operatorname{Im}\mathcal{G}_1(z)A\mathbf{w}_k,$$

where in the last step, we used the rigidity of λ_k given by (2.22). On the other hand, with the spectral decomposition of \mathcal{G}_1 , we obtain that on the event \mathscr{E}_1 , with high probability,

$$\eta^{2} \|\mathcal{G}_{1}(\lambda_{k})A\mathbf{w}_{k}\|^{2} = \sum_{j} \frac{\eta^{2} |(\mathbf{u}_{j}^{(1)})^{*}A\mathbf{w}_{k}|^{2}}{(\lambda_{j}^{(1)} - \lambda_{k})^{2}} \lesssim N^{2(c+\delta)} \sum_{j} \frac{\eta^{2} |(\mathbf{u}_{j}^{(1)})^{*}A\mathbf{w}_{k}|^{2}}{(\lambda_{j}^{(1)} - \lambda_{k})^{2} + \eta^{2}}$$
$$\lesssim N^{2(c+\delta)} \sum_{j} \frac{\eta^{2} |(\mathbf{u}_{j}^{(1)})^{*}A\mathbf{w}_{k}|^{2}}{(\lambda_{j}^{(1)} - \gamma_{k})^{2} + \eta^{2}} = N^{2(c+\delta)} \cdot \eta \mathbf{w}_{k}^{*}A^{*} \operatorname{Im} \mathcal{G}_{1}(z)A\mathbf{w}_{k}$$

where $\mathbf{u}_{j}^{(1)}$, $j \in [N]$, denote the eigenvectors of H_1 , and we used the definition of \mathscr{E}_1 in the second step and the rigidity of λ_k in the third step. The above two estimates together conclude (7.5).

Using (7.5) and the trivial bound

$$\mathbb{E}\operatorname{Tr}\left[\begin{pmatrix} 0 & 0\\ 0 & A^*\operatorname{Im}\mathcal{G}_1(z)A \end{pmatrix}\operatorname{Im}G(z)\right] \le \mathbb{E}\operatorname{Tr}\left[\operatorname{Im}G_0(z)\Lambda\operatorname{Im}G(z)\Lambda\right] \quad \text{with} \quad G_0(z) := (H-z)^{-1},$$

we obtain that

$$\mathbb{E}\left(\|\mathcal{G}_1(\lambda_k)A\mathbf{w}_k\|^2;\mathscr{E}_1\right) \lesssim N^{2(c+\delta)}\mathbb{E}\operatorname{Tr}\left[\operatorname{Im} G_0(z)\Lambda\operatorname{Im} G(z)\Lambda\right].$$
(7.6)

The estimate (7.4) then follows from the next lemma together with Markov's inequality.

Lemma 7.3. Under the setting of Theorem 2.4 (without requiring that D = 2), let $z = E + i\eta$ with $E = \gamma_k$ and $\eta = N^{-1+c}$. Then, we have that for any constant C > 0,

$$\mathbb{E}\operatorname{Tr}\left[\operatorname{Im} G_0(z)\Lambda \operatorname{Im} G(z)\Lambda\right] \prec \|A\|_{\mathrm{HS}}^2 + N^{-C}.$$
(7.7)

The proof of this lemma is the same as that for Lemma 8.2 below, which will be presented in full detail in Section 8. Hence, we omit the proof of Lemma 7.3 here.

Combining (7.7) with (7.6) gives that

$$\mathbb{E}\left(\|\mathcal{G}_1(\lambda_k)A\mathbf{w}_k\|^2;\mathscr{E}_1\right) \prec N^{2(c+\delta)}(\|A\|_{\mathrm{HS}}^2 + N^{-C}).$$

Then, applying Markov's inequality concludes (7.4) since c is arbitrarily small.

Now, we are ready to conclude Theorem 2.4.

Proof of Theorem 2.4. When
$$D = 2$$
, by Lemma 7.1, we have that for any constant $\varepsilon > 0$,

$$\mathbb{P}\left(\|\mathbf{u}_k\| \ge N^{-\varepsilon_A + \delta + \varepsilon}; \mathscr{E}_1\right) \le N^{-\varepsilon}.$$

By symmetry, a similar estimate holds for \mathbf{w}_k on \mathscr{E}_2 . Together with (7.1), these estimates imply that

 $\mathbb{P}\left(\min\{\|\mathbf{u}_k\|, \|\mathbf{w}_k\|\} \ge N^{-\varepsilon_A + \delta + \varepsilon}\right) \lesssim N^{-\varepsilon} + N^{-\delta/2},$

which concludes (2.8) as long as we choose $\delta + \varepsilon < \varepsilon_A$.

For the general case of $D \ge 2$, the proof is similar. More precisely, define events $\mathscr{E}_{(a)}, a \in \llbracket D \rrbracket$, as

$$\mathscr{E}_{(a)} := \left\{ \operatorname{dist} \left(\lambda_k, \bigcup_{k \in \llbracket D \rrbracket \setminus \{a\}} \operatorname{spec}(H_k) \right) \ge N^{-1-\delta} \right\}$$

Again, using the eigenvalue rigidity and bulk universality for Wigner matrices, we get that $\mathbb{P}(\bigcup_{a=1}^{D} \mathscr{E}_{(a)}) = 1 - O(N^{-\delta/2})$. Then, similar to (7.6), we can show that

$$\sum_{k \in [D] \setminus \{a\}} \mathbb{E} \left(\|E_k \mathbf{v}_k\|^2; \mathscr{E}_{(a)} \right) \lesssim N^{2(c+\delta)} \mathbb{E} \operatorname{Tr} \left[\operatorname{Im} G_0(z) \Lambda \operatorname{Im} G(z) \Lambda \right]$$

Applying Lemma 7.3 and Markov's inequality, we obtain that for any constant $\varepsilon > 0$,

$$\mathbb{P}\left(1 - \|E_a \mathbf{v}_k\|^2 \ge N^{-2\varepsilon_A + 2\delta + \varepsilon}; \mathscr{E}_{(a)}\right) \lesssim N^{-\varepsilon/2} + N^{-\delta/2}.$$

Finally, taking the union bound over $a \in \llbracket D \rrbracket$ concludes the proof.

8. INTEGRABLE REGIME: EIGENVALUES

Finally, in this section, we give the proof of Theorem 2.5. Recall the interpolating matrices defined in (2.24). Denote the eigenvalues and corresponding eigenvectors of $H_{\Lambda}(\theta)$ by $\lambda_i(\theta)$ and $\mathbf{v}_i(\theta)$, $i \in \mathcal{I}$. Then, we have that for any $k \in \mathcal{I}$,

$$\lambda_k(1) - \lambda_k(0) = \int_0^1 \frac{\mathrm{d}}{\mathrm{d}\theta} \lambda_k(\theta) \mathrm{d}\theta = \int_0^1 \mathbf{v}_k(\theta)^* \Lambda \mathbf{v}_k(\theta) \mathrm{d}\theta, \qquad (8.1)$$

from which we derive that

$$\mathbb{E}\left|\lambda_{k}(1)-\lambda_{k}(0)\right| \leq \int_{0}^{1} \mathbb{E}\left|\mathbf{v}_{k}(\theta)^{*}\Lambda\mathbf{v}_{k}(\theta)\right| \mathrm{d}\theta \leq \int_{0}^{1} \left(\mathbb{E}\left|\mathbf{v}_{k}(\theta)^{*}\Lambda\mathbf{v}_{k}(\theta)\right|^{2}\right)^{1/2} \mathrm{d}\theta.$$
(8.2)

Remark. Rigorously speaking, $d\lambda_k(\theta)/d\theta = \mathbf{v}_k(\theta)^* \Lambda \mathbf{v}_k(\theta)$ holds when $\lambda_k(\theta)$ is not a degenerate eigenvalue of $H_{\Lambda}(\theta)$. To handle this issue, we can add a small Gaussian component to H so that each entry of the diagonal blocks has a continuous distribution. For example, we can take a matrix H'_{Λ} whose blocks satisfy the law (4.8) with $t - t_0 = e^{-N}$. Then, almost surely, all the eigenvalues of H'_{Λ} are non-degenerate, which, together with Fubini's theorem, implies that

$$\mathbb{E}[\lambda_k(1) - \lambda_k(0)] = \int_0^1 \mathbb{E}\mathbf{v}_k(\theta)^* \Lambda \mathbf{v}_k(\theta) \mathrm{d}\theta.$$

Hence, (8.2) remains valid for the ensemble H'_{Λ} . After showing (2.9) for H'_{Λ} , we then remove the small Gaussian component, which leads to a negligible error of order $O_{\prec}(e^{-N})$ to both λ_k and $\lambda_k(H)$. For clarity and without loss of generality, it suffices to assume that (8.2) holds throughout the following proof.

Given (8.2), we will prove that there exists a constant $\varepsilon'_A > 0$ (depending on ε_A) such that for any $\theta \in [0, 1]$ and $k \in [\kappa DN, (1 - \kappa)DN]$,

$$\left(\mathbb{E}\left|\mathbf{v}_{k}(\theta)^{*}\Lambda\mathbf{v}_{k}(\theta)\right|^{2}\right)^{1/2} \leq N^{-1-\varepsilon_{A}'}.$$
(8.3)

Plugging this bound into (8.2) and applying Markov's inequality, we get that

$$\mathbb{P}\left(|\lambda_k(1) - \lambda_k(0)| \ge N^{-1-\varepsilon_A'/2}\right) \le N^{-\varepsilon_A'/2},$$

which concludes the proof.

Remark 8.1. We now explain heuristically why the estimate (8.3) should hold. Fix any $\theta \in [0, 1]$, suppose $\mathbf{v}_k \equiv \mathbf{v}_k(\theta)$ is decomposed into D sub-vectors $\mathbf{v}_{k,i} := (v_k(x) : x \in \mathcal{I}_i), i \in [D]$. Then, $\mathbf{v}_k^* \Lambda \mathbf{v}_k$ involves inner products $\mathbf{v}_{k,i}^* A \mathbf{v}_{k,j}, j \neq i$, and their complex conjugates. If we believe that the eigenvectors of different blocks $H_i, i \in [D]$, do not mix (which is partially suggested by Theorem 2.4), then $\mathbf{v}_{k,i}$ and $\mathbf{v}_{k,j}$ should be approximately independent random vectors, which implies that $\mathbf{v}_{k,i}^* A \mathbf{v}_{k,j} \prec N^{-1} \|A\|_{\mathrm{HS}} \leq N^{-1-\varepsilon_A}$.

The rest of this section is devoted to the proof of (8.3). Without loss of generality, we can take $\theta = 1$ in the following proof—for a general θ , we can just take $\theta \Lambda$ as our new Λ . With the spectral decomposition of Im G, we get that

$$\operatorname{Tr}\left[(\operatorname{Im} G)\Lambda(\operatorname{Im} G)\Lambda\right] = \sum_{k,l\in\mathcal{I}} \frac{\eta^2 |\mathbf{v}_k^*\Lambda\mathbf{v}_l|^2}{|z-\lambda_k|^2|z-\lambda_l|^2}, \quad z = E + \mathrm{i}\eta.$$

Let $E = \gamma_k$ and $\eta = N^{-1+c}$ for a small constant c > 0. Then, we obtain from the above equation that

$$\mathbb{E}|\mathbf{v}_k^*\Lambda\mathbf{v}_k|^2 \le \mathbb{E}\frac{\left[(\lambda_k - \gamma_k)^2 + \eta^2\right]^2}{\eta^2} \operatorname{Tr}\left[(\operatorname{Im} G)\Lambda(\operatorname{Im} G)\Lambda\right] \prec \eta^2 \mathbb{E}\operatorname{Tr}\left[(\operatorname{Im} G)\Lambda(\operatorname{Im} G)\Lambda\right],$$
(8.4)

where we used the eigenvalue rigidity (2.22) in the second step. Now, to conclude (8.3), it suffices to show that for $z = \gamma_k + i N^{-1+c}$,

 $\mathbb{E}\operatorname{Tr}\left[(\operatorname{Im} G)\Lambda(\operatorname{Im} G)\Lambda\right]\prec \|A\|_{\mathrm{HS}}^2+N^{-1}.$

Writing $\operatorname{Im} G = (G - G^*)/(2i)$, it suffices to prove the following lemma.

Lemma 8.2. Under the setting of Theorem 2.5, let $c \in (0,1)$ be an arbitrary small constant. Fix any $z = E + i\eta$ with $|E| \le 2 - \kappa$ and $\eta = N^{-1+c}$. Then, for $z_1, z_2 \in \{z, \bar{z}\}$ and any constant C > 0, we have that $\mathbb{E} \langle G_1 \Lambda G_2 \Lambda \rangle \prec N^{-1} ||A||_{\text{HS}}^2 + N^{-C}$. (8.5)

Again, the proof of this estimate relies on (4.23) and the cumulant expansion formula, Lemma 4.10. We first consider a simpler case where the entries of H are complex Gaussian. In this case, Lemma 4.10 can be replaced by the Gaussian integration by parts.

8.1. Proof of Lemma 8.2: Gaussian case. We recall the definitions of the scalar m and the matrix Min (2.15) and (2.16). In the following proof, we denote $\widetilde{m} := -(m+z)^{-1}$ and

$$\widetilde{M} := M - \widetilde{m} = -\widetilde{m}\Lambda M = -\sum_{k=1}^{\infty} (m+z)^{-(k+1)}\Lambda^k.$$
(8.6)

When $z = z_i$, $i \in \{1, 2\}$, we then use the notations \widetilde{m}_i and M_i .

By the averaged local law (2.21), we have that

$$\langle M_1 \Lambda G_2 \Lambda \rangle = \langle \Lambda^2 M_1 G_2 \rangle \prec N^{-1} \|A\|_{\mathrm{HS}}^2$$

Thus, we only need to bound $\langle (G_1 - M_1)\Lambda G_2\Lambda \rangle$. With (4.23) and Gaussian integration by parts, we get that

$$\mathbb{E}\langle (G_1 - M_1)\Lambda G_2\Lambda \rangle = -\mathbb{E}\langle G_1(H + m_1)\widetilde{M}_1\Lambda G_2\Lambda \rangle - \widetilde{m}_1\mathbb{E}\langle G_1(H + m_1)\Lambda G\Lambda \rangle = Y_1, \tag{8.7}$$

where Y_1 is defined as

$$Y_{1} = D\mathbb{E}\sum_{a=1}^{D} \langle G_{1}E_{a}\widetilde{M}_{1}\Lambda G_{2}\Lambda \rangle \langle (G_{1} - M_{1})E_{a} \rangle + D\mathbb{E}\sum_{a=1}^{D} \langle G_{1}E_{a}G_{2}\Lambda \rangle \langle \widetilde{M}_{1}\Lambda G_{2}E_{a} \rangle + D\widetilde{m}_{1}\mathbb{E}\sum_{a=1}^{D} \langle G_{1}E_{a}\Lambda G_{2}\Lambda \rangle \langle (G_{1} - M_{1})E_{a} \rangle + D\widetilde{m}_{1}\mathbb{E}\sum_{a=1}^{D} \langle G_{1}E_{a}G_{2}\Lambda \rangle \langle \Lambda G_{2}E_{a} \rangle.$$

$$(8.8)$$

We can keep expanding Y_1 using (4.23) and Gaussian integration by parts. During the expansions, all our expressions can be written into certain forms with some loop structures, which we now describe in more detail. In the expansions, each term only contains a deterministic coefficient and the following four types of loops (i.e., terms of the form $\langle \cdot \rangle$):

- (i) Light weights: $\langle (G M)E_{a_i} \rangle$ for some $a_i \in \llbracket D \rrbracket$.
- (ii) *G*-loops: $\langle \mathcal{G}^{(k)} \rangle$ for some $k \geq 2$, where $\mathcal{G}^{(k)}$ represents an expression of the form

$$\mathcal{G}^{(k)} = E_{a_0} \prod_{i=1}^k G_{s_i} E_{a_i}$$
(8.9)

for some sequences $a_i \in \llbracket D \rrbracket$ and $s_i \in \{1, 2\}$.

(iii) (G, Λ) -loops: $\langle \mathcal{G}^{(k)} \Lambda^{(n)} \rangle$ for some $k \geq 0$ and $n \geq 1$, where $\mathcal{G}^{(k)}$ is of the form (8.9) (with the convention that $\mathcal{G}^{(0)} = 1$ and $\Lambda^{(n)}$ represents an expression of the form

$$\Lambda^{(n)} \sim E_{a_0} \prod_{i=1}^n \Lambda E_{a_i} \tag{8.10}$$

for some sequence $a_i \in \llbracket D \rrbracket$. (iv) (G, Λ, G, Λ) -loops: $\langle \mathcal{G}^{(k_1)} \Lambda^{(n_1)} \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle$ for some $k_1, k_2, n_1, n_2 \geq 1$.

Let $\mathcal{W}^{(k)}$ denote a product of light weights of the form $\mathcal{W}^{(k)} = \prod_{i=1}^{k} \langle (G-M) E_{a_i} \rangle$, and let $\Gamma_n^{(m)}$ denote a product of G-loops of the form

$$\Gamma_n^{(m)} = \prod_{i=1}^n \langle \mathcal{G}^{(k_i)} \rangle \quad \text{with} \quad k_i \ge 2, \ \sum_{i=1}^n k_i = m.$$

Hereafter, we slightly abuse the notation and use m to denote an integer instead of m(z) defined in (2.15).

We will show that by following a specific expansion strategy, we can continue expanding Y_1 until each expression is bounded by $O_{\prec}(N^{-1}||A||_{\rm HS}^2 + N^{-C})$ as required. Before introducing the formal expansion strategy, we will examine some low-order expansions as illustrative examples. In equation (8.8), the first two terms have "smaller sizes" compared to the third and fourth terms, owing to the additional M_1 factor. Utilizing the local laws (2.21) and (2.23) along with the estimates (8.25)-(8.29) below, we can show that

$$\mathbb{E}\langle G_1 E_a \widetilde{M}_1 \Lambda G_2 \Lambda \rangle \langle (G_1 - M_1) E_a \rangle + \mathbb{E}\langle G_1 E_a G_2 \Lambda \rangle \langle \widetilde{M}_1 \Lambda G_2 E_a \rangle \prec \frac{\|A\|_{\mathrm{HS}}}{\sqrt{N\eta}} \frac{\|A\|_{\mathrm{HS}}^2}{N},$$
(8.11)

$$\mathbb{E}\langle G_1 E_a \Lambda G_2 \Lambda \rangle \langle (G_1 - M_1) E_a \rangle + \mathbb{E} \langle G_1 E_a G_2 \Lambda \rangle \langle \Lambda G_2 E_a \rangle \prec \frac{\|A\|_{\mathrm{HS}}^2}{N\eta}.$$
(8.12)

Therefore, we will focus on expanding the two terms in (8.12). Using (4.23), we can expand them as

$$\mathbb{E}\langle G_1 E_a \Lambda M_2 \Lambda \rangle \langle (G_1 - M_1) E_a \rangle - \mathbb{E} \langle G_1 E_a \Lambda G_2 (H + m_2) M_2 \Lambda \rangle \langle (G_1 - M_1) E_a \rangle + \mathbb{E} \langle G_1 E_a M_2 \Lambda \rangle \langle \Lambda G_2 E_a \rangle - \mathbb{E} \langle G_1 E_a G_2 (H + m_2) M_2 \Lambda \rangle \langle \Lambda G_2 E_a \rangle.$$
(8.13)

With the estimate (8.28) below, we can already bound the first and third terms by $O_{\prec}(N^{-1}||A||_{HS}^2)$. For the second and fourth terms in (8.13), by applying Gaussian integration by parts, we find:

$$D\mathbb{E}\sum_{x=1}^{D} \langle G_2 E_x M_2 \Lambda G_1 E_a \Lambda \rangle \langle (G_1 - M_1) E_a \rangle \langle (G_2 - M_2) E_x \rangle$$
(8.14)

+
$$D\mathbb{E}\sum_{x=1}^{D} \langle G_2 E_x G_1 E_a \Lambda \rangle \langle G_1 E_x M_2 \Lambda \rangle \langle (G_1 - M_1) E_a \rangle$$
 (8.15)

$$+ \frac{1}{DN^2} \mathbb{E} \sum_{x=1}^{D} \langle G_2 E_x G_1 E_a G_1 E_x M_2 \Lambda G_1 E_a \Lambda \rangle$$
(8.16)

$$+ D\mathbb{E}\sum_{x=1}^{D} \langle G_1 E_a G_2 E_x M_2 \Lambda \rangle \langle (G_2 - M_2) E_x \rangle \langle \Lambda G_2 E_a \rangle$$
(8.17)

$$+ D\mathbb{E}\sum_{x=1}^{D} \langle G_1 E_x M_2 \Lambda \rangle \langle G_1 E_a G_2 E_x \rangle \langle \Lambda G_2 E_a \rangle$$
(8.18)

$$+ \frac{1}{DN^2} \mathbb{E} \sum_{x=1}^{D} \langle G_1 E_a G_2 E_x G_2 E_a \Lambda G_2 E_x M_2 \Lambda \rangle.$$
(8.19)

Note the expressions (8.14) and (8.17) exhibit similar loop structures to the two terms in (8.12), albeit with an additional light weight. Consequently, their sizes are smaller by an extra factor of $(N\eta)^{-1}$:

$$(8.14) + (8.17) \prec ||A||_{\text{HS}}^2 / (N\eta)^2.$$

Moving from the two terms in (8.12) to (8.16) and (8.19), the two loops combine into a single loop while gaining an N^{-2} coefficient. Utilizing the estimate (8.29) below, we can bound them by

$$(8.16) + (8.19) \prec ||A||_{\rm HS}^2 / (N\eta)^2,$$

which is also better than (8.12) by a factor of $(N\eta)^{-1}$. Furthermore, from the first term in (8.12) to the expression (8.15), a (G, Λ, G, Λ) -loop is split into two (G, Λ) -loops. Using the estimate (8.28) below, we can show that (8.15) also satisfies a better bound than (8.12):

$$(8.15) \prec ||A||_{\text{HS}}^2 / (N\eta)^2.$$

It remains to address the expression (8.18). Utilizing the estimate (8.28), we can show that

$$(8.18) \prec \|A\|_{\rm HS}^2 / (N\eta). \tag{8.20}$$

While there is no improvement compared to the bound (8.12), we have a change in the loop structure. Specifically, two (G, Λ) -loops, denoted as $\langle \mathcal{G}^{(k_1)} \Lambda^{(n_1)} \rangle \langle \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle$ with $k_1 = 2$, $k_2 = 1$, and $n_1 = n_2 = 1$, transform into a $\langle \mathcal{G}^{(2)} \rangle$ loop along with two (G, Λ) -loops with $k_1 = k_2 = 1$ and $n_1 = n_2 = 1$. The key feature of this change is that $k_1 + k_2$ is decreased by 1, indicating that the two (G, Λ) -loops become "more deterministic". To proceed, we further expand (8.18) (without the summation over x and the D factor) as

$$\mathbb{E}\langle M_1 E_x M_2 \Lambda \rangle \langle G_2 E_a \Lambda \rangle \langle G_1 E_a G_2 E_x \rangle - \mathbb{E} \langle G_1 (H+m_1) M_1 E_x M_2 \Lambda \rangle \langle G_2 E_a \Lambda \rangle \langle G_1 E_a G_2 E_x \rangle.$$
(8.21)

Note that the first term contains two (G, Λ) -loops $\langle \mathcal{G}^{(k_1)} \Lambda^{(n_1)} \rangle \langle \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle$ with $k_1 = 0$, $k_2 = 1$, and $n_1 = n_2 = 1$, making it even "more deterministic" than (8.18). Moreover, recalling (8.6), the factor $\langle M_1 E_x M_2 \Lambda \rangle$ can be bounded by

$$\langle M_1 E_x M_2 \Lambda \rangle = \widetilde{m}_1 \langle E_x M_2 \Lambda \rangle + \widetilde{m}_2 \langle M_1 E_x \Lambda \rangle + \langle M_1 E_x M_2 \Lambda \rangle \lesssim N^{-1} \|A\|_{\mathrm{HS}}^2$$

This estimate, together with (2.23) and the estimate (8.28) below, implies that

$$\mathbb{E}\langle M_1 E_x M_2 \Lambda \rangle \langle G_2 E_a \Lambda \rangle \langle G_1 E_a G_2 E_x \rangle \prec \frac{\|A\|_{\mathrm{HS}}}{\sqrt{N}\eta} \cdot \frac{\|A\|_{\mathrm{HS}}^2}{N}$$

an improvement over (8.20) by a factor of $N^{-1/2} ||A||_{\text{HS}}$. Subsequently, by applying Gaussian integration by parts to the second term in (8.21), we obtain that

$$\begin{split} D\mathbb{E}\sum_{y=1}^{D} &\langle G_{1}E_{y}M_{1}E_{x}M_{2}\Lambda\rangle\langle G_{2}E_{a}\Lambda\rangle\langle (G_{1}-M_{1})E_{y}\rangle\langle G_{1}E_{a}G_{2}E_{x}\rangle\\ &+\frac{1}{DN^{2}}\mathbb{E}\sum_{y=1}^{D} \langle G_{2}E_{y}M_{1}E_{x}M_{2}\Lambda G_{1}E_{y}G_{2}E_{a}\Lambda\rangle\langle G_{1}E_{a}G_{2}E_{x}\rangle\\ &+\frac{1}{DN^{2}}\mathbb{E}\sum_{y=1}^{D} \langle G_{1}E_{y}M_{1}E_{x}M_{2}\Lambda G_{1}E_{y}G_{1}E_{a}G_{2}E_{x}\rangle\langle G_{2}E_{a}\Lambda\rangle\\ &+\frac{1}{DN^{2}}\mathbb{E}\sum_{y=1}^{D} \langle G_{1}E_{a}G_{2}E_{y}M_{1}E_{x}M_{2}\Lambda G_{1}E_{y}G_{2}E_{x}\rangle\langle G_{2}E_{a}\Lambda\rangle. \end{split}$$

By using the estimates (8.28) and (8.29) below, we can show that the first two expressions are bounded by $O_{\prec}(\|A\|_{\text{HS}}^2/(N\eta)^2)$, while the latter two expressions are bounded by $O_{\prec}(\|A\|_{\text{HS}}^2/(N\eta)^3)$. These bounds are better than (8.20) by at least a factor of $(N\eta)^{-1}$.

Inspired by the above calculations, we define the following two types of expressions with $n_1, n_2 \ge 1$:

• Type I expressions: $\mathcal{W}^{(k)}\Gamma_n^{(m)}\langle \mathcal{G}^{(k_1)}\Lambda^{(n_1)}\rangle\langle \mathcal{G}^{(k_2)}\Lambda^{(n_2)}\rangle$ with a deterministic coefficient of order

$$O\left(N^{-(m-n+k_1+k_2-3)-\mathbf{1}(k_1=0)-\mathbf{1}(k_2=0)}\right).$$
(8.22)

• Type II expressions: $\mathcal{W}^{(k)}\Gamma_n^{(m)}\langle \mathcal{G}^{(k_1)}\Lambda^{(n_1)}\mathcal{G}^{(k_2)}\Lambda^{(n_2)}\rangle$ with a deterministic coefficient of order

$$O\left(N^{-(m-n+k_1+k_2-2)-\mathbf{1}(k_1=0)-\mathbf{1}(k_2=0)}\right).$$
(8.23)

In our expansions, Y_1 can be expanded (in the sense of equal expectation) into a linear combination of these two types of expressions. The coefficients take the forms (8.22) and (8.23) due to the following reasons. In (8.8), every term has three G factors and two N^{-1} factors associated with the two loops, so the number of G factors is greater than the number of N^{-1} factors by 1. For each application of Gaussian integration by parts, we gain one more G factor and one more N^{-1} factor. Hence, we should have $(m-n) + (k_1-1) + (k_2-1) - 1$ many N^{-1} factors for type I expressions and $(m-n) + (k_1 + k_2 - 1) - 1$ many N^{-1} factors for type II expressions if $k_1 \neq 0$ and $k_2 \neq 0$. The case of $k_1 = 0$ or $k_2 = 0$ happens when we have replaced (at least) one or two G_i , $i \in \{1, 2\}$, factors with M_i , so we should gain one or two more N^{-1} factors in those cases.

We are now prepared to describe our expansion strategy and a systematic approach to bounding all expressions that appear during the expansion process. Prior to that, let us provide a brief overview of our proof. We will see that for a type I expression (resp. type II expression), if $k_1 = k_2 = 0$ (resp. $k_1 = 0$ or $k_2 = 0$), it can already be bounded by $O_{\prec}(N^{-1}||A||_{HS}^2)$. Otherwise, suppose $k_1 > 0$ without loss of generality. Then, we select the first G to the right of $\Lambda^{(n_1)}$, denoted as G_s with $s \in \{1, 2\}$. We expand this G_s with (4.23), i.e., $G_s = M_s - G_s(H + m_s)M_s$, and subsequently apply Gaussian integration by parts. Every resulting expression from this expansion satisfies one of the following properties:

- It can be bounded by $O_{\prec}(N^{-1}||A||_{\text{HS}}^2)$.
- It satisfies a better bound compared to the original expression by at least a factor of $(N\eta)^{-1}$.
- It is "more deterministic" than the original expression in the sense that $k_1 + k_2$ becomes strictly smaller.

This expansion process can be iterated until every resulting expression can be bounded by $O_{\prec}(N^{-1}||A||_{\text{HS}}^2 + N^{-C})$. The number of iterations is at most O(1).

With the above idea in mind, we now begin with the formal proof. With the local laws in Lemma 2.10 and Lemma 2.11, we immediately obtain that

$$\mathcal{W}^{(k)} \prec (N\eta)^{-k}, \quad \Gamma_n^{(m)} \prec \eta^{-(m-n)}. \tag{8.24}$$

Furthermore, let $G^{(k)}$ be an $N \times N$ block of the matrix $B_0 \prod_{i=1}^k G_i B_i$, where $(B_i)_{0 \leq i \leq k}$ is an arbitrary sequence of deterministic matrices satisfying $||B_i|| \leq 1$. Then, using Lemma 2.11 and the singular value decomposition $A = \sum_i \lambda_i^a \mathbf{u}_i^a (\mathbf{v}_i^a)^*$, we obtain that

$$Tr(G^{(k)}A) = \sum_{i} \lambda_{i}^{a}(\mathbf{v}_{i}^{a})^{*} G^{(k)} \mathbf{u}_{i}^{a} \prec \eta^{-(k-1)} \sqrt{N} \|A\|_{HS},$$
(8.25)

$$\operatorname{Tr}(G^{(k)}AA^*) = \sum_{i} (\lambda_i^a)^2 (\mathbf{u}_i^a)^* G^{(k)} \mathbf{u}_i^a \prec \eta^{-(k-1)} \|A\|_{\operatorname{HS}}^2,$$
(8.26)

$$\operatorname{Tr}(G^{(k)}A^2) = \sum_{i,j} \lambda_i^a \lambda_j^a (\mathbf{v}_j^a)^* G^{(k)} \mathbf{u}_i^a (\mathbf{v}_i^a)^* \mathbf{u}_j^a \prec \eta^{-(k-1)} \sum_j \lambda_j^a \left\| \sum_i \lambda_i^a \mathbf{u}_i^a (\mathbf{v}_i^a)^* \mathbf{u}_j^a \right\| \prec \eta^{-(k-1)} \|A\|_{\operatorname{HS}}^2.$$
(8.27)

Similar bounds also hold if we switch A and A^* . With these bounds, we can readily derive that

$$\langle \mathcal{G}^{(k)} \Lambda^{(n)} \rangle \prec \eta^{-(k-1)} \|A\|^{n-[1+1(n\geq 2)]} \cdot (N^{-1/2} \|A\|_{\mathrm{HS}})^{1+1(n\geq 2)}$$
(8.28)

for any $\mathcal{G}^{(k)}$ and $\Lambda^{(n)}$ with $k, n \geq 1$. Following a similar argument based on Lemma 2.11, we also obtain that

$$\langle \mathcal{G}^{(k_1)} \Lambda^{(n_1)} \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle \prec \frac{\|A\|^{n_1+n_2-2-(\mathbf{1}(n_1 \ge 2)+\mathbf{1}(n_2 \ge 2))}}{\eta^{k_1+k_2-2}} \cdot (N^{-1/2} \|A\|_{\mathrm{HS}})^{\mathbf{1}(n_1 \ge 2)+\mathbf{1}(n_2 \ge 2)} \|A\|_{\mathrm{HS}}^2$$
(8.29)

for any $\mathcal{G}^{(k_1)}, \mathcal{G}^{(k_2)}, \Lambda^{(n_1)}, \Lambda^{(n_2)}$ with $k_1, k_2, n_1, n_2 \ge 1$. When k = 0 and n = 1, we trivially have $\langle \mathcal{G}^{(0)} \Lambda^{(n)} \rangle = 0$; when k = 0 and $n \ge 2$, we have

$$\langle \mathcal{G}^{(0)} \Lambda^{(n)} \rangle \prec \|A\|^{n-2} \cdot N^{-1} \|A\|_{\mathrm{HS}}^2.$$
 (8.30)

When $k_1 = 0$ and $k_2, n_1, n_2 \ge 1$, we have

$$\langle \mathcal{G}^{(0)} \Lambda^{(n_1)} \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle \prec \eta^{-(k_2 - 1)} \|A\|^{n_1 + n_2 - 2} \cdot N^{-1} \|A\|^2_{\mathrm{HS}}.$$
 (8.31)

For simplicity of presentation, we use notations $\mathcal{T}_1(k, m, n; k_1, k_2, n_1, n_2)$ and $\mathcal{T}_2(k, m, n; k_1, k_2, n_1, n_2)$ to denote generic type I and type II expressions, together with their deterministic coefficients satisfying (8.22) and (8.23). Their exact forms may change from one line to the next. Using the above bounds (8.24)–(8.31), we readily obtain some "rough bounds" on type I and type II expressions: for $i \in \{1, 2\}$ and $n_1, n_2 \ge 1$,

$$\mathcal{T}_{i}(k,m,n;k_{1},k_{2},n_{1},n_{2}) \prec \left(\frac{1}{N\eta}\right)^{k+m-n+k_{1}+k_{2}-2} \cdot (N^{-1/2})^{\mathbf{1}(n_{1}\geq2)+\mathbf{1}(n_{2}\geq2)} \|A\|_{\mathrm{HS}}^{n_{1}+n_{2}}.$$
(8.32)

(Note we allow that $k_1 = 0$ or $k_2 = 0$ in this bound.) When $k_1 = 0$, we have better bounds:

$$\mathcal{T}_{1}(k,m,n;0,k_{2},n_{1},n_{2}) \prec \left(\frac{1}{N\eta}\right)^{k+m-n+k_{2}-1} \cdot (N^{-1/2})^{1+1(n_{2}\geq2)} \|A\|_{\mathrm{HS}}^{n_{1}+n_{2}}, \quad k_{2}\geq1,$$
(8.33)

$$\mathcal{T}_{2}(k,m,n;0,k_{2},n_{1},n_{2}) \prec \left(\frac{1}{N\eta}\right)^{k+m-n+k_{2}-1} \cdot N^{-1} \|A\|_{\mathrm{HS}}^{n_{1}+n_{2}}, \quad k_{2} \ge 1,$$
(8.34)

$$\mathcal{T}_{i}(k,m,n;0,0,n_{1},n_{2}) \prec \left(\frac{1}{N\eta}\right)^{k+m-n} \cdot N^{-1} \|A\|_{\mathrm{HS}}^{n_{1}+n_{2}}, \quad i \in \{1,2\}.$$
(8.35)

Similar bounds also hold when $k_2 = 0$. The bounds (8.32)–(8.35) show that the following expressions can be bounded by $O_{\prec}(N^{-1}||A||_{\text{HS}}^2)$ as desired:

- (E1) $\mathcal{T}_i(k, m, n; k_1, k_2, n_1, n_2), i \in \{1, 2\}, \text{ with } n_1, n_2 \ge 2;$
- (E2) $\mathcal{T}_i(k, m, n; k_1, k_2, n_1, n_2), i \in \{1, 2\}, \text{ with } k_1 = k_2 = 0;$
- (E3) $\mathcal{T}_2(k, m, n; k_1, k_2, n_1, n_2)$ with $k_1 = 0$ or $k_2 = 0$;
- (E4) $\mathcal{T}_1(k, m, n; k_1, k_2, n_1, n_2)$ with $k_1 = 0, n_2 \ge 2$ or $k_2 = 0, n_1 \ge 2$;
- (E5) $N^{-1}\mathcal{T}_i(k,m,n;k_1,k_2,n_1,n_2), i \in \{1,2\}, \text{ with } n_1,n_2 \ge 1.$

Now, the estimate (8.5) in the Gaussian case follows immediately from the next claim.

Claim 8.3. For any constant $C_0 > 0$, we can expand Y_1 in (8.8) as the expectation of a linear combination of O(1) many expressions satisfying (E1)–(E5) and some error expressions of order $O_{\prec}(N^{-C_0})$.

Proof. In the following proof, corresponding to (8.32)–(8.35), we define the *sizes* of the expressions on the LHS by their RHS. For example, if $k_1, k_2 \ge 1$, we define

$$\operatorname{size}\left[\mathcal{T}_{i}(k,m,n;k_{1},k_{2},n_{1},n_{2})\right] := \left(\frac{1}{N\eta}\right)^{k+m-n+k_{1}+k_{2}-2} \cdot (N^{-1/2})^{\mathbf{1}(n_{1}\geq2)+\mathbf{1}(n_{2}\geq2)} \|A\|_{\operatorname{HS}}^{n_{1}+n_{2}}.$$

If $k_1 = 0$ and $k_2 \ge 1$ (or $k_2 = 0$ and $k_1 \ge 1$), we define the sizes of $\mathcal{T}_i(k, m, n; k_1, k_2, n_1, n_2)$, $i \in \{1, 2\}$, through (8.33) and (8.34); if $k_1 = k_2 = 0$, we define the sizes through (8.35). If we have an expression, say $a_N \mathcal{T} + b_N \mathcal{T}'$, for some deterministic coefficients a_N, b_N and type I or type II expressions $\mathcal{T}, \mathcal{T}'$, then

$$\mathsf{size}(a_N\mathcal{T} + b_N\mathcal{T}') := a_N\mathsf{size}(\mathcal{T}) + b_N\mathsf{size}(\mathcal{T}')$$

We set the stopping rule of our expansions as follows: we stop expanding an expression $\mathbb{E}\mathcal{T}$ if it satisfies (E1)-(E5) or size $(\mathcal{T}) \prec N^{-C_0}$.

Given Y_1 in (8.8), we first expand M_1 using the Taylor expansion (8.6):

$$\widetilde{M}_1 = -\sum_{k=1}^{n_0} (m_1 + z_1)^{-(k+1)} \Lambda^k + (m_1 + z_1)^{-(n_0+1)} \Lambda^{n_0+1} M_1,$$
(8.36)

where n_0 is chosen sufficiently large such that $||A||^{n_0} \leq N^{-C_0-10}$. In every resulting expression, if it contains the error term $(m_1 + z_1)^{-(n_0+1)} \Lambda^{n_0+1} M_1$, then we can easily check that it is of order $O_{\prec}(N^{-C_0})$; otherwise, between each product of two matrices, we decompose the identity matrix as $I = \sum_{a=1}^{D} E_a$, i.e., for every matrix product B_1B_2 , we write it as $\sum_a B_1E_aB_2$. In this way, we can expand (8.8) into the expectation of a linear combination of O(1) many expressions in $\mathcal{T}_1(k = 0, m = 0, n = 0; k_1 = 2, k_2 = 1, n_1 = 1, n_2)$ and $\mathcal{T}_2(k = 1, m = 0, n = 0; k_1 = 1, k_2 = 1, n_1 = 1, n_2)$ with $n_2 \geq 1$ plus error expressions of size $O_{\prec}(N^{-C_0})$.

Now, given any type I or type II expression that is not an error of size $O_{\prec}(N^{-C_0})$ and does not satisfy (E1)–(E4), say $\mathcal{T}(k, m, n; k_1, k_2, n_1, n_2)$ with $n_1, n_2 \geq 1$, we expand it according to the following strategy.

Step 1: In this expression, we have either $n_1 = 1$ or $n_2 = 1$ since it does not satisfy (E1). Without loss of generality, suppose $n_1 = 1$. Then, we must have $k_1 \ge 1$. Otherwise, if $k_1 = 0$ and \mathcal{T} is a type I expression, then we trivially have $\langle \Lambda^{(1)} \rangle = 0$; if $k_1 = 0$ and \mathcal{T} is a type II expression, then \mathcal{T} satisfies (E3). Now, we pick the first G to the right of this $\Lambda^{(1)}$, say G_s with $s \in \{1, 2\}$, i.e., there is a factor $G_s E_{a_0} \Lambda E_{a_1}$ in the loop containing $\Lambda^{(1)} = E_{a_0} \Lambda E_{a_1}$, $a_0, a_1 \in [\![D]\!]$. We then expand G_s with (4.23), i.e., $G_s = M_s - G_s(H + m_s)M_s$.

Step 2: In a new expression from step 1 that does not satisfy the stopping rule, suppose we have replaced this G_s with M_s . By expanding \widetilde{M}_s with (8.36) and adding $I = \sum_a E_a$ between matrix products, we get some error expressions of size $O_{\prec}(N^{-C_0})$ plus expressions of the form $\mathcal{T}' = (N^{-1})^{\mathbf{1}(k_1 \geq 2)} \cdot \mathcal{T}_i(k, m, n; k_1 - 1, k_2, n'_1, n_2), i \in \{1, 2\}$, with $n'_1 \geq n_1 = 1$. Then:

- If $k_1 \geq 2$, then \mathcal{T}' satisfies (E5).
- If $k_1 = 1$ and i = 2, then \mathcal{T}' satisfies (E3).
- If $k_1 = 1$, i = 1, and $n_2 \ge 2$, then \mathcal{T}' satisfies (E4).
- If $k_1 = 1$, i = 1, $n_2 = 1$, and $k_2 = 0$, then \mathcal{T}' satisfies (E2).

In the remaining cases with i = 1, $k_1 = 1$, $k_2 \ge 1$, $n'_1 \ge n_1 = 1$, and $n_2 = 1$, \mathcal{T}' is a type I expression $\mathcal{T}_1(k, m, n; 0, k_2, n'_1, 1)$ with $\mathsf{size}(\mathcal{T}') \le N^{-1/2}\mathsf{size}(\mathcal{T})$. We send this \mathcal{T}' back to Step 1.

Step 3: In a new expression from step 1 that does not satisfy the stopping rule, suppose we have replaced the G_s with $-G_s(H+m_s)M_s$. Again, by expanding \widetilde{M}_s with (8.36) and adding $I = \sum_a E_a$ between matrix products, we get errors of size $O_{\prec}(N^{-C_0})$ plus expressions of the form $\mathbb{E}\mathcal{T}_1$ or $\mathbb{E}\mathcal{T}_2$, where for some $\widetilde{n}_1 \ge n_1$,

$$\mathcal{T}_1 = c_N \mathcal{W}^{(k)} \Gamma_n^{(m)} \langle \mathcal{G}^{(k_1 - 1)} G_s(H + m_s) \Lambda^{(\tilde{n}_1)} \rangle \langle \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle$$

with a deterministic coefficient $c_N = O(N^{-(m-n+k_1+k_2-3)-1(k_1=0)-1(k_2=0)})$ or

$$\mathcal{T}_2 = c_N \mathcal{W}^{(k)} \Gamma_n^{(m)} \langle \mathcal{G}^{(k_1-1)} G_s(H+m_s) \Lambda^{(\tilde{n}_1)} \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle$$

with a deterministic coefficient $c_N = O(N^{-(m-n+k_1+k_2-2)-1(k_1=0)-1(k_2=0)}).$

First, we apply the Gaussian integration by parts to $\mathbb{E}\mathcal{T}_1$:

$$\mathbb{E}\mathcal{T}_{1} = \frac{c_{N}}{DN^{2}} \mathbb{E}\sum_{x=1}^{D} \sum_{\alpha,\beta \in I_{x}} \partial_{\beta\alpha} \left[\left(\Lambda^{(\tilde{n}_{1})} \mathcal{G}^{(k_{1}-1)} G_{s} \right)_{\beta\alpha} \mathcal{W}^{(k)} \Gamma_{n}^{(m)} \langle \mathcal{G}^{(k_{2})} \Lambda^{(n_{2})} \rangle \right]$$

$$+ c_N m_s \mathbb{E} \mathcal{W}^{(k)} \Gamma_n^{(m)} \langle \mathcal{G}^{(k_1 - 1)} G_s \Lambda^{(\tilde{n}_1)} \rangle \langle \mathcal{G}^{(k_2)} \Lambda^{(n_2)} \rangle.$$

$$(8.37)$$

Taking the derivative $\partial_{\beta\alpha}$, we get the following cases.

- If $\partial_{\beta\alpha}$ acts on G_s , then the resulting expression together with the second term on the RHS of (8.37) gives a sum of expressions in $\mathcal{T}_1(k+1, m, n; k_1, k_2, \tilde{n}_1, n_2)$. These new expressions have sizes $\leq (N\eta)^{-1} \operatorname{size}(\mathcal{T}_1)$.
- If $\partial_{\beta\alpha}$ acts on a light weight in $\mathcal{W}^{(k)}$, then we get a sum of expressions in $\mathcal{T}_1(k-1, m, n; k_1 + 2, k_2, \tilde{n}_1, n_2)$. These new expressions have sizes $\leq (N\eta)^{-1} \operatorname{size}(\mathcal{T}_1)$.
- If $\partial_{\beta\alpha}$ acts on a *G*-loop in $\Gamma_n^{(m)}$, then we get a sum of expressions in $\mathcal{T}_1(k, m k', n 1; k_1 + k' + 1, k_2, \tilde{n}_1, n_2)$ for some $2 \leq k' \leq m$. These new expressions have sizes $\leq (N\eta)^{-2} \operatorname{size}(\mathcal{T}_1)$.
- If $\partial_{\beta\alpha}$ acts on $\mathcal{G}^{(k_1-1)}$, then we create a new *G*-loop and get a sum of expressions in $\mathcal{T}_1(k, m+k', n+1; k_1 k' + 1, k_2, \tilde{n}_1, n_2)$ for some $2 \leq k' \leq k_1$. These new expressions have sizes $\leq \mathsf{size}(\mathcal{T}_1)$, but the value of k_1 is decreased at least by 1.
- If $\partial_{\beta\alpha}$ acts on $\mathcal{G}^{(k_2)}$, then we create a (G, Λ, G, Λ) -loop and get a sum of expressions in $\mathcal{T}_2(k, m, n; k'_1, k_1 + k_2 k'_1 + 1, \tilde{n}_1, n_2)$ for some $1 \leq k'_1 \leq k_2$. These new expressions have sizes $\leq (N\eta)^{-1}$ size (\mathcal{T}_1) .

Second, we apply the Gaussian integration by parts to $\mathbb{E}\mathcal{T}_2$:

$$\mathbb{E}\mathcal{T}_{2} = \frac{c_{N}}{DN^{2}} \mathbb{E}\sum_{x=1}^{D} \sum_{\alpha,\beta \in I_{x}} \partial_{\beta\alpha} \left[\left(\Lambda^{(\tilde{n}_{1})} \mathcal{G}^{(k_{2})} \Lambda^{(n_{2})} \mathcal{G}^{(k_{1}-1)} G_{s} \right)_{\beta\alpha} \mathcal{W}^{(k)} \Gamma_{n}^{(m)} \right] \\ + c_{N} m_{s} \mathbb{E}\mathcal{W}^{(k)} \Gamma_{n}^{(m)} \langle \mathcal{G}^{(k_{1}-1)} G_{s} \Lambda^{(\tilde{n}_{1})} \mathcal{G}^{(k_{2})} \Lambda^{(n_{2})} \rangle.$$

$$(8.38)$$

Taking the derivative $\partial_{\beta\alpha}$, we get the following cases.

- If $\partial_{\beta\alpha}$ acts on G_s , then the resulting expression together with the second term on the RHS of (8.38) gives a sum of expressions in $\mathcal{T}_2(k+1, m, n; k_1, k_2, \tilde{n}_1, n_2)$. These new expressions have sizes $\leq (N\eta)^{-1} \operatorname{size}(\mathcal{T}_2)$.
- If $\partial_{\beta\alpha}$ acts on a light weight in $\mathcal{W}^{(k)}$, then we get a sum of expressions in $\mathcal{T}_2(k-1, m, n; k_1 + 2, k_2, \tilde{n}_1, n_2)$. These new expressions have sizes $\leq (N\eta)^{-1} \operatorname{size}(\mathcal{T}_2)$.
- If $\partial_{\beta\alpha}$ acts on a *G*-loop in $\Gamma_n^{(m)}$, then we get a sum of expressions in $\mathcal{T}_2(k, m k', n 1; k_1 + k' + 1, k_2, \tilde{n}_1, n_2)$ for some $2 \leq k' \leq m$. These new expressions have sizes $\leq (N\eta)^{-2} \operatorname{size}(\mathcal{T}_2)$.
- If $\partial_{\beta\alpha}$ acts on $\mathcal{G}^{(k_1-1)}$, then we create a new *G*-loop and get a sum of expressions in $\mathcal{T}_2(k, m+k', n+1; k_1 k' + 1, k_2, \tilde{n}_1, n_2)$ for some $2 \leq k' \leq k_1$. These new expressions have sizes $\leq \text{size}(\mathcal{T}_2)$, but the value of k_1 is decreased at least by 1.
- If $\partial_{\beta\alpha}$ acts on $\mathcal{G}^{(k_2)}$, then we create two (G, Λ) -loops and get a sum of expressions in $\mathcal{T}_1(k, m, n; k'_1, k_1 + k_2 k'_1 + 1, \tilde{n}_1, n_2)$ for some $1 \le k'_1 \le k_2$. These new expressions have sizes $\lesssim (N\eta)^{-1}$ size (\mathcal{T}_2) .

For every new expression from Step 3 that does not satisfy the stopping rule, we send it back to Step 1.

Now, we iterate the above expansion strategy, Steps 1–3, until all resulting expressions satisfy the stopping rule. It then suffices to show that this expansion process will finally stop after O(1) many iterations. In fact, we see that after one iteration of Steps 1–3, for each new expression, either it has a strictly smaller size than the input graph by a factor $(N\eta)^{-1} \leq N^{-c}$, or the value of $k_1 + k_2$ decreases at least by 1. With this observation, we can show that all expressions will satisfy the stopping rule after at most $(\lceil C_0/c \rceil + 10)^2$ many iterations. This concludes the proof of Lemma 8.2 in the Gaussian case.

8.2. **Proof of Lemma 8.2: non-Gaussian case.** In the non-Gaussian case, we apply the cumulant expansion, Lemma 4.10, to (8.7) and get some new terms containing higher-order cumulants of the H entries plus an error term \mathcal{R}_{l+1} . Again, this error term is negligible if we take l sufficiently large. We now need to estimate Y_1 in (8.8) and the following terms:

$$Y_{n,m} := -\frac{1}{DN} \sum_{\alpha,\beta \in \mathcal{I}} \frac{1}{n!m!} \mathcal{C}^{(n,m+1)}_{\beta\alpha} \mathbb{E} \partial^m_{\alpha\beta} \partial^n_{\beta\alpha} (M_1 \Lambda G_2 \Lambda G_1)_{\beta\alpha}, \quad 2 \le m+n \le l.$$
(8.39)

Taking the derivatives, we can write (8.39) into a linear combination of O(1) many terms of the following forms for some $s, t \in \mathbb{N}$ with s + t = n + m and $(x(i), y(i)), (\tilde{x}(j), \tilde{y}(j)) \in \{(\alpha, \beta), (\beta, \alpha)\}, i \in [\![s]\!], j \in [\![t]\!]$:

$$\frac{1}{DN} \sum_{\alpha,\beta \in \mathcal{I}} \mathcal{C}^{(n,m+1)}_{\beta\alpha} \mathbb{E}(M_1 \Lambda G_2)_{\beta x(1)} (G_2)_{y(1)x(2)} \cdots (G_2 \Lambda G_1)_{y(s)\tilde{x}(1)} (G_1)_{\tilde{y}(1)\tilde{x}(2)} \cdots (G_1)_{\tilde{y}(t)\alpha}, \quad s \ge 1; \quad (8.40)$$

$$\frac{1}{DN} \sum_{\alpha,\beta \in \mathcal{I}} \mathcal{C}^{(n,m+1)}_{\beta\alpha} \mathbb{E}(M_1 \Lambda G_2 \Lambda G_1)_{\beta \widetilde{x}(1)} (G_1)_{\widetilde{y}(1)\widetilde{x}(2)} \cdots (G_1)_{\widetilde{y}(t)\alpha}, \quad s = 0.$$
(8.41)

By the anisotropic local law (2.20), we have that

 $(M_1\Lambda G_2)_{\beta x(1)} \prec \|\mathbf{e}_{\beta}^*M_1\Lambda\|, \quad (G_2\Lambda G_1)_{y(s)\widetilde{x}(1)} \prec \sqrt{N}\|A\|_{\mathrm{HS}}, \quad (M_1\Lambda G_2\Lambda G_1)_{\beta\widetilde{x}(1)} \prec \|\mathbf{e}_{\beta}^*M_1\Lambda\| \cdot \sqrt{N}\|A\|_{\mathrm{HS}},$ where \mathbf{e}_{β} denotes the standard unit vector along the β -th direction. So, we can bound (8.40) and (8.41) by

$$|(8.40)| + |(8.41)| \prec \frac{N^{3/2} \|A\|_{\mathrm{HS}}}{N^{(n+m+1)/2+1}} \sum_{\beta} \|\mathbf{e}_{\beta}^* M_1 \Lambda\| \prec N^{-(m+n-3)/2} \cdot N^{-1} \|A\|_{\mathrm{HS}}^2$$

which is sufficiently small if $m + n \ge 3$.

When m + n = 2, the above bound has an extra $N^{1/2}$ factor, so we need to further expand the terms in $Y_{n,m}$. Take two terms in $Y_{1,1}$ as examples:

$$-\frac{1}{DN}\sum_{x=1}^{D}\sum_{\alpha,\beta\in\mathcal{I}_{x}}\mathcal{C}_{\beta\alpha}^{(1,2)}\mathbb{E}\left[(M_{1}\Lambda G_{2}\Lambda G_{1})_{\beta\beta}(G_{1})_{\alpha\alpha}+(M_{1}\Lambda G_{2})_{\beta\beta}(G_{2}\Lambda G_{1})_{\alpha\alpha}\right](G_{1})_{\beta\alpha}.$$

In the above expression, we can expand either G_2 or G_1 in the factor $G_2\Lambda G_1$ with (4.23). If we replace G_1 with M_1 , by the anisotropic local law (2.20), the resulting expression can be bounded by

$$\frac{1}{N^{5/2}} \sum_{x=1}^{D} \sum_{\alpha,\beta \in \mathcal{I}_x} \left[\|\mathbf{e}_{\beta}^* M_1 \Lambda\| \|\Lambda M_1 \mathbf{e}_{\beta}\| + \|\mathbf{e}_{\beta}^* M_1 \Lambda\| \|\Lambda M_1 \mathbf{e}_{\alpha}\| \right] \lesssim N^{-3/2} \|A\|_{\mathrm{HS}}^2$$

(Note that if we have replaced G_2 with M_2 instead, the resulting expression can be bounded by $N^{-1}||A||_{\text{HS}}^2$, which is still good enough.) Now, for the term with G_1 replaced by $-M_1(m_1 + H)G_1$, applying cumulant expansions we get a sum of small enough errors, Gaussian integration by parts terms, and terms involving third or higher-order cumulants: for some $2 \le m + n \le l$,

$$Y_{1,1;n,m} = \frac{1}{DN} \sum_{\alpha,\beta\in\mathcal{I}} \mathcal{C}_{\beta\alpha}^{(1,2)} \sum_{i,j\in\mathcal{I}} \mathcal{C}_{ji}^{(n,m+1)} \mathbb{E}\partial_{ij}^{m} \partial_{ji}^{n} \left[(M_{1}\Lambda G_{2}\Lambda M_{1})_{\beta i} (G_{1})_{j\beta} (G_{1})_{\alpha\alpha} (G_{1})_{\beta\alpha} \right] \\ + \frac{1}{DN} \sum_{\alpha,\beta\in\mathcal{I}} \mathcal{C}_{\beta\alpha}^{(1,2)} \sum_{i,j\in\mathcal{I}} \mathcal{C}_{ji}^{(n,m+1)} \mathbb{E}\partial_{ij}^{m} \partial_{ji}^{n} \left[(M_{1}\Lambda G_{2})_{\beta\beta} (G_{2}\Lambda M_{1})_{\alpha i} (G_{1})_{j\alpha} (G_{1})_{\beta\alpha} \right].$$

With the anisotropic local law (2.20), we can bound this term by

$$Y_{1,1;n,m} \prec \frac{1}{N^{3+\frac{n+m}{2}}} \sum_{\alpha,\beta,i,j} \|\mathbf{e}_{\beta}^* M_1 \Lambda\| \left(\|\Lambda M_1 \mathbf{e}_i\| + \|\Lambda M_1 \mathbf{e}_j\| \right) \lesssim N^{-\frac{n+m}{2}} \|A\|_{\mathrm{HS}}^2 \leq N^{-1} \|A\|_{\mathrm{HS}}^2.$$

Hence, we only need to further expand the Gaussian integration by parts terms.

We can estimate all the terms $Y_{n,m}$ with n + m = 2 using a similar argument as above. In this way, we obtain that

$$\mathbb{E}\langle (G_1 - M_1)\Lambda G_2\Lambda \rangle = Y_1 + \sum_{m+n=2} Y_{n,m;1} + \mathcal{O}_{\prec} (N^{-1} ||A||_{\mathrm{HS}}^2),$$

where Y_1 is given in (8.8) and $Y_{n,m;1}$ denotes the Gaussian integration by parts terms obtained from the expansions of $Y_{n,m}$. To bound the RHS, we still need to further expand Y_1 and $Y_{n,m;1}$. Since the proof closely resembles that of Claim 8.3, we will provide only a concise overview of the proof by outlining the main expansion strategy, while omitting some of the finer details.

In the expansions of $Y_{n,m;1}$, if an expression involves third or higher-order cumulants of the H entries, then it can be bounded by $O_{\prec}(N^{-1}||A||_{\text{HS}}^2)$. Hence, we only need to consider expressions obtained from repeated applications of (4.23) and Gaussian integration by parts. In this process, we either get a sufficiently small error of size $O_{\prec}(N^{-C_0})$ or expressions consisting of light weights, G-loops, at most one (G, Λ) -loop $\langle \mathcal{G}^{(k)}\Lambda^{(n)} \rangle$ for some $k \geq 0$ and $n \geq 1$, and terms of the form

$$(\Lambda^{(n_1)}\mathcal{G}^{(k_1)}\Lambda^{(n_1')})_{\beta\mathfrak{a}}(\Lambda^{(n_2')}\mathcal{G}^{(k_2)}\Lambda^{(n_2)}\mathcal{G}^{(k_2')}\Lambda^{(n_2'')})_{\mathfrak{bc}}, \quad \text{or} \quad (\Lambda^{(n_1)}\mathcal{G}^{(k_1)}\Lambda^{(n_2)}\mathcal{G}^{(k_2)}\Lambda^{(n_2')})_{\beta\mathfrak{a}}$$

for some $n_1, n_2 \ge 1, k_1, k_2, n'_1, n'_2, n''_2 \ge 0$, and $\mathfrak{a}, \mathfrak{b}, \mathfrak{c} \in \{\alpha, \beta\}$. One can check that if (i) there is a (G, Λ) -loop, (ii) at least one of n'_1, n'_2, n''_2 is non-zero, (iii) at least one of n_1, n_2 is at least 2, or (iv) $k_2 = 0$ or $k'_2 = 0$,

then the expressions is bounded by $O_{\prec}(N^{-1}||A||_{\text{HS}}^2)$. Hence, we only need to handle expressions containing the following factors:

$$(\Lambda \mathcal{G}^{(k_1)})_{\beta \mathfrak{a}} (\mathcal{G}^{(k_2)} \Lambda \mathcal{G}^{(k_2)})_{\mathfrak{b}\mathfrak{c}}, \quad \text{or} \quad (\Lambda \mathcal{G}^{(k_1)} \Lambda \mathcal{G}^{(k_2)})_{\beta \mathfrak{a}}, \quad k_1 \ge 0, \ k_2, k_2' \ge 1.$$

$$(8.42)$$

Then, similar to the expansion strategy (Steps 1–3) in the proof of Claim 8.3, we choose the G in $\mathcal{G}^{(k_2)}$ right next to Λ , use (4.23) to expand it, and then apply Gaussian integration by parts. (We will also need to use (8.36) and add $I = \sum_a E_a$ between matrix products in this process.) For each new expression, at least one of the following cases holds: (1) it is a small error $O_{\prec}(N^{-C_0})$; (2) it can be bounded by $O_{\prec}(N^{-1}||A||^2_{\text{HS}})$; (3) its size is smaller than the original expression by $O((N\eta)^{-1})$; (4) the value of $k_2 + k'_2$ is decreased at least by 1 (we let $k'_2 = 0$ for $(\Lambda \mathcal{G}^{(k_1)} \Lambda \mathcal{G}^{(k_2)})_{\beta \mathfrak{a}}$ as a convention). We can show that after at most $(\lceil C_0/c \rceil + 10)^2$ many iterations of this expansion strategy, all the resulting expressions is bounded by $O_{\prec}(N^{-1}||A||^2_{\text{HS}} + N^{-C_0})$.

Now, we look at the expansions of Y_1 . If an expression involves (1) a fourth or higher-order cumulant or (2) at least two third-order cumulants of the H entries, then it can be bounded by $O_{\prec}(N^{-1}||A||_{HS}^2)$. If we only apply Gaussian integration by parts in the whole expansion process, then the relevant expressions have been handled in Section 8.1. Finally, we need to address the scenario in which we employ the third-order cumulant expansion only once, while utilizing Gaussian integration for the remaining expansions. Before the third-order cumulant expansion, we follow the expansion strategy in the proof Claim 8.3, while after that, we follow the strategy below (8.42). Here we omit the relevant details. This concludes the proof Lemma 8.2, which further concludes Theorem 2.5.

APPENDIX A. SOME DETERMINISTIC ESTIMATES

In this section, we provide some deterministic estimates about M and \widehat{M} that have been used in the main proofs.

Lemma A.1 (Basic properties of M and \widehat{M}). Let A be an arbitrary deterministic matrix with ||A|| = o(1). For any constant $\tau > 0$, the following estimates hold uniformly for all $z = E + i\eta$ with $|z| \le \tau^{-1}$ and $\eta > 0$.

• We have that

$$|m(z) - m_{sc}(z)| \lesssim ||A||^{1/2}, ||M(z) - m(z)I|| \lesssim ||A||^{1/2}.$$
 (A.1)

Furthermore, when $|E| \leq 2 - \kappa$ for a constant $\kappa > 0$, we have the better bounds

$$|m(z) - m_{sc}(z)| \lesssim ||A||, ||M(z) - m(z)I|| \lesssim ||A||.$$
 (A.2)

As a consequence of (A.2), we have

$$|(m(z)+z)^{-1}| \lesssim 1$$
, and $|(1-m(z)^2)^{-1}| \lesssim 1$ if $|E| \le 2-\kappa$. (A.3)

- \widehat{M} is translation invariant, i.e., $\widehat{M}_{ab} = \widehat{M}_{a'b'}$ whenever $a b = a' b' \mod D$.
- For $z_1 = \overline{z}_2 \in \{z, \overline{z}\}$ and $a \in \llbracket D \rrbracket$, we have that

$$\sum_{b=1}^{D} \widehat{M}(z_1, z_2)_{ab} = \frac{\operatorname{Im} m(z)}{\operatorname{Im} m(z) + \eta} =: d_1,$$
(A.4)

where d_1 denotes the Perron-Frobenius eigenvalue of $\widehat{M}(z_1, z_2)$ with $(1, \ldots, 1)^{\top}$ being the corresponding eigenvector. The other eigenvalues of $\widehat{M}(z_1, z_2)$ satisfy

$$d_k = d_1 - a_k - ib_k, \quad k = 2, 3, \dots, D,$$
 (A.5)

where $a_k, b_k \in \mathbb{R}$ satisfy that

$$a_k \ge 0, \quad a_k \sim N^{-1} ||A||_{\text{HS}}^2, \quad |b_k| = o(N^{-1} ||A||_{\text{HS}}^2).$$
 (A.6)

• For $z_1 = \overline{z}_2 \in \{z, \overline{z}\}$, we have that

$$\left\| [1 - \widehat{M}(z_1, z_2)]^{-1} \right\| = \frac{\operatorname{Im} m(z) + \eta}{\eta} \lesssim \eta^{-1}.$$
(A.7)

• When $|E| \leq 2 - \kappa$ for a constant $\kappa > 0$, there exists a constant $C_{\kappa} > 0$ such that

$$\left\| [1 - \widehat{M}(z, z)]^{-1} \right\| \le C_{\kappa}.$$
(A.8)

Moreover, for $z_1, z_2 \in \{z, \overline{z}\}$, we have that

$$\max_{a,b,a',b' \in \llbracket D \rrbracket} \left| \left[(1 - \widehat{M}_{(1,2)})^{-1} \widehat{M}_{(1,2)} \right]_{ab} - \left[(1 - \widehat{M}_{(1,2)})^{-1} \widehat{M}_{(1,2)} \right]_{a'b'} \right| \lesssim \frac{N}{\|A\|_{\mathrm{HS}}^2}.$$
 (A.9)

Proof. The first estimates in (A.1) and (A.2) follow by comparing the self-consistent equation (2.15) with the equation $m_{sc}(z) = -[z + m_{sc}(z)]^{-1}$ and by utilizing the stability of the equation. The second estimates in (A.1) and (A.2) then follow from the Taylor expansion

$$M = -\sum_{k=0}^{\infty} (m+z)^{-k-1} \Lambda^k.$$
 (A.10)

The estimate (A.3) holds due to the simple fact that $m_{sc}(z)$ satisfies the corresponding bounds.

The translation invariance of M follows from the block translation invariance of M. As a consequence, its eigenvectors are given by $\mathbf{u}_k, k \in [\![D]\!]$, with

$$u_k(a) = D^{-1/2} \exp(i \cdot 2\pi (k-1)(a-1)/D).$$
(A.11)

The corresponding eigenvalues are

$$d_k = \sum_{a=1}^{D} \widehat{M}_{1a}(z_1, z_2) e^{i2\pi(k-1)(a-1)/D}.$$
(A.12)

In particular, when $z_1 = \bar{z}_2 \in \{z, \bar{z}\}$, d_1 is the Perron–Frobenius eigenvalue of $\widehat{M}(z_1, z_2)$ and the identity (A.4) follows by applying Ward's identity (C.11) below to M(z) and utilizing the identity

$$\sum_{b=1}^{D}\widehat{M}(z_1, z_2)_{ab} = \frac{1}{D}\sum_{a, b=1}^{D}\widehat{M}(z_1, z_2)_{ab} = \frac{1}{DN}\sum_{i, j \in \mathcal{I}} |M_{ij}(z)|^2.$$

With the Taylor expansion (A.10), we can derive that when $z_1 = \overline{z}_2 \in \{z, \overline{z}\}$,

$$\widehat{M}(z_1, z_2)_{11} = \frac{1}{|m+z|^2} + \frac{2(1+\mathbf{1}_{D>2})}{|m+z|^2} \operatorname{Re}\left[(m+z)^{-2}\right] \cdot N^{-1} \|A\|_{\operatorname{HS}}^2 + o(N^{-1}\|A\|_{\operatorname{HS}}^2),
\widehat{M}(z_1, z_2)_{12} = |m+z|^{-4}N^{-1}\|A\|_{\operatorname{HS}}^2 + o(N^{-1}\|A\|_{\operatorname{HS}}^2),
\widehat{M}(z_1, z_2)_{1D} = |m+z|^{-4}N^{-1}\|A\|_{\operatorname{HS}}^2 + o(N^{-1}\|A\|_{\operatorname{HS}}^2),
\widehat{M}(z_1, z_2)_{1k} = o(N^{-1}\|A\|_{\operatorname{HS}}^2), \quad 3 \le k \le D - 1.$$
(A.13)

Plugging these estimates into (A.12), we immediately obtain (A.5) and (A.6). Finally, using the eigendecomposition of $\widehat{M}(z_1, z_2)$, we can derive (A.7) from with (A.5).

When $|E| \leq 2 - \kappa$, we have

$$|1 - m_{sc}(z)^2| \ge c_\kappa \tag{A.14}$$

for a constant $c_{\kappa} > 0$. Combining this estimate with (A.1), we can obtain (A.8). Now, with the eigendecomposition of $\widehat{M}(z_1, z_2)$, we get

$$\left[(1 - \widehat{M}_{(1,2)})^{-1} \widehat{M}_{(1,2)} \right]_{ab} = \frac{1}{D} \sum_{k=2}^{D} \frac{d_k}{1 - d_k} e^{\mathbf{i} \cdot 2\pi (k-1)(a-b)/D} + \frac{1}{D} \frac{d_1}{1 - d_1},$$

which gives that

$$\left| \left[(1 - \widehat{M}_{(1,2)})^{-1} \widehat{M}_{(1,2)} \right]_{ab} - \frac{1}{D} \frac{d_1}{1 - d_1} \right| \lesssim \max_{k=2}^{D} \left| (1 - d_k)^{-1} \right|.$$
(A.15)

If $z_1 = z_2$, then combining (A.14) with (A.1), we obtain that $|1 - d_k| \gtrsim 1$, which gives (A.9). If $z_1 = \bar{z}_2$, applying (A.5) and (A.6), we obtain that $|1 - d_k| \gtrsim N^{-1} ||A||_{\text{HS}}^2$ for $k \in [\![2, D]\!]$. Together with (A.15), it concludes (A.9).

APPENDIX B. GREEN'S FUNCTION COMPARISON

In this section, we complete the proof of Lemma 3.4 using a standard Green's function comparison argument. We adopt the continuous comparison method introduced in [78], which is based on the following interpolation.

Definition B.1 (Interpolating matrices). Introduce the notations $H^0 := \tilde{H}$ and $H^1 := H$. Let ρ_{ij}^0 and ρ_{ij}^1 be the laws of \tilde{H}_{ij} and H_{ij} , respectively. For $\theta \in [0, 1]$, we define the interpolated laws $\rho_{ij}^{\theta} := (1 - \theta)\rho_{ij}^0 + \theta\rho_{ij}^1$. (Note that $\rho_{ij}^0 = \rho_{ij}^1 = \delta_0$ if i and j are not in the same \mathcal{I}_a .) Let $\{H^{\theta} : \theta \in (0, 1)\}$ be a collection of random matrices such that the followings hold. For any fixed $\theta \in (0, 1)$, (H^0, H^{θ}, H^1) is a triple of independent $DN \times DN$ random matrices, and the matrix $H^{\theta} = (H_{ij}^{\theta})$ has law

$$\prod_{i \le j \in \mathcal{I}} \rho_{ij}^{\theta} (\mathrm{d}H_{ij}^{\theta}). \tag{B.1}$$

(Note that we do not require H^{θ_1} to be independent of H^{θ_2} for $\theta_1 \neq \theta_2 \in (0,1)$.) For $\lambda \in \mathbb{C}$, $i, j \in \mathcal{I}$, we define the matrix $H^{\theta,\lambda}_{(ij)}$ as

$$\left(H_{(ij)}^{\theta,\lambda} \right)_{kl} := \begin{cases} H_{ij}^{\theta}, & \text{if } \{k,l\} \neq \{i,j\}, \\ \lambda, & \text{if } (k,l) = (i,j), \\ \bar{\lambda}, & \text{if } (k,l) = (j,i). \end{cases}$$
 (B.2)

Correspondingly, we define the resolvents

$$G^{\theta}(z) := G\left(z, H^{\theta}, \Lambda\right), \quad G^{\theta, \lambda}_{(ij)}(z) := G\left(z, H^{\theta, \lambda}_{(ij)}, \Lambda\right)$$

With (B.1), taking the derivative with respect to θ , we get the following interpolation formula: for any differentiable function $F : \mathbb{R}^{DN \times DN} \to \mathbb{C}$,

$$\frac{\mathrm{d}}{\mathrm{d}\theta}\mathbb{E}F(H^{\theta}) = \sum_{i\leq j\in\mathcal{I}} \left[\mathbb{E}F\left(H_{(ij)}^{\theta,H_{ij}^{1}}\right) - \mathbb{E}F\left(H_{(ij)}^{\theta,H_{ij}^{0}}\right)\right],\tag{B.3}$$

provided all the expectations exist. Then, Lemma 3.4 follows from the next estimate on the RHS of (B.3) with $F(H^{\theta}) = \langle G_1^{\theta} E_a G_2^{\theta} E_b \rangle$.

Lemma B.2. Under the assumptions of Lemma 3.4, we have

$$\sum_{i \le j \in \mathcal{I}} \left[\mathbb{E}F\left(H_{(ij)}^{\theta, H_{ij}^1}\right) - \mathbb{E}F\left(H_{(ij)}^{\theta, H_{ij}^0}\right) \right] \prec N^{-1-\delta} \eta^{-2}$$
(B.4)

for all $\theta \in [0,1]$, where $F(H^{\theta}) := \langle G_1^{\theta} E_a G_2^{\theta} E_b \rangle$.

Proof. The proof of (B.4) uses the moment matching conditions (3.11) and (3.12) and the following resolvent expansion: for any $\lambda, \lambda' \in \mathbb{C}$ and $K \in \mathbb{N}$,

$$G_{(ij)}^{\theta,\lambda'} = G_{(ij)}^{\theta,\lambda} + \sum_{k=1}^{K} G_{(ij)}^{\theta,\lambda} \left\{ \left[\operatorname{Re}(\lambda - \lambda')\Delta_{ij} + i\operatorname{Im}(\lambda - \lambda')\widetilde{\Delta}_{ij} \right] G_{(ij)}^{\theta,\lambda} \right\}^{k} + G_{(ij)}^{\theta,\lambda'} \left\{ \left[\operatorname{Re}(\lambda - \lambda')\Delta_{ij} + i\operatorname{Im}(\lambda - \lambda')\widetilde{\Delta}_{ij} \right] G_{(ij)}^{\theta,\lambda} \right\}^{K+1},$$
(B.5)

where the matrices Δ_{ij} and $\widetilde{\Delta}_{ji}$ are $DN \times DN$ matrices defined as $(\Delta_{ij})_{kl} := \delta_{ki}\delta_{lj} + \delta_{kj}\delta_{li}$ and $(\widetilde{\Delta}_{ij})_{kl} := \delta_{ki}\delta_{lj} - \delta_{kj}\delta_{li}$. Since H^{θ} and $G^{\theta,H^a_{ij}}$, $a \in \{0,1\}$, also satisfy the setting of Lemma 2.10, we have that for each $\theta \in [0,1]$, G^{θ} and $G^{\theta,H^a_{ij}}$ satisfy the local laws (2.20) and (2.23). Furthermore, let ξ be an arbitrary random variable satisfying $|\xi| \prec N^{-1/2}$. With the expansion (B.5), we can readily show that the local laws (2.20) and (2.23) hold also for the resolvent $G^{\theta,\xi}_{(ij)}$. We will use these estimates tacitly in the following proof.

By changing from $H_{ij}^0 := X_{ij}^0 + iY_{ij}^{(0)}$ to $H_{ij}^1 := X_{ij}^1 + iY_{ij}^1$, we can break down the replacement into two distinct steps. We first modify the real part X_{ij}^0 to X_{ij}^1 , followed by replacing the imaginary part Y_{ij}^0 with Y_{ij}^1 . Hence, without loss of generality, we assume for the rest of the proof that H_{ij}^0 and H_{ij}^1 have the same

imaginary part, i.e., $Y_{ij}^0 = Y_{ij}^1$, while the case of $X_{ij}^0 = X_{ij}^1$ can be handled in the same way. Moreover, with a slight abuse of notation, we denote

$$H_{(ij)}^{\theta, X_{ij}^{\gamma}} := H_{(ij)}^{\theta, H_{ij}^{\gamma}}, \quad G_{(ij)}^{\theta, X_{ij}^{\gamma}} := G_{(ij)}^{\theta, H_{ij}^{\gamma}}, \quad \gamma \in \{0, 1\}.$$

Using (B.5) with $\lambda = 0$ and K = 7 and applying the local law (2.20), we get that

$$\left[(G_s)_{(ij)}^{\theta, X_{ij}^{\gamma}} \right]_{xy} = \left[(G_s)_{(ij)}^{\theta, 0} \right]_{xy} + \sum_{k=1}^{\ell} \mathcal{X}_{xy}^{(ij)}(s, \gamma, k) + \mathcal{O}_{\prec}(N^{-4}), \quad x, y \in \mathcal{I}, \ s \in \{1, 2\}, \ \gamma \in \{0, 1\},$$
(B.6)

where the random matrices $\mathcal{X}^{(ij)}(s,\gamma,k)$ are defined by

$$\mathcal{X}^{(ij)}(s,\gamma,k) := (-X_{ij}^{\gamma})^k (G_s)_{(ij)}^{\theta,0} \left[\Delta_{ij} (G_s)_{(ij)}^{\theta,0} \right]^k, \quad \text{with} \quad \max_{x,y \in \mathcal{I}} |\mathcal{X}_{xy}^{(ij)}(s,\gamma,k)| \prec N^{-k/2}. \tag{B.7}$$

Note that $(G_s)_{(ij)}^{\theta,0}$ is independent of X_{ij}^{γ} , $\gamma \in \{0,1\}$. Then, with (B.6) and (B.7), using the moment matching condition (3.11), we obtain that

$$\sum_{i \leq j \in \mathcal{I}} \left[\mathbb{E}F\left(H_{(ij)}^{\theta, X_{ij}^1}\right) - \mathbb{E}F\left(H_{(ij)}^{\theta, X_{ij}^0}\right) \right]$$

$$= \sum_{i \leq j \in \mathcal{I}} \sum_{1 \leq k, l \leq 7, k+l \geq 4} \mathbb{E}\left[\langle \mathcal{X}^{(ij)}(1, 1, k) E_a \mathcal{X}^{(ij)}(2, 1, l) E_b \rangle - \langle \mathcal{X}^{(ij)}(1, 0, k) E_a \mathcal{X}^{(ij)}(2, 0, l) E_b \rangle \right]$$

$$+ \mathcal{O}_{\prec}(N^{-3/2} \eta^{-1/2}), \tag{B.8}$$

where we have applied the anisotropic local law for $(G_s)_{(ij)}^{\theta,0}$ to control the terms containing the error $O_{\prec}(N^{-4})$ in (B.6). When $k + l \ge 4$, using the moment matching condition (3.12), we get that

$$\sum_{i \le j \in \mathcal{I}} \mathbb{E} \left[\langle \mathcal{X}^{(ij)}(1, 1, k) E_a \mathcal{X}^{(ij)}(2, 1, l) E_b \rangle - \langle \mathcal{X}^{(ij)}(1, 0, k) E_a \mathcal{X}^{(ij)}(2, 0, l) E_b \rangle \right] \prec \left(N^{-2-\delta} \wedge N^{-(k+l)/2} \right) \sum_{i \le j \in \mathcal{I}} \mathbb{E} \left| \left\langle (G_1)_{(ij)}^{\theta, 0} \left[\Delta_{ij} (G_1)_{(ij)}^{\theta, 0} \right]^k E_a \left[(G_2)_{(ij)}^{\theta, 0} \Delta_{ij} \right]^l (G_2)_{(ij)}^{\theta, 0} E_b \rangle \right| \prec N^{-3-\delta} \sum_{i \le j \in \mathcal{I}} \mathbb{E} \left| \operatorname{Tr} \left[(G_2)_{(ij)}^{\theta, 0} E_b (G_1)_{(ij)}^{\theta, 0} \left[\Delta_{ij} (G_1)_{(ij)}^{\theta, 0} \right]^k E_a \left[(G_2)_{(ij)}^{\theta, 0} \Delta_{ij} \right]^l \right] \right| \prec N^{-3-\delta} \cdot \left(N^2 \eta^{-2} \right) = N^{-1-\delta} \eta^{-2},$$
(B.9)

where in the third step we have applied the estimates (2.20) and (2.23) to $(G_s)_{(ij)}^{\theta,0}$. Plugging (B.9) into (B.8) concludes the proof.

Appendix C. Proof of local laws

C.1. **Proof of Lemma 2.10.** In this subsection, we briefly outline the proof of Lemma 2.10, which is divided into two parts: a probabilistic part that focuses on deriving the self-consistent equation, and a deterministic part concerning the stability of this equation.

The probabilistic part of the proof has been addressed by the arguments presented in [72]. Following the notation therein, we define the functions $\Pi \equiv \Pi(\cdot, z) : \mathbb{C}^{DN \times DN} \to \mathbb{C}^{DN \times DN}$ and $R \equiv R(\cdot, z) : \mathbb{C} \to \mathbb{C}$ as follows:

$$\Pi(X) := I + zX + \mathcal{S}(X)X - \Lambda X, \quad R_{\xi} \equiv R(\xi) := (\Lambda - \xi - z)^{-1}$$

Then, we can write M(z) and G(z) as

 g_k

 $M = R_m, \quad G = R_{\mathcal{S}(G)} - R_{\mathcal{S}(G)} \Pi(G).$

Subtracting the above two equations yields

$$G - M = R_{\mathcal{S}(G)} - R_m - R_{\mathcal{S}(G)}\Pi(G)$$

Multiplying both sides with $E_k, k \in [D]$, and taking the averaged trace, we obtain that

$$-m - D\langle (R_{\mathcal{S}(G)} - R_m)E_k \rangle = -D\langle R_{\mathcal{S}(G)}\Pi(G)E_k \rangle, \quad \text{with} \quad g_k := D\langle GE_k \rangle. \tag{C.1}$$

Using the arguments in [72], one can show that $\Pi(G)$ is indeed a small error, and $\langle \Pi(G)B \rangle$ satisfies a better bound for any deterministic matrix B with $||B|| \leq 1$. Furthermore, $\mathcal{S}(G)$ can be written as $\mathcal{S}(G) = \sum_{k=1}^{D} g_k E_k$. Hence, (C.1) leads to a self-consistent vector equation for $\mathbf{g} = (g_1, \ldots, g_D) \in \mathbb{C}^D$:

$$g_k - m - D\langle (R_{\mathbf{g}\cdot\mathbf{E}} - R_m)E_k \rangle = \mathcal{E}_k, \quad k \in \llbracket D \rrbracket,$$
(C.2)

where $\mathbf{E} := (E_1, \ldots, E_D)$, $\mathbf{g} \cdot \mathbf{E} := \sum_{k=1}^{D} g_k E_k$, and \mathcal{E}_k denotes an error term. When $\mathcal{E}_k = 0$, this equation admits a solution $\mathbf{m} := (m(z), \ldots, m(z)) \in \mathbb{C}^D$. By combining the stability of the self-consistent equation (C.2) as presented in Lemma C.1 below with the error estimates established for $\Pi(G)$ and $\langle \Pi(G)B \rangle$, we can deduce the local laws (2.20) and (2.21) employing the arguments in Section 4 of [72]. Let $\kappa_E :=$ $|E - b_N| \wedge |E + a_N|$ (recall that $-a_N$ and b_N represent the spectral edges). In the proof of the averaged local law (2.21), we can also derive a stronger estimate outside the spectrum:

$$\langle G(z) \rangle - m(z) \prec \frac{1}{N(\kappa_E + \eta)} + \frac{1}{(N\eta)^2 \sqrt{\kappa_E + \eta}},$$

uniformly in $z = E + i\eta$ with $|z| \le \tau^{-1}$, $\eta \ge N^{-1+\tau}$, and $N\eta\sqrt{\kappa_E + \eta} \ge N^{\tau}$. By combining this estimate with the local law (2.21) when B = I, we can derive (2.22) using the argument in [67].

It remains to establish the stability of the self-consistent equation (C.2). Our framework closely aligns with the general settings outlined in [9,58], except that our model does not satisfy the flatness assumption in them (see Assumption A1 of [9] or Assumption E of [58]). This flatness assumption is required by [9,58] to guarantee the stability of the self-consistent equations. Nevertheless, the absence of this assumption can be compensated for by the simple form of our self-consistent equation, which is a perturbation of the quadratic self-consistent equation for semicircle law, due to our assumption that ||A|| = o(1). Consider the self-consistent equation $\mathbf{f}(z, \mathbf{x}) = 0$, where the vector-valued function $\mathbf{f} : \mathbb{C}_+ \times \mathbb{C}^D \to \mathbb{C}^D$ is defined as

$$f_k(z, \mathbf{x}) := x_k - m(z) - D\langle (R_{\mathbf{x} \cdot \mathbf{E}}(z) - R_{m(z)}(z))E_k \rangle, \quad k \in \llbracket D \rrbracket.$$

Define the domain $\mathbf{D}(\tau) := \{z = E + i\eta : |z| \le \tau^{-1}, \eta \ge N^{-1+\tau}\}$. For each $z \in \mathbf{D}(\tau)$, introduce the set

$$L(z) := \{z\} \cup \{z' \in \mathbf{D}(\tau) : \operatorname{Re} z' = \operatorname{Re} z, \operatorname{Im} z' \in [\operatorname{Im} z, \infty) \cap (N^{-10}\mathbb{N})\}.$$

In other words, L(z) is a 1-dimensional lattice with spacing N^{-10} along with the point z. The stability estimate we aim to establish for the equation $\mathbf{f}(z, \mathbf{x}) = 0$ on $\mathbf{D}(\tau)$ is as follows:

Lemma C.1. Let $c_0 > 0$ be a sufficiently small constant. The self-consistent equation $\mathbf{f}(z, \mathbf{x}) = 0$ is stable on $\mathbf{D}(\tau)$ in the following sense. Suppose the z-dependent function δ satisfies $N^{-2} \leq \delta(z) \leq (\log N)^{-1}$ for $z \in \mathbf{D}(\tau)$ and is Lipschitz continuous with a Lipschitz constant $\leq N^3$. Additionally, assume that for each fixed E, the function $\eta \mapsto \delta(E + i\eta)$ is non-increasing for $\eta > 0$. Suppose every component u_k of a vector function $\mathbf{u} : \mathbf{D}(\tau) \to \mathbb{C}$ is the Stieltjes transform of a probability measure on \mathbb{R} . For any $z \in \mathbf{D}(\tau)$, if for every $z' \in L(z)$ the bound

$$\|\mathbf{f}(z',\mathbf{u}(z'))\|_{\infty} \le \delta(z') \tag{C.3}$$

holds, then we have that

$$\|\mathbf{u}(z) - \mathbf{m}(z)\|_{\infty} \le \frac{C\delta(z)}{\operatorname{Im} m(z) + \sqrt{\delta(z)}},\tag{C.4}$$

where C > 0 is a constant independent of both z and N.

Proof. A delicate stability analysis of a general class of quadratic self-consistent equations has been carried out carefully in [7,9,20]. Following the ideas from these works, we will now outline the proof of the stability estimate (C.4) without giving all the details.

Given an arbitrary $z = E + i\eta \in \mathbf{D}(\tau)$, suppose we have proved that (C.4) holds for all $z \in L(z')$ with $\operatorname{Im} z' > \operatorname{Im} z$. Then, using the Lipschitz continuity of the functions $\delta(z)$, $\mathbf{u}(z)$, and $\mathbf{m}(z)$, we obtain that

$$\|\mathbf{u}(z) - \mathbf{m}(z)\|_{\infty} \le \frac{2C\delta(z)}{\operatorname{Im} m(z) + \sqrt{\delta(z)}}.$$
(C.5)

By assumption, $\mathbf{u}(z)$ satisfies the system of equations $f_k(z, \mathbf{u}(z)) = \varepsilon_k(z)$, $k \in \llbracket D \rrbracket$, for some quantities $\varepsilon_k(z)$ with $\max_k |\varepsilon_k| \le \delta(z)$. Under condition (C.5), we can perform a Taylor expansion of $R_{\mathbf{u}\cdot\mathbf{E}}$ to express these equations as

$$v_k - D\langle M(\mathbf{v} \cdot \mathbf{E})ME_k \rangle - D\langle M(\mathbf{v} \cdot \mathbf{E})M(\mathbf{v} \cdot \mathbf{E})ME_k \rangle = \varepsilon_k + O(\|\mathbf{v}\|_{\infty}^3), \quad k \in \llbracket D \rrbracket$$

where we denote $\mathbf{v} \equiv \mathbf{u} - \mathbf{m}$. Recalling \widehat{M} defined in (3.1), we can rewrite the above equation as

$$v_k - \sum_{l=1}^{D} \widehat{M}_{kl}(z, z) v_l - m(z)^3 v_k^2 = \varepsilon_k + \mathcal{O}(\|\mathbf{v}\|_{\infty}^3 + \|A\|^{1/2} \|\mathbf{v}\|_{\infty}^2).$$
(C.6)

In the derivation of (C.6), we also utilized the fact that

$$D\langle M(\mathbf{v}\cdot\mathbf{E})M(\mathbf{v}\cdot\mathbf{E})ME_k\rangle = \sum_{l,s} D\langle ME_lME_sME_k\rangle v_lv_s = m(z)^3 v_k^2 + \mathcal{O}(\|A\|^{1/2}\|\mathbf{v}\|_{\infty}^2),$$

where the second step stems from (A.1). Let $k \in [\![D]\!]$ such that $|v_k| = |\!|\mathbf{v}|\!|_{\infty}$. Then, we can write (C.6) as

$$a(z)v_k - m(z)^3 v_k^2 = \varepsilon_k + \mathcal{O}(\|\mathbf{v}\|_{\infty}^3 + \|A\|^{1/2} \|\mathbf{v}\|_{\infty}^2),$$
(C.7)

where $a(z) := 1 - \sum_{l=1}^{D} \widehat{M}_{kl}(z, z) v_l / v_k$ satisfies that

$$|a(z)| \gtrsim \sqrt{\kappa_E + \eta} \sim \operatorname{Im} m(z) \tag{C.8}$$

by using Lemma A.1 and a similar expansion as in (A.13) for $\widehat{M}_{kl}(z_1, z_2)$ with $z_1 = z_2 = z$. By using (C.8) and (C.5), along with an analysis of equation (C.7) similar to the reasoning in the proof of [20, Lemma 4.5], we can conclude the estimate (C.4). Detailed specifics are omitted here.

It remains to establish the stability estimate (C.4) for a specific $z_0 \in L(z)$. Combining this initial stability estimate with the continuity argument outlined above, we can conclude the proof of Lemma C.1 by induction. For this purpose, we choose $z_0 = E + iC_0$ for a sufficiently large constant $C_0 > 0$. Since both m(z) and the components of **u** are Stieltjes transforms, we have the a priori bounds

$$|m(z_0)|_{\infty} \le C_0^{-1}, \quad ||\mathbf{u}(z_0)||_{\infty} \le C_0^{-1}.$$
 (C.9)

As long as C_0 is chosen sufficiently large, $R_{\mathbf{u}\cdot\mathbf{E}}$ behaves well, allowing us to perform a Taylor expansion. This expansion leads us back to equation (C.6) with $|a(z_0)| \ge 1/2$. Analyzing this equation reveals that

$$\|\mathbf{u}(z_0) - \mathbf{m}(z_0)\|_{\infty} \le C_1 \delta(z_0) \text{ or } \|\mathbf{u}(z_0) - \mathbf{m}(z_0)\|_{\infty} \ge C_1^{-1}$$

for a constant $C_1 > 0$ independent of C_0 . However, the latter scenario contradicts (C.9) if we choose $C_0 > 2C_1$. Therefore, we must have $\|\mathbf{u}(z_0) - \mathbf{m}(z_0)\|_{\infty} \leq C\delta(z)$, establishing the initial stability estimate at z_0 and concluding the proof of Lemma C.1.

C.2. **Proof of Lemma 2.11.** The proof of Lemma 2.11 uses an induction argument based on the Cauchy-Schwarz inequality and the following classical Ward's identity, which follows from a simple algebraic calculation.

Lemma C.2 (Ward's identity). Let \mathcal{A} be a Hermitian matrix. Define its resolvent as $R(z) := (\mathcal{A} - z)^{-1}$ for any $z = E + i\eta \in \mathbb{C}_+$. Then, we have

$$\sum_{x} \overline{R_{xy'}} R_{xy} = \frac{R_{y'y} - \overline{R_{yy'}}}{2i\eta}, \quad \sum_{x} \overline{R_{y'x}} R_{yx} = \frac{R_{yy'} - \overline{R_{y'y}}}{2i\eta}.$$
 (C.10)

As a special case, if y = y', we have

$$\sum_{x} |R_{xy}(z)|^2 = \sum_{x} |R_{yx}(z)|^2 = \frac{\operatorname{Im} R_{yy}(z)}{\eta}.$$
 (C.11)

The case of p = 1 follows directly from the anisotropic local law (2.20). Suppose now that $p \ge 2$ and the estimate (2.23) holds for products of k many $G_i B_i$'s when $1 \le k \le p - 1$.

First, if p = 2q is even, then we have

$$\begin{aligned} |(G_1B_1\cdots G_pB_p)_{\mathbf{u}\mathbf{v}}| &= |(\mathbf{u}^*G_1B_1\cdots G_q)B_q(G_{q+1}B_{q+1}\cdots G_pB_p\mathbf{v})| \\ &\leqslant \Big(\sum_j |(G_1B_1\cdots G_q)_{\mathbf{u}j}|^2\Big)^{1/2} \Big(\sum_j |(G_{q+1}B_{q+1}\cdots G_pB_p)_{j\mathbf{v}}|^2\Big)^{1/2}. \end{aligned}$$

Using Ward's identity, we get

$$\sum_{j} |(G_1 B_1 \cdots G_q)_{\mathbf{v}j}|^2 = \eta^{-1} [G_1 B_1 \cdots G_{q-1} B_{q-1} (\operatorname{Im} G_q) B_{q-1}^* G_{q-1}^* \cdots B_1^* G_1^*]_{\mathbf{v}\mathbf{v}}$$
(C.12)

$$=\frac{(G_1B_1\cdots G_q\cdots B_1^*G_1^*)_{vv}-(G_1B_1\cdots G_q^*\cdots B_1^*G_1^*)_{vv}}{2i\eta}\prec \frac{1}{\eta^{2q-1}},$$

where we used the induction hypothesis in the last step with the observation that the two terms in the numerator have (2q-1) many G's. The term $\sum_{j} |(G_{q+1}B_{q+1}\cdots G_pB_p)_{j\mathbf{v}}|^2$ can be proved in the same way. This gives (2.23) for the even p case.

On the other hand, if p = 2q + 1 is odd, we find that

$$|(G_{1}B_{1}\cdots G_{p}B_{p})_{\mathbf{uv}}| \leq \left(\sum_{j} |(G_{1}B_{1}\cdots G_{q+1})_{\mathbf{uj}}|^{2}\right)^{1/2} \left(\sum_{j} |(G_{q+2}B_{q+2}\cdots G_{p}B_{p})_{j\mathbf{v}}|^{2}\right)^{1/2} \qquad (C.13)$$
$$\prec \eta^{-(2q-1)/2} \left(\sum_{j} |(G_{1}B_{1}\cdots G_{q+1})_{\mathbf{uj}}|^{2}\right)^{1/2},$$

where we have bounded $\sum_{j} |(G_{q+2}B_{q+2}\cdots G_{p}B_{p})_{j\mathbf{v}}|^{2}$ as in (C.12). Now, we use the induction hypothesis and the trivial bound $||G_{i}|| \leq \eta^{-1}$ to get that

$$\sum_{j} |(G_1 B_1 \cdots G_{q+1})_{\mathbf{u}j}|^2 = \eta^{-1} (G_1 B_1 \cdots G_q B_q (\operatorname{Im} G_{q+1}) B_q^* G_q^* \cdots B_1^* G_1^*)_{\mathbf{u}\mathbf{u}}$$
$$\leqslant \eta^{-2} (G_1 B_1 \cdots G_q G_q^* \cdots B_1^* G_1^*)_{\mathbf{u}\mathbf{u}} \prec \eta^{-(2q+1)}.$$

Plugging it into (C.13), we get (2.23) for the odd p case.

Finally, Lemma 2.11 follows by induction in p.

References

- A. Adhikari, S. Dubova, C. Xu, and J. Yin. Eigenstate thermalization hypothesis for generalized Wigner matrices. arXiv preprint arXiv:2302.00157, 2023.
- [2] A. Adhikari and J. Huang. Dyson Brownian motion for general β and potential at the edge. Probability Theory and Related Fields, 178(3):893–950, 2020.
- [3] A. Adhikari and B. Landon. Local law and rigidity for unitary Brownian motion. Probability Theory and Related Fields, 187(3):753–815, 2023.
- [4] A. Aggarwal, C. Bordenave, and P. Lopatto. Mobility edge for Lévy matrices. arXiv preprint arXiv:2210.09458, 2022.
- [5] A. Aggarwal, P. Lopatto, and J. Marcinek. Eigenvector statistics of Lévy matrices. The Annals of Probability, 49(4):1778– 1846, 2021.
- [6] A. Aggarwal, P. Lopatto, and H.-T. Yau. GOE statistics for Lévy matrices. J. Eur. Math. Soc., 23(11):3707-3800, 2021.
- [7] O. Ajanki, L. Erdős, and T. Krüger. Quadratic vector equations on complex upper half-plane. Mem. Amer. Math. Soc., 261(1261):v+133, 2019.
- [8] O. H. Ajanki, L. Erdős, and T. Krüger. Universality for general Wigner-type matrices. Probability Theory and Related Fields, 169:667–727, 2017.
- [9] O. H. Ajanki, L. Erdős, and T. Krüger. Stability of the matrix Dyson equation and random matrices with correlations. Probability Theory and Related Fields, 173(1):293–373, 2019.
- [10] J. Alt, R. Ducatez, and A. Knowles. Delocalization transition for critical Erdős-Rényi graphs. Communications in Mathematical Physics, 388(1):507–579, 2021.
- [11] N. Anantharaman and E. Le Masson. Quantum ergodicity on large regular graphs. Duke Math. J., 164(4):723–765, 2015.
- [12] P. W. Anderson. Absence of diffusion in certain random lattices. *Phys. Rev.*, 109:1492–1505, 1958.
- [13] Z. Bao and L. Erdős. Delocalization for a class of random block band matrices. Probab. Theory Related Fields, 167(3):673– 776, 2017.
- [14] R. Bauerschmidt, J. Huang, A. Knowles, and H.-T. Yau. Bulk eigenvalue statistics for random regular graphs. The Annals of Probability, 45(6A):3626 – 3663, 2017.
- [15] R. Bauerschmidt, J. Huang, and H.-T. Yau. Local Kesten-Mckay law for random regular graphs. Communications in Mathematical Physics, 369(2):523-636, 2019.
- [16] R. Bauerschmidt, A. Knowles, and H.-T. Yau. Local semicircle law for random regular graphs. Communications on Pure and Applied Mathematics, 70(10):1898–1960, 2017.
- [17] L. Benigni and P. Lopatto. Fluctuations in local quantum unique ergodicity for generalized Wigner matrices. Communications in Mathematical Physics, 391(2):401–454, 2022.
- [18] M. V. Berry and M. Robnik. Semiclassical level spacings when regular and chaotic orbits coexist. Journal of Physics A: Mathematical and General, 17(12):2413–2421, aug 1984.
- [19] M. V. Berry and M. Tabor. Level clustering in the regular spectrum. Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences, 356(1686):375–394, 1977.
- [20] A. Bloemendal, L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Isotropic local laws for sample covariance and generalized Wigner matrices. *Electron. J. Probab.*, 19(33):1–53, 2014.
- [21] O. Bohigas. Random matrix theories and chaotic dynamics. In M.-J. Giannoni, A. Voros, and J. Zinn-Justin, editors, Chaos et Physique Quantique/Chaos and Quantum Physics, page 87–199. Elsevier Science Publishers, 1991.

- [22] O. Bohigas and M.-J. Giannoni. Chaotic motion and random matrix theories. In J. S. Dehesa, J. M. G. Gomez, and A. Polls, editors, *Mathematical and Computational Methods in Nuclear Physics*, pages 1–99, Berlin, Heidelberg, 1984. Springer Berlin Heidelberg.
- [23] O. Bohigas, M. J. Giannoni, and C. Schmit. Characterization of chaotic quantum spectra and universality of level fluctuation laws. Phys. Rev. Lett., 52:1–4, 1984.
- [24] O. Bohigas, M. J. Giannoni, and C. Schmit. Spectral properties of the Laplacian and random matrix theories. J. Physique Lett., 45:1015–1022, 1984.
- [25] P. Bourgade. Random band matrices. In Proceedings of the International Congress of Mathematicians. Rio de Janeiro 2018. Vol. IV. Invited lectures, pages 2759–2784. World Sci. Publ., Hackensack, NJ, 2018.
- [26] P. Bourgade. Extreme gaps between eigenvalues of Wigner matrices. Journal of the European Mathematical Society, 24(8):2823–2873, 2022.
- [27] P. Bourgade, L. Erdős, H.-T. Yau, and J. Yin. Fixed energy universality for generalized Wigner matrices. Communications on Pure and Applied Mathematics, 69(10):1815–1881, 2016.
- [28] P. Bourgade, L. Erdos, H.-T. Yau, and J. Yin. Universality for a class of random band matrices. Advances in Theoretical and Mathematical Physics, 21(3):739–800, 2017.
- [29] P. Bourgade and H. Falconet. Liouville quantum gravity from random matrix dynamics. arXiv:2206.03029, 2022.
- [30] P. Bourgade, J. Huang, and H.-T. Yau. Eigenvector statistics of sparse random matrices. *Electron. J. Probab.*, 22:38 pp., 2017.
- [31] P. Bourgade and J. P. Keating. Quantum chaos, random matrix theory, and the Riemann ζ-function. In B. Duplantier, S. Nonnenmacher, and V. Rivasseau, editors, *Chaos: Poincaré Seminar 2010*, pages 125–168. Springer Basel, Basel, 2013.
- [32] P. Bourgade, F. Yang, H.-T. Yau, and J. Yin. Random band matrices in the delocalized phase, II: Generalized resolvent estimates. Journal of Statistical Physics, 174(6):1189–1221, 2019.
- [33] P. Bourgade and H.-T. Yau. The eigenvector moment flow and local quantum unique ergodicity. Communications in Mathematical Physics, pages 1–48, 2016.
- [34] P. Bourgade, H.-T. Yau, and J. Yin. Random band matrices in the delocalized phase, I: Quantum unique ergodicity and universality. Communications on Pure and Applied Mathematics, 73(7):1526–1596, 2020.
- [35] G. Casati, I. Guarneri, F. Izrailev, and R. Scharf. Scaling behavior of localization in quantum chaos. Phys. Rev. Lett., 64:5–8, Jan 1990.
- [36] G. Casati, L. Molinari, and F. Izrailev. Scaling properties of band random matrices. Phys. Rev. Lett., 64:1851–1854, Apr 1990.
- [37] N. Chen and C. K. Smart. Random band matrix localization by scalar fluctuations. arXiv:2206.06439, 2022.
- [38] Z. Cheng and J. L. Lebowitz. Statistics of energy levels in integrable quantum systems. Phys. Rev. A, 44:R3399–R3402, 1991.
- [39] Z. Cheng, J. L. Lebowitz, and P. Major. On the number of lattice points between two enlarged and randomly shifted, copies of an oval. *Probability Theory and Related Fields*, 100(2):253–268, Jun 1994.
- [40] G. Cipolloni, L. Erdős, and J. Henheik. Eigenstate thermalisation at the edge for Wigner matrices. arXiv preprint arXiv:2309.05488, 2023.
- [41] G. Cipolloni, L. Erdős, J. Henheik, and O. Kolupaiev. Gaussian fluctuations in the equipartition principle for Wigner matrices. Forum of Mathematics, Sigma, 11:e74, 2023.
- [42] G. Cipolloni, L. Erdős, and D. Schröder. Eigenstate thermalization hypothesis for Wigner matrices. Communications in Mathematical Physics, 388(2):1005–1048, Dec 2021.
- [43] G. Cipolloni, L. Erdős, and D. Schröder. Normal fluctuation in quantum ergodicity for Wigner matrices. The Annals of Probability, 50(3):984 – 1012, 2022.
- [44] G. Cipolloni, L. Erdős, and D. Schröder. Optimal multi-resolvent local laws for Wigner matrices. Electronic Journal of Probability, 27(none):1 – 38, 2022.
- [45] G. Cipolloni, L. Erdős, and D. Schröder. Thermalisation for Wigner matrices. Journal of Functional Analysis, 282(8):109394, 2022.
- [46] G. Cipolloni, L. Erdős, and D. Schröder. Mesoscopic central limit theorem for non-Hermitian random matrices. Probability Theory and Related Fields, 2023.
- [47] G. Cipolloni, L. Erdős, and D. Schröder. Rank-uniform local law for Wigner matrices. Forum of Mathematics, Sigma, 10:e96, 2022.
- [48] G. Cipolloni, L. Erdős, and Y. Xu. Universality of extremal eigenvalues of large random matrices. arXiv preprint arXiv:2312.08325, 2023.
- [49] G. Cipolloni, L. Erdős, and D. Schröder. Functional central limit theorems for Wigner matrices. The Annals of Applied Probability, 33(1):447 – 489, 2023.
- [50] G. Cipolloni, R. Peled, J. Schenker, and J. Shapiro. Dynamical localization for random band matrices up to $W \ll N^{1/4}$. arXiv:2206.05545, 2022.
- [51] Y. Colin de Verdiere. Ergodicité et fonctions propres du laplacien. Communications in Mathematical Physics, 102(3):497– 502, 1985.
- [52] L. Erdős and A. Knowles. Quantum diffusion and delocalization for band matrices with general distribution. Annales Henri Poincaré, 12(7):1227, 2011.
- [53] L. Erdős and A. Knowles. Quantum diffusion and eigenfunction delocalization in a random band matrix model. Communications in Mathematical Physics, 303(2):509–554, 2011.
- [54] L. Erdős, A. Knowles, and H.-T. Yau. Averaging fluctuations in resolvents of random band matrices. Ann. Henri Poincaré, 14:1837–1926, 2013.

- [55] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Spectral statistics of Erdős-Rényi graphs II: Eigenvalue spacing and the extreme eigenvalues. Communications in Mathematical Physics, 314(3):587–640, 2012.
- [56] L. Erdos, A. Knowles, H.-T. Yau, and J. Yin. Delocalization and diffusion profile for random band matrices. Communications in Mathematical Physics, 323(1):367–416, 2013.
- [57] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Spectral statistics of Erdős–Rényi graphs I: local semicircle law. The Annals of Probability, 41(3B):2279–2375, 2013.
- [58] L. Erdős, T. Krüger, and D. Schröder. Random matrices with slow correlation decay. Forum of Mathematics, Sigma, 7:e8, 2019.
- [59] L. Erdős, S. Péché, J. A. Ramírez, B. Schlein, and H.-T. Yau. Bulk universality for Wigner matrices. Comm. Pure Appl. Math., 63(7):895–925, 2010.
- [60] L. Erdős, B. Schlein, and H.-T. Yau. Local semicircle law and complete delocalization for Wigner random matrices. Commun. Math. Phys., 287(2):641–655, 2008.
- [61] L. Erdős, B. Schlein, and H.-T. Yau. Semicircle law on short scales and delocalization of eigenvectors for Wigner random matrices. Annals of Probability, 37:815–852, 2009.
- [62] L. Erdős, B. Schlein, and H.-T. Yau. Universality of random matrices and local relaxation flow. Invent. Math., 185(1):75– 119, 2011.
- [63] L. Erdős and H.-T. Yau. Universality of local spectral statistics of random matrices. Bull. Amer. Math. Soc., 49(3):377–414, 2012.
- [64] L. Erdős and H.-T. Yau. A dynamical approach to random matrix theory, volume 28. American Mathematical Soc., 2017.
 [65] L. Erdős, H.-T. Yau, and J. Yin. Universality for generalized Wigner matrices with Bernoulli distribution. Journal of

Combinatorics, 2(1):15-82, 2011.

- [66] L. Erdős, H.-T. Yau, and J. Yin. Bulk universality for generalized Wigner matrices. Probability Theory and Related Fields, 154(1):341–407, 2012.
- [67] L. Erdős, H.-T. Yau, and J. Yin. Rigidity of eigenvalues of generalized Wigner matrices. Advances in Mathematics, 229(3):1435–1515, 2012.
- [68] M. Feingold, D. M. Leitner, and M. Wilkinson. Spectral statistics in semiclassical random-matrix ensembles. Phys. Rev. Lett., 66:986–989, Feb 1991.
- [69] Y. V. Fyodorov and A. D. Mirlin. Scaling properties of localization in random band matrices: A σ -model approach. *Phys. Rev. Lett.*, 67:2405–2409, Oct 1991.
- [70] M. Goldstein. Fluctuations and localization length for random band GOE matrix. arXiv:2210.04346, 2022.
- [71] Y. He and A. Knowles. Mesoscopic eigenvalue statistics of Wigner matrices. The Annals of Applied Probability, 27(3):1510– 1550, 6 2017.
- [72] Y. He, A. Knowles, and R. Rosenthal. Isotropic self-consistent equations for mean-field random matrices. Probability Theory and Related Fields, 171(1):203–249, Jun 2018.
- [73] P. D. Hislop and M. Krishna. On the local eigenvalue statistics for random band matrices in the localization regime. Journal of Statistical Physics, 187(3):26, 2022.
- [74] R. Holowinsky and K. Soundararajan. Mass equidistribution for Hecke eigenforms. Annals of Mathematics, 172(2):1517– 1528, 2010.
- [75] J. Huang and B. Landon. Rigidity and a mesoscopic central limit theorem for Dyson Brownian motion for general β and potentials. *Probability Theory and Related Fields*, 175(1):209–253, 2019.
- [76] J. Huang, B. Landon, and H.-T. Yau. Bulk universality of sparse random matrices. Journal of Mathematical Physics, 56(12):123301, 12 2015.
- [77] A. Khorunzhy, B. Khoruzhenko, and L. Pastur. Asymptotic properties of large random matrices with independent entries. *Journal of Mathematical Physics*, 37(10):5033–5060, 1996.
- [78] A. Knowles and J. Yin. Anisotropic local laws for random matrices. Probability Theory and Related Fields, 169(1-2):pp-257, 2017.
- [79] B. Landon, P. Lopatto, and P. Sosoe. Single eigenvalue fluctuations of general Wigner-type matrices. Probability Theory and Related Fields, 188(1):1–62, 2024.
- [80] B. Landon and P. Sosoe. Almost-optimal bulk regularity conditions in the CLT for Wigner matrices. arXiv:2204.03419, 2022.
- [81] B. Landon, P. Sosoe, and H.-T. Yau. Fixed energy universality of Dyson Brownian motion. Advances in Mathematics, 346:1137–1332, 2019.
- [82] B. Landon and H.-T. Yau. Convergence of local statistics of Dyson Brownian motion. Comm. Math. Phys., 355(3):949– 1000, 2017.
- [83] J. O. Lee and K. Schnelli. Edge universality for deformed Wigner matrices. Reviews in Mathematical Physics, 27(08):1550018, 2015.
- [84] J.-O. Lee, K. Schnelli, B. Stetler, and H.-T. Yau. Bulk universality for deformed Wigner matrices. Ann. Probab., 44(3):2349-2425, 2016.
- [85] E. Lindenstrauss. Invariant measures and arithmetic quantum unique ergodicity. Annals of Mathematics, 163(1):165–219, 2006.
- [86] J. Marcinek and H.-T. Yau. High dimensional normality of noisy eigenvectors. Communications in Mathematical Physics, 395(3):1007–1096, 2022.
- [87] J. Marklof. The Berry-Tabor conjecture. In C. Casacuberta, R. M. Miró-Roig, J. Verdera, and S. Xambó-Descamps, editors, *European Congress of Mathematics*, pages 421–427, Basel, 2001. Birkhäuser Basel.

- [88] J. Marklof. Pair correlation densities of inhomogeneous quadratic forms, II. Duke Mathematical Journal, 115(3):409 434, 2002.
- [89] J. Marklof. Pair correlation densities of inhomogeneous quadratic forms. Annals of Mathematics, 158(2):419–471, 2003.
- [90] R. Oppermann and F. Wegner. Disordered system withn orbitals per site: 1/n expansion. Zeitschrift für Physik B Condensed Matter, 34(4):327–348, 1979.
- [91] R. Peled, J. Schenker, M. Shamis, and S. Sodin. On the Wegner Orbital Model. International Mathematics Research Notices, 2019(4):1030–1058, 2017.
- [92] Z. Rudnick and P. Sarnak. The behaviour of eigenstates of arithmetic hyperbolic manifolds. Comm. Math. Phys., 161(1):195-213, 1994.
- [93] L. Schäfer and F. J. Wegner. Disordered system with n orbitals per site: Lagrange formulation, hyperbolic symmetry, and goldstone modes. Zeitschrift für Physik B Condensed Matter, 38:113–126, 1980.
- [94] J. Schenker. Eigenvector localization for random band matrices with power law band width. Comm. Math. Phys., 290:1065–1097, 2009.
- [95] M. Shcherbina and T. Shcherbina. Characteristic polynomials for 1d random band matrices from the localization side. Communications in Mathematical Physics, 351(3):1009–1044, 2017.
- [96] M. Shcherbina and T. Shcherbina. Universality for 1d random band matrices: Sigma-model approximation. Journal of Statistical Physics, 172(2):627-664, 2018.
- [97] M. Shcherbina and T. Shcherbina. Universality for 1d random band matrices. Communications in Mathematical Physics, 385(2):667-716, 2021.
- [98] T. Shcherbina. On the second mixed moment of the characteristic polynomials of 1d band matrices. Comm. Math. Phys., 328:45–82, 2014.
- [99] T. Shcherbina. Universality of the local regime for the block band matrices with a finite number of blocks. J. Stat. Phys., 155:466-499, 2014.
- [100] T. Shcherbina. Universality of the second mixed moment of the characteristic polynomials of the 1d band matrices: Real symmetric case. J. Math. Phys., 56, 2015.
- [101] A. I. Shnirel'man. Ergodic properties of eigenfunctions. Uspekhi Mat. Nauk, 29:181-182, 1974.
- [102] T. Spencer. Random banded and sparse matrices. In G. Akemann, J. Baik, and P. D. Francesco, editors, Oxford Handbook of Random Matrix Theory, chapter 23. Oxford University Press, New York, 2011.
- [103] T. Tao and V. Vu. Random matrices: universality of local eigenvalue statistics. Acta Math., 206(1), 2011.
- [104] J. M. VanderKam. Pair correlation of four-dimensional flat tori. Duke Mathematical Journal, 97(2):413 438, 1999.
- [105] F. J. Wegner. Disordered system with n orbitals per site: $n = \infty$ limit. Phys. Rev. B, 19:783–792, Jan 1979.
- [106] E. P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. Annals of Mathematics, 62(3):548–564, 1955.
- [107] C. Xu, F. Yang, H.-T. Yau, and J. Yin. Bulk universality and quantum unique ergodicity for random band matrices in high dimensions. To appear in Ann. Probab., arXiv preprint arXiv:2207.14533, 2022.
- [108] F. Yang, H.-T. Yau, and J. Yin. Delocalization and quantum diffusion of random band matrices in high dimensions I: Self-energy renormalization. arXiv preprint arXiv:2104.12048, 2021.
- [109] F. Yang, H.-T. Yau, and J. Yin. Delocalization and quantum diffusion of random band matrices in high dimensions II: T-expansion. Communications in Mathematical Physics, pages 1–96, 2022.
- [110] F. Yang and J. Yin. Delocalization of a general class of random block schrödinger operators. In preparation.
- [111] F. Yang and J. Yin. Random band matrices in the delocalized phase, III: averaging fluctuations. Probability Theory and Related Fields, 179(1):451–540, 2021.
- [112] S. Zelditch. Uniform distribution of eigenfunctions on compact hyperbolic surfaces. Duke Mathematical Journal, 55(4):919– 941, 1987.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, LOS ANGELES, CA, USA. *Email address*: bertrand.stone@math.ucla.edu

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, AND BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS, BEIJING, CHINA.

Email address: fyangmath@mail.tsinghua.edu.cn

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, LOS ANGELES, CA, USA. *Email address:* jyin@math.ucla.edu