AN EXAMPLE OF AN INFINITE AMENABLE GROUP WITH THE ISR PROPERTY

YONGLE JIANG AND XIAOYAN ZHOU*

ABSTRACT. Let G be $S_{\mathbb{N}}$, the finitary permutation (i.e., permutations with finite support) group on the set of positive integers \mathbb{N} . We prove that G has the invariant von Neumann subalgebras rigidity (ISR, for short) property as introduced in Amrutam-Jiang's work. More precisely, every G-invariant von Neumann subalgebra $P \subseteq L(G)$ is of the form L(H) for some normal sugbroup $H \triangleleft G$ and in this case, $H = \{e\}, A_{\mathbb{N}}$ or G, where $A_{\mathbb{N}}$ denotes the finitary alternating group on \mathbb{N} , i.e., the subgroup of all even permutations in $S_{\mathbb{N}}$. This gives the first known example of an infinite amenable group with the ISR property.

1. INTRODUCTION

Let G be a countable discrete group and L(G) be the group von Neumann algebra generated by G. Note that G acts on L(G) naturally by conjugation. We say a von Neumann subalgebra $P \subseteq L(G)$ is *invariant* if it is invariant as a set under the G-conjugation action. For example, if H is a normal subgroup in G, then L(H) is an invariant von Neumann subalgebra in L(G).

It is a natural question to classify all invariant von Neumann subalgebras in L(G). In [1, Corollary 4.8], Alekseev and Brugger proved that for a lattice subgroup G in a higher rank simple real Lie group with trivial center, every invariant subfactor is either \mathbb{C} or has finite Jones's index in L(G) using character rigidity techniques introduced in [15,23]. For other classes of groups complementary to lattices in higher rank groups, including all non-abelian free groups, Chifan and Das made further progress (see [7, Theorem 3.15, Theorem 3.16 and Corollary 3.17] for precise statements) on describing invariant subfactors but by deformation/rigidity techniques for array/quasi-cocycles on groups as introduced and studied in [9–12].

Concerning the above question, the simplest answer one can hope is that every invariant von Neumann subalgebra is of the form L(H) for some normal subgroup $H \triangleleft G$. In [3], Amrutam and the first named author introduced "the invariant von Neumann subalgebras rigidity" (ISR for short) property for a group G once this best situation occurs, i.e., every invariant von Neumann subalgebra of L(G) arises

Date: April 4, 2024.

²⁰¹⁰ Mathematics Subject Classification. Primary 46L10, Secondary 22D25 22D10 43A35.

Key words and phrases. invariant von Neumann subalgebras, characters, finitary permutation groups, finitary alternating groups.

as the group von Neumann algebra of a normal subgroup of G. It is not hard to see that to search for groups with the ISR property, we may assume that G is infinite (see Proposition 2.5). The first known class of groups with this property is due to Kalantar and Panagopoulos [20]. They proved that any irreducible lattice inside a connected semisimple Lie group with trivial center, no non-trivial compact factors, and such that all its simple factors have real rank at least two has the ISR property. In fact, this result is the motivation for introducing the ISR property in [3]. In [3], Amrutam and the first named author found other classes of groups with the ISR property, including all torsion-free hyperbolic groups with trivial amenable radical, torsion-free non-amenable groups with zero first L^2 -Betti number under a mild assumption and finite direct product of groups from these classes. Subsequently, Chifan, Das and Sun [8] extended some results in [3] to much wider classes, e.g., all acylindricially hyperbolic groups with trivial amenable radical. Moreover, they conjectured that all non-amenable groups with trivial amenable radical should have the ISR property. Quite recently, in the remarkable work [2], Amrutam, Hartman and Oppelmayer made important progress on this conjecture by proving that for any non-amenable group G, $L(G_a)$ is the unique maximal invariant amenable von Neumann subalgebra in L(G), where G_a denotes the amenable radical of G. Therefore, for a non-amenable group G with trivial G_a , the only amenable invariant von Neumann subalgebra in L(G) is the trivial subalgebra \mathbb{C} .

However, despite all these results, there are still two fundamental questions unanswered. One is whether the ISR property could be shared by any infinite amenable groups. Note that techniques in [3, 8, 20] are mainly developed to deal with nonamenable groups with trivial amenable radical. The other question is whether the ISR property is preserved under W^{*}-equivalence.

In this paper, we answer the above mentioned two questions. For the first question, we show that the answer is positive by proving the following theorem.

Theorem 1.1. Let $S_{\mathbb{N}}$ be the finitary permutation group on the set of positive integers \mathbb{N} . Let $P \subseteq L(S_{\mathbb{N}})$ be an invariant von Neumann subalgebra in $L(S_{\mathbb{N}})$. Then $P = \mathbb{C}$, $L(A_{\mathbb{N}})$ or $L(S_{\mathbb{N}})$, where $A_{\mathbb{N}}$ denotes the finitary alternating group on \mathbb{N} .

Combining it with [3, Example 3.5], we show that the second question has a negative answer.

Corollary 1.2. The ISR property is not preserved under W^* -equivalence, i.e., there exist two countable discrete groups G and H such that $L(G) \cong L(H)$, but exactly one of these groups has the ISR property.

Let us briefly sketch the proof of Theorem 1.1.

Let $G = S_{\mathbb{N}}$ and P be an invariant von Neumann subalgebra in L(G). Denote by $E : (L(G), \tau) \to P$ the τ -preserving conditional expectation onto P, where τ denotes the canonical trace on L(G). Note that $\tau(x) = \langle x \delta_e, \delta_e \rangle$ for all $x \in L(G)$ and we set $\langle a, b \rangle_{\tau} := \tau(b^*a)$ for any $a, b \in L(G)$. As explained in [3], to show that Ghas the ISR property, we need to argue that $E(u_g) \in \mathbb{C}u_g$ for all $g \in G$. Hereafter we denote by g the canonical unitary $u_g \in L(G)$ for simplicity. In other words, we need to argue that the trace (in the sense of Definition 2.2) $G \ni g \stackrel{\phi}{\mapsto} \tau(E(g)g^{-1}) \in \mathbb{C}$ is the characteristic function supported on a normal subgroup of G.

First, we observe that for any $g \in G$, if we write $E(g) = \sum_{s \in G} c_s s$ for its Fourier expansion as an element in L(G), then $\operatorname{supp}(s) \subseteq \operatorname{supp}(g)$ for all $c_s \neq 0$. Here, for any $g \in G$, we write $\operatorname{supp}(g) = \{i \in \mathbb{N} : gi \neq i\}$ for the support of g. Then, it is not hard to see that for the transposition (1 2), we have E((1 2)) = (1 2) or E((1 2)) = 0. In the first case, we show that E(g) = g for all $g \in G$; equivalently, the trace ϕ defined above is the constant function one on G and P = L(G). So we may assume that E((1 2)) = 0.

The next step is to argue that $E((1\ 2\ 3)) = 0$ or $E((1\ 2\ 3)) = (1\ 2\ 3)$. In the case $E((1\ 2\ 3)) = (1\ 2\ 3)$, we argue that $P = L(A_{\mathbb{N}})$ since $A_{\mathbb{N}}$ is normally generated by all 3-cycles and has index two in G (see Proposition 2.1). Finally, for the case $E((1\ 2\ 3)) = 0$, we observe that the classification of characters on G by Thoma [28], or more precisely, Corollary 2.4 actually implies E(g) = 0 for all $e \neq g \in G$. This new approach to study the ISR property via the classification of characters is the main innovation in this paper.

Structure of the paper. In Section 2, we record the classification of all normal subgroups of $S_{\mathbb{N}}$ and review basic facts on the notion of traces and characters on groups, then we prove a key observation on traces on $S_{\mathbb{N}}$ based on Thoma's classification of characters on $S_{\mathbb{N}}$, i.e., Corollary 2.4. In Section 3, we prove the main theorem and its corollary.

Notations and conventions.

- Denote by \mathbb{N} the collection of all positive integers.
- For any $A \subseteq \mathbb{N}$, we write $S_A = \{s \in S_{\mathbb{N}} : \operatorname{supp}(s) \subseteq A\}$. Note that S_A is a subgroup in $S_{\mathbb{N}}$.
- Let $n \ge 1$. Set $[n] = \{1, 2, \dots, n\}$.
- Let n_1, \ldots, n_k be k-many distinct elements in N. We write $s = (n_1 \cdots n_k) \in S_{\mathbb{N}}$ to mean the k-cycle defined by $sn_i = n_{i+1 \mod k}$ and si = i for all $i \notin \{n_1, \ldots, n_k\}$.
- Let $k \geq 2$. For any two tuples $(n_1, n_2, \ldots, n_k), (m_1, m_2, \ldots, m_k) \in \mathbb{N}^k$, both of which have distinct entries, then we write $s = \begin{pmatrix} n_1 & n_2 & \cdots & n_k \\ m_1 & m_2 & \cdots & m_k \end{pmatrix}$ to mean any element $s \in S_{\mathbb{N}}$ such that $sn_i = m_i$ for all $1 \leq i \leq k$. Since we only use the fact $s(n_1 n_2 \cdots n_k)s^{-1} = (m_1 m_2 \cdots m_k)$ in this paper, the ambiguity of swould not cause problem.

2. Preliminaries

The classification of all normal subgroups of $S_{\mathbb{N}}$ is well-known, see e.g., [22, 26]. We are grateful to Prof. Mikael de la Salle for pointing out these references. For the sake of completion, we add a proof below.

Proposition 2.1. Let $N \triangleleft S_{\mathbb{N}}$ be a normal subgroup. Then either $N = \{e\}$, $N = A_{\mathbb{N}}$ or $N = S_{\mathbb{N}}$. Moreover, the subgroup index $[S_{\mathbb{N}} : A_{\mathbb{N}}] = 2$.

Proof. To see the subgroup index is two, we just note that $S_{\mathbb{N}} = A_{\mathbb{N}} \sqcup (1 \ 2)A_{\mathbb{N}}$. Indeed, recall that $[n] := \{1, \ldots, n\}$, then we may take the union over $n \ge 5$ on both sides of the equality $S_{[n]} = A_{[n]} \sqcup (1 \ 2)A_{[n]}$ for all $n \ge 5$.

Note that $S_{\mathbb{N}} = \bigcup_{n \geq 5} S_{[n]}$ and $A_{\mathbb{N}} = \bigcup_{n \geq 5} A_{[n]}$. Moreover, $A_{[n]}$ is a simple group and also the unique proper non-trivial normal subgroup in $S_{[n]}$ for $n \geq 5$, see e.g., [25, Theorem 3.11 and Exercise 3.21, Chapter 3].

Since $N \cap S_{[n]} \triangleleft S_{[n]}$, we deduce that $N \cap S_{[n]}$ is either $\{e\}$, $A_{[n]}$ or $S_{[n]}$ for any $n \ge 5$. If there exits some $n_0 \ge 5$ such that $N \cap S_{[n_0]} = S_{[n_0]}$, then $S_{[n_0]} \subseteq N$. Hence for all $m > n_0$, we have $S_{[n_0]} \subseteq N \cap S_{[m]} = S_{[m]}$ since $S_{[n_0]} \not\subseteq A_{[m]}$. Therefore, $S_{[m]} \subseteq N$ and hence $N = N \cap S_{\mathbb{N}} = N \cap (\bigcup_{m \ge n_0} S_{[m]}) = \bigcup_{m \ge n_0} (N \cap S_{[m]}) = \bigcup_{m \ge n_0} S_{[m]} = S_{\mathbb{N}}$. From now on, we may assume $N \cap S_{[n]} = A_{[n]}$ or $\{e\}$ for all $n \ge 5$.

Case 1: $N \cap S_{[n]} = A_{[n]}$ for all $n \ge 5$.

Then $A_{[n]} \subseteq N$ for all $n \geq 5$ and hence $A_{\mathbb{N}} = \bigcup_{n \geq 5} A_{[n]} \subseteq N \subseteq S_{\mathbb{N}}$. Clearly, $N \neq S_{\mathbb{N}}$, thus the fact that $A_{\mathbb{N}}$ has index two in $S_{\mathbb{N}}$ implies $N = A_{\mathbb{N}}$.

Case 2: there exists some $n_0 \ge 5$ such that $N \cap S_{[n_0]} = \{e\}$.

Take any $m > n_0$, then by our assumption, $N \cap S_{[m]} = \{e\}$ or $A_{[m]}$. Assume that $N \cap S_{[m]} = A_{[m]}$ for some $m > n_0$, then $\{e\} = N \cap S_{[n_0]} = (N \cap S_{[m]}) \cap S_{[n_0]} = A_{[m]} \cap S_{[n_0]} = A_{[n_0]}$, a contradiction. Hence $N \cap S_{[m]} = \{e\}$ for all $m > n_0$. Thus, $N = \bigcup_{m \ge n_0} (N \cap S_{[m]}) = \bigcup_{m \ge n_0} \{e\} = \{e\}$.

Next, we recall some basic facts on the notion of traces on groups.

Definition 2.2. A trace on a group G is a function $\phi : G \to \mathbb{C}$ so that

(1) ϕ is positive definite, i.e.,

$$\sum_{i=1}^{n} \sum_{j=1}^{n} \overline{\alpha_i} \alpha_j \phi(s_i^{-1} s_j) \ge 0$$

for all $n \in \mathbb{N}$ and any choice of elements $s_1, \ldots, s_n \in G$ and $\alpha_1, \ldots, \alpha_n \in \mathbb{C}$, (2) ϕ is conjugation invariant, i.e., $\phi(sts^{-1}) = \phi(t)$ for all elements $s, t \in G$ and (3) ϕ is normalized, i.e., $\phi(e) = 1$.

A character on G is an extreme point in the set of all traces on G.

We remind the reader that some researchers prefer calling characters instead of traces defined here (extremal or indecomposable characters instead of characters defined here respectively), see e.g., [16,17,19,23]. Here, we have followed the terminology used in [18,21].

Let Tr(G) be the space of traces on G equipped with the topology of pointwise convergence. Then Tr(G) is a compact convex subset of $\ell^{\infty}(G)$. Let Ch(G) be the set of characters on G. It is known that Tr(G) is a metrizable Choquet simplex [27]. This means that the barycenter map

$$Prob(Ch(G)) \to Tr(G), \ \mu \mapsto \phi_{\mu} := \int_{Ch(G)} \chi d\mu(\chi).$$

is a continuous affine bijection, see e.g., [5, Theorem 4.5, Chapter 4] and [24] for an overview of Choquet theory.

The classification of characters on various classes of groups have been an active topic, see [4] for an overview. In particular, the following famous classical theorem due to Thoma is the main ingredient in our proof. See [5, Chapter 4] for an exposition on this theorem and also [19] for an operator algebraic proof.

Theorem 2.3 (Thoma, [28]). A character of the group $S_{\mathbb{N}}$ is of the form

$$\chi(s) = \prod_{k=2}^{\infty} (\sum_{i=1}^{\infty} a_i^k + (-1)^{k-1} \sum_{j=1}^{\infty} b_j^k)^{m_k(s)}.$$

Here, $m_k(s)$ is the number of k-cycles in the permutation s and the two sequences $(a_i)_{i=1}^{\infty}, (b_j)_{j=1}^{\infty}$ satisfy

$$a_1 \ge a_2 \ge \dots \ge 0, \ b_1 \ge b_2 \ge \dots \ge 0, \ \sum_{i=1}^{\infty} a_i + \sum_{j=1}^{\infty} b_j \le 1.$$

Recall that each permutation $s \in S_{\mathbb{N}}$ can be decomposed into disjoint cycles $s_1, s_2, \ldots, s_n \in S_{\mathbb{N}}$ for some $n \in \mathbb{N}$ such that $s = s_1 s_2 \cdots s_n$. A cycle $(n_1 \ n_2 \ \cdots \ n_k)$ has the length k and is referred to as a k-cycle.

We observe the following fact holds.

Corollary 2.4. Let ϕ be a trace on $S_{\mathbb{N}}$. Assume that $\phi((1\ 2\ 3)) = 0$, then $\phi(s) = 0$ for all $e \neq s \in S_{\mathbb{N}}$, i.e., ϕ is the regular character δ_e .

Proof. Note that $m_k((1\ 2\ 3)) = 1$ if k = 3 and 0 for all $2 \le k \ne 3$. Then by Theorem 2.3, we deduce that

(1)
$$\chi((1\ 2\ 3)) = \sum_{i=1}^{\infty} a_i^3 + \sum_{j=1}^{\infty} b_j^3, \ \forall \ \chi \in Ch(S_{\mathbb{N}}).$$

Note that since $a_i \ge 0$ and $b_j \ge 0$ for all $i, j \ge 1$, the above formula implies that

(2)
$$\chi((1\ 2\ 3)) \ge 0, \ \forall \ \chi \in Ch(S_{\mathbb{N}}).$$

First, we assume that ϕ is a character, i.e., $\phi \in Ch(S_{\mathbb{N}})$. Since $\phi((1\ 2\ 3)) = 0$, we deduce that $a_i = b_j = 0$ for all $i, j \ge 1$ from (1) and (2). Given any $e \ne s$, there exists some $k \ge 2$ such that $m_k(s) > 0$, hence the k-th term in the product expression of $\phi(s)$ is 0, and therefore $\phi(s) = 0$.

Now, let ϕ be a general trace. Since Tr(G) is a Choquet simplex, we may write $\phi = \int_{Ch(G)} \chi d\mu(\chi)$ for some $\mu \in Prob(Ch(G))$. Then, from $0 = \phi((1\ 2\ 3)) = \int_{Ch(G)} \chi((1\ 2\ 3))d\mu(\chi)$ and (2), we deduce that for μ -a.e. $\chi \in Ch(G)$, we have

 $\chi((1\ 2\ 3)) = 0$. Thus $\chi(s) = 0$ for all $e \neq s \in G$ and μ -a.e. $\chi \in Ch(G)$ by what we have shown above. Thus, $\phi(s) = 0$ for all $e \neq s \in G$.

The following proposition explains why we focus on infinite groups when discussing the ISR property. In fact, it also follows directly from the proof of [3, Proposition 3.1], and we decide to include the proof here.

Proposition 2.5. Let G be a finite group. Then G has the ISR property if and only if G is trivial or $G = C_2$, the cyclic group of order two.

Proof. To see the "if" direction holds, observe that for $G = C_2$, $L(G) \cong \mathbb{C} \oplus \mathbb{C}$, which contains only two von Neumann subalgebras, i.e., the diagonal subalgebra $P = \{(c, c) : \in \mathbb{C}\} \cong \mathbb{C} = L(\{e\})$ and L(G) itself.

Next, we prove the "only if" direction holds. Assume that the finite group G has the ISR property, then the center $P := \mathcal{Z}(L(G)) = L(H)$ for some subgroup $H \lhd G$ since P is clearly G-invariant. Notice that $p = \sum_{g \in G} g \in P$, we deduce that G = H and thus, $L(G) = \mathcal{Z}(LG)$, which is abelian. Hence G is abelian.

Since G is abelian, every von Neumann subalgebra in L(G) is invariant. Moreover, $L(G) \cong \ell^{\infty}([n])$, where $n = \sharp G$. Assume that $n \ge 2$. Notice that $P := \{(x_i) \in \ell^{\infty}([n]) : x_1 = x_2\}$ is a von Neumann subalgebra and hence P = L(H) for some $H \triangleleft G$. Notice that $\sharp H = n - 1$, then from the Lagrange theorem, we get that $(n-1) \mid n$. Thus, n = 2. Clearly, this finishes the proof. \Box

3. Proofs

We will frequently use the simple fact that st = ts for any $s, t \in S_{\mathbb{N}}$ with disjoint supports.

Lemma 3.1. Let $A \subseteq \mathbb{N}$ be an infinite subset. Then $L(S_A)' \cap L(S_{\mathbb{N}}) = L(S_{\mathbb{N}\setminus A})$.

Proof. " \supseteq " holds trivially. To see " \subseteq " holds, it suffices to check that for any $s \in S_{\mathbb{N}} \setminus S_{\mathbb{N}\setminus A}$, we have $\sharp\{tst^{-1} : t \in S_A\} = \infty$.

Take any $s \in S_{\mathbb{N}} \setminus S_{\mathbb{N}\setminus A}$, then $\operatorname{supp}(s) \cap A \neq \emptyset$. Fix any $i \in \operatorname{supp}(s) \cap A$. Since $\sharp A = \infty$ and $\sharp \operatorname{supp}(s) < \infty$, we may find an infinite strictly increasing sequence $\{j_k\}_{k\geq 1}$ in A such that $j_1 > \max\{j : j \in \operatorname{supp}(s)\}$. Then define the transpositions $t_k = (i \ j_k) \in S_A$ for all $k \geq 1$. We are left to check that $t_k s t_k^{-1} \neq t_\ell s t_\ell^{-1}$ if $k \neq \ell$. Indeed, observe that $\operatorname{supp}(t_k s t_k^{-1}) = t_k(\operatorname{supp}(s)) = t_k((\operatorname{supp}(s) \setminus \{i\}) \sqcup \{i\}) =$ $(\operatorname{supp}(s) \setminus \{i\}) \sqcup \{j_k\}$. It is clear that $\operatorname{supp}(t_k s t_k^{-1}) \neq \operatorname{supp}(t_\ell s t_\ell^{-1})$ if $k \neq \ell$ by our choice of $\{j_k\}_{k\geq 1}$. Hence $\sharp\{t_k s t_k^{-1} : k \geq 1\} = \infty$. \Box

Let G be any countable discrete group. Let $P \subseteq L(G)$ be a G-invariant von Neumann subalgebra. Denote by $E : (L(G), \tau) \to P$ the canonical τ -preserving conditional expectation. Then set $\phi(g) = \tau(E(g)g^{-1}) = \tau(E(g)E(g)^{-1}) = ||E(g)||_2^2$. It is well-known that ϕ is a trace on G, which has been used in [1]. For convenience of the reader, we repeat the proof below.

Proposition 3.2. Under the above notations, then $\phi : G \to \mathbb{C}$ is a trace on G.

Proof. It is clear that $\phi(e) = \tau(e) = 1$.

Next, we show that ϕ is conjugation invariant. Since P is G-invariant, we deduce that $gE(x)g^{-1} = E(gxg^{-1})$ for all $x \in L(G)$ and $g \in G$. Hence, $\phi(sts^{-1}) = \tau(E(sts^{-1})st^{-1}s^{-1}) = \tau(sE(t)s^{-1}st^{-1}s^{-1}) = \tau(E(t)t^{-1}) = \phi(t)$.

Finally, we show that ϕ is positive definite. The proof has appeared in the proof of [6, Theorem 12.2.15] or [13, Lemma 1]. Note that E is a unital completely positive map by [6, Theorem 1.5.10]. Thus, for any $s_1, \ldots, s_n \in G$ and $\alpha_1, \ldots, \alpha_n \in \mathbb{C}$, we have

$$\begin{split} \sum_{i=1}^{n} \sum_{j=1}^{n} \overline{\alpha_i} \alpha_j \phi(s_i^{-1} s_j) &= \sum_{i=1}^{n} \sum_{j=1}^{n} \overline{\alpha_i} \alpha_j \tau(E(s_i^{-1} s_j) s_j^{-1} s_i) \\ &= \sum_{i=1}^{n} \sum_{j=1}^{n} \overline{\alpha_i} \alpha_j \tau(s_i E(s_i^{-1} s_j) s_j^{-1}) \\ &= \sum_{i=1}^{n} \sum_{j=1}^{n} \overline{\alpha_i} \alpha_j \langle E(s_i^{-1} s_j) s_j^{-1} \delta_e, s_i^{-1} \delta_e \rangle \\ &= \sum_{i=1}^{n} \sum_{j=1}^{n} \langle E(s_i^{-1} s_j) \alpha_j \delta_{s_j^{-1}}, \alpha_i \delta_{s_i^{-1}} \rangle \\ &= \langle [E(s_i^{-1} s_j)]_{1 \le i, j \le n} (\alpha_j \delta_{s_j^{-1}})_{1 \le j \le n}, (\alpha_i \delta_{s_i^{-1}})_{1 \le i \le n} \rangle \ge 0, \end{split}$$

where $(\alpha_j \delta_{s_j^{-1}})_{1 \leq j \leq n} \in \ell^2(G)^{\oplus n}$ and $[E(s_i^{-1}s_j)]_{1 \leq i,j \leq n} \in M_n(L(G))^+$ since E is a unital completely positive map and $[(s_i^{-1}s_j)]_{1 \leq i,j \leq n} \in M_n(L(G))^+$.

From now on, we write G for $S_{\mathbb{N}}$. The following proposition would be needed for the proof of Theorem 1.1.

Proposition 3.3. Let $P \subseteq L(G)$ be a *G*-invariant von Neumann subalgebra. Let $E: (L(G), \tau) \to P$ be the canonical τ -preserving conditional expectation. Suppose that $E((1\ 2\ 3)) = \mu(1\ 2\ 3) + \theta(1\ 3\ 2)$ for some real numbers $\mu, \theta \in \mathbb{R}$, then $(\mu, \theta) \neq (\frac{1}{2}, \pm \frac{1}{2})$.

Proof. First, we prove that $(\mu, \theta) \neq (\frac{1}{2}, \frac{1}{2})$.

Assume this does not hold, then $\tilde{E}((1\ 2\ 3)) = \frac{1}{2}((1\ 2\ 3) + (1\ 3\ 2)) \in P$ and $E((1\ 2\ 4)) = (3\ 4)E(1\ 2\ 3)(3\ 4) = \frac{1}{2}((1\ 2\ 4) + (1\ 4\ 2)) \in P$. In general, we have $E(i\ j\ k) = \frac{1}{2}((i\ j\ k) + (i\ k\ j)) \in P$ for any three distinct natural numbers i, j, k. Hence,

(3)
$$P \ni [(1\ 2\ 3) + (1\ 3\ 2)][(1\ 2\ 4) + (1\ 4\ 2)] \\= (1\ 3)(2\ 4) + (1\ 4\ 3) + (2\ 4\ 3) + (1\ 4)(2\ 3).$$

Thus, taking the conjugation of the RHS in (3) by the transposition (3 4) and noticing that P is G-invariant, we get that

(4)
$$P \ni (1 \ 4)(2 \ 3) + (1 \ 3 \ 4) + (2 \ 3 \ 4) + (1 \ 3)(2 \ 4).$$

Adding the RHS of (3) to (4) and noticing that $(1 4 3) + (1 3 4), (2 3 4) + (2 4 3) \in P$, we get that

(5)
$$(1\ 3)(2\ 4) + (1\ 4)(2\ 3) \in P.$$

Taking the conjugacy of the LHS of (5) by the transposition (2 3), we get that

 $(1\ 2)(3\ 4) + (1\ 4)(2\ 3) \in P.$ (6)

Thus, by subtracting (6) from (5), we get that

(7)
$$(1\ 3)(2\ 4) - (1\ 2)(3\ 4) \in P.$$

Taking the conjugation of the LHS of (5) by the transposition (24), we get that

$$(8) (1 3)(2 4) + (1 2)(3 4) \in P.$$

Then by adding (7) to (8), we deduce that $(1 \ 3)(2 \ 4) \in P$. Thus by taking the conjugation of $(1 \ 3)(2 \ 4)$ by $(3 \ 4)$, we also have $(1 \ 4)(2 \ 3) \in P$. In view of (3), $(5 4 3) \in P$. So $(2 4 3) + (5 4 3) \in P$ after taking conjugation by (1 2). Hence, $a - (25)a(25) = (243) - (543) \in P$. Therefore, $(243) \in P$. Thus, $(123) \in P$ and $E((1\ 2\ 3)) = (1\ 2\ 3)$, a contradiction.

Next, let us show that $(\mu, \theta) \neq (\frac{1}{2}, \frac{-1}{2})$. Assume that $(\mu, \theta) = (\frac{1}{2}, \frac{-1}{2})$ instead, then $E((1\ 2\ 3)) = \frac{1}{2}((1\ 2\ 3) - (1\ 3\ 2)) \in P$. Hence, $E((1\ 2\ 4)) = \frac{1}{2}((1\ 2\ 4) - (1\ 4\ 2)) \in P$ and more generally, $E((i\ j\ k)) = \frac{1}{2}((1\ 2\ 4)) =$ $\frac{1}{2}((i \ j \ k) - (i \ k \ j)) \in \overline{P}$ for any three distinct natural numbers i, j, k. Then,

(9)
$$P \ni [(1\ 2\ 3) - (1\ 3\ 2)][(1\ 2\ 4) - (1\ 4\ 2)] \\= (1\ 3)(2\ 4) - (1\ 4\ 3) - (2\ 4\ 3) + (1\ 4)(2\ 3)$$

Taking the conjugation of the RHS of (9) by the transposition (23), we deduce that

(10)
$$P \ni (1\ 2)(3\ 4) - (1\ 4\ 2) - (3\ 4\ 2) + (1\ 4)(2\ 3)$$

Notice that $-(2\ 4\ 3) + (3\ 4\ 2) = -(2\ 4\ 3) + (2\ 3\ 4) \in P$, we may subtract (10) from (9) to get that

$$(11) (13)(24) - (12)(34) - (143) + (142) \in P.$$

Taking the conjugation of the LHS of (11) by (13), we get that

$$(12) (13)(24) - (23)(14) - (134) + (342) \in P$$

Subtracting (12) from (11) and then noticing that $-(1 4 3) + (1 3 4) \in P$, we deduce that

(13)
$$-(1\ 2)(3\ 4) + (2\ 3)(1\ 4) + (1\ 4\ 2) - (3\ 4\ 2) \in P.$$

Adding (10) to (13), we get that

 $(2\ 3)(1\ 4) - (3\ 4\ 2) \in P.$ (14)

In view of (10), this implies that $(1\ 2)(3\ 4) - (1\ 4\ 2) \in P$.

Then

$$0 = \langle (1 \ 4 \ 2) - E((1 \ 4 \ 2)), (1 \ 2)(3 \ 4) - (1 \ 4 \ 2)) \rangle_{\tau}$$

= $\langle (1 \ 4 \ 2) - \frac{1}{2}((1 \ 4 \ 2) - (1 \ 2 \ 4)), (1 \ 2)(3 \ 4) - (1 \ 4 \ 2) \rangle_{\tau} = \frac{-1}{2}.$

This yields a contradiction.

We are ready for the proof of Theorem 1.1.

Proof of Theorem 1.1. We assume the above notations. To start the proof, we observe the following simple fact.

(15)
$$\forall e \neq s \in G$$
, we have $E(s) \in L(S_{\text{supp}(s)})$.

Indeed, $\forall t \in S_{\mathbb{N}\setminus \text{supp}(s)}$, we have $tst^{-1} = s$. Hence, $E(s) = E(tst^{-1}) = tE(s)t^{-1}$ implies that $E(s) \in L(S_{\mathbb{N}\setminus \text{supp}(s)})' \cap G = L(S_{\text{supp}(s)})$, where the last equality follows by applying Lemma 3.1 to the infinite set $A = \mathbb{N} \setminus \text{supp}(s)$.

From (15), we may write $E((1\ 2)) = c + \mu(1\ 2)$ for some $c, \mu \in \mathbb{C}$. Hence,

$$c + \mu(1 \ 2) = E((1 \ 2)) = E(E((1 \ 2)))$$
$$= E(c + \mu(1 \ 2)) = c + \mu E((1 \ 2)) = c + \mu(c + \mu(1 \ 2)).$$

Equivalently, $c = c + c\mu$ and $\mu = \mu^2$. Hence, we get that either $\mu = 0$ or $(\mu, c) = (1, 0)$.

Case 1: $(\mu, c) = (1, 0)$.

In this case, we have $E((1\ 2)) = (1\ 2)$. Thus, for any $i \neq j \in \mathbb{N}$, set $s = \begin{pmatrix} 1 & 2 \\ i & j \end{pmatrix}$, then $E((i\ j)) = E(s(1\ 2)s^{-1}) = sE((1\ 2))s^{-1} = (i\ j)$. Thus, all transpositions belong to P. Since $S_{\mathbb{N}}$ is generated by transpositions, we deduce that $P = L(S_{\mathbb{N}})$. Case 2: $\mu = 0$.

We may write $E((1\ 2)) = c$. Hence $c = \tau(E((1\ 2))) = \tau((1\ 2)) = 0$, i.e.,

 $E((1\ 2)) = 0.$

Next, we show that $E((1\ 2\ 3)) = (1\ 2\ 3)$ or 0. We split the proof into several steps.

Step 1: we show that $E((1\ 2\ 3)) = \mu(1\ 2\ 3) + \theta(1\ 3\ 2)$ for some $\mu, \theta \in \mathbb{C}$. To see this, we observe the following facts.

- the three transpositions (1 2), (1 3) and (2 3) belong to the same orbit for the conjugation action of the subgroup $\langle (1 2 3) \rangle \curvearrowright S_{[3]}$. Indeed, (1 2 3)(1 3)(1 3 2) = (1 2), (1 2 3)(1 2)(1 3 2) = (2 3).
- $E((1\ 2\ 3))$ is invariant under the conjugation of $(1\ 2\ 3)$ and $(1\ 3\ 2)$.

•
$$\tau(E((1\ 2\ 3))) = 0.$$

Combining the above facts with (15), we may write

$$E((1\ 2\ 3)) = c((1\ 2) + (1\ 3) + (2\ 3)) + \mu(1\ 2\ 3) + \theta(1\ 3\ 2)$$

for some $c, \mu, \theta \in \mathbb{C}$. Then, we observe that $E((1\ 2)) = 0$ implies $\langle (1\ 2), x \rangle_{\tau} = 0$ for all $x \in P$. In particular, plugging in $x = E((1 \ 2 \ 3))$, we deduce that c = 0. Therefore,

$$E((1\ 2\ 3)) = \mu(1\ 2\ 3) + \theta(1\ 3\ 2)$$

Step 2: we show that the pair $(\mu, \theta) \in \{(0, 0), (1, 0), (\frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, \frac{-1}{2})\}.$ First, we observe that $\mu, \theta \in \mathbb{R}$. Indeed, on the one hand,

 $E((1\ 3\ 2)) = E((2\ 3)(1\ 2\ 3)(2\ 3)) = (2\ 3)E((1\ 2\ 3))(2\ 3) = \mu(1\ 3\ 2) + \theta(1\ 2\ 3);$

on the other hand, $E((1\ 3\ 2)) = E((1\ 2\ 3)^{-1}) = E((1\ 2\ 3))^* = \overline{\mu}(1\ 3\ 2) + \overline{\theta}(1\ 2\ 3).$ Hence, we get that $\mu, \theta \in \mathbb{R}$.

Then, we have

$$E((1\ 2\ 3)) = E(E((1\ 2\ 3))) = E(\mu(1\ 2\ 3) + \theta(1\ 3\ 2)) = \mu E((1\ 2\ 3)) + \theta E((1\ 3\ 2))$$
$$= \mu(\mu(1\ 2\ 3) + \theta(1\ 3\ 2)) + \theta(\mu(1\ 3\ 2) + \theta(1\ 2\ 3))$$
$$= (\mu^2 + \theta^2)(1\ 2\ 3) + 2\mu\theta(1\ 3\ 2).$$

By comparing it with the expression for $E((1\ 2\ 3))$, we deduce that $\mu = \mu^2 + \theta^2$ and $\theta = 2\mu\theta$ hold true. Hence, we get that either

- (i) $\theta = 0, \mu \in \{0, 1\}$ or, (ii) $\mu = \frac{1}{2}, \ \theta = \pm \frac{1}{2}.$

In other words, the pair $(\mu, \theta) \in \{(0, 0), (1, 0), (\frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, \frac{-1}{2})\}.$

Therefore, we deduce that $(\mu, \theta) \in \{(0, 0), (1, 0)\}$ from Proposition 3.3. **Step 3**: we show that if $(\mu, \theta) = (0, 0)$, then $P = \mathbb{C}$; if $(\mu, \theta) = (1, 0)$, then $P = L(A_{\mathbb{N}}).$

Case 1: $(\mu, \theta) = (1, 0).$

Then $E((1\ 2\ 3)) = (1\ 2\ 3) \in P$. For any three distinct natural numbers i, j, k, set $s = \begin{pmatrix} 1 & 2 & 3 \\ i & j & k \end{pmatrix}$, then we have $E((i & j & k)) = E(s(1 & 2 & 3)s^{-1}) = sE((1 & 2 & 3))s^{-1} = s(1 & 2 & 3)s^{-1} = (i & j & k) \in P$. Recall that $[n] := \{1, \ldots, n\}$. Since $A_{\mathbb{N}} = \bigcup_{n \ge 5} A_{[n]}$, it equals the normal subgroup of $S_{\mathbb{N}}$ generated by all 3-cycles [25, Lemma 3.9, Chapter 3], we deduce that $L(A_{\mathbb{N}}) \subseteq P \subseteq L(S_{\mathbb{N}})$. Then notice that $E((1 \ 2)) = 0$ implies $E((1\ 2)s) = E((1\ 2))s = 0$ for all $s \in A_{\mathbb{N}}$, hence $P = L(A_{\mathbb{N}})$ since $S_{\mathbb{N}} = A_{\mathbb{N}} \sqcup (1\ 2)A_{\mathbb{N}}$. Case 2: $(\mu, \theta) = (0, 0).$

In this case, we have $E((1\ 2\ 3)) = 0$. Hence $\phi((1\ 2\ 3)) = 0$ for the trace ϕ defined by $\phi(s) = \tau(E(s)s^{-1})$. Then $\phi(s) = 0$ for all $e \neq s \in S_{\mathbb{N}}$ by Corollary 2.4. Hence $||E(s)||_2^2 = \tau(E(s)E(s)^*) = \tau(E(s)E(s^{-1})) = \tau(E(s)s^{-1}) = \phi(s) = 0$. Thus, E(s) = 0 for all $e \neq s \in S_{\mathbb{N}}$. Therefore, $P = \mathbb{C}$.

Finally, we present the proof of Corollary 1.2.

Proof of Corollary 1.2. Recall that we have proved in [3, Example 3.5] that the group $H := \mathbb{Z} \wr \mathbb{Z}$ does not have the ISR property. Observe that both H and $G = S_{\mathbb{N}}$ are i.c.c. amenable groups, hence $L(G) \cong L(H) \cong \mathbb{R}$, the hyperfinite II₁ factor by Connes's theorem [14]. The proof is done since $S_{\mathbb{N}}$ has the ISR property by Theorem 1.1.

Acknowledgements. Y. J. is partially supported by National Natural Science Foundation of China (Grant No. 12001081) and the Fundamental Research Funds for the Central Universities (Grant No.DUT19RC(3)075). X. Z. is partially supported by National Natural Science Foundation of China (Grant No. 12001085) and the Scientific Research Fund of Liaoning Provincial Education Department (Grant No. LJKZ1047). We are grateful to Prof. Adam Skalski and Dr. Amrutam Tattwamasi for helpful comments. We also thank the anonymous referee for excellent comments which help improving the readability of the paper greatly.

References

- V. Alekseev and R. Brugger, A rigidity result for normalized subfactors, J. Operator Theory 86 (2021), no. 1, 3–15.
- [2] T. Amrutam, Y. Hartman, and H. Oppelmayer, On the amenable subalgebras of group von Neumann algebras (2023), arXiv preprint, arXiv: 2309.10494.
- [3] T. Amrutam and Y. Jiang, On invariant von Neumann subalgebras rigidity property, J. Funct. Anal. 284 (2023), no. 5, Paper No. 109804.
- [4] B. Bekka and P. de la Harpe, Unitary representations of groups, duals, and characters, Mathematical Surveys and Monographs, vol. 250, American Mathematical Society, Providence, RI, [2020] ©2020.
- [5] A. Borodin and G. Olshanski, Representations of the infinite symmetric group, Cambridge Studies in Advanced Mathematics, vol. 160, Cambridge University Press, Cambridge, 2017.
- [6] N. P. Brown and N. Ozawa, C^{*}-algebras and finite-dimensional approximations, Graduate Studies in Mathematics, vol. 88, American Mathematical Society, Providence, RI, 2008.
- [7] I. Chifan and S. Das, Rigidity results for von Neumann algebras arising from mixing extensions of profinite actions of groups on probability spaces, Math. Ann. 378 (2020), no. 3-4, 907–950.
- [8] I. Chifan, S. Das, and B. Sun, Invariant subalgebras of von Neumann algebras arising from negatively curved groups, J. Funct. Anal. 285 (2023), no. 9, Paper No. 110098.
- [9] I. Chifan, Y. Kida, and S. Pant, Primeness results for von Neumann algebras associated with surface braid groups, Int. Math. Res. Not. IMRN 16 (2016), 4807–4848.
- [10] I. Chifan and T. Sinclair, On the structural theory of II₁ factors of negatively curved groups, Ann. Sci. Éc. Norm. Supér. (4) 46 (2013), no. 1, 1–33.
- [11] I. Chifan, T. Sinclair, and B. Udrea, On the structural theory of II₁ factors of negatively curved groups, II: Actions by product groups, Adv. Math. 245 (2013), 208–236.
- [12] _____, Inner amenability for groups and central sequences in factors, Ergodic Theory Dynam. Systems **36** (2016), no. 4, 1106–1129.
- [13] M. Choda, Group factors of the Haagerup type, Proc. Japan Acad. Ser. A Math. Sci. 59 (1983), no. 5, 174–177.
- [14] A. Connes, Classification of injective factors. Cases II_1 , II_{∞} , III_{λ} , $\lambda \neq 1$, Ann. of Math. (2) **104** (1976), no. 1, 73–115.
- [15] D. Creutz and J. Peterson, Character rigidity for lattices and commensurators, Amer. J. Math., to appear, preprint available at arXiv: 1311.4513.
- [16] A. Dudko and K. Medynets, On characters of inductive limits of symmetric groups, J. Funct. Anal. 264 (2013), no. 7, 1565–1598.
- [17] _____, Finite factor representations of Higman-Thompson groups, Groups Geom. Dyn. 8 (2014), no. 2, 375–389.
- [18] C. Eckhardt and T. Shulman, On amenable Hilbert-Schmidt stable groups, J. Funct. Anal. 285 (2023), no. 3, Paper No. 109954.

- [19] R. Gohm and C. Köstler, Noncommutative independence from characters of the infinite symmetric group S_{∞} (2010), arXiv preprint, arXiv: 1005.5726v1.
- [20] M. Kalantar and N. Panagopoulos, On invariant subalgebras of group C* and von Neumann algebras, Ergodic Theory Dynam. Systems 43 (2023), no. 10, 3341–3353.
- [21] A. Levit and I. Vigdorovich, *Characters of solvable groups, Hilbert-Schmidt stability and dense periodic measures* (2022), Math. Ann. to appear, preprint available at arXiv: 2206.02268.
- [22] L. Onofri, Teoria delle sostituzioni che operano su una infinità numerabile di elementi, Ann. Mat. Pura Appl. 7 (1929), no. 1, 103–130.
- [23] J. Peterson, *Character rigidity for lattices in higher-rank groups* (2014), preprint, available at www.math.vanderbilt.edu/peters10/rigidity.pdf.
- [24] Robert R. Phelps, Lectures on Choquet's theorem, 2nd ed., Lecture Notes in Mathematics, vol. 1757, Springer-Verlag, Berlin, 2001.
- [25] J. J. Rotman, An introduction to the theory of groups, 4th ed., Graduate Texts in Mathematics, vol. 148, Springer-Verlag, New York, 1995.
- [26] J. Schreier and S. Ulam, Uber die Permutationsgruppe der natürlichen Zahlenfolge, Studia Math. 4 (1933), no. 1, 134–141.
- [27] E. Thoma, Uber positiv-definite Klassenfunctionen abzählbarer Gruppen, Math. Z. 84 (1964), 389–402.
- [28] _____, Die unzerlegbaren, positiv-definiten Klassenfunktionen der abzählbar unendlichen, symmetrischen Gruppe, Math. Z. 85 (1964), 40–61.

Y.J., School of Mathematical Sciences, Dalian University of Technology, Dalian, Liaoning, 116024, China

Email address: yonglejiang@dlut.edu.cn

X.Z., School of Date Science and Artificial Intelligence, Dongbei University of Finance & Economics, Dalian, Liaoning, 116025, China

Email address: xyzhou@dufe.edu.cn