

## Universality of extremal eigenvalues of large random matrices.

Giorgio Cipolloni\*  
Princeton University  
gc4233@princeton.edu

László Erdős†  
IST Austria  
lerdos@ist.ac.at

Yuanyuan Xu†  
AMSS, CAS  
yyxu2023@amss.ac.cn

**Abstract.** We prove that the spectral radius of a large random matrix  $X$  with independent, identically distributed complex entries follows the Gumbel law irrespective of the distribution of the matrix elements. This solves a long-standing conjecture of Bordenave and Chafaï and it establishes the first universality result for one of the most prominent extremal spectral statistics in random matrix theory. Furthermore, we also prove that the argument of the largest eigenvalue is uniform on the unit circle and that the extremal eigenvalues of  $X$  form a Poisson point process.

*Keywords:* Extremal statistics, Gumbel distribution, Ginibre ensemble, Dyson Brownian motion

*MSC number:* 60B20, 60G55, 60G70

*Date:* February 2, 2024

### 1. INTRODUCTION

**1.1. The main result.** Let  $X$  be a large  $n \times n$  matrix with independent, identically distributed (i.i.d.) complex entries that have zero expectation and variance  $1/n$ . It is well known by the *Circular Law* [73, 21, 77, 101, 119] that most eigenvalues lie in the unit disc, in fact their density becomes uniform as  $n$  tends to infinity. Denote by  $\rho(X)$  the spectral radius of  $X$ , i.e. the modulus of its largest eigenvalue. Our main result is that the fluctuation of  $\rho(X)$ , after appropriate rescaling, follows the *Gumbel Law*, in particular it is universal. More precisely, we show that

$$\rho(X) = 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{\mathcal{G}_n}{\sqrt{4n\gamma_n}}, \quad \gamma_n := \log n - 2 \log \log n - \log 2\pi, \quad (1.1)$$

where  $\mathcal{G}_n$  is an asymptotically Gumbel random variable, i.e.  $\mathbf{P}(\mathcal{G}_n \leq r) \rightarrow \exp(-e^{-r})$  for any  $r \in \mathbb{R}$  as  $n \rightarrow \infty$ . This solves a long-standing conjecture by Bordenave and Chafaï [28], [42], see also [110] and [43] for the log-gas counterpart of this question. We also show that the argument of the largest eigenvalue of  $X$  is uniform on  $[0, 2\pi]$ , and that its few largest (in modulus) eigenvalues form a *Poisson point process*.

**1.2. Related universality results.** We now place this result in the broader context of random matrix theory. Since the revolutionary discovery of E. Wigner [125] on the universal behaviour of local eigenvalue statistics for large random matrices, rigorous proofs aiming at verifying Wigner's vision have been in the forefront of mathematical research. Motivated by quantum mechanical Hamiltonians, Wigner considered Hermitian random matrices and predicted that the local correlation functions depend only on the basic symmetry type but otherwise are insensitive to the details of the matrix ensemble. In particular, real symmetric and complex Hermitian matrices exhibit somewhat different but universal eigenvalue statistics. Mathematically this was formalised as the Wigner-Dyson-Gaudin-Mehta universality conjecture in the 1960's [98]. After important partial results on the most prominent *Wigner matrices* [85] and *sample covariance matrices* [24], the conjecture was proven in full generality in a series of papers by L. Erdős, H.-T. Yau, and their collaborators, after they developed the theory of *Dyson Brownian motion (DBM)* for spectral universality in the *bulk* of the spectrum, see [65, 66] for a summary. In a parallel development, around the same time, T. Tao and V. Vu [120] also proved bulk universality for the complex symmetry class by a different method based upon the *four moment matching* idea. The eigenvalue distribution

†Partially supported by ERC Advanced Grant "RMTBeyond" No. 101020331.

near the *spectral edges* has been identified in the early 1990's by C. Tracy and H. Widom [122, 123], and its universality was proven shortly afterwards by A. Soshnikov [116]. The third, and last type of local spectral universality for Hermitian matrices, described by the *Pearcey process*, occurring at the possible cusp singularities of the limiting eigenvalue density, was proven recently in [64, 50]. This largely completes the theory of Hermitian universality for the simplest mean field ensembles. However, much more work has been done and yet to be done on more general Hermitian ensembles, especially random band matrices [113, 114, 45, 46, 22, 33, 36, 39, 63, 76, 109, 115, 126, 127, 128, 129], on adjacency matrices of sparse random graphs [82, 83, 23, 84, 10, 11, 12, 13, 80], and on Lévy matrices [4, 5, 6, 20, 25, 29, 30], that we will not discuss here.

Non-Hermitian matrices exhibit similar universality patterns; although their mathematical theory is much less developed, the progress spectacularly accelerated in the last years. The local spectral universality for matrices with i.i.d. entries at the spectral edge (the non-Hermitian analogue of Tracy-Widom regime) was proven in [54] both for real and complex matrices. The universality in the bulk (analogue of the Hermitian Wigner-Dyson statistics) for complex symmetry class was solved very recently in [96], the real case is still open. The existence of the critical edge (analogue of the Hermitian cusp regime) was proven in [62] for both symmetry classes and the presumedly universal local statistics was computed in [94] for the complex case but its universality is still open.

In contrast to local spectral statistics, our main result (1.1) is the prime example of a universal *extremal* statistics arising in non-integrable random matrix theory. It has no natural Hermitian counterpart: the largest eigenvalue of a Hermitian matrix is always found around the endpoint of the support of the density and its distribution has already been identified by the Tracy-Widom edge universality. In contrast, the spectrum of a non-Hermitian i.i.d matrix is essentially supported on the unit disc by the Circular Law, there are about  $\sqrt{n}$  eigenvalues outside but very close, and around each reference point on the unit circle the eigenvalues locally form a strongly correlated determinantal process [72, 32, 54]. Thus the spectral radius is realised as the maximum of  $\sqrt{n}$  different eigenvalue moduli that themselves are correlated, albeit on a short scale. Their tails decay fast but they are not compactly supported hence, if they were independent, their extreme value distribution would naturally fall into the Gumbel class among the three possible limiting extreme value distributions given by the Fisher-Tippett-Gnedenko theorem [69, 74] for extrema of i.i.d. random variables. The correlation, however, prohibits us from using standard extreme value theory directly. Although our final conclusion coincides with the naive prediction based upon independence, in fact we even prove that the few largest eigenvalues form a Poisson point process, for the proof we need to follow a completely different path than these heuristics.

For completeness, we mention that another very prominent extremal statistics problem arises in the celebrated Fyodorov-Hiary-Keating (FHK) conjecture [71] that connects extrema of log-determinants of Haar unitary random matrices with the Riemann  $\zeta$ -function. In particular, the precise distribution of the maximum of the logarithm of the characteristic polynomial of the circular  $\beta$ -ensemble was identified very recently [104] (see previous papers [17, 44, 103]), proving the random matrix side of the FHK-conjecture. For the currently best result on the number theory side of FHK-conjecture, see [18, 19] and references therein. Superficially, these results are reminiscent to (1.1), but the underlying point process is much more correlated than our extreme eigenvalues, in particular the limiting law is not given directly by the Gumbel distribution. On the other hand, the structure of the underlying randomness is very different, and, as consequence, so are the proof methods. For the circular  $\beta$ -ensemble the joint density of all eigenvalues is explicitly known, this serves as the fundamental starting point for the analysis.

Returning to i.i.d. matrices, if the original random matrix has independent Gaussian entries (called *Ginibre ensemble*; see recent survey papers [40, 41]), then the limiting distribution of the spectral radius can be explicitly computed. The Gumbel law was obtained in this way for the complex Ginibre ensemble in [108] and for the real Ginibre ensemble in [110] (see also [26, 9] for similar calculations for the *elliptic* Ginibre matrices). Therefore our task is to prove that this distribution for any i.i.d. matrix is universal, hence it coincides with the Gumbel law for the Ginibre ensemble. In this sense our proof is modelled by all previous universality proofs; the core of the work is to show universality, then the limiting distribution is identified via explicit calculations performed on the distinguished Gaussian ensemble. In this paper we prove universality for the complex ensemble. The core of our method also applies to the real case but the technical details are somewhat more tedious and will be left to future work.

**1.3. New ideas and methods.** Our proof relies on two fundamentally new ideas compared with any previous work on spectral statistics of non-Hermitian matrices, in particular compared with our previous

papers [60, 61] that identified the correct order of magnitude of the spectral radius but failed to identify its distribution itself. Both ideas are expected to be influential for other spectral problems on non-Hermitian random matrices hence we now briefly outline them.

Almost all mathematical works on the spectrum of non-Hermitian random matrices (with the notable exception of [27, 31, 70]) are based upon *Girko's formula* [73], see e.g. [21, 121, 28, 37, 16, 60]. Modulo an irrelevant cutoff at infinity (see (6.2)), this identity, for any sufficiently regular test function  $f : \mathbb{C} \rightarrow \mathbb{R}$ , asserts that

$$\sum_{\sigma \in \text{Spec}(X)} f(\sigma) = -\frac{1}{4\pi} \int_{\mathbb{C}} \Delta f(z) A(z) d^2 z, \quad A(z) := \int_0^\infty \text{Im Tr} G^z(i\eta) d\eta, \quad (1.2)$$

where  $G^z$  is a family of Hermitized resolvents, parametrized by a family of spectral parameters  $z \in \mathbb{C}$ :

$$G^z(w) := (H^z - w)^{-1}, \quad H^z := \begin{pmatrix} 0 & X - z \\ X^* - \bar{z} & 0 \end{pmatrix}, \quad w \in \mathbb{C} \setminus \mathbb{R}. \quad (1.3)$$

The advantage of (1.2) is that linear eigenvalue statistics on the original non-Hermitian problem are translated into a Hermitian problem, which is technically much more tractable since Hermitian matrices tend to be much more stable against perturbations. In particular the resolvent  $G^z$  tends to concentrate around its deterministic approximation. There are two main disadvantages of this formula: i) it requires the understanding of several  $H^z$ 's simultaneously, ii) the Laplacian  $\Delta f$  is very large if  $f$  is chosen strongly localized to capture single eigenvalues. Moreover, the spectral parameter  $i\eta$  must be integrated down to the real axis which makes the  $\eta$ -integral logarithmically divergent and it requires an extra probabilistic regularisation via an independently obtained lower tail bound on the lowest non-negative eigenvalue  $\lambda_1^z$  of  $H^z$ , noticing that  $\|G^z(i\eta)\| \lesssim |\lambda_1^z + \eta|^{-1}$ . A meticulous analysis of (1.2) along these ideas were behind all previous results on eigenvalue statistics of  $X$  on fine scales [37, 38, 121, 14, 52, 58, 60, 61].

Catching the spectral radius via Girko's formula is especially difficult. In fact, the universality of size and fluctuation of the spectral radius was not known even under the additional four moment matching assumption [121], showing the difficulty of grasping this extremal spectral statistics. The main difficulty lies in the fact that the test function  $f$  needs to be a very *anisotropic* function; it is essentially a smoothed characteristic function of a narrow annulus of radius  $1 + \sqrt{\gamma_n/4n} \approx 1$  and width of order  $1/\sqrt{n \log n}$  – a domain where the eigenvalue with largest modulus is typically located. For such function  $\int |\Delta f|$  is especially large compared with its integral  $\int f$  thus the effect of any error term in estimating  $A(z)$  in (1.2) will be drastically magnified when integrated against the absolute value of  $\Delta f$ . As our previous work [61] on the size of  $\rho(X)$  demonstrated, all additional technical refinements along these ideas were barely not sufficient to identify the distribution of  $\rho(X)$ , so we need to change the strategy. Nevertheless, we point out that several estimates developed in [61] are essential in the current paper as well.

The *first key new idea* is to perform two integration by parts in (1.2) and study

$$\sum_{\sigma \in \text{Spec}(X)} f(\sigma) = -\frac{1}{4\pi} \int_{\mathbb{C}} f(z) \int_0^\infty \text{Im Tr} \Delta_z G^z(i\eta) d\eta d^2 z \quad (1.4)$$

exploiting the chiral symmetry of  $H^z$ . This innocently looking step is a paradigm change in the analysis of Girko's formula. The  $z$ -integration now becomes quite harmless, but

$$\Delta_z G^z = G^z F G^z F^* G^z + G^z F^* G^z F G^z, \quad \text{with } F := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix},$$

looks prohibitively singular: it involves the product of three resolvents. Thus the  $\eta$ -integration near 0 becomes quadratically divergent as  $\int d\eta/\eta^3$  and it looks unregularisable<sup>1</sup> even with the most optimal estimate on the tail of  $\lambda_1^z$ . The main observation is that the *chiral symmetry* of  $H^z$  implies  $G^z F G^z F^* = 0$  exactly at  $\eta = 0$  if  $\lambda_1^z \neq 0$ . This gives hope that (1.4) is still controllable if the precise behaviour of  $G^z F G^z F^* G^z$  for  $\eta$  near zero is better understood. Note that we are looking for the *distribution* of (1.4) not just its size, hence we will need to compute higher moments or even general functions of both sides. This would require to understand the joint distribution of  $(G^z F G^z F^* G^z)(i\eta)$  at least at a few different parameters  $z$  and in the entire range of the spectral parameter  $i\eta$  along the imaginary axis. The actual

<sup>1</sup>We remark that *one* integration by parts leads to  $\int \text{Tr}[G^z(i\eta)]^2 F d\eta = \text{Tr} G^z F$  which is still manageable without chiral symmetry, see, e.g. [78, Eq (1.9)].

proof is much more subtle due to the delicate condition  $\lambda_1^z \neq 0$ . In fact, if one naively ignores this condition, then

$$\int_0^\infty \text{Tr}(G^z F G^z F^* G^z)(i\eta) d\eta = \frac{1}{2i} \text{Tr}(G^z F G^z F^*)(i\eta) \Big|_{\eta=0}^{\eta=\infty} = 0, \quad \text{if } \lambda_1^z \neq 0,$$

would yield that (1.4) is always zero which is clearly wrong. The correct procedure is to regularise the condition  $\lambda_1^z \neq 0$  (see Section 6.1 for details) and consider the  $z$ -derivatives of this regularisation as well in the integration by parts (1.4); see (7.73)-(7.75) later. Along the way  $\text{Tr}G^z F G^z F^*$  instead of  $\text{Tr}G^z F G^z F^* G^z$  needs to be analysed at a small but non-zero  $\eta$  for a few different  $z$ -parameters.

Very crudely speaking, the behaviour of a single resolvent  $G^z(i\eta)$  is governed by two different mechanisms. If  $\eta$  is much smaller than the lowest eigenvalue  $\lambda_1^z$ , then  $G^z(i\eta)$  is strongly fluctuating, mainly depending on the fluctuation on  $\lambda_1^z$ . If  $\eta$  is much bigger than the typical size of  $\lambda_1^z$ , then  $G^z(i\eta)$  tends to be deterministic – such concentration results are commonly called *local laws*. In the intermediate regime both effects are present and contribute to the answer. So we need to tackle both regimes and in the much more complicated situations of several  $G^z F G^z F^*$  monomials with a few different  $z$ 's.

For the small  $\eta$  regime we prove that the lower tail distribution of the lowest eigenvalues  $\lambda_1^{z_1}, \lambda_1^{z_2}, \dots, \lambda_1^{z_k}$  are essentially independent whenever the pairwise distance of the  $z_p$  parameters is bigger than the typical eigenvalue spacing,  $|z_p - z_{p'}| \gg n^{-1/2}$  for any  $p \neq p'$ . This drastically improves our analogous previous result that proved a similar independence but only when  $|z_p - z_{p'}| \geq n^{-\epsilon}$  [61, Proposition 4.3]. Here we rely on the strong contraction effect of the coupled *Dyson Brownian motion* that allows us to mimic the joint distribution of several  $\lambda_1^z$  by a collection of *a priori* independent eigenvalues. The core of this idea has been exploited earlier [52, 58, 60, 61], but the current precision down to the scale  $|z_p - z_{p'}| \gg n^{-1/2}$  is unprecedented in the edge regime. For simplicity we do it only in the complex symmetry class, but extension to the real case requires only technical adjustments (see the corresponding analysis in [53]). The main input is that the eigenvectors of  $H^z$  belonging to  $\lambda_1^z$  and  $\lambda_1^{z'}$  are essentially orthogonal as long as  $|z - z'| \gg n^{-1/2}$ . This fact is proven by controlling products of resolvents  $G^z G^{z'}$  for spectral parameters  $\eta$  slightly above the eigenvalue spacing.

This leads us to the *second key new ingredient* of the current work: *multi-resolvent local law* involving  $G^z$  and  $G^{z'}$  with optimal  $|z - z'|$ -dependence. We point out that this is the first instance when such optimal  $|z - z'|$ -decay has been proven and this optimality is essential for obtaining the necessary eigenvector orthogonality. The simplest prototype of a multi-resolvent local law is to understand how  $\text{Tr}G^z(i\eta)[G^{z'}(i\eta)]^*$  behaves when  $|z - z'| \gg n^{-1/2}$  since this quantity governs the covariance of  $\text{Tr}G^z(i\eta)$  and  $\text{Tr}G^{z'}(i\eta)$ . When  $z = z'$ , then a simple resolvent identity (also called the *Ward-identity* in this context) shows that  $G^z(i\eta)[G^z(i\eta)]^* = \eta^{-1} \text{Im} G^z(i\eta)$ , i.e. a product of two resolvents can be reduced to one resolvent. Roughly speaking, a multi-resolvent local law for  $z = z'$  can be reduced to a single resolvent local law that is much better understood. However, for  $z \neq z'$  the spectral resolutions of  $H^z$  and  $H^{z'}$  are different and such reduction is not possible; a new idea is needed that we will explain in Section 1.4.

In fact, we will also need another multi-resolvent local law for  $G^z F G^z F^* G^z$  to handle the larger  $\eta$  regime in (1.4) with an improvement owing to the special choice of  $F$ . The key phenomenon that  $GAG$  is much smaller if the deterministic  $A$  is chosen in a special subspace has been first discovered in [54] for the resolvent  $G$  of a Wigner matrix. Later it has been extended to more general situations [55, 56, 57, 49, 47] by developing the method of (static) *master inequalities* for the fluctuations in the local law. However, here we need to exploit this phenomenon in the cusp regime for  $G^z$  which would be especially cumbersome via the master inequalities. Our new dynamical approach to multi-resolvent local laws also solves this problem in a short and elegant way.

**1.4. Dynamical approach to multi-resolvent local laws.** Multi-resolvent local laws have recently been derived in various situations; they are necessary to prove the central limit theorem for linear eigenvalue statistics [86, 95, 112, 79, 90, 89, 52, 55, 58, 91, 93, 117, 106, 107] and the Eigenstate Thermalisation Hypothesis [54, 47, 1]. Their canonical proof is to derive an approximate self-consistent equation, e.g. for  $\text{Tr}G^z[G^{z'}]^*$  for definiteness, then to compute the fluctuation error in this equation by a cumulant expansion and invert the equation. In certain cases, including  $\text{Tr}G^z[G^{z'}]^*$ , the self-consistent equation is very unstable which is manifested by the smallness of the explicitly computable smallest eigenvalue of its linear stability operator. This instability is offset by a cancellation mechanism on the fluctuation side that is effective exactly in the "worst direction" of the stability operator. Such cancellation is algebraically visible when Ward identity is available, but otherwise it remains very mysterious. One expects

an "effective Ward-like identity" operating in the background, but it has never been found in a sufficiently robust form. The interested reader is encouraged to work out the simplest instance of this phenomenon: prove the local law for  $G(z)$  and for  $G(z)G^*(z)$ , where  $G(z) = (W - z)^{-1}$  is the resolvent of the standard (Hermitian) Wigner matrix  $W$  following the cumulant expansion strategy, e.g. in [86, 79, 56]. The procedure applied directly for  $GG^*$  will give a bound that is off by a factor  $1/|\operatorname{Im} z| \gg 1$ . If, however, one first uses the Ward identity,  $GG^* = \frac{1}{\operatorname{Im} z} \operatorname{Im} G$ , and then applies the procedure for  $\operatorname{Im} G$ , then it gives the optimal result.

In this paper we present a different solution to this "missing Ward-identity" problem that does not require to invert the stability operator at all in its unstable direction: we apply the method of *characteristic flow* for a critical multi-resolvent situation. The basic idea to estimate resolvents *dynamically* appeared first in [92] in the context of edge universality. Here we consider a matrix valued Ornstein-Uhlenbeck process  $X_t$  with initial value  $X_0 = X$ , denote by  $G_t$  its Hermitized resolvent after a  $z$ -shift. We also let both spectral parameters  $z$  and  $\eta$  evolve in time in such a way that the time derivative of  $\operatorname{Tr} G_t^{z_t}(i\eta_t)[G_t^{z_t}(i\eta_t)]^*$  has an algebraic cancellation among its leading terms. The corresponding equation for the spectral parameters is called the characteristic flow, the name originates from the fact that in the simplest case of the single resolvent of a Wigner matrix this equation describes the characteristics of the underlying complex Burgers equation [34, 81, 105, 124]. Along this flow  $i\eta_t$  approaches the real axis, thus one can effectively compare the resolvents at small  $\eta$  with resolvents at large  $\eta$  where local laws are much easier to prove since the stability operator has no unstable direction. The price one pays for this convenience is the added Gaussian component to  $X_t$ , but this can later be removed by fairly standard *Green function comparison theorems (GFT)*.

The method of characteristic flow has been effectively used for local laws and closely related quantities in various models in [81, 2, 89, 3, 91, 35, 34, 92] but only for a single resolvent situation where it eventually leads to a linear equation thanks to the Ward identity. Two or more resolvents with dynamical method were considered only in [58, Theorem 3.3], [35, Proposition 4.5], [47, Proposition 3.3] and very recently in [118, Lemma 4.2]. The latter three works focused on a different, less delicate aspects and exhibited no correlation decay that would have been the analogue of our  $|z - z'|$ -decay. In [58] a local law for  $\operatorname{Tr} G^z(i\eta)G^{z'}(i\eta')$  was proven in the bulk regime (i.e. when  $|z|, |z'| \leq 1 - \epsilon$ ) with an error term that exhibited a small improvement when  $|z - z'| \gg n^{-1/2}$ . The decay of the error bound in  $|z - z'|$  was very far from being optimal. Our key result handles the much more critical edge regime (i.e. when  $|z|, |z'| \approx 1$ ) and it proves the optimal decay in  $|z - z'|$ ; mastering both features that are essential for proving the Gumbel distribution. It has been observed earlier [16] that in the edge regime,  $|z| \approx 1$ , the eigenvalue density of the Hermitised matrix  $H^z$  has a cusp like singularity near the origin and the stability operator has *two* small eigenvalues, hence two "bad directions". Therefore our current analysis requires to monitor all four quantities of the form  $\operatorname{Tr} G^z(i\eta)A_1 G^{z'}(i\eta')A_2$ , where the deterministic matrices  $A_1, A_2$  lie in one or the other bad direction.

There are two key reduction steps in this analysis. First, the evolution equation for a two-resolvent quantity, schematically like  $\operatorname{Tr} GAGAGA$ , naturally involves a three- and four-resolvent quantities of the form  $\operatorname{Tr} GAGAGAGA$  (see (5.14) and (5.28) later), whose own evolution equation involves eight resolvents etc. This would lead to an infinite hierarchy of equations, reminiscent to the well-known BBGKY-hierarchy from statistical physics for correlation functions of interacting particle systems. Such hierarchy is uncontrollable without a *closure* step, e.g.  $\operatorname{Tr} GAGAGAGA$  needs to be estimated by the square of  $\operatorname{Tr} GAGAGA$ . For this purpose we rely on a *reduction inequality* (see (5.30) below), this leads to a closed coupled system of four *nonlinear* differential equations plus an error term (its linearised form is given in (5.47) later).

Second, the linearised part of this system blows up exactly when the characteristic flow reaches the real axis and we need to compute the precise rate of the blow-up as it gives the optimal estimate in the local law, which is in fact necessary to obtain the desired  $|z - z'|$ -gain. We then need to control the higher order nonlinear terms and show that they remain negligible in the entire range of  $\eta$ 's above the eigenvalue spacing, i.e. where local laws are expected to hold. Eventually the nonlinear terms would prevail, so with the terminology of nonlinear evolution equations our problem is strictly speaking *supercritical*, but with an effective tiny cutoff provided by the underlying discreteness of spectrum.

In summary, the key methodological novelty of our work on multi-resolvent local laws is to identify the nonlinear system of equations that governs the evolution of the multi-resolvent local laws along the characteristic flow in the critical cusp regime. For brevity, we present its analysis in our special situation

of two resolvents of the Hermitisation of a complex i.i.d. matrix that is needed to prove the Gumbel law, but the method is very general; it is applicable to prove any multi-resolvent local law for any mean-field matrix ensemble, which will be fundamental in the analysis of eigenvalues and eigenvectors of both Hermitian and non-Hermitian matrices. Special cases of multi-resolvent local laws have already played a key role in several recent works with broader scope [99, 96, 100] and we expect their impact will continue to grow as they become available for more general ensembles. In particular, the necessary precise explicit calculations near the blow-up point can be done systematically based upon the underlying *Dyson equation* (3.3) and the corresponding characteristic flow (5.3), which hold for very general random matrix ensembles.

**Acknowledgement.** We thank Oleksii Kolupaiev for several critical remarks about the draft version of this paper.

**Conventions and notations.** For integers  $k, l \in \mathbb{N}$ , with  $k < l$ , we use the notations  $[k, l] := \{k, \dots, l\}$  and  $[k] := [1, k]$ . For any  $z \in \mathbb{C}$  we use  $d^2z := \frac{1}{2}i(dz \wedge d\bar{z})$  for the two dimensional area form on  $\mathbb{C}$  and  $\Delta_z := 4\partial_z\partial_{\bar{z}}$  denotes the Laplace operator on  $\mathbb{C}$ . For positive quantities  $f, g$  we write  $f \lesssim g$  and  $f \sim g$  if  $f \leq Cg$  or  $cg \leq f \leq Cg$ , respectively, for some constants  $c, C > 0$  which depend only on the constants appearing in (2.1). We set  $f \wedge g = \min\{f, g\}$  and  $f \vee g = \max\{f, g\}$ . Furthermore, for  $n$ -dependent positive quantities  $f_n, g_n$  we use the notation  $f_n \ll g_n$  to denote that  $\lim_{n \rightarrow \infty} (f_n/g_n) = 0$ . Throughout the paper  $c, C > 0$  denote small and large constants, respectively, which may change from line to line. We denote vectors by bold-faced lower case Roman letters  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^d$ , for some  $d \in \mathbb{N}$ . Vector and matrix norms,  $\|\mathbf{x}\|$  and  $\|A\|$ , indicate the usual Euclidean norm and the corresponding induced matrix norm. For any  $d \times d$  matrix  $A$  we use  $\langle A \rangle := d^{-1}\text{Tr}A$  to denote the normalized trace of  $A$ . Moreover, for vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^d$  we define the usual scalar product  $\langle \mathbf{x}, \mathbf{y} \rangle := \sum_{i=1}^d \bar{x}_i y_i$ . Furthermore,  $\|f\|_\infty$  and  $\|f\|_1$  denote the  $L^\infty$  and  $L^1$ -norm of a function  $f$ , respectively. Throughout the paper we will use the notations  $\mathbf{P}^{\text{Gin}}$  and  $\mathbf{E}^{\text{Gin}}$  to denote the corresponding probability and expectation for  $X$  being a complex Ginibre matrix.

We define the following special  $2n \times 2n$  matrices

$$F := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad E_1 := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad E_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad E_\pm := E_1 \pm E_2 = \begin{pmatrix} 1 & 0 \\ 0 & \pm 1 \end{pmatrix}, \quad (1.5)$$

in particular,  $E_+$  is the  $2n$ -dimensional identity matrix.

We will often use the concept of “with very high probability” for an  $n$ -dependent event, meaning that for any fixed  $D > 0$  the probability of the event is bigger than  $1 - n^{-D}$  if  $n \geq n_0(D)$ . Moreover, we use the convention that  $\xi > 0$  denotes an arbitrary small positive exponent which is independent of  $n$ . We recall the standard notion of *stochastic domination*: given two families of non-negative random variables

$$X = \left( X^{(n)}(u) : n \in \mathbb{N}, u \in U^{(n)} \right) \quad \text{and} \quad Y = \left( Y^{(n)}(u) : n \in \mathbb{N}, u \in U^{(n)} \right)$$

indexed by  $n$  (and possibly some parameter  $u$  in some parameter space  $U^{(n)}$ ), we say that  $X$  is *stochastically dominated* by  $Y$ , if for any  $\xi, D > 0$  we have

$$\sup_{u \in U^{(n)}} \mathbf{P} \left[ X^{(n)}(u) > n^\xi Y^{(n)}(u) \right] \leq n^{-D} \quad (1.6)$$

for large enough  $n \geq n_0(\xi, D)$ . In this case we use the notation  $X \prec Y$  or  $X = O_\prec(Y)$  and we say that  $X(u) \prec Y(u)$  holds uniformly in  $u \in U^{(n)}$ .

## 2. MAIN RESULTS

We consider *complex i.i.d. matrices*  $X \in \mathbb{C}^{n \times n}$ , i.e.  $n \times n$  matrices with independent and identically distributed (i.i.d.) entries  $x_{ab} \stackrel{d}{=} n^{-1/2}\chi$ , for some complex random variable  $\chi$ . If  $\chi$  is a standard complex Gaussian random variable, then  $X$  is called a (*complex*) *Ginibre matrix*. In general, we make the following assumptions on  $\chi$ :

**Assumption 2.1.** We assume that  $\mathbf{E}\chi = 0$ ,  $\mathbf{E}|\chi|^2 = 1$  and  $\mathbf{E}\chi^2 = 0$ . Furthermore, we require that higher moments exist<sup>2</sup>, i.e. that for any  $p \in \mathbb{N}$  there exists a constant  $C_p > 0$  such that

$$\mathbf{E}|\chi|^p \leq C_p. \quad (2.1)$$

<sup>2</sup>Here, for simplicity, we assume that all moment exists, but inspecting our proof it is clear that we need to formulate this assumption only for any  $p \leq p_0$ , for a certain fixed  $p_0 \in \mathbb{N}$ .

We denote the eigenvalues of  $X$  by  $\{\sigma_i\}_{i \in [n]}$ , and label them so that<sup>3</sup>

$$|\sigma_1| \geq |\sigma_2| \geq \cdots \geq |\sigma_n|. \quad (2.2)$$

**2.1. Spectral radius.** The spectral radius of  $X$  is given by  $\rho(X) = |\sigma_1|$ . Note that  $|\sigma_1| > |\sigma_2|$  with high probability (see Lemma 4.4 later), so the spectral radius comes from a unique eigenvalue. If  $\chi$  has a density function, then  $|\sigma_1| > |\sigma_2|$  holds even almost surely.

Our first main result is the universality of the joint fluctuations of  $|\sigma_1|$  and its argument:

**Theorem 2.2** (Spectral radius). *Let  $X$  be a complex i.i.d. matrix satisfying Assumption 2.1. Set*

$$\gamma_n := \log n - 2 \log \log n - \log 2\pi. \quad (2.3)$$

Then for any fixed  $r \in \mathbb{R}$  and  $\theta \in [0, 2\pi)$ , we have<sup>4</sup>

$$\lim_{n \rightarrow \infty} \mathbf{P}\left(|\sigma_1| \leq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4n\gamma_n}}, \quad 0 \leq \arg(\sigma_1) \leq \theta\right) = \frac{\theta}{2\pi} e^{-e^{-r}}. \quad (2.4)$$

Note that (2.4) proves the universality of the joint distribution of the largest eigenvalue (in absolute value) and its argument. In particular, by (2.4) it follows that  $\rho(X) = |\sigma_1|$  has Gumbel fluctuation:

$$\lim_{n \rightarrow \infty} \mathbf{P}\left(|\sigma_1| \leq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4n\gamma_n}}\right) = e^{-e^{-r}}. \quad (2.5)$$

Furthermore, (2.4) also implies that asymptotically the law of  $\arg(\sigma_1)$  is uniform and independent of  $|\sigma_1|$ :

$$\lim_{n \rightarrow \infty} \mathbf{P}\left(\arg(\sigma_1) \in [0, \theta]\right) = \frac{\theta}{2\pi}. \quad (2.6)$$

**Remark 2.3.** For simplicity we stated (2.2) only for the largest eigenvalue in absolute value, however, inspecting our proof in Section 4, it is clear that we can also prove universality of the joint distribution of the largest  $k$  eigenvalues and their arguments for any finite  $k \in \mathbb{N}$ .

**Remark 2.4** (Convergence of moments). *Our method proves not only the convergence in distribution to Gumbel, but also the convergence of all moments. More precisely, define the rescaled quantity*

$$Y_n := \sqrt{4n\gamma_n} \left[ |\sigma_1| - 1 - \sqrt{\frac{\gamma_n}{4n}} \right],$$

then, for any fixed  $p$ , we have that  $\mathbf{E}|Y_n|^p$  converges to the  $p$ -th moment of a standard Gumbel random variable. A similar result applies to all the joint moments of the first  $k$  largest eigenvalues (in absolute value) and their arguments. More details will be given in Remark 4.3 and in Section 7.3.

Our second main result is to prove that the eigenvalues (in modulus) around  $1 + \sqrt{\gamma_n/4n}$  with the scaling  $\sqrt{4n\gamma_n}$  form an inhomogeneous Poisson point process in the complex plane in the  $n \rightarrow \infty$  limit, and we also identify its intensity function.

**Theorem 2.5** (Poisson Point Process). *Under the conditions of Theorem 2.2, we rescale the eigenvalues*

$$\sigma_j = e^{i\theta_j} \left( 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r_j}{\sqrt{4\gamma_n n}} \right), \quad r_j \in \mathbb{R}, \quad \theta_j \in [0, 2\pi). \quad (2.7)$$

Fix any  $t \in \mathbb{R}$  and any non-negative function  $g \in C_c^2(\mathbb{C})$  supported on  $\{z = re^{i\theta} : r \in [t, \infty), \theta \in [0, 2\pi)\}$ . Then, we have

$$\lim_{n \rightarrow \infty} \mathbf{E} \left[ e^{-\sum_{j=1}^n g(r_j e^{i\theta_j})} \right] = \exp \left( - \int_0^{2\pi} \int_{\mathbb{R}} (1 - e^{-g(re^{i\theta})}) \frac{e^{-r}}{2\pi} dr d\theta \right). \quad (2.8)$$

<sup>3</sup>We chose this labelling purely for notational convenience. All our results are insensitive to the labelling; in fact, we think of the eigenvalues as an unlabelled point process.

<sup>4</sup>We formulated the convergence as a limiting statement, but inspecting our proof we can also obtain an effective speed of convergence  $O((\log \log n)^2 / (\log n))$  in (2.4) uniformly in  $(r, \theta)$  similarly to [59, Theorem 1].

**2.2. Rightmost eigenvalue.** Motivated by the long time behavior of the linear system of ODE's  $\dot{\mathbf{u}} = X\mathbf{u}$ , in [60] we computed the precise size of the real part of the rightmost eigenvalue of  $X$ . However, in [60] we were not able to identify the fluctuations of this quantity around its deterministic limit. Similarly to Theorems 2.2 and 2.5 now we also prove the following universality result:

**Theorem 2.6** (Rightmost eigenvalue). *Under the conditions of Theorem 2.2, for any fixed  $x \in \mathbb{R}$  we have*

$$\lim_{n \rightarrow \infty} \mathbf{P} \left( \max_{j \in [n]} \operatorname{Re} \sigma_j \leq 1 + \sqrt{\frac{\gamma'_n}{4n}} + \frac{x}{\sqrt{4n\gamma'_n}} \right) = e^{-e^{-x}}, \quad (2.9)$$

where  $\gamma'_n$  is given by

$$\gamma'_n := \frac{\log n - 5 \log \log n - \log(2\pi)^4}{2}. \quad (2.10)$$

Furthermore, rescaling the eigenvalues

$$\sigma_j = 1 + \sqrt{\frac{\gamma'_n}{4n}} + \frac{x_j}{\sqrt{4\gamma'_n n}} + \frac{iy_j}{(\gamma'_n n)^{1/4}}, \quad x_j, y_j \in \mathbb{R},$$

and fixing any  $t \in \mathbb{R}$  and any non-negative function  $g \in C_c^2(\mathbb{C})$  supported on  $[t, \infty) \times i\mathbb{R}$ , we have

$$\lim_{n \rightarrow \infty} \mathbf{E} \left[ e^{-\sum_{j=1}^n g(x_j + iy_j)} \right] = \exp \left( - \int_{\mathbb{R}} \int_{\mathbb{R}} (1 - e^{-g(x+iy)}) \frac{e^{-x-y^2}}{\sqrt{\pi}} dy dx \right). \quad (2.11)$$

We point out that a similar result holds for the largest eigenvalue in any given fixed direction, we stated only the result for the eigenvalue with largest real part because it is the most relevant direction for the application to the stability theory of the ODE  $\dot{\mathbf{u}} = X\mathbf{u}$ .

### 3. MAIN TECHNICAL RESULTS: MULTI-RESOLVENT LOCAL LAWS AT THE SPECTRAL EDGE OF $X$

In this section we present the main technical results which are needed to prove the results in Section 2. For any  $z \in \mathbb{C}$ , recall the definition of the resolvent  $G^z$  of the matrix  $H^z$  at spectral parameter  $w$  from (1.3). The  $2n \times 2n$  matrix  $H^z$  is known as the *Hermitization* of  $X - z$ . The  $2 \times 2$  block structure (*chiral symmetry*) of  $H^z$  yields a spectrum that is symmetric with respect to zero, i.e. the eigenvalues of  $H^z$  can be labelled as  $\{\lambda_{\pm i}^z\}_{i \in [n]}$  such that  $0 \leq \lambda_1^z \leq \lambda_2^z \leq \dots \leq \lambda_n^z$  and  $\lambda_{-i}^z = -\lambda_i^z$ . Note that  $\{\lambda_i^z\}_{i \in [n]}$  are the singular values of  $X - z$ . Denote by  $\mathbf{w}_i^z$  and  $\mathbf{w}_{-i}^z$  the orthonormal eigenvector of  $H^z$  corresponding to the eigenvalue  $\lambda_i^z$  and  $\lambda_{-i}^z$ , then, as a consequence of the chiral symmetry, the eigenvectors naturally split into two  $n$ -vectors in the form

$$\mathbf{w}_{\pm i}^z = \begin{pmatrix} \mathbf{u}_i^z \\ \pm \mathbf{v}_i^z \end{pmatrix}, \quad i \in [n],$$

where  $\mathbf{u}_i^z, \mathbf{v}_i^z$  are the left and right singular vectors of  $X - z$ , respectively, normalized so that  $\|\mathbf{u}_i^z\|^2 = \|\mathbf{v}_i^z\|^2 = 1/2$ . Furthermore, for spectral parameter  $w = i\eta$ ,  $\eta \in \mathbb{R} \setminus \{0\}$ , on the imaginary axis, we have

$$G_{vv}^z(i\eta) = i \operatorname{Im} G_{vv}^z(i\eta), \quad \eta \operatorname{Im} G_{vv}^z(i\eta) > 0, \quad v \in [1, 2n].$$

In this section we prove *local laws*, i.e. concentration statements asserting that single resolvents and certain products of them become deterministic in the large  $n$  limit even if the spectral parameter  $\eta$  is very small. All results are formulated in the *cusp regime*, i.e. for  $|z| \approx 1$ , where the limiting density of states of  $H^z$  has a cusp-like singularity at the origin. The cusp is technically the most complicated regime but this is necessary to analyse the spectral properties of  $X$  near its spectral edge, the unit circle. In Section 3.1 we recall the *averaged* and *isotropic* single resolvent laws for  $G^z$  from [61, Theorem 3.1]. In Section 3.2 we pick  $A_1, A_2 \in \{E_+, E_-\}$ , with  $E_{\pm}$  being defined in (1.5), and we prove a new local law for  $\langle G^{z_1} A_1 G^{z_2} A_2 \rangle$  with spectral parameters on the imaginary axis, exhibiting that the size of  $\langle G^{z_1} A_1 G^{z_2} A_2 \rangle$  decays as  $|z_1 - z_2|$  increases. Then in Section 3.3 we consider another mechanism to improve local laws: we show that if  $A_1, A_2 \in \{F, F^*\}$ , with  $F$  defined in (1.5), then  $\langle G^z A_1 \rangle$  and  $\langle G^z A_1 G^z A_2 \rangle$  are considerably smaller than for general  $A_1, A_2$ . The fact that the resolvent tested against certain matrices is smaller than expected was first observed in [54] for Wigner matrices and later extended to more general models in the bulk of the spectrum (see e.g. [1, 48, 49]) and at the spectral edges (see [47]). However, Theorem 3.5 below is the first result when we exhibit this effect at the cusp singularity of the spectral density. Finally, in Section 3.4 we first show that the smallest singular values of  $X - z_1, X - z_2, \dots, X - z_k$ , with a fix  $k \in \mathbb{N}$ , are asymptotically independent as long as  $|z_i - z_j| \gg n^{-1/2}$ , and then we recall a probabilistic



tail bound for the smallest singular value of  $X - z$  when  $z$  is outside of the spectrum (i.e.  $|z| > 1$ ) from [61].

**3.1. Local law for  $G^z$ .** We first define the deterministic approximation of  $G^z(w)$  as follows:

$$M^z(w) := \begin{pmatrix} m^z(w) & -zu^z(w) \\ -\bar{z}u^z(w) & m^z(w) \end{pmatrix}, \quad \text{with} \quad u^z(w) := \frac{m^z(w)}{w + m^z(w)}, \quad (3.1)$$

where  $m^z(w)$  is the unique solution of the scalar cubic equation with side condition

$$-\frac{1}{m^z(w)} = w + m^z(w) - \frac{|z|^2}{w + m^z(w)}, \quad \text{Im}[m^z(w)]\text{Im}w > 0. \quad (3.2)$$

We consider  $M^z(w)$  as a  $(2n) \times (2n)$  matrix which is block constant, i.e. it consists of four  $n \times n$  blocks and each of them is a constant multiple of the  $n \times n$  identity matrix. In fact,  $M^z$  is the unique solution of the *Matrix Dyson equation* (see e.g. [8])

$$-[M^z(w)]^{-1} = w + Z + \mathcal{S}[M^z(w)], \quad Z := \begin{pmatrix} 0 & z \\ \bar{z} & 0 \end{pmatrix}, \quad (3.3)$$

with the side condition  $\text{Im} M^z(w) \cdot \text{Im} w > 0$ . Here

$$\mathcal{S}[\cdot] := \langle \cdot E_+ \rangle E_+ - \langle \cdot E_- \rangle E_-, \quad \mathcal{S} : \mathbb{C}^{2n \times 2n} \rightarrow \mathbb{C}^{2n \times 2n}, \quad (3.4)$$

is the *covariance operator* of the Hermitization of  $X$ . Additionally, we define the *self-consistent eigenvalue density* of  $H^z$  by (see e.g. [14])

$$\rho^z(E) := \lim_{\eta \rightarrow 0^+} \frac{1}{\pi} \text{Im} \langle M^z(E + i\eta) \rangle, \quad E \in \mathbb{R}. \quad (3.5)$$

In the following we collect properties of the density  $\rho^z(E)$  as summarized in [61, Section 3.1] (based upon results from [64, 16, 15, 8]):

- 1) if  $|z| < 1$ , then the density  $\rho^z(E)$  has a local minimum at  $E = 0$  of height  $\rho^z(0) \sim (1 - |z|^2)^{1/2}$ ;
- 2) if  $|z| = 1$ , then the density has an exact cubic cusp singularity at  $E = 0$ ;
- 3) if  $|z| > 1$ , then there exists a small gap in the support of the symmetric density  $\rho^z(E)$ , denoted by  $[-\frac{\Delta}{2}, \frac{\Delta}{2}]$  with  $\Delta \sim (|z|^2 - 1)^{3/2}$ ; see Figure 1.

Moreover, we also recall the following quantitative properties of the density  $\rho^z(w) := \pi^{-1} |\text{Im} \langle M^z(w) \rangle|$  extended to the complex plane,  $w \in \mathbb{C}$ . If  $|z| \leq 1$ , then for any  $w = E + i\eta$ , we have

$$\rho^z(w) \sim (1 - |z|^2)^{1/2} + (|E| + \eta)^{1/3}, \quad |E| \leq c, \quad 0 \leq \eta \leq 1,$$

for some small  $c > 0$ , while in the complementary regime  $|z| > 1$ , for any  $w = \frac{\Delta}{2} + \kappa + i\eta$ , we have

$$\rho^z(w) \sim \begin{cases} (|\kappa| + \eta)^{1/2} (\Delta + |\kappa| + \eta)^{-1/6}, & \kappa \in [0, c] \\ \frac{\eta}{(\Delta + |\kappa| + \eta)^{1/6} (|\kappa| + \eta)^{1/2}}, & \kappa \in [-\Delta/2, 0], \end{cases} \quad 0 \leq \eta \leq 1. \quad (3.6)$$

Furthermore, on the imaginary axis  $m^z(i\eta) = \langle M^z(i\eta) \rangle$  is purely imaginary and  $u^z(i\eta)$  defined in (3.1) is real. We may often drop the superscript  $z$  and the argument  $w = i\eta$  for brevity, i.e., we write  $m = m^z(i\eta)$ , and

$$\rho := \frac{|\text{Im} m|}{\pi} \sim \begin{cases} \eta^{1/3} + |1 - |z|^2|^{1/2}, & |z| \leq 1 \\ \frac{\eta}{|1 - |z|^2| + \eta^{2/3}}, & |z| > 1, \end{cases} \quad 0 \leq \eta \leq 1. \quad (3.7)$$

We are now ready to recall the following optimal local law for the resolvent  $G^z$  near the cusp.

**Theorem 3.1** (Theorem 3.1 in [61]). *There are sufficiently small constants  $c, c' > 0$  such that for any  $z \in \mathbb{C}$  with  $||z| - 1| \leq c$  and for any  $w = E + i\eta$  with  $|E| \leq c'$  we have*

$$|\langle \mathbf{x}, (G^z(w) - M^z(w)) \mathbf{y} \rangle| \prec \|\mathbf{x}\| \|\mathbf{y}\| \left( \sqrt{\frac{\rho^z(w)}{n|\eta|}} + \frac{1}{n|\eta|} \right), \quad (3.8)$$

$$|\langle (G^z(w) - M^z(w)) A \rangle| \prec \frac{\|A\|}{n|\eta|}, \quad (3.9)$$

uniformly in spectral parameter  $|\eta| > 0$ , deterministic vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^{2n}$ , and matrices  $A \in \mathbb{C}^{2n \times 2n}$ .




FIGURE 1. For  $z$  in the regime  $|z| > 1$  slightly beyond the unit disk, the picture on the right shows the corresponding density  $\rho^z$  of  $H^z$ , which exhibits a small gap around zero of size  $\Delta \sim (|z|^2 - 1)^{3/2}$ ; see the paragraph below [16, Eq. (5.2)].

By standard arguments, see e.g. [7, Corollary 1.10-1.11], using the local law in (3.9) as an input, we readily conclude the following rigidity estimate for the eigenvalues  $\lambda_i^z$ . For this purpose we define the *quantiles*  $\gamma_i^z$  of  $\rho^z$  implicitly by the relation

$$\int_0^{\gamma_i^z} \rho^z(x) dx = \frac{i}{2n}, \quad \text{for } i \in [n]. \quad (3.10)$$

For negative indices we set  $\gamma_{-i}^z := -\gamma_i^z$ , with  $i \in [n]$ .

**Corollary 3.2** (Corollary 3.2 in [61]). *Fix a small constant  $c > 0$  and pick  $z$  such that  $0 \leq |z| - 1 \leq c$ . Then there exists a small  $c' > 0$  such that*

$$|\lambda_i^z - \gamma_i^z| < \max \left\{ \frac{1}{n^{3/4}|i|^{1/4}}, \frac{\Delta^{1/9}}{n^{2/3}|i|^{1/3}} \right\}, \quad \text{for } |i| \leq c'n, \quad (3.11)$$

with  $\Delta$  denoting the size of the gap around zero in the support of  $\rho^z$ . In addition there exists a small  $c'' > 0$  such that, for any  $E_1 < E_2$  with  $\max\{|E_1|, |E_2|\} \leq c''$ , we have

$$\left| \#\{j : E_1 \leq \lambda_j^z \leq E_2\} - 2n \int_{E_1}^{E_2} \rho^z(x) dx \right| < 1. \quad (3.12)$$

**3.2. Local law for  $G^{z_1} A G^{z_2}$  near the cusp.** We now consider the product of two resolvents  $G^{z_1} A G^{z_2}$ , with a deterministic matrix  $A \in \mathbb{C}^{2n \times 2n}$  in between, for  $z_1 \neq z_2$  close to but outside of the unit disk and with spectral parameters on the imaginary axis. This is the situation used in Girko's formula. Proving a local law for this quantity is quite challenging, because  $G^{z_1}$  and  $G^{z_2}$  belong to two different spectral families, i.e. no resolvent identity is available to linearize their product even if  $A$  were the identity. Before stating the local law for  $G^{z_1} A G^{z_2}$ , we define its deterministic approximation, which is *not* simply the product  $M^{z_1} A M^{z_2}$  as one might guess at first. The correct formula is given by

$$M_{12}^A = M_{12}^A(\eta_1, z_1, \eta_2, z_2) := \mathcal{B}_{12}^{-1} \left[ M^{z_1}(i\eta_1) A M^{z_2}(i\eta_2) \right], \quad (3.13)$$

where the *two-body stability operator*  $\mathcal{B}_{12} : \mathbb{C}^{2n \times 2n} \rightarrow \mathbb{C}^{2n \times 2n}$  is defined by

$$\mathcal{B}_{12}[\cdot] := 1 - M^{z_1}(i\eta_1) \mathcal{S}[\cdot] M^{z_2}(i\eta_2). \quad (3.14)$$

Furthermore, we have the following optimal bound (proven in Appendix A)

$$\left\| M_{12}^A(\eta_1, z_1, \eta_2, z_2) \right\| \lesssim \frac{\|A\|}{|z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}}, \quad (3.15)$$

using the short-hand notation  $m_i := m^{z_i}(i\eta_i)$  and  $\rho_i := \pi^{-1}|\text{Im } m_i|$  for any  $\eta_1, \eta_2 \neq 0$  and  $z_1, z_2 \in \mathbb{C}$ . We stress that we work in the regime where  $0 \leq |z_i| - 1 \ll 1$  and thus the eigenvalue density  $\rho^{z_i}$  has a cusp-like singularity. This fact strongly influences the form of (3.15) and its optimality. With our current method, a very similar (in fact easier) analysis would yield an optimal two-resolvent local law in the non-Hermitian bulk regime,  $|z_i| \leq 1 - \epsilon$ , as it was done non-optimally in [52, 58, 53]. In this paper, however, we focus on the regime  $|z_i| \geq 1$  needed for the results in Section 2.

We are now ready to formulate our first multi-resolvent local law, which has two parts. The first part is for random matrices with a substantial Gaussian component. The second part is for arbitrary random i.i.d. matrices, but instead of  $G^{z_1}, G^{z_2}$  we consider only their imaginary parts and we test  $\text{Im } G^{z_1} A \text{Im } G^{z_2}$  only against  $A$ . In both parts the test matrices are  $A \in \{E_+, E_-\}$ . This will be sufficient for our purposes and it simplifies the proof. In Section 3.2.1 we will also comment on the optimality and extensions of this theorem.

**Theorem 3.3** (Local law with  $|z_1 - z_2|$ -decorrelation). *Fix any small  $c, \epsilon > 0$ . Then, for any  $z_1, z_2$  with  $0 \leq |z_i| - 1 \leq c$ , the following statements hold uniformly in spectral parameters  $\eta_1, \eta_2$  satisfying  $n\ell \geq n^\epsilon$ , where  $\ell := \min_i(|\eta_i|\rho_i)$ .*

**Part A:** *Let  $X$  be a random matrix satisfying Assumption 2.1, and let  $X^{\text{Gin}}$  be a complex Ginibre matrix independent of  $X$ . Fix any small  $\mathfrak{s} > 0$ , consider a matrix of the form  $X_{\mathfrak{s}} := \sqrt{1 - \mathfrak{s}^2}X + \mathfrak{s}X^{\text{Gin}}$ , and denote by  $G_{\mathfrak{s}}^z(w)$  the resolvent of the Hermitization of  $X_{\mathfrak{s}} - z$  defined as in (1.3). Then we have*

$$\left| \left\langle (G_{\mathfrak{s}}^{z_1}(i\eta_1)A_1G_{\mathfrak{s}}^{z_2}(i\eta_2) - M_{12}^{A_1})A_2 \right\rangle \right| \prec \left( \frac{1}{(n\ell)^{1/2}} \frac{1}{|z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}} \right) \wedge \left( \frac{1}{n|\eta_1\eta_2|} \right), \quad (3.16)$$

for deterministic matrices  $A_1, A_2 \in \{E_+, E_-\}$ , with  $M_{12}^{A_1} = M_{12}^{A_1}(\eta_1, z_1, \eta_2, z_2)$  defined in (3.13).

**Part B:** *Consider any i.i.d. random matrix  $X$  satisfying Assumption 2.1. Then we have*

$$\left| \left\langle (\text{Im } G^{z_1}(i\eta_1)A \text{Im } G^{z_2}(i\eta_2) - \widehat{M}_{12}^A)A \right\rangle \right| \prec \left( \frac{1}{(n\ell)^{1/2}} \frac{1}{|z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}} \right) \wedge \left( \frac{1}{n|\eta_1\eta_2|} \right), \quad (3.17)$$

uniformly in comparable spectral parameters  $|\eta_1| \sim |\eta_2|$ ,  $\rho_1 \sim \rho_2$ , and deterministic matrix  $A \in \{E_+, E_-\}$ . Here the deterministic approximation  $\widehat{M}_{12}^A$  is naturally defined as a fourfold linear combination of  $M_{12}^A(\pm\eta_1, z_1, \pm\eta_2, z_2)$ , using the identity  $\text{Im } G = \frac{1}{2i}(G - G^*)$  twice.

In particular, the error term in the local law (3.16) for  $G_{\mathfrak{s}}^{z_1}A_1G_{\mathfrak{s}}^{z_2}$  is always smaller than the bound (3.15) for the deterministic term  $M_{12}^{A_1}$  since  $n\ell \gg 1$ . Note that  $n|\eta_i|\rho_i$  is roughly the number of eigenvalues of  $H^{z_i}$  in a window of size  $\eta_i$  about the origin, thus the condition  $n\ell \gg 1$  guarantees that we are in the local law regime for both resolvents  $G_i = G^{z_i}(i\eta_i)$ , i.e. when concentration of  $G_1AG_2$  around a deterministic quantity is expected. Note that both estimates (3.16)–(3.17) involve the minimum of two terms; the first one exhibits a  $|z_1 - z_2|$ -decay, the second one is independent of  $z_1, z_2$ .

As a consequence of Theorem 3.3, we obtain the following upper bound on the overlaps between the eigenvectors  $\mathbf{w}_i^z = (\mathbf{u}_i^z, \mathbf{v}_i^z)$  for different  $z_1, z_2$ ; for simplicity we state this result only for  $z_1, z_2$  in the regime  $0 \leq |z_i| - 1 \leq n^{-1/2+\tau}$ .

**Corollary 3.4.** *Fix any small  $\tau > 0$ . Then there exists a small  $0 < \omega_c < \tau/10$  such that for any  $z_1, z_2$  with  $0 \leq |z_i| - 1 \leq n^{-1/2+\tau}$  we have*

$$|\langle \mathbf{u}_i^{z_1}, \mathbf{u}_j^{z_2} \rangle|^2 + |\langle \mathbf{v}_i^{z_1}, \mathbf{v}_j^{z_2} \rangle|^2 \prec \frac{n^{2\tau}}{\sqrt{n}[|z_1 - z_2| + n^{-1/2}]}, \quad |i|, |j| \leq n^{\omega_c}. \quad (3.18)$$

*Proof.* By the rigidity estimate of the eigenvalues in (3.11) and  $\Delta \sim (|z| - 1)^{3/2} \leq n^{-3/4+3\tau/2}$ , we have

$$|\lambda_i^z - \gamma_i^z| \prec \max \left\{ \frac{1}{n^{3/4}|i|^{1/4}}, \frac{\Delta^{1/9}}{n^{2/3}|i|^{1/3}} \right\} \lesssim \frac{n^{\tau/6}}{n^{3/4}|i|^{1/4}}. \quad (3.19)$$

Choose  $\eta = n^{-3/4+\tau}$ , then there exists  $0 < \omega_c < \tau/10$  such that  $|\lambda_i^z| \lesssim \eta$  for any  $|i| \leq n^{\omega_c}$  using (3.19) and the elementary bound  $|\gamma_i^z| \leq \eta$  from (3.6) and (3.10). Additionally, by the spectral decomposition,

we have

$$\langle \text{Im } G^{z_1}(i\eta) \text{Im } G^{z_2}(i\eta) \rangle = \frac{1}{n} \sum_{i,j} \frac{\eta^2 (|\langle \mathbf{u}_i^{z_1}, \mathbf{u}_j^{z_2} \rangle|^2 + |\langle \mathbf{v}_i^{z_1}, \mathbf{v}_j^{z_2} \rangle|^2)}{((\lambda_i^{z_1})^2 + \eta^2)((\lambda_j^{z_2})^2 + \eta^2)}, \quad (3.20)$$

where we used that  $\text{Im } G^z$  is diagonal. Thus for any  $|i|, |j| \leq n^{\omega_\epsilon}$ , we have

$$|\langle \mathbf{u}_i^{z_1}, \mathbf{u}_j^{z_2} \rangle|^2 + |\langle \mathbf{v}_i^{z_1}, \mathbf{v}_j^{z_2} \rangle|^2 \lesssim n\eta^2 \langle \text{Im } G^{z_1}(i\eta) \text{Im } G^{z_2}(i\eta) \rangle \prec \frac{n^{-1/2+2\tau}}{|z_1 - z_2|}, \quad (3.21)$$

where we used the local law in (3.17) with  $A = E_+$  and the deterministic bound in (3.15). This proves (3.18) for  $|z_1 - z_2| \geq n^{-1/2}$ . For the complementary regime  $|z_1 - z_2| \leq n^{-1/2}$ , we use the trivial bound  $|\langle \mathbf{u}_i^{z_1}, \mathbf{u}_j^{z_2} \rangle|^2 + |\langle \mathbf{v}_i^{z_1}, \mathbf{v}_j^{z_2} \rangle|^2 \leq 1$ , yielding (3.18).  $\square$

We conclude Section 3.2 by commenting on the optimality and restrictions in Theorem 3.3.

**3.2.1. Optimality of the  $\langle G^{z_1} A_1 G^{z_2} A_2 \rangle$  local law.** First, in Theorem 3.3 we stated the local law only for  $A_1, A_2 \in \{E_+, E_-\}$  as we need only this case for our application and its proof is shorter but with a similar method we can prove exactly the same bound for any deterministic  $A_1, A_2 \in \mathbb{C}^{2n \times 2n}$ . Actually, the proof for  $A_1, A_2 \in \{E_-, E_+\}$  is the most delicate since the stability operator  $\mathcal{B}_{12}$  has two small eigenvalues in  $\text{Span}\{E_-, E_+\}$ , while  $\mathcal{B}_{12}$  has eigenvalues equal to 1 on the complement of  $\text{Span}\{E_-, E_+\}$ .

Second, in the regime where we gain from the  $|z_1 - z_2|$ -factor, we only prove that the error term in the local laws (3.16)–(3.17) is smaller than the deterministic approximation by a factor  $(n\ell)^{-1/2}$ , however the optimal factor would be  $(n\ell)^{-1}$ . We do not pursue this improvement here since we will use the estimate in the regime  $n\ell \sim n^\epsilon$ , for some very small  $\epsilon > 0$ , where practically there is no difference between the two estimates. More importantly, however, the  $|z_1 - z_2|^{-1}$  decay in (3.16) is optimal.

In particular, we expect that the optimal form of (3.16) is

$$\left| \left\langle (G^{z_1}(i\eta_1) A_1 G^{z_2}(i\eta_2) - M_{12}^{A_1}) A_2 \right\rangle \right| \prec \|A_1\| \|A_2\| \left( \frac{1}{n\ell |z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}} \right) \quad (3.22)$$

in the regime  $0 \leq |z_i| - 1 \leq n^{-1/2+\tau}$ , and it holds for general  $A_1, A_2$  and for any i.i.d. matrix  $X$ . Note that once we replace  $(n\ell)^{-1/2}$  with  $(n\ell)^{-1}$  then the factor  $1/(n|\eta_1\eta_2|)$  in (3.16) becomes obsolete. The same bound also holds even if  $0 \leq |z_i| - 1 \leq c$ , but it is optimal only in the most relevant regime  $|z_i| \approx 1$ . If  $|z_i| - 1 > n^{-1/2+\tau}$  then the error becomes smaller since in this regime there are typically no eigenvalues close to zero [16, Theorems 2.1–2.2] as the limiting eigenvalue density has a gap  $\Delta$  of size  $\Delta \sim (|z|^2 - 1)^{3/2}$ . In fact, the basic estimate (3.15) on the deterministic term also improves for larger  $\Delta$ , but here we will not investigate the regime far outside of the spectrum as it is irrelevant for our main results in Section 2.

For technical convenience we stated (3.17) only for the special case  $|\eta_1| \sim |\eta_2|$  and  $\rho_1 \sim \rho_2$ , because this regime is enough for our main application in (3.18) and it allows for a simpler *Green function comparison (GFT)* argument in Appendix D to remove the Gaussian component in (3.16). This restriction can easily be relaxed using the stronger Lindeberg replacement approach for the GFT by finding a matrix, with a large Gaussian component, which matches the third moment and (almost) matches the fourth moment of  $X$ . The same strengthening of GFT is necessary to achieve (3.22) for a general i.i.d.  $X$  without Gaussian component.

Furthermore, we mention that even relaxing the assumption  $|\eta_1| \sim |\eta_2|$  and  $\rho_1 \sim \rho_2$  in (3.17) would not give an optimal bound (modulo the  $(n\ell)^{-1}$  improvement) as the optimal local law for the product of two  $\text{Im } G$ 's carries an additional smallness involving the local density  $\rho$  that appears because of considering  $\text{Im } G$  instead of  $G$  (see [47, Theorem 2.4] for a similar effect). We expect that

$$\left| \left\langle (\text{Im } G^{z_1}(i\eta_1) A_1 \text{Im } G^{z_2}(i\eta_2) - \widehat{M}_{12}^{A_1}) A_2 \right\rangle \right| \prec \frac{\rho_1 \rho_2}{n\ell} \frac{\|A_1\| \|A_2\|}{[|z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}]^2} \quad (3.23)$$

holds in the regime  $0 \leq |z_i| - 1 \leq c$  for general  $A_1, A_2$  and for any i.i.d. matrix  $X$ , and we expect this bound to be optimal when  $0 \leq |z_i| - 1 \leq n^{-1/2+\tau}$ . In particular the local law (3.23) has a  $|z_1 - z_2|^{-2}$  decay, this also reflects the fact that for the deterministic approximation we have

$$\|\widehat{M}_{12}^A\| \lesssim \frac{\|A\| \rho_1 \rho_2}{[|z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}]^2}. \quad (3.24)$$

We stress that our method to prove multi-resolvent local laws dynamically would enable us to prove (3.22) and (3.23) as well. For this purpose we would need to study the flow not only for traces  $\langle G^{z_1} A_1 G^{z_2} A_2 \rangle$  as we do now, but also for *isotropic* quantities  $\langle \mathbf{x}, G^{z_1} A_1 G^{z_2} \mathbf{y} \rangle$ . For conciseness we do not pursue these improvements here since we do not need them for our main results in Section 2.

Lastly, we note that the optimal local law (3.23) would imply the optimal overlap bound (cf. (3.18))

$$|\langle \mathbf{u}_i^{z_1}, \mathbf{u}_j^{z_2} \rangle|^2 + |\langle \mathbf{v}_i^{z_1}, \mathbf{v}_j^{z_2} \rangle|^2 \prec \frac{n^{2\tau}}{n[|z_1 - z_2| + n^{-1/2}]^2}, \quad |i|, |j| \leq n^{\omega_c}. \quad (3.25)$$

**3.3. Local law for  $G^z F$  and  $G^z F G^z F^*$  near the cusp.** We now explore a different mechanism to improve the single and multi-resolvent local laws, (3.9) and (3.16). We show that the size of the resolvent  $G^z(i\eta)$  (in tracial sense) is smaller by  $\rho$ -factors when tested against the special matrices  $F, F^*$  from (1.5).

**Theorem 3.5** (Local laws with  $F$ -improvement). *Fix any small  $c, \epsilon > 0$ . For any  $z \in \mathbb{C}$  with  $0 \leq |z| - 1 \leq c$ , we have*

$$\left| \langle (G^z(i\eta) - M^z(i\eta)) F^{(*)} \rangle \right| \prec \frac{\rho}{n|\eta|}, \quad (3.26)$$

$$\left| \langle G^z(i\eta) F G^z(i\eta) - M_{12}^F(i\eta) \rangle \right| \prec \frac{\rho}{n\eta^2}, \quad (3.27)$$

$$\left| \langle (G^z(i\eta) F G^z(i\eta) - M_{12}^F(i\eta)) F^* \rangle \right| \prec \frac{\rho^2}{n\eta^2} \sqrt{n|\eta|\rho}, \quad (3.28)$$

uniformly in spectral parameters  $n|\eta|\rho \geq n^\epsilon$ . Here  $\rho = \rho^z(i\eta)$  is given in (3.7), the deterministic matrices  $M^z$  and  $M_{12}^F$  are defined in (3.1) and (3.13) with  $z_1 = z_2 = z$ ,  $\eta_1 = \eta_2 = \eta$ , and  $A = F$ , respectively.

On the deterministic approximations  $\langle M_{12}^F(i\eta) \rangle$ ,  $\langle M_{12}^F(i\eta) F^* \rangle$  in (3.27) and (3.28), respectively, we have the bounds (see Lemma 5.9)

$$|\langle M_{12}^F(i\eta) \rangle| \lesssim \frac{\rho^2}{|\eta|} \quad \left| \langle M_{12}^F(i\eta) F^* \rangle \right| \lesssim \frac{\rho^3}{|\eta|}, \quad (3.29)$$

i.e. the error terms in (3.27)–(3.28) are smaller than the bounds (3.29) by a factor  $(n|\eta|\rho)^{-1}$  and  $(n|\eta|\rho)^{-1/2}$ , respectively. We point out that (3.26)–(3.27) are optimal, while (3.28) is off by a factor  $\sqrt{n|\eta|\rho}$ . Our proof method in Section 5 can clearly improve (3.28) to be optimal, but we omit this for brevity as we will not need it.

As a corollary of Theorem 3.5, we have the following optimal upper bound on the overlaps between the left and right singular vectors of  $X - z$ .

**Corollary 3.6.** *Fix any small  $\tau > 0$ . Then there exists a small  $0 < \omega_c < \tau/10$  such that, for any  $0 \leq |z| - 1 \leq n^{-1/2+\tau}$ , we have<sup>5</sup>*

$$|\langle \mathbf{u}_i^z, \mathbf{v}_j^z \rangle|^2 \prec \frac{n^{2\tau}}{\sqrt{n}}, \quad |i|, |j| \leq n^{\omega_c}. \quad (3.30)$$

We mention that this bound for  $i = j$  is optimal for  $z$  very close to the edge. In the bulk regime,  $|z| \leq 1 - \epsilon$ , the optimal bound is  $\prec 1/n$ , see [49, Theorem 2.7]. In other words, in the bulk regime the overlap of the left and right singular vectors behaves as if they were independent random vectors, while at the edge  $|\langle \mathbf{u}_i^z, \mathbf{v}_i^z \rangle|^2 \prec n^{-1/2}$  indicates a "half-way" between the full independence and the substantial alignment  $|\langle \mathbf{u}_i^z, \mathbf{v}_i^z \rangle|^2 \sim 1$  in the regime  $|z| \geq 1 + \epsilon$ .

*Proof.* Following the proof of Corollary 3.4, choose  $\eta = n^{-3/4+\tau}$  and recall that there exists  $\omega_c > 0$  such that  $|\lambda_i^z|, |\lambda_j^z| \lesssim \eta$ , for any  $|i|, |j| \leq n^{\omega_c}$ . Using the spectral decomposition of  $\text{Im } G^z$ , we have

$$|\langle \mathbf{u}_i^z, \mathbf{v}_j^z \rangle|^2 \lesssim n\eta^2 \langle G^z(i\eta) F G^z(i\eta) F^* \rangle \prec n\eta^2 \frac{\rho^3}{\eta} \lesssim n^{-1/2+2\tau}, \quad (3.31)$$

where we used the local law in (3.28) and the deterministic bound in (3.29) and (3.7) in the last step.  $\square$

<sup>5</sup>Similarly to [49, Theorem 2.4], the optimal bound (3.30) for singular vector overlap can be turned into an optimal lower bound on the overlap of non-Hermitian eigenvectors. More precisely, let  $\{L_i, R_i\}_{i \in [n]}$  be the left and right bi-orthogonal eigenvectors of  $X$  with corresponding eigenvalue  $\sigma_i$ . Then, for the diagonal overlap  $O_{ii} := \|L_i\|^2 \|R_i\|^2$  we have

$$O_{ii} \geq n^{1/2-\xi},$$

with very high probability for any small  $\xi > 0$ , where the index  $i \in [n]$  is such that  $|\sigma_i| \geq 1$ .

We conclude this section with several useful identities which will often be used throughout the paper. They follow from the chiral symmetry of  $H^z$ . Their elementary proof is presented in Appendix A.

**Lemma 3.7.** *Let  $H^z$  be defined as in (1.3), let  $G^z = (H^z - i\eta)^{-1}$ , and let  $M^z, M_{12}^F$  be defined as in (3.1) and (3.13) with  $z = z_1 = z_2 = z$ ,  $\eta_1 = \eta_2 = \eta$ , and  $A = F$ , respectively. Then we have*

$$\langle G^z F G^z E_- \rangle = 0, \quad \langle M_{12}^F E_- \rangle = 0, \quad \langle G^z E_- \rangle = 0, \quad \langle M^z E_- \rangle = 0, \quad F G^z F^* = i F \text{Im} G F^*. \quad (3.32)$$

**3.4. Smallest singular value.** In this section, we show that the lower tails of the smallest singular values of  $X - z_1, X - z_2, \dots, X - z_k$  are asymptotical independent for any finite collection  $\{z_j\}_{j=1}^k$  with mesoscopically distant values. This will follow by the almost orthogonality of the overlaps in (3.18) for  $|z_i - z_j| \gg n^{-1/2}$  together with the *Dyson Brownian Motion* argument from [61, Section 7]. A similar result was obtained in [61, Proposition 4.3] with  $|z_i - z_j| \geq n^{-\gamma}$  for a small  $\gamma > 0$  (that result was stated for  $k = 2$  only, the proof can be easily extended to a general  $k \geq 2$ ). The main difference is that there we required a sufficiently large, almost macroscopic distance between  $z_i$  and  $z_j$ . Now the improved eigenvector overlap estimates in Corollary 3.4 allow us to extend the statement to the optimal regime  $|z_i - z_j| \geq n^{-1/2+\omega}$  for any small  $\omega > 0$ . This is formulated in the following proposition whose proof is postponed to Appendix C.

**Proposition 3.8.** *Let  $X$  be a complex i.i.d. random matrix satisfying Assumption 2.1, and let  $\lambda_1^z$  be the smallest singular value of  $X - z$ . Fix any  $k \in \mathbb{N}$  and any sufficiently small  $\tau, \omega, \gamma > 0$  such that  $\tau \leq \gamma\omega/10$ . Then for any  $\{E_j\}_{j \in [k]}$  with  $0 < E_j \lesssim n^{-3/4}$ , for any  $\{z_j\}_{j \in [k]}$  with  $0 \leq |z_j| - 1 \leq n^{-1/2+\tau}$  and any  $|z_i - z_j| \geq n^{-1/2+\omega}$  ( $\forall i, j \in [k]$ ), we have*

$$\begin{aligned} \prod_{j=1}^k \mathbf{P}\left(\lambda_1^{z_j} \leq E_j - 4n^{-3/4-\gamma\omega+\tau}\right) - O(n^{-D}) &\lesssim \mathbf{P}\left(\lambda_1^{z_j} \leq E_j, \forall j \in [k]\right) \\ &\lesssim \prod_{j=1}^k \mathbf{P}\left(\lambda_1^{z_j} \leq E_j + 4n^{-3/4-\gamma\omega+\tau}\right) + O(n^{-D}), \end{aligned} \quad (3.33)$$

for any fixed (large)  $D > 0$  if  $n$  is sufficiently large.

Note that the lower bound above is effective only when  $E_j > n^{-3/4-\gamma\omega+\tau}$ . To estimate the upper and lower bounds in (3.33) precisely, we recall from [60, 61] the following tail bound for the smallest singular value of  $X - z$  for  $z$  being outside of the unit disk.

**Proposition 3.9** (Proposition 5.2 in [61]). *Let  $X$  be a complex i.i.d. random matrix satisfying Assumption 2.1, and let  $\lambda_1^z$  be the smallest singular value of  $X - z$ . Fix any sufficiently small  $\epsilon, \epsilon', \tau > 0$  with  $0 < \tau/2 < \epsilon'$ . For any  $E$  with  $n^{-1+\epsilon} \leq E \leq n^{-3/4-\epsilon'}$ , and any  $z \in \mathbb{C}$  with  $n^{-1/2} \ll |z| - 1 \leq n^{-1/2+\tau}$ , we have<sup>6</sup>*

$$\mathbf{P}(\lambda_1^z \leq E) \lesssim n^{3/2} E^2 e^{-n\delta^2/2} + n^{-D}, \quad \delta := |z|^2 - 1, \quad (3.34)$$

for any fixed (large)  $D > 0$  if  $n$  is sufficiently large.

#### 4. PROOFS OF THEOREMS 2.2, 2.5 AND 2.6

For the purpose of proving our main universality results, Theorems 2.2, 2.5 and 2.6, we will compare the statistics of the spectral radius and of the eigenvalue with the largest real part of general i.i.d. matrices directly with the ones of the complex Ginibre ensemble, which are explicitly known. The key technical input is the following Green function comparison theorem. Its proof will be given in Sections 6-7.

**Theorem 4.1** (Green function comparison). *Fix any integer  $k \geq 1$  and any sufficiently small  $\epsilon, \tau, \nu > 0$  with  $\max\{\tau, \nu\} < \epsilon/100$ . For each  $1 \leq p \leq k$ , let  $f_p(z) : \mathbb{C} \rightarrow \mathbb{R}$  be a  $C^2$ -function such that*

$$\|f_p\|_\infty \lesssim 1, \quad \|\partial_z f_p(z)\|_\infty + \|\partial_{\bar{z}} f_p(z)\|_\infty \lesssim n^{1/2+\nu}, \quad \|\Delta_z f_p(z)\|_\infty \lesssim n^{1+2\nu}, \quad (4.1)$$

and assume that  $f_p$  is compactly supported on the annulus

$$\sqrt{\frac{\gamma_n}{4n}} - \frac{C_n}{\sqrt{4n\gamma_n}} \leq |z| - 1 \leq \frac{n^\tau}{\sqrt{n}}, \quad (4.2)$$

<sup>6</sup>The restriction  $|z| - 1 \gg n^{-1/2}$  can be relaxed to  $|z| - 1 \gtrsim n^{-1/2}$ ; see [60, Remark A.3].

with  $\gamma_n$  given in (2.3) and  $C_n/\sqrt{\log n} \rightarrow 0$  as  $n \rightarrow \infty$ . Furthermore, let  $\mathcal{F}(w_1, \dots, w_k) : \mathbb{R}^k \rightarrow \mathbb{R}$  be any smooth (but not necessarily bounded) function with uniformly bounded partial derivatives, i.e., for some constants  $C_\alpha > 0$ ,

$$|\partial^\alpha \mathcal{F}(w_1, \dots, w_k)| \leq C_\alpha, \quad \alpha := (\alpha_1, \dots, \alpha_k), \quad 1 \leq |\alpha| \leq 10. \quad (4.3)$$

Then, we have

$$\left| \mathbf{E} \left[ \mathcal{F} \left( \sum_{i=1}^n f_1(\sigma_i), \dots, \sum_{i=1}^n f_k(\sigma_i) \right) \right] - \mathbf{E}^{\text{Gin}} \left[ \mathcal{F} \left( \sum_{i=1}^n f_1(\sigma_i), \dots, \sum_{i=1}^n f_k(\sigma_i) \right) \right] \right| = O(n^{-\epsilon}). \quad (4.4)$$

**Remark 4.2.** Theorem 4.1 also holds true if instead of (4.2),  $\{f_p\}_{p=1}^k$  are supported on the set

$$\sqrt{\frac{\gamma'_n}{4n}} - \frac{C_n}{\sqrt{4n\gamma'_n}} \leq |z| - 1 \leq \frac{n^\tau}{\sqrt{n}}, \quad |\arg(z)| \leq \frac{n^{\tau/2}}{n^{1/4}}, \quad (4.5)$$

with  $\gamma'_n$  given in (2.10) and  $C_n/\sqrt{\log n} \rightarrow 0$  as  $n \rightarrow \infty$ . The proof is analogous and will be omitted. The support (4.5) is tailored for proving the universality of the eigenvalue with the largest real part in Theorem 2.6. In both cases,  $\gamma_n$  and  $\gamma'_n$  are chosen such that if  $C_n = O(1)$  then typically there are finitely ( $n$ -independent) many eigenvalues in the sets (4.2) and (4.5), respectively, hence  $\sum_{i=1}^n f(\sigma_i)$  in the argument of  $\mathcal{F}$  is typically finite.

Note that in this Green function comparison theorem we use only that the first and second moments of the two ensembles match, no higher moment matching is used. Furthermore, we stated Theorem 4.1 for the class of functions  $\mathcal{F}$  in (4.3) to keep the proof simpler and because this is enough for proving convergence of the (shifted and rescaled) spectral radius to Gumbel in distribution. However, the same result holds for a larger class of test functions, that, in particular, allows us to conclude the convergence in any finite moment sense (Remark 2.4).

**Remark 4.3.** Theorem 4.1 holds true for any  $\mathcal{F}$  with derivatives of polynomial growth, i.e.

$$|\partial^\alpha \mathcal{F}(w_1, \dots, w_k)| \leq C_\alpha \prod_{p=1}^k (1 + |w_p|)^{C\alpha_p}, \quad 1 \leq |\alpha| \leq 10 \quad (4.6)$$

with some  $C > 0$ . Later in Section 7.3 we will briefly discuss the modifications needed to extend Theorem 4.1 to this larger class of functions.

Using Theorem 4.1, we now prove the main results stated in Section 2, as well as the fact that with high probability there is a unique eigenvalue such that  $|\sigma_1|$  is equal to the spectral radius (i.e., Lemma 4.4 below). For pedagogical reasons, we first present the proof of (2.5) and (2.6), and then explain the minor modifications needed to prove Theorem 2.2 in full generality. Then we prove Lemma 4.4 and Theorem 2.5. In this way we can first present the main ideas of the proofs without obfuscating them with technical complications. Lastly, at the end of this section, using Remark 4.2 we prove Theorem 2.6.

**4.1. Proof of (2.5).** Recall the labelling of the eigenvalues  $|\sigma_1| \geq |\sigma_2| \geq \dots \geq |\sigma_n|$ . From [16, Theorem 2.1], we have a strong concentration estimate for the spectral radius  $|\sigma_1|$ , i.e., for any small  $\tau > 0$

$$1 - \frac{n^\tau}{\sqrt{n}} \leq |\sigma_1| \leq 1 + \frac{n^\tau}{\sqrt{n}}, \quad (4.7)$$

with a very high probability. For any set  $\Omega \subset \mathbb{C}$ , we let

$$\mathcal{N}_\Omega := \#\{\sigma_i : \sigma_i \in \Omega, \quad i \in [n]\} = \sum_{i=1}^n \mathbf{1}_\Omega(\sigma_i) \quad (4.8)$$

denote the number of eigenvalues within  $\Omega$ , where  $\mathbf{1}_\Omega$  is the indicator function of  $\Omega$ . Then for any fixed<sup>7</sup>  $r \in \mathbb{R}$ , using the upper bound in (4.7), we have

$$\mathbf{P} \left( |\sigma_1| \geq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4n\gamma_n}} \right) = \mathbf{P}(\mathcal{N}_{\Omega_r} \geq 1) + O(n^{-D}) = \mathbf{E}[\Phi(\mathcal{N}_{\Omega_r})] + O(n^{-D}), \quad (4.9)$$

<sup>7</sup>Inspecting the proof, the same result holds for any  $|r| \ll \sqrt{\log n}$ .

where  $\Omega_r \subset \mathbb{C}$  is a centered annulus given by

$$\Omega_r := \left\{ z : 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4n\gamma_n}} \leq |z| \leq 1 + \frac{n^\tau}{n^{1/2}} \right\}, \quad (4.10)$$

and  $\Phi : \mathbb{R}_+ \rightarrow [0, 1]$  is a smooth and non-decreasing cut-off function such that

$$\Phi(w) = 0, \quad \text{if } 0 \leq w \leq 1/9; \quad \Phi(w) = 1, \quad \text{if } w \geq 2/9. \quad (4.11)$$

Next, we regularize the indicator function of  $\Omega_r$  as follows. For any small  $\nu > 0$ , choose two test functions  $f_r^\pm \in \mathcal{C}_c^2(\mathbb{C})$  such that

$$\mathbf{1}_{\Omega_{r+n^{-\nu}}} \leq f_r^- \leq \mathbf{1}_{\Omega_r} \leq f_r^+ \leq \mathbf{1}_{\Omega_{r-n^{-\nu}}}, \quad (4.12)$$

and they both satisfy the  $L^\infty$  norm bounds in (4.1) as well as the support condition (4.2). Thus, using Theorem 4.1 for  $k = 1$ ,  $\mathcal{F}(w) = \Phi(w)$  and  $f(z) = f_r^\pm(z)$ , we have

$$\left| \mathbf{E} \left[ \Phi \left( \sum_{i=1}^n f_r^\pm(\sigma_i) \right) \right] - \mathbf{E}^{\text{Gin}} \left[ \Phi \left( \sum_{i=1}^n f_r^\pm(\sigma_i) \right) \right] \right| = O(n^{-\epsilon}). \quad (4.13)$$

Since  $\Phi$  is a non-decreasing function, the last two inequalities in (4.12) implies that

$$\mathbf{E} \left[ \Phi \left( \mathcal{N}_{\Omega_r} \right) \right] \leq \mathbf{E} \left[ \Phi \left( \sum_{i=1}^n f_r^+(\sigma_i) \right) \right] \leq \mathbf{E} \left[ \Phi \left( \mathcal{N}_{\Omega_{r-n^{-\nu}}} \right) \right] \quad (4.14)$$

and the same holds for  $\mathbf{E}$  replaced with  $\mathbf{E}^{\text{Gin}}$ .

Next, using the first inequality in (4.14), the GFT estimate in (4.13) for  $f_r^+$ , and the second inequality in (4.14) for the Ginibre ensemble, we have

$$\mathbf{E} \left[ \Phi \left( \mathcal{N}_{\Omega_r} \right) \right] \leq \mathbf{E} \left[ \Phi \left( \sum_{i=1}^n f_r^+(\sigma_i) \right) \right] \leq \mathbf{E}^{\text{Gin}} \left[ \Phi \left( \sum_{i=1}^n f_r^+(\sigma_i) \right) \right] + O(n^{-\epsilon}) \leq \mathbf{E}^{\text{Gin}} \left[ \Phi \left( \mathcal{N}_{\Omega_{r-n^{-\nu}}} \right) \right] + O(n^{-\epsilon}). \quad (4.15)$$

One can also obtain a similar lower bound using the first two inequalities in (4.12) and (4.13) for  $f_r^-$ , *i.e.*,

$$\mathbf{E} \left[ \Phi \left( \mathcal{N}_{\Omega_r} \right) \right] \geq \mathbf{E}^{\text{Gin}} \left[ \Phi \left( \mathcal{N}_{\Omega_{r+n^{-\nu}}} \right) \right] - O(n^{-\epsilon}). \quad (4.16)$$

Combining (4.15) and (4.16) with (4.9), we hence obtain that

$$\begin{aligned} \mathbf{P}^{\text{Gin}} \left( |\sigma_1| \geq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r+n^{-\nu}}{\sqrt{4n\gamma_n}} \right) - Cn^{-\epsilon} &\leq \mathbf{P} \left( |\sigma_1| \geq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4n\gamma_n}} \right) \\ &\leq \mathbf{P}^{\text{Gin}} \left( |\sigma_1| \geq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r-n^{-\nu}}{\sqrt{4n\gamma_n}} \right) + Cn^{-\epsilon}, \end{aligned} \quad (4.17)$$

for some constant  $C > 0$ . From the explicit Ginibre calculation in Corollary E.3 (or [108, Theorem 1]) for the complex Ginibre ensemble, for any  $r \in \mathbb{R}$  we have

$$\lim_{n \rightarrow \infty} \mathbf{P}^{\text{Gin}} \left( |\sigma_1| \leq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4n\gamma_n}} \right) = e^{-e^{-r}}. \quad (4.18)$$

Since the limiting function in (4.18) is bounded and Lipschitz continuous on  $[-C, \infty)$ , we hence finish the proof of (2.5) from (4.17).

**4.2. Proof of (2.6).** In this section we prove the uniform law (2.6) for the argument of  $\sigma_1$ . For any integer  $k \geq 1$ , we divide  $\mathbb{C}$  into  $k$  congruent angular sectors

$$\Theta_p := \left\{ z \in \mathbb{C} : \frac{2\pi(p-1)}{k} \leq \arg(z) < \frac{2\pi p}{k} \right\}, \quad p \in [k].$$

Then for any fixed  $(r_1, \dots, r_k) \in \mathbb{R}^k$ , we have

$$\mathbf{P} \left( \max_{\sigma_i \in \Theta_p} |\sigma_i| \geq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r_p}{\sqrt{4n\gamma_n}}, \quad \forall p \in [k] \right) = \mathbf{E} \left[ \prod_{p=1}^k \Phi \left( \mathcal{N}_{\Omega_{r_p} \cap \Theta_p} \right) \right] + O(n^{-D}). \quad (4.19)$$



with  $\Omega_{r_p}$  is defined as in (4.10) and  $\Phi$  is given in (4.11). We define the regularisation of the indicator function of  $\Omega_{r_p} \cap \Theta_p$  ( $p \in [k]$ ) as in (4.12), denoted by  $f_{r_p}^\pm$ , so that the conditions in (4.1) and (4.2) are both satisfied. Therefore for any  $k$ , using Theorem 4.1 for

$$\mathcal{F}(w_1, \dots, w_k) = \prod_{p=1}^k \Phi(w_p), \quad f_p = f_{r_p}^\pm, \quad p \in [k], \quad (4.20)$$

with  $\Phi$  defined in (4.11), we conclude that

$$\lim_{n \rightarrow \infty} \mathbf{P} \left( \max_{\sigma_i \in \Theta_p} |\sigma_i| \geq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r_p}{\sqrt{4n\gamma_n}}, \quad \forall p \in [k] \right) \quad (4.21)$$

coincides with its Ginibre counterpart. Note that all correlation functions of the Ginibre eigenvalues are rotational invariant (see (E.1) later), which implies that the probability in (4.21), even before the limit, is a symmetric function in  $(r_1, r_2, \dots, r_k)$  under cyclic permutations<sup>8</sup>. We also note that

$$|\sigma_1| = \max_{p \in [k]} \left\{ \max_{\sigma_i \in \Theta_p} |\sigma_i| \right\}.$$

This implies that  $\mathbf{P}(\sigma_1 \in \Theta_p)$  is independent of the index  $p$ , hence  $\mathbf{P}(\sigma_1 \in \Theta_p) = \frac{1}{k}$  for each  $p \in [k]$ . Repeating this argument for any integer  $k \geq 1$ , we obtain that the argument of  $\sigma_1$  is uniformly distributed on  $[0, 2\pi)$  in the limit as  $k$  increases. This proves (2.6).

Now we are ready to prove Theorem 2.2 in full generality.

*Proof of Theorem 2.2.* For any fixed  $-\infty < r_1 < r_2 < \infty$  and  $0 \leq \theta_1 < \theta_2 \leq 2\pi$ , we aim to show that the limit of the joint distribution

$$\mathbf{P} \left( \sqrt{4n\gamma_n} (|\sigma_1| - 1 - \sqrt{\frac{\gamma_n}{4n}}) \in [r_1, r_2), \quad \arg(\sigma_1) \in [\theta_1, \theta_2) \right) \quad (4.22)$$

can be identified similarly to the proof of (2.5)–(2.6) using Theorem 4.1 by choosing an appropriate  $\mathcal{F}$  function and test functions  $\{f_p\}$  as in (4.20), together with an inclusion-exclusion argument. This argument is standard and follows the same spirit of the above proofs, so we omit the details. In short, Theorem 4.1 implies that the limit of the joint distribution in (4.22) coincides with its Ginibre counterpart. This, together with Corollary E.3 in the Appendix, concludes the proof.  $\square$

We conclude this section with the following lemma:

**Lemma 4.4.** *Let  $X$  be a complex i.i.d. random matrix satisfying Assumption 2.1. Then for any (small)  $\nu > 0$ ,*

$$\lim_{n \rightarrow \infty} \mathbf{P}(\sqrt{4n\gamma_n} (|\sigma_1| - |\sigma_2|) \leq n^{-\nu}) = 0.$$

*Proof of Lemma 4.4.* Fix small  $\epsilon, \nu > 0$  with  $\nu \leq \epsilon/100$ . Normalize the moduli of eigenvalues as

$$r_i := \sqrt{4n\gamma_n} (|\sigma_i| - 1 - \sqrt{\frac{\gamma_n}{4n}}). \quad (4.23)$$

Then from [61, Theorem 2.2], for any sequence  $C_n \rightarrow \infty$ , we have

$$\lim_{n \rightarrow \infty} \mathbf{P}(|r_1| \geq C_n) = 0.$$

Thus we have

$$\begin{aligned} \mathbf{P}(\sqrt{4n\gamma_n} (|\sigma_1| - |\sigma_2|) \leq n^{-\nu}) &= \mathbf{P}(|r_1 - r_2| \leq n^{-\nu}) = \mathbf{P}(|r_1 - r_2| \leq n^{-\nu}, |r_1| \leq C_n) + o_n(1) \\ &\leq \sum_{k=0}^{2\lceil C_n \rceil n^\nu} \mathbf{P}(r_1 \in [a_k, a_{k+1}), r_2 \in [a_k - n^{-\nu}, a_{k+1} + n^{-\nu})) + o_n(1), \end{aligned} \quad (4.24)$$

with  $a_k := -C_n + kn^{-\nu}$ , for  $0 \leq k \leq 2\lceil C_n \rceil n^\nu$ , and  $[-C_n, C_n] \subset \bigcup_k [a_k, a_{k+1}]$ . Choose the sequence  $C_n \ll \sqrt{\log n}$  sufficiently slowly diverging and choose an appropriate function  $\mathcal{F}$  and functions  $f^\pm$  as in

<sup>8</sup>In fact, more information is available about (4.21). First, for each fixed  $1 \leq p \leq k$ , the limiting law of the largest eigenvalue in the  $p$ -th angular sector  $\Theta_p$  is given by  $\exp(-\frac{1}{k}e^{-r_p})$  with the rescaling in (4.21), see (E.16 later). By further computations using (E.1) and Proposition E.2, the joint limiting distribution in (4.21) is indeed a symmetric function given by  $\prod_{p=1}^k \exp(-\frac{1}{k}e^{-r_p})$ .

(4.20) satisfying the conditions in (4.1)-(4.3). By Theorem 4.1, the right side of (4.24) coincides with its Ginibre counterpart up to an error term  $n^{-\epsilon}$ . Note that for Ginibre matrices Kostlan's observation [88] (see Remark E.4 below) implies that, for any  $1 \leq k \leq n$ ,

$$(r_1, r_2, \dots, r_k) \stackrel{d}{=} (y_1, y_2, \dots, y_k), \quad y_k := \sqrt{4n\gamma_n} \left( x_k - 1 - \sqrt{\frac{\gamma_n}{4n}} \right), \quad (4.25)$$

where  $x_1 \geq x_2 \geq \dots \geq x_n$  is the ordered statistics of  $n$  independent chi-distributed random variables  $\{\chi_{2k}\}_{k \in [n]}$ . By a direct computation we have

$$\begin{aligned} & \mathbf{P}^{\text{Gin}} \left( r_1 \in [a_k, a_{k+1}), r_2 \in [a_k - n^{-\nu}, a_{k+1} + n^{-\nu}) \right) \\ &= \mathbf{P} \left( y_1 + C_n \in [kn^{-\nu}, (k+1)n^{-\nu}), y_2 + C_n \in [(k-1)n^{-\nu}, (k+2)n^{-\nu}) \right) \lesssim n^{-2\nu}, \end{aligned} \quad (4.26)$$

with  $C_n \ll \sqrt{\log n}$ . Summing up the terms with  $0 \leq k \leq 2\lceil C_n \rceil n^\nu$  in (4.24), the RHS of (4.24) goes to zero, hence we conclude Lemma 4.4.  $\square$

**4.3. Proof of Theorem 2.5.** By the rescaling of the eigenvalues as in (2.7), we have

$$r_i = \sqrt{4\gamma_n n} \left( |\sigma_i| - 1 - \sqrt{\frac{\gamma_n}{4n}} \right), \quad \theta_i = \arg(\sigma_i). \quad (4.27)$$

Then the test function  $f(z) := g(h(z))$  with  $g \in C_c^2(\mathbb{C})$  and  $h(z) = \sqrt{4n\gamma_n} (|z| - 1 - \sqrt{\frac{\gamma_n}{4n}}) \frac{z}{|z|}$  (which is chosen so that  $f(\sigma_i) = g(r_i e^{i\theta_i})$ ) satisfies the  $L^\infty$  norm bounds in (4.1) and the support condition in (4.2). Hence Theorem 2.5 follows directly by using Theorem 4.1 for  $k = 1$ , choosing  $\mathcal{F}(w) = e^{-w}$  for  $w \in \mathbb{R}_+$ , in combination with the explicit Ginibre computations in Corollary E.3 in the Appendix.  $\square$

**4.4. Proof of Theorem 2.6.** In this section we present the minor modifications needed to prove the universality of the rightmost eigenvalue. The local laws in [38, Theorem 1.2] and [16, Theorem 2.1] imply that, for any small  $\tau > 0$  there exists an eigenvalue in the rectangle

$$\left[ 1 - \frac{n^\tau}{n^{1/2}}, 1 + \frac{n^\tau}{n^{1/2}} \right] \times \left[ -\frac{n^{\tau/2}}{n^{1/4}}, \frac{n^{\tau/2}}{n^{1/4}} \right] \quad (4.28)$$

with very high probability, according to the curvature of the boundary. Combining this with the strong concentration estimate in (4.7), we know that the rightmost eigenvalue is indeed located in the rectangle given in (4.28). Thus, instead of the annulus  $\Omega_r$  in (4.10), we choose a rectangle given by

$$\Omega'_x := \left[ 1 + \sqrt{\frac{\gamma'_n}{4n}} + \frac{x}{\sqrt{4n\gamma'_n}}, 1 + n^{-1/2+\tau} \right] \times \left[ -\frac{n^{\tau/2}}{n^{1/4}}, \frac{n^{\tau/2}}{n^{1/4}} \right], \quad (4.29)$$

with  $\gamma'_n$  in (2.10). Choosing similar test functions as in (4.12), they both satisfy the  $L^\infty$  norm bounds in (4.1) and the support condition in (4.5). Thus using Remark 4.2 in combination with [59, Theorem 1] for the Ginibre ensemble, one can prove the Gumbel law in (2.9) for the rightmost eigenvalue as in Section 4.1. The proof of the Poisson point process in (2.11) is analogous to Section 4.3, but with the different scaling in (2.10), in combination with [59, Theorem 2] for the Ginibre ensemble, and so omitted.

## 5. PROOF OF THE LOCAL LAWS, THEOREMS 3.3 AND 3.5

Consider the following matrix valued stochastic differential equation, the Ornstein–Uhlenbeck flow,

$$dX_t = -\frac{1}{2}X_t dt + \frac{dB_t}{\sqrt{n}}, \quad (5.1)$$

with  $B_t$  being an  $n \times n$  matrix whose entries are i.i.d. standard complex Brownian motions. Set

$$W_t := \begin{pmatrix} 0 & X_t \\ X_t^* & 0 \end{pmatrix}$$

to be the Hermitization of  $X_t$ . Define a matrix  $\Lambda \in \mathbb{C}^{2n \times 2n}$  containing the spectral parameters, written in a  $2 \times 2$  block form as follows

$$\Lambda := Z + i\eta, \quad Z = \begin{pmatrix} 0 & z \\ \bar{z} & 0 \end{pmatrix}, \quad \eta \in \mathbb{R}, \quad z \in \mathbb{C}. \quad (5.2)$$

The main object of interest of this section is the evolution of the resolvent  $(W_t - \Lambda_t)^{-1}$  (and of certain products of two such resolvents) along the *characteristics flow* of the spectral parameters (see also [58, Eq. (5.3)])

$$\partial_t \Lambda_t = -\mathcal{S}[M(\Lambda_t)] - \frac{\Lambda_t}{2}, \quad t \geq 0, \quad (5.3)$$

where  $\mathcal{S}[\cdot]$  is the covariance operator (3.4) and  $M(\Lambda_t) := M^{z_t}(i\eta_t)$ . We point out that if the initial condition  $\Lambda_{i,0}$  of (5.3) is constant diagonal in each of its four  $n \times n$  blocks, as in our case from (3.1), then this structure persists for  $\Lambda_t$  for any  $t \geq 0$  and (5.3) simplifies to two coupled scalar equations:

$$\partial_t \eta_t = -\text{Im } m^{z_t}(i\eta_t) - \frac{\eta_t}{2}, \quad \partial_t z_t = -\frac{z_t}{2}. \quad (5.4)$$

Notice that the evolution of  $\eta_t$  depends on  $z_t$  through  $\text{Im } m^{z_t}$ , but the evolution of  $z_t$  is independent of  $\eta_t$ . We will often use the fact that  $\Lambda_t$  and the pair  $(z_t, \eta_t)$  carry the same information. We note that both the matrix Dyson equation (3.3) and the characteristic flow (5.3) literally hold true for arbitrary Hermitian *deformation matrix*  $Z$ , but then neither  $\Lambda_t$  nor  $M(\Lambda_t)$  is constant block diagonal.

Throughout this section we will always consider the characteristics (5.3) for times smaller than the maximal time

$$T^*(\eta_0, z_0) := \sup\{t : \text{sgn}(\eta_t) = \text{sgn}(\eta_0)\}, \quad (5.5)$$

for which the diagonal part of the characteristics  $i\eta_{T^*}$  reaches the real axis. In this section we will always consider initial conditions  $\eta_0, z_0$  such that  $|\eta_0| + |z_0| \lesssim 1$  (even if not stated explicitly); this implies that  $\|\Lambda_0\| \lesssim 1$ . Note that  $T^* \lesssim 1$  as a consequence of  $\|\Lambda_0\| \lesssim 1$ . Furthermore, since  $\text{Im } m^{z_t}(i\eta_t) \cdot \eta_t > 0$  (see (3.2)),  $t \rightarrow |\eta_t|$  is strictly decreasing, while  $z_t$  moves towards the origin of  $\mathbb{C}$  by (5.4). In the following lemma we summarize several trivially checkable properties of the characteristics (5.4):

**Lemma 5.1.** *Let  $\Lambda_t$  be the solution of (5.3) with some initial condition  $\Lambda_0$  and let  $M_t := M^{z_t}(i\eta_t)$ ,  $\rho_t := \pi^{-1}|\text{Im } m_t|$ , then for any  $0 \leq s \leq t \leq T^*(\eta_0, z_0)$  we have*

$$z_t = e^{-(t-s)/2} z_s, \quad M_t = e^{(t-s)/2} M_s, \quad \frac{|\eta_t|}{\rho_t} = e^{s-t} \frac{|\eta_s|}{\rho_s} - \pi(1 - e^{s-t}); \quad (5.6)$$

in particular  $t \mapsto |\eta_t|/\rho_t$  is a decreasing function. Additionally, for  $0 \leq s \leq t \leq T^*(\eta_0, z_0)$ , we have

$$|\eta_t| \rho_t = |\eta_s| \rho_s - \pi(1 - e^{s-t}) \rho_t^2. \quad (5.7)$$

Furthermore, for any  $\alpha > 1$  there is a  $C_\alpha$  such that

$$\int_s^t \frac{1}{|\eta_r|^\alpha} dr \leq \frac{C_\alpha}{|\eta_t|^{\alpha-1} \rho_t}, \quad \text{and} \quad \int_s^t \frac{\rho_r}{|\eta_r|} dr = \frac{1}{\pi} \left[ \log \left( \frac{\eta_s}{\eta_t} \right) - \frac{t-s}{2} \right]. \quad (5.8)$$

Additionally, by standard ODE theory we have:

**Lemma 5.2.** *Fix  $0 < T \leq 1$ , and pick  $\Lambda$  of the form (5.2) with  $|\eta| > 0$ ,  $|z| \geq 1$  and  $\|\Lambda\| \lesssim 1$ . Then there exists an initial condition  $\Lambda_0$ , with  $T^*(\eta_0, z_0) > T$ ,  $\|\Lambda_0\| \lesssim 1$ , such that the solution  $\Lambda_t$  of (5.3) with this initial condition at time  $T$  satisfies  $\Lambda_T = \Lambda$ . Furthermore, there exist constants  $c_1, c_2 > 0$  such that  $|z_0|^2 - 1 \geq c_1 T$  and  $|\eta_0|/\rho_0 \geq c_2 T$ .*

From now on we will consider the evolution of two different characteristics  $\Lambda_{1,t}, \Lambda_{2,t}$  with maximal times  $T_i^* := T^*(\eta_{i,0}, z_{i,0})$ ,  $i = 1, 2$ . Note that the first two relations in (5.6) imply that

$$|z_{1,s} - z_{2,s}| \sim |z_{1,t} - z_{2,t}| \quad \text{and} \quad \rho_{i,s} \sim \rho_{i,t} \quad \text{for} \quad s, t \in [0, \min_i T_i^*]. \quad (5.9)$$

For  $i = 1, 2$ , define the "resolvent" of  $W_t$  with the generalised spectral parameter  $\Lambda_{i,t}$ :

$$G_{i,t} := G_t^{z_{i,t}}(i\eta_{i,t}) = (W_t - \Lambda_{i,t})^{-1}. \quad (5.10)$$

We now describe the time evolution of  $\langle G_{1,t} A_1 G_{2,t} A_2 \rangle$  for general, time independent deterministic matrices  $A_1, A_2 \in \mathbb{C}^{2n \times 2n}$ . By Itô's formula and (5.10), we have

$$\begin{aligned} d\langle G_{1,t} A_1 G_{2,t} A_2 \rangle &= \sum_{a,b=1}^{2n} \partial_{ab} \langle G_{1,t} A_1 G_{2,t} A_2 \rangle \frac{dB_{ab,t}}{\sqrt{n}} + \langle G_{1,t} A_1 G_{2,t} A_2 \rangle dt \\ &\quad + 2 \sum_{i \neq j} \langle G_{1,t} A_1 G_{2,t} E_i \rangle \langle G_{2,t} A_2 G_{1,t} E_j \rangle dt \\ &\quad + \langle (G_{1,t} - M_{1,t}) \rangle \langle G_{1,t}^2 A_1 G_{2,t} A_2 \rangle dt + \langle (G_{2,t} - M_{2,t}) \rangle \langle G_{2,t}^2 A_2 G_{1,t} A_1 \rangle dt, \end{aligned} \quad (5.11)$$

where the sum in the second line indicates a summation over the matrix pairs  $(E_i, E_j) \in \{(E_1, E_2), (E_2, E_1)\}$  (this notation will be often used in this section even if not stated explicitly). Here  $\partial_{ab}$  denotes the directional derivative  $\partial_{w_{ab}}$ , where  $w_{ab}$  are the entries of the Hermitized matrix  $W_t$ .

The deterministic approximation of  $G_{1,t}AG_{2,t}$ , for any deterministic  $A \in \mathbb{C}^{2n \times 2n}$ , is given by

$$M_{12,t}^A := M^A(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}) := \mathcal{B}_{12,t}^{-1}[M_{1,t}AM_{2,t}]. \quad (5.12)$$

Here the two-body stability operator  $\mathcal{B}_{12,t}$  is defined exactly as in (3.14) with  $M^{z_i}(i\eta_i)$  replaced with its time-dependent version  $M^{z_{i,t}}(i\eta_{i,t})$ . The following lemma borrowed from [58] shows that  $\partial_t \langle M_{12,t}^{A_1} A_2 \rangle$  exactly cancels the deterministic approximations of the second and third terms in the rhs. of (5.11):

**Lemma 5.3** (Lemma 5.5 [58]). *For any matrices  $A_1, A_2 \in \mathbb{C}^{2n \times 2n}$ , we have*

$$\partial_t \langle M_{12,t}^{A_1} A_2 \rangle = \langle M_{12,t}^{A_1} A_2 \rangle + \langle \mathcal{S}[M_{12,t}^{A_1}] M_{21,t}^{A_2} \rangle. \quad (5.13)$$

Combining (5.11) and (5.13), we just derived a stochastic differential equation which describes the time evolution of  $\langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})A_2 \rangle$ :

$$\begin{aligned} d\langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})A_2 \rangle &= \sum_{a,b=1}^{2n} \partial_{ab} \langle G_{1,t}A_1G_{2,t}A_2 \rangle \frac{dB_{ab,t}}{\sqrt{n}} + \langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})A_2 \rangle dt \\ &\quad + 2 \sum_{i \neq j} \langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})E_i \rangle \langle M_{21,t}^{A_2} E_j \rangle dt \\ &\quad + 2 \sum_{i \neq j} \langle M_{12,t}^{A_1} E_i \rangle \langle (G_{2,t}A_2G_{1,t} - M_{21,t}^{A_2})E_j \rangle dt \\ &\quad + 2 \sum_{i \neq j} \langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})E_i \rangle \langle (G_{2,t}A_2G_{1,t} - M_{21,t}^{A_2})E_j \rangle dt \\ &\quad + \langle (G_{1,t} - M_{1,t}) \rangle \langle G_{1,t}^2 A_1 G_{2,t} A_2 \rangle dt + \langle (G_{2,t} - M_{2,t}) \rangle \langle G_{2,t}^2 A_2 G_{1,t} A_1 \rangle dt. \end{aligned} \quad (5.14)$$

This equation is the starting point of our analysis to propagate a two-resolvent local law from the global regime, where  $|\eta_{i,0}| \sim 1$ , to the local regime, where  $i\eta_{i,t}$  is very close to the real axis.

In the next two sections we focus on two different effects that improve the two resolvent local law and eventually yield Theorems 3.3 and 3.5. In Section 5.1 we consider the case  $z_1 \neq z_2$ , i.e. we want exploit the gain in the two resolvent local law when  $|z_1 - z_2|$  is fairly large, whilst in Section 5.2 we consider  $z_1 = z_2 = z$  and  $\eta_{1,t} = \eta_{2,t} = \eta_t$  and exploit the additional smallness of  $\langle (G_t A_1 G_t - M_t^{A_1}) A_2 \rangle$ , with  $M_t^A = M_{12,t}^A(\eta_t, z_t, \eta_t, z_t)$ , due to the special choice  $A_1 = F, A_2 = F^*$ . We remark that gaining from both effects simultaneously is not possible.

**5.1. Gain from  $|z_1 - z_2|$  in the local law for  $G^{z_1}AG^{z_2}$  (Theorem 3.3).** The main result of this section shows that if initially we have a good control of  $G_{1,0}AG_{2,0}$  in terms of  $|z_1 - z_2|$  then this is preserved for  $G_{1,t}AG_{2,t}$  along the characteristics (5.3).

**Proposition 5.4.** *Fix sufficiently small  $n$ -independent constants  $\epsilon, \omega_d, \omega_1, c > 0$ , and for  $i = 1, 2$  fix spectral parameters*

$$\Lambda_{i,0} := \begin{pmatrix} i\eta_{i,0} & z_{i,0} \\ z_{i,0} & i\eta_{i,0} \end{pmatrix}, \quad (5.15)$$

with  $0 \leq |z_{i,0}| - 1 \leq c$ ,  $|z_{1,0} - z_{2,0}| \leq \omega_d$ , and  $0 < |\eta_{i,0}| \leq \omega_1$ . For  $0 \leq t < T^*(\eta_{i,0}, z_{i,0})$ , let  $\Lambda_{i,t}$  be the solution of (5.3) with initial condition  $\Lambda_{i,0}$ . Recall the resolvent  $G_{i,t} = (W_t - \Lambda_{i,t})^{-1}$  and  $M_{12,t}^A$ , the deterministic approximation of  $G_{1,t}AG_{2,t}$ , from (5.12). Define the following control parameters and their time dependent versions:

$$\begin{aligned} \ell &= \ell(\eta_1, z_1, \eta_2, z_2) := \min_{i=1,2} |\eta_i| \rho^{z_i}(i\eta_i), & \ell_t &= \ell(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}), \\ \gamma &= \gamma(\eta_1, z_1, \eta_2, z_2) := |z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}, & \gamma_t &= \gamma(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}). \end{aligned} \quad (5.16)$$

Assume that for some small  $0 < \xi \leq \epsilon/10$ , with very high probability, it holds

$$\left| \langle (G_0^{z_1,0}(i\eta_1)A_1G_0^{z_2,0}(i\eta_2) - M_{12,t}^{A_1}(\eta_1, z_{1,0}, \eta_2, z_{2,0}))A_2 \rangle \right| \lesssim \frac{n^\xi}{\sqrt{n}\ell_0\gamma_0} \wedge \frac{n^\xi}{n|\eta_{1,0}\eta_{2,0}|}, \quad (5.17)$$

for any choice  $A_1, A_2 \in \{E_+, E_-\}$  and for any  $\eta_i$  with  $|\eta_i| \in [|\eta_{i,0}|, n^{100}]$ , where  $\ell_0$  and  $\gamma_0$  are evaluated at the spectral parameters  $(\eta_0, z_{1,0}, \eta_2, z_{2,0})$ . Then

$$|\langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})A_2 \rangle| \lesssim \frac{n^{2\xi}}{\sqrt{nl_t\gamma_t}} \wedge \frac{n^{2\xi}}{n|\eta_{1,t}\eta_{2,t}|}, \quad (5.18)$$

with very high probability uniformly in all  $t \leq \min_i T^*(\eta_{i,0}, z_{i,0})$  such that  $nl_t \geq n^\epsilon$ .

Note that by the strict monotonicity of  $t \rightarrow \eta_{i,t}/\rho_{i,t}$ , and by  $|z_{1,t} - z_{2,t}| = e^{-(t-s)/2}|z_{1,0} - z_{2,0}|$ , it follows that  $\gamma_s \geq \gamma_t$  for any  $0 \leq s \leq t \leq T^*$ , i.e.  $t \mapsto \gamma_t$  is strictly decreasing.

Before presenting the proof of Proposition 5.4 we first conclude the proof of Part A of Theorem 3.3 for matrices with a large Gaussian component. The proof for general i.i.d. matrices (Part B) will be given in Appendix D.

*Proof of Theorem 3.3 (Part A) with Gaussian component.* For any small  $c^* > 0$ , by the global law outside of the non-Hermitian spectrum [52, Theorem 5.2], for any small  $\xi > 0$  we have

$$|\langle (G^{z_1}(i\eta_1)A_1G^{z_2}(i\eta_2) - M_{12}^{A_1})A_2 \rangle| \lesssim \frac{n^\xi}{n}, \quad (5.19)$$

with very high probability, uniformly in spectral parameters such that  $1 + c^* \leq |z_i| \leq 10$ ,  $|\eta_i| \leq n^{100}$ , and deterministic matrices  $\|A_1\| + \|A_2\| \lesssim 1$ . Define  $t = t(\mathfrak{s})$  such that  $\mathfrak{s} = (1 - e^{-t/2})^{1/2}$ , then by the property of the Ornstein-Uhlenbeck process (5.1),  $G_t$  from (5.10) and  $G_{\mathfrak{s}}^z$  defined in Theorem 3.3 have the same distribution. Fix any small  $\omega_1, \omega_d, c$ , and pick any  $z_1, z_2$  such that  $1 \leq |z_i| \leq 1 + c/2$  and  $|z_1 - z_2| \leq \omega_d/2$ , and pick any  $\eta_i$  such that  $|\eta_i| \leq \omega_1/2$  and  $nl \geq n^\epsilon$ . Then, by Lemma 5.2 and that fact that  $\mathfrak{s}$  can be chosen small in Part A of Theorem 3.3, there exist  $z_{i,0}$  and  $\eta_{i,0}$  with  $|z_{1,0} - z_{2,0}| \leq \omega_d$ ,  $|\eta_{i,0}| \leq \omega_1$ , such that  $z_{i,t} = z_i$ ,  $\eta_{i,t} = \eta_i$ ,  $c_* \leq |z_{i,0}| - 1 \leq c$ , for some small  $c_* = c_*(\mathfrak{s}) > 0$ . Choosing these parameters sufficiently small so that (5.19) holds, and noticing that  $1/n \leq 1/(nl\gamma)$ , this guarantees that the assumption (5.17) is satisfied. We thus conclude from (5.18), using  $z_{i,t} = z_i$ ,  $\eta_{i,t} = \eta_i$ , hence  $\ell = \ell_t$  and  $\gamma = \gamma_t$ , that (here  $A_1, A_2 \in \{E_+, E_-\}$ )

$$|\langle (G_t^{z_1}(i\eta_1)A_1G_t^{z_2}(i\eta_2) - M_{12}^{A_1})A_2 \rangle| \lesssim \frac{n^{2\xi}}{\sqrt{nl\gamma}} + \frac{n^{2\xi}}{n|\eta_1\eta_2|}. \quad (5.20)$$

Since  $G_t \stackrel{d}{=} G_{\mathfrak{s}}^z$ , by the choice of  $t$ , we concluded (3.16).  $\square$

*Proof of Proposition 5.4.* By [61, Theorem 3.1] we have  $|\langle G_{i,t} - M_{i,t} \rangle| \lesssim n^\xi/(n|\eta_{i,t}|)$  with very high probability for any fix  $t$ , but in fact also implies  $\sup_{0 \leq t \leq T^*} (n|\eta_{i,t}|)|\langle G_{i,t} - M_{i,t} \rangle| \lesssim n^\xi$ , i.e. that the local law holds with very high probability simultaneously for any  $0 \leq t \leq T^*$ . This follows by a simple procedure called the *grid argument*. It consists of choosing equidistant intermediate times  $t_1, t_2, \dots, t_{K_n} \in [0, T^*]$ , for some large enough  $K_n$  (say  $K_n = n^{10}$ ), using the fact that for each  $t = t_i$  the local law holds and then extending the bound to any intermediate  $t \in [0, T^*]$  by the Lipschitz continuity of  $G_{i,t} - M_{i,t}$  (with Lipschitz parameter  $n$ ). The probability of the exceptional sets where the bound does not hold for some  $t_j$  can be estimated by a standard union bound since there are only polynomially many of them. We will use a similar procedure also for local laws for products of different resolvents. We also point out that throughout this proof, even if not stated explicitly, all the inequalities hold with very high probability. Additionally, to keep the notation simpler we assume that  $\eta_{1,t}$  and  $\eta_{2,t}$  are both positive, all the other cases are analogous and so omitted.

Fix any  $\epsilon > 0$ , and define the time,

$$T_i^\epsilon = T^\epsilon(\eta_{i,0}, z_{i,0}) := \min\{t > 0 : n\eta_{i,t}\rho_{i,t} = n^\epsilon\}, \quad (5.21)$$

such that  $\eta_{i,t}\rho_{i,t}$  reaches the critical threshold  $n\eta_{i,T_i^\epsilon}\rho_{i,T_i^\epsilon} = n^\epsilon$ . Notice that, by (5.4) and by taking the trace in the second relation in (5.6), it follows that

$$\partial_t(\eta_{i,t}\rho_{i,t}) = -\pi\rho_{i,t}^2.$$

Then, by the uniqueness of the solution of this simple ODE, together with the strict monotonicity<sup>9</sup> of the map  $\eta \mapsto \eta\rho^{z_{i,0}}(i\eta)$ , it follows that for a fixed  $z_{i,0}$  the map  $\eta_{i,0} \mapsto T^\epsilon(\eta_{i,0}, z_{i,0})$  is increasing.

<sup>9</sup>This monotonicity follows from a direct calculation of the derivative of  $m = m^{z_{i,0}}(i\eta)$

$$\partial_\eta \text{Im } m = \text{Im } m \frac{1 - 2\text{Im } m(\eta + \text{Im } m)}{2(\text{Im } m)^2(\eta + \text{Im } m) + \eta} > 0,$$

We start defining

$$Y_{\sigma,\sigma',t}(\eta_1, z_1, \eta_2, z_2) := \langle (G_t^{z_1}(i\eta_1)E_\sigma G_t^{z_2}(i\eta_2) - M_{12}^{E_\sigma}(\eta_1, z_1, \eta_2, z_2))E_{\sigma'} \rangle, \quad \sigma, \sigma' \in \{+, -\}. \quad (5.22)$$

Additionally, to keep the notation short we introduce the control parameter

$$\alpha(\eta_1, z_1, \eta_2, z_2) := \frac{1}{\sqrt{n\ell(\eta_1, z_1, \eta_2, z_2)}\gamma(\eta_1, z_1, \eta_2, z_2)} \wedge \frac{1}{n|\eta_1\eta_2|}, \quad \alpha_t := \alpha(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}), \quad (5.23)$$

with  $\ell, \gamma$  being defined in (5.16). Within the whole proof, for any given  $\eta_{i,0}, z_{i,0}$ ,  $i = 1, 2$ , we will consider the flow (5.14) up to times  $t \leq \tau$ , with  $\tau = \tau(\eta_{1,0}, z_{1,0}, \eta_{2,0}, z_{2,0})$  being the stopping time defined by<sup>10</sup>

$$\tau := \inf \left\{ t \geq 0 : \max_{i=1,2} \sup_{\tilde{\eta}_{i,0}} \max_{\sigma,\sigma' \in \{+,-\}} \alpha(\tilde{\eta}_{1,t}, z_{1,t}, \tilde{\eta}_{2,t}, z_{2,t})^{-1} |Y_{\sigma,\sigma',t}(\tilde{\eta}_{1,t}, z_{1,t}, \tilde{\eta}_{2,t}, z_{2,t})| = n^{2\xi} \right\} \wedge T_1^\epsilon \wedge T_2^\epsilon, \quad (5.24)$$

for some  $0 < \xi \leq \epsilon/10$ . The supremum  $\sup^*$  is taken over all  $\tilde{\eta}_{i,0}$  satisfying  $\tilde{\eta}_{i,0} \in [\eta_{i,0}, \omega_1]$ , with  $\omega_1$  given in Proposition 5.4. Here  $\tilde{\eta}_{i,t}$  is the solution to the diagonal part of (5.3) with initial conditions  $z_{i,0}, \tilde{\eta}_{i,0}$ . Note that the initial condition  $z_{i,0}$ , hence  $z_{i,t} = e^{-t/2}z_{i,0}$  along the tilde-flow, is always the same, irrespective of  $\tilde{\eta}_{i,0}$ . Furthermore, by (5.17), together with a standard grid argument, it follows that  $\tau > 0$  with very high probability.

Additionally, by the definition of the stopping time  $\tau$ , it follows that for any  $t \leq \tau$  we have

$$\sup_{\eta_{i,t} \leq \hat{\eta}_i \leq \omega_1/2} \max_{\sigma,\sigma' \in \{+,-\}} \alpha(\hat{\eta}_1, z_{1,t}, \hat{\eta}_2, z_{2,t})^{-1} |Y_{\sigma,\sigma',t}(\hat{\eta}_1, z_{1,t}, \hat{\eta}_2, z_{2,t})| \leq n^{2\xi}, \quad (5.25)$$

with very high probability. We remark that to obtain (5.25) we also used that for any fixed  $t \leq T_1^\epsilon \wedge T_2^\epsilon$ , by Lemma 5.2, any  $\hat{\eta}_i \in [\eta_{i,t}, \omega_1/2]$  can be achieved as the solution  $\tilde{\eta}_{i,t}$  of (5.4) with an appropriately chosen initial condition  $\tilde{\eta}_{i,0} \in [\eta_{i,0}, \omega_1]$ , i.e.  $\hat{\eta}_i = \tilde{\eta}_{i,t}$ . Note that in (5.25) the supremum is taken only over  $\hat{\eta}_i$ 's, whilst the  $z_{i,t}$ 's are fixed.

The supremum over a broader set of initial conditions  $\tilde{\eta}_{i,0}$  in (5.24), as well as over  $\hat{\eta}_i$  in (5.25), needs to be considered for a minor technical reason: at some point we will also need to consider quantities of the form  $GA|G|$  instead of  $GAG$ , and we use an integral representation of  $|G|$  in terms of  $G$  that requires integration over an entire segment of spectral parameters (see later in Appendix A after the end of the proof of Lemma A.1). The reader can safely ignore all tildes and hats at first reading and focus on the special case  $\tilde{\eta}_{i,t} = \eta_{i,t}$  and  $\hat{\eta}_i = \eta_{i,t}$  in these suprema.

Before proceeding with the estimate of the various terms in (5.14), we state some bounds for the functions  $\langle M_{12,t}^{A_1} A_2 \rangle$  in the special case when  $A_1, A_2 \in \{E_+, E_-\}$ . The proof of this lemma is postponed to Appendix A.

**Lemma 5.5.** *Let  $\gamma(\eta_1, z_1, \eta_2, z_2)$  be defined in (5.16). There exists an explicit real function  $b(\eta_1, z_1, \eta_2, z_2) \in \mathbb{R}$  such that*

$$\begin{aligned} |\langle M_{12}^{E_+} E_+ \rangle| &\lesssim \frac{1}{\gamma(\eta_1, z_1, \eta_2, z_2)}, & |\langle M_{12}^{E_-} E_- \rangle| &\lesssim \frac{1}{\gamma(\eta_1, z_1, \eta_2, z_2)}, \\ \langle M_{12}^{E_+} E_- \rangle = -\langle M_{12}^{E_-} E_+ \rangle &= ib(\eta_1, z_1, \eta_2, z_2), & |b(\eta_1, z_1, \eta_2, z_2)| &\lesssim \frac{1}{\gamma(\eta_1, z_1, \eta_2, z_2)}. \end{aligned} \quad (5.26)$$

Additionally, we have

$$\begin{aligned} \langle M_{12}^{E_+} E_+ \rangle &\leq -\langle M_{12}^{E_-} E_- \rangle && \text{for } \eta_1 \eta_2 > 0, \\ \langle M_{12}^{E_+} E_+ \rangle &\geq -\langle M_{12}^{E_-} E_- \rangle && \text{for } \eta_1 \eta_2 < 0. \end{aligned} \quad (5.27)$$

where we used that  $2\text{Im } m(\eta + \text{Im } m) \leq C\omega_1^{2/3}$  and that  $\omega_1$  is chosen sufficiently small.

<sup>10</sup>Here the stopping time  $\tau$  should not be confused with the small (deterministic) exponent used often starting from Section 3 to denote  $|z| \leq 1 + n^{-1/2+\tau}$ .

We now proceed with the estimate of the quadratic variation of the stochastic term in (5.14):

$$\begin{aligned} \frac{1}{n^2} \sum_{i \neq j} \left[ \langle G_{1,t} A_1 G_{2,t} A_2 G_{1,t} E_i G_{1,t}^* A_2 G_{2,t}^* A_2 G_{1,t}^* E_j \rangle + \langle G_{2,t} A_2 G_{1,t} A_1 G_{2,t} E_i G_{2,t}^* A_1 G_{1,t}^* A_2 G_{2,t}^* E_j \rangle \right. \\ \left. + 2\text{Re} \langle G_{1,t} A_1 G_{2,t} A_2 G_{1,t} E_i G_{2,t}^* A_1 G_{1,t}^* A_2 G_{2,t}^* E_j \rangle \right] dt. \end{aligned} \quad (5.28)$$

Here we used that  $A_1, A_2$  are Hermitian, since  $A_1, A_2 \in \{E_+, E_-\}$ . We present the estimate only for the first term in (5.28), the the second and third terms are handled completely analogously. To estimate this term we use

$$\langle G_{1,t} A_1 G_{2,t} A_2 G_{1,t} E_i G_{1,t}^* A_2 G_{2,t}^* A_2 G_{1,t}^* E_j \rangle \lesssim \frac{1}{\eta_{1,t}^2} \langle \text{Im} G_{1,t} A_1 G_{2,t} A_2 \text{Im} G_{1,t} A_2 G_{2,t}^* A_2 \rangle, \quad (5.29)$$

where we used a Schwarz inequality followed by the Ward identity  $GG^* = \text{Im}G/\eta$ . We now face the problem that even if we are studying the evolution of the product of two resolvents the quadratic variation consists of traces of products of four resolvents that needs to be reduced to traces with two resolvents in order to have a closed equation. For this purpose we will give two different estimates for (5.29). One estimate uses the deterministic bound  $\|G\| \leq 1/\eta$  to reduce the number of resolvents at the price of  $1/\eta$  factor, reminiscent to the Ward identity. This procedure loses the natural factor  $1/\gamma_t$  originating from every instance a factor  $G_{1,t} A G_{2,t}$  appears. Note that the factor  $1/\gamma_t$  is much better than  $1/\eta_t$ , in particular the former carries the optimal  $|z_1 - z_2|$  decay. The second estimate, instead, will preserve the number of  $1/\gamma_t$ 's (hence the  $|z_1 - z_2|$ -decay) at the price of losing a factor  $\sqrt{n\ell_t}$ .

We start with this latter estimate. We rely on the following *reduction inequality*

$$\langle \text{Im} G_{1,t} A_1 G_{2,t} A_2 \text{Im} G_{1,t} A_2 G_{2,t}^* A_1 \rangle \leq n \langle \text{Im} G_{1,t} A_1 | G_{2,t} | A_1 \rangle \langle \text{Im} G_{1,t} A_2 | G_{2,t} | A_2 \rangle, \quad (5.30)$$

which easily follows by spectral decomposition

$$\begin{aligned} \langle \text{Im} G_{1,t} A_1 G_{2,t} A_2 \text{Im} G_{1,t} A_2 G_{2,t}^* A_1 \rangle \\ = \frac{\eta_1^2}{n} \sum_{ijkl} \frac{\langle \mathbf{u}_i^{z_1}(t), A_1 \mathbf{u}_j^{z_2}(t) \rangle \langle \mathbf{u}_j^{z_2}(t), A_2 \mathbf{u}_k^{z_1}(t) \rangle \langle \mathbf{u}_k^{z_1}(t), A_2 \mathbf{u}_l^{z_2}(t) \rangle \langle \mathbf{u}_l^{z_2}(t), A_1 \mathbf{u}_i^{z_1}(t) \rangle}{[(\lambda_i^{z_1}(t))^2 + \eta_{1,t}^2] (\lambda_j^{z_2}(t) - i\eta_{2,t}) [(\lambda_k^{z_1}(t))^2 + \eta_{1,t}^2] (\lambda_l^{z_2}(t) - i\eta_{2,t})} \\ \lesssim n \langle \text{Im} G_{1,t} A_1 | G_{2,t} | A_1 \rangle \langle \text{Im} G_{1,t} A_2 | G_{2,t} | A_2 \rangle, \end{aligned} \quad (5.31)$$

where to go from the first to the second line we used the Schwarz inequality in the form (omitting the time dependence of the eigenvectors)

$$|\langle \mathbf{u}_i^{z_1}, A_1 \mathbf{u}_j^{z_2} \rangle \langle \mathbf{u}_j^{z_2}, A_2 \mathbf{u}_k^{z_1} \rangle \langle \mathbf{u}_k^{z_1}, A_2 \mathbf{u}_l^{z_2} \rangle \langle \mathbf{u}_l^{z_2}, A_1 \mathbf{u}_i^{z_1} \rangle| \leq |\langle \mathbf{u}_i^{z_1}, A_1 \mathbf{u}_j^{z_2} \rangle \langle \mathbf{u}_k^{z_1}, A_2 \mathbf{u}_l^{z_2} \rangle|^2 + |\langle \mathbf{u}_j^{z_2}, A_2 \mathbf{u}_k^{z_1} \rangle \langle \mathbf{u}_l^{z_2}, A_1 \mathbf{u}_i^{z_1} \rangle|^2.$$

To estimate the rhs. of (5.30), we claim that

$$\langle \text{Im} G_{1,t} A_1 | G_{2,t} | A_1 \rangle \lesssim \frac{\log n}{\gamma_t} + \frac{n^{2\xi} \log n}{\sqrt{n\ell_t} \gamma_t}, \quad (5.32)$$

i.e. that the bound (5.25) with the choice  $\hat{\eta}_t = \tilde{\eta}_{i,t}$  still essentially holds when one of the  $G$ 's is replaced by its absolute value. The proof of (5.32) is presented in Appendix A. For notational simplicity, in the remainder of the proof we will always omit  $(\log n)$ -factors.

For  $t \leq \tau$ , we thus estimate the integral of the quadratic variation of stochastic term in (5.14) as follows

$$\begin{aligned} \int_0^t \frac{1}{n^2 \eta_{1,s}^2} \langle \text{Im} G_{1,s} A_1 G_{2,s} A_2 \text{Im} G_{1,s} A_2 G_{2,s}^* A_1 \rangle ds &\leq \int_0^t \frac{1}{n \eta_{1,s}^2} \langle \text{Im} G_{1,s} A_1 | G_{2,s} | A_1 \rangle \langle \text{Im} G_{1,s} A_2 | G_{2,s} | A_2 \rangle ds \\ &\lesssim \int_0^t \frac{1}{n \eta_{1,s}^2} \left( \frac{1}{\gamma_s} + \frac{n^{2\xi}}{\sqrt{n\ell_s} \gamma_s} \right)^2 ds \\ &\lesssim \int_0^t \frac{1}{n \eta_{1,s}^2 \gamma_s^2} ds \lesssim \frac{1}{n \ell_t \gamma_t^2} \end{aligned} \quad (5.33)$$

where in the first inequality we used (5.30), in the second inequality (5.32), in the third inequality we used that  $n^{4\xi} \leq \ell_s$ , since  $\xi \leq \epsilon/10$  (we will often use these facts in the remainder of this section even if not stated explicitly), and in the last inequality we used the first relation in (5.8) together with  $\gamma_s \gtrsim \gamma_t$ .

We now present the other estimate for the integral of the quadratic variation which loses any  $1/\gamma_t$  but it is better in terms of  $1/n$ -powers:

$$\begin{aligned} \int_0^t \frac{1}{n^2 \eta_{1,s}^2} \langle \text{Im } G_{1,s} A_1 G_{2,s} A_2 \text{Im } G_{1,s} A_2 G_{2,s}^* A_1 \rangle ds &\leq \int_0^t \frac{1}{n^2 \eta_{1,s}^2} \|A_1 G_{2,s} A_2 \text{Im } G_{1,s} A_2 G_{2,s}^* A_1\| \langle \text{Im } G_{1,s} \rangle ds \\ &\lesssim \int_0^t \frac{\rho_{1,s}}{n^2 \eta_{1,s}^3 \eta_{2,s}^2} ds \lesssim \frac{1}{(n \eta_{1,t} \eta_{2,t})^2}, \end{aligned} \quad (5.34)$$

where in the first inequality we tacitly used the cyclicity of the trace and the positivity of  $\text{Im } G$  to write  $\langle \text{Im } G_{1,s} A_1 G_{2,s} A_2 \text{Im } G_{1,s} A_2 G_{2,s}^* A_1 \rangle = \langle [\text{Im } G_{1,s}]^{1/2} A_1 G_{2,s} A_2 \text{Im } G_{1,s} A_2 G_{2,s}^* A_1 [\text{Im } G_{1,s}]^{1/2} \rangle$  before the operator inequalities  $\|A_i\| \lesssim 1$ ,  $\|G_i\| \leq 1/\eta_i$  were applied. Furthermore, in the second inequality we also used the single resolvent local law  $|\langle G_{i,s} - M_{i,s} \rangle| \lesssim n^\xi / (n \eta_{i,s})$  in the form  $\langle \text{Im } G_{i,s} \rangle \lesssim \rho_{i,s} + n^\xi / (n \eta_{i,s}) \lesssim \rho_{i,s}$  since  $n \ell_s \geq n^\epsilon$ . Here we set  $\rho_{i,s} := \pi^{-1} |\text{Im } m^{z_{i,s}}(\eta_{i,s})|$ . In the last step we used  $\eta_{2,t} \leq \eta_{2,s}$  by monotonicity of  $t \rightarrow \eta_t$  and the first relation in (5.8).

Then, combining the two inequalities (5.33) and (5.34), we conclude the following very high probability bound for the quadratic variation of the Ito stochastic term in (5.14):

$$\left( \left[ \int_0^\cdot \partial_{ab} \langle G_{1,s} A_1 G_{2,s} A_2 \rangle \frac{dB_{ab,s}}{\sqrt{n}} \right]_{t \wedge \tau} \right)^{1/2} \lesssim \frac{n^\xi}{\sqrt{n \ell_{t \wedge \tau} \gamma_{t \wedge \tau}}} \wedge \frac{n^\xi}{n \eta_{1,t \wedge \tau} \eta_{2,t \wedge \tau}} = n^\xi \alpha_{t \wedge \tau}. \quad (5.35)$$

We now continue the estimates of the other terms in (5.14). The second term in the first line of (5.14) can be incorporated in the lhs. of (5.14) by considering the evolution of  $e^{-t} \langle (G_{1,t} A_1 G_{2,t} - M_{12,t}^{A_1}) A_2 \rangle$ ; we thus neglect this term from the analysis.

Next, we consider the terms in the last line of (5.14). These terms, using the single resolvent local law  $|\langle G_{1,t} - M_{1,t} \rangle| \lesssim n^\xi / (n \eta_{1,t})$ , for  $t \leq T_1^\epsilon \wedge T_2^\epsilon$ , are bounded by (we present the bound for only one of them)

$$\begin{aligned} &\int_0^t |\langle G_{1,s} - M_{1,s} \rangle| |\langle G_{1,s}^2 A_1 G_{2,s} A_2 \rangle| ds \\ &\lesssim \int_0^t |\langle (G_{1,s} - M_{1,s}) \rangle| \frac{1}{\eta_{1,s}} \langle \text{Im } G_{1,s} A_1 | G_{2,s} | A_1 \rangle^{1/2} \langle \text{Im } G_{1,s} A_2 | G_{2,s} | A_2 \rangle^{1/2} ds \\ &\lesssim \int_0^t \frac{n^\xi}{n \eta_{1,s}^2} \left( \frac{1}{\gamma_s} + \frac{n^\xi}{\sqrt{n \ell_s} \gamma_s} \right) ds \lesssim \frac{n^\xi}{n \ell_t \gamma_t}. \end{aligned} \quad (5.36)$$

Here in the first inequality we used the bound (easily checkable by spectral decomposition)

$$|\langle G_{1,s}^2 A_1 G_{2,s} A_2 \rangle| \lesssim \frac{1}{\eta_1} \langle \text{Im } G_{1,s} A_1 | G_{2,s} | A_1 \rangle^{1/2} \langle \text{Im } G_{1,s} A_2 | G_{2,s} | A_2 \rangle^{1/2}, \quad (5.37)$$

while in second inequality we used (5.32). In the last step, we used  $n^\xi \leq \sqrt{n \ell_s}$ ,  $\gamma_t \leq \gamma_s$ , and (5.8). Similarly, to the estimate of the stochastic term in (5.14), we now give another estimate for the term in (5.36):

$$\begin{aligned} &\int_0^t |\langle G_{1,s} - M_{1,s} \rangle| |\langle G_{1,s}^2 A_1 G_{2,s} A_2 \rangle| ds \\ &\lesssim \int_0^t |\langle (G_{1,s} - M_{1,s}) \rangle| \frac{1}{\eta_{1,s}} \langle \text{Im } G_{1,s} \rangle^{1/2} \langle \text{Im } G_{1,s} A_1 G_{2,s} A_2 G_{2,s}^* A_1 \rangle^{1/2} ds \\ &\lesssim \int_0^t \frac{n^\xi}{n \eta_{1,s}^2} \cdot \frac{\rho_{1,s}}{\eta_{2,s}} ds \lesssim \frac{n^\xi}{n \eta_{1,t} \eta_{2,t}}, \end{aligned} \quad (5.38)$$

where in the first inequality we used a simple Schwarz inequality, and in the second inequality we used a norm bound, as in (5.34), to estimate the last trace in the second line.



Then, for  $t \leq \tau$ , writing  $E_i, E_j$  as a linear combination of  $E_+, E_-$  (using the definitions in (1.5)) so that we could use (5.24), we estimate the term in the fourth line of (5.14) by

$$\begin{aligned} \int_0^t | \langle (G_{1,s}A_1G_{2,s} - M_{12,s}^{A_1})E_i \rangle | | \langle (G_{2,s}A_2G_{1,s} - M_{21,s}^{A_2})E_j \rangle | ds &\lesssim \int_0^t n^{4\xi} \left( \frac{1}{n\ell_s\gamma_s^2} \wedge \frac{1}{n^2\eta_{1,s}^2\eta_{2,s}^2} \right) ds \\ &\lesssim n^{4\xi} \left( \frac{1}{n\ell_t\gamma_t} \wedge \frac{1}{(n\ell_t)(n\eta_{1,t}\eta_{2,t})} \right). \end{aligned} \quad (5.39)$$

In the last step, to estimate the term  $1/(n\ell_s\gamma_s^2)$ , we used again  $\gamma_s \geq \gamma_t$  and the second formula of (5.8) for the bound

$$\frac{1}{\gamma_t} \int_0^t \frac{1}{\ell_s} ds \leq \frac{1}{\gamma_t} \int_0^t \left( \frac{1}{\eta_{1,s}\rho_{1,s}} + \frac{1}{\eta_{2,s}\rho_{2,s}} \right) ds \lesssim \frac{1}{\gamma_t} \left( \frac{1}{\rho_{1,s}^2} + \frac{1}{\rho_{2,s}^2} \right) \lesssim \frac{1}{\ell_t}, \quad (5.40)$$

where in the last inequality we used that  $\gamma_t \geq \max_i(\eta_{i,t}/\rho_{i,t})$ . The estimate for  $1/(n\eta_{1,t}\eta_{2,t})^2$  is simpler, and it follows by the monotonicity  $\eta_{i,s} \geq \eta_{i,t}$ , together with  $\int 1/\eta_{i,s}^2 \leq 1/\ell_t$ .

Collecting the bounds (5.35), (5.36) and (5.39) and writing again  $E_1, E_2$  in terms of  $E_+, E_-$ , we are thus left with

$$\begin{aligned} &d \langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})A_2 \rangle \\ &= \langle G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1} \rangle \langle M_{21,t}^{A_2} \rangle dt - \langle (G_{1,t}A_1G_{2,t} - M_{12,t}^{A_1})E_- \rangle \langle M_{21,t}^{A_2}E_- \rangle dt \\ &\quad + \langle M_{12,t}^{A_1} \rangle \langle G_{2,t}A_2G_{1,t} - M_{21,t}^{A_2} \rangle dt - \langle M_{12,t}^{A_1}E_- \rangle \langle (G_{2,t}A_2G_{1,t} - M_{21,t}^{A_2})E_- \rangle dt + h_t dt + de_t, \end{aligned} \quad (5.41)$$

with  $h_t$  being a forcing term, and  $e_t$  being a martingale term such that

$$\int_0^{t \wedge \tau} |h_s| ds + \left[ \int_0^\cdot de_s \right]_{t \wedge \tau}^{1/2} \lesssim n^\xi \alpha_{t \wedge \tau},$$

with very high probability. We also recall that by  $[\cdot]_t$  we denote the quadratic variation process.

With the four choices  $A_1 = E_\pm, A_2 = E_\pm$ , we view (5.41) a coupled system of four ODE's modulo a small error. Recalling the definition (5.22), we introduce the short-hand notations

$$Y_{\sigma,\sigma',t} := Y_{\sigma,\sigma',t}(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}), \quad \sigma, \sigma' \in \{+, -\}, \quad (5.42)$$

for these four components. Furthermore, we define the 4-dimensional column vector

$$\mathcal{Y}_t := \begin{pmatrix} Y_{+,+,t} \\ Y_{+,-,t} \\ Y_{-,+,t} \\ Y_{-,-,t} \end{pmatrix} \in \mathbb{C}^4, \quad (5.43)$$

and the 4 by 4 matrices

$$\begin{aligned} \widehat{\mathcal{M}}_t &:= \begin{pmatrix} 2a_{+,t} & ib_t & -ib_t & 0 \\ ib_t & a_{+,t} + a_{-,t} & 0 & -ib_t \\ -ib_t & 0 & a_{+,t} + a_{-,t} & ib_t \\ 0 & -ib_t & ib_t & 2a_{-,t} \end{pmatrix}, \\ \mathcal{D}_t &:= \begin{pmatrix} 2(a_t - a_{+,t}) & 0 & 0 & 0 \\ 0 & 2a_t - a_{+,t} - a_{-,t} & 0 & 0 \\ 0 & 0 & 2a_t - a_{+,t} - a_{-,t} & 0 \\ 0 & 0 & 0 & 2(a_t - a_{-,t}) \end{pmatrix} \end{aligned} \quad (5.44)$$

with

$$a_{+,t} = a_+(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}) := \langle M_{12,t}^{E_+} E_+ \rangle, \quad a_{-,t} = a_-(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}) := -\langle M_{12,t}^{E_-} E_- \rangle$$

and<sup>11</sup>

$$a_t := \begin{cases} a_{+,t} & \text{if } \eta_{1,t}\eta_{2,t} < 0, \\ a_{-,t} & \text{if } \eta_{1,t}\eta_{2,t} > 0, \end{cases} \quad b_t := b(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t}). \quad (5.45)$$

<sup>11</sup>In this proof we assumed that  $\eta_{i,t} > 0$  for simplicity, but here we give the definition of  $a_t$  for all cases.

Notice that by (5.27)  $\mathcal{D}_t$  is a positive matrix. Additionally, define

$$\mathcal{M}_t := \begin{pmatrix} 2a_t & ib_t & -ib_t & 0 \\ ib_t & 2a_t & 0 & -ib_t \\ -ib_t & 0 & 2a_t & ib_t \\ 0 & -ib_t & ib_t & 2a_t \end{pmatrix}, \quad (5.46)$$

so that  $\widehat{\mathcal{M}}_t = \mathcal{M}_t - \mathcal{D}_t$ . We point out that the four eigenvalues of  $\mathcal{M}_t$  are given by  $2a_t, 2a_t, 2a_t \pm 2ib_t$ .

Then, using these notations, by (5.41), it follows that

$$d\mathcal{Y}_t = (\mathcal{M}_t - \mathcal{D}_t)\mathcal{Y}_t dt + \mathcal{F}_t dt + d\mathcal{E}_t, \quad (5.47)$$

with  $\mathcal{F}_t$  being a forcing term, and  $\mathcal{E}_t$  being a  $4 \times 4$  matrix valued martingale such that

$$\int_0^{t \wedge \tau} \|\mathcal{F}_s\| ds + \left( \left\| \left[ \int_0^{\cdot} d\mathcal{E}_s \right]_{t \wedge \tau} \right\| \right)^{1/2} \lesssim n^\xi \alpha_{t \wedge \tau}. \quad (5.48)$$

In order to estimate  $\mathcal{Y}_t$ , the solution of (5.47), we will rely on the following simple lemma which is a stochastic matrix-valued Gronwall inequality:

**Lemma 5.6.** *Fix  $d \in \mathbb{N}$ , and let  $\mathcal{X}_t \in \mathbb{C}^d$  be the solution of the following matrix valued SDE:*

$$d\mathcal{X}_t = \mathcal{P}_t \mathcal{X}_t dt + \mathcal{F}_t dt + d\mathcal{E}_t,$$

with given initial condition  $\mathcal{X}_0$ , where the forcing term  $\mathcal{F}_t \in \mathbb{C}^d$  and the coefficient matrix  $\mathcal{P}_t \in \mathbb{C}^{d \times d}$  are adapted to continuous family of  $\sigma$ -algebras and  $\mathcal{E}_t \in \mathbb{C}^d$  is a martingale. We also assume that  $\|\mathcal{F}_t\| + \|\mathcal{E}_t\| \leq n^C$  for some large parameter  $n$  and for some constant  $C > 0$ . Furthermore, assume that  $\text{Re Spec}(\mathcal{P}_t) \subset (-\infty, f_t]$ , for some non-negative real valued function  $f_t \geq 0$ . Let  $\mathcal{C}_t dt \in \mathbb{C}^{d \times d}$  be the covariation process of  $d\mathcal{E}_t$ , let  $\tau$  be a stopping time, and assume that

$$\left( \int_0^{t \wedge \tau} \|\mathcal{F}_s\| ds \right)^2 + \int_0^{t \wedge \tau} \|\mathcal{C}_s\| ds \lesssim \alpha_{t \wedge \tau}^2, \quad (5.49)$$

for some positive deterministic function  $\alpha_t$ . Then, with very high probability, for any arbitrary small  $\zeta > 0$  and for any  $t \geq 0$ , we have

$$\sup_{0 \leq s \leq t \wedge \tau} \|\mathcal{X}_s\|^2 \lesssim \|\mathcal{X}_0\|^2 + n^{3\zeta} \alpha_t^2 + \int_0^{t \wedge \tau} (\|\mathcal{X}_0\|^2 + n^{3\zeta} \alpha_s^2) f_s \exp\left(2(1 + n^{-\zeta}) \int_s^{t \wedge \tau} f_r dr\right) ds. \quad (5.50)$$

*Proof.* We consider the evolution of  $\|\mathcal{X}_t\|^2$ :

$$d\|\mathcal{X}_t\|^2 = 2\langle \mathcal{X}_t, (\text{Re } \mathcal{P}_t) \mathcal{X}_t \rangle dt + 2\text{Re} \langle \mathcal{F}_t, \mathcal{X}_t \rangle dt + 2\text{Re} \langle d\mathcal{E}_t, \mathcal{X}_t \rangle + \text{Tr}[\mathcal{C}_t] dt.$$

Integrating this in time, we get

$$\|\mathcal{X}_t\|^2 \leq \|\mathcal{X}_0\|^2 + 2 \int_0^t f_s \|\mathcal{X}_s\|^2 ds + C \int_0^t \|\mathcal{F}_s\| \cdot \|\mathcal{X}_s\| ds + 2 \int_0^t \text{Re} \langle d\mathcal{E}_s, \mathcal{X}_s \rangle + \int_0^t \|\mathcal{C}_s\| ds \quad (5.51)$$

for some  $C = C(d) > 0$ . Define  $\mathcal{Z}_t := \sup_{0 \leq s \leq t} \|\mathcal{X}_s\|^2$ , then, to estimate the stochastic term in (5.51) we use the martingale inequality (see [102, Appendix B.6, Eq. (18)] with  $c = 0$  for continuous martingales)

$$\mathbf{P} \left( \sup_{0 \leq s \leq t} \left| \int_0^{s \wedge \tau} \text{Re} \langle d\mathcal{E}_r, \mathcal{X}_r \rangle \right| \geq \lambda, \quad \left[ \int_0^{\cdot} \text{Re} \langle d\mathcal{E}_s, \mathcal{X}_s \rangle \right]_{t \wedge \tau} \leq \mu \right) \leq 2e^{-\lambda^2/2\mu}, \quad (5.52)$$

for any fixed  $\lambda, \mu > 0$ . Using the boundedness of  $\mathcal{E}_t, \mathcal{X}_t$  and a dyadic argument, (5.52) implies

$$\sup_{0 \leq s \leq t} \left| \int_0^{s \wedge \tau} \text{Re} \langle d\mathcal{E}_r, \mathcal{X}_r \rangle \right| \leq \log n \left[ \int_0^{\cdot} \text{Re} \langle d\mathcal{E}_s, \mathcal{X}_s \rangle \right]_{t \wedge \tau}^{1/2} + n^{-100}, \quad (5.53)$$

with very high probability. Then, using that by (5.49) we have

$$\left[ \int_0^{\cdot} \text{Re} \langle d\mathcal{E}_s, \mathcal{X}_s \rangle \right]_{t \wedge \tau} \lesssim \int_0^{t \wedge \tau} \langle \mathcal{X}_s, \mathcal{C}_s \mathcal{X}_s \rangle ds \lesssim \int_0^{t \wedge \tau} \|\mathcal{C}_s\| \cdot \|\mathcal{X}_s\|^2 ds \lesssim \mathcal{Z}_{t \wedge \tau} \alpha_{t \wedge \tau}^2,$$

the estimate (5.53), plugged into (5.51), implies

$$\mathcal{Z}_{t \wedge \tau} \leq \|\mathcal{X}_0\|^2 + 2 \int_0^{t \wedge \tau} f_s \mathcal{Z}_s ds + \alpha_{t \wedge \tau} \mathcal{Z}_{t \wedge \tau}^{1/2} \log n + C \alpha_{t \wedge \tau}^2, \quad (5.54)$$

with very high probability. Using  $\alpha_{t \wedge \tau} \mathcal{Z}_{t \wedge \tau}^{1/2} \leq n^{2\zeta} \alpha_{t \wedge \tau}^2 + n^{-\zeta} \mathcal{Z}_{t \wedge \tau}$ , we thus conclude (neglecting  $\log n$ -factors)

$$\mathcal{Z}_{t \wedge \tau} \leq (1 + n^{-\zeta}) \|\mathcal{X}_0\|^2 + 2(1 + n^{-\zeta}) \int_0^{t \wedge \tau} f_s \mathcal{Z}_s ds + n^{3\zeta} \alpha_{t \wedge \tau}^2.$$

Then by a standard Gronwall inequality we conclude the proof.  $\square$

In order to apply this lemma, we now rewrite  $a_t$  from (5.45) as a time derivative of a certain quantity plus a lower order term (its proof is given in Appendix A):

**Lemma 5.7.** *Let  $a_t$  from (5.45), and define<sup>12</sup>*

$$\beta_{\pm, t} := 1 - \operatorname{Re} [z_{1, t} \overline{z_{2, t}}] u_{1, t} u_{2, t} \pm \sqrt{m_{1, t}^2 m_{2, t}^2 - (\operatorname{Im} [z_{1, t} \overline{z_{2, t}}])^2 u_{1, t}^2 u_{2, t}^2}. \quad (5.55)$$

Then, we have

$$a_t = -\frac{1}{2} \partial_t \log |\beta_{+, t} \beta_{-, t}| + \frac{|m_{1, t} m_{2, t}|}{|\beta_{+, t} \beta_{-, t}|}. \quad (5.56)$$

Additionally, we have

$$|\beta_{\pm, t}| \sim \gamma_t = |z_{1, t} - z_{2, t}| + \frac{\eta_{1, t}}{\rho_{1, t}} + \frac{\eta_{2, t}}{\rho_{2, t}}. \quad (5.57)$$

We are now ready to conclude the proof of Proposition 5.4. Using that the largest real part of the eigenvalues of  $\mathcal{M}_t$  is  $2a_t$  and that  $\mathcal{D}_t$  is positive, the propagator of (5.47) has operator norm bounded by

$$\exp \left( \int_s^t 2a_r dr \right) = \exp \left[ - \int_s^t \left( \partial_r \log |\beta_{+, r} \beta_{-, r}| + \frac{|m_{1, r} m_{2, r}|}{|\beta_{+, r} \beta_{-, r}|} \right) dr \right] \lesssim \frac{|\beta_{+, s} \beta_{-, s}|}{|\beta_{+, t} \beta_{-, t}|} \sim \frac{\gamma_s^2}{\gamma_t^2}. \quad (5.58)$$

Here, by  $|\beta_{\pm, r}| \sim \gamma_r \gtrsim \eta_{i, r} / \rho_{i, r}$  from (5.57), we used that

$$\int_s^t \frac{|m_{1, r} m_{2, r}|}{|\beta_{+, r} \beta_{-, r}|} dr \lesssim \int_s^t \frac{\rho_{1, r}^2 \rho_{2, r}^2}{\eta_{1, r} \eta_{2, r}} dr \leq \int_s^t \left( \frac{\rho_{1, r}^4}{\eta_{1, r}^2} + \frac{\rho_{2, r}^4}{\eta_{2, r}^2} \right) dr \lesssim \frac{\rho_{1, t}^3}{\eta_{1, t}} + \frac{\rho_{2, t}^3}{\eta_{2, t}} \lesssim 1,$$

where in the last inequality we used that  $\rho_{i, t} \lesssim \eta_{i, t}^{1/3}$  by (3.7). The leading term  $\partial_r \log |\beta_{+, r} \beta_{-, r}|$  has been explicitly integrated out in (5.58).

We now conclude the proof of Proposition 5.4 applying Lemma 5.6 for the equation (5.47). The fact that  $\mathcal{F}_t, d\mathcal{E}_t$  in (5.47) satisfy (5.49), with  $\alpha_t$  defined in (5.23), follows from (5.48). Using (5.58), we can thus apply Lemma 5.6 (with  $3\zeta = \xi$ ), for  $f_t = 2a_t \lesssim 1/\gamma_t$ , to obtain

$$\|\mathcal{Y}_t\|^2 \lesssim n^{3\xi} \alpha_t^2 + n^{3\xi} \int_0^t \frac{\alpha_s^2}{\gamma_s} \exp \left( \int_s^t 4a_r dr \right) ds \lesssim n^{3\xi} \alpha_t^2 + n^{3\xi} \int_0^t \frac{\alpha_s^2 \gamma_s^3}{\gamma_t^4} ds \lesssim n^{3\xi} \alpha_t^2, \quad (5.59)$$

with very high probability, where we used that  $\|\mathcal{Y}_0\| \lesssim \alpha_0 \leq \alpha_s$ , by the monotonicity of  $t \mapsto \alpha_t$ . The estimate of the integral in the last step will be explained later, first we conclude the main proof. This proves the desired bound for the initial condition  $\eta_{i, 0}$ . We point out that a completely analogous proof holds if we consider any other initial condition  $\tilde{\eta}_{i, 0} \in [\eta_{i, 0}, \omega_1]$ , obtaining exactly the same bound as in the last line of (5.59) but with a tilde on  $\eta$ 's and on all quantities depending on them. This together, with the definition of  $\tau$  (5.24) (note the  $n^{2\xi}$  threshold!) shows that  $\tau = T_1^\epsilon \wedge T_2^\epsilon$ , with very high probability. In order to conclude the proof we are now left only with the proof of the last inequality in (5.59).

We close this proof proving the last inequality of (5.59). Here we used (recall the definition of  $\alpha_t$  from (5.23))

$$\int_0^t \frac{\alpha_s^2 \gamma_s^3}{\gamma_t^4} ds = \int_0^t \left( \frac{1}{\sqrt{n} \ell_s \gamma_s} \wedge \frac{1}{n \eta_{1, s} \eta_{2, s}} \right)^2 \frac{\gamma_s^3}{\gamma_t^4} ds \leq \frac{1}{n \ell_t \gamma_t^2} \wedge \frac{1}{n^2 \eta_{1, t}^2 \eta_{2, t}^2} = \alpha_t^2. \quad (5.60)$$

Finally, we prove the last inequality (5.60). Let  $s_*$  be such that for any  $s \geq s_*$  we have  $\gamma_s \sim |z_{1, s} - z_{2, s}|$  and  $\gamma_s \sim (\eta_{1, s} / \rho_{1, s}) + (\eta_{2, s} / \rho_{2, s})$  for  $s < s_*$  (by the monotonicity of  $t \rightarrow \eta_{i, t} / \rho_{i, t}$  it follows that at least one such  $s_*$  exists, maybe  $s_* = 0$  or  $s_* = t$ ). We thus have

$$\int_{s_*}^t \left( \frac{1}{\sqrt{n} \ell_s \gamma_s} \wedge \frac{1}{n \eta_{1, s} \eta_{2, s}} \right)^2 \frac{\gamma_s^3}{\gamma_t^4} ds \lesssim \frac{1}{n^2 \gamma_t} \int_0^t \frac{1}{\eta_{1, s}^2 \eta_{2, s}^2} ds \leq \frac{1}{n^2 \gamma_t \eta_{2, t}^2} \int_0^t \frac{1}{\eta_{1, s}^2} ds \lesssim \frac{1}{n^2 \eta_{1, t}^2 \eta_{2, t}^2},$$

<sup>12</sup>Here  $\beta_{\pm, t}$  are the eigenvalues of the time-dependent two-body stability operator defined around (5.12) (see also (A.1) below). We use the standard convention that the complex square root  $\sqrt{\cdot}$  is defined using the branch cut  $\mathbb{C} \setminus (-\infty, 0)$ .

where in the first inequality we used that  $\gamma_{s_*} \sim |z_{1,s} - z_{2,s}| \sim |z_{1,0} - z_{2,0}|$  implies  $\gamma_s \sim |z_{1,0} - z_{2,0}|$  for any  $s_* \leq s \leq t$ , and so that  $\gamma_s/\gamma_t \sim 1$ , and in the last inequality we used the first relation in (5.8) together with the bound  $\gamma_t \geq \eta_{1,t}/\rho_{1,t}$ . Additionally, we also have the estimate

$$\int_{s_*}^t \left( \frac{1}{\sqrt{n\ell_s}\gamma_s} \wedge \frac{1}{n\eta_{1,s}\eta_{2,s}} \right)^2 \frac{\gamma_s^3}{\gamma_t^4} ds \lesssim \frac{1}{n\gamma_t^3} \int_0^t \frac{1}{\ell_s} ds \leq \frac{1}{n\ell_t\gamma_t^2},$$

where in the last inequality we used (5.40). For the complementary regime, we have

$$\begin{aligned} \int_0^{s_*} \left( \frac{1}{\sqrt{n\ell_s}\gamma_s} \wedge \frac{1}{n\eta_{1,s}\eta_{2,s}} \right)^2 \frac{\gamma_s^3}{\gamma_t^4} ds &\lesssim \frac{1}{n^2\gamma_t^4} \int_0^{s_*} \frac{\gamma_s^3}{\eta_{1,s}^2\eta_{2,s}^2} ds \\ &\sim \frac{1}{n^2\gamma_t^4} \int_0^t \left[ \frac{\eta_{2,s}}{\eta_{1,s}^2\rho_{2,s}^3} + \frac{\eta_{1,s}}{\eta_{2,s}^2\rho_{1,s}^3} \right] ds \\ &\lesssim \frac{1}{n^2\gamma_t^4} \int_0^t \left[ \frac{\eta_{2,t}}{\eta_{1,s}\eta_{1,t}\rho_{2,t}^3} + \frac{\eta_{1,t}}{\eta_{2,s}\eta_{2,t}\rho_{1,t}^3} \right] ds \\ &\quad + \frac{1}{n^2\gamma_t^4\rho_{1,t}\rho_{2,t}} \int_0^t \left[ \frac{1}{\eta_{1,s}\rho_{2,t}} + \frac{1}{\eta_{2,s}\rho_{1,t}} \right] ds \\ &\lesssim \frac{1}{(n\gamma_t)^2\rho_{1,t}\rho_{2,t}\eta_{1,t}\eta_{2,t}} \\ &\lesssim \frac{1}{(n\ell_t\gamma_t)^2} \wedge \frac{1}{(n\eta_{1,t}\eta_{2,t})^2}, \end{aligned}$$

where in the first  $\sim$  relation we used that  $\gamma_s \sim (\eta_{1,s}/\rho_{1,s}) + (\eta_{2,s}/\rho_{2,s})$  in this regime. In the second inequality we used  $\rho_{i,s} \sim \rho_{i,t}$  and we estimated  $\eta_{i,s}/\rho_{i,s}$  by using

$$\frac{\eta_{i,s}/\rho_{i,s}}{\eta_{j,s}/\rho_{j,s}} \lesssim 1 + \frac{\eta_{i,t}/\rho_{i,t}}{\eta_{j,t}/\rho_{j,t}},$$

that follows from the third relation in (5.6). Finally, in the last two inequalities we used (5.8) (ignoring log  $n$ -factors) and  $\rho_{1,t}\rho_{2,t}\gamma_t^2 \geq \eta_{1,t}\eta_{2,t}$ .  $\square$

**5.2. Improved local law for  $G^zFG^zF^*$  (Theorem 3.5).** As we mentioned after Proposition 5.4, its Part 1 would hold for any observables  $A_1, A_2$ , giving a bound of order  $1/(n\eta^2)$ , considering  $\eta_1 = \eta_2 = \eta$  for simplicity. Now we aim to improve this bound for the specific observables  $F, F^*$  to gain an extra  $\rho^2$ , i.e. we gain a factor  $\rho$  per  $F$  or  $F^*$  matrix. Moreover, along the way we also need an improved single-resolvent local law with  $F$ , and a two-resolvent local law when one observable is  $F$  and the other is the identity as these quantities naturally emerge. These three local laws will be proven simultaneously. This plan justifies the following short-hand notations.

Recall the definition of the characteristics  $\Lambda_t$  in (5.3), define the resolvent  $G_t = G^{z_t}(i\eta_t) := (W_t - \Lambda_t)^{-1}$ , and let

$$M_t^A := M_{12}^A(\eta_t, z_t, \eta_t, z_t) \quad (5.61)$$

be the deterministic approximation of  $G_tAG_t$  as defined in (5.12). Note that unlike in the previous section here we only consider resolvents evaluated at the same spectral parameters since this is simpler and enough for the current application, in particular in this section we drop the 12 indices from

$$M^A := \mathcal{B}^{-1}[MAM] := (1 - M\mathcal{S}[\cdot]M)^{-1}[MAM]. \quad (5.62)$$

We define

$$\begin{aligned} \Phi_1(t) &:= \frac{n|\eta_t|}{\rho_t} |\langle (G_t - M_t)F \rangle|, \\ \Phi_F(t) &:= \frac{n|\eta_t|^2}{\rho_t} |\langle G_tFG_t - M_t^F \rangle|, \\ \Phi_2(t) &:= \frac{\sqrt{n}|\eta_t|^{3/2}}{\rho_t^{5/2}} |\langle (G_tFG_t - M_t^F)F^* \rangle|, \end{aligned} \quad (5.63)$$

where we recall that  $\rho_t = \pi^{-1}|\text{Im } m^{z_t}(i\eta_t)|$ . We will show that the bound  $\Phi_1 + \Phi_F + \Phi_2 < 1$  propagates along the characteristic. We point that that the bound  $\Phi_1(t) + \Phi_F(t) < 1$  is optimal, but the bound

$\Phi_2(t) \prec 1$  is not, i.e. the optimal bound for  $\langle (G_t F G_t - M_t^F) F^* \rangle$  would be better by a factor  $\sqrt{n|\eta|\rho}$ . We do not pursue this additional gain here as it is not needed for the purpose of this paper. Furthermore, we can treat  $F$  and  $F^*$  on the same footing, in particular it is not necessary to define  $\Phi_F(t)$  for  $F^*$  instead of  $F$  as  $\langle (G_t F G_t)^* \rangle = \langle G_t^* F G_t^* \rangle$ . We also point out that by the first two equalities in (3.32) it follows that it is sufficient to define  $\Phi_F$  only when  $G_t F G_t$  is tested against the identity and not against  $E_-$ . We are now ready to state the main result of this section.

**Proposition 5.8.** *Fix  $n$ -independent constants  $\epsilon, \omega_1 > 0$ , and define the spectral parameter*

$$\Lambda_0 := \begin{pmatrix} i\eta_0 & z_0 \\ z_0 & i\eta_0 \end{pmatrix}, \quad (5.64)$$

with  $|z_0| \geq 1$ , and  $\eta_{i,0} \neq 0$ . For  $0 \leq t < T^*(\eta_0, z_0)$ , let  $\Lambda_t$  be the solution of (5.3) with initial condition  $\Lambda_0$ . Let  $\Phi_1(t), \Phi_2(t), \Phi_F(t)$  be defined as in (5.63), choose  $|\eta_0| \leq \omega_1$  and assume that for a small  $0 < \xi \leq \epsilon/10$  it holds

$$\Phi_1(0) + \Phi_F(0) + \Phi_2(0) \lesssim n^\xi, \quad (5.65)$$

with very high probability. Then

$$\Phi_1(t) + \Phi_F(t) + \Phi_2(t) \lesssim n^{2\xi}, \quad (5.66)$$

with very high probability uniformly in  $t \leq T^*(z_0, \eta_0)$  and spectral parameters satisfying  $n\eta_t \rho_t \geq n^\epsilon$ .

We are now ready to conclude the proof of Theorem 3.5, and then we present the proof of Proposition 5.8.

*Proof of Theorem 3.5 for Gaussian component.* Using the global law (5.19) as an input, the proof of this theorem for matrices with an order one Gaussian component is very similar (in fact simpler) to the proof of Theorem 3.3. The fact that the Gaussian component can be removed is proven in Section D.4.  $\square$

*Proof of Proposition 5.8.* Some estimates in this proof are similar to the ones presented in the proof of Proposition 5.4. We thus only focus on the main differences. For notation simplicity, from now on we assume that  $\eta_t > 0$ .

We now state some bounds on the deterministic approximations appearing in (5.63) (the proof is postponed to Appendix A):

**Lemma 5.9.** *Let  $H^z$  be defined as in (1.3), let  $G^z = (H^z - i\eta)^{-1}$ , and let  $M^z, M_{12}^F$  be defined as in (3.1) and (3.13) with  $z = z_1 = z_2 = z$ ,  $\eta_1 = \eta_2 = \eta$ , and  $A = F$ , respectively. Then there exists  $C > 0$  such that*

$$|\langle M_{12}^F F^* \rangle| \leq C \frac{\rho^3}{\eta}, \quad |\langle M_{12}^F \rangle| \leq C \frac{\rho^2}{\eta}, \quad 0 \leq \langle M_{12}^{E^+} \rangle \leq \frac{\text{Im } m}{\eta}. \quad (5.67)$$

Note that the last bound in (5.67) needs to be sharp; no multiplicative constant  $C$  could be afforded.

We will consider the time evolution of the quantities  $\Phi_1(t), \Phi_2(t), \Phi_F(t)$  for times  $t \leq \tilde{\tau}$ , with  $\tilde{\tau}$  the stopping time defined by

$$\tilde{\tau} := \inf \{ t \geq 0 : \max\{\Phi_1(t), \Phi_2(t), \Phi_F(t)\} = n^{2\xi} \} \wedge T^\epsilon, \quad (5.68)$$

for some  $0 < \xi \leq \epsilon/10$ , and  $T^\epsilon := \min\{t > 0 : n\ell_t = n^\epsilon\}$  is defined analogously to (5.21). Note that  $\tilde{\tau} > 0$  with very high probability as a consequence of our assumption (5.65).

We start considering  $\Phi_1(t)$ . By Itô's formula we immediately see that

$$d\langle (G_t - M_t)F \rangle = \frac{1}{\sqrt{n}} \sum_{a,b=1}^{2n} \langle G_t F \rangle dB_{ab,t} + \frac{1}{2} \langle (G_t - M_t)F \rangle + \frac{1}{2} \langle G_t - M_t \rangle \langle G_t^2 F \rangle, \quad (5.69)$$

where we used (1.5) and that  $\langle G_t E_- \rangle = 0$  by the third equality in (3.32). The second term in the rhs. of (5.69) can be incorporated into the lhs. by considering the evolution of  $e^{-t/2} \langle (G_t - M_t)F \rangle$  and thus neglected (recall that  $t \ll 1$ ).

The quadratic variation of the stochastic term in (5.69) is equal to

$$\frac{1}{n^2 \eta_t^2} \langle \text{Im } G_t F \text{Im } G_t F^* \rangle dt.$$

Integrating (5.69) in time, for  $t \leq \tilde{\tau}$ , and using the BDG inequality, we thus estimate the stochastic term by (recall  $\ell_t = \eta_t \rho_t$ )

$$\frac{n\eta_t}{\rho_t} \left( \int_0^t \frac{1}{n^2\eta_s^2} \langle \text{Im } G_s F \text{Im } G_s F^* \rangle ds \right)^{1/2} \lesssim \frac{n\eta_t}{\rho_t} \left( \int_0^t \left( \frac{\rho_s^3}{n^2\eta_s^3} + \frac{n^{2\xi}\rho_s^{5/2}}{n^{5/2}\eta_s^{7/2}} \right) ds \right)^{1/2} \lesssim 1 \quad (5.70)$$

where in the first inequality we used Lemma 5.9, together with the definition of the stopping time (5.68) to control  $\Phi_2(s) \leq n^{2\xi}$ , while in the second step we used (5.8) as before, and that  $n^{10\xi} \leq n\ell_s$ , for any  $0 \leq s \leq \tilde{\tau}$ .

Similarly, for the last term in the rhs. of (5.69) we estimate

$$\frac{n\eta_t}{\rho_t} \int_0^t |\langle G_s - M_s \rangle| |\langle G_s^2 F \rangle| ds \lesssim \frac{n\eta_t}{\rho_t} \int_0^t \frac{n^\xi}{n\eta_s} \left( \frac{\rho_s^2}{\eta_s} + \frac{n^{2\xi}\rho_s}{n\eta_s^2} \right) ds \lesssim 1, \quad (5.71)$$

with very high probability.

Using (5.65) to estimate  $\Phi_1(0) \lesssim n^\xi$ , and integrating (5.69), we thus obtain

$$\Phi_1(t) \lesssim n^\xi, \quad \text{with very high probability, for any } t \leq \tilde{\tau}. \quad (5.72)$$

Next, we consider  $\Phi_2(t)$ . The evolution of  $\langle (G_{1,t} F G_{2,t} - M_{12,t}^F) F^* \rangle$  is given as in (5.14) choosing  $A_1 = F$  and  $A_2 = F^*$ . In the following bounds we will repeatedly use Lemma 5.9 to estimate the deterministic terms, and the definition of the stopping time  $\tilde{\tau}$  for the fluctuation even if not stated explicitly. All the following estimates are performed for  $t \leq \tilde{\tau}$ , even if not stated explicitly. We estimate the quadratic variation by (we only write the bound for a representative term, the rest is analogous)

$$\frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \left( \int_0^t \frac{1}{n^2\eta_s^2} \langle \text{Im } G_s F G_s F^* \text{Im } G_s F G_s^* F^* \rangle ds \right)^{1/2} \lesssim \frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \left( \int_0^t \frac{1}{n\eta_s^2} \left( \frac{\rho_s^6}{\eta_s^2} + \frac{n^{2\xi}\rho_s^5}{n\eta_s^3} \right) ds \right)^{1/2} \lesssim 1. \quad (5.73)$$

We point out that in the first inequality we used that  $F G_s F^* = i F \text{Im } G_s F^*$  (see last identity in (3.32)) to obtain

$$\langle \text{Im } G_s F G_s F^* \text{Im } G_s F G_s^* F^* \rangle = \langle \text{Im } G_s F \text{Im } G_s F^* \text{Im } G_s F \text{Im } G_s^* F^* \rangle \leq n \langle \text{Im } G_s F \text{Im } G_s F^* \rangle^2 \quad (5.74)$$

to reduce the trace of the product of four resolvents in terms of traces of two resolvents. The second equality in (5.74) follows analogously to (5.30) and we thus omit the proof.

For the two terms in the last line of (5.14) we use the estimate (we only write the bound for one term)

$$\frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \int_0^t |\langle G_s - M_s \rangle| |\langle G_s^2 F G_s F^* \rangle| ds \lesssim \frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \int_0^t \frac{n^\xi}{n\eta_s^2} \left( \frac{\rho_s^3}{\eta_s} + \frac{n^{2\xi}\rho_s^{5/2}\phi_2}{\sqrt{n\eta_s^{3/2}}} \right) ds \lesssim \frac{n^\xi}{\sqrt{n\ell_t}}, \quad (5.75)$$

where in the first inequality we used that

$$|\langle G_s^2 F G_s F^* \rangle| = |\langle G_s^2 F \text{Im } G_s F^* \rangle| \leq \frac{1}{\eta_t} \langle \text{Im } G_s F \text{Im } G_s F^* \rangle.$$

Next, by

$$\langle (G_s F G_s - M_s^F) E_1 \rangle = \frac{1}{2} \langle G_s F G_s - M_s^F \rangle + \frac{1}{2} \langle (G_s F G_s - M_s^F) E_- \rangle = \frac{1}{2} \langle G_s F G_s - M_s^F \rangle,$$

where in the last equality we used (A.24), we estimate

$$\frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \int_0^t |\langle (G_s F G_s - M_s^F) E_1 \rangle| |\langle M_s^{F^*} E_2 \rangle| ds \lesssim \frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \int_0^t \frac{n^{2\xi}\rho_s}{n\eta_s^2} \frac{\rho_s^2}{\eta_s} ds \lesssim \frac{n^{2\xi}}{\sqrt{n\ell_t}} \quad (5.76)$$

and

$$\frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \int_0^t |\langle (G_s F G_s - M_s^F) E_1 \rangle| |\langle (G_s F^* G_s - M_s^{F^*}) E_2 \rangle| ds \lesssim \frac{\sqrt{n\eta_t}^{3/2}}{\rho_t^{5/2}} \int_0^t \frac{n^{4\xi}\rho_s^2}{(n\eta_s^2)^2} ds \lesssim \frac{n^{4\xi}}{(n\ell_t)^{3/2}}. \quad (5.77)$$

Using (5.65) to estimate

$$\frac{\sqrt{n\eta_t}^{3/2}}{\sqrt{n\eta_0}^{3/2}} \frac{\rho_0^{5/2}}{\rho_t^{5/2}} \Phi_2(0) \lesssim n^\xi \sqrt{n\ell_t}, \quad (5.78)$$

where we used that  $\rho_0 \sim \rho_t$ , we thus conclude

$$\Phi_2(t) \lesssim n^\xi, \quad \text{with very high probability, for any } t \leq \tilde{\tau}. \quad (5.79)$$

We are now only left with the derivation of the bound for  $\Phi_F$ . The estimates of the various terms in the evolution of  $\langle G_t F G_t - M_t^F \rangle$  are very similar to those for  $\langle (G_t F G_t - M_t^F) F^* \rangle$ , so we only highlight the main differences.

Using the BDG inequality, we estimate of the stochastic term in (5.14):

$$\begin{aligned} \frac{n\eta_t^2}{\rho_t} \left( \int_0^t \frac{1}{n^2\eta_s^2} \langle \text{Im } G_s F G_s \text{Im } G_s G_s^* F^* \rangle ds \right)^{1/2} &\lesssim \frac{n\eta_t^2}{\rho_t} \left( \int_0^t \frac{1}{n^2\eta_s^4} \langle \text{Im } G_s F \text{Im } G_s F^* \rangle ds \right)^{1/2} \\ &\lesssim \frac{n\eta_t^2}{\rho_t} \left( \int_0^t \frac{1}{n^2\eta_s^4} \left( \frac{\rho_s^3}{\eta_s} + \frac{n^{2\xi}\rho_s^{5/2}}{\sqrt{n}\eta_s^{3/2}} \right) ds \right)^{1/2} \\ &\lesssim 1. \end{aligned} \quad (5.80)$$

Note that in (5.80), compared to (5.73)–(5.74), we do not lose an  $n$ -factor to reduce a trace with four  $G$ 's to traces with two  $G$ 's. This is because in the current case the quadratic variation consists of a trace containing only two  $F$ 's that we need to preserve rather than four as in (5.73). This makes the bound (5.80) easier and stronger as we could estimated two of the four  $G$ 's just by  $G_s \text{Im } G_s G_s^* \leq \text{Im } G_s / \eta_s^2$  in the first inequality. Here we also tacitly used the cyclicity of the trace and the positivity of  $\text{Im } G$  to write  $\langle \text{Im } G F G \text{Im } G G^* F^* \rangle = \langle (\text{Im } G)^{1/2} F (G \text{Im } G G^*) F^* (\text{Im } G)^{1/2} \rangle$  before the operator inequality was applied.

Next, we estimate

$$\begin{aligned} \frac{n\eta_t^2}{\rho_t} \int_0^t |\langle G_s - M_s \rangle| |\langle G_s^3 F \rangle| ds &\lesssim \frac{n\eta_t^2}{\rho_t} \int_0^t \frac{1}{\eta_s^{3/2}} |\langle G_s - M_s \rangle| \langle \text{Im } G_s F \text{Im } G_s F^* \rangle^{1/2} \langle \text{Im } G_s \rangle^{1/2} ds \\ &\lesssim \frac{n\eta_t^2}{\rho_t} \int_0^t \frac{n^\xi \rho_s^{1/2}}{n\eta_s^{5/2}} \left( \frac{\rho_s^{3/2}}{\eta_s^{1/2}} + \frac{n^\xi \rho_s^{5/4}}{(n\eta_s^{3/2})^{1/4}} \right) ds \lesssim n^\xi. \end{aligned} \quad (5.81)$$

Using the first equality in (3.32) to turn  $\langle G_s F G_s E_1 \rangle$  into  $\frac{1}{2} \langle G_s F G_s \rangle$  in order to be able to use  $\phi_F$ , we estimate the term in the fifth line of (5.14) by

$$\frac{n\eta_t^2}{\rho_t} \int_0^t |\langle (G_s F G_s - M_s^F) E_1 \rangle| |\langle (G_s E_1 G_s - M_s^{E_1}) E_2 \rangle| ds \lesssim \frac{n\eta_t^2}{\rho_t} \int_0^t \frac{n^{4\xi} \rho_s}{(n\eta_s^2)^2} ds \lesssim \frac{n^{4\xi}}{n\ell_t}, \quad (5.82)$$

where we used

$$|\langle G_s A G_s - M_s^A \rangle B| \lesssim \frac{n^\xi}{n\eta_s^2},$$

for any deterministic  $A, B$  to estimate the second trace in the lhs. of (5.82). This local law follows follows by (3.16). Actually, in this case the proof would be much easier as we only consider the case  $z_1 = z_2$  and  $\eta_1 = \eta_2$ . Furthermore, we estimate the terms in the third line of (5.14) by

$$\frac{n\eta_t^2}{\rho_t} \int_0^t |\langle M_s^F E_1 \rangle| |\langle (G_s E_1 G_s - M_s^{E_1}) E_2 \rangle| ds \lesssim \frac{n\eta_t^2}{\rho_t} \int_0^t \frac{n^\xi \rho_s^2}{n\eta_s^3} ds \lesssim n^\xi. \quad (5.83)$$

For the terms in the second line of (5.14) we write

$$2\langle (G_t F G_t - M_t^F) E_1 \rangle \langle M_t^{E_+} E_2 \rangle + 2\langle (G_t F G_t - M_t^F) E_2 \rangle \langle M_t^{E_+} E_1 \rangle = \langle G_t F G_t - M_t^F \rangle \langle M_t^{E_+} \rangle,$$

where we used the definition (1.5). Then, combining, (5.80)–(5.83), we are left with

$$d\langle G_t F G_t - M_t^F \rangle = \langle M_t^{E_+} \rangle \langle G_t F G_t - M_t^F \rangle dt + d\widehat{e}_t, \quad (5.84)$$

with  $d\widehat{e}_t$  an error term such that

$$\sup_{0 \leq t \leq T} \frac{n\eta_t^2}{\rho_t} \left| \int_0^t d\widehat{e}_s \right| \lesssim n^\xi, \quad (5.85)$$

with very high probability. Using the last inequality of (5.67) and (5.8), the propagator of (5.84) is bounded by

$$\exp \left( \int_0^t \langle M_s^{E_+} \rangle ds \right) \leq \exp \left( \int_0^t \frac{\text{Im } m_s}{\eta_s} ds \right) \sim \frac{\eta_s}{\eta_t}. \quad (5.86)$$

Finally, using a Gronwall inequality and (5.85), we conclude that

$$\Phi_F(t) \lesssim n^\xi \quad \text{with very high probability, for any } t \leq \widetilde{\tau}, \quad (5.87)$$

where we proceeded similarly to (5.78) to estimate the initial condition. Finally, combining (5.72), (5.79), and (5.87), we have shown that  $\widetilde{\tau} = T^\epsilon$  with very high probability, and so that (5.66) holds.  $\square$

## 6. PROOF OF THE GFT: THEOREM 4.1

This section is divided into two parts: in Section 6.1 we present the main technical inputs to prove Theorem 4.1 (Proposition 6.1 and Theorem 6.2 below), informally sketch the main steps of their proofs, and conclude the proof of Theorem 4.1. Then, in Section 6.2 we present the proof of Proposition 6.1; the proof of Theorem 6.2 is postponed to Section 7.

To simplify the presentation of the proof, we may assume, additionally to Assumption 2.1, that there exist  $\alpha, \beta > 0$  such that the probability density of  $\chi$ , denoted by  $\varphi$ , satisfies

$$\varphi \in L^{1+\alpha}(\mathbb{C}), \quad \|\varphi\|_{1+\alpha} \leq n^\beta. \quad (6.1)$$

This condition is used to control the rare event that there is a tiny singular value of  $X - z$  in a simple way (see (6.16) below). However, this restriction can easily be removed using the argument in [121, Section 6.1] (see also [87, Section 2.2]). More precisely, if we consider a matrix  $X$  which does not satisfy (6.1), then we can instead study  $X + n^{-\gamma} X^{\text{Gin}}$ , with  $X^{\text{Gin}}$  a Ginibre matrix independent of  $X$  and  $\gamma > 0$  large, which does satisfy (6.1). The tiny Gaussian component  $n^{-\gamma} X^{\text{Gin}}$  can then be easily removed following the proof of [121, Theorem 23] (or its refinement [87, Lemma 4]) which combines a sampling idea with a standard (sufficiently high order) moment matching technique (see also [60, Remark 2.2]).

**6.1. Proof strategy of Theorem 4.1.** We will sketch only the proof for  $k = 1$ , as the proof for a general  $k \geq 1$  is analogous. By Girko's formula, for any test function  $f \in C_c^2(\mathbb{C})$  and any  $T > 0$ , we have

$$\mathcal{L}_f := \sum_{i=1}^n f(\sigma_i) = -\frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \int_0^T \text{Im Tr } G^z(i\eta) d\eta d^2z + \frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \log |\det(H^z - iT)| d^2z. \quad (6.2)$$

We now summarize the main steps to prove Theorem 4.1 (for  $k = 1$ ):

- Step 1:** We pick a very large  $T$ , e.g.,  $T := n^{100}$  to ensure the last term in (6.2) is very small with very high probability, as stated in (6.14) below.
- Step 2:** Under the density condition in (6.1), a very tiny  $\eta$ -integral over  $[0, \eta_c)$ , with  $\eta_c := n^{-L}$ , for a large  $L > 100$ , can be neglected in the first absolute moment, i.e.,  $L^1(d\mathbf{P})$ -sense; see (6.19) below.
- Step 3:** In order to use the local law in Theorem 3.1 effectively, we subtract a deterministic term  $M^z$  from  $G^z$  on the right side of (6.2) using that for  $|z| > 1$  this term vanishes, see (6.21) below.
- Step 4:** We truncate the smallest eigenvalue at the level  $E_0 := n^{-3/4-\epsilon}$  for a fixed small  $\epsilon > 0$ . Using the precise tail bound of the smallest eigenvalue in Proposition 3.9, the part with  $\mathbb{1}_{\lambda_1^z \leq E_0}$  is negligible in expectation (see (6.23)). Thus we obtain, in the first absolute moment sense, that

$$\mathcal{L}_f \approx -\frac{1}{4\pi} \int \Delta_z f(z) \left( \int_{\eta_c}^T \text{Im Tr}(G^z(i\eta) - M^z(i\eta)) d\eta \right) \mathbb{1}_{\lambda_1^z > E_0} d^2z. \quad (6.3)$$

- Step 5:** We regularize the truncating function  $\mathbb{1}_{\lambda_1^z > E_0}$  in terms of resolvents using Lemma 6.3 below:

$$q_z := q \left( \int_{-E_0}^{E_0} \text{Im Tr } G^z(y + i\eta_0) dy \right), \quad E_0 = n^{-3/4-\epsilon}, \quad \eta_0 := n^{-3\zeta} E_0, \quad (6.4)$$

for some small  $\zeta > 0$ , where  $q : \mathbb{R}_+ \rightarrow [0, 1]$  is a smooth and non-increasing cut-off function with

$$q(y) = 1, \quad \text{if } 0 \leq y \leq 1/9; \quad q(y) = 0, \quad \text{if } y \geq 2/9. \quad (6.5)$$

Thus, using Proposition 3.9, we have  $\mathbf{E}|\mathcal{L}_f - \widehat{\mathcal{L}}_f| = o(1)$ , where

$$\widehat{\mathcal{L}}_f := -\frac{1}{4\pi} \int \Delta_z f(z) \left( \int_{\eta_c}^T \text{Im Tr}(G^z(i\eta) - M^z(i\eta)) d\eta \right) q_z d^2z, \quad (6.6)$$

is obtained from (6.3) with  $\mathbb{1}_{\lambda_1^z > E_0}$  replaced by  $q_z$ . Hence a simple Taylor expansion of  $\mathcal{F}$  yields

$$\left| \mathbf{E}[\mathcal{F}(\mathcal{L}_f)] - \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f)] \right| \leq \|\mathcal{F}'\|_\infty \mathbf{E}|\mathcal{L}_f - \widehat{\mathcal{L}}_f| = o(1). \quad (6.7)$$

The formal statement is presented in Proposition 6.1 below even for a general  $k \geq 1$ .

- Step 6:** Finally it suffices to study  $\mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f)]$ . Using a GFT argument along a dynamical interpolating matrix flow we show that

$$\left| \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f)] - \mathbf{E}^{\text{Gin}}[\mathcal{F}(\widehat{\mathcal{L}}_f)] \right| = o(1), \quad (6.8)$$



see the formal statement in Theorem 6.2 for a general  $k \geq 1$ . The detailed proof of this step, which fundamentally relies on the independence estimates in Proposition 3.8 and the improved local laws in Theorem 3.5, will be presented in Section 7.

We point out that the different approximation steps hold in different senses. The large  $T$  cutoff (Step 1) holds with very high probability, the small  $\eta$  cutoff (Step 2) and the small  $\lambda_1^z$  cutoff (Step 4) work in first moment sense, while the regularisation of the cutoff (Step 5) and the actual comparison with the Ginibre ensemble (Step 6) are only done by comparing expectations.

The proof of Theorem 4.1 then follows directly from the following two results. The proof of the first result, Proposition 6.1 below, will be presented in the next subsection (as summarized in Step 1–5 for  $k = 1$ ).

**Proposition 6.1.** *Under the same assumption of Theorem 4.1, we consider linear statistics  $\mathcal{L}_{f_p} := \sum_{i=1}^n f_p(\sigma_i)$ ,  $p \in [k]$ , for some test functions  $f_p \in C_c^2(\mathbb{C})$  satisfying (4.1) and (4.2) (or alternatively (4.5)). Then there exists a sufficiently large  $L > 100$  (depending on  $\alpha, \beta$  in (6.1)) such that*

$$\left| \mathbf{E}[\mathcal{F}(\mathcal{L}_{f_1}, \dots, \mathcal{L}_{f_k})] - \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_{f_1}, \dots, \widehat{\mathcal{L}}_{f_k})] \right| = O(n^{-\epsilon}), \quad (6.9)$$

where for any  $p \in [k]$  we defined

$$\widehat{\mathcal{L}}_{f_p} := -\frac{1}{4\pi} \int \Delta_z f_p(z) \left( \int_{\eta_c}^T \text{Im Tr}(G^z(i\eta) - M^z(i\eta)) d\eta \right) q_z d^2 z, \quad (6.10)$$

with  $\eta_c = n^{-L}$ ,  $T = n^{100}$ , and  $q_z$  given in (6.4) with the smooth cut-off  $q$  from (6.5).

Once we translated  $\mathcal{L}_f$  to  $\widehat{\mathcal{L}}_f$ , we use the following Green function comparison theorem, whose proof will be presented in Section 7 (as sketched in Step 6 for  $k = 1$ ).

**Theorem 6.2.** *Under the same conditions of Theorem 4.1, we have*

$$\left| \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_{f_1}, \dots, \widehat{\mathcal{L}}_{f_k})] - \mathbf{E}^{\text{Gin}}[\mathcal{F}(\widehat{\mathcal{L}}_{f_1}, \dots, \widehat{\mathcal{L}}_{f_k})] \right| = O(n^{-1/4+C\epsilon}),$$

for some constant  $C > 0$  independent of  $\epsilon > 0$ .

Finally, combining Proposition 6.1 with Theorem 6.2, we conclude the proof Theorem 4.1.

**6.2. Proof of Proposition 6.1.** We follow Steps 1–5 in Section 6.1. Recall the Girko's formula

$$\mathcal{L}_f = -\frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \int_0^T \text{Im Tr} G^z(i\eta) d\eta d^2 z + \frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \log |\det(H^z - iT)| d^2 z, \quad (6.11)$$

where we dropped the subscript index  $p \in [k]$  of  $f_p$  and  $\mathcal{L}_{f_p}$  for notational simplicity.

**Step 1:** We start with showing that, choosing a large  $T = n^{100}$ , the last term in (6.11) is very small with very high probability. This follows directly from

$$\log |\det(H^z - iT)| = 2n \log T + \sum_j \log \left( 1 + \left( \frac{\lambda_j^z}{T} \right)^2 \right) = 2n \log T + O_{\prec} \left( \frac{n^2}{T^2} \right), \quad (6.12)$$

together with the  $L^1$ -bound

$$\int_{\mathbb{C}} |\Delta_z f(z)| d^2 z \lesssim n^{1/2+2\nu+\tau} \leq n^{1/2+\epsilon/20}, \quad (6.13)$$

which directly follows from the  $L^\infty$ -norm bound of  $\Delta f$  in (4.1) and the area of its support (4.2). We thus have

$$\mathcal{L}_f = -\frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \int_0^T \text{Im Tr} G^z(i\eta) d\eta d^2 z + O_{\prec}(n^{-100}), \quad T = n^{100}. \quad (6.14)$$

**Step 2:** We next show that the  $\eta$ -integral over  $[0, \eta_c]$ , with  $\eta_c = n^{-L}$  for some very large  $L > 100$ , is negligible in the first absolute moment sense. By a direct computation we have

$$\int_0^{\eta_c} \text{Im Tr} G^z(i\eta) d\eta = \frac{1}{2} \left( \sum_{|\lambda_i^z| \leq n^{-L}} + \sum_{n^{-L} < |\lambda_i^z| \leq E_0} + \sum_{|\lambda_i^z| > E_0} \right) \log \left( 1 + \frac{n^{-2L}}{(\lambda_i^z)^2} \right), \quad (6.15)$$

where  $E_0$  is slightly below the typical eigenvalue spacing near zero, *i.e.*,

$$E_0 = n^{-3/4-\epsilon}, \quad \text{for a fixed small } \epsilon > 0.$$

Note that the last sum in (6.15) can be trivially bounded by  $n^{1-2L}/E_0^2 \leq n^{-100}$ . To estimate the first sum in (6.15), we recall [14, Proposition 5.7], *i.e.*, that under the density condition (6.1), there exists  $C_\alpha > 0$  such that

$$\mathbb{P}\left(|\lambda_1^z| \leq \frac{u}{n}\right) \leq C_\alpha u^{\frac{2\alpha}{1+\alpha}} n^{\beta+1}, \quad z \in \mathbb{C}, u > 0, \quad (6.16)$$

with  $\alpha, \beta$  given in (6.1). Then, following [14, Eq. (5.34)-(5.35)], the first sum in (6.15) can be bounded by

$$\mathbf{E}\left[\sum_{|\lambda_i^z| \leq n^{-L}} \log\left(1 + \frac{n^{-2L}}{(\lambda_i^z)^2}\right)\right] \lesssim n \mathbf{E}[\log \lambda_1^z \mathbf{1}_{\lambda_1^z \lesssim n^{-L}}] \lesssim n^{-100},$$

with  $L > 100$  large enough depending on  $\alpha, \beta$ . The second sum in (6.15) can be bounded by

$$\mathbf{E}\left[\sum_{n^{-L} < |\lambda_i^z| \leq E_0} \log\left(1 + \frac{n^{-2L}}{(\lambda_i^z)^2}\right)\right] \prec (\log n) \mathbf{P}(\lambda_1^z \leq E_0), \quad (6.17)$$

using the rigidity estimate of eigenvalues in (3.12) (*i.e.*,  $\#\{|\lambda_i^z| \leq E_0\} \prec 1$ ). Using the support condition (4.2) or (4.5) and Proposition 3.9, we have

$$\mathbf{P}(\lambda_1^z \leq E_0) \lesssim n^{3/2} E_0^2 e^{-n\delta^2/2} \lesssim \begin{cases} n^{-1/2-2\epsilon}, & \text{if } z \text{ satisfies (4.2),} \\ n^{-1/4-2\epsilon}, & \text{if } z \text{ satisfies (4.5).} \end{cases} \quad (6.18)$$

Combining with the  $L^1$ -norm of  $\Delta f$  in (6.13), we know that the very tiny  $\eta$ -integral is negligible, *i.e.*, there exists a large  $L > 100$  such that

$$\mathbf{E}\left|\int_{\mathbb{C}} \Delta_z f(z) \int_0^{\eta_c} \text{Im Tr } G^z(i\eta) d\eta d^2 z\right| \lesssim n^{-\epsilon}, \quad \eta_c = n^{-L}. \quad (6.19)$$

**Step 3:** By Steps 1 and 2 it suffices to focus on the integral over  $\eta \in [\eta_c, T]$  in (6.11). In order to use the local law in (3.9), we subtract the deterministic term  $M^z(i\eta)$  from  $G^z(i\eta)$  on the right side of (6.14). By a direct computation using (3.1) and (3.2), we have (see also [52, Eq. (3.6)])

$$\partial_\eta m^z(i\eta) = \frac{1-\beta}{\beta}, \quad \partial_\eta u^z(i\eta) = \frac{2u^z m^z}{\beta}, \quad \beta := 1 - (m^z)^2 - (u^z)^2 |z|^2,$$

and thus  $m^z = m^z(i\eta) = \partial_\eta (\log u^z - (m^z)^2)/2$ . For any  $|z| > 1$ , we thus have

$$\int_{\eta_c}^T \langle M^z(i\eta) \rangle d\eta = \int_{\eta_c}^T m^z(i\eta) d\eta = \frac{\log u^z(i\eta) - (m^z(i\eta))^2}{2} \Big|_{\eta=\eta_c}^{\eta=T} = \log |z|^2 + O(n^{-100}),$$

where we used (3.7) and that  $m^z(0) = 0, u^z(0) = \frac{1}{|z|^2}$  for  $|z| > 1$ . Using that  $\log |z|^2$  is harmonic away from zero, *i.e.* on the support of  $f$ , after integration by parts we have

$$\int_{\mathbb{C}} \Delta_z f(z) \int_{\eta_c}^T \text{Tr } M^z(i\eta) d\eta d^2 z = O(n^{-98}). \quad (6.20)$$

Therefore, we conclude that

$$\mathbf{E}\left|\mathcal{L}_f + \frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \left(\int_{\eta_c}^T \text{Im Tr}(G^z(i\eta) - M^z(i\eta)) d\eta\right) d^2 z\right| = O(n^{-\epsilon}). \quad (6.21)$$

**Step 4:** We further truncate the smallest eigenvalue at the level  $E_0 = n^{-3/4-\epsilon}$  as in (6.15), *i.e.*,

$$\mathcal{L}_{\leq E_0} + \mathcal{L}_{> E_0} := -\frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \left(\int_{\eta_c}^T \text{Im Tr}(G^z(i\eta) - M^z(i\eta)) d\eta\right) \left(\mathbf{1}_{\lambda_1^z \leq E_0} + \mathbf{1}_{\lambda_1^z > E_0}\right) d^2 z. \quad (6.22)$$

Using the  $L^1$ -norm of  $\Delta_z f$  in (6.13), the local law in (3.9) and the tail bounds in (6.18), we have the bound

$$\mathbf{E}|\mathcal{L}_{\leq E_0}| \prec (\log n) \int_{\mathbb{C}} |\Delta_z f(z)| \mathbf{P}(\lambda_1^z \leq E_0) d^2 z \lesssim n^{-\epsilon}. \quad (6.23)$$

**Step 5:** For  $\mathcal{L}_{>E_0}$  in (6.22), we regularize the truncating function  $\mathbb{1}_{\lambda_i^z > E_0}$  by showing that

$$\mathbf{E}|\mathcal{L}_{>E_0} - \widehat{\mathcal{L}}_f| = O(n^{-\epsilon}), \quad (6.24)$$

where  $\widehat{\mathcal{L}}_f$  is given in (6.6). The proof of (6.24) relies on the following lemma which links the eigenvalue truncating function  $\mathbb{1}_{\lambda_i^z > E_0}$  with  $q_z$  in terms of the resolvent. We also remark the lemma below is in the same spirit of [67, Lemma 6.1-6.2] designed originally to prove the edge universality for Wigner matrices; we sketch the proof in Appendix B for the convenience of the reader.

**Lemma 6.3.** *Let  $0 \leq |z| - 1 \leq c$  for a small constant  $c > 0$ . Fix any small  $\epsilon' > 0$  and  $0 < \zeta < \epsilon'/10$ . Choose any spectral parameters  $E \gg l \gg l' \gg \eta \gg n^{-1}$  such that*

$$n^{-1+\epsilon'} \leq n^{6\zeta}\eta \leq n^{3\zeta}l' \leq n^\zeta l \leq E \leq Cn^{-3/4}. \quad (6.25)$$

Then we have, with very high probability,

$$\left| \#\{|\lambda_i^z| \leq E\} - \int_{-E}^E \text{Im Tr} G^z(y + i\eta) dy \right| \lesssim \#\{|\lambda_i^z| \in [E - l', E + l']\} + O(n^{-\zeta}), \quad (6.26)$$

and thus with very high probability,

$$\begin{aligned} \int_{|y| \leq E-l} \text{Im Tr} G^z(y + i\eta) dy - O(n^{-\zeta}) &\leq \#\{|\lambda_i^z| \leq E\} \\ &\leq \int_{|y| \leq E+l} \text{Im Tr} G^z(y + i\eta) dy + O(n^{-\zeta}). \end{aligned} \quad (6.27)$$

Recall the definitions of  $q$  and  $q_z$  in (6.5) and (6.4). Using that  $\mathbb{1}_{\lambda_i^z > E_0} = q(\#\{|\lambda_i^z| \leq E_0\})$  and (6.26) for  $E = E_0$ , we have, with very high probability,

$$\left| \mathbb{1}_{\lambda_i^z > E_0} - q_z \right| \lesssim \mathbb{1}_{\#\{|\lambda_i^z| \in [(1-n^{-3\zeta})E_0, (1+n^{-3\zeta})E_0]\} \geq 1} \lesssim \mathbb{1}_{\lambda_i^z \leq 3E_0/2}, \quad (6.28)$$

for some small  $\zeta > 0$ . Using (6.18) with  $3E_0/2$  instead of  $E_0$  and the  $L^1$ -norm of  $\Delta f$  in (6.13), we have

$$\mathbf{E}|\mathcal{L}_{>E_0} - \widehat{\mathcal{L}}_f| \lesssim (\log n) \int_{\mathbb{C}} |\Delta_z f(z)| \mathbf{P}(\lambda_1^z \leq 3E_0/2) d^2 z = O(n^{-\epsilon}). \quad (6.29)$$

Hence putting the above estimates in (6.21), (6.23) and (6.29) together, we obtain

$$\mathbf{E}|\mathcal{L}_f - \widehat{\mathcal{L}}_f| = O(n^{-\epsilon}). \quad (6.30)$$

This holds true for any test function  $f = f_p \in C_c^2(\mathbb{C})$  ( $p \in [k]$ ) satisfying (4.1) and (4.2) (or alternatively (4.5)). Then, since  $\mathcal{F}(w_1, \dots, w_k)$  is a smooth function satisfying (4.3), we obtain

$$\left| \mathbf{E}[\mathcal{F}(\mathcal{L}_{f_1}, \dots, \mathcal{L}_{f_k})] - \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_{f_1}, \dots, \widehat{\mathcal{L}}_{f_k})] \right| \lesssim \max_{|\alpha|=1} \{\|\partial^\alpha \mathcal{F}\|_\infty\} \sum_{p=1}^k \mathbf{E}|\mathcal{L}_{f_p} - \widehat{\mathcal{L}}_{f_p}| = O(n^{-\epsilon}). \quad (6.31)$$

This concludes the proof of Proposition 6.1.  $\square$

## 7. PROOF OF THEOREM 6.2

In this section, we prove Theorem 6.2 via a continuous interpolating flow. Given the initial ensemble  $H^z$  in (1.3) with  $X$  satisfying Assumption 2.1, we consider the following Ornstein-Uhlenbeck matrix flow

$$dH_t^z = -\frac{1}{2}(H_t^z + Z)dt + \frac{1}{\sqrt{n}}d\mathcal{B}_t, \quad Z := \begin{pmatrix} 0 & zI \\ \bar{z}I & 0 \end{pmatrix}, \quad \mathcal{B}_t := \begin{pmatrix} 0 & B_t \\ B_t^* & 0 \end{pmatrix}, \quad (7.1)$$

with initial condition  $H_{t=0}^z = H^z$ , where  $B_t$  is an  $n \times n$  matrix with i.i.d. standard complex valued Brownian motion entries. The matrix flow  $H_t^z$  interpolates between the initial matrix  $H^z$  and the same matrix with  $X$  replaced with an independent complex Ginibre ensemble. The Green function of  $H_t^z$  is denoted by  $G_t^z(i\eta) := (H_t^z - i\eta)^{-1}$ . Now we define  $\widehat{\mathcal{L}}_f(t)$  as in (6.10) but with the time dependent  $G_t^z$ . For notational brevity, we define, for any  $b \in \mathbb{N}$ ,

$$\mathcal{X}^z = \mathcal{X}_t^z := \int_{\eta_c}^T \text{Im Tr}(G_t^z(i\eta) - M^z(i\eta)) d\eta, \quad q_z^{(b)} = q_{z,t}^{(b)} := q^{(b)} \left( \int_{-E_0}^{E_0} \text{Im Tr} G_t^z(y + i\eta_0) dy \right), \quad (7.2)$$

with  $\eta_c$  and  $T$  chosen as in Proposition 6.1, where  $q^{(b)}$  is the  $b$ -th derivative of the smooth and uniformly bounded cut-off function  $q$  in (6.5) with  $E_0$  and  $\eta_0$  chosen as in (6.4). We also use the convention that  $q^{(0)} = q$ ,  $q' = q^{(1)}$  and  $q'' = q^{(2)}$ . Then  $\widehat{\mathcal{L}}_f(t)$  from (6.10) with  $f = f_p$  is given by

$$\widehat{\mathcal{L}}_f(t) := -\frac{1}{4\pi} \int \Delta_z f(z) \mathcal{X}_t^z q_{z,t} d^2 z. \quad (7.3)$$

Since the flow in (7.1) is stochastically Hölder continuous in time, the local law for the Green function in Theorem 3.1 also holds true for the time dependent  $G_t^z(i\eta)$  simultaneously for all  $t \geq 0$ , *i.e.*,

$$\sup_{t \in \mathbb{R}_+} \sup_{z \in \text{supp}(f)} \sup_{\eta_c \leq \eta \leq T} (n\eta) |\langle G_t^z(i\eta) - M^z(i\eta) \rangle| \prec 1. \quad (7.4)$$

The proof of (7.4) uses a standard grid argument together with the Hölder regularity of  $G_t^z(i\eta)$  in all three variables for any  $0 \leq t \leq n^{100}$ ,  $z \in \text{supp}(f)$  and  $\eta_c \leq \eta \leq T$ . For large times  $t \geq n^{100}$  a simple perturbation argument applies; see similar arguments at the beginning of the proof of Proposition 5.4 and around [60, Eq. (4.3)]. Thus the local law in (7.4) implies that  $|\mathcal{X}_t^z| \prec \log n$  for any  $t \in \mathbb{R}_+$ . In the following we often drop the  $t$ -dependence in the above notations, and all the estimates throughout this section hold true uniformly in  $t \in \mathbb{R}_+$  even if not mentioned explicitly.

We next compute how the time-dependent expectation  $\mathbf{E}[F(\widehat{\mathcal{L}}_f(t))]$  evolves along the flow in (7.1). Set  $W_t = H_t^z + Z$ , then  $dH_t^z = -\frac{1}{2}W_t dt + \frac{1}{\sqrt{n}}d\mathcal{B}_t$  and

$$W = W_t = \begin{pmatrix} 0 & X_t \\ X_t^* & 0 \end{pmatrix}, \quad X_t \stackrel{d}{=} e^{-\frac{t}{2}} X + \sqrt{1 - e^{-t}} X^{\text{Gin}}, \quad (7.5)$$

where  $X_\infty \stackrel{d}{=} X^{\text{Gin}}$  is a complex Ginibre matrix which is independent of  $X$ . Due to the above  $2 \times 2$  block structure, in the following we use lower case letters (*e.g.*,  $a, b$ ) to denote the indices taking values in  $[1, n]$  and upper case letters (*e.g.*,  $A, B$ ) to denote the indices taking values in  $[n+1, 2n]$ . We also use Fraktur letters, *e.g.*,  $\mathbf{u}, \mathbf{v}$  to denote the indices ranging fully from 1 to  $2n$ .

We apply Itô's formula to  $\mathcal{F}(\widehat{\mathcal{L}}_f)$  using that  $\partial/\partial w_{aB} = \partial/\partial h_{aB}$  and perform the cumulant expansion formula on the expectation (see [79, Lemma 7.1]). Since the variances of the matrix entries of  $X_t$  are preserved along the flow in (7.5), the second order cumulant terms are exactly cancelled and the following sum starts from the third order terms ( $p+q+1=3$ ), *i.e.*,

$$\begin{aligned} \frac{d}{dt} \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f)] &= \sum_{p+q+1 \geq 3}^{10} \frac{1}{n^{\frac{p+q+1}{2}}} \sum_{a=1}^n \sum_{B=n+1}^{2n} \left( \frac{\kappa^{(p+1,q)}}{p!q!} \mathbf{E} \left[ \frac{\partial^{p+q+1} \mathcal{F}(\widehat{\mathcal{L}}_f)}{\partial h_{aB}^{p+1} \partial h_{Ba}^q} \right] + \frac{\kappa^{(q,p+1)}}{p!q!} \mathbf{E} \left[ \frac{\partial^{p+q+1} \mathcal{F}(\widehat{\mathcal{L}}_f)}{\partial h_{aB}^q \partial h_{Ba}^{p+1}} \right] \right) \\ &\quad + O_{\prec}(n^{-1/4+C\epsilon}), \end{aligned} \quad (7.6)$$

where  $\kappa^{(p,q)}$  are the normalized  $(p, q)$ -cumulants of the time-dependent entries of  $\sqrt{n}X_t$ , in particular  $|\kappa^{(p,q)}| \lesssim 1$  by (2.1). Here we stopped the cumulant expansion at  $p+q+1=10$  so that the truncating error is bounded by  $O_{\prec}(n^{-1/4+C\epsilon})$ . Here we used the differentiation rules in Lemma 7.1 below, the  $L^1$ -norm bound of  $\Delta f$  in (6.13), together with the local law estimates in (7.26) and (7.32) later. This truncation argument is rather standard and has been used many times (see *e.g.* [54, 60, 61]), so we omit the details.

To study the terms on the right side of (7.6), we establish the following differentiation rules (these apply in the complex case). The proof is postponed till Section 7.2.

**Lemma 7.1.** *For any  $\mathbf{u}, \mathbf{v} \in [1, 2n]$ ,  $a \in [1, n]$  and  $B \in [n+1, 2n]$ , we have*

$$\frac{\partial G_{\mathbf{u}\mathbf{v}}^z}{\partial h_{aB}} = -G_{\mathbf{u}a}^z G_{B\mathbf{v}}^z, \quad \frac{\partial G_{\mathbf{u}\mathbf{v}}^z}{\partial h_{Ba}} = -G_{\mathbf{u}B}^z G_{a\mathbf{v}}^z. \quad (7.7)$$

Then for  $\mathcal{X}^z$  and  $q_z^{(b)}$  ( $b \in \mathbb{N}$ ) defined in (7.2), we have the following differentiation rules

$$\frac{\partial \mathcal{X}^z}{\partial h_{Ba}} = -G_{aB}^z(i\eta_c) + O_{\prec}(n^{-100}), \quad \frac{\partial q_z^{(b)}}{\partial h_{Ba}} = -q_z^{(b+1)} [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}] (G_{aB}^z), \quad (7.8)$$

where we used the short-hand notations for any function  $g, h : \mathbb{C} \setminus \mathbb{R} \rightarrow \mathbb{C}$ ,

$$\widetilde{\text{Im}} g(w) := \frac{1}{2i} (g(w) - g(\bar{w})), \quad \mathfrak{D}_{E_0, \eta_0} h := h(E_0 + i\eta_0) - h(-E_0 + i\eta_0), \quad (7.9)$$

with fixed  $E_0$  and  $\eta_0$  chosen in (6.4). The same rules above also hold when we interchange  $a$  and  $B$ . Moreover, both operations  $\widetilde{\text{Im}}$  and  $\mathfrak{D}_{E_0, \eta_0}$  in (7.9) commute with the derivatives  $\partial/\partial h_{Ba}$  or  $\partial/\partial h_{aB}$ .

By direct computations using Lemma 7.1, we have the following estimates on the partial derivatives of  $\widehat{\mathcal{L}}_f$  given by (7.3), whose proof is presented in Section 7.1 below.

**Lemma 7.2.** *For any  $a \in [1, n]$  and  $B \in [n+1, 2n]$ , we have*

$$\left| \frac{\partial \widehat{\mathcal{L}}_f}{\partial h_{Ba}} \right| \prec \left( n^{\frac{3}{4} + c\epsilon} + n^{1 + c\epsilon} \mathbb{1}_{a=B-n} \right) \int_{\text{supp}(f)} \mathbb{1}_{\lambda_{\bar{1}} \leq 3E_0/2} d^2 z, \quad (7.10)$$

for some constant  $c > 0$ . In general for any  $p, q \in \mathbb{N}$ , we have,

$$\left| \frac{\partial^{p+q+1} \widehat{\mathcal{L}}_f}{\partial h_{Ba}^{p+1} \partial h_{aB}^q} \right| \prec \left( n^{\frac{3}{4} + (-\frac{1}{4} + \epsilon)(p+q) + c\epsilon} + n^{1 + c\epsilon} \mathbb{1}_{a=B-n} \right) \int_{\text{supp}(f)} \mathbb{1}_{\lambda_{\bar{1}} \leq 3E_0/2} d^2 z, \quad (7.11)$$

and the same results holds true if we interchange  $a$  with  $B$ .

Using Lemma 7.2, we are now ready to estimate the terms on the right side of (7.6) starting from the third order  $p+q+1=3$ . As an example, we first consider one of the most critical third order terms when  $\partial h_{Ba}$  acts on the function  $\mathcal{F}$  for three times, *i.e.*,

$$\begin{aligned} \left| \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \mathcal{F}^{(3)}(\widehat{\mathcal{L}}_f) \left( \frac{\partial \widehat{\mathcal{L}}_f}{\partial h_{Ba}} \right)^3 \right] \right| &\lesssim \frac{1}{n^{3/2}} \sum_{a \neq B-n} \mathbf{E} \left| \frac{\partial \widehat{\mathcal{L}}_f}{\partial h_{Ba}} \right|^3 + \frac{1}{n^{3/2}} \sum_a \mathbf{E} \left| \frac{\partial \widehat{\mathcal{L}}_f}{\partial h_{a+n,a}} \right|^3 \\ &\lesssim n^{\frac{11}{4} + c\epsilon} \mathbf{E} \left[ \left( \int_{\text{supp}(f)} \mathbb{1}_{\lambda_{\bar{1}} \leq 3E_0/2} d^2 z \right)^3 \right], \end{aligned} \quad (7.12)$$

where we used that  $\mathcal{F}$  has uniformly bounded derivatives and the estimate in (7.10) in the last step. Then we split the triple  $z$ -integrals into three regimes

$$\mathbf{E} \left[ \left( \int_{\text{supp}(f)} \mathbb{1}_{\lambda_{\bar{1}} \leq 3E_0/2} d^2 z \right)^3 \right] = \left( \iiint_{(1)} + \iiint_{(2)} + \iiint_{(3)} \right) \mathbf{P} \left( \lambda_1^{z_j} \leq 3E_0/2, j \in [3] \right) d^2 z_1 d^2 z_2 d^2 z_3,$$

where the regimes are given as follows:

- (1) all  $z_1, z_2, z_3 \in \text{supp}(f)$  are sufficiently far from each other, *i.e.*,  $\min\{|z_1 - z_2|, |z_1 - z_3|, |z_2 - z_3|\} \geq n^{-1/2+\omega}$  for some  $\omega > 0$  to be chosen;
- (2) two of them are close while the remaining one is separated from them, *e.g.*,  $|z_1 - z_2| \leq n^{-1/2+\omega}$  and  $\min\{|z_1 - z_3|, |z_2 - z_3|\} \geq n^{-1/2+\omega}$  (and similarly for other two index combinations);
- (3) all  $z_1, z_2, z_3 \in \text{supp}(f)$  are close to each other, *i.e.*,  $\max\{|z_1 - z_2|, |z_1 - z_3|, |z_2 - z_3|\} \leq n^{-1/2+\omega}$ .

For any  $z_j \in \text{supp}(f)$  satisfying (4.2) with  $\min_{j \neq j'}\{|z_j - z_{j'}|\} \geq n^{-1/2+\omega}$ , choosing  $\omega = 2\epsilon/\gamma$  with the small constant  $\gamma > 0$  from Proposition 3.8, the independence estimate in (3.33) implies that

$$\mathbf{P} \left( \lambda_1^{z_1} \leq 3E_0/2, \dots, \lambda_1^{z_k} \leq 3E_0/2 \right) \lesssim \prod_{j=1}^k \mathbf{P} \left( \lambda_1^{z_j} \leq 2E_0 \right) + n^{-D}, \quad E_0 = n^{-3/4-\epsilon}, \quad (7.13)$$

where we also used that  $0 < \tau < \epsilon/100$ . Combining this with (6.18), we obtain that

$$\mathbf{P} \left( \lambda_1^{z_1} \leq 3E_0/2, \dots, \lambda_1^{z_k} \leq 3E_0/2 \right) \lesssim (n^{-1/2-2\epsilon})^k + n^{-D}. \quad (7.14)$$

Thus, using  $|\text{supp}(f)| \lesssim n^{-1/2+\tau}$ , the integrals over the above three regimes are bounded by

$$\begin{aligned} \mathbf{E} \left[ \left( \int_{\text{supp}(f)} \mathbb{1}_{\lambda_{\bar{1}} \leq 3E_0/2} d^2 z \right)^3 \right] &\lesssim (n^{-1/2+\tau})^3 (n^{-1/2-2\epsilon})^3 + n^{-1+2\omega} (n^{-1/2+\tau})^2 (n^{-1/2-2\epsilon})^2 \\ &\quad + (n^{-1+2\omega})^2 (n^{-1/2+\tau}) (n^{-1/2-2\epsilon}) = O(n^{-3+C\epsilon}), \end{aligned} \quad (7.15)$$

for some constant  $C > 0$  depending on  $\gamma$  from Proposition 3.8. We mention that if  $f$  satisfies the alternative support condition in (4.5), then the tail bound (7.14) becomes weaker  $(n^{-1/4-2\epsilon})^k$ , which is exactly compensated by the smaller volume of the support,  $|\text{supp}(f)| \lesssim n^{-3/4+\tau}$ , hence we obtain the same upper bound as in (7.15). Plugging the above estimates into (7.12), we obtain

$$\frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \mathcal{F}^{(3)}(\widehat{\mathcal{L}}_f) \left( \frac{\partial \widehat{\mathcal{L}}_f}{\partial h_{Ba}} \right)^3 \right] = O_{\prec}(n^{-1/4+C\epsilon}).$$

All the other third order terms in (7.6) can be estimated similarly using Lemma 7.2 and (7.14), e.g.,

$$\begin{aligned} \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \mathcal{F}''(\widehat{\mathcal{L}}_f) \left( \frac{\partial \widehat{\mathcal{L}}_f}{\partial h_{aB}} \right) \left( \frac{\partial^2 \widehat{\mathcal{L}}_f}{\partial h_{Ba}^2} \right) \right] &= O_{\prec}(n^{-1/4+C\epsilon}), \\ \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \mathcal{F}'(\widehat{\mathcal{L}}_f) \left( \frac{\partial^3 \widehat{\mathcal{L}}_f}{\partial h_{Ba}^3} \right) \right] &= O_{\prec}(n^{-1/4+C\epsilon}). \end{aligned} \quad (7.16)$$

Due to the moment condition in (2.1), the higher order terms ( $p+q+1 > 3$ ) can be estimated similarly with much less efforts. Therefore, we conclude from (7.6) that

$$\left| \frac{d}{dt} \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f(t))] \right| = O_{\prec}(n^{-1/4+C\epsilon}). \quad (7.17)$$

Integrating both sides of (7.17) over  $t \in [0, t_0]$  with a sufficiently large  $t_0 = O(\log n)$  to be fixed later, we then obtain that

$$\left| \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f(0))] - \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f(t_0))] \right| = O_{\prec}(n^{-1/4+C\epsilon}). \quad (7.18)$$

Recall that  $H_t^z$  is defined as in (1.3) with the time dependent  $X_t$  interpolating between the initial matrix  $X_0 = X$  and the independent Ginibre ensemble  $X_\infty = X^{\text{Gin}}$  as in (7.5). Owing to the exponential relaxation of  $X_t$  to the Ginibre ensemble, fixing  $t_0 := 10L \log n$  with a large  $L > 100$  from  $\eta_c = n^{-L}$  chosen in (6.19), we have

$$\|G_{t_0}^z(i\eta) - G_\infty^z(i\eta)\| \leq \|G_{t_0}^z(i\eta)\| \|G_\infty^z(i\eta)\| \|X_{t_0} - X_\infty\| \prec n^{-2L}, \quad \eta \geq \eta_c = n^{-L}, \quad (7.19)$$

where we used that  $\|G^z(i\eta)\| \leq \eta^{-1}$  and that  $|X_{ij}| \prec n^{-1/2}$  from the moment assumption in (2.1). Using the  $L^1$ -norm bound of  $\Delta f$  in (6.13), we have

$$\left| \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f(t_0))] - \mathbf{E}[\mathcal{F}(\widehat{\mathcal{L}}_f(\infty))] \right| = O_{\prec}(n^{-10}). \quad (7.20)$$

Combining (7.18) with (7.20), we hence finish the proof of Theorem 6.2.

**7.1. Proof of Lemma 7.2.** The proof relies on the following useful lemmas, whose proofs are postponed to Section 7.2 below.

Due to the  $2 \times 2$  block structure of  $H^z$  in (1.3), we observe the following (almost) precise algebraic cancellations between the eigenvalues and eigenvectors. Their origin also lies in the chiral symmetry of  $H^z$ , similarly to Lemma 3.7.

**Lemma 7.3.** *For any  $a \in [1, n]$ ,  $B \in [n+1, 2n]$ , we have*

$$\left| (G^z F G^z(i\eta_c))_{aB} \mathbf{1}_{\lambda_i^z \gtrsim E_0} \right| = O_{\prec}(n^{-100}), \quad \left| (G^z F^* G^z(i\eta_c))_{Ba} \mathbf{1}_{\lambda_i^z \gtrsim E_0} \right| = O_{\prec}(n^{-100}), \quad (7.21)$$

with  $E_0 = n^{-3/4-\epsilon}$  and  $\eta_c = n^{-L}$  for a sufficiently large  $L > 100$ . In particular,

$$[G^z F G^z F^*(i\eta_c)] \mathbf{1}_{\lambda_i^z \gtrsim E_0} = O_{\prec}(n^{-100}). \quad (7.22)$$

Moreover, for any bounded deterministic vector  $\mathbf{x} \in \mathbb{C}^{2n}$ , we have

$$\langle \mathbf{x}, G^z(i\eta_c) \mathbf{x} \rangle \mathbf{1}_{\lambda_i^z \gtrsim E_0} = O_{\prec}(n^{-100}). \quad (7.23)$$

Moreover, we state the following estimates on the Green function entries.

**Lemma 7.4.** *Fix a small  $\tau > 0$ . For any  $0 \leq |z|^2 - 1 \lesssim n^{-1/2+\tau}$  and  $w = E + i\eta \in \mathbb{C}^+$  with  $|E| + \eta \lesssim n^{-3/4}$ , the following hold*

$$\langle G^z(w) F^{(*)} \rangle = \langle G^z(i\tilde{\eta}) F^{(*)} \rangle + O_{\prec} \left( \frac{n^{-1/4+\xi+\tau}}{n\eta} \right), \quad \tilde{\eta} := n^{-3/4+\xi}, \quad (7.24)$$

$$\left| \langle G^z(w) F G^z(w) F^* \rangle \right| \prec \frac{n^{-1/2+2\xi+2\tau}}{n\eta^2}, \quad (7.25)$$

for a sufficiently small  $\xi > 0$ . Moreover, for any  $0 \leq |z|^2 - 1 \lesssim n^{-1/2+\tau}$  and any bounded deterministic vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^{2n}$  and deterministic matrices  $F, F^* \in \mathbb{C}^{2n \times 2n}$  from (1.5), we have

$$\left| \langle \mathbf{x}, (G^z(i\eta_c) - M^z(i\eta_c)) \mathbf{y} \rangle \right| \mathbf{1}_{\lambda_i^z \gtrsim E_0} \prec \frac{1}{nE_0}, \quad (7.26)$$

$$\left| \langle (G^z(i\eta_c) - M^z(i\eta_c))F^{(*)} \rangle \right| \mathbf{1}_{\lambda_1^z \gtrsim E_0} \prec \frac{n^{-1/4+\xi+\tau}}{nE_0}, \quad (7.27)$$

for a sufficiently small  $\xi > 0$ , with  $E_0 = n^{-3/4-\epsilon}$  and  $\eta_c = n^{-L}$ , for a large  $L > 100$ .

These bounds are modifications of the local laws in Theorem 3.1 and Theorem 3.5. First, the local law in (3.26) was stated only for  $w = i\eta$  with  $\eta$  bigger than the typical eigenvalue spacing  $n^{-3/4}$ . Now in (7.24) we have extended the estimate to a small neighbourhood of the imaginary axis below  $n^{-3/4}$ . Second, note that while the error terms in the local laws stated in (3.8) and (3.26) blow up with a very tiny  $\eta_c$ , we have obtained a much better estimate in (7.26)-(7.27) using that  $\lambda_1^z \gtrsim E_0$ .

In Section 7.2 we will also prove the following lemma on  $q$ -related functions.

**Lemma 7.5.** *Recall  $q_z$  given in (6.4) and  $q_z^{(b)}$  ( $b \geq 1$ ) in (7.2). Fix a small  $\tau > 0$ . For any  $0 \leq |z|^2 - 1 \lesssim n^{-1/2+\tau}$ , the following estimates hold*

$$|q_z| \lesssim \mathbf{1}_{\lambda_1^z \geq E_0/2}, \quad |q_z^{(b)}| \lesssim \mathbf{1}_{E_0/2 \leq \lambda_1^z \leq 3E_0/2} \quad (b \geq 1), \quad (7.28)$$

with very high probability. Moreover we have

$$\partial_z q_z = q'_z [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}] (\text{Tr} G^z F) = O_{\prec}(n^{1/2+\epsilon+\tau}) \mathbf{1}_{E_0/2 \leq \lambda_1^z \leq 3E_0/2}, \quad (7.29)$$

with  $\widetilde{\text{Im}}$  and  $\mathfrak{D}_{E_0, \eta_0}$  defined in (7.9). The same result also holds for  $\partial_{\bar{z}} q_z$  if we replace  $F$  with  $F^*$ . Finally,

$$\Delta_z q_z = O_{\prec}(n^{1+2\epsilon+2\tau}) \mathbf{1}_{E_0/2 \leq \lambda_1^z \leq 3E_0/2}. \quad (7.30)$$

Armed with the above lemmas, we are now ready to prove Lemma 7.2.

*Proof of Lemma 7.2.* We start with the simplest case in (7.10) (i.e., a special case of (7.11) with  $p = q = 0$ ). Recalling the definition of  $\widehat{\mathcal{L}}_f$  in (7.3) and using the differentiation rules in (7.8), we have

$$\frac{\partial \widehat{\mathcal{L}}_f}{\partial h_{Ba}} = \frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) G_{aB}^z(i\eta_c) q_z d^2 z + \frac{1}{4\pi} \int_{\mathbb{C}} \Delta_z f(z) \mathcal{X}^z q'_z [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}] (G_{aB}^z) d^2 z + O_{\prec}(n^{-99}). \quad (7.31)$$

We start with the second term in (7.31). From the definition of  $\widetilde{\text{Im}}$  and  $\mathfrak{D}_{E_0, \eta_0}$ , given in (7.9), using the local law in Theorem 3.1, the properties of  $M^z$  in (3.1) and (3.6), we have for any  $\mathbf{u}, \mathbf{v} \in [1, 2n]$

$$\left| [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}] (G_{\mathbf{u}\mathbf{v}}^z) \right| \prec n^{-1/4+c\epsilon}, \quad |\mathcal{X}^z| = \left| \int_{\eta_c}^T \text{Im} \text{Tr} G^z(i\eta) d\eta \right| \prec \log n. \quad (7.32)$$

Combining these with (7.28) and the  $L^1$ -norm bound of  $\Delta f$  from (6.13), the second term in (7.31) is bounded by

$$\left| \int_{\mathbb{C}} \Delta_z f(z) \mathcal{X}^z q'_z [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}] (G_{aB}^z) d^2 z \right| \prec n^{3/4+c\epsilon} \int_{\text{supp}(f)} \mathbf{1}_{\lambda_1^z \leq 3E_0/2} d^2 z. \quad (7.33)$$

We next look at the first term in (7.31). Applying integration by parts with respect to  $\partial_z$ , using that

$$\partial_z G^z = GFG, \quad \partial_{\bar{z}} G^z = GF^*G, \quad (7.34)$$

and  $\Delta_z = 4\partial_z \partial_{\bar{z}}$ , we obtain

$$\int_{\mathbb{C}} \Delta_z f(z) G_{aB}^z(i\eta_c) q_z d^2 z = 4 \int_{\mathbb{C}} \partial_{\bar{z}} f(z) (G^z F G^z(i\eta_c))_{aB} q_z d^2 z + 4 \int_{\mathbb{C}} \partial_{\bar{z}} f(z) G_{aB}^z(i\eta_c) \partial_z q_z d^2 z. \quad (7.35)$$

Note that the first term above almost vanishes using (7.28) and (7.21). Thus using (7.29) and (7.26), together with the  $L^\infty$ -bound of  $\partial_{\bar{z}} f$  in (4.1), we have

$$\left| \int_{\mathbb{C}} \Delta_z f(z) G_{aB}^z(i\eta_c) q_z d^2 z \right| \prec (n^{3/4+c\epsilon} + n^{1+c\epsilon} \mathbf{1}_{a=B-n}) \int_{\text{supp}(f)} \mathbf{1}_{\lambda_1^z \leq 3E_0/2} d^2 z. \quad (7.36)$$

Hence we have obtained (7.10) from (7.31) using (7.33) and (7.36).

Next, we prove (7.11) for a general  $p + q + 1 \geq 2$ . Recalling  $\widehat{\mathcal{L}}_f$  in (7.3) and using Lemma 7.1, the partial derivatives of  $\widehat{\mathcal{L}}_f$  consist of a sum of two types of terms:

- 1) Terms from acting by all the partial derivatives  $\partial/\partial h_{aB}, \partial/\partial h_{Ba}$  on  $\mathcal{X}^z$  and thus leaving  $q_z$  untouched. From (7.7) and (7.31), these terms are given by

$$\int_{\mathbb{C}} \Delta_z f(z) \frac{\partial^{p+q}(G_{aB}(i\eta_c))}{\partial h_{aB}^p \partial h_{Ba}^q} q_z d^2 z = \int_{\mathbb{C}} \Delta_z f(z) (G_{aB}^z(i\eta_c))^{p+q+1} q_z d^2 z + O_{\prec}(n^{-100}), \quad (7.37)$$

where the terms with at least one  $G_{aa}^z$  or  $G_{BB}^z$  factor are bounded by  $O_{\prec}(n^{-100})$  using (7.23) and (7.28). Using integration by parts as in (7.35), we have

$$\begin{aligned} \int_{\mathbb{C}} \Delta_z f(z) (G_{aB}^z(i\eta_c))^{p+q+1} q_z d^2 z &= \int_{\mathbb{C}} \partial_{\bar{z}} f(z) (G_{aB}^z(i\eta_c))^{p+q+1} \partial_z q_z d^2 z + O_{\prec}(n^{-100}) \\ &= O_{\prec}\left(n^{\frac{3}{4} + (-\frac{1}{4} + \epsilon)(p+q) + c\epsilon} + n^{1+c\epsilon} \mathbb{1}_{a=B-n}\right) \int_{\text{supp}(f)} \mathbb{1}_{\lambda_1^z \leq 3E_0/2} d^2 z, \end{aligned} \quad (7.38)$$

where we used (7.21) in the first line, and in the last line we used (7.26) and (7.29). Note that we gained additional  $n^{(-\frac{1}{4} + \epsilon)(p+q)}$  for  $a \neq B-n$  using (7.26) compared to (7.36).

- 2) Terms from acting by at least one partial derivative (either  $\partial/\partial h_{aB}$  or  $\partial/\partial h_{Ba}$ ) on  $q_z$ . Such terms are denoted by  $\int \Delta_z f(z) (\dots) q_z^{(b)} d^2 z$  with  $b \geq 1$ , where  $(\dots)$  contains  $p+q+1$  Green function entries whose row and column index are assigned either  $a$  or  $B$  according to (7.7)-(7.8). Hence using (7.26), (7.28), (7.32), and (4.1), these terms can be bounded by

$$\left| \int_{\mathbb{C}} \Delta_z f(z) (\dots) q_z^{(b)} d^2 z \right| \prec \left( n^{\frac{3}{4} + (-\frac{1}{4} + \epsilon)(p+q) + c\epsilon} + n^{1+c\epsilon} \mathbb{1}_{a=B-n} \right) \int_{\text{supp}(f)} \mathbb{1}_{\lambda_1^z \leq 3E_0/2} d^2 z. \quad (7.39)$$

Compared to (7.33) for  $p+q+1=1$ , we have gained additional  $n^{(-\frac{1}{4} + \epsilon)(p+q)}$  for  $a \neq B-n$  since we get  $p+q$  more off-diagonal Green function entries.

Summing up the above estimates in (7.38) and (7.39), we have proved (7.11) for any  $p+q+1 \geq 2$ . The same proof applies if we switch  $a$  with  $B$ , hence we finished the proof of Lemma 7.2.  $\square$

## 7.2. Proof of the auxiliary lemmas.

*Proof of Lemma 7.1.* The differentiation rule in (7.7) follows directly from the definition of  $G^z$  in (1.3). Using (7.7) and that  $\text{Im Tr}G(i\eta) = -i\text{Tr}G(i\eta)$ , we have

$$\frac{\partial \mathcal{X}^z}{\partial h_{Ba}} = i \int_{\eta_c}^T (G^z)_{Ba}^2(i\eta) d\eta = -G_{aB}^z(i\eta_c) + O_{\prec}(n^{-100}), \quad (7.40)$$

where we also used the identity  $G^2(i\eta) = -i\partial_{\eta}G(i\eta)$ . This proves the second rule in (7.8). Moreover, using that  $\text{Im Tr}G(w) = \frac{1}{2i}(\text{Tr}G(w) - \text{Tr}G(\bar{w}))$  and  $G^2(x+i\eta) = \partial_x G(x+i\eta)$ , we have

$$\frac{\partial q_z^{(b)}}{\partial h_{Ba}} = -q_z^{(b+1)} \int_{-E_0}^{E_0} \widetilde{\text{Im}}(G^2)_{aB}(x+i\eta_0) dx = -q_z^{(b+1)} [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}](G_{aB}^z), \quad (7.41)$$

with  $\widetilde{\text{Im}}$  and  $\mathfrak{D}_{E_0, \eta_0}$  defined in (7.9). This proves the first rule in (7.8), and hence finishes the proof.  $\square$

*Proof of Lemma 7.3.* We first prove (7.21). Using the spectral decomposition of  $H^z$  and denoting by  $\mathbf{e}_a$  the unit standard vector  $\mathbf{e}_a(b) = \delta_{ab}$ , we have

$$(G^z F G^z(i\eta_c))_{aB} = \sum_{j,k=-n}^n \frac{\langle \mathbf{e}_a, \mathbf{w}_j^z \rangle \langle \mathbf{w}_j^z, F \mathbf{w}_k^z \rangle \langle \mathbf{w}_k^z, \mathbf{e}_B \rangle}{(\lambda_j^z - i\eta_c)(\lambda_k^z - i\eta_c)}. \quad (7.42)$$

Recalling that  $\lambda_{-j}^z = -\lambda_j^z$  and the corresponding eigenvectors are related by  $\mathbf{w}_{\pm j}^z = (\mathbf{u}_j^z, \pm \mathbf{v}_j^z)$ , we thus have

$$(G^z F G^z(i\eta_c))_{aB} = \sum_{j,k=-n}^n \frac{\eta_c^2 \langle \mathbf{e}_a, \mathbf{u}_j^z \rangle \langle \mathbf{u}_j^z, F \mathbf{v}_k^z \rangle \langle \mathbf{v}_k^z, \mathbf{e}_B \rangle}{((\lambda_j^z)^2 + \eta_c^2)((\lambda_k^z)^2 + \eta_c^2)}. \quad (7.43)$$

Working on the event  $\lambda_1^z \gtrsim E_0$ , we have

$$\left| (G^z F G^z(i\eta_c))_{aB} \right| \mathbb{1}_{\lambda_1^z \gtrsim E_0} \lesssim \frac{n^2 \eta_c^2}{E_0^4} \lesssim n^{-100}, \quad (7.44)$$



for  $\eta_c = n^{-L}$  with a large  $L > 100$ . The same bound also applies to  $(G^z F^* G^z(i\eta_c))_{Ba}$  and  $G(i\eta_c) F G(i\eta_c) F^*$  entrywise. Similarly, for any bounded deterministic vector  $\mathbf{x} \in \mathbb{C}^{2n}$ , we have

$$\langle \mathbf{x}, G^z(i\eta_c) \mathbf{x} \rangle \mathbb{1}_{\lambda_1^z \gtrsim E_0} \lesssim \sum_j \frac{\eta_c |\langle \mathbf{w}_j, \mathbf{x} \rangle|^2}{\lambda_j^2 + \eta_c^2} \mathbb{1}_{\lambda_1^z \gtrsim E_0} \lesssim n^{-100}, \quad (7.45)$$

for  $\eta_c = n^{-L}$  with a large  $L > 100$ . We hence finished the proof of Lemma 7.3.  $\square$

*Proof of Lemma 7.4.* We start with the proof of (7.24). By the spectral decomposition of  $H^z$ , using the symmetry that  $\lambda_{-j}^z = -\lambda_j^z$ ,  $\mathbf{w}_j^z = (\mathbf{u}_j^z, \mathbf{v}_j^z)$ ,  $\mathbf{u}_{-j}^z = \mathbf{u}_j^z$  and  $\mathbf{v}_{-j}^z = -\mathbf{v}_j^z$ , we have

$$\langle G^z(w) F \rangle = \frac{1}{n} \sum_{j=1}^n \frac{\lambda_j^z \langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle}{(\lambda_j^z - E - i\eta)(\lambda_j^z + E + i\eta)}, \quad w = E + i\eta \in \mathbb{C}^+. \quad (7.46)$$

Given the fixed constant  $\omega_c > 0$  from Corollary 3.6, set  $\tilde{\eta} := n^{-3/4+\xi}$  for a sufficiently small  $\xi > 0$  such that, if  $0 \leq \lambda_j^z \leq \tilde{\eta}$ , then  $1 \leq j \leq n^{\omega_c}$ . Thus using the overlap estimate of the singular vectors in (3.30) corresponding to small indices, hence small eigenvalues  $\lambda_j^z \leq \tilde{\eta}$ , we have

$$\begin{aligned} \langle G^z(w) F \rangle &= \frac{1}{n} \left( \sum_{0 \leq \lambda_j^z \leq \tilde{\eta}} + \sum_{\lambda_j^z \geq \tilde{\eta}} \right) \frac{\lambda_j^z \langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle}{(\lambda_j^z - E - i\eta)(\lambda_j^z + E + i\eta)} \\ &= \frac{1}{n} \sum_{\lambda_j^z \geq \tilde{\eta}} \frac{\lambda_j^z \langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle}{(\lambda_j^z - E - i\eta)(\lambda_j^z + E + i\eta)} + O_{\prec} \left( \frac{n^{-1/4+\xi+\tau}}{n\eta} \right), \end{aligned} \quad (7.47)$$

where we also used that  $\#\{0 \leq \lambda_j^z \leq \tilde{\eta}\} \prec n^\xi$  from the rigidity estimate of the eigenvalues in (3.12).

Next we estimate the error to replace  $w = E + i\eta$  in (7.47), for any  $|E| + \eta \lesssim n^{-3/4}$ , with  $w = i\tilde{\eta} = in^{-3/4+\xi}$ , above the local eigenvalue spacing where the local laws are effective. By a direct computation, we have

$$\begin{aligned} \left| \frac{1}{n} \sum_{\lambda_j^z \geq \tilde{\eta}} \frac{\lambda_j^z \langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle}{(\lambda_j^z - E - i\eta)(\lambda_j^z + E + i\eta)} - \frac{1}{n} \sum_{\lambda_j^z \geq \tilde{\eta}} \frac{\lambda_j^z \langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle}{(\lambda_j^z - i\tilde{\eta})(\lambda_j^z + i\tilde{\eta})} \right| &\lesssim \frac{1}{n} \sum_{j=1}^n \frac{\tilde{\eta}^2 \lambda_j^z |\langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle|}{((\lambda_j^z)^2 + \tilde{\eta}^2)^2} \\ &\lesssim \sqrt{\frac{1}{n} \sum_{j=1}^n \frac{\tilde{\eta}^2}{(\lambda_j^z)^2 + \tilde{\eta}^2}} \sqrt{\frac{1}{n} \sum_{j=1}^n \frac{\tilde{\eta}^2 |\langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle|^2}{((\lambda_j^z)^2 + \tilde{\eta}^2)^2}} \\ &\lesssim \sqrt{\tilde{\eta} \langle \text{Im } G^z(i\tilde{\eta}) \rangle} \sqrt{\langle G^z F G^z F^*(i\tilde{\eta}) \rangle} \prec n^{-1/2+\xi}, \end{aligned} \quad (7.48)$$

where we also used the local laws in (3.9) with (3.7) and (3.28) with (3.29). Combining (7.48) with (7.47), we obtain that

$$\langle G^z(w) F \rangle = \frac{1}{n} \sum_{\lambda_j^z \geq \tilde{\eta}} \frac{\lambda_j^z \langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle}{(\lambda_j^z - i\tilde{\eta})(\lambda_j^z + i\tilde{\eta})} + O_{\prec} \left( \frac{n^{-1/4+\xi+\tau}}{n\eta} \right) = \langle G^z(i\tilde{\eta}) F \rangle + O_{\prec} \left( \frac{n^{-1/4+\xi+\tau}}{n\eta} \right), \quad (7.49)$$

where we also used (7.47) with  $w = i\tilde{\eta}$ . Thus we have proved (7.24).

The proof of (7.25) is similar. Using the symmetry among eigenvalues and eigenvectors, for any  $w = E + i\eta \in \mathbb{C}^+$ , we have

$$\langle G^z(w) F G^z(w) F^* \rangle = \frac{2}{n} \sum_{j,k=1}^n \frac{(E + i\eta)^2 |\langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle|^2}{((\lambda_j^z)^2 - (E + i\eta)^2)((\lambda_k^z)^2 - (E + i\eta)^2)}. \quad (7.50)$$

Similarly to (7.47), we use (3.12) and (3.30) to estimate small eigenvalues  $0 \leq \lambda_j^z, \lambda_k^z \leq \tilde{\eta}$  with small indices and obtain that

$$\langle G^z(w) F G^z(w) F^* \rangle = \frac{2}{n} \sum_{\lambda_j^z, \lambda_k^z \geq \tilde{\eta}} \frac{(E + i\eta)^2 |\langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle|^2}{((\lambda_j^z)^2 - (E + i\eta)^2)((\lambda_k^z)^2 - (E + i\eta)^2)} + O_{\prec} \left( \frac{n^{-1/2+2\xi+2\tau}}{n\eta^2} \right). \quad (7.51)$$

We remark that the above also holds if we replace  $w = E + i\eta$  with  $w = i\tilde{\eta} = in^{-3/4+\xi}$ . Thus for any  $w = E + i\eta$  with  $|E| + \eta \lesssim n^{-3/4}$ , we have

$$\begin{aligned} |\langle G^z(w)FG^z(w)F^* \rangle| &\lesssim \frac{2}{n} \sum_{\lambda_j^z, \lambda_k^z \geq \tilde{\eta}} \frac{|E + i\eta|^2 |\langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle|^2}{((\lambda_j^z)^2 + \tilde{\eta}^2)((\lambda_k^z)^2 + \tilde{\eta}^2)} + O_{\prec} \left( \frac{n^{-1/2+2\xi+2\tau}}{n\eta^2} \right) \\ &\lesssim \langle G^z(i\tilde{\eta})FG^z(i\tilde{\eta})F^* \rangle + O_{\prec} \left( \frac{n^{-1/2+2\xi+2\tau}}{n\eta^2} \right) \end{aligned} \quad (7.52)$$

where we also used that  $|E| + \eta \lesssim n^{-3/4} \ll \tilde{\eta} = n^{-3/4+\xi}$ . Since  $\tilde{\eta} = n^{-3/4+\xi}$ , we are above the local eigenvalue spacing in  $\langle G^z(i\tilde{\eta})FG^z(i\tilde{\eta})F^* \rangle$ , thus using the local law in (3.28) and the deterministic upper bound in (3.29), we have proved (7.25)

Next we prove (7.26). Using the spectral decomposition of  $H^z$  and the Cauchy-Schwarz inequality, for any bounded deterministic vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^{2n}$ , we have

$$\begin{aligned} \left| \langle \mathbf{x}, (G^z(i\eta_c) - G^z(iE_0))\mathbf{y} \rangle \mathbf{1}_{\lambda_{\tilde{1}}^z \gtrsim E_0} \right| &\lesssim \left| \sum_j \frac{E_0 \langle \mathbf{x}, \mathbf{w}_j^z \rangle \langle \mathbf{w}_j^z, \mathbf{y} \rangle}{(\lambda_j^z - i\eta_c)(\lambda_j^z - iE_0)} \right| \mathbf{1}_{\lambda_{\tilde{1}}^z \gtrsim E_0} \\ &\lesssim \sqrt{\langle \mathbf{x}, \text{Im } G(iE_0)\mathbf{x} \rangle \langle \mathbf{y}, \text{Im } G(iE_0)\mathbf{y} \rangle} \prec \frac{1}{nE_0}, \end{aligned} \quad (7.53)$$

where we also used the local law in (3.8) for  $w = iE_0$  and the property of  $M^z$  in (3.7). Moreover, from (3.1) and (3.7), we have

$$\left| \langle M^z(i\eta_c) - M^z(iE_0) \rangle \right| \lesssim (E_0)^{1/3}, \quad \left| \langle (M^z(i\eta_c) - M^z(iE_0))F^{(*)} \rangle \right| \lesssim (E_0)^{2/3}. \quad (7.54)$$

This, together with (7.53), proves (7.26).

Finally, we prove (7.27) similarly. Using (7.46) and a similar argument as in (7.48), we have

$$\begin{aligned} \left| \langle (G^z(i\eta_c) - G^z(iE_0))F \rangle \right| \mathbf{1}_{\lambda_{\tilde{1}}^z \gtrsim E_0} &\lesssim \frac{1}{n} \sum_{j=1}^n \frac{(E_0)^2 \lambda_j^z |\langle \mathbf{u}_j^z, \mathbf{v}_j^z \rangle|}{((\lambda_j^z)^2 + E_0^2)^2} \\ &\lesssim \sqrt{E_0 \langle \text{Im } G^z(iE_0) \rangle} \sqrt{\langle G^z F G^z F^*(iE_0) \rangle} \prec \frac{n^{-1/4+\xi+\tau}}{nE_0}, \end{aligned} \quad (7.55)$$

where we also used the local laws in (3.9) and (7.25) for  $w = iE_0$ . Together with (7.54), we have proved (7.27) and hence finished the proof of Lemma 7.4.  $\square$

*Proof of Lemma 7.5.* We first prove the estimates in (7.28). The former one follows directly from (6.26) with  $E = E_0$ , hence we focus on proving the latter one. Recall the definition of  $q$  given in (6.5) with uniformly bounded derivatives and  $q_z^{(b)}$  defined in (7.2). Recall  $E_0 = n^{-3/4-\epsilon}$  and choose the parameters  $l_0 \gg l'_0 \gg \eta_0$  as in Lemma 6.3, i.e.,

$$l_0 = n^{-\zeta} E_0, \quad l'_0 = n^{-3\zeta} E_0, \quad \eta_0 = n^{-6\zeta} E_0, \quad (7.56)$$

for some small  $\zeta > 0$ . Note that if  $q_z^{(b)} \neq 0$  ( $b \geq 1$ ), then it implies that

$$\int_{-E_0}^{E_0} \text{Im } \text{Tr} G^z(y + i\eta_0) dy \in [1/9, 2/9]. \quad (7.57)$$

Using the first inequality in (6.27) with  $E = E_0 + l_0$  and the second inequality with  $E = E_0 - l_0$ , we have

$$\#\{|\lambda_i^z| \leq E_0 - l_0\} - O(n^{-\zeta}) \leq \int_{-E_0}^{E_0} \text{Im } \text{Tr} G^z(y + i\eta_0) dy \leq \#\{|\lambda_i^z| \leq E_0 + l_0\} + O(n^{-\zeta}). \quad (7.58)$$

Combining this with (7.57) and using that both  $\#\{|\lambda_i^z| \leq E_0 \pm l_0\}$  are integer valued, we obtain

$$\#\{|\lambda_i^z| \leq E_0 - l_0\} = 0, \quad \#\{|\lambda_i^z| \leq E_0 + l_0\} \geq 1. \quad (7.59)$$

Hence we conclude that

$$|q_z^{(b)}| = \left| q^{(b)} \left( \int_{-E_0}^{E_0} \text{Im } \text{Tr} G^z(y + i\eta_0) dy \right) \right| \lesssim \mathbf{1}_{E_0/2 \leq \lambda_{\tilde{1}}^z \leq 3E_0/2}.$$

Next, we prove (7.29). By a direct computation using (7.34), we have

$$\partial_z q_z = q'_z \int_{-E_0}^{E_0} \widetilde{\text{Im}}(\text{Tr} G^z F G^z (x + i\eta_0)) dx = q'_z [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}] (\text{Tr} G^z F), \quad (7.60)$$

with  $\widetilde{\text{Im}}$  and  $\mathfrak{D}_{E_0, \eta_0}$  defined in (7.9), and  $\partial_z q_z$  can be computed similarly. Using (7.24) for any small  $\xi > 0$  and that  $E_0, \eta_0$  are chosen as in (7.56) for any small  $\zeta > 0$ , we obtain

$$\left| [\mathfrak{D}_{E_0, \eta_0} \widetilde{\text{Im}}] (\langle G^z F^* \rangle) \right| = O_{\prec}(n^{-1/2+\epsilon+\tau}). \quad (7.61)$$

Then (7.29) follows directly from (7.60) using (7.28) and (7.61). The proof of (7.30) is similar using additionally (7.25). This concludes the proof of Lemma 7.5.  $\square$

**7.3. Extension of Theorem 4.1 to general  $\mathcal{F}$ .** We sketch how to extend Theorem 4.1 to any function  $\mathcal{F}$  with derivatives of polynomial growth as in (4.6). We follow the same scheme as in Steps 1–6 of Section 6.1. More precisely, we repeat Steps 1–5 with modifications to generalise Proposition 6.1 to any function  $\mathcal{F}$  with polynomial growth as in (4.6), *i.e.*, for any fixed  $p \in \mathbb{N}$ , there exists a large  $L > 100$ , depending on  $p$  and  $\alpha, \beta$  in (6.1), such that

$$\mathbf{E} |\mathcal{L}_f - \widehat{\mathcal{L}}_f|^p = O(n^{-c(p)\epsilon}), \quad (7.62)$$

for some constant  $c(p) > 0$ , where  $\widehat{\mathcal{L}}_f$  is defined in (6.10) with  $\eta_c = n^{-L}$ . After these, in Step 6, we extend Theorem 6.2 to a general function  $\mathcal{F}$  in (4.6) using similar GFT arguments as in Section 7.

We first note that, the estimates in Step 1 and Step 3 of Section 6 also hold true in any finite moment sense. So we only focus on the modifications needed in Step 2 and Steps 4–6.

**Modifications in Steps 2, 4, 5:** We mainly focus on Step 4, *i.e.*, to show that the size of  $\mathcal{L}_{\leq E_0}$  in (6.23) is negligible in any finite moment sense, using additionally the independence estimates on the smallest eigenvalues in (3.33). The proof of (6.19) in Step 2 and (6.29) in Step 5 in any finite moment sense is exactly the same, so we omit the details.

Recall that  $E_0 = n^{-3/4-\epsilon}$ , for some small fixed  $\epsilon > 0$ , and that  $\|\Delta f\|_{\infty} \lesssim n^{1+2\nu}$ ,  $\|\Delta f\|_1 \lesssim n^{1/2+\tau+2\nu}$ , for some arbitrary small  $\tau, \nu > 0$ , in particular  $\tau, \nu$  will be chosen much smaller than  $\epsilon$ . We now show that for any  $p \in \mathbb{N}$  there exist constants  $c(p) > 0$  such that

$$\mathbf{E} |\mathcal{L}_{\leq E_0}|^p \lesssim (\log n)^p \int \dots \int \mathbf{P}(\lambda_1^{z_1} \leq E_0, \dots, \lambda_1^{z_p} \leq E_0) \prod_{i=1}^p |\Delta f(z_i)| d^2 z_i \leq n^{-c(p)\epsilon}. \quad (7.63)$$

To keep the notation simpler, we set  $\tau = \nu = 0$ , as any term of the form  $n^{C_p(\tau+\nu)}$  (with some large  $C_p > 0$ ) accumulated along the proof can be made smaller than  $n^{-c(p)\epsilon}$  from (7.63), by choosing  $\tau = \tau(\epsilon)$ ,  $\nu = \nu(\epsilon)$  sufficiently small. We will prove (7.63) by induction on  $p$ ; in particular, we will see that the choice  $c(p) = \gamma^p$  works, with  $\gamma$  being the small constant from Proposition 3.8.

The fact that (7.63) holds for  $p = 1$ , even with  $c(1) = 1$ , immediately follows from Proposition 3.9 for  $E = E_0$  (*i.e.*, (6.18)). We now assume that (7.63) holds for any  $l \leq p - 1$ , for some  $p \geq 2$ , and we show that (7.63) holds for  $l = p$  as well. For this purpose we define the set

$$Z := \left\{ |z_i - z_j| \geq n^{-1/2+\omega}, \forall i \neq j \right\} \subset \mathbb{C}^p, \quad (7.64)$$

for some  $\omega = \omega(p, \epsilon) > 0$  which we will choose later in the proof, and split the integral in (7.63) as

$$I_Z + I_{Z^c} := \left( \int \dots \int_Z + \int \dots \int_{Z^c} \right) \mathbf{P}(\lambda_1^{z_1} \leq E_0, \dots, \lambda_1^{z_p} \leq E_0) \prod_{i=1}^p |\Delta f(z_i)| d^2 z_i. \quad (7.65)$$

On  $Z$  we can use the upper bound in Proposition 3.8, followed by (3.34), to show that the probability essentially factorizes:

$$I_Z \lesssim \int \dots \int_Z \mathbf{P}(\lambda_1^{z_i} \leq E_0 + n^{-3/4-\gamma\omega}) \prod_{i=1}^p |\Delta f(z_i)| d^2 z_i \lesssim n^{-2p\gamma\omega} + n^{-2p\epsilon}. \quad (7.66)$$

Here we used that on the support of  $f$ , by (4.2), we have

$$\mathbf{P}(\lambda_i \leq E_0 + n^{-3/4-\gamma\omega}) \lesssim n^{3/2} (E_0 + n^{-3/4-\gamma\omega})^2 e^{-\gamma n/2(1+O(C_n/\gamma n))} \lesssim n^{-1/2} (n^{-2\gamma\omega} + n^{-2\epsilon}), \quad (7.67)$$

where we used the explicit form of  $\gamma_n$  from (2.3) and that  $C_n/\sqrt{\log n} \rightarrow 0$  as  $n \rightarrow \infty$ . On  $Z^c$  we may assume, without loss of generality that  $|z_p - z_{p-1}| \leq n^{-1/2+\omega}$ , then we have

$$I_{Z^c} \lesssim n^{2\omega} \int \dots \int \mathbf{P}(\lambda_1^{z_1} \leq E_0, \dots, \lambda_1^{z_{p-1}} \leq E_0) \prod_{i=1}^{p-1} |\Delta f(z_i)| d^2 z_i \lesssim n^{2\omega - c(p-1)\epsilon}, \quad (7.68)$$

where in the first inequality we integrated out  $z_p$  by using that the volume of the region  $|z_p - z_{p-1}| \leq n^{-1/2+\omega}$  is of order  $n^{-1+2\omega}$  and that  $\|\Delta f\|_\infty \lesssim n$ . In the last inequality we used the induction hypothesis. Putting (7.66)–(7.68) together, we thus obtain

$$\int \dots \int \mathbf{P}(\lambda_1^{z_1} \leq E_0, \dots, \lambda_1^{z_p} \leq E_0) \prod_{i=1}^p |\Delta f(z_i)| d^2 z_i \lesssim n^{-2p\gamma\omega} + n^{-2p\epsilon} + n^{2\omega - c(p-1)\epsilon}. \quad (7.69)$$

We now show that we can choose  $c(p)$  and  $\omega = \omega(p, \epsilon)$  such that

$$\begin{cases} 2p\gamma\omega > c(p)\epsilon, \\ c(p-1)\epsilon - 2\omega > c(p)\epsilon, \end{cases} \quad (7.70)$$

are both satisfied; this would complete the induction step of proving (7.63). Choosing  $c(p) = \gamma^p$ , we need to choose  $\omega = \omega(p, \epsilon)$  so that

$$\frac{\gamma^{p-1}}{2p} < \frac{\omega}{\epsilon} < \frac{\gamma^{p-1} - \gamma^p}{2}, \quad (7.71)$$

which is possible since  $\frac{1}{2p}\gamma^{p-1} < \frac{1}{2}(\gamma^{p-1} - \gamma^p)$ , recalling  $p \geq 2$ . This concludes the proof of (7.63).

**Modifications in Step 6:** Even though the derivatives of  $\mathcal{F}$  have polynomial growth, the following finite moment bound

$$\mathbf{E}|\widehat{\mathcal{L}}_f|^p \lesssim n^{c\epsilon p}, \quad (7.72)$$

for some fixed constant  $c > 0$  independent of  $p$ , still allows us to perform a similar GFT argument as in Section 7. To prove (7.72), we apply integration by parts as in (7.35) but in both  $\partial_z$  and  $\partial_{\bar{z}}$ , *i.e.*,

$$\begin{aligned} \widehat{\mathcal{L}}_f &= -\frac{1}{4\pi} \int f(z) \Delta_z \left[ \left( \int_{\eta_c}^T \text{Im Tr}(G^z(i\eta) - M^z(i\eta)) d\eta \right) q_z \right] d^2 z \\ &= \frac{2i}{\pi} \int f(z) \int_{\eta_c}^T \left( \text{Tr} G^z F G^z F^* G(i\eta) \right) q_z d^2 z + O_{\prec}(n^{1+c\epsilon}) \int_{\text{supp}(f)} \mathbb{1}_{E_0/2 \leq \lambda_1^z \leq 3E_0/2}, \end{aligned} \quad (7.73)$$

for some constant  $c > 0$ , where we used (6.20), the  $z$ -derivative bounds of  $q_z$  in (7.29)–(7.30) in combination with the local laws for  $GF$  and  $GFGF^*$  in (7.24)–(7.25) and (7.27). Note that  $\text{Tr}(G^z F G^z F^* G^z(i\eta)) = \frac{1}{2i} \frac{d}{d\eta} \text{Tr}(G^z F G^z F^*(i\eta))$ . Thus the first term in (7.73) almost vanishes, *i.e.*,

$$\frac{2i}{\pi} \int f(z) \int_{\eta_c}^T \left( \text{Tr} G^z F G^z F^* G(i\eta) \right) q_z d^2 z = \frac{1}{\pi} \int f(z) \left( \text{Tr} G^z F G^z F^*(i\eta) \Big|_{\eta=\eta_c}^{\eta=T} \right) q_z d^2 z = O_{\prec}(n^{-100}), \quad (7.74)$$

with  $\eta_c = n^{-L}$  for some large  $L > 100$ , using the precise cancellation in (7.22) due to the chiral symmetry of  $H^z$ . Moreover, the second term in (7.73) can be controlled effectively by a similar argument as in (7.63), using the precise tails bound of  $\lambda_1^z$  in (3.34) together with the independence estimates in (3.33). That is, we obtain from (7.73) and (7.74) that

$$\mathbf{E}|\widehat{\mathcal{L}}_f|^p \lesssim (n^{1+c\epsilon})^p \int_{\text{supp}(f)} \dots \int_{\text{supp}(f)} \mathbf{P}(\lambda_1^{z_1} \leq 2E_0, \dots, \lambda_1^{z_p} \leq 2E_0) \prod_{i=1}^p d^2 z_i + n^{-100p} \lesssim n^{c\epsilon p}, \quad (7.75)$$

where we also used (7.63) with the choice  $c(p) = \gamma^p$  with small  $\gamma > 0$  from (3.33). This, together with (7.73), proves the finite moment bound in (7.72). Given this size bound, a similar GFT argument as in Theorem 6.2 applies to any function  $\mathcal{F}$  in (4.6) with derivatives of polynomial growth, concluding the proof of Theorem 4.1 for such test functions.

## APPENDIX A. TECHNICAL RESULTS

In this appendix we present the proof of several additional technical results used within the proofs of Section 5. Recall the definition of the two-body stability operator from (3.14). The stability operator  $\mathcal{B}_{12}$  has two non-trivial small eigenvalues<sup>13</sup> (see [58, Appendix B] for more details about the eigendecomposition of  $\mathcal{B}_{12}$ ):

$$\beta_{\pm} = \beta_{\pm}(\eta_1, z_1, \eta_2, z_2) := 1 - u_1 u_2 \operatorname{Re}[z_1 \bar{z}_2] \pm \sqrt{s}, \quad s := m_1^2 m_2^2 - u_1^2 u_2^2 (\operatorname{Im}[z_1 \bar{z}_2])^2; \quad (\text{A.1})$$

the other eigenvalues are all ones. Note that the eigenvalues in (A.1) coincide with  $\beta_{\pm,t}$  from (5.55) when evaluated along time-dependent spectral parameters, i.e.  $\beta_{\pm,t} = \beta_{\pm}(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t})$ . The main input for most of the proofs in this section is the following size relation for  $\beta_{\pm}$ :

**Lemma A.1.** *For any sufficiently small  $\epsilon_1, \epsilon_2 > 0$  and a large  $C > 0$ , and fix  $z_1, z_2 \in \mathbb{C}$ ,  $\eta_1, \eta_2 \neq 0$  such that  $1 \leq |z_i| \leq C$ ,  $|z_1 - z_2| \leq \epsilon_1$ , and  $|\eta_i| \leq \epsilon_2$ . Let  $\beta_{\pm}$  be defined as in (A.1), then we have*

$$|\beta_{\pm}| \sim \gamma = |z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}. \quad (\text{A.2})$$

*Proof of Lemma A.1.* Note that by  $-u_i = m_i^2 - |z_i|^2 u_i^2$  (which is equivalent to (3.2)) it readily follows (recall  $\rho_i := \pi^{-1} |\operatorname{Im} m_i|$ )

$$u_i = \frac{1}{|z_i|^2} - \pi^2 \rho_i^2 + O(\rho_i^4). \quad (\text{A.3})$$

Additionally, we note that

$$\operatorname{Re}[z_1 \bar{z}_2] = \frac{|z_1|^2 + |z_2|^2 - |z_1 - z_2|^2}{2}. \quad (\text{A.4})$$

Using (A.3) in the first equality and (A.4), with  $||z_1| - |z_2|| \leq |z_1 - z_2|$ , in the second one, we then compute

$$\begin{aligned} 1 - u_1 u_2 \operatorname{Re}[z_1 \bar{z}_2] &= 1 - \frac{\operatorname{Re}[z_1 \bar{z}_2]}{|z_1 z_2|^2} + \frac{\pi^2 \rho_1^2}{|z_2|^2} \operatorname{Re}[z_1 \bar{z}_2] + \frac{\pi^2 \rho_2^2}{|z_1|^2} \operatorname{Re}[z_1 \bar{z}_2] + O(\rho_1^4 + \rho_2^4) \\ &= 1 - \frac{|z_1|^2 + |z_2|^2}{2|z_1 z_2|^2} + [\pi^2 \rho_1^2 + \pi^2 \rho_2^2] (1 + O(|z_1 - z_2|)) + \frac{|z_1 - z_2|^2}{2|z_1 z_2|^2} + O(\rho_1^4 + \rho_2^4). \end{aligned} \quad (\text{A.5})$$

Then, using that  $1 \leq |z_i| \leq C$ ,  $u_i \sim 1$ , and that

$$1 - \frac{|z_1|^2 + |z_2|^2}{2|z_1 z_2|^2} = \frac{|z_1|^2(|z_2|^2 - 1) + |z_2|^2(|z_1|^2 - 1)}{2|z_1 z_2|^2} \sim |z_1|^2 + |z_2|^2 - 2, \quad (\text{A.6})$$

this implies

$$\begin{aligned} \beta_+ \beta_- &= [1 - u_1 u_2 \operatorname{Re}[z_1 \bar{z}_2]]^2 - m_1^2 m_2^2 + u_1^2 u_2^2 (\operatorname{Im}[z_1 \bar{z}_2])^2 \\ &= [1 - u_1 u_2 \operatorname{Re}[z_1 \bar{z}_2] - \pi^2 \rho_1 \rho_2] [1 - u_1 u_2 \operatorname{Re}[z_1 \bar{z}_2] + \pi^2 \rho_1 \rho_2] + u_1^2 u_2^2 (\operatorname{Im}[z_1 \bar{z}_2])^2 \\ &\gtrsim (|z_1|^2 + |z_2|^2 - 2 + \rho_1^2 + \rho_2^2)^2 + (\operatorname{Im}[z_1 \bar{z}_2])^2 \\ &\gtrsim (|z_1|^2 + |z_2|^2 - 2 + \rho_1^2 + \rho_2^2)^2 + |z_1 - z_2|^2. \end{aligned} \quad (\text{A.7})$$

In the last inequality we used that

$$(|z_1|^2 + |z_2|^2 - 2 + \rho_1^2 + \rho_2^2)^2 + (\operatorname{Im}[z_1 \bar{z}_2])^2 \gtrsim |z_1 - z_2|^2. \quad (\text{A.8})$$

whose proof is postponed to the end.

On the other hand, using again (A.5)–(A.6) and that  $|\operatorname{Im}[z_1 \bar{z}_2]| \lesssim |z_1 - z_2|$ , we readily obtain

$$|\beta_{\pm}| \lesssim |z_1|^2 + |z_2|^2 - 2 + \rho_1^2 + \rho_2^2 + |z_1 - z_2|. \quad (\text{A.9})$$

Combining (A.7)–(A.9) we finally obtain

$$|\beta_{\pm}| \gtrsim |z_1|^2 + |z_2|^2 - 2 + |z_1 - z_2| + \rho_1^2 + \rho_2^2. \quad (\text{A.10})$$

<sup>13</sup>We use the standard convention that the complex square root  $\sqrt{\cdot}$  is defined using the branch cut  $\mathbb{C} \setminus (-\infty, 0)$ .

Noticing that by (3.2) it readily follows

$$\frac{\eta_i}{\operatorname{Im} m_i} = |z_i|^2 - 1 + \pi^2 \rho_i^2 |z_i|^4 + \mathcal{O}(\rho_i^4), \quad (\text{A.11})$$

and combining (A.9)–(A.10), we conclude the proof of Lemma A.1 modulo the proof of (A.8) that we present now.

Assume that  $1 \leq |z_1| \leq |z_2|$ . By rotational symmetry we can assume that  $z_1 = r \in \mathbb{R}_+$  and write  $z_2 = r + \varepsilon e^{i\theta}$ , with  $\varepsilon \geq 0$  and  $\theta \in [0, 2\pi)$ . Then we have

$$(|z_1|^2 + |z_2|^2 - 2 + \rho_1^2 + \rho_2^2)^2 + (\operatorname{Im} [z_1 \bar{z}_2])^2 \geq (2r^2 - 2 + \varepsilon^2 + 2r\varepsilon \cos \theta)^2 + (\varepsilon r \sin \theta)^2. \quad (\text{A.12})$$

We now distinguish two cases: i)  $|\sin \theta| \geq \delta$ , ii)  $|\sin \theta| < \delta$ , for some small  $\delta > 0$  we will choose shortly. In case i) we readily conclude

$$(\varepsilon r \sin \theta)^2 \gtrsim \varepsilon^2.$$

For case ii), we notice that  $|\sin \theta| < \delta$  implies  $|\cos \theta| \geq 1 - \delta^2$ . Additionally, since we assume that  $|z_1| \leq |z_2|$  and  $|z_1 - z_2| \leq \varepsilon$ , if  $\delta$  is chosen so that  $10\varepsilon \leq \delta \leq 1/10$  then this also implies that  $\cos \theta > 0$ . We thus have

$$(2r^2 - 2 + \varepsilon^2 + 2r\varepsilon \cos \theta)^2 \geq \varepsilon^2 (\cos \theta)^2 \gtrsim \varepsilon^2.$$

Recalling that  $\varepsilon^2 = |z_1 - z_2|^2$ , this concludes the proof of (A.8), hence Lemma A.1.  $\square$

*Proof of Lemma 5.5.* By explicit computations we find that

$$((\mathcal{B}_{12}^{-1})^* [E_{\pm}])^* = \frac{1}{1 + |z_1 z_2|^2 u_1^2 u_2^2 - m_1^2 m_2^2 - 2u_1 u_2 \operatorname{Re} [z_1 \bar{z}_2]} \begin{pmatrix} 1 - \bar{z}_1 z_2 u_1 u_2 \pm m_1 m_2 & 0 \\ 0 & m_1 m_2 \pm (1 - z_1 \bar{z}_2 u_1 u_2) \end{pmatrix}. \quad (\text{A.13})$$

Using that (here  $*$  denotes some explicit term that we do not need to compute)

$$M_1 E_{\pm} M_2 = \begin{pmatrix} m_1 m_2 \pm z_1 \bar{z}_2 u_1 u_2 & * \\ * & \pm m_1 m_2 + \bar{z}_1 z_2 u_1 u_2 \end{pmatrix},$$

we thus obtain

$$-\langle M_{12}^{E_-} E_+ \rangle = \langle M_{12}^{E_+} E_- \rangle = \frac{i \operatorname{Im} [z_1 \bar{z}_2] u_1 u_2}{1 + |z_1 z_2|^2 u_1^2 u_2^2 - m_1^2 m_2^2 - 2u_1 u_2 \operatorname{Re} [z_1 \bar{z}_2]} =: i b(\eta_1, z_1, \eta_2, z_2). \quad (\text{A.14})$$

Notice that  $b(\eta_1, z_1, \eta_2, z_2)$  is real by definition. The denominator above can also be written as

$$\beta_+ \beta_- = 1 + |z_1 z_2|^2 u_1^2 u_2^2 - m_1^2 m_2^2 - 2u_1 u_2 \operatorname{Re} [z_1 \bar{z}_2] > 0, \quad (\text{A.15})$$

where the last inequality follows by (A.7). In particular, (A.15) shows that even if  $\beta_+, \beta_-$  may not be real their product is always real and positive. This proves the first relation in the second line of (5.26).

By explicit calculation and using (A.15), we also obtain

$$\pm \langle M_{12}^{E_{\pm}} E_{\pm} \rangle = -1 + \frac{\pm m_1 m_2 + 1 - \operatorname{Re} [\bar{z}_1 z_2] u_1 u_2}{1 + |z_1 z_2|^2 u_1^2 u_2^2 - m_1^2 m_2^2 - 2u_1 u_2 \operatorname{Re} [z_1 \bar{z}_2]} = -1 + \frac{\pm m_1 m_2 + (\beta_+ + \beta_-)/2}{\beta_+ \beta_-} \quad (\text{A.16})$$

hence, recalling that  $|\beta_{\pm}| \sim \gamma$  by Lemma A.1, we have

$$|\langle M_{12}^{E_{\pm}} E_{\pm} \rangle| \lesssim \frac{\rho_1 \rho_2}{\beta_+ \beta_-} + \frac{1}{\gamma} \lesssim \frac{1}{\gamma}. \quad (\text{A.17})$$

In the last inequality we used

$$\frac{\rho_1 \rho_2}{\beta_+ \beta_-} \sim \frac{\rho_1 \rho_2}{\gamma^2} \leq \frac{1}{\gamma} \cdot \frac{(\rho_1 \rho_2)^{3/2}}{|\eta_1 \eta_2|^{1/2}} \lesssim \frac{1}{\gamma},$$

where we also used the lower bound  $\gamma \geq \sqrt{|\eta_1 \eta_2| / (\rho_1 \rho_2)}$  in the second inequality, and that  $\rho_i \lesssim |\eta_i|^{1/3}$  for  $|z_i| \geq 1$  in the last inequality. This proves the bounds in the first line of (5.26). Note that (A.16) also shows (5.27) since  $-\operatorname{sgn}(m_1 m_2) = \operatorname{sgn}(\operatorname{Im} m_1 \operatorname{Im} m_2) = \operatorname{sgn}(\eta_1 \eta_2)$ .

In order to conclude the remaining bound in the second line of (5.26) we estimate

$$|b(\eta_1, z_1, \eta_2, z_2)| = \left| \frac{\operatorname{Im} [z_1 \bar{z}_2] u_1 u_2}{\beta_+ \beta_-} \right| \lesssim \frac{|z_1 - z_2|}{\gamma^2} \leq \frac{1}{\gamma},$$

using  $|\operatorname{Im} [z_1 \bar{z}_2]| \lesssim |z_1 - z_2|$  and then  $\gamma \geq |z_1 - z_2|$  in the last step.  $\square$

*Proof of (3.15).* In Lemma 5.5 we already prove bounds  $|\langle M_{12}^A E_{\pm} \rangle| \lesssim 1/\gamma$  for  $A \in \{E_+, E_-\}$ . Completely analogous calculations give the same bound for  $|\langle M_{12}^A F^{(*)} \rangle|$ . Since  $M_{12}^{E_{\pm}}$  is block-constant matrix and  $\{E_+, E_-, F, F^*\}$  is a basis in the  $2 \times 2$  matrices, this proves

$$\|M_{12}^A\| \lesssim \frac{1}{|\beta_+|} + \frac{1}{|\beta_-|} \sim \frac{1}{\gamma} \quad (\text{A.18})$$

for  $A \in \{E_+, E_-\}$ . Note that so far we proved (A.18) only when  $A \in \{E_+, E_-\}$  which is the case needed in this paper. For completeness we mention that to prove the bound for general deterministic matrices  $A \in \mathbb{C}^{2n \times 2n}$  we need to rely on the spectral decomposition of the non-Hermitian operator  $\mathcal{B}_{12}$  similarly to what it was done in [58, Eq. (5.20)] for the bulk regime; the only difference is that now we would need to project the general matrix  $A$  into the two-dimensional subspace corresponding to the eigenvalues  $\beta_{\pm}$ , whilst in the bulk regime only  $\beta_-$  matters (i.e. we had projection into a one-dimensional subspace). We do not write the details of this elementary proof as it is not needed for our analysis.  $\square$

*Proof of (5.32).* Recall that  $G_{i,t} = G_t^{z_{i,t}}(\eta_{i,t})$  and that  $t \leq \tau$ , with  $\tau$  being the stopping time defined in (5.24). The main input to prove (5.32) is the following integral representation which expresses  $|G^z|$  in terms of its imaginary parts (see e.g. [56, Eq. (5.4)]):

$$|G^z(i\eta)| = \frac{2}{\pi} \int_0^{\infty} \frac{\text{Im } G^z(i\sqrt{v^2 + \eta^2})}{\sqrt{v^2 + \eta^2}} dv = \frac{2}{\pi} \int_0^{n^{100}} \frac{\text{Im } G^z(i\sqrt{v^2 + \eta^2})}{\sqrt{v^2 + \eta^2}} dv + O(n^{-10}). \quad (\text{A.19})$$

We point out that the same integral representation holds true for the deterministic approximation of  $|G^z(i\eta)|$ . The integration regime  $v \geq n^{100}$  can be easily estimated by an error  $n^{-10}$  in operator norm; we omit this cutoff for brevity, but in the application below it is important that the  $s$ -integration is always on a finite regime  $v \leq n^{100}$ .

In order to estimate all the  $v$ -regimes in our application of (A.19) below, we need to ensure a bound for  $G_t^{z_{1,t}}(i\eta_1)G_t^{z_{2,t}}(i\eta_2)$  for all  $\eta_i \in [\eta_{i,t}, n^{100}]$  and not just  $\eta_i = \eta_{i,t}$ . The more delicate regime, when both  $\eta_i$ 's are such that  $\eta_i \leq \omega_1/2$  for some small fixed constant  $\omega_1$ , has been covered in (5.25). If at least for one  $\eta_i$  we have  $\eta_i \gtrsim 1$  then it is easy to see (similarly to [56, Appendix B]) that we get the bound  $1/(n\eta_*)$ , with  $\eta_* := \eta_1 \wedge \eta_2$ , which is even stronger than our target  $1/(\sqrt{n\ell_t}\gamma_t)$ . We thus obtain

$$\sup_{\eta_{i,t} \leq \hat{\eta}_i \leq n^{100}} (n\hat{\ell}_t)^{1/2} \hat{\gamma}_t | \langle (G_t^{z_{1,t}}(i\hat{\eta}_1)A_1 G_t^{z_{2,t}}(i\hat{\eta}_2) - M_{12}^{A_1}(\hat{\eta}_1, z_{1,t}, \hat{\eta}_2, z_{2,t}))A_2 \rangle | \leq n^{2\xi}. \quad (\text{A.20})$$

Define  $\eta_{2,t,v} := \sqrt{\eta_{2,t}^2 + v^2}$ , then we estimate

$$\begin{aligned} \langle \text{Im } G_{1,t}A_1 | G_{2,t} | A_1^* \rangle &= \frac{2}{\pi} \int_0^{n^{100}} \langle \text{Im } G_{1,t}A_1 \text{Im } G_t^{z_{2,t}}(i\eta_{2,t,v})A_1^* \rangle \frac{dv}{\eta_{2,t,v}} \\ &\lesssim \int_0^{n^{100}} \left( \frac{1}{\gamma_t} + \frac{n^{2\xi}}{\sqrt{n\ell_t}\gamma_t} \right) \frac{dv}{\eta_{2,t,v}} \\ &\lesssim \frac{\log n}{\gamma_t} + \frac{n^{2\xi} \log n}{\sqrt{n\ell_t}\gamma_t}, \end{aligned} \quad (\text{A.21})$$

where in the first inequality we used that  $(\eta_{1,t}\rho_{1,t}) \wedge (\eta_{2,t,v}/\rho^{z_{2,t}}(i\eta_{2,t,v})) \geq \ell_t$  and that

$$\frac{1}{|z_{1,t} - z_{2,t}| + (\eta_{1,t}/\rho_{1,t}) + (\eta_{2,t,v}/\rho^{z_{2,t}}(i\eta_{2,t,v}))} \lesssim \frac{1}{|z_{1,t} - z_{2,t}| + (\eta_{1,t}/\rho_{1,t}) + (\eta_{2,t}/\rho_{2,t})} \sim \frac{1}{\gamma_t}$$

as a consequence of  $\eta_{2,t,v}/\rho^{z_{2,t}}(i\eta_{2,t,v}) \geq \eta_{2,t}/\rho_{2,t}$ .  $\square$

*Proof of Lemma 5.7.* From the definition of  $a_t$  in (5.45), it follows (for both cases  $\eta_{1,t}\eta_{2,t} < 0$  and  $\eta_{1,t}\eta_{2,t} > 0$ )

$$\begin{aligned} a_t &= \frac{|m_{1,t}m_{2,t}| + m_{1,t}^2 m_{2,t}^2 + \text{Re}[z_{1,t}\overline{z_{2,t}}]u_{1,t}u_{2,t} - |z_{1,t}z_{2,t}|^2 u_{1,t}^2 u_{2,t}^2}{1 + |z_{1,t}z_{2,t}|^2 u_{1,t}^2 u_{2,t}^2 - m_{1,t}^2 m_{2,t}^2 - 2u_{1,t}u_{2,t}\text{Re}[z_{1,t}\overline{z_{2,t}}]} \\ &= \frac{|m_{1,t}m_{2,t}|}{|\beta_{+,t}\beta_{-,t}|} - \frac{1}{2} \partial_t \log |\beta_{+,t}\beta_{-,t}|, \end{aligned} \quad (\text{A.22})$$

where we used (A.15) and that by Lemma 5.1 we have

$$m_{i,t} = e^{t/2} m_{i,0}, \quad u_{i,t} = e^t u_{i,0}, \quad z_{i,t} = e^{-t/2} z_{i,0}. \quad (\text{A.23})$$

This concludes the proof of (5.56). The asymptotics in (5.57) immediately follows from  $\beta_{\pm,t} = \beta_{\pm}(\eta_{1,t}, z_{1,t}, \eta_{2,t}, z_{2,t})$  and Lemma A.1.  $\square$

*Proof of Lemma 3.7.* We present the proof of the first two equalities in (3.32), the proof of the remaining ones is analogous and so omitted. In this proof we will omit the  $z, \eta$ -dependence of the various quantities, as in Lemma 3.7 all the quantities are evaluated at  $\eta_{1,t} = \eta_{2,t} = \eta_t$  and  $z_{1,t} = z_{2,t} = z_t$ ; in particular, we use  $\mathcal{B}_{12} = \mathcal{B}$ . By the chiral symmetry of  $W - \Lambda$  we have  $GE_- = -E_-G^*$ . This follows by spectral decomposition

$$E_-GE_- = \sum_i \frac{E_- \mathbf{w}_i \mathbf{w}_i^* E_-}{\lambda_i - i\eta} = \sum_i \frac{\mathbf{w}_{-i} \mathbf{w}_{-i}^*}{\lambda_i - i\eta} = \sum_i \frac{\mathbf{w}_i \mathbf{w}_i^*}{\lambda_{-i} - i\eta} = - \sum_i \frac{E_- \mathbf{w}_i \mathbf{w}_i^* E_-}{\lambda_i + i\eta} = -G^*,$$

where we used that  $E_- \mathbf{w}_i = \mathbf{w}_{-i}$  and that  $\lambda_{-i} = -\lambda_i$ . Noticing that  $E_-^2 = E_+$  this shows  $GE_- = -E_-G^*$ . Then we compute

$$\langle G_t F G_t E_- \rangle = -\langle G_t F E_- G_t^* \rangle = \frac{1}{\eta_t} \langle \text{Im } G_t F \rangle = 0, \quad (\text{A.24})$$

where we used  $G_t E_- = -E_- G_t^*$  in the first equality and that  $\text{Im } G_t$  is a diagonal matrix, which again follows by the chiral symmetry, in the last equality.

Next, we compute

$$MFM = \begin{pmatrix} -\bar{z}um & m^2 \\ -\bar{z}^2 u^2 & -\bar{z}um \end{pmatrix} \quad (\text{A.25})$$

using (3.1). Since the diagonal blocks are the same, and the same remains true once we compute  $\mathcal{B}^{-1}[MFM]$  this shows that  $\langle M^F E_- \rangle = 0$ .  $\square$

*Proof of Lemma 5.9.* We start noticing that

$$(\mathcal{B}^{-1})^*[F] = \begin{pmatrix} \frac{m(1-|z|^2 u^2) \bar{z} u + m^3 \bar{z} u}{\beta_+ \beta_-} & 1 \\ 0 & \frac{m(1-|z|^2 u^2) \bar{z} + m^3 \bar{z} u}{\beta_+ \beta_-} \end{pmatrix}. \quad (\text{A.26})$$

Recall that  $m = i\pi\rho$ , and from (3.7) that

$$\rho \sim \frac{\eta}{|z|^2 - 1 + \eta^{2/3}}, \quad 1 - u = \frac{\eta}{\eta + \text{Im } m}.$$

Then, this implies

$$1 - |z|^2 u^2 + m^2 = 1 - u = \frac{\eta}{\text{Im } m} \left( 1 + \mathcal{O}\left(\frac{\eta}{\rho}\right) \right),$$

where we used  $-u = m^2 - |z|^2 u^2$  (this follows by (3.2)) in the first equality. Using the definition of  $\beta_{\pm}$  in (A.1), by similar computations, this implies that

$$\beta_{\pm} \geq \frac{\eta}{\text{Im } m} \left( 1 - C \frac{\eta}{\rho} \right),$$

for some  $C > 0$ . We thus conclude that taking the absolute value entry-wise we have

$$(\mathcal{B}^{-1})^*[F] \leq \begin{pmatrix} \frac{\text{Im } m^2}{\eta} + C_1 \rho & C_2 \\ 0 & \frac{\text{Im } m^2}{\eta} + C_1 \rho \end{pmatrix}, \quad (\text{A.27})$$

for some constants  $C_1, C_2 > 0$ , where we used that  $|zu| \leq 1$ . Here the notation  $\leq$  denotes that the inequality holds entry-wise. This concludes the proof of the first two bounds in (5.67) using (A.25) and that

$$M^2 = \begin{pmatrix} m^2 + |z|^2 u^2 & -2zum \\ -2\bar{z}um & m^2 + |z|^2 u^2 \end{pmatrix},$$

together with  $\rho^3/\eta \gtrsim \rho^2$ . The third (and last) bound in (5.67) is a special case of (3.15) as  $z_1 = z_2$  and  $\eta_1 = \eta_2$ , which also imply  $\text{Im } m_1 = \text{Im } m_2$ .  $\square$



## APPENDIX B. PROOF OF LEMMA 6.3

The proof of Lemma 6.3 follows the one of [67, Lemma 6.1-6.2] or [111, Lemma 2.4-2.5], and relies on the rigidity bound of eigenvalues in Corollary 3.2.

*Proof of Lemma 6.3.* For any  $E > 0$ , we define  $\chi_E := \mathbf{1}_{[-E, E]}$ , then  $\#\{j : |\lambda_j^z| \leq E\} = \text{Tr} \chi_E(H^z)$ . For  $\eta > 0$ , we define the mollifier

$$\theta_\eta(x) := \frac{\eta}{\pi(x^2 + \eta^2)} = \frac{1}{\pi} \text{Im} \frac{1}{x - i\eta}. \quad (\text{B.1})$$

We can thus relate  $\text{Tr} \chi_E \star \theta_\eta(H)$  to the Green function by the following identity:

$$\text{Tr} \chi_E \star \theta_\eta(H^z) = \frac{1}{\pi} \int \chi_E(y) \text{Im} \text{Tr} G^z(y + i\eta) dy = \frac{1}{\pi} \int_{-E}^E \text{Im} \text{Tr} G^z(y + i\eta) dy. \quad (\text{B.2})$$

Choose the spectral parameters  $E \gg l \gg l' \gg \eta$  as in (6.25). Using [111, Eq.(A.2)-(A.4)], we have

$$\left| \text{Tr}(\chi_E - \chi_E \star \theta_\eta)(H^z) \right| \lesssim \#\{|\lambda_j^z| \in [E - l', E + l']\} + \frac{\eta}{l'} \#\{|\lambda_j^z| \leq E - l'\} + \text{Tr} g(H^z), \quad (\text{B.3})$$

with  $g(x) := \text{Tr} \chi_E \star \theta_\eta(x) \mathbf{1}_{|x| \geq E + l'}$ . Using the rigidity estimate of eigenvalues in (3.12) and the properties of  $\rho^z$  in (3.6), we have

$$\left| \text{Tr}(\chi_E - \chi_E \star \theta_\eta)(H^z) \right| \lesssim \#\{|\lambda_j^z| \in [E - l', E + l']\} + \frac{n^\xi \eta}{l'} + \text{Tr} g(H^z), \quad (\text{B.4})$$

for any small  $\xi > 0$  with very high probability. It is then sufficient to estimate  $\text{Tr} g(H^z)$ . Similarly to [111, Eq.(A.6)-(A.7)], if  $x \geq E + l'$  with  $E$  and  $l'$  chosen in (6.25), then  $x \pm E \geq l' \gg \eta$ , thus we have

$$g(x) = \int_{-E-x}^{E-x} \theta_\eta(y) dy \lesssim \min \left\{ \frac{\eta E}{(x-E)^2}, \frac{\eta}{x-E} \right\}. \quad (\text{B.5})$$

Choose a sufficiently small  $c > 0$  and we write

$$\text{Tr} g(H^z) = 2 \sum_{|\lambda_i^z| \geq E + l'} g(\lambda_i^z) = 2 \left( \sum_{k=0}^{K_0} \sum_{\lambda_i^z \in \mathcal{I}_k} + \sum_{\lambda_i^z \geq c} \right) g(\lambda_i^z), \quad \mathcal{I}_k := [E + 3^k l', E + 3^{k+1} l'], \quad (\text{B.6})$$

where  $[E + l', c] = \bigcup_{k=0}^{K_0} \mathcal{I}_k$  with  $K_0 = O(\log n)$ , using that  $g$  is an even function and the spectrum  $\{\lambda_{\pm i}^z\}_{i \in [n]}$  is symmetric. Combining with the rigidity estimate of eigenvalues in (3.12) and (B.5), we have

$$\text{Tr} g(H^z) \lesssim \sum_{k=0}^{K_0} \min \left\{ \frac{\eta E}{(3^k l')^2}, \frac{\eta}{3^k l'} \right\} \left( \#\{i : \lambda_i^z \in \mathcal{I}_k\} \right) + n\eta E \lesssim \frac{n^\xi \eta}{l'}, \quad (\text{B.7})$$

for any small  $\xi > 0$  with very high probability. Combining this with (B.2) and (B.4), we have proved

$$\left| \text{Tr}(\chi_E - \chi_E \star \theta_\eta)(H^z) \right| \lesssim \#\{|\lambda_j^z| \in [E - l', E + l']\} + \frac{n^\xi \eta}{l'}, \quad (\text{B.8})$$

for any small  $\xi > 0$  with very high probability and hence (6.26). Moreover, using that  $E \mapsto \chi_E$  is increasing, together with (B.8), we obtain that

$$\begin{aligned} \text{Tr} \chi_E(H^z) &\leq \frac{1}{l} \int_E^{E+l} \text{Tr} \chi_{E'}(H^z) dE' \\ &\leq \frac{1}{l} \int_E^{E+l} \text{Tr} \chi_E \star \theta_\eta(H^z) dE' + \frac{C}{l} \int_E^{E+l} \left( \#\{|\lambda_i^z| \in [E' - l', E' + l']\} + \frac{n^\xi \eta}{l'} \right) dE' \\ &\leq \text{Tr} \chi_{E+l} \star \theta_\eta(H^z) + Cn^\xi \left( \frac{\eta}{l'} + \frac{l'}{l} \right), \end{aligned} \quad (\text{B.9})$$

for any small  $\xi > 0$  with very high probability. One can obtain a lower bound similarly. With the choices of parameters in (6.25) and (B.2), we hence finished the proof of (6.27).  $\square$

## APPENDIX C. PROOF OF PROPOSITION 3.8

The proof of this proposition consists of two parts: 1) we first prove Proposition 3.8 for the matrix with a small ( $\sim n^{-1/2+\omega_1}$ ) Gaussian component, 2) we remove the Gaussian component using a GFT argument. In particular, Part 1) already proves Proposition 3.8 for Ginibre matrices.

We will only prove Proposition 3.8 for any  $\{E_j\}_{j \in [k]}$  with  $n^{-3/4-\gamma\omega+\tau} < E_j \lesssim n^{-3/4}$  and  $\tau \leq (\gamma\omega)/10$ , while the same upper bound applies to the complementary regime  $0 < E_j \leq n^{-3/4-\gamma\omega+\tau}$  using a simple monotonicity argument (the lower bound is a completely trivial in this case).

*Proof of Part 1) of Proposition 3.8.* This proof is similar to the proof of [61, Proposition 4.3], so we will be brief. For simplicity of notation we only present the proof for  $k = 2$ . Fix a time  $t_1 := n^{-1/2+\omega_1}$ , for some small  $\omega_1 > 0$ . Consider the matrix valued flow<sup>14</sup>

$$dX_t = \frac{dB_t}{\sqrt{n}}, \quad X_0 = e^{-t_1/2} X, \quad t \geq 0, \quad (\text{C.1})$$

where  $X$  is an i.i.d. complex matrix satisfying Assumption 2.1, and  $B_t$  is an  $n \times n$  matrix whose entries are i.i.d. standard complex Brownian motions. Hence,  $X_{\widehat{c}t_1}$ , with  $\widehat{c} = \widehat{c}(t_1) := \sqrt{1 - e^{-t_1}}/t_1$ , is an i.i.d. matrix which satisfies Assumption 2.1, and also has a Gaussian component of size  $\sim \sqrt{t_1}$ . By simple second order perturbation theory (see e.g. [52, Appendix B]) the eigenvalues  $\lambda_i^z(t)$  of the Hermitization of  $X_t - z$  are the unique strong solution of the following Dyson Brownian motion (DBM)

$$d\lambda_i^z(t) = \frac{db_i^z(t)}{\sqrt{2n}} + \frac{1}{2n} \sum_{j \neq i} \frac{1}{\lambda_i^z(t) - \lambda_j^z(t)} dt. \quad (\text{C.2})$$

Here  $b_i^z(t)$  are i.i.d. standard real Brownian motions for  $i \in [n]$ , and  $b_{-i}^z = -b_i^z$  for  $i \in [n]$ . Note that the collections  $\{b_i^z(t) : i \in [n]\}$  are not independent for different  $z$ 's.

To prove the (asymptotic) independence of  $\lambda_1^{z_j}(t)$  for different  $z_j$ 's, we couple the evolution (C.2) with the evolution of fully independent points  $\{\mu_i\}$  and show that after short time  $\sim n^{-1/2+\omega_1}$  these two processes are very close. More precisely, for  $l = 1, 2$ , define  $\{\mu_i^{(l)}(t)\}$  as the solution to

$$d\mu_i^{(l)}(t) = \frac{d\beta_i^{(l)}(t)}{\sqrt{2n}} + \frac{1}{2n} \sum_{j \neq i} \frac{1}{\mu_i^{(l)}(t) - \mu_j^{(l)}(t)} dt, \quad (\text{C.3})$$

with the singular values of two independent complex Ginibre matrices as initial conditions. Here the family  $\{\beta_i^{(l)} : i \in [n], l = 1, 2\}$  is a  $2n$ -dimensional standard Brownian motion and  $\beta_{-i}^{(l)} = -\beta_i^{(l)}$ . For any  $z_l \in \mathbb{C}$  such that  $0 \leq |z_l| - 1 \leq n^{-1/2+\tau}$  and  $|z_1 - z_2| \gtrsim n^{-1/2+\omega}$  for any sufficiently small  $\tau, \omega$  with  $\tau < \gamma\omega/10$ , Corollary 3.4 implies, by a standard grid argument, that

$$|\langle \mathbf{u}_i^{z_1}(t), \mathbf{u}_j^{z_2}(t) \rangle|^2 + |\langle \mathbf{v}_i^{z_1}(t), \mathbf{v}_j^{z_2}(t) \rangle|^2 \prec n^{-\omega+2\tau} =: n^{-\omega_E}, \quad |i|, |j| \leq n^{\omega_c}, \quad (\text{C.4})$$

holds simultaneously in  $0 \leq t \leq \widehat{c}t_1$ , for any small constants  $\omega_c, \omega_E$  such that  $10\omega_1 \leq 10\omega_c \leq \omega_E \leq \omega/2$ . Thus the assumptions of [61, Theorem 7.2] are verified, there hence exists a specific coupling between the pair of correlated Brownian motions  $\{b_i^{z_1}, b_i^{z_2} : i \in [n]\}$  and the independent ones  $\{\beta_i^{(1)}, \beta_i^{(2)} : i \in [n]\}$  such that, for  $l = 1, 2$ , we have

$$|\lambda_i^{z_l}(\widehat{c}t_1) - \mu_i^{(l)}(\widehat{c}t_1)| \prec n^{-3/4-\gamma\omega+\tau}, \quad |i| \leq n^{\widehat{\omega}}, \quad (\text{C.5})$$

for any sufficiently small constants  $\gamma, \omega, \widehat{\omega}, \tau$  satisfying  $\widehat{\omega} \leq (\gamma\omega)/10 \leq \omega_1/100$  and  $\tau \leq (\gamma\omega)/10$ . The bound (C.5) holds with very high probability in the joint probability space of the  $\lambda_i^{z_l}(\widehat{c}t_1)$ 's and  $\mu_i^{(l)}(\widehat{c}t_1)$ 's. We thus obtain

$$\begin{aligned} \mathbf{P}\left(\lambda_1^{z_l}(\widehat{c}t_1) \leq E_l, l \in [2]\right) &\leq \mathbf{P}\left(\mu_1^{(l)}(\widehat{c}t_1) \leq E_l + n^{-3/4-\gamma\omega+\tau}, l \in [2]\right) + O(n^{-D}) \\ &= \prod_{l=1}^2 \mathbf{P}\left(\mu_1^{(l)}(\widehat{c}t_1) \leq E_l + n^{-3/4-\gamma\omega+\tau}\right) + O(n^{-D}) \\ &\leq \prod_{l=1}^2 \mathbf{P}\left(\lambda_1^{z_l}(\widehat{c}t_1) \leq E_l + 2n^{-3/4-\gamma\omega+\tau}\right) + O(n^{-D}) \end{aligned} \quad (\text{C.6})$$

<sup>14</sup>We warn the reader that within Appendix C, to keep the same notation of [61, Section 7], by  $X_t$  we denote the solution of the flow (C.1), and not the solution of the Ornstein–Uhlenbeck flow from (7.5).

for any large  $D > 0$ . In the second line of (C.6), we used the fact that  $\mu_1^{(1)}(t)$  and  $\mu_1^{(2)}(t)$  are independent processes by construction. Similarly we obtain the desired lower bound as well. This proves Proposition 3.8 with  $k = 2$  for  $X_{\widehat{c}t_1}$ , i.e. for a matrix with a small ( $\sim n^{-1/2+\omega_1}$ ) Gaussian component, and the case of general  $k$  is analogous.  $\square$

The proof of Part 2) follows the strategy used in [61, Proposition 5.2]. In particular, we will remove the small  $\sim n^{-1/2+\omega_1}$  Gaussian component of  $X_{\widehat{c}t_1}$  from (C.6) that has been obtained in Part 1). Again we give the proof only for any  $\{E_j\}_{j \in [k]}$  with  $n^{-3/4-\gamma\omega+\tau} < E_j \lesssim n^{-3/4}$  and  $\tau \leq (\gamma\omega)/10$ .

*Proof of Part 2) of Proposition 3.8.* Consider  $\widehat{X}_t$  to be the solution of the Ornstein–Uhlenbeck flow

$$d\widehat{X}_t = -\frac{1}{2}\widehat{X}_t dt + \frac{d\widehat{B}_t}{\sqrt{n}}, \quad \widehat{X}_0 = X, \quad (\text{C.7})$$

where  $X$  is an i.i.d. complex matrix satisfying Assumption 2.1, and the entries of  $\widehat{B}_t$  are standard complex i.i.d. Brownian motions. The corresponding probability and expectation for the matrix flow  $\widehat{X}_t$  ( $t \geq 0$ ) are then denoted by  $\widehat{\mathbf{P}}_t$  and  $\widehat{\mathbf{E}}_t$ , respectively. We will now show that

$$\widehat{\mathbf{P}}_0\left(\lambda_1^{z_j} \leq E_j, j \in [k]\right) \lesssim \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1}\left(\lambda_1^{z_j} \leq E_j + n^{-3/4-\gamma\omega+\tau}\right) + n^{-D}$$

and the similar lower bound for  $t_1 = n^{-1/2+\omega_1}$ . This will conclude the proof of Proposition 3.8 noticing the following relation

$$\widehat{X}_{t_1} \stackrel{d}{=} e^{-t_1/2}X + \sqrt{1-e^{-t_1}}X^{\text{Gin}} \stackrel{d}{=} X_{\widehat{c}t_1}, \quad (\text{C.8})$$

where  $X^{\text{Gin}}$  is a Ginibre matrix independent of  $X$  and  $X_{\widehat{c}t_1}$  is the solution of (C.1) from Part 1) and for such matrix we already proved the asymptotic factorisation in (C.6).

Let  $F : \mathbb{R}_+^k \rightarrow [0, 1]$  be a smooth cut-off function given by  $F(w_1, \dots, w_k) := \Phi(w_1) \cdots \Phi(w_k)$  with the non-decreasing function  $\Phi$  defined in (4.11). Then for any fixed  $E_j \in \mathbb{R}^+$  and any  $t \geq 0$ ,

$$\widehat{\mathbf{P}}_t\left(\lambda_1^{z_j} \leq E_j, j \in [k]\right) = \widehat{\mathbf{E}}_t\left[F\left(\#\{|\lambda_i^{z_1}| \leq E_1\}, \dots, \#\{|\lambda_i^{z_k}| \leq E_k\}\right)\right].$$

For notational brevity in the proof below, we define the following short hand

$$\mathcal{I}_E^{z, \eta} := \int_{|y| \leq E} \text{Im Tr} \widehat{G}_t^z(y + i\eta) dy, \quad \forall E, \eta \in \mathbb{R}^+, z \in \mathbb{C}, \quad (\text{C.9})$$

where  $\widehat{G}_t^z$  is the corresponding resolvent of the Hermitised matrix defined as in (7.1) with  $\widehat{X}_t$  given by (C.7). Clearly  $\mathcal{I}_E^{z, \eta}$  is an increasing function in  $E \in \mathbb{R}^+$ . In this section we set  $\epsilon_* := \gamma\omega - \tau > 0$  for notational simplicity, hence  $n^{-3/4-\epsilon_*} \leq E_j \lesssim n^{-3/4}$ . Then using the inequalities in (6.27) in Lemma 6.3 for  $E = E_j, l = l_j = n^{-\epsilon_*}E_j, \eta = \eta_j = n^{-3\epsilon_*}E_j$  chosen as in (6.25) with  $\zeta = \epsilon_*$ , we have

$$\widehat{\mathbf{E}}_t\left[F\left(\mathcal{I}_{E_1-l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k-l_k}^{z_k, \eta_k}\right)\right] - n^{-D} \leq \widehat{\mathbf{P}}_t\left(\lambda_1^{z_j} \leq E_j, j \in [k]\right) \leq \widehat{\mathbf{E}}_t\left[F\left(\mathcal{I}_{E_1+l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+l_k}^{z_k, \eta_k}\right)\right] + n^{-D}, \quad (\text{C.10})$$

for any  $t \geq 0$  and any large constant  $D > 0$ . We remark that the inequalities in (C.10) hold true if we replace  $E_j$  with  $E_j + ml_j$  (for any fixed  $m \in \mathbb{Z}$ ). It then suffices to show that, there exists some constant  $c_0 > 0$  such that

$$\left|(\widehat{\mathbf{E}}_0 - \widehat{\mathbf{E}}_{t_1})\left[F\left(\mathcal{I}_{E_1 \pm l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k \pm l_k}^{z_k, \eta_k}\right)\right]\right| \lesssim n^{-c_0} \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1}\left(\lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*}\right) + n^{-D}. \quad (\text{C.11})$$

Once we prove (C.11), using with the inequalities in (C.10) for  $t = 0$  and  $t = t_1$ , we obtain that

$$\begin{aligned} \widehat{\mathbf{P}}_0\left(\lambda_1^{z_j} \leq E_j, j \in [k]\right) &\leq \widehat{\mathbf{E}}_0\left[F\left(\mathcal{I}_{E_1+l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+l_k}^{z_k, \eta_k}\right)\right] + n^{-D} \\ &\lesssim \widehat{\mathbf{E}}_{t_1}\left[F\left(\mathcal{I}_{E_1+l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+l_k}^{z_k, \eta_k}\right)\right] + n^{-c_0} \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1}\left(\lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*}\right) + n^{-D} \end{aligned}$$

$$\begin{aligned}
&\lesssim \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + 2l_j, j \in [k] \right) + n^{-c_0} \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*} \right) + n^{-D} \\
&\lesssim \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + 2n^{-3/4-\epsilon_*} \right) + n^{-D}, \tag{C.12}
\end{aligned}$$

where in the last line we applied (C.6) from Part 1) for  $t = \widehat{c}t_1$ , noticing the relation in (C.8). One obtains a similar lower bound with  $n^{-3/4-\epsilon_*}$  replaced with  $-n^{-3/4-\epsilon_*}$ . In particular for  $k = 1$

$$\widehat{\mathbf{P}}_{t_1} \left( \lambda_1^z \leq E - 2n^{-3/4-\epsilon_*} \right) - n^{-D} \lesssim \widehat{\mathbf{P}}_0 \left( \lambda_1^z \leq E \right) \lesssim \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^z \leq E + 2n^{-3/4-\epsilon_*} \right) + n^{-D}. \tag{C.13}$$

Combining (C.12) for a general  $k \geq 2$  and (C.13) for  $E = E_j + 2n^{-3/4+2\epsilon_*}$ , we prove Proposition 3.8 for the initial matrix  $\widehat{X}_0 = X$ , hence finish the proof of Proposition 3.8 for any i.i.d. matrix.

In the following, we will focus on proving (C.11). More precisely, we claim that, for any fixed  $m \in \mathbb{Z}$ ,

$$\frac{d}{dt} \widehat{\mathbf{E}}_t \left[ F \left( \mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k} \right) \right] \lesssim n^{-1/4+c\epsilon_*} \widehat{\mathbf{E}}_t \left[ F \left( \mathcal{I}_{E_1+(m+2)l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+(m+2)l_k}^{z_k, \eta_k} \right) \right] + n^{-D}, \tag{C.14}$$

uniformly in  $0 \leq t \leq t_1$ . Assuming we have proved (C.14), integrating (C.14) up to time  $t = t_1$ , we have

$$\left| \left( \widehat{\mathbf{E}}_t - \widehat{\mathbf{E}}_{t_1} \right) \left[ F \left( \mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k} \right) \right] \right| \lesssim n^{-1/4+c\epsilon_*} \int_t^{t_1} \left| \widehat{\mathbf{E}}_t \left[ F \left( \mathcal{I}_{E_1+(m+2)l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+(m+2)l_k}^{z_k, \eta_k} \right) \right] \right| + n^{-D}, \tag{C.15}$$

uniformly for any  $0 \leq t \leq t_1$ . Using (C.10), we know that, for any fixed  $m \in \mathbb{Z}$ ,

$$\begin{aligned}
\widehat{\mathbf{E}}_{t_1} \left[ F \left( \mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k} \right) \right] &\lesssim \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + (m+1)l_j, j \in [k] \right) + n^{-D} \\
&\lesssim \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*} \right) + n^{-D}, \tag{C.16}
\end{aligned}$$

where we also used (C.6) from Part 1) and the relation in (C.8). From (C.15) with (C.16) and using that  $\mathcal{F}$  is uniformly bounded, we have an initial estimate

$$\widehat{\mathbf{E}}_t \left[ F \left( \mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k} \right) \right] \lesssim n^{-1/4+c\epsilon_*} + \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*} \right) + n^{-D}, \tag{C.17}$$

for any fixed  $m \in \mathbb{Z}$  and  $0 \leq t \leq t_1$  uniformly. Plugging (C.17) (replacing  $m$  with  $m+2$ ) into the right side of (C.15), we then obtain

$$\left| \left( \widehat{\mathbf{E}}_t - \widehat{\mathbf{E}}_{t_1} \right) \left[ F \left( \mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k} \right) \right] \right| \lesssim \left( n^{-1/4+c\epsilon_*} \right)^2 + n^{-1/4+c\epsilon_*} \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*} \right) + n^{-D}, \tag{C.18}$$

uniformly in  $0 \leq t \leq t_1$ . Combining again with (C.16), we obtain that, for any fixed  $m \in \mathbb{Z}$  and  $0 \leq t \leq t_1$ ,

$$\left| \widehat{\mathbf{E}}_t \left[ F \left( \mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k} \right) \right] \right| \lesssim \left( n^{-1/4+c\epsilon_*} \right)^2 + \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*} \right) + n^{-D}. \tag{C.19}$$

In this way we have improved the first error term in (C.17) by an additional factor  $n^{-1/4+c\epsilon_*}$ . We further plug the above estimate into (C.14) and iterate the above process for sufficiently many times. In general for any  $s$ -th iteration step, the following holds true for any  $m \in \mathbb{Z}$  and  $0 \leq t \leq t_1$ ,

$$\left| \left( \widehat{\mathbf{E}}_t - \widehat{\mathbf{E}}_{t_1} \right) \left[ F \left( \mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k} \right) \right] \right| \lesssim \left( n^{-1/4+c\epsilon_*} \right)^s + n^{-1/4+c\epsilon_*} \prod_{j=1}^k \widehat{\mathbf{P}}_{t_1} \left( \lambda_1^{z_j} \leq E_j + n^{-3/4-\epsilon_*} \right) + n^{-D}. \tag{C.20}$$

We hence stop the iteration at  $s = O(D)$  to improve the first error in (C.20) to  $n^{-D}$  for any large  $D > 0$ . This finishes the proof of (C.11) with  $t = 0$ .

It still remains to prove (C.14). Recall that the Hermitized matrix flow  $\widehat{H}_t^z = (\widehat{h}_{ab})_{a,b \in [2n]}$  is defined as in (7.1) with  $\widehat{X}_t$  given by (C.7). Applying the Itô's formula on the corresponding  $F(\mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k})$  and using the cumulant expansion formula as in (7.6), we have

$$\begin{aligned} & \frac{d}{dt} \widehat{\mathbf{E}}_t \left[ F(\mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k}) \right] \\ &= \sum_{p+q+1 \geq 3} \frac{M_D}{n^{\frac{p+q+1}{2}}} \sum_{a=1}^n \sum_{B=n+1}^{2n} \left( \frac{\kappa^{(p+1, q)}}{p!q!} \widehat{\mathbf{E}}_t \left[ \frac{\partial^{p+q+1} F(\dots)}{\partial \widehat{h}_{aB}^{p+1} \partial \widehat{h}_{Ba}^q} \right] + \frac{\kappa^{(q, p+1)}}{p!q!} \widehat{\mathbf{E}}_t \left[ \frac{\partial^{p+q+1} F(\dots)}{\partial \widehat{h}_{aB}^q \partial \widehat{h}_{Ba}^{p+1}} \right] \right) + O_{\prec}(n^{-D}), \end{aligned} \quad (\text{C.21})$$

where we truncated the cumulant expansion at a sufficiently large order  $M_D$  such that the error is bounded by  $O_{\prec}(n^{-D})$ , using the moment condition in (2.1) and that  $\|G\| \leq |\eta|^{-1}$ . Using the differentiation rule as in (7.7), for any  $1 \leq j \leq k$ , we have

$$\frac{\partial \mathcal{I}_{E_j+ml_j}^{z_j, \eta_j}}{\partial \widehat{h}_{aB}} = \widetilde{\text{Im}} \widehat{G}_{Ba}^{z_p} (y + i\eta_j) \Big|_{y=-(E_j+ml_j)}^{y=E_j+ml_j} = O_{\prec}(n^{-1/4+c\epsilon_*}), \quad (\text{C.22})$$

with  $\widetilde{\text{Im}}$  defined in (7.9), and the last estimate follows from the local law in (3.8) and the choices of  $l_j, \eta_j$  above (C.10). In general, for any integers  $p, q \geq 1$ , we have

$$\left| \frac{\partial^{p+q} \mathcal{I}_{E_j+ml_j}^{z_j, \eta_j}}{\partial \widehat{h}_{aB}^p \partial \widehat{h}_{Ba}^q} \right| = O_{\prec} \left( (n^{-1/4+c\epsilon_*})^{p+q} \right). \quad (\text{C.23})$$

Therefore we obtain from (C.21) and that  $F(w_1, \dots, w_k) = \prod_{i=1}^k \Phi(w_i)$

$$\left| \frac{d}{dt} \widehat{\mathbf{E}}_t \left[ F(\mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k}) \right] \right| \prec n^{-1/4+c\epsilon_*} \sum_{|\alpha|=1}^{M_D} \sum_{\alpha_1, \dots, \alpha_k} \widehat{\mathbf{E}}_t \left| \prod_{j=1}^k \Phi^{(\alpha_j)}(\mathcal{I}_{E_j+ml_j}^{z_j, \eta_j}) \right|, \quad (\text{C.24})$$

where  $\alpha = (\alpha_1, \dots, \alpha_k)$  is a multi-index. Since  $\Phi$  given in (4.11) has uniformly bounded derivatives, then we have

$$\left| \Phi^{(\alpha_j)}(\mathcal{I}_{E_j+ml_j}^{z_j, \eta_j}) \right| \lesssim \mathbf{1}_{\mathcal{I}_{E_j+ml_j}^{z_j, \eta_j} \in [1/9, 2/9]} \lesssim \mathbf{1}_{\lambda_1^{z_j} \leq E_j + (m+1)l_j}, \quad \forall \alpha_j \geq 1$$

and a similar upper bound also applies to  $\alpha_j = 0$ . Thus we have

$$\begin{aligned} \left| \frac{d}{dt} \widehat{\mathbf{E}}_t \left[ F(\mathcal{I}_{E_1+ml_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+ml_k}^{z_k, \eta_k}) \right] \right| &\prec n^{-1/4+c\epsilon_*} \widehat{\mathbf{P}}_t \left( \lambda_1^{z_j} \leq E_j + (m+1)l_j, j \in [k] \right) \\ &\lesssim n^{-1/4+c\epsilon_*} \widehat{\mathbf{E}}_t \left[ F(\mathcal{I}_{E_1+(m+2)l_1}^{z_1, \eta_1}, \dots, \mathcal{I}_{E_k+(m+2)l_k}^{z_k, \eta_k}) \right] + n^{-D}, \end{aligned} \quad (\text{C.25})$$

where in the last step we used the second inequality (C.10) with  $E_j$  replaced by  $E_j + (m+1)l_j$ . Therefore we have proved (C.14) and hence finished the proof of Proposition 3.8.  $\square$

#### APPENDIX D. GFT: REMOVE GAUSSIAN COMPONENT IN PROPOSITION 5.4 AND 5.8

In this section, we will remove the Gaussian component in Part A of Theorem 3.3 using a standard GFT argument and hence prove Part B of Theorem 3.3 for any i.i.d. matrix. The same proof also applies to Theorem 3.5 using a similar GFT argument combining with Proposition 5.8.

We organize this section as follows. In the first two sections, we split the GFT argument used to prove Theorem 3.3 (Part B) for any i.i.d. matrix into two steps.

- Step 1:** In Section D.1, we first establish a standard GFT for  $\langle G^{z_1} A_1 G^{z_2} A_2 \rangle$  (see Proposition D.1 below) to remove any Gaussian component of size of order one with an error term bounded by  $(n|\eta_1 \eta_2|)^{-1}$ . This will be used to prove the local law with canonical error bound  $(n|\eta_1 \eta_2|)^{-1}$ .
- Step 2:** In Section D.2, we derive an improved GFT for quantities of the special form  $\langle \text{Im} G^{z_1} A \text{Im} G^{z_2} A^* \rangle$  for a large  $|z_1 - z_2| \gtrsim \eta_i / \rho_i$  (see Proposition D.3 below). This will be used to prove the local law in (3.17) with the  $|z_1 - z_2|$  decorrelation effect.

Moreover, in Section D.3 we show a refined GFT with  $\rho$ -improvement for  $\langle GFGF^* \rangle$  in order to prove the local laws in Theorem 3.5. Note that  $\langle GFGF^* \rangle = \langle \text{Im} GF \text{Im} GF^* \rangle$  from Lemma 3.7 due to the chiral symmetry of  $H^z$ , hence the GFT argument is completely analogous to Section D.2; for this reason we only sketch its proof. Lastly, in Section D.4, we combine Proposition 5.4 and Proposition 5.8 with the above GFT arguments, and present the proof of Theorem 3.3 (Part B) and Theorem 3.5, respectively, for any i.i.d. matrix.

**D.1. GFT for  $\langle G_1 A_1 G_2 A_2 \rangle$  with error  $(n|\eta_1 \eta_2|)^{-1}$ .** Recall the Ornstein-Uhlenbeck matrix flow  $H_t^z$  in (7.1) and its time-dependent resolvent  $G_t^z$  defined as in (1.3). For notational simplicity, we define

$$R_t := \langle (G_1 A_1 G_2 - M_{12}^{A_1}) A_2 \rangle, \quad G_i := G_t^{z_i}(i\eta_i), \quad i = 1, 2, \quad (\text{D.1})$$

with the deterministic matrix  $M_{12}^{A_1} := M_{12}^{A_1}(\eta_1, z_1, \eta_2, z_2)$  given by (3.13), where both  $z_i \in \mathbb{C}$  and  $\eta_i \in \mathbb{R}$  ( $i = 1, 2$ ) are time independent parameters. Then we establish the following GFT to remove the Gaussian component in Proposition 5.4 with the error bound  $(n|\eta_1 \eta_2|)^{-1}$ .

**Proposition D.1.** *Fix any  $m \geq 2$ . Then for any spectral parameters  $|\eta_1| \sim |\eta_2| \sim \eta$ ,  $\rho_1 \sim \rho_2 \sim \rho$ , the following holds*

$$\left| \frac{d\mathbf{E}|R_t|^{2m}}{dt} \right| \prec (1 + \eta^{1/3}) \left( \mathbf{E}|R_t|^{2m} + \left( \frac{1}{n|\eta_1 \eta_2|} \right)^{2m} \right). \quad (\text{D.2})$$

uniformly for any  $t \geq 0$ ,  $\|A_i\| \lesssim 1$  and  $n|\eta_i|\rho_i \gtrsim 1$ .

*Proof of Proposition D.1.* Applying Itô's formula to  $|R_t|^{2m} = (R_t)^m (R_t^*)^m$  and performing the cumulant expansions as in (7.6), we obtain

$$\begin{aligned} \frac{d\mathbf{E}|R_t|^{2m}}{dt} &= \sum_{p+q+1 \geq 3}^{M_0} \frac{1}{n^{\frac{p+q+1}{2}}} \sum_{a=1}^n \sum_{B=n+1}^{2n} \left( \frac{\kappa^{(p+1, q)}}{p!q!} \mathbf{E} \left[ \frac{\partial^{p+q+1} (R_t)^m (R_t^*)^m}{\partial h_{aB}^{p+1} \partial h_{Ba}^q} \right] + \frac{\kappa^{(q, p+1)}}{p!q!} \mathbf{E} \left[ \frac{\partial^{p+q+1} (R_t)^m (R_t^*)^m}{\partial h_{aB}^q \partial h_{Ba}^{p+1}} \right] \right) \\ &\quad + O_{\prec}(n^{-100m}), \end{aligned} \quad (\text{D.3})$$

where we stopped the cumulant expansions at a sufficiently high power  $M_0 = O(m)$  such that the error is bounded by  $O_{\prec}(n^{-100m})$  using the moment condition in (2.1) and that  $\|G\| \leq |\eta|^{-1}$ . To estimate the terms on the right side of (D.3) we introduce the following lemma, whose proof is postponed till the end of this section.

**Lemma D.2.** *For any  $\mathbf{u}, \mathbf{v} \in [1, 2n]$ , we have*

$$|(G_1 A_1 G_2)_{\mathbf{u}\mathbf{v}}| \prec \sqrt{\frac{\rho_1 \rho_2}{|\eta_1 \eta_2|}}, \quad |(G_1 A_1 G_2 A_2 G_1)_{\mathbf{u}\mathbf{v}}| \prec \frac{\rho_1}{|\eta_1 \eta_2|}, \quad (\text{D.4})$$

for any  $\|A_1\| + \|A_2\| \lesssim 1$  and  $n|\eta_i|\rho_i \gtrsim 1$ . Thus for any  $a \in [1, n]$  and  $B \in [n+1, 2n]$ , we have

$$\frac{\partial \langle G_1 A_1 G_2 A_2 \rangle}{\partial h_{aB}} = -\frac{1}{n} \left( (G_1 A_1 G_2 A_2 G_1)_{Ba} + (G_2 A_2 G_1 A_1 G_2)_{Ba} \right) = O_{\prec} \left( \frac{\rho_1 + \rho_2}{n|\eta_1 \eta_2|} \right), \quad (\text{D.5})$$

and the same holds if we switch the index  $a$  with  $B$ . In general for any  $p, q \in \mathbb{N}$  with  $p+q \geq 1$ , we have

$$\left| \frac{\partial^{p+q} \langle G_1 A_1 G_2 A_2 \rangle}{\partial h_{aB}^p \partial h_{Ba}^q} \right| \prec \frac{\rho_1 + \rho_2}{n|\eta_1 \eta_2|} (\rho_1 + \rho_2 + \mathbb{1}_{a=B-n})^{p+q-1}, \quad (\text{D.6})$$

for any  $\|A_1\| + \|A_2\| \lesssim 1$  and  $n|\eta_i|\rho_i \gtrsim 1$ .

Thus by a naive power counting using (D.6), the terms on the right side of (D.3) are bounded by

$$\sum_{p+q+1 \geq 3}^{K_0} \frac{n^2}{n^{\frac{p+q+1}{2}}} \sum_{s=1}^{2m} \left( \frac{\rho_1 + \rho_2}{n|\eta_1 \eta_2|} \right)^s \mathbf{E}|R_t|^{2m-s}. \quad (\text{D.7})$$

Note that the above naive estimate is already enough to show (D.2) as long as  $p+q+1 \geq 4$ . It then suffices to estimate the most critical third order terms with  $p+q+1 = 3$  in (D.3), i.e.,

$$\frac{1}{n^{3/2}} \sum_{a, B} \mathbf{E} \left[ (\partial R_t)^3 R_t^{2m-3} + \partial R_t (\partial^2 R_t) R_t^{2m-2} + (\partial^3 R_t) R_t^{2m-1} \right], \quad (\text{D.8})$$

where each  $\partial$  represents either  $\partial/\partial h_{aB}$  or  $\partial/\partial h_{Ba}$  and each  $R_t$  represents  $R_t$  itself or its conjugate  $R_t^*$ . By a direct computation, using (D.5), the first part in (D.8) is bounded by

$$\begin{aligned} \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ (\partial R_t)^3 R_t^{2m-3} \right] &\lesssim \frac{1}{n^{9/2}} \sum_{a,B} \mathbf{E} \left[ \left| (G_1 A_1 G_2 A_2 G_1)_{Ba} + (G_2 A_2 G_1 A_1 G_2)_{Ba} \right|^3 |R_t|^{2m-3} \right] \\ &\prec \frac{1}{n^{7/2}} \mathbf{E} \left[ \left( \frac{\rho_1 \langle \text{Im } G_1 A_1 G_2 A_2 \text{Im } G_1 A_2^* G_2^* A_1^* \rangle}{|\eta_1^3 \eta_2|} + \frac{\rho_2 \langle \text{Im } G_2 A_2 G_1 A_1 \text{Im } G_2 A_1^* G_1^* A_2^* \rangle}{|\eta_1 \eta_2^3|} \right) |R_t|^{2m-3} \right] \\ &\prec \frac{(\rho^*)^2}{n^{7/2} |\eta_1 \eta_2|^{5/2} (\eta_*)^2} \mathbf{E} |R_t|^{2m-3}, \end{aligned} \quad (\text{D.9})$$

with  $\rho^* := \rho_1 \vee \rho_2$  and  $\eta_* := |\eta_1| \wedge |\eta_2|$ , where we used the Ward identity in the second line, and in the last step we used the following inequality

$$\langle G_1 A_1 G_2 A_2 G_1^* A_2^* G_2^* A_1^* \rangle \leq \|A_1\| \|A_2\| \sqrt{\frac{\langle \text{Im } G_1 A_1 \text{Im } G_2 A_1^* \rangle \langle \text{Im } G_2 A_2^* \text{Im } G_1 A_2 \rangle}{\eta_1^2 \eta_2^2}} \prec \sqrt{\frac{\rho_1 \rho_2}{|\eta_1 \eta_2|^3}}. \quad (\text{D.10})$$

Note that in (D.9) we have gained an additional  $1/(n\eta_*\rho^*)$  compared to the naive size in (D.7) by power counting, since we are summing up the products of two off-diagonal resolvent entries.

We next compute the second term in (D.8). Using (D.5) and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \partial R_t (\partial^2 R_t) R_t^{2m-2} \right] &\prec \frac{(\rho_1 + \rho_2)^2}{n^{7/2} |\eta_1 \eta_2|} \sum_{a,B} \mathbf{E} \left[ \left( |(G_1 A_1 G_2 A_2 G_1)_{Ba}| + |(G_2 A_2 G_1 A_1 G_2)_{Ba}| \right) |R_t|^{2m-2} \right] \\ &\prec \frac{1}{n^{3/2}} \frac{(\rho_1 + \rho_2)^2}{|\eta_1 \eta_2|} \mathbf{E} \left[ \sqrt{\frac{1}{n^2} \sum_{a,B} |(G_1 A_1 G_2 A_2 G_1)_{Ba}|^2} |R_t|^{2m-2}} \right] \\ &\prec \frac{(\rho^*)^{5/2}}{n^2 |\eta_1 \eta_2|^{9/4}} \mathbf{E} |R_t|^{2m-2}, \end{aligned} \quad (\text{D.11})$$

where we gained an additional  $1/\sqrt{n\eta_*\rho^*}$  compared to the naive size in (D.7), since there exists one off-diagonal resolvent factor in the summation. Similarly, using (D.5), we see that the last part in (D.8) also contains at least one off-diagonal resolvent factor and so we have an additional smallness:

$$\frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ (\partial^3 R_t) R_t^{2m-1} \right] \lesssim \frac{(\rho^*)^{5/2}}{n |\eta_1 \eta_2| \sqrt{\eta_*}} \mathbf{E} |R_t|^{2m-1}. \quad (\text{D.12})$$

Combining (D.9), (D.11)–(D.12) with (D.7), we obtain that

$$\begin{aligned} \left| \frac{d\mathbf{E}|R_t|^{2m}}{dt} \right| &\prec \frac{(\rho^*)^{5/2}}{n |\eta_1 \eta_2| \sqrt{\eta_*}} \mathbf{E} |R_t|^{2m-1} + \frac{(\rho^*)^{5/2}}{n^2 |\eta_1 \eta_2|^{9/4}} \mathbf{E} |R_t|^{2m-2} + \frac{(\rho^*)^2}{n^{7/2} |\eta_1 \eta_2|^{5/2} \eta_*^2} \mathbf{E} |R_t|^{2m-3} \\ &\quad + \sum_{k \geq 4}^{2m} \frac{n^2}{n^{k/2}} \left( \frac{\rho^*}{n |\eta_1 \eta_2|} \right)^k \mathbf{E} |R_t|^{2m-k}. \end{aligned} \quad (\text{D.13})$$

Further assuming that  $|\eta_1| \sim |\eta_2| \sim \eta$ ,  $\rho_1 \sim \rho_2 \sim \rho$  and using Young's inequality, we obtain that

$$\left| \frac{d\mathbf{E}|R_t|^{2m}}{dt} \right| \prec \left( 1 + \frac{\rho^{5/2}}{\eta^{1/2}} + \frac{\rho^2}{\sqrt{n\eta}} \right) \left( \mathbf{E} |R_t|^{2m} + \left( \frac{1}{n |\eta_1 \eta_2|} \right)^{2m} \right). \quad (\text{D.14})$$

Thus (D.2) follows directly from (D.14) using that  $\rho \lesssim \eta^{1/3}$  and  $n\eta\rho \gtrsim 1$ . We hence finish the proof of Proposition D.1.  $\square$

We conclude this section with the proof of Lemma D.2.

*Proof of Lemma D.2.* Using the Cauchy-Schwarz inequality and the Ward identity, we have

$$\left| (G_1 A G_2)_{uv} \right| \leq \frac{\|A\| \sqrt{(\text{Im } G_1)_{uu} (\text{Im } G_2)_{vv}}}{\sqrt{|\eta_1 \eta_2|}} \prec \frac{1}{\sqrt{|\eta_1 \eta_2|}} \left( \sqrt{\rho_1 \rho_2} + \frac{1}{n \sqrt{|\eta_1 \eta_2|}} \right), \quad (\text{D.15})$$

where in the last step we used the local law in (3.8). This proves the first estimate in (D.4). Similarly to (D.15), we further have

$$\begin{aligned} |(G_1 A_1 G_2 A_2 G_1)_{\text{uv}}| &\leq \sqrt{(G_1 A_1 G_2 A_2 A_2^* G_2^* A_1^* G_1^*)_{\text{uu}} (G_1 G_1^*)_{\text{vv}}} \leq \frac{\|A_1\| \|A_2\| \sqrt{(\text{Im } G_1)_{\text{vv}} (\text{Im } G_1)_{\text{uu}}}}{|\eta_1 \eta_2|} \\ &\prec \frac{1}{|\eta_1 \eta_2|} \left( \rho_1 + \frac{1}{n|\eta_1|} \right), \end{aligned} \quad (\text{D.16})$$

where we also used (3.9) and that  $\|G(\eta)\| \leq |\eta|^{-1}$ . This proves the second estimate in (D.4).

Note that the differentiation rule in (D.5) follows directly from (7.7). Combining this with (D.16) we have proved the last estimate in (D.5). This also proves (D.6) for  $p + q = 1$ . By further computations using (7.7), the corresponding partial derivatives of  $\langle G_1 A_1 G_2 A_2 \rangle$  with  $p + q = 2$  are given by linear combinations of the following terms

$$\begin{aligned} &\frac{1}{n} (G_1 A_1 G_2 A_2 G_1)_{Ba} (G_1)_{Ba}, \quad \frac{1}{n} (G_1 A_1 G_2 A_2 G_1)_{aa} (G_1)_{BB}, \quad \frac{1}{n} (G_1 A_1 G_2 A_2 G_1)_{BB} (G_1)_{aa}, \\ &\frac{1}{n} (G_1 A_1 G_2)_{Ba} (G_2 A_2 G_1)_{Ba}, \quad \frac{1}{n} (G_1 A_1 G_2)_{aa} (G_2 A_2 G_1)_{BB}, \quad \frac{1}{n} (G_1 A_1 G_2)_{BB} (G_2 A_2 G_1)_{aa}, \end{aligned} \quad (\text{D.17})$$

and the terms obtained by switching the subscript 1 with 2. Thus for  $p + q = 2$ , (D.6) follows from (D.15), (D.16) and the local law in (3.8). For a general  $p + q \geq 2$ , one proves (D.6) similarly by induction using (D.15) and (D.16). We hence finish the proof of Lemma D.2.  $\square$

**D.2. GFT for  $\text{Im } G_1 A \text{Im } G_2 A^*$  with  $|z_1 - z_2|$ -decorrelation.** In this section, we consider a special form in terms of (D.1) and establish an improved GFT to remove the Gaussian component in Part A of Theorem 3.3 for a large  $|z_1 - z_2| \gtrsim |\eta_i|/\rho_i$ . Again for simplicity, we do it only for the regime  $|\eta_1| \sim |\eta_2|$  and  $\rho_1 \sim \rho_2$ .

**Proposition D.3.** *Define*

$$\widehat{R}_t := \langle (\text{Im } G_1 A \text{Im } G_2 - \widehat{M}_{12}^A) A^* \rangle, \quad G_i := G_t^{z_i}(i\eta_i), \quad i = 1, 2, \quad (\text{D.18})$$

where  $\widehat{M}_{12}^A = \widehat{M}_{12}^A(\eta_1, z_1, \eta_2, z_2)$  is the corresponding deterministic limit defined as in (3.13) using that  $\text{Im } G = (G - G^*)/(2i)$ . Fix any  $m > 1$ . Then for any spectral parameters  $|\eta_1| \sim |\eta_2| \sim \eta$  and  $\rho_1 \sim \rho_2 \sim \rho$ , the following holds

$$\left| \frac{d\mathbf{E}|\widehat{R}_t|^{2m}}{dt} \right| \prec \left( 1 + \frac{1}{\sqrt{n\eta}} \right) \left( \mathbf{E}|\widehat{R}_t|^{2m} + \left( \frac{1}{\sqrt{n\eta\rho}(|z_1 - z_2| + \frac{\eta}{\rho})} \right)^{2m} \right), \quad (\text{D.19})$$

uniformly for any  $t \geq 0$ ,  $\|A\| \lesssim 1$  and  $n\eta\rho \gtrsim 1$ .

*Proof of Proposition D.3.* Applying Itô's formula to  $|\widehat{R}_t|^{2m} = (\widehat{R}_t)^{2m}$  and performing the cumulant expansions, we obtain the analogue of (D.3) for  $\widehat{R}_t$  using that  $\text{Im } G = (G - G^*)/(2i)$ . The most critical third order terms with  $p + q + 1 = 3$  are then given by (D.8) in terms of  $\widehat{R}_t$ . Note that (D.5) implies

$$\begin{aligned} \frac{\partial \widehat{R}_t}{\partial a_B} &= -\frac{1}{2ni} \left( (G_1 A \text{Im } G_2 A^* G_1)_{Ba} - (G_1^* A \text{Im } G_2 A^* G_1^*)_{Ba} \right. \\ &\quad \left. + (G_2 A^* G_1 A G_2)_{Ba} - (G_2^* A^* \text{Im } G_1 A G_2^*)_{Ba} \right). \end{aligned} \quad (\text{D.20})$$

By a direct computation, using (D.20), the first part of (D.8) is bounded by (c.f., (D.9))

$$\begin{aligned} \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ (\partial \widehat{R}_t)^3 (\widehat{R}_t)^{2m-3} \right] &\lesssim \frac{1}{n^{9/2}} \sum_{a,B} \mathbf{E} \left[ \left| (G_1 A \text{Im } G_2 A^* G_1)_{Ba} + (G_2 A^* \text{Im } G_1 A G_2)_{Ba} \right|^3 |\widehat{R}_t|^{2m-3} \right] \\ &\lesssim \frac{\rho}{n^{7/2} \eta^2} \mathbf{E} \left[ \left( \frac{\langle \text{Im } G_1 A \text{Im } G_2 A^* \text{Im } G_1 A \text{Im } G_2 A^* \rangle}{\eta_1^2} + \frac{\langle \text{Im } G_2 A^* \text{Im } G_1 A \text{Im } G_2 A^* \text{Im } G_1 A \rangle}{\eta_2^2} \right) |\widehat{R}_t|^{2m-3} \right] \\ &\lesssim \frac{\rho}{n^{5/2} \eta^4} \mathbf{E} \left[ (\langle \text{Im } G_1 A \text{Im } G_2 A^* \rangle)^2 |\widehat{R}_t|^{2m-3} \right] \\ &\lesssim \frac{\rho}{n^{5/2} \eta^4} \left( \mathbf{E} |\widehat{R}_t|^{2m-1} + \frac{1}{(|z_1 - z_2| + \frac{\eta}{\rho})^2} \mathbf{E} |\widehat{R}_t|^{2m-3} \right), \end{aligned} \quad (\text{D.21})$$



where in the penultimate line we used

$$\langle \text{Im } G_1 A \text{Im } G_2 A^* \text{Im } G_1 A \text{Im } G_2 A^* \rangle \leq n \langle \text{Im } G_1 A \text{Im } G_2 A^* \rangle^2, \quad (\text{D.22})$$

and in the last line we used the upper bound in (3.15) with  $|\eta_1| \sim |\eta_2| \sim \eta$  and  $\rho_1 \sim \rho_2 \sim \rho$ . A similar upper bound also applies to the second part of (D.8) (c.f., (D.11)), i.e.,

$$\begin{aligned} \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \partial \widehat{R}_t (\partial^2 \widehat{R}_t) (\widehat{R}_t)^{2m-2} \right] &\prec \frac{\rho^2}{n^{7/2} \eta^2} \sum_{a,B} \mathbf{E} \left[ \left( |(G_1 A \text{Im } G_2 A^* G_1)_{Ba}| + |(G_2 A \text{Im } G_1 A^* G_2)_{Ba}| \right) |\widehat{R}_t|^{2m-2} \right] \\ &\lesssim \frac{\rho^2}{n^{3/2} \eta^2} \mathbf{E} \left[ \sqrt{\frac{\langle \text{Im } G_1 A \text{Im } G_2 A^* \text{Im } G_1 A \text{Im } G_2 A^* \rangle}{n \eta^2}} |\widehat{R}_t|^{2m-2} \right] \\ &\lesssim \frac{\rho^2}{n^{3/2} \eta^3} \left( \mathbf{E} |\widehat{R}_t|^{2m-1} + \frac{1}{|z_1 - z_2| + \frac{\eta}{\rho}} \mathbf{E} |\widehat{R}_t|^{2m-2} \right). \end{aligned} \quad (\text{D.23})$$

We use the same naive bound as in (D.12) to estimate the last part  $n^{-3/2} \sum_{a,B} \mathbf{E} [(\partial^3 \widehat{R}_t) (\widehat{R}_t)^{2m-1}]$ . Combining this with (D.21) and (D.23), the third order terms with  $p+q+1=3$  are bounded by

$$\begin{aligned} &\frac{\rho^{5/2}}{n \eta^{5/2}} \mathbf{E} |\widehat{R}_t|^{2m-1} + \frac{\rho^2}{n^{3/2} \eta^3} \frac{1}{|z_1 - z_2| + \frac{\eta}{\rho}} |\widehat{R}_t|^{2m-2} + \frac{\rho}{n^{5/2} \eta^4} \frac{1}{(|z_1 - z_2| + \frac{\eta}{\rho})^2} \mathbf{E} |\widehat{R}_t|^{2m-3} \\ &\lesssim \frac{1}{\sqrt{n} \eta} \left( \mathbf{E} |\widehat{R}_t|^{2m} + \left( \frac{\rho^2}{n \eta^2} \sqrt{n \eta \rho} \right)^{2m} + \left( \frac{1}{\sqrt{n \eta \rho} (|z_1 - z_2| + \frac{\eta}{\rho})} \right)^{2m} \right). \end{aligned} \quad (\text{D.24})$$

All the higher order terms with  $p+q+1 \geq 4$  can be bounded similarly with less efforts due to the moment condition in (2.1). More precisely, for general  $p+q+1 \geq 4$ , they are given by

$$\frac{1}{n^{\frac{p+q+1}{2}}} \sum_{a,B} \left( \sum_{1 \leq k \leq \lfloor \frac{p+q+1}{2} \rfloor} + \sum_{\lfloor \frac{p+q+1}{2} \rfloor < k \leq p+q+1} \right) \sum_{\substack{d_1 + \dots + d_k = p+q+1, \\ d_i \geq 1}} \mathbf{E} \left[ \prod_{i=1}^k (\partial^{d_i} \widehat{R}_t) (\widehat{R}_t)^{2m-k} \right] =: \Pi_1 + \Pi_2, \quad (\text{D.25})$$

where we split the above sum using the integer  $k$ , which is the number of times that  $\partial_{aB}$  or  $\partial_{Ba}$  acts on  $\widehat{R}_t$ . For any  $\lfloor \frac{p+q+1}{2} \rfloor < k \leq p+q+1$ , since  $\sum_{i=1}^k d_i = p+q+1$ , then there exists at least one  $d_i = 1$ , which implies that there exists at least one off-diagonal entry of  $G_1 A \text{Im } G_2 A^* G_1$  from (D.20). Using similar arguments as in (D.23) and the upper bound in (3.15), the second part in (D.25) is bounded by

$$\begin{aligned} |\Pi_2| &\prec \frac{\rho^{p+q}}{n^{\frac{p+q+3}{2}}} \sum_{a,B} \sum_{\lfloor \frac{p+q+1}{2} \rfloor < k \leq p+q+1} \left( \frac{1}{n \eta^2} \right)^{k-1} \mathbf{E} \left[ \left( |(G_1 A \text{Im } G_2 A^* G_1)_{Ba}| + |(G_2 A \text{Im } G_1 A^* G_2)_{Ba}| \right) (\widehat{R}_t)^{2m-k} \right] \\ &\lesssim \frac{\rho^{p+q}}{n^{\frac{p+q-1}{2}}} \sum_{\lfloor \frac{p+q+1}{2} \rfloor < k \leq p+q+1} \frac{1}{\rho^{2(k-1)} \eta} \left( \frac{\rho^2}{n \eta^2} \right)^{k-1} \mathbf{E} \left[ \langle \text{Im } G_1 A \text{Im } G_2 A^* \rangle (\widehat{R}_t)^{2m-k} \right] \\ &\lesssim \frac{1}{\sqrt{n} \eta} \left( \mathbf{E} |\widehat{R}_t|^{2m} + \left( \frac{\rho^2}{n \eta^2} \sqrt{n \eta \rho} + \frac{1}{\sqrt{n \eta \rho} (|z_1 - z_2| + \frac{\eta}{\rho})} \right)^{2m} \right), \end{aligned} \quad (\text{D.26})$$

where we also used that  $p+q+1 \geq 4$  and  $\rho \gtrsim n^{-1/4}$  for any  $n \eta \rho \gtrsim 1$  from (3.7) with  $|z| > 1$ . For any  $1 \leq k \leq \lfloor \frac{p+q+1}{2} \rfloor$  with  $p+q+1 \geq 4$ , we use a naive power counting by (D.6) to obtain that

$$|\Pi_1| \prec \frac{n^2}{n^{\frac{p+q+1}{2}}} \sum_{1 \leq k \leq \lfloor \frac{p+q+1}{2} \rfloor} \rho^{p+q+1} \left( \frac{1}{n \eta^2} \right)^k \mathbf{E} \left[ |\widehat{R}_t|^{2m-k} \right] \lesssim \mathbf{E} |\widehat{R}_t|^{2m} + \left( \frac{\rho^2}{n \eta^2} \right)^{2m}. \quad (\text{D.27})$$

Combining with (D.24) and (D.26), using that  $\rho \lesssim \eta^{1/3}$  and  $|z_1 - z_2| + \eta/\rho \lesssim 1$ , we hence finish the proof of Proposition D.3.  $\square$

**D.3. GFT for  $\langle GFGF^* \rangle$  with  $\rho$ -improvement.** To prove Theorem 3.5 with the  $\rho$ -improvements, we establish the following refined GFTs to remove the Gaussian component in Proposition 5.8.

**Proposition D.4.** *Let*

$$R_t^F := \langle (G_t^z(i\eta) F G_t^z(i\eta) - M_{12}^F) F^* \rangle, \quad \eta \in \mathbb{R}_+, \quad z \in \mathbb{C}, \quad (\text{D.28})$$

where  $M_{12}^F = M_{12}^F(\eta, z, \eta, z)$  is the corresponding deterministic limits given by (3.13) and we also define

$$S_t^F := \langle (G_t^z(\imath\eta) - M^z)F \rangle, \quad S_t^{2,F} := \langle G_t^z(\imath\eta)FG_t^z(\imath\eta) - M_{12}^F \rangle,$$

with  $M^z = M^z(\imath\eta)$  given by (3.1). Fix any  $m > 1$ . Then for any  $t \geq 0$  and  $n\eta\rho \gtrsim 1$  we have

$$\left| \frac{d\mathbf{E}|R_t^F|^{2m}}{dt} \right| \prec \left( 1 + \frac{1}{\sqrt{n\eta}} \right) \left( \mathbf{E}|R_t^F|^{2m} + \left( \frac{\rho^2}{n\eta^2} \sqrt{n\eta\rho} \right)^{2m} \right). \quad (\text{D.29})$$

Assuming that the following

$$|R_t^F| \prec \frac{\rho^2 \sqrt{n\eta\rho}}{n\eta^2}, \quad (\text{D.30})$$

holds uniformly for any  $t \geq 0$  and  $n\eta\rho \gtrsim 1$ , then we have

$$\left| \frac{d\mathbf{E}|S_t^F|^{2m}}{dt} \right| \lesssim \left( 1 + \frac{\rho}{\sqrt{n\eta}} \right) \left( \mathbf{E}|S_t^F|^{2m} + \left( \frac{\rho}{n\eta} \right)^{2m} \right), \quad (\text{D.31})$$

$$\left| \frac{d\mathbf{E}|S_t^{2,F}|^{2m}}{dt} \right| \lesssim \left( 1 + \frac{\rho}{\sqrt{n\eta}} \right) \left( \mathbf{E}|S_t^{2,F}|^{2m} + \left( \frac{\rho}{n\eta^2} \right)^{2m} \right). \quad (\text{D.32})$$

*Proof of Proposition D.4.* Note that that  $\langle GFGF^* \rangle = \langle \text{Im} GF \text{Im} GF^* \rangle$  from Lemma 3.7. The proof of the first GFT in (D.29) is completely analogous to the proof of Proposition D.3 with  $A = F$ ,  $\eta_1 = \eta_2 = \eta$  and  $z_1 = z_2 = z$ . The only difference is to replace the upper bound in (3.15) with the first upper bound in (5.67). Repeating the same arguments as in (D.21)-(D.24), the most critical third order terms are thus bounded by

$$\frac{\rho^{5/2}}{n\eta^{5/2}} \mathbf{E}|R_t^F|^{2m-1} + \frac{\rho^5}{n^{3/2}\eta^4} \mathbf{E}|R_t^F|^{2m-2} + \frac{\rho^7}{n^{5/2}\eta^6} \mathbf{E}|R_t^F|^{2m-3} \lesssim \frac{1}{\sqrt{n\eta}} \left( \mathbf{E}|R_t^F|^{2m} + \left( \frac{\rho^2}{n\eta^2} \sqrt{n\eta\rho} \right)^{2m} \right), \quad (\text{D.33})$$

where we also used that  $n\eta\rho \gtrsim 1$ . Repeating the same arguments as in (D.25)-(D.27), we obtain similar upper bounds for the higher order terms with  $p + q + 1 \geq 4$ , *i.e.*,

$$\frac{1}{\sqrt{n\eta}} \left( \mathbf{E}|\widehat{R}_t|^{2m} + \left( \frac{\rho^2}{n\eta^2} \sqrt{n\eta\rho} \right)^{2m} \right), \quad (\text{D.34})$$

where we used the first upper bound in (5.67) to replace the upper bound in (3.15). This, together with (D.33), prove the first GFT in (D.29) of Proposition D.1

The other two GFTs in (D.31)-(D.32) with less  $\rho$ -improvements can be proved with much less effort, so we only sketch their proofs. We will only consider  $S_t^F$  for simplicity and the other one can be proved exactly in the same way. Applying Ito's formula to  $|S_t^F|^{2m}$  and performing cumulant expansions, we obtain the analogue of (D.3). We next estimate the most critical third order terms with  $p + q + 1 = 3$ , which are splitted into three parts as in (D.8). Using (7.7) and the first upper bound in (D.4), the first part as in (D.8) is bounded by

$$\begin{aligned} \frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ (\partial S_t^F)^3 (S_t^F)^{2m-3} \right] &\lesssim \frac{1}{n^{9/2}} \sum_{a,B} \mathbf{E} \left[ \left| (GFG)_{Ba} + (GF^*G)_{Ba} \right|^3 |S_t^F|^{2m-3} \right] \\ &\prec \frac{\rho}{n^{7/2}\eta} \mathbf{E} \left[ \frac{\langle \text{Im} GF \text{Im} GF^* \rangle}{\eta^2} |S_t^F|^{2m-3} \right] \lesssim \frac{\rho^4}{n^{7/2}\eta^4} \mathbf{E}|S_t^F|^{2m-3}, \end{aligned} \quad (\text{D.35})$$

where we also used the assumption (D.30) and the first upper bound in (5.67). A similar upper bound also applies to the second part, *i.e.*,

$$\frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \partial S_t^F (\partial^2 S_t^F) (S_t^F)^{2m-2} \right] \prec \frac{\rho^{7/2}}{n^2 \eta^{5/2}} \mathbf{E}|S_t^F|^{2m-2}.$$

We use the following naive bound from (D.4) to estimate the last part of third order terms:

$$\frac{1}{n^{3/2}} \sum_{a,B} \mathbf{E} \left[ \partial^3 S_t^F (S_t^F)^{2m-1} \right] \prec \frac{\rho^{5/2}}{n\eta^{3/2}} \mathbf{E}|S_t^F|^{2m-1}.$$

All the higher order terms with  $p + q + 1 \geq 4$  can be bounded similarly with much less efforts, so we omit the details. Therefore, using that  $\rho \lesssim \eta^{1/3}$  from (3.7), we have proved (D.31). The last GFT in (D.32) can be proved similarly. We hence complete the proof of Proposition D.4.  $\square$

**D.4. Proof of the local laws.** We start with proving Part B of Theorem 3.3 for any i.i.d. matrix, while Part A of Theorem 3.3 with a Gaussian component was already proved in Section 5.1. And the proof of Theorem 3.5 is similar, so we only sketch its proof at the end of this section with some minor modifications.

*Proof of Theorem 3.3 (Part B) for i.i.d. matrix.* The proof of Theorem 3.3 (Part B) is split into two steps. In Step 1, we prove (3.17) without the  $|z_1 - z_2|$ -gain. Indeed we prove a more general result, *i.e.*,

$$|\langle (G^{z_1}(i\eta_1)A_1G^{z_2}(i\eta_2) - M_{12}^{A_1})A_2 \rangle| \prec \frac{1}{n|\eta_1\eta_2|}, \quad (\text{D.36})$$

uniformly in any spectral parameters in  $n\ell \geq n^\epsilon$  with  $\ell = \eta\rho$ , and  $\eta \sim |\eta_1| \sim |\eta_2|$ ,  $\rho \sim \rho_1 \sim \rho_2$ . In Step 2, based on the result from Step 1, we further prove (3.17) with the  $|z_1 - z_2|$ -decorrelation, *i.e.*,

$$|\langle (\text{Im } G^{z_1}(i\eta_1)A\text{Im } G^{z_2}(i\eta_2) - \widehat{M}_{12}^A)A^* \rangle| \prec \frac{1}{\sqrt{n\ell}(|z_1 - z_2| + \frac{\eta}{\rho})}, \quad (\text{D.37})$$

uniformly in  $n\ell \geq n^\epsilon$ , with  $\ell = \eta\rho$ , and  $\eta \sim |\eta_1| \sim |\eta_2|$ ,  $\rho \sim \rho_1 \sim \rho_2$ . This, together with (D.36) from Step 1, complete the proof of Theorem 3.3 (Part B).

**Step 1: Proof of (D.36).** Recall Part A of Theorem 3.3 that has already been proved in Section 5.1. Fixing  $\mathfrak{s} > 0$ , for any matrix of the form  $X_{\mathfrak{s}} := \sqrt{1 - \mathfrak{s}^2}X + \mathfrak{s}X^{\text{Gin}}$ , we have

$$|\langle (G_{\mathfrak{s}}^{z_1}(i\eta_1)A_1G_{\mathfrak{s}}^{z_2}(i\eta_2) - M_{12}^{A_1})A_2 \rangle| \prec \left( \frac{1}{(n\ell)^{1/2} |z_1 - z_2| + \frac{|\eta_1|}{\rho_1} + \frac{|\eta_2|}{\rho_2}} \right) \wedge \left( \frac{1}{n\eta_1\eta_2} \right). \quad (\text{D.38})$$

Fix  $z_i, \eta_i$  in (D.38) and recall the definition of  $R_t$  in (D.1). Using Proposition D.1 together with a Gronwall argument, we know that the Gaussian component of size  $\mathfrak{s} \sim 1$  in (D.38) can be removed with an error  $1/(n\eta_1\eta_2)$ . We hence prove (D.36).

**Step 2: Proof of (D.37).** We first assume that  $|\eta_i| \sim \eta \gtrsim n^{-1/2}$ . Similarly to Step 1 above, Proposition D.3 implies that the Gaussian component added in Part A of Theorem 3.3 can be removed by a Gronwall argument, with an error  $(\sqrt{n\ell}(|z_1 - z_2| + \eta/\rho))^{-1}$ . This, together with (D.38) from Step 1, prove (D.37) for any  $|\eta_i| \sim \eta \gtrsim n^{-1/2}$ .

We next extend this result to the complementary regime, *i.e.*,  $|\eta_i| \sim \eta \lesssim n^{-1/2}$ . Recall  $\eta \sim |\eta_1| \sim |\eta_2|$  and  $\rho \sim \rho_1 \sim \rho_2$  for simplicity, with  $n\eta\rho \gtrsim 1$ , and choose  $T \sim \sqrt{\eta/\rho} \lesssim 1$ . Then from Lemma 5.2, there exists  $\eta_{i,0}$  and  $z_{i,0}$  such that  $\eta_{i,T} = \eta_i$  and  $z_{i,T} = z_i$  with  $\rho_{i,0} \sim \rho$ ,  $\eta_{i,0} \gtrsim \rho T \gtrsim n^{-1/2}$  and  $n\ell_0 \geq n\ell \geq n^\epsilon$  with  $\ell_0$  defined in (5.16) at  $t = 0$ . As shown in Step 1 and at the beginning of Step 2 for  $\eta_{i,0} \gtrsim n^{-1/2}$ , we know that the initial condition (5.17) of Proposition 5.4 is satisfied. Thus Proposition 5.4 implies that

$$|\langle (\text{Im } G_T^{z_1}(i\eta_1)A\text{Im } G_T^{z_2}(i\eta_2) - \widehat{M}_{12}^A)A^* \rangle| \prec \frac{1}{\sqrt{n\ell}(|z_1 - z_2| + \frac{\eta}{\rho})} \wedge \frac{1}{n\eta^2}, \quad T \sim \sqrt{\eta/\rho}. \quad (\text{D.39})$$

Combining this with Proposition D.3 and using that  $n\ell \gtrsim n^\epsilon$ , we remove the Gaussian component of size  $T \sim \sqrt{\eta/\rho}$  in (D.39) and hence prove (D.37) for  $\eta \lesssim n^{-1/2}$ . This completes the proof of Part B of Theorem 3.3.  $\square$

*Proof of Theorem 3.5.* The proof of Theorem 3.5, in particular the local law in (3.28), is exactly the same as the proof of (D.37) presented above, using Proposition 5.8 and Proposition D.4 as inputs. Additionally, assuming the condition in (D.30) is satisfied (*i.e.*, the local law in (3.28) holds), one obtains the other two local laws in (3.26)-(3.27) (with less  $\rho$ -improvement) using the GFT arguments in (D.31)-(D.32) with much less efforts, so we omit the details. The fact that the assumptions on the initial conditions

$$\langle (G^{z_0}FG^{z_0}(i\eta_0) - M_{12}^F)F^* \rangle \prec \frac{\rho_0^2 \sqrt{n\eta_0\rho_0}}{n\eta_0^2}, \quad (\text{D.40})$$

$$\langle (G^{z_0}(i\eta_0) - M^{z_0})F \rangle \prec \frac{\rho_0}{n\eta_0}, \quad \langle (G^{z_0}FG^{z_0}(i\eta_0) - M_{12}^F) \rangle \prec \frac{\rho_0}{n\eta_0^2}. \quad (\text{D.41})$$

in Proposition 5.8 are satisfied immediately follows from (5.19) with  $|z_0| - 1 \gtrsim 1$ .  $\square$

## APPENDIX E. GINIBRE COMPUTATIONS

In this section we consider the complex Ginibre ensemble, which is a special case of  $X$  with i.i.d. Gaussian entries. Recall that the joint probability density of the eigenvalues  $\{\sigma_i\}_{i=1}^n$  of a complex Ginibre matrix is given by (see *e.g.*, [59, Lemma 3] and references therein)

$$\begin{aligned}\rho_n(z_1, \dots, z_n) &:= \frac{n^n}{\pi^n n! \dots n!} \exp\left(-n \sum_i |z_i|^2\right) \prod_{i < j} (n|z_i - z_j|^2) \\ &= \frac{n^n}{\pi^n n!} e^{-n|z|^2} \det\left(K_n(z_i, z_j)\right)_{i,j=1}^n,\end{aligned}\tag{E.1}$$

which forms a *determinantal point process* with the kernel

$$K_n(z, w) := \sum_{l=0}^{n-1} \frac{(nz\bar{w})^l}{l!}.$$

Recalling [59, Eq. (15)-(16)], then for any function  $\mathfrak{g} : \mathbb{C} \rightarrow [0, 1]$  we have

$$\mathbf{E}^{\text{Gin}} \prod_{i=1}^n (1 - \mathfrak{g}(\sigma_i)) = \det(1 - \sqrt{\mathfrak{g}} \tilde{K}_n \sqrt{\mathfrak{g}}),\tag{E.2}$$

where  $\det(\cdot)$  is the Fredholm determinant (see *e.g.*, [59, Definition 4] for definition) and the rescaled kernel  $\tilde{K}_n$  is given by

$$\tilde{K}_n(z, w) := \frac{n}{\pi} e^{-n(|z|^2 + |w|^2)/2} K_n(z, w) = \frac{n}{\pi} e^{-n(|z|^2 + |w|^2 - 2z\bar{w})/2} \frac{\Gamma(n, nz\bar{w})}{\Gamma(n)}.\tag{E.3}$$

Here  $\Gamma(\cdot, \cdot)$  denotes the incomplete Gamma function:

$$\Gamma(s, z) := \int_z^\infty t^{s-1} e^{-t} dt, \quad s \in \mathbb{N}, \quad z \in \mathbb{C},\tag{E.4}$$

with the integral contour going from  $z \in \mathbb{C}$  to the real infinity. Using [75, IV.(7.8)-(7.11)], or more precisely [59, Eq.(20)-(21)], we have

$$|\det(1 - K) - \exp(-\text{Tr}K)| \leq \|K\|_2 e^{(\|K\|_2 + 1)^2/2 - \text{Tr}K},\tag{E.5}$$

for any finite-rank kernel  $K$ , where

$$\text{Tr}K = \int_\Omega K(x, x) d\mu(x), \quad \|K\|_2^2 = \int_{\Omega^2} |K(x, y)|^2 d\mu(x) d\mu(y).\tag{E.6}$$

Then we state the following proposition as the analogue of [59, Proposition 5] (where we considered the rightmost eigenvalue as in Theorem 2.6).

**Proposition E.1.** *For any  $|t| \leq \sqrt{\log n}/2$  and  $0 \leq a < b \leq 2\pi$ , define*

$$A(t, a, b) := \left\{ z \in \mathbb{C} : |z| \geq 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{t}{\sqrt{4n\gamma_n}}, \quad \arg(z) \in [a, b] \right\}.\tag{E.7}$$

*Then for any function  $\mathfrak{g} : \mathbb{C} \rightarrow [0, 1]$  supported on  $A(t, a, b)$ , we obtain that*

$$\text{Tr} \sqrt{\mathfrak{g}} \tilde{K}_n \sqrt{\mathfrak{g}} = \int_t^\infty \int_a^b \mathfrak{g}(z) \frac{e^{-r}}{2\pi} d\theta dr + O\left(e^{-t} \frac{(\log \log n)^2 + |t|^2}{\log n}\right),\tag{E.8}$$

*with  $z = e^{i\theta} \left(1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4n\gamma_n}}\right)$ , and that*

$$\|\sqrt{\mathfrak{g}} \tilde{K}_n \sqrt{\mathfrak{g}}\|_2 \lesssim e^{-\sqrt{\log n}/32}.\tag{E.9}$$

The proof of Proposition E.1 is completely analogous to the proof of [59, Proposition 5] using the following lemma. The proof of Lemma E.2 is postponed to the end of this section.

**Lemma E.2.** *Recall  $\gamma_n$  in (2.3) and rescale the kernel variable as*

$$z = e^{i\theta} \left(1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4\gamma_n n}}\right), \quad \theta \in [0, 2\pi), \quad r \in \mathbb{R}.\tag{E.10}$$

Then for any  $|r| \leq \sqrt{\log n}/2$ , we have

$$\frac{\tilde{K}_n(z, z)}{\sqrt{4n\gamma_n}} = \frac{e^{-r}}{2\pi} \left( 1 + O\left(\frac{\log \log n + r^2}{\log n}\right) \right). \quad (\text{E.11})$$

Choose different  $z_1, z_2 \in \mathbb{C}$ , and rescale them as in (E.10). Then for any  $|r_i| \leq \sqrt{\log n}/2$  and  $|\theta_1 - \theta_2| \lesssim n^{-1/4}(\gamma_n)^{-1/2}$ , we have

$$\frac{|\tilde{K}_n(z_1, z_2)|}{\sqrt{n\gamma_n}} \lesssim e^{-\frac{1}{2}(r_1+r_2)} \left( 1 + O\left(\frac{\log \log n}{\log n}\right) \right), \quad (\text{E.12})$$

while for any  $|r_i| \leq \sqrt{\log n}/2$  and  $|\theta_1 - \theta_2| \gtrsim n^{-1/2+w}$  with a small  $w > 0$ , we have

$$\frac{|\tilde{K}_n(z_1, z_2)|}{\sqrt{n\gamma_n}} \lesssim \frac{\sqrt{\gamma_n}}{\sqrt{n}|\theta_1 - \theta_2|} e^{-\frac{1}{2}(r_1+r_2)} \left( 1 + O\left(\frac{\log \log n}{\log n}\right) \right). \quad (\text{E.13})$$

In addition, for any  $r_i \geq 0$ , we have the following uniform bound

$$\frac{|\tilde{K}_n(z_1, z_2)|}{\sqrt{n\gamma_n}} \lesssim |z_1 z_2| e^{-\frac{1}{6}(r_1+r_2)}. \quad (\text{E.14})$$

*Proof of Proposition E.1.* Note that from (E.6) and changing the variables as in (E.10), we have

$$\begin{aligned} \text{Tr} \sqrt{\mathfrak{g}} \tilde{K}_n \sqrt{\mathfrak{g}} &= \int_{A(t,a,b)} \mathfrak{g}(z) \tilde{K}_n(z, z) d^2z = \int_t^\infty \int_a^b \mathfrak{g}(z) \frac{\tilde{K}_n(z, z)}{\sqrt{4n\gamma_n}} \left( 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4\gamma_n n}} \right) d\theta dr \\ &= \left( \int_t^{r_0} + \int_{r_0}^\infty \right) \int_a^b \mathfrak{g}(z) \frac{\tilde{K}_n(z, z)}{\sqrt{4n\gamma_n}} \left( 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{r}{\sqrt{4\gamma_n n}} \right) d\theta dr \\ &= \int_t^\infty \int_a^b \mathfrak{g}(z) \frac{e^{-r}}{2\pi} d\theta dr + O\left( e^{-t} \frac{(\log \log n)^2 + |t|^2}{\log n} \right), \end{aligned} \quad (\text{E.15})$$

where we chose  $r_0 = 4(\log \log n + |t|)$  and we used (E.11) for the regime  $t \leq r \leq r_0$  and used (E.14) for the complementary regime  $r > r_0$ . This proves (E.8).

For any function  $\mathfrak{g} : \mathbb{C} \rightarrow [0, 1]$  supported on  $A(t, a, b)$  given by (E.7), then from (E.6) we have

$$\|\sqrt{\mathfrak{g}} \tilde{K}_n \sqrt{\mathfrak{g}}\|_2^2 = \text{Tr}(\sqrt{\mathfrak{g}} \tilde{K}_n \sqrt{\mathfrak{g}})^2 \leq \iint_{A(t,a,b)} |\tilde{K}_n(z_1, z_2)|^2 d^2z_1 d^2z_2.$$

Changing the variables as in (E.10), we use (E.12) to estimate the regime  $|r_i| \leq \sqrt{\log n}/2$ ,  $|\theta_1 - \theta_2| \lesssim n^{-1/4}(\gamma_n)^{-1/2}$ , and (E.12) to estimate the regime  $|r_i| \leq \sqrt{\log n}/2$ ,  $|\theta_1 - \theta_2| \gtrsim n^{-1/4}(\gamma_n)^{-1/2}$ , together with (E.14) to estimate the complementary regime  $|r_i| \geq \sqrt{\log n}/2$ . Thus we prove (E.9) and hence finish the proof of Proof of Proposition E.1.  $\square$

As a direct application of Proposition E.1, we obtain the following corollary.

**Corollary E.3.** *Both Theorem 2.2 and Theorem 2.5 hold true for the complex Ginibre ensemble.*

*Proof of Corollary E.3.* We start with proving Theorem 2.2 for the complex Ginibre ensemble. Note that

$$\begin{aligned} \mathbf{P}^{\text{Gin}} \left( \max_{\arg(\sigma_i) \in [a,b]} \{|\sigma_i|\} < 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{t}{\sqrt{4\gamma_n n}} \right) &= \mathbf{E}^{\text{Gin}} \prod_{i=1}^n \left( 1 - \mathbb{1}_{A(t,a,b)}(\sigma_i) \right) = \det(1 - \mathbb{1}_{A(t,a,b)} \tilde{K}_n \mathbb{1}_{A(t,a,b)}) \\ &\xrightarrow{n \rightarrow \infty} \exp\left(-\int_a^b \int_t^\infty \frac{e^{-r}}{2\pi} dr d\theta\right) = e^{-\frac{b-a}{2\pi} e^{-t}}, \end{aligned} \quad (\text{E.16})$$

with  $A(t, a, b)$  given by (E.7), where we also used (E.2) with  $\mathfrak{g}(z) = \mathbb{1}_{A(t,a,b)}(z)$ , together with (E.5) and Proposition E.1. Choose  $[a, b] = [0, 2\pi)$ , then this proves the Gumbel law for the spectral radius  $|\sigma_1|$  in (4.18).

Since the kernel function  $K_n$  in (E.1) is invariant under rotations, *i.e.*,  $K_n(z, w) = K_n(e^{i\theta}z, e^{i\theta}w)$ , clearly  $\arg(\sigma_1)$  is uniformly distributed on  $[0, 2\pi)$ . Moreover, fix any  $t \in \mathbb{R}$  and any integer  $k \geq 1$ , then the following holds uniformly in any  $1 \leq p \leq k$ :

$$\mathbf{P}^{\text{Gin}} \left( |\sigma_1| < 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{t}{\sqrt{4\gamma_n n}}, \frac{2\pi(p-1)}{k} \leq \arg(\sigma_1) < \frac{2\pi p}{k} \right)$$

$$\begin{aligned}
&= \frac{1}{k} \mathbf{P}^{\text{Gin}} \left( |\sigma_1| < 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{t}{\sqrt{4\gamma_n n}} \right) \\
&= \mathbf{P}^{\text{Gin}} \left( \frac{2\pi(p-1)}{k} \leq \arg(\sigma_1) < \frac{2\pi p}{k} \right) \mathbf{P}^{\text{Gin}} \left( |\sigma_1| < 1 + \sqrt{\frac{\gamma_n}{4n}} + \frac{t}{\sqrt{4\gamma_n n}} \right), \quad (\text{E.17})
\end{aligned}$$

where the first equality follows from the fact that the LHS is independent of  $p$  by rotation invariance. In the second equality we also used the uniform distribution of  $\arg(\sigma_1)$ . We can now extend the relation (E.17) involving events  $\arg(\sigma_1) \in [a, b]$  for any interval  $[a, b]$  by approximating it with intervals of rational endpoints. This implies that  $\arg(\sigma_1)$  is independent of  $|\sigma_1|$  and hence proves Theorem 2.2 for the complex Ginibre ensemble.

We continue to prove Theorem 2.5 for the complex Ginibre ensemble. For any function  $g : \mathbb{C} \rightarrow [0, \infty)$  supported on  $A(t, a, b)$ , using (E.2) with  $\mathbf{g}(z) = 1 - e^{-g(z)}$ , (E.5), and Proposition E.1, we have

$$\begin{aligned}
\mathbf{E}^{\text{Gin}} \exp \left( - \sum_{i=1}^n g(\sigma_i) \right) &= \det \left( 1 - \sqrt{1 - e^{-g}} \tilde{K}_n \sqrt{1 - e^{-g}} \right) \\
&\xrightarrow{n \rightarrow \infty} \exp \left( - \int_0^{2\pi} \int_t^\infty (1 - e^{-g(z)}) \frac{e^{-r}}{2\pi} dr d\theta \right), \quad (\text{E.18})
\end{aligned}$$

where we changed the variable  $z$  into  $(r, \theta)$  as in (E.10). We hence finished the proof of Corollary E.3.  $\square$

**Remark E.4.** *Alternatively, the explicit joint distribution of the largest  $k$  Ginibre eigenvalues (in modulus) can be directly computed due to Kostlan's observation [88]. That is, the collection of moduli of the eigenvalues  $\{|\sigma_i|\}_{i=1}^n$  of a complex Ginibre matrix has the same distribution as the collection of independent chi-distributed random variables  $\{\chi_{2k}\}_{k=1}^n$ , i.e.,*

$$\{|\sigma_1|, |\sigma_2|, \dots, |\sigma_n|\} \stackrel{d}{=} \{\chi_2, \chi_4, \dots, \chi_{2n}\}. \quad (\text{E.19})$$

*In other words the distribution of the decreasingly labelled eigenvalues  $|\sigma_1| \geq |\sigma_2| \geq \dots \geq |\sigma_n|$  coincides with the decreasingly ordered statistics of independent chi-distributed variables with even parameters.*

Finally it still remains to prove the key technical input Lemma E.2.

*Proof of Lemma E.2.* Note that the first estimate in (E.11) has already been proven in [59, Lemma E.1], while the last estimate in (E.14) follows directly from [59, Eq. (E.5)] using the Cauchy-Schwarz inequality. In the following we focus on proving (E.12) and (E.13). Recall that

$$\tilde{K}_n(z_1, z_2) = \frac{n}{\pi} e^{-n(|z_1|^2 + |z_2|^2 - 2z_1\bar{z}_2)/2} \frac{\Gamma(n, nz_1\bar{z}_2)}{\Gamma(n)}. \quad (\text{E.20})$$

Note that

$$z_1\bar{z}_2 = e^{i(\theta_1 - \theta_2)} \left( 1 + \sqrt{\frac{\gamma_n}{n}} + \frac{r_1 + r_2}{\sqrt{4\gamma_n n}} + \frac{r_1 + r_2 + \gamma_n}{4n} + \frac{r_1 r_2}{4\gamma_n n} \right), \quad (\text{E.21})$$

and hence for any  $|r_i| \leq \sqrt{\log n}/2$ ,

$$|1 - z_1\bar{z}_2| \gtrsim |\theta_1 - \theta_2| + \sqrt{\gamma_n/n}, \quad |z_i|^2 = 1 + \frac{\gamma_n + r_i}{\sqrt{\gamma_n n}} + O\left(\frac{\gamma_n}{n}\right). \quad (\text{E.22})$$

For  $|\theta_1 - \theta_2| \lesssim n^{-1/4}(\gamma_n)^{-1/2}$ , we use [110, Lemma 3.3] (or [59, Eq. (47)]) i.e.,

$$\frac{\Gamma(n, nz_1\bar{z}_2)}{\Gamma(n)} = \frac{e^{-n(z_1\bar{z}_2 - 1)^2/2}}{\sqrt{2\pi n}(z_1\bar{z}_2 - 1)} \left( 1 + O\left(|z_1\bar{z}_2 - 1| + n|z_1\bar{z}_2 - 1|^3 + \frac{1}{n|z_1\bar{z}_2 - 1|^2}\right) \right). \quad (\text{E.23})$$

Combining with (E.20)-(E.22) and using that  $|1 - z_1\bar{z}_2| \lesssim n^{-1/4}(\gamma_n)^{-1/2}$ , we obtain that

$$|\tilde{K}_n(z_1, z_2)| \lesssim \frac{\sqrt{n}}{|\theta_1 - \theta_2| + \sqrt{\gamma_n/n}} e^{-\frac{1}{2}(r_1 + r_2 + \gamma_n)} \quad (\text{E.24})$$

$$\lesssim \sqrt{n\gamma_n} e^{-\frac{1}{2}(r_1 + r_2)} \left( 1 + O\left(\frac{\log \log n}{\log n}\right) \right), \quad (\text{E.25})$$

where we also used that

$$e^{-\frac{1}{2}\gamma_n} = \exp\left(-\frac{1}{2}\log\frac{n}{(\log n)^{2\pi}}\right) = \frac{(2\pi)^{1/2}\gamma_n}{\sqrt{n}}\left(1 + O\left(\frac{\log\log n}{\log n}\right)\right). \quad (\text{E.26})$$

This proves (E.12).

For  $|\theta_1 - \theta_2| \gtrsim n^{-1/2+w}$ , we use [110, Lemma 3.4] (or [59, Eq. (42)])

$$\frac{\Gamma(n, nz_1\bar{z}_2)}{\Gamma(n)} = e^{n(1-z_1\bar{z}_2)} \frac{(z_1\bar{z}_2)^n}{\sqrt{2\pi n}(1-z_1\bar{z}_2)} \left(1 + O\left(\frac{1}{n|1-z_1\bar{z}_2|^2}\right)\right). \quad (\text{E.27})$$

Combining with (E.20), we obtain that

$$\begin{aligned} |\tilde{K}_n(z_1, z_2)| &= \frac{n e^{-\frac{n}{2}(|z_1|^2 + |z_2|^2 - 2\log|z_1\bar{z}_2|)}}{\pi \sqrt{2\pi n}(1-z_1\bar{z}_2)} \left(1 + O\left(\frac{1}{n|1-z_1\bar{z}_2|^2}\right)\right) \\ &\lesssim \frac{\sqrt{n}}{|\theta_1 - \theta_2| + \sqrt{\gamma_n/n}} e^{-\frac{1}{2}(r_1+r_2+\gamma_n)} \left(1 + O\left(\frac{1}{\sqrt{n}|\theta_1 - \theta_2| + \sqrt{\gamma_n}}\right)^2\right) \\ &\lesssim \frac{\gamma_n}{|\theta_1 - \theta_2|} e^{-\frac{1}{2}(r_1+r_2)} \left(1 + O\left(\frac{\log\log n}{\log n}\right)\right), \end{aligned} \quad (\text{E.28})$$

where we also used (E.21), (E.26) and that for any  $|r_i| \leq \sqrt{\log n}/2$ ,

$$|z_i|^2 - 1 - 2\log|z_i| = 2(|z_i| - 1)^2 + O((|z_i| - 1)^3) = \frac{(\gamma_n + r_i)^2}{2\gamma_n n} + O\left(\frac{\gamma_n^{3/2}}{n^{3/2}}\right).$$

We hence proved (E.13) and completed the proof of Lemma E.2.  $\square$

#### REFERENCES

- [1] Adhikari, A., Dubova, S., Xu, C., Yin, J.: Eigenstate Thermalisation Hypothesis for Generalised Wigner Matrices. Preprint (2023). arXiv:2302.00157
- [2] Adhikari, A., Huang, J.: Dyson Brownian motion for general  $\beta$  and potential at the edge. Probab. Theory Related Fields. **178**, 893–950 (2020)
- [3] Adhikari, A., Landon, B.: Local law and rigidity for unitary Brownian motion. Probab. Theory Related Fields. **187**, 753–815 (2023)
- [4] Aggarwal, A., Bordenave, C., Lopatto, P.: Mobility edge for Lévy matrices. Preprint (2022). arXiv:2210.09458.
- [5] Aggarwal, A., Lopatto P., Marcinek J.: Eigenvector statistics of Lévy matrices. Ann. Probab. **49**(4): 1778–1846 (2021).
- [6] Aggarwal, A., Lopatto, P. Yau, H.T.: GOE statistics for Lévy matrices. J. Eur. Math. Soc., **23**(11):3707–3800 (2021).
- [7] Ajanki O. H., Erdős, L., Krüger, T.: Universality for general Wigner-type matrices. Probab. Theory Related Fields. **169**, 667–727 (2017)
- [8] Ajanki O. H., Erdős, L., Krüger, T.: Stability of the Matrix Dyson Equation and Random Matrices with Correlations. Probab. Theory Related Fields. **173**, 293–373 (2019)
- [9] Akemann, G., Phillips, M.J.: The Interpolating Airy Kernels for the  $\beta = 1$  and  $\beta = 4$  Elliptic Ginibre Ensembles. J. Stat. Phys. **155**, 421–465 (2014)
- [10] Alt, J., Ducatez, R., Knowles, A.: Extremal eigenvalues of critical Erdős–Rényi graphs. Ann. Probab. **49**(3), 1347–1401 (2021)
- [11] Alt, J., Ducatez, R., Knowles, A.: Delocalization transition for critical Erdős–Rényi graphs. Comm. Math. Phys. **388**(1), 507–579 (2021)
- [12] Alt, J., Ducatez, R., Knowles, A.: The completely delocalized region of the Erdős–Rényi graph. Electron. Commun. Probab. **27**(10), 9 pp (2022)
- [13] Alt, J., Ducatez, R., Knowles, A.: Localized phase for the Erdős–Rényi graph graph. Preprint (2023). arXiv:2305.16294
- [14] Alt, J., Erdős, L., Krüger, T.: Local inhomogeneous circular law. Ann. Appl. Probab. **28**(1), 148–203 (2018)
- [15] Alt, J., Erdős, L., Krüger, T.: The Dyson equation with linear self-energy: spectral bands, edges and cusps. Doc. Math. **25**, 1421–1539 (2020)
- [16] Alt, J., Erdős, L., Krüger, T.: Spectral radius of random matrices with independent entries. Probability and Mathematical Physics. **2**(2), 221–280 (2021)
- [17] Arguin, L.P., Belius, D., Bourgade, P.: Maximum of the characteristic polynomial of random unitary matrices. Comm. Math. Phys. **349**, 703–751 (2017)
- [18] Arguin, L.P., Bourgade, P., Radziwiłł, M.: The Fyodorov-Hiary-Keating conjecture I. Preprint (2020). arXiv:2007.00988
- [19] Arguin, L.P., Bourgade, P., Radziwiłł, M.: The Fyodorov-Hiary-Keating Conjecture II. Preprint (2023). arXiv:2307.00982
- [20] Auffinger, A., Ben Arous, G., Péché S.: Poisson convergence for the largest eigenvalues of heavy tailed random matrices. Annales de l’IHP Probabilités et statistiques **45**, 589–610 (2009).
- [21] Bai, Z.D.: Circular law. Ann. Probab. **25**(1), 494–529 (1997)

- [22] Bao, Z.G., Erdős, L.: Delocalization for a class of random block band matrices. *Probab. Theory Related Fields.* **167**(3), 673–776 (2017)
- [23] Bauerschmidt, R., Huang, J., Knowles, A. and Yau, H.T.: Bulk eigenvalue statistics for random regular graphs. *Ann. Probab.* **45**(6A), 3626–3663 (2017)
- [24] Ben Arous, G., Péché, S.: Universality of local eigenvalue statistics for some sample covariance matrices. *Comm. Pure Appl. Math.* **58** 1316–1357 (2005)
- [25] Benaych-Georges, F., Guionnet, A.: Central limit theorem for eigenvectors of heavy tailed matrices. *Electronic Journal of Probability* **19**, 1–27 (2014).
- [26] Bender, M.: Edge scaling limits for a family of non-Hermitian random matrix ensembles. *Probab. Theory Related Fields.* **147**, 241–271 (2010)
- [27] Bordenave, C., Caputo, P., Chafaï, D., Tikhomirov, K.: On the spectral radius of a random matrix: An upper bound without fourth moment. *Ann. Probab.* **46**(4), 2268–2286 (2018)
- [28] Bordenave, C., Chafaï, D.: Around the circular law. *Probab. Surv.* **9**, 1–89 (2012)
- [29] Bordenave C., Guionnet, A.: Localization and delocalization of eigenvectors for heavy-tailed random matrices. *Probability Theory and Related Fields* **157**(3-4), 885–953 (2013).
- [30] Bordenave C., Guionnet, A.: Delocalization at small energy for heavy-tailed random matrices. *Communications in Mathematical Physics* **354**(1), 115–159 (2017).
- [31] Bordenave, C., Chafaï, D. García-Zelada, D.: Convergence of the spectral radius of a random matrix through its characteristic polynomial. *Probab. Theory Related Fields.* **182**, 1163–1181 (2022)
- [32] Borodin, A., Sinclair, C.D.: The Ginibre Ensemble of Real Random Matrices and its Scaling Limits. *Commun. Math. Phys.* **291**, 177–224 (2009)
- [33] Bourgade, P., Erdős, L., Yau, H.T., Yin, J.: Universality for a class of random band matrices. *Adv. Theor. Math. Phys.* **21**, 739–800 (2017)
- [34] Bourgade, P.: Extreme gaps between eigenvalues of Wigner matrices. *J. Eur. Math. Soc. (JEMS)* **24**, 2823–2873 (2022)
- [35] Bourgade, P., Falconet, H.: Liouville quantum gravity from random matrix dynamics. Preprint (2022). arXiv:2206.03029
- [36] Bourgade, P., Yau, H.-T., Yang, F., J. Yin.: Random band matrices in the delocalized phase, II: Generalized resolvent estimates. *J. Stat. Phys.* **174**, 1–33 (2019)
- [37] Bourgade, P., Yau, H.-T., Yin, J.: Local circular law for random matrices. *Probab. Theory Related Fields* **159**, 545–595 (2014)
- [38] Bourgade, P., Yau, H.-T., Yin, J.: The local circular law II: the edge case. *Probab. Theory Related Fields*, **159**, 619–660 (2014)
- [39] Bourgade, P., Yau, H.-T., Yin, J.: Random band matrices in the delocalized phase I: Quantum unique ergodicity and universality. *Comm. Pure Appl. Math.* **73**, 1526–1596 (2020)
- [40] Byun, S. S., Forrester, P. J.: Progress on the study of the Ginibre ensembles I: GinUE. Preprint (2022). arXiv:2211.16223
- [41] Byun, S. S., Forrester, P. J.: Progress on the study of the Ginibre ensembles II: GinOE and GinSE. Preprint (2023). arXiv:2301.05022
- [42] Chafaï, D.: Around the circular law: an update. (Webblog)  
<https://djalil.chafai.net/blog/2018/11/04/around-the-circular-law-an-update>. Version 2018-11-04
- [43] Chafaï, D., Péché, S.: A note on the second order universality at the edge of Coulomb gases on the plane. *J. Stat. Phys.* **156**, 368–383 (2014)
- [44] Chhaibi, R., Madaule, T., Najnudel, J.: On the maximum of the  $C\beta E$  field. *Duke Math.J.* **167**, 2243–2345 (2018)
- [45] Chen, N., Smart, C.K.: Random band matrix localization by scalar fluctuations. Preprint (2022). arXiv:2206.06439.
- [46] Cipolloni, G., Peled, R., Schenker, J., Shapiro, J.: Dynamical localization for random band matrices up to  $W \ll N^{1/4}$ . Preprint (2022). arXiv:2206.05545.
- [47] Cipolloni, G., Erdős, L., Henheik, J.: Eigenstate thermalisation at the edge for Wigner matrices. Preprint (2023). arXiv:2309.05488
- [48] Cipolloni, G., Erdős, L., Henheik, J., Kolupaiev, O.: Gaussian fluctuations in the Equipartition Principle for Wigner matrices. *Forum Math. Sigma.* **11**, (2024)
- [49] Cipolloni, G., Erdős, L., Henheik, J., Schröder, D.: Optimal Lower Bound on Eigenvector Overlaps for non-Hermitian Random Matrices. Preprint (2023). arXiv:2301.03549
- [50] Cipolloni, G., Erdős, L., Krüger, T., Schröder, D.: Cusp universality for random matrices II: the real symmetric case. *Pure Appl. Anal.* **1**(4), 615–707 (2019)
- [51] Cipolloni, G., Erdős, L., Schröder, D.: Edge universality for non-Hermitian random matrices. *Probability Theory and Related Fields*, **179**(1), 1–28 (2021)
- [52] Cipolloni, G., Erdős, L., Schröder, D.: Central limit theorem for linear eigenvalue statistics of non-Hermitian random matrices. *Comm. Pure Appl. Math.* **76**, 946–1034 (2023)
- [53] Cipolloni, G., Erdős, L., Schröder, D.: Fluctuation around the circular law for random matrices with real entries. *Electron. J. Probab.* **26**, 1-61 (2021)
- [54] Cipolloni, G., Erdős, L., Schröder, D.: Eigenstate thermalization hypothesis for Wigner matrices. *Commun. Math. Phys.* **388**, 1005–1048 (2021)
- [55] Cipolloni, G., Erdős, L., Schröder, D.: Functional Central Limit Theorems for Wigner Matrices. *Ann. Appl. Probab.* **33**(1), 447–489 (2023)
- [56] Cipolloni, G., Erdős, L., Schröder, D.: Optimal multi-resolvent local laws for Wigner matrices. *Electron. J. Probab.* **27**, 1–38 (2022)



- [57] Cipolloni, G., Erdős, L., Schröder, D.: Rank-uniform local law for Wigner matrices. *Forum Math. Sigma*. **10** (2022)
- [58] Cipolloni, G., Erdős, L., Schröder, D.: Mesoscopic Central Limit Theorem for non-Hermitian Random Matrices. *Probab. Theory Relat. Fields* (2023)
- [59] Cipolloni, G., Erdős, L., Schröder, D., Xu, Y.: Directional extremal statistics for Ginibre eigenvalues. *J. Math. Phys.* **63**(11), 103303, (2022)
- [60] Cipolloni, G., Erdős, L., Schröder, D., Xu, Y.: On the rightmost eigenvalue of non-Hermitian random matrices. *Ann. Probab.* **51**(6), 2192–2242 (2023)
- [61] Cipolloni, G., Erdős, L., Xu, Y.: Precise asymptotics for the spectral radius of a large random matrix. Preprint (2022). arXiv:2210.15643
- [62] Erdős, L., Ji, H. C.: Density of Brown measure of free circular Brownian motion. Preprint (2023). arXiv:2307.08626
- [63] Erdős, L., Knowles, A., Yau, H.-T., Yin, J.: Delocalization and diffusion profile for random band matrices. *Comm. Math. Phys.* **323**(1), 367–416 (2013)
- [64] Erdős, L., Krüger, T., Schröder, D.: Cusp universality for random matrices I: Local law and the complex Hermitian case. Preprint (2018). arXiv:1809.03971
- [65] Erdős, L., Yau, H.-T.: Universality of local spectral statistics of random matrices. *Bull. Amer. Math. Soc.* **49**(3), 377–414 (2012)
- [66] Erdős, L., Yau, H.-T.: A dynamical approach to random matrix theory. *Courant Lecture Notes* **28**, American Mathematical Soc. (2017)
- [67] Erdős, L., Yau, H. T., Yin, J.: Rigidity of eigenvalues of generalized Wigner matrices. *Adv. Math.* **229**(3), 1435–1515 (2012)
- [68] Erdős, L., Xu, Y.: Small deviation estimates for the largest eigenvalue of Wigner matrices. *Bernoulli* **29**(2), 1063–1079 (2023)
- [69] Fisher, R.A., Tippett, L.H.C.: Limiting forms of the frequency distribution of the largest and smallest member of a sample. *Proc. Camb. Phil. Soc.* **24**(2), 180–190 (1928)
- [70] François, Q., García-Zelada, D.: Asymptotic analysis of the characteristic polynomial for the Elliptic Ginibre Ensemble. Preprint (2023). arXiv:2306.16720
- [71] Fyodorov, Y.V., Hiary, G. A., Keating, J. P.: Freezing transition, characteristic polynomials of random matrices, and the Riemann zeta function. *Phys. Rev. Lett.* **108**, 170601 (2012)
- [72] Ginibre, J.: Statistical Ensembles of Complex, Quaternion, and Real Matrices. *J. Math. Phys.* **6**, 440–449 (1965)
- [73] Girko, V. L.: Circular law. *Teor. Veroyatnost. i Primenen.* **29**, 669–679 (1984)
- [74] Gnedenko, B.V.: Sur la distribution limite du terme maximum d’une série aléatoire. *Ann. Math.* **44**(3), 423–453 (1943)
- [75] Gohberg, I., Goldberg, S. and Krupnik, N.: *Traces and Determinants of Linear Operators. Operator Theory: Advances and Applications* **116**, Birkhäuser Basel, IX, 258 (2000)
- [76] Goldstein, M.: Fluctuations and localization length for random band GOE matrix. Preprint (2022). arXiv:2210.04346.
- [77] Götze, F., Tikhomirov, A.: The circular law for random matrices, *Ann. Probab.* **38**, 1444–1491 (2010)
- [78] He, Y.: Edge universality of sparse Erdős-Rényi digraphs. Preprint (2023). arXiv:2304.04723
- [79] He, Y., Knowles, A.: Mesoscopic eigenvalue statistics of Wigner matrices. *Ann. Appl. Probab.* **27** 1510–1550 (2017)
- [80] Hiesmayr, E. and McKenzie, T.: The Spectral Edge of Constant Degree Erdős-Rényi Graphs, Preprint (2023). arXiv:2309.11007
- [81] Huang, J., Landon, B.: Rigidity and a mesoscopic central limit theorem for Dyson Brownian motion for general  $\beta$  and potentials. *Probab. Theory Related Fields* **175**, 209–253 (2019)
- [82] Huang, J., Landon, B., Yau, H.-T.: Bulk universality of sparse random matrices. *J. Math. Phys.* **56**, 123301 (2015)
- [83] Huang, J., Yau, H.-T.: Edge Universality of Sparse Random Matrices. Preprint (2022). arXiv:2206.06580
- [84] Huang, J., Yau, H.-T.: Edge Universality of Random Regular Graphs of Growing Degrees. Preprint (2023). arXiv:2305.01428
- [85] Johansson, K.: On fluctuations of eigenvalues of random Hermitian matrices. *Duke Math. J.* **91**(1), 151–204 (1998)
- [86] Khorunzhy, A.M., Khoruzhenko, B.A., Pastur, L.A.: Asymptotic properties of large random matrices with independent entries, *J. Math. Phys.* **37**, 5033–5060 (1996)
- [87] Kopel, P.: Linear statistics of non-Hermitian matrices matching the real or complex Ginibre ensemble to four moments. Preprint (2015). arXiv:1510.02987
- [88] Kostlan, E.: On the spectra of Gaussian matrices. *Linear Algebra Appl.* **162/164**, Directions in matrix theory (Auburn, AL, 1990), 385–388 (1992)
- [89] Landon, B., Lopatto, P., Sosoe, P.: Single eigenvalue fluctuations of general Wigner-type matrices. *Probab. Theory Relat. Fields* (2023).
- [90] Landon, B., Sosoe, P.: Applications of mesoscopic CLTs in random matrix theory. *Ann. Appl. Probab.* **30**(6), 2769–2795 (2020)
- [91] Landon, B., Sosoe, P.: Almost-optimal bulk regularity conditions in the clt for Wigner matrices. Preprint (2022). arXiv:2204.03419
- [92] Lee, J.-O., Schnelli, K.: Edge universality for deformed Wigner matrices. *Rev. Math. Phys.* **27**(08), 1550018 (2015)
- [93] Li, Y., Schnelli, K., Xu, Y.: Central limit theorem for mesoscopic eigenvalue statistics of deformed Wigner matrices and sample covariance matrices. *Ann. Inst. Henri Poincaré Probab. Stat.* **57**, 506–546 (2021)
- [94] Liu, D.-Z., Zhang, L.: Critical edge statistics for deformed GinUEs. Preprint (2023). arXiv:2311.13227
- [95] Lytova, A., Pastur, L.: Central limit theorem for linear eigenvalue statistics of random matrices with independent entries. *Ann. Probab.* **37**, 1778–1840 (2009)
- [96] Maltsev, A., Osman, M.: Bulk universality for complex non-Hermitian matrices with independent and identically distributed entries. Preprint (2023). arXiv:2310.11429

- [97] May, R. M.: Will a large complex system be stable? *Nature* **238**, 413–414 (1972)
- [98] Mehta, M.: *Random Matrices*. Pure and Applied Mathematics **142**, Third Edition, Academic Press (2004)
- [99] Narayanan, H., Sheffield, S., Tao, T.: Sums of GUE matrices and concentration of hives from correlation decay of eigengaps. Preprint (2023). arXiv:2306.11514
- [100] Pacco, A., Ros, V.: Overlaps between eigenvectors of spiked, correlated random matrices: From matrix principal component analysis to random Gaussian landscapes. *Phys. Rev. E* **108**, 024145 (2023)
- [101] Pan, G., Zhou, W.: Circular law, extreme singular values and potential theory. *J. Multivar. Anal.* **101**(3), 645–656 (2010)
- [102] Shorack, G. R., Wellner, J. A.: *Empirical processes with applications to statistics*. Society for Industrial and Applied Mathematics (2009)
- [103] Paquette, E. and Zeitouni, O.: The maximum of the CUE field. *Int. Math. Res. Not. IMRN* **16**, 5028–5119 (2018)
- [104] Paquette, E. and Zeitouni, O.: The extremal landscape for the  $C\beta E$  ensemble. Preprint (2022). arXiv:2209.06743
- [105] Pastur, L. A.: On the spectrum of random matrices. *Teoreticheskaya i Matematicheskaya Fizika* **10.1**: 102–112 (1972).
- [106] Riabov, V.: Mesoscopic eigenvalue statistics for Wigner-type matrices. Preprint (2023). arXiv:2301.01712
- [107] Riabov, V.: Linear eigenvalue statistics at the cusp. Preprint (2023). arXiv:2307.07432
- [108] Rider, B.: A limit theorem at the edge of a non-Hermitian random matrix ensemble. *J. Phys. A* **36**(12), 3401–3409 (2003)
- [109] Schenker, J.: Eigenvector localization for random band matrices with power law band width. *Comm. Math. Phys.*, **290**:1065–1097 (2009).
- [110] Rider, B., Sinclair, C. D.: Extremal laws for the real Ginibre ensemble. *Ann. Appl. Probab.* **24**(4), 1621–1651 (2014)
- [111] Schnell, K., Xu, Y.: Convergence rate to the Tracy–Widom laws for the largest eigenvalue of Wigner matrices. *Comm. Math. Phys.* **393**, 839–907 (2022)
- [112] Shcherbina, M.: Central limit theorem for linear eigenvalue statistics of the Wigner and sample covariance random matrices. *Zh. Mat. Fiz. Anal. Geom.* **7**, 176–192 (2011)
- [113] Shcherbina, M., Shcherbina, T.: Characteristic polynomials for 1D random band matrices from the localization side. *Comm. Math. Phys.* **351** (2017)
- [114] Shcherbina, T.: On the Second Mixed Moment of the Characteristic Polynomials of 1D Band Matrices, *Comm. Math. Phys.* **328**, 45–82 (2014)
- [115] Sodin, S.: The spectral edge of some random band matrices. *Ann. Math.* **173**(3), 2223–2251 (2010)
- [116] Soshnikov, A.: Universality at the Edge of the Spectrum in Wigner Random Matrices. *Commun. Math. Phys.* **207**, 697–733 (1999)
- [117] Sosoe, P., Wong, P.: Regularity conditions in the CLT for linear eigenvalue statistics of Wigner matrices. *Advances in Mathematics* **249**, 37–87 (2013)
- [118] Stone, B., Yang, F., Yin, J.: A random matrix model towards the quantum chaos transition conjecture. Preprint (2023). arXiv:2312.07297
- [119] Tao, T., Vu, V.: Random matrices: universality of ESDs and the circular law. *Ann. Probab.* **38**(5), 2023–2065 (2010)
- [120] Tao, T., Vu, V.: Random matrices: Universality of local eigenvalue statistic. *Acta Math.* **206**, 127–204 (2011)
- [121] Tao, T., Vu, V.: Random matrices: universality of local spectral statistics of non-Hermitian matrices. *Ann. Probab.* **43**(2), 782–874 (2015)
- [122] Tracy, C., Widom, H.: Level-Spacing Distributions and the Airy Kernel. *Commun. Math. Phys.* **159**, 151–174 (1994)
- [123] Tracy, C., Widom, H.: On Orthogonal and Symplectic Matrix Ensembles, *Commun. Math. Phys.* **177**, 727–754 (1996)
- [124] von Soosten, P., Warzel, S.: Random characteristics for Wigner matrices. *Electron. Commun. Probab.* **24**: 1–12 (2019).
- [125] Wigner, E. P.: Characteristic vectors of bordered matrices with infinite dimensions. *Ann. Math.* **62**(3), 548–564 (1955)
- [126] Xu, C., Yang, F., Yau, H.T. and Yin, J.: Bulk universality and quantum unique ergodicity for random band matrices in high dimensions. Preprint (2022). arXiv:2207.14533
- [127] Yang, F., Yau, H.T., Yin, J.: Delocalization and quantum diffusion of random band matrices in high dimensions I: Self-energy renormalization. Preprint (2021). arXiv:2104.12048
- [128] Yang, F., Yau, H.T., Yin, J.: Delocalization and Quantum Diffusion of Random Band Matrices in High Dimensions II: T-expansion. *Commun. Math. Phys.* **396**, 527–622 (2022)
- [129] Yang, F., Yin, J.: Random band matrices in the delocalized phase, III: averaging fluctuations, *Probab. Theory Related Fields* **179**, 451–540 (2021)