

RELATING THE ROE ALGEBRA OF A SPACE TO THE UNIFORM ROE ALGEBRAS OF ITS DISCRETIZATIONS

V. MANUILOV

ABSTRACT. The Roe algebra $C^*(X)$ is a non-commutative C^* -algebra reflecting metric properties of a space X , and it is interesting to understand relation between the Roe algebra of X and the uniform Roe algebra of its discretizations. Here we construct, for a simplicial space X , a continuous field of C^* -algebras over $\mathbb{N} \cup \{\infty\}$ with the fibers over finite points the uniform C^* -algebras of discretizations of X , and the fiber over ∞ the Roe algebra of X . We also construct the direct limit of the uniform Roe algebras of discretizations and its embedding into the Roe algebra of X .

1. INTRODUCTION

Roe algebras play an increasingly important role in the index theory of elliptic operators on noncompact manifolds and their generalizations [10, 6, 8]. Following the ideology of noncommutative geometry [2], they provide an interplay between metric spaces (e.g. manifolds) and (noncommutative) C^* -algebras.

Recall the definition of the Roe algebra of a metric space X [10]. Let H_X be a Hilbert space with an action of the algebra $C_0(X)$ of continuous functions on X vanishing at infinity (i.e. a $*$ -homomorphism $\pi : C_0(X) \rightarrow \mathbb{B}(H_X)$). For a Hilbert space H we write $\mathbb{B}(H)$ (resp., $\mathbb{K}(H)$) for the algebra of all bounded (resp., all compact) operators on H . We will assume that

$$\{\pi(f)\xi : f \in C_0(X), \xi \in H_X\} \text{ is dense in } H_X \quad (1)$$

and that

$$\pi(f) \in \mathbb{K}(H_X) \text{ implies that } f = 0. \quad (2)$$

An operator $T \in \mathbb{B}(H_X)$ is *locally compact* if the operators $T\pi(f)$ and $\pi(f)T$ are compact for any $f \in C_0(X)$. It has *finite propagation* if there exists some $R > 0$ such that $\pi(f)T\pi(g) = 0$ whenever the distance between the supports of $f, g \in C_0(X)$ is greater than R . The *Roe algebra* $C^*(X, H_X)$ is the norm completion of the $*$ -algebra of locally compact, finite propagation operators on H_X . As it does not depend on the choice of H_X satisfying (1) and (2), it is usually denoted by $C^*(X)$. It is convenient to have a Borel measure on X . When the measure μ on X has no atoms, the simplest choice $H_X = L^2(X)$ with the standard action of $C_0(X)$ satisfies the properties (1) and (2), so it suffices to use $L^2(X)$ to define the Roe algebra.

When X is discrete (with an atomic measure), the choice $H_X = l^2(X)$ does not satisfy the condition (2). In order to fix this, one may take $H_X = l^2(X) \otimes H$ with an infinite-dimensional Hilbert space H to obtain the Roe algebra. But there is also another option: still to use $H_X = l^2(X)$. The resulting C^* -algebra is called the *uniform Roe algebra* of X , and is denoted by $C_u^*(X)$. This C^* -algebra differs from the Roe algebra $C^*(X)$. It is more tractable, but has less relations with elliptic theory.

Manifolds and some other spaces X are often endowed with discrete subspaces $D \subset X$ that are ε -dense for some ε , e.g. lattices in Lie groups, or, more generally, Delone sets in metric spaces [1]. Some problems related to X may become simpler when reduced to such discretization D . In particular, it would be interesting to understand relation

between the Roe algebra of X and the uniform Roe algebras of its discretizations D . The latter are embedded into $C^*(X)$, and in [7] we have shown that a large class of non-trivial projections in $C_u^*(D)$ vanish in the K -theory of $C^*(X)$. Delone sets on general metric spaces are too vague to establish such relation, so we restrict ourselves to the case when X is a simplicial complex. The set of vertices is a discretization of X , and in order to get more dense discretizations we can divide the simplices of X into smaller simplices.

In this paper we construct a continuous field of C^* -algebras over $\mathbb{N} \cup \{\infty\}$ such that the fiber over ∞ is the Roe algebra of X , while the fiber over any finite point is the uniform Roe algebra of a certain discretization $D(n)$, $n \in \mathbb{N}$, of X . Such non locally trivial continuous fields of C^* -algebras are interesting because they provide relations between fibers over different points. In particular, they may provide a map from the K -theory group of the fiber over ∞ to the K -theory group of the fibers over finite points (cf. [4]). In [9] we have implemented a similar construction of a continuous field in the case $X = \mathbb{R}$ using the group structure on \mathbb{R} .

We also construct the direct limit of the uniform Roe algebras of discretizations $D(n)$ of X that become more and more dense, and an embedding of this direct limit into $C^*(X)$. Such direct limits are good candidates for what could be the *uniform* Roe algebra of a non-discrete space.

Let X be a countable simplicial complex consisting of simplices Δ_α^k of dimension k . We assume that X is uniformly locally bounded, i.e. that each simplex has a uniformly bounded number of adjacent simplices, and that X is finitedimensional, i.e. that all dimensions of simplices are uniformly bounded. We say that a simplex has maximal dimension if it is not a subset of another simplex (X may have simplices of different maximal dimensions). We endow X with a metric d (resp., a Borel measure μ) such that the restriction of d (resp., of μ) onto each simplex Δ_α^k (resp. onto each simplex of maximal dimension) coincides with the standard metric (resp., standard measure) on the standard simplex

$$\Delta = \Delta(1) = \{(x_0, x_1, \dots, x_k) \in \mathbb{R}^{k+1} : \sum_{j=0}^k x_j = 1; x_j \geq 0, j = 0, 1, \dots, k\}. \quad (3)$$

If $x, y \in X$ do not lie in one simplex then $d(x, y)$ is defined as the infimum of lengths of paths connecting x and y . After fixing the isometric homeomorphisms of the simplices of X with the standard simplex we shall use the coordinates on the standard simplex as local coordinates on X .

2. DISCRETIZATIONS OF X AND PARTITIONS OF UNITY

For the given simplicial structure of X , let $D = D(1)$ denote the discrete metric space consisting of all the vertices Δ_α^0 , $\alpha \in A$. It is a 1-net for X . The assumption that X is finitedimensional and uniformly locally bounded implies that $D(1)$ has bounded geometry, i.e. the number of points of any ball of radius R in $D(1)$ is bounded by some number $b(R)$ which depends only on R .

Consider a subdivision of the simplicial structure $\{\Delta_\alpha^k\}_{\alpha \in A} = \{\Delta_{\alpha_1}^k(1)\}_{\alpha_1 \in A_1}$ of X into smaller simplices $\Delta_{\alpha_2}^k(2)$, $\alpha_2 \in A_2$, such that each simplex $\Delta_{\alpha_1}^k$ is the union of 2^k simplices $\Delta_{\alpha_2}^k(2)$, $\alpha_2 \in A_2$, isometric to the standard simplex

$$\Delta(2) = \{(x_0, x_1, \dots, x_k) \in \mathbb{R}^{k+1} : \sum_{j=0}^k x_j = \frac{1}{2}; x_j \geq 0, j = 0, 1, \dots, k\}.$$

Let $D(2)$ be the metric space of all vertices of the new simplicial structure $\{\Delta_{\alpha_2}^k(2)\}_{\alpha_2 \in A_2}$.

Proceeding, we obtain simplicial structures $\{\Delta_{\alpha_n}^k(n)\}_{\alpha_n \in A_n}$ such that each $\Delta_{\alpha_n}^k(n)$ is isometric to the standard simplex

$$\Delta(n) = \{(x_0, x_1, \dots, x_k) \in \mathbb{R}^{k+1} : \sum_{j=0}^k x_j = \frac{1}{2^n}; x_j \geq 0, j = 0, 1, \dots, k\}.$$

We also obtain a sequence of discretizations $D(1) \subset D(2) \subset \dots \subset X$ such that $D(n)$ is an $\frac{1}{2^n}$ -net in X . The metric spaces $D(n)$ are also of bounded geometry for each $n \in \mathbb{N}$.

For each vertex $u = \Delta_{\alpha_n}^0(n) \in D(n)$ let $\text{St}_u \subset X$ denote the open star of u . Then $\{\text{St}_u : u \in D(n)\}$ is an open cover of X . Let $\{\varphi_u^n : u \in D(n)\}$ be the piecewise linear partition of unity for this cover. If $u \in \Delta_{\alpha_n}^k(n) \subset D(n)$ has coordinates $(\frac{1}{2^n}, 0, \dots, 0)$ in the standard simplex then $\varphi_u^n(x) = x_0$ for any $x \in \Delta(n)$.

Let $G = (g_{uv})_{u,v \in D(n)}$, where $g_{uv} = \langle \varphi_u^n, \varphi_v^n \rangle$, be the Gram matrix for the set of functions $\{\varphi_u^n\}_{u \in D(n)}$. With an abuse of notation we shall use the same G to denote the operator on the Hilbert space $l^2(D(n))$ with the matrix G with respect to the standard basis of $l^2(D(n))$, which consists of the functions $\delta_u, u \in D(n)$, given by $\delta_u(v) = \begin{cases} 1, & \text{if } v = u; \\ 0, & \text{if } v \neq u. \end{cases}$

Clearly, the operator G is positive.

Theorem 1. *There exists $C > 0$ such that the operator G on $l^2(D(n))$ is bounded from below by $\frac{C}{2^n}$.*

Proof. It suffices to prove the claim for $n = 1$, as the general case can be obtained by scaling. We write Δ for $\Delta(1)$ and D for $D(1)$ to simplify the notation.

Let $\{\Delta_\beta\}$ be the set of simplices of maximal dimension. Note that

$$g_{uv} = \langle \varphi_u, \varphi_v \rangle = \int_X \bar{\varphi}_u(x) \varphi_v(x) d\mu(x) = \sum_\beta \int_{\Delta_\beta} \bar{\varphi}_u(x) \varphi_v(x) d\mu(x) = \sum_\beta g_{uv}^\beta,$$

$u, v \in D$. Let G^β be the operator on $l^2(D)$ with the matrix $(g_{uv}^\beta)_{u,v \in D}$ with respect to the standard basis. Then $G = \sum_\beta G^\beta$, where the sum is $*$ -strongly convergent due to finite propagation of G . For Δ_β let u_0, u_1, \dots, u_i be the set of its vertices, and let q_β denote the projection in $l^2(D)$ onto the linear span of $\delta_j, j = 0, 1, \dots, i$.

We claim that there exists $C > 0$ such that $G^\beta \geq Cq_\beta$. Assuming that this claim holds, it remains to note that each vertex lies in at least one simplex of maximal dimension, hence $\sum_\beta q_\beta \geq 1$, from which it follows that $G = \sum_\beta G^\beta \geq C \sum_\beta q_\beta \geq C \cdot 1$.

Fix a simplex Δ_β^k of maximal dimension k , with vertices u_0, u_1, \dots, u_k . After identifying the simplex with the standard simplex (3), these vertices have coordinates $(1, 0, \dots, 0), \dots, (0, 0, \dots, 1)$, and the functions φ_{u_i} are identified with the functions x_i on the standard simplex. Therefore,

$$g_{u_i u_j}^\beta = \int_{\Delta_\beta^k} x_i x_j d\mu(x).$$

By the definition of the projection q_β , the finite rank operator $q_\beta G^\beta q_\beta$ has $(g_{u_i u_j}^\beta)_{i,j=0}^k$ as its matrix.

It remains to show that the positive operator $q_\beta G^\beta q_\beta$ is separated from zero. Fix a k -dimensional cube K with the sides of length a , centered at the center $\frac{1}{k+1}(1, 1, \dots, 1)$ of the simplex Δ^k . If a is sufficiently small, $K \subset \Delta^k$. Then

$$g_{u_i u_j}^\beta = \int_K x_i x_j d\mu(x) + \int_{\Delta^k \setminus K} x_i x_j d\mu(x) = g'_{u_i u_j} + g''_{u_i u_j},$$

and both matrices, $(g'_{u_i u_j})_{i,j=0}^k$ and $(g''_{u_i u_j})_{i,j=0}^k$ are positive, so it would suffice to prove that the first one is separated from zero.

Let us calculate $g'_{u_i u_j} = \int_K x_i x_j d\mu(x)$. Let U be an orthogonal operator on \mathbb{R}^{k+1} that moves the standard simplex to a subset of the hyperplane $x_0 = \frac{1}{\sqrt{k+1}}$ (and the center of this simplex to the point $(\frac{1}{\sqrt{k+1}}, 0, \dots, 0)$). Changing variables $Ux = y$, we get

$$(g'_{u_i u_j})_{i,j=0}^k = \int_K (x_i x_j)_{i,j=0}^k d\mu(x) = U \left(\int_{U(K)} (y_p y_q)_{p,q=0}^k d\mu(y) \right) U^*,$$

so it remains to separate the matrix

$$\int_{U(K)} (y_p y_q)_{p,q=0}^k d\mu(y) \tag{4}$$

from zero.

If $p \neq q$ then

$$\int_{U(K)} y_p y_q d\mu(y) = \int_{-a}^a \cdots \int_{-a}^a y_p y_q dy_1 \cdots dy_k = 0,$$

so the matrix (4) is diagonal, and it remains to calculate

$$\int_{U(K)} y_p^2 d\mu(y) = \int_{-a}^a \cdots \int_{-a}^a y_p^2 dy_1 \cdots dy_k,$$

which, for $p = 0$, equals $(2a)^k \frac{1}{k+1}$, while for $p > 0$, it equals $(2a)^{k-1} \frac{2a^3}{3}$. If C is smaller than both these two numbers then $\int_{U(K)} (y_p y_q)_{p,q=0}^k d\mu(y) \geq CI$, where I denotes the unit matrix. \square

Corollary 2. *The operator G is bounded, invertible and has finite propagation.*

Proof. Invertibility follows from Theorem 1. Finite propagation follows from the fact that if $u, v \in D(n)$ and if $\text{St}_u \cap \text{St}_v = \emptyset$ then $\langle \varphi_u, \varphi_v \rangle = 0$, so uniform local finiteness of X implies that G has finite propagation. Finally, as all matrix entries are uniformly bounded, an operator of finite propagation is bounded. \square

3. TWO MAPS

In this section we construct the maps $\alpha_n : C_u^*(D(n)) \rightarrow C^*(X)$ and $\beta_n : C^*(X) \rightarrow C_u^*(D(n))$, $n \in \mathbb{N}$. Let p_n denote the projection, in $L^2(X)$, onto the closure H_n of the linear span of φ_x^n , $x \in D(n)$. The construction is the same for each n , so we shall skip n from the notation and write D for $D(n)$, H for H_n , and φ_u for φ_u^n .

Set $C = G^{-1/2}$, $C = (c_{uv})_{u,v \in D}$, where G is the operator determined by the Gram matrix for $\{\varphi_u\}$, $u \in D$. By functional calculus, C can be approximated by polynomials in G , hence C lies in the norm closure of operators of finite propagation, i.e. $C \in C_u^*(D)$.

Set $\psi_v = \sum_{u \in D} c_{uv} \varphi_u$. Then

$$\begin{aligned} \langle \psi_u, \psi_v \rangle &= \left\langle \sum_{x \in D} c_{xu} \varphi_x, \sum_{y \in D} c_{yv} \varphi_y \right\rangle = \sum_{x,y \in D} \bar{c}_{xu} c_{yv} \langle \varphi_x, \varphi_y \rangle \\ &= \sum_{x,y \in D} \bar{c}_{xu} c_{yv} g_{xy} = (CGC)_{uv} = \delta_{uv}, \end{aligned}$$

hence $\{\psi_u\}_{u \in D}$ is an orthonormal system. Invertibility of C implies that the closures of the linear spans of $\{\varphi_u\}_{u \in D}$ and of $\{\psi_u\}_{u \in D}$ coincide. The advantage of this orthonormal system with respect to the system obtained from $\{\varphi_u\}_{u \in D}$ by the Gram–Schmidt orthogonalization is that it is obtained from the original non-orthogonal system by using an operator from $C_u^*(\mathbb{Z})$.

Consider one more basis for H . By construction of C , for any $\varepsilon > 0$ there exists an operator $C_\varepsilon \in \mathbb{B}(l^2(D))$ of finite propagation M_ε , with the matrix $C_\varepsilon = (c_{uv}^\varepsilon)$, such that

$\|C - C_\varepsilon\| < \varepsilon$. Set $\psi_v^\varepsilon = \sum_{u \in D} c_{uv}^\varepsilon \varphi_u$. The basis $\{\psi_v^\varepsilon\}_{v \in D}$ is only almost orthonormal, but, unlike $\{\psi_v\}_{v \in D}$, it is compactly localized in D .

Let $U : l^2(D) \rightarrow L^2(X)$ be the isometry determined by $U(\delta_u) = \psi_u$, $u \in D$. Then the range of U is $H \subset L^2(X)$. Define a map $\alpha : C_u^*(D) \rightarrow C^*(X)$ by $\alpha(T)(\xi) = UTU^*(p\xi)$, where $\xi \in L^2(X)$. Note that if $T = (t_{uv})_{u,v \in D}$ is the matrix of $T \in \mathbb{B}(l^2(D))$ then

$$\alpha(T)(\xi) = \sum_{u,v \in D} t_{uv} \psi_u \langle \psi_v, \xi \rangle, \quad \xi \in L^2(X).$$

Clearly, α is a $*$ -homomorphism, in particular, it is isometric. As T is bounded, $\alpha(T)$ is bounded as well.

Let also $U_\varepsilon : l^2(D) \rightarrow L^2(X)$ be an operator given by $U_\varepsilon \delta_u = \psi_u^\varepsilon$, $u \in D$.

For a matrix $A = (a_{uv})_{u,v \in D}$ define an operator $\gamma(A)$ on H by $\gamma(A)\varphi_v = \sum_{u \in D} a_{uv} \varphi_u$. Then one can write $\alpha(T)$ as $\alpha(T) = \gamma(C^{-1}TC)$.

Lemma 3. *The map γ is bounded.*

Proof. Let $f = \sum_{u \in D} f_u \varphi_u \in H$. Then, by Corollary 2, the norm $\|f\|_{L^2(X)}^2 = \sum_{u,v \in D} g_{uv} \bar{f}_u f_v$ is equivalent to the norm $\|f\|_2^2 = \sum_{u \in D} |f_u|^2$. Therefore, there exists $K > 0$ such that

$$\|\gamma(A)f\|_{L^2(X)} \leq K \|\gamma(A)f\|_2 = K \|A\xi\|_2 \leq K \|A\| \|\xi\|_2 \leq K^2 \|A\| \|f\|_{L^2(X)},$$

where $\xi = \sum_{u \in D} f_u \delta_u \in l^2(D)$. □

Set $\alpha_\varepsilon(T) = \gamma(C_\varepsilon^{-1}TC_\varepsilon)$.

Lemma 4. *For sufficiently small ε there exists $M > 0$ such that $\|\alpha(T) - \alpha_\varepsilon(T)\| < M\varepsilon$.*

Proof. One should take ε small enough to provide invertibility of $\gamma(C_\varepsilon)$. Then

$$\begin{aligned} \|\alpha(T) - \alpha_\varepsilon(T)\| &\leq \|\gamma(C^{-1}) - \gamma(C_\varepsilon^{-1})\| \cdot \|\gamma(T)\| \cdot \|\gamma(C)\| \\ &\quad + \|\gamma(C_\varepsilon^{-1})\| \cdot \|\gamma(T)\| \cdot \|\gamma(C) - \gamma(C_\varepsilon)\| \\ &\leq \|\gamma(C - C_\varepsilon)\| \cdot \|\gamma(C^{-1})\| \cdot \|\gamma(C_\varepsilon^{-1})\| \cdot \|\gamma(T)\| \cdot \|\gamma(C)\| \\ &\quad + \|\gamma(C_\varepsilon^{-1})\| \cdot \|\gamma(T)\| \cdot \|\gamma(C - C_\varepsilon)\| \\ &< K^2 \varepsilon (\|\gamma(C^{-1})\| \cdot \|\gamma(C_\varepsilon^{-1})\| \cdot \|\gamma(C)\| + \|\gamma(C_\varepsilon^{-1})\|) \|\gamma(T)\|. \end{aligned}$$

□

Lemma 5. $\alpha(T) \in C^*(X)$ for any $T \in C_u^*(D)$.

Proof. As $T \in C_u^*(D)$, it can be approximated by finite propagation operators T^N , $N \in \mathbb{N}$, with propagation N . This means that the matrix of T^N has the property that $t_{uv}^N = 0$ when $d(u, v) > c$ for some $c > 0$.

As $\alpha(T^N)$ can be approximated by operators of the form $\alpha_\varepsilon(T^N)$, it suffices to show that $\alpha_\varepsilon(T^N) \in C^*(X)$.

Let $f \in C_0(X)$ has compact support $K \subset X$. Then

$$\alpha_\varepsilon(T^N)\pi(f)(g) = \sum_{u,v \in D} t_{uv}^N \langle \psi_v^\varepsilon, fg \rangle \psi_u^\varepsilon = \sum_{u,v \in D} t_{uv}^N \langle \gamma(C_\varepsilon)\varphi_v, fg \rangle \gamma(C_\varepsilon)\varphi_u \quad (5)$$

As $\text{supp}(fg) \subset K$ and as propagation of $C_\varepsilon \leq M_\varepsilon$, the set $\text{supp}(\gamma(C_\varepsilon)\varphi_u)$ intersects only finitely many simplices. Therefore, $\langle \gamma(C_\varepsilon)\varphi_u, fg \rangle \neq 0$ only for finitely many $u \in D$. Together with finite propagation of T^N , this shows that the sum (5) contains only finite number of non-zero summands, i.e. $\text{Ran } \alpha_\varepsilon(T^N)\pi(f)$ is finitedimensional. Similarly, $\text{Ran } \pi(f)\alpha_\varepsilon(T^N)$ is finitedimensional. Thus $\alpha(T)\pi(f)$ and $\pi(f)\alpha(T)$ are compact. Approximation of functions in $C_0(X)$ by functions f with finite support proves that $\alpha(T)$ is locally compact.

Similarly one can show that $\alpha(T^N)$ is of finite propagation. Indeed, let $f, g \in C_0(X)$ are such that the distance between their supports is greater than R . Then

$$\pi(f)\alpha_\varepsilon(T^N)\pi(g)(h) = \sum_{u,v \in D} t_{uv}^N \langle \gamma(C_\varepsilon)\varphi_v, gh \rangle f\gamma(C_\varepsilon)\varphi_u.$$

There exists $L > 0$ such that if $d(v, \text{supp } g) > L$ then $\langle \gamma(C_\varepsilon)\varphi_v, gh \rangle = 0$, and if $d(u, \text{supp } f) > L$ then $f\gamma(C_\varepsilon)\varphi_u = 0$. Also $t_{uv}^N = 0$ when $d(u, v) > c$. Thus, if $R > 2L + c$ then $\pi(f)\alpha_\varepsilon(T^N)\pi(g) = 0$. \square

For $S \in C^*(X)$ set $(\beta(S))_{uv} = \langle \psi_u, S\psi_v \rangle$, $u, v \in D$. Then the operator $\beta(S) \in \mathbb{B}(l^2(D))$ can be written as $\beta(S)(\delta_v) = \sum_{u \in D} \langle \psi_u, S\psi_v \rangle \delta_u$. It can be also written as $\beta(S) = U^*SU$, where $U : l^2(D) \rightarrow L^2(X)$ is the defined above isometry. In particular, this implies that $\beta(S)$ is bounded for any bounded operator S . Note that β is not a homomorphism.

Lemma 6. *Let $S \in C^*(X)$. Then $\beta(S) \in C_u^*(D)$.*

Proof. It suffices to show that $\beta(S) \in C_u^*(D)$ for operators of finite propagation. For an operator S of finite propagation set $\tilde{S} = U^*F^*SFU$, where $F = C_\varepsilon C^{-1}$. As $\|1 - F\| < \varepsilon\|C^{-1}\|$, $\beta(S)$ can be approximated by operators of the form \tilde{S} . Let us show that \tilde{S} has finite propagation. We have

$$\tilde{S}_{uv} = \langle \psi_u, F^*SF\psi_v \rangle = \langle F\psi_u, SF\psi_v \rangle = \langle \psi_u^\varepsilon, S\psi_v^\varepsilon \rangle.$$

As $\text{supp } \psi_u^\varepsilon$ is compactly localized at u , and as S has finite propagation, $\tilde{S}_{uv} = 0$ when $d(u, v)$ is sufficiently great. \square

Note that $\beta \circ \alpha(S) = pS|_H$.

As the constructions of the maps α and β work for each $D(n)$, we have two sequences of maps: $\alpha_n : C_u^*(D(n)) \rightarrow C^*(X)$ and $\beta_n : C^*(X) \rightarrow C_u^*(D(n))$. We also write $U_n : l^2(D(n)) \rightarrow L^2(X)$ for U , $U_{\varepsilon,n}$ for U_ε , $\psi_u^{\varepsilon;n}$ for ψ_u^ε , etc.

4. CONTINUOUS FIELD OF ROE ALGEBRAS

Continuous fields of C^* -algebras (aka bundles of C^* -algebras or $C(T)$ - C^* -algebras) were introduced by Fell ([5], see also Dixmier ([3], Section 10)). Recall that a continuous field of C^* -algebras over a locally compact Hausdorff space T is a triple $(T, A, \pi_t : A \rightarrow A_t)$, where A and A_t , $t \in T$, are C^* -algebras, the $*$ -homomorphisms π_t are surjective, the family $\{\pi_t\}_{t \in T}$ is faithful, and the map $t \mapsto \|\pi_t(a)\|$ is continuous for any $a \in A$.

Set $T = \mathbb{N} \cup \{\infty\}$, $A_n = C_u^*(D(n))$, $A_\infty = C^*(X)$. Let $F(T) = \prod_{t \in T} A_t$ denote the C^* -algebra of families $(a_t)_{t \in T}$ such that $a_t \in A_t$ for each $t \in T$, and $\sup_{t \in T} \|a_t\| < \infty$. Consider two C^* -subalgebras in $F(T)$: let

$$J = \{(a_t)_{t \in T} \in F(T) : a_\infty = \lim_{t \rightarrow \infty} \|a_t\| = 0\} \cong \bigoplus_{t \in \mathbb{N}} A_t$$

and

$$B = \{(\beta_t(S))_{t \in T} : S \in C^*(X)\} \subset F(T),$$

where $\beta_\infty(S) = S$ (as β_n is a contraction for any $n \in \mathbb{N}$, $(\beta(S))_{t \in T} \in F(T)$).

Let $A = J + B \subset F(T)$ be the $*$ -algebra of the sums of elements of the two C^* -subalgebras.

Lemma 7. *The $*$ -algebra A is norm closed.*

Proof. First, let us show that

$$\sup_{n \in \mathbb{N}} \|\beta_n(S)\| = \|S\| = \lim_{n \rightarrow \infty} \|\beta_n(S)\|. \quad (6)$$

Consider the projections $p_n = U_n U_n^*$ in $L^2(X)$, where U_n is given by $U_n(\delta_u) = \psi_u^n$. Note that $\|U_n^* S U_n\| = \|p_n S p_n\|$. Let $f \in C_0(X)$ be a Lipschitz function with compact support $K \subset X$, and let L be the Lipschitz constant for f . Set $g_n = \sum_{x \in D(n)} f(x) \varphi_x^n$ be a piecewise linear function such that $g_n(x) = f(x)$ for any $x \in D$. Then $|f(x) - g_n(x)| \leq L \operatorname{diam}(\Delta^k(n))$ for any $x \in X$, where $\Delta^k(n)$ is a simplex of maximal dimension from the n -th subdivision of the initial simplicial structure of X , hence $\|f - g_n\|_{L^2(X)} \leq L \operatorname{diam}(\Delta^k(n)) \mu(K)$. As $\operatorname{diam}(\Delta^k(n))$ vanishes as $n \rightarrow \infty$, $\|f - g_n\|_{L^2(X)}$ vanishes as well. Note that g_n lies in the linear span of the functions φ_x^n , $x \in D$, so $g_n = p_n g_n$. As $\|f - p_n f\|_{L^2(X)} \leq \|f - p_n g_n\|_{L^2(X)}$, we have $\lim_{n \rightarrow \infty} f - p_n f = 0$. As Lipschitz functions with compact support are dense in $L^2(X)$, we conclude that the $*$ -strong limit of p_n is the identity operator.

Consider the norm closure \overline{A} of A in $F(T)$. Then J is a closed ideal in \overline{A} . Let $\{f_i + \beta(S_i)\}_{i \in \mathbb{N}}$, $f_i \in J$, $S_i \in A_\infty$, be a Cauchy sequence in A . Passing to the quotient C^* -algebra \overline{A}/J , the sequence $\{f_i + \beta(S_i) + J\} = \{\beta(S_i) + J\}$ is also a Cauchy sequence, as the quotient $*$ -homomorphisms have norm 1 ([3], Section 1.8). Note that

$$\|\beta(S) + J\| = \inf_{f \in J} \|f + \beta(S)\| \geq \lim_{n \rightarrow \infty} \|f(n) + \beta_n(S)\| = \lim_{n \rightarrow \infty} \|\beta_n(S)\| = \|S\|,$$

hence $\{S_i\}_{i \in \mathbb{N}}$ is also a Cauchy sequence. As A_∞ is norm closed, this sequence has a limit in A_∞ . But then $\{f_i\}_{i \in \mathbb{N}}$ is also a Cauchy sequence, and as J is norm closed, its limit lies in J . Thus $\{f_i + \beta(S_i)\}_{i \in \mathbb{N}}$ converges in A , hence $\overline{A} = A$. \square

Define $\pi_t : A \rightarrow A_t$ by $\pi_t(f + \beta(S)) = f(t) + \beta_t(S)$, $t \in T$. These maps are well defined as $f_1 + \beta(S_1) = f_2 + \beta(S_2)$ implies that $f_1 = f_2$ and $S_1 = S_2$.

Theorem 8. *The triple $(T, A, \pi_t : A \rightarrow A_t)$ is a continuous field of C^* -algebras.*

Proof. Each π_t is clearly surjective. If $\pi_t(f_1 + \beta(S_1)) = \pi_t(f_2 + \beta(S_2))$ for any $t \in T$ then, taking $t = \infty$, we conclude that $S_1 = S_2$. Then we see that $f_1(t) = f_2(t)$ for any $t \in \mathbb{N}$, hence $f_1 = f_2$. Finally, we have to check that the map $t \mapsto \|\pi_t(a)\|$ is continuous at $t = \infty$, which follows from (6). \square

5. DIRECT LIMIT OF UNIFORM ROE ALGEBRAS

The main observation of this section is existence of a $*$ -homomorphism $j_n : C_u^*(D(n)) \rightarrow C_u^*(D(n+1))$ such that the diagram

$$\begin{array}{ccc} C_u^*(D(n)) & \xrightarrow{j_n} & C_u^*(D(n+1)) \\ & \searrow \alpha_n & \swarrow \alpha_{n+1} \\ & & C^*(X) \end{array} \quad (7)$$

commutes.

There is an obvious inclusion of $C_u^*(D(n))$ into $C_u^*(D(n+1))$ induced by the inclusion of $D(n)$ into $D(n+1)$, but this inclusion does not make the diagram (7) commute, so we cannot pass to the direct limit.

Recall that H_n is the closed subspace of $L^2(X)$ spanned by the partition of unity $\{\varphi_u^n\}_{u \in D(n)}$, which is a (non-orthonormal) basis for H_n . Our observation is the inclusion $H_n \subset H_{n+1}$. Indeed, it is easy to see that

$$\varphi_u^n = \varphi_u^{n+1} + \frac{1}{2} \sum_{x \in D(n+1) \setminus \{u\}; x \sim u} \varphi_x^{n+1},$$

where $u \in D(n)$, and $x \sim u$ denotes that u and x lie in the same simplex $\Delta_\alpha(n+1)$. Denote the inclusion $H_n \subset H_{n+1}$ by I_n , and recall that we have constructed isometric

isomorphisms $U_n : l^2(D(n)) \rightarrow H_n$. Define $W_n : l^2(D(n)) \rightarrow l^2(D(n+1))$ by $W_n = U_{n+1}^* I_n U_n$. Then W_n is an isometry, different from the one induced by the inclusion of $D(n)$ into $D(n+1)$. Set $j_n(T) = W_n T W_n^*$. Then $j_n : C_u^*(D(n)) \rightarrow \mathbb{B}(l^2(D(n+1)))$ is a $*$ -homomorphism.

Lemma 9. $j_n(T) \in C_u^*(D(n+1))$ for any $T \in C_u^*(D(n))$.

Proof. Recall that besides the maps U_n , we have constructed maps $U_{\varepsilon;n} : l^2(D(n)) \rightarrow H_n$ by setting $U_{\varepsilon;n}(\delta_u) = \psi_u^{\varepsilon;n}$ such that for any $\varepsilon > 0$ there exists $L > 0$ such that $\text{supp } \psi_u^{\varepsilon;n}$ lies in the ball of radius L centered at $u \in D(n)$, and $\lim_{\varepsilon \rightarrow 0} \|U_n - U_{\varepsilon;n}\| = 0$. As $\langle U_{\varepsilon;n}(\delta_u), \delta_v \rangle = \langle \delta_u, U_{\varepsilon;n}^*(\delta_v) \rangle$, the support $\text{supp } U_{\varepsilon;n}(\delta_u)$ also lies in the same ball. Set $W_{\varepsilon;n} = U_{\varepsilon;n+1}^* I_n U_{\varepsilon;n}$. Then $\lim_{\varepsilon \rightarrow 0} \|W_{\varepsilon;n} - W_n\| = 0$. As all the three operators $U_{\varepsilon;n+1}^*$, I_n and $U_{\varepsilon;n}$ have finite propagation, so $W_{\varepsilon;n}$ has finite propagation too, and if T has finite propagation then $W_{\varepsilon;n} T W_{\varepsilon;n}^*$ also has finite propagation, hence lies in $C_u^*(D(n+1))$. Thus we have approximated $j_n(T)$ by finite propagation operators. \square

Let $C_u^*(X)$ be the direct limit C^* -algebra for the sequence of C^* -algebras $C_u^*(D(n))$ and of $*$ -homomorphisms j_n .

Corollary 10. *The sequence of maps $\alpha_n : C^*(D(n)) \rightarrow C^*(X)$ induces an injective $*$ -homomorphism $\alpha : C_u^*(X) \rightarrow C^*(X)$.*

Proof. Commutativity of the diagram (7) implies that the direct limit map α is well defined. As each α_n is injective, hence an isometry, so is the limit map. \square

Let us describe $\alpha(C_u^*(X)) \subset C^*(X)$ in more detail. For each $u \in D(1)$ choose a closed set $F_u \subset X$ with the following properties:

- there exist $r, R > 0$ such that $B_r(u) \subset F_u \subset B_R(u)$ for any $u \in D(1)$;
- $X = \cup_{u \in D(1)} F_u$;
- $\mu(F_u \cap F_v) = 0$ for any $u, v \in D(1)$, $u \neq v$.

Then

$$L^2(X) = \oplus_{u \in D(1)} L^2(F_u), \quad (8)$$

and it is shown in [10] that if we write an operator T on $L^2(X)$ as a matrix $(T_{uv})_{u,v \in D(1)}$ with respect to the decomposition (8) then $T \in C^*(X)$ iff $T_{uv} : L^2(F_v) \rightarrow L^2(F_u)$ is compact for any $u, v \in D(n)$ and T lies in the closure of the operators of finite propagation. Consider the subset $R_{m,l}(X) \subset C^*(X)$ of operators $T \in C^*(X)$ of propagation $\leq l$ such that the rank of T_{uv} does not exceed m for any $u, v \in D(1)$. If $T, S \in R_{m,l}(X)$ then $TS \in R_{mb(l),b(l)}(X)$, where $b(l)$ denotes the maximal number of points in a ball of radius l in $D(1)$ (recall that $D(1)$ is assumed to have bounded geometry), hence $\cup_{m,l \in \mathbb{N}} R_{m,l}(X)$ is a $*$ -subalgebra in $C^*(X)$. Denote its norm closure by $R(X)$.

Lemma 11. *One has $\alpha(C_u^*(X)) \subset R(X)$.*

Proof. By definition, $T_{uv} = P_{F_v} T P_{F_u}$, $u, v \in D(1)$, where P_{F_u} denotes the projection onto $L^2(F_u)$. If T lies in the range of α_n then $T = p_n T p_n$, where p_n is the projection onto $H_n \subset L^2(X)$, so $T_{uv} = P_{F_v} p_n T p_n P_{F_u}$, and it suffices to estimate the rank of $P_{F_v} p_n$. Let $f \in H_n$, then $f = \sum_{w \in D(n)} f_w \varphi_w^n$, $f_w \in \mathbb{C}$. Let χ_F denote the characteristic function of the set $F \subset X$. Then the projection of $f \in L^2(X)$ onto $L^2(F_u)$ equals $\chi_{F_u} f$. Thus, $P_{F_v} f = \sum_{w \in D(n)} f_w \chi_{F_v} \varphi_w^n$. The product $\chi_{F_v} \varphi_w^n$ is non-zero only when $\text{supp } \varphi_w^n \cap F_v \neq \emptyset$, and there exists $M(n) > 0$ such that $\#\{w \in D(n) : \chi_{F_v} \varphi_w^n \neq 0\} \leq M(n)$ for any $v \in D(1)$. Thus, $\text{rk}((\alpha_n(T))_{uv}) \leq M(n)$ for any $u, v \in D(1)$. \square

It would be interesting to find the range of the map α at least for the case when X is a manifold and the sets F_u can be the Voronoi cells isometric to each other.

6. REMARK ON K -THEORY

Recall that a continuous field of C^* -algebras induces a homomorphism in K -theory. Let $S \in C^*(X)$, $(\beta_t(S))_{t \in T} \in A$, hence $(\beta_t(S)^2)_{t \in T} \in A$. If $S \in C^*(X)$ is a projection, i.e. satisfies $S = S^* = S^2$, then $\lim_{n \rightarrow \infty} \|\beta_n(S)^2 - \beta_n(S)\| = \|\beta_\infty(S)^2 - \beta_\infty(S)\| = \|S^2 - S\| = 0$, hence $\beta_n(S)$ is an almost projection for sufficiently great $n \in \mathbb{N}$. This also works for projections and for unitaries in matrix algebras, and equivalent projections give equivalent almost projections, so this gives a map from $K_*(C^*(X))$ to $K_*(C_u^*(X))$ (different projections give almost projections in $C_u^*(D(n))$ for different n , but all of them can be moved to the direct limit algebra $C_u^*(X)$ using the maps j_n). The inclusion $C_u^*(X) \subset C^*(X)$ induces a map in K -theory in the opposite direction. It is interesting to evaluate the compositions of these two maps for general spaces. Here we show that non-trivial elements of $K_*(C^*(X))$ are mapped to non-trivial elements in $K_*(C_u^*(D(n)))$ for the simplest non-trivial case $X = \mathbb{R}$.

Recall that the generator of $K_1(C^*(\mathbb{R}))$ can be given by the unitary V in the unitalization of $C^*(\mathbb{R})$ given by $V = PU + (1 - P)$, where $(Uf)(s) = f(s - 1)$ is the shift on \mathbb{R} , and P is the projection onto the closure of the linear span of the characteristic functions $\chi_k = U^k(\chi_{[-\frac{1}{2}, \frac{1}{2}]})$, $k \in \mathbb{Z}$, where $\chi_{[a,b]}$ denotes the characteristic function of the segment $[a, b] \subset \mathbb{R}$ [10]. (In fact, any function in $L^2([-\frac{1}{2}, \frac{1}{2}])$ would do instead of χ_0 .)

Theorem 12. $[\alpha_n(\beta_n(V))] = [V]$ in $K_1(C^*(\mathbb{R}))$ for sufficiently great n .

Proof. Note that $D(n) = \frac{1}{2^{n-1}}\mathbb{Z} \subset \mathbb{R}$. Therefore, $\beta_n(U)$ is a shift in $l^2(D(n))$, $\beta_n(U)(\delta_u) = \delta_{u+2^{n-1}}$, $u \in D(n)$. Let us find a projection in $C_u^*(D(n))$ close to $\beta_n(P)$. Set $\tilde{\omega}_k^n = \sum \varphi_u^n$, where this sum is taken over those $u \in D(n)$, for which $\text{supp } \varphi_k^n \subset [k - \frac{1}{2}, k + \frac{1}{2}]$; $\omega_k^n = \frac{\tilde{\omega}_k^n}{\|\tilde{\omega}_k^n\|}$. Then $\{\omega_k^n\}_{k \in \mathbb{Z}}$ is an orthonormal system in $L^2(\mathbb{R})$, and $\lim_{n \rightarrow \infty} \|\omega_k^n - \chi_k\| = 0$. Let q_k^n be the projection onto ω_k^n in $L^2([k - \frac{1}{2}, k + \frac{1}{2}])$, $Q^n(f) = \bigoplus_{k \in \mathbb{Z}} q_k^n$. Then

$$\|P - Q^n\| = \sup_{k \in \mathbb{Z}} \|p_k - q_k^n\| \leq 2\|\omega_0^n - \chi_0\|,$$

where p_k is the projection onto χ_k , vanishes as $n \rightarrow \infty$. Then $\lim_{n \rightarrow \infty} \|\beta_n(P) - \beta_n(Q^n)\| = 0$.

Recall that the bases $\{\varphi_u^n\}$ and $\{\psi_u^n\}$, $u \in D(n)$, are related by $\psi_v^n = \sum_{u \in D(n)} c_{uv}^n \varphi_u$, where the operator C_n with the matrix $(c_{uv}^n)_{u,v \in D(n)}$ is invertible and lies in $C_u^*(\mathbb{Z})$, hence C_n^{-1} can be approximated by operators with band matrices. Let $(d_{uv}^n)_{u,v \in D(n)}$ be the matrix of C_n^{-1} . Then we can write $\varphi_v^n = \sum_{u \in D(n)} d_{uv}^n \psi_u^n$, hence $\tilde{\omega}_k^n = \sum_v \sum_{u \in D(n)} d_{uv}^n \psi_u^n$, where the sum over v is taken for $\text{supp } \varphi_v^n \subset [k - \frac{1}{2}, k + \frac{1}{2}]$.

Then $\beta_n(Q^n)$ is the projection onto the linear span of $g_k^n = \sum_v \sum_{u \in D(n)} d_{uv}^n \delta_u \in l^2(D(n))$ (with the same condition on the sum over v). Approximating C_n^{-1} by matrices with finite number N of diagonals, we may approximate g_k^n by $\tilde{g}_k^n = \sum_v \sum_{u \in D(n); |v-u| \leq \frac{N}{2^{n-1}}} d_{uv}^n \delta_u$. As we have $d_{u+\frac{1}{2^{n-1}}, v+\frac{1}{2^{n-1}}}^n = d_{uv}^n$, it is easy to see that $\lim_{n \rightarrow \infty} \|\tilde{g}_k^n - f_k^n\| = 0$, where $f_k^n = \frac{1}{2^{n-1}} \sum_{u \in (k-\frac{1}{2}, k+\frac{1}{2})} \delta_u$. Therefore, $\lim_{n \rightarrow \infty} \|\beta_n(Q^n) - R_n\| = 0$, where $R_n = \bigoplus_{k \in \mathbb{Z}} r_k^n$, and r_k^n denotes the projection onto f_k^n . Now we have $\lim_{n \rightarrow \infty} \|\beta_n(P) - R_n\| = 0$, and $[\beta_n(V)] = [\beta_n(U)R_n + (1 - R_n)]$ in $K_1(C_u^*(D(n)))$, which is non-trivial.

As r_k^n is a rank one projection, it is equivalent to the projection s_k^n onto $\delta_{2^{n-1}k}$. Then $\alpha_n(R_n)$ is equivalent to the projection $\alpha_n(S_n)$ onto the linear span of the functions $\psi_{2^{n-1}k}$, $k \in \mathbb{N}$. Also $\alpha_n(\beta_n(U))\psi_u = \psi_{u+2^{n-1}}$, thus $[\alpha_n(\beta_n(V))] = [\alpha_n(S_n)W + (1 - \alpha_n(S_n))]$, where $W \in \mathbb{B}(L^2(\mathbb{R}))$ is given by $(Wf)(y) = y - 2^{n-1}$, and, clearly, $[\alpha_n(S_n)W + (1 - \alpha_n(S_n))]$ is equivalent to V . \square

REFERENCES

- [1] J. V. Bellissard. *Delone Sets and Material Science: a Program*. in Mathematics of Aperiodic Order. Progress in Mathematics, **309**, 405–428, Birkhäuser, 2015.
- [2] A. Connes. *Noncommutative Geometry*. Academic Press, 1995.
- [3] J. Dixmier. *C^* -Algebras*. North-Holland, 1977.
- [4] G. A. Elliott, T. Natsume, R. Nest. *The Heisenberg group and K -theory*. *K-Theory*, **7** (1991), 409–428.
- [5] J. M. G. Fell. *The structure of algebras of operator fields*. *Acta Math.* **106** (1961), 233–280.
- [6] N. Higson, J. Roe. *Analytic K -Homology*. Oxford Mathematical Monographs, OUP Oxford, 2000.
- [7] Yu. Kordyukov, V. Manuilov. *On topological obstructions to the existence of non-periodic Wannier bases*. ArXiv.org: 2304.00125.
- [8] M. Ludewig, G. C. Thiang. *Large-scale geometry obstructs localization*. *J. Math. Physics*, **63** (2022), 091902.
- [9] V. Manuilov. *An example of a continuous field of Roe algebras*. ArXiv.org: 2311.12951.
- [10] J. Roe. *Index theory, coarse geometry, and topology of manifolds*. CBMS Regional Conference Series in Mathematics, **90**. Amer. Math. Soc., Providence, RI, 1996.

MOSCOW CENTER FOR FUNDAMENTAL AND APPLIED MATHEMATICS, MOSCOW STATE UNIVERSITY,
LENINSKIE GORY 1, MOSCOW, 119991, RUSSIA
Email address: manuilov@mech.math.msu.su