Gradient flows for empirical Bayes in high-dimensional linear models

Zhou Fan, Leying Guan, Yandi Shen, Yihong Wu*

Abstract

Empirical Bayes provides a powerful approach to learning and adapting to latent structure in data. Theory and algorithms for empirical Bayes have a rich literature for sequence models, but are less understood in settings where latent variables and data interact through more complex designs.

In this work, we study empirical Bayes estimation of an i.i.d. prior in Bayesian linear models, via the nonparametric maximum likelihood estimator (NPMLE). We introduce and study a system of gradient flow equations for optimizing the marginal log-likelihood, jointly over the prior and posterior measures in its Gibbs variational representation using a smoothed reparametrization of the regression coefficients. A diffusion-based implementation yields a Langevin dynamics MCEM algorithm, where the prior law evolves continuously over time to optimize a sequence-model log-likelihood defined by the coordinates of the current Langevin iterate.

We show consistency of the NPMLE as $n, p \to \infty$ under mild conditions, including settings of random sub-Gaussian designs when $n \simeq p$. In high noise, we prove a uniform log-Sobolev inequality for the mixing of Langevin dynamics, for possibly misspecified priors and non-logconcave posteriors. We then establish polynomial-time convergence of the joint gradient flow to a near-NPMLE if the marginal negative log-likelihood is convex in a sub-level set of the initialization.

Contents

1	Introduction								
	1.1	Related work	5						
	1.2	Notational conventions	7						
2	Met	thod	7						
	2.1	Fisher-Rao gradient flow for the Gaussian sequence model: a review	$\overline{7}$						
	2.2	Variable reparametrization and joint gradient flow	8						
	2.3	Discretization and particle implementation	10						
	2.4	Posterior inference	11						
	2.5	Smoothness regularization	12						

*Department of Statistics and Data Science, Yale University

zhou.fan@yale.edu, leying.guan@yale.edu, yandi.shen@yale.edu, yihong.wu@yale.edu

This work is supported in part by NSF DMS-2142476, NSF DMS-2310836, NSF CCF-1900507.

[†]Department of Biostatistics, Yale University

3	The	eoretical guarantees	13
	3.1	Consistency of near-NPMLEs	13
	3.2	Convergence of gradient flows	14
		3.2.1 Univariate g -flow	15
		3.2.2 Univariate q -flow	16
		3.2.3 Joint flow	18
		3.2.4 Local convexity	20
4	Exp	periments	21
	4.1	Parameter choices	22
	4.2	Method comparison	24
5	Disc	cussion	32
\mathbf{A}	Pro	oof of consistency	34
	A.1	TV lower bound	34
	A.2	Covering net	37
	A.3	Completing the proof	40
	A.4	Assumption A for random designs	40
в	Ana	alysis of algorithm dynamics	41
	B.1	Fisher-Rao flow for the sequence model	41
	B.2	LSI for Langevin dynamics in high noise	43
	B.3	Analysis of the joint gradient flow	47
	B.4	Proof of (3.17)	50
	B.5	Proof of Proposition 3.8	50
\mathbf{C}	Solu	ution properties of gradient flows	57
	C.1	Solution properties of the g-flow	58
	C.2	Solution properties of the q-flow	62
	C.3	Completing the proof	67
	C.4	Proof of Lemma B.4	69
D	Der	rivations of gradient flows	71
	D.1	General background	71
	D.2	Derivatives of $F_n(g)$ and $F_n(q,g)$	73
	D.3	Non-convexity	74
\mathbf{E}	Det	tails of algorithms	74

1 Introduction

Introduced by Robbins [Rob51, Rob56] in the 1950s, empirical Bayes (EB) is a powerful framework for large-scale inference that learns and adapts to latent structure in data. In the most commonly studied setting, one or several independent samples are observed corresponding to each latent parameter, for example modeled via a Gaussian sequence model

$$y_i = \theta_i + \varepsilon_i$$
 for $i = 1, \dots, n.$ (1.1)

Treating $\{\theta_i\}_{i=1}^n$ as i.i.d. draws from a prior distribution g_* , empirical Bayes methods proceed by estimating g_* from the noisy observations $\{y_i\}_{i=1}^n$ and using the estimated prior g_* to perform downstream Bayesian inference. Such methods have found fruitful applications in numerous areas of statistics and the applied sciences, and we refer to the review articles [Cas85, Zha03, Efr21] and the monographs [ML89, CL96, Efr10b] for systematic treatments of this broad subject.

In many applications, one may instead wish to model the data $\mathbf{y} \in \mathbb{R}^n$ as having a more complex joint dependence on latent parameters $\boldsymbol{\theta} \in \mathbb{R}^p$. In this paper, we consider a canonical setting of a linear model with an i.i.d. prior,

$$\mathbf{y} = \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\varepsilon}, \qquad \theta_j \stackrel{iid}{\sim} g_* \text{ for } j = 1, \dots, p.$$
 (1.2)

Assuming that the sample size n and dimension p are both large, we study the analogous question of estimating g_* from the data (\mathbf{X}, \mathbf{y}) . One specific application in which this problem arises is in estimating effect size distributions and genetic architectures of complex traits, and we refer to [ZQPC18, O'C21, ZZ21, SSAAP22, MCW⁺23] for examples of methods developed in this context. From the perspective of statistical theory, basic questions such as conditions allowing for consistent estimation of g_* are, at present, only partially understood [MSS23].

We adopt a similar setting as in the recent work [MSS23], and focus on nonparametric estimation of a prior g_* supported on a fixed and known bounded domain $[-M, M] \subset \mathbb{R}$, via the nonparametric maximum likelihood estimator (NPMLE)

$$\widehat{g} = \arg \min_{g \in \mathcal{P}(M)} \overline{F}_n(g) \tag{1.3}$$

or its regularized variants. Here $\mathcal{P}(M)$ is the space of all probability distributions on [-M, M], and

$$\bar{F}_n(g) := -\frac{1}{p} \log P_g(\mathbf{y}) = -\frac{1}{p} \log \int_{[-M,M]^p} \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}{2\sigma^2}\right) \prod_{j=1}^p \mathrm{d}g(\theta_j) \tag{1.4}$$

is the (normalized) negative marginal log-likelihood of **y** under the prior distribution g. The NPMLE was introduced in [Rob50, KW56] and has been extensively studied in sequence models of the form (1.1). For the linear model (1.2), in contrast to the sequence model setting of (1.1), $\bar{F}_n(g)$ does not decompose across coordinates $j = 1, \ldots, p$ and is non-convex in the prior g for general regression designs. This renders optimization (or even the evaluation) of the log-likelihood $\bar{F}_n(g)$ a substantially more challenging task.

Inspired by and building upon a line of work on optimizing functionals of probability measures using gradient flows [JKO98, CB18, MMN18, LCB⁺22, YWR23], in this work we propose and study a system of bivariate gradient flow equations for optimizing $\bar{F}_n(g)$. Briefly, our approach consists of reparametrizing¹ the linear model (1.2) using an auxiliary variable $\varphi \in \mathbb{R}^p$ as

$$\mathbf{y} = \mathbf{X}\boldsymbol{\varphi} + \widetilde{\boldsymbol{\varepsilon}},\tag{1.5}$$

where φ has i.i.d. coordinates with a prior $\mathcal{N}_{\tau} * g$ that is a Gaussian convolution of g, and $\tilde{\varepsilon}$ is a (correlated) Gaussian residual defined so that (1.5) expresses the same model as (1.2). We write the Gibbs variational representation of $\bar{F}_n(g)$ in this parametrization as

$$\bar{F}_n(g) = \min_q F_n(q,g) = \min_q -\frac{1}{p} \int \left[\log P_g(\mathbf{y}, \boldsymbol{\varphi}) - \log q(\boldsymbol{\varphi}) \right] \mathrm{d}q(\boldsymbol{\varphi}) \tag{1.6}$$

¹This reparameterization is introduced for multiple reasons, such as ensuring the φ variable has a smooth density supported on all of \mathbb{R}^p which lends itself to Langevin-type sampling algorithms of the posterior. See Section 2.3 for a detailed discussion.

where the minimization is over all probability distributions q on \mathbb{R}^p . We then minimize $F_n(q,g)$ jointly over (q,g) via the simultaneous gradient flow equations

$$\begin{cases} \frac{\mathrm{d}}{\mathrm{d}t}q_t(\boldsymbol{\varphi}) &= -p \cdot \operatorname{grad}_q^{W_2} F_n(q_t, g_t)[\boldsymbol{\varphi}], \\ \frac{\mathrm{d}}{\mathrm{d}t}g_t(\boldsymbol{\theta}) &= -\alpha \cdot \operatorname{grad}_g^{\mathrm{FR}} F_n(q_t, g_t)[\boldsymbol{\theta}]. \end{cases}$$
(1.7)

Here, $\operatorname{grad}_q^{W_2}$ denotes a gradient with respect to the Wasserstein-2 geometry in q, $\operatorname{grad}_g^{\operatorname{FR}}$ denotes a gradient with respect to the Fisher-Rao geometry in g, and $\alpha > 0$ is a parameter for the relative learning rate of g versus q. Our use of the Fisher-Rao geometry for g is inspired by [YWR23], who studied also hybrid flows under combined Fisher-Rao and Wasserstein-2 geometries for computing the NPMLE in the sequence model (1.1).

Following the pioneering insights of [JKO98], the q-flow in (1.7) may be simulated via a sample $\varphi_t \sim q_t$ evolving in \mathbb{R}^p according to a Langevin diffusion process, in our setting with time-evolving drift whose form depends on g_t . The Fisher-Rao g-flow of (1.7) can be seen as a continuous-time version of the M-step in an expectation-maximization (EM) algorithm. Consequently, the resulting method may be understood as a continuous-time variant of a Monte-Carlo expectation-maximization (MCEM) procedure [WT90] with latent parameter φ . We provide a more detailed description and discussion of this method in Section 2. We discuss some issues of practical implementation and parameter tuning in Section 4, where we also present numerical comparisons with alternative MCEM and variational inference approaches. We note that related methods have also been proposed recently for general latent variable models with parametric priors in [KLJ23, ACG⁺23], and analyzed in contexts where the full-data log-likelihood is jointly convex in the latent variable and prior parameters. This joint convexity does not hold in our studied setting of linear models, requiring the development of new ideas to analyze the gradient flow dynamics. We discuss this distinction and related literature further in Section 1.1.

Our work takes a step towards understanding the theoretical properties of the NPMLE in the linear model, as well as of the gradient flow procedure (1.7) for its optimization. We summarize these contributions as follows:

1. In an asymptotic setting where $n, p \to \infty$, and under a certain deterministic condition on the design **X**, we establish consistency (in the sense of weak convergence) for any approximate minimizer \hat{g} of the negative log-likelihood $\bar{F}_n(g)$.

Our condition for **X** requires the existence of test vectors $\{\mathbf{z}_j\}_{j=1}^p$ that, roughly speaking, can decorrelate a sufficient number of variables $\{\mathbf{x}_j\}_{j=1}^p$ of the design. This builds upon and extends a recent result of [MSS23] that showed consistency for well-conditioned design matrices **X** when $n \ge p$. Our assumption is sufficiently general to encompass also settings where n < p, and in particular, settings where n and p are of a similar order and **X** is a random design having i.i.d. sub-Gaussian rows with well-conditioned population covariance.

- 2. For suitable initial conditions (q_0, g_0) , we show existence, uniqueness, and smoothness for the solution $\{(q_t, g_t)\}_{t\geq 0}$ of the coupled differential equations corresponding to (1.7), where g_t evolves in the space of densities on [-M, M] and q_t describes the law of a Langevin-type diffusion for a sample $\varphi_t \in \mathbb{R}^p$.
- 3. For fixed $q_t = q$ (or its approximation via a sample $\varphi_t = \varphi$), the "univariate" evolution of g_t in (1.7) is a Fisher-Rao gradient flow for optimizing the marginal log-likelihood in a sequence model (1.1). For any initialization g_0 fully supported on [-M, M], we prove the global convergence of this univariate flow $\{g_t\}_{t\geq 0}$ as $t \to \infty$ to the sequence model NPMLE, at the rate O(1/t).

4. For fixed $g_t = g$, the "univariate" evolution of q_t in (1.7) describes a standard Langevin diffusion with fixed drift. We show that for sufficiently large (but dimension-independent) noise variance $\sigma^2 > 0$, its stationary posterior law satisfies a log-Sobolev inequality (LSI), uniformly over all priors g supported on [-M, M]. Such a result is known to imply an exponential rate of convergence of $\{q_t\}_{t\geq 0}$, in KL-divergence as $t \to \infty$, to its stationary distribution.

Our proof of this result leverages the breakthrough insight of [BB19], who established a similar log-Sobolev inequality for high-temperature spin systems.

5. Finally, for the joint evolution of $\{(q_t, g_t)\}_{t\geq 0}$ following (1.7), under an assumption of a uniform LSI as guaranteed above, we show that if the marginal objective $\bar{F}_n(g)$ is convex in the sublevel set defined by the initial value $F_n(q_0, g_0)$, then $\{g_t\}_{t\geq 0}$ reaches, in polynomial time, an approximate NPMLE that has the same statistical consistency properties as \hat{g} .

These results are presented and discussed in Section 3.

We note that while the convergence guarantees in this paper are stated for the continuous-time flow (1.7), we believe that they also yield useful insights that pave the way to a discrete-time analysis. We discuss this and a few other interesting directions for future research in Section 5.

1.1 Related work

Empirical Bayes. The NPMLE was proposed in [Rob50, KW56] for nonparametric estimation of the unknown prior in sequence-type models. The monograph [Lin95] provides a systematic treatment of structural and geometric properties, and we refer to [PW20] for more recent results. Early consistency results on the NPMLE include [Jew82, Lin83b, LT84, Pfa88]; see also [Che17] for a recent survey. We refer to [GW00, GvdV01, Zha09, JZ09, SG20, PW21] for work on related problems of marginal density estimation and EB-compound estimation. Various algorithms have been proposed and studied for computing the NPMLE in sequence models, including EM-type methods [DLR77, Lai78, Cov84, VL93, JZ09], general constrained convex optimization techniques (e.g. the interior-point method) [LZ07, KM14], and vertex direction methods [LK92, Böh95, Wan07, GJW08]. We also mention the recent work [ZCST22] which proposed an EM-type algorithm for the sequence model (1.1) under general dimensions without a fixed support grid.

EB estimation in linear models is closely related to that in the sequence model with correlated errors, which was discussed in [Efr10a, Sch10, SS18] in a context of multiple testing and false discovery rate control. For designs \mathbf{X} having $n \ge p$ and full column rank, the sufficient z-score statistics follow such a correlated-errors model,² and the two models are equivalent. They are no longer equivalent in settings of n < p, where observations \mathbf{y} depend on $\boldsymbol{\theta}$ only via its projection onto the row span of \mathbf{X} . Some early and representative works studying EB in Bayesian linear models include [NS86, GF00, YL05] for parametric priors. A body of work in statistical genetics has developed methods in more semiparametric and nonparametric contexts, for example based on EM or hierarchical Bayes sampling algorithms [ZQPC18, ZZ21], mean-field variational approximations [SSAAP22, MCW⁺23], or moment-matching procedures in the Fourier domain [O'C21]. Nonparametric EB via a discretized computation of the NPMLE was also suggested in [WZ21].

Theoretical results on nonparametric EB in the linear model seem more scarce, and we are only aware of [JG21] who studied a mixture of (low-dimensional) regression models that is more akin to the setting of (1.1), and the recent work [MSS23] that analyzed the NPMLE and variational approximations in high-dimensional regression models. This latter work [MSS23] showed that the

²In this case, $(\mathbf{X}^{\top}\mathbf{X})^{-1}\mathbf{X}^{\top}\mathbf{y} = \boldsymbol{\theta} + \boldsymbol{\varepsilon}'$, where $\boldsymbol{\varepsilon}' \sim \mathcal{N}(0, \sigma^2(\mathbf{X}^{\top}\mathbf{X})^{-1})$.

NPMLE is consistent as $n, p \to \infty$ when $n \ge p$ and \mathbf{X}/σ has all singular values of constant order, and established conditions under which consistency holds also for the optimizer of a mean-field variational approximation to the log-likelihood objective. Our current work was motivated by applications where strong correlations between variables of \mathbf{X} may render mean-field variational methods inaccurate, and by the goal of designing a computationally efficient method for estimating the prior g_* in such contexts.

Gradient flows and Bayesian inference. Gradient flows of probability measures play a central role in our work, and we refer to [San17] for an introduction and [AGS08, Vil09] for fuller and systematic treatments. The connection between gradient-based optimization of entropy in the Wasserstein-2 geometry and Langevin diffusions was recognized in [JKO98, Ott01], and has enabled a fruitful line of work developing new analyses of Langevin dynamics as well as new Bayesian sampling algorithms [Per16, Wib18, DMM19, AC21, MCC⁺21]. Wasserstein gradient flows have also been applied to understand training dynamics of shallow neural networks [NS17, MMN18, SS20, RVE18, CB18] and to develop new methods for variational inference [LCB⁺22, YY22].

Our diffusion-based implementation of the gradient flow equations leads to a continuous-time Monte Carlo EM method [WT90, BH99, LC01], where an intractable E-step is simulated using an Unadjusted Langevin Algorithm [RT96, Dal17, DM17]. Classical theoretical treatments of MCEM include [CL95, FM03, Nea13], but typically assume an asymptotically growing number of MCMC iterations between M-step updates. Our analyses of the gradient flow equations may be understood as studying MCEM convergence in an opposite continuous-time regime, where the prior is continuously updated before the equilibration of the MCMC samples to their stationary measure.

Closest to our current work are the recent papers [KLJ23, ACG⁺23], which also proposed and studied EB methods for general latent variable models based on joint optimization or sampling of the prior and posterior in a Gibbs variational representation of the log-likelihood. These works focus on a setting where the prior q belongs to a fixed parametric family $\{q_{\alpha}\}$, and the analyses therein assume that the full-data log-likelihood log $P(\theta, \mathbf{y} \mid \boldsymbol{\alpha})$ of the latent variable θ and observation \mathbf{y} is jointly strongly concave in (θ, α) . Such a condition is sufficient to ensure that the marginal loglikelihood log $P(\mathbf{y} \mid \boldsymbol{\alpha})$ has a unique local maximizer $\hat{\boldsymbol{\alpha}}$, and that furthermore the Gibbs variational objective is Wasserstein-2 convex in q for each fixed α . It is then shown in [KLJ23] that a bivariate gradient flow in α and q converges at an exponential rate to $\hat{\alpha}$ and its corresponding posterior. It may be checked that this joint convexity does not hold in our problem, and furthermore the Gibbs variational objective may not be Wasserstein-2 convex in q under priors q where the posterior is non-log-concave. These issues motivate our choice of writing the variational objective using the reparametrized regression variable in (1.5), and our development of different ideas to analyze the gradient flow dynamics. We mention also the related works [Chi22, NWS22] that established quantitative convergence of McVean-Vlasov Langevin diffusions for optimizing entropy-regularized functionals F(q) under uniform LSI assumptions. However, these works also assume that F(q) is globally convex in the linear geometry. Such an assumption again does not hold in our setting, where the "entropy sandwich" arguments of [Chi22, NWS22] are not available.

For performing nonparametric optimization over the prior g, we will use (for simplicity) the Fisher-Rao geometry. The ideas of the Fisher-Rao geometry date back to the classical work [Hel09, Hot30, RR45, Kak48]. More recently, together with the Wasserstein geometry, it has played an important role in the study of unbalanced optimal transport and the associated Wasserstein-Fisher-Rao flow [CPSV18b, CPSV18a, KMV16, LMS18, LMS16, YWR23]. For sampling, recent work [LLN19, LSW23, DEP23] showed that dynamics resulting from the combined Wasserstein-Fisher-Rao gradient flow can accelerate the convergence of Langevin dynamics, especially when the target prior is not log-concave.

1.2 Notational conventions

 $\mathcal{P}(M), \mathcal{P}(\mathbb{R}), \mathcal{P}(\mathbb{R}^p)$ are the spaces of all probability distributions on [-M, M], \mathbb{R} , and \mathbb{R}^p , and $\mathcal{P}_*(M), \mathcal{P}_*(\mathbb{R}), \mathcal{P}_*(\mathbb{R}^p)$ are the spaces of probability densities on these domains with respect to Lebesgue measure.

For two measures P, Q such that P is absolutely continuous to Q, the Kullback-Leibler divergence is $D_{\text{KL}}(P||Q) = \int \log \frac{\mathrm{d}P}{\mathrm{d}Q} \mathrm{d}P$. The total variation distance is $d_{\text{TV}}(P,Q) = \frac{1}{2} \int |\mathrm{d}P - \mathrm{d}Q|$.

We use \mathcal{N}_{σ} as shorthand for the Gaussian law $\mathcal{N}(0, \sigma^2)$, and $\mathcal{N}_{\sigma}(x)$ for its density function at $x \in \mathbb{R}$. We use P * Q for the convolution of two measures, so $[\mathcal{N}_{\sigma} * g](x)$ is the density of the convolution of \mathcal{N}_{σ} with g.

Let ∇f , Δf , and $\nabla \cdot f$ denote the gradient, Laplacian, and divergence of a function $f : \mathbb{R}^d \to \mathbb{R}$. A function $f : \Omega \to \mathbb{R}$ on an open domain Ω belongs to $C^r(\Omega)$ (resp. $C^{\infty}(\Omega)$) if it is *r*-times (resp. infinitely-times) continuously differentiable on Ω . We use $\|\mathbf{A}\|_{op}$ for the operator norm of a matrix \mathbf{A} , and $\lambda_{\min}(\mathbf{A})$ (resp. $\lambda_{\max}(\mathbf{A})$) for its smallest (resp. largest) eigenvalue when \mathbf{A} is symmetric.

2 Method

In our exposition, we treat \mathbf{X} as deterministic, i.e. all results are conditional on \mathbf{X} . We assume throughout that g_* is a prior distribution supported on a fixed and known interval [-M, M], and we seek to estimate g_* from (\mathbf{X}, \mathbf{y}) following the linear model (1.2).

2.1 Fisher-Rao gradient flow for the Gaussian sequence model: a review

Consider first the sequence model (1.1) with Gaussian errors of variance σ^2 , corresponding to the identity design matrix $\mathbf{X} = \text{Id}$ with p = n. The NPMLE for priors supported on [-M, M] is

$$\widehat{g} = \arg\min_{g \in \mathcal{P}(M)} \overline{F}_n(g), \quad \text{where } \overline{F}_n(g) = -\frac{1}{n} \sum_{i=1}^n \log[\mathcal{N}_\sigma * g](y_i)$$
(2.1)

and $\mathcal{N}_{\sigma} * g$ denotes the convolution of $\mathcal{N}_{\sigma} \equiv \mathcal{N}(0, \sigma^2)$ with g.

Gradient flows for optimizing $\overline{F}_n(g)$ were studied in [YWR23], and we review here a simple such flow operating in the Fisher-Rao geometry for g: For any functional F(g) of a positive density gon [-M, M], we define its first variation $\delta F[g] : [-M, M] \to \mathbb{R}$ as the unique (up to an additive constant) function for which

$$\frac{\mathrm{d}}{\mathrm{d}\varepsilon}\Big|_{\varepsilon=0}F(g+\varepsilon\chi) = \int_{-M}^{M} \delta F[g](x)\chi(x)\mathrm{d}x$$
(2.2)

for every bounded function $\chi : [-M, M] \to \mathbb{R}$ satisfying $\int_{-M}^{M} \chi(x) dx = 0$. In particular, the first variation of $\bar{F}_n(g)$ in (2.1) may be computed to be

$$\delta \bar{F}_n[g](\theta) = -\frac{1}{n} \sum_{i=1}^n \frac{\mathcal{N}_\sigma(y_i - \theta)}{\mathcal{N}_\sigma * g(y_i)} + 1, \qquad (2.3)$$

where we choose the constant +1 so that it has mean 0 under g. The Fisher-Rao gradient flow is then

$$\frac{\mathrm{d}}{\mathrm{d}t}g_t(\theta) = -g_t(\theta) \cdot \delta \bar{F}_n[g_t](\theta) = g_t(\theta) \left[\frac{1}{n} \sum_{i=1}^n \frac{\mathcal{N}_\sigma(y_i - \theta)}{\mathcal{N}_\sigma * g_t(y_i)} - 1 \right].$$
(2.4)

This may be understood as an infinitesimal version of the widely studied EM algorithm [DLR77]

$$g_{t+1}(\theta) = \frac{1}{n} \sum_{i=1}^{n} \frac{g_t(\theta) \mathcal{N}_{\sigma}(y_i - \theta)}{\mathcal{N}_{\sigma} * g_t(y_i)},$$
(2.5)

and we refer to [YWR23] for other interpretations and a more detailed discussion. Both the Fisher-Rao flow (2.4) and EM algorithm (2.5) may be approximated numerically by restricting to $g = \sum_{k=1}^{K} w_k \delta_{b_k}$ supported on a fixed grid $b_1, \ldots, b_K \in [-M, M]$ and updating the weights (w_1, \ldots, w_K) .

For more general designs \mathbf{X} , the Fisher-Rao gradient flow may be written as (c.f. Appendix D.2 for a derivation)

$$\frac{\mathrm{d}}{\mathrm{d}t}g_t(\theta) = \bar{\mu}[g_t](\theta) - g_t(\theta) \tag{2.6}$$

where

$$\bar{\mu}[g](x) = \frac{1}{p} \sum_{j=1}^{p} \int_{\mathbb{R}^{p-1}} P_g(\theta_1, \dots, \theta_{j-1}, x, \theta_{j+1}, \dots, \theta_p \mid \mathbf{y}) \prod_{i: i \neq j} \mathrm{d}\theta_i$$
(2.7)

is the average marginal density of coordinates $\theta_1, \ldots, \theta_p$ under the posterior law $\mu[g](\boldsymbol{\theta}) := P_g(\boldsymbol{\theta} \mid \mathbf{y})$. Exact computation of $\bar{\mu}[g]$ is intractable for general regression designs. As mentioned in Section 1, we will address this challenge by performing the gradient flow instead on a Gibbs variational representation (1.6) of $\bar{F}_n(g)$, coupled with a gradient flow over the space of high-dimensional posterior measures on \mathbb{R}^p .

2.2 Variable reparametrization and joint gradient flow

Consider now the linear model with i.i.d. prior g_* and noise variance $\sigma^2 > 0$, which we restate here for clarity:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\varepsilon}, \qquad \theta_j \stackrel{iid}{\sim} g_* \text{ for } j = 1, \dots, p, \qquad \boldsymbol{\varepsilon} \sim \mathcal{N}(0, \sigma^2 \operatorname{Id}) \text{ independent of } \boldsymbol{\theta}.$$
 (2.8)

To describe the reparametrization (1.5), fix any parameter $\tau^2 > 0$ satisfying $\tau^2 < \sigma^2 / \lambda_{\max}(\mathbf{X}\mathbf{X}^{\top})$, where $\lambda_{\max}(\cdot)$ denotes the largest eigenvalue. Decompose the Gaussian noise $\boldsymbol{\varepsilon}$ as $\boldsymbol{\varepsilon} = \boldsymbol{\widetilde{\varepsilon}} + \mathbf{X}\mathbf{z}$ where $\mathbf{z} \sim \mathcal{N}(0, \tau^2 \operatorname{Id})$ and $\boldsymbol{\widetilde{\varepsilon}} \sim \mathcal{N}(0, \boldsymbol{\Sigma})$ with

$$\boldsymbol{\Sigma} = \sigma^2 \operatorname{Id} - \tau^2 \mathbf{X} \mathbf{X}^\top \succ \mathbf{0}.$$

Setting $\varphi = \theta + \mathbf{z}$, the marginal law of \mathbf{y} is equivalently expressed as

$$\mathbf{y} = \mathbf{X}\boldsymbol{\varphi} + \widetilde{\boldsymbol{\varepsilon}}, \qquad \varphi_j \stackrel{iid}{\sim} \mathcal{N}_\tau * g_*, \qquad \widetilde{\boldsymbol{\varepsilon}} \sim \mathcal{N}(0, \mathbf{\Sigma}) \text{ independent of } \boldsymbol{\varphi}, \tag{2.9}$$

where we recall that $\mathcal{N}_{\tau} * g$ denotes the convolution of the $\mathcal{N}(0, \tau^2)$ density with g. Then the negative log-likelihood $\bar{F}_n(g) = -\frac{1}{p} \log P_g(\mathbf{y})$ takes the form (equivalent to (1.4))

$$\bar{F}_n(g) = -\frac{1}{p} \log \int_{\mathbb{R}^p} \frac{1}{(2\pi)^{n/2} (\det \mathbf{\Sigma})^{1/2}} \exp\left(-\frac{1}{2} (\mathbf{y} - \mathbf{X}\boldsymbol{\varphi})^\top \mathbf{\Sigma}^{-1} (\mathbf{y} - \mathbf{X}\boldsymbol{\varphi})\right) \prod_{j=1}^p [\mathcal{N}_\tau * g](\varphi_j) \mathrm{d}\varphi_j.$$
(2.10)

This functional $\overline{F}_n(g)$ has the Gibbs variational representation,³ up to an additive constant not depending on g,

$$\bar{F}_n(g) = \min_{q \in \mathcal{P}_*(\mathbb{R}^p)} F_n(q,g),$$

³This representation follows from rearranging the identity $0 = \min_q D_{\text{KL}}(q(\boldsymbol{\varphi}) || P(\boldsymbol{\varphi} | \mathbf{y}))$, noting that the minimization may be restricted to probability densities $q(\cdot)$ since the posterior law $P(\boldsymbol{\varphi} | \mathbf{y})$ admits a density on \mathbb{R}^p .

$$F_n(q,g) := \frac{1}{p} \int \left[\frac{1}{2} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi}) - \sum_{j=1}^p \log[\mathcal{N}_\tau * g](\varphi_j) + \log q(\boldsymbol{\varphi}) \right] q(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} + \frac{n}{2p} \log(2\pi) + \frac{1}{2p} \log \det \boldsymbol{\Sigma}.$$
(2.11)

The minimization is over the space of all probability densities on \mathbb{R}^p , and the minimizer for each fixed prior g is given by the posterior law

$$\nu[g](\boldsymbol{\varphi}) := P_g(\boldsymbol{\varphi} \mid \mathbf{y}). \tag{2.12}$$

Thus the NPMLE $\hat{g} := \arg \min_g \bar{F}_n(g)$ may be computed by joint minimization of $F_n(q,g)$. This representation is also the basis of empirical Bayes learning using variational inference approaches [BBB⁺03], where the domain of the candidate posterior law $q(\cdot)$ is restricted to a variational subclass of densities on \mathbb{R}^p , as well as using expectation-maximization (EM) algorithms which may be understood as optimizing $F_n(q,g)$ using alternating updates of q and g [NH98].

Here, we study optimization of $F_n(q,g)$ over densities q on \mathbb{R}^p and g on [-M, M] via the simultaneous gradient flows (1.7), under a Wasserstein-2 geometry for q [JKO98, Vil09] and a Fisher-Rao geometry for g [KMV16, YWR23]. The first variations of $F_n(q,g)$ are (up to additive constants)

$$\delta_q F_n(q,g)[\boldsymbol{\varphi}] = \frac{1}{p} \left[\frac{1}{2} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi}) - \sum_{j=1}^p \log[\mathcal{N}_\tau * g](\varphi_j) + \log q(\boldsymbol{\varphi}) \right],$$

$$\delta_g F_n(q,g)[\theta] = - \left[\mathcal{N}_\tau * \frac{\bar{q}}{\mathcal{N}_\tau * g} \right] (\theta) + 1,$$
(2.13)

where, corresponding to any density $q(\cdot)$ on \mathbb{R}^p , we define its coordinate-wise average marginal density $\bar{q}(\cdot)$ on \mathbb{R} by

$$\bar{q}(x) = \frac{1}{p} \sum_{j=1}^{p} \int_{\mathbb{R}^{p-1}} q(\varphi_1, \dots, \varphi_{j-1}, x, \varphi_{j+1}, \dots, \varphi_p) \prod_{i: i \neq j} \mathrm{d}\varphi_i.$$
(2.14)

The gradient flow equations (1.7) then take the explicit form of the coupled system of differential equations

$$\frac{\mathrm{d}}{\mathrm{d}t}q_t(\boldsymbol{\varphi}) = \nabla \cdot \left[q_t(\boldsymbol{\varphi}) \left(\mathbf{X}^\top \boldsymbol{\Sigma}^{-1} (\mathbf{X} \boldsymbol{\varphi} - \mathbf{y}) - \left(\frac{[\mathcal{N}_\tau * g_t]'(\varphi_j)}{[\mathcal{N}_\tau * g_t](\varphi_j)} \right)_{j=1}^p \right) \right] + \Delta q_t(\boldsymbol{\varphi}), \quad (2.15)$$

$$\frac{\mathrm{d}}{\mathrm{d}t}g_t(\theta) = \alpha \, g_t(\theta) \left(\left[\mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * g_t} \right](\theta) - 1 \right), \tag{2.16}$$

where $\nabla \cdot f(\varphi)$ and $\Delta f(\varphi)$ denote the divergence and Laplacian in φ . We provide a heuristic derivation of (2.13) and (2.15–2.16) in Appendix D.2.⁴ We emphasize that the Fisher-Rao gradient in (2.16) is computed with respect to the unsmoothed prior law g for θ , not the smoothed prior law $\mathcal{N}_{\tau} * g$ for φ , as performing a gradient flow in the latter would not directly yield an estimate for g_* and, more importantly, would require a method of constraining the density to the space of Gaussian convolutions.

⁴We will henceforth treat the coupled system (2.15-2.16) as the definition of the gradient flow equations (1.7), and not formally justify their equivalence with a theory of gradient flows on metric spaces as developed in [AGS08].

2.3 Discretization and particle implementation

As first recognized in [JKO98], the evolution (2.15) resulting from the Wasserstein-2 gradient of $F_n(q,g)$ in q is a Fokker-Planck equation with q_t being the density of a sample $\varphi_t \in \mathbb{R}^p$ evolving according to the following Langevin diffusion:

$$d\boldsymbol{\varphi}_t = \left(-\mathbf{X}^{\top} \mathbf{\Sigma}^{-1} (\mathbf{X} \boldsymbol{\varphi}_t - \mathbf{y}) + \left(\frac{[\mathcal{N}_\tau * g_t]'(\boldsymbol{\varphi}_{t,j})}{[\mathcal{N}_\tau * g_t](\boldsymbol{\varphi}_{t,j})} \right)_{j=1}^p \right) dt + \sqrt{2} \, \mathrm{d}\mathbf{B}_t, \tag{2.17}$$

where $\{\mathbf{B}_t\}_{t\geq 0}$ is a standard Brownian motion in \mathbb{R}^p . In our setting, this diffusion is a McVean-Vlasov process in the sense that the first drift term is coupled to the evolution (2.16) for g_t , which in turn depends on the law q_t of its solution φ_t via the average marginal density \bar{q}_t .

We remark that estimating the high-dimensional density q_t on \mathbb{R}^p may be challenging, but estimating its average marginal \bar{q}_t on \mathbb{R} is conceivably easier. In this work, we will simulate the coupled system (2.16) and (2.17) using an estimate of \bar{q}_t in (2.16) by the empirical distribution of coordinates of φ_t ,

$$\bar{q}_t \approx \frac{1}{p} \sum_{j=1}^p \delta_{\varphi_{t,j}}.$$
(2.18)

To reduce the influence of Monte Carlo variance of this estimate on the prior g_t , we will set a small relative learning rate parameter α in (1.7) in our numerical implementations, so that g_t depends in aggregate on estimates of $\{\bar{q}_t\}$ over a longer effective history of past time. We conjecture that for high dimensions p and design matrices **X** that have limited long-range dependence across variables, the estimate on the right side of (2.18) may concentrate well around \bar{q}_t without needing to average across multiple parallel samples of φ_t . We discuss this further in Section 5, and leave the theoretical exploration of this to future work.

For simplicity, we opt for a discretization of (2.16) and (2.17) in time using a forward Euler scheme, and also discretize the prior distribution g in space as

$$g = \sum_{k=1}^{K} w_k \delta_{b_k}, \qquad (2.19)$$

where we fix an equally-spaced grid of K support points $b_1, \ldots, b_K \in [-M, M]$ with corresponding probability weights $w = (w_1, \ldots, w_K)$. Together with the approximation (2.18), this yields

$$\begin{bmatrix} \mathcal{N}_{\tau} * \frac{\bar{q}_t}{\mathcal{N}_{\tau} * g_t} \end{bmatrix} (\theta) = \int \frac{\mathcal{N}_{\tau}(\theta - \varphi)}{\sum_{k=1}^{K} w_k \mathcal{N}_{\tau}(b_k - \varphi)} \bar{q}_t(\varphi) \mathrm{d}\varphi \approx \frac{1}{p} \sum_{j=1}^{p} \frac{\mathcal{N}_{\tau}(\theta - \varphi_{t,j})}{\sum_{k=1}^{K} w_k \mathcal{N}_{\tau}(b_k - \varphi_{t,j})},$$
$$\frac{[\mathcal{N}_{\tau} * g]'(\varphi)}{[\mathcal{N}_{\tau} * g](\varphi)} = -\frac{1}{\tau^2} \cdot \frac{\sum_{k=1}^{K} (\varphi - b_k) w_k \mathcal{N}_{\tau}(\varphi - b_k)}{\sum_{k=1}^{K} w_k \mathcal{N}_{\tau}(\varphi - b_k)}.$$

Applying these discretizations to (2.16) and (2.17) leads to the discretized "EBflow" algorithm

$$\boldsymbol{\varphi}_{t+1} = \boldsymbol{\varphi}_t - \eta_t^{\varphi} \left[\mathbf{X}^\top \boldsymbol{\Sigma}^{-1} (\mathbf{X} \boldsymbol{\varphi}_t - \mathbf{y}) + \frac{1}{\tau^2} \left(\frac{\sum_{k=1}^K (\varphi_{t,j} - b_k) w_{t,k} \, \mathcal{N}_\tau (\varphi_{t,j} - b_k)}{\sum_{k=1}^K w_{t,k} \, \mathcal{N}_\tau (\varphi_{t,j} - b_k)} \right)_{j=1}^p \right] + \sqrt{2\eta_t^{\varphi}} \, \mathbf{b}_t,$$
(2.20)

$$w_{t+1,k} = w_{t,k} + \eta_t^w w_{t,k} \left[\frac{1}{p} \sum_{j=1}^p \frac{\mathcal{N}_\tau(b_k - \varphi_{t+1,j})}{\sum_{i=1}^K w_{t,i} \, \mathcal{N}_\tau(b_i - \varphi_{t+1,j})} - 1 \right] \text{ for } k = 1, \dots, K,$$
(2.21)

where $\mathbf{b}_t \sim \mathcal{N}(0, \mathrm{Id})$ is a standard Gaussian vector in \mathbb{R}^p generated independently in each iteration, $w_t := (w_{t,1}, \ldots, w_{t,K})$ are the probability weights representing $g_t = \sum_{k=1}^{K} w_{t,k} \delta_{b_k}$, and $\eta_t^{\varphi}, \eta_t^{w} > 0$ are step sizes for the updates of φ_t and w_t respectively, with $\eta_t^w / \eta_t^{\varphi}$ representing the relative learning rate $\alpha > 0$ in (1.7) and (2.16). (In our experiments, we will allow η_t^w and η_t^{φ} to vary with the iteration t, fixing their ratio $\eta_t^w / \eta_t^{\varphi}$.)

Note that the iterations (2.20) represent steps of an Unadjusted Langevin Algorithm (ULA) [RT96] with time-varying prior. The prior is updated via (2.21), which has the equivalent form

$$w_{t+1,k} = (1 - \eta_t^w) w_{t,k} + \eta_t^w \cdot \frac{1}{p} \sum_{j=1}^p \frac{w_{t,k} \,\mathcal{N}_\tau(b_k - \varphi_{t+1,j})}{\sum_{i=1}^K w_{t,i} \,\mathcal{N}_\tau(b_i - \varphi_{t+1,j})}$$

As such, as long as $\eta_t^w \leq 1$, we have $w_t \geq 0$ for all iterations $t \geq 1$, an appealing property resulting from the form of the Fisher-Rao gradient $\operatorname{grad}_q^{\operatorname{FR}} F_n(q,q)$.

Let us now briefly discuss the role of the initial variable reparametrization $\theta \mapsto \varphi$ in (2.9), and the interpretation of this algorithm as a Monte-Carlo EM (MCEM) procedure. If the true prior g_* has disconnected intervals of support, then the support of its posterior $\mu[g_*](\theta)$ would be similarly disconnected, and it is clear that a Langevin diffusion for sampling from this posterior would not be ergodic. Even if the true prior g_* were such that this Langevin diffusion is rapidly mixing, such a property may not hold under the priors $\{g_t\}$ evolving according to the gradient flow (2.15–2.16). Indeed, for identity design (i.e. the sequence model (1.1)) the exact NPMLE \hat{g} is known to be supported on a finite set of atoms [Lin83a, PW20], and we expect for more general designs that $\{g_t\}$ may also converge to a discrete distribution in the limit $t \to \infty$, if no additional regularization is imposed. In contrast, the prior $\mathcal{N}_{\tau} * g_t$ induces a continuous posterior distribution $\nu[g_t](\varphi)$ for φ even if g_t is discrete. For sufficiently large noise variance $\sigma^2 > 0$, we will further verify in Section 3 that this posterior $\nu[g_t](\varphi)$ satisfies a log-Sobolev inequality, which holds irrespective of the connectedness of the support of g_t or a lower bound for its density. For the discretized algorithm (2.20–2.21), reparametrization by φ likewise enables posterior sampling using a continuous Langevin diffusion even if the support of $g_t(\cdot)$ is discretized to a grid.

The two steps (2.20–2.21) of EBflow may be understood as "incremental" versions of the E-step and M-step in an EM algorithm with latent variable φ . Indeed, an exact EM algorithm in this parametrization would minimize $F_n(q, g)$ via the following alternating updates of q and g:

(E-step)
$$q_t = \nu[g_t]$$
 (the posterior law of φ under the prior $\mathcal{N}_{\tau} * g_t$),
(M-step) $g_{t+1} = \arg\min_g \int -\log[\mathcal{N}_{\tau} * g](\varphi)\bar{q}_t(\varphi)\mathrm{d}\varphi.$
(2.22)

Both steps have computational difficulties in practice: The E-step requires computation of the posterior $\nu[g_t]$, which is intractable for general design matrices **X**. The update (2.20) of EBflow replaces this E-step by a single step of ULA, and then approximates \bar{q}_t as needed in the M-step by the empirical distribution of coordinates of the sampled $\varphi_t = (\varphi_{t,j})_{j=1}^p$ cf. (2.18). Under this approximation, g_{t+1} in the M-step is precisely the solution of an NPMLE problem in the Gaussian sequence model (1.1) with observations $(\varphi_{t,1}, \ldots, \varphi_{t,p})$. Computing g_{t+1} exactly may be expensive to carry out in every iteration, and the update (2.21) of EBflow replaces this M-step by a single step of Fisher-Rao gradient descent for solving this sequence model NPMLE problem.

2.4 Posterior inference

For downstream tasks of posterior inference, one may perform Monte Carlo approximation using samples $\varphi_t \in \mathbb{R}^p$ of (2.20–2.21) for a set of final iterations $t \in \{T+1, \ldots, T+T'\}$ after fixing the final

estimated weights $w_T = (w_{T,1}, \ldots, w_{T,K})$. Typically one is interested in computing expectations under the posterior law of the original parameter $\boldsymbol{\theta}$, rather than under the reparametrized variable $\boldsymbol{\varphi}$. Such expectations may often be estimated using samples $\{\boldsymbol{\varphi}_t\}$ via the relation

$$\mathbb{E}[f(\boldsymbol{\theta}) \mid \mathbf{y}] = \int_{\mathbb{R}^p} \int_{[-M,M]^p} f(\boldsymbol{\theta}) P(\boldsymbol{\theta} \mid \boldsymbol{\varphi}) P(\boldsymbol{\varphi} \mid \mathbf{y}) \mathrm{d}\boldsymbol{\theta} \mathrm{d}\boldsymbol{\varphi}$$
$$\approx \frac{1}{T'} \sum_{t=T+1}^{T+T'} \int_{[-M,M]^p} f(\boldsymbol{\theta}) P(\boldsymbol{\theta} \mid \boldsymbol{\varphi}_t) \mathrm{d}\boldsymbol{\theta}.$$

Here, coordinates of $\boldsymbol{\theta}$ are independent conditional on $\boldsymbol{\varphi}$, with $P(\boldsymbol{\theta} \mid \boldsymbol{\varphi}) = \prod_{j=1}^{p} P(\theta_j \mid \varphi_j) \propto \prod_{j=1}^{p} g(\theta_j) \mathcal{N}_{\tau}(\varphi_j - \theta_j)$. This enables Monte Carlo approximation of $\mathbb{E}[f(\boldsymbol{\theta}) \mid \mathbf{y}]$ for functions f that depend only on low-dimensional marginals of $\boldsymbol{\theta}$.

For example, the posterior mean $\mathbb{E}[\boldsymbol{\theta} \mid \mathbf{y}] \in \mathbb{R}^p$ may be estimated as

$$\mathbb{E}[\boldsymbol{\theta} \mid \mathbf{y}] \approx \frac{1}{T'} \sum_{t=T+1}^{T+T'} \int_{[-M,M]^p} \boldsymbol{\theta} P(\boldsymbol{\theta} \mid \boldsymbol{\varphi}_t) \mathrm{d}\boldsymbol{\theta}$$
$$= \frac{1}{T'} \sum_{t=T+1}^{T+T'} \left(\frac{\int_{-M}^{M} \boldsymbol{\theta} \cdot g(\boldsymbol{\theta}) \, \mathcal{N}_{\tau}(\boldsymbol{\theta} - \boldsymbol{\varphi}_{t,j}) \mathrm{d}\boldsymbol{\theta}}{\int_{-M}^{M} g(\boldsymbol{\theta}) \mathcal{N}_{\tau}(\boldsymbol{\theta} - \boldsymbol{\varphi}_{t,j}) \mathrm{d}\boldsymbol{\theta}} \right)_{j=1}^{p}$$
$$= \frac{1}{T'} \sum_{t=T+1}^{T+T'} \left(\frac{\sum_{k=1}^{K} b_k w_{T,k} \, \mathcal{N}_{\tau}(b_k - \boldsymbol{\varphi}_{t,j})}{\sum_{k=1}^{K} w_{T,k} \, \mathcal{N}_{\tau}(b_k - \boldsymbol{\varphi}_{t,j})} \right)_{j=1}^{p}, \qquad (2.23)$$

where the last step holds under the space discretization (2.19) of g. This estimate (2.23) for $\mathbb{E}[\theta \mid \mathbf{y}]$ may then be used for downstream tasks such as predicting the response for new samples.

2.5 Smoothness regularization

The perspective of minimizing $F_n(q, g)$ using a joint gradient flow enables easy extensions and variations upon this basic algorithm, for example to use stochastic or preconditioned gradient updates for φ , or to incorporate regularization for the prior g.

The unregularized NPMLE \hat{g} may be adequate for purposes of prediction on new samples, but does not share the smoothness properties of g_* if the latter admits a smooth density. We describe here one simple approach for smoothness regularization, via addition of a smoothing-spline penalty to the log-likelihood objective:

$$\bar{F}_{n,\lambda}(g) = -\frac{1}{p}\log P_g(\mathbf{y}) + \frac{\lambda}{2} \int_{-M}^{M} g''(\theta)^2 d\theta.$$
(2.24)

On the discrete grid $b_1, \ldots, b_K \in [-M, M]$ with equal spacing $\Delta = b_{k+1} - b_k$, approximating g with weights $\{w_k\}_{k=1}^K$ as in (2.19) and applying $g(b_k) \approx w_k/\Delta$, we may then approximate

$$\frac{\lambda}{2} \int_{-M}^{M} g''(\theta)^2 d\theta \approx \frac{\lambda \Delta}{2} \sum_{k=1}^{K} \left[D\left(\frac{w}{\Delta}\right) \right]_k^2, \tag{2.25}$$

where D is a differential quadrature matrix approximating the second derivative, for example $D \in \mathbb{R}^{(K-2)\times K}$ having entries $D[i, i+1] = -2/\Delta^2$, $D[i, i] = D[i, i+2] = 1/\Delta^2$, and remaining

entries 0. This leads to the discretized algorithm where (2.21) is replaced by

$$w_{t+1,k} = w_{t,k} + \eta_t^w w_{t,k} \left[\frac{1}{p} \sum_{j=1}^p \frac{\mathcal{N}_\tau(b_k - \varphi_{t+1,j})}{\sum_{i=1}^K w_{t,i} \, \mathcal{N}_\tau(b_i - \varphi_{t+1,j})} - \lambda \left(\frac{D^\top D w_t}{\Delta} \right)_k - 1 + \lambda \sum_{i=1}^K w_{t,i} \left(\frac{D^\top D w_t}{\Delta} \right)_i \right].$$

$$(2.26)$$

Here, $\lambda(D^{\top}Dw)/\Delta$ represents the gradient of the right side of (2.25), and the last term centers the update so that $\sum_k w_{t,k} = 1$ remains constant. Noting that $D^{\top}D[w/\Delta]$ is an approximation to the fourth derivative $g^{(4)}(\theta)$, this term $\lambda(D^{\top}Dw)/\Delta$ may also be understood as a discrete approximation to the first variation of the continuous penalty on the left side of (2.25).

3 Theoretical guarantees

3.1 Consistency of near-NPMLEs

Formalizing (1.3), we may define any (constrained) NPMLE over $\mathcal{P}(M)$ by

$$\widehat{g} \in \arg \min_{g \in \mathcal{P}(M)} \overline{F}_n(g).$$

We note that the existence of a minimizer \hat{g} follows from the weak compactness of $\mathcal{P}(M)$ and weak continuity of $g \mapsto \bar{F}_n(g)$, although to our knowledge, uniqueness is not resolved except for identity designs [Lin83a]. As such, our consistency result applies to any such \hat{g} , and more generally, for any near-minimizer of $\bar{F}_n(g)$.

We start with the following assumption on the design matrix \mathbf{X} .

Assumption A. Let $\mathbf{x}_1, \ldots, \mathbf{x}_p$ denote the columns of \mathbf{X} , and let the noise variance $\sigma^2 > 0$ be a fixed constant. There are constants $C_*, c_*, \alpha, \beta > 0$ such that the following hold as $n, p \to \infty$:

- 1. $\|\mathbf{X}\|_{\text{op}} \leq C_*$ and $\operatorname{Tr} \mathbf{X}^\top \mathbf{X} \geq c_* p$.
- 2. Let $\mathbf{1} \in \mathbb{R}^p$ be the all-1's vector. Then $\|\mathbf{X}\mathbf{1}\|_2^2 \ge c_*p$.
- 3. For a subset $S \subseteq \{1, \ldots, p\}$ of size $|S| \ge \alpha p$ and for each $j \in S$, there exists a vector $\mathbf{z}_j \in \mathbb{R}^n$ (deterministic conditional on \mathbf{X}) such that $c_* \le \|\mathbf{z}_j\|_2^2 \le C_*$, $\|\mathbf{z}_j^\top \mathbf{X} \mathbf{1}\| \le C_*$, and

$$\max_{j \in \mathcal{S}} |\mathbf{z}_j^\top \mathbf{x}_j - 1| \to 0, \qquad \max_{j \in \mathcal{S}} \sum_{k: k \neq j} |\mathbf{z}_j^\top \mathbf{x}_k|^{2+\beta} \to 0.$$
(3.1)

4. For any $\{a_j : j \in \mathcal{S}\}$ satisfying $|a_j| \leq 1$,

$$\left\|\sum_{j\in\mathcal{S}}a_j\mathbf{z}_j\right\|_2^2 \le C_*p.$$

Let us give a brief discussion of Assumption A. Conditions (1) and (2) are mild, where (1) fixes a scaling of **X** to have typical singular values of constant order, and (2) rules out cases such as $\mathbf{X1} = 0$ where the mean of g_* would not be identifiable. Conditions (3) and (4) are key conditions which posit the existence of a sufficient number of "test vectors" $\{\mathbf{z}_j\}_{j \in S}$ such that:

- 1. each \mathbf{z}_i is sufficiently aligned with \mathbf{x}_i ;
- 2. \mathbf{z}_j has small alignment with the remaining $\{\mathbf{x}_k\}_{k\neq j}$.

The role of $\{\mathbf{z}_j\}$ is easy to understand: The distribution of $\mathbf{z}_j^{\top}\mathbf{y}$ for each $j \in S$ will "isolate" the j-th coordinate θ_j of $\boldsymbol{\theta}$, and be close to an observation from a Gaussian sequence model (1.1) with a larger error variance. This assumption is reminiscent of the construction of the debiased Lasso [ZZ14], but an important difference is that here $\{\mathbf{z}_j\}_{j\in S}$ is used only as a proof device for analyzing the NPMLE, and plays no role in the eventual algorithm.

Under Assumption A, the following result establishes the consistency of near optimizers of F_n , including any exact NPMLE \hat{g} . Its proof is presented in Appendix A.

Theorem 3.1. Let $g_* \in \mathcal{P}(M)$, and suppose Assumption A holds as $n, p \to \infty$. Let $\tilde{g} \in \mathcal{P}(M)$ be any estimator of g_* such that, almost surely,

$$\bar{F}_n(g_*) - \bar{F}_n(\tilde{g}) = \frac{1}{p} \Big(\log P_{\tilde{g}}(\mathbf{y}) - \log P_{g_*}(\mathbf{y}) \Big) \ge -\delta_n$$

for some $\delta_n \downarrow 0$ as $n, p \to \infty$. Then $\tilde{g} \Rightarrow g_*$ almost surely, in the sense of weak convergence.

A related consistency result was shown in the recent work [MSS23] studying variational inference, in settings where $p \leq n$ and **X** has singular values satisfying $c \leq \sigma_{\min}(\mathbf{X}) \leq \sigma_{\max}(\mathbf{X}) \leq C$. Under such conditions, the 'test vectors' in Assumption A can be constructed as $\mathbf{z}_j = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1}\mathbf{e}_j$ with \mathbf{e}_j denoting the *j*-th basis vector in \mathbb{R}^p , so that $\{\mathbf{z}_j^\top \mathbf{y}\}_{j=1}^p$ are the usual linear regression zscores. Our Assumption A can be seen as a relaxation of these conditions in [MSS23], allowing for a more general (n, p)-ratio, including settings where the row span of **X** is not all of \mathbb{R}^p .

The following result shows that Assumption A is satisfied for random sub-Gaussian designs with well-conditioned population covariance. We present its proof in Appendix A.4.

Proposition 3.2. Suppose $n/p \ge \gamma$ for any constant $\gamma > 0$. Suppose $\mathbf{X} \in \mathbb{R}^{n \times p}$ has i.i.d. sub-Gaussian rows $\{\frac{1}{\sqrt{n}}x^{(i)}\}_{i=1}^{n}$ where $x^{(i)}$ has mean 0, covariance $\Sigma_X \in \mathbb{R}^{p \times p}$ satisfying

$$c \le \lambda_{\min}(\mathbf{\Sigma}_X) \le \lambda_{\max}(\mathbf{\Sigma}_X) \le C$$

and sub-Gaussian norm⁵ bounded above by a constant K > 0. Then almost surely as $n, p \to \infty$, Assumption A is satisfied with $\alpha = 1$, any $\beta > 0$, some c_*, C_* depending only on (K, γ, C, c) , and

$$\mathbf{z}_j = \Pi \mathbf{X} \mathbf{\Sigma}_X^{-1} \mathbf{e}_j,$$

where $\Pi \in \mathbb{R}^{n \times n}$ is the projection matrix orthogonal to **X1**, and $\mathbf{e}_j \in \mathbb{R}^p$ is the *j*-th standard basis vector.

As mentioned above, the vectors \mathbf{z}_j are used only as a proof device to analyze the NPMLE, and they may depend on the covariance Σ_X even when Σ_X is unknown and cannot be consistently estimated in operator norm.

3.2 Convergence of gradient flows

In Section 3.2.1, we study convergence of the Fisher-Rao flow $\{g_t\}$ in (2.16) where q_t is replaced by a fixed $q \in \mathcal{P}(\mathbb{R}^p)$, corresponding to a sequence model NPMLE problem. In Section 3.2.2, we study the Langevin diffusion corresponding to a Wasserstein-2 flow $\{q_t\}$ in (2.15) where g_t is fixed at $g \in \mathcal{P}(M)$, showing a uniform log-Sobolev inequality for its stationary posterior measure under large noise variance $\sigma^2 > 0$. Lastly in Section 3.2.3, we combine the previous two ingredients to study the joint gradient flow (1.7).

⁵The sub-Gaussian norm of $U \in \mathbb{R}^p$ is defined as $||U||_{\psi_2} \triangleq \sup\{t > 0 : \sup_{a \in \mathbb{R}^p : ||a||_2 \le 1} \mathbb{E}[\exp(t(a^\top U)^2)] < \infty$. We say U is K-sub-Gaussian if $||U||_{\psi_2} \le K$.

3.2.1 Univariate g-flow

First consider the univariate Fisher-Rao gradient flow

$$\frac{\mathrm{d}}{\mathrm{d}t}g_t(\theta) = g_t(\theta) \left[\mathcal{N}_\tau * \frac{\bar{q}}{\mathcal{N}_\tau * g_t}(\theta) - 1 \right] := g_t(\theta) \left[\int \frac{\mathcal{N}_\tau(\theta - \varphi)}{\mathcal{N}_\tau * g_t(\varphi)} \mathrm{d}\bar{q}(\varphi) - 1 \right]$$
(3.2)

for $\theta \in [-M, M]$ and a fixed probability distribution $\bar{q} \in \mathcal{P}(\mathbb{R})$. This corresponds to the evolution of g_t alone in (2.16), which is a gradient flow for minimizing the objective

$$\bar{F}(g \mid \bar{q}) := \int -\log[\mathcal{N}_{\tau} * g](\varphi) \mathrm{d}\bar{q}(\varphi).$$

If $\bar{q} = \frac{1}{n} \sum_{i=1}^{n} \delta_{y_i}$ is an empirical distribution of n samples, then this objective is exactly the marginal log-likelihood $\bar{F}_n(g)$ in the Gaussian sequence model (1.1) with noise variance τ^2 , and (3.2) is exactly the gradient flow (2.4) for computing the sequence model NPMLE.

Corollary C.2 in Appendix C.1 shows that, for any $\bar{q} \in \mathcal{P}(\mathbb{R})$ with finite moment-generating function, if (3.2) is initialized at $g_0 \in \mathcal{P}_*(M)$ having strictly positive density on all of [-M, M], then it has a unique solution $\{g_t\}_{t\geq 0}$ in the space of positive densities. The following (deterministic) result establishes convergence of this solution as $t \to \infty$.

Theorem 3.3. Let $\bar{q} \in \mathcal{P}(\mathbb{R})$ with $\int e^{\lambda \varphi} \bar{q}(\varphi) d\varphi < \infty$ for all $\lambda \in \mathbb{R}$. Suppose that (3.2) is initialized from a density $g_0 \in \mathcal{P}_*(M)$ that is strictly positive on [-M, M]. Then the solution $\{g_t\}_{t\geq 0}$ to (3.2) satisfies, for any $h \in \mathcal{P}_*(M)$,

$$\bar{F}(g_t \mid \bar{q}) - \bar{F}(h \mid \bar{q}) \le \frac{D_{\mathrm{KL}}(h \parallel g_0)}{t}.$$
 (3.3)

If $\bar{q} = \mathcal{N}_{\tau} * g_*$ is a Gaussian mixture for some true prior with density $g_* \in \mathcal{P}_*(M)$, then the minimizer of $\bar{F}(g \mid \bar{q})$ is exactly g_* , and applying Theorem 3.3 with $h = g_*$ bounds directly the suboptimality gap of g_t . If instead \bar{q} is the empirical distribution of a sample (y_1, \ldots, y_n) and \hat{g} is the minimizer of $\bar{F}(g \mid \bar{q}) = \bar{F}_n(g)$ given by the NPMLE in the sequence model (1.1), then we may apply Theorem 3.3 with h being a smoothed approximation of \hat{g} , to show

$$\bar{F}_n(g_t) - \bar{F}_n(\widehat{g}) \le \frac{D_{\mathrm{KL}}(h || g_0)}{t} + \bar{F}_n(h) - \bar{F}_n(\widehat{g}).$$

In particular, taking $t \to \infty$ first, followed by $h \Rightarrow \hat{g}$ weakly, and noting that $\bar{F}_n(g)$ is weakly continuous in g, we have $\lim_{t\to\infty} \bar{F}_n(g_t) = \bar{F}_n(\hat{g})$, so g_t is a near-NPMLE for large t.

The proof of Theorem 3.3 can be found in Appendix B.1. Motivated by viewing (3.2) as a mirror flow [NY83], the analysis exploits the convexity of $\bar{F}(g) \equiv \bar{F}(g \mid \bar{q})$ in the standard linear geometry for any fixed \bar{q} , namely

$$\overline{F}(\lambda g_0 + (1 - \lambda)g_1) \le \lambda \overline{F}(g_0) + (1 - \lambda)\overline{F}(g_1), \tag{3.4}$$

to bound $\bar{F}(g_t) - \bar{F}(h)$ by the integrated first variation $\int_{-M}^{M} -\delta \bar{F}[g_t](\theta)h(\theta)d\theta$. Some related results include [LLN19, LSW23, DEP23] which established exponential convergence of the Fisher-Rao gradient flow for optimizing a different objective $g \mapsto D_{\mathrm{KL}}(g||\pi)$ with π denoting the measure to sample from, and [YWR23, Theorem 4] which showed a conditional convergence result for a discrete-time version of (3.2). Complementary to this result, Theorem 3.3 provides a quantitative and unconditional convergence guarantee for the continuous-time flow.

3.2.2 Univariate q-flow

Next, define the negative log-posterior-density for prior $g(\cdot)$ (up to a normalizing constant),

$$U_g(\boldsymbol{\varphi}) = \frac{1}{2} (\mathbf{y} - \mathbf{X}\boldsymbol{\varphi})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \mathbf{X}\boldsymbol{\varphi}) - \sum_{j=1}^p \log \mathcal{N}_\tau * g(\varphi_j).$$
(3.5)

We consider the Langevin diffusion

$$\mathrm{d}\boldsymbol{\varphi}_t = -\nabla U_g(\boldsymbol{\varphi}_t) + \sqrt{2} \,\mathrm{d}\mathbf{B}_t \tag{3.6}$$

with a fixed prior $g \in \mathcal{P}(M)$, whose law q_t satisfies the Fokker-Planck evolution

$$\frac{\mathrm{d}}{\mathrm{d}t}q_t(\boldsymbol{\varphi}) = \nabla \cdot [q_t(\boldsymbol{\varphi})\nabla U_g(\boldsymbol{\varphi})] + \Delta q_t(\boldsymbol{\varphi}).$$
(3.7)

This corresponds to the evolution of q_t alone in (2.15).

We prove a uniform log-Sobolev inequality (LSI) for the stationary measure $\nu[g](\varphi) = P_g(\varphi | \mathbf{y})$ of the diffusion (3.6), for any prior $g \in \mathcal{P}(M)$ and any noise level $\sigma^2 > M^2 ||\mathbf{X}||_{op}^2$. As this type of result may be of independent interest in other contexts, we also provide a LSI for the posterior measure $\mu[g](\theta) = P_g(\theta | \mathbf{y})$ in the original θ parametrization when $g \in \mathcal{P}_*(M)$ has a density bounded from above and below.

For any $g \in \mathcal{P}(M)$, let us denote the posterior density of φ given (\mathbf{X}, \mathbf{y}) , under its i.i.d. prior $\mathcal{N}_{\tau} * g$, by

$$\nu[g](\boldsymbol{\varphi}) := P_g(\boldsymbol{\varphi} \mid \mathbf{y}) = \frac{\exp(-U_g(\boldsymbol{\varphi}))}{\int \exp(-U_g(\boldsymbol{\varphi}')) \mathrm{d}\boldsymbol{\varphi}'}.$$
(3.8)

When the prior has a density $g \in \mathcal{P}_*(M)$, we denote the posterior density of $\boldsymbol{\theta}$ given (\mathbf{X}, \mathbf{y}) in the original parametrization of the linear model (2.8) by $\mu[g](\boldsymbol{\theta}) = P_g(\boldsymbol{\theta} \mid \mathbf{y})$. Following [Led99, Eq. (2.18)], we say that a distribution ρ on an open domain $\Omega \subseteq \mathbb{R}^p$ satisfies a log-Sobolev inequality (LSI) with constant C > 0 if,⁶ for any bounded, Lipschitz, and continuously differentiable function $f: \Omega \to \mathbb{R}$,

$$\operatorname{ent}_{\rho}(f^2) := \mathbb{E}_{\rho} f^2 \log f^2 - \mathbb{E}_{\rho} f^2 \cdot \log \mathbb{E}_{\rho} f^2 \le 2C \cdot \mathbb{E}_{\rho} \|\nabla f\|^2.$$
(3.9)

The following (deterministic) result, whose proof is given in Section B.2, establishes a LSI for the posterior distribution in both parametrizations.

Theorem 3.4. Let $g \in \mathcal{P}(M)$ and fix a constant $\delta > 0$. Suppose that $\sigma^2 / \|\mathbf{X}\|_{op}^2 > M^2 + \delta$. Then:

- 1. If $\tau^2 \in (0, \sigma^2/\|\mathbf{X}\|_{op}^2 \delta)$, then $\nu[g](\boldsymbol{\varphi})$ satisfies a LSI with constant $C = C(M, \tau, \delta) > 0$.
- 2. Suppose that $g \in \mathcal{P}_*(M)$ and $\kappa(g) \equiv \frac{\sup_{\theta \in [-M,M]} g(\theta)}{\inf_{\theta \in [-M,M]} g(\theta)} < \infty$. Then $\mu[g](\theta)$ satisfies a LSI on $(-M,M)^p$ with constant $C \cdot \kappa(g)^2$ for some $C = C(M,\delta) > 0$.

A direct and well-known consequence of Theorem 3.4-(1) (see [Vil09, p. 288]) is that the Langevin diffusion (3.6) converges exponentially fast in KL-divergence to $\nu[g]$.

⁶The inequality (3.9) corresponds to "LSI(λ)" in the formulation of [Vil09] with $\lambda = 1/C$.

Corollary 3.5. Under the assumption of Theorem 3.4-(1), let $q_0 \in \mathcal{P}(\mathbb{R}^p)$, and let $\{q_t\}_{t\geq 0}$ be the marginal laws of $\{\varphi_t\}_{t\geq 0}$ in (3.6). Then for any $t\geq 0$,

$$D_{\mathrm{KL}}(q_t \| \nu[g]) \le \exp(-2t/C) \cdot D_{\mathrm{KL}}(q_0 \| \nu[g]),$$

where C is the same constant as in Theorem 3.4-(1).

We mention in passing that Theorem 3.4-(2) likewise implies exponential convergence in KL for a Langevin diffusion in θ reflected at the boundaries of $[-M, M]^p$ [BEL18].

The posterior laws $\nu[g](\varphi) = P_g(\varphi \mid \mathbf{y})$ and $\mu[g](\theta) = P_g(\theta \mid \mathbf{y})$ are, in general, not logconcave and also not close to product measures, and a dimension-free LSI cannot be obtained by a naive application of Holley-Stroock perturbation in \mathbb{R}^p . Our proof of Theorem 3.4 uses instead the idea of [BB19] to express these posterior laws as log-concave mixtures of product distributions. In our setting of a Bayesian linear model, this idea has an intuitive interpretation related to the reparametrization of θ by φ : For the posterior of θ , we may consider the generative process $\theta \mapsto \varphi' \mapsto \mathbf{y}$ where $\varphi' = \theta + \mathbf{z}$ is given as in (2.9) for some $\tau'^2 > 0$ to be specified, and write

$$P_g(\boldsymbol{\theta} \mid \mathbf{y}) = \int P_g(\boldsymbol{\theta} \mid \boldsymbol{\varphi}') P_g(\boldsymbol{\varphi}' \mid \mathbf{y}) \mathrm{d}\boldsymbol{\varphi}'.$$

Then a LSI for $P_g(\boldsymbol{\theta} \mid \mathbf{y})$ follows from those for $P_g(\boldsymbol{\theta} \mid \boldsymbol{\varphi}')$ (uniform over $\boldsymbol{\varphi}'$) and $P_g(\boldsymbol{\varphi}' \mid \mathbf{y})$ (uniform over \mathbf{y}), which are established as follows:

- Since $P_g(\theta \mid \varphi')$ is a product measure, by tensorization it suffices to establish a uniform LSI over its univariate marginals. We do this using a Holley-Stroock perturbation argument and the characterization of LSI for one-dimensional measures [BG99] (c.f. Lemma B.2).
- The smoothed prior $\mathcal{N}_{\tau'} * g$, and hence also the posterior measure $P_g(\varphi' | \mathbf{y})$, become strongly log-concave when $\tau'^2 > M^2$, and a LSI follows from the Bakry-Emery criterion [BE85].

The LSI for $P_g(\varphi \mid \mathbf{y})$ follows from a similar argument, where we consider instead the generative process $\boldsymbol{\theta} \mapsto \varphi \mapsto \varphi' \mapsto \mathbf{y}$ to represent

$$P_g(\varphi \mid \mathbf{y}) = \int P_g(\varphi \mid \varphi') P_g(\varphi' \mid \mathbf{y}) \mathrm{d}\varphi'.$$

We remark that if $\tau^2 > M^2$ in our starting reparametrization by φ , then we may simply take $\varphi' = \varphi$ and the LSI follows directly from log-concavity. However, choosing a large value of τ^2 leads to a difficult sequence-model NPMLE problem to be solved by the evolution of $\{g_t\}$ in the joint flow, which we observe in practice can lead to unstable behavior of the discretized algorithm and slow convergence. Theorem 3.4 allows for any value $\tau^2 > 0$, for which the prior $\mathcal{N}_{\tau} * g$ is not necessarily log-concave.

The mixing of Langevin diffusions for posterior sampling in Bayesian regression models has been studied before, for example in [DT12] using a Meyn-Tweedie approach and in [Dal17] for log-concave priors. We mention also the recent results of [NW22] that show rapid mixing of Langevin diffusions in settings where the posterior concentrates in a locally log-concave region of the parameter space. However, to our knowledge, these results do not seem to directly imply polynomial-time mixing in many settings of interest in our application, such as that of Proposition 3.2 where p scales linearly with n and the posterior $P_g(\varphi | \mathbf{y})$ may not be log-concave or contract around the true regression coefficient parameter.

3.2.3 Joint flow

Equipped with the understanding of its two components, we now study the joint flow (2.15–2.16), recorded here for the reader's convenience:

$$\frac{\mathrm{d}}{\mathrm{d}t}q_t(\boldsymbol{\varphi}) = \nabla \cdot \left[q_t(\boldsymbol{\varphi})\nabla U_{g_t}(\boldsymbol{\varphi})\right] + \Delta q_t(\boldsymbol{\varphi})
\frac{\mathrm{d}}{\mathrm{d}t}g_t(\theta) = \alpha \cdot g_t(\theta) \left(\left[\mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * g_t} \right](\theta) - 1 \right).$$
(3.10)

The following first verifies the existence and uniqueness of a solution to this joint gradient flow in $\mathcal{P}_*(\mathbb{R}^p) \times \mathcal{P}_*(M)$.

Theorem 3.6. Fix any T > 0. Suppose the initialization $(q_0, g_0) \in \mathcal{P}_*(\mathbb{R}^p) \times \mathcal{P}_*(M)$ satisfies:

- (i) The density q_0 is continuous and upper bounded, and for any $\boldsymbol{\lambda} \in \mathbb{R}^p$ we have $\mathbb{E}_{\boldsymbol{\varphi} \sim q_0} e^{\boldsymbol{\lambda}^\top \boldsymbol{\varphi}} < \infty$.
- (ii) The density g_0 is strictly positive on all of [-M, M].

Then there exists a unique classical solution $\{(q_t, g_t)\}_{t \in [0,T]}$ to (2.15–2.16) such that $(q_t, g_t) \in \mathcal{P}_*(\mathbb{R}^p) \times \mathcal{P}_*(M)$ for all $t \in [0,T]$. This solution $\{(q_t, g_t)\}_{t \in [0,T]}$ satisfies:

- 1. $t \mapsto g_t(\theta)$ is C^{∞} on (0,T) for each fixed $\theta \in [-M,M]$. Furthermore, $g_t(\theta) \ge g_0(\theta)e^{-t}$ for each $\theta \in [-M,M]$ and $t \in [0,T]$
- 2. $\{q_t\}_{t\in[0,T]}$ are the densities of the marginal laws of $\{\varphi_t\}_{t\in[0,T]}$ solving the SDE (2.17) with initialization $\varphi_0 \sim q_0$. For any $\lambda \in \mathbb{R}^p$,

$$\sup_{t \in [0,T]} \mathbb{E}_{\varphi \sim q_t} e^{\lambda^\top \varphi} < \infty.$$
(3.11)

Furthermore, $(t, \varphi) \mapsto q_t(\varphi)$ is C^{∞} on $(0, T) \times \mathbb{R}^p$.

We prove Theorem 3.6 in Appendix C.3, by identifying (2.15–2.16) as the fixed point of a contractive mapping from a suitably defined Banach space to itself.

The following (deterministic) result shows, under the joint gradient flow (2.15–2.16), that $\{g_t\}$ reaches a near-NPMLE in polynomial time if the marginal log-likelihood objective \bar{F}_n is convex in a sub-level set defined by the initial objective value. In the absence of such a local convexity guarantee, we show more generally that $\{g_t\}$ reaches a density that approximately satisfies an averaged first-order optimality condition for $\bar{F}_n(g)$, defined in terms of its (centered) first variation integrated against any fixed test density $h \in \mathcal{P}_*(M)$. For any density $g \in \mathcal{P}_*(M)$ with full support, this first variation takes the form (c.f. Appendix D.2)

$$\delta \bar{F}_n[g](\theta) = -\frac{\bar{\mu}[g]}{g}(\theta) + 1 = -\mathcal{N}_\tau * \frac{\bar{\nu}[g]}{\mathcal{N}_\tau * g}(\theta) + 1$$
(3.12)

where $\bar{\mu}[g]$ (resp. $\bar{\nu}[g]$) is the average marginal density of $\mu[g]$ (resp. $\nu[g]$) as defined by (2.7) and (2.14). (As explained in Appendix D.2, the first variation does not depend on τ , and the last expression in (3.12) holds for any $\tau > 0$ and more generally for $g \in \mathcal{P}(M)$.)

Theorem 3.7. Suppose that $\nu[g](\varphi)$ satisfies a LSI with some constant $C = C(M, \tau) > 0$, uniformly over all priors $g \in \mathcal{P}(M)$. Suppose that $(q_0, g_0) \in \mathcal{P}_*(\mathbb{R}^p) \times \mathcal{P}_*(M)$ satisfy the conditions of Theorem 3.6, and let $\{(q_t, g_t)\}_{t\geq 0}$ be the solution to (2.15–2.16). Let $F_{n,*} = \min_g \bar{F}_n(g) = \min_{(q,q)} F_n(q,g)$, and define $\delta \bar{F}_n[g]$ by (3.12).

Then there exist (M, τ) -dependent constants C, c > 0 such that for any density $h \in \mathcal{P}_*(M)$, any $\varepsilon \in (0, \min(c, \frac{1}{16\alpha}))$, and some time

$$t \le T(h,\varepsilon) := \frac{C}{\varepsilon} D_{\mathrm{KL}}(h \| g_0) + \frac{Cp \cdot e^{C\sqrt{\log(1/\varepsilon)}}}{\varepsilon^2} (F_n(q_0, g_0) - F_{n,*}),$$
(3.13)

it holds that

$$\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta) h(\theta) \mathrm{d}\theta \le \varepsilon.$$
(3.14)

If \overline{F}_n is convex (in the usual sense of (3.4)) over the sub-level set $\mathcal{K}_n \equiv \{g : \overline{F}_n(g) \leq F_n(q_0, g_0)\}$, and if $h \in \mathcal{K}_n$, then there exists some $t \leq T(h, \varepsilon)$ such that

$$\bar{F}_n(g_t) - \bar{F}_n(h) \le \varepsilon. \tag{3.15}$$

Let us make several remarks regarding the conditions and interpretation of this theorem.

- 1. The required uniform LSI condition is provided, for example, by Theorem 3.4 in high-noise settings where $\sigma^2 > M^2 \|\mathbf{X}\|_{op}^2$. Note that under Assumption A, this represents a constant and dimension-independent requirement for σ^2 , for which consistency of the NPMLE as guaranteed by Theorem 3.1 can still hold. Reparametrization by $\boldsymbol{\varphi}$ is important for this LSI to hold for the posterior laws $\nu[g_t]$ even if $\{g_t\}$ converges to a discrete measure.
- 2. The resulting guarantee (3.14) may be understood as an averaged version of a first-order optimality condition for g_t to locally minimize $\bar{F}_n(g)$. Indeed, any local minimizer \tilde{g} of $\bar{F}_n(g)$ must satisfy $\frac{\mathrm{d}}{\mathrm{d}t}\bar{F}_n((1-t)\tilde{g}+t\delta_\theta)|_{t=0+} \geq 0$ for all $\theta \in [-M, M]$, which implies that

$$-\delta \bar{F}_n(\tilde{g})(\theta) \le 0 \quad \text{for all } \theta \in [-M, M].$$
(3.16)

The statement (3.14) averages this inequality with respect to a fixed (arbitrary) reference density h. If \bar{F}_n is convex on the sub-level set \mathcal{K}_n containing h, then

$$\bar{F}_n(h) - \bar{F}_n(g) \ge \int_{-M}^M \delta \bar{F}_n[g](\theta) \big(h(\theta) - g(\theta) \big) \mathrm{d}\theta = \int_{-M}^M \delta \bar{F}_n[g](\theta) h(\theta) \mathrm{d}\theta,$$

so in particular, the statement (3.14) is sufficient to imply the bound (3.15) on the suboptimality gap.

3. Under Assumption A and any choice $\tau^2 < \sigma^2 / \|\mathbf{X}\|_{op}^2 - \delta$, one may check (see Appendix B.4) that on an event of probability approaching 1,

$$F_n(q_0, g_0) - F_{n,*} \le C \left(1 + \frac{n + \mathbb{E}_{q_0}[\|\varphi\|_2^2]}{p} \right) + \frac{1}{p} D_{\mathrm{KL}} \left(q_0 \| (\mathcal{N}_\tau * g_0)^{\otimes p} \right)$$
(3.17)

for a constant $C := C(\delta, \sigma^2, M) > 0$. If $g_* \in \mathcal{P}_*(M)$ admits a density on [-M, M], then choosing $h = g_*$ in (3.15) verifies that, under the local convexity assumption of the theorem and the mild initial conditions $D_{\mathrm{KL}}(g_*||g_0) = O(1)$, $D_{\mathrm{KL}}(q_0||(\mathcal{N}_\tau * g_0)^{\otimes p}) = O(p)$, and $\mathbb{E}_{q_0}[||\boldsymbol{\varphi}||_2^2] = O(p)$, the gradient flow reaches in time $t = O(\varepsilon^{-2+o(1)}(n+p))$ a near-NPMLE g_t (in the sense of Theorem 3.1) satisfying $\bar{F}_n(g_t) \leq \bar{F}_n(g_*) + \varepsilon$.

If g_* does not admit a density, then the theorem may be applied with h being a smoothed approximation of g_* similar to the discussion after Theorem 3.3 for the univariate g-flow.

4. We emphasize that the local convexity assumption needed for (3.15) pertains to the marginal objective \overline{F}_n , and the theorem does not require $F_n(q,g)$ to be jointly convex on the sub-level set $\{(q,g): F_n(q,g) \leq F_n(q_0,g_0)\}$.

Fixing any initial prior g_0 and any $\delta > 0$, under the assumed LSI, one may first run a Langevin diffusion for a dimension-independent time $T_0(\delta)$ to obtain an initialization q_0 that satisfies $F_n(q_0, g_0) < F_n(\nu[g_0], g_0) + \delta = \bar{F}_n(g_0) + \delta$. The theorem may then be applied with this (q_0, g_0) , under the condition that \bar{F}_n is convex over $\{g : \bar{F}_n(g_0) < \bar{F}_n(g_0) + \delta\}$.

In certain asymptotic settings as $n, p \to \infty$, we expect $\bar{F}_n(g)$ and its functional derivatives to converge to those of a deterministic and dimension-independent functional $\bar{F}(g)$. In such settings, $\bar{F}_n(g)$ for all large n, p would be convex in a fixed sub-level set \mathcal{K} of $\bar{F}(g)$ containing g_* , whose definition is dimension-free. We formalize such a statement in Section 3.2.4 below.

The proof of Theorem 3.7 is given in Appendix B.3. On a high level, the proof analyzes the integrated first variation $\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta)h(\theta)d\theta$ as in Theorem 3.3 on convergence of the *g*flow, and shows that under the assumed LSI, replacing the posterior law $\bar{\nu}[g_t]$ in (D.2) by \bar{q}_t can inflate the value of $\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta)h(\theta)d\theta$ by at most a constant factor plus a quantity that is controllable in terms of $D_{\text{KL}}(q_t || \nu[g_t])$. The divergence $D_{\text{KL}}(q_t || \nu[g_t])$ may in turn be bounded by the time-derivative of the objective value $F_n(q_t, g_t)$ via a second application of the assumed LSI.

As mentioned in the introduction, convergence of MCEM methods to fixed points of the EM algorithm has been classically studied in [CL95, FM03, Nea13], in settings of asymptotically growing numbers of EM iterations between discrete M-step updates. Theorem 3.7 may be understood as a convergence result for a continuous-time version of MCEM, in a regime where the prior g_t is continuously updated before the law q_t of the posterior sample reaches the stationary law $\nu[g_t]$.

3.2.4 Local convexity

Finally, let us formalize an asymptotic condition under which the sub-level set \mathcal{K}_n of Theorem 3.7 may be taken to contain a fixed, dimension-free neighborhood of the true prior g_* , for all large n, p.

For any strictly positive density $g \in \mathcal{P}_*(M)$ and bounded function $\chi : [-M, M] \to \mathbb{R}$ satisfying $\int_{-M}^{M} \chi(\theta) d\theta = 0$, we may define a functional Hessian of \bar{F}_n in the direction of χ by

$$\bar{H}_n[g](\chi) = \frac{\mathrm{d}^2}{\mathrm{d}^2\varepsilon}\Big|_{\varepsilon=0}\bar{F}_n(g+\varepsilon\chi).$$

Explicitly, writing $\langle f \rangle_g = \int f(\varphi) \nu[g](\varphi) d\varphi$ as shorthand for the mean of $f(\varphi)$ under the posterior law $\nu[g](\varphi) = P_g(\varphi \mid \mathbf{y})$, we may check by direct differentiation of the form (2.10) for $\bar{F}_n(g)$ (c.f. Lemma B.5) that

$$\bar{H}_{n}[g](\chi) = \frac{1}{p} \left[\left\langle \sum_{j=1}^{p} \left(\frac{\mathcal{N}_{\tau} * \chi}{\mathcal{N}_{\tau} * g}(\varphi_{j}) \right)^{2} \right\rangle_{g} - \left\langle \left(\sum_{j=1}^{p} \frac{\mathcal{N}_{\tau} * \chi}{\mathcal{N}_{\tau} * g}(\varphi_{j}) \right)^{2} \right\rangle_{g} + \left\langle \sum_{j=1}^{p} \frac{\mathcal{N}_{\tau} * \chi}{\mathcal{N}_{\tau} * g}(\varphi_{j}) \right\rangle_{g}^{2} \right].$$
(3.18)

We may extend this definition (3.18) of $\overline{H}_n[g](\chi)$ to all probability measures $g \in \mathcal{P}(M)$ and signed measures χ on [-M, M] such that $\int_{-M}^{M} \chi(\mathrm{d}\theta) = 0$.

We consider a general setting where there exist limiting functionals $\bar{F}(g)$ and $\bar{H}[g](\chi)$ to which $\bar{F}_n(g)$ and $\bar{H}_n[g](\chi)$ converge uniformly, as $n, p \to \infty$. Let $\|\chi\|_{\text{TV}} = \int_{-M}^{M} |\chi(d\theta)|$ be the total variation norm of the signed measure χ , and define

$$\Gamma = \left\{ \chi : \chi \text{ is a signed measure supported on } [-M, M], \int_{-M}^{M} \chi(\mathrm{d}\theta) = 0, \ \|\chi\|_{\mathrm{TV}} = 1 \right\}.$$

Proposition 3.8. Define $\overline{H}_n[g]$ by (3.18), and suppose that there exist deterministic functionals $\overline{F} : \mathcal{P}(M) \to \mathbb{R}$ and $\overline{H} : \mathcal{P}(M) \times \Gamma \to \mathbb{R}$ (not depending on n, p) such that, as $n, p \to \infty$,

$$\sup_{g \in \mathcal{P}(M)} \left| \bar{F}_n(g) - \bar{F}(g) \right| \xrightarrow{P} 0, \qquad \sup_{g \in \mathcal{P}(M), \, \chi \in \Gamma} \left| \bar{H}_n[g](\chi) - \bar{H}[g](\chi) \right| \xrightarrow{P} 0.$$
(3.19)

Suppose furthermore that the true prior g_* is the unique minimizer of \overline{F} , and \overline{H} satisfies

$$\inf_{\chi \in \Gamma} \bar{H}[g_*](\chi) > 0. \tag{3.20}$$

Let $F_{n,*} = \min_g \bar{F}_n(g)$. Then there exist a weak neighborhood \mathcal{K} of g_* and a constant $c_0 > 0$, both independent of (n, p), such that with probability approaching 1, $\bar{F}_n(g)$ is convex over the sub-level set $\mathcal{K}_n = \{g : \bar{F}_n(g) \leq F_{n,*} + c_0\}$ which contains \mathcal{K} .

The proof of Proposition 3.8 is given in Appendix B.5. One simple type of design under which these asymptotic conditions may be verified is a block design where **X** is block-diagonal with Nequal blocks $\bar{\mathbf{X}} \in \mathbb{R}^{B_1 \times B_2}$ independent of (n, p), and $(n, p) = (NB_1, NB_2)$. Then we may simply take $\bar{F} = \mathbb{E}\bar{F}_n$ and $\bar{H} = \mathbb{E}\bar{H}_n$, noting that these functionals are the same for all $N \ge 1$. We verify in Proposition B.8 of Appendix B.5 that the conditions of Proposition 3.8 indeed hold for this example.

We believe that the conditions of Proposition 3.8 may hold also for random designs **X** with i.i.d. entries, and more generally for random sub-Gaussian designs of the form of Proposition 3.2 under certain asymptotic conditions for the design covariance Σ_X , in the proportional asymptotics limit as $n, p \to \infty$ with $n/p \to \gamma \in (0, \infty)$. Here, $\bar{F}(g)$ would be the limit evidence or free energy of the model under a misspecified prior g, and $\bar{H}[g]$ its functional Hessian. Establishing such a convergence result is outside the scope of the current paper, although we refer to [GKKZ23] for recent related results in a different problem of low-rank matrix estimation and [BCPS21] in a linear regression model where the misspecified prior g has a parametric Gaussian form.

4 Experiments

We compare the discretized method of Section 2—hereafter referred to as EBflow—with several more "vanilla" versions of Monte-Carlo expectation maximization (MCEM) and with an approach based on mean-field variational inference using coordinate ascent updates (CAVI).

For MCEM, the most direct comparison is with an implementation using ULA to approximate the E-step, with the same latent parameter $\varphi = \theta + \mathbf{z}$ (with $\mathbf{z} \sim \mathcal{N}(0, \tau^2 \operatorname{Id})$) as in EBflow. We perform T_{iter} ULA iterations (2.20) with fixed step size η^{φ} and prior weights w between M-steps, and update w in each M-step by computing the NPMLE in the sequence model (1.1) from the coordinates $(\varphi_{t,j})_{j=1}^{p}$ in the past T_{iter} iterations. This update is a convex program, which we solve using CVXR. We refer to this method as Langevin-MCEM. We also compare to a Gibbs-MCEM method and to CAVI in the original parametrization by θ . For Gibbs-MCEM, we perform T_{iter} Gibbs sampling iterations between updates of prior weights w, each iteration resampling $\{\theta_j\}_{j\in[p]}$ in sequence. For CAVI, we optimize the variational representation of the marginal log-likelihood over priors and product posteriors for θ , by updating the posterior distribution for each coordinate $\{\theta_j\}_{j\in[p]}$ in sequence, followed by the prior. As a short burn-in period, we perform the first 200 iterations of EBflow, Langevin-MCEM, and Gibbs-MCEM without updating prior weights w, under the initialization $w = \text{Uniform}(\{b_1, \ldots, b_K\})$. Further details of the methods are discussed in Appendix E. In Section 4.1 we first discuss some theoretical considerations on parameter tuning, such as step sizes and the reparametrization parameter τ . We focus on a simple setting with Gaussian prior and i.i.d. Gaussian design as a benchmark to choose the parameters of EBflow and other methods, so that they all have reasonable performance in this pilot example. In Section 4.2, we then compare these methods across other priors and designs.

4.1 Parameter choices

Variable reparametrization. It is understood that the approximation accuracy of ULA for continuous-time Langevin dynamics improves with the smoothness of the log-density of its stationary distribution [VW19, BCE⁺22]. In the parametrization by φ , the negative Hessian of the log-posterior-density is (by Tweedie's formula)

$$-\nabla^2 \log P(\boldsymbol{\varphi} \mid \mathbf{y}) = \mathbf{X}^\top \boldsymbol{\Sigma}^{-1} \mathbf{X} + \frac{1}{\tau^2} \operatorname{Id} - \frac{1}{\tau^4} \operatorname{diag} \left(\operatorname{Var}[\theta_j \mid \varphi_j] \right)_{j=1}^p$$
(4.1)

where we recall $\Sigma = (\sigma^2 \operatorname{Id} - \tau^2 \mathbf{X} \mathbf{X}^{\top})^{-1}$. Ignoring the last φ -dependent term, which becomes negligible for bounded θ_j and large τ , and treating the largest eigenvalue

$$\lambda_{\max} := \lambda_{\max} (\mathbf{X}^{\top} \boldsymbol{\Sigma}^{-1} \mathbf{X} + \tau^{-2} \operatorname{Id})$$

of the first two terms as a proxy for the smoothness of this log-posterior, we will fix the choice

$$\tau^2 = 0.5 \,\sigma^2 / \lambda_{\max}(\mathbf{X}\mathbf{X}^{\top}) \tag{4.2}$$

in the reparametrization of θ by φ , which is easily checked to be the value that minimizes λ_{max} . Theoretical results of [VW19, BCE⁺22] on ULA suggest a step size inversely proportional to the smoothness parameter. As such, we will rescale the step size η_t^{φ} in EBflow and Langevin-MCEM relative to $1/\lambda_{\text{max}}$, i.e. all further reference to η_t^{φ} will be in the following reparametrized form of (2.20):

$$\boldsymbol{\varphi}_{t+1} = \boldsymbol{\varphi}_t - \frac{\eta_t^{\varphi}}{\lambda_{\max}} \left[\mathbf{X}^\top \boldsymbol{\Sigma}^{-1} (\mathbf{X} \boldsymbol{\varphi}_t - \mathbf{y}) + \frac{1}{\tau^2} \left(\frac{\sum_{k=1}^K w_{t,k} (\varphi_{t,j} - b_k) \mathcal{N}_\tau (\varphi - b_k)}{\sum_{k=1}^K w_{t,k} \mathcal{N}_\tau (\varphi_{t,j} - b_k)} \right)_{j=1}^p \right] + \sqrt{\frac{2\eta_t^{\varphi}}{\lambda_{\max}}} \mathbf{b}_t.$$
(4.3)

Step sizes. A larger step size η_t^{φ} in EBflow leads to faster mixing, but can result in a higher discretization bias for ULA. The step size η_t^w determines an exponential weighting of past iterates when updating the prior. An overly large η_t^w can lead to high Monte Carlo variance in the estimated prior, while an overly small η_t^w can lead to slow convergence.

To illustrate these trade-offs, we generate data $\mathbf{y} = \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\varepsilon}$ with dimensions n = 500 and p = 1000, with standard Gaussian prior $g_* = \mathcal{N}(0, 1)$, i.i.d. Gaussian design \mathbf{X} , and noise variance σ^2 chosen⁷ so that $\boldsymbol{\varepsilon}$ and $\mathbf{X}\boldsymbol{\theta}$ each explain 50% of the variance of \mathbf{y} . Let $w_{*,1}, \ldots, w_{*,K}$ denote weights representing g_* discretized to the grid $b_1, \ldots, b_K \in [-3, 3]$. Figure 1(a) displays the total-variation (TV) error $d_{\mathrm{TV}}(g_t, g_*) \approx \frac{1}{2} \sum_{k=1}^{K} |w_{t,k} - w_{*,k}|$ of the density estimate for 10,000 iterations of EBflow, over 10 independent trials on the same data (\mathbf{X}, \mathbf{y}) . To produce a smooth density estimate, we apply all methods to minimize the negative log-likelihood objective with an additional smoothing spline regularizer as in (2.24), with common fixed regularization $\lambda = 0.003$.

In this example, fixing larger step sizes $\eta_t^{\varphi} = 1.0$ and $\eta_t^w = 0.01$ results in reasonable performance, with smaller $(\eta_t^w, \eta_t^{\varphi})$ leading to slower convergence and larger $\eta_t^w/\eta_t^{\varphi}$ leading to higher

⁷In our experiment, we first generate **X** and θ , and then set σ^2 equal to the sample variance of the entries of **X** θ .



Figure 1: TV distance $d_{\text{TV}}(g_t, g_*)$ across 10000 iterations for estimating a standard Gaussian prior $g_* = \mathcal{N}(0, 1)$ via optimization of a smoothed log-likelihood objective, using various parameter tunings for EBflow, Langevin-MCEM, and Gibbs-MCEM (each with 10 independent runs).

Monte Carlo variance in the estimated prior. There is a small amount of ULA bias incurred by the large step size choice $\eta_t^{\varphi} = 1.0$, and Figure 1(b) displays the TV error under an alternative log-linear step size decay over 10,000 iterations,

$$\eta_t^{\varphi} = a \cdot b^t$$
 with $\eta_t^{\varphi} = 1.0$ at $t = 1$ and $\eta_t^{\varphi} = 0.1$ at $t = 10000$, $\eta_t^w = 0.01 \eta_t^{\varphi}$. (4.4)

The reduction of ULA bias at the final step size $\eta_t^{\varphi} = 0.1$ yields a reduction of the average TV error of the estimated density to 0.046, from 0.075 for the fixed choice $\eta_t^{\varphi} = 1.0$. We recommend using this log-linear decay (4.4) as the default option for problems of this size, with the possibility of increasing η_t^{ψ} relative to η_t^{φ} for problems with higher dimension p.

Analogous trade-offs apply for the other MCEM approaches when determining the number of iterations T_{iter} between prior updates. Figure 1(c-d) displays TV error for Langevin-MCEM with $\eta_t^{\varphi} \in \{1.0, 0.1\}$ and $T_{\text{iter}} \in \{100, 1000\}$, and for Gibbs-MCEM with $T_{\text{iter}} \in \{10, 100, 1000\}$, both across 10 independent trials for the same data (**X**, **y**). We will henceforth fix $T_{\text{iter}} = 100$ for both methods to achieve a reasonable balance between Monte Carlo variance and convergence speed.

4.2 Method comparison

Priors and designs. We generate data $\mathbf{y} = \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\varepsilon}$ where $\{\theta_j\}_{j=1}^p \overset{iid}{\sim} g_*$, and the prior distribution g_* is one of four smooth densities:

- (Gaussian) $g_* = \mathcal{N}(0, 1)$
- (Cauchy) $g_* = \text{Cauchy}(0, 0.6)$
- (Skew) $g_* = \frac{1}{3}\mathcal{N}(-2, 0.5^2) + \frac{1}{3}\mathcal{N}(-1.5, 1^2) + \frac{1}{3}\mathcal{N}(0, 2^2)$
- (Bimodal) $g_* = \frac{1}{2}\mathcal{N}(-1.5, 0.5^2) + \frac{1}{2}\mathcal{N}(1.5, 0.5^2)$

For consistency, each prior is truncated to [-3,3] and discretized over a grid of K = 61 points b_1, \ldots, b_K with equal spacing 0.1. We generate **X** to be one of the following designs:

- (identity) $\mathbf{X} = \text{Id and } n = p$.
- (iid) X has i.i.d. Gaussian entries.
- (block02corr0.9) **X** has multivariate Gaussian rows, consisting of p/2 independent pairs of variables with correlation 0.9.
- (block10corr0.5) **X** has multivariate Gaussian rows, consisting of p/10 independent blocks of variables with pairwise correlation 0.5.

We fix the dimension p = 1000 for all designs. For the latter three random designs, we vary the sample sizes $n \in \{500, 1000, 2000\}$. The noise variance σ^2 is set so that ε and $\mathbf{X}\boldsymbol{\theta}$ each explain 50% of the variance of \mathbf{y} for the Gaussian prior, and ε explains 20% of the variance of \mathbf{y} for the Cauchy, skew, and bimodal priors. To study the effects of Monte Carlo variance of the MCEM methods on the estimated prior, we fix a single generated instance (\mathbf{X}, \mathbf{y}) for each prior and design, and then evaluate 10 independent runs of each algorithm (except for CAVI which is deterministic).

Experimental setup. We fix the prior support for all methods to be the grid b_1, \ldots, b_K over [-3,3]. EBflow is run with τ^2 set as in (4.2), and the log-linear step size decay (4.4) over 10,000

iterations. We tested also a preconditioned variant [RS02] of EBflow with the φ update in (4.3) replaced by

$$\boldsymbol{\varphi}_{t+1} = \boldsymbol{\varphi}_t - \eta_t^{\varphi} \mathbf{Q}^{-1} \left[\mathbf{X}^{\top} \boldsymbol{\Sigma}^{-1} (\mathbf{X} \boldsymbol{\varphi}_t - \mathbf{y}) + \frac{1}{\tau^2} \left(\frac{\sum_{k=1}^K w_{t,k} (\varphi_{t,j} - b_k) \mathcal{N}_\tau (\varphi - b_k)}{\sum_{k=1}^K w_{t,k} \mathcal{N}_\tau (\varphi_{t,j} - b_k)} \right)_{j=1}^p \right] + \sqrt{2\eta_t^{\varphi}} \mathbf{Q}^{-1/2} \mathbf{g}_t$$

$$(4.5)$$

fixing $\mathbf{Q} = \mathbf{X}^{\top} \mathbf{\Sigma}^{-1} \mathbf{X} + \tau^{-2}$ Id as a rough proxy for the Hessian (4.1). Related forms of preconditioning have been suggested, for example, for posterior sampling in logistic regression models in [Dal17] to accelerate mixing of Langevin dynamics with ill-conditioned designs. One may check that this preconditioning corresponds to the original gradient flow (1.7) with an alternative choice of inner-product on \mathbb{R}^p underlying the Wasserstein-2 geometry. Langevin-MCEM and Gibbs-MCEM are run fixing $T_{\text{iter}} = 100$ Monte Carlo iterations between M-step updates, also over 10,000 total Monte Carlo iterations. CAVI is run for 1000 iterations, as it converges much faster than the MCEM methods.

We compare prior density estimates and computation time for all methods applied to minimize the negative log-likelihood with a small smoothing spline regularization (2.24), fixing $\lambda = 0.003$ for the Gaussian prior and $\lambda = 0.001$ for the other three priors. We report also the achieved marginal log-likelihood for identity design and mean-squared prediction error for random designs, and verify that these are similar with and without smoothness regularization.

Results. Figure 2 displays the estimated priors in an illustrative example of the block02corr0.9 design with n = p = 1000, which highlights some of the main trends across other priors and designs.

In general, the estimates of EBflow, Langevin-MCEM with large step size $\eta_t^{\varphi} = 1.0$, and Gibbs-MCEM are more accurate than the remaining two methods. Preconditioning EBflow can yield more stable estimates, due to faster mixing of the ULA chain. Langevin-MCEM at the smaller step size $\eta_t^{\varphi} = 0.1$ suffers from high Monte Carlo variance. CAVI produces density estimates that tend to underestimate the prior variance, due to inaccuracy of the variational approximation for this type of design that has strong pairwise correlations between variables.

Tables 1–4 summarize estimation accuracy and runtime across all tested methods, priors, and designs. We report the mean and standard deviation of the TV error of the estimated density, across 10 independent runs for all methods except CAVI. As rough proxies for convergence speed, we report the median number of iterations (rounded to the nearest 100) used to reach a TV error less than 0.2, or NA if the final TV error is larger than this value.⁸ We report also the attained marginal log-likelihood value \bar{F}_n (up to an additive constant) for identity design,⁹ and the relative mean-squared prediction error

$$MSE = \frac{\|\mathbf{X}_{new}(\boldsymbol{\theta}_* - \mathbb{E}[\boldsymbol{\theta} \mid \mathbf{X}, \mathbf{y}])\|_2^2}{\|\mathbf{X}_{new}\boldsymbol{\theta}_*\|_2^2}$$

across $n_{\text{new}} = 1000$ test samples \mathbf{X}_{new} for random designs, where $\boldsymbol{\theta}_*$ is the true regression parameter. The posterior mean $\mathbb{E}[\boldsymbol{\theta} \mid \mathbf{X}, \mathbf{y}]$ is estimated in EBflow and Langevin-MCEM from T' = 50,000 additional iterates of $\boldsymbol{\varphi}_t$ using (2.23), T' = 50,000 additional iterates of $\boldsymbol{\theta}_t$ in Gibbs-MCEM, and the average of the final posterior marginals estimated in CAVI.

⁸The real compute times of Langevin-MCEM and Gibbs-MCEM are slower than EBflow in our straightforward implementations of these methods. For Langevin-MCEM, this is due to the complexity of solving the sequence-model NPMLE problem in each M-step (which is not reflected in the iteration count). For Gibbs sampling, it is due to a loop over coordinates as compared to a vectorized matrix-vector computation in EBflow. In our experiments at this problem size, EBflow was about 2–3 times faster than Gibbs sampling, and 5–10 times faster than Langevin-MCEM, although these comparisons are necessarily implementation dependent.

⁹For general design, it is difficult to evaluate the marginal log-likelihood which involves a p-dimensional integral.



Figure 2: Estimated prior densities across 6 methods, and 10 independent runs for each method except CAVI. Priors left to right: Gaussian, Cauchy, skew, bimodal. All settings have sample size n = 1000 and dimension p = 1000. Design is block02corr0.9.

Design	n/p	Method	TV	TV.sd	Time (iters)	Log-likelihood
		EBflow	0.063	0.0016	500	0.511
		EBflow.preconditioned	0.063	0.0016	500	0.511
identity	1.0	Langevin.eta1.0	0.068	0.0097	200	0.496
Identity	1.0	Langevin.eta0.1	0.081	0.0142	200	0.514
		Gibbs	0.084	0.0033	300	0.514
		CAVI	0.084	NA	2	0.514
Design	n/p	Method	TV	TV.sd	Time (iters)	Prediction MSE
		EBflow	0.052	0.0059	400	0.409
		EBflow.preconditioned	0.050	0.0038	400	0.407
iid	2.0	Langevin.eta1.0	0.044	0.0063	200	0.405
110		Langevin.eta0.1	0.069	0.0149	200	0.408
		Gibbs	0.057	0.0019	300	0.408
		CAVI	0.076	NA	1	0.419
		EBflow	0.044	0.0075	700	0.585
		EBflow.preconditioned	0.045	0.0052	600	0.583
iid	1.0	Langevin.eta1.0	0.065	0.0085	400	0.580
		Langevin.eta0.1	0.091	0.0191	700	0.586
		Gibbs	0.073	0.0038	500	0.584
		CAVI	0.078	NA	2	0.611
		EBflow	0.046	0.0069	1000	0.789
		EBnow.preconditioned	0.052	0.0034	900	0.788
iid	0.5	Langevin.eta1.0	0.075	0.0072	600	0.790
		Cithe	0.000	0.0228	1100	0.787
		GIDDS	0.062	0.0065 N A	800	0.780
		EDA	0.110	NA 0.0102	2	0.766
		EBROW EDdow proceeditioned	0.030	0.0123	600	0.303
	2.0	Langevin etal 0	0.031	0.0095	200	0.302
block02corr0.9		Langevin.eta1.0	0.041	0.0140	200	0.301
		Cibba	0.075	0.0226	200	0.303
		CAVI	0.038	0.0041 N 4	400	0.303
		EBflow	0.155	0.0070	800	0.555
		EBflow preconditioned	0.004	0.0070	700	0.525
	1.0	Langevin etal 0	0.071	0.0002	400	0.521
block02corr0.9		Langevin etal 1	0.097	0.0001	1150	0.520
		Gibbs	0.083	0.0021	600	0.525
		CAVI	0.161	NA	1	0.545
		EBflow	0.032	0.0120	1100	0.646
		EBflow.preconditioned	0.037	0.0104	950	0.646
		Langevin.eta1.0	0.053	0.0099	650	0.647
block02corr0.9	0.5	Langevin.eta0.1	0.062	0.0171	1750	0.644
		Gibbs	0.038	0.0036	800	0.640
		CAVI	0.157	NA	2	0.670
		EBflow	0.081	0.0141	600	0.304
		EBflow.preconditioned	0.079	0.0102	600	0.302
hlash10aann0 5	2.0	Langevin.eta1.0	0.074	0.0213	200	0.300
DIOCK10COIT0.5	2.0	Langevin.eta0.1	0.109	0.0344	200	0.305
		Gibbs	0.088	0.0052	400	0.299
		CAVI	0.349	NA	1	0.449
		EBflow	0.031	0.0072	900	0.478
		EBflow.preconditioned	0.026	0.0080	800	0.475
block10corr0 5	10	Langevin.eta1.0	0.049	0.0076	400	0.473
510081000110.0	1.0	Langevin.eta0.1	0.077	0.0211	1650	0.477
		Gibbs	0.051	0.0054	600	0.471
		CAVI	0.283	NA	1	0.513
		EBflow	0.076	0.0134	1500	0.650
		EBflow.preconditioned	0.075	0.0125	1300	0.645
block10corr0.5	0.5	Langevin.eta1.0	0.082	0.0123	800	0.642
	0.0	Langevin.eta0.1	0.100	0.0377	2750	0.645
		Gibbs	0.068	0.0073	1000	0.636
		CAVI	0.156	NA	2	0.639

Table 1: Comparison of methods: Gaussian prior.

Design	n/p	Method	TV	TV.sd	Time (iters)	Log-likelihood
		EBflow	0.066	0.0015	400	0.340
		EBflow.preconditioned	0.066	0.0015	400	0.340
identity	1.0	Langevin.eta1.0	0.072	0.0068	200	0.334
Identity	1.0	Langevin.eta0.1	0.080	0.0077	200	0.341
		Gibbs	0.077	0.0026	200	0.342
		CAVI	0.077	NA	1	0.342
Design	n/p	Method	TV	TV.sd	Time (iters)	Prediction MSE
		EBflow	0.082	0.0062	400	0.167
		EBflow.preconditioned	0.080	0.0043	400	0.166
iid	20	Langevin.eta1.0	0.081	0.0084	200	0.166
nu	2.0	Langevin.eta0.1	0.109	0.0176	200	0.167
		Gibbs	0.087	0.0025	200	0.166
		CAVI	0.206	NA	1	0.191
	EBflow 0.089 (0.0086	600	0.397	
		EBflow.preconditioned	0.087	0.0068	550	0.393
iid	1.0	Langevin.eta1.0	0.107	0.0158	200	0.392
nu	1.0	Langevin.eta0.1	0.137	0.0376	200	0.396
		Gibbs	0.125	0.0050	400	0.388
		CAVI	0.203	NA	1	0.565
		EBflow	0.153	0.0163	1150	0.661
		EBflow.preconditioned	0.154	0.0104	1050	0.656
iid	0.5	Langevin.eta1.0	0.166	0.0161	600	0.654
nu	0.5	Langevin.eta0.1	0.204	0.0367	6750	0.669
		Gibbs	0.165	0.0089	700	0.654
		CAVI	0.304	NA	1	0.915
		EBflow	0.069	0.0132	500	0.108
	2.0	EBflow.preconditioned	0.071	0.0083	500	0.107
block02corr0 0		Langevin.eta1.0	0.083	0.0174	200	0.107
DIOCK02C0110.9		Langevin.eta0.1	0.109	0.0338	200	0.111
		Gibbs	0.117	0.0056	400	0.107
		CAVI	0.263	NA	1	0.172
		EBflow	0.105	0.0259	850	0.258
		EBflow.preconditioned	0.102	0.0164	850	0.255
block02corr0 9	1.0	Langevin.eta1.0	0.114	0.0192	200	0.254
510CR02C0110.5	1.0	Langevin.eta0.1	0.155	0.0379	200	0.262
		Gibbs	0.096	0.0054	500	0.250
		CAVI	0.284	NA	1	0.355
		EBflow	0.132	0.0212	1350	0.431
		EBflow.preconditioned	0.132	0.0153	1250	0.425
block02corr0 9	0.5	Langevin.eta1.0	0.132	0.0135	600	0.418
510011020011010	0.0	Langevin.eta0.1	0.177	0.0405	4500	0.430
		Gibbs	0.113	0.0078	800	0.412
		CAVI	0.292	NA	1	0.520
		EBflow	0.118	0.0169	400	0.135
		EBflow.preconditioned	0.108	0.0160	400	0.131
block10corr0.5	2.0	Langevin.eta1.0	0.125	0.0137	200	0.130
		Langevin.eta0.1	0.130	0.0248	200	0.137
		Gibbs	0.121	0.0023	300	0.129
		CAVI	0.236	NA	1	0.159
		EBflow	0.130	0.0181	700	0.283
		EBflow.preconditioned	0.118	0.0207	750	0.275
block10corr0.5	1.0	Langevin.eta1.0	0.114	0.0138	200	0.272
		Langevin.eta0.1	0.166	0.0467	200	0.299
		Gibbs	0.119	0.0055	500	0.268
		UAVI	0.259	INA 0.00002	1050	0.390
		EBflow	0.170	0.0336	1250	0.469
		EBHOW.preconditioned	0.163	0.0194	1200	0.460
block10corr0.5	0.5	Langevin.eta1.0	0.172	0.0315	200	0.453
	1	Langevin.eta0.1	0.192	0.0655	200	0.473
		GIDDS	0.164	0.0059 N 4	800	0.450
		UAVI	0.378	INA	1	0.068

Table 2: Comparison of methods: Cauchy prior.

Design	n/p	Method	TV	TV.sd	Time (iters)	Log-likelihood
		EBflow	0.060	0.0016	300	0.360
		EBflow.preconditioned	0.060	0.0016	300	0.360
identity	1.0	Langevin.eta1.0	0.089	0.0105	200	0.354
Identity	1.0	Langevin.eta0.1	0.076	0.0145	200	0.361
		Gibbs	0.072	0.0014	200	0.361
		CAVI	0.073	NA	1	0.361
Design	n/p	Method	TV	TV.sd	Time (iters)	Prediction MSE
		EBflow	0.097	0.0070	300	0.120
		EBflow.preconditioned	0.094	0.0047	300	0.120
iid	20	Langevin.eta1.0	0.088	0.0133	200	0.118
nu	2.0	Langevin.eta0.1	0.125	0.0225	200	0.122
		Gibbs	0.120	0.0029	200	0.121
		CAVI	0.278	NA	1	0.159
		EBflow	0.077	0.0115	400	0.233
		EBflow.preconditioned	0.082	0.0081	400	0.232
iid	1.0	Langevin.eta1.0	0.107	0.0130	300	0.232
nu	1.0	Langevin.eta0.1	0.110	0.0298	300	0.234
		Gibbs	0.056	0.0073	300	0.230
		CAVI	0.357	NA	2	0.318
		EBflow	0.114	0.0143	600	0.299
		EBflow.preconditioned	0.118	0.0094	600	0.299
iid	0.5	Langevin.eta1.0	0.128	0.0166	500	0.302
nu	0.5	Langevin.eta0.1	0.156	0.0596	1700	0.304
		Gibbs	0.079	0.0028	400	0.296
		CAVI	0.382	NA	5	0.459
		EBflow	0.057	0.0121	400	0.070
		EBflow.preconditioned	0.061	0.0074	400	0.070
block02corr0.0	20	Langevin.eta1.0	0.067	0.0130	200	0.070
DIOCK02C0110.9	2.0	Langevin.eta0.1	0.096	0.0305	200	0.071
		Gibbs	0.082	0.0051	200	0.070
		CAVI	0.348	NA	4	0.094
		EBflow	0.106	0.0152	400	0.117
	1.0	EBflow.preconditioned	0.105	0.0102	400	0.117
block02corr0.0		Langevin.eta1.0	0.116	0.0209	250	0.118
biocko2corro.5	1.0	Langevin.eta0.1	0.162	0.0584	250	0.117
		Gibbs	0.145	0.0095	200	0.116
		CAVI	0.372	NA	2	0.158
		EBflow	0.165	0.0151	500	0.220
		EBflow.preconditioned	0.165	0.0114	500	0.220
block02corr0 9	0.5	Langevin.eta1.0	0.163	0.0131	400	0.220
STOCINO	0.0	Langevin.eta0.1	0.190	0.0525	1100	0.224
		Gibbs	0.208	0.0132	300	0.221
		CAVI	0.291	NA	2	0.239
		EBflow	0.088	0.0132	400	0.061
		EBflow.preconditioned	0.087	0.0101	400	0.060
block10corr0.5	2.0	Langevin.eta1.0	0.092	0.0149	200	0.060
		Langevin.eta0.1	0.107	0.0380	200	0.061
		Gibbs	0.075	0.0068	200	0.060
		CAVI	0.406	NA	4	0.089
		EBflow	0.131	0.0138	400	0.076
		EBflow.preconditioned	0.129	0.0105	400	0.076
block10corr0.5	1.0	Langevin.eta1.0	0.123	0.0102	200	0.076
		Langevin.eta0.1	0.144	0.0025	200	0.077
		GIDDS	0.129	U.U116	200	0.076
		UAVI ED4	0.197	INA 0.0105	2	0.082
		EDHOW proceeditions	0.145	0.0105	500	0.138
		Langevin etc. 1.0	0.140	0.0000	200	0.137
block10corr0.5	0.5	Langevin.eta1.0	0.150	0.0106	300	0.130
		Cibbs	0.208	0.0020	300	0.139
		CAVI	0.130	0.0097 N A	300 n	0.137
	1	UNVI	0.204	11/1	4	0.147

Table 3: Comparison of methods: Skew prior.

Design	n/p	Method	TV	TV.sd	Time (iters)	Log-likelihood
		EBflow	0.079	0.0007	1400	0.320
		EBflow.preconditioned	0.079	0.0007	1400	0.320
identity	1.0	Langevin.eta1.0	0.189	0.0076	400	0.310
Identity	1.0	Langevin.eta0.1	0.074	0.0121	300	0.323
		Gibbs	0.058	0.0024	700	0.325
		CAVI	0.061	NA	6	0.325
Design	n/p	Method	TV	TV.sd	Time (iters)	Prediction MSE
		EBflow	0.076	0.0079	800	0.121
		EBflow.preconditioned	0.079	0.0057	800	0.121
iid	2.0	Langevin.eta1.0	0.103	0.0143	600	0.122
na	2.0	Langevin.eta0.1	0.088	0.0182	900	0.123
		Gibbs	0.070	0.0031	600	0.121
		CAVI	0.096	NA	4	0.146
	EBflow 0.226 0.0193 N		NA	0.325		
		EBflow.preconditioned	0.235	0.0148	NA	0.323
iid	1.0	Langevin.eta1.0	0.260	0.0226	NA	0.325
na	1.0	Langevin.eta0.1	0.203	0.0516	6250	0.324
		Gibbs	0.137	0.0046	3800	0.312
		CAVI	0.206	NA	14	0.518
		EBflow	0.353	0.0106	NA	0.679
		EBflow.preconditioned	0.346	0.0091	NA	0.673 0.669 0.683 0.669 0.910
iid	0.5	Langevin.eta1.0	0.346	0.0188	NA	0.669
	0.0	Langevin.eta0.1	0.331	0.0699	NA	0.683
		Gibbs	0.333	0.0117	NA	0.669
		CAVI	0.688	NA	NA	0.910
		EBflow	0.117	0.0222	2050	0.092
	2.0	EBflow.preconditioned	0.118	0.0156	2150	0.091
block02corr0.9		Langevin.eta1.0	0.129	0.0213	1750	0.090
		Langevin.eta0.1	0.161	0.0589	5450	0.094
		Gibbs	0.080	0.0060	1400	0.087
		CAVI	0.715	NA	NA	0.170
		EBflow	0.285	0.0276	NA	0.235
	1.0	EBflow.preconditioned	0.289	0.0150	NA	0.233
block02corr0.9		Langevin.eta1.0	0.289	0.0171	NA	0.232
		Langevin.eta0.1	0.275	0.0985	NA	0.234
		GIDDS	0.232	0.0109 N A	NA 20	0.228
		CAVI	0.207	NA	32	0.331
		EBflow	0.342	0.0168	NA	0.431
		EBnow.preconditioned	0.332	0.0120	INA	0.429
block02corr0.9	0.5	Langevin.eta1.0	0.324	0.0147	INA NA	0.429
		Cibbs	0.303	0.1045	INA	0.440
		CAVI	0.319	0.0122 NA	INA NA	0.428
		EDflow	0.445	0.0224	7750	0.472
		EBflow proconditioned	0.189	0.0224 0.0205	6300	0.121
		Langovin etal 0	0.175	0.0200	3100	0.113
block10corr0.5	2.0	Langevin etal.0	0.175	0.0131	8450	0.114
		Gibbs	0.210	0.0015	2150	0.121
		CAVI	0.140	NA	8	0.162
		EBflow	0.201	0.0330	NA	0.224
		EBflow preconditioned	0.296	0.0000 0.0251	NA	0.224
		Langevin.eta1.0	0.293	0.0209	NA	0.216
block10corr0.5	1.0	Langevin eta0.1	0.329	0.1018	NA	0.229
		Gibbs	0.246	0.0092	NA	0.213
		CAVI	0.240	NA	NA	0.326
		EBflow	0.351	0.0292	NA	0.525
		EBflow.preconditioned	0.339	0.0153	NA	0.518
11 110 07	0-	Langevin.eta1.0	0.341	0.0248	NA	0.515
block10corr0.5	0.5	Langevin.eta0.1	0.372	0.0749	NA	0.532
		Gibbs	0.377	0.0087	NA	0.515
		CAVI	0.836	NA	NA	0.659

Table 4: Comparison of methods: Bimodal prior.

Design	Method	gaussian		skew		cauchy		bimodal	
		Log-lik		Log-lik		Log-lik		Log-lik	
	EBflow	0.511	0.512	0.359	0.360	0.341	0.342	0.320	0.321
	EBflow.preconditioned	0.511	0.512	0.359	0.360	0.341	0.342	0.320	0.321
identity	Langevin.eta1.0	0.496	0.496	0.354	0.353	0.336	0.335	0.309	0.310
Identity	Langevin.eta0.1	0.514	0.516	0.361	0.362	0.342	0.337	0.321	0.325
	Gibbs	0.515	0.515	0.361	0.362	0.342	0.343	0.324	0.327
	CAVI	0.514	0.516	0.361	0.362	0.342	0.343	0.325	0.328
		Μ	SE	М	SE	M	SE	M	SE
	EBflow	0.587	0.588	0.237	0.237	0.407	0.407	0.338	0.337
	EBflow.preconditioned	0.584	0.585	0.235	0.235	0.400	0.400	0.332	0.331
iid	Langevin.eta1.0	0.582	0.583	0.233	0.233	0.392	0.392	0.326	0.327
nu	Langevin.eta0.1	0.581	0.580	0.233	0.233	0.398	0.401	0.337	0.327
	Gibbs	0.584	0.584	0.230	0.229	0.389	0.385	0.312	0.309
	CAVI	0.611	0.615	0.318	0.353	0.565	0.615	0.518	0.584
	EBflow	0.532	0.532	0.118	0.118	0.262	0.262	0.244	0.244
	EBflow.preconditioned	0.529	0.529	0.117	0.117	0.259	0.259	0.240	0.240
block02corr0 0	Langevin.eta1.0	0.525	0.525	0.118	0.118	0.252	0.253	0.233	0.233
DIOCK02C0110.3	Langevin.eta0.1	0.520	0.521	0.116	0.117	0.268	0.261	0.234	0.228
	Gibbs	0.526	0.527	0.116	0.116	0.252	0.250	0.228	0.226
	CAVI	0.545	0.551	0.158	0.165	0.355	0.393	0.331	0.397
	EBflow	0.474	0.474	0.077	0.077	0.276	0.273	0.233	0.233
	EBflow.preconditioned	0.472	0.472	0.077	0.077	0.272	0.271	0.228	0.228
block10corr0 5	Langevin.eta1.0	0.478	0.476	0.076	0.075	0.269	0.266	0.219	0.219
DIOCKTOCOTTO.5	Langevin.eta0.1	0.489	0.486	0.076	0.077	0.304	0.283	0.228	0.233
	Gibbs	0.470	0.472	0.076	0.075	0.267	0.267	0.213	0.212
	CAVI	0.513	0.645	0.082	0.081	0.390	0.488	0.326	0.324

Table 5: Comparison of results with (left) and without (right) spline smoothing

Table 5 compares results with and without spline smoothing, for a single run of each method at dimensions n = p = 1000. The smoothing regularization (2.24) is needed to obtain accurate prior density estimates, but we observe that marginal log-likelihood and prediction MSE are nearly identical with and without smoothing for all methods except CAVI (where the prediction MSE without smoothing is sometimes worse).

We summarize here a few high-level trends and take-aways from these results, many of which are also visually apparent in Figure 2:

- Mean TV-accuracy of the prior density is similar for EBflow with and without preconditioning. However, preconditioning can reduce Monte Carlo variance across independent trials, especially for more correlated designs.
- Mean TV-accuracy of the prior density for EBflow is competitive with the best tunings of Langevin-MCEM and Gibbs-MCEM for the Gaussian, Cauchy, and skew priors, and in many settings EBflow yields the lowest error.
- For the bimodal prior and many of the tested designs, no method obtains an accurate prior density estimate with TV error less than 0.2. In settings where an accurate estimate is obtained, Gibbs-MCEM produces the best estimates, likely because it more rapidly mixes between the modes of the posterior distribution in comparison to Langevin dynamics, even when the latter is carried out in the smoothed parameter φ .
- In terms of iteration count, EBflow has competitive runtime with Langevin-MCEM and Gibbs-MCEM.
- CAVI is the fastest method, and is accurate under the identity design $\mathbf{X} = \text{Id}$ (where the variational approximation is exact) as well as several settings of i.i.d. Gaussian design. However,

its accuracy in terms of both the prior density estimate and prediction MSE can substantially degrade under correlated designs.

• All methods achieve similar marginal log-likelihood under identity design, and the MCEM methods achieve similar prediction MSE under random designs. We do note that Langevin-MCEM and EBflow with final step size $\eta_t^{\varphi} = 0.1$ have higher per-variable autocorrelation than with step size $\eta_t^{\varphi} = 1.0$ and than Gibbs sampling, resulting in the need to use more posterior samples to obtain an accurate estimate of $\mathbb{E}[\boldsymbol{\theta} \mid \mathbf{X}, \mathbf{y}]$.

5 Discussion

We have studied empirical Bayes estimation of an unknown prior distribution for the regression coefficients of a linear model, introducing a system of bivariate gradient flow equations for optimizing the marginal log-likelihood via its Gibbs variational representation. This gradient flow system may be simulated using a Langevin diffusion, where the prior is continuously updated based on an estimate of the average marginal distribution of coordinates of the Langevin sample. Although the regression model and Langevin sample are both high-dimensional, the one-dimensional nature of the estimand g_* enables its estimation using a fully nonparametric approach. Similar ideas can be explored in models beyond linear regression, where a high-dimensional latent parameter is comprised of independent draws from an unknown univariate or low-dimensional prior law.

We believe there are many interesting directions for future work, and we discuss here two of these directions:

1. (Global landscape analysis) For general regression designs, we believe the optimization landscape of the marginal log-likelihood $\bar{F}_n(g)$ may have multiple local minimizers, and we find empirically that the prior density estimated by EBflow is sometimes sensitive to the choice of initialization. In such settings, we have only provided a theoretical guarantee of convergence to a prior g that satisfies an approximate first-order optimality condition for $\bar{F}_n(g)$.

It is an open question to obtain a better understanding of the global landscape geometry of $\bar{F}_n(g)$, and to characterize possible conditions under which the local minimizer of $\bar{F}_n(g)$ may be unique. One interesting setting is that of random "mean-field" designs, in which $\bar{F}_n(g)$ may asymptotically converge to a deterministic limit functional $\bar{F}(g)$ in the sense of Proposition 3.8. Although simpler algorithms may be available for empirical Bayes estimation in mean-field contexts, we believe that nonetheless this is a natural setting to study questions about the optimization landscape, as they may be studied via the limit functional $\bar{F}(g)$.

2. (Time discretization and propagation-of-chaos) In this work we have studied the dynamics and convergence of the idealized continuous-time gradient flow, but have not yet taken the steps to translate this to convergence guarantees for the time-discretized EBflow algorithm. We believe that a careful analysis of the EBflow algorithm may lead to improved understanding of issues related to parameter tuning, such as the choice of variance τ^2 in our reparametrization by the smoothed regression variable φ .

One particular question of interest is to understand conditions under which the true law \bar{q}_t may be consistently estimated by the empirical average of coordinates (2.18) over a finite time horizon [0, T], using only one realization $\{\varphi_t\}_{t\in[0,T]}$ of the Langevin diffusion rather than an asymptotically growing number of independent parallel chains. We note that this type of "single-chain propagation-of-chaos" result is only possible in settings of growing dimensions $p \to \infty$, and we believe that its study may lead to improved understanding of functional inequalities and concentration of measure for the law $\{q_t\}_{t\in[0,T]}$ as it evolves along the gradient flow.

Our numerical simulations suggest that EBflow is competitive in many settings with alternative approaches such as MCEM with coordinate-wise Gibbs sampling, in terms of both estimation accuracy and runtime. More importantly, we hope that this EBflow approach can make available a broader toolbox of ideas related to Langevin-based algorithms for empirical Bayes estimation, including ideas around adaptive preconditioning [GC11] and underdamped/Hamiltonian dynamics [Nea11, CCBJ18] to handle more ill-conditioned designs, and stochastic gradient approximations [WT11, MCF15] to scale to larger-(n, p) problems. In extensions to more complex models where latent parameters are organized in multi-layer architectures, such gradient-based ideas rather than coordinate-based computations may become even more advantageous computationally, due to the lower complexity of computing gradients using a back-propagation procedure. We believe that such extensions to more complex models and sampling algorithms would be interesting to explore also in future work.

A Proof of consistency

In this appendix, we prove Theorem 3.1 and Proposition 3.2.

A.1 TV lower bound

For any $\eta > 0$ and $g \in \mathcal{P}(M)$, we write as shorthand $g_{\eta} = \mathcal{N}_{\eta} * g = \mathcal{N}(0, \eta^2) * g$. Let

$$d_{\eta}(g,g') = W_1(g_{\eta},g'_{\eta}) \tag{A.1}$$

be the smoothed Wasserstein-1 distance on $\mathcal{P}(M)$.

All constants in the lemmas and proofs may depend on the noise variance $\sigma^2 > 0$, on M defining the common support of \mathcal{P} , on the bound γ for n/p, and on C_*, c_*, α, β in Assumption A.

Lemma A.1. There exists a sufficiently large constant v > 0, constants c, c' > 0, and a sequence $\delta_n \to 0$ such that for all large n, p and all $g, g' \in \mathcal{P}(M)$,

$$d_{\rm TV}(P_g(\mathbf{y}), P_{g'}(\mathbf{y})) \ge 1 - 2e^{-cp(d_v(g,g') - \delta_n)_+^2}.$$

Proof. Fixing any constant $\eta > 0$, the smoothed distribution g_{η} satisfies an LSI with uniformly bounded LSI constant for all $g \in \mathcal{P}(M)$ [BGMZ18, Theorem 1.1]. Note that convolving g with \mathcal{N}_{η} corresponds to adding an independent $\mathcal{N}(0, \eta^2 \mathbf{X} \mathbf{X}^{\top})$ noise vector to \mathbf{y} . Hence by the data processing inequality,

$$d_{\mathrm{TV}}(P_g(\mathbf{y}), P_{g'}(\mathbf{y})) \ge d_{\mathrm{TV}}(P_{g_{\eta}}(\mathbf{y}), P_{g'_{\eta}}(\mathbf{y})),$$

and it suffices to lower bound the TV distance on the right side for priors g_{η} and g'_{η} .

Noting that g_{η} has variance at most $M^2 + \eta^2$ for every $g \in \mathcal{P}(M)$, we pick a constant v > 0 satisfying

$$v^{2} \geq \max_{j \in \mathcal{S}} \left(\eta^{2} + v_{j} \cdot (M^{2} + \eta^{2}) + w_{j} \right), \qquad \text{where } v_{j} = \sum_{k:k \neq j} (\mathbf{z}_{j}^{\top} \mathbf{x}_{k})^{2}, \qquad w_{j} = \sigma^{2} \|\mathbf{z}_{j}\|_{2}^{2}, \quad (A.2)$$

and $\{\mathbf{z}_j\}_{j\in\mathcal{S}}$ are given by Assumption A. We will check below that v_j, w_j are bounded above by a constant uniformly over $j \in \mathcal{S}$, so that we may indeed pick v here as a constant independent of n, p. Denote $\Delta = d_v(g, g') = W_1(g_v, g'_v)$. To establish the desired TV lower bound, it suffices to exhibit an explicit test statistic $T(\mathbf{y})$ with rejection threshold $t = (\mathbb{E}_{g_\eta}[T(\mathbf{y})] + \mathbb{E}_{g'_\eta}[T(\mathbf{y})])/2$ such that

$$\max(\mathbb{P}_{g_{\eta}}[T(\mathbf{y}) < t], \mathbb{P}_{g'_{\eta}}[T(\mathbf{y}) \ge t]) \le e^{-cp(\Delta - \delta_n)^2_+}.$$
(A.3)

Indeed, this immediately implies the conclusion of the lemma,

$$d_{\mathrm{TV}}(P_{g_{\eta}}(\mathbf{y}), P_{g'_{\eta}}(\mathbf{y})) \ge \left| \mathbb{P}_{g_{\eta}}[T(\mathbf{y}) \ge t] - \mathbb{P}_{g'_{\eta}}[T(\mathbf{y}) \ge t] \right| \ge 1 - 2e^{-cp(\Delta - \delta_n)^2_+}.$$

Let κ_1, κ_2 and κ'_1, κ'_2 denote the means and variances of g_η, g'_η respectively. To construct $T(\mathbf{y})$ satisfying (A.3), we consider three cases. The constants C, c > 0 below will change from instance to instance.

Case 1: $|\kappa_1 - \kappa'_1| > c_1 \Delta$ for a constant $c_1 > 0$ to be specified. Suppose without loss of generality $\kappa_1 > \kappa'_1$, and consider the test statistic

$$T(\mathbf{y}) = \mathbf{1}^{\top} \mathbf{X}^{\top} \mathbf{y} = \mathbf{1}^{\top} \mathbf{X}^{\top} \mathbf{X} \boldsymbol{\theta} + \mathbf{1}^{\top} \mathbf{X}^{\top} \boldsymbol{\varepsilon}.$$

Under Assumption A.2, we have

$$\mathbb{E}_{g_{\eta}}[T(\mathbf{y})] - \mathbb{E}_{g'_{\eta}}[T(\mathbf{y})] = \mathbf{1}^{\top} \mathbf{X}^{\top} \mathbf{X} \mathbf{1} \cdot (\kappa_1 - \kappa'_1) \ge c \Delta p.$$
(A.4)

Computing the gradient of $T(\mathbf{y})$ in $(\boldsymbol{\theta}, \boldsymbol{\varepsilon})$, we observe that $T(\mathbf{y})$ is *L*-Lipschitz in $(\boldsymbol{\theta}, \boldsymbol{\varepsilon})$ with respect to the ℓ_2 -norm, where $L^2 = \mathbf{1}^{\top} (\mathbf{X}^{\top} \mathbf{X})^2 \mathbf{1} + \mathbf{1}^{\top} \mathbf{X}^{\top} \mathbf{X} \mathbf{1} \leq Cp$ under Assumption A.1–2. Then by the LSI for $(\boldsymbol{\theta}, \boldsymbol{\varepsilon})$ under both g_{η} and g'_{η} , for any s > 0, $T(\mathbf{y})$ has the sub-Gaussian concentration arounds its mean

$$\max\left(\mathbb{P}_{g_{\eta}}[T(\mathbf{y}) - \mathbb{E}_{g_{\eta}}T(\mathbf{y}) \le -s\Delta p], \ \mathbb{P}_{g'_{\eta}}[T(\mathbf{y}) - \mathbb{E}_{g'_{\eta}}T(\mathbf{y}) \ge s\Delta p]\right) \le e^{-cs^2\Delta^2 p}.$$

Choosing a constant s > 0 small enough and combining with (A.4), this shows that (A.3) holds with $\delta_n = 0$.

Case 2: $|\kappa_2 - \kappa'_2| \ge c_2 \max(\Delta, \Delta^2)$ for a constant $c_2 > 0$ to be specified. Suppose without loss of generality $\kappa_2 > \kappa'_2$, let Π denote the projection orthogonal to **X1**, and consider the test statistic

$$T(\mathbf{y}) = \|\mathbf{X}^{\top}\Pi\mathbf{y}\|^{2} = \|\mathbf{X}^{\top}\Pi(\mathbf{X}\boldsymbol{\theta} + \boldsymbol{\varepsilon})\|^{2} = \begin{pmatrix} \boldsymbol{\theta} \\ \boldsymbol{\varepsilon} \end{pmatrix}^{\top} \underbrace{\begin{pmatrix} \mathbf{X}^{\top}\Pi\mathbf{X}\mathbf{X}^{\top}\Pi\mathbf{X} & \mathbf{X}^{\top}\Pi\mathbf{X}\mathbf{X}^{\top}\Pi \\ \Pi\mathbf{X}\mathbf{X}^{\top}\Pi\mathbf{X} & \Pi\mathbf{X}\mathbf{X}^{\top}\Pi \\ \vdots = M(\mathbf{X},\Pi) \end{pmatrix}}_{:=M(\mathbf{X},\Pi)} \begin{pmatrix} \boldsymbol{\theta} \\ \boldsymbol{\varepsilon} \end{pmatrix}$$

Under Assumptions A.1–2, the matrix $\mathbf{X}^{\top}\Pi\mathbf{X}$ has at least cp eigenvalues greater than c, for some sufficiently small constant c > 0. Then, irrespective of the means κ_1, κ'_1 , we have

$$\mathbb{E}_{G_{\eta}}[T(\mathbf{y})] - \mathbb{E}_{G'_{\eta}}[T(\mathbf{y})] = \operatorname{Tr} \mathbf{X}^{\top} \Pi \mathbf{X} \mathbf{X}^{\top} \Pi \mathbf{X} \cdot (\kappa_{2} - \kappa'_{2}) \ge c \max(\Delta, \Delta^{2}) p.$$

We have also $||M(\mathbf{X}, \Pi)||_{\text{op}} \leq C$ and $||M(\mathbf{X}, \Pi)||_{\text{F}}^2 \leq Cp$ for the above matrix $M(\mathbf{X}, \Pi)$, the latter holding because each block of $M(\mathbf{X}, \Pi)$ has rank at most p. Then by the Hanson-Wright inequality [RV13, Theorem 1.1], viewing $T(\mathbf{y})$ as a quadratic form in $(\boldsymbol{\theta}, \boldsymbol{\varepsilon})$ where $\boldsymbol{\theta}$ is sub-Gaussian under both g_{η} and g'_{η} , for any s > 0,

$$\mathbb{P}_{g_{\eta}}[T(\mathbf{y}) - \mathbb{E}_{g_{\eta}}T(\mathbf{y}) \le -s\max(\Delta, \Delta^2)p] \le e^{-c\min(s,s^2)\Delta^2 p}$$
$$\mathbb{P}_{g'_{\eta}}[T(\mathbf{y}) - \mathbb{E}_{g'_{\eta}}T(\mathbf{y}) \ge s\max(\Delta, \Delta^2)p] \le e^{-c\min(s,s^2)\Delta^2 p}.$$

Then (A.3) holds also in this case with $\delta_n = 0$.

Case 3: $|\kappa_1 - \kappa'_1| < c_1 \Delta$ and $|\kappa_2 - \kappa'_2| < c_2 \max(\Delta, \Delta^2)$. Let $S \subseteq \{1, \ldots, p\}$ and $\{\mathbf{z}_j : j \in S\}$ be as in Assumption A. For each $j \in S$, consider

$$s_j = \mathbf{z}_j^\top \mathbf{y} = \underbrace{\mathbf{z}_j^\top \mathbf{x}_j \cdot \theta_j}_{\equiv a_j} + \underbrace{\sum_{k:k \neq j} \mathbf{z}_j^\top \mathbf{x}_k \cdot \theta_k}_{\equiv b_j} + \underbrace{\mathbf{z}_j^\top \varepsilon}_{\equiv e_j},$$

where these three terms a_j, b_j, e_j are independent. Let $\mathcal{L}(f(\theta))$ denote the law of $f(\theta)$ under g_{η} . Then by Assumption A.3,

$$W_1(\mathcal{L}(a_j), g_\eta) = W_1(\mathcal{L}(a_j), \mathcal{L}(\theta_j)) \le |\mathbf{z}_j^\top \mathbf{x}_j - 1| \cdot \mathbb{E}_{g_\eta}[|\theta_j|] \to 0.$$

For b_j , we have

$$\mathbb{E}_{g_{\eta}}[b_j] = m_j \kappa_1, \qquad \operatorname{Var}_{g_{\eta}}[b_j] = v_j \kappa_2, \qquad \text{where } m_j = \sum_{k:k \neq j} \mathbf{z}_j^\top \mathbf{x}_k, \qquad v_j = \sum_{k:k \neq j} (\mathbf{z}_j^\top \mathbf{x}_k)^2.$$

The Wasserstein- $(2 + \beta)$ CLT of [Rio09, Eq. (4.1)] shows that for some $c = c(\beta) > 0$,

$$W_{2+\beta}\Big(\mathcal{L}(b_j), \mathcal{N}(m_j\kappa_1, v_j\kappa_2)\Big)^{2+\beta} \le c \sum_{k:k\neq j} |\mathbf{z}_j^\top \mathbf{x}_k|^{2+\beta} \cdot \mathbb{E}_{g_\eta}[|\theta_k|^{2+\beta}].$$

Assumption A.3 provides the Lyapunov condition ensuring that the right side converges to 0 as $n, p \to \infty$, uniformly over $j \in S$. Then also

$$W_1\Big(\mathcal{L}(b_j), \mathcal{N}(m_j\kappa_1, v_j\kappa_2)\Big) \leq W_{2+\beta}\Big(\mathcal{L}(b_j), \mathcal{N}(m_j\kappa_1, v_j\kappa_2)\Big) \to 0.$$

Finally, $e_j \sim \mathcal{N}(0, w_j)$ with variance $w_j = \sigma^2 \|\mathbf{z}_j\|_2^2$. Putting this together,

$$W_1\Big(\mathcal{L}(s_j), \, g_\eta * \mathcal{N}(m_j\kappa_1, v_j\kappa_2 + w_j)\Big) \le \delta_n/4 \tag{A.5}$$

for some $\delta_n \to 0$. Similarly, writing $\mathcal{L}'(\cdot)$ for the law under g'_{η} ,

$$W_1\Big(\mathcal{L}'(s_j), \, g'_\eta * \mathcal{N}(m_j\kappa'_1, v_j\kappa'_2 + w_j)\Big) \le \delta_n/4. \tag{A.6}$$

We use $|\kappa_1 - \kappa'_1| < c_1 \Delta$ and $|\kappa_2 - \kappa'_2| < c_2 \max(\Delta, \Delta^2)$ for some sufficiently small $c_1, c_2 > 0$ to compare these normal distributions. Note that for any two univariate normal laws, by the coupling $(\mu + \sigma Z, \mu' + \sigma' Z)$ for $Z \sim \mathcal{N}(0, 1)$,

$$W_{1}(\mathcal{N}(\mu, \sigma^{2}), \mathcal{N}(\mu', {\sigma'}^{2})) \leq |\mu - \mu'| + |\sigma - \sigma'| \cdot \mathbb{E}[|Z|] \\ \leq |\mu - \mu'| + C \min\left(|\sigma^{2} - {\sigma'}^{2}|, |\sigma^{2} - {\sigma'}^{2}|^{1/2}\right),$$

the second inequality holding when σ^2 , ${\sigma'}^2$ are lower bounded by a constant. By the definitions of m_j, v_j and under Assumption A.3, we have $|m_j| = |\mathbf{z}_j^\top \mathbf{X} \mathbf{1} - \mathbf{z}_j^\top \mathbf{x}_j| \le C$, $v_j \le \mathbf{z}_j^\top \mathbf{X} \mathbf{X}^\top \mathbf{z}_j \le C$, and $w_j = \sigma^2 ||\mathbf{z}_j||_2^2 \in [c, C]$ for some constants C, c > 0. Then for a sufficiently small choice of constants $c_1, c_2 > 0$, we have

$$W_1(\mathcal{N}(m_j\kappa_1, v_j\kappa_2 + w_j), \mathcal{N}(m_j\kappa_1', v_j\kappa_2' + w_j)) < \Delta/2.$$

Combining with (A.6), this implies

$$W_1\Big(\mathcal{L}'(s_j), \, g'_\eta * \mathcal{N}(m_j\kappa_1, v_j\kappa_2 + w_j)\Big) \le \delta_n/4 + \Delta/2. \tag{A.7}$$

Then by (A.5), (A.7), and the triangle inequality and translation invariance for W_1 ,

$$W_1(\mathcal{L}(s_j), \mathcal{L}'(s_j)) \ge W_1\left(g_\eta * \mathcal{N}(m_j\kappa_1, v_j\kappa_2 + w_j), g'_\eta * \mathcal{N}(m_j\kappa_1, v_j\kappa_2 + w_j)\right) - \delta_n/2 - \Delta/2$$
$$= W_1\left(g_{\sqrt{\eta^2 + v_j\kappa_2 + w_j}}, g'_{\sqrt{\eta^2 + v_j\kappa_2 + w_j}}\right) - \delta_n/2 - \Delta/2.$$

Recalling the choice of v in (A.2) and applying $\kappa_2 \leq M^2 + \eta^2$, we have $\eta^2 + v_j \kappa_2 + w_j \leq v^2$. Then by monotonicity of the smoothed Wasserstein distance in the smoothing parameter [GG20, Theorem 3], $W_1(g_{\sqrt{\eta^2 + v_j \kappa_2 + w_j}}, g'_{\sqrt{\eta^2 + v_j \kappa_2 + w_j}}) \geq W_1(g_v, g'_v) = \Delta$, so

$$W_1(\mathcal{L}(s_j), \mathcal{L}'(s_j)) \ge \Delta/2 - \delta_n/2$$
Finally, this implies by the Kantorovich duality for W_1 that there exists for each $j \in \mathcal{S}$ a smooth 1-Lipschitz test function $f_j : \mathbb{R} \to \mathbb{R}$ such that $\mathbb{E}_{g_\eta}[f_j(s_j)] - \mathbb{E}_{g'_\eta}[f_j(s_j)] \ge (\Delta - \delta_n)_+/3$. We consider the test statistic

$$T(\mathbf{y}) = \sum_{j \in \mathcal{S}} f_j(s_j) = \sum_{j \in \mathcal{S}} f_j(\mathbf{z}_j^\top (\mathbf{X} \boldsymbol{\theta} + \boldsymbol{\varepsilon})),$$

which then satisfies $\mathbb{E}_{g_{\eta}}[T(\mathbf{y})] - \mathbb{E}_{g'_{\eta}}[T(\mathbf{y})] \ge cp(\Delta - \delta_n)_+$ because $|\mathcal{S}| \ge \alpha p$. Computing the gradient in $(\boldsymbol{\theta}, \boldsymbol{\varepsilon})$, we see that $T(\mathbf{y})$ is *L*-Lipschitz in $(\boldsymbol{\theta}, \boldsymbol{\varepsilon})$ for

$$L^{2} = \left\| \mathbf{X}^{\top} \sum_{j \in \mathcal{S}} \mathbf{z}_{j} \cdot f_{j}'(s_{j}) \right\|^{2} + \left\| \sum_{j \in \mathcal{S}} \mathbf{z}_{j} \cdot f_{j}'(s_{j}) \right\|^{2}.$$

Applying $|f'_j(s_j)| \leq 1$ for all $j \in S$, $\|\mathbf{X}\|_{\text{op}} \leq C_*$, and the final condition of Assumption A.4, this gives $L^2 \leq Cp$. Then (A.3) holds by the LSI for $(\boldsymbol{\theta}, \boldsymbol{\varepsilon})$, similarly to Case 1.

A direct consequence of Lemma A.1 is the following pointwise comparison of the likelihood objective.

Corollary A.2. Let v > 0 and the sequence $\delta_n \downarrow 0$ be as in Lemma A.1. Fix any constant $\delta > 0$. Then there exists some constants $c = c(\delta)$ and $\tau = \tau(\delta)$ such that for all large n, p and any $g \in \mathcal{P}(M)$ with $d_v(g, g_*) \ge \delta$,

$$\mathbb{P}_{g_*}\left[\frac{1}{p}\left(\log P_g(\mathbf{y}) - \log P_{g_*}(\mathbf{y})\right) \ge -\tau\right] \le e^{-cp}.$$

Proof. Writing as shorthand $P_g = P_g(\mathbf{y})$, Lemma A.1 shows that for some constant c > 0, all large n, p, and any $g \in \mathcal{P}(M)$ with $d_v(g, g_*) \ge \delta$, we have

$$e^{-cp\delta^{2}} \ge 1 - d_{\mathrm{TV}}(P_{g_{*}}, P_{g}) = \int \min(P_{g_{*}}, P_{g}) \mathrm{d}\mathbf{y}$$
$$\ge \int e^{-p\tau} P_{g_{*}} \mathbf{1}\{P_{g} \ge e^{-p\tau} P_{g_{*}}\} \mathrm{d}\mathbf{y} = e^{-p\tau} \mathbb{P}_{g_{*}}\left[p^{-1}\left(\log P_{g} - \log P_{g_{*}}\right) \ge -\tau\right]$$

for any $\tau > 0$. The corollary follows upon choosing $\tau = c\delta^2/2$ and adjusting constants.

A.2 Covering net

We now define the following pseudo-metric $\tilde{d}_{\eta,B}$ over $\mathcal{P}(M)$: Let $G_{\eta}(\cdot), G'_{\eta}(\cdot)$ denote the CDFs of $g_{\eta} = \mathcal{N}_{\eta} * g$ and $g'_{\eta} = \mathcal{N}_{\eta} * g'$. (Here G'_{η} is not to be confused with a derivative of G_{η} .) We set

$$\widetilde{d}_{\eta,B}(g,g') = \max_{x:|x| \le B} |G_{\eta}(x) - G'_{\eta}(x)|.$$

Lemma A.3. Fix any constants $C_0, \iota > 0$. Let Π_X be the projection onto the column span of **X**. Then there exist constants $\eta, B, L > 0$ such that, on the event where $\|\Pi_X \mathbf{y}\|_2^2 \leq C_0 p$, for any $g, g' \in \mathcal{P}(M)$ we have

$$\frac{1}{p} \Big| \log P_g(\mathbf{y}) - \log P_{g'}(\mathbf{y}) \Big| \le L \,\widetilde{d}_{\eta,B}(g,g') + \iota.$$

Proof. Let $[\mathbf{V}_X | \mathbf{W}_X] \in \mathbb{R}^{n \times n}$ be an orthogonal matrix where $\Pi_X = \mathbf{V}_X \mathbf{V}_X^{\top}$ and the columns of \mathbf{V}_X form an orthonormal basis for the column span of \mathbf{X} . Write $\tilde{\mathbf{y}} = \mathbf{V}_X^{\top} \mathbf{y}$ and $\tilde{\mathbf{X}} = \mathbf{V}_X^{\top} \mathbf{X}$. Then $\tilde{\mathbf{y}} = \tilde{\mathbf{X}} \boldsymbol{\theta} + \mathbf{V}_X^{\top} \boldsymbol{\varepsilon}$ is independent of $\mathbf{W}_X^{\top} \mathbf{y} = \mathbf{W}_X^{\top} \boldsymbol{\varepsilon}$, and the law of the latter does not depend on the prior g. Thus

$$\log P_g(\mathbf{y}) = \log P_g(\widetilde{\mathbf{y}}) + \text{constant}$$

for a constant independent of g, where $P_g(\tilde{\mathbf{y}})$ denotes the marginal density of $\tilde{\mathbf{y}}$ under g. Observe that under Assumption A.1, $\|\tilde{\mathbf{X}}\tilde{\mathbf{X}}^{\top}\|_{\text{op}} = \|\mathbf{X}\mathbf{X}^{\top}\|_{\text{op}} < C$ for a constant C > 0. Then we may choose a constant $\eta > 0$ small enough such that $\eta \cdot \|\tilde{\mathbf{X}}\tilde{\mathbf{X}}^{\top}\|_{\text{op}} < \sigma^2/2$, and reparametrize the law of $\tilde{\mathbf{y}}$ equivalently as

$$\widetilde{\mathbf{y}} = \widetilde{\mathbf{X}} \boldsymbol{\varphi} + \widetilde{\boldsymbol{\varepsilon}}, \qquad \varphi_j \overset{iid}{\sim} g_{\eta}, \qquad \widetilde{\boldsymbol{\varepsilon}} \sim \mathcal{N}(0, \boldsymbol{\Sigma}),$$

where $\Sigma = \sigma^2 \operatorname{Id} - \eta \widetilde{\mathbf{X}} \widetilde{\mathbf{X}}^{\top}$ satisfies $(\sigma^2/2) \operatorname{Id} \preceq \Sigma \preceq \sigma^2 \operatorname{Id}$. The marginal log-likelihood log $P_g(\mathbf{y})$ then has the equivalent form

$$\log P_g(\mathbf{y}) = \log \int \exp\left(-\frac{1}{2}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}\boldsymbol{\varphi})^{\top} \boldsymbol{\Sigma}^{-1}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}\boldsymbol{\varphi})\right) \prod_{j=1}^p g_\eta(\varphi_j) \mathrm{d}\boldsymbol{\varphi} + \text{constant}$$
(A.8)

for a constant independent of g.

We now perform a change-of-measure from G_{η} to G'_{η} by introducing $F_{\eta}(x) = {G'_{\eta}}^{-1}(G_{\eta}(x))$ where ${G'_{\eta}}^{-1}(\cdot)$ is the functional inverse of the CDF $G'_{\eta}(\cdot)$. (This is the univariate optimal transport map from G_{η} to G'_{η} .) Let us write $F_{\eta}(\varphi)$ for the coordinate-wise application of F_{η} to φ . Then the marginal log-likelihood log $P_{g'}(\mathbf{y})$ may also be written as

$$\log P_{g'}(\mathbf{y}) = \log \int \exp\left(-\frac{1}{2}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}F_{\eta}(\boldsymbol{\varphi}))^{\top} \mathbf{\Sigma}^{-1}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}F_{\eta}(\boldsymbol{\varphi}))\right) \prod_{j=1}^{p} g_{\eta}(\varphi_{j}) \mathrm{d}\boldsymbol{\varphi} + \text{constant.} \quad (A.9)$$

Taking the difference of (A.8) and (A.9),

$$\log P_{g'}(\mathbf{y}) - \log P_g(\mathbf{y}) = \log \left\langle e^{-\frac{1}{2}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} F_{\eta}(\boldsymbol{\varphi}))^{\top} \mathbf{\Sigma}^{-1}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} F_{\eta}(\boldsymbol{\varphi})) + \frac{1}{2}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} \boldsymbol{\varphi})^{\top} \mathbf{\Sigma}^{-1}(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} \boldsymbol{\varphi})} \right\rangle,$$

where we introduce the shorthand

$$\langle f(\boldsymbol{\varphi}) \rangle = \frac{\int f(\boldsymbol{\varphi}) e^{-H(\boldsymbol{\varphi})} \mathrm{d}G_{\eta}(\boldsymbol{\varphi})}{\int e^{-H(\boldsymbol{\varphi})} \mathrm{d}G_{\eta}(\boldsymbol{\varphi})}, \quad H(\boldsymbol{\varphi}) = \frac{1}{2} (\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}\boldsymbol{\varphi})^{\top} \boldsymbol{\Sigma}^{-1} (\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}\boldsymbol{\varphi}), \quad \mathrm{d}G_{\eta}(\boldsymbol{\varphi}) = \prod_{j=1}^{p} g_{\eta}(\varphi_{j}) \mathrm{d}\boldsymbol{\varphi}$$

for the expectation with respect to the posterior law of φ under g_{η} given $(\mathbf{X}, \mathbf{\tilde{y}})$.

Let C, c > 0 denote constants changing from instance to instance, which may depend on η . By Jensen's inequality,

$$\log P_{g'}(\mathbf{y}) - \log P_g(\mathbf{y}) \geq \frac{1}{2} \left\langle -(\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} F_{\eta}(\boldsymbol{\varphi}))^{\top} \boldsymbol{\Sigma}^{-1} (\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} F_{\eta}(\boldsymbol{\varphi})) + (\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} \boldsymbol{\varphi})^{\top} \boldsymbol{\Sigma}^{-1} (\widetilde{\mathbf{y}} - \widetilde{\mathbf{X}} \boldsymbol{\varphi}) \right\rangle.$$

Setting $\mathbf{u} = \widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}\boldsymbol{\varphi}$ and $\mathbf{v} = \widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}F_{\eta}(\boldsymbol{\varphi})$, we apply

$$\langle -\mathbf{v}^{\top} \boldsymbol{\Sigma}^{-1} \mathbf{v} + \mathbf{u}^{\top} \boldsymbol{\Sigma}^{-1} \mathbf{u} \rangle = \langle -(\mathbf{u} - \mathbf{v})^{\top} \boldsymbol{\Sigma}^{-1} (\mathbf{u} - \mathbf{v}) + 2(\mathbf{u} - \mathbf{v})^{\top} \boldsymbol{\Sigma}^{-1} \mathbf{u} \rangle$$

$$\geq -\langle (\mathbf{u} - \mathbf{v})^{\top} \boldsymbol{\Sigma}^{-1} (\mathbf{u} - \mathbf{v}) \rangle - 2\langle (\mathbf{u} - \mathbf{v})^{\top} \boldsymbol{\Sigma}^{-1} (\mathbf{u} - \mathbf{v}) \rangle^{1/2} \langle \mathbf{u}^{\top} \boldsymbol{\Sigma}^{-1} \mathbf{u} \rangle^{1/2}.$$

Together with the conditions $\Sigma^{-1} \preceq (2/\sigma^2)$ Id, $\|\widetilde{\mathbf{X}}\|_{\text{op}} \leq C$ under Assumption A.1, and Cauchy-Schwarz, this gives

$$\log P_{g'}(\mathbf{y}) - \log P_g(\mathbf{y}) \ge -C\left(\left\langle \|F_{\eta}(\boldsymbol{\varphi}) - \boldsymbol{\varphi}\|_2^2 \right\rangle + \left\langle \|F_{\eta}(\boldsymbol{\varphi}) - \boldsymbol{\varphi}\|_2^2 \right\rangle^{1/2} \left\langle H(\boldsymbol{\varphi}) \right\rangle^{1/2} \right).$$
(A.10)

To lower bound (A.10), we may first bound $\langle H(\boldsymbol{\varphi}) \rangle$ as follows: Introduce the partition function $Z = \int e^{-H(\boldsymbol{\varphi})} dG_{\eta}(\boldsymbol{\varphi})$. Applying $H(\boldsymbol{\varphi}) \geq 0$, $e^{-H(\boldsymbol{\varphi})}/Z \leq 1$ on the event $e^{-H(\boldsymbol{\varphi})} \leq Z$, and $H(\boldsymbol{\varphi}) < -\log Z$ on the complementary event $e^{-H(\boldsymbol{\varphi})} > Z$, we have

$$\langle H(\boldsymbol{\varphi})\rangle = \int H(\boldsymbol{\varphi}) \frac{e^{-H(\boldsymbol{\varphi})}}{Z} \mathrm{d}G_{\eta}(\boldsymbol{\varphi}) \leq \int H(\boldsymbol{\varphi}) \mathrm{d}G_{\eta}(\boldsymbol{\varphi}) + \max(0, -\log Z) \leq 2 \int H(\boldsymbol{\varphi}) \mathrm{d}G_{\eta}(\boldsymbol{\varphi}),$$

the last step applying the Jensen's inequality lower bound $\log Z \ge -\int H(\varphi) dG_{\eta}(\varphi)$. Thus we have bounded the mean of $H(\varphi)$ under the posterior by its mean under the prior. On the event $\|\widetilde{\mathbf{y}}\|_2^2 = \|\Pi_X \mathbf{y}\|_2^2 \le C_0 p$, applying $\Sigma^{-1} \preceq (2/\sigma^2) \operatorname{Id}$, $\|\widetilde{\mathbf{X}}\|_{\operatorname{op}} \le C$, and that g_{η} is sub-Gaussian (since g has bounded support), we have $\int H(\varphi) dG_{\eta}(\varphi) \le Cp$. This implies

$$\langle H(\boldsymbol{\varphi}) \rangle \le Cp.$$
 (A.11)

To bound $\langle ||F_{\eta}(\varphi) - \varphi||_{2}^{2} \rangle$, let us fix a small constant t > 0 to be determined, pick B = B(t) > 0sufficiently large, and define $N_{B}(\varphi) = |\{j : |\varphi_{j}| > B\}|$. Applying again $\log Z \ge -\int H(\varphi) dG_{\eta}(\varphi) \ge -Cp$ and $H(\varphi) \ge 0$, we have

$$\langle \mathbf{1}\{N_B(\boldsymbol{\varphi}) > tp\} \rangle = \int \mathbf{1}\{N_B(\boldsymbol{\varphi}) > tp\} \frac{e^{-H(\boldsymbol{\varphi})}}{Z} \mathrm{d}G_{\eta}(\boldsymbol{\varphi}) \le e^{Cp} \int \mathbf{1}\{N_B(\boldsymbol{\varphi}) > tp\} \mathrm{d}G_{\eta}(\boldsymbol{\varphi}).$$

Under the product prior law $\prod_{j=1}^{p} g_{\eta}(\varphi_j)$, this quantity $N_B(\varphi)$ is a binomial random variable whose success probability is at most $q = 2e^{-cB^2}$ for a constant $c := c(M, \eta)$, by sub-Gaussianity of g_{η} . For any fixed constant t > 0, by choosing B = B(t) > 0 large enough, we have $q \le t/2$, so by the Chernoff bound for the binomial distribution [BLM13, pp. 24]

$$\int \mathbf{1}\{N_B(\boldsymbol{\varphi}) > tp\} \mathrm{d}G_{\eta}(\boldsymbol{\varphi}) \le \mathbb{P}_{g_{\eta}}\left(\mathrm{Bin}(p,q) - pq \ge \frac{t}{2}p\right) \le e^{-ph_q(q+\frac{t}{2})},$$

where $h_q(a) = \operatorname{KL}(\operatorname{Ber}(a)||\operatorname{Ber}(q))$. Since $h_q(q+t/2) \to \infty$ as $q \downarrow 0$, we may then choose B = B(t)large enough such that $\int \mathbf{1}\{N_B(\boldsymbol{\varphi}) > tp\} dG_\eta(\boldsymbol{\varphi}) \leq e^{-2Cp}$. Consequently, $\langle \mathbf{1}\{N_B(\boldsymbol{\varphi}) > tp\} \rangle \leq e^{Cp} \cdot e^{-2Cp} \leq e^{-Cp}$, and hence

$$\langle N_B(\boldsymbol{\varphi}) \rangle \le tp + p \cdot \langle \mathbf{1}\{N_B(\boldsymbol{\varphi}) > tp\} \rangle \le 2tp.$$
 (A.12)

Now note that for all $g' \in \mathcal{P}$, the smoothed distributions g'_{η} have densities lower-bounded by a η -dependent constant on any fixed compact interval, and hence the inverse CDFs ${G'_{\eta}}^{-1}(\cdot)$ are uniformly Lipschitz-continuous over any fixed compact sub-interval of (0, 1). Furthermore, for the above choice of B = B(t) > 0, the interval $[G_{\eta}(-B), G_{\eta}(B)]$ is contained in a common compact sub-interval of (0, 1) across all $g \in \mathcal{P}(M)$. Then there is a constant $C_1 = C_1(t) > 0$ such that for all $g, g' \in \mathcal{P}, {G'_{\eta}}^{-1}$ is C_1 -Lipschitz over the interval $[G_{\eta}(-B), G_{\eta}(B)]$. This shows that if $x \in [-B, B]$,

$$|F_{\eta}(x) - x| = |G'_{\eta}^{-1}(G_{\eta}(x)) - G'_{\eta}^{-1}(G'_{\eta}(x))| \le C_1|G_{\eta}(x) - G'_{\eta}(x)| \le C_1\widetilde{d}_{\eta,B}(g,g').$$
(A.13)

Note that if $\theta \sim g$, $\theta' \sim g'$ with $g, g' \in \mathcal{P}(M)$ and $\varepsilon \sim \mathcal{N}(0, \eta^2)$ are independent, then also $\mathbb{P}[\theta' + \varepsilon \leq x - 2M] \leq \mathbb{P}[\theta + \varepsilon \leq x] \leq \mathbb{P}[\theta' + \varepsilon \leq x + 2M]$ for every $x \in \mathbb{R}$ because $|\theta - \theta'| \leq 2M$. Then $G'_{\eta}(x - 2M) \leq G_{\eta}(x) \leq G'_{\eta}(x + 2M)$, implying that

$$|F_{\eta}(x) - x| = |G'_{\eta}^{-1}(G_{\eta}(x)) - x| \le 2M$$
(A.14)

for all $x \in \mathbb{R}$. Combining these bounds (A.13) and (A.14) and recalling $N_B(\varphi) = |\{j : |\varphi_j| > B\}|$ which satisfies (A.12),

$$\left\langle \|F_{\eta}(\boldsymbol{\varphi}) - \boldsymbol{\varphi}\|_{2}^{2} \right\rangle \leq \left\langle C_{1}^{2} \widetilde{d}_{\eta,B}(g,g')^{2} \cdot p + (2M)^{2} \cdot N_{B}(\boldsymbol{\varphi}) \right\rangle \leq (C_{1}^{2} \widetilde{d}_{\eta,B}(g,g')^{2} + 8M^{2}t)p. \quad (A.15)$$

We may apply (A.11) and (A.15) back to (A.10), now choosing the constant t > 0 sufficiently small based on the value of ι given in the lemma, and then B = B(t), $C_1 = C_1(t)$, and L = L(t)sufficiently large, to obtain

$$\log P_{g'}(\mathbf{y}) - \log P_g(\mathbf{y}) \ge -p\Big(L \cdot \max(\widetilde{d}_{\eta,B}(g,g')^2, \widetilde{d}_{\eta,B}(g,g')) + \iota\Big).$$

This is simplified by observing that $\tilde{d}_{\eta,B}(G,G') \leq 1$ by definition. Then, applying the same lower bound with g, g' interchanged, we obtain as desired $|\log P_{g'}(\mathbf{y}) - \log P_g(\mathbf{y})| \leq p(L \cdot \tilde{d}_{\eta,B}(g,g') + \iota)$. \Box

A.3 Completing the proof

Corollary A.4. Fix any $\delta > 0$. Then there exist constants $v, \tau > 0$ and C, c > 0 such that

$$\mathbb{P}_{g_*}\left[\sup_{g\in\mathcal{P}(M):d_v(g,g_*)\geq\delta}\frac{1}{p}\left(\log P_g(\mathbf{y})-\log P_{g_*}(\mathbf{y})\right)\geq-\tau\right]\leq Ce^{-cp}.$$

Proof. In Lemma A.3, set $\iota = \tau/3$, and set $C_0 > 0$ large enough such that $\mathbb{P}_{g_*}[\|\Pi_X \mathbf{y}\|_2^2 > C_0 p] \leq e^{-cp}$. (This is possible since $\|\mathbf{X}\|_{op} \leq C$, $\boldsymbol{\theta} \in \mathbb{R}^p$ has sub-Gaussian coordinates under g_* , and $\|\Pi_X \boldsymbol{\varepsilon}\|_2^2 \sim \chi_k^2$ where $k = \operatorname{rank}(\mathbf{X}) \leq p$.) Let η, B, L be as in the lemma. Take a $(\tau/3L)$ -covering net of $\{g \in \mathcal{P}(M) : d_v(g, g_*) \geq \delta\}$ in the pseudo-metric $\tilde{d}_{\eta,B}(\cdot, \cdot)$, which by a standard packing argument has some finite cardinality $C(\eta, B, L, \tau)$ because all CDFs G_η for $g \in \mathcal{P}(M)$ have a constant lower bound for their derivatives over [-B, B]. The result then follows from Lemma A.3, combined with Corollary A.2 applied to each point in this net with $-\tau/3$ in place of $-\tau$.

Proof of Theorem 3.1. By the Borel-Cantelli lemma, the above corollary implies that for any fixed constant $\delta > 0$, the estimator \tilde{g} of Theorem 3.1 satisfies $d_v(\tilde{g}, g_*) < \delta$ almost surely for all large n, p. Since $\delta > 0$ is arbitrary and this smoothed Wasserstein distance $d_v(\cdot, \cdot)$ metrizes weak convergence, Theorem 3.1 follows.

A.4 Assumption A for random designs

Proof of Proposition 3.2. Recall that $\Pi \in \mathbb{R}^{n \times n}$ is the projection orthogonal to $\mathbf{X1} \in \mathbb{R}^n$. For any $j \in [p]$, let

$$\mathbf{z}_j = \Pi \mathbf{X} \mathbf{\Sigma}_X^{-1} \mathbf{e}_j = \mathbf{b}_j - \mathbf{r}_j, \qquad \mathbf{b}_j = \mathbf{X} \mathbf{\Sigma}_X^{-1} \mathbf{e}_j, \qquad \mathbf{r}_j = \frac{\mathbf{X} \mathbf{1} \mathbf{1}^\top \mathbf{X}^\top}{\|\mathbf{X}\mathbf{1}\|^2} \mathbf{X} \mathbf{\Sigma}_X^{-1} \mathbf{e}_j.$$

Define the event

$$E = \left\{ \|\mathbf{X}\|_{\text{op}} \le C_0, \text{ Tr } \mathbf{X}^\top \mathbf{X} \ge c_0 p, \|\mathbf{X}\mathbf{1}\|^2 \ge c_0 p, c_0 \le \min_{j \in [p]} \|\mathbf{b}_j\|^2 \le \max_{j \in [p]} \|\mathbf{b}_j\|^2 \le C_0, \max_{j \in [p]} \|$$

On this event E, from these conditions and the decomposition $\mathbf{z}_j = \mathbf{b}_j - \mathbf{r}_j$, for a constant $C_\beta > 0$ we have

$$\max_{j \in [p]} \sum_{k: k \neq j} |\mathbf{z}_j^\top \mathbf{x}_k|^{2+\beta} \le C_\beta p \Big(\sqrt{\frac{\log p}{n}} + \frac{1}{\sqrt{p}} \Big)^{2+\beta}.$$

This approaches 0 as $n, p \to \infty$ for $n/p \ge \gamma$ and any fixed $\beta, \gamma > 0$. We have also $\mathbf{z}_j^\top \mathbf{X} \mathbf{1} = 0$ since $\Pi \mathbf{X} \mathbf{1} = 0$ by definition of Π , and it is direct to check the remaining conditions of Assumption A.1–3 on this event E. For Assumption A.4, observe that

$$\left\|\sum_{j\in\mathcal{S}}a_j\mathbf{z}_j\right\|^2 = \mathbf{a}^\top \boldsymbol{\Sigma}_X^{-1} \mathbf{X}^\top \Pi \mathbf{X} \boldsymbol{\Sigma}_X^{-1} \mathbf{a} \le \|\mathbf{a}\|^2 \cdot \left\|\boldsymbol{\Sigma}_X^{-1} \mathbf{X}^\top \Pi \mathbf{X} \boldsymbol{\Sigma}_X^{-1}\right\|_{\text{op}} \le Cp,$$

when $|a_j| \leq 1$ for all $j \in [p]$ and $\|\mathbf{X}\|_{\text{op}} \leq C_0$. Thus Assumption A.4 holds also on E.

Next we show that for some sufficiently large $C_0 > 0$ and small $c_0 > 0$, the event E holds with probability at least $1 - p^{-10}$ for all large n, p. In the following, C, C', c, c' > 0 denote constants changing from instance to instance. We have $\|\sqrt{n}\mathbf{X}\mathbf{\Sigma}_X^{-1/2}\|_{op} \leq C(\sqrt{n} + \sqrt{p})$ with probability $1 - e^{-cp}$, by [Ver18, Theorem 4.6.1]. Then, since $n/p \geq \gamma$ and $\|\mathbf{\Sigma}_X\|_{op} \leq C$, this implies $\|\mathbf{X}\|_{op} \leq C'$. For the remaining statements, observe that for any (deterministic) unit vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^p$, each of the products $\mathbf{X}\mathbf{u}, \mathbf{X}\mathbf{v} \in \mathbb{R}^n$ has i.i.d. (C/\sqrt{n}) -sub-Gaussian entries. Then $\mathbf{u}^\top \mathbf{X}^\top \mathbf{X}\mathbf{v}$ is a sum of ni.i.d. (C'/n)-sub-exponential random variables, and Bernstein's inequality [Ver18, Theorem 2.8.1] implies

$$\mathbb{P}\Big[\Big|\mathbf{u}^{\top}\mathbf{X}^{\top}\mathbf{X}\mathbf{v}-\mathbf{u}^{\top}\boldsymbol{\Sigma}_{X}\mathbf{v}\Big| \geq t\Big] = \mathbb{P}\Big[\Big|\mathbf{u}^{\top}\mathbf{X}^{\top}\mathbf{X}\mathbf{v}-\mathbb{E}\mathbf{u}^{\top}\mathbf{X}^{\top}\mathbf{X}\mathbf{v}\Big| \geq t\Big] \leq 2e^{-cn\min(t^{2},t)}.$$

Applying this with $\mathbf{u} = \mathbf{v} = \mathbf{e}_j$ and taking a union bound over $j \in [p]$ shows $\operatorname{Tr} \mathbf{X}^\top \mathbf{X} \ge cp$. Applying this with $\mathbf{u} = \mathbf{v} = \mathbf{1}/\sqrt{p}$ shows $\|\mathbf{X}\mathbf{1}\|^2 \ge cp$. Applying this with $\mathbf{u} = \mathbf{v} = \mathbf{\Sigma}_X^{-1}\mathbf{e}_j/\|\mathbf{\Sigma}_X^{-1}\mathbf{e}_j\|$ shows $c \le \|\mathbf{b}_j\|^2 \le C$. These statements all hold with probability at least $1 - e^{-cn} \ge 1 - p^{-20}$. Furthermore, applying the above with $\mathbf{u} = \mathbf{\Sigma}_X^{-1}\mathbf{e}_j/\|\mathbf{\Sigma}_X^{-1}\mathbf{e}_j\|$ and $\mathbf{v} = \mathbf{e}_j$ or $\mathbf{v} = \mathbf{e}_k$ shows $|\mathbf{b}_j^\top \mathbf{x}_j - 1| \le C_0\sqrt{\log p/n}$ and $|\mathbf{b}_j^\top \mathbf{x}_k| \le C_0\sqrt{\log p/n}$ for any $j \ne k \in [p]$, with probability $1 - c^{2(n-1)/2}$.

Furthermore, applying the above with $\mathbf{u} = \mathbf{\Sigma}_X^{-1} \mathbf{e}_j / \|\mathbf{\Sigma}_X^{-1} \mathbf{e}_j\|$ and $\mathbf{v} = \mathbf{e}_j$ or $\mathbf{v} = \mathbf{e}_k$ shows $|\mathbf{b}_j^\top \mathbf{x}_j - 1| \leq C_0 \sqrt{\log p/n}$ and $|\mathbf{b}_j^\top \mathbf{x}_k| \leq C_0 \sqrt{\log p/n}$ for any $j \neq k \in [p]$, with probability $1 - e^{-cn \cdot C_0^2 (\log p)/n} \geq 1 - p^{-20}$ for sufficiently large $C_0 > 0$. Applying $\|\mathbf{X}\mathbf{1}\|^2 \geq cp$ already shown, we have $\|\mathbf{r}_j\| \leq (cp)^{-1/2} |\mathbf{1}^\top \mathbf{X}^\top \mathbf{X} \mathbf{\Sigma}_X^{-1} \mathbf{e}_j|$. Then applying the above with $\mathbf{u} = \mathbf{1}/\sqrt{p}$ and $\mathbf{v} = \mathbf{\Sigma}_X^{-1} \mathbf{e}_j / \|\mathbf{\Sigma}_X^{-1} \mathbf{e}_j\|$ shows $\|\mathbf{r}_j\| \leq C/\sqrt{p} + C_0 \sqrt{\log p/n}$ also with probability $1 - e^{-cn \cdot C_0^2 (\log p)/n} \geq 1 - p^{-20}$. Taking a union bound shows that for some constants $C_0, c_0 > 0$, the event E holds with probability at least $1 - p^{-10}$, as claimed. The proposition then follows from the Borel-Cantelli lemma.

B Analysis of algorithm dynamics

B.1 Fisher-Rao flow for the sequence model

Proposition B.1. For any probability distributions g, h on [-M, M],

$$\frac{\mathcal{N}_{\tau} * g(\varphi)}{\mathcal{N}_{\tau} * h(\varphi)} \le e^{(M^2/2\tau^2)} e^{(2M/\tau^2)|\varphi|}$$

Proof. Observe that for any $\theta, \theta' \in [-M, M]$,

$$\frac{\mathcal{N}_{\tau}(\theta-\varphi)}{\mathcal{N}_{\tau}(\theta'-\varphi)} = \frac{e^{-\frac{1}{2\tau^2}(\theta-\varphi)^2}}{e^{-\frac{1}{2\tau^2}(\theta'-\varphi)^2}} \in \left[e^{-\frac{M^2}{2\tau^2}-\frac{2M}{\tau^2}|\varphi|}, \ e^{\frac{M^2}{2\tau^2}+\frac{2M}{\tau^2}|\varphi|}\right].$$

Then also

$$\frac{\mathcal{N}_{\tau}(\theta - \varphi)}{\mathcal{N}_{\tau} * h(\varphi)} = \left(\int_{-M}^{M} \frac{\mathcal{N}_{\tau}(\theta' - \varphi)}{\mathcal{N}_{\tau}(\theta - \varphi)} \mathrm{d}h(\theta')\right)^{-1} \le e^{\frac{M^2}{2\tau^2} + \frac{2M}{\tau^2}|\varphi|},$$

and the result follows from further integrating this with respect to $dg(\theta)$.

Proof of Theorem 3.3. Assume that $D_{\mathrm{KL}}(h||g_0) < \infty$ for otherwise there is nothing to prove. By Corollary C.2, $\inf_{t \in [0,T]} \inf_{\theta \in [-M,M]} g_t(\theta)/g_0(\theta) \ge e^{-t}$. As such, $D_{\mathrm{KL}}(h||g_t) = D_{\mathrm{KL}}(h||g_0) + \int h \log \frac{g_0}{g_t} \le D_{\mathrm{KL}}(h||g_0) + t$ which is finite for all t. Together with the weak lower semicontinuity of the KL divergence, this implies

$$\lim_{t \to 0} D_{\text{KL}}(h \| g_t) = D_{\text{KL}}(h \| g_0).$$
(B.1)

Next, write for simplicity $\bar{F}(g) = \bar{F}(g \mid \bar{q})$, and define its first variation

$$\delta \bar{F}[g](\theta) = -\mathcal{N}_{\tau} * \frac{\bar{q}}{\mathcal{N}_{\tau} * g}(\theta) + 1$$

Applying Proposition B.1, $\mathcal{N}_{\tau} * \frac{\bar{q}}{\mathcal{N}_{\tau}*g}(\theta) \leq \int C e^{C|\varphi|} d\bar{q}(\varphi) < \infty$, so $\delta \bar{F}[g]$ is uniformly bounded over $\theta \in [-M, M]$. Let *h* be any density on [-M, M]. Applying the definition of the flow (3.2) and using this boundedness to differentiate under the integral by the dominated convergence theorem,

$$\int_{-M}^{M} -\delta \bar{F}[g_t](\theta)h(\theta)d\theta = \int_{-M}^{M} \frac{\mathrm{d}}{\mathrm{d}t} \left(\log \frac{g_t(\theta)}{h(\theta)}\right)h(\theta)d\theta = -\frac{\mathrm{d}}{\mathrm{d}t}D_{\mathrm{KL}}(h||g_t). \tag{B.2}$$

By convexity of $x \mapsto -\log x$, we have

$$\begin{split} \bar{F}(g) - \bar{F}(h) &= \int \left(-\log[\mathcal{N}_{\tau} * g](\varphi) + \log[\mathcal{N}_{\tau} * h](\varphi) \right) \mathrm{d}\bar{q}(\varphi) \\ &\leq \int \frac{\mathrm{d}}{\mathrm{d}\varepsilon} \Big[\log[\mathcal{N}_{\tau} * (g + \varepsilon(h - g))](\varphi) \Big]_{\varepsilon = 0} \mathrm{d}\bar{q}(\varphi) \\ &= \int \frac{[\mathcal{N}_{\tau} * (h - g)](\varphi)}{[\mathcal{N}_{\tau} * g](\varphi)} \mathrm{d}\bar{q}(\varphi) \\ &= \int \frac{\mathcal{N}_{\tau}(\theta - \varphi)h(\theta)}{\mathcal{N}_{\tau} * g(\varphi)} \mathrm{d}\bar{q}(\varphi) - 1 = \int_{-M}^{M} -\delta\bar{F}[g](\theta)h(\theta)\mathrm{d}\theta. \end{split}$$

Combining the above two displays and integrating from time t to time T yields

$$\int_t^T \left[\bar{F}(g_s) - \bar{F}(h) \right] \mathrm{d}s \le \int_t^T -\frac{\mathrm{d}}{\mathrm{d}t} D_{\mathrm{KL}}(h \| g_t) \Big|_{t=s} \mathrm{d}s = D_{\mathrm{KL}}(h \| g_t) - D_{\mathrm{KL}}(h \| g_T).$$

So taking the limit $t \to 0$ above gives

$$\int_{0}^{T} \left[\bar{F}(g_{s}) - \bar{F}(h) \right] \mathrm{d}s \le D_{\mathrm{KL}}(h \| g_{0}) - D_{\mathrm{KL}}(h \| g_{T}) \le D_{\mathrm{KL}}(h \| g_{T}).$$
(B.3)

Now observe that

$$\left|\frac{\mathrm{d}}{\mathrm{d}t}\int_{-M}^{M}\mathcal{N}_{\tau}(\varphi-\theta)g_{t}(\theta)\mathrm{d}\theta\right| = \left|\int_{-M}^{M}\mathcal{N}_{\tau}(\varphi-\theta)g_{t}(\theta)\delta\bar{F}[g_{t}](\theta)\mathrm{d}\theta\right| \le C\max_{\theta\in[-M,M]}\mathcal{N}_{\tau}(\varphi-\theta)$$

for a constant C > 0, using boundedness of $|\delta \bar{F}[g]|$ as shown above to differentiate under the integral. Then

$$\left|\frac{\mathrm{d}}{\mathrm{d}t}\log[\mathcal{N}_{\tau}*g_{t}](\varphi)\right| = \left|\frac{\frac{\mathrm{d}}{\mathrm{d}t}\int_{-M}^{M}\mathcal{N}_{\tau}(\varphi-\theta)g_{t}(\theta)\mathrm{d}\theta}{\mathcal{N}_{\tau}*g_{t}(\varphi)}\right| \le Ce^{C|\varphi|}$$

by Proposition B.1, so again differentiating under the integral by the dominated convergence theorem and applying $\int g(\theta) \delta \bar{F}[g](\theta) d\theta = 0$,

$$\frac{\mathrm{d}}{\mathrm{d}t}\bar{F}(g_t) = \int -\frac{\mathrm{d}}{\mathrm{d}t}\log[\mathcal{N}_{\tau} * g_t](\varphi)\mathrm{d}\bar{q}(\varphi)
= \int -\frac{\mathcal{N}_{\tau}(\varphi - \theta)g_t(\theta)\delta\bar{F}[g_t](\theta)}{\mathcal{N}_{\tau} * g_t(\varphi)}\mathrm{d}\bar{q}(\varphi) = -\int \left(\delta\bar{F}[g_t](\theta)\right)^2 g_t(\theta)\mathrm{d}\theta \le 0, \quad (B.4)$$

i.e. \bar{F} is non-increasing along the gradient flow path. Thus $T(\bar{F}(g_T) - \bar{F}(h)) \leq \int_0^T [\bar{F}(g_s) - \bar{F}(h)] ds \leq D_{\text{KL}}(h \| g_0)$ and rearranging yields the theorem.

B.2 LSI for Langevin dynamics in high noise

Proof of Theorem 3.4-(1). Recall that $\varphi = \theta + \mathbf{z}$ where $\mathbf{z} \sim \mathcal{N}(0, \tau^2 \operatorname{Id})$ and $\tau^2 + \delta < \sigma^2 / \|\mathbf{X}\|_{\operatorname{op}}^2$. Using also $M^2 + \delta < \sigma^2 / \|\mathbf{X}\|_{\operatorname{op}}^2$, we may pick some constant $\tau'^2 \in (\min(M^2, \tau^2), \sigma^2 / \|\mathbf{X}\|_{\operatorname{op}}^2)$ depending only on (M, τ, δ) . Then, defining $\eta^2 = \tau'^2 - \tau^2 > 0$ and $\varphi' = \varphi + \mathbf{z}' = \theta + \mathbf{z} + \mathbf{z}'$ for $\mathbf{z}' \sim \mathcal{N}(0, \eta^2 \operatorname{Id})$ independent of \mathbf{z} , we have the equivalent representation $\mathbf{y} = \mathbf{X}\varphi' + \varepsilon'$ with $\varepsilon' \sim \mathcal{N}(0, \mathbf{\Sigma}')$ independent of φ' such that $\mathbf{\Sigma}' = \sigma^2 \operatorname{Id} - {\tau'}^2 \mathbf{X} \mathbf{X}^\top$ is positive definite.

Write as shorthand $\nu(\varphi) = P_g(\varphi \mid \mathbf{y}), \ \nu_{\varphi'}(\varphi) = P_g(\varphi \mid \varphi')$, and $\mu(\varphi') = P_g(\varphi' \mid \mathbf{y})$. Following the decomposition of [BB19], for any bounded, Lipschitz, and continuously differentiable function $f : \mathbb{R}^p \to \mathbb{R}$, by the tower property

$$\operatorname{ent}_{\nu}(f^{2}) = \mathbb{E}_{\nu}f^{2}\log f^{2} - \mathbb{E}_{\nu}f^{2} \cdot \log \mathbb{E}_{\nu}f^{2}$$

$$= \mathbb{E}_{\varphi' \sim \mu} \Big[\mathbb{E}_{\nu_{\varphi'}}f^{2}\log f^{2} - \mathbb{E}_{\nu_{\varphi'}}f^{2} \cdot \log \mathbb{E}_{\nu_{\varphi'}}f^{2} \Big]$$

$$+ \mathbb{E}_{\varphi' \sim \mu} \Big[\mathbb{E}_{\nu_{\varphi'}}f^{2} \cdot \log \mathbb{E}_{\nu_{\varphi'}}f^{2} \Big] - \mathbb{E}_{\varphi' \sim \mu} [\mathbb{E}_{\nu_{\varphi'}}f^{2}] \cdot \log \mathbb{E}_{\varphi' \sim \mu} [\mathbb{E}_{\nu_{\varphi'}}f^{2}]$$

$$= \mathbb{E}_{\mu}\operatorname{ent}_{\nu_{\varphi'}}(f^{2}) + \operatorname{ent}_{\mu} \mathbb{E}_{\nu_{\varphi'}}(f^{2}). \tag{B.5}$$

For the first term, note that $\nu_{\varphi'}(\varphi)$ is a product measure with marginals

$$\nu_{\varphi',j}(\varphi_j) \propto P_g(\varphi_j \mid \varphi'_j) \propto e^{-\frac{(\varphi'_j - \varphi_j)^2}{2\eta^2}} \cdot \int e^{-\frac{(\varphi_j - \theta)^2}{2\tau^2}} \mathrm{d}g(\theta) \propto \int e^{-\frac{(\varphi_j - c(\theta))^2}{2\gamma^2}} \mathrm{d}\bar{g}(\theta), \qquad (B.6)$$

where

$$\gamma^{2} = \frac{\eta^{2}\tau^{2}}{\eta^{2} + \tau^{2}}, \quad c(\theta) = \frac{\tau^{2}\varphi_{j}' + \eta^{2}\theta}{\eta^{2} + \tau^{2}}, \quad \mathrm{d}\bar{g}(\theta) = \frac{e^{-\frac{(\theta - \varphi_{j}')^{2}}{2(\eta^{2} + \tau^{2})}}\mathrm{d}g(\theta)}{\int e^{-\frac{(u - \varphi_{j}')^{2}}{2(\eta^{2} + \tau^{2})}}\mathrm{d}g(u)}.$$
(B.7)

In other words, $\nu_{\varphi',j}$ is the convolution between $\mathcal{N}(0,\gamma^2)$ and the pushforward of \bar{g} under c, which is supported on an interval of length at most 2M. By [Zim16, Theorem 1], $\nu_{\varphi',j}$ satisfies a onedimensional LSI with constant $C = C(\tau, \tau', M)$:

$$\operatorname{ent}_{\nu_{\varphi',j}}(f^2) \le C \cdot \mathbb{E}_{\nu_{\varphi',j}} |\partial_j f(\varphi)|^2 \tag{B.8}$$

where we write $\operatorname{ent}_{\nu_{\varphi',j}}$ and $\mathbb{E}_{\nu_{\varphi',j}}$ for the partial entropy and expectation over the coordinate φ_j of $\varphi \sim \nu_{\varphi'}$. Finally, by tensorization of LSI [Vil09, Theorem 9.9(i)], the product measure $\nu_{\varphi'}$ satisfies a LSI with the same C, namely,

$$\operatorname{ent}_{\nu_{\varphi'}}(f^2) \le C \cdot \mathbb{E}_{\nu_{\varphi'}} \|\nabla f(\varphi)\|^2.$$
(B.9)

Taking further expectation under μ yields

$$\mathbb{E}_{\mu} \operatorname{ent}_{\nu_{\varphi'}}(f^2) \le C \cdot \mathbb{E}_{\nu} \|\nabla f(\varphi)\|^2.$$
(B.10)

To bound the second term in (B.5), note that $\mu(\varphi') \propto \exp\left(-\frac{1}{2}(\mathbf{y} - \mathbf{X}\varphi')^{\top} \mathbf{\Sigma}'^{-1}(\mathbf{y} - \mathbf{X}\varphi') - \sum_{j=1}^{p} V(\varphi'_{j})\right)$, where

$$V(\varphi') = -\log \int e^{-\frac{1}{2\tau'^2}(\varphi'-\theta)^2} \mathrm{d}g(\theta).$$

We have

$$V''(\varphi') = \frac{1}{{\tau'}^2} - \partial_{\varphi'}^2 \log \int e^{\frac{\varphi'\theta}{{\tau'}^2} - \frac{\theta^2}{2{\tau'}^2}} \mathrm{d}g(\theta) = \frac{1}{{\tau'}^2} - \frac{\mathrm{Var}_g(\theta \mid \varphi')}{{\tau'}^4} \ge \frac{1}{{\tau'}^2} - \frac{M^2}{{\tau'}^4} =: c(\tau', M) > 0,$$

using the fact that $P(\theta \mid \varphi')$ is supported on [-M, M] and ${\tau'}^2 > M^2$. Hence the density of $\mu(\varphi')$ is strictly log-concave, so by the Bakry-Emery criterion [Vil09, Theorem 9.9(iii)], $\mu(\varphi')$ satisfies a LSI with constant $c(\tau', M)^{-1}$. Let $G(\varphi') \equiv [\mathbb{E}_{\nu_{\varphi'}}(f^2)]^{1/2}$. Then

$$\operatorname{ent}_{\mu} \mathbb{E}_{\nu_{\varphi'}}(f^2) = \operatorname{ent}_{\mu}(G^2(\varphi')) \le c(\tau', M)^{-1} \cdot \mathbb{E}_{\mu} \|\nabla G(\varphi')\|^2.$$
(B.11)

We have

$$\partial_{\varphi_i'} G(\varphi') = \frac{\partial_{\varphi_i'} G^2(\varphi')}{2G(\varphi')} = \frac{1}{2\eta^2} \frac{\operatorname{Cov}_{\nu_{\varphi'}}(f^2, \varphi_i)}{[\mathbb{E}_{\nu_{\varphi'}}(f^2)]^{1/2}} = \frac{1}{2\eta^2} \frac{\mathbb{E}_{\nu_{\varphi',-i}} \operatorname{Cov}_{\nu_{\varphi',i}}(f^2, \varphi_i)}{[\mathbb{E}_{\nu_{\varphi'}}(f^2)]^{1/2}},$$
(B.12)

where the second equality follows from $\partial_{\varphi'_i} G^2(\varphi') = \mathbb{E}_{\nu_{\varphi'}} [f^2 \partial_{\varphi'_i} \log P_g(\varphi_i | \varphi'_i)]$ and the fact that $\partial_{\varphi'_i} \log P_g(\varphi_i | \varphi'_i) = \frac{1}{\eta^2} (\varphi_i - \mathbb{E}[\varphi_i | \varphi'_i])$ from (B.6), and the last equality uses the fact that $\nu_{\varphi'}$ is a product measure. Shorthand $\operatorname{Cov}_{\nu_{\varphi',i}}(f^2, \varphi_i)$ as $\operatorname{Cov}_i(f^2, \varphi_i | \varphi'_i)$. Note that (B.6) yields the representation of the conditional law $(\varphi_i | \varphi'_i) \stackrel{d}{=} c(\theta) + z$ where $c(\theta)$ is given in (B.7), and $\theta \sim \overline{g}(\cdot)$ and $z \sim \mathcal{N}(0, \gamma^2)$ are independent (conditional on φ'_i). Consequently,

$$\begin{aligned} \operatorname{Cov}_{i}(f^{2}(\boldsymbol{\varphi}),\varphi_{i} \mid \varphi_{i}') &= \operatorname{Cov}_{i}(f^{2}(\boldsymbol{\varphi}_{-i},c(\theta)+z),c(\theta)+z \mid \varphi_{i}') \\ &= \operatorname{Cov}_{i}(f^{2}(\boldsymbol{\varphi}_{-i},c(\theta)+z),c(\theta) \mid \varphi_{i}') + \operatorname{Cov}_{i}(f^{2}(\boldsymbol{\varphi}_{-i},c(\theta)+z),z \mid \varphi_{i}') \equiv (I) + (II). \end{aligned}$$

To bound (I), let $\theta_1, \theta_2 \sim \bar{g}(\cdot)$ and $z_1, z_2 \sim \mathcal{N}(0, \gamma^2)$ be independent. Then

$$2 \cdot (I) = \mathbb{E} \Big[(f^{2}(\varphi_{-i}, c(\theta_{1}) + z_{1}) - f^{2}(\varphi_{-i}, c(\theta_{2}) + z_{2}))(c(\theta_{1}) - c(\theta_{2})) \mid \varphi_{i}' \Big] \\ \leq \mathbb{E} \Big[\Big(f(\varphi_{-i}, c(\theta_{1}) + z_{1}) - f(\varphi_{-i}, c(\theta_{2}) + z_{2}) \Big)^{2} \mid \varphi_{i}' \Big]^{1/2} \\ \cdot \mathbb{E} \Big[\Big(f(\varphi_{-i}, c(\theta_{1}) + z_{1}) + f(\varphi_{-i}, c(\theta_{2}) + z_{2}) \Big)^{2} (c(\theta_{1}) - c(\theta_{2}))^{2} \mid \varphi_{i}' \Big]^{1/2} \\ = (2 \operatorname{Var}_{\nu_{\varphi',i}} [f])^{1/2} \cdot \frac{\eta^{2}}{\tau^{\prime 2}} \cdot \mathbb{E} \Big[\Big(f(\varphi_{-i}, c(\theta_{1}) + z_{1}) + f(\varphi_{-i}, c(\theta_{2}) + z_{2}) \Big)^{2} (\theta_{1} - \theta_{2})^{2} \mid \varphi_{i}' \Big]^{1/2} \Big]^{1/2}$$

$$\leq C(M,\tau,\tau') \cdot \left[\mathbb{E}_{\nu_{\varphi',i}}(\partial_i f)^2\right]^{1/2} \cdot \left[\mathbb{E}_{\nu_{\varphi',i}}(f^2)\right]^{1/2},$$

where the last step applies the Poincaré inequality for $\nu_{\varphi',i}$ (as implied by its LSI (B.8) – see [Vil09, (9.32)]) to $\operatorname{Var}_{\nu_{\varphi',i}}[f]$, and the fact that $\theta_1, \theta_2 \in [-M, M]$. To bound (II), using the independence of z and θ and applying Stein's lemma,

$$(II) = \mathbb{E} \Big[f^2(\varphi_{-i}, c(\theta) + z) z \mid \varphi_i' \Big] = 2\gamma^2 \cdot \mathbb{E} \Big[f(\varphi_{-i}, c(\theta) + z) \partial_i f(\varphi_{-i}, c(\theta) + z) \mid \varphi_i' \Big] \\ \leq C(\tau, \tau') \cdot \big[\mathbb{E}_{\nu_{\varphi',i}}(\partial_i f)^2 \big]^{1/2} \cdot \big[\mathbb{E}_{\nu_{\varphi',i}}(f^2) \big]^{1/2}.$$

Combining the two bounds and applying Cauchy-Schwarz and the tower property for $\mathbb{E}_{\nu_{\omega',-i}}$ yields

$$\mathbb{E}_{\nu_{\varphi',-i}} \operatorname{Cov}_i(f^2(\varphi), \varphi_i \mid \varphi'_i) \le C \cdot \left[\mathbb{E}_{\nu_{\varphi'}}(\partial_i f)^2\right]^{1/2} \cdot \left[\mathbb{E}_{\nu_{\varphi'}}(f^2)\right]^{1/2}$$

Substituting this into (B.11) and (B.12), the second term in (B.5) is bounded by

$$\operatorname{ent}_{\mu} \mathbb{E}_{\nu_{\varphi'}}(f^2) \le C' \cdot \mathbb{E}_{\mu} \sum_{i=1}^{p} \mathbb{E}_{\nu_{\varphi'}}(\partial_i f)^2 = C' \cdot \mathbb{E}_{\nu} \|\nabla f\|^2$$
(B.13)

for some $C' = C'(\tau, \tau', M)$. Combining (B.10) and (B.13) concludes the proof.

The proof of Theorem 3.4-(2) requires the following additional lemma, whose proof we present after the main proof.

Lemma B.2. Fix any M > 0. Uniformly over $h \in \mathbb{R}$, the measure $\mu_h(x) \propto e^{hx} \mathbf{1}_{|x| < M}$ satisfies a LSI over (-M, M) with constant C = C(M) > 0.

Proof of Theorem 3.4-(2). The proof is similar to Theorem 3.4-(1), and we skip some of the common steps. Pick ${\tau'}^2 \in (M^2, \sigma^2/||\mathbf{X}||_{op}^2)$ depending only on (M, δ) . Let $\boldsymbol{\varphi}' = \boldsymbol{\theta} + \mathbf{z}$ with $\mathbf{z} \sim \mathcal{N}(0, {\tau'}^2 \operatorname{Id})$, and represent $\mathbf{y} = \mathbf{X} \boldsymbol{\varphi}' + \boldsymbol{\varepsilon}'$ with $\boldsymbol{\varepsilon}' \sim \mathcal{N}(0, \boldsymbol{\Sigma}')$ and $\boldsymbol{\Sigma}' = \sigma^2 - {\tau'}^2 \mathbf{X} \mathbf{X}^{\top}$. Then, denoting $\nu = P_g(\boldsymbol{\theta} \mid \mathbf{y})$, $\nu_{\boldsymbol{\varphi}'} = P_g(\boldsymbol{\theta} \mid \boldsymbol{\varphi}')$ and $\mu = P_g(\boldsymbol{\varphi}' \mid \mathbf{y})$, we have, analogous to (B.5),

$$\operatorname{ent}_{\nu}(f^2) = \mathbb{E}_{\mu} \operatorname{ent}_{\nu_{\varphi'}}(f^2) + \operatorname{ent}_{\mu} \mathbb{E}_{\nu_{\varphi'}}(f^2).$$

Here $\nu_{\varphi'}$ is a product measure with marginals

$$u_{\varphi',j}(\theta_j) \propto e^{-\frac{\varphi'_j}{\tau^2}\theta_j} \cdot e^{-\frac{\theta_j^2}{2\tau^2}}g(\theta_j).$$

By Lemma B.2, the density proportional to $e^{-\frac{\varphi'_j}{\tau^2}\theta_j}$ on (-M, M) satisfies a LSI with constant C = C(M) > 0 uniformly over φ'_j . Then by the Holley-Stroock perturbation principle [Vil09, Theorem 9.9(ii)], $\nu_{\varphi',j}$ also satisfies a LSI with constant $C = C'(M, \tau)\kappa(g)^2$. Applying this and tensorization shows

$$\mathbb{E}_{\mu} \operatorname{ent}_{\nu_{\boldsymbol{\omega}'}}(f^2) \leq C \cdot \mathbb{E}_{\nu} \|\nabla f(\boldsymbol{\theta})\|^2.$$

For the second term, we have again that μ is strongly log-concave and hence satisfies a LSI with some constant $c(\tau', M)^{-1} < \infty$. Let $G(\varphi') \equiv [\mathbb{E}_{\nu_{\alpha'}}(f^2)]^{1/2}$. Then as before,

$$\operatorname{ent}_{\mu} \mathbb{E}_{\nu_{\varphi'}}(f^2) = \operatorname{ent}_{\mu}(G^2(\varphi')) \le c(\tau', M)^{-1} \cdot \mathbb{E}_{\mu} \|\nabla G(\varphi')\|^2,$$

where

$$\partial_{\varphi_i'} G(\varphi') = \frac{\partial_{\varphi_i'} G^2(\varphi')}{2G(\varphi')} = \frac{1}{2\tau'^2} \frac{\mathbb{E}_{\nu_{\varphi',-i}} \operatorname{Cov}_{\nu_{\varphi',i}}(f^2, \theta_i)}{[\mathbb{E}_{\nu_{\varphi'}}(f^2)]^{1/2}}.$$

Here, since $\theta_i \in [-M, M]$ is bounded, we have simply

$$\operatorname{Cov}_{\nu_{\varphi',i}}(f^2,\theta_i) \le C(M) \cdot \operatorname{Var}_{\nu_{\varphi',i}}(f^2) \le C'(M) \cdot \left[\operatorname{Var}_{\nu_{\varphi',i}}(f)\right]^{1/2} \cdot \left[\mathbb{E}_{\nu_{\varphi',i}}(f^2)\right]^{1/2}.$$

Then, applying again the Poincaré inequality for $\nu_{\varphi',i}$ gives

$$\operatorname{ent}_{\mu}\nu_{\varphi'}(f^2) \le C \cdot \mathbb{E}_{\mu} \sum_{i=1}^{p} \mathbb{E}_{\nu_{\varphi'}}(\partial_i f)^2 = C \cdot \mathbb{E}_{\nu} \|\nabla f\|^2$$

which completes the proof.

To prove Lemma B.2, we use the following characterization of the LSI constant due to [BG99, Theorem 5.3]. We note that this result has also been applied in [ZQM11] to prove an analogue of Lemma B.2 under spherical priors and general dimensions.

Lemma B.3 ([BG99]). Let μ be a (Borel) probability measure on \mathbb{R} and let $\rho(x)$ be the density of its absolutely continuous part with respect to Lebesgue measure. Let m be a median of μ . Then the optimal constant C_* in the LSI

$$ent_{\mu}(f^2) \le C_* \mathbb{E}_{\mu}[{f'}^2]$$

satisfies $c_1 \max(B_-, B_+) \leq C_* \leq c_2 \max(B_-, B_+)$ for some universal constants c_1, c_2 , where

$$B_{+} = \sup_{x > m} \mu([x, \infty)) \cdot \log \frac{1}{\mu([x, \infty))} \cdot \int_{m}^{x} \frac{1}{\rho(t)} dt,$$

$$B_{-} = \sup_{x > m} \mu((-\infty, x]) \cdot \log \frac{1}{\mu((-\infty, x])} \cdot \int_{x}^{m} \frac{1}{\rho(t)} dt$$

This result does not require μ to have full support on all of \mathbb{R} , and B_+ or B_- is defined as 0 if $\mu((m,\infty)) = 0$ or $\mu((-\infty,m)) = 0$ respectively.

Proof of Lemma B.2. The uniform density on (-M, M) satisfies a LSI with constant C(M) > 0by [GMW14]. Then by the Holley-Stroock perturbation principle, it suffices to show the result for $|h| > h_0(M)$ and any fixed and large constant $h_0(M) > 0$. By symmetry, we may assume h > 0. For presentational simplicity, we only prove the case M = 1, and the case of general M follows similarly.

Using Lemma B.3, it suffices to control the quantities B_+, B_- for μ_h . Let us write $a(h) \simeq b(h)$ if there exist constants C, c > 0 for which $ca(h) \le b(h) \le Ca(h)$ for all large h. Let $m \equiv m(h)$ be the median of μ_h , and let $\rho_h(t) = e^{ht}/(h^{-1}(e^h - e^{-h}))$ be the density of μ_h . We claim that $1 - m \simeq \frac{1}{h}$. To see this, note that for any $x \in (-1, 1)$,

$$\mu_h([-1,x]) = \int_{-1}^x \frac{e^{ht}}{h^{-1}(e^h - e^{-h})} dt = \frac{e^{hx} - e^{-h}}{e^h - e^{-h}},$$

so for sufficiently large h, $\inf\{x : \mu_h([-1, x]) \ge 1/2\} \in [1 - C/h, 1 - c/h].$

We first bound B_+ . Using the boundedness of $t \mapsto t \log(1/t)$ and

$$\int_{m}^{x} \frac{1}{\rho_{h}(t)} dt = \frac{1}{h^{2}} (e^{h} - e^{-h})(e^{-hm} - e^{-hx}) \approx \frac{1}{h^{2}} (e^{h(1-m)} - e^{h(1-x)}),$$

we have for $1 - m \approx \frac{1}{h}$ and any $x \in (m, 1)$ that

$$\mu_h([x,1]) \cdot \log \frac{1}{\mu_h([x,1])} \cdot \int_m^x \frac{1}{\rho_h(t)} \mathrm{d}t \le \frac{C}{h^2}$$

for a constant C > 0 independent of x. Then $B_+ \leq C/h^2$.

Next we bound B_{-} . Using

$$\mu_h\big([-1,x]\big) = \frac{e^{hx} - e^{-h}}{e^h - e^{-h}} \asymp e^{-h(1-x)}(1 - e^{-h(1+x)}), \quad \int_x^m \frac{1}{\rho_h(t)} \mathrm{d}t \asymp \frac{e^{h(1-x)}}{h^2}(1 - e^{h(x-m)}),$$

we have for any $x \in (-1, m)$ and a constant C > 0 that

$$\mu_h([-1,x]) \cdot \log \frac{1}{\mu_h([-1,x]))} \cdot \int_x^m \frac{1}{\rho_h(t)} dt \le \frac{C}{h^2} (1 - e^{-h(1+x)}) \log \frac{e^h - e^{-h}}{e^{hx} - e^{-h}} \le \frac{C}{h^2} (1 - e^{-h(1+x)}) \log \frac{e^{h(1-x)}}{1 - e^{-h(1+x)}}.$$

Applying again boundedness of $t \mapsto t \log(1/t)$, we have in this case $B_{-} \leq C'/h$ for a constant C' > 0. In summary, $\max(B_{+}, B_{-})$ is at most a constant, which in light of Lemma B.3 completes the proof.

B.3 Analysis of the joint gradient flow

We isolate the following intuitive but somewhat technical lemma, which verifies the statement

$$\frac{\mathrm{d}}{\mathrm{d}t}F_n(q_t, g_t) = -p \|\operatorname{grad}_q^{W_2} F_n(q_t, g_t)\|_{q_t}^2 - \alpha \|\operatorname{grad}_g^{\mathrm{FR}} F_n(q_t, g_t)\|_{g_t}^2$$

where $\|\cdot\|_{q_t}^2$, $\|\cdot\|_{g_t}^2$ are the squared norms under the Wasserstein-2 metric tensor at q_t and Fisher-Rao metric tensor at g_t , respectively. (For background on these metric tensors, see Section D.1.)

Lemma B.4. Under the conditions of Theorem 3.7,

$$\limsup_{t \to 0} F_n(q_t, g_t) \le F_n(q_0, g_0).$$

Furthermore $t \mapsto F_n(q_t, g_t)$ is differentiable at every t > 0, and

$$\frac{\mathrm{d}}{\mathrm{d}t}F_n(q_t, g_t) = -\frac{1}{p} \int \left\| \nabla \log \frac{\mathrm{d}q_t}{\mathrm{d}\nu[g_t]} \right\|_2^2 q_t(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} - \alpha \int \left(\left[\mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * g_t} \right](\boldsymbol{\theta}) - 1 \right)^2 g_t(\boldsymbol{\theta}) \mathrm{d}\boldsymbol{\theta}.$$
(B.14)

The proof of Lemma B.4 is deferred to Appendix C.4.

Proof of Theorem 3.7. Assume $D_{\mathrm{KL}}(h||g_0) < \infty$, as otherwise there is nothing to prove. Then by Theorem 3.6-(1) and the same argument leading to (B.1), $D_{\mathrm{KL}}(h||g_t)$ is finite for all t and continuous

at t = 0. Next, recall that $-\delta \bar{F}_n[g_t](\theta) = \left(\mathcal{N}_\tau * \frac{\bar{\nu}[g_t]}{\mathcal{N}_\tau * g_t}\right)(\theta) - 1$, so for any density h supported on [-M, M],

$$\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta) \cdot h(\theta) = \int_{\mathbb{R}} \frac{\mathcal{N}_\tau * h}{\mathcal{N}_\tau * g_t}(\varphi) \bar{\nu}[g_t](\varphi) - 1 = \frac{1}{p} \sum_{j=1}^p \int_{\mathbb{R}^p} \frac{\mathcal{N}_\tau * h}{\mathcal{N}_\tau * g_t}(\varphi_j) \nu[g_t](\varphi) - 1.$$
(B.15)

Denote

$$f_t(\varphi) = \log \frac{\mathcal{N}_{\tau} * h}{\mathcal{N}_{\tau} * g_t}(\varphi), \qquad \nu_j[g_t](f_t) = \int f_t(\varphi_j) \nu[g_t](\varphi).$$

Then

$$\int \frac{\mathcal{N}_{\tau} * h}{\mathcal{N}_{\tau} * g_t} (\varphi_j) \nu[g_t](\varphi) = \int e^{f_t(\varphi_j)} \nu[g_t](\varphi)$$
$$= \int e^{f_t(\varphi_j)} q_t(\varphi) + e^{\nu_j [g_t](f_t)} \int e^{f_t(\varphi_j) - \nu_j [g_t](f_t)} (\nu[g_t](\varphi) - q_t(\varphi)). \quad (B.16)$$

Observe that $f_t(\varphi_j)$ is 2*M*-Lipschitz in φ , because

$$\frac{\mathrm{d}}{\mathrm{d}\varphi}f_t(\varphi) = \frac{(\mathcal{N}_\tau * h)'}{\mathcal{N}_\tau * h}(\varphi) - \frac{(\mathcal{N}_\tau * g_t)'}{\mathcal{N}_\tau * g_t}(\varphi) = \mathbb{E}_h[\theta \mid \theta + \tau Z = \varphi] - \mathbb{E}_{g_t}[\theta \mid \theta + \tau Z = \varphi] \in [-2M, 2M]$$

with $Z \sim \mathcal{N}(0,1)$. Then for some (M,τ) -dependent constants $C, C', c > 0, f_t(\varphi_j)$ is C'-subgaussian under the law $\nu[g_t](\varphi)$ by the given LSI assumption (cf. [Vil09, Theorem 9.9(iv)]), so for any L > 0,

$$\int_{f_t(\varphi_j)-\nu_j[g_t](f_t)>L} e^{f_t(\varphi_j)-\nu_j[g_t](f_t)} (\nu[g_t](\varphi) - q_t(\varphi)) \\
\leq \int_{f_t(\varphi_j)-\nu_j[g_t](f_t)>L} e^{f_t(\varphi_j)-\nu_j[g_t](f_t)} \nu[g_t](\varphi) \leq C e^{-cL^2}.$$
(B.17)

Next, applying $d_{\text{TV}}(\nu[g_t], q_t) = \frac{1}{2} \int |\nu[g_t](\varphi) - q_t(\varphi)|$ and Pinsker's inequality $d_{\text{TV}}(\nu[g_t], q_t) \leq \sqrt{\frac{1}{2}D_{\text{KL}}(q_t\|\nu[g_t])}$, we have

$$\int_{f_t(\varphi_j) - \nu_j[g_t](f_t) \le L} e^{f_t(\varphi_j) - \nu_j[g_t](f_t)} (\nu[g_t](\varphi) - q_t(\varphi)) \le 2e^L d_{\mathrm{TV}}(\nu[g_t], q_t) \le e^L \sqrt{2D_{\mathrm{KL}}(q_t \| \nu[g_t])}.$$
(B.18)

Applying (B.14) without the second term as an upper bound, and applying again the LSI for $\nu[g_t](\varphi)$ with $f = \sqrt{\mathrm{d}q_t/\mathrm{d}\nu[g_t]}$ in its definition (3.9), we have for a constant $c := c(M, \tau) > 0$,

$$\frac{\mathrm{d}}{\mathrm{d}t}F_n(q_t, g_t) \le -\frac{1}{p} \int \left\| \nabla \log \frac{\mathrm{d}q_t}{\mathrm{d}\nu[g_t]} \right\|_2^2 q_t(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \le -\frac{c}{p} D_{\mathrm{KL}}(q_t \| \nu[g_t]) \le 0.$$
(B.19)

Applying (B.19) to (B.18) and combining with (B.17), we obtain for (M, τ) -dependent constants $C_0, c_0 > 0$ that

$$\int e^{f_t(\varphi_j) - \nu_j[g_t](f_t)}(\nu[g_t](\varphi) - q_t(\varphi)) \le C_0 \left(e^{-c_0 L^2} + e^L \sqrt{-p \cdot \frac{\mathrm{d}}{\mathrm{d}t} F_n(q_t, g_t)} \right) := H_t(L) \quad (B.20)$$

where we define $H_t(L)$ to be this (non-negative) upper bound.

Now by Jensen's inequality, $e^{\nu_j[g_t](f_t)} \leq \int e^{f_t(\varphi_j)} \nu[g_t](\varphi)$. Applying this and (B.20) back to (B.16),

$$\int e^{f_t(\varphi_j)} \nu[g_t](\varphi) - 1 \le \int e^{f_t(\varphi_j)} q_t(\varphi) - 1 + H_t(L) \int e^{f_t(\varphi_j)} \nu[g_t](\varphi).$$

Then, averaging over j = 1, ..., p, applying the identity (B.15), and using

$$\frac{1}{p}\sum_{j=1}^{p}e^{f_t(\varphi_j)}q_t(\varphi) - 1 = \int \frac{\mathcal{N}*h}{\mathcal{N}*g_t}(\varphi)\bar{q}_t(\varphi)\mathrm{d}\varphi - 1 = \frac{\mathrm{d}}{\mathrm{d}t}[-D_{\mathrm{KL}}(h\|g_t)]$$

(where we may differentiate under the integral by the same argument as in (B.2)), we get

$$\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta) \cdot h(\theta) \le \frac{\mathrm{d}}{\mathrm{d}t} [-D_{\mathrm{KL}}(h||g_t)] + H_t(L) \Big(\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta) \cdot h(\theta) + 1\Big).$$

This yields the implication

$$\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta) \cdot h(\theta) \ge 0 \text{ and } H_t(L) \le 1/2 \Longrightarrow \int -\delta \bar{F}_n[g_t](\theta) \cdot h(\theta) \le 2H_t(L) + 2\frac{\mathrm{d}}{\mathrm{d}t}[-D_{\mathrm{KL}}(h||g_t)]$$

Now, suppose by contradiction that $\int -\delta \bar{F}_n[g_t](\theta) \cdot h(\theta) \geq \varepsilon > 0$ for all $t \in [0,T]$, where $\varepsilon < \min(1/2, 1/(4C_0), 1/(16\alpha))$. Fix L such that $C_0 e^{-c_0 L^2} = \varepsilon/4$, i.e. $L = \sqrt{(-1/c_0) \log[\varepsilon/(4C_0)]}$, and denote

$$K = C_0 e^L \sqrt{p}, \qquad A(t) = -D_{\mathrm{KL}}(h \| g_t), \qquad B(t) = -F_n(q_t, g_t)$$

Then $H_t(L) = K\sqrt{B'(t)} + \varepsilon/4$. Then the above implies that for each $t \in [0, T]$,

either
$$K\sqrt{B'(t)} \ge 1/2 - \varepsilon/4 \ge 1/4$$
 or $\varepsilon \le 2K\sqrt{B'(t)} + \varepsilon/2 + 2A'(t)$.

In the latter case $\frac{\varepsilon}{4} \leq K\sqrt{B'(t)} + A'(t)$, so $\frac{\varepsilon^2}{8} \leq \frac{\varepsilon}{2}A'(t) + \frac{\varepsilon}{2}K\sqrt{B'(t)} \leq \frac{\varepsilon}{2}A'(t) + \frac{\varepsilon^2}{16} + K^2B'(t)$, the latter inequality applying Cauchy-Schwarz. This implies

$$K^2 B'(t) + A'(t) \cdot \varepsilon/2 \ge \varepsilon^2/8$$

This holds also in the former case because $K^2B'(t) + A'(t)\varepsilon/2 \ge 1/16 + A'(t)\varepsilon/2 \ge \varepsilon^2/8$, the latter inequality using $\varepsilon < \min(1/2, 1/(16\alpha))$ and

$$A'(t) = \frac{\mathrm{d}}{\mathrm{d}t} \int_{-M}^{M} h(\theta) \log g_t(\theta) = \alpha \cdot \int_{-M}^{M} \left(\left[\mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * g_t} \right](\theta) - 1 \right) h(\theta) \ge \alpha \cdot \int_{-M}^{M} -h(\theta) = -\alpha.$$

Then, recalling the definitions of A(t), B(t) and integrating from time t to time T gives

$$K^{2}(F_{n}(q_{t},g_{t})-F_{n}(q_{T},g_{T}))+\frac{\varepsilon}{2}(D_{\mathrm{KL}}(h||g_{t})-D_{\mathrm{KL}}(h||g_{T}))\geq\frac{(T-t)\varepsilon^{2}}{8}.$$

We have $\limsup_{t\to 0} F_n(q_t, g_t) \leq F_n(q_0, g_0)$ by Lemma B.4, and $\lim_{t\to 0} D_{\mathrm{KL}}(h||g_t) = D_{\mathrm{KL}}(h||g_0)$. Then

$$K^{2}(F_{n}(q_{0},g_{0})-F_{n,*})+\frac{\varepsilon}{2}D_{\mathrm{KL}}(h||g_{0})\geq\frac{T\varepsilon^{2}}{8}$$

This is a contradiction for

$$T \ge C \left[\frac{p \cdot e^{C\sqrt{-\log \varepsilon}}}{\varepsilon^2} (F_n(q_0, g_0) - F_{n,*}) + \frac{1}{\varepsilon} D_{\mathrm{KL}}(h \| g_0) \right]$$

and some large enough constant $C := C(M, \tau) > 0$, which proves (3.14).

For the second claim (3.15), since $\bar{F}_n(g_t) \leq F_n(q_t, g_t) \leq F_n(q_0, g_0)$ where the second inequality follows from (B.19), we have that g_t belongs to \mathcal{K} for all $t \geq 0$. Let $t \leq T$ be as in (3.14) such that $\int_{-M}^{M} -\delta \bar{F}_n[g_t](\theta)h(\theta) \leq \varepsilon$, and define $H(\lambda) = \bar{F}_n(\lambda h + (1 - \lambda)g_t)$. Since also $h \in \mathcal{K}$, by the assumed convexity of \bar{F}_n on \mathcal{K} we have $H''(\lambda) \geq 0$ for all $\lambda \in [0, 1]$ so $H(1) \geq H(0) + H'(0)$, which is equivalent to

$$\bar{F}_n(g_t) - \bar{F}_n(h) \le \int_{-M}^M -\delta \bar{F}_n[g_t](\theta) h(\theta) \mathrm{d}\theta \le \varepsilon,$$

as desired.

B.4 Proof of (3.17)

We verify (3.17) under Assumption A and the condition $\tau^2 < \sigma^2 / \|\mathbf{X}\|_{op}^2 - \delta$. Recall that $F_{n,*} = \min_{g \in \mathcal{P}(M)} -\frac{1}{p} \log P_g(\mathbf{y})$. From the form (1.4), we have $P_g(\mathbf{y}) \leq (2\pi\sigma^2)^{-n/2}$, hence $F_{n,*} \geq \frac{n}{2p} \log(2\pi\sigma^2)$. Next, recall from (2.11) that we have

$$F_n(q_0, g_0) = \frac{1}{2p} \mathbb{E}_{\boldsymbol{\varphi} \sim q_0} [(\mathbf{y} - \mathbf{X} \boldsymbol{\varphi})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi})] + \frac{1}{p} D_{\mathrm{KL}}(q_0 \| (\mathcal{N}_{\tau} * g_0)^{\otimes p}) + \frac{n}{2p} \log(2\pi) + \frac{1}{2p} \log \det(\boldsymbol{\Sigma}).$$
(B.21)

Note that

$$\mathbb{E}_{\boldsymbol{\varphi} \sim q_0}[(\mathbf{y} - \mathbf{X}\boldsymbol{\varphi})^{\top} \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \mathbf{X}\boldsymbol{\varphi})] \leq \|\boldsymbol{\Sigma}^{-1}\|_{\mathrm{op}} \mathbb{E}_{\boldsymbol{\varphi} \sim q_0}[\|\mathbf{y} - \mathbf{X}\boldsymbol{\varphi}\|_2^2] \leq 2\|\boldsymbol{\Sigma}^{-1}\|_{\mathrm{op}}(\|\mathbf{y}\|_2^2 + \mathbb{E}_{q_0}[\|\boldsymbol{\varphi}\|_2^2]\|\mathbf{X}\|_{\mathrm{op}}^2),$$

where $\|\mathbf{\Sigma}^{-1}\|_{\text{op}} = (\sigma^2 - \tau^2 \|\mathbf{X}\|_{\text{op}}^2)^{-1} < 1/(\delta \|\mathbf{X}\|_{\text{op}}^2) \le 1/(C_*^2 \delta)$ and $\|\mathbf{X}\|_{\text{op}}^2 \le C_*^2$ by Assumption A. Furthermore, in view of (1.2), $\|\mathbf{y}\|_2^2 \le 2(\|\mathbf{X}\|_{\text{op}}^2 \|\boldsymbol{\theta}\|_2^2 + \|\boldsymbol{\varepsilon}\|_2^2)$, where $\|\mathbf{X}\|_{\text{op}}^2 \|\boldsymbol{\theta}\|_2^2 \le C_*^2 M^2 p$ with probability one, and $\|\boldsymbol{\varepsilon}\|_2^2 \le 2\sigma^2 n$ with probability approaching 1 as $n \to \infty$. Thus the first term in (B.21) is at most $C(1 + \frac{n + \mathbb{E}_{q_0}[\|\boldsymbol{\varphi}\|_2^2)}{p})$ for some $C = C(\sigma, \delta, M, C_*)$. For the last term, we have $\log \det(\mathbf{\Sigma}) \le n \log \|\mathbf{\Sigma}\|_{\text{op}} \le n \log \sigma^2$, and combining these statements proves the desired (3.17).

B.5 Proof of Proposition 3.8

We need two lemmas before proving Proposition 3.8. The first lemma computes the Hessian of \bar{F}_n .

Lemma B.5. Let $g \in \mathcal{P}(M)$ and $\chi \in \Gamma$ be such that $g_t \equiv g + t\chi \in \mathcal{P}(M)$ for $t \in [0, 1]$. Then

$$\frac{\mathrm{d}^2}{\mathrm{d}t^2}\bar{F}_n(g_t) = \bar{H}_n[g_t](\chi)$$

and

$$\frac{\mathrm{d}^2}{\mathrm{d}t^2} \mathbb{E}_g \bar{F}_n(g_t) \Big|_{t=0} = \frac{1}{p} \int \frac{\left(\sum_{j=1}^p \int P(\mathbf{y}|\boldsymbol{\varphi}) \prod_{k:k \neq j} \mathcal{N}_\tau * g(\varphi_k) \cdot \mathcal{N}_\tau * \chi(\varphi_j) \mathrm{d}\boldsymbol{\varphi}\right)^2}{\int P(\mathbf{y}|\boldsymbol{\varphi}) \prod_{k=1}^p \mathcal{N}_\tau * g(\varphi_k) \mathrm{d}\boldsymbol{\varphi}} \mathrm{d}\mathbf{y}$$

Proof. Let $\langle \cdot \rangle_g$ denote the posterior average under $P_g(\varphi \mid \mathbf{y})$, i.e., for any $f : \mathbb{R}^p \to \mathbb{R}$,

$$\langle f \rangle_g = \frac{\int e^{-\frac{1}{2} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi}) + \sum_{j=1}^p \log \mathcal{N}_\tau * g(\varphi_j)} f(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi}}{\int e^{-\frac{1}{2} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \mathbf{X} \boldsymbol{\varphi}) + \sum_{j=1}^p \log \mathcal{N}_\tau * g(\varphi_j)} \mathrm{d}\boldsymbol{\varphi}}.$$

Then differentiating (2.10) yields that

$$\frac{\mathrm{d}}{\mathrm{d}t}\bar{F}(g_t) = -\frac{1}{p} \left\langle \sum_{j=1}^p \frac{\mathcal{N}_\tau * \chi}{\mathcal{N}_\tau * g_t}(\varphi_j) \right\rangle_{g_t},$$

and

$$\frac{\mathrm{d}^2}{\mathrm{d}t^2}\bar{F}_n(g_t) = \frac{1}{p} \left[\left\langle \sum_{j=1}^p \frac{\mathcal{N}_\tau * \chi}{\mathcal{N}_\tau * g_t}(\varphi_j) \right\rangle_{g_t}^2 + \left\langle \sum_{j=1}^p \left(\frac{\mathcal{N}_\tau * \chi}{\mathcal{N}_\tau * g_t}\right)^2(\varphi_j) \right\rangle_{g_t} - \left\langle \left(\sum_{j=1}^p \frac{\mathcal{N}_\tau * \chi}{\mathcal{N}_\tau * g_t}(\varphi_j)\right)^2 \right\rangle_{g_t} \right],$$

as desired. At t = 0 and $g_0 = g$, we have

$$\mathbb{E}_{g}\left[\left\langle\sum_{j=1}^{p}\frac{\mathcal{N}_{\tau}*\chi}{\mathcal{N}_{\tau}*g}(\theta_{j})\right\rangle_{g}^{2}\right] = \mathbb{E}_{g}\left[\left(\sum_{j=1}^{p}\frac{\int P(\mathbf{y}|\boldsymbol{\varphi})\prod_{k:k\neq j}\mathcal{N}_{\tau}*g(\varphi_{k})\cdot\mathcal{N}_{\tau}*\chi(\varphi_{j})\mathrm{d}\boldsymbol{\varphi}}{\int P(\mathbf{y}|\boldsymbol{\varphi})\prod_{k=1}^{p}\mathcal{N}_{\tau}*g(\varphi_{k})\mathrm{d}\boldsymbol{\varphi}}\right)^{2}\right]$$
$$=\int\frac{\left(\sum_{j=1}^{p}\int P(\mathbf{y}|\boldsymbol{\varphi})\prod_{k:k\neq j}\mathcal{N}_{\tau}*g(\varphi_{k})\cdot\mathcal{N}_{\tau}*\chi(\varphi_{j})\mathrm{d}\boldsymbol{\varphi}\right)^{2}}{\int P(\mathbf{y}|\boldsymbol{\varphi})\prod_{k=1}^{p}\mathcal{N}_{\tau}*g(\varphi_{k})\mathrm{d}\boldsymbol{\varphi}}\mathrm{d}\mathbf{y},$$

and, by the tower property of conditional expectation and independence of $(\varphi_1, \ldots, \varphi_p)$ under the prior,

$$\mathbb{E}_{g}\left\langle\sum_{j=1}^{p}\left(\frac{\mathcal{N}_{\tau}*\chi}{\mathcal{N}_{\tau}*g}\right)^{2}(\varphi_{j})\right\rangle_{g} = \mathbb{E}_{g}\left[\sum_{j=1}^{p}\left(\frac{\mathcal{N}_{\tau}*\chi}{\mathcal{N}_{\tau}*g}\right)^{2}(\varphi_{j})\right] = \mathbb{E}_{g}\left\langle\left(\sum_{j=1}^{p}\frac{\mathcal{N}_{\tau}*\chi}{\mathcal{N}_{\tau}*g}(\varphi_{j})\right)^{2}\right\rangle_{g},$$

where $\mathbb{E}_{g}\left(\sum_{j=1}^{p}\frac{\mathcal{N}_{\tau}*\chi}{\mathcal{N}_{\tau}*g}(\varphi_{j})\right)^{2}\right\rangle_{g}$

as desired.

We write $\bar{H}_n[g](\chi)(\mathbf{y})$ to make explicit the dependence of $\bar{H}_n[g](\chi)$ on the observed data \mathbf{y} . The following lemma establishes regularity of the function $\bar{H}_n[g](\chi)$. We present its proof after the main proof.

Lemma B.6. 1. The envelope function $\mathsf{F}(\mathbf{y}) = \sup_{g \in \mathcal{P}(M), \chi \in \Gamma} \overline{H}_n[g](\chi)(\mathbf{y})$ satisfies $\mathbb{E}_{g_*}\mathsf{F}(\mathbf{y}) < \infty$.

2. For any \mathbf{y} , the map $(g, \chi) \mapsto \overline{H}_n[g](\chi)(\mathbf{y})$ is jointly continuous on $\mathcal{P}(M) \times \Gamma$, with respect to the weak topology for $g \in \mathcal{P}(M)$ and the TV norm $\|\chi\|_{TV} = \int_{-M}^M |\chi(\mathrm{d}\theta)|$ for $\chi \in \Gamma$.

Proof of Proposition 3.8. Let $H_n(g) \equiv \inf_{\chi \in \Gamma} \overline{H}_n[g](\chi)$ and H(g) be defined similarly. Since Γ is compact, condition (3.20) yields that $H(g_*) > 0$. Since $\overline{H}_n[g](\chi)$ is jointly continuous in (g, χ) by Lemma B.6-(2), $\overline{H}[g](\chi)$ is also jointly continuous by the uniform convergence of $\{\overline{H}_n[g](\chi)\}_n$ in (3.19). Then since Γ is compact, H(g) is also continuous in g. Recall the definition of d_v in (A.1), which metrizes weak convergence [GG20]. Then, fixing a constant v > 0 to be specified later, there exists some (n, p)-independent $\delta > 0$ such that H(g) > 0 whenever $g \in B_v(\delta; g_*) = \{\overline{g} \in \mathcal{P}(M) :$ $d_v(\overline{g}, g_*) \leq \delta\}$. For any $g \in B_v(\delta; g_*)$, if $\chi_0 = \arg \min_{\chi \in \Gamma} \overline{H}[g](\chi)$, then

$$H_n(g) - H(g) = \min_{\chi \in \Gamma} \bar{H}_n[g](\chi) - \bar{H}[g](\chi_0) \le \bar{H}_n[g](\chi_0) - \bar{H}[g](\chi_0).$$

An analogous lower bound concludes

$$\sup_{g\in B_v(\delta;g_*)} \left| H_n(g) - H(g) \right| \le \sup_{g\in B_v(\delta;g_*),\chi\in\Gamma} \left| \bar{H}_n[g](\chi) - \bar{H}[g](\chi) \right|.$$

Hence by (3.19),

$$\sup_{g \in B_v(\delta;g_*)} |H_n(g) - H(g)| \stackrel{P}{\to} 0$$

so there exists an event with probability converging to 1 on which

$$H_n(g) \ge H(g)/2 > 0$$
 for all $g \in B_v(\delta; g_*).$ (B.22)

Now observe that $B_v(\delta; g_*)$ is a convex set, by definition of d_v in (A.1) and convexity of the map $g \mapsto W_1(\mathcal{N}_v * g_*, \mathcal{N}_v * g)$. Fix any such $g_0, g_1 \in B_v(\delta; g_*)$ and let $s(t) \equiv \overline{F}_n((1-t)g_0 + tg_1)$. Then by (B.22) and Lemma B.5,

$$s''(t) = \bar{H}_n[g_t](g_1 - g_0) = \bar{H}_n[g_t] \left(\frac{g_1 - g_0}{\|g_1 - g_0\|_{\mathrm{TV}}}\right) \cdot \|g_1 - g_0\|_{\mathrm{TV}}^2 \ge 0,$$

which establishes the desired convexity for all $g \in B_v(\delta; g_*)$.

Lastly, note that since g_* is the unique minimizer of F(g), there exists an (n, p)-independent constant $c_0 > 0$ such that

$$\left\{g: \bar{F}(g) \leq \bar{F}(g_*) + c_0\right\} \subset B_v(\delta; g_*)$$

Indeed, otherwise there would exist a sequence $\{g_t\}$ outside $B_v(\delta; g_*)$ satisfying $\bar{F}(g_t) \leq \bar{F}(g_*) + t^{-1}$. By weak compactness of $\mathcal{P}(M)$, this sequence has a limit point g', which by weak continuity of \bar{F} must satisfy $\bar{F}(g') \leq \bar{F}(g_*)$, contradicting uniqueness of g_* as the minimizer of \bar{F} . By the uniform convergence of $\{\bar{F}_n(g)\}_n$ in (3.19), for all large n, p it holds

$$\mathcal{K}_n := \left\{ g : \bar{F}_n(g) \le \bar{F}_n(g_*) + c_0/2 \right\} \subset \left\{ g : \bar{F}(g) \le \bar{F}(g_*) + c_0 \right\} \subset B_v(\delta; g_*).$$

Similarly, $\mathcal{K} := \{g : \overline{F}(g) < \overline{F}(g_*) + c_0/4\} \subset \mathcal{K}_n$, where \mathcal{K} is a (n, p)-independent weak neighborhood of g_* by weak continuity of \overline{F} , as desired.

Proof of Lemma B.6. To prove claim (1), note that for any $\chi \in \Gamma$ and $g \in \mathcal{P}(M)$,

$$\left|\frac{\mathcal{N}_{\tau} * \chi}{\mathcal{N}_{\tau} * g}(\varphi_{j})\right| = \frac{\left|\int_{-M}^{M} e^{-\frac{u^{2}}{2\tau^{2}} + \frac{\varphi_{j}u}{\tau^{2}}} \chi(u) \mathrm{d}u\right|}{\int_{-M}^{M} e^{-\frac{u^{2}}{2\tau^{2}} + \frac{\varphi_{j}u}{\tau^{2}}} g(u) \mathrm{d}u} \le C e^{C|\varphi_{j}|} \cdot \|\chi\|_{\mathrm{TV}} \le C e^{C\|\varphi\|}.$$
 (B.23)

Hence by Lemma B.7 to follow (with Γ and $c(\theta)$ as defined therein),

$$\left\langle e^{C\|\boldsymbol{\varphi}\|} \right\rangle_{g} = \int_{[-M,M]^{p}} P_{g}(\boldsymbol{\theta} \mid \mathbf{y}) \mathbb{E}_{\boldsymbol{\varphi} \sim \mathcal{N}(c(\boldsymbol{\theta}), \boldsymbol{\Gamma})} e^{C\|\boldsymbol{\varphi}\|} \mathrm{d}\boldsymbol{\theta} \le C \sup_{\boldsymbol{\theta} \in [-M,M]^{p}} e^{C\|c(\boldsymbol{\theta})\|} \le C' e^{C'\|\mathbf{y}\|}, \quad (B.24)$$

where C, C' may depend on $(\|\mathbf{X}\|_{op}, \sigma^2, \tau^2, p, M)$. Since $\mathbf{y} \stackrel{d}{=} \mathbf{X}\boldsymbol{\theta} + \boldsymbol{\varepsilon}$ with $\theta_j \stackrel{iid}{\sim} g_*$ and $\varepsilon_i \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2)$, $\sup_{g \in \mathcal{P}(M), \chi \in \Gamma} \left| \frac{\mathcal{N}_{\tau} * \chi}{\mathcal{N}_{\tau} * g} (\varphi_j) \right|^2$ and hence $\mathsf{F}(\mathbf{y})$ is integrable, establishing claim (1). Next we establish claim (2). For presentational simplicity, we only prove the joint continuity of

Next we establish claim (2). For presentational simplicity, we only prove the joint continuity of the first term of $\bar{H}_n[g](\chi)(\mathbf{y})$ in (3.18), and that of the other two terms follows similarly. It suffices to prove the continuity for the first summand there, which is

$$H(g,\chi;\mathbf{y}) \equiv \left\langle \left(\frac{\mathcal{N}_{\tau} * \chi}{\mathcal{N}_{\tau} * g}\right)^2 (\varphi_1) \right\rangle_g = \frac{\int P(\mathbf{y}|\boldsymbol{\varphi}) \prod_{j=1}^p \mathcal{N}_{\tau} * g(\varphi_j) \cdot \left(\frac{\mathcal{N}_{\tau} * \chi}{\mathcal{N}_{\tau} * g}\right)^2 (\varphi_1) \mathrm{d}\boldsymbol{\varphi}}{P_g(\mathbf{y})}.$$

For two pairs $(g_i, \chi_i), i = 1, 2, |H(g_1, \chi_1) - H(g_2, \chi_2)| \le \sum_{i=1}^4 (I_i)$, where

$$(I_{1}) = \left(\frac{1}{P_{g_{1}}(\mathbf{y})} - \frac{1}{P_{g_{2}}(\mathbf{y})}\right) \int P(\mathbf{y}|\boldsymbol{\varphi}) \prod_{j=1}^{p} \mathcal{N}_{\tau} * g_{1}(\varphi_{j}) \cdot \left(\frac{\mathcal{N}_{\tau} * \chi_{1}}{\mathcal{N}_{\tau} * g_{1}}\right)^{2} (\varphi_{1}) \mathrm{d}\boldsymbol{\varphi},$$

$$(I_{2}) = P_{g_{2}}(\mathbf{y})^{-1} \int P(\mathbf{y}|\boldsymbol{\varphi}) \left(\prod_{j=1}^{p} \mathcal{N}_{\tau} * g_{1}(\varphi_{j}) - \prod_{j=1}^{p} \mathcal{N}_{\tau} * g_{2}(\varphi_{j})\right) \cdot \left(\frac{\mathcal{N}_{\tau} * \chi_{1}}{\mathcal{N}_{\tau} * g_{1}}\right)^{2} (\varphi_{1}) \mathrm{d}\boldsymbol{\varphi},$$

$$(I_{3}) = P_{g_{2}}(\mathbf{y})^{-1} \int \prod_{j=1}^{p} \mathcal{N}_{\tau} * g_{2}(\varphi_{j}) \cdot \left[\left(\frac{\mathcal{N}_{\tau} * \chi_{1}}{\mathcal{N}_{\tau} * g_{1}}\right)^{2} (\varphi_{1}) - \left(\frac{\mathcal{N}_{\tau} * \chi_{1}}{\mathcal{N}_{\tau} * g_{2}}\right)^{2} (\varphi_{1})\right] \mathrm{d}\boldsymbol{\varphi},$$

$$(I_{4}) = P_{g_{2}}(\mathbf{y})^{-1} \int P(\mathbf{y}|\boldsymbol{\varphi}) \prod_{j=1}^{p} \mathcal{N}_{\tau} * g_{2}(\varphi_{j}) \cdot \frac{\left(\mathcal{N}_{\tau} * \chi_{1}\right)^{2} (\varphi_{1}) - \left(\mathcal{N}_{\tau} * \chi_{2}\right)^{2} (\varphi_{1})}{\left(\mathcal{N}_{\tau} * g_{2}\right)^{2} (\varphi_{1})} \mathrm{d}\boldsymbol{\varphi}.$$

We will show that (I_1) - (I_4) converge to 0 when (g_1, χ_1) converges to (g_2, χ_2) under the weak topology for g and TV norm for χ .

Bound on (I_1) For i = 1, 2, let μ_i denote the law of $\mathbf{X}\boldsymbol{\theta}$ on \mathbb{R}^n with $\theta_j \stackrel{iid}{\sim} g_i$, then μ_i is a measure supported in the Euclidean ball $B_n(0, \sqrt{p} || \mathbf{X} ||_{\text{op}} M)$. It is clear that μ_1 converges weakly to μ_2 if g_1 converges weakly to g_2 . Using

$$P_{g_1}(\mathbf{y}) - P_{g_2}(\mathbf{y}) = (2\pi\sigma^2)^{-\frac{n}{2}} \int e^{-\frac{\|\mathbf{y}-\mathbf{u}\|^2}{2\sigma^2}} (\mu_1 - \mu_2) (\mathrm{d}\mathbf{u}),$$

we have

$$(I_1) = \frac{P_{g_2}(\mathbf{y}) - P_{g_1}(\mathbf{y})}{P_{g_2}(\mathbf{y})} \left\langle \left(\frac{\mathcal{N}_{\tau} * \chi_1}{\mathcal{N}_{\tau} * g_1}\right)^2 (\varphi_1) \right\rangle_{g_1} = (2\pi\sigma^2)^{-\frac{n}{2}} \int f(\mathbf{u})(\mu_2 - \mu_1)(\mathrm{d}\mathbf{u}),$$

where

$$f(\mathbf{u}) = \frac{\left\langle \left(\frac{\mathcal{N}_{\tau} * \chi_1}{\mathcal{N}_{\tau} * g_1}\right)^2 (\varphi_1) \right\rangle_{g_1}}{P_{g_2}(\mathbf{y})} e^{-\frac{\|\mathbf{y} - \mathbf{u}\|^2}{2\sigma^2}}.$$

By (B.23)-(B.24), $\left|\left\langle \left(\frac{N_{\tau}*\chi_1}{N_{\tau}*g_1}\right)^2(\varphi_1)\right\rangle_{g_1}\right| \leq C$, and similarly $P_{g_2}(\mathbf{y}) \geq C$, where the constant C may depend on \mathbf{y} but does not depend on (g_i, χ_i) . Hence $f(\mathbf{u})$ admits an upper bound independent of (g_i, χ_i) , and $(I_1) \to 0$ as (g_1, χ_1) converges to (g_2, χ_2) .

Bound on (I_2) We have by telescoping

$$(I_2) \leq \frac{1}{P_{g_2}(\mathbf{y})} \cdot \sum_{j=1}^p \int P(\mathbf{y}|\boldsymbol{\varphi}) \mathcal{N}_{\tau} * (g_1 - g_2)(\varphi_j) \prod_{k:k \neq j} \mathcal{N}_{\tau} * \widetilde{g}(\varphi_k) \cdot \left(\frac{\mathcal{N}_{\tau} * \chi_1}{\mathcal{N}_{\tau} * g_1}\right)^2 (\varphi_1) \mathrm{d}\boldsymbol{\varphi},$$

where $\tilde{g} \in \{g_1, g_2\}$. Fix any $j \in [p]$, and note that the *j*th term equals some constant multiple of $\int f(u)(g_1 - g_2)(du)$, where

$$f(u) = \int P(\mathbf{y}|\boldsymbol{\varphi}) e^{-\frac{(\varphi_j - u)^2}{2\tau^2}} \prod_{k:k \neq j} \mathcal{N}_{\tau} * \widetilde{g}(\varphi_k) \cdot \left(\frac{\mathcal{N}_{\tau} * \chi_1}{\mathcal{N}_{\tau} * g_1}\right)^2 (\varphi_1) d\boldsymbol{\varphi}$$

$$\leq C \int e^{C|\varphi_1|} \prod_{k:k \neq j} \mathcal{N}_{\tau} * \widetilde{g}(\varphi_k) \cdot e^{-\frac{(\varphi_j - u)^2}{2\tau^2}} \mathrm{d}\varphi \leq C e^{C|u|},$$

using (B.23)-(B.24). This and the lower bound of $P_{g_2}(\mathbf{y})$ above imply f(u) is bounded on [-M, M] independently of (g_i, χ_i) , and hence $(I_2) \to 0$ as (g_1, χ_1) converges to (g_2, χ_2) .

Bound on (I₃) Note that (I₃) equals a constant multiple of $\int f(u)(g_1 - g_2)(du)$, where

$$\begin{aligned} \mathsf{f}(u) &= \frac{1}{P_{g_2}(\mathbf{y})} \int P(\mathbf{y}|\boldsymbol{\varphi}) \prod_{j=1}^p \mathcal{N}_\tau * g_2(\varphi_j) \frac{(\mathcal{N}_\tau * \chi)^2(\varphi_1) \cdot [\mathcal{N}_\tau * (g_1 + g_2)](\varphi_1)}{(\mathcal{N}_\tau * g_1)^2(\varphi_1)(\mathcal{N}_\tau * g_2)^2(\varphi_1)} e^{-\frac{(\varphi_1 - u)^2}{2\tau^2}} \mathrm{d}\boldsymbol{\varphi} \\ &\leq \int C e^{C|\varphi_1|} e^{-\frac{(\varphi_1 - u)^2}{2\tau^2}} \mathrm{d}\varphi_1 \leq C e^{C|u|}, \end{aligned}$$

yielding $(I_3) \to 0$ when (g_1, χ_1) converges to (g_2, χ_2) .

Bound on (I_4) Note that (I_4) is bounded by a constant multiple of $\int |f(u)||(\chi_1 - \chi_2)(du)|$, where

$$\mathsf{f}(u) = \frac{1}{P_{g_2}(\mathbf{y})} \int P(\mathbf{y}|\boldsymbol{\varphi}) \prod_{j=1}^p \mathcal{N}_\tau * g_2(\varphi_j) \cdot \frac{\left[\mathcal{N}_\tau * (\chi_1 + \chi_2)\right](\varphi_1)}{\left(\mathcal{N}_\tau * g_2\right)^2(\varphi_1)} e^{-\frac{(\varphi_1 - u)^2}{2\tau^2}} \mathrm{d}\boldsymbol{\varphi} \le C e^{C|u|},$$

using the same argument as (I_3) . Combining the bounds for (I_1) - (I_4) completes the proof for claim (2).

Lemma B.7. For any density $g(\cdot)$ on [-M, M], it holds that

$$P_g(\boldsymbol{\varphi} \mid \mathbf{y}) = \int_{[-M,M]^p} \phi_{\Gamma}(\boldsymbol{\varphi} - c(\boldsymbol{\theta})) P_g(\boldsymbol{\theta} \mid \mathbf{y}) \mathrm{d}\boldsymbol{\theta},$$

where $\phi_{\Gamma}(\cdot)$ denotes the density of $\mathcal{N}(0,\Gamma)$ on \mathbb{R}^p , and

$$\Gamma = \frac{\tau^2}{\sigma^2} (\sigma^2 \operatorname{Id} - \tau^2 \mathbf{X}^\top \mathbf{X}), \quad c(\boldsymbol{\theta}) = \frac{\tau^2}{\sigma^2} \mathbf{X}^\top \mathbf{y} + \boldsymbol{\theta} - \frac{\tau^2}{\sigma^2} \mathbf{X}^\top \mathbf{X} \boldsymbol{\theta}.$$

Proof. We have

$$P_g(\boldsymbol{\varphi} \mid \mathbf{y}) = \int P_g(\boldsymbol{\varphi}, \boldsymbol{\theta} \mid \mathbf{y}) \mathrm{d}\boldsymbol{\theta} = \int P_g(\boldsymbol{\varphi} \mid \boldsymbol{\theta}, \mathbf{y}) P_g(\boldsymbol{\theta} \mid \mathbf{y}) \mathrm{d}\boldsymbol{\theta},$$

and direct calculation yields that $P_g(\boldsymbol{\varphi} \mid \boldsymbol{\theta}, \mathbf{y}) = \mathcal{N}(c(\boldsymbol{\theta}), \boldsymbol{\Gamma})$ with $\boldsymbol{\Gamma} = (\mathbf{X}^{\top} \boldsymbol{\Sigma}^{-1} \mathbf{X} + \tau^{-2} \operatorname{Id})^{-1}$ and $c(\boldsymbol{\theta}) = \boldsymbol{\Gamma} (\mathbf{X}^{\top} \boldsymbol{\Sigma}^{-1} \mathbf{y} + \tau^{-2} \boldsymbol{\theta})$. To simplify the expression for $\boldsymbol{\Gamma}$, recall that the Woodbury matrix identity reads $(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)^{-1}VA^{-1}$, provided that A is invertible. Apply this formula with $A = \tau^{-2} \operatorname{Id}, U = \mathbf{X}^{\top}, C = \mathbf{\Sigma}^{-1}, V = \mathbf{X}$ yields that

$$\boldsymbol{\Gamma} = \tau^2 \operatorname{Id} - \tau^4 \mathbf{X}^\top (\boldsymbol{\Sigma} + \tau^2 \mathbf{X} \mathbf{X}^\top)^{-1} \mathbf{X} = \tau^2 \operatorname{Id} - \frac{\tau^4}{\sigma^2} \mathbf{X}^\top \mathbf{X} = \frac{\tau^2}{\sigma^2} (\sigma^2 \operatorname{Id} - \tau^2 \mathbf{X}^\top \mathbf{X}),$$

as desired. We can then simplify $c(\boldsymbol{\theta})$ as

$$c(\boldsymbol{\theta}) = \frac{\tau^2}{\sigma^2} (\sigma^2 \operatorname{Id} - \tau^2 \mathbf{X}^\top \mathbf{X}) \mathbf{X}^\top \boldsymbol{\Sigma}^{-1} \mathbf{y} + \frac{1}{\sigma^2} (\sigma^2 - \tau^2 \mathbf{X}^\top \mathbf{X}) \boldsymbol{\theta}$$

$$= \frac{\tau^2}{\sigma^2} \mathbf{X}^\top (\sigma^2 \operatorname{Id} - \tau^2 \mathbf{X} \mathbf{X}^\top) \boldsymbol{\Sigma}^{-1} \mathbf{y} + \boldsymbol{\theta} - \frac{\tau^2}{\sigma^2} \mathbf{X}^\top \mathbf{X} \boldsymbol{\theta} = \frac{\tau^2}{\sigma^2} \mathbf{X}^\top \mathbf{y} + \boldsymbol{\theta} - \frac{\tau^2}{\sigma^2} \mathbf{X}^\top \mathbf{X} \boldsymbol{\theta}.$$

54

Finally we illustrate a simple setting where the assumptions of Proposition 3.8 hold, where \mathbf{X} is a block design.

Proposition B.8. Suppose $\mathbf{X} \in \mathbb{R}^{n \times p}$ is block diagonal with N identical blocks $\bar{\mathbf{X}} \in \mathbb{R}^{B_1 \times B_2}$, where $\bar{\mathbf{X}}$ and (B_1, B_2) are fixed independent of (n, p), $\bar{\mathbf{X}} \neq 0$, and $(n, p) = (NB_1, NB_2)$. Then there exists a set in $\mathbb{R}^{B_1 \times B_2}$ of full Lebesgue measure such that for every $\bar{\mathbf{X}}$ in this set, the assumptions of Proposition 3.8 are satisfied with $\bar{F} = \mathbb{E}\bar{F}_n$ and $\bar{H} = \mathbb{E}\bar{H}_n$.

Proof. We verify uniform convergence of \overline{H}_n to \overline{H} ; a similar argument holds for \overline{F}_n , which we omit for brevity. Partition the observation $\mathbf{y} = (\overline{\mathbf{y}}^1, \dots, \overline{\mathbf{y}}^N)$ according to the blocks, with $\overline{\mathbf{y}}^i \in \mathbb{R}^{B_1}$, so that each block is generated by

$$\bar{\mathbf{y}}^i \stackrel{d}{=} \bar{\mathbf{X}} \bar{\boldsymbol{\varphi}}^i + \bar{\boldsymbol{\varepsilon}}^i, \quad (\bar{\varphi}^i_1, \dots, \bar{\varphi}^i_{B_2}) \stackrel{iid}{\sim} g_*, \quad \bar{\boldsymbol{\varepsilon}}^i \sim \mathcal{N}(0, \bar{\mathbf{\Sigma}}) \text{ with } \bar{\mathbf{\Sigma}} = \sigma^2 \operatorname{Id}_{B_1} - \tau^2 \bar{\mathbf{X}} \bar{\mathbf{X}}^\top.$$
 (B.25)

Note that under this block design,

$$\bar{H}_n[g](\chi) = \frac{1}{N} \sum_{i=1}^N \mathsf{h}_{g,\chi}(\bar{\mathbf{y}}^i) \quad \text{and} \quad \bar{H}[g](\chi) = \mathbb{E}_{g_*}\bar{H}_n[g](\chi) = \mathbb{E}_{g_*}\mathsf{h}_{g,\chi}(\bar{\mathbf{y}}), \tag{B.26}$$

where, with the notation

$$\langle f \rangle_g = \frac{\int f(\bar{\varphi}) e^{-\frac{1}{2}(\bar{\mathbf{y}} - \bar{\mathbf{X}}\bar{\varphi})^\top \bar{\boldsymbol{\Sigma}}^{-1}(\bar{\mathbf{y}} - \bar{\mathbf{X}}\bar{\varphi}) + \sum_{j=1}^{B_2} \log \mathcal{N}_\tau * g(\bar{\varphi}_j)}{\int e^{-\frac{1}{2}(\bar{y} - \bar{\mathbf{X}}\bar{\varphi})^\top \bar{\boldsymbol{\Sigma}}^{-1}(\bar{\mathbf{y}} - \bar{\mathbf{X}}\bar{\varphi}) + \sum_{j=1}^{B_2} \log \mathcal{N}_\tau * g(\bar{\varphi}_j)} \mathrm{d}\bar{\varphi}}, \quad f: \mathbb{R}^{B_2} \to \mathbb{R},$$

we define

$$\mathsf{h}_{g,\chi}(\bar{\mathbf{y}}) = \frac{1}{B_2} \left[\left\langle \sum_{j=1}^{B_2} \left(\frac{\mathcal{N}_\tau * \chi}{\mathcal{N}_\tau * g}(\bar{\varphi}_j) \right)^2 \right\rangle_g - \left\langle \left(\sum_{j=1}^{B_2} \frac{\mathcal{N}_\tau * \chi}{\mathcal{N}_\tau * g}(\bar{\varphi}_j) \right)^2 \right\rangle_g + \left\langle \sum_{j=1}^{B_2} \frac{\mathcal{N}_\tau * \chi}{\mathcal{N}_\tau * g}(\bar{\varphi}_j) \right\rangle_g^2 \right].$$
(B.27)

It is clear that $\bar{H}[g](\chi)$ is an (n, p)-independent function. To show uniform convergence of $\bar{H}_n[g](\chi)$, it suffices to establish that

$$\mathcal{H} \equiv \left\{ \bar{\mathbf{y}} \mapsto \mathsf{h}_{g,\chi}(\bar{\mathbf{y}}) : g \in \mathcal{P}(M), \chi \in \Gamma \right\}$$

is Glivenko-Cantelli with respect to the measure $P_{g_*}(\bar{\mathbf{y}})$. Since the parameter space $\Theta \equiv \{(g, \chi) : g \in \mathcal{P}(M), \chi \in \Gamma\}$ is compact, a sufficient condition (see [Wel05, Lemma 6.1] and [VW23, Theorem 2.4.1]) is to show: (i) For every $\bar{\mathbf{y}}, (g, \chi) \mapsto \mathsf{h}_{g,\chi}(\bar{\mathbf{y}})$ is jointly continuous; (ii) the envelope function $\mathsf{F}(\bar{\mathbf{y}}) \equiv \sup_{(g,\chi)\in\Theta} \mathsf{h}_{g,\chi}(\bar{\mathbf{y}})$ is P_{g_*} -integrable. These conditions are established by Lemma B.6, so uniform convergence follows.

Next we show g_* is the unique global minimizer of $\overline{F}(g)$, whenever $\overline{\mathbf{X}}$ contains a row $\mathbf{x}_{i_0} \in \mathbb{R}^{B_2}$ such that

$$\sum_{j=1}^{B_2} x_{i_0,j}^q \neq 0, \quad \text{for all } q = 1, 2, 3, \dots$$
(B.28)

It is clear that the set of such $\bar{\mathbf{X}}$ has full Lebesgue measure. Since $\bar{F}(g) = D_{\mathrm{KL}}(P_{g_*}(\bar{\mathbf{y}})||P_g(\bar{\mathbf{y}}))$, it suffices to show that the map $g \mapsto P_g(\bar{\mathbf{y}})$ (the image as a distribution) is injective. Since the reduced model (B.25) is equivalent to $\bar{\mathbf{y}} = \bar{\mathbf{X}}\bar{\boldsymbol{\theta}} + \mathcal{N}(0, \sigma^2 \operatorname{Id}_{B_1})$, it suffices to show $g \mapsto P_g(\bar{\mathbf{X}}\bar{\boldsymbol{\theta}})$ is injective.

We prove by contradiction. Suppose there are g, \tilde{g} such that $\bar{\mathbf{X}}\bar{\boldsymbol{\theta}}$ has the same distribution when the components of $\bar{\boldsymbol{\theta}}$ are drawn i.i.d. from g, \tilde{g} , then $\mathbf{x}_{i_0}^{\top} \bar{\boldsymbol{\theta}}$ also has the same distribution. Let $m_q(g)$ and $m_q(\tilde{g})$ denote the *q*th moment of g, \tilde{g} respectively. We show by induction that $m_q(g) = m_q(\tilde{g})$ for all $q \geq 1$. The base case q = 1 holds because

$$\mathbb{E}_{g}\mathbf{x}_{i_{0}}^{\top}\bar{\boldsymbol{\theta}} = m_{1}(g) \cdot \sum_{j=1}^{B_{2}} x_{i_{0},j}, \quad \mathbb{E}_{\widetilde{g}}\mathbf{x}_{i_{0}}^{\top}\bar{\boldsymbol{\theta}} = m_{1}(\widetilde{g}) \cdot \sum_{j=1}^{B_{2}} x_{i_{0},j},$$

and the common factor $\sum_{j=1}^{B_2} x_{i_0,j}$ is not zero by (B.28). Suppose the moments match up to order q, then

$$\mathbb{E}_g \left(\mathbf{x}_{i_0}^\top \bar{\boldsymbol{\theta}} \right)^{q+1} = m_{q+1}(g) \cdot \sum_{j=1}^p x_{i_0,j}^{q+1} + R_q,$$

where R_q is some term that only depends on the first q moments of g. In view of (B.28), it holds $m_{q+1}(g) = m_{q+1}(\tilde{g})$, completing the induction. The conclusion now follows because boundedness of $\mathbf{x}_{i_0}^{\top} \bar{\boldsymbol{\theta}}$ implies that its distribution is completely determined by its moments ([Bil95, Theorem 30.1]).

Finally let us check the lower bound (3.20). Note that $\bar{H}_n[g](\chi) \ge 0$ by definition, so also $\bar{H}[g](\chi) \ge 0$, and it suffices to show $\bar{H}[g_*](\chi) \ne 0$ for all $\chi \in \Gamma$. Suppose by contradiction that $\bar{H}[g_*](\chi) = 0$ for some $\chi \ne 0$. Then Lemma B.5 implies that with probability 1 over $\bar{\mathbf{y}}$,

$$\sum_{j=1}^{B_2} \int P(\bar{\mathbf{y}}|\bar{\boldsymbol{\varphi}}) \prod_{k:k\neq j} \mathcal{N}_\tau * g_*(\bar{\varphi}_k) \cdot \mathcal{N}_\tau * \chi(\bar{\varphi}_j) \mathrm{d}\bar{\boldsymbol{\varphi}} = 0.$$

Take the same i_0 as in (B.28). Then, integrating over the variables $\{y_i : i \neq i_0\}$, the above identity yields that

$$\frac{1}{B_2}\sum_{j=1}^{B_2}\int P(\bar{y}_{i_0}|\bar{\varphi})\prod_{k:k\neq j}\mathcal{N}_{\tau}*g_*(\bar{\varphi}_k)\cdot\mathcal{N}_{\tau}*\chi(\bar{\varphi}_j)\mathrm{d}\bar{\varphi}=0,$$

or equivalently,

$$\frac{1}{B_2}\sum_{j=1}^{B_2}\int P(\bar{y}_{i_0}|\bar{\boldsymbol{\theta}})\prod_{k:k\neq j}g_*(\bar{\theta}_k)\cdot\chi(\bar{\theta}_j)\mathrm{d}\bar{\boldsymbol{\theta}}=0$$

Let h_+ and h_- be positive measures such that $\chi = h_+ - h_-$. Since $\int \chi = 0$, we have $\int h_+ = \int h_-$. Let $\tilde{h}_+ = h_+ / \int h_+$ and $\tilde{h}_- = h_- / \int h_-$ be the normalized probability measures. Then the above identity yields that

$$\int \frac{1}{B_2} \sum_{j=1}^{B_2} P(\bar{y}_{i_0}|\bar{\boldsymbol{\theta}}) \prod_{k:k\neq j} g_*(\bar{\theta}_k) \cdot \tilde{h}_+(\bar{\theta}_j) \mathrm{d}\bar{\boldsymbol{\theta}} = \int \frac{1}{B_2} \sum_{j=1}^{B_2} P(\bar{y}_{i_0}|\bar{\boldsymbol{\theta}}) \prod_{k:k\neq j} g_*(\bar{\theta}_k) \cdot \tilde{h}_-(\bar{\theta}_j) \mathrm{d}\bar{\boldsymbol{\theta}}.$$

Note that (B.25) is equivalent to $y_{i_0}|\bar{\boldsymbol{\theta}} \stackrel{d}{=} \mathbf{x}_{i_0}^\top \bar{\boldsymbol{\theta}} + \mathcal{N}(0, \sigma^2)$, so this implies that the distribution of $\mathbf{x}_{i_0}^\top \bar{\boldsymbol{\theta}}$ is the same under the following two priors on $\bar{\boldsymbol{\theta}}$:

$$P: J \sim \text{Unif}([B_2]), \quad (\bar{\theta}|J=j) \sim \prod_{k:k \neq j} g_*(\bar{\theta}_k) \cdot \tilde{h}_+(\theta_k),$$

$$Q: J \sim \text{Unif}([B_2]), \quad (\bar{\theta}|J=j) \sim \prod_{k:k \neq j} g_*(\bar{\theta}_k) \cdot \tilde{h}_-(\theta_k).$$

Let $m_q(\tilde{h}_+)$ and $m_q(\tilde{h}_-)$ denote the q^{th} moment of \tilde{h}_+ and \tilde{h}_- . Then induction again implies $m_q(\tilde{h}_+) = m_q(\tilde{h}_-)$ under (B.28). Indeed, the base case follows from the same argument as above, and for the induction step,

$$\mathbb{E}_{P}(\mathbf{x}_{i_{0}}^{\top}\bar{\boldsymbol{\theta}})^{q+1} = m_{q+1}(\widetilde{h}_{+}) \cdot \frac{1}{B_{2}} \sum_{j=1}^{p} x_{i_{0},j}^{q+1} + R_{q},$$

where R_q is again some term that only depends on the first q moments of \tilde{h}_+ , concluding the induction. Since the distributions of \tilde{h}_+, \tilde{h}_- are determined by their moments ([Bil95, Theorem 30.1]), the above arguments yield $h_+ = h_-$ and hence $\chi \equiv 0$, leading to a contradiction.

C Solution properties of gradient flows

In this appendix, we prove Theorem 3.6. Throughout this appendix, $n, p, \mathbf{X}, \mathbf{y}$ are fixed, and C, C', C_0, K etc. denote constants that may depend on $n, p, \mathbf{X}, \mathbf{y}$ unless otherwise specified.

By a solution $\{q_t(\cdot), g_t(\cdot)\}_{t \in [0,T]}$ to (2.15-2.16), we mean functions $t \mapsto g_t(\theta)$ in $C^1([0,T])$ for each $\theta \in [-M, M]$ and $(t, \varphi) \mapsto q_t(\varphi)$ in $C^{1,2}([0,T] \times \mathbb{R}^p)$, such that q_t, g_t are probability densities for all $t \in [0,T]$ and (2.15-2.16) hold in the classical sense of pointwise equality. We call the solution unique if for any other such solution $\{\tilde{q}_t, \tilde{g}_t\}_{t \in [0,T]}$, we have $q_t(\varphi) = \tilde{q}_t(\varphi)$ and $g_t(\theta) = \tilde{g}_t(\theta)$ for all $t \in [0,T], \theta \in [-M, M]$, and $\varphi \in \mathbb{R}^p$.

We equip $\mathcal{P}(M)$ with the total variation metric $d_{\mathrm{TV}}(g, \tilde{g}) = \sup_{A \subseteq [-M,M]} |g(A) - \tilde{g}(A)|$, or equivalently $d_{\mathrm{TV}}(g, \tilde{g}) = \frac{1}{2} \int_{-M}^{M} |g(\theta) - \tilde{g}(\theta)| \mathrm{d}\theta$ when $g, \tilde{g} \in \mathcal{P}_*(M)$ both admit densities. Then $\mathcal{P}(M)$ is complete under d_{TV} . For any T > 0, let

$$\mathcal{L}_g \equiv \mathcal{L}_g(T) = C([0,T], \mathcal{P}(M)),$$

the space of continuous (with respect to d_{TV}) $\mathcal{P}(M)$ -valued processes $\{g_t\}_{t\in[0,T]}$, equipped with the metric $d_{\infty,\text{TV}}(g,\tilde{g}) = \sup_{t\in[0,T]} d_{\text{TV}}(g_t,\tilde{g}_t)$. Then also \mathcal{L}_g is complete under $d_{\infty,\text{TV}}$.

Next, let

$$E_q \equiv E_q(T) = C([0, T], \mathbb{R}^p),$$

the space of continuous \mathbb{R}^p -valued sample paths $\{\varphi_t\}_{t\in[0,T]}$, equipped analogously with the norm $\|\varphi\|_{\infty,2} = \sup_{t\in[0,T]} \|\varphi_t\|_2$. Let $\mathcal{L}_q \equiv \mathcal{L}_q(T)$ be the space of probability distributions on E_q for which $\mathbb{E}[\|\varphi\|_{\infty,2}^2] < \infty$, equipped with the Wasserstein-2 distance induced by this norm,

$$d_W(q, \widetilde{q}) = \inf_{\boldsymbol{\varphi} = \{\boldsymbol{\varphi}_t\}_{t \in [0,T]} \sim q, \, \widetilde{\boldsymbol{\varphi}} = \{\widetilde{\boldsymbol{\varphi}}_t\}_{t \in [0,T]} \sim \widetilde{q}} \left(\mathbb{E} \| \boldsymbol{\varphi} - \widetilde{\boldsymbol{\varphi}} \|_{\infty,2}^2 \right)^{1/2}$$
$$= \inf_{\boldsymbol{\varphi} = \{\boldsymbol{\varphi}_t\}_{t \in [0,T]} \sim q, \, \widetilde{\boldsymbol{\varphi}} = \{\widetilde{\boldsymbol{\varphi}}_t\}_{t \in [0,T]} \sim \widetilde{q}} \left(\mathbb{E} \sup_{t \in [0,T]} \| \boldsymbol{\varphi}_t - \widetilde{\boldsymbol{\varphi}}_t \|_2^2 \right)^{1/2}$$
(C.1)

where the infimum is taken over all couplings of $q, \tilde{q} \in \mathcal{L}_q$. Then E_q is complete, and hence \mathcal{L}_q is also complete by [Vil09, Theorem 6.18]. We will write $\{q_t\}_{t \in [0,T]}$ for the marginal laws of $q \in \mathcal{L}_q$.

C.1 Solution properties of the g-flow

The following result gives the solution properties of the Fisher-Rao g-flow for a fixed law $q \in \mathcal{L}_q(T)$.

Lemma C.1. 1. Fix any T > 0. Let $q \in \mathcal{L}_q(T)$ be such that there exists some $K < \infty$ for which its marginal laws $\{q_t\}_{t \in [0,T]}$ satisfy

$$\sup_{t \in [0,T]} \sup_{j \in [p]} \mathbb{E}_{\varphi_t \sim q_t} e^{(8M/\tau^2)|\varphi_{t,j}|} < K.$$
(C.2)

Then for any initialization $g_0 \in \mathcal{P}_*(M)$ that is strictly positive on [-M, M], there exists a unique solution $\{g_t\}_{t \in [0,T]}$ to

$$\frac{\mathrm{d}}{\mathrm{d}t}g_t(\theta) = g_t(\theta) \left(\mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * g_t}(\theta) - 1\right)$$
(C.3)

such that $g_t \in \mathcal{P}_*(M)$ for all $t \in [0,T]$. Furthermore, $g_t(\theta) \ge g_0(\theta)e^{-t}$ for all $t \in [0,T]$ and $\theta \in [-M,M]$.

2. For any K > 0, there exists $C_0 > 0$ depending only on (M, τ, K) such that the following holds: Fix any $T \leq 1/C_0$. For any $q, \tilde{q} \in \mathcal{L}_q(T)$ both satisfying the property (C.2), and for any $g_0 \in \mathcal{P}_*(M)$ satisfying $g_0(\theta) > 0$ for all $\theta \in [-M, M]$, if g, \tilde{g} are the solutions of (C.3) associated to q, \tilde{q} given by Part (1), then

$$d_{\infty,\mathrm{TV}}(g,\widetilde{g}) \le 2C_0T \cdot d_W(q,\widetilde{q}).$$

Proof of Lemma C.1-Part (1). Fix $g_0 \in \mathcal{P}_*(M)$ with $\inf_{\theta \in [-M,M]} g_0(\theta) > 0$ for all $\theta \in [-M,M]$. Define

$$B(g_0) = \{g \in L^1([-M, M]) : g(\theta) \in [g_0(\theta)/2, 2g_0(\theta)] \text{ for Lebesgue-a.e. } \theta \in [-M, M]\}$$

equipped with the L^1 metric $||g - \tilde{g}||_{L^1} = \int_{-M}^{M} |g(\theta) - \tilde{g}(\theta)|$. Note that $L^1([-M, M])$ is complete, and if $g_n \in B(g_0)$ converges in L^1 to some $g_* \in L^1([-M, M])$, then also $g_n(\theta) - g_*(\theta) \to 0$ for Lebesgue-a.e. $\theta \in [-M, M]$. Then $g_*(\theta) \in [g_0(\theta)/2, 2g_0(\theta)]$ for Lebesgue-a.e. $\theta \in [-M, M]$, implying that $B(g_0)$ is closed in $L^1([-M, M])$ and hence complete.

For some $t_0 > 0$ to be specified later, define $E(g_0) = C([0, t_0], B(g_0))$ equipped with the metric $d_{\infty,L^1}(g, \tilde{g}) = \sup_{t \in [0, t_0]} \|g_t - \tilde{g}_t\|_{L^1}$. Then $E(g_0)$ is also complete, by the above completeness of $B(g_0)$. Define a map $\Gamma : E(g_0) \to E(g_0)$ with $\{g_t\}_{t \in [0, t_0]} \mapsto \{\Gamma g\}_{t \in [0, t_0]}$, by

$$(\Gamma g)_t(\theta) = g_0(\theta) \cdot \exp\left(\int_0^t \left[\mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s}(\theta) - 1\right] \mathrm{d}s\right).$$
(C.4)

We claim that for small enough t_0 , this definition of Γ indeed maps $E(g_0)$ to itself, and furthermore is contractive under d_{∞,L^1} . To see that Γ maps $E(g_0)$ to itself, note that for any $g \in E(g_0)$ and $t \in [0, t_0]$,

$$\log (\Gamma g)_t(\theta) = \log g_0(\theta) + \int_0^t \left[\mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s}(\theta) - 1 \right] \mathrm{d}s \ge \log g_0(\theta) - t.$$
(C.5)

Observe that condition (C.2) implies also $\int e^{(8M/\tau^2)|\varphi|} \bar{q}_s(\varphi) < K$. Applying this, $\frac{N_{\tau}(\varphi-\theta)}{N_{\tau}(\varphi-\theta')} \leq Ce^{(2M/\tau^2)|\varphi|}$ for all $\theta, \theta' \in [-M, M]$ by Proposition B.1, and $\int_{-M}^{M} g_s \geq \int_{-M}^{M} g_0/2 = 1/2$,

$$\mathcal{N}_{\tau} * \frac{\bar{q}_s}{\mathcal{N}_{\tau} * g_s}(\theta) = \int \frac{\mathcal{N}_{\tau}(\varphi - \theta)}{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta') g_s(\theta') \mathrm{d}\theta'} \bar{q}_s(\varphi) \mathrm{d}\varphi$$

$$\leq \int C e^{(2M/\tau^2)|\varphi|} \bar{q}_s(\varphi) \mathrm{d}\varphi \cdot \frac{1}{\int_{-M}^{M} g_s(\theta') \mathrm{d}\theta'} < C'$$
(C.6)

for some C' > 0. Here and below, C, C' etc. are constants depending only on (M, τ, K) . So

$$\log (\Gamma g)_t(\theta) = \log g_0(\theta) + \int_0^t \left[\mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s}(\theta) - 1 \right] \mathrm{d}s \le \log g_0(\theta) + C't.$$
(C.7)

Choosing $t_0 < (\log 2) / \max(C', 1)$, this shows that for all $t \leq t_0$ and $\theta \in [-M, M]$, we have $(\Gamma g)_t(\theta) \in (g_0(\theta)/2, 2g_0(\theta))$, so $(\Gamma g)_t \in B(g_0)$. Furthermore, it is clear from the definition of Γ that if $s \to t$, then $(\Gamma g)_s(\theta) \to (\Gamma g)_t(\theta)$ pointwise over $\theta \in [-M, M]$. Then also $\int_{-M}^M |(\Gamma g)_s - (\Gamma g)_t| \to 0$ by (C.5), (C.7), and the dominated convergence theorem, so $t \mapsto (\Gamma g)_t$ is continuous in L^1 . Thus Γ is a well-defined map from $E(g_0)$ to itself.

To see that Γ is contractive, observe that for any $g, \tilde{g} \in E(g_0)$ and $t \in [0, t_0]$,

$$\begin{split} &\int_{-M}^{M} \left| (\Gamma g)_{t}(\theta) - (\Gamma \widetilde{g})_{t}(\theta) \right| \mathrm{d}\theta \\ &= \int_{-M}^{M} g_{0}(\theta) \Big| \exp \Big(\int_{0}^{t} \left[\mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * g_{s}}(\theta) - 1 \right] \mathrm{d}s \Big) - \exp \Big(\int_{0}^{t} \left[\mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * \widetilde{g}_{s}}(\theta) - 1 \right] \mathrm{d}s \Big) \Big| \mathrm{d}\theta \\ &\leq \sup_{\theta \in [-M,M]} \Big| \exp \Big(\int_{0}^{t} \mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * g_{s}}(\theta) \mathrm{d}s \Big) - \exp \Big(\int_{0}^{t} \mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * \widetilde{g}_{s}}(\theta) \mathrm{d}s \Big) \Big| \\ &\leq t \cdot \sup_{\theta \in [-M,M]} e^{\xi(\theta)} \cdot \sup_{s \in [0,t]} \Big| \mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * g_{s}}(\theta) - \mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * \widetilde{g}_{s}}(\theta) \Big|, \end{split}$$

where $\xi(\theta)$ is some value between $\int_0^t \mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s}(\theta) ds$ and $\int_0^t \mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * \tilde{g}_s}(\theta) ds$. Using (C.6), we have $\sup_{\theta \in [-M,M]} e^{\xi(\theta)} < 2$ because $t \le t_0 < (\log 2)/C'$. Moreover, using again Proposition B.1 and the lower bounds $\int_{-M}^M g_s(\theta) \ge 1/2$ and $\int_{-M}^M \tilde{g}_s(\theta) \ge 1/2$ for all $s \in [0, t_0]$, we have

$$\begin{split} \sup_{\theta \in [-M,M]} \sup_{s \in [0,t]} \left| \mathcal{N}_{\tau} * \frac{\bar{q}_{s}}{\mathcal{N}_{\tau} * g_{s}}(\theta) - \mathcal{N}_{\tau} * \frac{\bar{q}_{s}}{\mathcal{N}_{\tau} * \tilde{g}_{s}}(\theta) \right| \\ &\leq \sup_{\theta \in [-M,M]} \sup_{s \in [0,t]} \int_{-M}^{M} \int \frac{\mathcal{N}_{\tau}(\varphi - \theta) \mathcal{N}_{\tau}(\varphi - \theta') \bar{q}_{s}(\varphi)}{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta') g_{s}(\theta') \mathrm{d}\theta' \int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta') \tilde{g}_{s}(\theta') \mathrm{d}\theta'} \cdot |g_{s}(\theta') - \tilde{g}_{s}(\theta')| \mathrm{d}\varphi \mathrm{d}\theta' \\ &\leq \sup_{s \in [0,t]} \int C e^{(4M/\tau^{2})|\varphi|} \bar{q}_{s}(\varphi) \mathrm{d}\varphi \cdot \sup_{s \in [0,t]} \frac{1}{\int_{-M}^{M} g_{s}(\theta') \mathrm{d}\theta' \cdot \int_{-M}^{M} \tilde{g}_{s}(\theta') \mathrm{d}\theta'} \cdot \sup_{s \in [0,t]} \int_{-M}^{M} |g_{s}(\theta') - \tilde{g}_{s}(\theta')| \mathrm{d}\theta' \mathrm{d}\theta' \\ &\leq C'' \|g - \tilde{g}\|_{\infty,L^{1}}. \end{split}$$

For some C'' > 0. In summary,

$$\|\Gamma g - \Gamma \widetilde{g}\|_{\infty,L^1} = \sup_{t \in [0,t_0]} \int_{-M}^{M} \left| (\Gamma g)_t(\theta) - (\Gamma \widetilde{g})_t(\theta) \right| \mathrm{d}\theta \le 2C'' t_0 \|g - \widetilde{g}\|_{\infty,L^1}.$$

Choosing $t_0 \leq 1/4C''$, this establishes that $\Gamma: E(g_0) \to E(g_0)$ is a contraction.

By the Banach fixed point theorem, there exists a unique fixed point $g = \{g_t\}_{t \in [0,t_0]} \in E(g_0)$ of Γ . Choosing the version of g satisfying (C.4) with pointwise equality for all $t \in [0, t_0]$ and $\theta \in [-M, M]$, we must then have

$$\log g_t(\theta) = \log g_0(\theta) + \int_0^t \left[\mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s}(\theta) - 1 \right] \mathrm{d}s \tag{C.8}$$

for all $t \in [0, t_0]$ and $\theta \in [-M, M]$. The functions $\varphi \mapsto p^{-1} \sum_{j=1}^p \frac{\mathcal{N}_{\tau}(\varphi_j - \theta)}{\mathcal{N}_{\tau} * g_s(\varphi_j)} q_s(\varphi)$ are uniformly integrable over $s \in [0, t_0]$, by Proposition B.1 and the condition (C.2) for q_s . Thus $s \mapsto \mathcal{N}_{\tau} * \frac{\bar{q}_s}{\mathcal{N}_{\tau} * g_s}(\theta) = p^{-1} \sum_{j=1}^p \int \frac{\mathcal{N}_{\tau}(\varphi_j - \theta)}{\mathcal{N}_{\tau} * g_s(\varphi_j)} q_s(\varphi) d\varphi$ is continuous in s, so $g_t(\theta)$ is C^1 in t. Differentiating (C.8) in t shows that $\{g_t\}_{t \in [0, t_0]}$ solves (C.3). Furthermore, $g_t \in \mathcal{P}_*(M)$ for all $t \in [0, t_0]$, because

$$\partial_t \int_{-M}^{M} g_t = \int_{-M}^{M} g_t \cdot \partial_t \log g_t = \int_{-M}^{M} g_t \cdot \left[\mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * g_t} - 1 \right] = 1 - \int_{-M}^{M} g_t,$$

which has the unique solution $\int_{-M}^{M} g_t = 1$ with initialization $\int_{-M}^{M} g_0 = 1$.

Conversely, let $\{\tilde{g}_t\}_{t\in[0,t_0]}$ be any other solution of (C.3) where $\tilde{g}_t \in \mathcal{P}_*(M)$ for all $t \in [0,t_0]$. Suppose by contradiction that there exists some $\theta \in [-M, M]$ for which $\tilde{g}_t(\theta) \notin [g_0(\theta)/2, 2g_0(\theta)]$ for some $t \in [0,t_0]$. Fixing this θ , let t' be the infimum of all such t. Then $\tilde{g}_{t'}(\theta) = g_0(\theta)/2$, $2g_0(\theta)/2$ or $\tilde{g}_{t'} = 2g_0(\theta)$ by continuity of $t \mapsto \tilde{g}_t(\theta)$. On the other hand, we must have $\log \tilde{g}_t(\theta) \ge \log g_0(\theta) - t$ and $\log \tilde{g}_t(\theta) \le \log g_0(\theta) + C't$ for all $t \in [0,t']$, by the same arguments as (C.5) and (C.7) where we may use $\int_{-M}^{M} \tilde{g}_s = 1$ in place of the previously used condition $\int_{-M}^{M} g_s \ge 1/2$ to derive (C.7). Since $t' \le t_0 < (\log 2) / \max(C', 1)$, integrating these inequalities leads to the contradiction that $\tilde{g}_{t'}(\theta) > g_0(\theta)/2$ and $\tilde{g}_{t'}(\theta) < 2g_0(\theta)$ strictly. Thus, $\tilde{g}_t(\theta) \in [g_0(\theta)/2, 2g_0(\theta)]$ for all $t \in [0, t_0]$ and $\theta \in [-M, M]$. So $\{\tilde{g}_t\}_{t\in[0,t_0]}$ is also a fixed point of Γ in $E(g_0)$, implying by uniqueness of this fixed point that $g_t = \tilde{g}_t$ as distributions in $\mathcal{P}(M)$ for all $t \in [0, t_0]$. This implies that the right sides of (C.8) are equal for g_t and \tilde{g}_t , and consequently the left sides are equal pointwise, i.e. $g_t(\theta) = \tilde{g}_t(\theta)$ pointwise for all $\theta \in [-M, M]$ and $t \in [0, t_0]$. Thus the solution $\{g_t\}_{t\in[0,t_0]}$ to (C.3) is unique.

This solution satisfies that g_{t_0} is a strictly positive density in $\mathcal{P}_*(M)$. Then we may apply the above argument sequentially to $[t_0, 2t_0]$, $[2t_0, 3t_0]$ etc. with g_{t_0} , g_{2t_0} etc. in place of g_0 . This proves the existence and uniqueness of the solution $\{g_t\}_{t\in[0,T]}$ to (C.3). Finally, from (C.8) we get $g_t(\theta)/g_0(\theta) \ge e^{-t}$ for all $\theta \in [-M, M]$ and $t \in [0, T]$, completing the proof of Part (1).

Proof of Lemma C.1-Part (2). By definition,

$$g_t(\theta) = g_0(\theta) + \int_0^t g_s(\theta) \Big[\mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s}(\theta) - 1 \Big] \mathrm{d}s,$$

$$\tilde{g}_t(\theta) = g_0(\theta) + \int_0^t \tilde{g}_s(\theta) \Big[\mathcal{N}_\tau * \frac{\bar{\tilde{q}}_s}{\mathcal{N}_\tau * g_s}(\theta) - 1 \Big] \mathrm{d}s.$$

Then for any $t \in [0, T]$,

$$\int_{-M}^{M} |g_{t}(\theta) - \widetilde{g}_{t}(\theta)| \mathrm{d}\theta \leq \int_{-M}^{M} \left| \int_{0}^{t} \left(g_{s}(\theta) \left[\mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * g_{s}}(\theta) \right] - \widetilde{g}_{s}(\theta) \left[\mathcal{N}_{\tau} * \frac{\overline{\tilde{q}}_{s}}{\mathcal{N}_{\tau} * \widetilde{g}_{s}}(\theta) \right] \right) \mathrm{d}s \right| \mathrm{d}\theta \\
+ \int_{0}^{t} \int_{-M}^{M} |g_{s}(\theta) - \widetilde{g}_{s}(\theta)| \mathrm{d}\theta \mathrm{d}s \\
\leq (I) + (II) + (III) + 2T \cdot d_{\mathrm{TV}}(g, \widetilde{g}),$$
(C.9)

where

$$(I) = \int_{-M}^{M} \int_{0}^{t} \left| g_{s}(\theta) - \widetilde{g}_{s}(\theta) \right| \cdot \mathcal{N}_{\tau} * \frac{\overline{q}_{s}}{\mathcal{N}_{\tau} * g_{s}}(\theta) \mathrm{d}s \mathrm{d}\theta,$$

$$(II) = \int_{-M}^{M} \int_{0}^{t} \widetilde{g}_{s}(\theta) \cdot \left| \mathcal{N}_{\tau} * \frac{\overline{q}_{s} - \overline{\widetilde{q}}_{s}}{\mathcal{N}_{\tau} * g_{s}}(\theta) \right| \mathrm{d}s \mathrm{d}\theta,$$

$$(III) = \int_{-M}^{M} \int_{0}^{t} \widetilde{g}_{s}(\theta) \cdot \left| \mathcal{N}_{\tau} * \left(\frac{\overline{\widetilde{q}}_{s}}{\mathcal{N}_{\tau} * g_{s}} - \frac{\overline{\widetilde{q}}_{s}}{\mathcal{N}_{\tau} * \widetilde{g}_{s}} \right)(\theta) \right| \mathrm{d}s \mathrm{d}\theta$$

Let C, C' etc. denote constants depending only on (M, τ, K) and changing from instance to instance. Applying Proposition B.1, we have $\mathcal{N}_{\tau} * \frac{\bar{q}_s}{\mathcal{N}_{\tau} * g_s}(\theta) \leq C \int e^{(2M/\tau^2)|\varphi|} \bar{q}_s(\varphi) \mathrm{d}\varphi$ for any $\theta \in [-M, M]$, hence

$$(I) \leq Ct \sup_{s \in [0,t]} \left(\int_{-M}^{M} \left| g_s(\theta) - \widetilde{g}_s(\theta) \right| \mathrm{d}\theta \cdot \int e^{(2M/\tau^2)|\varphi|} \bar{q}_s(\varphi) \mathrm{d}\varphi \right) \leq C'T \cdot d_{\infty,\mathrm{TV}}(g,\widetilde{g}).$$

Similarly,

$$(III) \leq \int_{-M}^{M} \int_{0}^{t} \widetilde{g}_{s}(\theta) \cdot \mathcal{N}_{\tau}(\theta - \varphi) \overline{\widetilde{q}}_{s}(\varphi) \cdot \frac{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta') |g_{s}(\theta') - \widetilde{g}_{s}(\theta')| \mathrm{d}\theta'}{\mathcal{N}_{\tau} * g_{s}(\varphi) \cdot \mathcal{N}_{\tau} * \widetilde{g}_{s}(\varphi)} \mathrm{d}s\mathrm{d}\theta$$
$$\leq Ct \sup_{s \in [0,t]} \left(\int_{-M}^{M} \left| g_{s}(\theta') - \widetilde{g}_{s}(\theta') \right| \mathrm{d}\theta' \cdot \int e^{(4M/\tau^{2})|\varphi|} \overline{\widetilde{q}}_{s}(\varphi) \mathrm{d}\varphi \right) \leq C'T \cdot d_{\infty,\mathrm{TV}}(g, \widetilde{g}).$$

To bound (II), let $\varphi = \{\varphi_t\}_{t \in [0,T]}$ and $\tilde{\varphi} = \{\tilde{\varphi}_t\}_{t \in [0,T]}$ be two processes on the same probability space such that $\varphi \sim q$ and $\tilde{\varphi} \sim \tilde{q}$. Then

$$(II) = \int_{-M}^{M} \int_{0}^{t} \widetilde{g}_{s}(\theta) \cdot \left| \int \frac{\mathcal{N}_{\tau}(\theta - \varphi)}{\mathcal{N}_{\tau} * g_{s}(\varphi)} (\bar{q}_{s}(\varphi) - \bar{\widetilde{q}}_{s}(\varphi)) d\varphi \right| ds d\theta$$
$$= \int_{-M}^{M} \int_{0}^{t} \widetilde{g}_{s}(\theta) \cdot \left| \frac{1}{p} \sum_{j=1}^{p} \mathbb{E} f_{s}(\varphi_{s,j}; \theta) - \mathbb{E} f_{s}(\widetilde{\varphi}_{s,j}; \theta) \right| ds d\theta,$$

where $f_s(\varphi; \theta) = \frac{N_{\tau}(\varphi - \theta)}{N_{\tau} * g_s(\varphi)}$. For any $\theta \in [-M, M]$, applying Proposition B.1,

$$\left|\mathsf{f}_{s}'(\varphi;\theta)\right| = \frac{1}{\tau^{2}} \left| \frac{\mathcal{N}_{\tau}(\varphi-\theta)\cdot\theta}{\mathcal{N}_{\tau}*g_{s}(\varphi)} - \frac{\mathcal{N}_{\tau}(\varphi-\theta)\cdot\int\mathcal{N}_{\tau}(\varphi-\theta')\cdot\theta'g_{s}(\theta')\mathrm{d}\theta'}{\left(\mathcal{N}_{\tau}*g_{s}(\varphi)\right)^{2}} \right| \le Ce^{(4M/\tau^{2})|\varphi|},$$

so it holds that

$$\begin{aligned} \left| \mathbb{E} \mathsf{f}_{s}(\varphi_{s,j};\theta) - \mathbb{E} \mathsf{f}_{s}(\widetilde{\varphi}_{s,j};\theta) \right| &\leq C \,\mathbb{E} \left(e^{(4M/\tau^{2})|\varphi_{s,j}|} + e^{(4M/\tau^{2})|\widetilde{\varphi}_{s,j}|} \right) \cdot |\varphi_{s,j} - \widetilde{\varphi}_{s,j}| \\ &\leq C \big(\mathbb{E} e^{(8M/\tau^{2})|\varphi_{s,j}|} + e^{(8M/\tau^{2})|\widetilde{\varphi}_{s,j}|} \big)^{1/2} \cdot \big(\mathbb{E} (\varphi_{s,j} - \widetilde{\varphi}_{s,j})^{2} \big)^{1/2} \\ &\leq C' \big(\mathbb{E} (\varphi_{s,j} - \widetilde{\varphi}_{s,j})^{2} \big)^{1/2}, \end{aligned}$$

using the condition (C.2) for $\{q_t\}$ and $\{\widetilde{q}_t\}$. Therefore,

$$(II) \le C \int_0^t \frac{1}{p} \sum_{j=1}^p \left[\mathbb{E}(\varphi_{s,j} - \widetilde{\varphi}_{s,j})^2 \right]^{1/2} \mathrm{d}s \le C'T \cdot \left(\mathbb{E} \sup_{s \in [0,T]} \|\varphi_s - \widetilde{\varphi}_s\|_2^2 \right)^{1/2}.$$

Taking the infimum over couplings of $q, \tilde{q} \in \mathcal{L}_q$ yields $(II) \leq C'T \cdot d_W(q, \tilde{q})$. Plugging the bounds for (I), (II), (III) into (C.9) yields, for some $C_0 > 0$ (not depending on T),

$$d_{\infty,\mathrm{TV}}(g,\widetilde{g}) \le C_0 T \cdot (d_{\infty,\mathrm{TV}}(g,\widetilde{g}) + d_W(q,\widetilde{q}))$$

61

Then choosing $T \leq 1/(2C_0)$ and rearranging concludes the proof.

An immediate corollary is the existence and uniqueness of the univariate g-flow $\{g_t\}_{t\geq 0}$ in the setting of Theorem 3.3.

Corollary C.2. Fix any T > 0. Let $\bar{q} \in \mathcal{P}(\mathbb{R})$ satisfy $\int e^{\lambda \varphi} \bar{q}(\varphi) d\varphi < \infty$ for all $\lambda \in \mathbb{R}$. Then, given any initial density $g_0 \in \mathcal{P}_*(M)$ that is strictly positive on [-M, M], there exists a unique solution $\{g_t\}_{t\in[0,T]}$ to (3.2) where $g_t \in \mathcal{P}_*(M)$ for all $t \in [0,T]$. Furthermore, $g_t(\theta) \geq g_0(\theta)e^{-t}$ for all $\theta \in [-M, M]$ and $t \in [0,T]$.

Proof. Let $q \in \mathcal{L}_q$ be the law of a constant process $\varphi_t = \varphi_0$ for all $t \in [0, T]$, where coordinates of φ_0 are i.i.d. with distribution \bar{q} . Then the condition (C.2) for q is satisfied by the given moment generating function condition for \bar{q} , and $\bar{q}_t = \bar{q}$ for all $t \in [0, T]$, so the result follows from Lemma C.1-(1).

C.2 Solution properties of the q-flow

Recall the negative log-posterior-density $U_g(\cdot)$ of φ from (3.5). The following result gives the solution properties of the Wasserstein q-flow for any fixed $g = \{g_t\}_{t \in [0,T]} \in \mathcal{L}_g(T)$.

Lemma C.3. 1. Fix any T > 0, and let $g = \{g_t\}_{t \in [0,T]} \in \mathcal{L}_g(T)$. Then for any initialization $q_0 \in \mathcal{P}(\mathbb{R}^p)$, there is a strong solution $\varphi = \{\varphi_t\}_{t \in [0,T]}$ to the Langevin diffusion

$$\mathrm{d}\boldsymbol{\varphi}_t = -\nabla U_{q_t}(\boldsymbol{\varphi}_t)\mathrm{d}t + \sqrt{2}\,\mathrm{d}\mathbf{B}_t,\tag{C.10}$$

which is pathwise unique given the initialization $\varphi_0 \sim q_0$.

The marginal laws $\{q_t\}_{t\in[0,T]}$ of $\{\varphi_t\}_{t\in[0,T]}$ form the unique weak solution valued in $\mathcal{P}(\mathbb{R}^p)$ to the Fokker-Planck PDE

$$\partial_t q_t(\boldsymbol{\varphi}) = \nabla \cdot [q_t(\boldsymbol{\varphi}) \nabla U_{g_t}(\boldsymbol{\varphi})] + \Delta q_t(\boldsymbol{\varphi}), \qquad (C.11)$$

in the sense that $q_t \Rightarrow q_0$ weakly as $t \to 0$ and

$$0 = \int_0^t \int \left(\partial_s \zeta_s(\varphi) - \nabla U_{g_s}(\varphi)^\top \nabla \zeta_s(\varphi) + \Delta \zeta_s(\varphi) \right) q_s(\mathrm{d}\varphi) \mathrm{d}s \tag{C.12}$$

for any smooth function $(t, \varphi) \mapsto \zeta_t(\varphi)$ with compact support in $(0, T) \times \mathbb{R}^p$.

2. Let $q_0 \in \mathcal{P}(\mathbb{R}^p)$ satisfy $\mathbb{E}_{\varphi_0 \sim q_0} e^{\lambda^\top \varphi_0} < \infty$ for all $\lambda \in \mathbb{R}^p$. Then, for each $\lambda \in \mathbb{R}^p$, there exists some $C(q_0, \lambda) < \infty$ (depending also on $n, p, \mathbf{X}, \mathbf{y}, \sigma, \tau, M$) such that for any T > 0 and $g \in \mathcal{L}_g(T)$, the law $q \in \mathcal{L}_q(T)$ of the solution $\{\varphi_t\}_{t \in [0,T]}$ of Part (1) satisfies

$$\sup_{t \in [0,T]} \mathbb{E}_{\boldsymbol{\varphi}_t \sim q_t} e^{\boldsymbol{\lambda}^\top \boldsymbol{\varphi}_t} < C(q_0, \boldsymbol{\lambda}).$$
(C.13)

In particular, there exists some $K(q_0) < \infty$ such that for any T > 0 and $g \in \mathcal{L}_q(T)$,

$$\sup_{t \in [0,T]} \max_{j \in [p]} \mathbb{E}_{\varphi_t \sim q_t} e^{(8M/\tau^2)|\varphi_{t,j}|} \le K(q_0).$$
(C.14)

3. For any K > 0, there exist $T_0, C_0 > 0$ (depending on K and also $(n, p, \mathbf{X}, \mathbf{y}, \sigma, \tau, M)$) such that the following holds: Let $q_0 \in \mathcal{P}(\mathbb{R}^p)$ be any initialization as in Part (2), such that (C.14) holds with $K(q_0) = K$. Then for any $T \leq T_0$ and $g, \tilde{g} \in \mathcal{L}_g(T)$, if $q, \tilde{q} \in \mathcal{L}_q(T)$ are the associated solutions of (C.10) given by Part (1), then

$$d_W(q, \widetilde{q}) \le C_0 T \cdot d_{\infty, \mathrm{TV}}(g, \widetilde{g}).$$

Proof of Lemma C.3-Part (1). We show that $\nabla U_g : \mathbb{R}^p \to \mathbb{R}^p$ is Lipschitz uniformly over $g \in \mathcal{P}(M)$. Observe that by Tweedie's formula,

$$\nabla U_g(\boldsymbol{\varphi}) = -\mathbf{X}^{\top} \mathbf{\Sigma}^{-1}(\mathbf{y} - \mathbf{X}\boldsymbol{\varphi}) - \left(\frac{[\mathcal{N}_\tau * g]'(\varphi_j)}{[\mathcal{N}_\tau * g](\varphi_j)}\right)_{j=1}^p$$
(C.15)

$$= \left(\mathbf{X}^{\top} \mathbf{\Sigma}^{-1} \mathbf{X} + \tau^{-2} \operatorname{Id} \right) \boldsymbol{\varphi} - \mathbf{X}^{\top} \mathbf{\Sigma}^{-1} \mathbf{y} - \tau^{-2} \theta_g(\boldsymbol{\varphi})$$
(C.16)

where $\theta_q(\varphi) = \mathbb{E}_q[\theta \mid \varphi]$ is the posterior mean in the scalar model

$$\varphi = \theta + z, \qquad \theta \sim g, \qquad z \sim \mathcal{N}(0, \tau^2),$$

and $\theta_g(\varphi)$ is its entrywise application to $\varphi = (\varphi_1, \ldots, \varphi_p)$. We have $|\theta'_g(\varphi)| = |\tau^{-2} \operatorname{Var}[\theta | \varphi]| \le M^2/\tau^2$. Then $\|\nabla U_g(\varphi) - \nabla U_g(\widetilde{\varphi})\|_2 \le C \|\varphi - \widetilde{\varphi}\|_2$ for some C > 0 (depending on $\mathbf{X}, \mathbf{y}, \sigma, \tau, M$), uniformly over $g \in \mathcal{P}(M)$, as claimed.

From the weak continuity of $t \mapsto g_t$ and the form (C.15) for U_g , we see that $(t, \varphi) \mapsto U_{g_t}(\varphi)$ is jointly continuous. Then the existence of a pathwise unique strong solution $\varphi = \{\varphi_t\}_{t \in [0,T]}$ to the Langevin diffusion (C.10) follows from [LG16, Theorem 8.3]. By Ito's formula, for any smooth bounded function $\zeta_t(\varphi)$, we have

$$\mathbb{E}[\zeta_t(\boldsymbol{\varphi}_t) - \zeta_0(\boldsymbol{\varphi}_0)] = \mathbb{E}\left[\int_0^t \left(\partial_s \zeta_s(\boldsymbol{\varphi}_s) - \nabla U_{g_s}(\boldsymbol{\varphi}_s)^\top \nabla \zeta_t(\boldsymbol{\varphi}_s) + \Delta \zeta_s(\boldsymbol{\varphi}_s)\right) \mathrm{d}s\right]$$
(C.17)

and the statement (C.12) follows from applying the boundary condition $\zeta_t(\varphi_t) = \zeta_0(\varphi_0) = 0$. This is the unique weak solution to (C.11) for which every $q_t \in \mathcal{P}(\mathbb{R}^p)$, by [BRS11, Theorem 4.9].

Proof of Lemma C.3-Part (2). Let $\{\varphi_t\}_{t\in[0,T]}$ be a solution of the Langevin diffusion in Part (1), with initialization $\varphi_0 \sim q_0$. Let $\Gamma = \mathbf{X}^\top \mathbf{\Sigma}^{-1} \mathbf{X} + \tau^{-2} \operatorname{Id} \in \mathbb{R}^{p \times p}$, and let $\mathbf{u} \in \mathbb{R}^p$ be an eigenvector of Γ (having arbitrary ℓ_2 norm) with eigenvalue $\mu > 0$. Fix some B > 0 to be specified later. Then

$$\mathbb{E}e^{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}} = \mathbb{E}e^{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}}\mathbf{1}_{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}\leq B} + \mathbb{E}e^{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}}\mathbf{1}_{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}>B} \leq e^{B} + \mathbb{E}e^{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}}\mathbf{1}_{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}>B}$$

To bound the second term, let h be a twice differentiable function such that: (i) h(u) = 0 for $u \leq B-1$; (ii) h(u) = 1 for $u \geq B$; (iii) $h(u), h'(u) \geq 0$ for $u \in \mathbb{R}$; (iv) $\sup_{u \in \mathbb{R}} \max(h'(u), |h''(u)|) < C$ for some C > 0. Then with $f(\varphi) = e^{\mathbf{u}^{\top}\varphi}h(\mathbf{u}^{\top}\varphi)$,

$$\mathbb{E}e^{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}}\mathbf{1}_{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}>B} \leq \mathbb{E}e^{\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}}h(\mathbf{u}^{\top}\boldsymbol{\varphi}_{t}) = \mathbb{E}f(\boldsymbol{\varphi}_{t}).$$

Using Ito's formula, we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}f(\boldsymbol{\varphi}_t) = \mathbb{E}\left\langle \nabla f(\boldsymbol{\varphi}_t), \mathrm{d}\boldsymbol{\varphi}_t \right\rangle + \mathbb{E}\operatorname{Tr}(\nabla^2 f(\boldsymbol{\varphi}_t)) \\ = \mathbb{E}\left\langle \nabla f(\boldsymbol{\varphi}_t), -\nabla U_{g_t}(\boldsymbol{\varphi}_t) \right\rangle + \mathbb{E}\operatorname{Tr}(\nabla^2 f(\boldsymbol{\varphi}_t)).$$
(C.18)

For the first term, note from (C.16) that $-\nabla U_{g_t}(\varphi) = -\Gamma \varphi + \mathbf{s}$ where $\mathbf{s} = \mathbf{X}^\top \Sigma^{-1} \mathbf{y} + \tau^{-2} \theta_{g_t}(\varphi)$ Since g_t is supported on [-M, M], there exists some C > 0 (depending on $\mathbf{X}, \mathbf{y}, \sigma, \tau, M$) such that $\|\mathbf{s}\| < C$. Hence using

$$\nabla f(\boldsymbol{\varphi}) = \mathbf{u} e^{\mathbf{u}^{\top} \boldsymbol{\varphi}} h(\mathbf{u}^{\top} \boldsymbol{\varphi}) + \mathbf{u} e^{\mathbf{u}^{\top} \boldsymbol{\varphi}} h'(\mathbf{u}^{\top} \boldsymbol{\varphi}) = \mathbf{u} \cdot \left(f(\boldsymbol{\varphi}) + e^{\mathbf{u}^{\top} \boldsymbol{\varphi}} h'(\mathbf{u}^{\top} \boldsymbol{\varphi}) \right),$$

and applying $\Gamma \mathbf{u} = \mu \cdot \mathbf{u}, f(\cdot) \ge 0$, and $h'(\cdot) \ge 0$, we have

$$\langle \nabla f(\boldsymbol{\varphi}), -\nabla U_{g_t}(\boldsymbol{\varphi}) \rangle = -\left(f(\boldsymbol{\varphi}) + e^{\mathbf{u}^\top \boldsymbol{\varphi}} h'(\mathbf{u}^\top \boldsymbol{\varphi})\right) \cdot \mathbf{u}^\top \boldsymbol{\Gamma} \boldsymbol{\varphi} + \left(f(\boldsymbol{\varphi}) + e^{\mathbf{u}^\top \boldsymbol{\varphi}} h'(\mathbf{u}^\top \boldsymbol{\varphi})\right) \mathbf{u}^\top \mathbf{s}$$

$$\leq -(\mu \cdot \mathbf{u}^{\top} \varphi - \|\mathbf{u}\| \|\mathbf{s}\|) f(\varphi) - \mu e^{\mathbf{u}^{\top} \varphi} h'(\mathbf{u}^{\top} \varphi) \cdot \mathbf{u}^{\top} \varphi + \|\mathbf{u}\| \|\mathbf{s}\| \cdot e^{\mathbf{u}^{\top} \varphi} h'(\mathbf{u}^{\top} \varphi) \cdot \mathbf{u}^{\top} \varphi$$

Choose $B = B(||\mathbf{u}||, \mu) > 1$ large enough such that $\mu \cdot \mathbf{u}^{\top} \varphi - ||\mathbf{u}|| ||\mathbf{s}|| > B\mu/2$ whenever $\varphi \in \mathbb{R}^p$ satisfies $\mathbf{u}^{\top} \varphi > B - 1$. Then, the first term is bounded above by $-B\mu/2 \cdot f(\varphi)$ (which holds also when $\mathbf{u}^{\top} \varphi \leq B - 1$, in which case $f(\varphi) = 0$). The second term $-\mu e^{\mathbf{u}^{\top} \varphi} h'(\mathbf{u}^{\top} \varphi) \cdot \mathbf{u}^{\top} \varphi$ is non-positive, because $h'(\mathbf{u}^{\top} \varphi) \geq 0$ and equals 0 unless $\mathbf{u}^{\top} \varphi > B - 1 > 0$. For the third term, since also $h'(\mathbf{u}^{\top} \varphi) = 0$ unless $\mathbf{u}^{\top} \varphi \leq B$, it is bounded by some $C = C(||\mathbf{u}||, \mu) > 0$. In summary,

$$\mathbb{E} \left\langle \nabla f(\boldsymbol{\varphi}_t), -\nabla U_{g_t}(\boldsymbol{\varphi}_t) \right\rangle \leq -\frac{B\mu}{2} \mathbb{E} f(\boldsymbol{\varphi}_t) + C.$$

On the other hand, since

$$\nabla^2 f(\boldsymbol{\varphi}) = \mathbf{u} \mathbf{u}^\top \Big(e^{\mathbf{u}^\top \boldsymbol{\varphi}} h(\mathbf{u}^\top \boldsymbol{\varphi}) + 2e^{\mathbf{u}^\top \boldsymbol{\varphi}} h'(\mathbf{u}^\top \boldsymbol{\varphi}) + e^{\mathbf{u}^\top \boldsymbol{\varphi}} h''(\mathbf{u}^\top \boldsymbol{\varphi}) \Big),$$

we have

$$\operatorname{tr}\left(\nabla^{2} f(\boldsymbol{\varphi})\right) = \|\mathbf{u}\|^{2} \cdot f(\boldsymbol{\varphi}) + \|\mathbf{u}\|^{2} \left(2e^{\mathbf{u}^{\top}\boldsymbol{\varphi}} h'(\mathbf{u}^{\top}\boldsymbol{\varphi}) + e^{\mathbf{u}^{\top}\boldsymbol{\varphi}} h''(\mathbf{u}^{\top}\boldsymbol{\varphi})\right)$$

Using $h'(\mathbf{u}^{\top}\boldsymbol{\varphi}) = h''(\mathbf{u}^{\top}\boldsymbol{\varphi}) = 0$ unless $\mathbf{u}^{\top}\boldsymbol{\varphi} \leq B$, and choosing $B = B(\|\mathbf{u}\|, \mu)$ large enough so that $B\mu \geq 4\|\mathbf{u}\|^2$, for some $C = C(\|\mathbf{u}\|, \mu) > 0$ we have

$$\mathbb{E}\operatorname{tr}(\nabla^2 f(\boldsymbol{\varphi}_t)) \leq \frac{B\mu}{4} \mathbb{E}f(\boldsymbol{\varphi}_t) + C.$$

Plugging the above bounds into (C.18), we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}f(\boldsymbol{\varphi}_t) \leq -\frac{B\mu}{4}\mathbb{E}f(\boldsymbol{\varphi}_t) + C'.$$

This implies that $\mathbb{E}f(\varphi_t) \leq x(t)$ where $x(t) = (\mathbb{E}[f(\varphi_0)] + 4C'/B\mu)e^{-tB\mu/4} - 4C'/B\mu$ is the solution of $x'(t) = -(B\mu/4)x(t) + C'$ with initialization $x(0) = \mathbb{E}f(\varphi_0)$. If $\mathbb{E}e^{\mathbf{u}^\top \varphi_0} < A$, then also $\mathbb{E}f(\varphi_0) \leq \mathbb{E}e^{\mathbf{u}^\top \varphi_0} < A$ and $\mathbb{E}e^{\mathbf{u}^\top \varphi_t} \leq e^B + \mathbb{E}f(\varphi_t)$, so this shows that there exists some $C = C(A, \|\mathbf{u}\|, \mu) > 0$ for which $\mathbb{E}e^{\mathbf{u}^\top \varphi_t} < C$ and every $t \in [0, T]$.

Finally, for any $\lambda \in \mathbb{R}^p$, we may write $\lambda = \sum_{j=1}^p \mathbf{u}_j$ where $\mathbf{u}_1, \ldots, \mathbf{u}_p$ are eigenvectors of Γ . Under the given finite moment-generating-function condition for q_0 , there exists some $A = A(q_0, \lambda) > 0$ for which $\mathbb{E}_{\varphi_0 \sim q_0} e^{p\mathbf{u}_j^\top \varphi_0} < A$ for every $j = 1, \ldots, p$. Then for any $t \in [0, T]$, applying the above with $\mathbf{u} = p\mathbf{u}_j$ yields

$$\mathbb{E}e^{\boldsymbol{\lambda}^{\top}\boldsymbol{\varphi}_{t}} = \mathbb{E}\prod_{j=1}^{p}e^{\mathbf{u}_{j}^{\top}\boldsymbol{\varphi}_{t}} \leq \prod_{j=1}^{p}\left(\mathbb{E}e^{p\mathbf{u}_{j}^{\top}\boldsymbol{\varphi}_{t}}\right)^{1/p} < C(q_{0},\boldsymbol{\lambda})$$

for some $C(q_0, \lambda) < \infty$. This shows (C.13), and specializing to $\lambda \in \{\pm (8M/\tau^2)\mathbf{e}_j\}_{j\in[p]}$ for the standard basis vectors \mathbf{e}_j shows (C.14).

Proof of Lemma C.3-Part (3). On a probability space containing $\varphi_0 \sim q_0$ and a standard Brownian motion $\{\mathbf{B}_t\}_{t\in[0,T]}$, define $\varphi = \{\varphi_t\}_{t\in[0,T]}$ and $\tilde{\varphi} = \{\tilde{\varphi}_t\}_{t\in[0,T]}$ as the (pathwise unique) solutions to

$$\boldsymbol{\varphi}_t = \boldsymbol{\varphi}_0 + \int_0^t -\nabla U_{g_s}(\boldsymbol{\varphi}_s) \mathrm{d}s + \sqrt{2}\mathbf{B}_t,$$

$$\widetilde{\varphi}_t = \varphi_0 + \int_0^t -\nabla U_{\widetilde{g}_s}(\widetilde{\varphi}_s) \mathrm{d}s + \sqrt{2}\mathbf{B}_t.$$

Thus $\varphi \sim q$ and $\widetilde{\varphi} \sim \widetilde{q}$. For any $t \in [0, T]$,

$$\begin{split} \sup_{t\in[0,T]} \|\varphi_t - \widetilde{\varphi}_t\|^2 &= \sup_{t\in[0,T]} \left\| \int_0^t \left(\nabla U_{g_s}(\varphi_s) - \nabla U_{\widetilde{g}_s}(\widetilde{\varphi}_s) \right) \mathrm{d}s \right\|^2 \\ &\leq T \cdot \sup_{t\in[0,T]} \int_0^t \left\| \nabla U_{g_s}(\varphi_s) - \nabla U_{\widetilde{g}_s}(\widetilde{\varphi}_s) \right\|^2 \mathrm{d}s \\ &\leq 2T^2 \sup_{t\in[0,T]} \left\| \nabla U_{\widetilde{g}_t}(\varphi_t) - \nabla U_{\widetilde{g}_t}(\widetilde{\varphi}_t) \right\|^2 + 2T \int_0^T \left\| \nabla U_{g_t}(\varphi_t) - \nabla U_{\widetilde{g}_t}(\varphi_t) \right\|^2 \mathrm{d}t. \end{split}$$

As shown in the proof of Part (1), $\nabla U_g : \mathbb{R}^p \to \mathbb{R}^p$ is Lipschitz uniformly over $g \in \mathcal{P}(M)$, so the first term is at most $CT^2 \sup_{t \in [0,T]} \|\varphi_t - \widetilde{\varphi}_t\|^2 \leq \frac{1}{2} \sup_{t \in [0,T]} \|\varphi_t - \widetilde{\varphi}_t\|^2$ for some $C, T_0 > 0$ and any $T \leq T_0$. For the second term, using (C.16),

$$\|\nabla U_{g_t}(\varphi_t) - \nabla U_{\widetilde{g}_t}(\varphi_t)\|^2 \le C \sum_{j=1}^p \left(\theta_{g_t}(\varphi_{t,j}) - \theta_{\widetilde{g}_t}(\varphi_{t,j})\right)^2.$$

Applying Proposition B.1, we have

$$\begin{split} \left| \theta_{g}(\varphi) - \theta_{\widetilde{g}}(\varphi) \right| &= \left| \frac{\int_{-M}^{M} \theta \cdot \mathcal{N}_{\tau}(\varphi - \theta) g(\theta) \mathrm{d}\theta}{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) g(\theta) \mathrm{d}\theta} - \frac{\int_{-M}^{M} \theta \cdot \mathcal{N}_{\tau}(\varphi - \theta) \widetilde{g}(\theta) \mathrm{d}\theta}{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) |g(\theta) - \widetilde{g}(\theta)| \mathrm{d}\theta} \right| \\ &\leq \frac{\int_{-M}^{M} \theta \cdot \mathcal{N}_{\tau}(\varphi - \theta) |g(\theta) - \widetilde{g}(\theta)| \mathrm{d}\theta}{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) g(\theta) \mathrm{d}\theta} \\ &+ \int_{-M}^{M} \theta \cdot \mathcal{N}_{\tau}(\varphi - \theta) \widetilde{g}(\theta) \mathrm{d}\theta \cdot \frac{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) |g(\theta) - \widetilde{g}(\theta)| \mathrm{d}\theta}{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) g(\theta) \mathrm{d}\theta} \cdot \frac{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) |g(\theta) - \widetilde{g}(\theta)| \mathrm{d}\theta}{\int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) g(\theta) \mathrm{d}\theta \cdot \int_{-M}^{M} \mathcal{N}_{\tau}(\varphi - \theta) \widetilde{g}(\theta) \mathrm{d}\theta} \\ &\leq C e^{(4M/\tau^{2})|\varphi|} \cdot \int_{-M}^{M} |g(\theta) - \widetilde{g}(\theta)| \mathrm{d}\theta. \end{split}$$

In summary, there exist $C', T_0 > 0$ such that for any $T \leq T_0$,

$$\sup_{t\in[0,T]} \|\boldsymbol{\varphi}_t - \widetilde{\boldsymbol{\varphi}}_t\|^2 \le C'T \int_0^T \sum_{j=1}^p e^{(4M/\tau^2)|\boldsymbol{\varphi}_{t,j}|} \mathrm{d}t \cdot d_{\infty,\mathrm{TV}}(g,\widetilde{g})^2.$$

Taking expectation on both sides and using the bound (C.14) with $K(q_0) = K$ gives $d_W^2(q, \tilde{q}) \leq C_0^2 T^2 d_{\infty, \text{TV}}(q, \tilde{g})^2$, for some $C_0 > 0$ depending on K. This completes the proof.

We now provide a lemma that will allow us to inductively establish the smoothness of the preceding solutions $\{q_t, g_t\}$ in both space and time.

Lemma C.4. In the setting of Lemma C.3, suppose in addition that $g_t \in \mathcal{P}_*(M)$ admits a density for each $t \in [0,T]$, where $t \mapsto g_t(\theta)$ belongs to $C^r((0,T))$ for some $r \ge 0$ and every $\theta \in [-M,M]$, and $\sup_{t \in (0,T)} \int_{-M}^{M} |\frac{\mathrm{d}^a}{\mathrm{d}t^a} g_t(\theta)| \mathrm{d}\theta < \infty$ for all $a = 0, \ldots, r$. Then: 1. The map $t \mapsto f_t(\theta) := \mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * q_t}(\theta)$ belongs also to $C^r((0,T))$, and for all $a = 0, \ldots, r$,

$$\sup_{t \in (0,T)} \sup_{\theta \in [-M,M]} \left| \frac{\mathrm{d}^a}{\mathrm{d}t^a} f_t(\theta) \right| < \infty.$$
(C.19)

2. If $q_0 \in \mathcal{P}_*(\mathbb{R}^p)$ has a continuous bounded density, then the marginal laws $\{q_t\}_{t\in[0,T]}$ of $\{\varphi_t\}_{t\in[0,T]}$ admit continuous densities bounded uniformly over $t \in [0,T]$. Furthermore if $r \geq 1$, then $(t,\varphi) \mapsto q_t(\varphi)$ belongs to $C^{r-1,\infty}((0,T) \times \mathbb{R}^p)$.

Proof. For part (1), we claim by induction on $a \in \{0, 1, ..., r\}$ that $f_t(\theta)$ belongs to $C^a((0,T))$, the bound (C.19) holds for a, and $\frac{d^a}{dt^a} f_t(\theta) = \mathbb{E}_{\varphi \sim q_t}[F^a_{t,\theta}(\varphi)]$ for some $F^a_{t,\theta}(\varphi)$ that is a polynomial of the arguments

$$\varphi_k, \qquad \frac{\int_{-M}^{M} \frac{\mathrm{d}^i}{\mathrm{d}\varphi_k^i} \mathcal{N}_{\tau}(\varphi_k - \theta) \frac{\mathrm{d}^j}{\mathrm{d}t^j} g_t(\theta) \mathrm{d}\theta}{\mathcal{N}_{\tau} * g_t(\varphi_k)} \quad \text{and} \quad \frac{\frac{\mathrm{d}^i}{\mathrm{d}\varphi_k^i} \mathcal{N}_{\tau}(\varphi_k - \theta)}{\mathcal{N}_{\tau} * g_t(\varphi_k)} \tag{C.20}$$

for $k \in \{1, \dots, p\}, i \ge 0$, and $j \in \{0, \dots, a\}$.

For the base case a = 0, observe that

$$f_t(\theta) = \mathcal{N}_\tau * \frac{\bar{q}_t}{\mathcal{N}_\tau * g_t}(\theta) = \mathbb{E}_{\varphi \sim q_t}[F_{t,\theta}^0(\varphi)], \qquad F_{t,\theta}^0(\varphi) := \frac{1}{p} \sum_{k=1}^p \frac{\mathcal{N}_\tau(\varphi_k - \theta)}{\mathcal{N}_\tau * g_t(\varphi_k)}$$

By the bound $\sup_{t\in[0,T]} \sup_{\theta\in[-M,M]} |\mathcal{N}_{\tau}(\varphi_k - \theta)/\mathcal{N}_{\tau} * g_t(\varphi_k)| \leq Ce^{(2M/\tau^2)|\varphi|}$ from Proposition B.1 and Lemma C.3-(2), we see that $\sup_{t\in[0,T]} \sup_{\theta\in[-M,M]} |f_t(\theta)| < \infty$, and the maps $\varphi \mapsto F^0_{t,\theta}(\varphi)q_t(\varphi)$ for $t \in [0,T]$ are uniformly integrable, hence $f_t(\theta)$ is continuous in t. So the inductive claims hold for a = 0. Assuming they hold for a - 1 where $1 \leq a \leq r$, we have that $F^{a-1}_{t,\theta}(\varphi)$ belongs to $C^{1,\infty}((0,T) \times \mathbb{R}^p)$. Then by Ito's formula,

$$\frac{\mathrm{d}^{a}}{\mathrm{d}t^{a}}f_{t}(\theta) = \frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}[F_{t,\theta}^{a-1}(\varphi)] = \mathbb{E}_{\varphi \sim q_{t}}\left[\left(\frac{\mathrm{d}}{\mathrm{d}t}F_{t,\theta}^{a-1} - \nabla U_{g_{t}}^{\top}\nabla F_{t,\theta}^{a-1} + \Delta F_{t,\theta}^{a-1}\right)(\varphi)\right].$$

From the form (C.15), we have that each coordinate of $\nabla U_{g_t}(\varphi)$ is a polynomial of terms of the form (C.20), and it is easily verified that differentiating any term (C.20) in either t or φ_k yields again a polynomial of terms (C.20), where differentiating in t increases the maximum order j of the time derivative of $g_t(\theta)$ by at most 1. Thus the above may be written as $\frac{d^a}{dt^a} f_t(\theta) = \mathbb{E}_{\varphi \sim qt}[F^a_{t,\theta}(\varphi)]$ where $F^a_{t,\theta}(\varphi)$ is a polynomial of the terms (C.20) with $j \leq a$. By Proposition B.1, for each fixed $i \geq 0$ we have $\sup_{t \in [0,T]} \sup_{\theta \in [-M,M]} \left| \frac{d^i}{d\varphi_k^i} \mathcal{N}_{\tau}(\varphi_k - \theta) / \mathcal{N} * g_t(\varphi_k) \right| \leq Ce^{C|\varphi_k|}$ for some C > 0. Then, applying the given condition $\sup_{t \in (0,T)} \int_{-M}^{M} \left| \frac{d^a}{dt^a} g_t(\theta) \right| d\theta < \infty$ and Lemma C.3-(2), we get $\sup_{t \in (0,T)} \sup_{\theta \in [-M,M]} \left| \frac{d^a}{dt^a} f_t(\theta) \right| < \infty$. Furthermore, $\varphi \mapsto F^a_{t,\theta}(\varphi) q_t(\varphi)$ is uniformly integrable over $t \in (0,T)$, so $\frac{d^a}{dt^a} f_t(\theta)$ is continuous in t. This completes the induction, showing the inductive claim for all $a = 0, \ldots, r$, and hence establishes Part (1).

For Part (2), let us write $q_t(\varphi) \in H^{j,k} \equiv H^{j,k}((0,T) \times \mathbb{R}^p)$ if all components of the weak derivatives $D_t^a D_{\varphi}^b q_t(\varphi)$ for $a \leq j$ and $b \leq k$ exist and belong to $L^r((0,T) \times \mathbb{R}^p)$ (defined with respect to Lebesgue measure) for every $r < \infty$. We write $q_t(\varphi) \in H_{\text{loc}}^{j,k}$ if for any compact subset $S \subset (0,T) \times \mathbb{R}^p$, these derivatives belong to $L^r(S)$ for every $r < \infty$. By Holder's inequality and the Leibniz rule for weak differentiation, $H_{\text{loc}}^{j,k}$ is closed under sums and products. Lemma C.3-(2) implies that for any fixed $k \geq 0$, some C, C' > 0, and all $t \in [0,T]$,

$$\mathbb{E}_{\boldsymbol{\varphi} \sim q_t}[\|\boldsymbol{\varphi}\|_2^k] \le C \sum_{j=1}^p \mathbb{E}_{\boldsymbol{\varphi} \sim q_t}[|\varphi_j|^k] \le C'.$$
(C.21)

Then the expression (C.16) for ∇U_{g_t} implies, for any fixed k > 0, that $\mathbb{E}_{\varphi \sim q_t}[\|\nabla U_{g_t}(\varphi)\|^k] < C$ uniformly over $t \in [0, T]$. Then [BKRS22, Corollaries 6.4.3, 7.3.8] show that $\{q_t\}_{t \in [0,T]}$ admit continuous densities with respect to Lebesgue measure, uniformly bounded over $t \in [0, T]$, which further satisfy

$$q_t(\boldsymbol{\varphi}) \in H^{0,1}_{\text{loc}}.\tag{C.22}$$

To improve this smoothness to $H_{\text{loc}}^{r,\infty}$, we apply a similar argument as in [JKO98, Theorem 5.1]: Fix $(t, \varphi) \in (0, T) \times \mathbb{R}^p$ and apply the Ito formula (C.17) with $\zeta_s(\mathbf{x}) = \omega(s)\eta(\mathbf{x})\mathcal{N}_{2(\delta+t-s)}(\varphi - \mathbf{x})$, where $\delta > 0$, $\eta : \mathbb{R}^p \to [0, 1]$ is a smooth compactly supported function on \mathbb{R}^p , and $\omega : \mathbb{R} \to [0, 1]$ is a smooth compactly supported function satisfying $\omega(s) = 0$ for all $s \leq 0$ and $\omega(t) = 1$. Then, applying $\partial_s \mathcal{N}_{2(\delta+t-s)}(\varphi - \mathbf{x}) + \Delta \mathcal{N}_{2(\delta+t-s)}(\varphi - \mathbf{x}) = 0$ (the heat equation) and the chain rule, (C.17) yields

$$\int \eta(\mathbf{x})q_t(\mathbf{x})\mathcal{N}_{2\delta}(\boldsymbol{\varphi} - \mathbf{x})d\mathbf{x}$$

= $\int_{-\infty}^t \int \left[\omega'(s)\eta(\mathbf{x}) - \omega(s)\nabla U_{g_s}(\mathbf{x})^\top \nabla \eta(\mathbf{x}) + \omega(s)\Delta \eta(\mathbf{x}) \right] \mathcal{N}_{2(\delta+t-s)}(\boldsymbol{\varphi} - \mathbf{x})q_s(\mathbf{x})d\mathbf{x}ds$
+ $\int_{-\infty}^t \int \left[\omega(s)\eta(\mathbf{x})\nabla U_{g_s}(\mathbf{x}) - \omega(s)\nabla \eta(\mathbf{x}) \right]^\top \nabla \mathcal{N}_{2(\delta+t-s)}(\boldsymbol{\varphi} - \mathbf{x})q_s(\mathbf{x})d\mathbf{x}ds.$

Applying the weak differentiability in \mathbf{x} of $q_s(\mathbf{x})$ from (C.22) to integrate by parts in the second term, and taking the limit $\delta \to 0$, this yields

$$\eta(\boldsymbol{\varphi})q_t(\boldsymbol{\varphi}) = \int_{-\infty}^t \left[\left(q_s [\omega'(s)\eta - \omega(s)\nabla U_{g_s}^\top \nabla \eta + \omega(s)\Delta \eta] - \nabla \cdot \left(q_s [\omega(s)\eta\nabla U_{g_s} - \omega(s)\nabla \eta] \right) \right) * \mathcal{N}_{t-s} \right] (\boldsymbol{\varphi}) \mathrm{d}s.$$
(C.23)

The estimate of [LSU68, Chapter 4, (3.1)] shows the implication

$$f_t(\boldsymbol{\varphi}) \in H^{j,k}$$
 for all $2j + k \le 2m \Rightarrow \int_{-\infty}^t [f_s * \mathcal{N}_{t-s}](\boldsymbol{\varphi}) \mathrm{d}s \in H^{j,k}$ for all $2j + k \le 2m + 2$. (C.24)

From (C.15) and the given condition $\sup_{t \in (0,T)} \int_{-M}^{M} |\frac{d^{a}}{dt^{a}} g_{t}(\theta)| d\theta < \infty$ for each $a \leq r$, it is direct to check that each coordinate of $(t, \varphi) \mapsto \nabla U_{g_{t}}(\varphi)$ belongs to $H_{\text{loc}}^{r,\infty}$. Then (C.24) yields the implications $q_{t}(\varphi) \in H_{\text{loc}}^{0,1} \Rightarrow q_{t}(\varphi) \in H_{\text{loc}}^{0,2} \Rightarrow q_{t}(\varphi) \in H_{\text{loc}}^{0,3} \Rightarrow \dots$ where the initial condition comes from (C.22), hence $q_{t}(\varphi) \in H_{\text{loc}}^{0,\infty}$. Then (C.24) further yields the implications $q_{t}(\varphi) \in H_{\text{loc}}^{0,\infty} \Rightarrow q_{t}(\varphi) \in H_{\text{loc}}^{r,\infty}$. Finally, by the Sobolev embedding theorem (cf. [AF03, Theorem 4.2 Part II]) we have $H_{\text{loc}}^{r,\infty} \subset C^{r-1,\infty}((0,T) \times \mathbb{R}^{p})$.

C.3 Completing the proof

We are now ready to prove Theorem 3.6.

Proof of Theorem 3.6. We start by showing the existence and uniqueness of a solution of (2.15–2.16) on $[0, t_0]$ for some small t_0 to be specified later. For the given initialization $q_0 \in \mathcal{P}_*(\mathbb{R}^p)$, let K > 0 be such that (C.14) holds with $K(q_0) = K$, and define

$$\mathcal{L}'_{q}(t_{0}) = \{ q \in \mathcal{L}_{q}(t_{0}) : \max_{t \in [0, t_{0}]} \max_{j \in [p]} \mathbb{E}_{\varphi_{t} \sim q_{t}} e^{(8M/\tau^{2})|\varphi_{t,j}|} \le K \}.$$
 (C.25)

Observe that $\mathcal{L}'_q(t_0)$ is closed in $\mathcal{L}_q(t_0)$. Indeed, since d_W metrizes weak convergence, for any sequence $q_n \in \mathcal{L}'_q(t_0)$ converging to $q \in \mathcal{L}_q(t_0)$, it holds that $q_n \Rightarrow q$ weakly, and thus $q_{n,t} \Rightarrow q_t$ weakly for all $t \in [0, t_0]$. Then by Fatou's lemma, $\mathbb{E}_{\varphi_t \sim q_t} e^{(8M/\tau^2)|\varphi_{t,j}|} \leq \liminf_n \mathbb{E}_{\varphi_t \sim q_{n,t}} e^{(8M/\tau^2)|\varphi_{t,j}|} \leq K(q_0)$. This shows that $\mathcal{L}'_q(t_0)$ is closed, and hence also complete.

Define a map $\Gamma : \mathcal{L}'_q(t_0) \times \mathcal{L}_g(t_0) \to \mathcal{L}'_q(t_0) \times \mathcal{L}_g(t_0)$ where $\Gamma(q,g) = (\Gamma_q(g), \Gamma_g(q))$, as follows:

- $\Gamma_q(g)$ is the law of the unique strong solution to (C.10) defined by g, with initialization q_0 , as guaranteed by Lemma C.3-(1). Here, Lemma C.3-(2) ensures that $\Gamma_q(g) \in \mathcal{L}'_q(t_0)$.
- $\Gamma_g(q)$ is the unique solution to (C.3) defined by q, with initialization g_0 , as guaranteed by Lemma C.1-(1).

Equip $\mathcal{L}'_q(t_0) \times \mathcal{L}_g(t_0)$ with the metric

$$d_*((q,g),(\widetilde{q},\widetilde{g})) = d_W(q,\widetilde{q}) + d_{\infty,\mathrm{TV}}(g,\widetilde{g}).$$

Since both \mathcal{L}'_q and \mathcal{L}_g are complete metric spaces, $\mathcal{L}'_q \times \mathcal{L}_g$ is also complete. Moreover, by Lemmas C.1-(2) and C.3-(3), for any $q, \tilde{q} \in \mathcal{L}'_q(t_0)$ and $g, \tilde{g} \in \mathcal{L}_g(t_0)$, there exist $C_0, T_0 > 0$ depending on K for which

$$d_*(\Gamma(q,g),\Gamma(\widetilde{q},\widetilde{g})) \le C_0 t_0 \cdot d_*((q,g),(\widetilde{q},\widetilde{g}))$$

for any $t_0 \leq T_0$. Then, choosing t_0 to satisfy also $t_0 \leq 1/2C_0$, we obtain that Γ is a contraction in d_* . By the Banach fixed point theorem, there is a unique fixed point $(q^*, g^*) \in \mathcal{L}'_q(t_0) \times \mathcal{L}_g(t_0)$ of Γ .

By Lemma C.1-(1), g^* is a (classical) solution to (2.16) defined by q^* and the initial condition g_0 , and by Lemma C.3-(1), q^* is a weak solution to (C.11) defined by g^* with the initial condition q_0 , in the sense (C.12). Suppose inductively that $t \mapsto g_t^*(\theta)$ belongs to $C^r((0, t_0))$ for each fixed $\theta \in [-M, M]$ and that $\sup_{t \in (0, t_0)} \int_{-M}^{M} |\frac{\mathrm{d}^a}{\mathrm{d}t^a} g_t^*(\theta)| \mathrm{d}\theta < \infty$ for all $a = 0, \ldots, r$. Then Lemma C.4-(1) shows that $t \mapsto f_t(\theta) = \mathcal{N}_{\tau} * \frac{\bar{q}_t^*}{\mathcal{N}_{\tau} * g_t^*}(\theta)$ belongs also to $C^r((0, t_0))$ and $\sup_{t \in (0, t_0)} \sup_{\theta \in [-M, M]} |\frac{\mathrm{d}^a}{\mathrm{d}t^a} f_t(\theta)| < \infty$ for each $a = 0, \ldots, r$. Then $\frac{\mathrm{d}^{r+1}}{\mathrm{d}t^{r+1}} g_t(\theta) = \frac{\mathrm{d}^r}{\mathrm{d}t^r} [g_t(\theta)(f_t(\theta) - 1)]$ is continuous in t, and satisfies

$$\begin{split} \sup_{t \in (0,t_0)} \int_{-M}^{M} \left| \frac{\mathrm{d}^{r+1}}{\mathrm{d}t^{r+1}} g_t(\theta) \right| \mathrm{d}\theta &= \sup_{t \in (0,t_0)} \int_{-M}^{M} \left| \frac{\mathrm{d}^r}{\mathrm{d}t^r} [g_t(\theta)(f_t(\theta) - 1)] \right| \mathrm{d}\theta \\ &\leq \sup_{t \in (0,t_0)} \sum_{a=0}^r \binom{r}{a} \sup_{\theta \in [-M,M]} \left| \frac{\mathrm{d}^{r-a}}{\mathrm{d}t^{r-a}} (f_t(\theta) - 1) \right| \int_{-M}^{M} \left| \frac{\mathrm{d}^a}{\mathrm{d}t^a} g_t(\theta) \right| \mathrm{d}\theta < \infty. \end{split}$$

Thus the inductive claim holds for r+1. Then $t \mapsto g_t^*(\theta)$ belongs to $C^{\infty}((0,t_0))$ for each fixed θ , and $\sup_{t \in (0,t_0)} \int_{-M}^{M} |\frac{d^r}{dt^r} g_t^*(\theta)| d\theta < \infty$ for each fixed $r \ge 0$. Then Lemma C.4-(2) implies that $(t, \varphi) \mapsto q_t^*(\varphi)$ belongs to $C^{\infty}((0,t_0) \times \mathbb{R}^p)$. Applying this smoothness of $q_t^*(\varphi)$ and integrating (C.12) by parts, we obtain that q^* satisfies (2.15) pointwise at each $(t,\varphi) \in (0,t_0) \times \mathbb{R}^p$. Furthermore, the uniform boundedness and continuity of $q_t^*(\varphi)$ and weak convergence $q_t^* \Rightarrow q_0^*$ imply that $q_t^*(\varphi) \to q_0^*(\varphi)$ pointwise over $\varphi \in \mathbb{R}^p$, so $\{q_t^*\}_{t \in [0,t_0]}$ is a classical solution to (2.15).

Conversely, for any solution (q, g) to (2.16-2.15), Lemma C.3-(2) applied with this g shows that we must have $q \in \mathcal{L}'_q(t_0)$. Then $(q, g) \in \mathcal{L}'_q(t_0) \times \mathcal{L}_g(t_0)$ is also a fixed point of Γ , so by the uniqueness guarantees in Lemmas C.1-(1) and C.3-(1), $q = q^*$ and $g = g^*$ as equalities of elements in $\mathcal{L}'_q(t_0)$ and $\mathcal{L}_g(t_0)$. Since (q, g) are continuous, this implies $g_t(\theta) = g_t^*(\theta)$ and $q_t(\varphi) = q_t^*(\varphi)$ pointwise, so the solution (q^*, g^*) to (2.15-2.16) is unique.

We now extend this solution to [0, T], by applying this argument sequentially. Let $\{g_t, q_t\}_{t \in [0, t_0]}$ be the above solution over $[0, t_0]$. For any T > 0 and extension of $\{g_t\}_{t \in [0, t_0]}$ to $\{g_t\}_{t \in [0, t_0+T]} \in$ $\mathcal{L}_g(t_0+T)$, Lemma C.3 shows that there is a unique solution to the diffusion (C.10) over $[0, t_0+T]$ with initialization q_0 , which satisfies (C.14) for all $t \in [0, t_0 + T]$ with $K(q_0) = K$ for the above value K > 0 defining (C.25). The portion of this solution on $[t_0, t_0 + T]$ must coincide with the unique solution to (C.11) over [0, T] that is initialized at q_{t_0} and defined by $\{g_t\}_{t\in[t_0,t_0+T]}$. So for the initialization q_{t_0} , we must have that (C.14) holds with $K(q_{t_0}) = K$ and the same value Kas above. Then we may apply the above argument to $[t_0, 2t_0]$ with (q_{t_0}, g_{t_0}) in place of (q_0, g_0) , and then sequentially to $[2t_0, 3t_0]$, etc. This proves the existence and uniqueness of the solution $\{(q_t, g_t)\}_{t\in[0,T]}$ to (2.15–2.16), as claimed. The properties $g_t(\theta) > 0$ and $\sup_{t\in[0,T]} \mathbb{E}_{\varphi_t \sim q_t} e^{\lambda^\top \varphi_t} < \infty$ follow from the properties of the solutions guaranteed by Lemmas C.1-(1) and C.3-(2), and this concludes the proof.

C.4 Proof of Lemma B.4

Proof of Lemma B.4. Recall the form of F_n in (2.11). To show $\limsup_{t\to 0} F_n(q_t, g_t) \leq F_n(q_0, g_0)$, recall from (C.22) that for any fixed T > 0, $q_t(\varphi)$ is uniformly bounded over $t \in [0, T]$. Since $U_{g_t}(\varphi)$ is also uniformly bounded over $\|\varphi\|_2 \leq R$, for any R > 0,

$$\lim_{t\to 0} \int \mathbf{1}\{\|\boldsymbol{\varphi}\|_{\infty} \leq R\} U_{g_t}(\boldsymbol{\varphi}) q_t(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} = \int \mathbf{1}\{\|\boldsymbol{\varphi}\|_{\infty} \leq R\} U_{g_0}(\boldsymbol{\varphi}) q_0(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi}$$

by the bounded convergence theorem. Applying $U_{g_t}(\varphi) < C(\|\varphi\|_2^2 + 1)$ and Cauchy-Schwarz,

$$\int \mathbf{1}\{\|\boldsymbol{\varphi}\|_{\infty} > R\} |U_{g_t}(\boldsymbol{\varphi})| q_t(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \le \mathbb{P}_{q_t}[\|\boldsymbol{\varphi}_t\|_{\infty} > R]^{1/2} (\mathbb{E}_{q_t}[\|\boldsymbol{\varphi}_t\|^4] + 1)^{1/2}]$$

By Lemma C.3-(2), $\mathbb{E}_{q_t}[\|\varphi_t\|^4] < C$ and $\mathbb{P}_{q_t}[|\varphi_{t,j}| > R] \leq e^{-R}\mathbb{E}[e^{|\varphi_{t,j}|}] < Ce^{-R}$ for a constant C > 0, uniformly over $j = 1, \ldots, p$ and $t \in [0, T]$. Thus the above display converges to 0 uniformly over $t \in [0, T]$ as $R \to \infty$, so

$$\lim_{t\to 0} \int U_{g_t}(\boldsymbol{\varphi}) q_t(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} = \int U_{g_0}(\boldsymbol{\varphi}) q_0(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi}.$$

Applying the uniform upper bound $\log q_t(\varphi) \leq C_0$ and Fatou's lemma, we have also

$$\int [C_0 - \log q_0(\boldsymbol{\varphi})] q_0(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \le \liminf_{t \to 0} \int [C_0 - \log q_t(\boldsymbol{\varphi})] q_t(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi},$$

hence $\limsup_{t\to 0} \int (\log q_t) q_t \leq \int (\log q_0) q_0$. Combining these statements gives $\limsup_{t\to 0} F_n(q_t, g_t) \leq F_n(q_0, g_0)$.

To show the form (B.14) for the derivative of F_n , we will appeal to the following properties of $\{q_t\}_{t \in [0,T]}$: For any fixed 0 < t < T and compact subset $K \subset \mathbb{R}^p$,

$$\sup_{\in[t,T]} \sup_{\varphi \in \mathbb{R}^p} \frac{|\log q_s(\varphi)|}{\|\varphi\|^2 + 1} < \infty,$$
(C.26)

$$\sup_{s \in [t,T]} \sup_{\varphi \in K} \|\nabla q_s(\varphi)\|_2 < \infty, \tag{C.27}$$

$$\sup_{s \in [t,T]} \sup_{\varphi \in K} \|\nabla^2 q_s(\varphi)\|_{\mathcal{F}} < \infty,$$
(C.28)

$$\int_{t}^{T} \int_{\mathbb{R}^{p}} \|\nabla \log q_{s}(\boldsymbol{\varphi})\|_{2}^{2} q_{s}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \mathrm{d}s < \infty.$$
(C.29)

The upper bound for $\log q_s(\varphi)$ in (C.26) follows from (C.22), and the lower bound follows from [BKRS22, Example 8.2.3], where we may take $\beta = 1$ in light of the form (C.16) for the drift $\nabla U_{g_t}(\varphi)$. The next two statements (C.27–C.28) follow from the C^{∞} smoothness of $(t, \varphi) \mapsto q_t(\varphi)$. By Lemma C.3-(2), we must have $\mathbb{E}_{q_s}[\|\varphi_s\|_2^2] < C$ for some C > 0 uniformly over $s \in [0, T]$. Then (C.26) implies $\int |\log q_t(\varphi)| q_t(\varphi) d\varphi < \infty$, (C.16) implies

$$\int \|\nabla U_{g_s}(\varphi)\|_2^2 q_s(\varphi) \mathrm{d}\varphi < C \tag{C.30}$$

for some C > 0 uniformly over $s \in [0, T]$, and the last statement (C.29) then follows from [BKRS22, Theorem 7.4.1].

Let $\boldsymbol{\eta}(\boldsymbol{\varphi}) = \prod_{j=1}^{p} \eta(\varphi_j)$ where η is a smooth, compactly supported bump function on \mathbb{R} . Define a truncated version of F_n by

$$F_n^{\eta}(q_t, g_t) = \frac{1}{p} \int \boldsymbol{\eta}(\boldsymbol{\varphi}) [U_{g_t}(\boldsymbol{\varphi}) + \log q_t(\boldsymbol{\varphi})] q_t(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} + \frac{n}{2p} \log(2\pi) + \frac{1}{2p} \log \det \boldsymbol{\Sigma}.$$

Then for any t > 0, we may decompose

$$\frac{\mathrm{d}}{\mathrm{d}t}F_n^\eta(q_t, g_t) = \mathrm{I}^\eta(t) + \mathrm{II}^\eta(t)$$

Here, we set

$$\begin{split} \mathbf{I}^{\eta}(t) &= \frac{1}{p} \int \boldsymbol{\eta}(\boldsymbol{\varphi}) \cdot \frac{\mathrm{d}}{\mathrm{d}t} U_{g_{t}}(\boldsymbol{\varphi}) \cdot q_{t}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \\ &= \int \boldsymbol{\eta}(\boldsymbol{\varphi}) \cdot \frac{\mathrm{d}}{\mathrm{d}t} \Big(-\log[\mathcal{N}_{\tau} * g_{t}](\boldsymbol{\varphi}) \Big) \cdot \bar{q}_{t}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} = -\alpha \int \left(\left[\mathcal{N}_{\tau} * \frac{\eta \bar{q}_{t}}{\mathcal{N}_{\tau} * g_{t}} \right](\boldsymbol{\theta}) - 1 \right)^{2} g_{t}(\boldsymbol{\theta}) \mathrm{d}\boldsymbol{\theta}, \end{split}$$

where these calculations and argument for differentiating under the integral are the same as in (B.4). We also set

$$\begin{split} \mathrm{II}^{\eta}(t) &= \frac{1}{p} \int \boldsymbol{\eta}(\boldsymbol{\varphi}) [U_{g_{t}}(\boldsymbol{\varphi}) + \log q_{t}(\boldsymbol{\varphi}) + 1] \cdot \frac{\mathrm{d}}{\mathrm{d}t} q_{t}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \\ &= \frac{1}{p} \int \boldsymbol{\eta}(\boldsymbol{\varphi}) [U_{g_{t}}(\boldsymbol{\varphi}) + \log q_{t}(\boldsymbol{\varphi}) + 1] \cdot \left(\nabla \cdot [q_{t}(\boldsymbol{\varphi}) \nabla U_{g_{t}}(\boldsymbol{\varphi})] + \Delta q_{t}(\boldsymbol{\varphi}) \right) \mathrm{d}\boldsymbol{\varphi}, \end{split}$$

where we may differentiate under the integral in the first line using the dominated convergence theorem and boundedness of U_{g_t} , ∇U_{g_t} , $\log q_t$, $\|\nabla q_t\|$, $|\Delta q_t|$ in compact subsets of $(0, T) \times \mathbb{R}^p$ as follows from (C.26–C.28). Applying integration by parts in φ , we then have $\Pi^{\eta}(t) = \Pi\Pi^{\eta}(t) + IV^{\eta}(t)$ where

$$\begin{aligned} \mathrm{III}^{\eta}(t) &= -\frac{1}{p} \int \boldsymbol{\eta}(\boldsymbol{\varphi}) \left\| \nabla \left(U_{g_{t}}(\boldsymbol{\varphi}) + \log q_{t}(\boldsymbol{\varphi}) \right) \right\|_{2}^{2} q_{t}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi}, \\ \mathrm{IV}^{\eta}(t) &= -\frac{1}{p} \int [U_{g_{t}}(\boldsymbol{\varphi}) + \log q_{t}(\boldsymbol{\varphi}) + 1] \cdot \nabla \boldsymbol{\eta}(\boldsymbol{\varphi})^{\top} \nabla [U_{g_{t}}(\boldsymbol{\varphi}) + \log q_{t}(\boldsymbol{\varphi})] q_{t}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi}. \end{aligned}$$

Take $\eta \equiv \eta_R$ such that $\eta_R(\varphi) = 1$ for $|\varphi| \leq R$, $\eta_R(\varphi) = 0$ for $|\varphi| \geq R+1$, and $\eta'_R(\varphi)$ is bounded by a constant. Then, recalling from the arguments preceding (B.2) that $\mathcal{N}_{\tau} * \frac{q_t}{\mathcal{N}_{\tau}*g}$ is uniformly bounded, a direct application of the dominated convergence theorem yields

$$\lim_{R \to \infty} \int_t^T \mathbf{I}^{\eta_R}(s) \mathrm{d}s = -\alpha \int_t^T \int_{-M}^M \left(\left[\mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s} \right](\theta) - 1 \right)^2 g_s(\theta) \mathrm{d}s.$$

Applying the dominated convergence theorem using (C.30) and (C.29), we also have

$$\lim_{R \to \infty} \int_t^T \operatorname{III}^{\eta_R}(s) \mathrm{d}s = -\frac{1}{p} \int_t^T \int \left\| \nabla \left(U_{g_s}(\varphi) + \log q_s(\varphi) \right) \right\|_2^2 q_s(\varphi) \mathrm{d}\varphi \mathrm{d}s.$$

Finally, applying again (C.30) and (C.29), the bounds $|U_{g_s}(\varphi)|, |\log q_s(\varphi)| \leq C(||\varphi||_2^2+1)$ uniformly over $s \in [t, T]$, and Holder's inequality, for some C, C' > 0 we have

$$\begin{split} \int_{t}^{T} |\mathrm{IV}^{\eta_{R}}(s)| \mathrm{d}s &\leq C \left[\int_{t}^{T} \int \mathbf{1} \{ \nabla \boldsymbol{\eta}_{R}(\boldsymbol{\varphi}) \neq 0 \} q_{s}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \mathrm{d}s \right]^{1/4} \cdot \left[\int_{t}^{T} \int (\|\boldsymbol{\varphi}\|_{2}^{8} + 1) q_{s}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \mathrm{d}s \right]^{1/4} \\ & \cdot \left[\int_{t}^{T} \int \left\| \nabla \left(U_{g_{s}}(\boldsymbol{\varphi}) + \log q_{s}(\boldsymbol{\varphi}) \right) \right\|_{2}^{2} q_{s}(\boldsymbol{\varphi}) \mathrm{d}\boldsymbol{\varphi} \mathrm{d}s \right]^{1/2} \\ & \leq C' \sup_{s \in [t,T]} \mathbb{P}_{q_{s}} [\nabla \boldsymbol{\eta}_{R}(\boldsymbol{\varphi}_{s}) \neq 0]^{1/4} \cdot \sup_{s \in [t,T]} (\mathbb{E}_{q_{s}}[\|\boldsymbol{\varphi}_{s}\|_{2}^{8}] + 1)^{1/4}. \end{split}$$

By Lemma C.3-(2), $\mathbb{E}_{q_s}[\|\varphi_s\|^8] < C$ and $\mathbb{P}_{q_s}[|\varphi_{s,j}| > R] \leq e^{-R}\mathbb{E}[e^{|\varphi_{s,j}|}] < Ce^{-R}$ for a constant C > 0, uniformly over $j = 1, \ldots, p$ and $s \in [0, T]$. Thus

$$\lim_{R \to \infty} \int_t^T |\mathrm{IV}^{\eta_R}(s)| \mathrm{d}s = 0$$

Combining these statements,

$$\begin{aligned} F_n(q_T, g_T) &- F_n(q_t, g_t) \\ &= \lim_{R \to \infty} F_n^{\eta_R}(q_T, g_T) - F_n^{\eta_R}(q_t, g_t) \\ &= \lim_{R \to \infty} \int_t^T \left(I^{\eta_R}(s) + III^{\eta_R}(s) + IV^{\eta_R}(s) \right) \mathrm{d}s \\ &= \int_t^T \left(-\alpha \int \left(\left[\mathcal{N}_\tau * \frac{\bar{q}_s}{\mathcal{N}_\tau * g_s} \right](\theta) - 1 \right)^2 g_s(\theta) \mathrm{d}s - \frac{1}{p} \int \left\| \nabla \left(U_{g_s}(\varphi) + \log q_s(\varphi) \right) \right\|_2^2 q_s(\varphi) \mathrm{d}\varphi \right) \mathrm{d}s \end{aligned}$$

where the first equality follows from applying the dominated convergence theorem to $\lim_{R\to\infty} F_n^{\eta_R}$. Differentiating both sides in T shows that $T \mapsto F_n(q_T, g_T)$ is differentiable at any T > 0, with derivative given by (B.14).

D Derivations of gradient flows

We discuss the derivations of the Wasserstein-2 and Fisher-Rao gradient flow equations (2.15–2.16) used in this paper. This discussion is non-rigorous and only at the level of formal derivations, and is not used in our proofs.

D.1 General background

Let \mathcal{P} be the space of probability density functions on \mathcal{X} with finite second moment. For a smooth curve $\{\rho_t\}_{t\in[0,1]}$ in \mathcal{P} , we identify its tangent vector $\dot{\rho}_t = \partial_t \rho_t(\cdot)$ at ρ_t as a function $\dot{\rho}_t : \mathcal{X} \to \mathbb{R}$. We equip \mathcal{P} with the formal structure of a Riemannian manifold, associating to each $\rho \in \mathcal{P}$ a tangent space $\operatorname{Tan}_{\rho} \mathcal{P}$ of such tangent vectors, and a metric tensor $g_{\rho}(\cdot, \cdot) : \operatorname{Tan}_{\rho} \mathcal{P} \times \operatorname{Tan}_{\rho} \mathcal{P} \to \mathbb{R}$. This induces a metric, for any $\mu, \nu \in \mathcal{P}$,

$$\mathsf{d}(\mu,\nu) = \inf\left\{\int_{0}^{1} \|\dot{\rho}_{t}\|_{\rho_{t}} : \rho_{0} = \mu, \rho_{1} = \nu\right\}$$
(D.1)

where the infimum is over all smooth curves $\{\rho_t\}_{t\in[0,1]}$ in \mathcal{P} from μ to ν , and $\|\zeta\|_{\rho} = \sqrt{g_{\rho}(\zeta,\zeta)}$ for any $\zeta \in \operatorname{Tan}_{\rho} \mathcal{P}$. Given a functional $F: \mathcal{P} \to \mathbb{R}$, the gradient of F at $\rho \in \mathcal{P}$, denoted by grad $F[\rho]$, is the unique element in $\operatorname{Tan}_{\rho} \mathcal{P}$ such that, for all curves $\{\rho_t\}_{t\in[0,1]}$ with $\rho_0 = \rho$ and $\dot{\rho}_0 = \zeta \in \operatorname{Tan}_{\rho} \mathcal{P}$,

$$g_{\rho}\left(\operatorname{grad} F[\rho], \zeta\right) = \frac{\mathrm{d}}{\mathrm{d}t}\Big|_{t=0} F(\rho_t) = \int \delta F[\rho](x)\zeta(x)\mathrm{d}x \tag{D.2}$$

where $\delta F[\rho] : \mathbb{R}^d \to \mathbb{R}$ is the first variation of F. We may then define the gradient flow $\frac{\mathrm{d}}{\mathrm{d}t}\rho_t = -\operatorname{grad} F[\rho_t]$.

For any $\mu, \nu \in \mathcal{P}$ if the infimum (D.1) is achieved by a curve $\{\rho_t\}_{t\in[0,1]}$, then $\{\rho_t\}$ is a geodesic from μ to ν . We say the functional F is geodesically convex at ρ if for all geodesics $\{\rho_t\}_{t\in[0,1]}$ with $\rho_0 = \rho$, we have $\frac{d^2}{dt^2}|_{t=0}\mathcal{F}(\rho_t) \geq 0$.

We now specialize to the Wasserstein-2 and Fisher-Rao geometries, following closely the presentation in [YWR23].

Wasserstein-2 geometry. For the formal Riemannian structure underlying the Wasserstein-2 geometry on $\mathcal{P}_*(\mathbb{R}^p)$, the tangent space at $\rho \in \mathcal{P}_*(\mathbb{R}^p)$ is

$$\operatorname{Tan}_{\rho}^{W_2} \mathcal{P}_*(\mathbb{R}^p) = \Big\{ \zeta : \mathbb{R}^p \to \mathbb{R} \ \Big| \ \zeta = -\nabla \cdot (\rho \nabla u) \text{ for some } u \text{ satisfying } \int \|\nabla u\|^2 \mathrm{d}\rho < \infty \Big\},$$

equipped with the metric tensor $g_{\rho}^{W_2}(\zeta_1, \zeta_2) = \int \langle \nabla u_1, \nabla u_2 \rangle \rho(x) dx$. To compute the Wasserstein-2 gradient, let $\{\rho_t\}_{t \in [0,1]}$ be a curve with $\rho_0 = \rho$ and $\dot{\rho}_0 = \zeta = -\nabla \cdot (\rho \nabla v)$. Then (D.2) reads

$$g_{\rho}^{W_2}\big(\operatorname{grad}^{W_2} F[\rho],\zeta\big) = \int \delta F[\rho] \cdot \zeta \,\mathrm{d}x = \int -\delta F[\rho] \Big(\nabla \cdot (\rho \nabla v)\Big) \mathrm{d}x = \int \langle \nabla \delta F[\rho], \nabla v \rangle \,\rho \,\mathrm{d}x,$$

the last equality applying integration by parts. Comparing with the definition of $g_{\rho}^{W_2}$, this identifies the Wasserstein-2 gradient as

$$\operatorname{grad}^{W_2} F[\rho] = -\nabla \cdot (\rho \nabla \delta F[\rho]). \tag{D.3}$$

Fisher-Rao geometry. For the formal Riemannian structure underlying the Fisher-Rao geometry on $\mathcal{P}_*(M)$, the tangent space at $\rho \in \mathcal{P}_*(M)$ is

$$\operatorname{Tan}_{\rho}^{\operatorname{FR}} \mathcal{P} = \Big\{ \zeta : [-M, M] \to \mathbb{R} \ \Big| \ \zeta = \rho \Big(\alpha - \int_{-M}^{M} \alpha \mathrm{d}\rho \Big) \text{ for some } \alpha \text{ satisfying } \int_{-M}^{M} \alpha^2 \mathrm{d}\rho < \infty \Big\},$$

equipped with the metric tensor $g_{\rho}^{\text{FR}}(\zeta_1, \zeta_2) = \int \frac{\zeta_1 \cdot \zeta_2}{\rho^2} d\rho = \int (\alpha_1 - \int \alpha_1 d\rho)(\alpha_2 - \int \alpha_2 d\rho) d\rho$. To compute the Fisher-Rao gradient, let $\{\rho_t\}_{t\in[0,1]}$ be a curve with $\rho_0 = \rho$ and $\dot{\rho}_0 = \zeta = \rho(\alpha - \int \alpha d\rho)$. Then (D.2) reads

$$\begin{split} g_{\rho}^{\mathrm{FR}}(\mathrm{grad}^{\mathrm{FR}} F[\rho], \zeta) &= \int_{-M}^{M} \delta F[\rho] \cdot \zeta \, \mathrm{d}x = \int_{-M}^{M} \delta F[\rho] \cdot \left(\alpha - \int_{-M}^{M} \alpha \mathrm{d}\rho\right) \mathrm{d}\rho \\ &= \int_{-M}^{M} \left(\delta F[\rho] - \int_{-M}^{M} \delta F[\rho] \mathrm{d}\rho\right) \cdot \left(\alpha - \int_{-M}^{M} \alpha \mathrm{d}\rho\right) \mathrm{d}\rho. \end{split}$$

Comparing with the definition of g_{ρ}^{FR} , this identifies the Fisher-Rao gradient as

$$\operatorname{grad}^{\operatorname{FR}} F[\rho] = \rho \Big(\delta F[\rho] - \int \delta F[\rho] \mathrm{d}\rho \Big). \tag{D.4}$$
D.2 Derivatives of $\overline{F}_n(g)$ and $F_n(q,g)$

From the form (1.4) of $\overline{F}_n(g)$, for any positive density $g \in \mathcal{P}_*(M)$ and bounded $\chi : [-M, M] \to \mathbb{R}$ with $\int_{-M}^M \chi(\theta) d\theta = 0$, we have

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}\varepsilon}\Big|_{\varepsilon=0}\bar{F}_n(g+\varepsilon\chi) &= -\frac{1}{p}\sum_{j=1}^p \frac{\int_{[-M,M]^p} \exp(-\frac{\|\mathbf{y}-\mathbf{X}\boldsymbol{\theta}\|^2}{2\sigma^2})\frac{\chi(\theta_j)}{g(\theta_j)}\prod_{i=1}^p g(\theta_i)\mathrm{d}\boldsymbol{\theta}}{\int_{[-M,M]^p} \exp(-\frac{\|\mathbf{y}-\mathbf{X}\boldsymbol{\theta}\|^2}{2\sigma^2})\prod_{i=1}^p g(\theta_i)\mathrm{d}\boldsymbol{\theta}} \\ &= -\frac{1}{p}\sum_{j=1}^p \int_{-M}^M \frac{\chi(\theta_j)}{g(\theta_j)}P_g(\boldsymbol{\theta} \mid \mathbf{y}) = \int_{-M}^M -\frac{\chi(\theta)}{g(\theta)}\bar{\mu}[g](\theta)\mathrm{d}\theta = \int_{-M}^M -\frac{\bar{\mu}[g](\theta)}{g(\theta)}\chi(\theta)\mathrm{d}\theta, \end{aligned}$$

where $\bar{\mu}[g]$ is the average marginal density of $\mu[g](\boldsymbol{\theta}) = P_g(\boldsymbol{\theta} \mid \mathbf{y})$ as defined in (2.7). Similarly, from the equivalent form (2.10), we have

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}\varepsilon}\Big|_{\varepsilon=0}\bar{F}_n(g+\varepsilon\chi) &= -\frac{1}{p}\sum_{j=1}^p \int \frac{[\mathcal{N}_\tau * \chi](\varphi_j)}{[\mathcal{N}_\tau * g](\varphi_j)} P_g(\varphi \mid \mathbf{y}) \\ &= \int -\frac{[\mathcal{N}_\tau * \chi](\varphi)}{[\mathcal{N}_\tau * g](\varphi)} \bar{\nu}[g](\varphi) \mathrm{d}\varphi = \int_{-M}^M - \left[\mathcal{N}_\tau * \frac{\bar{\nu}[g]}{\mathcal{N}_\tau * g}\right](\theta)\chi(\theta) \mathrm{d}\theta, \end{aligned}$$

where the last equality uses the identity $\int f(\varphi) [\mathcal{N}_{\tau} * \chi](\varphi) d\varphi = \int_{-M}^{M} [\mathcal{N}_{\tau} * f](\theta) \chi(\theta) d\theta$. Thus we identify two equivalent forms for the first variation of $\bar{F}_n(g)$,

$$\delta \bar{F}_n[g](\theta) = -\frac{\bar{\nu}[g](\theta)}{g(\theta)} + 1 = -\mathcal{N}_\tau * \frac{\bar{\nu}[g]}{\mathcal{N}_\tau * g}(\theta) + 1,$$
(D.5)

where the constant +1 is chosen to center $\delta \bar{F}_n[g]$ to have mean 0 under g. The second form shows (3.12), as well as (2.3) when specialized to $\mathbf{X} = \text{Id}$ with $\sigma^2 = \tau^2$ and $\bar{\nu}[g] = n^{-1} \sum_{i=1}^n \delta_{y_i}$. The first form shows (2.6), in light of the general definition (D.4) for the Fisher-Rao gradient. More generally, for $g \in \mathcal{P}(M)$, the last expression of (D.5) still holds and equals to

$$\delta \bar{F}_n[g](\theta) = -\frac{1}{p} \sum_{j=1}^p \frac{\int_{[-M,M]^{p-1}} \exp(-\frac{\|\mathbf{y} - \mathbf{X}(\theta, \theta_{-j})\|^2}{2\sigma^2}) \prod_{i:i \neq j} g(\mathrm{d}\theta_i)}{\int_{[-M,M]^p} \exp(-\frac{\|\mathbf{y} - \mathbf{X}\theta\|^2}{2\sigma^2}) \prod_{i=1}^p g(\mathrm{d}\theta_i)} + 1,$$

which satisfies $\frac{\mathrm{d}}{\mathrm{d}\varepsilon}\Big|_{\varepsilon=0} \bar{F}_n(g+\varepsilon\chi) = \int_{[-M,M]} \delta \bar{F}_n[g](\theta)\chi(\theta)\mathrm{d}\theta$ for any χ on [-M,M] such that $g+\varepsilon\chi$ remains a valid distribution for ε in a neighborhood of 0.

Similarly, for the functional $F_n(q, g)$ in (2.11), we have

$$\frac{\mathrm{d}}{\mathrm{d}\varepsilon}\Big|_{\varepsilon=0}F_n(q,g+\varepsilon\chi) = -\frac{1}{p}\sum_{j=1}^p \int \frac{[\mathcal{N}_\tau *\chi](\varphi_j)}{[\mathcal{N}_\tau *g](\varphi_j)}q(\varphi)\mathrm{d}\varphi = \int_{-M}^M -\left[\mathcal{N}_\tau *\frac{\bar{q}}{\mathcal{N}_\tau *g}\right](\theta)\chi(\theta)\mathrm{d}\theta,$$
$$\frac{\mathrm{d}}{\mathrm{d}\varepsilon}\Big|_{\varepsilon=0}F_n(q+\varepsilon\chi,g) = \frac{1}{p}\int \left[U_g(\varphi) + \log q(\varphi) + 1\right]\chi(\varphi)\mathrm{d}\varphi,$$

where $U_g(\varphi)$ is the negative log-posterior-density (3.5). Then, applying $\int_{-M}^{M} \chi(\theta) d\theta = 0$ and $\int \chi(\varphi) d\varphi = 0$, the first variations of F_n are

$$\delta_g F_n[q,g](\theta) = -\mathcal{N}_\tau * \frac{\bar{q}}{\mathcal{N}_\tau * g}(\theta) + 1, \qquad \delta_q F_n[q,g](\varphi) = \frac{1}{p} \big[U_g(\varphi) + \log q(\varphi) \big],$$

where, for notational convenience, we have centered this first quantity $\delta_g F_n[q,g](\theta)$ to have mean 0 under g. This leads to the forms (2.15–2.16) for the gradient flows (1.7), in light of the general definitions (D.3) and (D.4) for the Wasserstein-2 and Fisher-Rao gradients.

D.3 Non-convexity

We provide a concrete example to show that even for the identity design $\mathbf{X} = \text{Id}$, the functional $\bar{F}_n(g)$ is not geodesically convex under either the Wasserstein-2 geometry or the Fisher-Rao geometry, even though it is convex in the sense (3.4) in the linear geometry.

For the Wasserstein-2 geometry, consider $g_0 = g_* = \delta_0$, $g_1 = \frac{1}{2}\delta_{-1} + \frac{1}{2}\delta_1$, and $y_1, \ldots, y_n \stackrel{iid}{\sim} \mathcal{N}(0,1) * g_*$. The constant-speed Wasserstein-2 geodesic connecting g_0 and g_1 is $g_t = \frac{1}{2}\delta_{-t} + \frac{1}{2}\delta_t$ for $t \in [0,1]$. Explicit computation shows

$$\frac{\mathrm{d}^2}{\mathrm{d}t^2}\bar{F}_n(g_t) = 1 - \frac{1}{n}\sum_{i=1}^n \frac{y_i^2}{\cosh^2(ty_i)},$$

yielding that $\mathbb{P}_{g_*}(\frac{\mathrm{d}^2}{\mathrm{d}^2 t}|_{t=0}\bar{F}_n(g_t) \ge 0) = \mathbb{P}_{g_*}(n^{-1}\sum_{i=1}^n Y_i^2 \le 1) \approx 1/2$ for large n. For the Fisher-Rao geometry, let $g_* = \operatorname{Bernoulli}(1/2), g_0 = \operatorname{Bernoulli}(1/4), g_1 = \operatorname{Bernoulli}(3/4),$

For the Fisher-Rao geometry, let $g_* = \text{Bernoull}(1/2), g_0 = \text{Bernoull}(1/4), g_1 = \text{Bernoull}(3/4),$ and $y_1, \ldots, y_n \stackrel{iid}{\sim} \mathcal{N}(0, 1) * g_*$. Using the fact [MMC23] that the geodesic distance in (D.1) is $d_{\text{FR}}(\text{Bernoulli}(\theta_1), \text{Bernoulli}(\theta_2)) = 2| \operatorname{arcsin}(\theta_1) - \operatorname{arcsin}(\theta_2)|$ for $\theta_1, \theta_2 \in (0, 1)$, we have $d_{\text{FR}}(g_0, g_1) = \pi/3$. The constant-speed Fisher-Rao geodesic connecting g_0 and g_1 is $g_t = \text{Bernoulli}(p_t)$ with $p_t = \sin^2(\frac{\pi}{6}(t+1))$, from equating $d_{\text{FR}}(g_0, g_t) = t \cdot \pi/3$. Explicit computation shows that

$$\mathbb{E}_{g_*}[\bar{F}_n(g_t)] = -\frac{1}{2} \int \left(\mathcal{N}(x) + \mathcal{N}(x-1) \right) \cdot \log \left[\cos^2 \left(\frac{\pi}{6} (t+1) \right) \mathcal{N}(x) + \sin^2 \left(\frac{\pi}{6} (t+1) \right) \mathcal{N}(x-1) \right] \mathrm{d}x$$

is not convex in t, where $\mathcal{N}(x)$ is the standard $\mathcal{N}(0,1)$ density at x.

E Details of algorithms

CAVI. We consider the Gibbs variational representation

$$\bar{F}_n(g) = \min_q \tilde{F}_n(q,g) := \min_q \frac{1}{p} \int_{[-M,M]^p} \left[\frac{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}{2\sigma^2} - \sum_{j=1}^p \log g(\theta_j) + \log q(\boldsymbol{\theta}) \right] q(\boldsymbol{\theta}) \mathrm{d}\boldsymbol{\theta}$$

in the parametrization by $\boldsymbol{\theta}$; in other words, $\widetilde{F}_n(q,g)$ coincides with $F_n(q,g)$ in (2.11) for $\tau = 0$. Mean-field variational inference optimizes $\widetilde{F}_n(q,g)$ over product distributions $q(\boldsymbol{\theta}) = \prod_{j=1}^p q_j(\theta_j)$. It is easy to check that, for fixed g and $q_{-j} = (q_1, \ldots, q_{j-1}, q_{j+1}, \ldots, q_p)$, the minimizer over q_j has the explicit form

$$q_j(\theta_j) \propto \exp\left(-\frac{\|\mathbf{x}_j\|^2}{2\sigma^2}\theta_j^2 + \frac{1}{\sigma^2}(\mathbf{y} - \mathbf{X}_{-j}\mathbb{E}_{q-j}[\boldsymbol{\theta}_{-j}])^\top \mathbf{x}_j \cdot \theta_j\right) g(\theta_j)$$
(E.1)

where $\mathbf{X}_{-j} \in \mathbb{R}^{n \times (p-1)}$ is \mathbf{X} without the *j*-th column, and $\boldsymbol{\theta}_{-j} \in \mathbb{R}^{p-1}$ is $\boldsymbol{\theta}$ without the *j*-th coordinate. Over a discrete support $b_1, \ldots, b_K \in [-M, M]$, we represent the prior *g* by probability weights $w = (w_1, \ldots, w_K)$ and each q_j by weights $w_j = (w_{j,1}, \ldots, w_{j,K})$. Then w_j is updated as

$$w_{j,k} = \frac{w_k \exp\left(-\frac{\|\mathbf{x}_j\|^2}{2\sigma^2} b_k^2 + \frac{1}{\sigma^2} (\mathbf{y} - \mathbf{X}_{-j} \mathbb{E}_{q_{-j}}[\boldsymbol{\theta}_{-j}])^\top \mathbf{x}_j \cdot b_k\right)}{\sum_{i=1}^K w_i \exp\left(-\frac{\|\mathbf{x}_j\|^2}{2\sigma^2} b_i^2 + \frac{1}{\sigma^2} (\mathbf{y} - \mathbf{X}_{-j} \mathbb{E}_{q_{-j}}[\boldsymbol{\theta}_{-j}])^\top \mathbf{x}_j \cdot b_i\right)} \quad \text{for all } k = 1, \dots, K, \quad (E.2)$$

where $\mathbb{E}_{q_{-j}}[\boldsymbol{\theta}_{-j}] = (\sum_{k=1}^{K} b_k w_{j',k})_{j' \neq j}.$

Fixing $q = \prod_{j=1}^{p} q_j$, we update g to minimize

$$\widetilde{F}_{n}(q,g) + \frac{\lambda}{2} \int_{-M}^{M} g''(\theta)^{2} \mathrm{d}\theta = \int_{-M}^{M} [-\log g(\theta)] \bar{q}(\theta) \mathrm{d}\theta + \frac{\lambda}{2} \int_{-M}^{M} g''(\theta)^{2} \mathrm{d}\theta + \text{constant}$$
(E.3)

where $\bar{q} = p^{-1} \sum_{j} q_{j}$. This is implemented by the weight update

$$w = \arg \min_{w \ge 0, \|w\|_{1} = 1} - \sum_{k=1}^{K} \frac{\sum_{j=1}^{p} w_{j,k}}{p} \log w_{k} + \frac{\lambda \Delta}{2} \sum_{k=1}^{K} \left[\frac{Dw}{\Delta} \right]_{k}^{2}$$

where D is the differential quadrature matrix described following (2.25). When $\lambda = 0$, this takes the closed form $w_k = \frac{1}{p} \sum_{j=1}^{p} w_{j,k}$; when $\lambda > 0$, we solve this convex program using CVXR [FNB20]. Each iteration of CAVI iteratively updates (q_1, \ldots, q_p, g) once in this order.

We initialize CAVI from $q_1 = \ldots = q_p = g = \text{Uniform}(\{b_1, \ldots, b_K\}).$

Gibbs-MCEM. Under the prior $g(\theta)$, the conditional posterior density of θ_j given θ_{-j} is

$$P_g(\theta_j \mid \mathbf{y}, \boldsymbol{\theta}_{-j}) \propto \exp\left(-\frac{\|\mathbf{x}_j\|^2}{2\sigma^2}\theta_j^2 + \frac{1}{\sigma^2}(\mathbf{y} - \mathbf{X}_{-j}\boldsymbol{\theta}_{-j})^\top \mathbf{x}_j \cdot \theta_j\right) g(\theta_j).$$
(E.4)

Over a discrete support $b_1, \ldots, b_K \in [-M, M]$, this is represented by the weight vector $w_j = (w_{j,1}, \ldots, w_{j,K})$ with weights

$$w_{j,k} = \frac{w_k \exp\left(-\frac{\|\mathbf{x}_j\|^2}{2\sigma^2} b_k^2 + \frac{1}{\sigma^2} (\mathbf{y} - \mathbf{X}_{-j} \boldsymbol{\theta}_{-j})^\top \mathbf{x}_j \cdot b_k\right)}{\sum_{i=1}^K w_i \exp\left(-\frac{\|\mathbf{x}_j\|^2}{2\sigma^2} b_i^2 + \frac{1}{\sigma^2} (\mathbf{y} - \mathbf{X}_{-j} \boldsymbol{\theta}_{-j})^\top \mathbf{x}_j \cdot b_i\right)} \quad \text{for all } k = 1, \dots, K$$
(E.5)

where $w = (w_1, \ldots, w_k)$ are weights for the prior.

Gibbs-MCEM iteratively resamples the coordinates $\theta_1, \ldots, \theta_p$ of $\boldsymbol{\theta}$ in sequence from the conditional law (E.5) over the support points b_1, \ldots, b_K , fixing the priors weights $w = (w_1, \ldots, w_K)$. Having drawn T_{iter} samples $\theta_j^{(1)}, \ldots, \theta_j^{(T_{\text{iter}})}$ of each *j*-th coordinate, it then approximates \bar{q} in (E.3) by the empirical distribution of these samples, yielding the update for prior weights

$$w = \arg \min_{w \ge 0, \|w\|_{1} = 1} - \frac{\sum_{j=1}^{p} \sum_{t=1}^{T_{\text{iter}}} \sum_{k=1}^{K} 1\{\theta_{j}^{(t)} = b_{k}\}}{pT_{\text{iter}}} \log w_{k} + \frac{\lambda \Delta}{2} \sum_{k=1}^{K} \left[\frac{Dw}{\Delta}\right]_{k}^{2}$$

We initialize Gibbs-MCEM from $\theta = 0$ and $g = \text{Uniform}(\{b_1, \ldots, b_K\})$, and run the first 200 sampling iterations without updating prior weights.

Langevin-MCEM. Langevin-MCEM applies the ULA iterations (4.3) to sample from $P_g(\varphi \mid \mathbf{y})$, fixing the prior weights $w = w_t$ between every T_{iter} iterations. After every T_{iter} iterations, denoting by $\varphi^{(1)}, \ldots, \varphi^{(T_{\text{iter}})}$ the sampled iterates, we sample S = 10000 coordinates $\varphi_1, \ldots, \varphi_S$ uniformly at random from the pT_{iter} total coordinates of $\varphi^{(1)}, \ldots, \varphi^{(T_{\text{iter}})}$ (for reasons of computational efficiency), and approximate the exact M-step update (2.22) in EM by the prior weight update

$$w = \arg \min_{w \ge 0, \|w\|_1 = 1} -\frac{1}{S} \sum_{s=1}^{S} \log \left(\sum_{k=1}^{K} w_k \mathcal{N}_\tau(b_k - \varphi_s) \right) + \frac{\lambda \Delta}{2} \sum_{k=1}^{K} \left[\frac{Dw}{\Delta} \right]_k^2$$

In other words, this is a discretized NPMLE with sample size S with a smoothness penalty. This convex program is solved using CVXR.

We initialize Langevin-MCEM from $\varphi = 0$ and $q = \text{Uniform}(\{b_1, \ldots, b_K\})$, and run the first 200 sampling iterations at large step size $\eta_t^{\varphi} = 1.0$ without updating prior weights, before transitioning to the fixed step size $\eta_t^{\varphi} \in \{0.1, 1.0\}$ with prior weight updates.

EBflow. EBflow alternately applies the ULA iteration (4.3) and the *w* iteration (2.26) with spline smoothing, or (4.5) and (2.26) in the preconditioned setting. We initialize EBflow also from $\varphi = 0$ and $g = \text{Uniform}(\{b_1, \ldots, b_K\})$, and again run the first 200 sampling iterations at large step size $\eta_t^{\varphi} = 1.0$ without updating prior weights, before transitioning to the log-linear step size decay schedule (4.4).

References

- [AC21] Kwangjun Ahn and Sinho Chewi, *Efficient constrained sampling via the mirror-Langevin algorithm*, Advances in Neural Information Processing Systems **34** (2021), 28405–28418.
- [ACG⁺23] O Deniz Akyildiz, Francesca Romana Crucinio, Mark Girolami, Tim Johnston, and Sotirios Sabanis, Interacting particle Langevin algorithm for maximum marginal likelihood estimation, arXiv preprint arXiv:2303.13429 (2023).
- [AF03] Robert A Adams and John JF Fournier, *Sobolev spaces*, 2nd ed., Elsevier, 2003.
- [AGS08] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré, *Gradient flows in metric spaces and in the space of probability measures*, 2nd ed., Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2008.
- [BB19] Roland Bauerschmidt and Thierry Bodineau, A very simple proof of the LSI for high temperature spin systems, J. Funct. Anal. **276** (2019), no. 8, 2582–2588.
- [BBB⁺03] JM Bernardo, MJ Bayarri, JO Berger, AP Dawid, D Heckerman, AFM Smith, and M West, The variational Bayesian EM algorithm for incomplete data: with application to scoring graphical model structures, Bayesian Statistics 7 (2003), no. 453-464, 210.
- [BCE+22] Krishna Balasubramanian, Sinho Chewi, Murat A Erdogdu, Adil Salim, and Shunshi Zhang, Towards a theory of non-log-concave sampling: first-order stationarity guarantees for Langevin Monte Carlo, Conference on Learning Theory, PMLR, 2022, pp. 2896–2923.
- [BCPS21] Jean Barbier, Wei-Kuo Chen, Dmitry Panchenko, and Manuel Sáenz, *Performance of Bayesian linear regression in a model with mismatch*, arXiv preprint arXiv:2107.06936 (2021).
- [BE85] D. Bakry and Michel Émery, *Diffusions hypercontractives*, Séminaire de probabilités, XIX, 1983/84, Lecture Notes in Math., vol. 1123, Springer, Berlin, 1985, pp. 177–206.
- [BEL18] Sébastien Bubeck, Ronen Eldan, and Joseph Lehec, Sampling from a log-concave distribution with projected Langevin Monte Carlo, Discrete Comput. Geom. **59** (2018), no. 4, 757–783.
- [BG99] S. G. Bobkov and F. Götze, Exponential integrability and transportation cost related to logarithmic Sobolev inequalities, J. Funct. Anal. 163 (1999), no. 1, 1–28.
- [BGMZ18] Jean-Baptiste Bardet, Nathaël Gozlan, Florent Malrieu, and Pierre-André Zitt, Functional inequalities for Gaussian convolutions of compactly supported measures: explicit bounds and dimension dependence.
- [BH99] James G Booth and James P Hobert, Maximizing generalized linear mixed model likelihoods with an automated Monte Carlo EM algorithm, Journal of the Royal Statistical Society Series B: Statistical Methodology 61 (1999), no. 1, 265–285.
- [Bil95] Patrick Billingsley, *Probability and measure*, 3rd ed., Wiley Series in Probability and Mathematical Statistics, John Wiley & Sons, Inc., New York, 1995, A Wiley-Interscience Publication.

- [BKRS22] Vladimir I Bogachev, Nicolai V Krylov, Michael Röckner, and Stanislav V Shaposhnikov, Fokker-Planck-Kolmogorov Equations, vol. 207, American Mathematical Society, 2022.
- [BLM13] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart, *Concentration inequalities: A nonasymptotic theory of independence*, Oxford University Press, Oxford, 2013.
- [Böh95] Dankmar Böhning, A review of reliable maximum likelihood algorithms for semiparametric mixture models, vol. 47, 1995, Statistical modelling (Leuven, 1993), pp. 5–28.
- [BRS11] VI Bogachev, M Röckner, and SV Shaposhnikov, On uniqueness problems related to the Fokker– Planck–Kolmogorov equation for measures, J. Math. Sci.(New York) 179 (2011), no. 1, 7–47.
- [Cas85] George Casella, An introduction to empirical Bayes data analysis, Amer. Statist. **39** (1985), no. 2, 83–87.
- [CB18] Lenaic Chizat and Francis Bach, On the global convergence of gradient descent for overparameterized models using optimal transport, Advances in neural information processing systems **31** (2018).
- [CCBJ18] Xiang Cheng, Niladri S Chatterji, Peter L Bartlett, and Michael I Jordan, Underdamped Langevin MCMC: A non-asymptotic analysis, Conference on learning theory, PMLR, 2018, pp. 300–323.
- [Che17] Jiahua Chen, Consistency of the MLE under mixture models, Statist. Sci. **32** (2017), no. 1, 47–63.
- [Chi22] Lénaïc Chizat, Mean-field Langevin dynamics: Exponential convergence and annealing, arXiv preprint arXiv:2202.01009 (2022).
- [CL95] KS Chan and Johannes Ledolter, Monte Carlo EM estimation for time series models involving counts, Journal of the American Statistical Association 90 (1995), no. 429, 242–252.
- [CL96] Bradley P. Carlin and Thomas A. Louis, *Bayes and empirical Bayes methods for data analysis*, Monographs on Statistics and Applied Probability, vol. 69, Chapman & Hall, London, 1996.
- [Cov84] Thomas M. Cover, An algorithm for maximizing expected log investment return, IEEE Trans. Inform. Theory **30** (1984), no. 2, 369–373.
- [CPSV18a] Lénaïc Chizat, Gabriel Peyré, Bernhard Schmitzer, and François-Xavier Vialard, An interpolating distance between optimal transport and Fisher-Rao metrics, Found. Comput. Math. 18 (2018), no. 1, 1–44.
- [CPSV18b] Lénaïc Chizat, Gabriel Peyré, Bernhard Schmitzer, and François-Xavier Vialard, Unbalanced optimal transport: Dynamic and Kantorovich formulations, Journal of Functional Analysis 274 (2018), no. 11, 3090–3123.
- [Dal17] Arnak S. Dalalyan, Theoretical guarantees for approximate sampling from smooth and logconcave densities, J. R. Stat. Soc. Ser. B. Stat. Methodol. 79 (2017), no. 3, 651–676.
- [DEP23] Carles Domingo-Enrich and Aram-Alexandre Pooladian, An explicit expansion of the Kullback-Leibler divergence along its Fisher-Rao gradient flow, arXiv preprint arXiv:2302.12229 (2023).
- [DLR77] A. P. Dempster, N. M. Laird, and D. B. Rubin, Maximum likelihood from incomplete data via the EM algorithm, J. Roy. Statist. Soc. Ser. B 39 (1977), no. 1, 1–38, With discussion.
- [DM17] Alain Durmus and Éric Moulines, Nonasymptotic convergence analysis for the unadjusted Langevin algorithm, Ann. Appl. Probab. **27** (2017), no. 3, 1551–1587.
- [DMM19] Alain Durmus, Szymon Majewski, and Błażej Miasojedow, Analysis of Langevin Monte Carlo via convex optimization, The Journal of Machine Learning Research **20** (2019), no. 1, 2666–2711.
- [DT12] Arnak S Dalalyan and Alexandre B Tsybakov, Sparse regression learning by aggregation and Langevin Monte-Carlo, Journal of Computer and System Sciences **78** (2012), no. 5, 1423–1443.

- [Efr10a] Bradley Efron, Correlated z-values and the accuracy of large-scale statistical estimates, Journal of the American Statistical Association **105** (2010), no. 491, 1042–1055.
- [Efr10b] _____, Large-scale inference, Institute of Mathematical Statistics (IMS) Monographs, vol. 1, Cambridge University Press, Cambridge, 2010, Empirical Bayes methods for estimation, testing, and prediction.
- [Efr21] _____, Empirical bayes: Concepts and methods, 2021.
- [FM03] Gersende Fort and Eric Moulines, Convergence of the Monte Carlo expectation maximization for curved exponential families, The Annals of Statistics 31 (2003), no. 4, 1220–1259.
- [FNB20] Anqi Fu, Balasubramanian Narasimhan, and Stephen Boyd, CVXR: An R package for disciplined convex optimization, Journal of Statistical Software 94 (2020), no. 14, 1–34.
- [GC11] Mark Girolami and Ben Calderhead, Riemann manifold Langevin and Hamiltonian Monte Carlo methods, Journal of the Royal Statistical Society Series B: Statistical Methodology 73 (2011), no. 2, 123–214.
- [GF00] Edward I. George and Dean P. Foster, *Calibration and empirical Bayes variable selection*, Biometrika **87** (2000), no. 4, 731–747.
- [GG20] Ziv Goldfeld and Kristjan Greenewald, Gaussian-smoothed optimal transport: Metric structure and statistical efficiency, International Conference on Artificial Intelligence and Statistics, PMLR, 2020, pp. 3327–3337.
- [GJW08] Piet Groeneboom, Geurt Jongbloed, and Jon A. Wellner, The support reduction algorithm for computing non-parametric function estimates in mixture models, Scand. J. Statist. 35 (2008), no. 3, 385–399.
- [GKKZ23] Alice Guionnet, Justin Ko, Florent Krzakala, and Lenka Zdeborová, *Estimating rank-one matrices with mismatched prior and noise: universality and large deviations*, arXiv preprint arXiv:2306.09283 (2023).
- [GMW14] Whan Ghang, Zane Martin, and Steven Waruhiu, The sharp log-Sobolev inequality on a compact interval, Involve 7 (2014), no. 2, 181–186.
- [GvdV01] Subhashis Ghosal and Aad W. van der Vaart, Entropies and rates of convergence for maximum likelihood and Bayes estimation for mixtures of normal densities, Ann. Statist. 29 (2001), no. 5, 1233–1263.
- [GW00] Christopher R. Genovese and Larry Wasserman, *Rates of convergence for the Gaussian mixture sieve*, Ann. Statist. **28** (2000), no. 4, 1105–1127.
- [Hel09] E. Hellinger, Neue Begründung der Theorie quadratischer Formen von unendlichvielen Veränderlichen, J. Reine Angew. Math. **136** (1909), 210–271.
- [Hot30] Harold Hotelling, Spaces of statistical parameters, Bull. Amer. Math. Soc **36** (1930), 191.
- [Jew82] Nicholas P. Jewell, *Mixtures of exponential distributions*, Ann. Statist. **10** (1982), no. 2, 479–484.
- [JG21] Hansheng Jiang and Adityanand Guntuboyina, A nonparametric maximum likelihood approach to mixture of regression, arXiv preprint arXiv:2108.09816 (2021).
- [JKO98] Richard Jordan, David Kinderlehrer, and Felix Otto, *The variational formulation of the Fokker-Planck equation*, SIAM J. Math. Anal. **29** (1998), no. 1, 1–17.
- [JZ09] Wenhua Jiang and Cun-Hui Zhang, General maximum likelihood empirical Bayes estimation of normal means, Ann. Statist. **37** (2009), no. 4, 1647–1684.
- [Kak48] Shizuo Kakutani, On equivalence of infinite product measures, Ann. of Math. (2) **49** (1948), 214–224.

- [KLJ23] Juan Kuntz, Jen Ning Lim, and Adam M Johansen, Particle algorithms for maximum likelihood training of latent variable models, International Conference on Artificial Intelligence and Statistics, PMLR, 2023, pp. 5134–5180.
- [KM14] Roger Koenker and Ivan Mizera, Convex optimization, shape constraints, compound decisions, and empirical Bayes rules, J. Amer. Statist. Assoc. **109** (2014), no. 506, 674–685.
- [KMV16] Stanislav Kondratyev, Léonard Monsaingeon, and Dmitry Vorotnikov, A new optimal transport distance on the space of finite Radon measures, Adv. Differential Equations 21 (2016), no. 11-12, 1117–1164.
- [KW56] J. Kiefer and J. Wolfowitz, Consistency of the maximum likelihood estimator in the presence of infinitely many incidental parameters, Ann. Math. Statist. 27 (1956), 887–906.
- [Lai78] Nan Laird, Nonparametric maximum likelihood estimation of a mixed distribution, J. Amer. Statist. Assoc. 73 (1978), no. 364, 805–811.
- [LC01] Richard A Levine and George Casella, *Implementations of the Monte Carlo EM algorithm*, Journal of Computational and Graphical Statistics **10** (2001), no. 3, 422–439.
- [LCB⁺22] Marc Lambert, Sinho Chewi, Francis Bach, Silvère Bonnabel, and Philippe Rigollet, Variational inference via Wasserstein gradient flows, Advances in Neural Information Processing Systems 35 (2022), 14434–14447.
- [Led99] Michel Ledoux, Concentration of measure and logarithmic Sobolev inequalities, Séminaire de probabilités de Strasbourg **33** (1999), 120–216 (en).
- [LG16] Jean-François Le Gall, Brownian motion, martingales, and stochastic calculus, french ed., Graduate Texts in Mathematics, vol. 274, Springer, [Cham], 2016.
- [Lin83a] Bruce G. Lindsay, The geometry of mixture likelihoods: a general theory, Ann. Statist. 11 (1983), no. 1, 86–94.
- [Lin83b] Bruce G Lindsay, The geometry of mixture likelihoods, part II: the exponential family, The Annals of Statistics 11 (1983), no. 3, 783–792.
- [Lin95] , Mixture models: Theory, geometry, and applications, Ims, 1995.
- [LK92] Mary L Lesperance and John D Kalbfleisch, An algorithm for computing the nonparametric MLE of a mixing distribution, Journal of the American Statistical Association 87 (1992), no. 417, 120– 126.
- [LLN19] Yulong Lu, Jianfeng Lu, and James Nolen, Accelerating Langevin sampling with birth-death, arXiv preprint arXiv:1905.09863 (2019).
- [LMS16] Matthias Liero, Alexander Mielke, and Giuseppe Savaré, Optimal transport in competition with reaction: the Hellinger-Kantorovich distance and geodesic curves, SIAM J. Math. Anal. 48 (2016), no. 4, 2869–2911.
- [LMS18] _____, Optimal entropy-transport problems and a new Hellinger-Kantorovich distance between positive measures, Invent. Math. **211** (2018), no. 3, 969–1117.
- [LSU68] Olga Aleksandrovna Ladyzhenskaia, Vsevolod Alekseevich Solonnikov, and Nina N Ural'tseva, Linear and quasi-linear equations of parabolic type, vol. 23, American Mathematical Soc., 1968.
- [LSW23] Yulong Lu, Dejan Slepvcev, and Lihan Wang, Birth-death dynamics for sampling: global convergence, approximations and their asymptotics, Nonlinearity 36 (2023), no. 11, 5731.
- [LT84] Diane Lambert and Luke Tierney, Asymptotic properties of maximum likelihood estimates in the mixed Poisson model, Ann. Statist. 12 (1984), no. 4, 1388–1399.
- [LZ07] Lei Liu and Yu Zhu, Partially projected gradient algorithms for computing nonparametric maximum likelihood estimates of mixing distributions, J. Statist. Plann. Inference 137 (2007), no. 7, 2509–2522.

- [MCC⁺21] Yi-An Ma, Niladri S Chatterji, Xiang Cheng, Nicolas Flammarion, Peter L Bartlett, and Michael I Jordan, *Is there an analog of Nesterov acceleration for gradient-based MCMC?*
- [MCF15] Yi-An Ma, Tianqi Chen, and Emily Fox, A complete recipe for stochastic gradient MCMC, Advances in neural information processing systems **28** (2015).
- [MCW⁺23] Fabio Morgante, Peter Carbonetto, Gao Wang, Yuxin Zou, Abhishek Sarkar, and Matthew Stephens, A flexible empirical Bayes approach to multivariate multiple regression, and its improved accuracy in predicting multi-tissue gene expression from genotypes, PLoS Genetics 19 (2023), no. 7, e1010539.
- [ML89] J. S. Maritz and T. Lwin, *Empirical Bayes methods*, 2nd ed., Monographs on Statistics and Applied Probability, vol. 35, Chapman & Hall, London, 1989.
- [MMC23] Henrique K Miyamoto, Fábio CC Meneghetti, and Sueli IR Costa, On closed-form expressions for the Fisher-Rao distance, arXiv preprint arXiv:2304.14885 (2023).
- [MMN18] Song Mei, Andrea Montanari, and Phan-Minh Nguyen, A mean field view of the landscape of two-layer neural networks, Proc. Natl. Acad. Sci. USA **115** (2018), no. 33, E7665–E7671.
- [MSS23] Sumit Mukherjee, Bodhisattva Sen, and Subhabrata Sen, A mean field approach to empirical bayes estimation in high-dimensional linear regression, arXiv preprint arXiv:2309.16843 (2023).
- [Nea11] Radford M Neal, *MCMC using Hamiltonian dynamics*, Handbook of Markov Chain Monte Carlo **2** (2011), no. 11, 2.
- [Nea13] Ronald C Neath, On convergence properties of the Monte Carlo EM algorithm, Advances in modern statistical theory and applications: a Festschrift in Honor of Morris L. Eaton **10** (2013), 43–63.
- [NH98] Radford M Neal and Geoffrey E Hinton, A view of the EM algorithm that justifies incremental, sparse, and other variants, Learning in graphical models, Springer, 1998, pp. 355–368.
- [NS86] Fassil Nebebe and T. W. F. Stroud, Bayes and empirical Bayes shrinkage estimation of regression coefficients, Canad. J. Statist. 14 (1986), no. 4, 267–280.
- [NS17] Atsushi Nitanda and Taiji Suzuki, Stochastic particle gradient descent for infinite ensembles, arXiv preprint arXiv:1712.05438 (2017).
- [NW22] Richard Nickl and Sven Wang, On polynomial-time computation of high-dimensional posterior measures by Langevin-type algorithms, Journal of the European Mathematical Society (2022).
- [NWS22] Atsushi Nitanda, Denny Wu, and Taiji Suzuki, *Convex analysis of the mean field langevin dynamics*, International Conference on Artificial Intelligence and Statistics, PMLR, 2022, pp. 9741– 9757.
- [NY83] A. S. Nemirovsky and D. B. Yudin, Problem complexity and method efficiency in optimization, Wiley-Interscience Series in Discrete Mathematics, John Wiley & Sons, Inc., New York, 1983, Translated from the Russian and with a preface by E. R. Dawson, A Wiley-Interscience Publication.
- [O'C21] Luke J O'Connor, The distribution of common-variant effect sizes, Nature Genetics 53 (2021), no. 8, 1243–1249.
- [Ott01] Felix Otto, The geometry of dissipative evolution equations: the porous medium equation, Comm. Partial Differential Equations **26** (2001), no. 1-2, 101–174.
- [Per16] Marcelo Pereyra, Proximal Markov Chain Monte Carlo algorithms, Statistics and Computing 26 (2016), 745–760.
- [Pfa88] J. Pfanzagl, Consistency of maximum likelihood estimators for certain nonparametric families, in particular: mixtures, J. Statist. Plann. Inference 19 (1988), no. 2, 137–158.

- [PW20] Yury Polyanskiy and Yihong Wu, Self-regularizing property of nonparametric maximum likelihood estimator in mixture models, arXiv preprint arXiv:2008.08244 (2020).
- [PW21] _____, Sharp regret bounds for empirical bayes and compound decision problems, arXiv preprint arXiv:2109.03943 (2021).
- [Rio09] Emmanuel Rio, Upper bounds for minimal distances in the central limit theorem, Annales de l'IHP Probabilités et statistiques, vol. 45, 2009, pp. 802–817.
- [Rob50] Herbert Robbins, A generalization of the method of maximum likelihood-estimating a mixing distribution, no. 2, 314–315.
- [Rob51] _____, Asymptotically subminimax solutions of compound statistical decision problems, Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability, The Regents of the University of California, 1951.
- [Rob56] _____, An empirical Bayes approach to statistics, Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics, The Regents of the University of California, 1956.
- [RR45] C. Radhakrishna Rao, Information and the accuracy attainable in the estimation of statistical parameters, Bull. Calcutta Math. Soc. 37 (1945), 81–91.
- [RS02] Gareth O Roberts and Osnat Stramer, Langevin diffusions and Metropolis-Hastings algorithms, Methodology and computing in applied probability 4 (2002), 337–357.
- [RT96] Gareth O Roberts and Richard L Tweedie, Exponential convergence of Langevin distributions and their discrete approximations, Bernoulli (1996), 341–363.
- [RV13] Mark Rudelson and Roman Vershynin, Hanson-Wright inequality and sub-gaussian concentration.
- [RVE18] Grant M Rotskoff and Eric Vanden-Eijnden, Neural networks as interacting particle systems: Asymptotic convexity of the loss landscape and universal scaling of the approximation error, Stat 1050 (2018), 22.
- [San17] Filippo Santambrogio, {Euclidean, metric, and Wasserstein} gradient flows: an overview, Bull. Math. Sci. 7 (2017), no. 1, 87–154.
- [Sch10] Armin Schwartzman, Comment on "Correlated z-values and the accuracy of large-scale statistical estimates" by Bradley Efron, Journal of the American Statistical Association 105 (2010), no. 491, 1059.
- [SG20] Sujayam Saha and Adityanand Guntuboyina, On the nonparametric maximum likelihood estimator for Gaussian location mixture densities with application to Gaussian denoising, Ann. Statist. 48 (2020), no. 2, 738–762.
- [SS18] Lei Sun and Matthew Stephens, Solving the empirical bayes normal means problem with correlated noise, arXiv preprint arXiv:1812.07488 (2018).
- [SS20] Justin Sirignano and Konstantinos Spiliopoulos, *Mean field analysis of neural networks: A law of large numbers*, SIAM Journal on Applied Mathematics **80** (2020), no. 2, 725–752.
- [SSAAP22] Jeffrey P Spence, Nasa Sinnott-Armstrong, Themistocles L Assimes, and Jonathan K Pritchard, A flexible modeling and inference framework for estimating variant effect sizes from GWAS summary statistics, BioRxiv (2022), 2022–04.
- [Ver18] Roman Vershynin, *High-dimensional probability: An introduction with applications in data sci*ence, vol. 47, Cambridge university press, 2018.
- [Vil09] Cédric Villani, Optimal transport: Old and new, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 338, Springer-Verlag, Berlin, 2009.

- [VL93] Y. Vardi and D. Lee, From image deblurring to optimal investments: maximum likelihood solutions for positive linear inverse problems, J. Roy. Statist. Soc. Ser. B 55 (1993), no. 3, 569–612, With discussion.
- [VW19] Santosh Vempala and Andre Wibisono, *Rapid convergence of the unadjusted Langevin algorithm: Isoperimetry suffices*, Advances in neural information processing systems **32** (2019).
- [VW23] AW van der Vaart and Jon A Wellner, Weak convergence and empirical processes: With applications to statistics, Springer, 2023.
- [Wan07] Yong Wang, On fast computation of the non-parametric maximum likelihood estimate of a mixing distribution, J. R. Stat. Soc. Ser. B Stat. Methodol. 69 (2007), no. 2, 185–198.
- [Wel05] Jon A Wellner, *Empirical processes: Theory and applications*, Notes for a course given at Delft University of Technology (2005), 17.
- [Wib18] Andre Wibisono, Sampling as optimization in the space of measures: The Langevin dynamics as a composite optimization problem, Conference on Learning Theory, PMLR, 2018, pp. 2093–3027.
- [WT90] Greg CG Wei and Martin A Tanner, A Monte Carlo implementation of the EM algorithm and the poor man's data augmentation algorithms, Journal of the American statistical Association 85 (1990), no. 411, 699–704.
- [WT11] Max Welling and Yee W Teh, Bayesian learning via stochastic gradient Langevin dynamics, Proceedings of the 28th international conference on machine learning (ICML-11), 2011, pp. 681– 688.
- [WZ21] Yihe Wang and Sihai Dave Zhao, A nonparametric empirical Bayes approach to large-scale multivariate regression, Computational Statistics & Data Analysis **156** (2021), 107130.
- [YL05] Ming Yuan and Yi Lin, Efficient empirical Bayes variable selection and estimation in linear models, J. Amer. Statist. Assoc. 100 (2005), no. 472, 1215–1225.
- [YWR23] Yuling Yan, Kaizheng Wang, and Philippe Rigollet, *Learning Gaussian mixtures using the Wasserstein-Fisher-Rao gradient flow*, arXiv preprint arXiv:2301.01766 (2023).
- [YY22] Rentian Yao and Yun Yang, *Mean field variational inference via wasserstein gradient flow*, arXiv preprint arXiv:2207.08074 (2022).
- [ZCST22] Yangjing Zhang, Ying Cui, Bodhisattva Sen, and Kim-Chuan Toh, On efficient and scalable computation of the nonparametric maximum likelihood estimator in mixture models, arXiv preprint arXiv:2208.07514 (2022).
- [Zha03] Cun-Hui Zhang, Compound decision theory and empirical Bayes methods, vol. 31, 2003, Dedicated to the memory of Herbert E. Robbins, pp. 379–390.
- [Zha09] _____, Generalized maximum likelihood estimation of normal mixture densities, Statist. Sinica 19 (2009), no. 3, 1297–1318.
- [Zim16] David Zimmermann, Elementary proof of logarithmic Sobolev inequalities for gaussian convolutions on ℝ, Annales mathématiques Blaise Pascal, vol. 23, 2016, pp. 129–140.
- [ZQM11] Zhengliang Zhang, Bin Qian, and Yutao Ma, Uniform logarithmic Sobolev inequality for Boltzmann measures with exterior magnetic field over spheres, Acta Appl. Math. 116 (2011), no. 3, 305–315.
- [ZQPC18] Yan Zhang, Guanghao Qi, Ju-Hyun Park, and Nilanjan Chatterjee, Estimation of complex effectsize distributions using summary-level statistics from genome-wide association studies across 32 complex traits, Nature Genetics 50 (2018), no. 9, 1318–1326.
- [ZZ14] Cun-Hui Zhang and Stephanie S Zhang, Confidence intervals for low dimensional parameters in high dimensional linear models, Journal of the Royal Statistical Society: Series B: Statistical Methodology (2014), 217–242.
- [ZZ21] Geyu Zhou and Hongyu Zhao, A fast and robust Bayesian nonparametric method for prediction of complex traits using summary statistics, PLoS genetics **17** (2021), no. 7, e1009697.