IRREDUCIBLE CHARACTERS AND BITRACE FOR THE q-ROOK MONOID

NAIHUAN JING, YU WU, AND NING LIU

ABSTRACT. This paper studies irreducible characters of the q-rook monoid algebra $R_n(q)$ using the vertex algebraic method. Based on the Frobenius formula for $R_n(q)$, a new iterative character formula is derived with the help of the vertex operator realization of the Schur symmetric function. The same idea also leads to a simple proof of the Murnaghan-Nakayama rule for $R_n(q)$. We also introduce the bitrace for the q-monoid and obtain a general combinatorial formula for the bitrace, which generalizes the counterpart for the Iwahori-Hecke algebra. The character table of $R_n(q)$ with $|\mu|=5$ is listed in the appendix

CONTENTS

²⁰¹⁰ *Mathematics Subject Classification.* Primary: 20C08; Secondary: 17B69, 05E10.

Key words and phrases. q-rook monoid algebra, vertex operators, Murnaghan-Nakayama rule, bitrace, Schur polynomials.

1. INTRODUCTION

The rook monoid R_n or symmetric inverse monoid is the monoid of partial permutations, i.e. $n \times n$ permutation matrices with some nonzero entries changed to 0. Solomon [\[16\]](#page-21-0) introduced a q -analogue $R_n(q)$ of the group algebra of R_n as an "Iwahori-Hecke algebra" for the monoid of $n \times n$ matrices $M_n(\mathbb{F}_q)$ over the finite field \mathbb{F}_q with q elements. Its "Borel subgroup" of invertible upper triangular matrices, i.e., the q-rook monoid $R_n(q)$ is a semisimple $\mathbb{C}(q)$ -algebra that specializes when $q \to 1$ to $\mathbb{C}[R_n]$. In [\[18\]](#page-21-1), Solomon also obtained a presentation of $R_n(q)$ by generators and relations and gave an action of $R_n(q)$ on $U^{\otimes n}$ where $U = L(\omega_1) \oplus L(0)$ is the $(r + 1)$ -dimensional representation of $U_q(gl_r)$, which is the direct sum of the r-dimensional fundamental representation $L(\omega_1)$ and a trivial representation $L(0)$. Halverson [\[3\]](#page-20-1) gave a different presentation of $R_n(q)$ and showed that as an $(R_n(q), U_q(gl_r))$ -bimodule

$$
U^{\otimes n} \cong \bigoplus_{\lambda \in \hat{R}_n} R_n^{\lambda}(q) \otimes L(\lambda)
$$

summed over all partitions λ with no more than n boxes and r rows. Here $R_n^{\lambda}(q)$ are simple $R_n(q)$ modules and $L(\lambda)$ are simple $U_q(gl_n)$ -modules. A third presentation provided by Halverson and Ram showed that $R_n(q)$ is a quotient of the affine Hecke algebra in type A. In [\[1\]](#page-20-2), the character theory of $R_n(q)$ was developed, which generalized the earlier work of Munn [\[13\]](#page-21-2). [\[1\]](#page-20-2) used a Schur-Weyl duality between $R_n(q)$ and the quantum general linear group $U_q(gl_r)$ to compute a Frobenius formula, in the ring of symmetric functions, for the irreducible characters of $R_n(q)$. A recursive Murnaghan-Nakayama rule for these characters $\chi^{\lambda}_{\mu}(q)$ was also derived in [\[1\]](#page-20-2), which expresses $\chi^{\lambda}_{\mu}(q)$ in terms of those whose lower partitions have shorter length.

The aim of this paper is to find more effective and computation friendly formulae for $\chi^{\lambda}_{\mu}(q)$. We use vertex algebraic techniques to give a new iterative formula for $\chi^{\lambda}_{\mu}(q)$. Different from the Murnaghan-Nakayama rule, our new formula express $\chi^{\lambda}_{\mu}(q)$ in terms of those whose upper partitions have shorter length. Many special cases are considered. In particular, the compact rules for $\chi^{\lambda}_{\mu}(q)$ with λ hook and two-row type are derived respectively. The bitrace can be viewed a deformed orthogonality relation for irreducible characters. We also introduce the bitrace of the q-rook monoid algebra as an analog of the bitrace of the Hecke algebra $[4, 9]$ $[4, 9]$. A parallel combinatorial formula is obtained.

The paper is organized as follows. In the next section, we will review briefly some essential preliminaries about symmetric functions, vertex operator realization of Schur functions and some structures of the q-Rook Monoid algebra. In Sec. 3, we will give our main results: a new recursive formula for $\chi^{\lambda}_{\mu}(q)$ (Theorem [3.3\)](#page-8-0) and the closed formulae for the cases of λ being hook and two-row type (Theorem

[3.10\)](#page-13-0). The Murnaghan-Nakayama rule is also redrived. Sec. 4 deals with the bitrace of the q-Rook Monoid algebra. A general combinatorial formula for it is deduced (Theorem [4.1\)](#page-19-1). In Sec. 5, we list a table of $\chi^{\lambda}_{\mu}(q)$ with $|\mu|=5$ using our formulae.

2. Preliminaries

In this section, first we present some well-known facts about symmetric functions [\[11\]](#page-21-4), which will be used in the subsequent sections. Then we will review briefly the vertex operator realisation of Schur functions and the Frobenius formula for the q-rook monoid algebra.

2.1. Partitions and symmetric functions. A partition λ of $n > 0$ is a sequence of non-negative integers $(\lambda_1, \lambda_2, \ldots)$ such that $\lambda_1 \geq \lambda_2 \geq \ldots$ and $\sum_i \lambda_i = |\lambda| = n$. We will call \emptyset a partition of 0. The nonzero λ_i are called parts of λ , and n is called the weight of λ and denoted by $\lambda \vdash n$. The number of nonzero parts is length $l(\lambda)$. Sometimes λ is arranged in the ascending order: $\lambda = (1^{m_1}2^{m_2} \dots)$ with m_i being the multiplicity of part i in λ . The conjugate λ' of a partition λ is the partition defined by $\lambda'_i = Card\{j : \lambda_j \geqslant i\}$ and $m_i = \lambda'_i - \lambda'_{i+1}$.

A composition τ of n is a sequence of non-negative integers (τ_1, \ldots, τ_l) satisfying $\sum_i \tau_i = n$, denoted by $\tau \models n$. Clearly, a partition is a composition. For any two compositions τ and ρ , we denote $\tau \subset \rho$ if $\tau_i \leqslant \rho_i$ for all $i \geqslant 1$.

A partition λ can be visualized by its Young diagram, which consists of left adjusted array of boxes (also called nodes) with λ_i boxes on the *i*th row. For two partitions λ, μ , the set-theoretic difference $\theta = \lambda/\mu$ is called a *skew diagram*. A skew diagram $\theta = \lambda/\mu$ is a *vertical* (resp. *horizontal*) m-strip if $|\theta| = m$ and $\theta_i = \lambda_i - \mu_i \leq 1$ (resp. $\theta'_i = \lambda'_i - \mu'_i \leq 1$) for each $i \geq 1$. In other words, a vertical (resp. horizontal) strip has at most one square in each row (resp. column).

Let Λ be the ring of symmetric functions in the x_n $(n \in \mathbb{N})$ over Q, and we will also study the ring $Λ_Z$ as a lattice of Λ. The ring Λ has several linear bases indexed by partitions. For each $r > 1$, let $p_r = \sum x_i^r$ be the rth power-sum. Then $p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_l}$, $\lambda \in \mathcal{P}$ form a Q-basis of Λ . We introduce the modified power sum functions $\hat{p}_n = p_n(1, x_1, x_2, \ldots) = 1 + p_n$.

The ring Λ also has various \mathbb{Z} -bases indexed by partitions: the set of monomial symmetric functions $m_{\lambda} = \sum x_{i_1}^{\lambda_1}$ $\frac{\lambda_1}{i_1}\cdots x_{i_k}^{\lambda_k}$ \hat{i}_k^k ; that of elementary symmetric functions $e_{\lambda} = e_{\lambda_1} \cdots e_{\lambda_k}$ with $e_n = m_{(1^n)}$; that of complete symmetric function $h_{\lambda} = h_{\lambda_1} \cdots h_{\lambda_k}$ with $h_n = \sum_{\lambda \vdash n} m_{\lambda}$, and that of Schur symmetric functions s_λ .

The space Λ is equipped with the standard inner product given by:

$$
\langle p_{\lambda}, p_{\mu} \rangle = \delta_{\lambda \mu} z_{\lambda},
$$

where $z_{\lambda} = \prod_{i \geq 1} \lambda_i m_i(\lambda)!$, under which the Schur functions s_{λ} 's form an orthonormal basis of Λ .

The function p_n acts on Λ as a multiplication operator, and its adjoint operator is the differential operator $p_n^* = n \frac{\partial}{\partial p}$ $\frac{\partial}{\partial p_n}$. Note that * is Q-linear and anti-involutive satisfying

(2.2)
$$
\langle p_n u, v \rangle = \langle u, p_n^* v \rangle
$$

for $u, v \in \Lambda$.

Using the degree gradation, Λ becomes a graded ring

(2.3)
$$
\Lambda = \bigoplus_{n=0}^{\infty} \Lambda_n.
$$

A linear operator A is of degree n if $A(\Lambda_m) \subset \Lambda_{m+n}$. Clearly, p_n (resp. p_n^*) is of degree n (resp. $-n$).

2.2. Vertex operator realization of Schur functions. We recall the vertex operator realization of Schur functions. Let $S(z)$ be the map: $\Lambda \longrightarrow \Lambda[[z, z^{-1}]]$ defined by (see [\[8\]](#page-21-5))

(2.4)
$$
S(z) = \exp\left(\sum_{n\geq 1} \frac{1}{n} p_n z^n\right) \exp\left(-\sum_{n\geq 1} \frac{\partial}{\partial p_n} z^{-n}\right) = \sum_{n\in \mathbb{Z}} S_n z^n.
$$

We use the convention to index the components by their degrees. The operators $S_n \in \text{End}(\Lambda)$ are the Bernstein vertex operators realizing the Schur functions.

For any composition $\mu = (\mu_1, \dots, \mu_k)$, the product $S_{\mu_1} \cdots S_{\mu_k}$. $1 = s_{\mu}$ is the Schur function labeled by μ . In general, $s_{\mu} = 0$ or $\pm s_{\lambda}$ for a partition λ such that $\lambda \in \mathfrak{S}_l(\mu+\delta) - \delta$. Here $\delta = (l-1, l-2, \ldots, 1, 0)$ and $l = l(\mu)$. \mathfrak{S}_l is the symmetric group on l letters.

Note that $p_n^* = n \frac{\partial}{\partial p}$ $\frac{\partial}{\partial p_n}$, we have the dual or adjoint operator of $S(z)$:

(2.5)
$$
S^*(z) = \exp\left(-\sum_{n\geqslant 1}^{\infty} \frac{1}{n} p_n z^n\right) \exp\left(\sum_{n\geqslant 1}^{\infty} \frac{\partial}{\partial p_n} z^{-n}\right) = \sum_{n\in\mathbb{Z}} S_n^* z^{-n}.
$$

Moreover, $S_{-n}.1 = \delta_{n,0}, S_n^* . 1 = \delta_{n,0}, (n \ge 0).$

The special Schur function $h_n = S_n.1$ is called the complete homogeneous symmetric function, whose generating function is given by

(2.6)
$$
\exp\left(\sum_{n\geq 1} \frac{1}{n} p_n z^n\right) = \prod_{i\geq 1} \frac{1}{1 - x_i z} = \sum_{n\geq 0} h_n z^n.
$$

Similarly, its dual operator can be written as

(2.7)
$$
\exp\left(\sum_{n\geq 1}\frac{\partial}{\partial p_n}z^{-n}\right) = \sum_{n\geq 0}h_n^*z^{-n}.
$$

Introduce the symmetric functions $q_n(t)$ defined by its generating function:

(2.8)
$$
\exp\left(\sum_{n=1}^{\infty}\frac{1-t^n}{n}p_nz^n\right) = \sum_{n=0}^{\infty}q_n(t)z^n = q(z;t)
$$

where $q_n(t)$ is the Hall-Littlewood polynomial of one-row partition (n) , and clearly

(2.9)
$$
q_n(t) = \sum_{\lambda \vdash n} \frac{1}{z_{\lambda}(t)} p_{\lambda}.
$$

Here $z_{\lambda}(t) = \frac{z_{\lambda}}{\prod_{i\geq 1}(1-t^{\lambda_i})}$. For each partition λ , denote $q_{\lambda}(t) = q_{\lambda_1}(t)q_{\lambda_2}(t)\cdots$, then $\{q_{\lambda}(t)\}$ forms a basis of $\Lambda \otimes \mathbb{Q}(t)$.

Recall that $\hat{p}_n = p_n(1, x_1, x_2, \dots)$. We introduce a modification of $q_n(t)$ denoted by $\hat{q}_n(t)$ as follows:

(2.10)
$$
\exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} \hat{p}_n z^n\right) = \sum_{n=0}^{\infty} \hat{q}_n(t) z^n = \hat{q}(z; t).
$$

Similarly, we have $\hat{q}_n(t) = \sum_{\lambda \vdash n} \frac{1}{z_{\lambda}(t)}$ $\frac{1}{z_{\lambda}(t)}\hat{p}_{\lambda}$. For a partition λ , denote $\hat{q}_{\lambda}(t) = \hat{q}_{\lambda_1}(t)\hat{q}_{\lambda_2}(t)\cdots$.

Lemma 2.1. The relation between $\hat{q}_{\lambda}(t)$ and $q_{\lambda}(t)$ is given by

(2.11)
$$
\hat{q}_{\lambda}(t) = \sum_{\tau} (1-t)^{l(\tau)} q_{\lambda-\tau}(t)
$$

summed over all compositions τ such that $\tau \subset \lambda$. Here $\lambda - \tau = (\lambda_1 - \tau_1, \lambda_2 - \tau_2, \cdots)$. In particular, $\hat{q}_n(t) = q_n(t) + (1-t) \sum_{i \geq 1}^n q_{n-i}(t).$

Proof. By definition it follows that

$$
\sum_{n=0}^{\infty} \hat{q}_n(t) z^n = \exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} \hat{p}_n z^n\right)
$$

=
$$
\exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} p_n z^n\right) \exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} z^n\right)
$$

=
$$
\left(\sum_{n=0}^{\infty} q_n(t) z^n\right) \left(1 + (1-t) \sum_{i \geq 1}^{\infty} z^i\right).
$$

Comparing the coefficient of z^n on both sides gives $\hat{q}_n(t) = q_n(t) + (1-t)\sum_{i\geq 1}^n q_{n-i}(t)$, which yields (2.11) by induction.

2.3. Irreducible characters for the q-rook monoid algebra. Let q be an indeterminate. For $n \geq$ 2, the q-rook monoid $R_n(q)$ is defined as the associative $\mathbb{C}(q)$ -algebra with generators $1, T_1, \cdots, T_{n-1}$, P_1, \cdots, P_n subject to the relations [\[3\]](#page-20-1):

$$
(T_i - q)(T_i + 1) = 0, \quad 1 \le i \le n - 1,
$$

$$
T_i T_j = T_j T_i, \quad 1 \le i, j \le n - 1, |i - j| > 1,
$$

$$
T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad 1 \le i \le n-2,
$$

\n
$$
T_i P_j = P_j T_i = q P_j, \quad 1 \le i < j \le n,
$$

\n
$$
T_i P_j = P_j T_i, \quad 1 \le j < i \le n-1,
$$

\n
$$
P_i^2 = P_i, \quad 1 \le i \le n,
$$

\n
$$
P_{i+1} = q P_i T_i^{-1} P_i, \quad 2 \le i \le n.
$$

Define $R_0(q) = \mathbb{C}(q)$, and $R_1(q)$ is the associative $\mathbb{C}(q)$ -algebra spanned by 1 and P_1 subject to $P_1^2 = P_1$. The subalgebra of $R_n(q)$ generated by T_1, \dots, T_{n-1} is isomorphic to the Iwahori-Hecke algebra $H_n(q)$ in type A_{n-1} . Here we adopt Halverson's presentation of $R_n(q)$ for our purposes.

Let $s_i \in \mathfrak{S}_n$ be the transposition of i and $i + 1$, we define $\gamma_1 = T_{\gamma_1} = 1$ and for $2 \le t \le n$,

$$
\gamma_t = s_1 s_2 \cdots s_{i-1},
$$

$$
T_{\gamma_t} = T_1 \cdots T_{t-1}.
$$

For a composition $\mu = (\mu_1, \ldots, \mu_l) \models k, 0 \leq k \leq n$, we define

$$
T_{\gamma_{\mu}} = T_{\gamma_{\mu_1}} \otimes \ldots \otimes T_{\gamma_{\mu_l}},
$$

$$
T_{\mu} = P_{n-k} \otimes T_{\gamma_{\mu}},
$$

where the tensor product means that the factors $T_{\gamma_{\mu_i}}$ are generated by T_j with $j = \sum_{a=1}^{j-1} \mu_a +$ $1, \sum_{a=1}^{j-1} \mu_a + 2, \ldots, \sum_{a=1}^{j} \mu_a$ (disjoint indices). Here $T_{\mu} \in R_{n-k}(q) \otimes R_{\mu_1}(q) \otimes \ldots \otimes R_{\mu_l}(q) \subset R_n(q)$ is called the *standard element*. It is clear that T_{μ} is the identity when $\mu = (1^{n})$.

If $K \subset \{1, 2, \ldots, n\}$ define the subgroup $S_K \subset S_n$ to be the group of permutations on the elements of K. For $1 \leq i \leq n$, we define $T_{i,i} = 1$, and define for $1 \leq i < j \leq n$

$$
(2.12) \t\t T_{i,j} = T_{j-1}T_{j-2}...T_i.
$$

Let $A = \{a_1, a_2, \ldots, a_k\} \subset \{1, 2, \ldots, n\}$ and assume that $a_1 < a_2 < \ldots < a_k$. Define

(2.13)
$$
T_A = T_{1,a_1} T_{2,a_2} \dots T_{k,a_k}.
$$

For $0 \leq k \leq n$, let Ω_k be the following set of triples,

$$
\Omega_k = \left\{ (A, B, \omega) \middle| \begin{array}{c} A, B \subset \{1, 2, \dots, n\}, |A| = |B| = k, \\ \omega \in S_{\{k+1, \dots, n\}} \end{array} \right\}
$$

and let $\Omega = \bigcup_{k=0}^{n} \Omega_k$. If we define the standard words

(2.14)
$$
T_{(A,B,\omega)} = T_A T_{\omega} P_k T_B^{-1}, (A,B,\omega) \in \Omega_k.
$$

Then the set $\{T_{(A,B,\omega)}|(A,B,\omega \in \Omega)\}\)$ forms a linear $\mathbb{C}(q)$ -basis of $R_n(q)$ [\[3\]](#page-20-1).

It is known that irreducible characters of $R_n(q)$ are parameterized by partitions of $k \leq n, k =$ $0, 1, \ldots, n$. Here a partition of 0 means \emptyset . The irreducible module V^{λ} indexed by $\lambda \vdash k \leq n$ have been studied in [\[2,](#page-20-4) [13,](#page-21-2) [17\]](#page-21-6), and the seminormal bases of V^{λ} were found in [\[3\]](#page-20-1) (which will be recalled in the next section). Dieng, Halverson and Poladian [\[1\]](#page-20-2) used the Schur-Weyl duality to prove the following Frobenius-type formula in the ring of symmetric functions $(\mu \vdash n)$

(2.15)
$$
\frac{q^{|\mu|}}{(q-1)^{l(\mu)}}\hat{q}_{\mu}(q^{-1}) = \sum_{k=0}^{n} \sum_{\lambda \vdash k} \chi^{\lambda}_{R_n(q)}(T_{\mu})s_{\lambda}
$$

where $\chi^{\lambda}_{R_n(q)}(T_\mu)$ is the value of the irreducible characters of $R_n(q)$ indexed by $\lambda \vdash k(\leq n)$ at the standard element T_{μ} . In the following we sometimes write $\chi^{\lambda}_{\mu}(q)$ for $\chi^{\lambda}_{R_n(q)}(T_{\mu})$ in short. Formula [\(2.15\)](#page-6-1) is a generalization of the Frobenius formula given by Ram [\[14\]](#page-21-7) for the Iwahori-Hecke algebra $H_n(q)$ in type A. The irreducible $R_n(q)$ -characters are completely determined by their values on the set of standard elements $T_{\mu}, \mu \vdash k, 0 \leq k \leq n$. Specifically, the character table of $R_n(q)$ is block upper triangular with the diagonal blocks being the character tables of the Iwahori-Hecke algebras $H_k(q)$ in type $A, k \leq n$.

By the orthonormality and the vertex operator realization of s_{λ} ,

$$
\chi^{\lambda}_{\mu}(q) = \frac{q^{|\mu|}}{(q-1)^{l(\mu)}} \langle \hat{q}_{\mu}(q^{-1}), s_{\lambda} \rangle = \frac{q^{|\mu|}}{(q-1)^{l(\mu)}} \langle \hat{q}_{\mu}(q^{-1}), S_{\lambda}.1 \rangle.
$$

Denote $X^{\lambda}_{\mu}(t) = \langle \hat{q}_{\mu}(t), S_{\lambda} \cdot 1 \rangle$, then $\chi^{\lambda}_{\mu}(q) = \frac{q^{|\mu|}}{(q-1)^{l(\mu)}} X^{\lambda}_{\mu}(q^{-1})$. We are going to compute $X^{\lambda}_{\mu}(t)$ in the sequel.

3. Irreducible characters

In this section, we will give a new iterative formula, which implies compact formulae for some special cases. Moreover, we rederive the Murnaghan-Nakayama rule based on the vertex operator realization of Schur functions and the Frobenius formula.

It is known that the Young seminormal representation [\[7\]](#page-21-8) of the Hecke algebra $H_n(q)$ can be extended to that of $R_n(q)$ [\[3\]](#page-20-1). For $\lambda \vdash k$, an n-standard tableau L of shape λ is a filling of the diagram of λ with numbers from $\{1, 2, \ldots, n\}$ such that (1) each number appears at most once, (2) the entries in each column strictly increase from top to bottom, and (3) the entries in each row strictly increase from left to right.

If b is a box of L in position $(i, j) \in \lambda$, the *content* of b is defined as

$$
c(b) = j - i.
$$

For $i \in \{1, \ldots, k\}$, we sometimes call $i \in L$ if i appears inside L, and refer $L(i)$ as the box of L where i stands.

The number f_λ of n-standard tableaux of shape $\lambda \vdash n$ is given by $f_\lambda = n! / \prod_{(i,j) \in \lambda} h_{ij}$. Subsequently the number of n–standard tableaux of shape $\lambda \vdash k$ ($k \leq n$) is given by $\binom{n}{k}$ $\binom{n}{k}$ f_{λ} .

Let T_n^{λ} be the set of *n*-standard tableaux of shape $\lambda \vdash k$ ($k \leq n$), and define V^{λ} as the $\mathbb{C}(q^{1/2})$ -vector space spanned by $v_L, L \in T_n^{\lambda}$. We define an action of $R_n(q)$ on V^{λ} as follows: $P_i v_L = v_i$ for $i \notin L$ otherwise $P_i v_L = 0$ and

(3.2)
$$
T_i v_L = \begin{cases} \frac{q-1}{1-q^{c(L(i)) - c(L(i+1))}} v_L + (1 + \frac{q-1}{1-q^{c(L(i)) - c(L(i+1))}}) v_{L'}, & \text{if } i, i+1 \in L, \\ (q-1)v_L + q^{1/2} v_{s_i L}, & \text{if } i \notin L, i+1 \in L, \\ q^{1/2} v_{s_i L}, & \text{if } i \in L, i+1 \notin L, \\ q v_L, & \text{if } i, i+1 \notin L, \end{cases}
$$

where

(3.3)
$$
v_{L'} = \begin{cases} v_{s_i L}, & \text{if } s_i L \text{ is } n-\text{standard,} \\ 0, & \text{otherwise.} \end{cases}
$$

Theorem 3.1. [\[3\]](#page-20-1) For each $\lambda \vdash k \leq n$, the above $R_n(q)$ -action on V^{λ} affords an irreducible representation of $R_n(q)$. Moveover, the set $V^{\lambda}, \lambda \vdash k \leq n$, is a complete set of irreducible, pairwise non-isomorphic $R_n(q)$ –modules.

In the following we will simply use χ^{λ} to denote the irreducible character of V^{λ} .

3.1. A new iterative formula. For each partition $\lambda = (\lambda_1, \dots, \lambda_l)$, we define that

(3.4)
$$
\lambda^{[i]} = (\lambda_{i+1}, \cdots, \lambda_l), \qquad i = 0, 1, \ldots, l
$$

So $\lambda^{[0]} = \lambda$ and $\lambda^{[l]} = \emptyset$.

In [\[10,](#page-21-9) Sec. 5], two of us obtained the following decomposition of Schur functions. For an arbitrary partition $\lambda = (\lambda_1, \lambda_2, \cdots, \lambda_l) \vdash n$, then

(3.5)
$$
s_{\lambda} = \sum_{\nu} (-1)^{n-|\nu| - \lambda_1} h_{n-|\nu|} s_{\nu}
$$

summed over all partitions $\nu \subset \lambda^{[1]}$ such that $\lambda^{[1]}/\nu$ is a vertical strip.

Proposition 3.2. Let k be a nonnegative integer and $\mu = (\mu_1, \mu_2, \dots, \mu_m) \vdash n$ be a partition, then we have

(3.6)
$$
h_k^* \hat{q}_{\mu} \cdot 1 = \sum_{\tau \in C(\mu;k)} (1-t)^{l(\tau)} \hat{q}_{\mu-\tau} \cdot 1,
$$

where $C(\mu; k) := \{ \tau \subset \mu \mid \tau \models k \}.$

Proof. Note that $[\frac{\partial}{\partial p_n}, \hat{p}_m] = \delta_{n,m}$. The usual vertex operator calculus gives that

$$
\exp\left(\sum_{n=1}^{\infty}\frac{\partial}{\partial p_n}z^{-n}\right)\exp\left(\sum_{n=1}^{\infty}\frac{1-t^n}{n}\hat{p}_nw^n\right)=\exp\left(\sum_{n=1}^{\infty}\frac{1-t^n}{n}\hat{p}_nw^n\right)\exp\left(\sum_{n=1}^{\infty}\frac{\partial}{\partial p_n}z^{-n}\right)\frac{z-tw}{z-w}.
$$

Using repeatedly the above commutative relation noting that $\frac{\partial}{\partial n}$ $\frac{\partial}{\partial p_n}$.1 = 0, we have

(3.7)
$$
\exp\left(\sum_{n=1}^{\infty} \frac{\partial}{\partial p_n} z^{-n}\right) \exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} \hat{p}_n w_1^n\right) \cdots \exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} \hat{p}_n w_m^n\right).1
$$

$$
=\exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} \hat{p}_n w_1^n\right) \cdots \exp\left(\sum_{n=1}^{\infty} \frac{1-t^n}{n} \hat{p}_n w_m^n\right) \exp\prod_{m\geq i\geq 1} \frac{z-tw_i}{z-w_i}.
$$

The coefficient of $z^{-k}w_1^{\mu_1}w_2^{\mu_2}\cdots w_m^{\mu_m}$ is $h_k^*q_\mu$ in [\(3.7\)](#page-8-1) equals $h_k^*q_\mu$, and the right-hand side in (3.7) equals to

$$
\prod_{j=1}^{m} \left(1 + (1-t) \sum_{l \ge 1} \left(\frac{w_j}{z} \right)^l \right) \prod_{j=1}^{m} \left(\sum_{n \ge 0} \hat{q}_n w_j^n \right) \left(\sum_{n \ge 0} h_n^* z^{-n} \right).
$$

=
$$
\prod_{j=1}^{m} \left(1 + (1-t) \sum_{l \ge 1} \left(\frac{w_j}{z} \right)^l \right) \prod_{j=1}^{m} \left(\sum_{n \ge 0} \hat{q}_n w_j^n \right).
$$

in which the corresponding coefficient of $z^{-k}w_1^{\mu_1}w_2^{\mu_2}\cdots w_m^{\mu_m}$ is exactly $\sum_{\tau \in C(\mu;k)} (1-t)^{l(\tau)} \hat{q}_{\mu-\tau}$. $1 \square$

Now we state our main result.

Theorem 3.3. Let μ be a partition of $n, \lambda = (\lambda_1, \ldots, \lambda_r)$ be a partition of $m \leq n$. We have that

(3.8)
$$
\chi^{\lambda}_{\mu}(q) = \sum_{\nu} \sum_{\tau \in C(\mu; m - |\nu|)} (-1)^{m - |\nu| - \lambda_1} \frac{q^{m - |\nu| - l(\tau)}}{(q - 1)^{l(\mu) - l(\tau) - l(\mu - \tau)}} \chi^{\nu}_{\mu - \tau}(q)
$$

summed over all partitions $\nu \subset \lambda^{[1]}$ such that $\lambda^{[1]}/\nu$ is a vertical strip.

Proof. Direct computation gives that

$$
X^{\lambda}_{\mu}(t) = \langle \hat{q}_{\mu}.1, S_{\lambda}.1 \rangle
$$

=
$$
\sum_{\nu} (-1)^{m-|\nu| - \lambda_1} \langle \hat{q}_{\mu}.1, h_{m-|\nu|} s_{\nu} \rangle \quad \text{(by (3.5))}
$$

$$
= \sum_{\nu} (-1)^{m-|\nu|-\lambda_1} \left\langle h_{m-|\nu|}^* \hat{q}_{\mu}.1, s_{\nu} \right\rangle
$$

= $\sum_{\nu} (-1)^{m-|\nu|-\lambda_1} \left\langle \sum_{\tau \in C(\mu; m-|\nu|)} (1-t)^{l(\tau)} \hat{q}_{\mu-\tau}.1, s_{\nu} \right\rangle$ (by (3.6))
= $\sum_{\nu} \sum_{\tau \in C(\mu; m-|\nu|)} (-1)^{m-|\nu|-\lambda_1} (1-t)^{l(\tau)} X_{\mu-\tau}^{\nu}(t)$

summed over all partitions $\nu \subset \lambda^{[1]}$ such that $\lambda^{[1]}/\nu$ is a vertical strip. Then [\(3.8\)](#page-8-3) follows from $\chi^\lambda_\mu(q)=\frac{q^{|\mu|}}{(q-1)^{l(\mu)}}X^\lambda_\mu(q^{-1})$). \Box

We remark that in [\(3.8\)](#page-8-3) $\mu - \tau$ is seen as a partition obtained by arranging parts of composition $\mu - \tau$ in descending order.

Example 3.4. Suppose $\lambda = (31^2)$, $\mu = (321)$. The possible chooses of ν are \emptyset , (1), (1, 1). By Theorem [3.3,](#page-8-0) we have

$$
\chi_{(321)}^{(31^2)}(q) = \sum_{\tau \in C(\mu;5)} \frac{q^{5-l(\tau)}}{(q-1)^{3-l(\tau)-l(\mu-\tau)}} \chi_{\mu-\tau}^{\emptyset}(q) - \sum_{\tau \in C(\mu;4)} \frac{q^{4-l(\tau)}}{(q-1)^{3-l(\tau)-l(\mu-\tau)}} \chi_{\mu-\tau}^{(1)}(q)
$$

+
$$
\sum_{\tau \in C(\mu;3)} \frac{q^{3-l(\tau)}}{(q-1)^{3-l(\tau)-l(\mu-\tau)}} \chi_{\mu-\tau}^{(1^2)}(q)
$$

=
$$
(3q^3 - 2q^2) \chi_{(1)}^{\emptyset}(q) - (2q^2 - q) \chi_{(2)}^{(1)}(q) - (3q^3 - 4q^2 + q) \chi_{(1^2)}^{(1)}(q)
$$

+
$$
q \chi_{(3)}^{(1^2)}(q) + (4q^2 - 4q + 1) \chi_{(21)}^{(1^2)}(q) + q(q-1)^2 \chi_{(1^3)}^{(1^2)}(q).
$$

So
$$
\chi_{(321)}^{(31^2)}(q) = (q-1)(2q^2 - 8q + 2).
$$

Remark 3.5. In [\(3.8\)](#page-8-3) there appears the character $\chi_{(1)}^{\emptyset}(q)$ which is not necessarily equal to 1. In fact, we have that

$$
\chi_{\mu}^{\emptyset}(q) = \frac{q^{|\mu|}}{(q-1)^{l(\mu)}} \langle \hat{q}_{\mu}(q^{-1}), 1 \rangle
$$

=
$$
\frac{q^{|\mu|}}{(q-1)^{l(\mu)}} \times constant \ term \ of \ \hat{q}_{\mu}(q^{-1})
$$

=
$$
q^{|\mu|-l(\mu)} \quad (by \ (2.11)).
$$

3.2. **Murnaghan-Nakayama rule.** A path in a skew diagram θ is a sequence x_0, x_1, \dots, x_n of squares in θ such that x_{i-1} and x_i have a common side, for $1 \leq i \leq n$. A subset ξ of θ is connected if any two squares in ξ are connected by a path in ξ . The connected components, themselves skew diagrams, are by definition the maximal connected subsets of θ . A skew diagram θ is a border strip if it is connected and contains no 2×2 blocks of squares (cf. [\[11,](#page-21-4) p. 5]). A skew diagram is a *generalized*

border strip (written as GBS in short) if it is a union of connected border strips. Any GBS is a union of its connected components, each of which is a border strip. A k -GBS is a GBS with k boxes. In the example below, $\lambda = (4, 3, 3, 1), \mu = (3, 2, 1)$, the GBS λ/μ has three border strips. It's a 5-GBS.

If a GBS $\theta = \lambda/\mu$ has m connected components $(\xi_1, \xi_2, \dots, \xi_m)$, we define the weight of θ by

$$
wt(\theta; t) = (t-1)^{m-1} \prod_{i=1}^m (-1)^{r(\xi_i)-1} t^{c(\xi_i)-1},
$$

where the $r(\xi_i)$ (resp. $c(\xi_i)$) denotes the number of rows (resp. columns) in the border strip ξ_i . We make the convention that $wt(\lambda/\mu;t) = 1$ if $|\lambda/\mu| = 0$.

More generally, we define

(3.9)
$$
wt(\lambda/\mu; k, t) := \begin{cases} t^{k-1}wt(\lambda/\mu; t), & |\lambda/\mu| = 0; \\ (t-1)t^{k-|\lambda/\mu|-1}wt(\lambda/\mu; t), & 0 < |\lambda/\mu| < k; \\ wt(\lambda/\mu; t), & |\lambda/\mu| = k; \\ 0, & |\lambda/\mu| > k. \end{cases}
$$

Due to $[9,$ Corollary 4.2], we have

(3.10)
$$
q_k^*(t)S_{\lambda}.1 = \sum_{\mu} t^{k-1}(1-t)wt(\lambda/\mu;t^{-1})S_{\mu}.1 \quad (k>0)
$$

summed over all partitions μ such that λ/μ is a $k\text{-GBS}.$

We rewrite (2.11) as

(3.11)
$$
\frac{t^k}{t-1}\hat{q}_k(t^{-1}) = \frac{t^k}{t-1}q_k(t^{-1}) + \sum_{i=1}^{k-1}t^{k-1}q_i(t^{-1}) + t^{k-1}.
$$

Now we give an alternative proof of the following Murnaghan-Nakayama rule obtained first in [\[1,](#page-20-2) Prop. 3.3].

Theorem 3.6. Let k ba a positive integer, then

(3.12)
$$
\frac{t^k}{t-1} \hat{q}_k^*(t^{-1}) S_\lambda . 1 = \sum_{\mu} wt(\lambda/\mu; k, t) S_\mu . 1
$$

summed over partitions μ such that λ/μ is a GBS with $0 \leq |\lambda/\mu| \leq k$. In terms of irreducible characters, the iterative Murnaghan-Nakayama rule takes the following form

(3.13)
$$
\chi^{\lambda}_{\mu}(q) = \sum_{\nu} wt(\lambda/\nu; \mu_1, q) \chi^{\nu}_{\mu^{[1]}}(q)
$$

summed over partitions ν such that $|\nu| \leq |\mu^{[1]}|$ and λ/ν is a GBS with $0 \leq |\lambda/\nu| \leq \mu_1$.

Proof. It follows from immediate calculation that

$$
\frac{t^k}{t-1} \hat{q}_k^*(t^{-1}) S_\lambda.1
$$
\n
$$
= \frac{t^k}{t-1} q_k^*(t^{-1}) S_\lambda.1 + \sum_{i=1}^{k-1} t^{k-1} q_i^*(t^{-1}) S_\lambda.1 + t^{k-1} S_\lambda.1 \quad \text{(by (3.11))}
$$
\n
$$
= \sum_{\mu \vdash |\lambda| = k} wt(\lambda/\mu; t) S_\mu.1 + \sum_{i=1}^{k-1} t^{k-1} \sum_{\mu \vdash |\lambda| = i} (t-1) t^{-i} wt(\lambda/\mu; t) S_\mu.1 + t^{k-1} S_\lambda.1 \quad \text{(by (3.10))}
$$
\n
$$
= \sum_{\mu} wt(\lambda/\mu; k, t) S_\mu.1 \quad \text{(by (3.9))}
$$

summed over partitions μ such that λ/μ is a GBS with $0 \le |\lambda/\mu| \le k$.

3.3. Compact formulae for hook and two-row types. For a Young diagram λ , the hook-length of λ at $(i, j) \in \lambda$ is defined as $h_{i,j} = \lambda_i + \lambda'_j - i - j + 1$. In this subsection, we will consider character formulae for special cases based on the new iterative formula derived in the previous subsection. We first consider the following simple cases.

Proposition 3.7. Let $\lambda \vdash m$, $\mu \vdash n$ and $m \leq n$,

(3.14)
$$
\chi^{\lambda}_{(1^n)}(q) = {n \choose m} \frac{m!}{\prod_{(i,j)\in\lambda} h_{ij}},
$$

(3.15)
$$
\chi_{\mu}^{(m)}(q) = \frac{q^n}{(q-1)^{l(\mu)}} \sum_{\tau \in C(\mu;m)} (1 - q^{-1})^{l(\tau) + l(\mu - \tau)},
$$

(3.16)
$$
\chi_{\mu}^{(1^m)}(q) = \frac{q^n}{(q-1)^{l(\mu)}} \sum_{\tau \in C(\mu; n-m)} (1 - q^{-1})^{l(\tau) + l(\mu - \tau)} (-q^{-1})^{m - l(\mu - \tau)},
$$

(3.17)
$$
\chi^{\lambda}_{(n)}(q) = \sum_{\nu} wt(\lambda/\nu; n, q)
$$

summed over partitions ν such that $|\nu| \leq |\mu^{[1]}|$ and λ/ν is a GBS with $0 \leq |\lambda/\nu| \leq \mu_1$.

Proof. (3.15) and (3.17) are immediate results of (3.8) and (3.13) respectively. For (3.14) , we have $\chi^\lambda_{(1^n)}(q)=\langle \frac{q^n}{(q-1)^n}\hat{q}_{(1^n)}(q^{-1}),s_\lambda\rangle=\langle (p_1+1)^n,s_\lambda\rangle=\binom{n}{m}$ $\binom{n}{m}\langle p_1^m, s_\lambda\rangle = \binom{n}{m}$ $\binom{n}{m} \frac{m!}{\prod h_{ij}}$, in which the last identity

follows from the well-known hook formula of the symmetric group \mathfrak{S}_n [\[15\]](#page-21-10). For [\(3.16\)](#page-11-5), one has

$$
\chi_{\mu}^{(1^{m})}(q) = \left\langle \frac{q^{n}}{(q-1)^{l(\mu)}} \hat{q}_{\mu}(q^{-1}), s_{(1^{m})} \right\rangle
$$

\n
$$
= \frac{q^{n}}{(q-1)^{l(\mu)}} \left\langle \sum_{\tau \subset \mu} (1 - q^{-1})^{l(\tau)} q_{\mu-\tau}(q^{-1}), s_{(1^{m})} \right\rangle \text{ (by (2.11))}
$$

\n
$$
= \frac{q^{n}}{(q-1)^{l(\mu)}} \sum_{\tau \in C(\mu; n-m)} (1 - q^{-1})^{l(\tau) + l(\mu-\tau)} (-q^{-1})^{m-l(\mu-\tau)} \text{ (by [9, (2.27)])}.
$$

 \Box

For $\mu \vdash n$, we define two families of polynomials in t associated with μ by

$$
a_{ij}(\mu; t) = \sum_{\tau \in C(\mu; i)} \sum_{\theta \in C(\mu - \tau; j)} (1 - t^{-1})^{l(\tau) + l(\theta)} (1 - t)^{l(\mu - \tau - \theta)};
$$

$$
b_{ij}(\mu; t) = \sum_{\tau \in C(\mu; i)} \sum_{\theta \in C(\mu - \tau; j)} (1 - t^{-1})^{l(\tau) + l(\theta) + l(\mu - \tau - \theta)},
$$

for $i + j \leq n$ and $a_{ij}(\mu; t) = b_{ij}(\mu; t) = 0$ for $i + j > n$.

Symmetry of τ and θ yields

(3.18)
$$
a_{ij}(\mu;t) = a_{ji}(\mu;t),
$$

(3.19)
$$
b_{ij}(\mu;t) = b_{ji}(\mu;t).
$$

It is clear that $a_{0,0}(\mu;t) = (1-t)^{l(\mu)}$ and $b_{0,0}(\mu;t) = (1-t^{-1})^{l(\mu)}$.

Proposition 3.8. We have the generating functions of a_{ij} and b_{ij} as follows:

$$
(3.20)
$$

$$
\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} a_{ij}(\mu; t) v^i u^j = \prod_{i=1}^{l(\mu)} \left(\sum_{\tau_i=0}^{\mu_i} \sum_{\theta_i=0}^{\mu_i - \tau_i} (1 - t^{-1})^{1 - \delta_{\tau_i,0}} (1 - t^{-1})^{1 - \delta_{\theta_i,0}} (1 - t)^{1 - \delta_{\mu_i, \tau_i + \theta_i}} v^{\tau_i} u^{\theta_i} \right),
$$
\n(3.21)
$$
\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} b_{ij}(\mu; t) v^i u^j = \prod_{i=1}^{l(\mu)} \left(\sum_{\tau_i=0}^{\mu_i} \sum_{\theta_i=0}^{\mu_i - \tau_i} (1 - t^{-1})^{1 - \delta_{\tau_i,0}} (1 - t^{-1})^{1 - \delta_{\theta_i,0}} (1 - t^{-1})^{1 - \delta_{\mu_i, \tau_i + \theta_i}} v^{\tau_i} u^{\theta_j} \right).
$$

Proof. It follows from immediate calculations that

$$
\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} a_{ij}(\mu; t) v^i u^j
$$
\n
$$
= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \sum_{\tau \in C(\mu; i)} \sum_{\theta \in C(\mu - \tau; j)} (1 - t^{-1})^{l(\tau) + l(\theta)} (1 - t)^{l(\mu - \tau - \theta)} v^i u^j
$$
\n
$$
= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \sum_{\tau_1 + \dots + \tau_l = i} \sum_{\theta_1 + \dots + \theta_l = j} (1 - t^{-1})^{2l - \sum_{s=1}^l \delta_{\tau_s, 0} - \sum_{s=1}^l \delta_{\theta_s, 0}} (1 - t)^{l - \sum_{s=1}^l \delta_{\mu_s, \tau_s + \theta_s} v^i u^j
$$

$$
= \prod_{i=1}^l \left(\sum_{\tau_i=0}^{\mu_i} \sum_{\theta_i=0}^{\mu_i-\tau_i} (1-t^{-1})^{1-\delta_{\tau_i,0}} (1-t^{-1})^{1-\delta_{\theta_i,0}} (1-t)^{1-\delta_{\mu_i,\tau_i+\theta_i}} v^{\tau_i} u^{\theta_i} \right).
$$

The second identity can be derived similarly.

Prop. [3.8](#page-12-0) gives the following special cases, which is essential to the subsequent contents.

Corollary 3.9. For a partition $\mu = (\mu_1, \dots, \mu_l) \vdash n$, we have

(3.22)
$$
\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} a_{ij}(\mu; q) q^{i+j} = \prod_{i=1}^{l} (q-1) q^{\mu_i-1},
$$

(3.23)
$$
\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} a_{ij}(\mu; q) (-q)^i q^j = \prod_{i=1}^l (1-q)(-q)^{\mu_i-1},
$$

(3.24)
$$
\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} a_{ij}(\mu; q)(-q)^i (-q)^j = \prod_{i=1}^l (1-q)A(\mu_i; q),
$$

$$
(3.25) \qquad \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} b_{ij}(\mu; q) = (1 - q^{-1})^l \prod_{i=1}^l \left(3\mu_i - 3q^{-1}(\mu_i - 1) + \frac{(\mu_i - 1)(\mu_i - 2)(1 - q^{-1})^2}{2} \right),
$$

where $A(\mu_i;q) = \left(\frac{(-q)^{\mu_i-1}(q^2+6q+1)+4}{(q+1)^2} + \frac{2(-q)^{\mu_i-1}(q-1)\mu_i}{q+1} \right)$.

Now we present our main results: compact formulae for $\chi^{\lambda}_{\mu}(q)$ with λ hook and two-row type.

Theorem 3.10. Let $\mu \vdash n$ and $1 \leq k \leq m \leq n$, we have

(3.26)
$$
\chi_{\mu}^{(k,1^{m-k})}(q) = \frac{q^{n-m}}{(q-1)^{l(\mu)}} \sum_{i=k}^{m} (-1)^{m-k} a_{i,n-m}(\mu;q) q^i,
$$

(3.27)
$$
\chi_{\mu}^{(k,m-k)}(q) = \frac{q^n}{(q-1)^{l(\mu)}} \left(b_{k,m-k}(\mu;q) - b_{k+1,m-k-1}(\mu;q) \right).
$$

Proof. For the first identity, by Theorem [3.3,](#page-8-0) all possible ν 's can be chosen from $\{(1^j) | 0 \le j \le m-k\}$, then

$$
\chi_{\mu}^{(k,1^{m-k})}(q) = \sum_{j=0}^{m-k} \sum_{\tau \in C(\mu;m-j)} (-1)^{m-k-j} \frac{q^{m-j-l(\tau)}}{(q-1)^{l(\mu)-l(\tau)-l(\mu-\tau)}} \chi_{\mu-\tau}^{(1^j)}(q)
$$

$$
= \sum_{j=0}^{m-k} \sum_{\tau \in C(\mu;m-j)} (-1)^{m-k-j} \frac{q^{m-j-l(\tau)}}{(q-1)^{l(\mu)-l(\tau)-l(\mu-\tau)}} \frac{q^{n-m+j}}{(q-1)^{l(\mu-\tau)}}
$$

$$
\sum_{\theta \in C(\mu-\tau;n-m)} (1-q^{-1})^{l(\theta)+l(\mu-\tau-\theta)} (-q^{-1})^{j-l(\mu-\tau-\theta)} \quad \text{(by (3.16))}.
$$

Replacing $j \to m - i$ and simplification give [\(3.26\)](#page-13-1). The second identity is obtained similarly. Just note $\nu = (m - k)$ or $(m - k - 1)$ in this case.

Example 3.11. The following two examples show effectiveness of our compact formulas.

(i) Suppose $\lambda = (31^2)$, $\mu = (321)$, we have

(3.28)
$$
\chi^{(31^2)}_{(321)}(q) = \frac{q}{(q-1)^3} \left(a_{31}(\mu;q) q^3 + a_{41}(\mu;q) q^4 + a_{51}(\mu;q) q^5 \right).
$$

Theorem [3.8](#page-12-0) gives

$$
\sum_{i=0}^{6} \sum_{i=0}^{6} a_{ij} v^i u^j = (1 - t^{-1})^3 \left[-t + (v + u) \right] \left[-t + (v^2 + u^2) + (1 - t)(u + v) + (1 - t^{-1})vu \right]
$$

$$
\left[-t + (v^3 + u^3) + (1 - t)(v + v^2 + u + u^2) + (1 - t^{-1})(v^2u + vu^2) + (1 - t^{-1})(1 - t)uv \right].
$$

By taking the coefficient of v^3u , v^4u , v^5u , we conclude that

$$
\begin{cases}\na_{31}(\mu;q) = (1 - q^{-1})^3 (7q^2 - 19q + 13 - 2q^{-1}) \\
a_{41}(\mu;q) = (1 - q^{-1})^3 (11 - 8q - 3q^{-1}) \\
a_{51}(\mu;q) = 2(1 - q^{-1})^4.\n\end{cases}
$$

Substituting them into [\(3.28\)](#page-14-0) yields $\chi^{(31^2)}_{(321)}(q) = q^3 - 10q^2 + 10q - 2$.

(ii) Suppose $\lambda = (32)$, $\mu = (2^3)$, we have

(3.29)
$$
\chi_{(2^3)}^{(32)}(q) = \frac{q^6}{(q-1)^3} (b_{32}(\mu;q) - b_{41}(\mu;q)).
$$

Theorem [3.8](#page-12-0) implies

$$
\sum_{i=0}^{6} \sum_{j=0}^{6} b_{ij} v^{i} u^{j} = (1 - t^{-1})^{3} [v^{2} + u^{2} + (1 - t^{-1})(vu + v + u) + 1]^{3}.
$$

By taking the coefficients of v^3u^2 , v^4u , we conclude

$$
\begin{cases}\nb_{32}(\mu;q) = 6(1-q^{-1})^5 + 6(1-q^{-1})^4 \\
b_{41}(\mu;q) = 6(1-q^{-1})^5 + 3(1-q^{-1})^4.\n\end{cases}
$$

Plugging them into (3.29) gives $\chi_{(23)}^{(32)}$ $\binom{(32)}{(2^3)}(q) = 3q^2(q-1).$

Now we study the sign q-permutation representation of the rook monoid $R_n(q)$. Let $V = V_{\bar{0}} \bigoplus V_{\bar{1}}$ be the Z₂-graded vector space over $\mathbb{C}(q)$ with basis $\{v_1, v_2, \cdots, v_{a+b}\}$, where $V_{\overline{0}} = \langle v_{a+1}, v_{a+2}, \cdots, v_{a+b} \rangle$ and $V_{\bar{1}} = \langle v_{a+1}, v_{a+2}, \cdots, v_{a+b} \rangle.$

 \Box

Define the endomorphism π of $V \otimes V$ by

(3.30)
$$
\pi(v_k \otimes v_l) = \begin{cases} (-1)^{|v_k||v_l|} v_l \otimes v_k + (q-1)v_k \otimes v_l, k < l \\ \frac{(-1)^{|v_k|}(q+1) + q - 1}{2} v_k \otimes v_l, k = l \\ (-1)^{|v_k||v_l|} q v_l \otimes v_k, k > l, \end{cases}
$$

and let π_i be the endomorphism of $V^{\otimes n}$ by letting π acting on the $(i, i+1)$ factor and the trivial identity action elsewhere, $i = 1, \dots, n-1$. The action π gives rise to the so-called q-permutation representation $\Phi_{a,b}^{q,n}$ of $H_n(q)$. In particular, $\varphi_{1,1}^n$ and $\varphi_{2,0}^n$ decomposes themselves [\[19\]](#page-21-11) into some special direct sums of irreducible representations of $H_n(q)$:

(3.31)
$$
\varphi_{1,1}^n = \sum_{k=0}^{n-1} \zeta^{(n-k,1^k)},
$$

(3.32)
$$
\varphi_{2,0}^n = \sum_{k=0}^{\lceil n/2 \rceil} (n - 2k + 1) \zeta^{(n-k,k)},
$$

where ζ^{λ} is the irreducible character of the Hecke algebra indexed by λ and $\lceil n/2 \rceil$ denotes the maximum integer number $\leq \frac{n}{2}$ $\frac{n}{2}$. In [\[9\]](#page-21-3) two compact formulas were given for $\varphi_{1,1}^n, \varphi_{2,0}^n$:

(3.33)
$$
\varphi_{1,1}^n(T_{\gamma\mu}) = \sum_{i=0}^{n-1} \zeta_{\mu}^{(n-i,1^i)}(q) = (-1)^{n-l(\mu)} 2^{l(\mu)-1} \prod_{i=1}^{l(\mu)} [\mu_i]_{-q},
$$

(3.34)
$$
\varphi_{2,0}^n(T_{\gamma\mu}) = \sum_{i=0}^{\lceil n/2 \rceil} (n-2i+1) \zeta_{\mu}^{(n-i,i)}(q) = q^{n-2l(\mu)} \prod_{i=1}^{l(\mu)} (1+q+\mu_i(q-1)),
$$

where $[n]_q$ is the q-binomial number defined by $[n]_q := \frac{q^{n-1}}{q-1}$ $rac{p^{n}-1}{q-1}$.

The following results seem analogs of (3.33) and (3.34) for the monoid $H_q(n)$.

Theorem 3.12. Under the same assumption of Theorem [3.10,](#page-13-0) one has

(3.35)
$$
\sum_{m=0}^{n} \sum_{k=0}^{m-1} \chi_{\mu}^{(m-k,1^{k})}(q) = \frac{(-1)^{n+l(\mu)}}{2} \left(\prod_{i=1}^{l(\mu)} A(\mu_{i};q) - (-q)^{n-l(\mu)} \right),
$$

(3.36)

$$
\sum_{m=0}^{n} \sum_{k=0}^{\lceil m/2 \rceil} (m-2k+1) \chi_{\mu}^{(m-k,k)}(q) = q^{n-l(\mu)} \prod_{i=1}^{l(\mu)} \left(3\mu_i - 3q^{-1}(\mu_i - 1) + \frac{(\mu_i - 1)(\mu_i - 2)(1 - q^{-1})^2}{2} \right).
$$

Proof. For the first identity, by (3.26) , we have

(3.37)
$$
\sum_{m=0}^{n} \sum_{k=0}^{m-1} \chi_{\mu}^{(m-k,1^k)}(q) = \sum_{m=0}^{n} \sum_{k=0}^{m-1} \frac{(-1)^k}{(q-1)^l} \sum_{i=m-k}^{m} a_{i,n-m}(\mu;q) q^i q^{n-m}.
$$

Note that

$$
\sum_{k=0}^{m-1} \sum_{i=m-k}^{m} (-1)^k a_{i,n-m}(\mu;q) q^i
$$

=
$$
\sum_{i=1}^{m} \sum_{k=m-i}^{m-1} (-1)^k a_{i,n-m}(\mu;q) q^i
$$
 (exchange the sums)
=
$$
\sum_{i=1}^{m} \frac{(-1)^m ((-1)^i - 1)}{2} a_{i,n-m}(\mu;q) q^i.
$$

Since $\frac{(-1)^m((-1)^i-1)}{2} = 0$ when $i = 0$ and $a_{i,j}(\mu; q) = 0$ when $i + j > |\mu|$,

$$
\sum_{k=0}^{m-1} \sum_{i=m-k}^{m} (-1)^{k} a_{i,n-m}(\mu;q) q^{i} = \sum_{i=0}^{n} \frac{(-1)^{m}((-1)^{i}-1)}{2} a_{i,n-m}(\mu;q) q^{i}.
$$

Substituting this to (3.37) gives

$$
\sum_{m=0}^{n} \sum_{k=0}^{m-1} \chi_{\mu}^{(m-k,1^{k})}(q)
$$
\n
$$
= \frac{(-1)^{n}}{(q-1)^{l(\mu)}} \sum_{m=0}^{n} \sum_{i=0}^{n} \frac{(-1)^{i}-1}{2} a_{i,n-m}(\mu;q) q^{i}(-q)^{n-m}
$$
\n
$$
= \frac{(-1)^{n}}{(q-1)^{l(\mu)}} \sum_{m=0}^{n} \sum_{i=0}^{n} \frac{(-1)^{i}-1}{2} a_{i,m}(\mu;q) q^{i}(-q)^{m} \quad \text{(replace } m \text{ by } n-m)
$$
\n
$$
= \frac{(-1)^{n}}{(q-1)^{l(\mu)}} \sum_{m=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-1)^{i}-1}{2} a_{i,m}(\mu;q) q^{i}(-q)^{m} \quad (a_{i,j}(\mu;q) = 0 \text{ if } i+j > n)
$$
\n
$$
= \frac{(-1)^{n}}{2(q-1)^{l(\mu)}} \left(\sum_{i=0}^{\infty} \sum_{m=0}^{\infty} a_{i,m}(\mu;q) (-q)^{i} (-q)^{m} - \sum_{m=0}^{\infty} \sum_{i=0}^{\infty} a_{m,i}(\mu;q) (-q)^{m} q^{i} \right) \quad (a_{i,j}(\mu;q) = a_{j,i}(\mu;q)).
$$

Then [\(3.35\)](#page-15-3) follows from Corollary [3.9.](#page-13-2)

For the second identity, we have

$$
\sum_{m=0}^{n} \sum_{k=0}^{\lfloor m/2 \rfloor} (m - 2k + 1) \chi_{\mu}^{(m-k,k)}(q)
$$

=
$$
\sum_{m=0}^{n} \frac{q^n}{(q-1)^{l(\mu)}} \sum_{k=0}^{\lfloor m/2 \rfloor} (m - 2k + 1) (b_{m-k,k}(\mu; q) - b_{m-k+1,k-1}(\mu; q))
$$
 (by (3.27))
=
$$
\frac{q^n}{(q-1)^{l(\mu)}} \sum_{m=0}^{n} \sum_{k=0}^{m} b_{m-k,k}(\mu; q) \quad (b_{i,j}(\mu; q) = b_{j,i}(\mu; q))
$$

=
$$
\frac{q^n}{(q-1)^{l(\mu)}} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} b_{m,k}(\mu; q) \quad (b_{i,j}(\mu; q) = 0 \text{ if } i+j > n).
$$

Then Corollary [3.9](#page-13-2) gives (3.36) .

It would be interesting to construct the representation of $H_q(n)$ afforded by Theorem [3.12.](#page-15-5)

4. The second orthogonal relation

In this section, we introduce the bitrace of $R_n(q)$ as an analogue of the bitrace of the Iwahori-Hecke algebra $H_n(q)$ in type A [\[4\]](#page-20-3).

Recall that $\{T_{(A,B,\omega)}|(A,B,\omega\in\Omega)\}$ forms a linear $\mathbb{C}(q)$ -basis of $R_n(q)$. Define the bitrace of elements $T_{(C,D,x)}$ and $T_{(E,F,y)}$ by

(4.1)
$$
btr(T_{(C,D,x)}, T_{(E,F,y)}):=\sum_{(A,B,\omega)\in\Omega} T_{(C,D,x)}T_{(A,B,\omega)}T_{(E,F,y)}|_{T_{(A,B,\omega)}}
$$

where $T_{(C,D,x)}T_{(A,B,\omega)}T_{(E,F,y)}|_{T_{(A,B,\omega)}}$ denotes the coefficient of the basis element $T_{(A,B,\omega)}$ in the linear expansion of $T_{(C,D,x)}T_{(A,B,\omega)}T_{(E,F,y)}$. If $(A, B, \omega) \in \Omega$, let $L_{(A,B,\omega)}$ (resp. $R_{(A,B,\omega)}$) denote the linear transformation of $R_n(q)$ induced by the action of (A, B, ω) on $R_n(q)$ by the left (resp. right) multiplication, then $L_{(A,B,\omega)}$ and $R_{(A,B,\omega)}$ are commutative each other and

(4.2)
$$
br(T_{(C,D,x)}, T_{(E,F,y)}) = Tr(L_{(C,D,x)}R_{(E,F,y)}).
$$

The algebra $R_n(q)$ becomes a $R_n(q)$ -bimodule under the left and right multiplication. Since $R_n(q)$ is semisimple, by double centralizer theory, we have that

(4.3)
$$
R_n(q) \cong \bigoplus_{k=0}^n \bigoplus_{\lambda \vdash k} R_{\lambda} \otimes R^{\lambda}
$$

where R_{λ} is the irreducible left $R_n(q)$ -module labeled by λ , R^{λ} is the irreducible right $R_n(q)$ -module labeled by λ , then

(4.4)
$$
btr(T_{(C,D,x)}, T_{(E,F,y)}) = \sum_{k=0}^{n} \sum_{\lambda \vdash k} \chi^{\lambda}_{R_n(q)}(T_{(C,D,x)}) \chi^{\lambda}_{R_n(q)}(T_{(E,F,y)}).
$$

Note that any irreducible character of $R_n(q)$ is completely determined by its values on the standard elements $T_{\mu}, \mu \vdash k, k \leq n$ [\[1,](#page-20-2) Theorem 5.5]. For this reason, we can define the bitrace of arbitrary two compositions $\mu, \nu \models n$ as

(4.5)
$$
btr(\mu, \nu) := br(T_{\mu}, T_{\nu}) = \sum_{k=0}^{n} \sum_{\lambda \vdash k} \chi_{\mu}^{\lambda}(q) \chi_{\nu}^{\lambda}(q).
$$

Using (2.15) and (2.11) respectively, one has

$$
btr(\mu, \nu) = \frac{q^{2n}}{(q-1)^{l(\mu)+l(\nu)}} \langle \hat{q}_{\mu}(x; q^{-1}), \hat{q}_{\nu}(x; q^{-1}) \rangle
$$

\n
$$
= \frac{q^{2n}}{(q-1)^{l(\mu)+l(\nu)}} \left\langle \sum_{\tau \subset \mu} (1 - q^{-1})^{l(\mu-\tau)} q_{\tau}(q^{-1}), \sum_{\theta \subset \nu} (1 - q^{-1})^{l(\nu-\theta)} q_{\theta}(q^{-1}) \right\rangle
$$

\n
$$
= \frac{q^{2n}}{(q-1)^{l(\mu)+l(\nu)}} \sum_{k=0}^{n} \sum_{\substack{\tau \in C(\mu;k) \\ \theta \in C(\nu;k)}} (1 - q^{-1})^{l(\mu-\tau)+l(\nu-\theta)} \langle q_{\tau}(q^{-1}), q_{\theta}(q^{-1}) \rangle
$$

\n
$$
= \frac{q^{2n}}{(q-1)^{l(\mu)+l(\nu)}} \sum_{k=0}^{n} \sum_{\substack{\tau \in C(\mu;k) \\ \theta \in C(\nu;k)}} (1 - q^{-1})^{l(\tau)+l(\theta)} \langle q_{\mu-\tau}(q^{-1}), q_{\nu-\theta}(q^{-1}) \rangle.
$$

For simplicity we denote for integer $r > 0$

$$
(r)_t = (t-1)\frac{t^{2r} - 1}{t+1}
$$

and make the convention that $(r)_t = 0$ for $r < 0$ and $(0)_t = 1$.

By $[9, (5.8)]$ we have that

(4.7)
$$
\langle q_{\mu-\tau}(q^{-1}), q_{\nu-\theta}(q^{-1}) \rangle = \frac{1}{q^{2n-2k}} \sum_{M} \prod_{m_{ij}} (m_{ij})_q.
$$

summed over all $l(\mu) \times l(\nu)$ nonnegative integer matrices $M = (m_{ij})$ such that the row sums are $\mu_1 - \tau_1, \ldots, \mu_{l(\mu)} - \tau_{l(\mu)}$ and the column sums are $\nu_1 - \theta_1, \ldots, \nu_{l(\nu)} - \theta_{l(\nu)}$.

Substituting (4.7) into (4.6) gives that

$$
btr(\mu, \nu)
$$
\n
$$
= \frac{1}{(q-1)^{l(\mu)+l(\nu)}} \sum_{k=0}^{n} \sum_{\substack{\tau \in C(\mu;k) \\ \theta \in C(\nu;k)}} (1-q^{-1})^{l(\tau)+l(\theta)} q^{2k} \sum_{M} \prod_{m_{ij}} (m_{ij})_q
$$
\n
$$
= \frac{1}{(q-1)^{l(\mu)+l(\nu)}} \sum_{k=0}^{n} \sum_{\substack{\tau \in C(\mu;k) \\ \theta \in C(\nu;k)}} \sum_{M} \prod_{i \geq 1}^{l(\mu)} (1-q^{-1})^{1-\delta_{0,\tau_i}} q^{\tau_i} \prod_{j \geq 1} (1-q^{-1})^{1-\delta_{0,\theta_i}} q^{\theta_i} \prod_{m_{ij}} (m_{ij})_q
$$

summed over all $l(\mu) \times l(\nu)$ nonnegative integer matrices $M = (m_{ij})$ such that the row sums are $\mu_1 - \tau_1, \ldots, \mu_{l(\mu)} - \tau_{l(\mu)}$ and the column sums are $\nu_1 - \theta_1, \ldots, \nu_{l(\nu)} - \theta_{l(\nu)}$.

For an arbitrary nonnegative integer matrix $M = (m_{ij})$, define the weight of element m_{ij} by

(4.9)
$$
wt(m_{ij}) := \begin{cases} (1 - q^{-1})^{1 - \delta_{0, m_{ij}}} q^{m_{ij}}, & i = 1 \text{ or } j = 1; \\ (m_{ij})_q, & i > 1 \text{ and } j > 1. \end{cases}
$$

Moreover, the weight of M is defined as $wt(M) := \prod_{m_{ij}} wt(m_{ij}).$

Then we can derive the following combinatorial formula for $btr(\mu, \nu)$.

Theorem 4.1. Let $\mu, \nu \models n$, then

(4.10)
$$
btr(\mu, \nu) = \frac{1}{(q-1)^{l(\mu)+l(\nu)}} \sum_{M} wt(M)
$$

summed over all $(l(\mu) + 1) \times (l(\nu) + 1)$ nonnegative integer matrices $M = (m_{ij})$ satisfying (i) $m_{11} = 0;$

(ii) the first row sum equals to the first column sum;

(iii) the *i*-th row sum is μ_i and the *j*-th column sum is ν_j , $2 \le i \le l(\mu)$, $2 \le j \le l(\nu)$.

Corollary 4.2. Let $T_{\mu} \in R_n(q)$ and $\mu \models n$, then the trace of regular representation of $R_n(q)$ at element T_{μ} is given by

(4.11)
$$
\chi^{reg}(T_{\mu}) = \frac{q^n}{(q-1)^{l(\mu)}} \sum_{i=0}^{k} \sum_{\tau \in C(\mu;i)} {n \choose i} (1-q^{-1})^{l(\mu-\tau)+i} \frac{i!}{\prod_{j} \tau_{j}!}.
$$

Proof. By definition, $\chi^{reg}(T_\mu) = \text{btr}(\mu, (1^n))$. Let us first fix the first row (column) sum in M to be i $(0 \leq i \leq n)$, then the first row in M has $\binom{n}{i}$ $\binom{n}{i}$ possibilities. If \tilde{M} denotes the submatrix of M obtained by removing the first row and the first column, then the entries in \tilde{M} are 0 or 1. Assume that we choose the first column to be $(0, \tau_1, \cdots, \tau_{l(\mu)}) \subset \mu$, then each column has at most one 1 and the *j*-th row has $\mu_j - \tau_j$ 1's in \tilde{M} . Thus the number of these \tilde{M} is $\frac{(n-i)!}{\prod_{i=1}^{n} n_i}$ $\frac{(h-i)!}{\prod_j (\mu_j - \tau_j)!}$. In this case, $wt(M) = (1 - q^{-1})^{l(\tau) + i} q^{2i} (q - \tau_j)$ 1)²ⁿ⁻²ⁱ. Hence $br(\mu, (1^n)) = \frac{1}{(q-1)^{l(\mu)+n}} \sum_{i=0}^{n} \sum_{\tau \in C(\mu; i)} {n \choose i}$ $\binom{n}{i} \frac{(n-i)!}{\prod (u - \tau)}$ $\frac{(h-i)!}{\prod_j(\mu_j-\tau_j)!}(1-q^{-1})^{l(\tau)+i}q^{2i}(q-1)^{2n-2i}.$ Substituting $i \to n - i$ and $\tau \to \mu - \tau$ gives [\(4.11\)](#page-19-2).

We remark that [\(4.11\)](#page-19-2) reduces to $\chi^{reg}(1) = \sum_{i=0}^{n} {n \choose i}$ $\left(\mu_{i}\right)^{2} i!$ when $\mu = (1^{n})$, which gives the dimension of $R_n(q)$.

5. Appendix

In [\[1\]](#page-20-2), a table of $\chi^{\lambda}_{\mu}(q)$ ($|\mu| \leq 4$) was presented. Here we list $\chi^{\lambda}_{\mu}(q)$ with $|\mu| = 5$. Since $\chi^{\lambda}_{\mu}(q)$ equals to the irreducible character of the Hecke algebra when $|\lambda| = |\mu|$, we just list those with $|\mu| = 5$ and $|\lambda| < 5$.

Acknowledgments

The paper is partially supported by Simons Foundation grant No. 523868 and NSFC grant No. 12171303. The third author is supported by China Scholarship Council.

REFERENCES

- [1] M. Dieng, T. Halverson, V. Poladian, *Character formulas for* q*-rook monoid algebras.* J. Algebraic Combin. 17 (2003), no. 2, 99-123.
- [2] C. Grood, *A Spect module analog for the rook monoid*, Electronic J. Combinatorics 9 (1) (2002).
- [3] T. Halverson, *Representations of the q-rook monoid.* J. Algebra 273 (2004), no. 1, 227-251.
- [4] T. Halverson, R. Leduc, A. Ram, *Iwahori-Hecke algebras of type A, bitraces and symmetric functions*, Int. Math. Res. Not. 9 (1997), 401-416.
- [5] T. Halverson, A. Ram, *Bitraces for* GLn(Fq) *and the Iwahori-Hecke algebra of type* Aⁿ−1, Indag. Math, 10(1999),247- 268.
- [6] T. Halverson, A. Ram, q*-rook monoid algebras, Hecke algebras, and Schur-Weyl duality.* Zap. Nauchn. Sem. S.- Peterburg. Otdel. Mat. Inst. Steklov. (POMI) 283 (2001), Teor. Predst. Din. Sist. Komb. i Algoritm. Metody. 6, 224-250, 262-263; reprinted in J. Math. Sci. (N.Y.) 121 (2004), no. 3, 2419-2436
- [7] P.N. Hoefsmit, *Representations of Hecke algebras of finite groups with* BN*-pairs of classical type*, Thesis, Univ. of British Columbia, 1974.
- [8] N. Jing, *Vertex operators, symmetric functions, and the spin group* Γn, J. Algebra 138:2 (1991), 340-398.
- [9] N. Jing, N. Liu, *On irreducible characters of the Iwahori-Hecke algebra in type* A, J. Algebra 598 (2022) 24-47.
- [10] N. Jing, N. Liu, *A multiparametric Murnaghan-Nakayama rule for Macdonald polynomials*, preprint, 2023, arXiv: 2310.15730.
- [11] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford university press, 1998.
- [12] H. Mitsuhashi, *Schur-Weyl reciprocity between the quantum superalgebra and the Iwahori-Hecke algebra*, Alg. Represen. Theory 9 (2006), 309-322.
- [13] W. Munn, *The characters of the symmetric inverse semigroup.* Proc. Cambridge Philos. Soc. 53 (1957), 13-18.
- [14] A. Ram, *A Frobenius formula for the characters of the Hecke algebras.* Invent. Math. 106 (1991), no. 3, 461-488.
- [15] B. E. Sagan, *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, 2nd ed., Springer-Verlag, New York, 2001.
- [16] L. Solomon, *The Bruhat decomposition, Tits system and Iwahori ring for the monoid of matrices over a finite field.* Geom. Dedicata 36 (1990), no. 1, 15-49.
- [17] L. Solomon, *Representations of the rook monoid*, J. Algebra, 256 (2002), 309-342.
- [18] L. Solomon, *The Iwahori algebra of* $M_n(\mathbb{F}_q)$. A presentation and a representation on tensor space. J. Algebra 273 (2004), no. 1, 206-226.
- [19] D. Zhao, *Characters of Iwahori-Hecke algebras*, Israel J. Math. 229 (2019), 67-83.

Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA *Email address*: jing@ncsu.edu

School of Mathematics, South China University of Technology, Guangzhou, Guangdong 510640, China *Email address*: wywymath@163.com

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090 VIENNA, AUSTRIA *Email address*: mathliu123@outlook.com