

Eigenvalues of quantum Gelfand invariants

Naihuan Jing, Ming Liu and Alexander Molev

Abstract

We consider the quantum Gelfand invariants which first appeared in a landmark paper by Reshetikhin, Takhtadzhyan and Faddeev (1989). We calculate the eigenvalues of the invariants acting in irreducible highest weight representations of the quantized enveloping algebra for \mathfrak{gl}_n . The calculation is based on Liouville-type formulas relating two families of central elements in the quantum affine algebras of type A .

1 Introduction

The *quantized enveloping algebras* and *quantum affine algebras* associated with simple Lie algebras comprise remarkable families of quantum groups, as introduced by V. Drinfeld [5] and M. Jimbo [13]. These algebras and their representations have since found numerous connections with many areas in mathematics and physics.

In this paper we will be concerned with those families associated with the general linear Lie algebras \mathfrak{gl}_n . Both the quantized enveloping algebra $U_q(\mathfrak{gl}_n)$ and the quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$ admit *R-matrix* (or *RTT*) *presentations* going back to the work of the Leningrad school headed by L. D. Faddeev; see e.g [19] and [27] for reviews of the foundations of the *R-matrix* approach originated in the quantum inverse scattering method.

Central elements in both $U_q(\mathfrak{gl}_n)$ and $U_q(\widehat{\mathfrak{gl}}_n)$ are constructed with the use of the *R-matrix* presentations and found as coefficients of the respective *quantum determinants*; see [3], [14], [19], and also [6] for more general constructions of central elements in the quantized enveloping algebras and quantum affine algebras. As pointed out in [27], the quantum traces of powers of generator matrices are central in $U_q(\mathfrak{gl}_n)$; see also [1]. By taking the limit $q \rightarrow 1$ one recovers the central elements of $U(\mathfrak{gl}_n)$ going back to [8], which are known as the *Gelfand invariants*. Note that a different generalization of the Gelfand invariants for \mathfrak{gl}_n as central elements in $U_q(\mathfrak{gl}_n)$, was given in [9], where their eigenvalues in irreducible highest weight representations were calculated.

A new family of central elements in $U_q(\widehat{\mathfrak{gl}}_n)$ was given in [2] and they were related to the quantum determinants by Liouville-type formulas, although they were not accompanied by proofs. This result is quite analogous to the corresponding quantum Liouville formulas for the Yangians originated in [22] and we give a complete proof in this paper.

By taking the images of the new central elements of $U_q(\widehat{\mathfrak{gl}}_n)$ under the evaluation homomorphism, we recover the quantum Gelfand invariants of [27]. The Liouville formulas will then

allow us to calculate the eigenvalues of the quantum Gelfand invariants in irreducible highest weight representations of $U_q(\mathfrak{gl}_n)$. We thus obtain q -analogues of the Perelomov–Popov formulas [24]. To recall the eigenvalue formulas from [24], consider the irreducible highest weight representation $L(\lambda)$ of \mathfrak{gl}_n with the highest weight $\lambda = (\lambda_1, \dots, \lambda_n)$ and combine the standard basis elements E_{ij} into the matrix $E = [E_{ij}]$. Then the eigenvalue of the Gelfand invariant $\text{tr } E^m$ in $L(\lambda)$ is found by

$$\text{tr } E^m \mapsto \sum_{k=1}^n \ell_k^m \frac{(\ell_1 - \ell_k + 1) \dots (\ell_n - \ell_k + 1)}{(\ell_1 - \ell_k) \dots \wedge \dots (\ell_n - \ell_k)}, \quad (1.1)$$

where $\ell_i = \lambda_i + n - i$ and the symbol \wedge indicates that the zero factor is skipped. Formula (1.1) can be derived with the use of R -matrix calculations in the Yangian $Y(\mathfrak{gl}_n)$ or in the universal enveloping algebra $U(\mathfrak{gl}_n)$; see [20, Sec. 7.1] and [21, Sec. 4.8], respectively.

Our main result concerning the quantum Gelfand invariants is the following theorem, where we calculate the eigenvalues of the quantum traces $\text{tr}_q M^m$ of the powers of the generator matrix $M = L^-(L^+)^{-1}$ in the representation $L_q(\lambda)$ of $U_q(\mathfrak{gl}_n)$ (see Section 3 for the definitions). We use a standard notation for the q -numbers

$$[k]_q = \frac{q^k - q^{-k}}{q - q^{-1}}, \quad k \in \mathbb{Z}.$$

Theorem 1.1. *The eigenvalue of the quantum Gelfand invariant $\text{tr}_q M^m$ in $L_q(\lambda)$ is found by*

$$\text{tr}_q M^m \mapsto \sum_{k=1}^n q^{2\ell_k m} \frac{[\ell_1 - \ell_k + 1]_q \dots [\ell_n - \ell_k + 1]_q}{[\ell_1 - \ell_k]_q \dots \wedge \dots [\ell_n - \ell_k]_q}. \quad (1.2)$$

We will prove Theorem 1.1 in Section 3 by deriving it from the Liouville formula given in Theorem 2.4 in a way similar to [20, Sec. 7.1]. We also consider three more families of central elements of $U_q(\mathfrak{gl}_n)$ which, however, turn out to coincide with the quantum Gelfand invariants, up to a possible replacement $q \mapsto q^{-1}$.

In the limit $q \rightarrow 1$ we have

$$\frac{M - 1}{q - q^{-1}} \rightarrow E \quad (1.3)$$

by (3.1) and (3.2) below. Therefore, the Gelfand invariants in $U(\mathfrak{gl}_n)$ are recovered from the elements $\text{tr}_q M^m$ in the limit, while the Perelomov–Popov formulas (1.1) follow from (1.2); see Remark 3.1 below.

We point out a related recent work [18], where explicit formulas for certain central elements in the reflection equation algebras were given and their relation with the quantum Gelfand invariants were reviewed. This includes the connection with an earlier construction of central elements in [25] and with the Cayley–Hamilton theorem and Newton identities of [10], [12] and [23].

This work was completed during the first and third named authors' visits to the South China University of Technology and to the Shanghai University. They are grateful to the Departments of Mathematics in both universities for the warm hospitality.

2 Liouville formulas

We will regard q as a nonzero complex number which is not a root of unity. Recall the R -matrix presentation of the quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$ as introduced in [26]. We follow [4] and use the same settings as in our earlier work [17]. Let $e_{ij} \in \text{End } \mathbb{C}^n$ denote the standard matrix units. Consider the R -matrix

$$R(x) = \frac{f(x)}{q - q^{-1}x}(R - x\tilde{R}), \quad (2.1)$$

where

$$R = q \sum_i e_{ii} \otimes e_{ii} + \sum_{i \neq j} e_{ii} \otimes e_{jj} + (q - q^{-1}) \sum_{i < j} e_{ij} \otimes e_{ji} \quad (2.2)$$

and

$$\tilde{R} = q^{-1} \sum_i e_{ii} \otimes e_{ii} + \sum_{i \neq j} e_{ii} \otimes e_{jj} - (q - q^{-1}) \sum_{i > j} e_{ij} \otimes e_{ji},$$

while the formal power series

$$f(x) = 1 + \sum_{k=1}^{\infty} f_k x^k, \quad f_k = f_k(q),$$

is uniquely determined by the relation

$$f(xq^{2n}) = f(x) \frac{(1 - xq^2)(1 - xq^{2n-2})}{(1 - x)(1 - xq^{2n})}.$$

The quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$ is generated by elements

$$l_{ij}^+[-r], \quad l_{ij}^-[r] \quad \text{with} \quad 1 \leq i, j \leq n, \quad r = 0, 1, \dots,$$

and the invertible central element q^c , subject to the defining relations

$$\begin{aligned} l_{ji}^+[0] &= l_{ij}^-[0] = 0 & \text{for} & \quad 1 \leq i < j \leq n, \\ l_{ii}^+[0] l_{ii}^-[0] &= l_{ii}^-[0] l_{ii}^+[0] = 1 & \text{for} & \quad i = 1, \dots, n, \end{aligned}$$

and

$$R(u/v)L_1^\pm(u)L_2^\pm(v) = L_2^\pm(v)L_1^\pm(u)R(u/v), \quad (2.3)$$

$$R(uq^{-c}/v)L_1^+(u)L_2^-(v) = L_2^-(v)L_1^+(u)R(uq^c/v). \quad (2.4)$$

In the last two formulas we consider the matrices $L^\pm(u) = [l_{ij}^\pm(u)]$, whose entries are formal power series in u and u^{-1} ,

$$l_{ij}^+(u) = \sum_{r=0}^{\infty} l_{ij}^+[-r]u^r, \quad l_{ij}^-(u) = \sum_{r=0}^{\infty} l_{ij}^-[r]u^{-r}.$$

Here and below we regard the matrices as elements

$$L^\pm(u) = \sum_{i,j=1}^n e_{ij} \otimes l_{ij}^\pm(u) \in \text{End } \mathbb{C}^n \otimes U_q(\widehat{\mathfrak{gl}}_n)[[u^{\pm 1}]]$$

and use a subscript to indicate a copy of the matrix in the multiple tensor product algebra

$$\underbrace{\text{End } \mathbb{C}^n \otimes \dots \otimes \text{End } \mathbb{C}^n}_k \otimes U_q(\widehat{\mathfrak{gl}}_n)[[u^{\pm 1}]] \quad (2.5)$$

so that

$$L_a^\pm(u) = \sum_{i,j=1}^n 1^{\otimes(a-1)} \otimes e_{ij} \otimes 1^{\otimes(k-a)} \otimes l_{ij}^\pm(u).$$

We regard the usual matrix transposition also as the linear map

$$t : \text{End } \mathbb{C}^n \rightarrow \text{End } \mathbb{C}^n, \quad e_{ij} \mapsto e_{ji}.$$

For any $a \in \{1, \dots, k\}$ we will denote by t_a the corresponding partial transposition on the algebra (2.5) which acts as t on the a -th copy of $\text{End } \mathbb{C}^n$ and as the identity map on all the other tensor factors.

The scalar factor in (2.1) is necessary for the R -matrix to satisfy the *crossing symmetry relations* [7]. We will use one of them given by

$$\left(R_{12}(x)^{-1} \right)^{t_2} D_2 R_{12}(xq^{2n})^{t_2} = D_2, \quad (2.6)$$

where D denotes the diagonal $n \times n$ matrix

$$D = \text{diag}[q^{n-1}, q^{n-3}, \dots, q^{-n+1}].$$

The following proposition was stated in [2, Eq. (4.28)] without proof.

Proposition 2.1. *There exist a series $z^+(u)$ in u and a series $z^-(u)$ in u^{-1} with coefficients in the algebra $U_q(\widehat{\mathfrak{gl}}_n)$ such that*

$$L^\pm(uq^{2n})^t D (L^\pm(u)^{-1})^t = z^\pm(u) D \quad (2.7)$$

and

$$(L^\pm(u)^{-1})^t D^{-1} L^\pm(uq^{2n})^t = z^\pm(u) D^{-1}. \quad (2.8)$$

Moreover, the coefficients of the series $z^\pm(u)$ belong to the center of the algebra $U_q(\widehat{\mathfrak{gl}}_n)$.

Proof. Multiply both sides of (2.3) by $L_2^\pm(v)^{-1}$ from the left and the right and apply the transposition t_2 to get

$$R(u/v)^{t_2} \left(L_2^\pm(v)^{-1} \right)^t L_1^\pm(u) = L_1^\pm(u) \left(L_2^\pm(v)^{-1} \right)^t R(u/v)^{t_2}$$

and hence

$$\left(R(u/v)^{t_2}\right)^{-1} L_1^\pm(u) \left(L_2^\pm(v)^{-1}\right)^t = \left(L_2^\pm(v)^{-1}\right)^t L_1^\pm(u) \left(R(u/v)^{t_2}\right)^{-1}. \quad (2.9)$$

Use the crossing symmetry relation (2.6) to replace the R -matrix by

$$\left(R(u/v)^{t_2}\right)^{-1} = D_2^{-1} \left(R(u/vq^{2n})^{-1}\right)^{t_2} D_2$$

and get

$$\left(R(u/vq^{2n})^{-1}\right)^{t_2} D_2 L_1^\pm(u) \left(L_2^\pm(v)^{-1}\right)^t D_2^{-1} = D_2 \left(L_2^\pm(v)^{-1}\right)^t L_1^\pm(u) D_2^{-1} \left(R(u/vq^{2n})^{-1}\right)^{t_2}.$$

Now cancel the scalar factors appearing in (2.1) on both sides of this relation and observe that the R -matrix $R - x\tilde{R}$ evaluated at $x = 1$ equals $(q - q^{-1})P$, where P is the permutation operator. Therefore,

$$\left((R - \tilde{R})^{-1}\right)^{t_2} = \frac{1}{q - q^{-1}} Q \quad \text{with} \quad Q = \sum_{i,j=1}^n e_{ij} \otimes e_{ij}.$$

Hence, by taking $u = vq^{2n}$ we get

$$Q D_2 L_1^\pm(vq^{2n}) \left(L_2^\pm(v)^{-1}\right)^t D_2^{-1} = D_2 \left(L_2^\pm(v)^{-1}\right)^t L_1^\pm(vq^{2n}) D_2^{-1} Q.$$

Since Q is an operator in $\text{End } \mathbb{C}^n \otimes \text{End } \mathbb{C}^n$ with a one-dimensional image, both sides must be equal to $Q z^\pm(v)$ for series $z^\pm(v)$ with coefficients in the quantum affine algebra. Using the relations $QX_1 = QX_2^t$ and $X_1Q = X_2^tQ$ which hold for an arbitrary matrix X , we can write the definition of $z^\pm(v)$ as

$$Q L_2^\pm(vq^{2n})^t D_2 \left(L_2^\pm(v)^{-1}\right)^t = Q D_2 z^\pm(v) \quad (2.10)$$

and

$$\left(L_2^\pm(v)^{-1}\right)^t D_2^{-1} L_2^\pm(vq^{2n})^t Q = D_2^{-1} Q z^\pm(v). \quad (2.11)$$

By taking trace over the first copy of $\text{End } \mathbb{C}^n$ on both sides of (2.10) and (2.11) we arrive at (2.7) and (2.8), respectively.

We will now use (2.8) to show that the series $z^-(v)$ commutes with $L^+(u)$. We have

$$L_1^+(u) z^-(v) = L_1^+(u) D_2 \left(L_2^-(v)^{-1}\right)^t D_2^{-1} L_2^-(vq^{2n})^t. \quad (2.12)$$

Transform relation (2.4) in the same way as we did for (2.3) in the beginning of the proof to get the following counterpart of (2.9):

$$\left(R(uq^{-c}/v)^{t_2}\right)^{-1} L_1^+(u) \left(L_2^-(v)^{-1}\right)^t = \left(L_2^-(v)^{-1}\right)^t L_1^+(u) \left(R(uq^c/v)^{t_2}\right)^{-1}.$$

Hence, the right hand side of (2.12) equals

$$D_2 R_{12}(uq^{-c}/v)^{t_2} \left(L_2^-(v)^{-1}\right)^t L_1^+(u) \left(R_{12}(uq^c/v)^{t_2}\right)^{-1} D_2^{-1} L_2^-(vq^{2n})^t. \quad (2.13)$$

Applying again (2.6), we can write

$$\left(R_{12}(uq^c/v)^{t_2}\right)^{-1} D_2^{-1} = D_2^{-1} \left(R_{12}(uq^c/vq^{2n})^{-1}\right)^{t_2}.$$

Continue transforming (2.13) by using the following consequence of (2.4):

$$L_1^+(u) \left(R_{12}(uq^c/vq^{2n})^{-1}\right)^{t_2} L_2^-(vq^{2n})^t = L_2^-(vq^{2n})^t \left(R_{12}(uq^{-c}/vq^{2n})^{-1}\right)^{t_2} L_2^-(vq^{2n})^t,$$

so that (2.13) becomes

$$D_2 R_{12}(uq^{-c}/v)^{t_2} \left(L_2^-(v)^{-1}\right)^t D_2^{-1} L_2^-(vq^{2n})^t \left(R_{12}(uq^{-c}/vq^{2n})^{-1}\right)^{t_2} L_1^+(u).$$

By (2.8) this simplifies to

$$D_2 R_{12}(uq^{-c}/v)^{t_2} D_2^{-1} \left(R_{12}(uq^{-c}/vq^{2n})^{-1}\right)^{t_2} z^-(v) L_1^+(u)$$

which equals $z^-(v) L_1^+(u)$ by (2.6). This proves that $L_1^+(u) z^-(v) = z^-(v) L_1^+(u)$. The relation $L_1^-(u) z^-(v) = z^-(v) L_1^-(u)$ and the centrality of $z^+(v)$ are verified in the same way. \square

Remark 2.2. The counterparts of the series $z^\pm(u)$ for the quantum affine algebras of types B , C and D appear in [15, Prop. 3.3] and [16, Prop. 3.3], where they were introduced by relations analogous to (2.7) and (2.8).

Corollary 2.3. *We have the formulas*

$$z^\pm(u) = \frac{1}{[n]_q} \operatorname{tr} DL^\pm(uq^{2n})L^\pm(u)^{-1} \quad (2.14)$$

and

$$z^\pm(u) = \frac{1}{[n]_q} \operatorname{tr} D^{-1} L^\pm(u)^{-1} L^\pm(uq^{2n}). \quad (2.15)$$

Proof. The formulas follow by taking trace on both sides of the respective matrix relations (2.7) and (2.8). \square

Recall that the *quantum determinants* $\operatorname{qdet} L^+(u)$ and $\operatorname{qdet} L^-(u)$ are series in u and u^{-1} , respectively, whose coefficients belong to the center of the quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$:

$$\operatorname{qdet} L^\pm(u) = \sum_{\sigma \in \mathfrak{S}_n} (-q)^{-l(\sigma)} l_{\sigma(1)1}^\pm(uq^{2n-2}) \cdots l_{\sigma(n)n}^\pm(u),$$

where $l(\sigma)$ denotes the length of the permutation σ .

The following is a q -analogue of the quantum Liouville formula of [22]. It was stated in [2, Eq. (4.32)].

Theorem 2.4. *We have the relations*

$$z^\pm(u) = \frac{\operatorname{qdet} L^\pm(uq^2)}{\operatorname{qdet} L^\pm(u)}. \quad (2.16)$$

Proof. We follow [20, Sec. 1.9] and use the *quantum comatrices* $\widehat{L}^\pm(u)$ introduced in [23] and [28]. They are defined by the relations

$$\widehat{L}^\pm(uq^2)L^\pm(u) = \text{qdet } L^\pm(u) 1, \quad (2.17)$$

where 1 denotes the identity matrix. We will derive formulas for the entries of the matrices $\widehat{L}^\pm(u)$ by using the quantum minor relations reviewed e.g. in [11] which we outline below.

Recall that the q -permutation operator $P^q \in \text{End } \mathbb{C}^n \otimes \text{End } \mathbb{C}^n$ is defined by

$$P^q = \sum_{i=1}^n e_{ii} \otimes e_{ii} + q \sum_{i>j} e_{ij} \otimes e_{ji} + q^{-1} \sum_{i<j} e_{ij} \otimes e_{ji}.$$

The action of the symmetric group \mathfrak{S}_k on the space $(\mathbb{C}^n)^{\otimes k}$ can be defined by setting $s_i \mapsto P_{s_i}^q := P_{i,i+1}^q$ for $i = 1, \dots, k-1$, where s_i denotes the transposition $(i, i+1)$. If $\sigma = s_{i_1} \cdots s_{i_l}$ is a reduced decomposition of an element $\sigma \in \mathfrak{S}_k$ then we set $P_\sigma^q = P_{s_{i_1}}^q \cdots P_{s_{i_l}}^q$. Denote by e_1, \dots, e_n the canonical basis vectors of \mathbb{C}^n . Then for any indices $a_1 < \cdots < a_k$ and any $\tau \in \mathfrak{S}_k$ we have

$$P_\sigma^q(e_{a_{\tau(1)}} \otimes \cdots \otimes e_{a_{\tau(k)}}) = q^{l(\sigma\tau^{-1})-l(\tau)} e_{a_{\tau\sigma^{-1}(1)}} \otimes \cdots \otimes e_{a_{\tau\sigma^{-1}(k)}}. \quad (2.18)$$

We denote by $A_q^{(k)}$ the q -antisymmetrizer

$$A_q^{(k)} = \sum_{\sigma \in \mathfrak{S}_k} \text{sgn } \sigma \cdot P_\sigma^q. \quad (2.19)$$

The defining relations (2.3) and the *fusion procedure* [3] for the quantum affine algebra imply the relations

$$A_q^{(k)} L_1^\pm(uq^{2k-2}) \cdots L_k^\pm(u) = L_k^\pm(u) \cdots L_1^\pm(uq^{2k-2}) A_q^{(k)}. \quad (2.20)$$

The quantum minors are the series $l_{b_1 \cdots b_k}^{\pm a_1 \cdots a_k}(u)$ with coefficients in $U_q(\widehat{\mathfrak{gl}}_n)$ defined by the expansion of the elements (2.20) as

$$\sum_{a_i, b_i} e_{a_1 b_1} \otimes \cdots \otimes e_{a_k b_k} \otimes l_{b_1 \cdots b_k}^{\pm a_1 \cdots a_k}(u).$$

Explicit formulas for the quantum minors have the form: if $a_1 < \cdots < a_k$ then

$$l_{b_1 \cdots b_k}^{\pm a_1 \cdots a_k}(u) = \sum_{\sigma \in \mathfrak{S}_k} (-q)^{-l(\sigma)} \cdot l_{a_{\sigma(1)} b_1}^\pm(uq^{2k-2}) \cdots l_{a_{\sigma(k)} b_k}^\pm(u). \quad (2.21)$$

If $b_1 < \cdots < b_k$ (and the a_i are arbitrary) then

$$l_{b_1 \cdots b_k}^{\pm a_1 \cdots a_k}(u) = \sum_{\sigma \in \mathfrak{S}_k} (-q)^{l(\sigma)} \cdot l_{a_k b_{\sigma(k)}}^\pm(u) \cdots l_{a_1 b_{\sigma(1)}}^\pm(uq^{2k-2})$$

and for any $\tau \in \mathfrak{S}_k$ we have

$$l_{b_{\tau(1)} \cdots b_{\tau(k)}}^{\pm a_1 \cdots a_k}(u) = (-q)^{-l(\tau)} l_{b_1 \cdots b_k}^{\pm a_1 \cdots a_k}(u).$$

The following lemma was pointed out in [23] and [28].

Lemma 2.5. *The (i, j) entry of the matrix $\widehat{L}^\pm(u)$ is given by*

$$\widehat{l}_{ij}^\pm(u) = (-q)^{j-i} l_{1 \dots \widehat{i} \dots n}^{\pm 1 \dots \widehat{j} \dots n}(u), \quad (2.22)$$

where the hats on the right-hand side indicate the indices to be omitted.

Proof. According to (2.21), the quantum determinants are defined by the relations

$$A_q^{(n)} L_1^\pm(uq^{2n-2}) \cdots L_n^\pm(u) = A_q^{(n)} \text{qdet } L^\pm(u).$$

Hence, the definition (2.17) of the quantum comatrices implies

$$A_q^{(n)} L_1^\pm(uq^{2n-4}) \cdots L_{n-1}^\pm(u) = A_q^{(n)} \widehat{L}_n^\pm(u).$$

Apply both sides to the vector $e_1 \otimes \cdots \otimes \widehat{e}_i \otimes \cdots \otimes e_n \otimes e_j$ and use (2.18) and (2.19) to equate the coefficients of the vector $A_q^{(n)}(e_1 \otimes \cdots \otimes e_n)$ to obtain (2.22). \square

We will also use the relations for the transposed comatrices; see [23] and [28].

Lemma 2.6. *We have the relations*

$$D \widehat{L}^\pm(u)^t D^{-1} L^\pm(uq^{2n-2})^t = \text{qdet } L^\pm(u) 1.$$

Proof. Since each relation only involves generators belonging to the subalgebra of $U_q(\widehat{\mathfrak{gl}}_n)$ generated by the coefficients of the series $l_{ij}^+(u)$ or $l_{ij}^-(u)$, we may assume that the central charge is specialized to zero, $c = 0$, and derive the desired relations in the quotient algebra $U_q^\circ(\widehat{\mathfrak{gl}}_n)$. The mapping

$$\theta : L^\pm(u) \rightarrow L^\mp(u^{-1})^t$$

defines an automorphism of the algebra $U_q^\circ(\widehat{\mathfrak{gl}}_n)$. By Lemma 2.5 and the quantum minor formulas, for the images under θ we have

$$\theta : \widehat{L}^\pm(u) \mapsto D \widehat{L}^\mp(u^{-1}q^{-2n+4})^t D^{-1}$$

and

$$\theta : \text{qdet } L^\pm(u) \mapsto \text{qdet } L^\mp(u^{-1}q^{-2n+2}).$$

It remains to apply θ to both sides of relations (2.17) and then replace u by $u^{-1}q^{-2n+2}$. \square

Now, using (2.17) and the centrality of $\text{qdet } L^\pm(u)$, we can write (2.7) as

$$z^\pm(u) = L^\pm(uq^{2n})^t D(L^\pm(u)^{-1})^t D^{-1} = (\text{qdet } L^\pm(u))^{-1} L^\pm(uq^{2n})^t D \widehat{L}^\pm(uq^2)^t D^{-1}.$$

By applying Lemma 2.6 with u replaced by uq^2 we find that this expression coincides with the right hand side of (2.16). \square

Remark 2.7. Since the second part of Proposition 2.1 was not used in the proof of Theorem 2.4, the fact that the coefficients of both series $z^+(u)$ and $z^-(u)$ belong to the center of the quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$ also follows from (2.16) due to the respective properties of the quantum determinants $\text{qdet } L^+(u)$ and $\text{qdet } L^-(u)$.

3 Quantum Gelfand invariants

We will follow [4] to define the *quantized enveloping algebra* $U_q(\mathfrak{gl}_n)$ in its R -matrix presentation [27] as the algebra generated by elements l_{ij}^+ and l_{ij}^- with $1 \leq i, j \leq n$ subject to the relations

$$\begin{aligned} l_{ij}^- &= l_{ji}^+ = 0, & 1 \leq i < j \leq n, \\ l_{ii}^- l_{ii}^+ &= l_{ii}^+ l_{ii}^- = 1, & 1 \leq i \leq n, \\ R L_1^\pm L_2^\pm &= L_2^\pm L_1^\pm R, & R L_1^+ L_2^- = L_2^- L_1^+ R, \end{aligned}$$

where the R -matrix R is defined in (2.2), while L^+ and L^- are the matrices

$$L^\pm = \sum_{i,j} e_{ij} \otimes l_{ij}^\pm \in \text{End } \mathbb{C}^n \otimes U_q(\mathfrak{gl}_n).$$

The universal enveloping algebra $U(\mathfrak{gl}_n)$ is recovered from $U_q(\mathfrak{gl}_n)$ in the limit $q \rightarrow 1$ by the formulas

$$\frac{l_{ij}^-}{q - q^{-1}} \rightarrow E_{ij}, \quad \frac{l_{ji}^+}{q - q^{-1}} \rightarrow -E_{ji} \quad \text{for } i > j, \quad (3.1)$$

and

$$\frac{l_{ii}^- - 1}{q - 1} \rightarrow E_{ii}, \quad \frac{l_{ii}^+ - 1}{q - 1} \rightarrow -E_{ii} \quad \text{for } i = 1, \dots, n. \quad (3.2)$$

Set $M = L^-(L^+)^{-1}$. By [27], the quantum traces defined by $\text{tr}_q M^m = \text{tr } D M^m$ belong to the center of the algebra $U_q(\mathfrak{gl}_n)$. The elements $\text{tr}_q M^m$ act by multiplication by scalars in the irreducible highest weight representations $L_q(\lambda)$. The representation $L_q(\lambda)$ of $U_q(\mathfrak{gl}_n)$ is generated by a nonzero vector ξ such that

$$\begin{aligned} l_{ij}^+ \xi &= 0 & \text{for } 1 \leq i < j \leq n, \\ l_{ii}^- \xi &= q^{\lambda_i} \xi & \text{for } 1 \leq i \leq n, \end{aligned}$$

for an n -tuple $\lambda = (\lambda_1, \dots, \lambda_n)$ of integers (or real numbers). This is a q -deformation of the irreducible \mathfrak{gl}_n -module $L(\lambda)$ with the highest weight λ .

Proof of Theorem 1.1. We will show that the eigenvalue of the quantum Gelfand invariant $\text{tr}_q M^m$ in $L_q(\lambda)$ is given by formula (1.2).

Recall the evaluation homomorphism $U_q(\widehat{\mathfrak{gl}}_n) \rightarrow U_q(\mathfrak{gl}_n)$ defined by

$$L^+(u) \mapsto L^+ - L^- u, \quad L^-(u) \mapsto L^- - L^+ u^{-1}, \quad q^c \mapsto 1, \quad (3.3)$$

and apply it to both sides of the Liouville formula

$$z^+(u) = \frac{\text{qdet } L^+(uq^2)}{\text{qdet } L^+(u)} \quad (3.4)$$

proved in Theorem 2.4. The image of the quantum determinant $\text{qdet } L^+(u)$ is found by

$$\sum_{\sigma \in \tilde{\mathfrak{S}}_n} (-q)^{-l(\sigma)} \left(l_{\sigma(1)1}^+ - l_{\sigma(1)1}^- u q^{2n-2} \right) \cdots \left(l_{\sigma(n)n}^+ - l_{\sigma(n)n}^- u \right).$$

By applying this element to the highest vector ξ of $L_q(\lambda)$, we find that its eigenvalue is given by

$$\left(q^{-\lambda_1} - q^{\lambda_1+2n-2} u \right) \cdots \left(q^{-\lambda_n} - q^{\lambda_n} u \right) = q^{n(n-1)/2} (q^{-\ell_1} - q^{\ell_1} u) \cdots (q^{-\ell_n} - q^{\ell_n} u)$$

with $\ell_i = \lambda_i + n - i$. Therefore, the eigenvalue of the image of the right hand side of (3.4) in $L_q(\lambda)$ is given by

$$\frac{(q^{-\ell_1} - q^{\ell_1+2} u) \cdots (q^{-\ell_n} - q^{\ell_n+2} u)}{(q^{-\ell_1} - q^{\ell_1} u) \cdots (q^{-\ell_n} - q^{\ell_n} u)}.$$

To expand this rational function into a series in u write it as

$$C + \frac{a_1}{1 - q^{2\ell_1} u} + \cdots + \frac{a_n}{1 - q^{2\ell_n} u}$$

to find that the constants a_k are given by

$$a_k = (q^{n-1} - q^{n+1}) \frac{[\ell_1 - \ell_k + 1]_q \cdots [\ell_n - \ell_k + 1]_q}{[\ell_1 - \ell_k]_q \cdots \wedge \cdots [\ell_n - \ell_k]_q}.$$

Then write

$$\frac{1}{1 - q^{2\ell_k} u} = \sum_{m=0}^{\infty} q^{2\ell_k m} u^m.$$

On the other hand, using (2.14), for the image of $z^+(u)$ under the evaluation homomorphism (3.3) we get

$$\frac{1}{[n]_q} \text{tr } D(L^+ - L^- u q^{2n}) (L^+ - L^- u)^{-1} = \frac{1}{[n]_q} \text{tr } D(1 - M u q^{2n}) (1 - M u)^{-1}$$

which equals

$$1 + (q^{n-1} - q^{n+1}) \sum_{m=1}^{\infty} \text{tr}_q M^m u^m.$$

Formula (1.2) now follows by equating the coefficients of the powers u^m on both sides of the power expansions. Note that the formula is also valid for $m = 0$. \square

Remark 3.1. Due to (1.3), the Gelfand invariant $\text{tr } E^m \in U(\mathfrak{gl}_n)$ is obtained as the limit value as $q \rightarrow 1$ of the expression

$$\frac{1}{(q - q^{-1})^m} \text{tr } D(M - 1)^m = \frac{1}{(q - q^{-1})^m} \sum_{r=0}^m \binom{m}{r} (-1)^{m-r} \text{tr}_q M^r.$$

We will look at the limit value of the corresponding linear combination of the eigenvalues given by (1.2) and note that the q -numbers specialize by the rule $[r]_q \rightarrow r$ as $q \rightarrow 1$. Hence, for each $k = 1, \dots, n$ it suffices to find the limit value of the expression

$$\frac{1}{(q - q^{-1})^m} \sum_{r=0}^m \binom{m}{r} (-1)^{m-r} q^{2\ell_k r} = \frac{(q^{2\ell_k} - 1)^m}{(q - q^{-1})^m},$$

which equals ℓ_k^m , thus yielding formula (1.1). \square

Since the evaluation homomorphism (3.3) is surjective, the centrality property of the elements $\text{tr}_q M^m$ also follows from the above calculations due to Proposition 2.1. Similarly, by using the three remaining formulas in (2.14) and (2.15), we get three more families of central elements in $U_q(\mathfrak{gl}_n)$ together with the relations between them given by

$$\text{tr } D^{-1}((L^+)^{-1}L^-)^m = \text{tr}_q M^m \quad \text{and} \quad \text{tr } D^{-1}((L^-)^{-1}L^+)^m = \text{tr } D(L^+(L^-)^{-1})^m.$$

Moreover, additional relations between the families are provided by the isomorphism

$$U_q(\mathfrak{gl}_n) \rightarrow U_{q^{-1}}(\mathfrak{gl}_n), \quad L^\pm \mapsto (L^\pm)^{-1}.$$

This implies that the formulas for the eigenvalues of the other two central elements of $U_q(\mathfrak{gl}_n)$ in $L_q(\lambda)$ are obtained by the replacement $q \mapsto q^{-1}$ in (1.2):

$$\text{tr } D(L^+(L^-)^{-1})^m \mapsto \sum_{k=1}^n q^{-2\ell_k m} \frac{[\ell_1 - \ell_k + 1]_q \cdots [\ell_n - \ell_k + 1]_q}{[\ell_1 - \ell_k]_q \cdots \wedge \cdots [\ell_n - \ell_k]_q}.$$

Alternatively, these eigenvalue formulas can be derived in the same way as in the proof of Theorem 1.1 by working with the other Liouville formula in (2.16) instead of (3.4).

Declarations

Competing interests

The authors have no competing interests to declare that are relevant to the content of this article.

Funding

This work was supported by the Australian Research Council, grant DP240101572.

Availability of data and materials

No data was used for the research described in the article.

References

- [1] P. Baumann, *On the center of quantized enveloping algebras*, J. Algebra **203** (1998), 244–260.
- [2] S. Belliard and E. Ragoucy, *The nested Bethe ansatz for ‘all’ open spin chains with diagonal boundary conditions*, J. Phys. A **42** (2009), no. 20, 205203, 35 pp.
- [3] I. V. Cherednik, *A new interpretation of Gelfand–Tsetlin bases*, Duke Math. J. **54** (1987), 563–577.

- [4] J. Ding and I. B. Frenkel, *Isomorphism of two realizations of quantum affine algebra $U_q(\widehat{\mathfrak{gl}}(n))$* , *Comm. Math. Phys.* **156** (1993), 277–300.
- [5] V. G. Drinfeld, *Hopf algebras and the quantum Yang–Baxter equation*, *Soviet Math. Dokl.* **32** (1985), 254–258.
- [6] P. I. Etingof, *Central elements for quantum affine algebras and affine Macdonald’s operators*, *Math. Res. Lett.* **2** (1995), 611–628.
- [7] I. B. Frenkel and N. Yu. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, *Comm. Math. Phys.* **146** (1992), 1–60.
- [8] I. M. Gelfand, *Center of the infinitesimal group ring* (Russian), *Mat. Sbornik* **26** (1950), 103–112. English transl. in: I. M. Gelfand, “Collected Papers”, Vol. II, Berlin: Springer-Verlag, 1988, pp. 22–30.
- [9] M. D. Gould, R. B. Zhang and A. J. Bracken, *Generalized Gel’fand invariants and characteristic identities for quantum groups*, *J. Math. Phys.* **32** (1991), 2298–2303.
- [10] D. I. Gurevich, P. N. Pyatov and P. A. Saponov, *Hecke symmetries and characteristic relations on reflection equation algebras*, *Lett. Math. Phys.* **41** (1997), 255–264.
- [11] M. J. Hopkins and A. I. Molev, *A q -analogue of the centralizer construction and skew representations of the quantum affine algebra*, *Symmetry, Integrability and Geometry: Methods and Applications* **2** (2006), paper 092, 29 pp.
- [12] A. Isaev, O. Ogievetsky and P. Pyatov, *On quantum matrix algebras satisfying the Cayley–Hamilton–Newton identities*, *J. Phys. A* **32** (1999), L115–L121.
- [13] M. Jimbo, *A q -difference analogue of $U(\mathfrak{g})$ and the Yang–Baxter equation*, *Lett. Math. Phys.* **10** (1985), 63–69.
- [14] M. Jimbo, *A q -analogue of $U_q(\mathfrak{gl}(N + 1))$, Hecke algebra and the Yang–Baxter equation*, *Lett. Math. Phys.* **11** (1986), 247–252.
- [15] N. Jing, M. Liu and A. Molev, *Isomorphism between the R -matrix and Drinfeld presentations of quantum affine algebra: type C* , *J. Math. Phys.* **61** (2020), no. 3, 031701, 41 pp.
- [16] N. Jing, M. Liu and A. Molev, *Isomorphism between the R -matrix and Drinfeld presentations of quantum affine algebra: types B and D* , *SIGMA* **16** (2020), 043, 49 pp.
- [17] N. Jing, M. Liu and A. Molev, *Quantum Sugawara operators in type A* , preprint [arXiv:2212.11435](https://arxiv.org/abs/2212.11435).
- [18] D. Jordan and N. White, *The center of the reflection equation algebra via quantum minors*, *J. Algebra* **542** (2020), 308–342.

- [19] P. P. Kulish and E. K. Sklyanin, *Quantum spectral transform method: recent developments*, in: “Integrable Quantum Field Theories”, Lecture Notes in Phys. **151**, Springer, Berlin, 1982, pp. 61–119.
- [20] A. Molev, *Yangians and classical Lie algebras*, Mathematical Surveys and Monographs, 143. American Mathematical Society, Providence, RI, 2007.
- [21] A. Molev, *Sugawara operators for classical Lie algebras*. Mathematical Surveys and Monographs 229. AMS, Providence, RI, 2018.
- [22] M. L. Nazarov, *Quantum Berezinian and the classical Capelli identity*, Lett. Math. Phys. **21** (1991), 123–131.
- [23] M. Nazarov and V. Tarasov, *Yangians and Gelfand–Zetlin bases*, Publ. Res. Inst. Math. Sci. Kyoto Univ. **30** (1994), 459–478.
- [24] A. M. Perelomov and V. S. Popov, *Casimir operators for semisimple Lie algebras*, Izv. AN SSSR Ser. Mat. **32** (1968), 1368–1390.
- [25] N. Yu. Reshetikhin, *Quasitriangular Hopf algebras and invariants of links* (Russian), Algebra i Analiz **1** (1989), no. 2, 169–188; translation in Leningrad Math. J. **1** (1990), no. 2, 491–513.
- [26] N. Yu. Reshetikhin and M. A. Semenov-Tian-Shansky, *Central extensions of quantum current groups*, Lett. Math. Phys. **19** (1990), 133–142.
- [27] N. Yu. Reshetikhin, L. A. Takhtadzhyan and L. D. Faddeev, *Quantization of Lie groups and Lie algebras* (Russian), Algebra i Analiz **1** (1989), no. 1, 178–206; translation in Leningrad Math. J. **1** (1990), no. 1, 193–225.
- [28] V. O. Tarasov, *Cyclic monodromy matrices for $sl(n)$ trigonometric R -matrices*, Comm. Math. Phys. **158** (1993), 459–484.

N.J.:

Department of Mathematics
North Carolina State University, Raleigh, NC 27695, USA
jing@math.ncsu.edu

M.L.:

School of Mathematical Sciences
South China University of Technology
Guangzhou, Guangdong 510640, China
mamliu@scut.edu.cn

A.M.:

School of Mathematics and Statistics
University of Sydney, NSW 2006, Australia
alexander.molev@sydney.edu.au