# SYMMETRIC FUNCTIONS FROM THE SIX-VERTEX MODEL IN HALF-SPACE

#### ALEXANDR GARBALI, JAN DE GIER, WILLIAM MEAD AND MICHAEL WHEELER

ABSTRACT. We study the stochastic six-vertex model in half-space with generic integrable boundary weights, and define two families of multivariate rational symmetric functions. Using commutation relations between double-row operators, we prove a skew Cauchy identity of these functions. In a certain degeneration of the right-hand side of the Cauchy identity we obtain the partition function of the six-vertex model in a half-quadrant, and give a Pfaffian formula for this quantity. The Pfaffian is a direct generalization of a formula obtained by Kuperberg in his work on symmetry classes of alternating-sign matrices.

One of our families of symmetric functions admits an integral (sum over residues) formula, and we use this to conjecture an orthogonality property of the dual family. We conclude by studying the reduction of our integral formula to transition probabilities of the (initially empty) asymmetric simple exclusion process on the half-line.

## Contents

1. Introduction	1
2. Vertex model with boundaries	10
3. Row operators and symmetric functions	14
4. Triangular partition function	29
5. Integral formula for initially empty symmetric function	36
6. Open ASEP on the half-line	43
Compliance with ethical standards	46
Appendix A. Properties of Pfaffians	46
References	47

# 1. INTRODUCTION

1.1. **Background.** The connection between symmetric functions, exactly solvable vertex models and stochastic processes is a fertile branch of integrable probability, and is motivated by the rigorous study of the asymptotic behaviour of certain random variables within the KPZ universality class. Cornerstone results are the Macdonald processes [BC14] and their half-space analogues [BBC20], as well as the construction of probability measures on full-space from Cauchy identities of multivariate partition functions within the stochastic six-vertex model [Bor17, BCG16]. Generalizations to the setting of higher-spin [CP16, BP18] and higher-rank [BW22, BW20] models are also known, and have been very topical in recent literature.

The study of stochastic vertex models in half-space is less developed and comes with additional complications due the presence of a boundary. Some of the first algebraic results in this setting were obtained in [BBCW18], with application to rigorous asymptotics of the asymmetric simple exclusion process (ASEP) on the half-line, albeit for a rather restricted case of hopping rates at the boundary. More recently, the results of [BBCW18] were upgraded to fully generic boundaries in [He23]. In each of these works the interplay with symmetric functions is always close to the foreground, as both [BBCW18, He23] rely on symmetric function identities obtained in [BWZJ15] and [IMS22] to match half-space vertex-model expectations with quantities in the half-space and free-boundary Schur processes, respectively.

A much wider body of work exists in the context of the ASEP (recovered as a continuous-time limit of the stochastic six-vertex model [Agg16]) with open boundaries. The integrable structure of the ASEP on a strip (with open left and right boundaries), arising from commuting doublerow transfer matrices [Skl88], was reviewed in [CRV14] with a successive exploration of various integrable boundary conditions in [CFRV16]. The dynamic phase diagram based on the Bethe ansatz solution for the spectral gap of the open ASEP was computed in [dGE05]. A more recent approach to spectral gap analysis for the open ASEP uses high genus Riemann surface analysis of the Bethe equations [GP20]. Integrability of the classical open ASEP is relevant to quantum stochastic systems described by integrable Lindbladians [EP20].

Substantial literature has also been devoted to the stationary properties of the ASEP with two open boundaries. In the context of symmetric polynomials, the ASEP stationary state is related to Askey–Wilson polynomials [USW04, CW06] and the multivariate Koornwinder polynomials in the case of multi-species, or coloured, models [CGdGW16, FV17].

In the setting of the half-line (one non-trivial boundary), distribution functions can be accessed following the initial results of [TW13] for reflecting boundary conditions. Recent works include results for transition probabilities, boundary current fluctuation analysis for special boundary conditions [BBC20, BKD22, IMS22, He23] and Markov duality analysis [BC24]. While analyses of current fluctuations both at the boundary and in the bulk has been studied for last passage percolation [BBCS18].

The goal of the present text is to further develop our understanding of stochastic vertex models in half-space, viewed through the lens of multivariate symmetric functions. Stated in the simplest possible terms, our aim is to construct half-space analogues of the symmetric functions introduced by Borodin in [Bor17], and to study their algebraic properties (Cauchy summation identities, integral formulas and orthogonality). As mentioned above, classical symmetric functions (specifically, Hall– Littlewood and Schur polynomials) have already played an important role in probabilistic results in the half-space setting. In the current paper, however, our rational symmetric functions are new, and they are a key result in their own right.

We proceed to a detailed summary of our main results.

1.2. Six-vertex model with a boundary. Our primary algebraic tool is the *stochastic six-vertex* model. The interpretation of the six-vertex model as a Markov process in the quadrant dates back to a number of earlier works; see, for example, [GS92b, BCG16, BP18]. In the current text we will follow the same conventions as [BW22, Chapter 1].

We shall assign weights to finite collections of *paths* drawn on the square lattice. Each edge of the lattice (whether horizontal or vertical) supports at most one path, and the vertices obey the *ice rule*, which enforces that the total number of paths entering a vertex is the same as the total number of paths exiting it. Imposing the ice rule, one obtains six possible types of vertices:



where we have indicated the weight of each vertex underneath its picture. Here q denotes the quantum deformation parameter arising from the underlying  $U_q(\widehat{\mathfrak{sl}}_2)$  algebra, while z denotes the spectral parameter associated to the vertex. The weight of a lattice configuration is defined as the product of the weights of the individual vertices which comprise it.

The weights (1.1) have a number of well known fundamental properties. The first is that they obey the Yang–Baxter equation; see Proposition 2.5 of the text. This property endows the model with its rich algebraic structure and facilitates the exact computation of observables. The second is that the weights have a sum-to-unity property; see Proposition 2.2. This property allows one to assign probability measures to collections of paths in the square grid, and is the key feature that in turn allows reductions to one-dimensional stochastic processes such as the *asymmetric simple exclusion process*.

It turns out to be possible to extend both of the above properties, namely integrability and Markovian dynamics, to the setting of *boundary vertices*. A boundary vertex consists of a single incoming and outgoing edge, such that the two edges join to form a right angle. Once again, each edge of a boundary vertex supports at most one path. However, in contrast to the bulk vertices (1.1), we no longer enforce any ice rule on boundary vertices; this means that the total flux of paths through a boundary vertex need not be conserved. In the absence of an ice rule, there are two possible path assignments to the incoming/outgoing edges, leading to four types of boundary vertices:



where we have defined

$$h(x) = \frac{ac(1-x^2)}{(a-x)(c-x)}.$$
(1.3)

The weights (1.2) depend on a spectral parameter x, but unlike their bulk counterparts, they no longer have any q dependence. Instead, they acquire dependence on two boundary parameters a and c, whose values are free but the same for all boundary vertices.

One may verify that the weights (1.2), together with (1.1), satisfy the *reflection equation*, which is the boundary-analogue of the Yang–Baxter equation introduced by Sklyanin [Skl88]. It is also clear that these weights have a sum-to-unity property and are non-negative for certain mild choices of x and a, c, allowing one to maintain links with probability.

1.3. Double-row operators. A standard algebraic device in the setting of lattice models with a boundary is the *double-row transfer matrix*. Motivated by this, we introduce a family of double-row operators whose matrix elements are computed as partition functions in the model (1.1)-(1.2).

In the sequel, let  $\mu, \nu \in \mathbb{N}$  be finite subsets of the natural numbers<sup>1</sup>, and for each  $i \in \mathbb{N}$  introduce the indicator function  $\eta_i^{\mu} = \mathbf{1}_{i \in \mu}$  which assigns a value of 1 if i is an element of  $\mu$ , and 0 otherwise. We then construct the following partition function on a semi-infinite lattice:

$$A_{\mu \to \nu}(x) \coloneqq \begin{array}{c} \eta_{1}^{\mu} & \eta_{2}^{\mu} & \eta_{3}^{\mu} & \dots & \dots \\ \eta_{1}^{\mu} & \eta_{2}^{\mu} & \eta_{3}^{\mu} & \dots & \dots \\ & & & & & \\ \eta_{1}^{\mu} & \eta_{2}^{\mu} & \eta_{3}^{\mu} & \dots & \dots \\ & & & & & \\ \eta_{1}^{\mu} & \eta_{2}^{\mu} & \eta_{3}^{\mu} & \dots & \dots \end{array}$$
(1.4)

<sup>&</sup>lt;sup>1</sup>That is, both  $\mu$  and  $\nu$  have finitely many elements of finite size.

where the assignment of 1 to any edge of the lattice means that a path is present there, while the assignment of 0 means that it is vacant. The vertices used in the top row of (1.4) are given by the table (1.1), where in the *j*-th vertex in the row (read from left to right) we set  $z \mapsto xy_j$ . Similarly, the vertices used in the bottom row of (1.4) are given by (1.1) under 90° counterclockwise rotation, where in the *j*-th vertex in the row we set  $z \mapsto x/y_j$ . The boundary vertex appearing in (1.4) has weights given by (1.2). Note that  $A_{\mu \to \nu}(x)$  depends implicitly on the alphabet  $Y = (y_1, y_2, \ldots)$ , but we suppress this dependence in our notation where there is no potential for confusion.

One evaluates  $A_{\mu\to\nu}(x)$  by computing the weighted sum over all possible path configurations of the picture (1.4); that is, by computing it as a statistical mechanical partition function. Although we have defined  $A_{\mu\to\nu}(x)$  on a semi-infinite lattice, since  $\mu$  and  $\nu$  are finite subsets of N it is easily verified that sufficiently far from the boundary vertex one sees only empty vertices (devoid of paths). The weight of such vertices is 1, meaning that the quantity  $A_{\mu\to\nu}(x)$  is a finite sum of rational functions in  $x, y_i, q, a, c$ , for any fixed  $\mu, \nu$ .

In view of the sum-to-unity property of the bulk weights (1.1) and boundary weights (1.2), one finds that  $A_{\mu\to\nu}(x)$  obeys a sum-to-unity property:

$$\sum_{\nu \in \mathbb{W}} A_{\mu \to \nu}(x) = 1, \quad \text{for any fixed } \mu \in \mathbb{W}, \tag{1.5}$$

where  $\mathbb{W}$  denotes the set of finite subsets of  $\mathbb{N}$ . This fact allows us to view  $A_{\mu \to \nu}(x)$  as a Markov kernel and to generate a discrete-time Markov process of paths in half-space, similarly to what has been done in the context of the stochastic six-vertex model in a quadrant [BCG16, BP18].

It is then natural to construct an infinite-dimensional Markov matrix A(x) with entries  $A_{\mu \to \nu}(x)$ , where  $\mu$  is the row index and  $\nu$  the column index, which acts in the vector space obtained by taking the formal linear span of all finite subsets of  $\mathbb{N}$ . Our first result is that these operators commute:

**Proposition 1.1** (Proposition 3.7 below). Fix  $x_1, x_2 \in \mathbb{C}$  and assume there exists  $\rho > 0$  such that

$$\left|\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{q(1 - x_j / y_k)}{1 - q x_j / y_k}\right| \le \rho < 1,\tag{1.6}$$

for all  $1 \le i \ne j \le 2$  and for all  $k \in \mathbb{N}$ . One then has

$$A(x_1)A(x_2) = A(x_2)A(x_1),$$
(1.7)

where the latter identity holds in End(Span W).

The proof of Proposition 1.1 follows from use of the Yang-Baxter and reflection equations for our model. It proceeds along similar lines to the proof that double-row transfer matrices commute (see, for example, [Skl88]), however it requires a careful adjustment to the current setting where our operators act in the infinite-dimensional space Span W. The condition (1.6) is an artefact of the proof, and ensures that the product of operators  $A(x_i)A(x_j)$  converges.

1.4. Symmetric functions and Cauchy identities. With our commuting double-row operators in hand, there is a natural passage to the definition of a family of symmetric functions. In particular, for all  $\mu, \nu \in \mathbb{N}$  and a fixed alphabet  $(x_1, \ldots, x_L)$ , we define



which is again interpreted as a statistical mechanical partition function, with vertex weights given by (1.1) and (1.2). From its definition, it is clear that  $G_{\nu/\mu}(x_1, \ldots, x_L)$  may be interpreted as the concatenation of L discrete-time Markov kernels (1.4); expressing this algebraically via the double-row operators  $A(x_i)$ , we have that

$$G_{\nu/\mu}(x_1,\ldots,x_L) = \langle \mu | A(x_1) \cdots A(x_L) | \nu \rangle.$$
(1.9)

The functions (1.9) are a key focal point of this work<sup>2</sup>. In view of the commutativity (1.7) of the double-row operators that are used to define them, it is clear that the functions  $G_{\nu/\mu}(x_1,\ldots,x_L)$  are symmetric in  $(x_1,\ldots,x_L)$  (but for generic subsets  $\mu$  and  $\nu$  they do not exhibit symmetry in the  $(y_1, y_2, \ldots)$  alphabet). They also satisfy a sum-to-unity property (see Proposition 3.13) that is a direct consequence of (1.5).

Given a family of symmetric functions, a general goal is to produce another family which is *dual* to the first. In practice, this means that the original family is orthogonal to the dual one with respect to a certain inner product, or alternatively, the two families should pair together to yield a *Cauchy summation identity*. We are able to solve the latter of these two problems. In particular, for any subsets  $\mu, \nu \in \mathbb{N}$  and a fixed alphabet  $(z_1, \ldots, z_M)$ , we construct a second family of symmetric functions

$$F_{\mu/\nu}(z_1,\ldots,z_M) = \langle \mu | \dot{B}(z_1) \cdots \dot{B}(z_M) | \nu \rangle$$
(1.10)

where  $\dot{B}(z_i)$  denotes another kind of double-row operator (see Definition 3.2). As with the functions (1.9),  $F_{\mu/\nu}(z_1, \ldots, z_M)$  also depends on the secondary alphabet  $(y_1, y_2, \ldots)$ , but we suppress this from our notation. We then observe the following Cauchy identity between the two families (1.9) and (1.10):

**Theorem 1.2** (Theorem 3.14 below). Fix alphabets of complex numbers  $(x_1 \ldots, x_L), (z_1, \ldots, z_M)$ and assume there exists  $\rho > 0$  such that

$$\left|\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{q(1 - z_j/y_k)}{1 - q z_j/y_k}\right| \le \rho < 1, \qquad \left|\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{1 - q z_j y_k}{1 - z_j y_k}\right| \le \rho < 1, \tag{1.11}$$

for all  $1 \le i \le L, 1 \le j \le M$  and  $k \in \mathbb{N}$ . Then the partition functions (1.9) and (1.10) satisfy the skew Cauchy identity

$$\sum_{\kappa} G_{\kappa/\mu}(x_1,\ldots,x_L) F_{\kappa/\nu}(z_1,\ldots,z_M)$$

<sup>&</sup>lt;sup>2</sup>There is at least one other known instance of symmetric functions appearing from stochastic vertex models with a boundary; see [Zho22]. However, the weights that we use in the present work (both in the bulk and in the boundary) are more general and our results are otherwise disjoint from the work performed in [Zho22].

$$= \prod_{i=1}^{M} \prod_{j=1}^{L} \left[ \frac{x_j - qz_i}{x_j - z_i} \frac{1 - z_i x_j}{1 - qz_i x_j} \right] \sum_{\lambda} F_{\mu/\lambda}(z_1, \dots, z_M) G_{\nu/\lambda}(x_1, \dots, x_L). \quad (1.12)$$

where the sum on the left is over all finite subsets  $\kappa \subset \mathbb{N}$ , while the sum on the right is over subsets  $\lambda \subset \mathbb{N}$  whose elements do not exceed the maximal element of  $\mu$ .

The proof of Theorem 1.12 is by application of a non-trivial commutation relation between the operators  $A(x_i)$  and  $\dot{B}(z_j)$ ; see Proposition 3.9. As (1.12) holds for any fixed  $\mu$  and  $\nu$ , one may examine specific choices of these subsets such that the right hand side summation simplifies. One such choice is to take  $\mu = \emptyset$ , when the sum on the right hand side of (1.12) collapses to a single term, yielding the following corollary:

**Corollary 1.3** (Corollary 3.15 below). With the same set of assumptions as in Theorem 1.2, one has the summation identity

$$\sum_{\kappa} G_{\kappa}(x_1, \dots, x_L) F_{\kappa/\nu}(z_1, \dots, z_M) = \prod_{i=1}^M h(z_i) \prod_{i=1}^M \prod_{j=1}^L \left[ \frac{x_j - qz_i}{x_j - z_i} \frac{1 - z_i x_j}{1 - qz_i x_j} \right] G_{\nu}(x_1, \dots, x_L), \quad (1.13)$$

where we have introduced the abbreviation  $G_{\kappa}(x_1, \ldots, x_L) = G_{\kappa/\emptyset}(x_1, \ldots, x_L)$ , and where  $h(z_i)$  is given by (1.3).

Equation (1.13) is likely to have probabilistic utility. Note that, in view of the stochasticity of  $G_{\kappa}(x_1,\ldots,x_L)$ , one may view the left hand side of (1.3) as an expectation value  $\mathbb{E}(F_{\kappa/\nu})$ . Here  $\kappa$  is a random variable and  $F_{\kappa/\nu}(z_1,\ldots,z_M)$  is a family of observables in which both M and  $\nu$  are free. It would be interesting to explore the full range of observables that one may access through such a scheme, and to examine the types of explicit formulas that one obtains for these averages, via the right hand side of (1.13); similar approaches have previously been successfully followed in [BP18, BW20].

1.5. A half-space analogue of the domain wall partition function. Motivated by the quest for further simplifications of the Cauchy identity (1.13), we were led to consider the partition function  $G_{\nu/\mu}(x_1, \ldots, x_L)$  in which both  $\mu = \nu = \emptyset$ ; we denote this quantity  $G_{\emptyset}(x_1, \ldots, x_L)$  in the sequel. In this situation, no paths enter or exit via the external vertical edges of the lattice (1.8). At first glance, it may seem that this renders  $G_{\emptyset}(x_1, \ldots, x_L)$  trivial; this turns out not to be the case, since paths may be created/destroyed by the boundary vertices, and may thus trace out non-trivial configurations within the bulk of the lattice. Nevertheless, we do find an unexpected simplification of the function  $G_{\emptyset}(x_1, \ldots, x_L)$  that brings us into the realm of the six-vertex model in the half-quadrant.

**Theorem 1.4** (Theorem 3.23 below). Fix an integer L and an alphabet  $(x_1, \ldots, x_L)$  of complex parameters. Introduce the following *triangular partition function*:



where the vertex in the *i*-th column (counted from the left) and *j*-th row (counted from the bottom) is given by (1.1) with  $z \mapsto x_i x_j$ , and the boundary vertices are given by (1.2). One then has that

$$G_{\varnothing}(x_1,\ldots,x_L) = Z_L(x_1,\ldots,x_L). \tag{1.15}$$

In particular,  $G_{\emptyset}(x_1, \ldots, x_L)$  does not depend on the collection of vertical spectral parameters Y that are implicit in its definition (1.8).

Theorem 1.4 is proved by a non-obvious application of the Yang–Baxter equation, and initially came as a complete surprise to us. There is also a version of it that applies to the more general object  $G_{\nu}(x_1, \ldots, x_L)$ ; see Lemma 3.21 in the text. One of the main points of interest in these results is that they connect our probability measures on the half-line to six-vertex measures in the half-quadrant. The latter have been quite topical in recent years; see, for example, [BBCW18,He23].

Another point of interest is that (1.14) provides a natural two-parameter generalization of the partition function of off diagonally symmetric alternating-sign matrices (OSASMs), as introduced by Kuperberg in [Kup02]. In particular, [Kup02] gave a Pfaffian evaluation of the partition function (1.14), in a limit where only the middle two boundary vertices in (1.2) survive. One result of the current text is that this Pfaffian structure is preserved in the presence of four non-trivial boundary vertices<sup>3</sup>:

**Theorem 1.5** (Theorem 4.8 below). When L is even, the triangular partition function (1.14) admits the Pfaffian formula

$$Z_L(x_1, \dots, x_L) = \prod_{1 \le i < j \le L} \frac{1 - x_i x_j}{x_i - x_j} \cdot \Pr\left(\frac{x_i - x_j}{1 - x_i x_j} Q(x_i, x_j)\right)_{1 \le i, j \le L}$$
(1.16)

where Q is a symmetric function in two variables given by

$$Q(x_i, x_j) = (1 - h(x_i))(1 - h(x_j)) - \frac{h(x_i)h(x_j)}{ac} \frac{(1 - q)x_i x_j}{1 - qx_i x_j}.$$
(1.17)

A Pfaffian formula is also available in the case where L is odd, but we omit it here as its presentation is less elegant. The reduction of (1.16) to the OSASM partition function of [Kup02] is obtained by taking  $a \to 1, c \to -1$ ; note that in this limit one has that  $h(x) \to 1$ .

The proof of Theorem 1.5 proceeds by establishing a set of conditions that uniquely determine  $Z_L(x_1, \ldots, x_L)$  by Lagrange interpolation, and then showing that the right hand side of (1.16) satisfies these conditions. For the verification step, we show that  $Z_L(x_1, \ldots, x_L)$  may be recovered from the *shuffle exponential* of a certain linear combination of the partition functions  $Z_1$  and  $Z_2$ , assuming a *shuffle product* that we define; see Section 4.3. The connection with shuffle algebras is non-essential for our proof, but expedites the verification of our recursion relations significantly.

Combining Theorem 1.5 with Corollary 1.3 in the case  $\nu = \emptyset$ , we then have the following result:

**Corollary 1.6** (Corollary 4.10 below). With the same set of assumptions as in Theorem 1.2, one has the summation identity

$$\sum_{\kappa} G_{\kappa}(x_{1}, \dots, x_{L}) F_{\kappa}(z_{1}, \dots, z_{M}) = \prod_{i=1}^{M} h(z_{i}) \prod_{i=1}^{M} \prod_{j=1}^{L} \left[ \frac{x_{j} - qz_{i}}{x_{j} - z_{i}} \frac{1 - z_{i}x_{j}}{1 - qz_{i}x_{j}} \right] \times \prod_{1 \le i < j \le L} \frac{1 - x_{i}x_{j}}{x_{i} - x_{j}} \cdot \operatorname{Pf}\left(\frac{x_{i} - x_{j}}{1 - x_{i}x_{j}}Q(x_{i}, x_{j})\right)_{1 \le i, j \le L}$$
(1.18)

where  $F_{\kappa}(z_1, \ldots, z_M) = F_{\kappa/\emptyset}(z_1, \ldots, z_M)$  and Q is given by (1.17).

Corollary 1.6 has a very similar flavour to the refined Cauchy and Littlewood identities obtained in [BWZJ15, WZJ16], in which infinite summations of (products of) symmetric functions are evaluated as partition functions in the six-vertex model. As we have already mentioned, the left hand side of (1.18) admits the interpretation of an average; the right hand side of (1.18) provides a Pfaffian

<sup>&</sup>lt;sup>3</sup>After completion of this work we became aware that an analogous result was very recently also presented in [BFK23]; we thank Roger Behrend for bringing this to our attention.

evaluation of this average, which is likely to be valuable for asymptotic purposes. A similar scheme, in the full-space setting, was recently elaborated in [ABW23, Appendix C].

1.6. Integral formula for transition probabilities. Our next result concerns the evaluation of  $G_{\nu}(x_1, \ldots, x_L)$ , where  $\nu$  is non-empty. In this more general setting we lose the Pfaffian structure of  $G_{\emptyset}(x_1, \ldots, x_L)$ , but it still turns out to be possible to obtain a compact multiple-integral formula for the object in question.

**Theorem 1.7** (Theorem 5.4 below). In the special case  $\mu = \emptyset$ ,  $\nu = \{\nu_1 > \cdots > \nu_n \ge 1\}$ , the partition function (1.8) can be expressed as the following *n*-fold integral:

$$G_{\nu}(x_{1},\ldots,x_{L}) = \oint_{\mathcal{C}_{1}} \frac{\mathrm{d}w_{1}}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{C}_{n}} \frac{\mathrm{d}w_{n}}{2\pi \mathrm{i}} Z_{L+n} \left(x_{1},\ldots,x_{L},w_{1}^{-1},\ldots,w_{n}^{-1}\right) \\ \times \prod_{i=1}^{n} \prod_{j=1}^{L} \left[ \frac{qw_{i}-x_{j}}{w_{i}-x_{j}} \frac{1-w_{i}x_{j}}{1-qw_{i}x_{j}} \right]_{1 \leq i < j \leq n} \left[ \frac{w_{j}-w_{i}}{qw_{j}-w_{i}} \frac{1-qw_{i}w_{j}}{1-w_{i}w_{j}} \right] \\ \times \prod_{i=1}^{n} \left[ \frac{ac\left(qw_{i}^{2}-1\right)}{(w_{i}-a)(w_{i}-c)} \frac{y_{\nu_{i}}}{1-qw_{i}y_{\nu_{i}}} \prod_{j=1}^{\nu_{i}-1} \frac{1-w_{i}y_{j}}{1-qw_{i}y_{j}} \right], \quad (1.19)$$

where  $Z_{L+n}$  denotes the partition function (1.14) in (L+n) variables and  $C_1, \ldots, C_n$  are a certain family of q-nested contours that surround the points  $(x_1, \ldots, x_L)$  (for the precise definition of these contours, see Definition 5.3 and Figure 2).

Originally, we constructed a sum-over-subsets formula for  $G_{\nu}(x_1, \ldots, x_L)$  (see Theorem 5.1 below) by the application of *Drinfeld twists* to the columns of the partition function (1.8), similarly to what was done in [WZJ16]. Once obtained, it was then straightforward to match this sum-over-subsets formula to a sum-over-residues produced by evaluating the integral (1.19). While this is a totally direct method, the technical details are quite unwieldy; for this reason, we present instead a simpler (though non-constructive) verification argument, that once again relies on checking a set of recursive properties that uniquely determine  $G_{\nu}(x_1, \ldots, x_L)$ .

The remainder of the text is then devoted to studying the special case  $c \to \infty$  of (1.19), when the integrand becomes fully factorized. While this loses some of the generality of boundary vertex weights (under this limit, the third vertex in (1.2) vanishes) it has the advantage of making our formulas much more tractable without sacrificing the non-trivial injection of paths into the lattice (1.8) that still occurs in this regime.

1.7. An orthogonality conjecture. Orthogonality with respect to an integral scalar product was a key feature of the multivariate rational functions studied in [BP18,Bor17,BW22]. It seems likely that the symmetric functions considered in the current text also have nice orthogonality properties, and we find the first evidence of this in the following conjecture.

**Conjecture 1.8** (Conjecture 5.5 below). Fix a finite subset  $\nu = \{\nu_1 > \cdots > \nu_n \ge 1\}$ , and a second finite subset  $\kappa$  whose cardinality satisfies  $|\kappa| \le n$ . Then in the limit  $c \to \infty$ , one has that

$$\oint_{\mathcal{C}} \frac{\mathrm{d}w_1}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{C}} \frac{\mathrm{d}w_n}{2\pi \mathrm{i}} \prod_{1 \le i < j \le n} \left[ \frac{w_j - w_i}{qw_j - w_i} \frac{1 - qw_i w_j}{1 - w_i w_j} \right] \\ \times \prod_{i=1}^n \left[ \frac{w_i - a}{w_i (1 - aw_i)} \frac{1 - qw_i^2}{1 - w_i^2} \frac{y_{\nu_i}}{1 - qw_i y_{\nu_i}} \prod_{j=1}^{\nu_i - 1} \frac{1 - w_i y_j}{1 - qw_i y_j} \right] F_{\kappa}(w_1, \dots, w_n) = \delta_{\kappa, \nu}, \quad (1.20)$$

where the contour  $\mathcal{C}$  is a small, positively oriented circle surrounding the points  $y_j^{-1}$ ,  $j \ge 1$  and no other singularities of the integrand, such that  $q \cdot \mathcal{C}$  is disjoint from the interior of  $\mathcal{C}$ .

This conjecture has been tested extensively on examples with  $n \leq 3$ . It was motivated by considering the integral (1.19) at  $c \to \infty$ . In that limit, one may show that the triangular partition

function (1.14) factorizes:

$$\lim_{c \to \infty} Z_L(x_1, \dots, x_L) = \lim_{c \to \infty} \prod_{i=1}^L (1 - h(x_i)) = \prod_{i=1}^L \frac{x_i(1 - ax_i)}{x_i - a}.$$

This has two implications. On the one hand, the right side of the Cauchy identity (1.18) factorizes at this point; on the other hand, the integrand of (1.19) is itself factorized and one may recognize the Cauchy kernel embedded within it. Expanding the Cauchy kernel via the left side of (1.18) and collecting coefficients of  $G_{\kappa}(x_1, \ldots, x_L)$  on both sides of the equation, one deduces (1.20). We note that Conjecture 1.8 is true assuming the linear independence of the functions  $G_{\kappa}(x_1, \ldots, x_L)$ , but the proof of such a statement is outside of the scope of the present work.

1.8. Reduction to the ASEP on the half-line. It is well known that the stochastic six-vertex model converges, in a certain continuous-time limit, to the asymmetric simple exclusion process (ASEP); for a rigorous proof of this convergence in the full-space setting see, for example, [Agg16]. Our final result is an explicit integral formula for transition probabilities of the ASEP on the half-line, obtained by taking an appropriate degeneration of the integral expression (1.19).

In what follows, we consider the ASEP on the half-line with bulk hopping rates q and 1 associated to left and right moves, respectively, and incoming/outgoing hopping rates parameterized as

$$\alpha = \frac{ac(1-q)}{(1-a)(1-c)}, \qquad \gamma = -\frac{1-q}{(1-a)(1-c)},$$

respectively. For more details on the definition of this system, see Section 6.

**Theorem 1.9.** Let  $\mathbb{P}_t(\emptyset \to \nu)$  denote the probability that the ASEP on the half-line has particles at positions  $\nu \in \mathbb{N}$  at time t, given that it is initially empty. Under the limit  $c \to \infty$  ( $\gamma \to 0$ ), and assuming that  $\alpha + q \neq 1$ , one has that

$$\lim_{\gamma \to 0} \mathbb{P}_{t}(\emptyset \to \nu) = \alpha^{n} e^{-\alpha t} \oint_{\mathcal{D}_{1}} \frac{\mathrm{d}w_{1}}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{D}_{n}} \frac{\mathrm{d}w_{n}}{2\pi \mathrm{i}} \prod_{1 \le i < j \le n} \left[ \frac{w_{j} - w_{i}}{qw_{j} - w_{i}} \frac{1 - qw_{i}w_{j}}{1 - w_{i}w_{j}} \right] \\ \times \prod_{i=1}^{n} \left[ \frac{1 - qw_{i}^{2}}{w_{i}(q + \alpha - 1 - \alpha w_{i})(1 - qw_{i})} \left( \frac{1 - w_{i}}{1 - qw_{i}} \right)^{\nu_{i} - 1} \exp\left( \frac{(1 - q)^{2}w_{i}t}{(1 - w_{i})(1 - qw_{i})} \right) \right], \quad (1.21)$$

where the contours  $\mathcal{D}_1, \ldots, \mathcal{D}_n$  surround the point 1 and satisfy the conditions of Definition 6.3.

In the case of closed (or hard-wall) boundary conditions, integral formulas for the half-line ASEP were previously obtained in [TW13]. Our formula (1.21) appears to be new, however, as we deal with more generic boundary hopping rates than were considered in [TW13]. In particular, (1.21) allows the non-trivial injection of particles into the half-line (indeed, this is necessary to make this probability non-zero), whereas in [TW13] no particle may ever enter or exit the system.

1.9. Future prospects. A number of natural directions are suggested by the current work, and we plan to pursue at least a few of the following topics in later texts:

• The stochastic six-vertex model has a *higher-spin* analogue [BP18, Bor17, CP16] that may be obtained via the *fusion procedure* of [KRS81]. In this more general model, vertical lattice lines may now accommodate any number of paths, while horizontal lines still permit at most one path.

It is quite straightforward to push many of the formulas in the current text through the machinery of fusion, yielding a theory of higher-spin double-row symmetric functions. We expect that these functions will have many interesting properties, including simpler Cauchy identities and links with known families of functions, such as *BC*-symmetric Hall–Littlewood polynomials.

• In another direction, the stochastic six-vertex model may also be generalized to ensembles of *coloured paths*, following [BW22]. Algebraically, this corresponds to lifting the underlying quantum group  $U_q(\widehat{\mathfrak{sl}}_2)$  to  $U_q(\widehat{\mathfrak{sl}}_{n+1})$ , with  $n \ge 1$ . We expect that, in this coloured lattice model, our double-row operators will satisfy non-trivial commutation relations and that one should observe a family of *non-symmetric* multivariate rational functions that transform nicely under the action of the *Hecke algebra*. Evidence of such a construction, albeit in a slightly different model, has already been obtained in [Zho22].

• It is interesting to further explore the combinatorial implications of the Pfaffian formula (1.16). In view of the generic boundary vertex weights (1.2), the summation set of the partition function (1.14) is equivalent to *diagonally symmetric* alternating-sign matrices (with a generic diagonal), whose enumeration was unknown for twenty years since Kuperberg's work [Kup02] but very recently resolved in [BFK23].

1.10. Acknowledgments. We warmly thank Roger Behrend, Vadim Gorin, Leonid Petrov, Jeremy Quastel, Travis Scrimshaw and Ole Warnaar for helpful discussions. The authors gratefully acknowledge financial support of the Australian Research Council. AG was partially supported by the ARC DECRA DE210101264. WM was supported by an Australian Government Research Training Program Scholarship. MW was partially supported by the ARC Future Fellowship FT200100981.

# 2. Vertex model with boundaries

In this section we will outline a stochastic vertex model which will be the focus of much of this text. Explicit constructions of multi-parameter symmetric functions will be provided using these vertex models on the square lattice with non-trivial boundaries. These constructions have generic boundary parameters which will allow for a relation to the half-line open ASEP in Section 6.

2.1. Bulk vertex weights. Here we define the weights and relations of the vertex models that will be used throughout this work. First, we provide a definition of the stochastic six-vertex model [GS92a] and its  $U_q(\widehat{\mathfrak{sl}}_2)$  *R*-matrix. We will follow the conventions of [BW22].

Definition 2.1. The stochastic six-vertex model is an assignment of a rational function to a vertex

$$R_{y/x}(i,j;k,l) = \begin{array}{c} k \\ x \to j & & \\ & & \\ & & \\ & & \\ i \\ & & \\ y \end{array}; \quad i,j,k,l \in \{0,1\}.$$

$$(2.1)$$

At any given edge a 1 indicates the presence of a path while a 0 indicates an absence thereof. The values of the weights (2.1) are given pictorially in the following table where z = y/x. Any weight which does not appear in the table is defined to be equal to zero.



**Proposition 2.2** (Stochasticity). The weights of the stochastic six-vertex model satisfy a sum to unity property

$$\sum_{k,l \in \{0,1\}} R_z(i,j;k,l) = 1,$$
(2.3)

for any fixed  $i, j \in \{0, 1\}$ .

**Definition 2.3.** The *R*-matrix of the stochastic six-vertex model is  $R_{12}(z) \in \text{End}(V_1 \otimes V_2)$  for  $V_1, V_2 \cong \mathbb{C}^2$  given by

$$R_{12}(z) = \sum_{i,j,k,l \in \{0,1\}} R_z(i,j;k,l) E_1^{(j,l)} \otimes E_2^{(i,k)}, \qquad (2.4)$$

where  $E_m^{(i,j)} \in \text{End}(V_m)$  is the 2 × 2 elementary matrix with a 1 in row *i* and column *j*.

We will be interested in *R*-matrices acting on (possibly infinite-dimensional) spaces  $V_1 \otimes V_2 \otimes \cdots$ where each  $V_i \cong \mathbb{C}^2$ . We will denote by  $R_{jk}$  the *R*-matrix which acts non-trivially in the space  $V_j$ aligned horizontally and  $V_k$  aligned vertically as in the picture (2.1).

**Proposition 2.4** (Factorization). At the special value z = 1, the *R*-matrix (2.4) satisfies the relation  $R_{z=1}(i, j; k, l) = \delta_{i,l}\delta_{j,k}$ . This has pictoral representation

$$R_{x/x}(i,j;k,l) = \begin{array}{c} k & k \\ \uparrow & \uparrow \\ k & l \\ k & l \\ k & k \\ k & l \\ k & k \\ k &$$

**Proposition 2.5** (Yang–Baxter equation). The *R*-matrix (2.4) satisfies the Yang–Baxter equation  $P_{n}(u|v)P_{n}(v|v)P_{n}(v|v) = P_{n}(v|v)P_{n}(v|v)P_{n}(v|v)P_{n}(v|v)$ (2.6)

$$R_{12}(y|x)R_{13}(z|x)R_{23}(z|y) = R_{23}(z|y)R_{13}(z|x)R_{12}(y|x).$$
(2.6)

For fixed  $i_1, i_2, i_3, j_1, j_2, j_3 \in \{0, 1\}$ , this is represented pictorially as

$$\sum_{\substack{k_1,k_2,k_3\in\{0,1\}\\ x \to i_1 \\ y \to i_2 \\ k_1 \\ \vdots_2 \\ k_1 \\ k_2 \\ k_1 \\ k_3 \\ k_1 \\ k_3 \\ j_1 \\ k_1 \\ k_2 \\ k_1 \\ k_2 \\ k_2 \\ k_1 \\ k_2 \\ k_2 \\ k_1 \\ k_2 \\ k_3 \\ k_2 \\ k_1 \\ k_2 \\ k_2 \\ k_1 \\ k_2 \\ k_2 \\ k_2 \\ k_2 \\ k_2 \\ k_2 \\ j_1 \\ k_2 \\ k_$$

**Proposition 2.6** (*R*-matrix unitarity). The *R*-matrix (2.4) satisfies the unitarity condition  $R_{21}(x/y) R_{12}(y/x) = id,$ 

where id is the identity within  $\operatorname{End}(V_1 \otimes V_2)$ . For fixed  $i_1, i_2, j_1, j_2 \in \{0, 1\}$ , this is represented pictorially as



(2.7)

We will also define a re-normalized version of the weights from Definition 2.1. These new weights inherit many of the algebraic properties of the stochastic ones.

**Definition 2.7.** A *re-normalized vertex* is represented pictorially

$$\dot{R}_{y/x}(i,j;k,l) = \begin{array}{c} x \to j & & & \\ & & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\$$

The values of these weights are given in the following table, where z = y/x. Any weight which does not appear in the table is defined to be equal to zero.



These weights are related to their stochastic counterparts (2.1) through the relation

$$\dot{R}_z(i,j;k,l) = \frac{1-qz}{1-z} R_z(i,j;k,l).$$
(2.10)

Since the re-normalized vertices differ from the stochastic ones only by an overall multiplicative factor, we can write versions of the Yang–Baxter equation (2.6) what incorporate mixtures of both R and re-normalized  $\dot{R}$ -matrices.

2.2. Boundary vertex weights. We also define weights of a boundary vertex. Such boundary vertices and their matrices were introduced in [Che84, Skl88] and in a more general non-diagonal case in [DVGR93]. We consider the non-diagonal case in which the boundary vertex weights depend on two free parameters.

**Definition 2.8.** A stochastic boundary weight is an assignment of a rational function to a half-vertex

$$K_x(i;j) = \underbrace{j \rightarrow x^{-1}}_{i \leftarrow x}; \quad i, j \in \{0,1\}.$$

$$(2.11)$$

As previously, a 0/1 at a given edge indicates the absence/presence of a path. With the use of the rational function

$$h(x) = \frac{ac(1-x^2)}{(a-x)(c-x)},$$
(2.12)

these weights are tabulated below.



**Proposition 2.9** (Stochasticity). The stochastic boundary weights also satisfy a sum to unity property

$$\sum_{j \in \{0,1\}} K_x(i;j) = 1, \tag{2.14}$$

for any fixed  $i \in \{0, 1\}$ .

**Definition 2.10.** The *K*-matrix of the stochastic boundary weights is  $K_1(x) \in \text{End}(V_1)$  for  $V_1 \cong \mathbb{C}^2$  given by

$$K_1(x) = \sum_{i,j \in \{0,1\}} K_x(i;j) E_1^{(i,j)}.$$
(2.15)

Just as with the *R*-matrix, while acting on spaces  $V_1 \otimes V_2 \otimes \cdots$  we will denote by  $K_j$  the *K*-matrix which acts non-trivially on the space  $V_j$ . The boundary vertices (2.11), together with the bulk vertices (2.1), combine to give the reflection equation.

**Proposition 2.11** (Sklyanin reflection equation). The boundary vertex weights from Definition 2.8 and the bulk weights from Definition 2.1 satisfy the *Sklyanin reflection equation* 

$$R_{21}\left(\frac{x}{y}\right)K_1(x)R_{12}(xy)K_2(y) = K_2(y)R_{21}(xy)K_1(x)R_{12}\left(\frac{x}{y}\right).$$
(2.16)

For fixed  $i_1, i_2, j_1, j_2 \in \{0, 1\}$  this can be represented pictorially as



This relation is sometimes referred to as the boundary Yang-Baxter equation.

**Proposition 2.12** (*K*-matrix unitarity). The *K*-matrix from Definition 2.8 satisfies its own unitarity relation

$$K_1(x)K_1(x^{-1}) = \mathrm{id},$$
 (2.17)

where id is the identity within  $End(V_1)$ . This can be represented pictorially as



for all  $i, j \in \{0, 1\}$ .

# 3. Row operators and symmetric functions

3.1. Space of states and row-operators. In this section we construct double-row operators which serve as our transfer matrices. Our operators will act on the vector space with basis elements indexed by configurations in the set

$$\mathbb{W} = \left\{ S \subset \mathbb{N} : \sum_{i} S_i \text{ is finite} \right\}.$$

We denote a configuration  $\mu \in \mathbb{W}$  with  $m \ge 0$  parts by  $\mu = (\mu_1, \ldots, \mu_m)$  where  $\mu_m < \cdots < \mu_2 < \mu_1$ . By agreement  $\mu = \emptyset$  is defined when m = 0. We will only consider finite configurations  $\mu, \nu \in \mathbb{W}$ . That means that there are only finitely-many occupations in these states and that these occur at finite positions. We also define an orthogonal inner product on  $\mathbb{W}$  by  $\langle \mu | \nu \rangle = \delta_{\mu,\nu}$ .

**Definition 3.1** (Occupation notation). For  $\mu \in \mathbb{W}$  we define the occupation at site  $i \in \mathbb{N}$  as

$$\eta_i^{\mu} = \begin{cases} 1 & \text{if } i \in \mu \\ 0 & \text{otherwise} \end{cases}$$
(3.1)

**Definition 3.2.** A *double-row operator* on  $\mathbb{W}$  is defined by its action on co-vectors:

where  $x, z \in \mathbb{C}$  are horizontal spectral parameters and  $Y = (y_1, y_2, ...)$  is an infinite collection of vertical spectral parameters. For conciseness, we will often omit the family of vertical parameters from our notation by writing A(x|Y) = A(x) and  $\dot{B}(z|Y) = \dot{B}(z)$ .

**Proposition 3.3.** The empty state  $\emptyset \in \mathbb{W}$  corresponds to a left-eigenvector of the double-row operator (3.3)

$$\langle \varnothing | \dot{B}(z) = h(z) \langle \varnothing |, \qquad (3.4)$$

where h(z) is given by (2.12).

*Proof.* When the bottom state is empty in (3.3) there is only one possible configuration on the double-row. This single state is depicted as



which propagates the empty state from below to above the double-row. The weight of all bulk vertices in this picture are 1, while the boundary vertex has weight h(z).

A crucial property of the double-row operators of Definition 3.2 is their algebra of commutation and exchange relations. In order to prove these relations, we must first define a version of the double-row operators with finitely many columns.

**Definition 3.4.** For some fixed  $N \in \mathbb{N}$ , we define the monodromy matrices  $\mathcal{T}^{(N)}(x|Y)$  and  $\dot{\mathcal{T}}^{(N)}(z|Y)$ . The elements of these matrices are double-row row transfer matrices with N columns indexed by  $i, j \in \{0, 1\}$ . For fixed states  $\mu, \nu \in \mathbb{N}$  with  $\mu_1, \nu_1 \leq N$  these matrices are represented as



where  $x, z \in \mathbb{C}$  are horizontal spectral parameters and  $Y = (y_1, \ldots, y_N)$  is a collection of vertical spectral parameters. As with the definition of the double-row operators, we will omit the family of vertical spectral parameters from our notation by writing  $\mathcal{T}^{(N)}(x|Y) = \mathcal{T}^{(N)}(x)$  and  $\dot{\mathcal{T}}^{(N)}(z|Y) = \dot{\mathcal{T}}^{(N)}(z)$ . The elements of these matrices are represented as

$$\mathcal{T}^{(N)}(x) = \begin{pmatrix} A^{(N)}(x) & B^{(N)}(x) \\ C^{(N)}(x) & D^{(N)}(x) \end{pmatrix}, \quad \dot{\mathcal{T}}^{(N)}(z) = \begin{pmatrix} \dot{A}^{(N)}(z) & \dot{B}^{(N)}(z) \\ \dot{C}^{(N)}(z) & \dot{D}^{(N)}(z) \end{pmatrix}$$
(3.7)

For states  $\mu, \nu \in \mathbb{W}$ , we can recover the infinite column double row operators by

$$\lim_{N \to \infty} \langle \mu | A^{(N)}(x|Y) | \nu \rangle = \langle \mu | A(x|Y) | \nu \rangle, \qquad (3.8)$$

$$\lim_{N \to \infty} \langle \mu | \dot{B}^{(N)}(z|Y) | \nu \rangle = \langle \mu | \dot{B}(z|Y) | \nu \rangle, \qquad (3.9)$$

where the infinite column row-double row operators depend on the infinite collection of vertical parameters  $Y = (y_1, y_2, ...)$  and we regard the finite column operators as having dependence on the first N elements of the collection, i.e.  $(y_1, ..., y_N)$ .

**Proposition 3.5.** Fix  $x_1, x_2 \in \mathbb{C}$  and assume that there exists  $\rho > 0$  such that

$$\left|\frac{1 - x_1 y_k}{1 - q x_1 y_k} \frac{q(1 - x_2/y_k)}{1 - q x_2/y_k}\right| \le \rho < 1,\tag{3.10}$$

for all  $k \in \mathbb{N}$ . Then the following limit holds for states  $\mu, \nu \in \mathbb{W}$ 

$$\lim_{N \to \infty} \sum_{p \in \{0,1\}} \langle \mu | \mathcal{T}_{0,p}^{(N)}(x_1) \mathcal{T}_{p,0}^{(N)}(x_2) | \nu \rangle R_{x_1 x_2}(0,p;p,0) = \langle \mu | A(x_1) A(x_2) | \nu \rangle.$$
(3.11)

Which has a graphical interpretation



*Proof.* Consider  $\mu, \nu \in \mathbb{W}$  to be finite states. We denote their maximum occupation by  $\tau = \max\{\mu_1, \nu_1\}$ . Let N be an integer satisfying  $N > \tau$  which is also independent of  $\tau$ . By expanding the sum on the left side of (3.11) as

$$\lim_{N \to \infty} \langle \mu | A^{(N)}(x_1) A^{(N)}(x_2) | \nu \rangle + \frac{x_1 x_2 (1-q)}{1-q x_1 x_2} \lim_{N \to \infty} \langle \mu | B^{(N)}(x_1) C^{(N)}(x_2) | \nu \rangle.$$
(3.13)

The first term, which corresponds to p = 0, has the limit

 $\lim_{N\to\infty} \langle \mu | A^{(N)}(x_1) A^{(N)}(x_2) | \nu \rangle = \langle \mu | A(x_1) A(x_2) | \nu \rangle,$ 

which is our final result. So all that remains to prove is that the second term (p = 1) in (3.13) vanishes under the limit.

Since we are interested in the large N limit, we consider for  $N > \tau$  the partition function

Since  $N > \tau$ , it follows that  $\eta_i^{\mu}, \eta_i^{\nu} = 0$  for all  $\tau < i \le N$ . This freezes the columns to the right of the rectangle as shown in (3.14). The rectangle itself can be identified as a double row partition with  $\tau$  columns. By evaluating the frozen section, (3.14) is reduced to

$$\langle \mu | B^{(N)}(x_1) C^{(N)}(x_2) | \nu \rangle = \prod_{k=\tau+1}^{N} \left[ \frac{1 - x_1 y_k}{1 - q x_1 y_k} \frac{q(1 - x_2/y_k)}{1 - q x_2/y_k} \right] \langle \mu | B^{(\tau)}(x_1) C^{(\tau)}(x_2) | \nu \rangle.$$
(3.15)

Then with condition (3.10), we can bound (3.15) as

$$\left| \langle \mu | B^{(N)}(x_1) C^{(N)}(x_2) | \nu \rangle \right| \le \rho^{N-\tau} \left| \langle \mu | B^{(\tau)}(x_1) C^{(\tau)}(x_2) | \nu \rangle \right|,$$
(3.16)  
m of (3.13) vanishes as  $N \to \infty$ .

so that the p = 1 term of (3.13) vanishes as  $N \to \infty$ .

**Proposition 3.6.** Fix  $x, z \in \mathbb{C}$  and assume that there exists  $\rho > 0$  such that

$$\frac{1 - xy_k}{1 - qxy_k} \frac{q(1 - z/y_k)}{1 - qz/y_k} \le \rho < 1, \qquad \left| \frac{1 - xy_k}{1 - qxy_k} \frac{1 - qzy_k}{1 - zy_k} \right| \le \rho < 1, \tag{3.17}$$

for all  $k \in \mathbb{N}$ . Then the following limit holds

$$\lim_{N \to \infty} \sum_{\substack{p_1, p_2 \in \{0,1\}\\ p_1 \ge p_2}} \langle \mu | \mathcal{T}_{0,p_1}^{(N)}(x) \dot{\mathcal{T}}_{p_1 - p_2, 1 - p_2}^{(N)}(z) | \nu \rangle R_{xz}(0, p_1; p_1 - p_2, p_2) R_{z/x}(p_2, 1 - p_2; 0, 1)$$
$$= \frac{x - z}{x - qz} \langle \mu | A(x) \dot{B}(z) | \nu \rangle. \quad (3.18)$$



Figure 1. The terms of the sum (3.18) corresponding to  $(p_1, p_2) = (0,0), (1,0), (1,1)$  respectively.

Which has graphical interpretation



*Proof.* In a similar manner to the proof of Proposition 3.5, we fix  $\mu, \nu \in \mathbb{W}$  with maximum occupation  $\tau = \max\{\mu_1, \nu_1\}$ . We may write the terms in the sum over  $p_1, p_2$  explicitly as

$$\lim_{N \to \infty} \langle \mu | \left[ \frac{x-z}{x-qz} A^{(N)}(x) \dot{B}^{(N)}(z) + \frac{xz(1-q)}{1-qxz} \frac{x-z}{x-qz} B^{(N)}(x) \dot{D}^{(N)}(z) + \frac{1-xz}{1-qxz} \frac{x(1-q)}{x-qz} B^{(N)}(x) \dot{A}^{(N)}(z) \right] | \nu \rangle, \quad (3.20)$$

where each of the terms corresponds to  $(p_1, p_2) = (0, 0), (1, 0), (1, 1)$  respectively. Each of these three terms are depicted in Figure 1. We will analyse each term individually.

Firstly, the  $p_1 = p_2 = 0$  term has the limit

$$\lim_{N \to \infty} \left\langle \mu \right| \frac{x - z}{x - qz} A^{(N)}(x) \dot{B}^{(N)}(z) \left| \nu \right\rangle = \frac{x - z}{x - qz} \left\langle \mu \right| A(x) \dot{B}(z) \left| \nu \right\rangle$$

which is the desired result. So it remains to show that the other terms vanish in the limit.

Consider now the term associated to  $p_1 = 1, p_2 = 0$ . By virtue of the configurations being finite, this term can be decomposed into two possible configuration types. For some integer  $N > \tau$  which is independent of  $\tau$ , these configurations are summed over





where the dotted rectangles can are identified as stacked double row partition functions with  $\tau$  columns. The columns attached to the right of these rectangles in (3.21) can be explicitly evaluated as

$$\langle \mu | B^{(N)}(x) \dot{D}^{(N)}(z) | \nu \rangle = \prod_{k=\tau+1}^{N} \left[ \frac{1 - xy_k}{1 - qxy_k} \frac{q(1 - z/y_k)}{1 - qz/y_k} \right] \langle \mu | B^{(\tau)}(x) \dot{D}^{(\tau)}(z) | \nu \rangle$$

$$+ \prod_{k=\tau+1}^{N} \frac{1 - xy_k}{1 - qxy_k} \sum_{\ell=\tau+1}^{N} \frac{1 - q}{1 - qz/y_\ell} \frac{1 - q}{1 - zy_\ell} \prod_{k=\tau+1}^{\ell-1} \frac{1 - qzy_k}{1 - zy_k} \prod_{k=\ell+1}^{N} \frac{q(1 - z/y_k)}{1 - qz/y_k} \langle \mu | B^{(\tau)}(x) \dot{A}^{(\tau)}(z) | \nu \rangle.$$
(3.22)

Using condition (3.18), we can then bound this term effectively as

$$\left| \left\langle \mu | B^{(N)}(x) \dot{D}^{(N)}(z) | \nu \right\rangle \right| \leq \rho^{N-\tau} \left| \left\langle \mu | B^{(\tau)}(x) \dot{D}^{(\tau)}(z) | \nu \right\rangle \right|$$
  
+  $\rho^{N-\tau-1} (N-\tau) \max_{\ell \in \{\tau+1,\dots,N\}} \left\{ \left| \frac{1-xy_{\ell}}{1-qxy_{\ell}} \right| \right\} \max_{\ell \in \{\tau+1,\dots,N\}} \left\{ \left| \frac{1-q}{1-qz/y_{\ell}} \frac{1-q}{1-zy_{\ell}} \right| \right\} \left| \left\langle \mu | B^{(\tau)}(x) \dot{A}^{(\tau)}(z) | \nu \right\rangle \right|.$ (3.23)

We note here that the conditions (3.17) imply that, for all  $N > \tau$ , the points  $qxy_{\ell}, qz/y_{\ell}, zy_{\ell}$  are all bounded uniformly away 1 for all  $\ell \in \{\tau + 1, \ldots, N\}$ , and hence the maxima in the second term in (3.23) remain finite as  $N \to \infty$ , and therefore both terms in (3.23) vanish as  $N \to \infty$ .

It remains to show that the third term in (3.20) vanishes. This follows similarly as

$$\langle \mu | B^{(N)}(x) \dot{A}^{(N)}(z) | \nu \rangle = \prod_{k=\tau+1}^{N} \left[ \frac{1 - xy_k}{1 - qxy_k} \frac{1 - qzy_k}{1 - zy_k} \right] \langle \mu | B^{(\tau)}(x) \dot{A}^{(\tau)}(z) | \nu \rangle, \qquad (3.24)$$

which can be bounded using condition (3.18) as

$$\left| \langle \mu | B^{(N)}(x) \dot{A}^{(N)}(z) | \nu \rangle \right| \le \rho^{N-\tau} \left| \langle \mu | B^{(\tau)}(x) \dot{A}^{(\tau)}(z) | \nu \rangle \right|.$$
(3.25)

The limit of this term vanishes also, so we may conclude that only the  $p_1 = p_2 = 0$  term remains in the limit  $N \to \infty$  which implies the result.

**Proposition 3.7.** Fix  $x_1, x_2 \in \mathbb{C}$  and assume there exists  $\rho > 0$  such that

$$\left|\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{q(1 - x_j/y_k)}{1 - q x_j/y_k}\right| \le \rho < 1,$$
(3.26)

for all  $i \neq j$  and  $k \in \mathbb{N}$ . Then the double-row operators from (3.2) commute:

$$A(x_1)A(x_2) = A(x_2)A(x_1).$$
(3.27)

*Proof.* Let  $\mu, \nu \in \mathbb{W}$  with maximum occupation  $\tau = \max\{\mu_1, \nu_1\}$ . Also let N be an integer satisfying  $N \ge \tau$  and consider the double row partition functions with N columns



which, due to Proposition 3.5, has limit  $\langle \mu | A(x_1)A(x_2) | \nu \rangle$  as  $N \to \infty$ .

We may then append an additional intertwining vertex to the lattice after the last column. The boundary conditions of this vertex mean that there is only one allowed vertex configuration on the intertwiner so that it can be added at no overall cost to the partition function. We have the (3.28) is equal to



We may then repeatedly apply the Yang–Baxter equation (2.6) to manipulate the diagram. This equation can be applied successively to each column of partition function leading to the relation



At this point the reflection equation (2.16) can be applied, followed by the Yang–Baxter equation to push the intertwining vertices back to the right edge of the partition function. This yields



At this point, the intertwiner at the bottom-right of the diagram can be removed at no cost the partition function due to the boundary conditions. This yields



which we recognize as (3.28) with  $x_1$  and  $x_2$  interchanged. The limit of (3.30) can then be evaluated as  $\langle \mu | A(x_2)A(x_1) | \nu \rangle$  as  $N \to \infty$  due to condition (3.26). Since (3.28) and (3.30) are equal for all  $N \ge \tau$ , we can conclude that their limits must be equal. This is the result (3.27).

**Proposition 3.8.** Given configurations  $\mu, \nu \in \mathbb{W}$  let N be an integer  $N \ge \max\{\mu_1, \nu_1\}$ . For  $z_1, z_2$  the double-row operators with N columns commute

$$\langle \mu | \dot{B}^{(N)}(z_1) \dot{B}^{(N)}(z_2) | \nu \rangle = \langle \mu | \dot{B}^{(N)}(z_2) \dot{B}^{(N)}(z_1) | \nu \rangle.$$
(3.31)

This can be extended to the case of infinite columns to obtain the commutation relation of the double-row operators (3.3)

$$\dot{B}(z_1)\dot{B}(z_2) = \dot{B}(z_2)\dot{B}(z_1).$$
 (3.32)

*Proof.* Let  $\mu, \nu \in W$  with maximum occupation  $\tau = \max\{\mu_1, \nu_1\}$ . Also let  $N \ge \tau$  and consider the following double row partition functions with N columns



with two intertwining vertices appended to the right of the diagram. We note that these intertwiners are frozen in their own right and can simply be evaluated. For any  $N \ge \tau$  this yields

$$f^{(N)}(z_1, z_2) = \frac{1 - z_1 z_2}{1 - q z_1 z_2} \langle \mu | \dot{B}^{(N)}(z_1) \dot{B}^{(N)}(z_2) | \nu \rangle.$$
(3.34)

It is important to note here that this holds for all  $N \ge \tau$  here rather than in just under the large N limit as with the proof of Proposition 3.7.

Then following the same procedure as the proof of Proposition 3.7, we may apply the Yang–Baxter equation (2.6) and reflection equation (2.16) to manipulate the diagram (3.33) to obtain



The intertwiners on the right side of (3.35) are also frozen in their own right and can be evaluated as

$$f^{(N)}(z_1, z_2) = \frac{1 - z_1 z_2}{1 - q z_1 z_2} \langle \mu | \dot{B}^{(N)}(z_2) \dot{B}^{(N)}(z_1) | \nu \rangle.$$
(3.36)

The result (3.31) is obtained by comparing (3.34) and (3.36). Taking the large N limit yields the result (3.32).  $\Box$ 

**Proposition 3.9.** Fix  $x, z \in \mathbb{C}$  and assume there exists  $\rho > 0$  such that

$$\left|\frac{1 - xy_k}{1 - qxy_k}\frac{q(1 - z/y_k)}{1 - qz/y_k}\right| \le \rho < 1, \quad \left|\frac{1 - xy_k}{1 - qxy_k}\frac{1 - qzy_k}{1 - zy_k}\right| \le \rho < 1, \tag{3.37}$$

for all  $k \in \mathbb{N}$ . Then the double-row operators from Definition 3.2 obey the exchange relation

$$A(x)\dot{B}(z) = \frac{x - qz}{x - z} \frac{1 - xz}{1 - qxz} \dot{B}(z)A(x).$$
(3.38)

*Proof.* Let  $\mu, \nu \in \mathbb{W}$  with maximum occupation  $\tau = \max\{\mu_1, \nu_1\}$ . Also let N be an integer satisfying  $N \ge \tau$  and consider the double row partition functions with N columns



Due to condition (3.37) and Proposition 3.6, the limit of (3.39) is

$$\lim_{N \to \infty} f^{(N)}(x, z) = \frac{x - z}{x - qz} \langle \mu | A(x) \dot{B}(z) | \nu \rangle$$
(3.40)

We may manipulate the diagram of (3.39) to obtain an exchange relation in a similar way to the proof of Proposition 3.7.

By following the same steps as in the proof of Proposition 3.7, we may arrive at the following diagram which is equal to (3.39) as a partition function



The last equality follows from noticing that the intertwining vertices are both frozen by the boundary conditions, so they may be evaluated as a factor and removed from the diagram. The limit of the right diagram yields

$$\lim_{N\to\infty} f^{(N)}(x,z) = \frac{1-xz}{1-qxz} \langle \mu | \dot{B}(z) A(x) | \nu \rangle,$$

which can be combined with (3.40) to give the result.

3.2. Multi-parameter symmetric functions. We will now define a partition function which will be central to much the remainder of this work.

**Definition 3.10.** Fix two alphabets  $(x_1, \ldots, x_L)$  and  $(z_1, \ldots, z_M)$ , and configurations  $\mu, \nu \in \mathbb{W}$ . We define

$$G_{\nu/\mu}(x_1,\ldots,x_L|Y) = \langle \mu | A(x_1|Y) \cdots A(x_L|Y) | \nu \rangle, \qquad (3.42)$$

$$F_{\mu/\nu}(z_1,\ldots,z_M|Y) = \langle \mu | \dot{B}(z_1|Y)\cdots\dot{B}(z_M|Y) | \nu \rangle.$$
(3.43)

The functions (3.42) and (3.43) can be represented diagrammatically by stacking double-row operators (3.2) and (3.3) appropriately. We find



and



We will also, where convenient, omit the family of parameters Y from our notation.

The primary focus for the remainder of this work will be the partition function depicted in (3.44). This will ultimately be shown to reduce to describing the behaviour of the ASEP on the half-line with generic open boundary conditions.

In many cases we will be interested in the partition function (3.44) whose bottom state is empty, so that  $\mu = \emptyset$ . While for the partition function (3.45) we will often be interested in cases when the top state is empty, so that  $\nu = \emptyset$ . In such cases we will write

$$G_{\nu/\varnothing}(x_1, \dots, x_L | Y) \coloneqq G_{\nu}(x_1, \dots, x_L)$$
  
$$F_{\mu/\varnothing}(z_1, \dots, z_M | Y) \coloneqq F_{\mu}(z_1, \dots, z_M)$$

**Corollary 3.11** (of Propositions 3.7 and 3.8). Fix  $\mu, \nu \in \mathbb{W}$ . Given parameters  $x_1, \ldots, x_L \in \mathbb{C}$  and  $Y \in \mathbb{C}^{\mathbb{N}}$  all satisfying the conditions (3.26), the partition function  $G_{\nu/\mu}(x_1, \ldots, x_L|Y)$  is symmetric under permuting its *x*-alphabet. Given  $z_1, \ldots, z_M \in \mathbb{C}$ , the partition function  $F_{\mu/\nu}(z_1, \ldots, z_M|Y)$  is symmetric in permuting its *z*-alphabet.

**Proposition 3.12** (Branching relations). The partition functions from Definition 3.10 obey the branching relations

$$G_{\nu/\mu}(x_1,\ldots,x_{L+M}) = \sum_{\kappa} G_{\nu/\kappa}(x_{M+1},\ldots,x_{L+M}) G_{\kappa/\mu}(x_1,\ldots,x_M), \qquad (3.46)$$

$$F_{\mu/\nu}(z_1, \dots, z_{L+M}) = \sum_{\lambda} F_{\lambda/\nu}(z_{M+1}, \dots, z_{L+M}) F_{\mu/\lambda}(z_1, \dots, z_M).$$
(3.47)

*Proof.* This can be seen by inserting a sum over a complete set of states between the double-row operators in (3.42) and (3.43).

**Proposition 3.13.** Fix a configuration  $\mu \in \mathbb{W}$ . The partition function (3.42) obeys the sum-to-unity property

$$\sum_{\nu} G_{\nu/\mu}(x_1, \dots, x_L) = 1.$$
(3.48)

*Proof.* This follows from the stochasticity of the bulk and boundary vertices, Propositions 2.2 and 2.9.

3.3. Cauchy summation identity. In this section, we use the exchange relation (3.38) to prove an infinite summation identity of Cauchy type between the functions (3.42) and (3.43). This identity is the hint of a deeper orthogonality theory behind these functions that we plan to explore in a future text.

**Theorem 3.14.** Fix the alphabets  $(x_1 \ldots, x_L), (z_1, \ldots, z_M)$  and assume there exists  $\rho > 0$  such that

$$\left|\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{q(1 - z_j / y_k)}{1 - q z_j / y_k}\right| \le \rho < 1, \qquad \left|\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{1 - q z_j y_k}{1 - z_j y_k}\right| \le \rho < 1, \tag{3.49}$$

for all  $1 \leq i \leq L, 1 \leq j \leq M$  and  $k \in \mathbb{N}$ . Then the partition functions from Definition 3.10 satisfy the skew Cauchy identity

$$\sum_{\kappa} G_{\kappa/\mu}(x_1, \dots, x_L) F_{\kappa/\nu}(z_1, \dots, z_M)$$
  
= 
$$\prod_{i=1}^{M} \prod_{j=1}^{L} \left[ \frac{x_j - qz_i}{x_j - z_i} \frac{1 - z_i x_j}{1 - qz_i x_j} \right] \sum_{\lambda} F_{\mu/\lambda}(z_1, \dots, z_M) G_{\nu/\lambda}(x_1, \dots, x_L). \quad (3.50)$$

where the sum on the left is an infinite sum over  $\kappa \in \mathbb{W}$  while the sum on the right is a finite sum is over  $\lambda \in \mathbb{W}$  contained within  $\mu$ . That is,  $\lambda_i < \mu_i$  for all *i* less than the lengths of both  $\lambda$  and  $\mu$ .

*Proof.* We begin by writing the left-hand side of (3.50) in double-row operator notation as

$$\sum_{\kappa} G_{\kappa/\mu}(x_1,\ldots,x_L) F_{\kappa/\nu}(z_1,\ldots,z_M) = \langle \mu | A(x_1) \cdots A(x_L) \dot{B}(z_1) \cdots \dot{B}(z_M) | \nu \rangle.$$

From here we may commute the A operators through the  $\dot{B}$  operators using Proposition 3.9. Each commutation will generate a multiplicative rational factor; collecting all of these, we have

$$\langle \mu | A(x_1) \cdots A(x_L) \dot{B}(z_1) \cdots \dot{B}(z_M) | \nu \rangle = \prod_{i=1}^{M} \prod_{j=1}^{L} \left[ \frac{x_j - qz_i}{x_j - z_i} \frac{1 - z_i x_j}{1 - qz_i x_j} \right] \langle \mu | \dot{B}(z_1) \cdots \dot{B}(z_M) A(x_1) \cdots A(x_L) | \nu \rangle.$$
(3.51)

The right side of this may be recognized as the right-hand side of (3.50)

**Corollary 3.15.** With the same set of assumptions as in Theorem 3.14, one has the following summation identity

$$\sum_{\kappa} G_{\kappa}(x_1, \dots, x_L) F_{\kappa/\nu}(z_1, \dots, z_M) = \prod_{i=1}^M h(z_i) \prod_{i=1}^M \prod_{j=1}^L \left[ \frac{x_j - qz_i}{x_j - z_i} \frac{1 - z_i x_j}{1 - qz_i x_j} \right] G_{\nu}(x_1, \dots, x_L).$$
(3.52)

*Proof.* This is the  $\mu = \emptyset$  case of Theorem 3.14. Indeed, following the same steps as in the previous proof we use the fact, due to Proposition 3.3, that

$$\langle \varnothing | \dot{B}(z_1) \cdots \dot{B}(z_M) = \prod_{i=1}^M h(z_i) \langle \varnothing \rangle$$

in (3.51). This recovers precisely (3.52).

**Remark 3.16.** A further specification of Corollary 3.15 appears later in the text in Section 4.2.1 where  $\nu = \emptyset$ .

**Remark 3.17.** Subject to the positivity, Propositions 3.12 and 3.13 motivate the understanding of  $G_{\nu/\mu}$  as the propagator of a discrete-time Markov process from initial state  $\mu$  to state  $\nu$ . In order to define an appropriate probability measure, the spectral parameters  $(x_1, \ldots, x_L)$  and  $Y = (y_1, y_2, \ldots)$  need to be fixed so that the weights from table (2.2) and (2.13) are all real, non-negative and less than or equal to 1.

We may also regard the symmetric function F as an observable of this Markov process; indeed, the left-side of (3.52) can be interpreted as the formal definition of the expectation value of the observable  $F_{\kappa/\nu}$  with respect to the discrete measure  $G_{\kappa}$ . Provided that the right-hand side of (3.52) can be evaluated explicitly, Corollary 3.15 then provides a systematic method for evaluating the expectation value of the observable  $F_{\kappa/\nu}$ .

3.4. Recursion relations. In this section we demonstrate a series of recursion relations for the symmetric function (3.42) which will prove important in deriving a formula for the function. These relations follow from the unitary of the R and K-matrices (Propositions 2.6 and 2.12) and the R-matrix factorization (Proposition 2.4).

**Proposition 3.18.** We have the following relations for the row-operator (3.2):

$$A(0) = 0, (3.53)$$

$$A(\pm 1) = \mathsf{id},\tag{3.54}$$

where id is the identity within  $\operatorname{End}(\operatorname{Span} \mathbb{W})$ . Further, for fixed  $x \in \mathbb{C}$ , if there exists  $\rho > 0$  such that

$$\left|\frac{1 - xy_k}{1 - qxy_k} \frac{q(xy_k - 1)}{xy_k - q}\right| \le \rho < 1 \tag{3.55}$$

for all  $j \in \mathbb{N}$ , then it holds that

$$A(x)A(x^{-1}) = \mathsf{id}.$$
 (3.56)

*Proof.* We will proceed with the proof of each identity separately. In each proof, we will consider arbitrary finite configurations  $\mu, \nu \in \mathbb{W}$ .

(i) **Proof of** (3.53). It is sufficient here to consider the partition function

where we will be interested in the case x = 0. We note here that all contributions from the weights which make up (3.57) are non-singular at x = 0. In this case, the vertex configuration with weight z(1-q)/(1-qz) = 0 from (2.2) cannot appear. This means that the lower row of (3.57) cannot carry any occupations on any horizontal edges.

This will mean that the only possible boundary vertex configuration will be that of weight h(x) = 1. Since the weight z(1-q)/(1-qz) = 0 will not appear on the upper row, each horizontal edge on the upper row must be occupied.

However, this is not permitted since the boundary conditions on the right edge must be vacant. This means that there is no possible path configuration in (3.57). It follows that f(0) = 0, which gives the result.

(ii) **Proof of** (3.54). We will again consider (3.57), this time with  $x = \pm 1$ . In this case the function  $h(\pm 1) = 0$  so that the boundary vertex must have both entry and exits either both occupied or both vacant. Both of these configurations carry weight 1.

This is equivalent to the fact that that  $K(\pm 1)$  is the identity matrix. We are then left with



where we have made repeated use of the unitarity condition of *R*-matrices (Proposition 2.6), to produce the final equality. From here we may conclude that  $f(\pm 1) = 1$  which gives the result.

(iii) **Proof of** (3.56). Let  $\tau = \max\{\mu_1, \nu_1\}$  and let N be an integer satisfying  $N \ge \tau$ . Then for  $x \ne 0$ , consider



which is equal to  $\langle \mu | A(x)A(x^{-1}) | \nu \rangle$  in the limit  $N \to \infty$  from Proposition 3.5. Due to the factorization property of the *R*-matrix (Proposition 2.4), we may recognize that the intertwining vertex in (3.58) is the identity. We may simplify (3.58) as



where we have use the unitarity property of the *R*-matrix (Proposition 2.6) over the *N* columns to obtain the second diagram. Within this diagram, we may use the unitarity property of the *K*-matrix (Proposition 2.12) to simplify (3.59) as



from which we may again use *R*-matrix unitarity to remove all vertices. This shows that (3.66) is equal to  $\langle \mu | \nu \rangle$ , which yields the result under the limit  $N \to \infty$ .

The relations in Proposition 3.18 for the row-operators lead to recursion relations for the partition function (3.42).

**Corollary 3.19.** The partition function  $G_{\nu/\mu}(x_1, \ldots, x_L|Y)$  from (3.42) satisfies the following recursion relations

$$G_{\nu/\mu}(x_1,\ldots,x_L)\Big|_{x_i=0} = 0,$$
 (3.61)

$$G_{\nu/\mu}(x_1,\ldots,x_L)\Big|_{x_i=\pm 1} = G_{\nu/\mu}(x_1,\ldots,\hat{x}_i,\ldots,x_L),$$
(3.62)

$$G_{\nu/\mu}(x_1,\ldots,x_L)\Big|_{x_j=1/x_k} = G_{\nu/\mu}(x_1,\ldots,\hat{x}_j,\ldots,\hat{x}_k,\ldots,x_L),$$
(3.63)

for all  $1 \le i \le L$  and  $1 \le j < k \le L$ . Here,  $\hat{x}_i$  means that  $x_i$  is omitted from the alphabet  $(x_1, \ldots, x_L)$ . *Proof.* These follow from the row-operator identities, (3.53), (3.54), (3.56), in Proposition 3.18 as well as the fact that  $G_{\nu/\mu}$  is symmetric in its alphabet. **Corollary 3.20.** Let  $\mu, \nu \in \mathbb{W}$  and  $x_1, \ldots, x_L \in \mathbb{C} \setminus \{0\}$ . The partition function (3.42) satisfies the following unitarity property:

$$\sum_{\kappa} G_{\nu/\kappa} \left( x_1^{-1}, \dots, x_L^{-1} \right) G_{\kappa/\mu} (x_1, \dots, x_L) = \delta_{\mu,\nu}.$$
(3.64)

*Proof.* The proof follows from the branching relation (3.46), the recursion relation (3.63), and the fact that  $G_{\nu/\mu}$  evaluated on an empty alphabet is equal to  $\delta_{\mu,\nu}$ .

### 3.5. Empty initial conditions.

**Lemma 3.21.** Let  $(x_1, \ldots, x_L)$  be an alphabet satisfying conditions (3.26) for all  $i \neq j$ . The evaluation of the symmetric function indexed by a single configuration  $G_{\nu} = G_{\nu/\emptyset}$  from (3.42) reduces to the following partition function



*Proof.* Given an integer  $N \ge \nu_1$ , consider the N column version of the double-row picture (3.44). We may append a triangular arrangement of R-matrices onto the right-hand side of the lattice to obtain the following object



where the sum is over all occupations of the edges of the appended triangle. Absorbing the sum into the notation of the partition function, we can apply the Yang–Baxter equation to move the

intertwiners to be adjacent to the boundary. This is depicted as



We note that the left-moving sector of the vertical columns is frozen with no occupations. As a result, these vertices may be evaluated to 1 and removed from the diagram. In the limit  $N \to \infty$  this results in the desired diagram (3.65).

Now directly consider the large N limit of (3.66). Due to the conditions (3.26) the only term which survives this limit is the one with  $p_1 = \cdots = p_{2L-2} = 0$ . This forces all  $p_i = 0$  so that the sum of (3.66) collapses into a single term where all intertwining vertices are equal to 1. This yields

$$\lim_{N \to \infty} g_{\nu}^{(N)}(x_1, \dots, x_L) = \lim_{N \to \infty} \langle \emptyset | A^{(N)}(x_1) \cdots A^{(N)}(x_L) | \nu \rangle = G_{\nu}(x_1, \dots, x_L).$$
(3.68)

The diagram in (3.65) leads us to the following definition and theorem.

**Definition 3.22.** We refer to the following partition function, with generic boundary parameters, as the *triangular partition function*:



**Theorem 3.23.** When both the initial and final configurations are empty, i.e.  $\mu = \nu = \emptyset$ , the symmetric function (3.42) reduces to the triangular partition function (3.69)

$$G_{\varnothing}(x_1,\ldots,x_L) = Z_L(x_1,\ldots,x_L). \tag{3.70}$$

In particular,  $G_{\emptyset}$  does not depend on the collection of vertical spectral parameters Y.

*Proof.* This result is immediate from diagram (3.65) in Lemma 3.21 when we set  $\nu$  to be empty. This causes there to be no occupations on any of columns or rows past the point of  $y_i$ -dependence.

This means that the entire bulk on the right evaluates to 1 and may be removed without effect on the partition function evaluation.  $\hfill \Box$ 

# 4. TRIANGULAR PARTITION FUNCTION

In this section we turn our attention to the triangular partition function (3.69). An important feature of the six-vertex model with generic open boundary conditions is that the partition function  $G_{\emptyset}(x_1, \ldots, x_L)$  is non-trivial. Theorem 3.23 tells us that this partition function is equal to the triangular partition function. This function plays an analogous role to the domain wall partition function [Ize87] of the six-vertex model on a square geometry. In fact, this function directly generalizes a partition function related to diagonally symmetric alternating-sign matrices introduced in [Kup02].

We first note that for any  $m \in \mathbb{N}$ , the partition function  $Z_m(x_1, \ldots, x_m)$  is symmetric and satisfies the recursion relations of Corollary 3.19. It is instructive to derive these properties directly from (3.69) and we will do so below. Moreover, the triangular partition function is a rational function in its alphabet where the degree of numerator and denominator can be easily established. All these properties together completely determine  $Z_m(x_1, \ldots, x_m)$  and we shall find several closed formulas for it.

# 4.1. Properties and recursion relations.

**Proposition 4.1.** The triangular partition function  $Z_m(x_1, \ldots, x_m)$  from (3.69) is symmetric in  $(x_1, \ldots, x_m)$ .

*Proof.* Considering  $Z_m(x_1, \ldots, x_m)$ , the symmetry can be seen by inserting an intertwining vertex at the bottom of the diagram (3.69) to cross the adjacent lines with spectral parameters  $x_i$  and  $x_{i+1}$ . This can be done at no overall cost to the partition function since the boundary conditions enforce that the only possible vertex configuration is the all-empty one, which carries weight 1.

Using repeated applications of the Yang-Baxter equation (2.6) and reflection equation (2.16), the intertwining *R*-matrix can be pulled through to the right hand side of the lattice between the *i*-th and (i + 1)-th lines of the lattice. In the process of shifting this intertwiner, the  $x_i$  and  $x_{i+1}$ spectral parameters swap positions. The intertwiner can then be removed from the right hand side of the lattice at no overall cost, again enforced by the boundary conditions, leaving us with

$$Z_m(x_1,\ldots,x_i,x_{i+1},\ldots,x_m)=Z_m(x_1,\ldots,x_{i+1},x_i,\ldots,x_m),$$

which generates symmetry over the whole alphabet.

**Proposition 4.2.** The function defined by

$$\widetilde{Z}_m(x_1,\ldots,x_m) \coloneqq \prod_{i=1}^m \left[ (a-x_i)(c-x_i) \right] \prod_{1 \le i < j \le m} (1-qx_ix_j) \cdot Z_m(x_1,\ldots,x_m)$$
(4.1)

is a symmetric polynomial of degree at most m + 1 in each of the variables  $x_1, \ldots, x_m$ .

*Proof.* The pre-factors remove all possible denominators of bulk and boundary vertex weights of any lattice configuration, so that we can conclude that  $\tilde{Z}_m$  is a polynomial in all  $x_1, \ldots, x_m$ . Once the denominators are removed, the vertex on the boundary with argument  $x_i$  contributes a factor of power 2. Each of the m-1 bulk vertices also contribute a factor of power of at most 1, giving the leading order power of m+1.

**Proposition 4.3.** The triangular partition function (3.69) both satisfies, and is completely determined by, the following recursion relations

$$Z_m(x_1, \dots, x_m)\Big|_{x_i=0} = 0, \tag{4.2}$$

$$Z_m(x_1, \dots, x_m)\Big|_{x_i=\pm 1} = Z_{m-1}(x_1, \dots, \hat{x}_i, \dots, x_m),$$
(4.3)

$$Z_m(x_1, \dots, x_m) \Big|_{x_i = 1/x_j} = Z_{m-2}(x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_m).$$
(4.4)

$$\square$$

*Proof.* Because of Proposition 4.2, the triangular partition function  $Z_m$  is defined by a polynomial of degree m + 1 in each variable. The m + 2 recursions for each variable in the statement of Proposition 4.3 therefore completely determine the rational function  $Z_m$ .

We now prove the recursion relations individually.

(i) **Proof of** (4.2). We make use of the symmetry of the partition function and consider

$$Z_m(x_i, x_1, \ldots, \hat{x}_i, \ldots, x_m)\Big|_{x_i=0}$$

so that the boundary vertex corresponding to the parameter  $x_i$  is at the bottom-left-most position. Setting  $x_i = 0$  forces the bottom-left boundary vertex to generate a path with weight h(0) = 1, where we recall the definition of h(x) in (2.12). Since  $x_i = 0$ , the bottomright weight in (2.2) vanishes and hence this path cannot turn to any vertical edge on any vertex on the bottom line in the diagram (3.69). And so this path must proceed to the right hand side of the lattice. However, due to the imposed boundary conditions there cannot be any occupations on the external boundary edges on the right hand side and so we conclude that there are no allowed configurations when  $x_i = 0$  and hence that the partition function is equal to zero.

(ii) **Proof of** (4.3). Again making use of symmetry we consider

$$Z_m(x_i, x_1, \ldots, \hat{x}_i, \ldots, x_m)\Big|_{x_i=\pm 1}$$

This forces the boundary vertex at the bottom-left of the diagram in (3.69) to generate no paths and be of weight  $1 - h(\pm 1) = 1$ . This causes there to be no occupations along any edges of the bottom line, and hence this line may be removed at no cost leaving us with  $Z_{m-1}(x_1, \ldots, \hat{x}_i, \ldots, x_m)$ .

(iii) **Proof of** (4.4). Again using symmetry we consider

$$Z_m(x_i, x_j, x_1, \ldots, \hat{x}_i, \ldots, \hat{x}_j, \ldots, x_m)\Big|_{x_i=1/x_j}.$$

Using the R-matrix factorization, Lemma 2.4, the partition function becomes



Here, we have used the unitarity of the K-matrix (Proposition 2.12) to obtain the second diagram. We note here that the two bottom rows in the second diagram are completely frozen with no occupations and total weight 1, so that they can be removed at no cost to the partition function. This yields the result.

**Remark 4.4.** Another set of recursions can be derived for the numerator (4.1) of  $Z_m$ ,

$$\begin{aligned} \widetilde{Z}_m(x_1, \dots, x_m)|_{x_i=1/qx_j} &= -ac(1-q)q^{m-3}(1-x_j^2)(1-1/q^2x_j^2) \prod_{\substack{k=1\\k\neq i,j}}^m (1-x_k/qx_j)(1-x_kx_j) \\ &\times \widetilde{Z}_{m-2}(x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_m). \end{aligned}$$
(4.5)

These relations follow from the observation that the bottom-right vertex in diagram (3.69) completely freezes when  $qx_1x_m = 1$ , and as a consequence so does the bottom row and right-most column, leaving a partition function of size m - 2 multiplied by the pre-factors in (4.5) that arise from the weights of the frozen vertices. By symmetry a similar result follows for  $qx_ix_j = 1$  for any i and j.

4.2. Solution to recursion relations. This section provides solutions to the recursion relations of Proposition 4.3 which in turn provide closed form solutions to the triangular partition function.

Let  $Z_m^K$  be the partition function corresponding to the off-diagonal boundary conditions, which can be realized by setting a = -c = 1, set

$$Z_m^K(x_1, \dots, x_m) \coloneqq Z_m(x_1, \dots, x_m)|_{a = -c = 1}.$$
(4.6)

This partition function admits a Pfaffian formula due to Kuperberg [Kup02]:

$$Z_m^K(x_1, \dots, x_m) = \prod_{i=1}^m x_i \prod_{1 \le i < j \le m} \frac{1 - x_i x_j}{x_i - x_j} \operatorname{Pf} \left( M(x_i, x_j) \right)_{1 \le i, j \le m},$$
(4.7)

$$M(x_i, x_j) = \frac{(1-q)(x_i - x_j)}{(1 - x_i x_j)(1 - q x_i x_j)}.$$
(4.8)

We note that when m is odd the partition function  $Z_m^K$  vanishes.

**Theorem 4.5.** The triangular partition function (3.69) with the general boundary weights can be expressed as,

$$Z_m(x_1, \dots, x_m) = H_m(x) \sum_{r=0}^{\lfloor m/2 \rfloor} \left(\frac{1}{-ac}\right)^r \sum_{\substack{S \in [1,m] \\ |S| = 2r}} \prod_{i \in S} \frac{h(x_i)}{1 - h(x_i)} \prod_{\substack{i \in S \\ j \in S^c}} \frac{1 - x_i x_j}{x_i - x_j} Z_{2r}^K(x_S),$$
(4.9)

where

$$H_m(x) \coloneqq \prod_{i=1}^m (1 - h(x_i)), \qquad h(x) = \frac{ac(1 - x^2)}{(a - x)(c - x)}.$$
(4.10)

The proof of Theorem 4.5 is presented in the next section using shuffle algebra techniques.

**Corollary 4.6.** Expression (4.9) for  $Z_m$  as a sum over subsets can routinely be converted to a contour integration over a family of contours  $\mathcal{L}_i$  which all enclose each pole at  $x_1, \ldots, x_m$  but omit all other singularities of the integrand

$$Z_{m}(x_{1},...,x_{m}) = H_{m}(x) \sum_{r=0}^{m} \frac{1}{r!} \left(\frac{1}{-ac}\right)^{r/2} \oint_{\mathcal{L}_{1}} \frac{\mathrm{d}v_{1}}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{L}_{r}} \frac{\mathrm{d}v_{r}}{2\pi \mathrm{i}} \prod_{1 \le i < j \le r} \frac{v_{j} - v_{i}}{1 - v_{i}v_{j}} \times \prod_{i=1}^{r} \left(\frac{v_{i}h(v_{i})}{(1 - v_{i}^{2})(1 - h(v_{i}))} \prod_{j=1}^{m} \frac{1 - v_{i}x_{j}}{v_{i} - x_{j}}\right) \operatorname{Pf}\left(M(v_{k}, v_{\ell})\right)_{1 \le k, \ell \le r}.$$
 (4.11)

**Corollary 4.7.** For all  $m \ge 1$ , one has that

$$\lim_{c \to \infty} Z_m(x_1, \dots, x_m) = \lim_{c \to \infty} \prod_{i=1}^m (1 - h(x_i)) = \prod_{i=1}^m \frac{x_i(1 - ax_i)}{x_i - a}.$$
(4.12)

*Proof.* Examining the sum-over-subsets formula (4.9), it is easily verified that the limit  $c \to \infty$  eliminates all terms in the sum over r except that corresponding to r = 0. The claim (4.12) is then immediate.

Alternatively, one may prove (4.12) directly from the definition of the partition function (3.69), by noting that the  $c \to \infty$  limit causes the third vertex in the table (2.13) to vanish. Since the boundary vertices then only have the option to inject (but never eject) paths, and no paths exit the partition function (3.69) via its right-outgoing edges, it follows that the whole partition function is frozen as a product of empty vertices. The factorization (4.12) follows trivially.

The sum over subsets (4.9) can be compactly written in terms of Pfaffians in various ways. A particularly elegant expression is the following single Pfaffian expression for  $Z_m$ .<sup>4</sup>

**Theorem 4.8.** When m is even, the triangular partition function (3.69) with the generic open boundary weights can be expressed in terms of a Pfaffian,

$$Z_m(x_1, \dots, x_m) = \prod_{1 \le i < j \le m} \frac{1 - x_i x_j}{x_i - x_j} \cdot \Pr\left(\frac{x_i - x_j}{1 - x_i x_j} Q(x_i, x_j)\right)_{1 \le i, j \le m},$$
(4.13)

where Q is a symmetric function in two variables given by

$$Q(x_i, x_j) = (1 - h(x_i))(1 - h(x_j)) - \frac{h(x_i)h(x_j)}{ac} \frac{(1 - q)x_i x_j}{1 - qx_i x_j}.$$
(4.14)

*Proof.* Theorem 4.8 follows from Theorem 4.5 and the Pfaffian summation identity (A.3) in Lemma A.1.

**Remark 4.9.** In order to obtain an odd-sized solution from (4.13) we would write, for  $m = 2\ell$ ,

$$Z_{2\ell-1}(x_1,\ldots,x_{2\ell-1}) \coloneqq Z_{2\ell}(x_1,\ldots,x_{2l-1},1),$$

which makes use of the recursion relation (4.3).

We note that the Pfaffian kernel (4.14) bears some resemblance to the one appearing in a refined Littlewood identity for spin Hall–Littlewood symmetric rational functions [Gav23], though is quite different due to the boundary factors.

### 4.2.1. Cauchy summation identity revisited.

**Corollary 4.10.** Fix alphabets  $(x_1, \ldots, x_L), (z_1, \ldots, z_M)$  and assume that there exists  $\rho > 0$  such that conditions (3.49) hold. Then the following Cauchy summation identity holds

$$\sum_{\kappa} G_{\kappa}(x_{1}, \dots, x_{L}) F_{\kappa}(z_{1}, \dots, z_{M}) = \prod_{i=1}^{M} h(z_{i}) \prod_{i=1}^{M} \prod_{j=1}^{L} \left[ \frac{x_{j} - qz_{i}}{x_{j} - z_{i}} \frac{1 - z_{i}x_{j}}{1 - qz_{i}x_{j}} \right] \\ \times \prod_{1 \le i < j \le L} \frac{1 - x_{i}x_{j}}{x_{i} - x_{j}} \cdot \operatorname{Pf}\left( \frac{x_{i} - x_{j}}{1 - x_{i}x_{j}} Q(x_{i}, x_{j}) \right)_{1 \le i, j \le L}, \quad (4.15)$$

where Q is given by (4.14).

*Proof.* The proof follows by using Theorem 4.8 in the Cauchy summation identity of Corollary 3.15.

We note here the parallel of (4.15) to the refined Cauchy identity of Macdonald polynomials from [KN96], which is expressed as the product of the Macdonald Cauchy kernel and the Izergin–Korepin determinant in [War08].

4.3. Shuffle-exponential generating function. The partition function  $Z_m$  and its generating function

$$Z(v) \coloneqq \sum_{m=0}^{\infty} v^m Z_m, \tag{4.16}$$

can both be conveniently written in terms of a shuffle product.

**Definition 4.11.** Let  $f(x_1 \dots x_k)$  and  $g(x_1 \dots x_\ell)$  be two symmetric rational functions. We define the *shuffle product* f \* g to be the symmetric rational function given by

$$f \star g = \sum_{\substack{S \subseteq [1, k+\ell] \\ |S|=k}} f(x_S)g(x_{S^c}) \prod_{\substack{i \in S \\ j \in S^c}} \frac{1 - x_i x_j}{x_i - x_j}.$$
(4.17)

<sup>&</sup>lt;sup>4</sup>After completion of this work we became aware that an analogous formula was very recently also presented in [BFK23].

The identity with respect to the shuffle product is the rational symmetric function 1 in zero number of arguments and f \* 1 = 1 \* f = f. Further, for any rational function  $f(x_1 \dots x_k)$  we define the shuffle power and the shuffle exponential, exp<sub>\*</sub>, by

$$f^{*j} := \underbrace{f * f * \dots * f}_{j \text{ times}}, \qquad \exp_*(f) := 1 + f + \frac{1}{2!} f^{*2} + \frac{1}{3!} f^{*3} + \dots.$$
(4.18)

From Definition 4.11 it follows that the shuffle product of  $f(x_1 \dots x_k)$  and  $g(x_1 \dots x_l)$  is commutative unless both k and l are odd

$$f * g = (-1)^{kl} g * f. \tag{4.19}$$

It can also be easily shown that this shuffle product is associative (f \* g) \* h = f \* (g \* h). This shuffle product can be used to construct an algebra of functions and constitutes a convenient notation.

**Proposition 4.12.** Consider the first three triangular partition functions  $Z_0, Z_1$  and  $Z_2$ . These can be explicitly calculated from the the diagram (3.69) as

$$Z_0 = 1, \qquad Z_1 = 1 - h(x_1), \qquad Z_2 = (1 - h(x_1))(1 - h(x_2)) - \frac{h(x_1)h(x_2)}{ac} \frac{(1 - q)x_1x_2}{1 - qx_1x_2}, \qquad (4.20)$$

which are rational symmetric functions in 0,1 and 2 arguments x respectively. The generating function Z(v) takes form

$$Z(v) = \exp_{*}(v^{2}Z_{2} + vZ_{1}).$$
(4.21)

This exponential formula is equivalent to

$$Z_{2m} = \frac{1}{m!} Z_2^{*m}, \qquad Z_{2m+1} = \frac{1}{m!} Z_1 * Z_2^{*m}.$$
(4.22)

*Proof.* First we note that the two terms in the exponent in (4.21) commute with each other and  $Z_1^{*k} = 0$  for k > 1 due to (4.19). Applying definitions (4.18) to the generating function (4.21) leads to (4.22). Let us examine the expression for  $Z_{2m}$  given by (4.22). Computing  $Z_2^{*m}$  produces a rational function with the (minimal) denominator

$$\prod_{i=1}^{2m} (a - x_i)(c - x_i) \prod_{1 \le i < j \le 2m} (1 - qx_i x_j).$$
(4.23)

This denominator is a polynomial of degree 2m + 1 in the individual  $x_l$ , l = 1, ..., 2m. Let us fix l and show that the limit  $x_l \to \infty$  of  $Z_2^{*m}$  exists. By writing  $Z_2^{*m}$  using (4.18) and (4.17) we can see that  $Z_2^{*m}$  is of the form

$$Z_2^{*m} = \sum \cdots Z_2(x_{i_1}, x_{i_2}) \cdots Z_2(x_{j_1}, x_{j_2}) \cdots \times \prod_{a, b=1, 2} \frac{1 - x_{i_a} x_{j_b}}{x_{i_a} - x_{j_b}} \times \cdots,$$
(4.24)

therefore in each summand the dependence on  $x_{\ell}$  is of the form

$$Z_2(x_\ell, x_a) \prod_b \frac{1 - x_\ell x_b}{x_\ell - x_b},$$

where a and b are some indices not equal to l. Computing  $x_{\ell} \to \infty$  in both of these factors shows that this limit exists. Therefore  $Z_2^{*m}$  is given by a ratio of a polynomial of degree at most 2m + 1in  $x_{\ell}$  and the polynomial in (4.23). This implies that in order to prove (4.22) for  $Z_{2m}$  we need to show that  $Z_2^{*m}/m!$  satisfies the recursion relations of Proposition 4.3.

The specializations given in (4.2) and (4.3) follow from

$$Z_1(0) = 0, \qquad Z_2(x,0) = Z_2(0,x) = 0, \qquad (4.25)$$

$$Z_1(1) = Z_0 = 1, \quad Z_2(x,1) = Z_2(1,x) = Z_1(x).$$
(4.26)

We have (4.22) satisfies (4.2) due to (4.25) and it satisfies (4.3) due to (4.26). Consider next (4.4) and set  $x_k = 1/x_\ell$  in  $Z_2^{*m}$  as written in (4.24) for any distinct  $k, \ell = 1...2m$ . In each term of the sum in (4.24) the arguments  $x_1 \ldots x_{2m}$  are distributed between various factors  $Z_2$ . Considering a generic summand we encounter two cases: either  $k \in \{i_1, i_2\}$  and  $\ell \in \{j_1, j_2\}$  or  $k, \ell \in \{i_1, i_2\}$ . In the

first case the contribution is zero because of the factor which is explicitly written in (4.24) and in the second case we compute

$$\frac{1}{m!} Z_2^{*m}|_{x_k=1/x_\ell} = \frac{1}{m!} \sum \cdots Z_2 \left( x_\ell, \frac{1}{x_\ell} \right) \cdots Z_2 (x_{j_1}, x_{j_2}) \cdots \times \prod_{b=1,2} \frac{1 - x_\ell x_{j_b}}{x_\ell - x_{j_b}} \frac{1 - \frac{1}{x_\ell} x_{j_b}}{\frac{1}{x_\ell} - x_{j_b}} \times \cdots \\ = \frac{1}{(m-1)!} \left( Z_2^{*(m-1)} \right) (\dots, \hat{x}_k, \dots, \hat{x}_\ell, \dots),$$

where we noted that  $Z_2(x_l, 1/x_l) = 1$  and the explicitly written rational function is also equal to 1. There are in total *m* different summands for which  $k, \ell \in \{i_1, i_2\}$ . All these summands are equal to each other and to the symmetric function  $Z_2^{*(m-1)}$  which depends on  $x_1, \ldots, x_{2m}$  with  $x_k, x_\ell$  omitted. These computations show that  $Z_{2m}$  given by (4.22) satisfies the conditions of Proposition 4.3. The case of  $Z_{2m+1}$  can be proven analogously.

**Corollary 4.13.** The summation formula for  $Z_m$  given in Theorem 4.5 holds as a consequence of (4.22).

*Proof.* We note that  $Z_2$  in (4.20) is given by a sum of two terms and therefore (4.22) can be expanded using the binomial theorem

$$Z_{2m} = \frac{1}{m!} \left( (1 - h(x_1))(1 - h(x_2)) + \frac{-1}{ac} h(x_1) h(x_2) Z_2^K(x_1, x_2) \right)^{*m}$$
  
=  $\sum_{r=0}^m \left( \frac{-1}{ac} \right)^r \frac{1}{(m-r)!r!} \left( (1 - h(x_1))(1 - h(x_2)) \right)^{*(m-r)} * \left( h(x_1) h(x_2) Z_2^K(x_1, x_2) \right)^{*r}.$ 

The two terms given by the shuffle powers \*(m-r) and \*r can be computed. For example, the second term is computed by observing that  $(Z_2^K)^{*r} = r!Z_{2r}^K$  as a consequence of (4.22) and (4.6). After this we can write the shuffle product of these two terms using (4.11) and match the outcome with (4.9).

4.4. Alternative form of solution to recursion relations. Theorem 4.14 below contains an alternative explicit expression for the triangular partition function  $Z_m(x_1, \ldots, x_m)$  in terms of subsetsums over factorized expressions and valid for both m even and odd. We first define S by

$$S(x_i, x_j) = \frac{x_i - x_j}{1 - x_i x_j},$$
(4.27)

and let

$$Q^{e}(x_{i}, x_{j}) = S(x_{i}, x_{j}) + \frac{u^{2}q^{1/2}}{ac} x_{i}x_{j}h(x_{i})h(x_{j}) S\left(q^{1/2}x_{i}, q^{1/2}x_{j}\right),$$

$$Q^{o}(x_{i}, x_{j}) = x_{i}x_{j}S(x_{i}, x_{j}) + \frac{u^{2}}{q^{1/2}ac}h(x_{i})h(x_{j})S\left(q^{1/2}x_{i}, q^{1/2}x_{j}\right),$$
(4.28)

where u is a generating parameter. Furthermore, we define the following functions in terms of Pfaffians

$$Z_{2m}^{e}(u;x_{1},\ldots,x_{2m}) = \prod_{1 \le i < j \le 2m} \frac{1-x_{i}x_{j}}{x_{i}-x_{j}} \cdot \operatorname{Pf}\left(Q_{2m}^{e}(x_{i},x_{j})\right)_{1 \le i,j \le 2m},\tag{4.29}$$

$$Z_{2m-1}^{o}(u; x_1, \dots, x_{2m-1}) = \prod_{1 \le i < j \le 2m-1} \frac{1 - x_i x_j}{x_i - x_j} \cdot \Pr\left( \begin{pmatrix} (Q_{2m-1}^{o}(x_i, x_j))_{1 \le i < j \le 2m-1} & (-uh(x_i))_{1 \le i \le 2m-1} \\ (uh(x_j))_{1 \le j \le 2m-1} & 0 \end{pmatrix} \right).$$

$$(4.30)$$

We also set

$$Z_{2m-1}^{e}(u; x_1, \dots, x_{2m-1}) = Z_{2m}^{e}(u; x_1, \dots, x_{2m-1}, 1),$$
  

$$Z_{2m}^{o}(u; x_1, \dots, x_{2m}) = Z_{2m+1}^{o}(u; x_1, \dots, x_{2m}, 1).$$
(4.31)

Using these definitions we can now state the following theorem and corollary.

**Theorem 4.14.** The triangular partition function (3.69) is recovered by  $Z_m(x_1, \ldots, x_m) = Z_m(u = 1; x_1, \ldots, x_m)$  for  $m \in \mathbb{N}$  with

$$Z_m(u; x_1, \dots, x_m) = Z_m^{\rm e}(u; x_1, \dots, x_m) + Z_m^{\rm o}(u; x_1, \dots, x_m).$$
(4.32)

Furthermore, the partition function with generating parameter,  $Z_m(u; x_1, \ldots, x_m)$ , can be written as

$$Z_m(u;x_1,\ldots,x_m) = \sum_{S \subseteq [1,m]} (-u)^{|S|} g_S(x) \prod_{i \in S} h(x_i) \prod_{i \in S} \prod_{j \in S^c} \frac{x_i x_j - 1}{x_i - x_j} \prod_{\substack{1 \le i < j \le m \\ i, j \in S}} \frac{1 - x_i x_j}{1 - q x_i x_j},$$
(4.33)

where  $r_S = \lfloor |S|/2 \rfloor$  and

$$g_S(x) = \frac{q^{r_S^2}}{(ac)^{r_S}} \begin{cases} \prod_{i \in S} x_i, & |S| \text{ is even} \\ \prod_{i \in S^c} x_i, & |S| \text{ is odd} \end{cases}$$

**Remark 4.15.** The recursions (4.5) appear in (4.33) as residues of the simple poles at  $x_k = 1/qx_\ell$ .

Proof of Theorem 4.14. The equivalence of (4.32) and (4.33) follows in a straightforward manner from the Pfaffian definitions of  $Z_m^{\rm e}$  and  $Z_m^{\rm o}$ , the Pfaffian identity (A.1) and from the fact that the Pfaffian of S factorizes [Ste90],

$$Pf(S(x_i, x_j))_{1 \le i, j \le 2m} = \prod_{1 \le i < j \le 2m} \frac{x_i - x_j}{1 - x_i x_j}.$$
(4.34)

Next we need to show that  $Z_m(x_1, \ldots, x_m) = Z_m(u = 1; x_1, \ldots, x_m)$ . We do that by computing the generating function

$$Z(u;v) \coloneqq \sum_{m=0}^{\infty} v^m Z_m(u;x_1,\ldots,x_m)$$

at u = 1 with  $Z_m(u = 1; x_1, ..., x_m)$  given by (4.33). We will show that this generating function is equal to the generating function of  $Z_m(x_1, ..., x_m)$  (4.21) from Proposition 4.12. Using the definition of the shuffle product (4.11) we rewrite (4.33) with u = 1 as

$$Z_{2m}(u = 1; x_1, \dots, x_{2m}) = \sum_{j=0}^{2m} V_{2m-j} * W_j^{e},$$

$$Z_{2m+1}(u = 1; x_1, \dots, x_{2m+1}) = \sum_{j=0}^{2m+1} V_{2m+1-j} * W_j^{o},$$
(4.35)

where we introduced symmetric functions  $V_m = V_m(x_1, \ldots, x_m)$ 

$$V_{2m} = \frac{q^{m^2}}{(ac)^m} \prod_{i=1}^{2m} x_i h(x_i) \prod_{1 \le i < j \le 2m} \frac{1 - x_i x_j}{1 - q x_i x_j},$$

$$V_{2m+1} = -\frac{q^{m^2}}{(ac)^m} \prod_{i=1}^{2m+1} h(x_i) \prod_{1 \le i < j \le 2m+1} \frac{1 - x_i x_j}{1 - q x_i x_j},$$
(4.36)

and  $W_m^{\mathrm{e/o}} = W_m^{\mathrm{e/o}}(x_1, \ldots, x_m)$  are defined by

$$W_{2m}^{e} = 1, \qquad W_{2m+1}^{e} = -x_1 \cdots x_{2m+1}, \qquad W_{2m}^{o} = x_1 \cdots x_{2m}, \qquad W_{2m+1}^{o} = 1.$$
 (4.37)

All of these functions also factorize with respect to the shuffle product

$$X_{2m} = \frac{1}{m!} X_2^{*m}, \qquad X_{2m+1} = \frac{1}{m!} X_1 * X_2^{*m}, \qquad \text{for} \quad X = V, W^e, W^o.$$
(4.38)

From these formulas it follows that the generating functions of  $V, W^e, W^o$  can be expressed in terms of the shuffle exponential (4.18). We compute the generating function Z(u = 1; v)

$$Z(u=1;v) = \sum_{m=0}^{\infty} v^{2m} Z_{2m}(u=1;x_1,\ldots,x_{2m}) + \sum_{m=0}^{\infty} v^{2m+1} Z_{2m+1}(u=1;x_1,\ldots,x_{2m+1}),$$
(4.39)

using (4.35) and by representing each function  $V_k, W_j^{e/o}$  in the form (4.38). The first summand in (4.39) is computed as follows

$$\sum_{m=0}^{\infty} v^{2m} Z_{2m} (u = 1; x_1, \dots, x_{2m}) = (1 + v^2 V_1 * W_1^e) * \exp_* (v^2 (V_2 + W_2^e))$$
$$= \exp_* (v^2 (V_2 + W_2^e + V_1 * W_1^e)) = \exp_* (v^2 Z_2), \quad (4.40)$$

where  $Z_2 = Z_2(x_1, x_2)$  in the last expression is the triangular partition function for two sites. In (4.40) the second equality is due to the nilpotency of the shuffle product and the third equality is a consequence of the identity

$$V_2 + W_2^{\rm e} + V_1 * W_1^{\rm e} = Z_2. \tag{4.41}$$

Let us remark that the numerator on the right hand side of (4.41) is a polynomial of degree 3 in each  $x_i$  while on the left hand side some terms have numerators which are polynomials of degree 4 in individual  $x_i$ . In the above equation it is easy to check that the degree 4 terms cancel on the left hand side. This phenomenon manifests itself if one tries to evaluate the degrees produced by the formula (4.33). The apparent degree is higher than expected and, in order to show the connection with  $Z_m$ , it is required to argue that (4.33) actually produces the correct degree.

In the next step we calculate the generating function of the second term in (4.39)

$$\sum_{m=0}^{\infty} v^{2m+1} Z_{2m+1} (u = 1; x_1, \dots, x_{2m+1}) = v \left( V_1 + W_1^o \right) * \exp_* \left( v^2 \left( V_2 + W_2^o \right) \right)$$
$$= v Z_1 * \exp_* \left( v^2 \left( Z_2 + Z_1 * x_1 \right) \right) = v Z_1 * \exp_* \left( v^2 Z_2 \right)$$
(4.42)

where  $Z_1 = Z_1(x_1)$  and  $Z_2 = Z_2(x_1, x_2)$  in the second line are the triangular partition functions for one and two sites. In the second equality in (4.42) we used

$$V_1 + W_1^{o} = Z_1$$
  $V_2 + W_2^{o} = Z_2 + Z_1 * x_1$ 

and the last equality of (4.42) is due to the nilpotency  $Z_1^{*n} = 0$ , n > 1. By combining (4.42) with (4.40) in (4.39) we obtain the full generating function

$$Z(u = 1; v) = \exp_*(v^2 Z_2 + v Z_1).$$

which coincides with (4.21) and therefore proves the statement of the Theorem.

### 5. INTEGRAL FORMULA FOR INITIALLY EMPTY SYMMETRIC FUNCTION

The central objects of this work are the two symmetric functions of Definition 3.10. Theorem 3.23 shows that the function  $G_{\nu/\mu}$  reduces to the triangular partition function when both bottom and top configurations are empty. The previous section demonstrates how even when both conditions are empty this symmetric function is highly non-trivial. In this section we provide more insight into this behaviour by providing two equivalent evaluations of  $G_{\nu/\mu}$  for arbitrary  $\nu$  from an empty  $\mu = \emptyset$ . The form of this function leads to a striking conjecture on the orthogonality of the dual family  $F_{\kappa}$ .

#### 5.1. Subset formula.

**Theorem 5.1.** Let the state on the bottom be empty while the arbitrary state on top  $\nu = (\nu_1, \ldots, \nu_n)$  consist of *n* occupations at positions finitely far from the origin. Assume that that  $L \ge n$  and there exists  $\rho > 0$  such that

$$\left|\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{q(1 - x_j/y_k)}{1 - q x_j/y_k}\right| \le \rho < 1,$$

for all  $1 \le i \ne j \le L, k \in \mathbb{N}$ . Then the partition function (3.42) is calculated explicitly as

$$G_{\nu}(x_{1},...,x_{L}|Y) = \sum_{\substack{K \subseteq [L] \\ |K|=n}} Z_{L-n}(x_{\bar{k}_{1}},...,x_{\bar{k}_{L-n}}) \prod_{i \in K} h(x_{i})$$

$$\times \prod_{i \in K} \prod_{j \in K^{c}} \left[ \frac{x_{j} - qx_{i}}{x_{j} - x_{i}} \frac{1 - x_{i}x_{j}}{1 - qx_{i}x_{j}} \right] \prod_{\substack{1 \leq i < j \leq L \\ i, j \in K}} \frac{1 - x_{i}x_{j}}{1 - qx_{i}x_{j}}$$

$$\times \sum_{\sigma \in S_{n}} \prod_{1 \leq i < j \leq n} \frac{x_{k_{\sigma(j)}} - qx_{k_{\sigma(i)}}}{x_{k_{\sigma(j)}} - x_{k_{\sigma(i)}}} \prod_{i=1}^{n} \left[ \frac{(1 - q)x_{k_{\sigma(i)}}y_{\nu_{i}}}{1 - qx_{k_{\sigma(i)}}y_{\nu_{i}}} \prod_{j=1}^{\nu_{i} - 1} \frac{1 - x_{k_{\sigma(i)}}y_{j}}{1 - qx_{k_{\sigma(i)}}y_{j}} \right].$$
(5.1)

The outer sum is over subsets  $K = \{k_1, \ldots, k_n\}$  of  $[L] = \{1, \ldots, L\}$  with *n* elements, whose complement is denoted  $K^c = \{\bar{k}_1, \ldots, \bar{k}_{L-n}\}$ .

We will now prepare for the proof of this important result. Instead of proving that the partition function with empty initial condition (3.65) is equal the rational function (5.1), it is convenient to invert the family of vertical spectral parameters, with  $Y^{-1} = (y_1^{-1}, y_2^{-1}, ...)$ , by considering the diagram from (3.65)



Which we will show is equal to the following rational function formula, which is equivalent to (5.1) with inverted vertical spectral parameters

$$\mathfrak{G}_{\nu}\left(x_{1},\ldots,x_{L}|Y^{-1}\right) = \sum_{\substack{K \in [L] \\ |K|=n}} Z_{L-n}\left(x_{\bar{k}_{1}},\ldots,x_{\bar{k}_{L-n}}\right) \prod_{i \in K} h(x_{i})$$

$$\times \prod_{i \in K} \prod_{j \in K^{c}} \left[\frac{x_{j} - qx_{i}}{x_{j} - x_{i}} \frac{1 - x_{i}x_{j}}{1 - qx_{i}x_{j}}\right] \prod_{\substack{1 \leq i < j \leq L \\ i,j \in K}} \frac{1 - x_{i}x_{j}}{1 - qx_{i}x_{j}}$$

$$\times \sum_{\sigma \in S_{n}} \prod_{1 \leq i < j \leq n} \frac{x_{k_{\sigma(j)}} - qx_{k_{\sigma(i)}}}{x_{k_{\sigma(j)}} - x_{k_{\sigma(i)}}} \prod_{i=1}^{n} \left[\frac{(1 - q)x_{k_{\sigma(i)}}/y_{\nu_{i}}}{1 - qx_{k_{\sigma(i)}}/y_{\nu_{i}}} \prod_{j=1}^{\nu_{i}-1} \frac{1 - x_{k_{\sigma(i)}}/y_{j}}{1 - qx_{k_{\sigma(i)}}/y_{j}}\right]. \quad (5.3)$$

Before we present the proof of Theorem 5.1, we will need some important properties of the partition function which largely follow from Lemma 3.21.

**Lemma 5.2.** The partition function from Definition 3.10 with empty initial condition satisfies the following properties. We note that for configuration  $\nu \in \mathbb{W}$  with at least one occupation, The coordinate  $\nu_1 \in \mathbb{N}$  denotes the right-most occupation in  $\nu$ .

- (i)  $G_{\nu}(x_1, \ldots, x_L|Y^{-1})$  is a meromorphic function in  $y_{\nu_1}$ . Its poles are all simple and occur at the points  $y_{\nu_1} = qx_i$  for  $1 \le i \le L$ .
- (ii)  $G_{\nu}(x_1,\ldots,x_L|Y^{-1})$  is symmetric in its alphabet  $(x_1,\ldots,x_L)$ .
- (iii) The residue of  $G_{\nu}(x_1, \ldots, x_L|Y^{-1})$  at its simple pole  $y_{\nu_1} = qx_1$  is given by

$$\operatorname{Res}_{y_{\nu_{1}}=qx_{1}}\left[G_{\nu}\left(x_{1},\ldots,x_{L}|Y^{-1}\right)\right] = (1-q)x_{1}h(x_{1})\prod_{j=2}^{L}\frac{x_{j}-qx_{1}}{x_{j}-x_{1}}\frac{1-x_{1}x_{j}}{1-qx_{1}x_{j}}$$

$$\times \prod_{j=1}^{\nu_1 - 1} \frac{y_j - x_1}{y_j - qx_1} G_{(\nu_2, \dots, \nu_n)} \left( x_2, \dots, x_L | Y^{-1} \right).$$
 (5.4)

(iv) The limit in  $y_{\nu_1}$  is

$$\lim_{y_{\nu_1} \to \infty} G_{\nu} \left( x_1, \dots, x_L | Y^{-1} \right) = 0.$$
(5.5)

(v) When the coordinate is empty

$$G_{\emptyset}(x_1, \dots, x_L | Y^{-1}) = Z_L(x_1, \dots, x_L).$$
 (5.6)

*Proof.* We will demonstrate the properties diagrammatically on the partition function (5.2).

(i) From the diagram, the only dependence on  $y_{\nu_1}$  is from the  $\nu_1$ 'th column. The weights which contribute to the partition function from this column will be from (2.2) with  $z = x_i/y_{\nu_1}$ , where i corresponds to the rows  $1 \le i \le L$ . All of the vertex configurations carry weights which are either entire functions of  $y_{\nu_1}$  or are analytic except at the isolated point  $y_{\nu_1} = qx_i$ . Since these weights contribute the only dependence on  $y_{\nu_1}$  we can conclude that  $G_{\nu}(x_1, \ldots, x_L|Y^{-1})$  is a meromorphic function for all  $y_{\nu_1}$  with possible singularities at the isolated points  $y_{\nu_1} = qx_i$  for  $1 \le i \le L$ .

Each global path configuration on (5.2) is will feature a weight from each vertex in the  $\nu_1$ 'th column at most once. Since each weight generates at most a simple pole at  $y_{\nu_1} = qx_i$ , we can conclude that the partition function will be a sum of rational functions with simple poles at  $y_{\nu_1} = qx_i$ . Therefore the poles at these points will be simple.

- (ii) This property follows from Corollary 3.11.
- (iii) Observing the boundary conditions, the vertex in the  $\nu_1$ 'th column and first row has two possible vertex configurations. these are shown in the table below.



Since the all-empty configuration on the left has weight 1, lattice configurations where this vertex is empty will have a partition function contribution which are analytic at  $y_{\nu_1} = qx_1$ . When the other weight is involved the contribution will have a simple pole at  $y_{\nu_1} = qx_1$ . By taking the residue of the whole partition function at the point  $y_{\nu_1} = qx_1$  we isolate contributions where this vertex is non-empty. Such configurations are depicted in the following diagram



By fixing the configuration at this vertex, we really freeze the contribution along the whole line associated with spectral parameter  $x_1$  and the line associated with vertical spectral

parameter  $y_{\nu_1}$ . This freezing passes on the empty boundary conditions below the first line the below the second line. Likewise it enforces the empty conditions on the right of the  $\nu_1$ 'th column to the  $(\nu_1 - 1)$ 'th column.

After removing the frozen contribution when taking the residue, what is left in the rectangle in (5.7) is that of the same partition function (5.2) with a n-1 coordinates  $(\nu_2, \ldots, \nu_n)$ and L-1 rows with spectral parameters  $(x_2, \ldots, x_L)$ .

Taking this residue can be written as

$$\operatorname{Res}_{y_{\nu_{1}}=qx_{1}}\left[G_{\nu}\left(x_{1},\ldots,x_{L}|Y^{-1}\right)\right] = h(x_{1})\prod_{j=2}^{n}\frac{1-x_{1}x_{j}}{1-qx_{1}x_{j}}\prod_{j=1}^{\nu_{1}-1}\frac{y_{j}-x_{1}}{y_{j}-qx_{1}}G_{(\nu_{2},\ldots,\nu_{n})}\left(x_{2},\ldots,x_{L}|Y^{-1}\right) \\ \times \lim_{y_{\nu_{1}}\to qx_{1}}\left[\left(y_{\nu_{1}}-qx_{1}\right)\frac{x_{1}(1-q)}{y_{\nu_{1}}-qx_{1}}\prod_{j=2}^{n}\frac{q(y_{\nu_{1}}-x_{j})}{y_{\nu_{1}}-qx_{j}}\right], \quad (5.8)$$

\_\_\_\_

which can be easily manipulated to take the form of (5.4). We note here we can include the entire inverted alphabet  $Y^{-1}$  and remove it from the evaluation of the limit since the smaller partition function will only have explicit dependence on  $y_j$  for  $1 \le j \le \nu_2$ .

(iv) From the table of weights (2.2), the configuration



However when observing the boundary conditions of (5.2), we note that there is a path exit through the top of the  $\nu_1$ 'th column while there are no occupations on the bottom entry or to the right of this column. Consequently, this weight must appear in the  $\nu_1$ 'th column exactly once for any configuration to provide a non-zero contribution.

Meanwhile, all other vertex configurations within (2.2) will not diverge under the same the limit. This is sufficient to conclude that the limit of the whole partition function will evaluate to zero.

(v) This property follows from Theorem 3.23.

We will now proceed with the proof of the theorem.

*Proof of Theorem 5.1.* Let us write (5.3) in the more compact form

$$\mathfrak{G}_{\nu}\left(x_{1},\ldots,x_{L}|Y^{-1}\right) = \sum_{\substack{K \subseteq [L] \\ |K|=n}} Z_{L-n}\left(x_{K^{c}}\right)h(x_{K})\Delta(x_{K}|x_{K^{c}})\Phi_{\nu}\left(x_{K}|Y^{-1}\right),$$
(5.9)

where we have defined

$$h(x_K) = \prod_{i \in K} h(x_i),$$
  
$$\Delta(x_K | x_{K^c}) = \prod_{i \in K} \prod_{j \in K^c} \left[ \frac{x_j - qx_i}{x_j - x_i} \frac{1 - x_i x_j}{1 - qx_i x_j} \right] \prod_{\substack{1 \le i < j \le L \\ i, j \in K}} \frac{1 - x_i x_j}{1 - qx_i x_j},$$

$$\Phi_{\nu}\left(x_{K}|Y^{-1}\right) = \sum_{\sigma \in S_{n}} \prod_{1 \le i < j \le n} \frac{x_{k_{\sigma(j)}} - qx_{k_{\sigma(i)}}}{x_{k_{\sigma(j)}} - x_{k_{\sigma(i)}}} \prod_{i=1}^{n} \left[\frac{(1-q)x_{k_{\sigma(i)}}/y_{\nu_{i}}}{1 - qx_{k_{\sigma(i)}}/y_{\nu_{i}}} \prod_{j=1}^{\nu_{i}-1} \frac{1 - x_{k_{\sigma(i)}}/y_{j}}{1 - qx_{k_{\sigma(i)}}/y_{j}}\right]$$

We shall begin by proving that (5.9) obeys the same set of properties as  $G_{\nu}(x_1, \ldots, x_L|Y^{-1})$  from (5.2). These are the properties in Lemma 5.2 which provide a recursive construction for the formula

(5.3) with an initial condition, and so completely define the formula for (5.2) through an inductive argument.

- (i)  $\mathfrak{G}_{\nu}(x_1, \ldots, x_L|Y^{-1})$  is a meromorphic function in  $y_{\nu_1}$ . Its poles are all simple and occur at the points  $y_{\nu_1} = qx_i$ ,  $1 \le i \le L$ . This property is immediate from the formula for  $\Phi_{\nu}(x_K|Y^{-1})$ , which is the only place where  $\mathfrak{G}_{\nu}(x_1, \ldots, x_L|Y^{-1})$  has dependence on the family Y.
- (ii)  $\mathfrak{G}_{\nu}(x_1,\ldots,x_L|Y^{-1})$  is symmetric in its alphabet  $(x_1,\ldots,x_L)$ . This is manifest from the form (5.9) of  $\mathfrak{G}_{\nu}(x_1,\ldots,x_L|Y^{-1})$ .
- (iii) The residue of  $\mathfrak{G}_{\nu}(x_1,\ldots,x_L|Y^{-1})$  at its simple pole  $y_{\nu_1} = qx_1$  is given by

$$\operatorname{Res}_{y_{\nu_{1}}=qx_{1}}\left[\mathfrak{G}_{\nu}\left(x_{1},\ldots,x_{L}|Y^{-1}\right)\right] = (1-q)x_{1}h(x_{1})\prod_{j=2}^{L}\frac{x_{j}-qx_{1}}{x_{j}-x_{1}}\frac{1-x_{1}x_{j}}{1-qx_{1}x_{j}} \times \prod_{j=1}^{\nu_{1}-1}\frac{y_{j}-x_{1}}{y_{j}-qx_{1}}\mathfrak{G}_{(\nu_{2},\ldots,\nu_{n})}\left(x_{2},\ldots,x_{L}|Y^{-1}\right).$$

This is easily seen by computing

$$\operatorname{Res}_{y_{\nu_{1}}=qx_{1}}\left[\Phi_{\nu}\left(x_{K}|Y^{-1}\right)\right] = \begin{cases} (1-q)x_{1}\prod_{j\in K\setminus\{1\}}\frac{x_{j}-qx_{1}}{x_{j}-x_{1}}\prod_{j=1}^{\nu_{1}-1}\frac{y_{j}-x_{1}}{y_{j}-qx_{1}}\Phi_{(\nu_{2},\dots,\nu_{n})}\left(x_{K\setminus\{1\}}|Y^{-1}\right) & \text{if } 1\in K\\ 0 & \text{if } 1\notin K \end{cases}$$

and noting that for  $1 \in K$ , we have

$$\Delta(x_K|x_{K^c}) = \prod_{j \in K^c} \frac{x_j - qx_1}{x_j - x_1} \prod_{j=2}^L \frac{1 - x_1 x_j}{1 - qx_1 x_j} \Delta(x_{K \setminus \{1\}} | x_{K^c}), \qquad h(x_K) = h(x_1) h(x_{K \setminus \{1\}})$$

- (iv)  $\mathfrak{G}_{\nu}(x_1,\ldots,x_L|Y^{-1}) \to 0$  as  $y_{\nu_1} \to \infty$ . This follows by computing this limit directly on  $\Phi_{\nu}(x_K|Y^{-1})$ .
- (v)  $\mathfrak{G}_{\emptyset}(x_1,\ldots,x_L|Y^{-1}) = Z_L(x_1,\ldots,x_L)$ . This is simply the n = 0 case of the formula (5.9).

We have shown that  $\mathfrak{G}_{\nu}(x_1,\ldots,x_L|Y^{-1})$  obeys the same set of properties as  $G_{\nu}(x_1,\ldots,x_L|Y^{-1})$  does according to Lemma 5.2. It remains to show that these properties imply the equality of the two objects; we do this by induction on the length of  $\nu$ . To that end, define the function

$$\mathfrak{Z}_{\nu}(x_1,\ldots,x_L|Y) = \mathfrak{G}_{\nu}(x_1,\ldots,x_L|Y^{-1}) - G_{\nu}(x_1,\ldots,x_L|Y^{-1}).$$

By construction,  $\mathfrak{Z}_{\varnothing}(x_1, \ldots, x_L|Y) = 0$ . It follows that there exists an integer  $m \ge 0$  such that  $\mathfrak{Z}_{\mu}(x_1, \ldots, x_L|Y) = 0$  for all strict partitions  $\mu = (\mu_1 > \cdots > \mu_m)$  of length m (with L being arbitrary); this is our inductive hypothesis.

Now let  $\lambda = (\lambda_1 > \cdots > \lambda_{m+1})$  be a strict partition of length m + 1. We know that  $\mathfrak{Z}_{\lambda}(x_1, \ldots, x_L|Y)$ is a meromorphic function in  $y_{\lambda_1}$ , its poles are all simple, and it vanishes as  $y_{\lambda_1} \to \infty$ . However, from the recursion relation obeyed by  $\mathfrak{G}_{\lambda}(x_1, \ldots, x_L|Y^{-1})$  and  $G_{\lambda}(x_1, \ldots, x_L|Y^{-1})$ , as well as the inductive hypothesis, all poles have vanishing residue. This means that  $\mathfrak{Z}_{\lambda}(x_1, \ldots, x_L|Y)$  is entire and bounded in  $y_{\lambda_1}$  and therefore constant. This constant must be zero in view of the known  $y_{\lambda_1} \to \infty$  behaviour. It follows that  $\mathfrak{Z}_{\lambda}(x_1, \ldots, x_L|Y) = 0$  for all strict partitions  $\lambda = (\lambda_1 > \cdots > \lambda_{m+1})$ of length m + 1, and the inductive step of the proof is complete.  $\Box$ 

5.2. Integral formula. Here we will present the sum over subset expression (5.1) as an equivalent nested integral formula.

**Definition 5.3.** Fix an alphabet  $(x_1, \ldots, x_L) \in \mathbb{C}^L$ . We denote by  $\mathcal{C}_1, \ldots, \mathcal{C}_n$  a collection of positively oriented closed complex contours satisfying

- For all  $1 \leq i < j \leq n$ , we have that  $qC_j$  lies completely outside the interior of  $C_i$ , where  $qC_j$  denotes the image of  $C_j$  under multiplication by q. In addition, if 1 is within the interior of  $C_i$  we also require that  $C_i$  is completely contained in the interior of  $C_j$ .
- For all  $1 \leq i \leq L$ , the contour  $C_i$  surrounds all points  $x_j$  and does not surround the points  $qx_j, q^{-1}x_j^{-1}, q^{-1}y_k^{-1}, a, c$  for all  $1 \leq j \leq L$  and  $k \in \mathbb{N}$ .



Figure 2. Diagrams depicting arrangements of contours allowed by Defintion 5.3.

Examples of contours satisfying the conditions of Definition 5.3 are shown in Figure 2. These conditions allow for some freedom with contour choice. In particular we may choose all contours to be equal,  $C_i =: C$  for all i, provided that the contours neither lie upon nor enclose  $1 \in \mathbb{C}$ .

Provided that the integrand considered has no singularity at 1, the contours may surround 1 if we choose that they are nested. That is, for all  $1 \le i \le n-1$  the contour  $C_i$  is completely contained within the interior of  $C_{i+1}$ . This choice is convenient as it allows us to choose elements of our alphabet,  $x_i$ , to be arbitrarily close to 1.

**Theorem 5.4.** The partition function expression (5.1) can be expressed as the following *n*-fold integral:

$$G_{\nu}(x_{1},\ldots,x_{L}|Y) = \oint_{\mathcal{C}_{1}} \frac{dw_{1}}{2\pi i} \cdots \oint_{\mathcal{C}_{n}} \frac{dw_{n}}{2\pi i} Z_{L+n}\left(x_{1},\ldots,x_{L},w_{1}^{-1},\ldots,w_{n}^{-1}\right) \\ \times \prod_{i=1}^{n} \prod_{j=1}^{L} \left[ \frac{qw_{i}-x_{j}}{w_{i}-x_{j}} \frac{1-w_{i}x_{j}}{1-qw_{i}x_{j}} \right]_{1 \le i < j \le n} \left[ \frac{w_{j}-w_{i}}{qw_{j}-w_{i}} \frac{1-qw_{i}w_{j}}{1-w_{i}w_{j}} \right] \\ \times \prod_{i=1}^{n} \left[ \frac{ac\left(qw_{i}^{2}-1\right)}{(w_{i}-a)(w_{i}-c)} \frac{y_{\nu_{i}}}{1-qw_{i}y_{\nu_{i}}} \prod_{j=1}^{\nu_{i}-1} \frac{1-w_{i}y_{j}}{1-qw_{i}y_{j}} \right], \quad (5.10)$$

where the contours  $C_1, \ldots, C_n$  satisfy the conditions of Definition 5.3.

*Proof.* The idea of the proof is to replace each  $x_i$  for  $i \in K$  in the sum in (5.1) by an auxiliary variable  $w_i$  that will be integrated over a contour surrounding simple poles at all  $x_1, \ldots, x_L$ . The sum over K then dictates which n of the L possible residues are evaluated, whilst the sum over  $\sigma \in S_n$  dictates the order in which the residues are evaluated for a given K.

For this to work, only the residues at  $w_i = x_j$  in (5.10) should be evaluated. All other poles in (5.10) therefore need to be excluded from the contours, and this is guaranteed by Definition 5.3. Firstly, it is obvious that the explicit poles in (5.10) at  $a, c, q^{-1}y_j^{-1}$  and  $q^{-1}x_j^{-1}$  need to lie outside each contour. Secondly, the poles at  $qx_j$  also need to be avoided because these cause singularities in  $Z_{L+n}$ , see below. Moreover, for  $1 \le i < j \le n$ , the factors of the form

$$\frac{w_j - w_i}{qw_j - w_i} \frac{1 - qw_i w_j}{1 - w_i w_j},$$

produce potential residues which will be avoided when  $qC_i$  lies outisde the interior of  $C_i$ .

In order to reproduce the triangular partition function  $Z_{L-n}$  that appears in the summand of (5.1) we extend this function to  $Z_{L+n}$  in the combined alphabet  $(x_1, \ldots, x_L, w_1^{-1}, \ldots, w_n^{-1})$ . During the evaluation of the residue of the simple pole at each  $w_i = x_j$ , the recursion relation (4.4) ensures that we re-obtain  $Z_{L-n}$  in the complement alphabet of x-variables of (5.1). Furthermore, according to Proposition 4.2, the rational function  $Z_{L+n}(x_1, \ldots, x_L, w_1^{-1}, \ldots, w_n^{-1})$  has poles at  $w_i = a^{-1}$ ,  $w_i = c^{-1}$  and  $w_i = qx_j$  and so has no singularities at  $w_i = x_j$  that could affect the residue evaluation.

In order to proceed, we rewrite the following factor that occurs in (5.1),

$$\prod_{i \in K} \prod_{j \in K^{c}} \left[ \frac{x_{j} - qx_{i}}{x_{j} - x_{i}} \frac{1 - x_{i}x_{j}}{1 - qx_{i}x_{j}} \right] = \prod_{i \in K} \prod_{j \in K^{c}} \frac{1}{x_{i} - x_{j}} \prod_{i \in K} \prod_{j=1}^{L} \frac{(qx_{i} - x_{j})(1 - x_{i}x_{j})}{1 - qx_{i}x_{j}} \\
\times \prod_{\substack{i \neq j \\ i, j \in K}} \frac{1 - qx_{i}x_{j}}{(qx_{i} - x_{j})(1 - x_{i}x_{j})} \prod_{i \in K} \frac{qx_{i}^{2} - 1}{(1 - q)x_{i}(1 - x_{i}^{2})}.$$
(5.11)

Incorporating this we notice that for each K the second line in (5.1) is manifestly symmetric in the variables  $x_i$  for  $i \in K$ . This allows us to replace each  $x_i$  for  $i \in K$  with  $w_i$ , i.e. the right hand side of (5.11) is replaced by

$$\prod_{i,j=1\atop i\neq j}^{n} (w_i - w_j) \prod_{i=1}^{n} \prod_{j=1}^{L} \frac{(qw_i - x_j)(1 - w_i x_j)}{(w_i - x_j)(1 - qw_i x_j)} \prod_{i,j=1\atop i\neq j}^{n} \frac{1 - qw_i w_j}{(qw_i - w_j)(1 - w_i w_j)} \prod_{i=1}^{n} \frac{qw_i^2 - 1}{(1 - q)w_i(1 - w_i^2)}, \quad (5.12)$$

and each  $x_{\sigma(i)}$  and  $x_{\sigma(j)}$  in (5.1) is similarly replaced by  $w_i$  and  $w_j$  respectively. Simplifying and cancelling common factors we thus obtain the integrand of (5.10). Finally we note that the factors  $w_j - w_i$  in (5.10) ensure that after evaluating the residue of  $w_i$  at  $x_k$  the singularity at  $w_j = x_k$  is removable for all  $j \neq i$ , and hence that the residue for each  $w_i$  is evaluated at a different simple pole.

5.3. An orthogonality conjecture. In this section we examine an interesting implication of the Cauchy identity (4.15) and integral formula (5.10) when  $c \to \infty$ . Our starting point is the observation made in Corollary 4.7 regarding the  $c \to \infty$  limit of the triangular partition function (3.69) which we recall now:

$$\lim_{c \to \infty} Z_m(x_1, \dots, x_m) = \lim_{c \to \infty} \prod_{i=1}^m (1 - h(x_i)) = \prod_{i=1}^m \frac{x_i(1 - ax_i)}{x_i - a}.$$
(5.13)

Making use of Corollary 4.7 in the Cauchy identity (4.15), it reads

$$\lim_{c \to \infty} \sum_{\kappa} G_{\kappa}(x_1, \dots, x_L) F_{\kappa}(z_1, \dots, z_M) = \prod_{i=1}^M \frac{a(1-z_i^2)}{a-z_i} \prod_{i=1}^M \prod_{j=1}^L \left[ \frac{x_j - qz_i}{x_j - z_i} \frac{1 - z_i x_j}{1 - qz_i x_j} \right] \prod_{i=1}^L \frac{x_i(1 - ax_i)}{x_i - a}.$$
 (5.14)

A similar application of Corollary 4.7 inside the integral formula (5.10) yields, after redistribution of factors in the integrand,

$$\lim_{c \to \infty} G_{\nu}(x_1, \dots, x_L) = \prod_{i=1}^{L} \frac{x_i(1 - ax_i)}{x_i - a} \oint_{\mathcal{C}_1} \frac{\mathrm{d}w_1}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{C}_n} \frac{\mathrm{d}w_n}{2\pi \mathrm{i}} \prod_{i=1}^n \prod_{j=1}^{L} \left[ \frac{qw_i - x_j}{w_i - x_j} \frac{1 - w_i x_j}{1 - qw_i x_j} \right] \\ \times \prod_{1 \le i < j \le n} \left[ \frac{w_j - w_i}{qw_j - w_i} \frac{1 - qw_i w_j}{1 - w_i w_j} \right] \prod_{i=1}^n \left[ \frac{a \left( 1 - qw_i^2 \right)}{w_i (1 - aw_i)} \frac{y_{\nu_i}}{1 - qw_i y_{\nu_i}} \prod_{j=1}^{\nu_i - 1} \frac{1 - w_i y_j}{1 - qw_i y_j} \right].$$
(5.15)

One may then recognize the right hand side of (5.14) (under the substitutions  $M \mapsto n$  and  $z_i \mapsto w_i$ ) as being embedded within the integrand of (5.15), leading to the equality

$$\lim_{c \to \infty} G_{\nu}(x_1, \dots, x_L) = \sum_{\kappa} \lim_{c \to \infty} G_{\kappa}(x_1, \dots, x_L) \oint_{\mathcal{C}_1} \frac{\mathrm{d}w_1}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{C}_n} \frac{\mathrm{d}w_n}{2\pi \mathrm{i}} \lim_{c \to \infty} F_{\kappa}(w_1, \dots, w_n) \\ \times \prod_{1 \le i < j \le n} \left[ \frac{w_j - w_i}{qw_j - w_i} \frac{1 - qw_i w_j}{1 - w_i w_j} \right] \prod_{i=1}^n \left[ \frac{(w_i - a)(1 - qw_i^2)}{w_i(1 - aw_i)(1 - w_i^2)} \frac{y_{\nu_i}}{1 - qw_i y_{\nu_i}} \prod_{j=1}^{\nu_i - 1} \frac{1 - w_i y_j}{1 - qw_i y_j} \right], \quad (5.16)$$

where it is necessary to assume the convergence constraints

$$\frac{1 - x_i y_k}{1 - q x_i y_k} \frac{q(1 - w_j / y_k)}{1 - q w_j / y_k} \le \rho < 1, \qquad \left| \frac{1 - x_i y_k}{1 - q x_i y_k} \frac{1 - q w_j y_k}{1 - w_j y_k} \right| \le \rho < 1$$
(5.17)

in order to introduce the infinite sum over  $\kappa^5$ . The contours  $\mathcal{C}_1, \ldots, \mathcal{C}_n$  are as previously, but we must now examine which poles of the function  $\lim_{c\to\infty} F_{\kappa}(w_1, \ldots, w_n|Y)$  they enclose (noting that the points  $w_i = x_i$  are no longer singularities of the integrand).

To deduce this, we make some assumptions concerning the parameters q and  $Y = (y_1, y_2, ...)^6$ . We note that one way to satisfy the constraints (5.17) is to assume that |q| is arbitrarily small. This renders the first constraint in (5.17) trivial, while the second one becomes equivalent to establishing the bound  $|x_i - y_k^{-1}| < \rho |w_j - y_k^{-1}|$  for all  $1 \le i \le L$ ,  $1 \le j \le n$ ,  $k \in \mathbb{N}$ ; the latter constraint is satisfied if the points  $y_k^{-1}$  are arbitrarily close to the points  $x_i$ , and are thereby enclosed by the integration contours.

In summary, we may replace the contours  $C_1, \ldots, C_n$  in (5.16) by a single contour C that encloses all points  $Y^{-1} = (y_1^{-1}, y_2^{-1}, \ldots)$  and no other singularities of the integrand, and such that  $q \cdot C$  is disjoint from the interior of C. Reading off the coefficient of  $\lim_{c\to\infty} G_{\nu}(x_1, \ldots, x_L)$  on both sides of the resulting equation, we arrive at the following conjecture:

**Conjecture 5.5.** Fix a finite subset  $\nu = \{\nu_1 > \cdots > \nu_n \ge 1\}$ , and a second finite subset  $\kappa$  whose cardinality satisfies  $|\kappa| \le n$ . Then one has that

$$\oint_{\mathcal{C}} \frac{\mathrm{d}w_1}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{C}} \frac{\mathrm{d}w_n}{2\pi \mathrm{i}} \prod_{1 \le i < j \le n} \left[ \frac{w_j - w_i}{qw_j - w_i} \frac{1 - qw_i w_j}{1 - w_i w_j} \right] \\
\times \prod_{i=1}^n \left[ \frac{w_i - a}{w_i (1 - aw_i)} \frac{1 - qw_i^2}{1 - w_i^2} \frac{y_{\nu_i}}{1 - qw_i y_{\nu_i}} \prod_{j=1}^{\nu_i - 1} \frac{1 - w_i y_j}{1 - qw_i y_j} \right] \lim_{c \to \infty} F_{\kappa}(w_1, \dots, w_n) = \delta_{\kappa, \nu}, \quad (5.18)$$

where the contour  $\mathcal{C}$  is a small, positively oriented circle surrounding the points  $y_j^{-1}$ ,  $j \ge 1$  and no other singularities of the integrand, such that  $q \cdot \mathcal{C}$  is disjoint from the interior of  $\mathcal{C}$ .

Although we have given an essentially complete formulation of (5.18), it remains conjectural as we have not established that the functions  $\lim_{c\to\infty} G_{\kappa}(x_1,\ldots,x_L)$  are linearly independent. Conjecture 5.5 has been extensively tested, and we plan to return to its proof in a later text.

# 6. Open ASEP on the half-line

In this section we demonstrate an important reduction of the partition function  $G_{\nu/\mu}$  defined by (3.42). In section 3 it was demonstrated that  $G_{\nu/\mu}$  can describe a discrete-time Markov process of interacting particles on a half-line with both creation and annihilation occurring at the origin. A continuous-time limit of this propagator will recover the dynamics of the *asymmetric simple exclusion process* (ASEP) on the half-line with open boundary conditions.

6.1. Markov generator. We will consider the continuous-time asymmetric simple exclusion process on the semi-infinite half-line  $\mathbb{N} := \mathbb{Z}_{>0}$ . As with the vertex model, we only consider configurations with a finite yet varying number of particles. Our conventions for ASEP coordinates follow that of Section 3.1. These configurations are indexed by the random variable  $\nu = (\nu_1(t), \ldots, \nu_n(t)) \in \mathbb{W}$ where  $\nu_1 > \nu_2 > \cdots > \nu_n$  for some finite  $n \ge 0$ . We may also regard the number of particles n as a random variable.

These configurations evolve according to the following bulk transition rates

$$\nu \mapsto (\nu_1, \dots, \nu_i + 1, \dots, \nu_n) \text{ at rate } 1 \text{ if } \nu_{i-1} > \nu_i + 1,$$
  
$$\nu \mapsto (\nu_1, \dots, \nu_i - 1, \dots, \nu_n) \text{ at rate } q \text{ if } \nu_{i+1} < \nu_i - 1.$$

<sup>&</sup>lt;sup>5</sup>More precisely, we have used the fact that this convergence is uniform with  $(x_1, \ldots, x_L)$  and  $(w_1, \ldots, w_n)$  ranging over compact subsets of  $\mathbb{C}$ , which is necessary to be able to switch the order of integration and summation.

<sup>&</sup>lt;sup>6</sup>It is later possible to relax these constraints, since we ultimately derive (5.18), which is an identity of rational functions in q and Y that holds when these parameters take values in certain compact subsets of  $\mathbb{C}$ ; it must therefore hold generally.



**Figure 3.** Dynamics of the ASEP on the half-line  $\mathbb{N}$ .

If we have the case where  $\nu_{i-1} = \nu_i + 1$  then this process is excluded (occurs at rate zero). We also have the boundary transition rates

 $(\nu_1, \dots, \nu_n) \mapsto (\nu_1, \dots, \nu_n, 1)$  at rate  $\alpha$  if  $\nu_n > 1$  or n = 0,  $(\nu_1, \dots, \nu_n) \mapsto (\nu_1, \dots, \nu_{n-1})$  at rate  $\gamma$  if  $\nu_n = 1$ .

The bulk and boundary dynamics are depicted in Figure 3. For fixed  $\nu \in \mathbb{W}$ , the Markov generator for this process is expressed as an operator  $\mathscr{L}$  which acts on functions  $f : \mathbb{W} \to \mathbb{C}$  given by

$$\mathscr{L}[f](\nu) = (1 - \eta_1^{\nu}) \left(\gamma f(\nu \cup \{1\}) - \alpha f(\nu)\right) + \eta_1^{\nu} \left(\alpha f(\nu \setminus \{1\}) - \gamma f(\nu)\right) \\ + \sum_{s=1}^{\infty} \left[\eta_s^{\nu} (1 - \eta_{s+1}^{\nu}) + q \eta_{s+1}^{\nu} (1 - \eta_s^{\nu})\right] \left(f(\nu^{s,s+1}) - f(\nu)\right), \quad (6.1)$$

where  $\nu^{s,s+1} \in \mathbb{W}$  is the configuration obtained by the interchange of occupations at sites s and s+1. We have also used the notation  $\eta_j^{\nu}$  as the occupation of  $\nu$  at site j defined in Definition 3.1. Using the Markov generator we can arrive at the evolution equation of the ASEP on the half-line

$$\frac{\mathrm{d}}{\mathrm{d}t}\psi_t(\nu) = \mathscr{L}[\psi_t](\nu), \tag{6.2}$$

whose solution,  $\psi_t : \mathbb{W} \to \mathbb{R}$ , is an eigenfunction of the Markov generator indexed by continuous-time parameter  $t \ge 0$ . The rest of this section is devoted to the presentation a solution to (6.2) for an initially empty configuration with partially open boundary parameters.

6.2. Transition probability from the vertex model. In this section we outline a method to obtain solutions to the half-line ASEP equations of motion (6.2) via a reduction of the symmetric function (3.42). An explicit evaluation is provided for the case of an empty initial configuration of particles following from the results of Section 5.

Let  $\mu, \nu \in \mathbb{W}$  be arbitrary particle configurations. We will denote  $\mathbb{P}_t(\mu \to \nu)$  as a solution to (6.2), indexed by a time parameter  $t \ge 0$ , subject to the initial condition

$$\mathbb{P}_0(\mu \to \nu) = \delta_{\mu,\nu}$$

That is, a probability of being in configuration  $\nu$  at time t after being initially in  $\mu$ . We refer to this as the *transition probability*. The transition probability can be expressed as a formal solution to the evolution equation in terms of the Markov generator. This is given as

$$\mathbb{P}_t(\mu \to \nu) = \langle \mu | \exp(t\mathcal{L}) | \nu \rangle, \qquad (6.3)$$

where the exponential is regarded as the formal operator exponential. We now proceed by outlining the specialization required to recover the ASEP transition probability from the half-space six-vertex model.

**Proposition 6.1.** Fix configurations  $\mu, \nu \in \mathbb{W}$  and time  $t \ge 0$ . The ASEP half-line transition probability is recovered as the limit of the symmetric function

$$\mathbb{P}_t(\mu \to \nu) = \lim_{L \to \infty} G_{\nu/\mu}(x_1 \dots, x_L | Y) \Big|_{y_j = 1, \ x_i = 1 - (1 - q)t/(2L)},\tag{6.4}$$

where we have specified the spectral parameters  $x_i = 1 - (1-q)t/(2L)$  and vertical parameters  $y_j = 1$  for all  $1 \le i \le L$  and  $j \in \mathbb{N}$  prior to taking the limit  $L \to \infty$ . The limit (6.4) holds provided that we choose boundary Markov rates as

$$\alpha = \frac{ac(1-q)}{(1-a)(1-c)}, \qquad \gamma = -\frac{1-q}{(1-a)(1-c)}, \tag{6.5}$$

where we are free to choose a, c so that both  $\alpha, \gamma \ge 0$  while we restrict  $q \ge 0$ .

We will first demonstrate a degeneration of the double-row

**Lemma 6.2.** Fix  $\epsilon > 0$  let  $x = 1 - (1 - q)\epsilon$  and  $y_j = 1$  for all  $j \in \mathbb{N}$ . Then the double-row operator (3.2) is given by

$$A(x|Y)\Big|_{x=1-(1-q)\epsilon, y_j=1} = 1 + 2\epsilon \mathscr{L} + O(\epsilon^2).$$
(6.6)

*Proof.* To demonstrate this, we choose  $x_i = 1 - (1 - q)\epsilon$ ,  $y_j = 1$  and observe that the weights from (2.2) with horizontal parameters  $x_i^{-1}$  and vertical parameters  $y_j$  acquire the following form.



This table represents the weights of the upper row in the double row of (3.2). The weights of the lower row can be obtained by rotating these vertices. The boundary weights have a similar form.



The double-row operator (6.6) can be determined by calculating for specific configurations  $\mu, \nu \in \mathbb{W}$  the partition function using the weights in (6.7),(6.8) and matching with the action of the ASEP generator (6.1).

Proof of Proposition 6.1. Using the double-row operator definition of the partition (3.42) and the scaling (6.6) with  $\epsilon = t/(2L)$ , we can arrive at the following expression for the partition function

$$G_{\nu/\mu}(x_1...,x_L|Y)\Big|_{y_j=1,\ x_i=1-(1-q)t/(2L)} = \langle \mu | \left(1 + \frac{t}{L}\mathscr{L} + O\left(\frac{1}{L^2}\right)\right)^L |\nu\rangle.$$
(6.9)

In order to take the limit we use the definition of the operator exponential. This yields

$$\lim_{L \to \infty} \left( 1 + \frac{t}{L} \mathscr{L} + O\left(\frac{1}{L^2}\right) \right)^L = \lim_{L \to \infty} \left( 1 + \frac{t}{L} \mathscr{L} \right)^L = \exp(t\mathscr{L}).$$
(6.10)

This recovers the formal solution for ASEP transition probability (6.3).

6.3. Particle injection case. In this section we provide an explicit expression for the ASEP transition probability (6.3) under the specialization  $\gamma = 0$ , whereby particles may enter from the boundary at rate  $\alpha$  but may not exit the system.

**Definition 6.3.** Fix an alphabet  $(x_1, \ldots, x_L)$  and let  $\{C_1, \ldots, C_n\}$  be a set of contours satisfying the conditions of Definition 5.3 whilst also surrounding 1. We define  $\{D_1, \ldots, D_n\}$  to be these nested contours satisfying the conditions of Definition 5.3 having taken  $x_i = 1$ .

**Theorem 6.4.** Under the limit  $c \to \infty$  ( $\gamma \to 0$ ) for  $\alpha + q \neq 1$ , the ASEP transition probability is given by

$$\mathbb{P}_{t}(\varnothing \to \nu) = \alpha^{n} \mathrm{e}^{-\alpha t} \oint_{\mathcal{D}_{1}} \frac{\mathrm{d}w_{1}}{2\pi \mathrm{i}} \cdots \oint_{\mathcal{D}_{n}} \frac{\mathrm{d}w_{n}}{2\pi \mathrm{i}} \prod_{1 \le i < j \le n} \left[ \frac{w_{j} - w_{i}}{qw_{j} - w_{i}} \frac{1 - qw_{i}w_{j}}{1 - w_{i}w_{j}} \right] \\ \times \prod_{i=1}^{n} \left[ \frac{1 - qw_{i}^{2}}{w_{i}(q + \alpha - 1 - \alpha w_{i})(1 - qw_{i})} \left( \frac{1 - w_{i}}{1 - qw_{i}} \right)^{\nu_{i} - 1} \exp\left( \frac{(1 - q)^{2}w_{i}t}{(1 - w_{i})(1 - qw_{i})} \right) \right], \quad (6.11)$$

where the contours satisfy the conditions of Definition 6.3. That is, they surround the essential singularity at  $w_i = 1$ .

*Proof.* Consider the integral formula (5.10) with contours  $\{C_1, \ldots, C_n\}$  taken to surround 1 as well as the points  $x_1 \ldots, x_L$ . The conditions of the contours from Definition 5.3 also means that the contours must be nested whilst they do not intersect.

In the limit  $c \to \infty$ , only the term r = 0 in the sum over subset formula (4.9) for  $Z_L$  survives, see Corollary 4.7. Hence  $Z_L$  trivializes and completely factorizes into factors of the form  $1 - h(x_i)$ .

Upon substitution in (5.10) and setting  $x_i = 1 - (1 - q)t/(2L)$ , the limit  $L \to \infty$  can be taken on the integrand in a straightforward manner. Note that in this limit we have written  $a = \alpha/(\alpha + q - 1)$  for  $\alpha + q \neq 1$ . We have also calculated under the ASEP limit

$$\lim_{L \to \infty} \left. \prod_{j=1}^{L} \left[ \frac{qw_i - x_j}{w_i - x_j} \frac{1 - w_i x_j}{1 - qw_i x_j} \right] \right|_{x_j = 1 - (1 - q)t/(2L)} = \exp\left(\frac{(1 - q)^2 w_i t}{(1 - w_i)(1 - qw_i)}\right).$$

In order to obtain the result of the theorem, the  $L \to \infty$  limit must must also be simultaneously applied to the contours. This will deform the contours  $C_i \mapsto D_i$  using the contours from Definitions 5.3 and 6.3. This deformation occurs without crossing over any other singularities of the integrand, thus yielding the result.

#### COMPLIANCE WITH ETHICAL STANDARDS

The authors are not aware of any potential conflict of interest.

# APPENDIX A. PROPERTIES OF PFAFFIANS

A Pfaffian is taken over a  $2m \times 2m$ -dimensional skew-symmetric matrix. It is defined as

$$PfA = \frac{1}{2^m m!} \sum_{\sigma \in S_m} (-1)^{|\sigma|} \prod_{i=1}^m a_{\sigma(2i-1),\sigma(2i)}.$$
 (A.1)

Importantly, the determinant of skew-symmetric matrix is the square of a polynomial in its entries. The Pfaffian can be identified as this polynomial. That is, so long as A is an even-dimensional skew-symmetric matrix then

$$\det A = (\operatorname{Pf} A)^2. \tag{A.2}$$

The following identity appears in [Ste90].

**Lemma A.1.** Let A and B be skew-symmetric  $m \times m$  matrices. Then we have

$$Pf(A+B) = \sum_{r=0}^{m} (-1)^{r/2} \sum_{\substack{S \subseteq [m] \\ |S|=r}} (-1)^{\sum_{i} S_{i}} Pf(A_{S}) Pf(B_{S^{c}}), \qquad (A.3)$$

where  $S^{c}$  denotes the set which is the complement to the set S w.r.t. [m].

#### References

- [ABW23] A. Aggarwal, A. Borodin, and M. Wheeler. Deformed polynuclear growth in (1 + 1) dimensions. *Int. Math. Res. Not.*, 2023(7):5728–5780, 2023. arXiv:2108.06018.
- [Agg16] A. Aggarwal. Convergence of the Stochastic Six-Vertex Model to the ASEP. J. Math. Phys. Anal. Geom., 20(2):3, 2016. arXiv:1607.08683.
- [BBC20] G. Barraquand, A. Borodin, and I. Corwin. Half-space Macdonald processes. *Forum Math. Pi*, 8:e11, 2020. arXiv:1802.08210.
- [BBCS18] J. Baik, G. Barraquand, I. Corwin, and T. Suidan. Pfaffian Schur processes and last passage percolation in a half-quadrant. Ann. Probab., 46(6), 2018. arXiv:1606.00525.
- [BBCW18] G. Barraquand, A. Borodin, I. Corwin, and M. Wheeler. Stochastic six-vertex model in a half-quadrant and half-line open ASEP. *Duke Math. J.*, 167(13), 2018. arXiv:1704.04309.
- [BC14] A. Borodin and I. Corwin. Macdonald processes. *Probab. Theory Relat. Fields*, 158, 2014. arXiv:1111.4408.
- [BC24] G. Barraquand and I. Corwin. Markov duality and Bethe ansatz formula for half-line open ASEP. Probab. Math. Phys., 5(1):89–129, 2024. arXiv:2212.07349.
- [BCG16] A. Borodin, I. Corwin, and V. Gorin. Stochastic six-vertex model. *Duke Math. J.*, 165(3), 2016. arXiv:1407.6729.
- [BFK23] R. E. Behrend, I. Fischer, and C. Koutschan. Diagonally symmetric alternating sign matrices. 2023. arXiv:2309.08446.
- [BKD22] G. Barraquand, A. Krajenbrink, and P. L. Doussal. Half-space stationary Kardar–Parisi–Zhang equation beyond the Brownian case. J. Phys. A: Math. Theor., 55(27):275004, 2022. arXiv:2202.10487.
- [Bor17] A. Borodin. On a family of symmetric rational functions. *Adv. Math.*, 306:973–1018, 2017. arXiv:1410.0976.
- [BP18] A. Borodin and L. Petrov. Higher spin six vertex model and symmetric rational functions. *Sel. Math. New Ser.*, 24(2):751–874, 2018. arXiv:1601.05770.
- [BW20] A. Borodin and M. Wheeler. Observables of coloured stochastic vertex models and their polymer limits. *Probab. Math. Phys.*, 1(1):205–265, 2020. arXiv:2001.04913.
- [BW22] A. Borodin and M. Wheeler. Coloured stochastic vertex models and their spectral theory. *Astérisque*, 437, 2022. arXiv:1808.01866.
- [BWZJ15] D. Betea, M. Wheeler, and P. Zinn-Justin. Refined Cauchy/Littlewood identities and six-vertex model partition functions: II. Proofs and new conjectures. J. Algebr. Comb., 42(2):555–603, 2015. arXiv:1405.7035.
- [CFRV16] N. Crampe, C. Finn, E. Ragoucy, and M. Vanicat. Integrable boundary conditions for multi-species ASEP. J. Phys. A Math. Theor., 49(37):375201, 2016. arXiv:1606.01018.
- [CGdGW16] L. Cantini, A. Garbali, J. de Gier, and M. Wheeler. Koornwinder polynomials and the stationary multispecies asymmetric exclusion process with open boundaries. J. Phys. A Math. Theor., 49(44):444002, 2016. arXiv:1607.00039.
- [Che84] I. Cherednik. Factorizing particles on a half-line and root systems. *Theor. Math. Phys.*, 61(1):35–44, 1984.
- [CP16] I. Corwin and L. Petrov. Stochastic higher spin vertex models on the line. *Commun. Math. Phys.*, 343:651–700, 2016. arXiv:1502.07374.
- [CRV14] N. Crampe, E. Ragoucy, and M. Vanicat. Integrable approach to simple exclusion processes with boundaries. Review and progress. J. Stat. Mech. Theory Exp., 2014(11):P11032, 2014. arXiv:1408.5357.
- [CW06] S. Corteel and L. K. Williams. Tableaux combinatorics for the asymmetric exclusion process. *Adv. Appl. Math.*, 39:293–310, 2006. arXiv:math/0602109.
- [dGE05] J. de Gier and F. H. L. Essler. Bethe ansatz solution of the asymmetric exclusion process with open boundaries. *Phys. Rev. Lett.*, 95:240601, 2005. arXiv:cond-mat/0508707.
- [DVGR93] H. De Vega and A. Gonzalez Ruiz. Boundary K-matrices for the six vertex and the  $n(2n-1) A_{n-1}$  vertex models. J. Phys. A: Math. Gen., 26(12):L519, 1993. arXiv:hep-th/9211114.
- [EP20] F. H. L. Essler and L. Piroli. Integrability of one-dimensional Lindbladians from operator-space fragmentation. *Phys. Rev. E*, 102:062210, 2020. arXiv:2009.11745.
- [FV17] C. Finn and M. Vanicat. Matrix product construction for Koornwinder polynomials and fluctuations of the current in the open ASEP. J. Stat. Mech. Theory Exp., 2017(2):023102, 2017. arXiv:1610.08320.
- [Gav23] S. Gavrilova. Refined Littlewood identity for spin Hall–Littlewood symmetric rational functions. *Algebr. Comb.*, 6(1):37–51, 2023. arXiv:2104.09755.
- [GP20] U. Godreau and S. Prolhac. Spectral gaps of open TASEP in the maximal current phase. J. Phys. A Math. Theor., 53(38):385006, 2020. arXiv:2005.04461.

ALEXANDR GARBALI, JAN DE GIER, WILLIAM MEAD AND MICHAEL WHEELER

- [GS92a] L.-H. Gwa and H. Spohn. Bethe solution for the dynamical-scaling exponent of the noisy Burgers equation. *Phys. Rev. A*, 46(2):844, 1992.
- [GS92b] L.-H. Gwa and H. Spohn. Six-vertex model, roughened surfaces, and an asymmetric spin Hamiltonian. *Phys. Rev. Lett.*, 68(6):725–728, 1992.
- [He23] J. He. Boundary current fluctuations for the half space ASEP and six vertex model, 2023. arXiv:2303.16335.
- [IMS22] T. Imamura, M. Mucciconi, and T. Sasamoto. Solvable models in the KPZ class: approach through periodic and free boundary Schur measures, 2022. arXiv:2204.08420.
- [Ize87] A. G. Izergin. Partition function of a six-vertex model in a finite volume. *Proc. USSR Acad. Sci.*, 297(2):331–333, 1987.
- [KN96] A. Kirillov and M. Noumi. q-difference raising operators for macdonald polynomials and the integrality of transition coefficients. *CRM Proc. Lect. Notes*, 22, 1996. arXiv:q-alg/9605005.
- [KRS81] P. P. Kulish, N. Y. Reshetikhin, and E. K. Sklyanin. Yang–Baxter equation and representation theory: I. Lett. Math. Phys., 5:393–403, 1981.
- [Kup02] G. Kuperberg. Symmetry Classes of Alternating-Sign Matrices under One Roof. Ann. Math., 156(3):835–866, 2002. arXiv:math/0008184.
- [Skl88] E. K. Sklyanin. Boundary conditions for integrable quantum systems. J. Phys. A: Math. Gen., 21(10):2375, 1988.
- [Ste90] J. R. Stembridge. Nonintersecting paths, pfaffians, and plane partitions. Adv. Math., 83(1):96–131, 1990.
- [TW13] C. A. Tracy and H. Widom. The asymmetric simple exclusion process with an open boundary. J. Math. Phys., 54(10):103301, 2013. arXiv:1304.0800.
- [USW04] M. Uchiyama, T. Sasamoto, and M. Wadati. Asymmetric simple exclusion process with open boundaries and Askey–Wilson polynomials. J. Phys. A: Math. Gen., 37(18):4985, 2004. arXiv:cond-mat/0312457.
- [War08] O. Warnaar. Bisymmetric functions, Macdonald polynomials and \$\$\mathbf{s}\$\mathbf{1}\$ basic hypergeometric series. Compos. Math., 144:271–303, 2008. arXiv:math/0511333.
- [WZJ16] M. Wheeler and P. Zinn-Justin. Refined Cauchy/Littlewood identities and six-vertex model partition functions: III. Deformed bosons. *Adv. Math.*, 299:543–600, 2016. arXiv:1508.02236.
- [Zho22] C. Zhong. Stochastic symplectic ice. Lett. Math. Phys., 112(3):55, 2022. arXiv:2102.00660.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VICTORIA 3010, AUSTRALIA Email address: alexandr.garbali@unimelb.edu.au, jdgier@unimelb.edu.au, wmead@student.unimelb.edu.au, wheelerm@unimelb.edu.au

48