Orbit determination from one position vector and a very short arc of optical observations

Erica Scantamburlo^{*1}, Giovanni F. Gronchi^{†2}, and Giulio Baù^{‡2}

¹Department of Mechanical and Aerospace Engineering, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129, Italy

²Dipartimento di Matematica, University of Pisa, Largo B. Pontecorvo 5, 56127, Italy

December 25, 2023

Abstract

In this paper we address the problem of computing a preliminary orbit of a celestial body from one topocentric position vector and a very short arc (VSA) of optical observations. Using the conservation laws of the two-body dynamics, we write the problem as a system of 8 polynomial equations in 6 unknowns. We prove that this system is generically consistent, namely it admits solutions at least in the complex field. From this system we derive a univariate polynomial v of degree 8 in the unknown topocentric distance at the mean epoch of the VSA. Through Gröbner bases theory, we show that the degree of v is minimum among the degrees of all the univariate polynomials solving this problem. The proposed method is relevant for different purposes, e.g. the computation of a preliminary orbit of an Earth satellite with radar and optical observations, the detection of maneuvres of an Earth satellite, and the recovery of asteroids which are lost due to a planetary close encounter. We also show some numerical tests in the case of asteroids undergoing a close encounter with the Earth.

Keywords: Orbit determination - Keplerian integrals - Algebraic methods

1 Introduction

The problem of computing the orbit of celestial bodies has attracted the interest of scientists since a long time, see Laplace (1780), Lagrange (1783), Gauss (1809). The recent improvements in the observation technology have posed new interesting mathematical problems in this field. The number of asteroids observations performed by modern telescopes is very large. Usually they can be grouped into very short arcs (VSAs) of optical observations, however it is not easy to determine whether VSAs collected in different nights belong to the same observed objects. In general, from a VSA we can not compute a reliable preliminary orbit, but we can try to put together different VSAs to perform this task, see e.g. Milani et al. (2005). Assuming that two VSAs belong to the same asteroid,

^{*}erica.scantamburlo@polito.it

[†]giovanni.federico.gronchi@unipi.it

[‡]giulio.bau@unipi.it

we can write polynomial equations to compute a preliminary orbit using the conservation laws of the two-body dynamics. Recently, different ways to combine these integrals of motion have been developed and tested, see Taff and Hall (1977); Gronchi et al. (2010, 2011, 2015, 2017).

In this paper we address the problem of computing the orbit of a celestial body (hereafter, the OD problem) from one topocentric position $\mathcal{P}_1 = (\alpha_1, \delta_1, \rho_1)$ at epoch t_1 and a VSA of optical observations, from which we can derive an attributable $\mathcal{A}_2 = (\alpha_2, \delta_2, \dot{\alpha}_2, \dot{\delta}_2)$ at the mean epoch \bar{t}_2 of the arc. Here, α , δ , ρ denote respectively right ascension, declination and topocentric distance of the body, while $\dot{\alpha}, \dot{\delta}$ stand for the angular rates.

Using the conservation of angular momentum, energy and Laplace-Lenz vector we write the OD problem as a system of polynomial equations in the unknowns $\dot{\rho}_1$, $\dot{\alpha}_1$, $\dot{\delta}_1$, ρ_2 , $\dot{\rho}_2$, z_2 , where z_2 is an auxiliary variable, and the other variables allow us to obtain an orbit in spherical coordinates at the two epochs.

We prove that this polynomial system is consistent, namely it generically admits solutions, at least in the complex field, and we obtain a univariate polynomial \boldsymbol{v} of degree 8 in the unknown range ρ_2 to solve the OD problem. Through a computer algebra software we are also able to show that the degree of \boldsymbol{v} is minimum among the degrees of all the univariate polynomials in ρ_2 solving this problem.

The proposed method is relevant for different purposes, such as the computation of a preliminary orbit of an Earth satellite with radar and optical observations, the detection of maneuvres of an Earth satellite, the recovery of asteroids which are lost due to a planetary encounter, if the latter can be modeled as an instantaneous change of direction of the velocity vector like in Öpik theory (Öpik, 1976). Here we show the results of the application of our algorithm to the orbits of some near-Earth asteroids, whose positions have been changed to increase the effect of the close encounter with the Earth.

The paper is organized as follows: after recalling the expressions of the Keplerian integrals in Section 2, we introduce the OD problem in Section 3 as an overdetermined polynomial system, whose consistency is shown in Section 4, where the univariate polynomial \mathfrak{v} is derived. The minimality of the degree of \mathfrak{v} is proved in Section 5. In Section 6 we discuss the selection of the solutions. Finally, in Section 7 we present the numerical tests.

2 Keplerian integrals

Let us consider a celestial body whose dynamics can be modeled by Kepler's problem

$$\ddot{\boldsymbol{r}} = -\mu \frac{\boldsymbol{r}}{|\boldsymbol{r}|^3},\tag{1}$$

where \boldsymbol{r} and μ denote respectively its position and the gravitational parameter. Equation (1) admits the first integrals

$$\boldsymbol{c} = \boldsymbol{r} \times \dot{\boldsymbol{r}}, \qquad \mathcal{E} = \frac{1}{2} |\dot{\boldsymbol{r}}|^2 - \frac{\mu}{|\boldsymbol{r}|}, \qquad \boldsymbol{L} = \frac{1}{\mu} \dot{\boldsymbol{r}} \times \boldsymbol{c} - \frac{\boldsymbol{r}}{|\boldsymbol{r}|},$$
(2)

corresponding to the angular momentum, the energy, and the Laplace-Lenz vector of the body, respectively. We call *Keplerian integrals* these constants of motion. Note that we can write

$$\mu L = \left(|\dot{\boldsymbol{r}}|^2 - rac{\mu}{|\boldsymbol{r}|}
ight) \boldsymbol{r} - (\dot{\boldsymbol{r}} \cdot \boldsymbol{r}) \dot{\boldsymbol{r}}.$$

3 The OD problem

Let us introduce a reference frame whose origin is the center of force of the Keplerian dynamics. We consider the Keplerian motion of a celestial body around the origin observed from a moving point of view. Assume the position \boldsymbol{q} and the velocity $\dot{\boldsymbol{q}}$ of the observer are known functions of time. The position of the observed body is given by

$$\boldsymbol{r} = \boldsymbol{q} + \rho \hat{\boldsymbol{e}}^{\rho},\tag{3}$$

where ρ represents the topocentric distance, and

$$\hat{\boldsymbol{e}}^{\rho} = (\cos\delta\cos\alpha, \cos\delta\sin\alpha, \sin\delta)$$

is the line of sight.

Let us assume that we know the position vector

$$\mathcal{P}_1 = (\alpha_1, \delta_1, \rho_1) \in [-\pi, \pi) \times (-\pi/2, \pi/2) \times \mathbb{R}^+$$

at epoch t_1 , and a very short arc of optical observations of the same object, from which we compute the attributable

$$\mathcal{A}_2 = (\alpha_2, \delta_2, \dot{\alpha}_2, \dot{\delta}_2) \in [-\pi, \pi) \times (-\pi/2, \pi/2) \times \mathbb{R}^2$$

at the mean epoch \bar{t}_2 . We can combine these data to compute the quantities $\dot{\rho}_1$, ξ_1 , ζ_1 , ρ_2 , $\dot{\rho}_2$, where

$$\xi_1 = \rho_1 \dot{\alpha}_1 \cos \delta_1, \qquad \zeta_1 = \rho_1 \dot{\delta}_1,$$

which are missing to have a complete set of orbital elements at both epochs. For this purpose, we use the conservation of the integrals listed in (2).

Hereafter, we shall use subscripts 1, 2 for all the quantities relative to epochs t_1 , \bar{t}_2 . The dependence of position and velocity on the unknowns is given by

$$\dot{\mathbf{r}}_{1} = \dot{\mathbf{q}}_{1} + \dot{\rho}_{1} \hat{\mathbf{e}}_{1}^{\rho} + \xi_{1} \hat{\mathbf{e}}_{1}^{\alpha} + \zeta_{1} \hat{\mathbf{e}}_{1}^{\delta},
\mathbf{r}_{2} = \mathbf{q}_{2} + \rho_{2} \hat{\mathbf{e}}_{2}^{\rho},
\dot{\mathbf{r}}_{2} = \dot{\mathbf{q}}_{2} + \dot{\rho}_{2} \hat{\mathbf{e}}_{2}^{\rho} + \rho_{2} \hat{\mathbf{e}}_{2}^{\perp},$$
(4)

where

$$\hat{oldsymbol{e}}_2^\perp=\dot{lpha}_2\cos\delta_2\hat{oldsymbol{e}}_2^lpha+\dot{\delta}_2\hat{oldsymbol{e}}_2^\delta$$

is a known vector. Note that $\boldsymbol{r}_1 = \boldsymbol{q}_1 + \rho_1 \hat{\boldsymbol{e}}_1^{\rho}$ is known.

To write a polynomial system, we replace the term $\frac{\mu}{|\mathbf{r}_2|}$ with an auxiliary variable z_2 , independent from ρ_2 , and set

$$egin{aligned} &\mu \widetilde{m{L}}_2 = \Big(|\dot{m{r}}_2|^2 - z_2 \Big) m{r}_2 - (\dot{m{r}}_2 \cdot m{r}_2) \dot{m{r}}_2, \ &\widetilde{\mathcal{E}}_2 = rac{1}{2} |\dot{m{r}}_2|^2 - z_2. \end{aligned}$$

Lemma 1. The following relations hold:

$$\mu^{2} |\boldsymbol{L}_{1}|^{2} = 2\mathcal{E}_{1} |\boldsymbol{c}_{1}|^{2} + \mu^{2}, \qquad \mu^{2} |\boldsymbol{\widetilde{L}}_{2}|^{2} = 2\boldsymbol{\widetilde{\mathcal{E}}}_{2} |\boldsymbol{c}_{2}|^{2} + z_{2}^{2} |\boldsymbol{r}_{2}|^{2}.$$
(5)

Proof. From the definitions of L_1 and \widetilde{L}_2 we have

$$\begin{split} \mu^{2} |\boldsymbol{L}_{1}|^{2} &= |\dot{\boldsymbol{r}}_{1} \times \boldsymbol{c}_{1}|^{2} - 2\frac{\mu}{|\boldsymbol{r}_{1}|}(\dot{\boldsymbol{r}}_{1} \times \boldsymbol{c}_{1} \cdot \boldsymbol{r}_{1}) + \mu^{2}, \\ \mu^{2} |\widetilde{\boldsymbol{L}}_{2}|^{2} &= |\dot{\boldsymbol{r}}_{2} \times \boldsymbol{c}_{2}|^{2} - 2z_{2}(\dot{\boldsymbol{r}}_{2} \times \boldsymbol{c}_{2} \cdot \boldsymbol{r}_{2}) + z_{2}^{2}|\boldsymbol{r}_{2}|^{2}. \end{split}$$

We note that, at both epochs,

$$|\dot{\boldsymbol{r}} imes \boldsymbol{c}|^2 = |\dot{\boldsymbol{r}}|^2 |\boldsymbol{c}|^2, \qquad \dot{\boldsymbol{r}} imes \boldsymbol{c} \cdot \boldsymbol{r} = |\boldsymbol{c}|^2,$$

so that

$$\mu^{2} |\mathbf{L}_{1}|^{2} = |\mathbf{c}_{1}|^{2} \left(|\dot{\mathbf{r}}_{1}|^{2} - 2\frac{\mu}{|\mathbf{r}_{1}|} \right) + \mu^{2},$$

$$\mu^{2} |\widetilde{\mathbf{L}}_{2}|^{2} = |\mathbf{c}_{2}|^{2} \left(|\dot{\mathbf{r}}_{2}|^{2} - 2z_{2} \right) + z_{2}^{2} |\mathbf{r}_{2}|^{2}.$$

Relations (5) immediately follow from the definitions of \mathcal{E}_1 and $\widetilde{\mathcal{E}}_2$.

We consider the overdetermined polynomial system

$$\boldsymbol{c}_1 = \boldsymbol{c}_2, \quad \boldsymbol{L}_1 = \widetilde{\boldsymbol{L}}_2, \quad \mathcal{E}_1 = \widetilde{\mathcal{E}}_2, \quad z_2^2 |\boldsymbol{r}_2|^2 = \mu^2,$$
 (6)

consisting of 8 equations in the 6 unknowns $\dot{\rho}_1$, ξ_1 , ζ_1 , ρ_2 , $\dot{\rho}_2$, z_2 . System (6) corresponds to the equations of our OD problem.

4 Consistency of the equations

We will show the following property:

Theorem 1. The overdetermined polynomial system (6) is generically consistent, i.e. it always has solutions in the complex field for a generic choice of the data \mathcal{P}_1 , \mathcal{A}_2 , \mathbf{q}_1 , $\dot{\mathbf{q}}_1$, \mathbf{q}_2 , $\dot{\mathbf{q}}_2$.

The proof makes use of the results presented in the following subsections.

We start by noting that, because of relations (5), equation

$$z_2^2 |m{r}_2|^2 = \mu^2$$

is a consequence of the reduced system

$$c_1 = c_2, \quad L_1 = \widetilde{L}_2, \quad \mathcal{E}_1 = \widetilde{\mathcal{E}}_2,$$
 (7)

of 7 equations in 6 unknowns. Therefore, to prove Theorem 1 we show that system (7) is generically consistent.

Moreover, for a generical choice of the data \mathcal{P}_1 , \mathcal{A}_2 , \mathbf{q}_1 , $\dot{\mathbf{q}}_1$, \mathbf{q}_2 , $\dot{\mathbf{q}}_2$, system (7) is equivalent to

$$\mathfrak{q}_1 = \mathfrak{q}_2 = \mathfrak{q}_3 = \mathfrak{q}_4 = \mathfrak{q}_5 = \mathfrak{q}_6 = \mathfrak{q}_7 = 0,$$

where

$$\begin{aligned}
\mathbf{q}_{1} &= (\mathbf{c}_{1} - \mathbf{c}_{2}) \cdot \mathbf{W}_{12}, \\
\mathbf{q}_{2} &= (\mathbf{c}_{1} - \mathbf{c}_{2}) \cdot \mathbf{D}_{1} \times \mathbf{W}_{12}, \\
\mathbf{q}_{3} &= (\mathbf{c}_{1} - \mathbf{c}_{2}) \cdot \mathbf{D}_{2} \times \mathbf{W}_{12}, \\
\mathbf{q}_{4} &= \mu(\mathbf{L}_{1} - \tilde{\mathbf{L}}_{2}) \cdot \mathbf{D}_{1}, \\
\mathbf{q}_{5} &= \mu(\mathbf{L}_{1} - \tilde{\mathbf{L}}_{2}) \cdot \mathbf{D}_{2}, \\
\mathbf{q}_{6} &= \mu(\mathbf{L}_{1} - \tilde{\mathbf{L}}_{2}) \cdot (\mathbf{r}_{1} \times \hat{\mathbf{e}}_{2}^{\rho}), \\
\mathbf{q}_{7} &= \mathcal{E}_{1} - \tilde{\mathcal{E}}_{2},
\end{aligned}$$
(8)

with

$$\boldsymbol{D}_j = \boldsymbol{q}_j \times \hat{\boldsymbol{e}}_j^{\rho}, \quad j = 1, 2, \tag{9}$$

and

$$\boldsymbol{W}_{12} = \boldsymbol{D}_1 \times \boldsymbol{D}_2. \tag{10}$$

4.1 The angular momentum

The angular momenta c_1 and c_2 , written in terms of the unknowns, become

$$m{c}_1 = m{D}_1 \dot{
ho}_1 + m{N}_1 \xi_1 + m{O}_1 \zeta_1 + m{P}_1,$$

 $m{c}_2 = m{D}_2 \dot{
ho}_2 + m{E}_2
ho_2^2 + m{F}_2
ho_2 + m{G}_2,$

where D_1 , D_2 are defined as in (9), and

$$egin{aligned} &oldsymbol{N}_1 = oldsymbol{r}_1 imes \hat{oldsymbol{e}}_1^lpha, &oldsymbol{E}_2 = \dot{lpha}_2\cos\delta_2\hat{oldsymbol{e}}_2^\delta - \dot{\delta}_2\hat{oldsymbol{e}}_2^lpha, \ &oldsymbol{O}_1 = oldsymbol{r}_1 imes \hat{oldsymbol{e}}_1^\delta, &oldsymbol{F}_2 = \dot{lpha}_2\cos\delta_2(oldsymbol{q}_2 imes \hat{oldsymbol{e}}_2^lpha) + \dot{\delta}_2(oldsymbol{q}_2 imes \hat{oldsymbol{e}}_2^\delta) + (\hat{oldsymbol{e}}_2^
ho imes \dot{oldsymbol{q}}_2), \ &oldsymbol{P}_1 = oldsymbol{r}_1 imes \dot{oldsymbol{q}}_1, &oldsymbol{G}_2 = oldsymbol{q}_2 imes \dot{oldsymbol{q}}_2. \end{aligned}$$

Therefore, we can write

$$c_1 - c_2 = D_1 \dot{\rho}_1 - D_2 \dot{\rho}_2 + J(\xi_1, \zeta_1, \rho_2), \qquad (11)$$

with

$$J(\xi_1, \zeta_1, \rho_2) = N_1 \xi_1 + O_1 \zeta_1 - E_2 \rho_2^2 - F_2 \rho_2 + P_1 - G_2.$$
(12)

4.2 Elimination of variables

Using relations (11), (12), the polynomials q_1 , q_2 , q_3 defined in (8) can be written as

$$\begin{aligned} \mathbf{q}_1 &= Q_{100}^{(1)} \xi_1 + Q_{010}^{(1)} \zeta_1 + Q_{002}^{(1)} \rho_2^2 + Q_{001}^{(1)} \rho_2 + Q_{000}^{(1)}, \\ \mathbf{q}_2 &= -|\mathbf{W}_{12}|^2 \dot{\rho}_2 + Q_{100}^{(2)} \xi_1 + Q_{010}^{(2)} \zeta_1 + Q_{002}^{(2)} \rho_2^2 + Q_{001}^{(2)} \rho_2 + Q_{000}^{(2)}, \\ \mathbf{q}_3 &= |\mathbf{W}_{12}|^2 \dot{\rho}_1 + Q_{100}^{(3)} \xi_1 + Q_{010}^{(3)} \zeta_1 + Q_{002}^{(3)} \rho_2^2 + Q_{001}^{(3)} \rho_2 + Q_{000}^{(3)}, \end{aligned}$$

where

$$\begin{aligned} Q_{100}^{(1)} &= \mathbf{N}_1 \cdot \mathbf{W}_{12}, & Q_{010}^{(1)} &= \mathbf{O}_1 \cdot \mathbf{W}_{12}, \\ Q_{002}^{(1)} &= -\mathbf{E}_2 \cdot \mathbf{W}_{12}, & Q_{001}^{(1)} &= -\mathbf{F}_2 \cdot \mathbf{W}_{12}, \\ Q_{000}^{(1)} &= (\mathbf{G}_2 - \mathbf{P}_1) \cdot \mathbf{W}_{12}, \\ Q_{100}^{(2)} &= \mathbf{N}_1 \cdot \mathbf{D}_1 \times \mathbf{W}_{12}, & Q_{010}^{(2)} &= \mathbf{O}_1 \cdot \mathbf{D}_1 \times \mathbf{W}_{12}, \\ Q_{002}^{(2)} &= -\mathbf{E}_2 \cdot \mathbf{D}_1 \times \mathbf{W}_{12}, & Q_{001}^{(2)} &= -\mathbf{F}_2 \cdot \mathbf{D}_1 \times \mathbf{W}_{12}, \\ Q_{000}^{(2)} &= (\mathbf{P}_1 - \mathbf{G}_2) \cdot \mathbf{D}_1 \times \mathbf{W}_{12}, \\ Q_{100}^{(3)} &= \mathbf{N}_1 \cdot \mathbf{D}_2 \times \mathbf{W}_{12}, & Q_{010}^{(3)} &= \mathbf{O}_1 \cdot \mathbf{D}_2 \times \mathbf{W}_{12}, \\ Q_{002}^{(3)} &= -\mathbf{E}_2 \cdot \mathbf{D}_2 \times \mathbf{W}_{12}, & Q_{001}^{(3)} &= -\mathbf{F}_2 \cdot \mathbf{D}_2 \times \mathbf{W}_{12}, \\ Q_{002}^{(3)} &= (\mathbf{P}_1 - \mathbf{G}_2) \cdot \mathbf{D}_2 \times \mathbf{W}_{12}. \end{aligned}$$

In particular, the occurence of the variables $\dot{\rho}_1$, ξ_1 , ζ_1 , $\dot{\rho}_2$ in \mathfrak{q}_1 , $\mathfrak{q}_2,\mathfrak{q}_3$ is at most linear. From the definitions above, we obtain

$$\begin{aligned} Q_{100}^{(1)} &= (\boldsymbol{N}_1 \times \boldsymbol{D}_1) \cdot \boldsymbol{D}_2 = \left[(\boldsymbol{r}_1 \times \hat{\boldsymbol{e}}_1^{\alpha}) \times (\boldsymbol{r}_1 \times \hat{\boldsymbol{e}}_1^{\rho}) \right] \cdot \boldsymbol{D}_2 \\ &= \left[(\boldsymbol{r}_1 \times \hat{\boldsymbol{e}}_1^{\alpha}) \cdot \hat{\boldsymbol{e}}_1^{\rho} \right] \boldsymbol{r}_1 \cdot \boldsymbol{D}_2 = -(\boldsymbol{r}_1 \cdot \hat{\boldsymbol{e}}_1^{\delta})(\boldsymbol{r}_1 \cdot \boldsymbol{D}_2), \\ Q_{010}^{(1)} &= (\boldsymbol{O}_1 \times \boldsymbol{D}_1) \cdot \boldsymbol{D}_2 = \left[(\boldsymbol{r}_1 \times \hat{\boldsymbol{e}}_1^{\delta}) \times (\boldsymbol{r}_1 \times \hat{\boldsymbol{e}}_1^{\rho}) \right] \cdot \boldsymbol{D}_2 \\ &= \left[(\boldsymbol{r}_1 \times \hat{\boldsymbol{e}}_1^{\delta}) \cdot \hat{\boldsymbol{e}}_1^{\rho} \right] \boldsymbol{r}_1 \cdot \boldsymbol{D}_2 = (\boldsymbol{r}_1 \cdot \hat{\boldsymbol{e}}_1^{\alpha})(\boldsymbol{r}_1 \cdot \boldsymbol{D}_2). \end{aligned}$$

Therefore, the coefficients $Q_{100}^{(1)}$ and $Q_{010}^{(1)}$ are both vanishing iff

$$\boldsymbol{r}_1 \cdot \boldsymbol{D}_2 = 0$$
 or $\boldsymbol{D}_1 = \boldsymbol{0}$.

In fact, the second condition in the alternative above is equivalent to

$$\boldsymbol{r}_1 \cdot \hat{\boldsymbol{e}}_1^{\alpha} = \boldsymbol{r}_1 \cdot \hat{\boldsymbol{e}}_1^{\delta} = 0$$

Thus, for a generic choice of the data, the angular momentum equations allow us to eliminate $\dot{\rho}_1$, $\dot{\rho}_2$ and one variable between ξ_1 and ζ_1 . Assuming $Q_{100}^{(1)} \neq 0$, we choose to eliminate ξ_1 , which can be written as

$$\xi_1 = -\frac{Q_{010}^{(1)}\zeta_1 + \sum_{h=0}^2 Q_{00h}^{(1)}\rho_2^h}{Q_{100}^{(1)}}.$$
(13)

Substituting (13) in equations $q_3 = q_2 = 0$ and assuming $W_{12} \neq 0$ we find

$$\dot{\rho}_{1} = -\frac{1}{|\boldsymbol{W}_{12}|^{2}Q_{100}^{(1)}} \left[\left(Q_{010}^{(3)}Q_{100}^{(1)} - Q_{010}^{(1)}Q_{100}^{(3)} \right) \zeta_{1} + \sum_{h=0}^{2} \left(Q_{00h}^{(3)}Q_{100}^{(1)} - Q_{00h}^{(1)}Q_{100}^{(3)} \right) \rho_{2}^{h} \right] \quad (14)$$

$$\dot{\rho}_{2} = \frac{1}{|\boldsymbol{W}_{12}|^{2}Q_{100}^{(1)}} \Big[\Big(Q_{010}^{(2)}Q_{100}^{(1)} - Q_{010}^{(1)}Q_{100}^{(2)} \Big) \zeta_{1} + \sum_{h=0}^{2} \Big(Q_{00h}^{(2)}Q_{100}^{(1)} - Q_{00h}^{(1)}Q_{100}^{(2)} \Big) \rho_{2}^{h} \Big].$$
(15)

Hence, by using relations (13), (14), and (15) we can eliminate the variables ξ_1 , $\dot{\rho}_1$, $\dot{\rho}_2$ in the generators \mathfrak{q}_4 , \mathfrak{q}_5 , \mathfrak{q}_6 , \mathfrak{q}_7 . The resulting polynomials, named $\tilde{\mathfrak{q}}_4$, $\tilde{\mathfrak{q}}_5$, $\tilde{\mathfrak{q}}_6$, $\tilde{\mathfrak{q}}_7$, can be written as

$$egin{aligned} & ilde{\mathfrak{q}}_4 = (m{r}_2 \cdot m{D}_1) z_2 + h_4(\zeta_1,
ho_2), \ & ilde{\mathfrak{q}}_5 = & ilde{\mathfrak{q}}_5(\zeta_1,
ho_2), \ & ilde{\mathfrak{q}}_6 = -(m{r}_1 \cdot m{D}_2) z_2 + h_6(\zeta_1,
ho_2), \ & ilde{\mathfrak{q}}_7 = z_2 + h_7(\zeta_1,
ho_2), \end{aligned}$$

for some bivariate polynomials h_4 , h_6 , h_7 . For later use we observe that

$$\tilde{\mathfrak{q}}_5 = P_{12}^{(5)}\zeta_1\rho_2^2 + P_{11}^{(5)}\zeta_1\rho_2 + P_{10}^{(5)}\zeta_1 + P_{04}^{(5)}\rho_2^4 + P_{03}^{(5)}\rho_2^3 + P_{02}^{(5)}\rho_2^2 + P_{01}^{(5)}\rho_2 + P_{00}^{(5)},$$

for some coefficients $P_{ij}^{(5)}$. Moreover, using equation $\tilde{\mathfrak{q}}_7 = 0$, we can eliminate z_2 from $\tilde{\mathfrak{q}}_4$ and $\tilde{\mathfrak{q}}_6$. In particular, with this elimination $\tilde{\mathfrak{q}}_6$ becomes

$$\mathfrak{p}_{6} = P_{20}^{(6)}\zeta_{1}^{2} + P_{12}^{(6)}\zeta_{1}\rho_{2}^{2} + P_{11}^{(6)}\zeta_{1}\rho_{2} + P_{10}^{(6)}\zeta_{1} + P_{04}^{(6)}\rho_{2}^{4} + P_{03}^{(6)}\rho_{2}^{3} + P_{02}^{(6)}\rho_{2}^{2} + P_{01}^{(6)}\rho_{2} + P_{00}^{(6)},$$

for some coefficients $P_{ij}^{(6)}$.

4.3 A redundant relation

Here, we prove that equation $q_4 = 0$ can be dropped from system (7) without losing any solution. In particular, we show:

Proposition 1. If $r_1 \cdot D_2 \neq 0$, the polynomial q_4 is generated by

$$\mathfrak{q}_1,\,\mathfrak{q}_2,\,\mathfrak{q}_3,\,\mathfrak{q}_6$$

Proof. Let us consider the polynomials $\tilde{\mathfrak{q}}_4$, $\tilde{\mathfrak{q}}_6$ obtained from \mathfrak{q}_4 , \mathfrak{q}_6 by eliminating $\dot{\rho}_1$, $\dot{\rho}_2$, ξ_1 through relations $\mathfrak{q}_1 = \mathfrak{q}_2 = \mathfrak{q}_3 = 0$. We note that

$$ilde{\mathfrak{q}}_4 = \sum_{j=1}^3 \mathfrak{a}_j \mathfrak{q}_j + \mathfrak{q}_4, \qquad ilde{\mathfrak{q}}_6 = \sum_{j=1}^3 \mathfrak{b}_j \mathfrak{q}_j + \mathfrak{q}_6,$$

for some polynomials \mathfrak{a}_j , \mathfrak{b}_j . We prove that $\tilde{\mathfrak{q}}_6$ divides $\tilde{\mathfrak{q}}_4$; in particular,

$$\tilde{\mathfrak{q}}_4 = -\frac{\boldsymbol{r}_2 \cdot \boldsymbol{D}_1}{\boldsymbol{r}_1 \cdot \boldsymbol{D}_2} \,\tilde{\mathfrak{q}}_6. \tag{16}$$

For this purpose, we first note that system $q_1 = q_2 = q_3 = 0$ is equivalent to $c_1 = c_2$ and, inserting the last relation in the expressions of q_4 , q_6 , after eliminating $\dot{\rho}_1$, $\dot{\rho}_2$, ξ_1 , we obtain

$$\begin{split} \widetilde{\mathfrak{q}}_4 &= (oldsymbol{c}_1 \cdot \hat{oldsymbol{e}}_1^
ho) \left[oldsymbol{r}_1 \cdot (\dot{oldsymbol{r}}_1 - \dot{oldsymbol{r}}_2)
ight] + z_2(oldsymbol{r}_2 \cdot oldsymbol{D}_1), \ \widetilde{\mathfrak{q}}_6 &= (oldsymbol{c}_2 \cdot \hat{oldsymbol{e}}_2^
ho) \left[oldsymbol{r}_1 \cdot (\dot{oldsymbol{r}}_1 - \dot{oldsymbol{r}}_2)
ight] - z_2(oldsymbol{r}_1 \cdot oldsymbol{D}_2), \end{split}$$

where c_1 , c_2 , r_2 , \dot{r}_1 , \dot{r}_2 are meant as functions of ζ_1 , ρ_2 only. More details about these computations can be found in Appendix A.

We have

$$oldsymbol{r}_2\cdotoldsymbol{D}_1=-(oldsymbol{r}_1 imesoldsymbol{r}_2)\cdot\hat{oldsymbol{e}}_1^
ho,\qquadoldsymbol{r}_1\cdotoldsymbol{D}_2=(oldsymbol{r}_1 imesoldsymbol{r}_2)\cdot\hat{oldsymbol{e}}_2^
ho.$$

Moreover, $c_1 = c_2$ implies that $r_1 \times r_2$ is parallel to c_1 and c_2 . Hence

$$(oldsymbol{c}_2\cdot\hat{oldsymbol{e}}_2^
ho)\left[(oldsymbol{r}_1 imesoldsymbol{r}_2)\cdot\hat{oldsymbol{e}}_1^
ho
ight]=(oldsymbol{c}_1\cdot\hat{oldsymbol{e}}_1^
ho)\left[(oldsymbol{r}_1 imesoldsymbol{r}_2)\cdot\hat{oldsymbol{e}}_2^
ho
ight],$$

that is

$$-(\boldsymbol{c}_2\cdot\hat{\boldsymbol{e}}_2^
ho)(\boldsymbol{r}_2\cdot\boldsymbol{D}_1)=(\boldsymbol{c}_1\cdot\hat{\boldsymbol{e}}_1^
ho)(\boldsymbol{r}_1\cdot\boldsymbol{D}_2).$$

The latter relation immediately yields (16).

Setting

$$A = -\frac{\boldsymbol{r}_2 \cdot \boldsymbol{D}_1}{\boldsymbol{r}_1 \cdot \boldsymbol{D}_2},$$

we can write

$$\mathfrak{q}_4 = \tilde{\mathfrak{q}}_4 - \sum_{j=1}^3 \mathfrak{a}_j \mathfrak{q}_j = A \tilde{\mathfrak{q}}_6 - \sum_{j=1}^3 \mathfrak{a}_j \mathfrak{q}_j = \sum_{j=1}^3 (A \mathfrak{b}_j - \mathfrak{a}_j) \mathfrak{q}_j + A \mathfrak{q}_6,$$

which concludes the proof.

4.4 The univariate polynomial

We compute a univariate polynomial $\mathfrak{u}(\rho_2)$ of degree 8, which is a consequence of the equations in (7). Let us consider the polynomial system

$$\mathfrak{q}_1 = \mathfrak{q}_2 = \mathfrak{q}_3 = \mathfrak{q}_5 = \mathfrak{q}_6 = \mathfrak{q}_7 = 0 \tag{17}$$

including all the polynomials in (8) except q_4 . To solve the OD problem, we solve the polynomial system defined by (17).

Now, we compute the resultant of $\tilde{\mathfrak{q}}_5$ and \mathfrak{p}_6 with respect to ζ_1 , which is denoted by $\mathfrak{v}(\rho_2)$. For this purpose, the terms in $\tilde{\mathfrak{q}}_5$ and \mathfrak{p}_6 are grouped in the following way:

$$\begin{split} \tilde{\mathfrak{q}}_5 &= a_1(\rho_2)\zeta_1 + a_0(\rho_2), \\ \mathfrak{p}_6 &= P_{20}^{(6)}\zeta_1^2 + b_1(\rho_2)\zeta_1 + b_0(\rho_2), \end{split}$$

where

$$a_{1}(\rho_{2}) = P_{12}^{(5)}\rho_{2}^{2} + P_{11}^{(5)}\rho_{2} + P_{10}^{(5)},$$

$$a_{0}(\rho_{2}) = P_{04}^{(5)}\rho_{2}^{4} + P_{03}^{(5)}\rho_{2}^{3} + P_{02}^{(5)}\rho_{2}^{2} + P_{01}^{(5)}\rho_{2} + P_{00}^{(5)},$$

$$b_{1}(\rho_{2}) = P_{12}^{(6)}\rho_{2}^{2} + P_{11}^{(6)}\rho_{2} + P_{10}^{(6)},$$

$$b_{0}(\rho_{2}) = P_{04}^{(6)}\rho_{2}^{4} + P_{03}^{(6)}\rho_{2}^{3} + P_{02}^{(6)}\rho_{2}^{2} + P_{01}^{(6)}\rho_{2} + P_{00}^{(6)}.$$

Therefore, we have

$$\mathfrak{v}(\rho_2) = a_1(\rho_2)a_0(\rho_2)b_1(\rho_2) - a_0^2(\rho_2)P_{20}^{(6)} - b_0(\rho_2)a_1^2(\rho_2),$$

which has degree 8.

We can conclude the proof of Theorem 1. In fact, $\mathfrak{v}(\rho_2)$ has (generically) 8 complex roots. For each of these roots, equation $\tilde{\mathfrak{q}}_5 = 0$ gives a unique value of ζ_1 . Then, equation $\tilde{\mathfrak{q}}_7 = 0$ and relations (13), (14), (15), respectively, allow us to compute unique values for the remaining components z_2 , ξ_1 , $\dot{\rho}_1$, $\dot{\rho}_2$ of the solutions.

5 An optimal property

We show that the univariate polynomial $\mathfrak{v}(\rho_2)$ has the minimal degree among all the univariate polynomials in the variable ρ_2 that are algebraic consequences of

$$c_1 - c_2, \ \mu(L_1 - \widetilde{L}_2), \ \mathcal{E}_1 - \widetilde{\mathcal{E}}_2.$$

More precisely, let us introduce the polynomial ideal

$$I = \langle \boldsymbol{c}_1 - \boldsymbol{c}_2, \ \mu(\boldsymbol{L}_1 - \widetilde{\boldsymbol{L}}_2), \ \mathcal{E}_1 - \widetilde{\mathcal{E}}_2 \rangle \subseteq \mathbb{R}[\dot{\rho}_1, \xi_1, \zeta_1, \rho_2, \dot{\rho}_2, z_2].$$

We will show the following result:

Theorem 2. For a generic choice of the data we can find a Gröbner basis \mathcal{G} in $\mathbb{R}[\dot{\rho}_1, \xi_1, \zeta_1, \rho_2, \dot{\rho}_2, z_2]$ of the ideal I for the lexicographic order

$$\dot{\rho}_1 \succ \dot{\rho}_2 \succ \xi_1 \succ z_2 \succ \zeta_1 \succ \rho_2.$$

such that $\mathfrak{v} \in \mathcal{G}$.

Proof. As a consequence of Proposition 1 we obtain

$$I = \langle \mathfrak{q}_1, \mathfrak{q}_2, \mathfrak{q}_3, \mathfrak{q}_5, \mathfrak{q}_6, \mathfrak{q}_7 \rangle,$$

where we dropped q_4 from the set of generators. Let us consider the elimination ideal

$$I' = I \cap \mathbb{R}[\zeta_1, \rho_2] = \langle \tilde{\mathfrak{q}}_5, \mathfrak{p}_6 \rangle$$

We compute a Gröbner basis \mathcal{G}' of the ideal I' for the lexicographic order $\zeta_1 \succ \rho_2$ using the software *Mathematica*¹: in this basis we have a univariate polynomial

$$\mathfrak{u}(\rho_2) = \sum_{i=0}^8 u_i \rho_2^i,$$

for some constant coefficients u_i . The roots of \mathfrak{u} are the only values of the ρ_2 components of the solutions of system (17).

We note that, since the resultant v is a univariate polynomial of degree 8, v is necessarily proportional to u, i.e.

$$\mathfrak{v} = \kappa \mathfrak{u}, \qquad \kappa \in \mathbb{R} \setminus \{0\}.$$

In place of the generators q_1 , q_2 , q_3 , we can consider

$$\begin{split} \tilde{\mathfrak{q}}_1 &= \xi_1 + h_1(\zeta_1, \rho_2), \\ \tilde{\mathfrak{q}}_2 &= \dot{\rho}_2 + h_2(\zeta_1, \rho_2), \\ \tilde{\mathfrak{q}}_3 &= \dot{\rho}_1 + h_3(\zeta_1, \rho_2), \end{split}$$

where the polynomials h_1 , h_2 , h_3 are defined by relations (13), (14), (15).

The set

$$\mathcal{G} = \{ \tilde{\mathfrak{q}}_1, \tilde{\mathfrak{q}}_2, \tilde{\mathfrak{q}}_3, \tilde{\mathfrak{q}}_7 \} \cup \mathcal{G}'$$

is a Gröbner basis of the ideal I. In fact, the leading term of every polynomial in I is divisible by the leading term of one polynomial in \mathcal{G} .

As a consequence of the previous theorem we obtain the following property:

Corollary 1. The polynomial v has the minimum degree among the univariate polynomials in ρ_2 belonging to I.

6 Selecting the solutions

From the solutions of (7), we first discard the ones with non-real components and the ones with $\rho_2, z_2 \leq 0$. From the remaining solutions we can compute Keplerian orbital elements at epochs

$$\tilde{t}_1 = t_1 - \frac{\rho_1}{c}, \qquad \tilde{t}_2^{(j)} = \bar{t}_2 - \frac{\rho_2^{(j)}}{c},$$

where c is the speed of light and $\rho_2^{(j)}$ is the value of ρ_2 for the j-th solution.

We emphasize that the accepted solutions of the system are such that the Keplerian elements at the two epochs are the same. In fact, from $c_1 - c_2 = 0$ we get

 $i_1 = i_2,$ $\Omega_1 = \Omega_2,$ $a_1(1 - e_1^2) = a_2(1 - e_2^2),$

¹Wolfram Research, Inc. Mathematica, Version 12.1, https://www.wolfram.com/mathematica.

where i, Ω , a and e denote the inclination, ascending node, semi-major axis and eccentricity. In particular, from $z_2^2 |\mathbf{r}_2|^2 - \mu^2 = 0$, which is a consequence of system (7), we get $z_2 = \pm \frac{\mu}{|\mathbf{r}_2|}$. Having discarded the solutions with $z_2 \leq 0$, we obtain

$$\mu(\boldsymbol{L}_1 - \widetilde{\boldsymbol{L}}_2) = \boldsymbol{0} \quad \Leftrightarrow \quad \mu(\boldsymbol{L}_1 - \boldsymbol{L}_2) = \boldsymbol{0},$$

which implies

$$e_1 = e_2, \qquad \omega_1 = \omega_2.$$

Since the eccentrities e_1 , e_2 are equal, then also

$$a_1 = a_2.$$

We observe that for the other orbit determination methods using the Keplerian integrals (Gronchi et al., 2010, 2011, 2015) the trajectories of the solutions at the two epochs are not necessarily the same.

In the following subsections we explain how to choose or discard the remaining solutions.

6.1 Selection without covariance

In case of multiple solutions, labeled with (j), we make our choice according to the following procedure. We propagate each of the computed orbits, referring to epoch $\tilde{t}_2^{(j)}$, backward to epoch \tilde{t}_1 in the framework of Kepler's dynamics, and consider the norm of the differences $\rho_1^{(j)} - \mathcal{P}_1$, where

$$\boldsymbol{\rho}_{1}^{(j)} = \boldsymbol{r}^{(j)}(\tilde{t}_{1}) - \boldsymbol{q}(t_{1})$$

is the topocentric position of the *j*-th solution propagated to epoch \tilde{t}_1 . Note that in general we have

$$|\boldsymbol{\rho}_1^{(j)}| \neq \rho_1.$$

We select the solution attaining the minimum value

$$\min_{j} | oldsymbol{
ho}_{1}^{(j)} - \mathcal{P}_{1} |.$$

6.2 Selection using covariance

Assume that the data

$$\mathcal{D} = (\mathcal{P}_1, \mathcal{A}_2),$$

where

$$\mathcal{P}_1 = (\alpha_1, \delta_1, \rho_1), \qquad \mathcal{A}_2 = (\alpha_2, \delta_2, \dot{\alpha}_2, \delta_2),$$

have a covariance matrix

$$\Gamma_{\mathcal{D}} = \left[\begin{array}{cc} \Gamma_{\mathcal{P}_1} & 0\\ 0 & \Gamma_{\mathcal{A}_2} \end{array} \right].$$

Let

$$S = S(D) = (V_1(D), \mathcal{R}_2(D)),$$

with

$$\mathcal{V}_1 = (\xi_1, \zeta_1, \dot{\rho}_1), \qquad \mathcal{R}_2 = (\rho_2, \dot{\rho}_2)$$

be a solution of

$$\boldsymbol{\Phi}(\mathcal{S};\mathcal{D}) = \begin{pmatrix} \boldsymbol{c}_1 - \boldsymbol{c}_2 \\ [\mu \boldsymbol{L}_1 + (\dot{\boldsymbol{r}}_2 \cdot \boldsymbol{r}_2)\dot{\boldsymbol{r}}_2] \cdot \boldsymbol{q}_2 \times \boldsymbol{r}_2 \\ [-(\dot{\boldsymbol{r}}_1 \cdot \boldsymbol{r}_1)\dot{\boldsymbol{r}}_1 - (\frac{1}{2}|\dot{\boldsymbol{r}}_2|^2 + \mathcal{E}_1)\boldsymbol{r}_2 + (\dot{\boldsymbol{r}}_2 \cdot \boldsymbol{r}_2)\dot{\boldsymbol{r}}_2] \cdot \boldsymbol{r}_1 \times (\boldsymbol{r}_2 - \boldsymbol{q}_2) \end{pmatrix} = \boldsymbol{0}.$$
(18)

Note that, generically, the solutions S of system (18) correspond to the S components of the solutions of system (17).

Let $\mathbf{E}_{car} = (\mathbf{E}_{car}^{(1)}, \mathbf{E}_{car}^{(2)})$ and $\mathbf{E}_{att} = (\mathbf{E}_{att}^{(1)}, \mathbf{E}_{att}^{(2)})$ be the vectors of the Cartesian coordinates and the attributable elements at epochs \tilde{t}_1 and \tilde{t}_2 .² Let us introduce the transformation $\mathcal{T}_{att}^{car} : \mathbf{E}_{att} \to \mathbf{E}_{car}$ by (3), (4) for both epochs, and consider the map Ψ defined by $\Phi = \Psi \circ \mathcal{T}_{att}^{car}$. Equation $\Phi = \mathbf{0}$ is equivalent to the system

$$c_1 - c_2 = 0, \quad \mu(L_1 - \tilde{L}_2) \cdot D_2 = 0, \quad \mu(L_1 - \tilde{L}_2) \cdot r_1 \times (r_2 - q_2) = 0.$$

We introduce the vector

$$\widetilde{\mathcal{V}}_1 = (\dot{\alpha}_1, \dot{\delta}_1, \dot{\rho}_1).$$

The covariance matrix of the Cartesian coordinates at epoch \tilde{t}_1 is

$$\Gamma_{car}^{(1)} = \frac{\partial \mathbf{E}_{car}^{(1)}}{\partial \mathcal{D}} \Gamma_{\mathcal{D}} \left[\frac{\partial \mathbf{E}_{car}^{(1)}}{\partial \mathcal{D}} \right]^T \,,$$

with

$$\frac{\partial \mathbf{E}_{car}^{(1)}}{\partial \mathcal{D}} = \frac{\partial \mathbf{E}_{car}^{(1)}}{\partial \mathbf{E}_{att}^{(1)}} \frac{\partial \mathbf{E}_{att}^{(1)}}{\partial \mathcal{D}}, \qquad \frac{\partial \mathbf{E}_{att}^{(1)}}{\partial \mathcal{D}} = M \begin{bmatrix} I_3 & O_{3 \times 4} \\ & \frac{\partial \widetilde{\mathcal{V}}_1}{\partial \mathcal{D}} \end{bmatrix},$$

where

$$M = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

is the matrix that exchanges the 3-rd, 4-th and 5-th lines, and

$$\frac{\partial \widetilde{\mathcal{V}}_1}{\partial \mathcal{D}} = \frac{\partial \widetilde{\mathcal{V}}_1}{\partial \mathcal{V}_1} \frac{\partial \mathcal{V}_1}{\partial \mathcal{D}}, \qquad \frac{\partial \widetilde{\mathcal{V}}_1}{\partial \mathcal{V}_1} = \begin{bmatrix} \frac{1}{\rho_1 \cos \delta_1} & 0 & 0\\ 0 & \frac{1}{\rho_1} & 0\\ 0 & 0 & 1 \end{bmatrix}.$$

From the implicit function theorem, we have

$$\begin{aligned} \frac{\partial \widetilde{\mathcal{S}}}{\partial \mathcal{D}}(\mathcal{D}) &= -\left[\frac{\partial \Phi}{\partial \widetilde{\mathcal{S}}}(\mathbf{E}_{att})\right]^{-1} \frac{\partial \Phi}{\partial \mathcal{D}}(\mathbf{E}_{att}),\\ \frac{\partial \Phi}{\partial \widetilde{\mathcal{S}}} &= \left(\frac{\partial \Psi}{\partial \mathbf{E}_{car}} \circ \mathcal{T}_{att}^{car}\right) \frac{\partial \mathcal{T}_{att}^{car}}{\partial \widetilde{\mathcal{S}}}, \qquad \frac{\partial \Phi}{\partial \mathcal{D}} = \left(\frac{\partial \Psi}{\partial \mathbf{E}_{car}} \circ \mathcal{T}_{att}^{car}\right) \frac{\partial \mathcal{T}_{att}^{car}}{\partial \mathcal{D}},\\ \widetilde{\mathcal{S}} &= (\widetilde{\mathcal{V}}_1, \mathcal{R}_2), \qquad \frac{\partial \widetilde{\mathcal{S}}}{\partial \mathcal{S}} = \left[\begin{array}{c} \frac{\partial \widetilde{\mathcal{V}}_1}{\partial \mathcal{V}_1} & O_{3\times 2}\\ O_{2\times 3} & I_2\end{array}\right]. \end{aligned}$$

with

² for simplicity, in this section we drop the label (j), referring to the possible multiple solutions.

The matrices $\partial \mathcal{T}_{att}^{car}/\partial \tilde{S}$ and $\partial \mathcal{T}_{att}^{car}/\partial \mathcal{D}$ are respectively made by columns 3, 4, 6, 11, 12 and by columns 1, 2, 5, 7, 8, 9, 10 of $\partial \mathbf{E}_{car}/\partial \mathbf{E}_{att}$. The covariance matrix of the Cartesian coordinates at epoch \tilde{t}_2 is given by

$$\Gamma_{car}^{(2)} = \frac{\partial \mathbf{E}_{car}^{(2)}}{\partial \mathcal{D}} \Gamma_{\mathcal{D}} \left[\frac{\partial \mathbf{E}_{car}^{(2)}}{\partial \mathcal{D}} \right]^{T},$$

with

$$\frac{\partial \mathbf{E}_{car}^{(2)}}{\partial \mathcal{D}} = \frac{\partial \mathbf{E}_{car}^{(2)}}{\partial \mathbf{E}_{att}^{(2)}} \frac{\partial \mathbf{E}_{att}^{(2)}}{\partial \mathcal{D}}, \qquad \frac{\partial \mathbf{E}_{att}^{(2)}}{\partial \mathcal{D}} = \begin{bmatrix} O_{4\times3} & I_4 \\ \frac{\partial \mathcal{R}_2}{\partial \mathcal{D}} \end{bmatrix}$$

and

$$\frac{\partial(\rho_2,\dot{\rho}_2)}{\partial\widetilde{\mathcal{S}}} = \frac{\partial(\rho_2,\dot{\rho}_2)}{\partial\widetilde{\mathcal{S}}}\frac{\partial\widetilde{\mathcal{S}}}{\partial\mathcal{D}}, \qquad \frac{\partial(\rho_2,\dot{\rho}_2)}{\partial\widetilde{\mathcal{S}}} = \begin{bmatrix} 0 & 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

For a given vector $\mathbf{u} \in \mathbb{R}^3$ we define the map

$$\mathbb{R}^{3} \ni (u_{1}, u_{2}, u_{3}) = \mathbf{u} \mapsto \widehat{\mathbf{u}} = \begin{bmatrix} 0 & -u_{3} & u_{2} \\ u_{3} & 0 & -u_{1} \\ -u_{2} & u_{1} & 0 \end{bmatrix} \in so(3).$$

Then, using $\hat{\mathbf{u}}^T = -\hat{\mathbf{u}}$, we have

$$\frac{\partial \Psi}{\partial \mathbf{E}_{car}} = \begin{bmatrix} -\hat{r_1} & \hat{r_1} & \hat{r_2} & -\hat{r_2} \\ \frac{\partial \Phi_4}{\partial r_1} & \frac{\partial \Phi_4}{\partial \dot{r_1}} & \frac{\partial \Phi_4}{\partial r_2} & \frac{\partial \Phi_4}{\partial \dot{r_2}} \\ \frac{\partial \Phi_5}{\partial r_1} & \frac{\partial \Phi_5}{\partial \dot{r_1}} & \frac{\partial \Phi_5}{\partial r_2} & \frac{\partial \Phi_5}{\partial \dot{r_2}} \end{bmatrix},$$

where

$$\begin{split} \Phi_4 &= \left[\mu \boldsymbol{L}_1 + (\dot{\boldsymbol{r}}_2 \cdot \boldsymbol{r}_2) \dot{\boldsymbol{r}}_2\right] \cdot \boldsymbol{q}_2 \times \boldsymbol{r}_2, \\ \Phi_5 &= \left[-(\dot{\boldsymbol{r}}_1 \cdot \boldsymbol{r}_1) \dot{\boldsymbol{r}}_1 - (\frac{1}{2} |\dot{\boldsymbol{r}}_2|^2 + \mathcal{E}_1) \boldsymbol{r}_2 + (\dot{\boldsymbol{r}}_2 \cdot \boldsymbol{r}_2) \dot{\boldsymbol{r}}_2 \right] \cdot \boldsymbol{r}_1 \times (\boldsymbol{r}_2 - \boldsymbol{q}_2), \end{split}$$

and

$$\begin{split} \frac{\partial \Phi_4}{\partial r_1} &= \left(|\dot{r}_1|^2 - \frac{\mu}{|r_1|} \right) (q_2 \times r_2) - \dot{r}_1 \left[\dot{r}_1 \cdot (q_2 \times r_2) \right] + \frac{\mu}{|r_1|^3} r_1 \left[r_1 \cdot (q_2 \times r_2) \right], \\ \frac{\partial \Phi_4}{\partial \dot{r}_1} &= 2 \dot{r}_1 \left[r_1 \cdot (q_2 \times r_2) \right] - r_1 \left[\dot{r}_1 \cdot (q_2 \times r_2) \right] - (\dot{r}_1 \cdot r_1) (q_2 \times r_2), \\ \frac{\partial \Phi_4}{\partial r_2} &= \dot{r}_2 \left[\dot{r}_2 \cdot (q_2 \times r_2) \right] + \left[\left(\left| \dot{r}_1 \right|^2 - \frac{\mu}{|r_1|} \right) r_1 - (\dot{r}_1 \cdot r_1) \dot{r}_1 + (\dot{r}_2 \cdot r_2) \dot{r}_2 \right] \times q_2, \\ \frac{\partial \Phi_4}{\partial \dot{r}_2} &= r_2 \left[\dot{r}_2 \cdot (q_2 \times r_2) \right] + (\dot{r}_2 \cdot r_2) (q_2 \times r_2), \\ \frac{\partial \Phi_5}{\partial r_1} &= -\dot{r}_1 \left[\dot{r}_1 \cdot r_1 \times (r_2 - q_2) \right] - \frac{\mu}{|r_1|^3} r_1 \left[r_2 \cdot r_1 \times (r_2 - q_2) \right] \\ &+ (r_2 - q_2) \times \left[- (\dot{r}_1 \cdot r_1) \dot{r}_1 - \left(\frac{|\dot{r}_2|^2}{2} - \frac{|\dot{r}_1|^2}{2} + \frac{\mu}{|r_1|} \right) r_2 + (\dot{r}_2 \cdot r_2) \dot{r}_2 \right], \\ \frac{\partial \Phi_5}{\partial \dot{r}_1} &= -r_1 \left[\dot{r}_1 \cdot r_1 \times (r_2 - q_2) \right] - (\dot{r}_1 \cdot r_1) \left[r_1 \times (r_2 - q_2) \right] + \dot{r}_1 \left[r_2 \cdot r_1 \times (r_2 - q_2) \right] \\ &+ \left[- (\dot{r}_1 \cdot r_1) \dot{r}_1 - \left(\frac{|\dot{r}_2|^2}{2} + \frac{|\dot{r}_1|^2}{2} - \frac{\mu}{|r_1|} \right) r_2 + (\dot{r}_2 \cdot r_2) \dot{r}_2 \right] \times r_1, \\ \frac{\partial \Phi_5}{\partial \dot{r}_2} &= -\dot{r}_2 \left[r_2 \cdot r_1 \times (r_2 - q_2) \right] + r_2 \left[\dot{r}_2 \cdot r_1 \times (r_2 - q_2) \right] + (\dot{r}_2 \cdot r_2) \dot{r}_2 \right] \times r_1, \end{aligned}$$

Equations (7) do not set any constraints on the mean anomalies ℓ_1 , ℓ_2 of the pairs of Keplerian orbits that we compute. To properly select a solution we integrate back each orbit $\mathbf{E}_{car}^{(2)}$ computed at epoch \tilde{t}_2 , to the epoch \tilde{t}_1 of the orbit $\mathbf{E}_{car}^{(1)}$, together with its covariance matrix $\Gamma^{(2)}$. Then, we compute a predicted position vector $\mathcal{P}_{1,p}$ and its 3×3 covariance matrix $\Gamma_{\mathcal{P}_{1,p}}$. To define a metric to choose/discard solutions we use a three-dimensional version of the attribution algorithm (Milani and Gronchi, 2010), and introduce the identification penalty

$$\chi_3^2 = \left(\mathcal{P}_1 - \mathcal{P}_{1,p}\right) \cdot \left[C_{\mathcal{P}_{1,p}} - C_{\mathcal{P}_{1,p}}\Gamma_0 C_{\mathcal{P}_{1,p}}\right] \left(\mathcal{P}_1 - \mathcal{P}_{1,p}\right),$$

where

$$C_{\mathcal{P}_{1,p}} = \Gamma_{\mathcal{P}_{1,p}}^{-1}, \qquad \Gamma_0 = C_0^{-1},$$

with

$$C_0 = C_{\mathcal{P}_{1,p}} + C_{\mathcal{P}_1}, \qquad C_{\mathcal{P}_1} = \Gamma_{\mathcal{P}_1}^{-1}.$$

We keep only the solutions with the lowest values of χ_3 or, if we wish to admit multiple solutions, we keep the ones whose value of χ_3 is below a certain threshold.

7 Numerical tests

In this section we show an application of our algorithm to asteroids undergoing a close approch with the Earth. We consider all the near-Earth asteroids present in the NEODyS database³ to the date of December 6, 2023, and select the 1305 asteroids with Earth

³http://newton.spacedys.com/neodys



Figure 1: Effects of the close approach on the semimajor axes (left) and on the whole trajectories (right).

MOID⁴ d_{\min} smaller than 10^{-3} au. For each selected asteroid we compute the points P_A and P_{\oplus} , lying on the osculating orbits of the asteroid and the Earth at epoch t_0 , where the minimum distance d_{\min} is attained. Then, we find the closest epoch $t_1 > t_0$ when the Earth arrives at P_{\oplus} and change the asteroid phase so that it would arrive at P_A at the same epoch t_1 in the framework of the Sun-asteroid two-body dynamics. In this way, we try to enhance the effect of the close approach with the Earth at t_1 . A full *n*-body propagation, from t_0 to t_1 , is used to compute the topocentric position vector \mathcal{P}_1 from the point of view of the Pan-STARRS 1 telescope (et al., 2019). Continuing the propagation from t_1 to a successive epoch t_2 , we derive an attributable \mathcal{A}_2 by coordinate change. As a result, the components α_2 , δ_2 , $\dot{\alpha}_2$, $\dot{\delta}_2$ of \mathcal{A}_2 are not affected by interpolation errors, which are usually introduced when an attributable is obtained from the astrometric observations.

Assume we can model a close approach at epoch t_1 by an instantaneous velocity change, like in Öpik's theory (Öpik, 1976). Then, our algorithm can be applied using $\mathcal{P}_1, \mathcal{A}_2$ as input data, because the values of the Keplerian integrals are conserved in $(t_1, t_2]$. At t_1 , when the velocity instantaneously changes, the position vector \mathcal{P}_1 remains unchanged.

In practice, the close encounter is not instantaneous, but takes a finite time: we select the epoch t_2 when the close approach is over, as described below. In a typical close encounter, the value of Tisserand's parameter T changes quickly, going back approximately to its pre-encounter value when the encounter is over. In Cavallari et al. (2023, Sec. 3), for a given value C of the Jacobi constant, the authors provide a value d_C of the geocentric distance attaining $\left|\frac{dT}{dt}\right| \leq \varepsilon$, for a fixed small parameter ε , so that we can think that the encounter is over when the asteroid reaches that distance from the Earth. Following this approach, we choose t_2 as the time when the value of the distance becomes d_C .

In Fig. 1 we show the effect of the close approach on these orbits. On the left, we plot the distribution of the relative differences between the semimajor axes a_1 , a_2 obtained by the known orbits propagated at epochs t_1 and t_2 using the software OrbFit⁵. On the right, we plot the distribution of the distance $\delta(\mathcal{T}_1, \mathcal{T}_2)$ between the propagated trajectories \mathcal{T}_1 ,

⁴Minimum Orbit Intersection Distance, see Bowell and Muinonen (1994), Gronchi et al. (2023)

⁵http::adams.dm.unipi.it/orbfit

 \mathcal{T}_2 , where

$$\delta(\mathcal{T}_1, \mathcal{T}_2) = \sqrt{\frac{(a_1 - a_2)^2}{a_1^2} + (e_1 - e_2)^2 + (i_1 - i_2)^2 + (\Omega_1 - \Omega_2)^2 + (\omega_1 - \omega_2)^2}.$$
 (19)

In (19) the subscripts 1, 2 of the Keplerian elements a, e, i, Ω, ω and the trajectories \mathcal{T} refer to the epochs t_1, t_2 . In most cases the relative change in semimajor axis is within 20%. However, there are a few cases where the change is larger, with one extreme case (Fig 1, left) where it passes from $a_1 \sim 2.84$ to $a_2 \sim 0.69$ au. The approximate values of the mean and the standard deviation of $(a_1 - a_2)/a_1$ are about 0.012 and 0.0478, respectively.

Next, we show the performance of our algorithm using \mathcal{P}_1 , \mathcal{A}_2 as input data, which have been computed for each of these orbits by a full *n*-body propagation with OrbFit. In case of multiple solutions, we select the best one according to the procedure explained in Section 6.1.

In 8 cases, out of 1305, we could not obtain an orbital solution. However, by increasing the time span $[t_1, t_2]$ by 10%, while keeping the same value of t_1 , thus changing only \mathcal{A}_2 , we recovered an orbital solution in all these cases. In one case the computation of the second epoch t_2 with Newton's method failed. However, employing the *starting guess* value for t_2 , computed with a geocentric two-body dynamics, we obtained an orbital solution also in this case.

In Fig. 2 we show the distribution of the differences between the Keplerian elements computed with our algorithm (with subscript c) and the same elements at t_2 obtained by propagation. In Fig. 2 (bottom right) we also show the distribution of the distances $\delta(\mathcal{T}_2, \mathcal{T}_c)$ between propagated (\mathcal{T}_2) and computed (\mathcal{T}_c) trajectories, where the function δ is the same as in (19).

The results of this preliminary statistical test are satisfactory: the mean and the standard deviation of the distributions displayed in Fig. 2 are shown in Table 1.

	mean	std
$(a_c - a_2)/a_2$	$-5.0803{ imes}10^{-4}$	0.0778
$e_c - e_2$	-0.003	0.0341
$i_c - i_2$	$-3.8397{\times}10^{-4}$	0.0045
$\Omega_c - \Omega_2$	0.0019	0.0232
$\omega_c - \omega_2$	-0.0021	0.0244
$\delta(\mathcal{T}_c,\mathcal{T}_2)$	0.0264	0.0877

Table 1: Mean and standard deviation of the distributions represented in Fig. 2.

Finally, we show the dependence of the results of our algorithm on the type of close encounter. Following Cavallari et al. (2023, Sect. 3), close encounters can be classified as *deep/shallow* and *fast/slow*, according to the values of the minimum distance q from the Earth center and the corresponding geocentric velocity v. Here, we compute an approximation of q and v, still denoted by these symbols, using the following procedure: qis the pericenter distance of the geocentric two-body orbit computed from the Keplerian elements at t_1 , and v is the corresponding two-body velocity. In Fig. 3 we plot the values of q and v using a color scale representing the values of the trajectory distance $\delta(\mathcal{T}_c, \mathcal{T}_2)$.

For values of q greater than 0.001 au, we obtain better results (e.g. lower values of the



Figure 2: Distribution of the differences in orbital elements a (top left), e (top right), i (center left), Ω (center right), ω (bottom left) between the propagated trajectory and the one computed with our algorithm at epoch t_2 . Distribution of the distances $\delta(\mathcal{T}_2, \mathcal{T}_c)$ between propagated and computed trajectories (bottom right).

trajectory distance) for higher values of the velocity v. For q smaller than 0.001 au, the results appear worse, and there is no apparent correlation with velocity in the figure.



Figure 3: Values of q and v in a log-log plot. The colors represent the values of the distance $\delta(\mathcal{T}_2, \mathcal{T}_c)$ between known and computed trajectories.

8 Conclusions

We have introduced a method to compute preliminary orbits with one topocentric position vector \mathcal{P}_1 and a very short arc of optical observations, from which we can derive an attributable \mathcal{A}_2 . This method is based on polynomial equations coming from the first integrals of Kepler's problem, i.e. angular momentum, energy, and Laplace-Lenz vector. Using the conservation laws of these integrals, after introducing the auxiliary variable z_2 , we obtain a polynomial system that always has solutions, at least in the complex field, even if \mathcal{P}_1 , \mathcal{A}_2 do not correspond to the same celestial object. There are some checks that can be performed to accept or reject solutions.

We applied this algorithm to the computation of 1305 NEA orbits, whose phase was changed in order to enhance the close encounter effect with the Earth: these preliminary results are satisfactory, see Section 7. The ideal situation for the application of the proposed algorithm in this context would be given by an instantaneous effect of the close encounter, which is not the case. The selected epoch t_1 does not exactly correspond to the time of passage at the MOID because we use a full *n*-body propagation starting from t_0 , so that at t_1 both the osculating trajectory and the time law along it may have changed. Moreover, from this preliminary test, we saw that also varying t_2 may affect the results. Therefore, a more detailed study is necessary to better understand the applicability in case of close encounters and the reliability of the computed orbits. On top of that, the sensitivity of the algorithm to astrometric errors has still to be investigated.

Applications of the method introduced in this work to the orbit computation of Earth satellites undergoing a maneuvre are also possible, and are worth to be investigated in a future work.

Acknowledgments The authors acknowledge the project MIUR-PRIN 20178CJA2AB "New Frontiers of Celestial Mechanics: theory and applications". G.F. Gronchi and G. Baù acknowledge the project MSCA-ITN Stardust-R, Grant Agreement n. 813644 under the H2020 research and innovation program. E. Scantamburlo acknowledges the project "Advanced Space System Engineering to Address Broad Societal Benefits – Starting Grant" funded by a contract between Politecnico di Torino and Compagnia di San Paolo (CSP) 2019/2021 within the call "Attrazione e retention di docenti di qualità".

A Appendix

Lemma 2. For $c_1 = c_2$, the generators q_4 , q_5 , q_6 defined in (8) assume the following form:

$$\begin{split} \tilde{\mathfrak{q}}_4 &= (\boldsymbol{c}_1 \cdot \hat{\boldsymbol{e}}_1^{\rho}) \left[\boldsymbol{r}_1 \cdot (\dot{\boldsymbol{r}}_1 - \dot{\boldsymbol{r}}_2) \right] + (\boldsymbol{r}_2 \cdot \boldsymbol{D}_1) z_2, \\ \tilde{\mathfrak{q}}_5 &= (\boldsymbol{c}_2 \cdot \hat{\boldsymbol{e}}_2^{\rho}) \left[\boldsymbol{r}_2 \cdot (\dot{\boldsymbol{r}}_1 - \dot{\boldsymbol{r}}_2) \right] - (\boldsymbol{r}_1 \cdot \boldsymbol{D}_2) \frac{\mu}{|\boldsymbol{r}_1|}, \\ \tilde{\mathfrak{q}}_6 &= (\boldsymbol{c}_2 \cdot \hat{\boldsymbol{e}}_2^{\rho}) \left[\boldsymbol{r}_1 \cdot (\dot{\boldsymbol{r}}_1 - \dot{\boldsymbol{r}}_2) \right] - (\boldsymbol{r}_1 \cdot \boldsymbol{D}_2) z_2. \end{split}$$

Proof. The result is given by a direct computation:

$$\begin{split} \tilde{\mathfrak{q}}_{4} &= -(\dot{r}_{1} \cdot r_{1}) \left[\dot{r}_{1} \cdot (r_{1} \times \hat{e}_{1}^{\rho}) \right] - \mu \tilde{L}_{2} \cdot D_{1} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) (\dot{r}_{1} \cdot r_{1}) - (\|r_{2}\|^{2} - z_{2}) \left[r_{2} \cdot (r_{1} \times \hat{e}_{1}^{\rho}) \right] - (\dot{r}_{2} \cdot r_{2}) \left[\dot{r}_{2} \cdot (r_{1} \times \hat{e}_{1}^{\rho}) \right] \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) (\dot{r}_{1} \cdot r_{1}) - (\dot{r}_{2} \times c_{2}) \cdot (r_{1} \times \hat{e}_{1}^{\rho}) + (r_{2} \cdot D_{1}) z_{2} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) (\dot{r}_{1} \cdot r_{1}) - [\dot{r}_{2} \times (r_{1} \times \dot{r}_{1})] \cdot (r_{1} \times \hat{e}_{1}^{\rho}) + (r_{2} \cdot D_{1}) z_{2} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) (\dot{r}_{1} \cdot r_{1}) - [r_{1} (\dot{r}_{1} \cdot \dot{r}_{2}) - \dot{r}_{1} (\dot{r}_{2} \cdot r_{1})] \cdot (r_{1} \times \hat{e}_{1}^{\rho}) + (r_{2} \cdot D_{1}) z_{2} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) (\dot{r}_{1} \cdot r_{1}) - [r_{1} (\dot{r}_{1} \cdot \dot{r}_{2}) - \dot{r}_{1} (\dot{r}_{2} \cdot r_{1})] \cdot (r_{1} \times \hat{e}_{1}^{\rho}) + (r_{2} \cdot D_{1}) z_{2} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) [\dot{r}_{1} \cdot (\dot{r}_{1} - \dot{r}_{2})] + (r_{2} \cdot D_{1}) z_{2} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) [r_{1} \cdot (\dot{r}_{1} - \dot{r}_{2})] + (r_{2} \cdot D_{1}) z_{2} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) [r_{1} \cdot (\dot{r}_{1} - \dot{r}_{2})] + (r_{2} \cdot D_{1}) z_{2} \\ &= (c_{1} \cdot \hat{e}_{1}^{\rho}) [r_{1} \cdot (\dot{r}_{1} - \dot{r}_{2})] + (r_{2} \cdot d_{2}^{\rho}) - (c_{2} \cdot \hat{e}_{2}^{\rho}) (\dot{r}_{2} \cdot r_{2}) \\ &= \left(\dot{r}_{1} \times c_{2} - \frac{\mu}{|r_{1}|} \right) \cdot (r_{2} \times \hat{e}_{2}^{\rho}) - (c_{2} \cdot \hat{e}_{2}^{\rho}) (\dot{r}_{2} \cdot r_{2}) \\ &= \left((\dot{r}_{1} \cdot \dot{r}_{2}) r_{2} - (r_{2} \cdot \dot{r}_{1}) \dot{r}_{2} - \frac{\mu}{|r_{1}|} \right) \cdot (r_{2} \times \hat{e}_{2}^{\rho}) - (c_{2} \cdot \hat{e}_{2}^{\rho}) (\dot{r}_{2} \cdot r_{2}) \\ &= -(r_{2} \cdot \dot{r}_{1}) \left[\dot{r}_{2} \cdot (r_{2} \times \hat{e}_{2}^{\rho}) \right] - \frac{\mu}{|r_{1}|} r_{1} \cdot (r_{2} \times \hat{e}_{2}^{\rho}) - (c_{2} \cdot \hat{e}_{2}^{\rho}) (\dot{r}_{2} \cdot r_{2}) \\ &= (c_{2} \cdot \hat{e}_{2}^{\rho}) \left[r_{2} \cdot (\dot{r}_{1} - \dot{r}_{2}) \right] - (r_{1} \cdot D_{2}) \frac{\mu}{|r_{1}|}, \\ \tilde{\mathfrak{q}}_{6} = - \left[(\dot{r}_{2} \times c_{2}) - z_{2} r_{2} \right] \cdot (r_{1} \times \hat{e}_{2}^{\rho}) + (c_{2} \cdot \hat{e}_{2}^{\rho}) (\dot{r}_{1} \cdot r_{1}) \\ &= - \left[(\dot{r}_{1} \cdot \dot{r}_{2}) r_{1} - (\dot{r}_{2} \cdot r_{1}) \dot{r}_{1} - z_{2} r_{2} \right] \cdot (r_{1} \times \hat{e}_{2}^{\rho}) + (c_{2} \cdot \hat{e}_{2}^{\rho}) (\dot{r}_{1} \cdot r_{1}) \\ &= (c_{1} \cdot \dot{r}_{2}) \left[r_{1}$$

References

E. Bowell and K. Muinonen. *Hazards due to comets and asteroids*. Tucson: The University of Arizona Press, 1994.

- I. Cavallari, C. Grassi, G. F. Gronchi, G. Baù, and G. B. Valsecchi. A dynamical definition of the sphere of influence of the Earth. *Communications in Nonlinear Science and Numerical Simulation*, 119:107091, 2023.
- K. C. Chambers et al. The Pan-STARRS1 surveys, 2019.
- C. F. Gauss. *Theoria motus corporum in sectionibus conicis solem ambientium*. reprinted by Dover publications in 1963, 1809.
- G. F. Gronchi, L. Dimare, and A. Milani. Orbit determination with the two-body integrals. Celestial Mechanics and Dynamical Astronomy, 107:299–318, 2010.
- G. F. Gronchi, D. Farnocchia, and L. Dimare. Orbit determination with the two-body integrals. II. *Celestial Mechanics and Dynamical Astronomy*, 110:257–270, 2011.
- G. F. Gronchi, G. Baù, and S. Marò. Orbit determination with the two-body integrals. III. Celestial Mechanics and Dynamical Astronomy, 123:105–122, 2015.
- G. F. Gronchi, G. Baù, and A. Milani. Keplerian integrals, elimination theory and identification of very short arcs in a large database of optical observations. *Celestial Mechanics* and Dynamical Astronomy, 127:211–232, 2017.
- G. F. Gronchi, G. Baù, and C. Grassi. Revisiting the computation of the critical points of the Keplerian distance. *Celestial Mechanics and Dynamical Astronomy*, 135, 2023.
- J. L. Lagrange. Sur le problème de la détermination des orbites des cométes d'aprés trois observations. Troisiè memémoire. Nouveaux mémoires de l'Académie royale des sciences et belles-lettres de Berlin, 1783. Reprinted in Œuvres de Lagrange, Gauthier-Villars et fils, Paris (1869), volume 4, pp. 496–532.
- P. S. Laplace. Mémoire sur la détermination des orbites des comètes. Mémoires de l'Académie royale des sciences de Paris, 1780. Reprinted in Œuvres complètes de Laplace, Gauthier-Villars et fils, Paris (1894), volume 10, pp. 93–146.
- A. Milani and G. F. Gronchi. Theory of Orbit Determination. Cambridge Univ. Press, 2010.
- A. Milani, G. F. Gronchi, Z. Knežević, M. E. Sansaturio, and O. Arratia. Orbit Determination with Very Short Arcs. II Identifications. *Icarus*, 179:350–364, 2005.
- E. J. Opik. Interplanetary encounters: close-range gravitational interactions. Elsevier, Amsterdam, 1976.
- L. G. Taff and D. L. Hall. The use of angles and angular rates. I Initial Orbit Determination. *Celestial Mechanics*, 16:481–488, 1977.