Family Puzzle, Framing Topology, $c_{-} = 24$ and $3(E_8)_1$ Conformal Field Theories: $\frac{48}{16} = \frac{45}{15} = \frac{24}{8} = 3$

Juven Wang^{1, *}

¹Center of Mathematical Sciences and Applications, Harvard University, MA 02138, USA

Family Puzzle or Generation Problem demands an explanation of why there are 3 families or generations of quarks and leptons in the Standard Model of particle physics. Here we propose a novel solution — the multiple of 3 families of 16 Weyl fermions (namely $(N_f = 3) \times 16$) in the 3+1d spacetime dimensions are topologically robust due to constraints rooted in profound mathematics (such as Hirzebruch signature and Rokhlin's theorems, and cobordism) and derivable in physics (such as chiral edge states, quantized thermal Hall conductance, and gravitational Chern-Simons theory), which holds true even forgetting or getting rid of any global symmetry or gauge structure of the Standard Model. By the dimensional reduction through a sequence of sign-reversing mass domain wall of domain wall and so on, we reduce the Standard Model fermions to obtain the $(N_f = 3) \times 16$ multiple of 1+1d Majorana-Weyl fermion with a total chiral central charge $c_{-} = 24$. Effectively via the fermionization-bosonization, the 1+1d theory becomes 3 copies of $c_{-} = 8$ of $(E_8)_1$ conformal field theory, living on the boundary of 3 copies of $2+1d E_8$ quantum Hall states. Based on the framing anomaly-free $c_{-} = 0 \mod 24$ modular invariance, the framed bordism and string bordism \mathbb{Z}_{24} class, the 2-framing and p_1 -structure, the w_1 - p_1 bordism \mathbb{Z}_3 class constraints, we derive the family number constraint $N_f \in (\frac{48}{16} = \frac{24}{8} = 3)\mathbb{Z}$. The dimensional reduction process, although not necessary, is sufficiently supported by the \mathbb{Z}_{16} class Smith homomorphism. We also comment on the $\frac{45}{15} = 3$ relation: the 3 families of 15 Weyl-fermion Standard Model vacuum where the absence of some starily relations is followed by the \mathbb{Z}_{16} class some starily relations in the source of start of the start of some sterile right-handed neutrinos is fulfilled by additional topological field theories or conformal field theories in Ultra Unification.

CONTENTS

I.	Introduction and Summary	2
	A. Family Puzzle	2
	B. Background Information	2
II.	Physics Model Setup	3
	A. General Arguments: $c_{-} = 24$ and $3(E_8)_1$ Conformal Field Theories	4
	B. Mass Domain Wall of Domain Wall Reduction Toy Model	6
III.	From 6d Anomaly Polynomial, 5d Convertible Field Theory, 4d Anomaly of the SM to 3d Gravitational	C
	Chern-Simons Term	8
	A. Spin $\times_{\mathbb{Z}_2^F} \mathbb{U}(1)_{\mathbf{Q}-N_c \mathbf{L}} \equiv \text{Spin}^\circ$ and $\text{Spin} \times_{\mathbb{Z}_2^F} \mathbb{Z}_{4,X}$ with a vector $\mathbb{U}(1)_{\mathbf{Q}-N_c \mathbf{L}}$ and a chiral X	g
	B. Another Spin $\times_{\mathbb{Z}_2^F} U(1) \equiv \text{Spin}^c$ with a chiral U(1): $3(E_8)_1$ quantum Hall states	10
	C. Index Theorem and (Gravitational) Chern-Simons: Framing, String, 2-Framing, and w_1 - p_1 structures	10
	D. Cobordism constraints of Framing, String, 2-Framing, and w_1 - p_1 structures	12
IV.	Conclusion and Comparison	12
	Acknowledgments	13
	References	14

^{*} jw@cmsa.fas.harvard.edu; http://sns.ias.edu/~juven/

I. INTRODUCTION AND SUMMARY

A. Family Puzzle

The 3 families or 3 generations of quarks and leptons of the Standard Model (SM) of particle physics had been advocated since the 1970s [1]. The Family Puzzle or Generation Problem demands an explanation of what dictates the SM family number $N_f = 3$. Any valid theoretical solution to this puzzle, unraveling the SM's underlying mysterious structure, can guide us further to explore Beyond the Standard Model physics (BSM) with elevated confidence, making new predictions. What is the evidence of $N_f = 3$?

On one hand, in the quark sector, the (C)KM matrix indicates that the charge-conjugation-parity CP symmetry violation of quarks via the weak interaction predicts the existence of at least 3 families of quarks in nature theoretically [2]. The discovery of the most massive known elementary particle, top quark t, in 1995 by the CDF and Dø experiments at the Fermilab [3, 4] confirmed the completion of 3 families of quarks experimentally. Moreover, if there is any gapped hypothetical 4th generation of quarks massive than t quark, say t' and b' quarks, the Higgs decay rate to two gluons $\phi_H \rightarrow gg$ through loop triangle Feynman diagrams of 3 types of virtual massive quarks (t, t', b') will be enhanced by a factor of $3^2 = 9$ times larger than the SM prediction, but which possibility has been excluded [5].

On the other hand, in the lepton sector, the Z boson (the third most massive known particle of about 91 GeV) decays into at most 3 families of light neutrinos (assuming light neutrinos are lighter than about 45 GeV, namely mass $m_{\nu} < m_Z/2$), verified at CERN's Large Electron Positron Collider (LEP) [6, 7]. In addition, the astrophysical data from the cosmic microwave background (CMB) or big bang nucleosythesis (BBN) also suggest 3 families of light neutrinos [8].

Here we show a novel theoretical solution to the Family Puzzle — the multiple of $N_f = 3$ of 16 Weyl fermions in the 3+1d spacetime dimensions are robust due to the almost purely topological constraints (without requiring internal symmetry, neither global symmetry nor gauge structure) rooted in the profound mathematics and derivable in physics.¹

B. Background Information

To understand and appreciate our solution better, some prior knowledge and familiarity with literature can help: • (1) Modular invariance of 2d conformal field theory (CFT) [12] and 3d Chern-Simons-Witten and gravitational Chern-Simons theories [13]: framing anomaly-free with chiral central charge $c_{-} = 0 \mod 24$.²

• (2) Mapping between fermions by domain-wall reduction (see Fig. 1 and Fig. 2): crossing different dimensions by the sign-reversing mass domain wall of domain wall and so on. We will introduce a toy model (following [14]) as the explicit mass domain wall m(x) reincarnation version of Jackiw-Rebbi [15] or Callan-Harvey [16] where they introduce instead some additional scalar field $\Phi(x)$. Our mass domain wall does not require extra scalar field and does not necessarily implement symmetry-breaking mass; in contrast, other related (similar but not exactly the same) recent models that explore the symmetry-breaking domain wall and anomaly inflow include [17–20]. A lattice version of our model can be implemented as the domain wall of the domain wall fermion of Kaplan's [21] and so on.

• (3) Cobordism theory [22, 23] and Atiyah-Patodi-Singer (APS) et a invariant η [24]. Given the spacetime and internal symmetry G structure of manifold³

$$G \equiv \left(\frac{G_{\text{spacetime}} \ltimes G_{\text{internal}}}{N_{\text{shared}}}\right) \equiv G_{\text{spacetime}} \ltimes_{N_{\text{shared}}} G_{\text{internal}},\tag{1}$$

we will apply the data of bordism group Ω_D^G that classifies the *G*-structure *D*-dimensional (*D*d) manifold M^D up to the cobordant relation (identified by bounding the (D+1)d manifold), and the Anderson dual version of the cobordism group TP_D^G that classifies the *D*d invertible field theories or *D*d Symmetry-Protected Topological states (SPTs)

¹ One inspiration of this work is noticing that the deep UV regularization of 2+1d E_8 quantum Hall state required more strict branchdependent structure on the triangulation of the manifold, while the 2+1d $3E_8$ quantum Hall state required less strict branch-independent structure on the triangulation of the manifold [9]. The branch structure seems to be related to the framing structure of the manifold in some way. The author realizes that the Hirzebruch signature [10], framing-anomaly, and modular invariance may give rise to the purely topological constraint on the Family Puzzle $N_f \in 3\mathbb{Z}$. Another inspiration comes from studying the anomalies of SM by the index theorem in [11].

 $^{^{2}}$ For spacetime dimensionality, we either denote the space + time dimension (e.g. 1+1d CFT) or the total dimension (e.g. 2d CFT).

³ The semi-direct product \ltimes specifies a group extension. The N_{shared} is the shared common normal subgroup symmetry between $G_{\text{spacetime}}$ and G_{internal} , e.g. N_{shared} can be the fermion parity symmetry $\mathbb{Z}_{2}^{\text{F}}$.

(2)

realized in quantum condensed matter, and their boundary's (D-1)d quantum anomalies (e.g., [25–29]).⁴ Later we will use freely these relation between the co/bordism and the classification of manifolds vs invertible field theories. The G_{internal} implies the structure group of the principal bundle and gauge connection. The $G_{\text{spacetime}} = \text{Spin}, \text{Pin}^{\pm}, \dots$ can contain the fermionic Lorentz group structure (rotation + boost) such as Spin group or the time-reversal or reflection-symmetry enhanced Pin^{\pm} group [22]. Moreover, we emphasize that our Family Puzzle solution require nothing of any specific G_{internal} of the SM nor any specific $G_{\text{spacetime}}$, except only the tangential structure of manifold (denoted Struct). We will apply the data of bordism group Ω_d^{Struct} and cobordism group $\mathrm{TP}_d^{\text{Struct}}$ where the tangential structure can include the characteristic class [34] defined by tangent bundles: the Steifel-Whitney class (e.g. w_1, w_2) or String structure, and the Pontryagin class (e.g. p_1), or Witten's framing structure [13], or Atiyah's 2-framing structure [35]. In particular, we will study the 3-manifold bounding the boundary of a 4-manifold, and we will use the relation between (a) framing provides the String structure, and (2) 2-framing provides the w_1 - p_1 -structure. For Family Puzzle, we will need the following structures:

Structure := $\begin{cases} \text{(a) framing (fr): trivialization of tangent bundle } TM \\ \simeq \text{String structure: trivialization of } w_1(TM), w_2(TM), \text{ and } \frac{1}{2}p_1(TM). \\ \text{(b) 2-framing (2-fr): trivialization of the spin bundle of 2 copies of the tangent bundle } TM \oplus TM \end{cases}$ (3) $\simeq w_1$ - p_1 -structure: trivialization of $w_1(TM)$ and $p_1(TM)$.

• (4) Atiyah-Singer index theorem (e.g.[36]), specifically Hirzebruch signature [10] and Rokhlin's [37] theorems.

PHYSICS MODEL SETUP II.

Standard Model (SM) [38] is a 4d chiral gauge theory with Yang-Mills spin-1 gauge fields of the Lie algebra $\mathcal{G}_{\mathrm{SM}} \equiv su(3) \times su(2) \times u(1)_{\tilde{Y}}$ (with four compatible Lie group $G_{\mathrm{SM}_{q}} \equiv \frac{\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)_{\tilde{Y}}}{\mathbb{Z}_{q}}$, with q = 1, 2, 3, 6) coupling to $N_f = 3$ families of 15 or 16 Weyl fermions (spin- $\frac{1}{2}$ Weyl spinor in the $\mathbf{2}_L^{\mathbb{C}}$ representation of the spacetime symmetry Spin(1,3), written as a left-handed 15- or 16-plet ψ_L) in the following $\mathcal{G}_{\mathrm{SM}}$ representation

$$(\psi_L)_{\rm I} = (\bar{d}_R \oplus l_L \oplus q_L \oplus \bar{u}_R \oplus \bar{e}_R)_{\rm I} \oplus n_{\nu_{{\rm I},R}} \bar{\nu}_{{\rm I},R} \sim ((\bar{\mathbf{3}}, \mathbf{1})_2 \oplus (\mathbf{1}, \mathbf{2})_{-3} \oplus (\mathbf{3}, \mathbf{2})_1 \oplus (\bar{\mathbf{3}}, \mathbf{1})_{-4} \oplus (\mathbf{1}, \mathbf{1})_6)_{\rm I} \oplus n_{\nu_{{\rm I},R}} (\mathbf{1}, \mathbf{1})_0$$

$$(4)$$

for each family label I = 1,2,3. The right-handed neutrino $\bar{\nu}_R$ is written as right-handed anti-particle so to be regarded as a left-handed particle. The $n_{\nu_{e,R}}, n_{\nu_{\pi,R}} \in \{0,1\}$ for I = 1,2,3 labels either the absence or presence of electron e, muon μ , or tauon τ types of sterile neutrinos (i.e., "right-handed" neutrinos sterile to \mathcal{G}_{SM} gauge forces). There are also additional Yukawa-Higgs terms. But none of these are crucial to our solution except the total number of Weyl fermions —

$$N_f = 3) \times 15 + n_{\nu_{e,R}} + n_{\nu_{\mu,R}} + n_{\nu_{\tau,R}} = 45, 46, 47, 48, \dots$$
(5)

For simplicity, we focus on the 48-Weyl-fermion SM first in the limit of massless fermions, turning off the SM gauge structure and SM Higgs mechanism while relegating those extra refined structures to later discussions.⁵ We will comment on other fermion numbers later.

⁴ Freed-Hopkins version of cobordism group $\operatorname{TP}_D^G = \Omega_{D+1}^{G; \text{free}} \oplus \Omega_D^{G; \text{torsion}}$ [23] consists of the integer free \mathbb{Z} class of $\Omega_{D+1}^{G; \text{torsion}}$ and the finite group torsion \mathbb{Z}_n class of $\Omega_D^{G; \text{torsion}}$. In summary,

	$\Omega_{D+1}^{G;\text{free}}$ classifies $(D+1)$ d anomaly polynomials, Dd Chern-Simons like invertible field
	theories, and $(D-1)d$ perturbative local anomalies [30] of G; those anomalies can be
	detected by infinitesimal or small gauge/diffeomorphism transformations and
$TPG = \Omega^{G; \text{free}} \oplus \Omega^{G; \text{torsion}} =$	perturbative Feynman diagrams.
$\prod_{D=M} D = M_{D+1} \oplus M_D = -$	$\Omega_D^{G;\text{torsion}}$ classifies Dd APS η invariants as invertible topological field theories (TFTs) as well as
	cobordism invariants, and $(D-1)d$ nonperturbative global anomalies [31–33] of G; those anomalies
	can only be detected by large gauge/diffeomorphism transformations
	(that are undeformable continuously from the identity [null transformation]).

 5 It may still be possible to have 0, 1, 2, 3, or more than 3 sterile neutrinos. Also, turning off gauge and Higgs fields is highly-motivated for the high-energy early universe scenario — the fermions become massless. In addition, any Grand Unification with an appropriate semi-simple compact Lie group and with these amounts of fermions exhibits asymptotic freedom like free quarks at higher energy. Here we also clarify the physical meanings of energy spectrum. Typically for particle physics, whenever the particle field description is

available, we have massless or massive spectrum. On the other hand, for interacting CFT, it is better to refer its spectrum as gapless; for interacting topological quantum field theory (TQFT), it is better to refer its spectrum as gapped above the (possibly degenerate) ground state(s). Thus, gapless vs gapped are more general than massless vs massive.

Moreover, the "heavy mass" concept in quantum matter context may mean something different: the inverse of the curvature of the energy dispersion E(p) of momentum p, namely $(\nabla_p^2 E(p))^{-1}$. This "heavy mass" can coincidentally coincide with the mass gap for Lorentz invariant particle theory, but the Lorentzian coincidence between these masses in general fail either in non-Lorentz invariant or in interacting many-body quantum systems. Regardless of these subtleties of massive or gapped systems, we only focus on Lorentz invariant theories in this work.

A. General Arguments: $c_{-} = 24$ and $3(E_8)_1$ Conformal Field Theories

What absolutely crucial is the dimension-reduction relationship between 3+1d's 48 Weyl fermions in (5) and the 1+1d 48 Majorana-Weyl fermion (each with chiral central charge $c_{-} = \frac{1}{2}$) with a combined chiral central charge

$$c_{-} \equiv c_{L} - c_{R} = \frac{1}{2}48 = 24. \tag{6}$$

Here are the steps of our arguments (here succinctly summarizing the main logic, later we will fill in more details):

- (1): Mass domain wall reduction maps between fermions as Lorentz Spin(d, 1) spinors crossing different di**mensions** (see Fig. 1 (a)):
 - 4+1d Dirac $\psi_{\rm D}^{5d} \rightarrow 3+1$ d Weyl $\psi_{\rm W}^{4d} \sim 3+1$ d Majorana $\psi_{\rm M}^{4d} \rightarrow 2+1$ d Majorana $\psi_{\rm M}^{3d} \rightarrow 1+1$ d Majorana-Weyl $\psi_{\rm MW}^{2d}$. (7)
 - (i): A single (d+1)d Dirac or Majorana fermion allows to pair themselves with Dirac or Majorana mass term respectively by the fermion bilinear $m\bar{\psi}\psi \equiv m\psi^{\dagger}\Gamma^{0}\psi$.
 - (ii): Whenever the massive fermion ψ^{d+1} is allowed in (d+1)d, with time and space coordinates $(t, x_1, x_2, \ldots, x_d)$, we may consider the spatial-dependent mass profile $m(x_d)\overline{\psi}\psi$ such that

$$m(x_d) = \begin{cases} |m|, \ x_d \gg 1. \\ 0, \ x_d = 0. \\ -|m|, \ x_d \ll 1. \end{cases}$$
(8)

The (d+1)d fermion obeys the Lagrangian $\mathcal{L}_{(d+1)d} = \bar{\psi}^{d+1}(i\Gamma^{\mu}\partial_{\mu} - m(x))\psi^{d+1}$ with the spacetime indices $\mu = 0, 1, \dots, d$, then there is an effective massless dd domain wall fermion theory ψ^d at $x_d = 0$, with its time and space coordinates $(t, x_1, x_2, \ldots, x_{d-1})$, which is obtained by the projection $P_{\pm} = \frac{1 \pm i \Gamma^d}{2}$, so⁶

$$\psi_{\pm}^{d} = \mathbf{P}_{\pm}\psi^{d+1} = \frac{1 \pm i\Gamma^{d}}{2}\psi^{d+1}.$$
(9)

We get a *d*d domain wall fermion Lagrangian, either + or - version of $\mathcal{L}_{dd,\pm} = \psi_{\pm}^{d\dagger} i(\partial_t - \Gamma^0 \Gamma^j \partial_j) \psi_{\pm}^d$ depending on the orientation of the kink of the domain wall, with the spatial indices $j = 1, \ldots, d-1$. Importantly, either of a single kink domain wall fermion ψ_{\pm}^d carries 1/2 of degrees of freedom of the bulk fermion ψ_{\pm}^{d+1} , while the combined $\psi_{\pm}^d + \psi_{\pm}^d$ reproduces the full bulk ψ^{d+1} degrees of freedom (DOF). This DOF counting combined with the spinor representation theory unambiguously suggests a unique domain wall reduction path in Fig. 1 (a) from the 4d Weyl fermion ψ_W^{4d} of the SM. This approach works in any dimension, but we focus on the reduction from 5d to 2d in Fig. 1 (a):

Dirac ψ_{D}^{5d} in $4^{\mathbb{C}}$, Weyl ψ_{W}^{4d} in $2_{L}^{\mathbb{C}}$ equivalently as Majorana ψ_{M}^{4d} in $4^{\mathbb{R}}$, Majorana ψ_{M}^{3d} in $2^{\mathbb{R}}$, Majorana-Weyl ψ_{MW}^{2d} in $1_{L}^{\mathbb{R}}$.

For N_f -family n-Weyl fermion SM (e.g. $N_f = 3\mathbb{Z}$ and n = 16), we obtain a reduction map following:

$$N_f n \psi_{\rm D}^{5d} \to N_f n \psi_{\rm M}^{4d} \sim N_f n \psi_{\rm M}^{4d} \to N_f n \psi_{\rm M}^{3d} \to N_f n \psi_{\rm MW}^{2d} .$$

$$\Rightarrow 48\mathbb{Z} \psi_{\rm D}^{5d} \to 48\mathbb{Z} \psi_{\rm W}^{4d} \sim 48\mathbb{Z} \psi_{\rm M}^{4d} \to 48\mathbb{Z} \psi_{\rm M}^{3d} \to 48\mathbb{Z} \psi_{\rm MW}^{2d} .$$
(11)

The reduced 1+1d theory has a total chiral central charge $c_{-} \in \frac{1}{2}48\mathbb{Z} = 24\mathbb{Z}$. We thus reduced the standard 3+1d left-handed Weyl fermion theory with Pauli's gamma matrices to 1+1d chiral Majoarna-Weyl theory:

$$\mathcal{L}_{4d} = \sum_{I=1}^{N_f n} \psi_{W,I}^{4d\dagger} i \bar{\sigma}^{\mu} \partial_{\mu} \psi_{W,I} \longmapsto \mathcal{L}_{2d} = \sum_{I=1}^{N_f n} \psi_{MW,I}^{2d} i (\partial_t - \partial_x) \psi_{MW,I}^{2d}.$$
(12)

⁶ Follow [14], for any dimension d + 1, we write down a set of Gamma matrices satisfy $\{\Gamma^{\mu}, \Gamma^{\nu}\} = 2\eta^{\mu\nu}$ for a proper Lorentzian Minkowski metric $\eta^{\mu\nu} = \text{diag}(+, -, \dots, -)$ in any dimension with spacetime indices $\mu, \nu = 0, 1, \dots, d-1, d$. Note that $P_{\pm}^2 = P_{\pm}$ and $\mathbf{P}_+\mathbf{P}_-=\mathbf{P}_-\mathbf{P}_+=0 \mbox{ and } \mathbf{i}\Gamma^d\mathbf{P}_\pm=\pm\mathbf{P}_\pm.$

[•] For even d, we can choose the chiral or Weyl representation, $\Gamma^0 = \begin{pmatrix} 0 & \mathbb{I} \\ \mathbb{I} & 0 \end{pmatrix}$, $\Gamma^j = \begin{pmatrix} 0 & \gamma^j \\ -\gamma^j & 0 \end{pmatrix}$ for $j = 1, \dots, d-1$ with $\{\gamma^i, \gamma^j\} = 2\delta^{ij}$,

 $[\]Gamma^d = \text{idiag}(-\mathbb{I},\mathbb{I}), \text{ so } P_{\pm} = \text{diag}(\mathbb{I},0) \text{ or } \text{diag}(0,\mathbb{I}).$ The Gamma matrices Γ^{μ} in dd are the same as those in (d+1)d for even d. The P_{\pm} projection maps a (d+1)d Dirac (or Majorana) fermion to a dd chiral Weyl (or Majorana-Weyl) fermion. The P_{\pm} decouples the left-handed and right-handed Weyl (Majorana-Weyl) fermion in dd at $x_d = 0$ with $m(x_d) = 0$. We get a dd domain wall fermion Lagrangian $\mathcal{L}_{d\,d} = \psi_{\pm}^{d\dagger} i(\partial_t \mp \gamma^j \partial_j) \psi_{\pm}^d$. • For odd *d*, the projection maps a (d+1)d Dirac (Majorana) fermion to a *dd* Dirac (Majorana) fermion. See Fig. 1 (a).



FIG. 1. (a) This subfigure is a key step in our argument. Here the left column d labels the total spacetime dimension of fermionic theories in the same row. For example, in d = 2, we have Majorana-Weyl (MW) ψ_{MW}^{2d} in $\mathbb{1}_{L}^{\mathbb{R}}$, Majorana (M) ψ_{M}^{2d} in $\mathbb{1}_{L}^{\mathbb{R}} \oplus \mathbb{1}_{R}^{\mathbb{R}}$, Weyl (W) in $\mathbb{1}_{L}^{\mathbb{C}}$, and Dirac (D) in $\mathbb{1}_{L}^{\mathbb{C}} \oplus \mathbb{1}_{R}^{\mathbb{C}}$. Here \mathbb{R} and \mathbb{C} for real and complex values, L and R for the left and right chirality. The key dimensional reduction path along the color path with arrows from 5d to 2d $(4^{\mathbb{C}} \to 2_{L}^{\mathbb{C}} \to 4^{\mathbb{R}} \to 2^{\mathbb{R}} \to \mathbb{1}_{L}^{\mathbb{R}})$ follows (7) and (10), its intentional colors match with the colors of bulks or domain walls in Fig. 2. (b) This subfigure is *not* a key step in our argument. This shows the cobordism (TP_d) and bordism (Ω_d) version of the Smith homomorphism [39]. This Smith homomorphism helps to group the minimal representation of fermions (as the (d-1)d free fermion boundary of the cobordism invariant of the TP_d for anomaly matching) in 0 mod 16 or 0 mod 8 manner, which is supportive and sufficient, but not necessarily required for solving the Family Puzzle.

(2): The 1+1d reduced theory has chiral central charge $c_{-} = 0 \mod 24$ thus framing anomaly free [12, 13]. The 3+1d SM and the 1+1d reduced theories all have the fermion parity symmetry $\mathbb{Z}_2^{\mathrm{F}}$, which acts on fermions by

 $\psi \mapsto -\psi$. Moreover, we can do fermionization-bosonization $(\psi_I \sim \exp(i\phi_I) : of \text{ compact boson } \phi_I \text{ with proper normal ordering) to map the (12)'s 2d 16N_f Majorana-Weyl fermions to 2d 8N_f Weyl fermions, to the bosonized version, and then by summing over all Spin structures of the 2-manifold of the 2d theory,⁷ which becomes N_f (e.g., 3Z) copies of 1+1d (E₈)₁ bosonic CFT or Wess-Zumino-Witten (WZW) model:⁸$

$$\mathcal{L}_{2d}^{B} = \sum_{I=1}^{N_{f}} \sum_{I,J}^{8} \frac{1}{4\pi} \left((K_{E_{8}})_{IJ} \partial_{t} \phi_{I} - V_{IJ} \partial_{x} \phi_{I} \right) \partial_{x} \phi_{J}. K_{E_{8}} \equiv \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 2 \end{pmatrix}, V = \mathbb{I}_{8}, \quad I, J \in \{1, \dots, 8\}$$

$$(13)$$

where K_{E_8} is the unimodular symmetric bilinear form's K matrix (corresponding to the gapless boundary of 2+1d abelian Chern-Simons theory as 2+1d invertible TFT that corresponds to the unimodular $|\det(K)| = 1$), here the rank-8 Cartan matrix of E_8 . When $N_f \in 3\mathbb{Z}$, we have in total \mathbb{Z} copies of 1+1d $3(\text{E}_8)_1$ CFT where the combined rank-24 $K_{3\text{E}_8}$ matrix is the 3 block diagonal copies of K_{E_8} :

$$K_{3\mathrm{E}_8} \equiv K_{\mathrm{E}_8} \oplus K_{\mathrm{E}_8} \oplus K_{\mathrm{E}_8}.\tag{14}$$

There are at least three different constructions of these 1+1d and 2+1d theories:

⁷ For standard terminology, the fermionic theory is called Spin that requires a spacetime manifold with Spin structure so to be compatible with fermions as spinors; the bosonic theory is called non-Spin that spacetime manifold requires no Spin structure. For example, on a 2-torus, we have periodic (P or Ramond [R]) and anti-periodic (AP or Neveu-Schwarz [NS]) boundary conditions along each 1-cycle, so that odd Spin structure includes P-P, while even Spin structure includes AP-AP, P-AP, and AP-P. Bosonization requires to sum over all Spin structures.

⁸ There is a boundary-bulk correspondence between 1+1d (E₈)₁ CFT and 2+1d (E₈)₁ TQFT.

The 1+1d (E₈)₁ CFT is a unique holomorphic Vertex Operator Algebra (VOA) with $c_{-} = 8$.

The 2+1d $(E_8)_1$ TQFT is also known as the E_8 quantum Hall state [40].

⁽¹⁾ $(E_8)_1$: based on the E_8 affine Lie algebra in WZW model and the level-1 non-abelian exceptional simple Lie E_8 gauge group Chern-Simons theory.

⁽²⁾ $SO(16)_1$ up to stack with a trivial $SO(0)_1$ Spin-TQFT: $SO(16)_1$ is Spin-TQFT, which $(E_8)_1$ non-Spin-TQFT stacked with an $SO(0)_1$ spin-TQFT [41].

⁽³⁾ $K_{\rm E_8}$ matrix: The 1+1d compact chiral boson with U(1)⁸ global symmetry, or the 2+1d abelian Chern-Simons theory with U(1)⁸ gauge group, which that the symmetric bilinear form pairing between fields is the Cartan E₈ matrix $K_{\rm E_8}$. The rank-8 $K_{\rm E_8}$ matrix is unimodular, positive definite, even and symmetric rank-8 matrix. The $K_{\rm E_8}$ is also the intersection form of the unique compact simply connected topological E₈ 4-manifold which has a signature $\sigma = 8$.

See further discussions about $3E_8$ and Leech lattices in Sec. IV [42]. To have the 2d theory well-defined on a generic orientable 2-manifold (a genus-g Riemann surface) imposed additional constraints. In particular, we can consider a genus-one 2-torus T^2 . Recall the 2-torus partition function of 2d CFT is

$$\mathbf{Z}(\tau) = \text{Tr}\left(\exp(2\pi i(\tau_1 P - \tau_2 H))\right) = \text{Tr}\left(e^{2\pi i\tau(L_0 - \frac{c}{24})}e^{-2\pi i\bar{\tau}(\bar{L}_0 - \frac{\bar{c}}{24})}\right) = \text{Tr}(q^{L_0 - \frac{c}{24}}\bar{q}^{\bar{L}_0 - \frac{\bar{c}}{24}})$$
(15)

in terms of holomorphic and anti-holomorphic Virasoro generators L_0 and \bar{L}_0 , the modular parameter $\tau = \tau_1 + i\tau_2$ identifies lattice vectors $w \simeq w + 2\pi \simeq w + 2\pi\tau$ as a 2-torus on the complex plane $w = \sigma_1 + i\sigma_2$, $q = e^{2\pi i\tau}$, and trace (Tr) over states in the Hilbert space. The central charges of holomorphic and anti-holomorphic sectors are denoted as $c = c_L$ (left-moving) and $\bar{c} = c_R$ (right-moving) respectively.⁹

Modular invariance demands the $\mathbf{Z}(\tau)$ is invariant under the modular (P)SL(2, \mathbb{Z}) transformations (i.e., the Mapping Class Goup of 2-torus MCG(T^2)) generated by $S: \tau \to -\tau^{-1}$ and the Dehn twist $\mathcal{T}: \tau \to \tau + 1$, so $\mathbf{Z}(-\tau^{-1}) = \mathbf{Z}(\tau)$ and $\mathbf{Z}(\tau+1) = \mathbf{Z}(\tau)$. Modular invariance happens when the chiral central charge

$$c_{-} = c - \bar{c} = c_{L} - c_{R} = 0 \mod 24.$$
⁽¹⁷⁾

thus also framing anomaly free [12, 13]. In our SM reduced model (11), we do have $c_L = 24\mathbb{Z}$ and $c_R = 0$ satisfy (17). Although a generic $c_- = 24\mathbb{Z} \neq 0$ 2d theory still suffers from a 2d perturbative gravitational anomaly that corresponds to 4d anomaly polynomial $\frac{c_-}{24}p_1$ with the 1st Pontryagin class p_1 , but the modular invariance demands the theory have quantized integer spin and angular momentum, which is nice quantum mechanically. We will fill in more details in Sec. IV.

- (3): The 1+1d reduced massless system can be attached to a 2+1d bulk of a trivial class 0 in the 3d framed cobordism and string cobordism: $0 \in TP_3^{fr} \cong TP_3^{String} \cong \mathbb{Z}_{24}$. The corresponding manifold generator is also in the trivial class of 3d framed bordism and string bordism: $0 \in \Omega_3^{fr} \cong \Omega_3^{String} \cong \mathbb{Z}_{24}$.¹⁰ Our SM reduced model (11) obeys this, detailed in Sec. IV.
- (4): The 1+1d reduced massless system can be attached to a 2+1d bulk of a trivial class 0 in the 3d w_1 - p_1 cobordism: $0 \in \operatorname{TP}_3^{w_1-p_1} \cong \mathbb{Z}_3$. The corresponding manifold generator is also in the trivial class of 3d w_1 - p_1 bordism: $0 \in \Omega_3^{w_1-p_1} \cong \mathbb{Z}_3$. Our SM reduced model (11) obeys this, detailed in Sec. IV.
- (5): Smith homomorphism [39]: This step is not necessary but only for further supporting our argument sufficiently. Only for the convenience of reducing from 3+1d to 1+1d, we may implement Wilczek-Zee's $\mathbb{Z}_{4,X}$ symmetry (more discussions in Sec. II B) in the SM [47, 48], so the following Smith homomorphism between bordism groups of Pin structure manifolds [49] (see Fig. 1 (b) and Fig. 2)

$$\dots \to \Omega_5^{\operatorname{Spin} \times_{\mathbb{Z}_2} \mathbb{Z}_4} = \mathbb{Z}_{16} \to \Omega_4^{\operatorname{Pin}^+} = \mathbb{Z}_{16} \to \Omega_3^{\operatorname{Spin} \times \mathbb{Z}_2} = \mathbb{Z}_8 \to \Omega_2^{\operatorname{Pin}^-} = \mathbb{Z}_8 \to \dots$$
(18)

can guide us to map between fermions in different dimensions as (11), in particular in a multiple of 16, detailed in Sec. II B.

We emphasize that our Family Puzzle solution is more topologically robust (depending mainly on (1), (2), (3), (4)) without really relying on any specific global symmetry or any specific gauge group (such as the $\mathbb{Z}_{4,X}$ of (5)). Even breaking $\mathbb{Z}_{4,X}$ symmetry in the SM, our argument still holds.

B. Mass Domain Wall of Domain Wall Reduction Toy Model

The purpose of this subsection is to introduce a concrete model filling the Argument (5) in Sec. II A. As emphasized a few times already, the additional discrete symmetries are not crucial (for solving Family Puzzle), nonetheless

⁹ Momentum $P = L_0 - \bar{L}_0$ generates the σ_1 translation, and Hamiltonian $H = L_0 + \bar{L}_0 - \frac{1}{24}(c+\bar{c})$ generates the σ_2 translation. Viasoro algebra for $m, n \in \mathbb{Z}$ satisfies

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12}(m^3 - m)\delta_{m+n,0}$$
(16)

while $[\bar{L}_m, \bar{L}_n]$ obeys the same but replaced with \bar{c} , and $[L_m, \bar{L}_n] = 0$. The factor of 12 or 24 here comes from Riemann zeta function regularization

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = 1^{-s} + 2^{-s} + 3^{-s} + \dots$$
 and $\zeta(-1) = \sum_{n=1}^{\infty} n = \frac{-1}{12}$

which appears in Casimir energy of the quantum vacuum.

¹⁰ Note that Pontryagin-Thom [43–46] construction bridges between geometric manifold and algebraic homotopy aspects of topology. Under Pontryagin-Thom isomorphism, the inherently geometric aspects of framed cobordism theory (such as *d*-dimensional framed bordism group Ω_d^{fr}) is mapped to the algebraic and topological structure of stable homotopy theory (such as $\lim_{n\to\infty} \pi_{d+n}S^n$). So we have $\Omega_d^{\mathrm{fr}} \cong \lim_{n\to\infty} \pi_{d+n}S^n$, here $\Omega_3^{\mathrm{fr}} \cong \lim_{n\to\infty} \pi_{3+n}S^n = \mathbb{Z}/24\mathbb{Z} = \mathbb{Z}_{24}$. See more in Sec. III and [42].

supportive to realize *not only* dimensional reduction of Fig. 1(a) *but also* the Smith map of Fig. 1(b) simultaneously in one model.

Only for the convenience of the Argument (5) (but not necessary for Argument (1)- (4)), we recall that the SM has an excellent discrete order-four finite abelian group $\mathbb{Z}_{4,X}$ unitary symmetry, where $X \equiv 5(\mathbf{B} - \mathbf{L}) - \frac{2}{3}\tilde{Y} = \frac{5}{N_c}(\mathbf{Q} - N_c\mathbf{L}) - \frac{2}{3}\tilde{Y}$ is a linear combination of the conventional baryon minus lepton $\mathbf{B} - \mathbf{L}$ charge (but more precisely the properly quantized quark number \mathbf{Q} minus N_c times lepton number \mathbf{L}) and the properly integer quantized hypercharge \tilde{Y} [47, 48]. All the quarks and leptons have a unit charge 1 under $\mathbb{Z}_{4,X}$, also $X^2 = (-1)^F$, so $\mathbb{Z}_{4,X} \supset \mathbb{Z}_2^F$. Thus this $\mathbb{Z}_{4,X}$ symmetry is not only preserved with or without SM gauge group G_{SM_q} , but also robust even including four-fermion BSM interaction deformation deviated from the SM fixed point. Given the family number N_f and the total number of sterile right-handed neutrino type $n_{\nu_R} = n_{\nu_{e,R}} + n_{\nu_{\mu,R}} + n_{\nu_{\tau,R}} + \dots$, there is an index $-N_f + n_{\nu_R} \mod 16$ nonperturbative global anomaly classified by \mathbb{Z}_{16} (from the bordism group $\Omega_5^{\mathrm{Spin} \times \mathbb{Z}_2\mathbb{Z}_4} \cong \mathbb{Z}_{16}$ in [17, 50–54], studied recently in the context of SM in [11, 51, 54–59]) captured by the large gauge-diffeomorphism transformations. We shall leave the more formal discussions and mathematical forms of anomaly derivations into Sec. III and in [42].



FIG. 2. Domain wall of domain wall reduction and so on. The extra discrete symmetries (namely, the extra \mathbb{Z}_2 quotient in Spin $\times_{\mathbb{Z}_2} \mathbb{Z}_4$, Pin⁺, Spin $\times \mathbb{Z}_2$, Pin⁻) are not necessary for solving the Family Puzzle, but they are supportive to realize *not only* dimensional reduction of Fig. 1(a) *but also* the Smith map of Fig. 1(b) simultaneously in this model. The colors (purple, blue, green, red) are intentionally meant to match the colors of the dimensional reduction path in Fig. 1(a).

Let us discuss Fig. 2 step by step.

(a): Step 1 Fig. 2 (a), 5d to 4d: We have a 5d bulk massive Dirac fermion of $m(x_4)$ profile (purple) given by (8). The 5d mass term $m\psi_D^5\psi_D^5$ preserves the $\mathbb{Z}_{4,X}$ symmetry: $\psi_D^5 \mapsto i\psi_D^5$. The $\mathbb{Z}_{4,X}$ symmetry projects to 4d massless Weyl fermion that also preserves the $\mathbb{Z}_{4,X}$: $\psi_W^{4d} \mapsto i\psi_W^{4d}$. So this Step 1 scenario is a 5d bulk $\mathbb{Z}_{4,X}$ -SPTs (5d $\mathbb{Z}_{4,X}$ -topological superconductor) classified by $\operatorname{TP}_5^{\operatorname{Spin} \times \mathbb{Z}_2 \mathbb{Z}_4} \cong \mathbb{Z}_{16}$ with a symmetry-preserving massless (gapless) 4d boundary (blue) at $x_4 = 0$. The 5d-4d bulk-boundary system thus far has unitary $X^2 = (-1)^F$ thus $\operatorname{Spin} \times_{\mathbb{Z}_2} \mathbb{Z}_4$ symmetry.

5d Dirac mass term breaks the charge conjugation C and reflection R symmetries. But the R₄ symmetry (spatial reflection along x_4) becomes an internal symmetry at x_4 , which turns out to be the same internal symmetry X we desired for: $R_4(x_4 = 0) = X$.

(b): Step 2 Fig. 2 (b), 4d to 3d: We then give the 4d Weyl fermion a 4d Majorana mass of $m(x_3)$ profile (blue) given by (8). We then obtain the domain wall inside the domain wall. The 4d Majorana mass $m\psi_W^{\intercal}(-i\sigma^2)\psi_W + h.c.$ breaks the $\mathbb{Z}_{4,X}$: $\psi_W^{4d} \mapsto i\psi_W^{4d}$. But there is an antiuntary \mathbb{Z}_4^T symmetry that both the 4d Majorana fermion with Majorana mass

preserves and the 3d massless Majorana fermion at $x_3 = 0$ preserves. So this Step 2 scenario is a 4d bulk $\mathbb{Z}_4^{\mathrm{T}}$ -SPTs (4d $\mathbb{Z}_4^{\mathrm{T}}$ -topological superconductor [60, 61] classified by $\mathrm{TP}_4^{\mathrm{Pin}^+} \cong \mathbb{Z}_{16}$ [23, 26]) with a symmetry-preserving massless (gapless) Majorana fermion 3d boundary (green) at $x_3 = 0$. The 4d-3d bulk-boundary system thus far has antiunitary $\mathrm{T}^2 = (-1)^{\mathrm{F}}$ thus Pin⁺ symmetry.

(c): Step 3 Fig. 2 (c), 3d to 2d: We then give the 3d Majorana fermion a Majorana mass of $m(x_2)$ profile (green) given by (8). We then obtain the domain wall inside the domain wall inside the domain wall. The 3d Majorana mass breaks the $\mathbb{Z}_4^{\mathrm{T}}$, while the 2d domain wall's Majorana-Weyl fermion (at $x_2 = 0$) also breaks $\mathbb{Z}_4^{\mathrm{T}}$ symmetry down to $\mathbb{Z}_2^{\mathrm{F}}$.

In summary of the above, we not only realize (7):

4+1d Dirac
$$\psi_{\rm D}^{5d} \rightarrow$$
 3+1d Weyl $\psi_{\rm W}^{4d} \sim$ 3+1d Majorana $\psi_{\rm M}^{4d} \rightarrow$ 2+1d Majorana $\psi_{\rm M}^{3d} \rightarrow$ 1+1d Majorana-Weyl $\psi_{\rm MW}^{2d}$,

but also enhanced it by cobordism version of Smith map (18):

$$\dots \to \operatorname{TP}_{3}^{\operatorname{Spin} \times \mathbb{Z}_{2}} = \mathbb{Z} \oplus \mathbb{Z}_{8} \to \operatorname{TP}_{4}^{\operatorname{Pin}^{+}} = \mathbb{Z}_{16} \to \operatorname{TP}_{5}^{\operatorname{Spin} \times_{\mathbb{Z}_{2}} \mathbb{Z}_{4}} = \mathbb{Z}_{16} \to \dots$$
(19)

The anomaly indices of (7) are mapped under (19) as

$$(\mathbf{\nu}' = 2(c_L - c_R) = 1, \mathbf{\nu}_2 = 0)|_{\in \mathbb{Z} \oplus \mathbb{Z}_8} \to \mathbf{\nu}_3 = 1|_{\in \mathbb{Z}_{16}} \to \mathbf{\nu}_4 = 1|_{\in \mathbb{Z}_{16}}.$$
(20)

This satisfies a constraint derived in [17]: $\nu' + 2\nu_2 = 2(c_L - c_R) + 2\nu_2 = \nu_3 \mod 16$. We also complete the Argument (5).¹¹

III. FROM 6D ANOMALY POLYNOMIAL, 5D CONVERTIBLE FIELD THEORY, 4D ANOMALY OF THE SM TO 3D GRAVITATIONAL CHERN-SIMONS TERM

The purpose of this subsection is to fill the anomaly perspectives of the Argument (1), (2), (3), and (4) in Sec. II A.

Follow the SM computation in [11], the anomaly polynomial of Weyl fermions follows the Atiyah-Singer index theorem. The 4d anomaly of a single Weyl fermion in 4d is the degree 6 part of the 6d anomaly polynomial from $\hat{A} \operatorname{ch}(\mathcal{E})$.¹² The explicit expression in terms of Pontryagin and Chern characteristic classes [62–64], p_j and c_j , can be obtained using the expansions of \hat{A} and $\operatorname{ch}(\mathcal{E})$:

$$\hat{A} = 1 - \frac{p_1}{24} + \frac{7p_1^2 - 4p_2}{5760} + \dots,$$
(21)

$$\operatorname{ch}(\mathcal{E}) = \operatorname{rank} \mathcal{E} + c_1(\mathcal{E}) + \frac{1}{2} \left(c_1^2(\mathcal{E}) - 2c_2(\mathcal{E}) \right) + \frac{1}{6} \left((c_1^3(\mathcal{E}) - 3c_1(\mathcal{E})c_2(\mathcal{E}) + 3c_3(\mathcal{E})) + \dots \right)$$
(22)

The explicit 6d anomaly polynomial for the gauge, global, and diffeomorphism symmetries of the 4d SM, $G = \text{Spin} \times \text{U}(1)_{\mathbf{Q}} \times \text{U}(1)_{\mathbf{L}} \times G_{\text{SM}_{q}}$ (or more properly $G = \text{Spin} \times_{\mathbb{Z}_{2}^{\text{F}}} \text{U}(1)_{\mathbf{Q}-N_{c}\mathbf{L}} \times_{\mathbb{Z}_{2}^{\text{F}}} \mathbb{Z}_{2N_{c}N_{f},\mathbf{Q}+N_{c}\mathbf{L}} \times G_{\text{SM}_{q}}$ concerning the extra U(1) to be mixed $G_{\text{SM}_{c}}$ -anomaly free), with the matter representation given in (4) becomes:¹³

$$I_{6} \equiv \left(N_{c}c_{1}(\mathrm{U}(1)_{\mathbf{Q}}) + c_{1}(\mathrm{U}(1)_{\mathbf{L}})\right)N_{f}\left(-18\frac{c_{1}(\mathrm{U}(1)_{\tilde{Y}})^{2}}{2} - c_{2}(\mathrm{SU}(2))\right) + \left(N_{f} - n_{\nu_{R}}\right)\left(\frac{c_{1}(\mathrm{U}(1)_{\mathbf{L}})^{3}}{6} - \frac{c_{1}(\mathrm{U}(1)_{\mathbf{L}})p_{1}(TM)}{24}\right), \quad (23)$$

When M^6 is a closed 6-manifold, then $\int_{M^6} I_6 \in \mathbb{Z}$, and there is a 6d *invertible* topological field theory (iTFT) with the partition function $\exp(i \int \theta I_6)$ where $\theta \in [0, 2\pi)$. When M^6 has a boundary $\partial M^6 = M^5$, we can consider this M^5 as a 5d interface between two 6d bulks with the Lagrangian density θI_6 such that $\theta = 0$ on one 6d side and $\theta = 2\pi$ on

¹¹ From this domain wall reduction discussion, we indeed physically "prove" the ratio of $\Omega_4^{SO} = \mathbb{Z}$ over $\Omega_4^{Spin} = 16\mathbb{Z}$ is isomorphic to $\mathbb{Z}/(16\mathbb{Z}) = \mathbb{Z}_{16} \cong \Omega_4^{Pin^+} \cong \Omega_5^{Spin \times \mathbb{Z}_2 \mathbb{Z}_4}$.

¹² Here \hat{A} is the A-roof genus of the spacetime tangent bundle TM over the base spacetime manifold M, expressed in terms of j-th Pontryagin classes p_j , while the ch is the total Chern character expressed in terms of j-th Chern classes c_j , and \mathcal{E} is the complex vector bundle associated with the representation of the fermion. We also use the properties $ch(\mathcal{E}_1 \oplus \mathcal{E}_2) = ch(\mathcal{E}_1) + ch(\mathcal{E}_2)$, and $ch(\mathcal{E}_1 \otimes \mathcal{E}_2) = ch(\mathcal{E}_1) ch(\mathcal{E}_2)$.

¹³ To obtain the polynomial coefficients correctly, here we use the convention such that every fermion in 4d is written as a left-handed Weyl spinor in 4d (left-handed particle ψ_L or right-handed anti-particle $i\sigma_2\psi_R^*$). Every particle contributes +1 (e.g., ψ_L) and every anti-particle contributes -1 (e.g., $i\sigma_2\psi_R^*$), to the quark **Q** or lepton **L** number, namely the integer charge representation of U(1)_{**Q**} or U(1)_{**L**}. We abbreviate $c_j(\mathcal{E}_G) \equiv c_j(\mathcal{G})$ the *j*-th Chern class of the vector bundle \mathcal{E}_G associated with the defining representation of G, and $p_j(TM)$ is the *j*-th Pontryagin class of the spacetime tangent bundle TM.

the other 6d side. On the M^5 interface, we have an iTFT with the action $S_5 = 2\pi \int_{M^5} I_5 \in 2\pi \mathbb{R}$ from $I_6 = dI_5$. The 5d iTFT partition function is $\exp(iS_5) \in U(1)$. The S_5 value modulo 2π is independent of the choice of M^6 . From the 6d anomaly polynomial (23), the explicit 5d iTFT is

$$S_5 \equiv \int_{M^5} (N_c A_{\mathbf{Q}} + A_{\mathbf{L}}) N_f \left(-18 \, \frac{c_1(\mathrm{U}(1)_{\tilde{Y}})^2}{2} - c_2(\mathrm{SU}(2)) \right) + (N_f - n_{\nu_R}) \, A_{\mathbf{L}} \left(\frac{c_1(\mathrm{U}(1)_{\mathbf{L}})^2}{6} - \frac{p_1(TM)}{24} \right). \tag{24}$$

Here $A_{\mathbf{Q}}$ and $A_{\mathbf{L}}$ are background fields for $U(1)_{\mathbf{Q}}$ and $U(1)_{\mathbf{L}}$ symmetries respectively. This 5d TQFT encodes the anomaly of the 4d SM by the anomaly inflow.

A. Spin $\times_{\mathbb{Z}_2^{\mathsf{F}}} \mathrm{U}(1)_{\mathbf{Q}-N_c \mathbf{L}} \equiv \mathrm{Spin}^c$ and $\mathrm{Spin} \times_{\mathbb{Z}_2^{\mathsf{F}}} \mathbb{Z}_{4,X}$ with a vector $\mathrm{U}(1)_{\mathbf{Q}-N_c \mathbf{L}}$ and a chiral X

Concerning the Spin $\times_{\mathbb{Z}_{2}^{F}} U(1) \equiv \text{Spin}^{c}$ structure with $A_{\mathbf{Q}-N_{c}\mathbf{L}}$ background field for $U(1)_{\mathbf{Q}-N_{c}\mathbf{L}}$, the (24) becomes

$$S_5 \equiv (-N_f + n_{\nu_R}) \int_{M^5} A_{\mathbf{Q}-N_c \mathbf{L}} \left(N_c^3 \frac{c_1 (\mathbf{U}(1)_{\mathbf{Q}-N_c \mathbf{L}})^2}{6} - N_c \frac{p_1(TM)}{24} \right).$$
(25)

In the familiar form, we have the relation between Pontryagin class p_1 and gravitational Chern-Simons (GCS) 3-form:

$$p_{1} \coloneqq -\frac{1}{8\pi^{2}} \operatorname{Tr}[R \wedge R] = -\operatorname{dGCS}/(2\pi),$$

$$\operatorname{GCS} \coloneqq \frac{1}{4\pi} \operatorname{Tr}[\omega \wedge \mathrm{d}\omega + \frac{2}{3}\omega \wedge \omega \wedge \omega],$$
(26)

(where $R = d\omega + \omega \wedge \omega$ is the curvature 2-form of the Levi-Civita spin-connection 1-form ω) and the relation between Chern class and Chern-Simons (CS) 3-form:

$$-c_{2} + \frac{1}{2}c_{1}^{2} \coloneqq \frac{1}{8\pi^{2}}\operatorname{Tr}(F \wedge F) = \frac{1}{2\pi}\operatorname{dCS},$$

$$\operatorname{CS} \coloneqq \frac{1}{4\pi}\operatorname{Tr}[A \wedge dA + \frac{2}{3}A \wedge A \wedge A]$$
(27)

 $(F = dA + A \land A$ is the curvature 2-form of gauge connection 1-form). Note however the topologically invariant data from a characteristic class is *not* captured by its local expression in a single patch or chart, but instead is typically captured by the transition functions between different overlapping patches. So in order to define characteristic classes differential-geometrically, we *cannot* just use differential forms *locally*, but we need to define them *globally*. We will also write down the more well-defined global expression later [42].

By restricting Spin $\times_{\mathbb{Z}_2^F} U(1) \equiv \operatorname{Spin}^{\widetilde{c}}$ to Spin $\times_{\mathbb{Z}_2} \mathbb{Z}_4$ (e.g. $U(1)_{\mathbf{Q}-N_c\mathbf{L}} \supset \mathbb{Z}_{4,X} \supset \mathbb{Z}_2^F$), the (25) becomes a \mathbb{Z}_{16} class 5d iTFT: ¹⁴

$$S_5 \equiv (-N_f + n_{\nu_R}) \frac{2\pi}{16} \eta_{4d} (\text{PD}(A_{\mathbb{Z}_{2,X}})) \big|_{M^5}.$$
(28)

Let us comments about the physics of the $(-N_f + n_{\nu_R})$ coefficient in (25) and (28):

- For $N_f = 3$ with 16 Weyl-fermion SM scenario, the $-N_f + n_{\nu_R} = -3 + 3 = 0$, so all these \mathbb{Z} and \mathbb{Z}_{16} anomalies in (25) and (28) cancel. There is no anomaly descendant to 2d theory that we can look for.
- However, when $-N_f + n_{\nu_R} \neq 0$, like the scenario of Ultra Unification [55–59], we do have a room to descend the U(1)³, U(1)-gravity, and \mathbb{Z}_4 -gravity anomalies in 4d to the U(1)² and gravity-gravity anomalies in 2d. This descended anomaly in 2d can give constraints on the dimensional reduced domain wall theory.

• Here the 5d Atiyah-Patodi-Singer (APS) eta-invariant $\eta_{5d} = \eta_{4d}(\text{PD}(A_{\mathbb{Z}_{2,X}}))$ is valued in $\mathbb{Z}_{16} \equiv \mathbb{Z}/(16\mathbb{Z})$ and is written as the 4d eta invariant $\eta_{4d} \in \mathbb{Z}_{16}$ on the 4d Pin⁺ submanifold representing Poincaré dual (PD) to $A_{\mathbb{Z}_{2,X}}$. The Pin⁺ structure is obtained from the 5d bulk Spin $\times_{\mathbb{Z}_{2}^{F}} \mathbb{Z}_{4,X}$ -structure by Smith isomorphism: $\Omega_{5}^{\text{Spin} \times_{\mathbb{Z}_{2}} \mathbb{Z}_{4}} \cong \Omega_{4}^{\text{Pin}^{+}} \cong \mathbb{Z}_{16}$ [17, 50, 52, 53]. The eta invariant $\eta_{4d} \in \mathbb{Z}_{16}$ is the effective topological action of the interacting fermionic time-reversal symmetric topological superconductor of condensed matter in three spatial dimensions with an *anti-unitary* time-reversal symmetry $\mathbb{Z}_{4}^{\text{TF}}$ such that the time-reversal symmetry generator T squares to the fermion parity operator, namely $T^{2} = (-1)^{\text{F}}$. The symmetry can be defined by the nontrivial group extension $1 \to \mathbb{Z}_{2}^{\text{F}} \to \mathbb{Z}_{4}^{\text{TF}} \to \mathbb{Z}$ (see a review [27, 65]). In contrast, in the SM, we have the *unitary* $\mathbf{B} - \mathbf{L}$ -like symmetry $\mathbb{Z}_{4,X} \to \mathbb{Z}_{2,X} \to 1$.

 $^{^{14}}$ Some explanation about the notation [11]:

[•] Because all the quarks and leptons have charge 1 under $\mathbb{Z}_{4,X}$, there is no N_c factor in this 5d iTFT.

[•] The background gauge field $A_{\mathbb{Z}_{2,X}} \in \mathrm{H}^1(M^5, \mathbb{Z}_2)$ is obtained by the quotient map down to $\mathbb{Z}_{2,X} \equiv \mathbb{Z}_{4,X}/\mathbb{Z}_2^{\mathrm{F}}$ from the Spin $\times_{\mathbb{Z}_2^{\mathrm{F}}} \mathbb{Z}_{4,X}$ -structure on the 5d spacetime manifold M^5 .

• The $U(1)_{\mathbf{Q}-N_c\mathbf{L}}$ is a vector symmetry with properly quantized charge of $\mathbf{B} - \mathbf{L}$. Due to the vector U(1)symmetry, the anomalies of left-handed and right-handed Weyl fermions in the SM cancel nicely (nearly, except that if $-N_f + n_{\nu_R} \neq 0$.¹⁵ However, we are motivated to resolve the Family Puzzle by thinking of more robust topological constraint without symmetry, thus we should forget the SM internal (gauge or global) symmetry structure. Once $G_{SM_{\alpha}}$ is removed, we can consider the chiral U(1) symmetry of the Weyl fermions, which has a stronger constraint of the modular invariance of 2d theory and the framing anomaly of the 3d TQFT (as we will show in Sec. III B).

B. Another $\operatorname{Spin} \times_{\mathbb{Z}_2^F} U(1) \equiv \operatorname{Spin}^c$ with a chiral U(1): $3(\operatorname{E}_8)_1$ quantum Hall states

For a single left-handed Weyl fermion of charge q in 4d, we take \mathcal{E} to be the complex line bundle associated with the corresponding representation of U(1). The fermionic 6d anomaly polynomial that captures the 4d anomaly is

$$I_{6,f} = [\hat{A} \operatorname{ch}(\mathcal{E})]_6 = q^3 \frac{c_1^3}{6} - q \frac{c_1 p_1}{24}, \qquad \int_{M^6} I_{6,f} \in \mathbb{Z}.$$
(29)

Consider a collection of left-handed Weyl fermions in 4d with the global U(1) symmetry charges $q_i = 1$ with $i = 1, \ldots, N_f n$ for N_f families of n fermions per family (e.g. $N_f = 3$ and n = 15, 16).

The dimensional reduced 2d theory potential has $U(1)^2$ and gravity-gravity anomalies (2-point function under oneloop diagram) in 2d. However, the domain wall construction given in Sec. II can break all U(1) symmetries, thus we are left with only the perturbative gravitational anomalies, captured by the gravity-gravity 2-point function under one-loop diagram, 3d GCS, 4d p_1 and the signature σ of 4-manifold with proper coefficients given in (31):¹⁶

$$\begin{array}{ll} p_{1} \in \frac{24}{c_{-}}\mathbb{Z}, & \frac{\sigma}{8}c_{-} = \frac{c_{-}}{24}p_{1}, & \frac{c_{-}}{24}\text{GCS} = \frac{c_{-}}{96\pi}\text{Tr}[\omega\,\mathrm{d}\omega + \frac{2}{3}\,\omega^{3}], & c_{-}, & \kappa_{xy} = 2c_{-}\kappa_{xy}^{\text{MW}}, & c_{-} \text{ Gapless/CFT} \\ \hline p_{1} \in 48\mathbb{Z}, & \frac{\sigma}{16} = \frac{1}{48}p_{1}, & \frac{1}{48}\text{GCS} = \frac{1}{192\pi}\text{Tr}[\omega\,\mathrm{d}\omega + \frac{2}{3}\,\omega^{3}], & c_{-} = \frac{1}{2}, & \kappa_{xy} = \kappa_{xy}^{\text{MW}}, & \text{Majorana-Weyl} \\ p_{1} \in 24\mathbb{Z}, & \frac{\sigma}{8} = \frac{1}{24}p_{1}, & \frac{1}{24}\text{GCS} = \frac{1}{96\pi}\text{Tr}[\omega\,\mathrm{d}\omega + \frac{2}{3}\,\omega^{3}], & c_{-} = 1, & \kappa_{xy} = 2\kappa_{xy}^{\text{MW}}, & \text{Weyl with U(1) symmetry} \\ p_{1} \in 3\mathbb{Z}, & \sigma = \frac{1}{3}p_{1}, & \frac{1}{3}\text{GCS} = \frac{1}{12\pi}\text{Tr}[\omega\,\mathrm{d}\omega + \frac{2}{3}\,\omega^{3}], & c_{-} = 8, & \kappa_{xy} = 16\kappa_{xy}^{\text{MW}}, & (\text{E}_{8})_{1} \text{ chiral boson} \\ & 3\sigma = p_{1}, & \text{GCS} = \frac{1}{4\pi}\text{Tr}[\omega\,\mathrm{d}\omega + \frac{2}{3}\,\omega^{3}], & c_{-} = 24, & \kappa_{xy} = 48\kappa_{xy}^{\text{MW}}, & 3(\text{E}_{8})_{1} \text{ or Leech chiral boson} \\ \end{array}$$

Under (7), the 4d Weyl ψ_{W}^{4d} is reduced to 2d Majorana-Weyl ψ_{MW}^{2d} with $c_{-} = 1/2$; the 48 of 4d Weyl ψ_{W}^{4d} is reduced to 48 of 2d Majorana-Weyl $\psi_{\rm MW}^{2d}$ with $c_{-} = 24$.

Index Theorem and (Gravitational) Chern-Simons: Framing, String, 2-Framing, and w_1 - p_1 structures С.

Eq. (31) also shows the information of the topology of 4-manifolds. Hirzebruch signature [10] writes in terms of the L genus: For the dimension $d = 0 \mod 4$, the signature $\sigma(M) \in \mathbb{Z}$ of manifold M^d is

$$\sigma(M^d) = \int_{M^d} L_n = \int_{M^d} L_{d/4} \equiv \langle L_{d/4}, [M^d] \rangle$$

$$L = L_0 + L_1 + L_2 + \dots = 1 + \frac{p_1}{3} + \frac{-p_1^2 + 7p_2}{45} + \dots$$
(32)

and the $[M^d]$ is the fundamental class of M^d . For example, 4-manifold $\sigma(M^4) = \langle \frac{p_1}{3}, [M^4] \rangle \equiv \frac{p_1}{3}$. Under the Special Orthogonal SO structure, the SO (non-Spin) 4-manifold has a signature $\sigma = \frac{p_1}{3} \in \mathbb{Z}$.

Under the Spin structure, the Rokhlin's theorem [37] says the $\sigma = \frac{p_1}{3} \in 16\mathbb{Z}$ for 4-manifolds.

By (7), the 4d Weyl ψ_{W}^{4d} is reduced to 2d Majorana-Weyl ψ_{MW}^{2d} with $c_{-} = 1/2$ which is attached to a 3d bulk of $\frac{1}{48}$ GCS. However, GCS written as Levi-Civita spin-connection 1-form ω is *metric dependent* thus *not* strictly mathematically topological.

Here are some issues:

¹⁶ For condensed matter related physical observable, we write the Thermal Hall conductance [66] in the unit

$$\kappa_{xy}^{\text{MW}} = \frac{\pi k_B^2 T}{12\hbar} = \frac{\pi^2 k_B^2 T}{6\hbar}.$$
(30)

 $^{^{15}}$ For a unit charge 1 of an axial U(1) symmetry in the Weyl fermion basis, we choose the left-handed particle and right-handed antiparticle to have q = 1. We choose the right-handed particle and left-handed anti-particle to have q = -1.

For a unit charge 1 of a vector U(1) symmetry in the Weyl fermion basis, we choose the left-handed particle and right-handed particle to have q = 1. We choose the left-handed anti-particle and right-handed anti-particle to have q = -1.

- 1. We like to address the issue of how to make the theory *metric independent* thus strictly mathematically topological.
- 2. We shall impose framing anomaly-free in 3d TQFT thus related to the modular invariance of 2d CFT (on the boundary).
- 3. We aim to define GCS globally instead of locally.

The answers to address the above issues are:

1. Metric independent and (mathematically) topological: Follow [13], we not only need the 3d GCS in (31), but also the 3d CS gauge theory with a CS action S(A) such as $(E_8)_1$ CS with the gauge field A. We also need gravitational eta invariant η_{grav} [13]. We write

$$S(A) + c_{-}\frac{1}{24}\text{GCS} = c_{-}(\frac{\pi}{2}\eta_{\text{grav}} + \frac{1}{24}\text{GCS}) + (S(A) - c_{-}\frac{\pi}{2}\eta_{\text{grav}})$$

$$= c_{-}(\frac{\pi}{2})(\eta_{\text{grav}} + \frac{1}{12\pi}\text{GCS}) + (S(A) - c_{-}\frac{\pi}{2}\eta_{\text{grav}}).$$
(33)

- S(A): 3d CS gauge theory action of the gauge field A: It is metric-dependent (not topological), but no framing is required.
- $\frac{\pi}{2}\eta_{\text{grav}}$: The gravitational eta invariant version of CS (e.g., of the (E₈)₁ CS). It is also metric-dependent (not topological), but no framing is required.
- $(S(A) c_{-\frac{\pi}{2}}\eta_{\text{grav}})$ is metric-independent (topological) and no framing is required.
- GCS is indeed the gravitational counter term (invertible field theory iFT) from the free part of Ω_4 bordism group. GCS is metric-dependent (non-topological), framing-dependent.
- $\eta_{\text{grav}} + \frac{1}{12\pi} \text{GCS}$ is metric-independent but framing-dependent.
- The combined $S(A) + c_{-\frac{1}{24}}$ GCS is generally metric-independent but framing-dependent.
- 2. Framing anomaly: The partition function / path integral on an oriented 3-manifold M^3 defined by the action (33) is in principle framing anomalous under the change of framing $f \in \pi_3(SO(3)) = \mathbb{Z}$ or $H^3(M^3, \mathbb{Z}) = \mathbb{Z}$,

$$\mathbf{Z} \mapsto \mathbf{Z} \exp(2\pi \mathrm{i} f \frac{c_{-}}{24}). \tag{34}$$

Only when $c_{-} = 0 \mod 24$, which we like to impose, then the theory is not only metric-independent (topological) but also framing anomaly-free. This theory is particularly good like the $3(E_8)_1$ quantum Hall state in Sec. III D.

3. Define GCS or CS globally instead of locally: Recall (3) we define GCS (or CS) globally by putting it on the 3-manifold $M^3 = \partial M^4$ bounding the boundary of a 4-manifold M^4 , and we will use the relation between (a) framing provides the String structure, and (2) 2-framing provides the w_1 - p_1 -structure. For Family Puzzle, we will need the following structures:

 $Structure \coloneqq \begin{cases} (a) \text{ (Fermionic) GCS on Spin manifold} & --\text{framing (fr): trivialization of tangent bundle } TM \\ \simeq String structure: trivialization of <math>w_1(TM), w_2(TM), \text{ and } \frac{1}{2}p_1(TM). \\ (b) \text{ (Bosonic) GCS on non-Spin manifold} & --2\text{-framing (2-fr): trivialization of the spin bundle of 2 copies of } \\ \simeq w_1\text{-}p_1\text{-structure: trivialization of } w_1(TM) \text{ and } p_1(TM). \end{cases}$

Follow [11],

• For 2-framing, the relative characteristic number $\frac{1}{2}p_1(2TM,\beta) \in \mathbb{Z}$ with a trivialization β of the spin bundle of 2 copies of the tangent bundle 2TM. Using this number, one can then define the value of gravitational Chern-Simons action in \mathbb{R} (instead of just in $\mathbb{R}/(2\pi\mathbb{Z})$):

$$\frac{c_{-}}{24} \int_{M^3} \text{GCS} = \frac{c_{-}}{24} 2\pi \left(\frac{1}{8\pi^2} \int_{M^4} \text{Tr}[R \wedge R] + \frac{1}{2} p_1(2TM^4, \beta) \right).$$
(36)

• For p_1 -structure β' on M^4 as a choice of the trivialization of the first Pontryagin class $p_1(TM^3) = 0$ (which vanishes exactly in 3d by dimensional reasons). Similarly extending M^3 to a 4-manifold M^3 , we have an integral

relative characteristic number $p_1(TM^4, \beta') \in \mathbb{Z}$, which can be used to define the value of the gravitational Chern-Simons action in \mathbb{R} (instead of just in $\mathbb{R}/(2\pi\mathbb{Z})$) as follows:

$$\frac{c_{-}}{24} \int_{M^3} \text{GCS} = \frac{c_{-}}{24} 2\pi \left(\frac{1}{8\pi^2} \int_{M^4} \text{Tr}[R \wedge R] + p_1(TM^4, \beta') \right).$$
(37)

Similarly, for CS, we also write (defined mod $\mathbb{R}/(2\pi\mathbb{Z})$) related to $2\pi \int_{M^4} (-c_2 + \frac{1}{2}c_1^2)$:

$$k \int_{M^3} \operatorname{CS}(A) = 2\pi \frac{k}{8\pi^2} \int_{M^4} \operatorname{Tr}(F \wedge F).$$
(38)

D. Cobordism constraints of Framing, String, 2-Framing, and w_1 - p_1 structures

Finally, we like to check the bordism and cobordism constraints from the given structures (Framing, 2-Framing, and w_1 - p_1 structures). We show that when we have $N_f = 3$ families or 48 Weyl fermions in 4d SM, or $c_- = 24$ in 2d, the theory nicely sits in the trivial cobordism class. This will completely prove Argument (3), and (4) in Sec. II A.

First recall, the bordism map of manifolds suggests by the Whitehead tower relation is

	Ω_d^{String} -	$\rightarrow \Omega_d^{\mathrm{Spin}} -$	$\rightarrow \Omega_d^{\mathrm{SC}}$
d = 3	\mathbb{Z}_{24}	0	0
d = 4	0	$\mathbb{Z}^{\times 1}$	$\xrightarrow{6} \mathbb{Z}$

Second, the cobordism map of iFT or iTQFT as Framing or String cobordism invariants becomes¹⁷

$$TP_{d}^{\text{String}} \leftarrow TP_{d}^{\text{Spin}} \leftarrow TP_{d}^{\text{SO}}$$

$$d = 3 \qquad \mathbb{Z}_{24} \xleftarrow{\text{mod } 24} \mathbb{Z} \xleftarrow{\times 16} \mathbb{Z}$$

$$16 \xleftarrow{\text{mod } 24} 16 \xleftarrow{\times 16} 1$$

$$0 \xleftarrow{\text{mod } 24} 48 \xleftarrow{\times 16} 3 \quad . \tag{39}$$

We show that $p_1/3$ maps to 16 mod 24 (thus $c_- = 0 \mod 8$) in the string cobordism $\text{TP}_3^{\text{String}}$ which is nonzero. But the p_1 maps to $3 \times 16 = 48 = 0 \mod 24$ (thus $c_- = 0 \mod 24$) in the $\text{TP}_3^{\text{String}}$ which vanishes!

Third, the cobordism map of iFT or iTQFT as 2-Framing or w_1 - p_1 cobordism invariants becomes¹⁸

$$TP_{d}^{w_{1},p_{1}} \leftarrow TP_{d}^{SO}$$

$$d = 3 \qquad \mathbb{Z}_{3} \xleftarrow{}^{\text{mod }3} \mathbb{Z}$$

$$1 \xleftarrow{}^{\text{mod }3} 1$$

$$0 \xleftarrow{}^{\text{mod }3} 3 \qquad (40)$$

We show that $p_1/3$ maps to 1 mod 3 (thus $c_- = 0 \mod 8$) in the w_1 - p_1 cobordism $\operatorname{TP}_3^{w_1,p_1}$ which is nonzero. But the p_1 maps to $3 = 0 \mod 3$ (thus $c_- = 0 \mod 24$) in the $\operatorname{TP}_3^{w_1,p_1}$ which vanishes!

IV. CONCLUSION AND COMPARISON

Above we have presented a theoretical solution as to why the family number $N_f = 3$ or its multiple $N_f \in 3\mathbb{Z}$ is favored due to modular invariant and framing anomaly-free. Especially with 16 Weyl fermion per generation in the SM, we map this $N_f \in 3\mathbb{Z}$ SM to a 1+1d chiral central charge $c_- = 24\mathbb{Z}$ CFT by the dimensional reduction.

We address additional refined questions about our proposed solution:

¹⁷ We derive $\operatorname{TP}_d^{\operatorname{String}}$ (trivialize $w_1, w_2, \frac{1}{2}p_1$ [framing]) from the ratio of the allowed relative $\frac{1}{2}p_1$ class: $\frac{\frac{1}{2}p_1 \in \mathbb{Z}}{\frac{1}{3}p_1 \in 16\mathbb{Z}} = \frac{p_1 \in \mathbb{Z}}{p_1 \in 48\mathbb{Z}} = \mathbb{Z}_{24}$. ¹⁸ We derive $\operatorname{TP}_d^{w_1, p_1}$ (trivialize w_1, p_1 [Atiyah's 2-framing]) from the ratio of the allowed relative p_1 class: $\frac{p_1 \in \mathbb{Z}}{\frac{1}{3}p_1 \in \mathbb{Z}} = \frac{p_1 \in \mathbb{Z}}{p_1 \in 3\mathbb{Z}} = \mathbb{Z}_3$.

1: The 4th family? More than $N_f = 3$ families?

I the hypothetical 4th family or more new families of quarks and leptons do not couple to the SM Higgs nor gain mass from the SM Higgs mechanism, and also if these new families are heavier than $m_Z/2$, then the contemporary experiments have not yet ruled out those possibilities — they are novel mass-generating mechanisms (without symmetry-breaking Higgs fields) known as Symmetric Mass Generation (SMG, see a recent overview [67]). The SMG gapping out the 4th family SM can still survive under the known experimental constraints above as a valid theoretical possibility [68–73], potentially relevant for the Strong CP problem as well [73, 74]. If $N_f \in 3\mathbb{Z}$ constraint still holds, it may suggests a group of 3 more families at much higher-energy above the low-energy SM. It seems more likely that $N_f \in 3\mathbb{Z}$ only constrains more on the lower energy spectrum. It will be interesting to sharpen the statement regarding the energy scales.

2: 16 Weyl fermions vs 15 Weyl fermions?

We can comment on the 3 families of 15 Weyl fermions, the absence of sterile right-handed neutrino replaced by topological field theory, and its $\frac{45}{15} = 3$ relation to Ultra Unification. Notice that the missing sterile neutrinos also appear to be 3 (so far). So the ratio $\frac{45}{15} = 3$ holds. For the reduced 2d CFT, we can obtain $c_{-} = 45/2$ but we miss $c_{-} = 3/2$, which suggests a possible extra Pfaffian-like non-abelian Quantum Hall states [75] with $c_{-} = 3/2$ in the dimensional reduced 3d bulk as well. This picture could match with the TQFT sector for Ultra Unification.

3: Our argument is topologically robust without requiring any global symmetry and gauge group structure: Even breaking or forgetting all SM gauge group structures or global symmetry structures, our constraints ((1), (2), (3), (4) in Sec. II A) still favor the N_f ∈ 3Z family.

In comparison to the literature, we requires no $\mathbb{Z}_{4,X}$ symmetry's \mathbb{Z}_{16} nonperturbative anomaly in 4d, nor additional \mathbb{Z}_3 symmetry's \mathbb{Z}_9 nonperturbative anomaly in 4d for baryon triality or proton hexality in [51], nor the homotopy group analysis of $G_{\rm SM_q}$ [76].¹⁹ Our solution is entirely distinct from the previous proposals [51, 76] that require specific global or gauge internal symmetry constraints.

4: E_8 , $3E_8$, and Leech lattices: Prediction of additional gapped sectors above the SM?

Along the discussion in (13), indeed there exists proper $SL(N, \mathbb{Z})$ transformations to map between the fermionic $K_f = \mathbb{I}_8$ matrix of 8 Weyl fermions to the bosonic K_{E_8} by enlarging the matrix along the diagonal block via introducing extra rank-2 canonical unimodular fermionic matrix $K_{f,2} \equiv \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}$. There exists also $SL(N,\mathbb{Z})$ transformations to map between $K_{3E_8} \equiv K_{E_8} \oplus K_{E_8} \oplus K_{E_8}$ to the other 23 out of the 24 unimodular rank-24 Niemeier lattices, by enlarging the matrix along the diagonal block via introducing extra rank-2 canonical unimodular bosonic matrix $K_{b,2} \equiv \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. One of the most famous kind of Niemeier lattices is the rank-24 Leech lattice K_{Leech} . (Beware that there are also other 1+1d holomorphic but also meromorphic $c_- = 24 \text{ CFT}$ [78].) Recall that both the E_8 lattice and Leech lattice (as a discrete subgroup of \mathbb{R}^8 and \mathbb{R}^{24} respectively) are the solutions to the Spherical Packing Problem [79] in 8 and 24 dimensions [80, 81]. Since the 48 Weyl-fermion version of the 3+1d SM can be mapped to 1+1d 3(E_8)_1 CFT of K_{3E_8} , one could ask what does the 1+1d Leech lattice CFT of K_{Leech} imply? — Does the rank-24 Leech lattice have any use of prediction about the 3+1d SM and BSM real-world physics? It implies that by adding additional gapped bosonic sectors from copies of 1+1d non-chiral bosonic or fermion CFT of $K_{b,2}$ or $K_{f,2}$, bringing these gapped sector down and allowing interactions between new sectors and the SM sector, we can deform the SM's $3(E_8)_1$ CFT to the Leech lattice CFT. The inverse map from the copies of 1+1d non-chiral bosonic CFT sector to the 3+1d sectors may predict other hidden BSM sectors [42].

5: Apparently since we refer to the E_8 lattice and Leech lattice in the Spherical Packing, potential relation to the Conformal Bootstrap, Moonshine (Monstrous moonshine, Mathieu moonshine), Sporadic group and Monster CFT may also help to connect to the Family $N_f \in 3\mathbb{Z}$ structure in the SM.

ACKNOWLEDGMENTS

JW appreciates the inspiring conversations (either discussions on the related topics or the feedback) with Meng Cheng, Dan Freed, Inaki Garcia-Etxebarria, Michael Hopkins, Zohar Komargodski, Seth Koren, Stephen McKean,

¹⁹ Although [76] uses the 6d homotopy group, $\pi_6(SU(2)) = \mathbb{Z}_{12}$, $\pi_6(SU(3)) = \mathbb{Z}_6$ and $\pi_6(G_2) = \mathbb{Z}_3$, to argue the nonperturbative global anomaly constraints in 6d, we find that the cobordism classification of nonperturbative global anomaly constraints showing them vanish thus these anomaly constraints may not really exist: $\Omega_7^{\text{Spin} \times G_{\text{SM}_q}} = 0$ [77] shows no 6d nonperturbative global anomalies.

Jake McNamara, Miguel Montero, Gregory Moore, Pavel Putrov, Matthew Reece, Clifford Taubes, Constantin Teleman, Zheyan Wan, Xiao-Gang Wen, Edward Witten, David Wu, Yizhuang You, and Yunqin Zheng. JW is supported by Harvard University CMSA.

- H. Harari, "Three generations of quarks and leptons," in Presented at 5th Intern. Conf. on Exptl. Meson Spectry (1977) pp. 29–30.
- [2] Makoto Kobayashi and Toshihide Maskawa, "CP Violation in the Renormalizable Theory of Weak Interaction," Prog. Theor. Phys. 49, 652–657 (1973).
- [3] F. Abe *et al.* (CDF), "Observation of top quark production in $\bar{p}p$ collisions," Phys. Rev. Lett. **74**, 2626–2631 (1995), arXiv:hep-ex/9503002.
- [4] S. Abachi et al. (D0), "Observation of the top quark," Phys. Rev. Lett. 74, 2632–2637 (1995), arXiv:hep-ex/9503003.
- [5] Otto Eberhardt, Geoffrey Herbert, Heiko Lacker, Alexander Lenz, Andreas Menzel, Ulrich Nierste, and Martin Wiebusch, "Impact of a Higgs boson at a mass of 126 GeV on the standard model with three and four fermion generations," Phys. Rev. Lett. 109, 241802 (2012), arXiv:1209.1101 [hep-ph].
- [6] D. Decamp et al. (ALEPH), "Determination of the Number of Light Neutrino Species," Phys. Lett. B 231, 519–529 (1989).
- [7] S. Schael *et al.* (ALEPH, DELPHI, L3, OPAL, SLD, LEP Electroweak Working Group, SLD Electroweak Group, SLD Heavy Flavour Group), "Precision electroweak measurements on the Z resonance," Phys. Rept. 427, 257–454 (2006), arXiv:hep-ex/0509008.
- [8] C. L. Bennett *et al.* (WMAP), "Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results," Astrophys. J. Suppl. 208, 20 (2013), arXiv:1212.5225 [astro-ph.CO].
- [9] Zheyan Wan, Juven Wang, and Xiao-Gang Wen, "(3+1)d boundaries with gravitational anomaly of (4+1)d invertible topological order for branch-independent bosonic systems," Phys. Rev. B 106, 045127 (2022), arXiv:2112.12148 [condmat.str-el].
- [10] F. Hirzebruch, Topological Methods in Algebraic Geometry, Classics in mathematics (Springer-Verlag, 1954, 1978).
- [11] Pavel Putrov and Juven Wang, "Categorical Symmetry of the Standard Model from Gravitational Anomaly," (2023), arXiv:2302.14862 [hep-th].
- [12] P. Di Francesco, P. Mathieu, and D. Senechal, *Conformal Field Theory*, Graduate Texts in Contemporary Physics (Springer-Verlag, New York, 1997).
- [13] Edward Witten, "Quantum Field Theory and the Jones Polynomial," Commun. Math. Phys. 121, 351–399 (1989).
- [14] Zheyan Wan, Juven Wang, Shing-Tung Yau, and Yi-Zhuang You, "C-R-T Fractionalization, Fermions, and Mod 8 Periodicity," to appear (to appear).
- [15] R. Jackiw and C. Rebbi, "Solitons with Fermion Number 1/2," Phys. Rev. D 13, 3398–3409 (1976).
- [16] Curtis G. Callan, Jr. and Jeffrey A. Harvey, "Anomalies and Fermion Zero Modes on Strings and Domain Walls," Nucl. Phys. B250, 427–436 (1985).
- [17] Itamar Hason, Zohar Komargodski, and Ryan Thorngren, "Anomaly Matching in the Symmetry Broken Phase: Domain Walls, CPT, and the Smith Isomorphism," SciPost Phys. 8, 062 (2020), arXiv:1910.14039 [hep-th].
- [18] Clay Córdova, Kantaro Ohmori, Shu-Heng Shao, and Fei Yan, "Decorated Z₂ symmetry defects and their time-reversal anomalies," Phys. Rev. D 102, 045019 (2020), arXiv:1910.14046 [hep-th].
- [19] Juven Wang, Yi-Zhuang You, and Yunqin Zheng, "Gauge enhanced quantum criticality and time reversal deconfined domain wall: SU(2) Yang-Mills dynamics with topological terms," Phys. Rev. Research. 2, 013189 (2020), arXiv:1910.14664 [cond-mat.str-el].
- [20] Arun Debray, Sanath K. Devalapurkar, Cameron Krulewski, Yu Leon Liu, Natalia Pacheco-Tallaj, and Ryan Thorngren, "A Long Exact Sequence in Symmetry Breaking: order parameter constraints, defect anomaly-matching, and higher Berry phases," (2023), arXiv:2309.16749 [hep-th].
- [21] David B. Kaplan, "A Method for simulating chiral fermions on the lattice," Phys. Lett. B 288, 342–347 (1992), arXiv:heplat/9206013.
- [22] Daniel S. Freed, "Bordism: Old and new," (2003), online: https://web.ma.utexas.edu/users/dafr/bordism.pdf.
- [23] Daniel S. Freed and Michael J. Hopkins, "Reflection positivity and invertible topological phases," arXiv e-prints, arXiv:1604.06527 (2016), arXiv:1604.06527 [hep-th].
- [24] M. F. Atiyah, V. K. Patodi, and I. M. Singer, "Spectral asymmetry and Riemannian geometry 1," Math. Proc. Cambridge Phil. Soc. 77, 43 (1975).
- [25] Anton Kapustin, "Symmetry Protected Topological Phases, Anomalies, and Cobordisms: Beyond Group Cohomology," (2014), arXiv:1403.1467 [cond-mat.str-el].
- [26] Anton Kapustin, Ryan Thorngren, Alex Turzillo, and Zitao Wang, "Fermionic Symmetry Protected Topological Phases and Cobordisms," JHEP 12, 052 (2015), [JHEP12,052(2015)], arXiv:1406.7329 [cond-mat.str-el].
- [27] Meng Guo, Pavel Putrov, and Juven Wang, "Time reversal, SU(N) Yang-Mills and cobordisms: Interacting topological superconductors/insulators and quantum spin liquids in 3 + 1 D," Annals of Physics 394, 244–293 (2018), arXiv:1711.11587 [cond-mat.str-el].
- [28] Zheyan Wan and Juven Wang, "Higher Anomalies, Higher Symmetries, and Cobordisms I: Classification of Higher-Symmetry-Protected Topological States and Their Boundary Fermionic/Bosonic Anomalies via a Generalized Cobordism

Theory," Ann. Math. Sci. Appl. 4, 107–311 (2019), arXiv:1812.11967 [hep-th].

- [29] Edward Witten and Kazuya Yonekura, "Anomaly Inflow and the η-Invariant," in The Shoucheng Zhang Memorial Workshop Stanford, CA, USA, May 2-4, 2019 (2019) arXiv:1909.08775 [hep-th].
- [30] Luis Alvarez-Gaume and Edward Witten, "Gravitational Anomalies," Nucl. Phys. B 234, 269 (1984).
- [31] Edward Witten, "Global Gravitational Anomalies," Commun. Math. Phys. 100, 197 (1985), [,197(1985)].
- [32] Edward Witten, "An SU(2) Anomaly," Phys. Lett. 117B, 324–328 (1982), [,230(1982)].
- [33] Juven Wang, Xiao-Gang Wen, and Edward Witten, "A New SU(2) Anomaly," J. Math. Phys. 60, 052301 (2019), arXiv:1810.00844 [hep-th].
- [34] John W. Milnor and James D. Stasheff, *Characteristic classes*, Annals of Mathematics Studies, No. 76 (Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974) pp. vii+331.
- [35] Michael Atiyah, "On framings of 3-manifolds," Topology 29, 1–7 (1990).
- [36] Daniel S. Freed, "The Atiyah-Singer index theorem," Bull. Am. Math. Soc. 58, 517–566 (2021), arXiv:2107.03557 [math.HO].
- [37] Vladimir Rokhlin, "New results in the theory of four-dimensional manifolds," Doklady Akademii Nauk SSSR 84, 221–224 (1952).
- [38] Steven Weinberg, The quantum theory of fields. Vol. 2: Modern applications (Cambridge University Press, 2013).
- [39] P. A. Smith, "New results and old problems in finite transformation groups," Bulletin of the American Mathematical Society 66, 401–415 (1960).
- [40] A. Kitaev, "Anyons in an exactly solved model and beyond," Annals of Physics **321**, 2–111 (2006), cond-mat/0506438.
- [41] Nathan Seiberg and Edward Witten, "Gapped Boundary Phases of Topological Insulators via Weak Coupling," PTEP 2016, 12C101 (2016), arXiv:1602.04251 [cond-mat.str-el].
- [42] Juven Wang and et al, "Generation Problem, Gravitational Chern-Simons, String Cobordism, and Modularity: $\frac{48}{16} = \frac{45}{15} = \frac{24}{2} = 3$," to appear (to appear).
- [43] Lev Pontryagin, "A classification of mappings of the three-dimensional complex into the two-dimensional sphere," Doklady Akademii Nauk SSSR 31, 129–132 (1941).
- [44] Lev Pontryagin, "Smooth manifolds and their applications in homotopy theory," American Mathematical Society Translations, Series 2 11, 1–114 (1950).
- [45] René Thom, Espaces fibrés en sphères et carrés de Steenrod, Ph.D. thesis (1952).
- [46] René Thom, "Quelques propriétés globales des variétés différentiables," Commentarii Mathematici Helvetici 28, 17–86 (1954).
- [47] Frank Wilczek and A. Zee, "Operator Analysis of Nucleon Decay," Phys. Rev. Lett. 43, 1571–1573 (1979).
- [48] Frank Wilczek and A. Zee, "Conservation or Violation of B L in Proton Decay," Phys. Lett. B 88, 311-314 (1979).
- [49] R. C. Kirby and L. R. Taylor, "Pin structures on low-dimensional manifolds," in Geometry of low-dimensional manifolds, 2 (Durham, 1989), London Math. Soc. Lecture Note Ser., Vol. 151 (Cambridge Univ. Press, Cambridge, 1990) pp. 177–242.
- [50] Yuji Tachikawa and Kazuya Yonekura, "Why are fractional charges of orientifolds compatible with Dirac quantization?" arXiv e-prints, arXiv:1805.02772 (2018), arXiv:1805.02772 [hep-th].
- [51] Inaki Garcia-Etxebarria and Miguel Montero, "Dai-Freed anomalies in particle physics," JHEP 08, 003 (2019), arXiv:1808.00009 [hep-th].
- [52] Chang-Tse Hsieh, "Discrete gauge anomalies revisited," (2018), arXiv:1808.02881 [hep-th].
- [53] Meng Guo, Kantaro Ohmori, Pavel Putrov, Zheyan Wan, and Juven Wang, "Fermionic Finite-Group Gauge Theories and Interacting Symmetric/Crystalline Orders via Cobordisms," Commun. Math. Phys. 376, 1073 (2020), arXiv:1812.11959 [hep-th].
- [54] Zheyan Wan and Juven Wang, "Beyond Standard Models and Grand Unifications: Anomalies, Topological Terms, and Dynamical Constraints via Cobordisms," JHEP 07, 062 (2020), arXiv:1910.14668 [hep-th].
- [55] Juven Wang, "Anomaly and Cobordism Constraints Beyond the Standard Model: Topological Force," (2020), arXiv:2006.16996 [hep-th].
- [56] Juven Wang, "Anomaly and Cobordism Constraints Beyond Grand Unification: Energy Hierarchy," (2020), arXiv:2008.06499 [hep-th].
- [57] Juven Wang, "Ultra Unification," Phys. Rev. D 103, 105024 (2021), arXiv:2012.15860 [hep-th].
- [58] Juven Wang, Zheyan Wan, and Yi-Zhuang You, "Cobordism and deformation class of the standard model," Phys. Rev. D 106, L041701 (2022), arXiv:2112.14765 [hep-th].
- [59] Juven Wang, Zheyan Wan, and Yi-Zhuang You, "Proton stability: From the standard model to beyond grand unification," Phys. Rev. D 106, 025016 (2022), arXiv:2204.08393 [hep-ph].
- [60] Lukasz Fidkowski, Xie Chen, and Ashvin Vishwanath, "Non-Abelian Topological Order on the Surface of a 3D Topological Superconductor from an Exactly Solved Model," Phys. Rev. X 3, 041016 (2013), arXiv:1305.5851 [cond-mat.str-el].
- [61] C. Wang and T. Senthil, "Interacting fermionic topological insulators/superconductors in three dimensions," Phys. Rev. B 89, 195124 (2014), arXiv:1401.1142 [cond-mat.str-el].
- [62] Shiing-Shen Chern, "Characteristic classes of hermitian manifolds," Annals of Mathematics 47, 85–121 (1946).
- [63] L.S. Pontryagin, "Characteristic classes of differentiable manifolds," Mat. Sb. 21, 233–284 (1947).
- [64] J.W. Milnor and J. Stasheff, Characteristic Classes, by Milnor and Stasheff, Annals of Mathematics Studies, No. 76 (Princeton University Press, 1974).
- [65] T. Senthil, "Symmetry-Protected Topological Phases of Quantum Matter," Annual Review of Condensed Matter Physics 6, 299–324 (2015), arXiv:1405.4015 [cond-mat.str-el].

- [66] C. L. Kane and Matthew P. A. Fisher, "Quantized thermal transport in the fractional quantum Hall effect," Phys. Rev. B 55, 15832–15837 (1997), arXiv:cond-mat/9603118.
- [67] Juven Wang and Yi-Zhuang You, "Symmetric Mass Generation," Symmetry 14, 1475 (2022), arXiv:2204.14271 [cond-mat.str-el].
- [68] Xiao-Gang Wen, "A lattice non-perturbative definition of an SO(10) chiral gauge theory and its induced standard model," Chin. Phys. Lett. 30, 111101 (2013), arXiv:1305.1045 [hep-lat].
- [69] Yi-Zhuang You and Cenke Xu, "Interacting topological insulator and emergent grand unified theory," Phys. Rev. B **91**, 125147 (2015), arXiv:1412.4784.
- [70] Yoshio Kikukawa, "On the gauge invariant path-integral measure for the overlap Weyl fermions in <u>16</u> of SO(10)," PTEP 2019, 113B03 (2019), arXiv:1710.11618 [hep-lat].
- [71] Juven Wang and Xiao-Gang Wen, "A Non-Perturbative Definition of the Standard Models," Phys. Rev. Res. 2, 023356 (2020), arXiv:1809.11171 [hep-th].
- [72] Shlomo S. Razamat and David Tong, "Gapped Chiral Fermions," Phys. Rev. X 11, 011063 (2021), arXiv:2009.05037 [hep-th].
- [73] Juven Wang, "Strong CP Problem and Symmetric Interacting Mass Solution," (2022), arXiv:2212.14036 [hep-ph].
- [74] Juven Wang, "CT or P problem and symmetric gapped fermion solution," Phys. Rev. D 106, 125007 (2022), arXiv:2207.14813 [hep-th].
- [75] Gregory W. Moore and N. Read, "Nonabelions in the fractional quantum Hall effect," Nucl. Phys. B 360, 362–396 (1991).
- [76] Bogdan A. Dobrescu and Erich Poppitz, "Number of fermion generations derived from anomaly cancellation," Phys. Rev. Lett. 87, 031801 (2001), arXiv:hep-ph/0102010.
- [77] Zheyan Wan, Juven Wang, and et al., "Higher anomalies, higher symmetries, and cobordisms IV: Beyond Standard Models, Grand Unification, and Ultra Unification (202n)," ("202n"), arXiv:2xxx.xxxxx [hep-th].
- [78] A. N. Schellekens, "Meromorphic c = 24 conformal field theories," Commun. Math. Phys. 153, 159–186 (1993), arXiv:hep-th/9205072.
- [79] J. H. Conway and N. J. A. Sloane, Sphere Packings, Lattices and Groups (Springer, 1988).
- [80] Maryna Viazovska, "The sphere packing problem in dimension 8," arXiv e-prints , arXiv:1603.04246 (2016), arXiv:1603.04246 [math.NT].
- [81] Henry Cohn, Abhinav Kumar, Stephen D. Miller, Danylo Radchenko, and Maryna Viazovska, "The sphere packing problem in dimension 24," arXiv e-prints, arXiv:1603.06518 (2016), arXiv:1603.06518 [math.NT].