# On orbital stability of solitons for 2D Maxwell–Lorentz equations

A. I. Komech<sup>1</sup>

Institute of Mathematics, BOKU University, Vienna alexander.komech@gmail.com

E.A. Kopylova<sup>2</sup>

Faculty of Mathematics, Vienna University

#### Abstract

We prove the orbital stability of soliton solutions for 2D Maxwell–Lorentz system with extended charged particle. The solitons corresponds to the uniform motion and rotation of the particle.

We reduce the corresponding Hamilton system by the canonical transformation via transition to a comoving frame. The solitons are the critical points of the reduced Hamiltonian. The key point of the proof is a lower bound for the Hamiltonian.

 $<sup>^1\</sup>mathrm{Supported}$  partly by Austrian Science Fund (FWF) PAT 3476224

<sup>&</sup>lt;sup>2</sup>Supported partly by Austrian Science Fund (FWF) P34177

### 1 Introduction

We consider the 2D Maxwell–Lorentz equations with extended charged spinning particle

$$\begin{cases} \dot{E}(x,t) = J\nabla B(x,t) - [\dot{q}(t) - \omega(t)J(x - q(t))]\rho(x - q) \\ \dot{B}(x,t) = -\nabla \cdot JE(x,t), \quad \nabla \cdot E(x,t) = \rho(x - q(t)), \\ m\ddot{q}(t) = \langle E(x,t) + [J\dot{q}(t) + \omega(t)(x - q(t))]B(x,t), \rho(x - q(t))\rangle \\ I\dot{\omega}(t) = \langle (x - q(t)) \cdot [JE(x,t) - \dot{q}(t)B(x,t)], \rho(x - q(t))\rangle \end{cases} , \quad x \in \mathbb{R}^{2}, \quad t \in \mathbb{R}.$$
(1.1)

Here  $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ ,  $E(x,t) \in \mathbb{R}^2$  is the electric field,  $B(x,t) \in \mathbb{R}$  is the magnetic field,  $\rho(x-q(t))$  is the charge distribution of the extended particle centered at a point  $q(t) \in \mathbb{R}^2$ , m is the mass of the particle, I > 0 is its moment of inertia, and  $\omega(t) = \dot{\varphi}(t)$  is the angular velocity of the particle rotation. The brackets  $\langle , \rangle$  denote the inner product in the Hilbert space  $L^2 := L^2(\mathbb{R}^2) \otimes \mathbb{R}^2$ . We choose the units where the speed of light is c = 1.

Note that the 2D Maxwell–Lorentz equations can be introduced directly from 3D Maxwell– Lorentz system [18, (2.39)–(2.41)] by substituting  $x = (x_1, x_2, 0)$  and

$$E(x,t) = (E_1(x,t), E_2(x,t), 0), \quad B(x,t) = (0,0, B_3(x,t)), \quad q(t) = (q_1(t), q_2(t), 0).$$

In the main body of the paper, we focus on the equations with a *nonrelativistic particle*, and at the end, we comment on the corresponding modifications for the case of a *relativistic particle*.

We assume that the charge density  $\rho(x)$  is smooth and rotation-invariant, i.e.,

$$\rho \in C_0^{\infty}(\mathbb{R}^2), \qquad \rho(x) = \rho_1(|x|). \tag{1.2}$$

Moreover, we assume that the total charge equals zero (neutrality of the particle):

$$\int \rho(x)dx = 0. \tag{1.3}$$

The Maxwell potentials  $A(x,t) = (A_1(x,t), A_2(x,t)), \Phi(x,t)$  are introduced by

$$B(x,t) = \nabla \cdot JA, \qquad E(x,t) = -\dot{A}(x,t) - \nabla \Phi(x,t). \tag{1.4}$$

We choose the Coulomb gauge  $\nabla \cdot A(x,t) = 0$ . The second line of (1.1) implies that  $\Delta \Phi(x,t) = -\rho(x-q(t))$ . Hence,

$$\Phi(x,t) = \Phi_0(x - q(t)), \text{ where } \Phi_0(y) = -\frac{1}{2\pi} \int \log|z - y|\rho(z)dz.$$
 (1.5)

In the Maxwell potentials, the system (1.1) becomes

$$\begin{cases} \ddot{A}(x,t) = \Delta A(x,t) + \mathcal{P}\left([\dot{q}(t) - \dot{\varphi}(t)J(x-q(t))]\rho(x-q)\right) \\ m\ddot{q}(t) = \left\langle -\dot{A}(x,t) + [J\dot{q}(t) + \dot{\varphi}(t)(x-q(t))]\nabla \cdot JA(x,t), \rho(x-q(t))\right\rangle \\ I\ddot{\varphi}(t) = \left\langle (x-q(t)) \cdot [-J\dot{A}(x,t) - \dot{q}(t)\nabla \cdot JA(x,t)], \rho(x-q(t))\right\rangle \end{cases}$$
(1.6)

Here  $\mathcal{P}$  is the projection onto the space of solenoidal (divergence-free) vector fields. In the Fourier space,

$$\widehat{\mathcal{P}a}(k) = \hat{a}(k) - \frac{\hat{a}(k) \cdot k}{k^2}k.$$
(1.7)

The system is invariant under the action of the group of translations and rotations of the space  $\mathbb{R}^2$ . Accordingly, the system admits solitons of the form

$$S_{v,\omega}(t) = (A_{v,\omega}(x - vt), \dot{A}_{v,\omega}(x - vt), vt, v, \omega)$$

$$(1.8)$$

moving with constant speed v and rotating with constant angular velocity  $\omega$ .

The main result of present paper is the orbital stability of the solitons. We establish the Lagrangian and the Hamiltonian structures of the system (1.1), and prove the conservations of energy, linear and angular momenta.

The proof of the orbital stability of solitons relies on the reduction of the system (1.6) by the canonical transformation via transition to a comoving frame. The Hamiltonian of the reduced system depends on two parameters: the total linear momentum P and the angular momentum M, which are invariant in time.

The key point is the proof of the lower bound for the reduced Hamiltonian with the fixed P and M. The bound implies that the soliton is a strict global minimizer of the reduced Hamiltonian. We show that this fact implies the orbital stability. Moreover, the lower bound implies that the reduced system admits a unique soliton for every  $P, M \in \mathbb{R}^2$ .

Let us comment on related results.

Low-dimensional models are widely exploited as useful models in the classical and quantum field theories. For example, 't Hooft [4] established the confining for quarks in the meson theory in 1+1 space-time. The 2D version of the Maxwell electrodynamics was considered by Schwinger in [17], to show the possibility of the nonzero mass particles in the gauge theories. In [3], the Maxwell–Lorentz equations in 1D space were considered in the context of plasma dynamics, when the charge and currents correspond to a smooth distribution of the charged dust subject to the Lorentz dynamics. The well-posedness is proved and the point charge asymptotics is analyzed. The solitons of 2D Maxwell–Lorentz equations and their stability were not consider previously.

In [2], Bambusi and Galgani proved the existence and orbital stability of solitons with velocities |v| < 1 for the 3D Maxwell–Lorentz system without spinning. The proof relies on the transition to a comoving frame and a lower bound for the reduced Hamiltonian. The global convergence to solitons for the same system was proved in [5]. In [13], the global attraction to stationary states was established for a similar system with a relativistic particle in presence of an external confining

potential.

The results [5, 14, 13] provide the first rigorous proof of the radiation damping in classical electrodynamics. All these results were obtained under the *Wiener-type condition* on the charge density  $\rho$ . For the corresponding surveys, see [8] and [9].

The global convergence to rotating solitons was established in [6] for solutions of the 3D Maxwell–Lorentz system with  $q(t) \equiv 0$  in the case of sufficiently small charge density  $\rho$ . This result was strengthened in [14] under a considerably weaker Wiener-type condition.

In [7], the asymptotic completeness was proved for the scattering of solutions to the Maxwell– Lorentz equations without spinning. The result refers to solutions which are close to a solitary manifold.

In [10, 12] the orbital stability of solitons was studied for the 3D Maxwell–Lorentz system with moving and spinning particle. The solitons move with a velocity  $v \in \mathbb{R}^3$ , |v| < 1 and rotate with an angular velocity  $\omega \in \mathbb{R}^3$ . The results are obtained for solitons either with  $\omega = 0$  or with v = 0. The proofs rely on the Hamiltonian–Poisson structure of the equations which has been constructed in [11].

### 2 Well-posedness

Let us introduce a phase space for the system (1.6). Denote by  $\dot{H}^1$  the closure of  $C_0^{\infty}(\mathbb{R}^2) \otimes \mathbb{R}^2$ with the respect to the norm  $||A||_{\dot{H}^1} = ||\nabla A||_{L^2}$ , where  $L^2 := L^2(\mathbb{R}^3) \otimes \mathbb{R}^2$ . Let  $\mathcal{F}^0$ ,  $\dot{\mathcal{F}}^1$  be the subspaces of solenoidal vector fields:

$$\mathcal{F}^{0} := \{ A \in L^{2} : \operatorname{div} A(x) \equiv 0 \}, \qquad \dot{\mathcal{F}}^{1} := \{ A \in \dot{H}^{1} : \operatorname{div} A(x) \equiv 0 \}.$$
(2.1)

Denote

$$\mathcal{F} := \dot{\mathcal{F}}^1 \oplus \mathcal{F}^0, \quad \mathbb{Y} := \dot{\mathcal{F}}^1 \oplus \mathcal{F}^0 \oplus \mathbb{R}^2 \oplus \mathbb{R}^2 \oplus \mathbb{R}^2, \quad \mathbb{W} := L^2 \oplus H^{-1} \oplus \mathbb{R}^2 \oplus \mathbb{R}^2 \oplus \mathbb{R}^2.$$
(2.2)

**Proposition 2.1.** i) For any initial state  $Y(0) = (A(x,0), \dot{A}(x,0), q(0), \dot{q}(0), \varphi(0), \dot{\varphi}(0)) \in \mathbb{Y}$ , there exists a unique solution of the system (1.6),

$$Y(t) = (A(x,t), \dot{A}(x,t), q(t), \dot{q}(t), \varphi(t), \dot{\varphi}(t)) \in C(\mathbb{R}, \mathbb{Y}) \cap C^{1}(\mathbb{R}, \mathbb{W}).$$

$$(2.3)$$

ii) The map

$$W(t): Y(0) \mapsto Y(t) \tag{2.4}$$

is continuous in  $\mathbb{Y}$  for every  $t \in \mathbb{R}$ .

iii) For  $Y(0) \in \mathbb{Y}$ , the energy  $\mathcal{E}(t)$ , linear momentum  $P(t) = (P_1(t), P_2(t))$  and angular momentum

M(t) are conserved:

$$\mathcal{E}(t): = \frac{1}{2} \int [\dot{A}^2(x,t) + |\nabla A(x,t)|^2] dx + \frac{1}{2} m \dot{q}^2(t) + \frac{1}{2} I \dot{\varphi}^2(t) = \text{const},$$
(2.5)

$$P_{j}(t) := -\langle \dot{A}(x,t), \nabla_{j}A(x,t) \rangle + m\dot{q}_{j}(t) + \langle \rho(x-q(t)), A_{j}(x,t) \rangle = \text{const}, \ j = 1, 2, \quad (2.6)$$

$$M(t) := I\dot{\varphi}(t) - \left\langle [J(x-q(t))] \cdot A(x,t), \rho(x-q(t)) \right\rangle = \text{const}.$$

$$(2.7)$$

iv) Let

$$\begin{cases} A(x,0) \in C^{3}(\mathbb{R}^{2}) \otimes \mathbb{R}^{2}, & \dot{A}(x,0) \in C^{2}(\mathbb{R}^{2}) \otimes \mathbb{R}^{2} \\ A(x,0) = 0, & \dot{A}(x,0) + \nabla \Phi_{0}(x - q(0)) = 0, & |x| > R_{0} \end{cases}$$
(2.8)

for some  $R_0 > 0$ . Then

$$\begin{cases} A(x,t) \in C^2(\mathbb{R}^2 \times \mathbb{R}) \otimes \mathbb{R}^2, \\ |\partial_x^{\alpha} A(x,t)| + |\partial_x^{\alpha} \dot{A}(x,t)| \leq C_{\alpha} (1+|x|)^{-2-|\alpha|}, \quad |x| > R(t) + 1, \qquad \forall \alpha \end{cases},$$
(2.9)

where  $R(t) = R + |q(0)| + \overline{v}t + R_{\rho}$  with  $\overline{v} > 0$  defined by the relation  $\mathcal{E} = \frac{1}{2}m\overline{v}^2$ .

Proof. The proof is similar to that of Proposition A.5 in [13]. Namely, consider at first the case of smooth initial functions  $A(x,0), \dot{A}(x,0)$  with compact supports. The elimination of the fields reduces the system (1.6) to the system of nonlinear integral equations for q(t) and  $\omega(t)$ . Then the existence and uniqueness of the solution  $(q(t), \omega(t))$  for small |t| follows by the application of the contraction mapping principle. The corresponding fields  $A(\cdot, t), \dot{A}(\cdot, t)$  are smooth and have compact supports, hence, the energy conservation (2.5) follows by standard integration by parts. Therefore, the solution Y(t) can be extended to all  $t \in \mathbb{R}$ .

The continuity of the map W(t), as constructed on the dense subspace of  $\mathbb{Y}$ , follows from the continuity of the map  $\mathbb{Y} \to C^2([0,t]; \mathbb{R}^2 \oplus \mathbb{R}^2)$  defined as  $Y(0) \mapsto (q(\cdot), \omega(\cdot))|_{[0,t]}$ . Now i) and (2.5) follow for general initial state  $Y(0) \in \mathbb{Y}$  by suitable approximations of Y(0). The conservation laws (2.6), and (2.7) will be proved later.

It remains to prove iv). Note that  $q(\cdot) \in C^3(\mathbb{R}) \otimes \mathbb{R}^2$  and  $\omega(\cdot) \in C^2(\mathbb{R}) \otimes \mathbb{R}^2$  by (2.3) and the last two equations of (1.6). This fact and the first line of (2.8) imply the first line of (2.9) due to the Kirchhoff integral representation of solutions of the first equation of (1.6). Further, applying  $-J\nabla$  to both sides of the first equation in (1.4), and using  $\nabla \cdot A = 0$ , we obtain that

$$\Delta A(x,t) = -J\nabla B(x,t).$$

The second line of (2.8) implies that

$$E(x,0) = B(x,0) = 0, \qquad |x| > R.$$
 (2.10)

Moreover, (2.5) implies that  $|\dot{q}(t)| \leq \overline{v}, t \in \mathbb{R}$ . Hence,  $|q(t)| \leq |q(0)| + \overline{v}|t|$ .

Therefore, (2.10) implies that E(x,t) = B(x,t) = 0, for |x| > R(t) by the 2D analogue of the

$$A(x,t) = -\frac{1}{2\pi} \int_{|y| < R(t)} J\nabla B(y,t) \log |x-y| dy = -\frac{1}{2\pi} J\nabla \int_{|y| < R(t)} B(y,t) \log |x-y| dy.$$
(2.11)

Now the second line of (2.9) follows.

### 3 The Lagrangian structure

Here we show that, under assumption (1.2), sufficiently smooth solutions  $X(t) = (A(t), q(t), \omega(t))$ of the 2D system (1.6) satisfy the Hamilton least action principle. The Lagrangian L is well known (cf. [15, (28.6)]:

$$L(A, \dot{A}, q, \dot{q}, \varphi, \dot{\varphi}) = \frac{1}{2} \int [\dot{A}^2 - (\nabla A)^2] dx + \frac{m\dot{q}^2}{2} + \frac{I\dot{\varphi}^2}{2} + \left\langle [\dot{q} - \dot{\varphi}J(x-q)] \cdot A, \rho(x-q) \right\rangle$$
(3.1)

up to an additive constant depending on  $\Phi$ . This Lagrangian is well defined and Fréchet differentiable on the extended Hilbert phase space  $\mathbb{Y}$ .

**Proposition 3.1.** Under the assumption (1.2), the system (1.6) with  $\omega(t) = \dot{\varphi}(t)$  is equivalent to the Euler-Lagrange equations

$$\dot{\Pi} = D_A L, \qquad \dot{p} = D_q L, \qquad \dot{M} = D_{\varphi} L, \tag{3.2}$$

where the canonical conjugate momenta are defined by

$$\Pi := D_{\dot{A}}L, \qquad p := D_{\dot{q}}L, \qquad M := D_{\dot{\varphi}}L. \tag{3.3}$$

*Proof.* All the calculations of variations are justified by  $C^2$ -approximations, which exist by Proposition 2.1, iv).

Step i) Differentiating (3.1), and using (1.4), we find the momenta

$$\Pi = \dot{A}, \qquad p = m\dot{q} + \left\langle A(q+y), \rho(y) \right\rangle, \qquad M = I\dot{\varphi} - \left\langle [Jy] \cdot A(q+y), \rho(y) \right\rangle. \tag{3.4}$$

It remains to calculate the derivative  $D_A L$  on the right-hand side of the first equation (3.2). For any  $h \in \dot{\mathcal{F}}^1$ , we have

$$\frac{1}{2} \langle D_A \int (\nabla A(x))^2 dx, h \rangle = \frac{1}{2} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \int |\nabla [A(x) + \varepsilon h(x)]|^2 dx = \int \nabla A(x) \nabla h(x) dx = -\langle \Delta A, h \rangle.$$

Hence,

$$D_A L = \Delta A(q+y) + \mathcal{P}([\dot{q} - \dot{\varphi} Jy]\rho(y)).$$
(3.5)

Now the first equation of (3.2) coincides with the Maxwell equation (the first line of (1.6)).

Step ii) Now we check the Newton equation (the second equation of (1.6)). Differentiating (3.1)

in  $q_k$ , we obtain

$$D_{q_k}L = D_{q_k} \langle [\dot{q} - \dot{\varphi}Jy] \cdot A(q+y), \rho(y) \rangle = \langle [\dot{q} - \dot{\varphi}Jy] \cdot \nabla_k A(q+y), \rho(y) \rangle, \quad k = 1, 2.$$
(3.6)

Hence, the corresponding Euler–Lagrange equation reads as the system

$$m\ddot{q}_{k} + \left\langle \dot{A}_{k}(q+y,t),\rho(y)\right\rangle + \left\langle (\dot{q}(t)\cdot\nabla)A_{k}(q+y,t),\rho(y)\right\rangle = \left\langle [\dot{q}-\dot{\varphi}Jy]\cdot\nabla_{k}A(q+y,t),\rho(y)\right\rangle.$$
(3.7)

We should check that this system coincides with the second equation of (1.6) with  $\omega(t) = \dot{\varphi}(t)$ :

$$m\ddot{q}_{k}(t) = \left\langle -\dot{A}_{k}(q+y,t) + [(J\dot{q})_{k}(t) + \dot{\varphi}(t)y_{k}]\nabla \cdot JA(q+y,t), \rho(y) \right\rangle.$$
(3.8)

It remains to prove that

$$-\langle (\dot{q} \cdot \nabla)A_k, \rho \rangle + \langle [\dot{q} - \dot{\varphi}Jy] \cdot \nabla_k A, \rho \rangle = \langle [(J\dot{q})_k + \dot{\varphi}y_k] \nabla \cdot JA, \rho \rangle, \quad k = 1, 2.$$
(3.9)

It suffices to consider the case k = 1. Now (3.9) becomes

$$-\left\langle (\dot{q}_1 \nabla_1 + \dot{q}_2 \nabla_2) A_1, \rho \right\rangle + \left\langle [\dot{q}_1 - \dot{\varphi} y_2] \nabla_1 A_1, \rho \right\rangle + [\dot{q}_2 + \dot{\varphi} y_1] \nabla_1 A_2, \rho \right\rangle = \left\langle [\dot{q}_2 + \dot{\varphi} y_1] \nabla \cdot J A, \rho \right\rangle.$$

After cancellations, this identity is equivalent to

$$\dot{\varphi} \langle [-y_2 \nabla_1 + y_1 \nabla_2] A_1, \rho \rangle = \dot{\varphi} \langle A_1, [y_2 \nabla_1 - y_1 \nabla_2] \rho \rangle = 0.$$
(3.10)

This identity follows by the rotational symmetry (1.2).

Step *iii*) Finally, let us check that the last equation of (3.2) is equivalent to the last (torque) equation of the system (1.6). The right-hand side of the last equation of (3.2) vanishes since the Lagrangian (3.1) does not depend on  $\varphi$ . Now, the expression for M from (3.4) implies

$$\dot{M}(t) = I\ddot{\varphi}(t) - \left\langle [Jy] \cdot [\dot{A}(q+y,t) + (\dot{q} \cdot \nabla)A(q+y,t)], \rho(y) \right\rangle = 0.$$
(3.11)

So, the angular momentum is conserved:

$$M(t) = M, \qquad t \in \mathbb{R}. \tag{3.12}$$

To prove the equivalence of the last equations of (3.2) and (1.6), we should check that

$$-\langle (y_1\dot{q}_1 + y_2\dot{q}_2)(\nabla_1 A_2 - \nabla_2 A_1), \rho \rangle = \langle y_2[(\dot{q}_1 \nabla_1 + \dot{q}_2 \nabla_2)A_1] - y_1[(\dot{q}_1 \nabla_1 + \dot{q}_2 \nabla_2)A_2], \rho \rangle.$$
(3.13)

After cancellations, we arrive at

$$\dot{q}_2 \langle (-y_2 \nabla_1 + y_1 \nabla_2) A_2, \rho \rangle = \dot{q}_1 \langle (-y_1 \nabla_2 + y_2 \nabla_1) A_1, \rho \rangle.$$

Integrating by parts, and using the rotational symmetry (1.2), we conclude that both parts of the last equation vanish.

# 4 The conservation of linear momentum

The Lagrangian (3.1) is invariant under translations  $x \mapsto x + a$  and rotations  $R(\theta) : x_1 + ix_2 \mapsto e^{i\theta}(x_1 + ix_2)$  of the plane  $\mathbb{R}^2$ . The rotation symmetry implies the conservation (3.12) of the angular momentum M. Let us calculate the conserved linear momentum P. Consider the action of the group of translations

$$T_a: X = (A, q, \varphi) \mapsto T_a X = (A(x - a), q + a, \varphi), \qquad a \in \mathbb{R}^2.$$

The derivative of this map is

$$T'_a: \dot{X} = (\dot{A}, \dot{q}, \dot{\varphi}) \mapsto T'_a \dot{X} = (\dot{A}(x-a), \dot{q}, \dot{\varphi}).$$

It is easy to check that  $L(T_aX, T'_a\dot{X}) = L(X, \dot{X})$ . Hence, we can apply the Noether theorem on invariants [1, p. 88] to the one-parametric subgroups  $T_{\varepsilon a}$  with any  $a \in \mathbb{R}^2$  and  $\varepsilon \in \mathbb{R}$ . It suffices to consider  $C^2$ -approximations, which exist by Proposition 2.1. Thus, we obtain the conservation of the projection of the linear momentum

$$P(Y, \dot{Y}) \cdot a = D_{\dot{Y}}L \cdot \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} T_{\varepsilon a}Y = (\dot{A}, m\dot{q} + \langle \rho(x-q), A(x) \rangle, D_{\dot{\varphi}}L) \cdot (-a \cdot \nabla A, a, 0)$$
  
$$= -\langle \dot{A}, a \cdot \nabla A \rangle + m\dot{q} \cdot a + \langle \rho(x-q), A(x) \rangle \cdot a.$$
(4.1)

Therefore, the conserved linear momentum is the vector

$$P = -\langle \Pi, \nabla_* A \rangle + p = -\langle \dot{A}, \nabla_* A \rangle + m \dot{q} + \langle \rho(x-q), A(x) \rangle, \tag{4.2}$$

where the expression with  $\nabla_*$  is defined by

$$\langle \Pi, \nabla_* A \rangle_j = \langle \Pi, \nabla_j A \rangle, \quad j = 1, 2.$$
 (4.3)

### 5 The Hamiltonian structure

The Hamiltonian structure is obtained through the Legendre transformation [1, 16]

$$(\dot{A}, \dot{q}, \dot{\varphi}) \mapsto (\Pi, p, M), \quad H(A, q, \varphi, \Pi, p, M) = \langle \Pi, \dot{A} \rangle + p\dot{q} + M\dot{\varphi} - L.$$
 (5.1)

The formulas (3.4) give

$$\dot{A} = \Pi, \qquad \dot{q} = \frac{1}{m} (p - \left\langle A(q+y), \rho(y) \right\rangle), \qquad \dot{\varphi} = \frac{1}{I} (M + \left\langle [Jy] \cdot A(q+y), \rho(y) \right\rangle). \tag{5.2}$$

Hence, we obtain the Hamiltonian

$$H = \frac{1}{2} \int [\Pi^2 + |\nabla A|^2] dy + [m\dot{q} + \langle A(q+y), \rho(y) \rangle] \dot{q} + [I\dot{\varphi} - \langle [Jy] \cdot A(q+y), \rho(y) \rangle] \dot{\varphi} - \frac{m\dot{q}^2}{2} - \frac{I\dot{\varphi}^2}{2} - \langle [\dot{q} - \dot{\varphi}Jy] \cdot A(q+y), \rho(y) \rangle = \frac{1}{2} \int [\Pi^2 + |\nabla A|^2] dy + \frac{m\dot{q}^2}{2} + \frac{I\dot{\varphi}^2}{2}, \quad (5.3)$$

which coincides with (2.5). Here  $\frac{m\dot{q}^2}{2}$  and  $\frac{I\dot{\varphi}^2}{2}$  are *defined* according to (5.2):

$$\frac{m\dot{q}^2}{2} = \frac{1}{2m} (p - \left\langle A(q+y), \rho(y) \right\rangle)^2, \qquad \frac{I\dot{\varphi}^2}{2} = \frac{1}{2I} (M + \left\langle [Jy] \cdot A(q+y), \rho(y) \right\rangle)^2. \tag{5.4}$$

The Hamiltonian (5.3) is well defined and Fréchet differentiable on the Hilbert phase space  $\mathbb{Y}$ . The system (1.6) is equivalent to the canonical Hamiltonian system

$$\dot{A} = D_{\Pi}H, \ \dot{\Pi} = -D_AH; \ \dot{q} = D_pH, \ \dot{p} = -D_qH; \ \dot{\varphi} = D_MH, \ \dot{M} = -D_{\varphi}H.$$
 (5.5)

Now the energy conservation (2.5) obviously follows.

# 6 The canonical transformation to a comoving frame

Let us denote the fields in the "comoving frame" as

$$\mathbf{A}(y) = A(q+y), \qquad \mathbf{\Pi}(y) = \mathbf{\Pi}(q+y), \tag{6.1}$$

and define the map of the phase space  $\mathbb{Y} \oplus \mathbb{R}$ ,

$$T: (A, \Pi, q, p, \varphi, M) \mapsto (\mathbf{A}, \Pi, q, P, \varphi, M), \tag{6.2}$$

where P is defined by (4.2). Let us show that T is a canonical transformation, i.e. it preserves the canonical form ([1, 16]):

$$\langle \mathbf{\Pi}, \dot{\mathbf{A}} \rangle + P\dot{q} + M\dot{\varphi} = \langle \Pi, \dot{A} \rangle + p\dot{q} + M\dot{\varphi}.$$
 (6.3)

Indeed, differentiating in time  $\mathbf{A}(y,t) := A(q(t) + y, t)$ , we obtain

$$\dot{\mathbf{A}}(y,t) = \dot{A}(q+y,t) + \dot{q}(t) \cdot \nabla A(q+y,t).$$
(6.4)

Hence, the identity (6.3) reduces to  $\langle \Pi, \dot{q} \cdot \nabla A \rangle + P \cdot \dot{q} = p\dot{q}$ , which holds due to (4.2).

As a result, the canonical Hamiltonian equations (5.5) in the variables  $(A, \Pi, q, P, \varphi, M)$  with the Hamiltonian (5.3) are equivalent to the similar equations in the variables  $(\mathbf{A}, \mathbf{\Pi}, q, P, \varphi, M)$  with the Hamiltonian

$$\mathbf{H}(\mathbf{A}, \mathbf{\Pi}, P, M) = \frac{1}{2} \int [|\mathbf{\Pi}|^2 + |\nabla \mathbf{A}|^2] dy + \frac{1}{2m} (P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle)^2 + \frac{1}{2I} (M + \langle [Jy] \cdot \mathbf{A}(y), \rho(y) \rangle)^2, \qquad (6.5)$$

where we used the expressions (5.4), (4.2) and (3.12). The equivalence follows from the fact that the identity of the canonical forms (6.3) implies that the corresponding Lagrangians and the Lagrangian actions for the both systems coincide for the trajectories related by the canonical transformation T. Hence, the solutions of these systems are also related by this transformation since they are critical trajectories of the action. In other words, in the new variables, the Hamiltonian equations (5.5) read similarly:

$$\dot{\mathbf{A}} = D_{\mathbf{\Pi}}\mathbf{H}, \ \dot{\mathbf{\Pi}} = -D_{\mathbf{A}}\mathbf{H}; \ \dot{q} = D_{P}\mathbf{H}, \ \dot{P} = -D_{q}\mathbf{H}; \ \dot{\varphi} = D_{M}\mathbf{H}, \ \dot{M} = -D_{\varphi}\mathbf{H}.$$
 (6.6)

Now the key point is that the Hamiltonian (6.5) does not depend on q and  $\varphi$ . Hence, the equations for  $q, P, \varphi, M$  reduce to

$$\dot{q} = D_P \mathbf{H} = \frac{1}{m} (P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle), \quad \dot{P} = 0,$$
 (6.7)

$$\dot{\varphi} = D_M \mathbf{H} = \frac{1}{I} (M + \langle [Jy] \cdot \mathbf{A}(y), \rho(y) \rangle), \qquad \dot{M} = 0.$$
(6.8)

which also coincide with the formulas (5.2), (4.2) and (3.4) and correspond to the conservation of P and M. As the result, the system (6.6) reduces to the following family of the Hamiltonian systems with parameters  $P \in \mathbb{R}^2$  and  $M \in \mathbb{R}$ :

$$\dot{\mathbf{A}} = D_{\mathbf{\Pi}} \mathbf{H}_{P,M}, \quad \dot{\mathbf{\Pi}} = -D_{\mathbf{A}} \mathbf{H}_{P,M}, \tag{6.9}$$

where the "reduced Hamiltonian" is given by

$$\mathbf{H}_{P,M}(\mathbf{A},\mathbf{\Pi}) = \mathbf{H}(\mathbf{A},\mathbf{\Pi},P,M).$$
(6.10)

Expressing the variational derivatives, we write (6.9) as

$$\begin{cases} \dot{\mathbf{A}} = \mathbf{\Pi} + (\dot{q} \cdot \nabla) \mathbf{A} \\ \dot{\mathbf{\Pi}} = \Delta \mathbf{A} + (\dot{q} \cdot \nabla) \mathbf{\Pi} + \mathcal{P}[\dot{q}\rho(y) - \dot{\varphi}Jy\rho(y)] \end{cases}.$$
(6.11)

Here  $\dot{q}$  and  $\dot{\varphi}$  are given by (6.7) and (6.8). By the way, these formulas allow us to reconstruct the entire solution of the system (6.6) knowing the solution of the reduced system (6.11).

### 7 The solitons

The solitons (1.8) correspond to stationary solutions  $\mathbf{S}_{v,\omega} = (\mathbf{A}_{v,\omega}, \mathbf{\Pi}_{v,\omega})$  of the reduced system (6.11) with the parameters  $P_{v,\omega}$  and  $M_{v,\omega}$  defined by (6.7)–(6.8):

$$mv = P_{v,\omega} + \left\langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \right\rangle - \left\langle \mathbf{A}_{v,\omega}, \rho \right\rangle, \qquad I\omega = M_{v,\omega} + \left\langle [Jy] \cdot \mathbf{A}_{v,\omega}(y), \rho(y) \right\rangle. \tag{7.1}$$

The stationary equations corresponding to (6.11) read as

$$0 = \mathbf{\Pi}_{v,\omega} + (v \cdot \nabla)\mathbf{A}_{v,\omega}, \qquad 0 = \Delta \mathbf{A}_{v,\omega} + (v \cdot \nabla)\mathbf{\Pi}_{v,\omega} + \mathcal{P}[v\rho(y) - \omega J y \rho(y)].$$
(7.2)

This means that the solitons  $\mathbf{S}_{v,\omega}$  are critical points of the reduced Hamiltonian  $\mathbf{H}_{P_{v,\omega},M_{v,\omega}}$ . We calculate  $\mathbf{A}_{v,\omega}$  from the first equation of the system (1.6) with  $A(x,t) = \mathbf{A}_{v,\omega}(x-vt)$ :

$$\Delta \mathbf{A}_{v,\omega}(y) - (v \cdot \nabla)^2 \mathbf{A}_{v,\omega}(y) = -\mathcal{P}[v\rho(y)] + \omega J y \rho(y).$$
(7.3)

The equation (7.3) is easy to solve in the case |v| < 1: in the Fourier transform

$$\hat{\mathbf{A}}_{v,\omega}(k) = \frac{(v - \frac{k(v \cdot k)}{k^2} + \omega J i \nabla) \hat{\rho}(k)}{k^2 - (v \cdot k)^2}$$
(7.4)

by (1.7). The conditions (1.2) and (1.3) imply that

$$\mathbf{A}_{v,\omega} \in \dot{H}^1, \quad \mathbf{\Pi}_{v,\omega} = -(v \cdot \nabla) \mathbf{A}_{v,\omega} \in L^2.$$
(7.5)

Finally, equations (6.7) and (6.8) with q(t) = vt and  $\omega(t) = \omega$  hold according to (7.1). Thus, the solitons  $\mathbf{S}_{v,\omega} \in \dot{H}^1 \oplus L^2$  exist for  $(v,\omega) \in \Sigma = \{(v,\omega) \in \mathbb{R}^2 \times \mathbb{R} : |v| < 1\}$ .

The following lemma is proved in Appendix A. It implies that the solitons  $\mathbf{S}_{v,\omega}$  can be locally parametrised by their total momenta  $P_{v,\omega}$  and angular momenta  $M_{v,\omega}$ .

Lemma 7.1. The following Jacobian is nondegenerate:

$$J_{v,\omega} = \det \frac{\partial(P_{v,\omega}, M_{v,\omega})}{\partial(v,\omega)} \neq 0, \qquad (v,\omega) \in \Sigma.$$
(7.6)

# 8 The Lyapunov function

The stationary equations (7.2) mean that

$$D_{\mathbf{A}}\mathbf{H}_{P,M}(\mathbf{S}_{v,\omega}) = 0, \quad D_{\mathbf{\Pi}}\mathbf{H}_{P,M}(\mathbf{S}_{v,\omega}) = 0.$$
(8.1)

Hence, each soliton  $\mathbf{S}_{v,\omega}$  is a critical point of the reduced Hamiltonian  $\mathbf{H}_{P,M}$  with the corresponding linear momentum  $P = P_{v,\omega}$  and angular momentum  $M = M_{v,\omega}$ . Moreover, it turns out that, the soliton is a strict global minimizer of  $\mathbf{H}_{P,M}$ , which implies the orbital stability. Lemma 8.1. The lower bound holds,

$$\delta \mathbf{H}_{P,M} := \mathbf{H}_{P,M}(\mathbf{A}_{v,\omega} + \delta \mathbf{A}, \mathbf{\Pi}_{v,\omega} + \delta \mathbf{\Pi}) - \mathbf{H}_{P,M}(\mathbf{A}_{v,\omega}, \mathbf{\Pi}_{v,\omega})$$
  

$$\geq \frac{1 - |v|}{2} (\|\delta \mathbf{A}\|_{\dot{\mathcal{F}}^{1}}^{2} + \|\delta \mathbf{\Pi}\|_{\mathcal{F}^{0}}^{2}), \qquad (\delta \mathbf{A}, \delta \mathbf{\Pi}) \in \mathcal{F}.$$
(8.2)

*Proof.* Denote  $\delta \mathbf{A} = \mathbf{a}, \, \delta \mathbf{\Pi} = \boldsymbol{\pi}$ . Then  $\nabla \cdot \mathbf{a} = 0$ , and so we have

$$\delta \mathbf{H}_{P,M} = \frac{1}{2} \int (|\mathbf{\Pi}_{v,\omega} + \boldsymbol{\pi}|^2 + |\nabla(\mathbf{A}_{v,\omega} + \mathbf{a})|^2) \, dy - \frac{1}{2} \int (|\mathbf{\Pi}_{v,\omega}|^2 + |\nabla\mathbf{A}_{v,\omega}|^2) \, dy$$
$$+ \frac{1}{2m} (P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle)^2 - \frac{1}{2m} (P + \langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \rangle - \langle \mathbf{A}_{v,\omega}, \rho \rangle)^2$$
$$+ \frac{1}{2I} (M + \langle [Jy] \cdot \mathbf{A}, \rho \rangle)^2 - \frac{1}{2I} (M + \langle [Jy] \cdot \mathbf{A}_{v,\omega}, \rho \rangle)^2. \tag{8.3}$$

Evaluating, we obtain

$$\begin{split} \delta \mathbf{H}_{P,M} &= \frac{1}{2} \int (|\boldsymbol{\pi}|^2 + |\nabla \mathbf{a}|^2) \, dy + \int (\boldsymbol{\Pi}_{v,\omega} \cdot \boldsymbol{\pi} + \nabla \mathbf{A}_{v,\omega} \cdot \nabla \mathbf{a}) \, dy \\ &+ \frac{1}{2m} \big( P + \big\langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \big\rangle + \big\langle \boldsymbol{\pi}, \nabla_* \mathbf{A}_{v,\omega} \big\rangle + \big\langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{a} \big\rangle + \big\langle \boldsymbol{\pi}, \nabla_* \mathbf{a} \big\rangle - \big\langle \mathbf{A}_{v,\omega}, \rho \big\rangle - \big\langle \mathbf{a}, \rho \big\rangle \big)^2 \\ &- \frac{1}{2m} \big( P + \big\langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \big\rangle - \big\langle \mathbf{A}_{v,\omega}, \rho \big\rangle \big)^2 \\ &+ \frac{1}{2I} \big( M + \big\langle [Jy] \cdot [\mathbf{A}_{v,\omega} + \mathbf{a}], \rho \big\rangle \big)^2 - \frac{1}{2I} \big( M + \big\langle [Jy] \cdot \mathbf{A}_{v,\omega}, \rho \big\rangle \big)^2. \end{split}$$

After a rearrangement, we get

$$\delta \mathbf{H}_{P,M} = \frac{1}{2} \int (|\boldsymbol{\pi}|^2 + |\nabla \mathbf{a}|^2) \, dy + \int (\boldsymbol{\Pi}_{v,\omega} \cdot \boldsymbol{\pi} + \nabla \mathbf{A}_{v,\omega} \cdot \nabla \mathbf{a}) \, dy + \frac{1}{2m} [(\mathbf{p}_{v,\omega} + \delta \mathbf{p})^2 - \mathbf{p}_{v,\omega}^2] + \frac{1}{2I} [(M_{v,\omega} + \delta M)^2 - M_{v,\omega}^2], \tag{8.4}$$

where

$$\begin{cases} \mathbf{p}_{v,\omega} := P + \langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \rangle - \langle \mathbf{A}_{v,\omega}(y), \rho(y) \rangle = mv \\ \delta \mathbf{p} := \langle \pi, \nabla_* \mathbf{A}_{v,\omega} \rangle + \langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{a} \rangle + \langle \pi, \nabla_* \mathbf{a} \rangle - \langle \mathbf{a}, \rho \rangle \\ M_{v,\omega} := M + \langle [Jy] \cdot \mathbf{A}_{v,\omega}, \rho \rangle = I\omega \\ \delta M := \langle [Jy] \cdot \mathbf{a}, \rho \rangle \end{cases}$$
(8.5)

It remains to calculate the integral

$$\mathcal{I} := \int (\mathbf{\Pi}_{v,\omega} \cdot \boldsymbol{\pi} + \nabla \mathbf{A}_{v,\omega} \cdot \nabla \mathbf{a}) \, dy = \langle \mathbf{\Pi}_{v,\omega}, \boldsymbol{\pi} \rangle - \langle \Delta \mathbf{A}_{v,\omega}, \mathbf{a} \rangle.$$

By (7.2),

$$\mathcal{I} = -\langle (v \cdot \nabla) \mathbf{A}_{v,\omega}, \boldsymbol{\pi} \rangle + \langle (v \cdot \nabla) \mathbf{\Pi}_{v,\omega}, \mathbf{a} \rangle + \langle v\rho - \omega J y\rho, \mathbf{a} \rangle$$
(8.6)

since  $\nabla \cdot \mathbf{a} = 0$ . Integrating by parts in (8.6), we obtain

$$\mathcal{I} = -\langle (v \cdot \nabla) \mathbf{A}_{v,\omega}, \boldsymbol{\pi} \rangle - \langle \mathbf{\Pi}_{v,\omega}, (v \cdot \nabla) \mathbf{a} \rangle + \langle v\rho - \omega J y\rho, \mathbf{a} \rangle = -v \cdot \delta \mathbf{p} - \omega \delta M + \langle \boldsymbol{\pi}, (v \cdot \nabla) \mathbf{a} \rangle.$$
(8.7)

Substituting into (8.4), we get

$$\delta \mathbf{H}_{P,M} = \frac{1}{2} \int (|\boldsymbol{\pi}|^2 + |\nabla \mathbf{a}|^2) \, dy + \left\langle \boldsymbol{\pi}, (v \cdot \nabla) \mathbf{a} \right\rangle \\ + \frac{1}{2m} [(\mathbf{p}_{v,\omega} + \delta \mathbf{p})^2 - \mathbf{p}_{v,\omega}^2] - v \cdot \delta \mathbf{p} + \frac{1}{2I} [(M_{v,\omega} + \delta M)^2 - M_{v,\omega}^2] - \omega \delta M.$$
(8.8)

The last line can be written as

$$\frac{m}{2} \left( (v + \delta \mathbf{p}/m)^2 - v^2 - 2v \cdot \delta \mathbf{p}/m \right) + \frac{I}{2} \left( (\omega + \delta M/I)^2 - \omega^2 - 2\omega \delta M/I \right)$$
$$= \frac{m}{2} (\delta \mathbf{p}/m)^2 + \frac{I}{2} (\delta M/I)^2 \ge 0.$$
(8.9)

We have  $|\langle \boldsymbol{\pi}, (v \cdot \nabla) \mathbf{a} \rangle| \leq |v| |\boldsymbol{\pi}| |\nabla \mathbf{a}|$ . Hence, (8.8) and (8.9) imply that

$$\begin{split} \delta \mathbf{H}_{P,M} &\geq \frac{1}{2} \int (|\boldsymbol{\pi}|^2 + |\nabla \mathbf{a}|^2) \, dy + \left\langle \boldsymbol{\pi}, (v \cdot \nabla) \mathbf{a} \right\rangle \geq \frac{1}{2} \int (|\boldsymbol{\pi}|^2 - 2|v| |\boldsymbol{\pi}| |\nabla \mathbf{a}| + |\nabla \mathbf{a}|^2) dy \\ &= \frac{1 - |v|}{2} \int (|\boldsymbol{\pi}| + |\nabla \mathbf{a}|)^2 dy + \frac{|v|}{2} \int (|\boldsymbol{\pi} - \nabla \mathbf{a}||^2 dy \geq \frac{1 - |v|}{2} \int (|\boldsymbol{\pi}| + |\nabla \mathbf{a}|)^2 dy. \end{split}$$

# 9 The orbital stability of solitons

Introduce the distance between two solutions  $\mathbf{F}_1(y,t)$  and  $\mathbf{F}_2(y,t)$  as

$$d(\mathbf{F}_{1}(t), \mathbf{F}_{2}(t)) := \|\mathbf{F}_{1}(t) - \mathbf{F}_{2}(t)\|_{\mathcal{F}} + |\dot{q}_{1}(t) - \dot{q}_{2}(t)| + |\dot{\varphi}_{1}(t) - \dot{\varphi}_{2}(t)|, \quad t \in \mathbb{R},$$
(9.1)

where  $\dot{q}_j(t)$  and  $\dot{\varphi}_j(t)$  are defined by (6.7)–(6.8). Recall that the reduced soliton (1.8) does not depend on time:

$$\mathbf{F}_{v,\omega}(y,t) = \mathbf{S}_{v,\omega}(y) := (\mathbf{A}_{v,\omega}, \mathbf{\Pi}_{v,\omega}).$$
(9.2)

**Definition 9.1.** The soliton  $\mathbf{S}_{v,\omega}$  is orbitally stable if for any  $\varepsilon > 0$  there is a  $\delta > 0$  such that for any solution  $\mathbf{F}$  of the system (6.11), the inequality

$$d(\mathbf{F}(0), \mathbf{S}_{v,\omega}) < \delta \tag{9.3}$$

implies

$$d(\mathbf{F}(t), \mathbf{S}_{v,\omega}) < \varepsilon, \qquad t \in \mathbb{R}.$$
(9.4)

**Theorem 9.2.** Let |v| < 1 and let  $\rho(x)$  satisfy the conditions (1.2) and (1.3). Then the soliton (1.8) is orbitally stable.

14

*Proof.* For the proof we develop the general scheme of [2, 5]. Namely, by Lemma 8.1, the soliton  $\mathbf{S}_{v,\omega}$  is a nondegenerate global minimizer of the reduced Hamiltonian  $\mathbf{H}_{P,M}$ . The inequality (9.3) with sufficiently small  $\delta > 0$  implies that the linear and angular momenta  $P^*$  and  $M^*$  of the solution  $\mathbf{F}(t)$  are close to P and M, respectively:

$$|P^* - P| + |M^* - M| \to 0, \qquad \delta \to 0.$$
 (9.5)

By Lemma 7.1, for sufficiently small  $\delta > 0$ , there exists a unique soliton  $\mathbf{S}_{v_*,\omega_*}$  with the total momentum  $P^*$  and angular momentum  $M^*$ , which is close to  $\mathbf{S}_{v,\omega}$ :

$$\|\mathbf{S}_{v_*,\omega_*} - \mathbf{S}_{v,\omega}\|_{\mathcal{F}} \to 0, \qquad \delta \to 0.$$
(9.6)

Combining with (9.3), we get

$$\|\mathbf{F}(0) - \mathbf{S}_{v_*,\omega_*}\|_{\mathcal{F}} \to 0, \qquad \delta \to 0.$$
(9.7)

Therefore,

$$|\mathbf{H}_{P_*,M_*}(\mathbf{F}(0)) - \mathbf{H}_{P_*,M_*}(\mathbf{S}_{v_*,\omega_*})| \le \delta_1, \qquad \delta_1 \to 0 \text{ as } \delta \to 0.$$
(9.8)

Now the conservation of the reduced Hamiltonian  $\mathbf{H}_{P_*,M_*}$  implies that

$$|\mathbf{H}_{P_*,M_*}(\mathbf{F}(t)) - \mathbf{H}_{P_*,M_*}(\mathbf{S}_{v_*,\omega_*})| \le \delta_1, \quad t \in \mathbb{R}.$$
(9.9)

Therefore, the bound (8.2) with  $v, \omega$  replaced by  $v_*, \omega_*$ , allows us to conclude that

$$\sup_{t \in \mathbb{R}} \|\mathbf{F}(t) - \mathbf{S}_{v_{*},\omega_{*}}\|_{\mathcal{F}} \to 0 \text{ as } \delta \to 0.$$
(9.10)

Finally, combined with (9.6), this inequality gives

$$\sup_{t \in \mathbb{R}} \|\mathbf{F}(t) - \mathbf{S}_{v,\omega}\|_{\mathcal{F}} \to 0 \text{ as } \delta \to 0.$$
(9.11)

It remains to show that the terms with  $\dot{q}$  and  $\dot{\varphi}$  in (9.1) are small uniformly in time:

$$|\dot{q}(t) - v| + |\dot{\varphi}(t) - \omega| \le |\dot{q}(t) - v_*| + |\dot{\varphi}(t) - \omega_*| + |v_* - v| + |\omega_* - \omega|.$$
(9.12)

Indeed, the last two terms are small by (7.1) together with (9.5) and (9.6). Similarly, the smallness of the first two terms follows from (6.7)–(6.8) together with (9.5) and (9.10).

# 10 The case of a relativistic particle

In this section we comment on the required modifications of our arguments in the case of a relativistic particle with kinetic momentum

$$\mathbf{p} = m\dot{q}/\sqrt{1-\dot{q}^2}.$$

Now the third equation of the system (1.1) should be replaced by

$$\dot{\mathbf{p}}(t) = \left\langle E(x,t) + [J\dot{q}(t) + \omega(t)(x-q(t))]B(x,t), \rho(x-q(t)) \right\rangle.$$

Respectively, the second equation of the system (1.6) now reads

$$\dot{\mathbf{p}}(t) = \left\langle -\dot{A}(x,t) + [J\dot{q}(t) + \omega(t)(x - q(t))), \rho(x - q(t)]\nabla \cdot JA(x,t) \right\rangle.$$

The Lagrangian (3.1) is replaced by

$$L(A, \dot{A}, q, \dot{q}, \varphi, \dot{\varphi}) = \frac{1}{2} \int [\dot{A}^2 - (\nabla A)^2] dx - m\sqrt{1 - \dot{q}^2} + \frac{I\dot{\varphi}^2}{2} + \left\langle [\dot{q} - \dot{\varphi}J(x - q)] \cdot A, \rho(x - q) \right\rangle.$$

Accordingly, the momentum in (3.4) changes to

$$p = \mathbf{p} + \langle \mathbf{A}, \rho \rangle. \tag{10.1}$$

The equations (3.7) and (3.8) become

$$\dot{\mathbf{p}}_{k} + \left\langle \dot{A}_{k}(q+y,t),\rho(y)\right\rangle + \left\langle (\dot{q}(t)\cdot\nabla)A_{k}(q+y,t),\rho(y)\right\rangle = \left\langle [\dot{q}-\dot{\varphi}Jy]\cdot\nabla_{k}A(q+y,t),\rho(y)\right\rangle,\\ \dot{\mathbf{p}}_{k}(t) = \left\langle -\dot{A}_{k}(q+y,t) + [(J\dot{q})_{k}(t)+\dot{\varphi}(t)y_{k}]\nabla\cdot JA(q+y,t),\rho(y)\right\rangle, \quad k = 1, 2.$$

Now the linear momentum (4.2) keeps its form with p given by (10.1), and hence the formula for  $\dot{q}$  in (5.2) is replaced by

$$\dot{q} = \frac{\mathbf{p}}{\sqrt{m^2 + \mathbf{p}^2}} = \frac{p - \langle \mathbf{A}, \rho \rangle}{\sqrt{m^2 + (p - \langle \mathbf{A}, \rho \rangle)^2}} = \frac{P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle}{\sqrt{m^2 + (P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle)^2}}.$$
(10.2)

The Hamiltonian (5.3) becomes

$$H = \frac{1}{2} \int [\Pi_A^2 + |\nabla A|^2] dx + \frac{m}{\sqrt{1 - \dot{q}^2}} + \frac{I\dot{\varphi}^2}{2}.$$
 (10.3)

Here  $m/\sqrt{1-\dot{q}^2}$  is expressed via (10.2):

$$\frac{m}{\sqrt{1-\dot{q}^2}} = \sqrt{m^2 + \mathbf{p}^2} := \sqrt{m^2 + (P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle)^2}, \tag{10.4}$$

while  $I\dot{\varphi}^2/2$  is given by (5.2) as for the nonrelativistic particle.

The system (6.11) remains the same, but now  $\dot{q}$  is defined by (10.2), while  $\dot{\varphi}$  remains (6.8). As a result, equations (7.2) for the soliton fields are the same. The difference is that in the relativistic case the inequality |v| < 1 holds automatically.

Finally, Lemma 8.1 remains valid in the case of a relativistic particle as well. The changes in the proof are as follows. The second line of (8.3) is replaced by

$$\sqrt{m^2 + \mathbf{p}^2} - \sqrt{m^2 + \mathbf{p}_v^2} = \sqrt{m^2 + (\mathbf{p}_v + \delta \mathbf{p})^2} - \sqrt{m^2 + \mathbf{p}_v^2}$$

Here  $\mathbf{p}_v := mv/\sqrt{1-v^2}$  and

$$\delta \mathbf{p} := \mathbf{p} - \mathbf{p}_v = (P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle) - (P + \langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \rangle - \langle \mathbf{A}_{v,\omega}, \rho \rangle),$$

which coincides with the second line of (8.5). Hence, the formula (8.7) remains valid, and so (8.8) becomes

$$\delta \mathbf{H} = \frac{1}{2} \int (|\boldsymbol{\pi}|^2 + |\nabla \mathbf{a}|^2) d^3 x + \langle \boldsymbol{\pi}, d_v \mathbf{a} \rangle + \sqrt{m^2 + (\mathbf{p}_v + \delta \mathbf{p})^2} - \sqrt{m^2 + \mathbf{p}_v^2} - v \cdot \delta \mathbf{p} + \frac{I}{2} (\delta M/I)^2.$$
(10.5)

Here

$$\sqrt{m^2 + (\mathbf{p}_v + \delta \mathbf{p})^2} - \sqrt{m^2 + \mathbf{p}_v^2} - v \cdot \delta \mathbf{p} \ge 0$$
(10.6)

since the function  $E(\mathbf{p}) := \sqrt{m^2 + \mathbf{p}^2}$  is convex (its graph is the hyperbola  $E^2 - \mathbf{p}^2 = m^2$ ) and  $\nabla E(\mathbf{p}_v) = v$ . Hence, (8.2) follows.

Finally, Theorem 9.2 and its proof remain unchanged.

# A On alternative paramerisation of solitons

Here we prove Lemma 7.1. The Jacobian (7.6) reads

$$J_{\nu,\omega} = \det \begin{pmatrix} \frac{\partial P_1}{\partial v_1} & \frac{\partial P_2}{\partial v_1} & \frac{\partial M}{\partial v_1} \\ \frac{\partial P_1}{\partial v_2} & \frac{\partial P_2}{\partial v_2} & \frac{\partial M}{\partial v_2} \\ \frac{\partial P_1}{\partial \omega} & \frac{\partial P_2}{\partial \omega} & \frac{\partial M}{\partial \omega} \end{pmatrix}; \quad M = M_{\nu,\omega}; \quad P_j = (P_{\nu,\omega})_j, \quad j = 1, 2.$$
(A.7)

By (7.1),

$$\frac{\partial M}{\partial \omega} = I + \int \frac{|\nabla \hat{\rho}|^2 dk}{k^2 - (v \cdot k)^2} > 0, \quad \frac{\partial M}{\partial v_l} = 2\omega \int \frac{(v \cdot k)k_l |\nabla \hat{\rho}|^2}{(k^2 - (v \cdot k)^2)^2} dk, \quad l = 1, 2.$$
(A.8)

Further, we split  $\mathbf{A} = \mathbf{A}_{v,\omega}$  into even "+" and odd "-" components. In the Fourier transform,  $\hat{\mathbf{A}} = \hat{\mathbf{A}}^+ + \hat{\mathbf{A}}^-$ , where, by (7.4),

$$\hat{\mathbf{A}}^{+}(k) = \frac{(v - \frac{k(v \cdot k)}{k^2})\hat{\rho}(k)}{k^2 - (v \cdot k)^2}, \qquad \hat{\mathbf{A}}^{-}(k) = \frac{i\omega J \nabla \hat{\rho}(k)}{k^2 - (v \cdot k)^2}$$

Respectively, using (7.1), we split P as  $P = P^+ + P^-$ , where

$$P_j^- = \langle (v \cdot \nabla) \mathbf{A}^-, \nabla_j \mathbf{A}^- \rangle = \langle (v \cdot k) \hat{\mathbf{A}}^-, k_j \hat{\mathbf{A}}^- \rangle = \omega^2 \int \frac{(v \cdot k) k_j |\nabla \hat{\rho}|^2 dk}{(k^2 - (v \cdot k)^2)^2},$$

$$P_{j}^{+} = mv_{j} + \langle \mathbf{A}_{j}^{+}, \rho \rangle + \langle (v \cdot \nabla) \mathbf{A}^{+}, \nabla_{j} \mathbf{A}^{+} \rangle = mv_{j} + \langle \hat{\mathbf{A}}_{j}^{+}, \hat{\rho} \rangle + \langle (v \cdot k) \hat{\mathbf{A}}^{+}, k_{j} \hat{\mathbf{A}}^{+} \rangle$$
  
$$= mv_{j} + v_{j} \int \frac{|\hat{\rho}|^{2} dk}{k^{2} - (v \cdot k)^{2}} - \int \frac{(v \cdot k)k_{j}|\hat{\rho}|^{2} dk}{k^{2} (k^{2} - (v \cdot k)^{2})} + \int \left(v^{2} - \frac{(v \cdot k)^{2}}{k^{2}}\right) \frac{(v \cdot k)k_{j}|\hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{2}}.$$

Differentiation with respect to  $v_l$  and  $\omega$  gives

$$\frac{\partial P_{j}^{-}}{\partial v_{l}} = \omega^{2} \int \frac{k_{j}k_{l} |\nabla \hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{2}} + \omega^{2} \int \frac{4(v \cdot k)^{2}k_{j}k_{l} |\nabla \hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{3}}$$
(A.9)
$$\frac{\partial P_{j}^{+}}{\partial v_{l}} = \delta_{jl} \left( m + \int \frac{|\hat{\rho}|^{2} dk}{k^{2} - (v \cdot k)^{2}} \right) + v_{j} \int \frac{2(v \cdot k)k_{l} |\hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{2}} - \int \frac{k_{j}k_{l} |\hat{\rho}|^{2} dk}{k^{2}(k^{2} - (v \cdot k)^{2})} \\
- \int \frac{2k_{j}k_{l}(v \cdot k)^{2} |\hat{\rho}|^{2}}{k^{2}(k^{2} - (v \cdot k)^{2})^{2}} dk + \int \left( v^{2} - \frac{(v \cdot k)^{2}}{k^{2}} \right) \frac{k_{j}k_{l} |\hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{2}} \\
+ \int \left( v_{l} - \frac{(v \cdot k)k_{l}}{k^{2}} \right) \frac{2(v \cdot k)k_{j} |\hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{2}} + \int \left( v^{2} - \frac{(v \cdot k)^{2}}{k^{2}} \right) \frac{4(v \cdot k)^{2}k_{j}k_{l} |\hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{3}}, \quad (A.10)$$

$$\frac{\partial P_{j}}{\partial \omega} = \frac{\partial P_{j}^{-}}{\partial \omega} = 2\omega \int \frac{(v \cdot k)k_{j} |\nabla \hat{\rho}|^{2} dk}{(k^{2} - (v \cdot k)^{2})^{2}}.$$

We can assume that v = (|v|, 0). In this case  $\frac{\partial P_2}{\partial \omega} = 0$  and  $\frac{\partial P_j^+}{\partial v_l} = \frac{\partial P_j^-}{\partial v_l} = 0$  for  $j \neq l$ . Hence, (A.7) becomes

$$J_{\nu,\omega} = \frac{\partial P_2}{\partial v_2} \Big( \frac{\partial P_1}{\partial v_1} \frac{\partial M}{\partial \omega} - \frac{\partial M}{\partial v_1} \frac{\partial P_1}{\partial \omega} \Big).$$
(A.12)

Denoting  $D = k^2 - v^2 k_1^2$ , (A.9) and (A.10) gives

$$\frac{\partial P_j^-}{\partial v_j} = \omega^2 \int \frac{k_j^2 |\nabla \hat{\rho}|^2 dk}{D^2} + \omega^2 \int \frac{4v^2 k_1^2 k_j^2 |\nabla \hat{\rho}|^2 dk}{D^3} \ge 0, \quad \frac{\partial P_1}{\partial \omega} = 2\omega |v| \int \frac{k_1^2 |\nabla \hat{\rho}|^2 dk}{D^2}, \tag{A.13}$$

$$\begin{aligned} \frac{\partial P_1^+}{\partial v_1} &= m + \int \frac{|\hat{\rho}|^2 dk}{D} + \int \frac{2v^2 k_1^2 |\hat{\rho}|^2}{D^2} dk - \int \frac{k_1^2 |\hat{\rho}|^2}{k^2 D} dk - \int \frac{2v^2 k_1^4 |\hat{\rho}|^2}{k^2 D^2} dk \\ &+ \int \frac{v^2 k_1^2 k_2^2 |\hat{\rho}|^2}{k^2 D^2} + \int \frac{2v^2 k_1^2 k_2^2 |\hat{\rho}|^2}{k^2 D^2} + \int \frac{4v^4 k_1^4 k_2^2 |\hat{\rho}|^2}{k^2 D^3} \\ &= m + \int \frac{k_2^2 |\hat{\rho}|^2 dk}{k^2 D} + \int \frac{5v^2 k_1^2 k_2^2 |\hat{\rho}|^2}{k^2 D^2} + \int \frac{4v^4 k_1^4 k_2^2 |\hat{\rho}|^2}{k^2 D^3} > 0. \end{aligned}$$
(A.14)

$$\begin{aligned} \frac{\partial P_2^+}{\partial v_2} &= m + \int \frac{|\hat{\rho}|^2 dk}{D} - \int \frac{k_2^2 |\hat{\rho}|^2 dk}{k^2 D} - \int \frac{2v^2 k_2^2 k_1^2 |\hat{\rho}|^2 dk}{k^2 D^2} \\ &+ \int \frac{v^2 k_2^4 |\hat{\rho}|^2 dk}{k^2 D^2} - \int \frac{2v^2 k_1^2 k_2^2 |\hat{\rho}|^2 dk}{k^2 D^2} + \int \frac{4v^4 k_1^2 k_2^4 |\hat{\rho}|^2 dk}{k^2 D^3} \\ &= m + \int \frac{v^2 k_2^4 |\hat{\rho}|^2 dk}{k^2 D^2} + \int \frac{k_1^2 |\hat{\rho}|^2}{k^2 D} \left(1 - \frac{4v^2 k_2^2}{D} + \frac{4v^4 k_2^4}{D^2}\right) dk > 0 \end{aligned}$$
(A.15)

since  $1 - \frac{4v^2k_2^2}{D} + \frac{4v^4k_2^4}{D^2} = \left(1 - \frac{2v^2k_2^2}{D}\right)^2$ . From (A.13) and (A.15) it follows that

$$\frac{\partial P_2}{\partial v_2} = \frac{\partial P_2^+}{\partial v_2} + \frac{\partial P_2^-}{\partial v_2} > 0. \tag{A.16}$$

Moreover, (A.8) and (A.13)–(A.14) imply that

$$\frac{\partial P_1}{\partial v_1} \frac{\partial M}{\partial \omega} - \frac{\partial M}{\partial v_1} \frac{\partial P_1}{\partial \omega} = \frac{\partial P_1^+}{\partial v_1} \frac{\partial M}{\partial \omega} + \frac{\partial P_1^-}{\partial v_1} \frac{\partial M}{\partial \omega} - \frac{\partial M}{\partial v_1} \frac{\partial P_1}{\partial \omega} = \frac{\partial P_1^+}{\partial v_1} \frac{\partial M}{\partial \omega} + \omega^2 I \left( \int \frac{k_1^2 |\nabla \hat{\rho}|^2 dk}{D^2} + \int \frac{4v^2 k_1^4 |\nabla \hat{\rho}|^2 dk}{D^3} \right) + \omega^2 \int \frac{|\nabla \hat{\rho}|^2 dk}{D} \cdot \int \frac{k_1^2 |\nabla \hat{\rho}|^2 dk}{D^2} + 4\omega^2 v^2 \left( \int \frac{|\nabla \hat{\rho}|^2 dk}{D} \cdot \int \frac{k_1^4 |\nabla \hat{\rho}|^2 dk}{D^3} - \left( \int \frac{k_1^2 |\nabla \hat{\rho}|^2 dk}{D^2} \right)^2 \right) > 0$$
(A.17)

since  $\frac{\partial P_1^+}{\partial v_1} \frac{\partial M}{\partial \omega} > 0$  by (A.8) and (A.14), and  $\left(\int \frac{k_1^2 |\nabla \hat{\rho}|^2 dk}{D^2}\right)^2 \leq \int \frac{|\nabla \hat{\rho}|^2 dk}{D} \cdot \int \frac{k_1^4 |\nabla \hat{\rho}|^2 dk}{D^3}$  by the Cauchy-Schwarz inequality. As the result,  $J_{v,\omega} > 0$  by (A.12), (A.16) and (A.17).

### References

- [1] V. I. Arnold, Mathematical Methods of Classical Mechanics, Springer, New York, 1989.
- [2] D. Bambusi, L. Galgani, Some rigorous results on the Pauli–Fierz model of classical electrodynamics, Ann. de l'I.H.P., section A 58 (1993), no. 2, 155–171.
- [3] G. Hoermann, M. Kunzinger, Regularized derivatives in a 2-dimensional model of selfinteracting fields with singular data, *Zeitschr. Anal. Anw.* 19 (2000) 1, 147–158.
- [4] G. 't Hooft, A two-dimensional model for mesons, Nuclear Physics B75 (1974), 461–470.

- [5] V. Imaykin, A. Komech, N. Mauser, Soliton-type asymptotics for the coupled Maxwell–Lorentz equations, Ann. Inst. Poincaré, Phys. Theor. 5 (2004) 1117–1135.
- [6] V. Imaykin, A. Komech, H. Spohn, Rotating charge coupled to the Maxwell field: scattering theory and adiabatic limit, *Monatsh. Math.* 142 (2004), no. 1–2, 143–156.
- [7] V. Imaykin, A. Komech, H. Spohn, Scattering asymptotics for a charged particle coupled to the Maxwell field, J. Math. Phys. 52 (2011), no. 4, 042701.
- [8] A. Komech, E. Kopylova, Attractors of nonlinear Hamiltonian partial differential equations, Russ. Math. Surv. 75 (2020), no. 1, 1–87.
- [9] A. Komech, E. Kopylova, Attractors of Hamiltonian Nonlinear Partial Differential Equations, Cambridge University Press, Cambridge, 2022.
- [10] A. Komech, E. Kopylova, On the stability of solitons for Maxwell-Lorentz equations with rotating particle, *Milan Journal of Mathematics* **91** (2023), 155–173.
- [11] A. Komech, E. Kopylova, On the Hamilton–Poisson structure and solitons for the Maxwell– Lorentz equations with spinning particle, J. Math. Anal. Appl. 522 (2023), no. 2, 126976.
- [12] A. Komech, E. Kopylova, On orbital stability of solitons for 3D Maxwell–Lorentz equations with spinning particle, arXiv:2306.00508 [math-ph].
- [13] A.I. Komech, H. Spohn, Long-time asymptotics for the coupled Maxwell–Lorentz equations, Comm. Partial Differential Equations 25 (2000), 559–584.
- [14] M. Kunze, On the absence of radiationless motion for a rotating classical charge Advances in Mathematics 223, no. 5, 1632–1665.
- [15] L.D. Landau, E.M. Lifshitz, The classical theory of fields, Pergamon, 1975.
- [16] J.E. Marsden, T. Ratiu, Introduction to Mechanics and Symmetry, Springer, New York, 2002.
- [17] J. Schwinger, Gauge invariance of mass II, Phys. Rev. 128 (1962), no. 5, 2425–2429.
- [18] H. Spohn, Dynamics of Charged Particles and Their Radiation Field, Cambridge University Press, Cambridge, 2004.