# Hidden Conformal Symmetry in $AdS_2 \times S^2$ Beyond Tree Level

P. J. Heslop<sup>*a*</sup>, A. E. Lipstein<sup>*a*</sup>, and M. Santagata<sup>*b*</sup>

<sup>a</sup>Department of Mathematical Sciences, Durham University, Durham, DH1 3LE, UK

<sup>b</sup>Department of Physics, National Taiwan University, Taipei 10617, Taiwan

Correlators of a certain one-dimensional superconformal field theory dual to hypermultiplets in  $AdS_2 \times S^2$  exhibit a hidden four-dimensional conformal symmetry which allows one to repackage all tree-level 4-point correlators into a single four-dimensional object corresponding to a contact diagram arising from a massless  $\phi^4$  theory in  $AdS_2 \times S^2$ . This theory serves as a toy model for IIB string theory in  $AdS_5 \times S^5$  and is interesting in its own right because  $AdS_2 \times S^2$  describes the near-horizon limit of extremal black holes in four dimensions. We argue that after acting with an  $SU(1, 1) \times SU(2)$  Casimir, all one-loop correlators can similarly be encoded by a four-dimensional function which arises from a one-loop scalar bubble diagram in  $AdS_2 \times S^2$ , explaining how the hidden conformal symmetry extends beyond tree level. Finally, we conjecture a scalar effective field theory with a derivative interaction in  $AdS_2 \times S^2$  whose Witten diagrams should directly reproduce 4-point correlators to all loops without acting with Casimirs.

### I. INTRODUCTION

The celebrated AdS/CFT correspondence [1] relates quantum gravity in Anti-de Sitter (AdS) background to a non-gravitational conformal field theory (CFT) living in the boundary. The basic observables of a CFT are its correlation functions, which encode the scattering of gravitons and other degrees of freedom in AdS and can be computed perturbatively using Witten diagrams. For certain backgrounds of the form AdS×S, where S is a sphere, simple formulae packaging tree-level correlation functions of all the modes on the sphere into a single generating function have been obtained, originating from a hidden higher dimensional conformal symmetry. From the point of view of the dual CFT, these correspond to correlation functions of an infinite number of certain protected operators known as 1/2-BPS operators.

Hidden conformal symmetry was first discovered in 4point tree-level amplitudes of type IIB supergravity in  $AdS_5 \times S^5$  [2], which is dual to maximally supersymmetric Yang-Mills theory in four dimensions ( $\mathcal{N} = 4$  SYM) at strong coupling, and the symmetry has also been observed at weak coupling [3, 4].<sup>1</sup> If one includes higherderivative corrections to supergravity coming from string theory, the higher dimensional conformal symmetry gets broken but there are still remnants of a higher dimensional organising principle [6–9]. Indeed it was discovered that the four-point correlators of  $\mathcal{N} = 4$  SYM theory arise from massless scalar contact Witten diagrams on  $AdS_5 \times S^5$  [10].

This higher dimensional symmetry was also found in  $AdS_3 \times S^3$  [11, 12],  $AdS_5 \times S^3$  [13] and  $AdS_2 \times S^2$  [14] backgrounds. The last case is of particular interest because the geometry arises in the near-horizon limit of fourdimensional extremal black holes [15, 16] and may therefore have actual relevance for quantum gravity in the real world. The 4-point tree-level amplitudes of hypermultiplets in this background (which are dual to correlators of a 1d CFT with SU(1, 1|2) superconformal symmetry) can be encoded by a massless  $\phi^4$  contact Witten diagram in the bulk [14] very similarly to the IIB/  $\mathcal{N} = 4$  SYM case.<sup>2</sup> Hence the scalar theory in  $AdS_2 \times S^2$  can be thought of as a toy model for IIB string theory on  $AdS_5 \times S^5$ .

The purpose of this work is to go beyond the classical theory and investigate this at loop level. In particular, we consider four-point functions at one-loop order and show that they can be obtained by acting with Casimir operators on a 4d generating function. We then show that, quite remarkably, this generating function can be derived from a one-loop bubble diagram of the  $\phi^4$  theory. We observe that the infinite sum over modes of the 2-sphere which flow through each edge of a Feynman diagram can be resummed into a single bulk-to-bulk propagator in  $AdS_2 \times S^2$  which takes the same form as a propagator in four-dimensional flat space.

Finally, we conjecture a full effective field theory in  $AdS_2 \times S^2$  whose Feynman rules should directly compute correlators of the boundary CFT to all loop-orders, without acting with Casimirs. The key insight that leads us to this action is that acting with a boundary Casimir on a contact diagram is equivalent to acting with a Laplacian in the bulk. We verify this proposal for free-theory and tree-level correlators and write down a one-loop formula whose explicit evaluation we leave for future work.

## II. BASICS OF SU(1, 1|2) CORRELATORS

The purpose of this section is to review what is currently known about four-point functions of half-BPS operators in the SU(1,1|2) SCFT.<sup>3</sup> We will denote the 1/2-BPS operators by  $\mathcal{O}_p(x,y)$ , where x is the spacetime coordinate and y an analogous internal coordinate dealing with the SU(2) R-symmetry. These operators have protected dimension, and SU(2) charge  $p + \frac{1}{2}$  with  $p = 0, 1, \ldots$ 

 $\mathcal{O}_p$  is the primary operator of the superfield  $\mathcal{O}_p$ :

$$O_p = \mathcal{O}_p + \theta \mathcal{L}_p + \bar{\theta} \bar{\mathcal{L}}_p + \cdots \tag{1}$$

and so the descendant  $\mathcal{L}_p$  has dimension p+1 and SU(2) charge p. It is extremely useful to combine operators for

<sup>&</sup>lt;sup>1</sup> See [5] for a review on the subject.

 $<sup>^{2}</sup>$  Note that a straightforward generalisation of this describes four-

point open string scattering in IIB string theory on  $AdS_5 \times S^5 / \mathbb{Z}_2$ [17].

<sup>&</sup>lt;sup>3</sup> We follow the conventions of [14].

all p into single operators  $\mathcal{O}, \mathcal{L}, \bar{\mathcal{L}}$ :

$$\mathcal{O} = \sum_{p=0}^{\infty} \mathcal{O}_p, \qquad \mathcal{L} = \sum_{p=0}^{\infty} \mathcal{L}_p, \qquad \bar{\mathcal{L}} = \sum_{p=0}^{\infty} \bar{\mathcal{L}}_p. \quad (2)$$

Correlators of these objects will be referred to as "master correlators" and they act as generating functions in the sense that correlators with given external charges  $p_i$  are obtained simply by Taylor-expanding in the y variables and projecting onto the component with the correct conformal weight.

We will be interested in the four-point function of both the primaries and their descendants and we will expand all correlators in a large central charge expansion, e.g.

$$\langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}}\rangle = \sum_{m=0}^{\infty} \frac{1}{c^m} \langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}}\rangle^{(m)}, \qquad (3)$$

where the first term corresponds to free theory, the second to tree level amplitudes in the bulk, and so on. The goal of this work is to compute the one-loop contribution  $\langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}}\rangle^{(2)}$  and understand its structure.

Superconformal symmetry implies that the four-point function of primaries and descendants are closely related, taking the form [14]

$$\langle \mathcal{O}\mathcal{O}\mathcal{O}\mathcal{O} \rangle = \langle \mathcal{O}\mathcal{O}\mathcal{O}\mathcal{O} \rangle^{(0)} + I \times \mathbf{G}(x_i, y_i; c) , \langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}} \rangle = \langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}} \rangle^{(0)} + \mathcal{C}_{12}\mathbf{G}(x_i, y_i; c) .$$
(4)

Here I is a certain kinematic factor given in (A4), and  $C_{12}$  is the quadratic Casimir of  $SU(1,1) \times SU(2)$  arising from the superconformal SU(1,1|2) Casimir [14] applied at points 1 and 2:

$$C_{12} = C_{12}^{x} - C_{12}^{y},$$
  

$$C_{12}^{x} = \frac{1}{2} (D_{x_{1}} + D_{x_{2}})_{AB} (D_{x_{1}} + D_{x_{2}})^{AB},$$
(5)

where

$$D_{xAB} = x_A \frac{\partial}{\partial x^B} - x_B \frac{\partial}{\partial x^A} \tag{6}$$

is a generator of the conformal or SU(2) R-symmetry group.<sup>4</sup>

All the correlators are functions of  $x_i, y_i$ , but the hidden 4-dimensional symmetry means that the master correlators can be expressed in terms of functions of 4d distances, obtained by summing conformal and internal distances:

$$\mathbf{x}_{ij}^2 = x_{ij}^2 + y_{ij}^2 \,, \tag{7}$$

with the (boldface) 4d cross ratios defined accordingly:

$$\mathbf{u} = \frac{\mathbf{x}_{12}^2 \mathbf{x}_{34}^2}{\mathbf{x}_{13}^2 \mathbf{x}_{24}^2} = \mathbf{z}\overline{\mathbf{z}}, \qquad \mathbf{v} = \frac{\mathbf{x}_{14}^2 \mathbf{x}_{23}^2}{\mathbf{x}_{13}^2 \mathbf{x}_{24}^2} = (1 - \mathbf{z})(1 - \overline{\mathbf{z}}).$$
(8)

**Free theory.** The 4d conformal symmetry in free theory is realised at the level of correlators of descendants which can be written entirely in terms of 4d distances [14]

$$\langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}}\rangle^{(0)} = \frac{1}{\mathbf{x}_{14}^2\mathbf{x}_{23}^2} + \frac{1}{\mathbf{x}_{13}^2\mathbf{x}_{24}^2}.$$
 (9)

**Tree level.** At tree level on the other hand, the 4d conformal symmetry is directly realised on the correlators of primaries and we have

$$\mathbf{G}^{(1)} = -D_{1111}(\mathbf{x}_{ij}^2), \tag{10}$$

where  $D_{1111}$  is the (normalised) one-loop box diagram:

$$D_{1111}(\mathbf{x}_{ij}^{2}) = \frac{1}{\mathbf{x}_{13}^{2}\mathbf{x}_{24}^{2}} \frac{\phi^{(1)}(\mathbf{z}, \bar{\mathbf{z}})}{\mathbf{z} - \bar{\mathbf{z}}} = \frac{1}{\mathbf{x}_{13}^{2}\mathbf{x}_{24}^{2}} \frac{2\left(\operatorname{Li}_{2}(\mathbf{z}) - \operatorname{Li}_{2}(\bar{\mathbf{z}})\right) + \log \mathbf{u} \log\left(\frac{1-\mathbf{z}}{1-\bar{\mathbf{z}}}\right)}{\mathbf{z} - \bar{\mathbf{z}}}.$$
(11)

### III. ONE-LOOP CORRELATORS

Let us now consider one-loop correlators. We will focus on the trascendental weight 2 part of the one-loop correlators, which can be uniquely fixed by requiring consistency of the double-discontinuity with the OPE and crossing symmetry. Now, the double-discontinuity is simply related to the tree-level discontinuity via the action of the Casimir

$$\mathbf{G}^{(2)}\big|_{\log^2 u} = -\frac{1}{2} \mathcal{C}_{12}(D_{1111}\big|_{\log \mathbf{x}_{12}^2}), \qquad (12)$$

where

$$(D_{1111})|_{\log \mathbf{x}_{12}^2} = \frac{1}{\mathbf{x}_{13}^2 \mathbf{x}_{24}^2 (\mathbf{z} - \bar{\mathbf{z}})} \log \left(\frac{1 - \mathbf{z}}{1 - \bar{\mathbf{z}}}\right) , \quad (13)$$

and  $u = x_{12}^2 x_{34}^2 / (x_{13}^2 x_{24}^2)$  is the 1d cross-ratio.<sup>5</sup> The simplicity of this relation<sup>6</sup> is really a consequence of the fact that the leading log is controlled by tree-level CFT data, which, in theories with hidden conformal symmetries, depend on the higher dimensional spin [2, 14, 18–20] and in this case the latter is truncated to spin 0 only.<sup>7</sup>

The crossing completion of the above doublediscontinuity then automatically yields the complete result for the the maximal trascendental weight part of the one-loop correlator:

$$\mathbf{G}^{(2)} = -\frac{1}{2} \Big( \mathcal{C}_{12} \left( (D_{1111}) |_{\log \mathbf{x}_{12}^2} \right) \log u \log \frac{u}{v} \\ + \mathcal{C}_{23} \left( (D_{1111}) |_{\log \mathbf{x}_{23}^2} \right) \log v \log \frac{v}{u} \\ + \mathcal{C}_{13} \left( (D_{1111}) |_{\log \mathbf{x}_{13}^2} \right) \log u \log v \Big), \quad (14)$$

where  $C_{12}$  is given in (5) and  $C_{23}$  and  $C_{13}$  are its crossing versions. Note that the solution dictated by OPE and

 $<sup>^5</sup>$  Recall that in 1d the cross ratios u,v are not independent.

<sup>&</sup>lt;sup>6</sup> In fact, this relation directly generalises to all loop orders  $\mathbf{G}^{(l)}|_{\log^l u} = -\frac{1}{ll} \mathcal{C}_{12}^{l-1}(D_{1111}|_{\log \mathbf{x}_{12}^2})$ . An analogous statement was also recently noted in a similar context [21].

<sup>&</sup>lt;sup>4</sup> Note that we will mainly use 3-component embedding space coordinates  $x^A, y^A$  throughout the paper. For more details on our conventions see appendix A and equation (D5).

<sup>&</sup>lt;sup>7</sup> For more details, see e.g. section 3 of [22].

crossing symmetry carves out a three-dimensional space in the four-dimensional space of harmonic polylogarithms of weight 2, given by the three orientations of the logs. In particular, note the absence of  $Li_2$ .

A remarkable 4d uplift. Despite being quite simple, the one-loop formula presented above is still not fully satisfactory, because it is in a hybrid form: the function on which the Casimir acts is manifestly 4d but the logs are not. Instead, we wish to find a higher-dimensional function where the only piece breaking the conformal symmetry is the Casimir. In other words, we seek a function  $f_s(\mathbf{z}, \overline{\mathbf{z}})$  of four-dimensional distances *only* such that

$$\mathbf{G}^{(2)} = -\frac{1}{2} \Big( \mathcal{C}_{12} \frac{\mathfrak{f}_s(\mathbf{z}, \bar{\mathbf{z}})}{\mathbf{x}_{13}^2 \mathbf{x}_{24}^2} + \mathcal{C}_{23} \frac{\mathfrak{f}_t(\mathbf{z}, \bar{\mathbf{z}})}{\mathbf{x}_{13}^2 \mathbf{x}_{24}^2} + \mathcal{C}_{13} \frac{\mathfrak{f}_u(\mathbf{z}, \bar{\mathbf{z}})}{\mathbf{x}_{13}^2 \mathbf{x}_{24}^2} \Big),$$
(15)

where

$$\mathfrak{f}_t(\mathbf{z}, \overline{\mathbf{z}}) = \mathfrak{f}_s(1 - \mathbf{z}, 1 - \overline{\mathbf{z}}), \qquad \mathfrak{f}_u(\mathbf{z}, \overline{\mathbf{z}}) = \mathfrak{f}_s\left(\frac{1}{\mathbf{z}}, \frac{1}{\overline{\mathbf{z}}}\right).$$
(16)

In particular, we expect the function  $f_s(\mathbf{z}, \overline{\mathbf{z}})$  to transform correctly under  $1 \leftrightarrow 2$  crossing and to be a weight-3 combination of single-valued multiple polylogarithms (SVMPL's) [23].<sup>8</sup>

Quite remarkably, it turns out that such a generating function exists and is given by

$$f_s(\mathbf{z}, \overline{\mathbf{z}}) = \frac{1}{3} \frac{1}{\mathbf{z} - \overline{\mathbf{z}}} \left( f^{(3)}(\mathbf{z}, \overline{\mathbf{z}}) + \left( \log \mathbf{u} + \log \frac{\mathbf{u}}{\mathbf{v}} \right) \phi^{(1)}(\mathbf{z}, \overline{\mathbf{z}}) \right)$$
(17)

where  $f^{(3)}$  is a weight-3 single-valued antysimmetric function which, in AdS context, recently made its first appearance in [24]. In fact,  $f_s$  is the *unique* linear combination of SVMPL's built out of the alphabet  $\{\mathbf{z}, 1-\mathbf{z}, \mathbf{\bar{z}}, 1-\mathbf{\bar{z}}, \mathbf{z}-\mathbf{\bar{z}}\}$ such that it correctly reproduces the double-discontinuity and, crucially, its crossing versions  $f_t$ ,  $f_u$  do not contain  $\log^2 \mathbf{u}$  terms. We refer to appendix B for a more precise definition and a list of some useful properties.

#### IV. GENERALISED WITTEN DIAGRAMS

One of the consequences of the existence of a hidden conformal symmetry is that tree-level correlators can be obtained from a massless  $\phi^4$  contact Witten diagram in  $AdS_2 \times S^2$  [14]. The purpose of the remaining part of the paper is to test the applicability of such an approach beyond tree level. Before doing so, let us first recall how to obtain the tree level result from a  $\phi^4$  Witten diagram.

Points in AdS and S are parametrised in (d+2)dimensional embedding space as  $-\hat{x}\cdot\hat{x} = \hat{y}\cdot\hat{y} = 1$ , where we set the radii of AdS and S to one. Analogously, boundary coordinates, denoted by un-hatted letters, satisfy  $x \cdot x = y \cdot y = 0$ . **Tree level.** The following interaction term in the bulk effective action

$$S_{\rm int} = \frac{1}{4!} \int_{\rm AdS_2 \times S^2} d^2 \hat{x}_0 d^2 \hat{y}_0 \, \phi(\hat{x}_0, \hat{y}_0)^4 \qquad (18)$$

computes all tree-level correlators of the supersymmetric CFT using generalised Witten diagrams which treat AdS and S on equal footing. In particular all half-BPS correlators are generated by the following  $AdS_2 \times S^2$  integral, which is consistent with the bootstrapped result (10):

$$\mathbf{G}^{(1)} = -\frac{1}{\pi^2} \int \frac{d^2 \hat{x}_0 d^2 \hat{y}_0}{\mathbf{x}_{10}^2 \mathbf{x}_{20}^2 \mathbf{x}_{30}^2 \mathbf{x}_{40}^2} = -D_{1111}(\mathbf{x}_{ij}^2), \quad (19)$$

where  $\mathbf{x}_{i0}^2 = -2x_i \cdot \hat{x}_0 - 2y_i \cdot \hat{y}_0$  is the (inverse of the) AdS×S bulk-to-boundary propagator, which satisfies

$$\nabla^2 \frac{1}{\mathbf{x}_{i0}^2} = \left(\nabla_{\hat{x}_0}^2 + \nabla_{\hat{y}_0}^2\right) \frac{1}{\mathbf{x}_{i0}^2} = 0, \qquad (20)$$

where  $\nabla_{\hat{x}_0}^2$  ( $\nabla_{\hat{y}_0}^2$ ) is the AdS (S) Laplacian, which is given in equation (26) below.

**One loop.** In order to test whether the effective action extends beyond tree level, we first need to find an expression for the  $AdS_2 \times S^2$  bulk-to-bulk propagator. Intriguingly, it turns out that the bulk-to-bulk propagator in the  $AdS_{d+1} \times S^{d+1}$  product geometry takes exactly the same functional form as the bulk-to-boundary propagator:

$$\frac{1}{\hat{\mathbf{x}}_{lr}^{2d}} = \frac{1}{(-2\hat{x}_l \cdot \hat{x}_r - 2\hat{y}_l \cdot \hat{y}_r)^d} \ . \tag{21}$$

It turns out that in  $AdS \times S$  this simple propagator satisfies (20). This is very different from pure AdS space where bulk-to-bulk propagators are hypergeometric functions. Remarkably, this propagator also encodes an infinite sum over the Kaluza-Klein modes of the sphere propagating in AdS:

$$\frac{1}{\hat{\mathbf{x}}_{lr}^{2d}} = \frac{(-2)^{-d}}{(d-1)!} \sum_{p=0}^{\infty} \left[ (-1)^{\tilde{p}} (p-d+1)_{d-1} \right] \\
\times \frac{1}{(\hat{x}_l \cdot \hat{x}_r)^p} {}_2F_1 \left( \frac{p+1}{2}, \frac{p}{2}, p+1-\frac{d}{2}; \frac{1}{(\hat{x}_l \cdot \hat{x}_r)^2} \right) \\
\times \frac{1}{(\hat{y}_l \cdot \hat{y}_r)^{\tilde{p}}} {}_2F_1 \left( \frac{\tilde{p}+1}{2}, \frac{\tilde{p}}{2}, \tilde{p}+1-\frac{d}{2}; \frac{1}{(\hat{y}_l \cdot \hat{y}_r)^2} \right) \right].$$
(22)

With the bulk-to-bulk propagator at hand, we can then compute a one-loop bubble diagram in  $AdS_{d+1} \times S^{d+1}$ which in the s-channel reads (up to an unimportant overall constant)

$$\int_{2}^{1} \frac{d^{d+1}\hat{x}_{l}d^{d+1}\hat{x}_{r}d^{d+1}\hat{y}_{l}d^{d+1}\hat{x}_{r}}{\mathbf{x}_{1l}^{2d}\mathbf{x}_{2l}^{2d}\mathbf{x}_{3r}^{2d}\mathbf{x}_{4r}^{2d}\hat{\mathbf{x}}_{lr}^{4d}}, \quad (23)$$

with a similar formula for the t- and u-channels.

The double integral over the sphere can be evaluated combinatorially for arbitrary d and  $p_i$ , see appendix C for more details. Here, we will focus on  $d = 1, p_i = 0$ , for which the integral reduces to

$$\mathcal{K}_{s}|_{\substack{d=1\\y_{i}=0}} = \int \frac{d^{2}\hat{x}_{l}d^{2}\hat{x}_{r} \times 2^{-6}}{(x_{1}\cdot\hat{x}_{l})(x_{2}\cdot\hat{x}_{l})(x_{3}\cdot\hat{x}_{r})(x_{4}\cdot\hat{x}_{r})\big((\hat{x}_{l}\cdot\hat{x}_{r})^{2}-1\big)}.$$
(24)

<sup>&</sup>lt;sup>8</sup> In fact, this is the expected weight for one-loop amplitudes in d > 1 originating from a four-point contact interaction in the bulk [22, 24–27].

The remaining integration over AdS produces divergences and needs to be regularised. This can be done, for example, following the procedure outlined in [27]. In short, the idea is that the integral can be mapped to a flat-space integral so that one can take advantage of standard flat space techniques, such as dimensional regularisation, to isolate the divergence and extract the finite part. We refer to appendix D for details on the computation. Here we just quote the result for the finite part, which reads

$$\mathcal{K}_{s}^{(\text{fin})}\Big|_{\substack{d=1\\y_{i}=0}} \propto \frac{\mathfrak{f}_{s}(\mathbf{z}, \overline{\mathbf{z}})}{\mathbf{x}_{13}^{2}\mathbf{x}_{24}^{2}}\Big|_{y_{i}=0}$$
(25)

and analogously for t- and u-channels. Quite remarkably, this is precisely the function in equation (15) appearing in the master correlator of the SU(1, 1|2) CFT! The computation of the bubble diagram beyond  $y_i = 0$  is much more complicated and we leave it to the future but it is natural to expect that the relation (25) will extend beyond the  $y_i = 0$  case.

At this point, it is important to remark on the difference compared to tree-level: the bubble diagram at one loop reproduces the correlator only upon acting with a Casimir in each crossing orientation (15). This leads to a natural question: is there a way to explain the presence of the Casimir from a bulk perspective and make a more precise statement? We will address this in the next section.

### V. AN ALL-LOOPS CONJECTURE

The key observation we will make use of is that acting with a boundary Casimir on a pair or bulk-to-boundary propagators is equivalent to acting with the AdS×S Laplacian in the bulk. This is a generalisation of similar observations in AdS [28, 29]. The AdS and S Laplacians are equal (up to an important sign) to the SO(d, 2) or SO(d+2) Casimir operators acting on bulk coordinates:

$$\nabla_{\hat{x}}^2 = -\frac{1}{2} D_{\hat{x}AB} D_{\hat{x}}^{AB} \qquad \nabla_{\hat{y}}^2 = +\frac{1}{2} D_{\hat{y}AB} D_{\hat{y}}^{AB}, \quad (26)$$

where  $D_{\hat{x}AB}$  is a generator, identical in form to the conformal generator on the boundary (6). When the bulk Laplacian acts on a pair of bulk-to-boundary propagators meeting at common point in the bulk, it can be traded with a Casimir acting on the boundary points:

$$\nabla^2 \left( \frac{1}{\mathbf{x}_{10}^2 \mathbf{x}_{20}^2} \right) = -\mathcal{C}_{12} \left( \frac{1}{\mathbf{x}_{10}^2 \mathbf{x}_{20}^2} \right) \,, \tag{27}$$

where  $\nabla^2 = \nabla_{\hat{x}}^2 + \nabla_{\hat{y}}^2$  is the AdS×S Laplacian, and  $C_{12} = C_{12}^x - C_{12}^y$  is the AdS<sub>2</sub>×S<sup>2</sup> Casimir (5).

Now note that the master correlator of descendants  $\langle \mathcal{L}\mathcal{L}\mathcal{L}\mathcal{L}\mathcal{L}\rangle$  enjoys the following properties: in free theory the higher dimensional symmetry is manifest (9); at tree level it takes the form of the Casimir  $C_{12}$  acting on an object with higher dimensional symmetry (4, 10); at one loop it can be written as the Casimir  $C_{12}$  acting on further Casimirs which themselves act on functions with higher dimensional symmetry (4, 15). Along with the observation that external Casimirs can arise from a bulk Laplacian operator, this suggests the following effective action AdS×S space (including kinetic terms):

$$S' = \int_{\mathrm{AdS}_2 \times \mathrm{S}^2} d^4 \hat{\mathbf{x}}_0 \left( \frac{1}{2} \bar{\phi} \nabla^2 \phi - \frac{G_N}{4} \bar{\phi}^2 \nabla^2 \phi^2 \right), \qquad (28)$$

where we identify the *complex* field  $\phi$  as the bulk field coupled to the descendant operator  $\mathcal{L}$ . Here the Newton's constant is proportional to the inverse of the central charge,  $G_N \propto 1/c$  and we employed the notation  $d^4 \hat{\mathbf{x}}_0 \equiv d^2 \hat{x}_0 d^2 \hat{y}_0$ . Let us now work out the predictions for the boundary correlators arising from this action.

**Free theory.** By performing Wick contraction, the free-theory four-point function reads

$$\langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}}\rangle^{(0)} = \frac{1}{2} \underbrace{\overset{3}{\overbrace{\phantom{a}}}_{4}^{3} + \overset{1}{2} \underbrace{\overset{3}{\overbrace{\phantom{a}}}_{4}^{3}}_{4} = \frac{1}{\mathbf{x}_{13}^{2}\mathbf{x}_{24}^{2}} + \frac{1}{\mathbf{x}_{14}^{2}\mathbf{x}_{23}^{2}},$$

$$(29)$$

which indeed reproduces the result for the free theory correlator [14], quoted in (9).

**Tree level.** Following the standard AdS/CFT procedure, we can readily find the tree level four-point function, which reads:

$$\langle \mathcal{L}\mathcal{L}\bar{\mathcal{L}}\bar{\mathcal{L}}\bar{\mathcal{L}}\rangle^{(1)} = \frac{1}{2} \int d^{4}\hat{\mathbf{x}}_{0} \nabla^{2} \left(\frac{1}{\mathbf{x}_{10}^{2}\mathbf{x}_{20}^{2}}\right) \frac{1}{\mathbf{x}_{30}^{2}\mathbf{x}_{40}^{2}}$$
$$= \frac{-1}{\pi^{2}} C_{12} \int \frac{d^{4}\hat{\mathbf{x}}_{0}}{\mathbf{x}_{10}^{2}\mathbf{x}_{20}^{2}\mathbf{x}_{30}^{2}\mathbf{x}_{40}^{2}} = -C_{12}D_{1111}(\mathbf{x}_{ij}^{2}).$$
(30)

The arrows on the vertex distinguish the pairs of legs on which the Laplacian acts. This indeed reproduces the result for the master correlator (4, 10).

**One loop.** Finally, let us consider the one-loop amplitude. Here we have to consider three orientations of the bubble diagram

These yield the corresponding expressions

$$\int d^{4} \hat{\mathbf{x}}_{l} d^{4} \hat{\mathbf{x}}_{r} \nabla_{l}^{2} \left( \frac{1}{\mathbf{x}_{1l}^{2}} \frac{1}{\mathbf{x}_{2l}^{2}} \right) \frac{1}{\hat{\mathbf{x}}_{lr}^{4}} \nabla_{r}^{2} \left( \frac{1}{\mathbf{x}_{3r}^{2}} \frac{1}{\mathbf{x}_{4r}^{2}} \right) ,$$
$$\int d^{4} \hat{\mathbf{x}}_{l} d^{4} \hat{\mathbf{x}}_{r} \nabla_{l}^{2} \left( \frac{1}{\mathbf{x}_{1l}^{2}} \frac{1}{\hat{\mathbf{x}}_{lr}^{2}} \right) \frac{1}{\mathbf{x}_{3l}^{2} \mathbf{x}_{4r}^{2}} \nabla_{r}^{2} \left( \frac{1}{\mathbf{x}_{2r}^{2} \hat{\mathbf{x}}_{lr}^{2}} \right) , \quad (31)$$

with the *u*-channel expression obtained by swapping  $(x_3, y_3) \leftrightarrow (x_4, y_4)$  in the *t*-channel expression.

The first expression is directly related (via (27)) to the bubble integral evaluated in the previous section:

$$\int d^4 \hat{\mathbf{x}}_l d^4 \hat{\mathbf{x}}_r \, \nabla_l^2 \left( \frac{1}{\mathbf{x}_{1l}^2} \frac{1}{\mathbf{x}_{2l}^2} \right) \frac{1}{\hat{\mathbf{x}}_{lr}^4} \nabla_r^2 \left( \frac{1}{\mathbf{x}_{3r}^2} \frac{1}{\mathbf{x}_{4r}^2} \right)$$
$$= \mathcal{C}_{12}^2 \mathcal{K}_s|_{d=1}, \qquad (32)$$

and thus it correctly yields the *s*-channel part of the correlator (4, 15) (at least for the  $y_i = 0$  component). The evaluation of the integral in the other two channels is unfortunately less trivial and we will reserve it for future studies.

#### VI. CONCLUSION

The study of quantum gravity in curved background is both technically and conceptually very challenging, but great progress has been made in AdS background because in this case there is a dual description in terms of a conformal field theory in the boundary whose correlation functions are strongly constrained by conformal and crossing symmetry. In recent years a new symmetry principle has emerged for certain AdS×S backgrounds known as hidden conformal symmetry. In this paper, we explored the implications of this symmetry for 4-point correlators in AdS<sub>2</sub>×S<sup>2</sup> at the quantum level.

This paper has two main results: first, we found that the weight-2 transcendental sector at one loop can be organised into a 4d master correlator, which precisely coincides with the  $y_i = 0$  component of the one-loop bubble diagram of a massless  $\phi^4$  theory in  $AdS_2 \times S^2$ , upon acting with a Casimir in each orientation of the bubble. Moreover, we conjectured a scalar effective field theory in  $AdS_2 \times S^2$  which gives rise to the aforementioned Casimir via a two-derivative interaction in the bulk, and should produce all 4-point correlators to all loops.

In this paper we have only dealt with the maximal transcendental weight part of the correlators. It would be interesting to see whether the generating function (15)automatically incorporates the full structure dictated by the OPE. Moreover, it would very useful to test our conjectured action in (28) for correlators with arbitrary charges as well as for higher loops, and to understand how the structures we found in this paper generalise to higher-dimensional theories. For example, does a oneloop massless scalar bubble diagram in  $AdS_5 \times S^5$  have anything to do with correlators in  $\mathcal{N} = 4$  SYM? In fact, while a (Mellin) formula for arbitrary charges at order  $\lambda^{-\frac{3}{2}}$  is known [26], it is still not clear if and how this can be written in terms of a generating function, which will most likely involve the sixth order differential operator found in [14].

While the correlators we considered have superconformal symmetry, the bulk effective theories we constructed are non-supersymmetric so it is conceivable that they can be adapted to describe less idealised situations. Indeed, since  $AdS_2 \times S^2$  arises in the near horizon limit of extremal black holes, the approach developed in this paper may ultimately provide powerful new tools for the study of semirealistic black holes using effective field theory. Moreover, 4-point correlators in asymptotically AdS backgrounds are known to encode bound-states, quasi-normal modes, and gravitational radiation [30]. It would therefore be very exciting if these ideas can be combined with the ones developed in this paper to compute real-world gravitational wave observables.

### ACKNOWLEDGMENTS

We thank James Drummond, Hynek Paul, Ivo Sachs, and Pierre Vanhove for useful discussions. AL and PH are supported by an STFC Consolidated Grant ST/T000708/1. MS is supported by the Ministry of Science and Technology (MOST) through the grant 110-2112-M-002-006-.

## Appendix A: Internal coordinate conventions

The formulae in this paper are all written in terms of x, y coordinates in embedding space (either for the boundary or bulk spaces) where they have three components. The operators  $\mathcal{O}_p$  are however defined in terms of the 2-component  $y_a$ , homogeneous coordinates of the Riemann sphere  $\mathbb{CP}^1$ . The two are related as follows

$$y_a y_b = y_A (\sigma^A \epsilon)_{ab} . \tag{A1}$$

with  $\sigma^A$  Pauli matrices. Choosing coordinates  $y^a = (1, y)$  this gives

$$y^{A} = \left(\frac{1}{2}(y^{2}-1), -\frac{i}{2}(y^{2}+1), y\right)$$
(A2)

which satisfies  $y \cdot y = 0$ . Then we define the invariant combinations of two of these

$$y_{ij}^2 := (y_i - y_j) \cdot (y_i - y_j) = -2y_i \cdot y_j = (y_i - y_j)^2$$
 (A3)

where the last expression is given in terms of the 1d Poincare patch  $y_i$  coordinates but has exactly the same form as the second expression where the dot indicates the dot product in 3d embedding space.

The bulk  $\hat{y}$ 's (we use the terms "boundary" and "bulk" for the internal coordinates by analogy with AdS even though a sphere has no boundary) for the S<sup>2</sup> sphere are also given as 3-component vectors satisfying  $\hat{y} \cdot \hat{y} = 1$  (we assume a unit radius for the sphere and AdS space).

A completely analogous story applies to the above applies for the external  $x, \hat{x}$  coordinates where the dot product assumes the (-, -, +) metric and  $\hat{x} \cdot \hat{x} = -1$ . More details can be found in Appendix D.

In these conventions, the expression for the kinematic factor I in (4) is given as

$$I = x_{12}x_{34}y_{13}y_{24} - y_{12}y_{34}x_{13}x_{24} .$$
 (A4)

This is most straightforwardly viewed in the Poincare patch with  $y_{ij} := y_i - y_j$ ,  $x_{ij} := x_i - x_j$ , but could equally be viewed in embedding space with  $y_{ij} := \sqrt{(y_i - y_j) \cdot (y_i - y_j)}$ ,  $x_{ij} := \sqrt{(x_i - x_j) \cdot (x_i - x_j)}$ .

Note that the 4d invariants defining boundary-toboundary, bulk-to-boundary and bulk-to-bulk propagators are written (7)

$$\begin{aligned} \mathbf{x}_{ij}^{2} &:= (x_{i} - x_{j}) \cdot (x_{i} - x_{j}) + (y_{i} - y_{j}) \cdot (y_{i} - y_{j}) = -2x_{i} \cdot x_{j} - 2y_{i} \cdot y_{j} \\ \mathbf{x}_{i0}^{2} &:= (x_{i} - \hat{x}_{0}) \cdot (x_{i} - \hat{x}_{0}) + (y_{i} - \hat{y}_{0}) \cdot (y_{i} - \hat{y}_{0}) = -2x_{i} \cdot \hat{x}_{0} - 2y_{i} \cdot \hat{y}_{0} \\ \mathbf{\hat{x}}_{lr}^{2} &:= (\hat{x}_{l} - \hat{x}_{r}) \cdot (\hat{x}_{l} - \hat{x}_{r}) + (\hat{y}_{l} - \hat{y}_{r}) \cdot (\hat{y}_{l} - \hat{y}_{r}) = -2\hat{x}_{l} \cdot \hat{x}_{r} - 2\hat{y}_{l} \cdot \hat{y}_{r} \\ (A5) \end{aligned}$$

with the equality on the right due to the cancellations of the form  $\hat{x}\cdot\hat{x} + \hat{y}\cdot\hat{y} = -1 + 1 = 0$ .

# Appendix B: The functions $f^{(3)}$ and $\mathfrak{f}_s$

In this Appendix we will give more details about the weight-3 function appearing in the one-loop correlators.

The function  $f^{(3)}$  can be simply defined in terms of its total derivative:

$$df^{(3)} = \left(4\phi^{(1)} - \log^2 \mathbf{v} + 2\log \mathbf{u}\log \mathbf{v}\right) d\log \mathbf{z} \\ + \left(4\phi^{(1)} + \log^2 \mathbf{v} - 2\log \mathbf{u}\log \mathbf{v}\right) d\log \mathbf{\bar{z}} \\ + \left(4\phi^{(1)} + \log^2 \mathbf{u} - 2\log \mathbf{u}\log \mathbf{v}\right) d\log(1 - \mathbf{z}) \\ + \left(4\phi^{(1)} - \log^2 \mathbf{u} + 2\log \mathbf{u}\log \mathbf{v}\right) d\log(1 - \mathbf{\bar{z}}) \\ + \left(-12\phi^{(1)}\right) d\log(\mathbf{z} - \mathbf{\bar{z}}).$$
(B1)

The interested reader can find the full integrated expression – given in terms of SVMPL's – in [24].<sup>9</sup> Note that  $f^{(3)}$  satisfies

$$f^{(3)}(\mathbf{z}, \bar{\mathbf{z}}) = -f^{(3)}(1 - \mathbf{z}, 1 - \bar{\mathbf{z}}) = -f^{(3)}\left(\frac{1}{\mathbf{z}}, \frac{1}{\bar{\mathbf{z}}}\right).$$
(B2)

The function  $f_s$ , which we recall here for convenience,

$$f_s(\mathbf{z}, \bar{\mathbf{z}}) = \frac{1}{3} \frac{1}{\mathbf{z} - \bar{\mathbf{z}}} \left( f^{(3)} + \left( \log \mathbf{u} + \log \frac{\mathbf{u}}{\mathbf{v}} \right) \phi^{(1)} \right) \quad (B3)$$

is nothing but the "s-channel version" of  $f^{(3)}/(\mathbf{z}-\bar{\mathbf{z}})$ . In fact, it only preserves a  $1 \rightarrow 2$  crossing transformation:

$$\mathfrak{f}_s(\mathbf{z}, \bar{\mathbf{z}}) = \frac{1}{\mathbf{v}^2} \mathfrak{f}_s\left(\frac{\mathbf{z}}{1-\mathbf{z}}, \frac{\bar{\mathbf{z}}}{1-\bar{\mathbf{z}}}\right) \,, \tag{B4}$$

and we have

$$\mathfrak{f}_s(\mathbf{z}, \bar{\mathbf{z}}) + \mathfrak{f}_t(\mathbf{z}, \bar{\mathbf{z}}) + \frac{1}{\mathbf{z}\,\bar{\mathbf{z}}}\mathfrak{f}_u(\mathbf{z}, \bar{\mathbf{z}}) = \frac{1}{\mathbf{z} - \bar{\mathbf{z}}}f^{(3)}(\mathbf{z}, \bar{\mathbf{z}})\,, \quad (B5)$$

where  $f_t$ ,  $f_u$  are the crossing versions defined in (16).

Finally, note that  $f_s$  admits the log **u** stratification

$$f_s(\mathbf{z}, \bar{\mathbf{z}}) = \frac{1}{\mathbf{z} - \bar{\mathbf{z}}} \log \frac{1 - \mathbf{z}}{1 - \bar{\mathbf{z}}} \times \log \mathbf{u} \log \frac{\mathbf{u}}{\mathbf{v}} + \cdots$$
(B6)

where the dots do not contain  $\log^2 \mathbf{u}$ , which implies

$$\mathfrak{f}_{s}\Big|_{\log^{2}\mathbf{u}} = \frac{\phi^{(1)}}{\mathbf{z} - \bar{\mathbf{z}}}\Big|_{\log\mathbf{u}} = \frac{1}{\mathbf{z} - \bar{\mathbf{z}}}\log\frac{1 - \mathbf{z}}{1 - \bar{\mathbf{z}}}.$$
 (B7)

This, together with the property that

$$\mathfrak{f}_t(\mathbf{z}, \overline{\mathbf{z}})\Big|_{\log^2 \mathbf{u}} = \mathfrak{f}_u(\mathbf{z}, \overline{\mathbf{z}})\Big|_{\log^2 \mathbf{u}} = 0, \qquad (B8)$$

imply that the uplift (15) manifestly reproduces the trascendental weight 2 part of the 1d correlators given by (14).

The function  $f^{(3)}$  is not new in this context, and it appears in the position-space representation of correlators with a weakly-coupled dual AdS description [22, 24, 31–33]. In particular, this function appears whenever there is a logarithmic divergence in the associated flat-space amplitude. In fact, an important feature of this function is that it contains the letter  $\mathbf{z} - \bar{\mathbf{z}}$  and, as pointed out in [32] (see also [22]), the presence of this letter signals a logarithmic divergence  $\sim \log(\mathbf{z} - \bar{\mathbf{z}})$  in the bulk-point limit. This is indeed the case for a 4d one-loop flat-space amplitude, which the AdS<sub>2</sub>×S<sup>2</sup> correlator should approach to in the bulk-point limit.

### Appendix C: Sphere integration

The sphere integration of the one-loop bubble diagram (23) can be performed explicitly for all charges and for generic dimensions. This is far more general than we need but we give it in full generality as it may be useful in the future. We start by expanding all the propagators  $\mathbf{x}_{ij}^2 = -2x_i \cdot x_j - 2y_i \cdot y_j$  in the  $y, \hat{y}$  coordinates:

$$\mathcal{K}_{s} = \int d^{d+1}\hat{x}_{r} d^{d+1}\hat{x}_{l} \sum_{p_{i},k} I_{p_{i};k}(y_{i}) \times \frac{(2d)_{k}}{k!} \left(\prod_{i} \frac{(d)_{p_{i}}}{p_{i}!}\right) \frac{2^{-6d-2\Sigma_{p}}}{(x_{1}\cdot\hat{x}_{l})^{p_{1}+d}(x_{2}\cdot\hat{x}_{l})^{p_{2}+d}(\hat{x}_{l}\cdot\hat{x}_{r})^{2d+k}(x_{3}\cdot\hat{x}_{r})^{p_{3}+d}(x_{4}\cdot\hat{x}_{r})^{p_{4}+d}}$$
(C1)

where

$$I_{p_i;k}(y_i) = 2^{2\Sigma_p} \int d^{d+1} \hat{y}_l d^{d+1} \hat{y}_r (y_1 \cdot \hat{y}_l)^{p_1} (y_2 \cdot \hat{y}_l)^{p_2} (\hat{y}_l \cdot \hat{y}_r)^k \times (y_3 \cdot \hat{y}_r)^{p_3} (y_4 \cdot \hat{y}_r)^{p_4} \quad (C2)$$

and  $\Sigma_p = (p_1+p_2+p_3+p_4)/2$ . We will now do the  $\hat{y}_{l,r}$  integrations. First consider the three-point function corresponding to the integration over  $y_l$ 

$$J = \int d^{d+1} y_l (y_1 \cdot \hat{y}_l)^{p_1} (y_2 \cdot \hat{y}_l)^{p_2} (\hat{y}_l \cdot \hat{y}_r)^k$$
  
=  $y_1^{A_1} \dots y_1^{A_{p_1}} y_2^{B_1} \dots y_2^{B_{p_2}} \hat{y}_r^{E_1} \dots \hat{y}_r^{E_k} \int d^{d+1} \hat{y}_l \hat{y}_l^{A_1} \dots \hat{y}_l^{E_k}$ .  
(C3)

The integral of the product of  $2L y_l$ 's gives a factor  $\mathcal{N}_L$  times a product of symmetrised delta functions

$$\int d^{d+1}\hat{y}_l\hat{y}_l^{A_1}\dots\hat{y}_l^{A_{2L}} = \mathcal{N}_L\delta^{(A_1A_2}\dots\delta^{A_{2L-1}A_{2L})} \quad (C4)$$

and zero if there are an odd number of  $\hat{y}_l$ 's (see e.g. [34]) with

$$\mathcal{N}_L = \operatorname{vol}(S^{d+1}) \frac{(2L)!}{2^{2L} L! (d/2 + 1)_L} .$$
 (C5)

We need to consider then the product of  $(p_1 + p_2 + k)/2$ Kronecker-delta's with indices completely symmetrised:

$$\delta^{(A_1A_2}\dots\delta^{A_{p_1-1}A_{p_1}}\delta^{B_1B_2}\dots\delta^{B_{p_2-1}A_{p_2}}\delta^{E_1E_2}\dots\delta^{E_{k-1}E_k)}$$
(C6)

and count the proportion of terms which have the schematic form:

$$(\delta^{AA})^a (\delta^{BB})^b (\delta^{EE})^e (\delta^{AB})^\epsilon (\delta^{AE})^\alpha (\delta^{BE})^\beta \tag{C7}$$

with

$$p_1 = 2a + \alpha + \epsilon$$
  $p_2 = 2b + \beta + \epsilon$   $k = 2e + \alpha + \beta$ .  
(C8)

Counting these is equivalent to counting the number of inequivalent graphs of the form



<sup>&</sup>lt;sup>9</sup> The expression here differs by an overall factor of -2 compared to the definition given in [24].

where each line represents a number of parallel edges given by the associated label. The number of such inequivalent graphs is

$$\frac{p_1!p_2!k!}{a!b!e!\alpha!\beta!\epsilon!2^{a+b+e}} \tag{C9}$$

whereas the total number of inequivalent terms of the form (C6) is  $(p_1 + p_2 + k - 1)!!$ .

With this information we can obtain the integral J in (C3). Terms from (C8) with a > 0 or b > 0 produce  $y_1 \cdot y_1 = 0$  or  $y_2 \cdot y_2 = 0$  in (C3) thus only terms with a = b = 0 survive. Thus

$$J = \mathcal{N}_{(p_1+p_2+k)/2} \sum_{e} \left( (y_1.y_2)^{\epsilon} (y_1.\hat{y}_r)^{\alpha} (y_2.\hat{y}_r)^{\beta} \frac{p_1! p_2! k!}{e! \alpha! \beta! \epsilon! 2^e (p_1+p_2+k-1)!!} \right)$$
(C10)

with

$$\alpha = \frac{1}{2} (-2e + k + p_1 - p_2)$$
  

$$\beta = \frac{1}{2} (-2e + k - p_1 + p_2),$$
  

$$\epsilon = \frac{1}{2} (2e - k + p_1 + p_2).$$
 (C11)

Inserting this into (C2) then gives the exchange diagram as a sum of contact sphere diagrams  $B_{p_1p_2p_3p_4}(y_i)$  (similarly to what happens for AdS exchange Witten diagrams):

$$I_{p_i,k} = \mathcal{N}_{\frac{p_1+p_2+k}{2}} \sum_{e} \left( \frac{p_1!p_2!k!}{e!\alpha!\beta!\epsilon!(p_1+p_2+k-1)!!2^e} \times \frac{B_{\alpha\beta p_3 p_4}(y_1, y_2, y_3, y_4)}{2^{\alpha+\beta+p_3+p_4}} \right)$$
(C12)

where

$$B_{p_1p_2p_3p_4} = 2^{2\Sigma_p} \int d^{d+1} \hat{y}(\hat{y}.y_1)^{p_1} (\hat{y}.y_2)^{p_2} (\hat{y}.y_3)^{p_3} (\hat{y}.y_4)^{p_4}$$
(C13)

is the tree-level function which can be explicitly integrated giving:

$$B_{p_1p_2p_3p_4}(y_i) = \operatorname{vol}(S^{d+1}) \frac{p_1!p_2!p_3!p_4!(d/2)!(-1)^{\Sigma_p}}{((p_1+p_2+p_3+p_4+d)/2)!} \times \sum_{\{d_{ij}\}} \prod_{i < j} \frac{(y_{ij}^2)^{d_{ij}}}{d_{ij}!}.$$
(C14)

with

$$\sum_{i} d_{ij} = p_j , \quad d_{ii} = 0 , \quad d_{ij} = d_{ji} .$$
 (C15)

We can thus obtain an explicit expression for the exchange diagram I by inserting this into (C12). Since it is  $B_{\alpha\beta p_3 p_4}$  rather than  $B_{p_1 p_2 p_3 p_4}$  we replace  $d_{12} \rightarrow d_{12} - \epsilon$ in (C14). After some simplification and performing the sum over e we get

$$I_{p_i;k} = (-2)^{\sum_p + k} \sum_{\{d_{ij}\}} c_{p_i;k}(d_{ij}) \prod_{i < j} (y_{ij}^2)^{d_{ij}}$$
(C16)

with

$$c_{p_{i};k}(d_{ij}) = \frac{\operatorname{vol}(S^{d+1})^{2} p_{1}! p_{2}! p_{3}! p_{4}! (d!!)^{2}}{d_{12}! d_{13}! d_{14}! d_{23}! d_{24}! d_{34}! \left(\frac{d}{2} + \Sigma_{p}\right)!} \times \frac{k! \left(\frac{1}{2} \left(k - d_{13} - d_{14} - d_{23} - d_{24}\right) + 1\right)_{\Sigma_{p} + d/2}}{(d + k + p_{1} + p_{2})!! \left(d + k + p_{3} + p_{4}\right)!!}.$$
(C17)

Plugging everything in (C1) after some work one can do the sum over k explicitly. The result is given in terms of a single  ${}_{3}F_{2}$  hypergeometric function:

$$\mathcal{K}_{s} = \sum_{d_{ij}=0}^{\infty} \left( \prod_{i < j} \frac{(y_{ij}^{2})^{d_{ij}}}{d_{ij}!} \right) \frac{(\prod_{i}(d)_{p_{i}})(2d)_{d_{4}}(d/2)!^{2}}{2^{d_{12}+d_{34}+d_{4}-6d+\Sigma_{p}}} \times \int \frac{d^{d+1}\hat{x}_{l}d^{d+1}\hat{x}_{r}}{(x_{1}\cdot\hat{x}_{l})^{p_{1}+d}(x_{2}\cdot\hat{x}_{l})^{p_{2}+d}(\hat{x}_{l}\cdot\hat{x}_{r})^{2d+d_{4}}(x_{3}\cdot\hat{x}_{r})^{p_{3}+d}(x_{4}\cdot\hat{x}_{r})^{p_{4}+d}}}{(C18)}$$

where

$$\begin{aligned} a_1 &= \frac{2d + d_4}{2} \,, \\ a_2 &= a_1 + 1/2 \,, \\ a_3 &= \frac{2 + d + 2d_{12} + 2d_{34} + 2d_4}{2} \\ a_4 &= \frac{2 + d + 2d_{12} + 2d_4}{2} \,, \\ a_5 &= \frac{2 + d + 2d_{34} + 2d_4}{2} \,, \end{aligned}$$

and  $d_4 = d_{13} + d_{14} + d_{34} + d_{24}$ . Note that in the case of  $p_i = 0, d = 1$ , the hypergeometric is

$$\frac{1}{(\hat{x}_l \cdot \hat{x}_r)^2} {}_3F_2\left(1, \frac{3}{2}, \frac{3}{2}; \frac{3}{2}, \frac{3}{2}, \frac{3}{2}, \frac{1}{(\hat{x}_l \cdot \hat{x}_r)^2}\right) = \frac{1}{(\hat{x}_l \cdot \hat{x}_r)^2 - 1}$$

This gives the result quoted in the main text, see formula (24).

#### Appendix D: AdS Integral

In this appendix we give some more details on the AdS integration of the bubble diagram. We will restrict to d = 1, which is the theory we are interested in, and focus on the simplest case, i.e.  $p_i = 0$ . Nicely, plugging d = 1 in (C18) and projecting onto the  $y_i = 0$  component, the hypergeometric simplifies to a geometric function. In the end, we are left with the following double integral:

$$\mathcal{K}_{s}\Big|_{\substack{d=1\\y_{i}=0}} = \int \frac{d^{2}\hat{x}_{l}d^{2}\hat{x}_{r} \times 2^{-6}}{(x_{1}\cdot x_{l})(x_{2}\cdot x_{l})(x_{3}\cdot x_{r})(x_{4}\cdot x_{r})((\hat{x}_{l}\cdot \hat{x}_{r})^{2}-1)},$$
(D1)

and its crossing orientations.<sup>10</sup> This integral can be evaluated following the logic outlined in [27]. The idea is to turn the AdS integral into a flat-space integral and evaluate it by borrowing known flat-space techniques.

<sup>&</sup>lt;sup>10</sup> From now on, we will always consider  $\mathcal{K}_s$  evaluated at  $d = y_i + 1 = 1$  and hence drop the notation  $|_{d=1,y_i=0}$ .

To this extent, let us map euclidean AdS to the upper half plane:

$$\mathcal{C}_2 = \{ Z = (x, z), \, x \in \mathbb{R}, \, z > 0 \}.$$
 (D2)

The bulk coordinates are then parametrised as follows

$$\hat{x} = \left(\hat{x}^{(-1)}, \hat{x}^{(0)}, \hat{x}^{(1)}\right) =$$
$$= \frac{1}{\sqrt{2}z} \left(1 + \frac{1}{2}|Z|^2, \sqrt{2}x, 1 - \frac{1}{2}|Z|^2\right), \quad (D3)$$

with  $\hat{x} \cdot \hat{x} = -(\hat{x}^{(-1)})^2 + (\hat{x}^{(0)})^2 + (\hat{x}^{(1)})^2 = -1$ , and we have defined the euclidean distance,

$$|Z|^2 \equiv x^2 + z^2. \tag{D4}$$

Analogously, the boundary coordinates are parametrised in terms of the 1d coordinate  $x_i$  as

$$x_i^A = \left(x_i^{(-1)}, x_i^{(0)}, x_i^{(1)}\right)$$
$$= \frac{1}{\sqrt{2}} \left(1 + \frac{1}{2}x_i^2, \sqrt{2}x_i, 1 - \frac{1}{2}x_i^2\right), \qquad (D5)$$

with  $x_i \cdot x_i = 0$ .

In these coordinates, we have

$$\hat{x}_l \cdot \hat{x}_r = -1 - \frac{|Z_l - Z_r|^2}{2(p \cdot Z_l)(p \cdot Z_r)},$$
(D6)

$$x_i \cdot \hat{x}_l = -\frac{|Z_i - Z_l|^2}{2(p \cdot Z_l)}, \qquad Z_i = (x_i, 0), \quad (D7)$$

where we introduced a constant vector  $p \in \mathbb{R}^2$ :<sup>11</sup>

$$p \cdot Z_{l,r} = z_{l,r}, \qquad p = (0,1),$$
 (D8)

and we also embedded the boundary coordinates in  $\mathbb{R}^2$ so that:

$$|Z_l - Z_i|^2 = (x_l - x_i)^2 + z_l^2, \qquad Z_i = (x_i, 0).$$
 (D9)

Now, note that the effective propagator satisfies the following identity:

$$\frac{2}{(\hat{x}_{l}\cdot\hat{x}_{r})^{2}-1} = \left(\frac{1}{\hat{x}_{l}\cdot\hat{x}_{r}-1} - \frac{1}{\hat{x}_{l}\cdot\hat{x}_{r}+1}\right) = \\
= \frac{2z_{l}z_{r}}{(z_{l}-z_{r})^{2}+(x_{l}-x_{r})^{2}} - \frac{2z_{l}z_{r}}{(z_{l}+z_{r})^{2}+(x_{l}-x_{r})^{2}} \\
= \frac{2(p\cdot Z_{l})(p\cdot Z_{r})}{|Z_{l}-Z_{r}|^{2}} + \frac{2(p\cdot Z_{l})(p\cdot\sigma(Z_{r}))}{|Z_{l}-\sigma(Z_{r})|^{2}} \tag{D10}$$

where  $\sigma(Z)$  is the antipodal map

$$\sigma(x,z) := (x,-z). \tag{D11}$$

The trick is then to use the identity (D10) to split the integral and then rewrite it as a single integral with the

domain of integration extended to the full plane, so that it effectively becomes a flat-space integral:

$$\mathcal{K}_{s} = \frac{1}{2} \int_{\mathbb{R}^{4}} \frac{d^{2}Z_{l} d^{2}Z_{r}}{(p \cdot Z_{l})^{2} (p \cdot Z_{r})^{2}} \times \frac{(p \cdot Z_{l})^{3} (p \cdot Z_{r})^{3}}{|Z_{l} - Z_{r}|^{2} |Z_{l} - Z_{1}|^{2} |Z_{l} - Z_{2}|^{2} |Z_{r} - Z_{3}|^{2} |Z_{r} - Z_{4}|^{2}} .$$
 (D12)

The integral contains divergences and it needs to be regularised. A way to do this is to change the dimension of integration without changing the pre-factor coming from the metric [27]:

$$\frac{d^2 Z}{(p \cdot Z)^2} \quad \to \quad \frac{d^D Z}{(p \cdot Z)^2}, \qquad D = 2 - 2\epsilon. \tag{D13}$$

The regularised integral becomes

$$\mathcal{K}_{s}^{(\epsilon)} \equiv \frac{1}{2} \int_{\mathbb{R}^{2D}} \frac{d^{D} Z_{l} d^{D} Z_{r}}{(p \cdot Z_{l})^{2} (p \cdot Z_{r})^{2}} \mathcal{I}$$
(D14)

where the integrand  $\mathcal{I}$  reads

$$\mathcal{I} \equiv \frac{(p \cdot Z_l)^3 (p \cdot Z_r)^3}{|Z_l - Z_r|^2 |Z_l - Z_1|^2 |Z_l - Z_2|^2 |Z_r - Z_3|^2 |Z_r - Z_4|^2}.$$
(D15)

Here,  $D = 2 - 2\epsilon$  and the superscript ( $\epsilon$ ) emphasises that the integral has been regularised.

To simplify the evaluation of the integral, we will take advantage of conformal symmetry to recast it in terms of conformal cross ratios. In doing so, we need to be careful as the dimensional regularisation scheme we are using breaks conformal invariance, as a consequence of the fact that we changed the dimension without changing the measure factor. Let us first take care of the integrand and then of the measure. Let us shift every boundary point by  $Z_4$  and then invert every bulk point. The inversion only changes the bulk-to-boundary propagator which maps to:

$$\frac{p \cdot Z_l}{|Z_l - Z_i|^2} \to \frac{p \cdot Z'_l}{|Z'_l - Z_i|^2} = \frac{1}{|Z_i|^2} \frac{p \cdot Z_l}{|Z_l - \frac{Z_i}{|Z_i|^2}|^2}$$
(D16)

Then, we shift every bulk point by  $Z_{l,r} \rightarrow Z_{l,r} + \tilde{Z}_{14}$ where we defined  $\tilde{Z}_{ij} = \frac{Z_{ij}}{Z_{ij}^2}$ ,  $Z_{ij} = Z_i - Z_j$  and then rescale every bulk point by  $Z_{l,r} \rightarrow \left(\tilde{Z}_{34} - \tilde{Z}_{14}\right) Z_{l,r}^{12}$ Thus, after these transformations the integrand reads<sup>13</sup>

$$\mathcal{I} = \frac{1}{x_{13}^2 x_{24}^2} \frac{(p \cdot Z_l)^3 (p \cdot Z_r)^3}{|Z_l - Z_r|^2 |Z_l|^2} \times \frac{1}{|Z_l - \frac{\tilde{Z}_{24} - \tilde{Z}_{14}}{|\tilde{Z}_{34} - \tilde{Z}_{14}|^2} |^2 |Z_r - \frac{\tilde{Z}_{34} - \tilde{Z}_{14}}{|\tilde{Z}_{34} - \tilde{Z}_{14}|}|^2}.$$
(D17)

We can finally use the conformal group acting on the boundary to map  $Z_3$  to infinity,  $Z_4$  to the origin,  $Z_1 =$ 

<sup>&</sup>lt;sup>11</sup> Note that this is the scalar product in  $\mathbb{R}^2$  equipped with the euclidean metric.

 $<sup>^{12}</sup>$  Note that in the case of one-dimensional boundary we could also simplify  $\frac{Z_{ij}}{Z_{ij}^2} = \frac{1}{Z_{ij}}$ . It is however convenient to keep the dimension generic.  $^{ij}_{13}$  Note that  $Z^2_{ij}=x^2_{ij}.$ 

(-1,0) and parametrise  $Z_2$  with the conformal ratio x:<sup>14</sup>

$$Z_2 = \left(-\frac{1}{1-x}, 0\right), \qquad x = \frac{x_{12}x_{34}}{x_{13}x_{24}}.$$
 (D19)

This gives:

$$\mathcal{I} = \frac{1}{x_{13}^2 x_{24}^2} \frac{(p \cdot Z_l)^3 (p \cdot Z_r)^3}{|Z_l - Z_r|^2 |Z_l|^2} \frac{1}{|Z_l - m_x|^2} \frac{1}{|Z_r - m_1|^2},$$
(D20)

with  $m_1 = (1, 0), m_x = (x, 0).$ 

On the other hand, the measure, after all the above conformal transformations, maps to

$$\int_{\mathbb{R}^{2D}} \frac{d^{D} Z_{l} d^{D} Z_{r}}{(p \cdot Z_{l})^{2} (p \cdot Z_{r})^{2}} \rightarrow \\
\int_{\mathbb{R}^{2D}} \frac{d^{D} Z_{l} d^{D} Z_{r}}{(p \cdot Z_{l})^{2} (p \cdot Z_{r})^{2}} \frac{1}{|Z_{l} - m_{1}|^{2(D-2)} |Z_{r} - m_{1}|^{2(D-2)}}.$$
(D21)

Putting all together we arrive at the following integral

$$\mathcal{K}_{s}^{(\epsilon)} = \frac{1}{2x_{13}^{2}x_{24}^{2}} \int_{\mathbb{R}^{2D}} d^{D}Z_{l} d^{D}Z_{r} \\ \times \frac{(p \cdot Z_{l})(p \cdot Z_{r})|Z_{l} - m_{1}|^{4\epsilon}}{|Z_{l} - Z_{r}|^{2}|Z_{l}|^{2}|Z_{l} - m_{x}|^{2}|Z_{r} - m_{1}|^{2-4\epsilon}}.$$
 (D22)

This regularised integral can be analytically evaluated with standard flat-space techniques. In short, one first introduces Feynman parameters, which allow to perform the spacetime integration. Then, the remaining parametric integral can be computed using e.g. the HyperInt package [35].

Feynman parameters and analytic evaluation of the integral. Let us recall a few useful formulae for the evaluation of the AdS integral. We begin with the familiar Feynman parametrisation:

$$\frac{1}{A_1^{\alpha_1} \cdots A_n^{\alpha_n}} = \frac{\Gamma(\alpha_1 + \cdots + \alpha_n)}{\Gamma(\alpha_1) \cdots \Gamma(\alpha_n)} \int_0^1 \frac{\prod_{i=1}^n dl_i l_i^{\alpha_i - 1} \delta\left(1 - \sum_{i=1}^n l_i\right)}{(\sum_{i=1}^n l_i A_i) \sum_{i=1}^n \alpha_i}.$$
(D23)

Let us also recall the *one-loop master formula* (see e.g. [36])

$$\int \frac{d^D k}{i\pi^{\frac{D}{2}}} \frac{(-k^2)^a}{(-Uk^2+V)^b} = \frac{\Gamma(\frac{D}{2}+a)\Gamma(b-\frac{D}{2}-a)}{\Gamma(\frac{D}{2})\Gamma(b)} \frac{U^{-\frac{D}{2}-a}}{V^{b-\frac{D}{2}-a}},$$
(D24)

and a very useful AdS-variation of it:

1

$$\frac{1}{\pi^{d+1}} \int d^d z \, dz_0 \, d^d w \, dw_0 \frac{(w_0^2)^{a_1} (z_0^2)^{a_2}}{(c_1 w^2 + c_2 w_0^2 + c_3 z^2 + c_4 z_0^2 + c_5)^b} \\
= \frac{1}{\pi \Gamma(b)} \Gamma \left( a_1 + \frac{1}{2} \right) \Gamma \left( a_2 + \frac{1}{2} \right) \Gamma \left( b - d - 1 - a_1 - a_2 \right) \times \\
\frac{c_1^{-\frac{d}{2}} c_2^{-\frac{1}{2} - a_1} c_3^{-\frac{d}{2}} c_4^{-\frac{1}{2} - a_2}}{c_5^{b-d-1 - a_1 - a_2}}.$$
(D25)

$$Z_2 = \left(\frac{\mathbf{z} + \overline{\mathbf{z}} - 2}{2(1 - \mathbf{z})(1 - \overline{\mathbf{z}})}, \frac{\mathbf{z} - \overline{\mathbf{z}}}{2i(1 - \mathbf{z})(1 - \overline{\mathbf{z}})}, 0, \dots, 0\right)^T.$$
(D18)

This is all we need to turn the AdS integrals involved in our one-loop bubble diagram into integrals over Feynman parameters. Let us first rewrite the integrand using the Feynman parametrisation (D23):

$$\frac{(p \cdot Z_l)(p \cdot Z_r)}{|Z_r - m_1|^{2-4\epsilon} |Z_l|^2 |Z_l - m_1|^{-4\epsilon} |Z_l - m_x|^2 |Z_l - Z_r|^2} = \frac{\Gamma(4-4\epsilon)}{\Gamma(1-2\epsilon)\Gamma(-2\epsilon)} \int \left(\prod_{i=1}^5 dl_i\right) l_1^{-2\epsilon} l_3^{-1-2\epsilon} \delta(1-\sum_i l_i) \times \frac{(p \cdot Z_l)(p \cdot Z_r)}{(\beta_1 |Z_r - Z'|^2 + \beta_2 |Z_l - Z''|^2 + \beta_3)^{4-4\epsilon}},$$
(D26)

where Z', Z'' are two-component vectors,  $\beta_i$  are constants and they do not depend on  $Z_r$ :

$$\begin{split} \beta_1 &= l_1 + l_5 \,, \\ \beta_2 &= -\frac{(l_2 + l_3 + l_4)l_5 + (l_2 + l_3 + l_4 + l_5)l_1}{l_1 + l_5} \,, \\ \beta_3 &= \frac{l_1(l_2(l_3 + l_4x^2 + l_5) + l_3l_4(1 - x)^2 + l_4l_5(1 - x)^2)}{(l_1 + l_5)\beta_2} \,, \\ + \frac{l_5(l_2(l_3 + l_4x^2) + l_3l_4(1 - x)^2)}{(l_1 + l_5)\beta_2} \,, \\ Z' &= \frac{l_1m_1 + l_5Z_l}{l_1 + l_5} \,, \\ Z'' &= \frac{l_1(l_3m_1 + l_4m_x + l_5m_1) + l_5(l_3m_1 + l_4m_x)}{(l_1 + l_5)\beta_2} \,, \end{split}$$
(D27)

where we used the fact that  $||m_1|| = 1, ||m_x|| = x$ . Now, since this expression will be integrated in  $Z_r$ , we can shift  $Z_r \to Z_r + Z'$ . Under this shift, the numerator maps to<sup>15</sup>

$$(p \cdot Z_l)(p \cdot Z_r) \to (p \cdot Z_l)(p \cdot Z_r) + \frac{l_5}{l_1 + l_5} (p \cdot Z_l)^2.$$
 (D28)

The first integrand is odd in  $Z_R$ , thus it vanishes upon integration. We are thus left with the following integrand

$$\frac{\Gamma(4-4\epsilon)}{\Gamma(1-2\epsilon)\Gamma(-2\epsilon)} \int \prod_{i=1}^{5} dl_{i} \ l_{1}^{-2\epsilon} l_{3}^{-1-2\epsilon} \delta\left(1-\sum_{i} l_{i}\right) \times \frac{l_{5}}{l_{1}+l_{5}} \frac{(p \cdot Z_{l})^{2}}{(\beta_{1}|Z_{r}|^{2}+\beta_{2}|Z_{l}-Z''|^{2}+\beta_{3})^{4-4\epsilon}}.$$
(D29)

After performing the shift  $Z_l \to Z_l + Z''$ , the integrand takes the form (D25) with  $c_1 = c_2 = \beta_1, c_3 = c_4 = \beta_2, c_5 = \beta_3, a_1 = 0, a_2 = 1, b = 4 - 4\epsilon$  and  $d = 1 - 2\epsilon$ . The double spacetime integration in  $Z_l, Z_r$  thus gives

$$\begin{split} \mathcal{K}_{s}^{(\epsilon)} &= \frac{1}{2x_{13}^{2}x_{24}^{2}} \frac{\Gamma(4-4\epsilon)}{\Gamma(1-2\epsilon)\Gamma(-2\epsilon)} \int_{\mathbb{R}^{2D}} d^{2D}Z_{l} d^{2D}Z_{r} \times \\ &\int \prod_{i=1}^{5} dl_{i} \; \frac{l_{1}^{-2\epsilon} l_{3}^{-1-2\epsilon} l_{5} \delta\left(1-\sum_{i} l_{i}\right) (p \cdot Z_{l})^{2}}{(l_{1}+l_{5})(\beta_{1}|Z_{r}|^{2}+\beta_{2}|Z_{l}|^{2}+\beta_{3})^{4-4\epsilon}} \\ &= \frac{1}{4x_{13}^{2}x_{24}^{2}} \frac{\pi^{2-2\epsilon}}{\Gamma(-2\epsilon)} \int \prod_{i=1}^{5} dl_{i} \frac{l_{1}^{-2\epsilon} l_{3}^{-1-2\epsilon} l_{5}((l_{1}+l_{5})\beta_{2})^{-2+\epsilon}}{\beta_{3}^{1-2\epsilon}} \end{split}$$
(D30)

<sup>15</sup> Note that  $p \cdot m_x = p \cdot m_1 = 0.$ 

<sup>&</sup>lt;sup>14</sup> In d > 1,  $Z_2$  is parametrised by two cross ratios:

which can be evaluted with the HyperInt package [35]. In fact, this job has actually already been done in [27]: upon inspection, it turns out that the parametric integral is precisely equal to (the diagonal limit of) their expressions (C.4)-(C.7). The result can be written as

$$\mathcal{K}_{s}^{(\epsilon)} = \frac{\pi^{2-2\epsilon} e^{-2\gamma_{\epsilon}}}{16\mathbf{x}_{13}^{2}\mathbf{x}_{24}^{2}} \times \left( -\frac{2}{\epsilon} \bar{D}_{1111} + \mathfrak{f}_{s} - 2\log \mathbf{v} \bar{D}_{1111} + \mathcal{O}(\epsilon) \right) \Big|_{y_{i}=0}$$
(D31)

where  $\bar{D}_{1111} = \frac{\phi^{(1)}}{\mathbf{z}-\bar{\mathbf{z}}}$ , and similarly for the other orientations. Let us conclude with two comments. First, while the result is really a function of the 1d cross ratio x, we have highlighted the fact it has a natural interpretation as the  $y_i = 0$  component of a higher dimensional function.<sup>16</sup> Second, note that, because of the term  $\log \mathbf{v}\bar{D}_{1111}$ , the expression is not invariant under  $1 \rightarrow 2$  crossing. This had to be expected as the regularisation scheme breaks conformal invariance. However, as explained in [27], the non-crossing symmetric part is exactly the order  $\epsilon$  of the tree-level contact diagram in  $D = 2 - 4\epsilon$ and thus it can be removed upon renormalising the bare four-point coupling, yielding a finite, crossing-symmetric, and conformally-covariant function. In other words, the one-loop bubble diagram can be rewritten as

$$\mathcal{K}_{s}^{(\epsilon)} = - \left. \frac{\pi^{2-2\epsilon} e^{-2\gamma_{\epsilon}}}{16\mathbf{x}_{13}^{2}\mathbf{x}_{24}^{2}} \left( \frac{2}{\epsilon} \bar{D}_{1111}^{(\epsilon)} + \mathfrak{f}_{s} + \mathcal{O}(\epsilon) \right) \right|_{y_{i}=0}, \tag{D32}$$

where  $\bar{D}_{1111}^{(\epsilon)}$  is precisely the  $\epsilon$ -expansion of the four-point contact Witten diagram,

$$\bar{D}_{1111}^{(\epsilon)} = \bar{D}_{1111} - \epsilon \left( \mathfrak{f}_s + \log \mathbf{v} \bar{D}_{1111} \right) + \mathcal{O}(\epsilon^2) \,. \quad (D33)$$

- J. M. Maldacena, "The Large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. 2 (1998), 231-252 doi:10.4310/ATMP.1998.v2.n2.a1 [arxiv:hep-th/9711200 [hep-th]].
- [2] S. Caron-Huot and A. K. Trinh, "All tree-level correlators in AdS<sub>5</sub>×S<sub>5</sub> supergravity: hidden tendimensional conformal symmetry," JHEP 01 (2019), 196 doi:10.1007/JHEP01(2019)196 [arXiv:1809.09173 [hepth]].
- [3] S. Caron-Huot and F. Coronado, "Ten dimensional symmetry of  $\mathcal{N} = 4$  SYM correlators," JHEP **03** (2022), 151 doi:10.1007/JHEP03(2022)151 [arXiv:2106.03892 [hep-th]].
- [4] S. Caron-Huot, F. Coronado and B. Mühlmann, "Determinants in self-dual  $\mathcal{N} = 4$  SYM and twistor space," JHEP **08** (2023), 008 doi:10.1007/JHEP08(2023)008 [arXiv:2304.12341 [hep-th]].
- [5] P. Heslop, "The SAGEX Review on Scattering Amplitudes, Chapter 8: Half BPS correlators," J. Phys. A 55 (2022) no.44, 443009 doi:10.1088/1751-8121/ac8c71 [arXiv:2203.13019 [hep-th]].

- [6] J. M. Drummond, D. Nandan, H. Paul and K. S. Rigatos, "String corrections to AdS amplitudes and the doubletrace spectrum of  $\mathcal{N} = 4$  SYM," JHEP **12** (2019), 173 doi:10.1007/JHEP12(2019)173 [arXiv:1907.00992 [hepth]].
- J. M. Drummond, H. Paul and M. Santagata, "Boot-strapping string theory on AdS5×S5," Phys. Rev. D 108 (2023) no.2, 026020 doi:10.1103/PhysRevD.108.026020 [arXiv:2004.07282 [hep-th]].
- [8] F. Aprile and P. Vieira, "Large p explorations. From SUGRA to big STRINGS in Mellin space," JHEP 12 (2020), 206 doi:10.1007/JHEP12(2020)206 [arXiv:2007.09176 [hep-th]].
- [9] F. Aprile, J. M. Drummond, H. Paul and M. Santagata, "The Virasoro-Shapiro amplitude in  $AdS_5 \times S^5$  and level splitting of 10d conformal symmetry," JHEP **11** (2021), 109 doi:10.1007/JHEP11(2021)109 [arXiv:2012.12092 [hep-th]].
- [10] T. Abl, P. Heslop and A. E. Lipstein, "Towards the Virasoro-Shapiro amplitude in  $AdS_5 \times S^5$ ," JHEP **04** (2021), 237 doi:10.1007/JHEP04(2021)237 [arXiv:2012.12091 [hep-th]].
- [11] L. Rastelli, K. Roumpedakis and X. Zhou, " $AdS_3 \times S^3$ Tree-Level Correlators: Hidden Six-Dimensional Conformal Symmetry," JHEP **10** (2019), 140 doi:10.1007/JHEP10(2019)140 [arXiv:1905.11983 [hepth]].
- [12] S. Giusto, R. Russo, A. Tyukov and C. Wen, "The CFT<sub>6</sub> origin of all tree-level 4-point correlators in  $AdS_3 \times S^3$ ," Eur. Phys. J. C **80** (2020) no.8, 736 doi:10.1140/epjc/s10052-020-8300-4 [arXiv:2005.08560 [hep-th]].
- [13] L. F. Alday, C. Behan, P. Ferrero and X. Zhou, "Gluon Scattering in AdS from CFT," JHEP 06 (2021), 020 doi:10.1007/JHEP06(2021)020 [arXiv:2103.15830 [hepth]].
- [14] T. Abl, P. Heslop and A. E. Lipstein, "Higher-dimensional symmetry of  $AdS_2 \times S^2$  correlators," JHEP **03** (2022), 076 doi:10.1007/JHEP03(2022)076 [arXiv:2112.09597 [hep-th]].
- [15] B. Bertotti, "Uniform electromagnetic field in the theory of general relativity," Phys. Rev. 116 (1959), 1331 doi:10.1103/PhysRev.116.1331
- [16] I. Robinson, "A Solution of the Maxwell-Einstein Equations," Bull. Acad. Pol. Sci. Ser. Sci. Math. Astron. Phys. 7 (1959), 351-352
- [17] R. Glew and M. Santagata, "The Veneziano amplitude in  $AdS_5 \times S^3$  from an 8-dimensional effective action," JHEP **08** (2023), 010 doi:10.1007/JHEP08(2023)010 [arXiv:2305.01013 [hep-th]].
- [18] F. Aprile, J. Drummond, P. Heslop and H. Paul, "Double-trace spectrum of N = 4 supersymmetric yang-Mills theory at strong coupling," Phys. Rev. D **98** (2018) no.12, 126008 doi:10.1103/PhysRevD.98.126008 [arXiv:1802.06889 [hep-th]].
- [19] F. Aprile and M. Santagata, "Two particle spectrum of tensor multiplets coupled to AdS3×S3 gravity," Phys. Rev. D 104 (2021) no.12, 126022 doi:10.1103/PhysRevD.104.126022 [arXiv:2104.00036 [hep-th]].
- [20] J. M. Drummond, R. Glew and M. Santagata, "Bern-Carrasco-Johansson relations in AdS5×S3 and the double-trace spectrum of super gluons," Phys. Rev. D 107 (2023) no.8, L081901 doi:10.1103/PhysRevD.107.L081901 [arXiv:2202.09837 [hep-th]].
- [21] P. Ferrero and C. Meneghelli, "Unmixing the Wilson line defect CFT. Part II: analytic bootstrap," [arXiv:2312.12551 [hep-th]].
- [22] H. Paul and M. Santagata, "Genus-one open string amplitudes on  $AdS_5 \times S^3$  from CFT," JHEP **12** (2023), 057 doi:10.1007/JHEP12(2023)057 [arXiv:2309.15506 [hep-

<sup>&</sup>lt;sup>16</sup> This expression is obtained by simply parametrising  $Z_2$  with the two cross ratios  $\mathbf{z}, \bar{\mathbf{z}}$ , as in equation (D18).

th]].

- [23] O. Schnetz, "Generalized single-valued hyperlogarithms," [arXiv:2111.11246 [math-ph]].
- [24] J. M. Drummond and H. Paul, "One-loop string corrections to AdS amplitudes from CFT," JHEP 03 (2021), 038 doi:10.1007/JHEP03(2021)038 [arXiv:1912.07632 [hep-th]].
- [25] J. M. Drummond, R. Glew and H. Paul, "One-loop string corrections for AdS Kaluza-Klein amplitudes," JHEP 12 (2021), 072 doi:10.1007/JHEP12(2021)072 [arXiv:2008.01109 [hep-th]].
- [26] F. Aprile, J. M. Drummond, R. Glew and M. Santagata, "One-loop string amplitudes in  $AdS_5 \times S^5$ : Mellin space and sphere splitting," JHEP **02** (2023), 087 doi:10.1007/JHEP02(2023)087 [arXiv:2207.13084 [hepth]].
- [27] T. Heckelbacher, I. Sachs, E. Skvortsov and P. Vanhove, "Analytical evaluation of AdS<sub>4</sub> Witten diagrams as flat space multi-loop Feynman integrals," JHEP 08 (2022), 052 doi:10.1007/JHEP08(2022)052 [arXiv:2201.09626 [hep-th]].
- [28] L. Eberhardt, S. Komatsu and S. Mizera, "Scattering equations in AdS: scalar correlators in arbitrary dimensions," JHEP 11 (2020), 158 doi:10.1007/JHEP11(2020)158 [arXiv:2007.06574 [hepth]].
- [29] A. Herderschee, R. Roiban and F. Teng, "On the differential representation and color-kinematics duality of AdS boundary correlators," JHEP 05 (2022), 026

doi:10.1007/JHEP05(2022)026 [arXiv:2201.05067 [hep-th]].

- [30] M. Dodelson and A. Zhiboedov, "Gravitational orbits, double-twist mirage, and many-body scars," JHEP 12 (2022), 163 doi:10.1007/JHEP12(2022)163 [arXiv:2204.09749 [hep-th]].
- [31] Z. Huang and E. Y. Yuan, "Graviton scattering in  $AdS_5 \times S^5$  at two loops," JHEP **04** (2023), 064 doi:10.1007/JHEP04(2023)064 [arXiv:2112.15174 [hep-th]].
- [32] J. M. Drummond and H. Paul, "Two-loop supergravity on  $AdS_5 \times S^5$  from CFT," JHEP **08** (2022), 275 doi:10.1007/JHEP08(2022)275 [arXiv:2204.01829 [hep-th]].
- [33] Z. Huang, B. Wang, E. Y. Yuan and X. Zhou, "AdS super gluon scattering up to two loops: a position space approach," JHEP 07 (2023), 053 doi:10.1007/JHEP07(2023)053 [arXiv:2301.13240 [hepth]].
- [34] H. Y. Chen and J. i. Sakamoto, "Superconformal Block from Holographic Geometry," JHEP 07 (2020), 028 doi:10.1007/JHEP07(2020)028 [arXiv:2003.13343 [hepth]].
- [35] E. Panzer, "Algorithms for the symbolic integration of hyperlogarithms with applications to Feynman integrals," Comput. Phys. Commun. 188 (2015), 148-166 doi:10.1016/j.cpc.2014.10.019 [arXiv:1403.3385 [hep-th]].
- [36] S. Weinzierl, "Feynman Integrals," doi:10.1007/978-3-030-99558-4 [arXiv:2201.03593 [hep-th]].