

On the rank of two-dimensional simplicial distributions

Cihan Okay*

Department of Mathematics, Bilkent University, Ankara, Turkey

December 27, 2023

Abstract

Simplicial distributions provide a framework for studying quantum contextuality, a generalization of Bell's non-locality. Understanding extremal simplicial distributions is of fundamental importance with applications to quantum computing. We introduce a rank formula for twisted simplicial distributions defined for 2-dimensional measurement spaces and provide a systematic approach for describing extremal distributions.

Contents

1	Introduction	1
2	Simplicial distributions	3
2.1	Two-dimensional simplicial sets	3
2.2	Simplicial distributions	5
2.3	Twisted distributions	7
2.4	Product of distributions	10
2.5	Cohomology exact sequence	11
3	Distributions on graphs	12
3.1	Distributions on graphs	12
3.2	Rank of a distribution	14
3.3	The rank formula	16
4	Examples	17
4.1	N -cycle scenario	18
4.2	Boundary of tetrahedron	19
4.3	Mermin torus	19

1 Introduction

The theory of simplicial distributions introduced in [1] is a framework for describing quantum contextuality, a fundamental feature of quantum theory generalizing Bell's non-locality [2, 3]. In this framework, measurements and outcomes are represented by spaces modeled by combinatorial objects called simplicial sets [4]. This framework generalizes the theory of non-signaling distributions formulated in the language of sheaf-theory [5] by elevating *sets* of measurements and outcomes to *spaces* of measurements and outcomes. Simplicial distributions constitute a polytope whose vertices (extreme distributions) are of fundamental importance in quantum foundations [6–9]. Non-contextual distributions are described by a subpolytope whose



Figure 1: Triangle σ with faces $x = d_2\sigma$, $y = d_0\sigma$ and $z = d_1\sigma$. (a) A twisted simplicial distribution on σ with $\beta(\sigma) = 0$. The d_0 -face is given by $p_y^0 = p^{00} + p^{10}$. (b) A twisted simplicial distribution with $\beta(\sigma) = 1$. The d_0 -face is given by $p_y^0 = p^{01} + p^{11}$.

facets are given by the Bell inequalities. In this paper, we introduce graph-theoretic methods to identify the contextual vertices of the polytope of twisted simplicial distributions.

A simplicial distribution on a scenario consisting of a space X of measurements and space Y of outcomes is a simplicial set map of the form

$$p : X \rightarrow D(Y)$$

where $D(Y)$ is a simplicial set that models the space of distributions on the outcome space. In this paper we will restrict to the case where X is a simplicial set generated by 2-dimensional simplices $\sigma_1, \sigma_2, \dots, \sigma_N$, and Y is the nerve space of the additive group $\mathbb{Z}_2 = \{0, 1\}$. Topologically X is a 2-dimensional space obtained by gluing the 2-simplices (triangles) along their faces and possibly collapsing some of them. Our choice of the outcomes space reflects the restriction that our measurements have binary outcomes. More concretely, a simplicial distribution p consists of a family $\{p_{\sigma_i} : i = 1, 2, \dots, N\}$ of probability distributions

$$\{p_{\sigma_i}^{ab} \in \mathbb{R}_{\geq 0} : a, b \in \mathbb{Z}_2, \sum_{a,b} p_{\sigma_i}^{ab} = 1\}$$

satisfying a compatibility condition induced by the face relations. Simplicial distributions on X define a polytope with finitely many vertices. The face relations impose conditions in the form of marginals, e.g., for the d_0 -face of a triangle σ , we have

$$p_{d_0\sigma}^0 = p_{\sigma}^{00} + p_{\sigma}^{10}.$$

In this paper, we extend our interest to twisted distributions. For a 2-cocycle $\beta : X_2 \rightarrow \mathbb{Z}_2$ we write $\text{sDist}_{\beta}(X)$ for the polytope of β -twisted simplicial distributions on X . The effect of the twisting appears only in the d_0 -face, and the marginalization formula above is generalized as follows:

$$p_{d_0\sigma}^0 = \begin{cases} p_{\sigma}^{00} + p_{\sigma}^{10} & \beta(\sigma) = 0 \\ p_{\sigma}^{11} + p_{\sigma}^{01} & \beta(\sigma) = 1. \end{cases}$$

Given a twisted simplicial distribution p , we define a simplicial subset $Z_p \subset X$ consisting of simplices on which the distribution is deterministic, i.e., given by a delta-distribution. The rank of p is defined to be the rank of the matrix consisting of the defining inequalities of the polytope $\text{sDist}_{\beta}(X)$ that are tight at p . Our main result provides a formula for the rank. For a simplicial set X , we will write X_n° to denote the subset of non-degenerate simplices. Given a triangle σ , the set of 1-simplices in its boundary is denoted by $\partial\sigma$.

Theorem. *Let X be a simplicial set generated by 2-simplices $\sigma_1, \sigma_2, \dots, \sigma_N$ such that each $\partial\sigma_i$ consists of either three distinct non-degenerate 1-simplices or two distinct non-degenerate 1-simplices and a remaining degenerate 1-simplex. Consider a twisted distribution $p \in \text{sDist}_{\beta}(X)$ satisfying the following conditions:*

- for each generating 2-simplex σ , $p_{\sigma}^{ab} = 0$ for at least one pair $(a, b) \in \mathbb{Z}_2^2$, and

*cihan.okay@bilkent.edu.tr

- every non-degenerate 1-simplex of $\bar{X} = X/Z_p$ belongs precisely to two generating 2-simplices.

Then we have

$$\text{rank}(p) = |(Z_p)_1^\circ| + |\bar{X}_2^\circ| - b(\bar{X}, \bar{p}).$$

The crucial component of the rank formula is $b(X, p)$, a natural number defined for a twisted simplicial distribution. Our graph-theoretic approach is based on constructing a signed graph associated with p and $b(X, p)$ is the number of balanced components in this graph. Balancedness is an important property for signed graphs. A connected graph is called balanced if the sign of every circle contained in it, which is simply given by the product of the signs of the edges in the circle, is positive. In the theorem, we apply this construction to the quotient space $\bar{X} = X/Z_p$ and the twisted distribution \bar{p} constructed in Proposition 2.20 using a cocycle obtained by a cohomology long exact sequence. In the last section, we demonstrate how to apply the rank formula to describe the vertices of $\text{sDist}(X)$ for the following important scenarios:

- In Section 4.1, we study cycle scenarios whose measurement space consists of a disk triangulated into N triangles. This scenario is the generalization of the famous Clauser–Horne–Shimony–Holt (CHSH) scenario [10].
- The scenario whose measurement space is the boundary of a tetrahedron is considered in Section 4.2. Describing these vertices was a key step in the topological proof of Fine’s theorem [1], that characterizes non-contextual distributions on the CHSH scenario.
- The Mermin scenario [11] provides contextual distributions that arise in quantum theory. In [12], a topological realization is provided where the underlying measurement space is a torus. The vertices of the polytope of twisted distributions for different cocycles are studied in [13]. We reproduce the vertices in Section 4.3.

As displayed in these examples, the rank formula provides a systematic study of the vertices of the polytope of twisted distributions. We expect this approach will also be useful in analyzing more complicated polytopes in the context of classical simulation algorithms for quantum computation [14]. Some ideas in this direction appear in [13, 15].

The paper is structured as follows. In Section 2, we introduce simplicial sets focusing on the 2-dimensional case. We introduce twisted simplicial distributions in this restricted setting. Our main result of this section, Proposition 2.20, is proved using ideas from cohomology of simplicial sets and products of twisted distributions. In Section 3, we develop our graph-theoretical methods and introduce distributions on graphs of interest. Our rank formula is proved in Theorem 3.15. Section 4 contains the examples of practical interest where we put the rank formula in work.

Acknowledgments. This work is supported by the US Air Force Office of Scientific Research under award number FA9550-21-1-0002 and the Digital Horizon Europe project FoQaCiA, GA no. 101070558. The author would like to thank Selman Ipek and Walker H. Stern for their helpful comments on an earlier version of this paper.

2 Simplicial distributions

In this section we recall some basic definitions from [1] on simplicial distributions and introduce a twisted version in the case of 2-dimensional simplicial sets based on the exposition in [16]. Our main result is Proposition 2.20, an important ingredient for the rank formula in Section 3.

2.1 Two-dimensional simplicial sets

A simplicial set X consists of a sequence of sets X_n for $n \geq 0$ together with the face maps $d_i : X_n \rightarrow X_{n-1}$ and the degeneracy maps $s_j : X_n \rightarrow X_{n+1}$ satisfying the simplicial identities [4, 17]. A simplicial set map $f : X \rightarrow Y$ consists of functions $f_n : X_n \rightarrow Y_n$ for $n \geq 0$ compatible with the face and the degeneracy maps.

We will write f_σ for the simplex $f_n(\sigma) \in Y_n$ for a given n -simplex $\sigma \in X_n$. With this notation compatibility with the simplicial structure can be expressed as

$$d_i f_\sigma = f_{d_i \sigma} \quad \text{and} \quad s_j f_\sigma = f_{s_j \sigma}.$$

A simplex is called non-degenerate if it does not lie in the image of a degeneracy map. A non-degenerate simplex is called generating if it does not lie in the image of a face map. For a set U of simplices we write U° to denote the subset of non-degenerate simplices.

In this paper we only consider simplicial sets where each X_n is a finite set. We will also have a restriction on the dimension of the simplicial set. A 1-dimensional simplicial set is the same as a directed graph with vertex set X_0 and edge set X_1 (allowing loops and parallel edges). A 1-simplex can be treated as an arrow

$$v \xrightarrow{x} w$$

with source $d_1(x) = v$ and target $d_0(x) = w$. The degenerate 1-simplex $s_0(v)$ have both source and target given by v . A 2-dimensional simplicial set consists of such a directed graph specified by $(X_0, X_1, d_0, d_1, s_0)$ together with a finite set of 2-simplices $\sigma_1, \dots, \sigma_N$ glued to the graph. The gluing is encoded by specifying the faces $d_i(\sigma_k) \in X_1$ for $i = 0, 1, 2$. For example, the standard 2-simplex Δ^2 has a single generating 2-simplex σ glued to the directed graph on the three vertices $\{0, 1, 2\}$ with edges $0 \rightarrow 1$, $1 \rightarrow 2$ and $0 \rightarrow 2$. The face maps are given by

$$d_i \sigma = \begin{cases} 1 \xrightarrow{y} 2 & i = 0 \\ 0 \xrightarrow{z} 2 & i = 1 \\ 0 \xrightarrow{x} 1 & i = 2. \end{cases}$$

The degenerate edges $0 \xrightarrow{s_0(0)} 0$, $1 \xrightarrow{s_0(1)} 1$ and $2 \xrightarrow{s_0(2)} 2$ are usually omitted. We will write $\partial\sigma = \{x, y, z\}$ for the set of edges in the boundary. Let $f : X \rightarrow Y$ be a simplicial set map where X is generated by the 2-simplices $\sigma_1, \sigma_2, \dots, \sigma_N$. Such a map assigns a 2-simplex $f_{\sigma_k} \in Y_2$ for each $k = 1, \dots, N$ such that

$$d_i f_{\sigma_k} = d_j f_{\sigma_l}$$

whenever $d_i \sigma_k = d_j \sigma_l$. Therefore to study such maps it suffices to understand the face maps of Y in dimension 2:

$$d_i : Y_2 \rightarrow Y_1, \quad i = 0, 1, 2.$$

For simplicity when we introduce a simplicial set which will appear only in the target of a simplicial set we will only specify these face maps.

As an important target space we will be considering the nerve of the additive group $\mathbb{Z}_2 = \{0, 1\}$ of integers modulo 2. This is a simplicial set denoted by $N\mathbb{Z}_2$ whose n -simplices are given by n -tuples of elements in \mathbb{Z}_2 . The face maps in dimension 2 are given by

$$d_i(a, b) = \begin{cases} b & i = 0 \\ a + b & i = 1 \\ a & i = 2. \end{cases} \quad (1)$$

Proposition 2.1. *Let X be a simplicial set with generating 2-simplices $\sigma_1, \sigma_2, \dots, \sigma_N$. There is a bijection between simplicial set maps $s : X \rightarrow N\mathbb{Z}_2$ and functions $\varphi : X_1 \rightarrow \mathbb{Z}_2$ satisfying*

$$\varphi(x) + \varphi(y) + \varphi(z) = 0 \pmod{2}$$

for $x, y, z \in \partial\sigma_i$ where $i = 1, 2, \dots, N$.

Proof. See [1, Proposition 3.13]. □

2.2 Simplicial distributions

We will write $D(U)$ for the set of probability distributions on a set U . It consists of functions $p : U \rightarrow \mathbb{R}_{\geq 0}$ with finite support, i.e., $|\{u \in U : p(u) > 0\}| < \infty$, such that

$$\sum_{u \in U} p(u) = 1.$$

Given a function $f : U \rightarrow V$ and a distribution $p \in D(U)$ we define $f_*(p) \in D(V)$ by the formula

$$f_*(p)(v) = \sum_{u \in f^{-1}(v)} p(u).$$

We will write δ^u for the delta-distribution peaked at an element $u \in U$. Given a simplicial set Y we can construct a new simplicial set $D(Y)$ whose set of n -simplices consists of $D(Y_n)$. The face and the degeneracy maps of this simplicial set are given by $d_i(p) = (d_i)_*(p)$ and $s_j(p) = (s_j)_*(p)$. We will write $\delta : Y \rightarrow D(Y)$ for the simplicial set map that sends a simplex to the delta-distribution peaked at that simplex.

Our interest is in the space $D(N\mathbb{Z}_2)$. The set of n -simplices consists of distributions of the form

$$p : \mathbb{Z}_2^n \rightarrow \mathbb{R}_{\geq 0}.$$

We write $p^{a_1 \dots a_n}$ for the value $p(a_1, \dots, a_n)$. With this notation the face maps in dimension 2 are given by

$$(d_i p)^0 = \begin{cases} p^{00} + p^{10} & i = 0 \\ p^{00} + p^{11} & i = 1 \\ p^{00} + p^{01} & i = 2. \end{cases} \quad (2)$$

See Figure (1a).

Definition 2.2. A (simplicial) scenario consists of a pair (X, Y) of simplicial sets representing the space of measurements and outcomes, respectively. A simplicial distribution on the scenario (X, Y) is a simplicial set map

$$p : X \rightarrow D(Y)$$

We write $\text{sDist}(X, Y)$ for the set of simplicial distributions.

When $Y = N\mathbb{Z}_2$ we will simplify the notation and write $\text{sDist}(X)$ for the set of simplicial distributions on X .

Proposition 2.3. Let X be a simplicial set with generating 2-simplices $\sigma_1, \sigma_2, \dots, \sigma_N$. Then a simplicial distribution $p : X \rightarrow D(N\mathbb{Z}_2)$ is given by a collection of distributions $p_{\sigma_k} \in D(\mathbb{Z}_2^2)$ satisfying

$$d_i p_{\sigma_k} = d_j p_{\sigma_l} \quad (3)$$

whenever $d_i \sigma_k = d_j \sigma_l$. In particular, $\text{sDist}(X)$ is a (convex) polytope with finitely many vertices.

Proof. A simplicial set map $p : X \rightarrow D(N\mathbb{Z}_2)$ is determined by its values, i.e., the distributions $p_{\sigma_i} \in D(\mathbb{Z}_2^2)$, on the generating simplices. Note that on the remaining simplices the image is determined by the simplicial structure maps. The only relations imposed on p_{σ_i} 's come from the face maps given in Equation (3). Therefore $\text{sDist}(X)$ is the subspace obtained by intersecting $[0, 1]^{4N} \subset \mathbb{R}^{4N}$ with the linear equations corresponding to normalization and identifications under the face maps. This subspace is convex and bounded. Since the set of equations involved is finite there are finitely many extreme points (vertices). \square

Example 2.4. The simplest case is when $X = \Delta^2$:

$$\text{sDist}(\Delta^2) = \{(p^{00}, p^{01}, p^{10}, p^{11}) \in [0, 1]^4 : p^{00} + p^{01} + p^{10} + p^{11} = 1\},$$

which defines a polytope in \mathbb{R}^3 , e.g., by retaining the coordinates (p^{01}, p^{10}, p^{11}) .

In general, X is obtained by gluing N of the Δ^2 's and $\text{sDist}(X)$ is a convex polytope in \mathbb{R}^{3N} cut out by the linear equations (3).

Definition 2.5. A simplicial distribution is called a deterministic distribution if it is of the form $\delta^s : X \xrightarrow{s} Y \xrightarrow{\delta} D(Y)$ where $s : X \rightarrow Y$ is a simplicial set map. We will write $\text{dDist}(X, Y)$ for the set of deterministic distributions. There is a natural map

$$\Theta : D(\text{dDist}(X, Y)) \rightarrow \text{sDist}(X, Y)$$

defined by sending $d = \sum_s d(s)\delta^s$ to the simplicial distribution p given by

$$p_\sigma(\theta) = \sum_{s:s_\sigma=\theta} d(s).$$

A simplicial distribution p is non-contextual if p is in the image of Θ . Otherwise, p is called contextual.

When $Y = N\mathbb{Z}_2$ we will simply write $\text{dDist}(X)$ for the set of deterministic distributions on X .

Example 2.6. Let us consider $X = \Delta^2$. By Proposition 2.1 we have

$$\text{dDist}(\Delta^2) \xrightarrow{\cong} \mathbb{Z}_2^2$$

obtained by sending $s : \Delta^2 \rightarrow N\mathbb{Z}_2$ to the pair $(s_{d_2\sigma}, s_{d_0\sigma})$. The corresponding deterministic distribution is given by

$$(\delta_\sigma^s)^{ab} = \begin{cases} 1 & (a, b) = (s_{d_2\sigma}, s_{d_0\sigma}) \\ 0 & \text{otherwise.} \end{cases}$$

For notational convenience we will write δ^{ab} when s is specified by (a, b) under the bijection above. Observe that any simplicial distribution p on the triangle can be written as

$$p = \sum_{a,b} p^{ab} \delta^{ab}.$$

Therefore every p on Δ^2 is non-contextual according to Definition 2.5.

Example 2.7. The simplest contextual example can be obtained by identifying any two edges of the triangle. Formally this is expressed as a push-out diagram

$$\begin{array}{ccc} \Delta^1 \sqcup \Delta^1 & \hookrightarrow & \Delta^2 \\ \downarrow & & \downarrow \\ \Delta^1 & \longrightarrow & \Delta^2/x \sim z \end{array}$$

where the top horizontal map sends the generating simplices of Δ^1 's to $x = d_2\sigma$ and $z = d_1\sigma$. Then we have

$$\Theta : D(\{\delta^{00}, \delta^{10}\}) \rightarrow \{(p^{00}, p^{01}, p^{10}, p^{11}) \in [0, 1]^4 : \sum_{a,b} p^{ab} = 1, p^{01} = p^{11}\}$$

and a simplicial distribution p is contextual if and only if $p^{11} > 0$. The analysis is similar for the case where the other pairs of edges are identified.

We can also identify all three edges to obtain $\Delta^2/x \sim y \sim z$. Then

$$\Theta : D(\{\delta^{00}\}) \rightarrow \{(p^{00}, p^{01}, p^{10}, p^{11}) \in [0, 1]^4 : \sum_{a,b} p^{ab} = 1, p^{01} = p^{10} = p^{11}\}$$

and again p is contextual if and only if $p^{11} > 0$.

Example 2.8. Another topological operation we can perform on a single triangle is to collapse an edge. This is given by a push-out diagram of the form

$$\begin{array}{ccc} \Delta^1 & \xrightarrow{\quad} & \Delta^2 \\ \downarrow & & \downarrow \\ \Delta[0] & \xrightarrow{\quad} & \Delta^2/z \sim * \end{array}$$

Then we have

$$\Theta : D(\{\delta^{00}, \delta^{11}\}) \xrightarrow{\cong} \{(p^{00}, 0, 0, p^{11}) \in [0, 1]^4 : p^{00} + p^{11} = 1\}.$$

In this case every p is non-contextual. Collapsing two of the edges to obtain $\Delta^2/y \sim z \sim *$ we obtain

$$\Theta : D(\{\delta^{00}\}) \xrightarrow{\cong} \{(1, 0, 0, 0)\}$$

and again every simplicial distribution is non-contextual. The situation is exactly the same when all three edges are collapsed.

2.3 Twisted distributions

We will use the more general notion of simplicial distributions introduced in [16].

Definition 2.9. A (simplicial) bundle scenario is a simplicial set map $\pi : E \rightarrow X$. A simplicial distribution on the bundle scenario f is a simplicial set map $p : X \rightarrow D(E)$ that makes the following diagram commute

$$\begin{array}{ccc} & & D(E) \\ & \nearrow p & \downarrow D(\pi) \\ X & \xrightarrow{\delta} & D(X) \end{array} \quad (4)$$

A deterministic distribution on f is a simplicial set map of the form $\delta^s : X \xrightarrow{s} E \xrightarrow{\delta} D(E)$ where $s : X \rightarrow E$ is a section of π .

The earlier notion given in Definition 2.2 can be recovered by considering simplicial distributions on the bundle scenario given by the projection map

$$\pi : Y \times X \rightarrow X$$

For twisted distributions we will need the notion of twisted products. Twisted products model principal bundles [18]. Let us write $H_n(X)$ and $H^n(X)$ for the n -th homology and cohomology groups with coefficients in \mathbb{Z}_2 , respectively. When X is 2-dimensional there is a well-known classification theorem: the set of isomorphism classes of principal bundles with fiber $N\mathbb{Z}_2$ is in bijective correspondence with the classes in the second cohomology group $H^2(X)$. Let us recall the definition of the n -th cohomology group of a simplicial set [4]. Given a simplicial set X let us write $C^n(X)$ for the set $\mathbb{Z}_2^{X_n}$ of functions $\alpha : X_n \rightarrow \mathbb{Z}_2$. The coboundary map $\delta_n : C^n(X) \rightarrow C^{n+1}(X)$ is defined by

$$\delta_n \alpha(x) = \sum_{i=0}^n (-1)^i \alpha(d_i x).$$

Then $H^n(X)$ is defined as the quotient of the kernel of δ_n by the image of δ_{n-1} . A function $\alpha : X_n \rightarrow \mathbb{Z}_2$ is called a cocycle if it belongs to the kernel of δ_n . It is called normalized if $\alpha(x) = 0$ for every degenerate n -simplex x .

Definition 2.10. Let $\beta : X_2 \rightarrow \mathbb{Z}_2$ be a normalized 2-cocycle. The twisted product $N\mathbb{Z}_2 \times_{\beta} X$ is the simplicial set whose set of n -simplices is given by $(N\mathbb{Z}_2)_n \times X_n$. The face map in dimension 2 is given by

$$d_i((a, b), \sigma) = \begin{cases} (b + \beta(\sigma), d_0 \sigma) & i = 0 \\ (a + b, d_1 \sigma) & i = 1 \\ (a, d_2 \sigma) & i = 2. \end{cases} \quad (5)$$

Note that the d_0 -face is twisted by the cocycle β . Projecting onto the second coordinate gives a principal $N\mathbb{Z}_2$ -bundle

$$\pi : N\mathbb{Z}_2 \times_{\beta} X \rightarrow X$$

We are interested in simplicial distributions on such maps.

Definition 2.11. A β -twisted distribution is a simplicial distribution on $\pi : N\mathbb{Z}_2 \times_{\beta} X \rightarrow X$. A β -twisted deterministic distribution is a deterministic distribution on π . We will write $\text{sDist}_{\beta}(X)$ and $\text{dDist}_{\beta}(X)$ for the sets of β -twisted simplicial and deterministic distributions on X , respectively.

Let us unravel the definition of a twisted distribution when X is 2-dimensional generated by $\sigma_1, \dots, \sigma_N$. A twisted simplicial distribution associates a distribution $p_{\sigma_i} \in D(\mathbb{Z}_2^2 \times X_2)$ with each 2-simplex σ_i . The commutativity of diagram (4) implies that the support of the distribution is contained in $\mathbb{Z}_2^2 \times \{\sigma_i\}$. Face maps can be worked out from Equation (5). Only the d_0 -face is twisted

$$(d_0 p_{\sigma})^0 = \begin{cases} p^{00} + p^{10} & \beta(\sigma) = 0 \\ p^{11} + p^{01} & \beta(\sigma) = 1. \end{cases} \quad (6)$$

See Figure (1b).

A version of Proposition 2.3 holds for $\text{sDist}_{\beta}(X)$ once the d_0 -face is twisted in this way. We will provide a more explicit description of the resulting polytope. We define the correlation function

$$c : X_1 \rightarrow \mathbb{R} \quad (7)$$

by $c(x) = p_x^0 - p_x^1$. The probabilities p_{σ}^{ab} for each 2-simplex $\sigma \in X_2$ can be recovered from the correlation function.

Lemma 2.12. *We have*

$$p_{\sigma}^{ab} = \frac{1}{4}(1 + (-1)^a c(x_2) + (-1)^{b+\beta(\sigma)} c(x_0) + (-1)^{a+b} c(x_1)). \quad (8)$$

Proof. This formula can be verified using the marginal relations (e.g., d_0 face as in Equation (6) and others) and the relation $p_{x_i}^1 = 1 - p_{x_i}^0$. For example, for $(a, b) = (0, 0)$ and $\beta(\sigma) = 1$ we have

$$\begin{aligned} \frac{1}{4}(1 + c(x_2) - c(x_0) + c(x_1)) &= \frac{1}{4}(1 + (2p_{x_2} - 1) - (2p_{x_0} - 1) + (2p_{x_1} - 1)) \\ &= \frac{1}{4}(1 + (2(p_{\sigma}^{00} + p_{\sigma}^{01}) - 1) - (2(p_{\sigma}^{11} + p_{\sigma}^{01}) - 1) + (2(p_{\sigma}^{00} + p_{\sigma}^{11}) - 1)) \\ &= p_{\sigma}^{00}. \end{aligned}$$

□

Using this result we can provide a more explicit description for the polytope of twisted simplicial distributions. For a $m \times d$ matrix M and a column vector b of size m we will write

$$P(M, b) = \{t \in \mathbb{R}^d : Mt \geq b\}$$

for the corresponding polytope in \mathbb{R}^d . Let $\mathbb{1}$ denote a column vector consisting of 1's.

Proposition 2.13. *Let $d = |X_1|$ and $m = |X_2 \times \mathbb{Z}_2^2|$. Then*

$$\text{sDist}_{\beta}(X) = P(M, -\mathbb{1})$$

where M is the $m \times d$ matrix defined by

$$M_{(\sigma, ab), x} = \begin{cases} (-1)^{b+\beta(\sigma)} & x = d_0\sigma \\ (-1)^{a+b} & x = d_1\sigma \\ (-1)^a & x = d_2\sigma. \end{cases}$$

Proof. This follows directly from Lemma 2.12. □

Proposition 2.14. *We have the following properties.*

1. The set $\text{dDist}_\beta(X)$ is non-empty if and only if $[\beta] = 0$.
2. There is a bijection

$$\text{sDist}_\beta(X) \xrightarrow{\cong} \text{sDist}_\alpha(X)$$

when $[\beta] = [\alpha]$.

Proof. Part (1): The cohomology class $[\beta]$ vanishes if and only if the principal bundle $N\mathbb{Z}_2 \times_\beta X \rightarrow X$ is trivial. The latter holds if and only if it admits a section. Part (2): The cohomology classes coincide if and only if the corresponding principal bundles are isomorphic (as principal bundles). This gives a commutative diagram where the top arrow is an isomorphism

$$\begin{array}{ccc} D(N\mathbb{Z}_2 \times_\beta X) & \xrightarrow{\cong} & D(N\mathbb{Z}_2 \times_\alpha X) \\ & \searrow & \swarrow \\ & D(X) & \end{array}$$

Therefore the resulting sets of simplicial distributions are in bijective correspondence. □

Example 2.15. Let D_N be a simplicial set obtained by gluing N many 2-simplices to obtain a disk, e.g., as in Figure (2). For a more general definition see [19, Definition 3.1]. Since the resulting space is contractible we have $H_2(D_N) = 0$. Therefore for any normalized cocycle $\beta : (D_N)_2 \rightarrow \mathbb{Z}_2$ we have $[\beta] = 0$. By Proposition 2.14 we have $\text{dDist}_\beta(D_N) \neq \emptyset$ and

$$\text{sDist}_\beta(D_N) \cong \text{sDist}(D_N).$$

Then the gluing lemma of [1] implies that every $p \in \text{sDist}_\beta(D_N)$ is non-contextual.




Figure 2

For a simplicial subset $Z \subset X$ and a simplicial distribution on X we write $p|_Z$ for the composite $Z \xrightarrow{i} X \xrightarrow{p} D(E)$ where i is the inclusion map. The following simple observation will be very useful in analyzing the vertices of the polytope of twisted distributions.

Lemma 2.16. ([13]) *Let $Z \subset X$ be a simplicial subset with a single generating 2-simplex σ . If two of the faces, say $x, y \in \partial\sigma$, satisfy $p_x = \delta^a$ and $p_y = \delta^b$ for some $a, b \in \mathbb{Z}_2$ then $p|_Z$ is a deterministic distribution.*

Proof. When the distribution is deterministic on two of the edges this forces three of the four parameters in $\text{sDist}_\beta(\Delta^2) \subset [0, 1]^4$ to be zero. Together with the normalization condition we obtain a unique solution, i.e., a deterministic distribution. □

Example 2.17. The measurement space \tilde{C}_4 of the Clauser–Horne–Shimony–Holt (CHSH) scenario consists of four triangles $\sigma_1, \sigma_2, \sigma_3, \sigma_4$ glued as in Figure (3a). This is a special case of cycle scenarios we will examine in Section 4.1. It is well-known that the polytope $\text{sDist}(\tilde{C}_4)$ has two kinds of vertices consisting of the deterministic distributions and contextual distributions known as the Popescu–Rohrlich (PR) boxes [7];

see also [19, Example 3]. PR boxes are characterized by the property that the restriction $p|_{\partial\tilde{C}_4}$ to the boundary, which consists of four edges x_1, x_2, x_3, x_4 , is a deterministic distribution δ^s such that

$$\sum_{i=1}^4 a_i = 1 \pmod{2}$$

where $\delta_{x_i}^s = \delta^{a_i}$ for $i = 1, 2, \dots, N$.



Figure 3: (a) The measurement space of the CHSH scenario. (b) PR box with $a_1 + a_2 + a_3 + a_4 = 1 \pmod{2}$. Green color indicates the deterministic edges.

2.4 Product of distributions

In [20] a product is introduced for simplicial distributions. We can extend this product to 2-dimensional twisted distributions. The convolution product of $p, q \in D(\mathbb{Z}_2^n)$ is the distribution $p * q$ defined by

$$p * q(c) = \sum_{a+b=c} p(a)q(b)$$

where the sum runs over $a, b \in \mathbb{Z}_2^n$ such that $a + b = c$. Using the convolution product we define a map

$$\text{sDist}_\alpha(X) \times \text{sDist}_\beta(X) \rightarrow \text{sDist}_{\alpha+\beta}(X)$$

by sending (p, q) to the distribution $p \cdot q$ given by

$$(p \cdot q)_\sigma = p_\sigma * q_\sigma.$$

Lemma 2.18. *The product $p \cdot q$ is an $(\alpha + \beta)$ -twisted distribution.*

Proof. We have

$$\begin{aligned} ((p \cdot q)_{d_0\sigma})^0 &= p_{d_0\sigma}^0 q_{d_0\sigma}^0 + p_{d_0\sigma}^1 q_{d_0\sigma}^1 \\ &= \sum_a p_\sigma^{a(\alpha(\sigma))} \sum_b q_\sigma^{b(\beta(\sigma))} + \sum_a p_\sigma^{a(\alpha(\sigma)+1)} \sum_b q_\sigma^{b(\beta(\sigma)+1)} \\ &= (p \cdot q)^{0(\alpha(\sigma)+\beta(\sigma))} + (p \cdot q)^{1(\alpha(\sigma)+\beta(\sigma))} \\ &= (d_0(p \cdot q)_\sigma)^0. \end{aligned}$$

Similarly one can verify that $p \cdot q$ is compatible with the remaining simplicial structure maps. Commutativity of diagram (4) follows from the observation that the support of $p \cdot q$ is contained in $\mathbb{Z}_2^2 \times \{\sigma_i\}$ by definition of the product. \square

It is instructive to consider the action of $\text{dDist}_\alpha(X)$ on $\text{sDist}_\beta(X)$ induced by this product. Note that because of part (1) of Proposition 2.14 we assume $[\alpha] = 0$, or for computational simplicity $\alpha = 0$. Then we have

$$(\delta^{ab} \cdot q)_\sigma^{cd} = q_\sigma^{(c+a)(b+d)} \quad (9)$$

where $\delta^{ab} \in \text{dDist}(X)$ and $q \in \text{sDist}_\beta(X)$.

2.5 Cohomology exact sequence

Let $Z \subset X$ be a simplicial subset. Given a normalized 2-cocycle $\beta : X_2 \rightarrow \mathbb{Z}_2$ we will write $\beta|_Z$ for the pull-back $i^*\beta$ along the inclusion map $i : Z \rightarrow X$. We assume $[\beta|_Z] = 0$ so that there exists a normalized 1-cochain $s : Z_1 \rightarrow \mathbb{Z}_2$ such that $\beta|_Z = \delta s$. We will write

$$\text{sDist}_\beta(X, s) = \{p \in \text{sDist}_\beta(X) : p|_Z = \delta^s\}.$$

Consider the cofiber sequence

$$Z \xrightarrow{i} X \xrightarrow{q} X/Z$$

and the associated cohomology long exact sequence

$$H^1(X/Z) \rightarrow H^1(X) \rightarrow H^1(Z) \xrightarrow{\zeta} H^2(X/Z)$$

Let $\zeta(s)$ denote the 2-cocycle obtained by the snake lemma [21], i.e., first extending s to X and applying the coboundary:

- Let $\tilde{s} : X_1 \rightarrow \mathbb{Z}_2$ denote the 2-cochain defined by

$$\tilde{s}(x) = \begin{cases} s(x) & x \in Z_1 \\ 0 & x \in X_1 - Z_1. \end{cases}$$

- By applying the coboundary map $\delta : C_1(X) \rightarrow C_2(X)$ we obtain

$$\delta\tilde{s}(\sigma) = \tilde{s}(d_0\sigma) - \tilde{s}(d_1\sigma) + \tilde{s}(d_2\sigma)$$

where $\sigma \in X_2$.

- Since $\delta\tilde{s}|_Z = 0$ it comes from a 2-cochain, which will be denoted by $\zeta(s) : (X/Z)_2 \rightarrow \mathbb{Z}_2$.

See also [1, Section 5]

Lemma 2.19. *The deterministic distribution $\delta^{\tilde{s}}$ on X defined by*

$$(\delta^{\tilde{s}})^{ab} = \begin{cases} 1 & (a, b) = (\tilde{s}(d_2\sigma), \tilde{s}(d_0\sigma)) \\ 0 & \text{otherwise} \end{cases}$$

is a $\delta^{\tilde{s}}$ -twisted distribution.

Proof. This is direct verification. □

Proposition 2.20. *We have a commutative diagram*

$$\begin{array}{ccc} D(\text{dDist}_\beta(X, s)) & \xrightarrow{\Theta} & \text{sDist}_\beta(X, s) \\ \cong \downarrow D(\phi) & & \cong \downarrow \phi \\ D(\text{dDist}_{\beta+\zeta(s)}(X/Z)) & \xrightarrow{\Theta} & \text{sDist}_{\beta+\zeta(s)}(X/Z) \end{array}$$

where $\phi(p) = \delta^{\tilde{s}} \cdot p$. Both vertical arrows are isomorphisms.

Proof. Let $p \in \text{sDist}_\beta(X)$ be such that $p|_Z = \delta^s$ for some s satisfying $\delta s = \beta|_Z$. By Lemma 2.18 and 2.19 the distribution $\delta^{\tilde{s}} \cdot p$ is $(\beta + \zeta(s))$ -twisted and

$$(\delta^{\tilde{s}} \cdot p)|_Z = \delta^{\tilde{s}}|_Z \cdot p|_Z = \delta^s \cdot \delta^s = \delta^0$$

where $\delta^0 : Z \rightarrow D(N\mathbb{Z}_2)$ is the constant map (with image the unique vertex). Note that in the last equation we used Equation (9). Therefore $\delta^{\tilde{s}} \cdot p$ factors through the quotient X/Z . The inverse of ϕ is given by the composite

$$\phi^{-1} : \text{sDist}_{\beta+\zeta(s)}(X/Z) \xrightarrow{q^*} \text{sDist}_{\beta+d\tilde{s}}(X) \xrightarrow{\delta^{\tilde{s}}} \text{sDist}_\beta(X)$$

The maps ϕ and ϕ^{-1} restrict to a bijection between $\text{dDist}_\beta(X, s)$ and $\text{dDist}_{\beta+\zeta(s)}(X/Z)$. □



Figure 4: Twisted simplicial distributions on a triangle where x is identified with z and y is collapsed: (a) $\beta(\sigma) = 0$ and (b) $\beta(\sigma) = 1$. The resulting polytope can be identified with the unit interval $[0, 1]$. Here $\bar{t} = 1 - t$.

Corollary 2.21. *If $\phi(p)$ is non-contextual then p is non-contextual.*

Example 2.22. Let us consider $\text{sDist}(X)$ where $X = \Delta^2/x \sim z$; see Example 2.7. We take Z to be the simplicial subset given by the edge y . If the restriction $p|_Z$ is deterministic then it is either δ^0 or δ^1 . Then by Proposition 2.20 we have an isomorphism

$$\text{sDist}(X, s) \xrightarrow{\cong} \text{sDist}_\beta(X/y \sim *)$$

Note that $\beta(\sigma) = a$ if $p_y^a = \delta^a$. See Figure (4).

3 Distributions on graphs

Our goal in this section is to describe twisted distributions as distributions on graphs. This is achieved by associating a graph with a simplicial set. Throughout the paper we only consider simple graphs, i.e., graphs with no loops and parallel edges. To land in simple graphs we impose some conditions on the simplicial sets that represent the measurements. We will restrict to a measurement space X given by a simplicial set

- generated by the 2-simplices $\sigma_1, \sigma_2, \dots, \sigma_N$, and
- each set $\partial\sigma_k$ of edges in the boundary consists of either three distinct non-degenerate edges $\{x_0, x_1, x_2\}$ or two distinct non-degenerate edges $\{x_i, x_j\}$, where $i > j$, and a remaining degenerate edge.

Therefore we have the situation depicted in Figure (5).




Figure 5

3.1 Distributions on graphs

We begin by constructing a bipartite graph associated to the simplicial set X . A bipartite graph Γ consists of a vertex set $V(\Gamma)$, partitioned into two sets $V^0(\Gamma) \sqcup V^1(\Gamma)$, and an edge set $E(\Gamma)$ connecting the vertices from these two sets. We will also consider graphs with a sign given by a function $\gamma : E(\Gamma) \rightarrow \{\pm 1\}$.

Definition 3.1. Given a finite 2-dimensional simplicial set X let $\Gamma(X)$ denote the bipartite graph with

- vertex set $V = X_1^\circ \sqcup X_2^\circ$,
- edge set E consisting of $\{x, \sigma\}$ where $x \in \partial\sigma^\circ$.

Next, we enlarge this graph by including the set of outcomes into the picture. The idea is to replace each vertex corresponding to σ by four vertices labeled by s_σ^{ab} where $a, b \in \mathbb{Z}_2$. The new vertices are also connected to the same vertices corresponding to the 1-simplices in $\partial\sigma^\circ$. These edges of the graph are also assigned a sign ± 1 indicating the outcomes. See Figure (6) for the local picture over a triangle with two kinds of boundaries, one with $|\partial\sigma^\circ| = 3$ and 2.

Definition 3.2. Given a normalized 2-cocycle $\beta : X_2 \rightarrow \mathbb{Z}_2$ we define a signed bipartite graph $\Gamma_\beta(X)$ with

- vertex set $X_1^\circ \sqcup (X_2^\circ \times \mathbb{Z}_2^2)$,
- edge set consisting of $\{x, s_\sigma^{ab}\}$ where $x \in \partial\sigma^\circ$,
- and sign given by

$$\gamma(x, s_\sigma^{ab}) = (-1)^{s_\sigma(x)} \quad (10)$$

where $s_\sigma : \partial\sigma \rightarrow \mathbb{Z}_2$ is defined by

$$s_\sigma(x) = \begin{cases} (-1)^{b+\beta(\sigma)} & x = d_0\sigma \\ (-1)^{a+b} & x = d_1\sigma \\ (-1)^a & x = d_2\sigma. \end{cases}$$

Next, we introduce the notion of a distribution on the bipartite graph associated to the pair (X, β) . The goal is to capture the notion of twisted simplicial distributions as distributions on graphs. The next definition is motivated by Proposition 2.13.

Definition 3.3. A distribution on $\Gamma_\beta(X)$ is a function $p : X_2^\circ \times \mathbb{Z}_2^2 \rightarrow \mathbb{R}_{\geq 0}$ such that

$$\sum_{a,b} p(s_\sigma^{ab}) = 1$$

for every $\sigma \in X_2^\circ$ and

$$\sum_{a,b} \gamma(x, s_\sigma^{ab}) p(s_\sigma^{ab}) = \sum_{a,b} \gamma(x, s_{\sigma'}^{ab}) p(s_{\sigma'}^{ab}) \quad (11)$$

for every $\sigma, \sigma' \in X_2^\circ$ such that $x \in \partial\sigma^\circ \cap \partial\sigma'^\circ$. In addition, for simplices $\sigma \in X_2^\circ$ whose boundary contains a single degenerate edge x' we also require that

$$\begin{cases} p(s_\sigma^{(0+\beta(\sigma))(0+\beta(\sigma))}) + p(s_\sigma^{(1+\beta(\sigma))(0+\beta(\sigma))}) = 1 & x' = d_0\sigma \\ p(s_\sigma^{00}) + p(s_\sigma^{11}) = 1 & x' = d_1\sigma \\ p(s_\sigma^{00}) + p(s_\sigma^{01}) = 1 & x' = d_2\sigma. \end{cases} \quad (12)$$

We write $\text{Dist}(\Gamma_\beta(X))$ for the set of distributions on the graph $\Gamma_\beta(X)$.

Equation (12) has the following interpretation. As in simplicial distributions we want $p_{x'} = \delta^0$ for each degenerate edge x' . This condition is imposed via this equation separately since the degenerate edges of X does not appear in the graph $\Gamma_\beta(X)$. See Figure (6) for the local picture over a simplex. Definition 3.3 is tailored so that the following identification can be done.

Proposition 3.4. Sending a twisted distribution $p \in \text{sDist}_\beta(X)$ to the distribution on the graph $\Gamma_\beta(X)$ defined by

$$p(s_\sigma^{ab}) = p_\sigma^{ab}$$

gives a bijection of convex sets

$$\text{sDist}_\beta(X) \xrightarrow{\cong} \text{Dist}(\Gamma_\beta(X)) \quad (13)$$

Proof. Follows from Proposition 2.13. □

We will identify twisted simplicial distributions and distributions on the associated enlarged graph.




Figure 6: Signs on the edges induced by s_σ^{ab} are indicated by blue and red color for $+1$ and -1 , respectively. The probabilities are given by $p^{ab} = p(s_\sigma^{ab})$. In this example $\beta(\sigma) = 0$.

3.2 Rank of a distribution

Definition 3.5. Given a distribution $p \in \text{Dist}(\Gamma_\beta(X))$ we consider the induced signed subgraph $\Gamma_\beta(X, p) \subset \Gamma_\beta(X)$ determined by the vertex set

$$X_1^\circ \sqcup \{s_\sigma^{ab} \in X_2^\circ \times \mathbb{Z}_2^2 : p(s_\sigma^{ab}) = 0\}.$$

The edge set is determined by the vertices, that is, $\{x, s_\sigma^{ab}\}$ is an edge if $p(s_\sigma^{ab}) = 0$ and $x \in \partial\sigma^\circ$.

Let $A(\Gamma_\beta(X, p))$ denote the adjacency matrix of the signed graph. Its rows and columns are indexed by the vertices of the graph and its entries are $0, \pm 1$ indicating whether there exists an edge connecting the pair of vertices (taking into account the sign). Since the graph is bipartite the adjacency matrix has the form

$$A(\Gamma_b) = \begin{pmatrix} \mathbb{0} & B(\Gamma_\beta(X, p)) \\ B(\Gamma_\beta(X, p))^T & \mathbb{0} \end{pmatrix}$$

where

$$B(\Gamma_\beta(X, p))_{s_\sigma^{ab}, x} = \begin{cases} \gamma(x, s_\sigma^{ab}) & \{x, s_\sigma^{ab}\} \in E(\Gamma_\beta(X, p)) \\ 0 & \text{otherwise.} \end{cases}$$

Definition 3.6. The rank of $p \in \text{Dist}(\Gamma_\beta(X))$ is defined to be the rank of the matrix $B(\Gamma_\beta(X, p))$. The rank of a twisted distribution $p \in \text{sDist}_\beta(X)$ is the rank of the associated distribution on the graph.

Corollary 3.7. A twisted distribution $p \in \text{sDist}_\beta(X) \subset \mathbb{R}^{|X_1^\circ|}$ is a vertex if and only if $\text{rank}(p) = |X_1^\circ|$.

Proof. This observation follows from the general theory of polytopes; see [22, Theorem 18.1]. \square

Note that the polytope $\text{sDist}_\beta(X)$ is not full-dimensional in $\mathbb{R}^{|X_1^\circ|}$ as its dimension is given by

$$d(X) = |X_1^\circ| - \sum_{\sigma \in X_2^\circ} |\partial\sigma - \partial\sigma^\circ|.$$

This dimension count follows from Example 2.4 and 2.8.

Definition 3.8. Let $p \in \text{sDist}_\beta(X)$ be a twisted distribution. A non-degenerate 1-simplex x is called a deterministic edge if $p_x = \delta^a$ for some $a \in \mathbb{Z}_2$. Similarly a non-degenerate 2-simplex σ is called a deterministic triangle if $p_\sigma = \delta^{ab}$ for some $(a, b) \in \mathbb{Z}_2^2$. We will write $Z_p \subset X$ for the simplicial subset generated by the deterministic edges and triangles with respect to p .

Recall the bijection in Proposition 2.20:

$$\phi : \mathbf{sDist}_\beta(X, s) \rightarrow \mathbf{sDist}_{\bar{\beta}}(\bar{X})$$

where $\bar{X} = X/Z_p$ and $\bar{\beta} = \beta + \zeta(s)$. In the next result we show that the rank of $\bar{p} = \phi(p)$ is related to the rank of p .

Lemma 3.9. *For $p \in \mathbf{sDist}_\beta(X)$ we have*

$$\text{rank}(p) = \text{rank}(\bar{p}) + |(Z_p^\circ)_1|. \quad (14)$$

Proof. For $\sigma \in X_2^\circ$ let us write $\Gamma_\beta(X, p)|_\sigma \subset \Gamma_\beta(X, p)$ for the induced subgraph on the vertices $\partial\sigma^\circ \sqcup (\{\sigma\} \times \mathbb{Z}_2^2)$. We will take the quotient in steps, that is, one deterministic simplex at a time. By Lemma 2.16 each $\sigma \in (Z_p)^\circ_2$ satisfies $|\partial\sigma^\circ| = 2$ or 3. Therefore we have

$$X = X^{(0)} \rightarrow X^{(1)} \rightarrow \dots \rightarrow X^{(i)} \rightarrow X^{(i+1)} \rightarrow \dots \rightarrow X^{(t-1)} \rightarrow X^{(t)} = X/Z_p$$

where $X^{(i+1)}$ is obtained from $X^{(i)}$ by killing a non-degenerate deterministic edge or a non-degenerate deterministic triangle. Let us consider these two types of quotients:

1. Killing a non-degenerate deterministic edge x which does not belong to a deterministic triangle: Let σ be a non-degenerate triangle whose boundary contains x . Since p_σ is not deterministic $|\partial\sigma^\circ| = 3$ (by Lemma 2.16). We have

$$B(\Gamma_\beta(X, p)|_\sigma) = \begin{bmatrix} (-1)^{s(x)} & 1 & (-1)^{s(x)+\beta(\sigma)} \\ (-1)^{s(x)} & -1 & (-1)^{s(x)+\beta(\sigma)+1} \end{bmatrix} \sim \begin{bmatrix} 0 & 1 & (-1)^{s(x)+\beta(\sigma)} \\ 1 & 0 & 0 \end{bmatrix}$$

where \sim indicates a sequence of elementary row operations. On the other hand, we have

$$B(\Gamma_{\bar{\beta}}(\bar{X}, \bar{p})|_\sigma) = \begin{bmatrix} 1 & (-1)^{\bar{\beta}(\sigma)} \\ -1 & (-1)^{\bar{\beta}(\sigma)+1} \end{bmatrix} = \begin{bmatrix} 1 & (-1)^{s(x)+\beta(\sigma)} \\ -1 & (-1)^{s(x)+\beta(\sigma)+1} \end{bmatrix} \sim \begin{bmatrix} 1 & (-1)^{s(x)+\beta(\sigma)} \\ 0 & 0 \end{bmatrix}$$

where $\bar{\beta}(\sigma) = \beta(\sigma) + \zeta(s)(\sigma)$. Therefore we have $\text{rank}(B(\Gamma_\beta(X, p)|_\sigma)) = \text{rank}(B(\Gamma_{\bar{\beta}}(\bar{X}, \bar{p})|_\sigma)) + 1$. That is, the rank at σ has two parts. First part comes from the new distribution on the quotient space and the second part from the deterministic edge. Therefore Equation (14) holds.

2. Killing a non-degenerate deterministic triangle σ : Since p_σ is deterministic $B(\Gamma_\beta(X, p)|_\sigma)$ is a 3×3 -matrix of rank 3. This time all the contribution comes from the deterministic edges and thus Equation (14) holds.

When a deterministic edge is killed, either as in (1) or part of a deterministic triangle as in (2), the corresponding column in $B(\Gamma_\beta(X, p))$ can be removed. Once all such columns are removed the remaining matrix has the same rank as $B(\Gamma_{\bar{\beta}}(\bar{X}, \bar{p}))$ by the local computations at (1) and (2). \square

By this lemma we can assume that p does not give rise to any deterministic simplex (edge or triangle). Otherwise, we can always take a quotient by the simplicial subset Z_p and use the formula in Lemma 3.9. If $\partial\sigma$ contains a degenerate edge x' then $p(s_\sigma^{ab}) = 0$ for precisely two of the pairs $(a, b) \in \mathbb{Z}_2^2$. Note that these pairs are of the form (a, b) and $(a+1, b+1)$. We remove one of these vertices for each such simplex keeping only s_σ^{ab} with $a = 0$.

Definition 3.10. Let $p \in \mathbf{sDist}_\beta(X)$ be a twisted distribution with no deterministic simplices (edges or triangles). Let $\Gamma_\beta^0(X, p)$ denote the induced signed subgraph of $\Gamma_\beta(X, p)$ where only the vertex s_σ^{0b} is kept when σ has a degenerate 1-simplex in its boundary together with the sign defined by

$$\gamma_p(x, \sigma) = \gamma(x, s_\sigma^{ab}) \quad (15)$$

where s_σ^{ab} is the unique vertex with $p(s_\sigma^{ab}) = 0$ if $|\partial\sigma^\circ| = 3$, or s_σ^{0b} if $|\partial\sigma^\circ| = 2$.

The sign γ_p satisfies

$$\prod_{x \in \partial\sigma^\circ} \gamma_p(x, \sigma) = \begin{cases} (-1)^{\beta(\sigma)} & |\partial\sigma^\circ| = 3 \\ (-1)^{a+\beta(\sigma)+1} & |\partial\sigma^\circ| = 2, x' = d_1\sigma \\ (-1)^{a+1} & |\partial\sigma^\circ| = 2, x' = d_0\sigma, \text{ or } d_2\sigma \end{cases} \quad (16)$$

where x' is the unique degenerate edge in the case $|\partial\sigma^\circ| = 2$.

Proposition 3.11. *Let $p \in \text{sDist}_\beta(X)$ be a twisted distribution with no deterministic simplices. Then*

$$\text{rank}(p) = \text{rank}(B(\Gamma_\beta^0(X, p))).$$

The choice we made in the definition of the graph does not affect the rank.



Figure 7: (a) The graph $\Gamma_{\bar{\beta}}(\bar{X}, \bar{p})$ where p is a PR box. (b) The subgraph $\Gamma_{\bar{\beta}}^0(\bar{X}, \bar{p})$.

3.3 The rank formula

We recall some basic definitions about signed graphs following [23]. Let $\Sigma = (V, E)$ be a simple (undirected) graph. A path is a sequence of edges $e_1 e_2 \cdots e_k$ with no repetition such that e_{i-1} and e_i has a common vertex. If v and w are the vertices of e_1 and e_k not common to e_2 and e_{k-1} , respectively, then we say the path is from v to w . A path is called closed if $v = w$. A circle is the graph determined by a closed path. If Σ comes with a sign $\gamma : E \rightarrow \{\pm 1\}$ then a circle C given by the sequence $e_1 e_2 \cdots e_k$ has sign

$$\gamma(C) = \gamma(e_1)\gamma(e_2) \cdots \gamma(e_k).$$

If $\gamma(C) = 1$ ($\gamma(C) = -1$) the circle is called positive (negative).

Definition 3.12. A signed graph Σ is called balanced if every circle in it is positive. We will write $b(\Sigma)$ for the number of components of the graph that are balanced.

A bidirected graph comes with a choice of sign $\eta(v, e) = \pm 1$ for each v incident to an edge e . Given a signed graph (Σ, γ) we can define a bidirected graph (Σ, η) such that

$$\gamma(e) = -\eta(v, e)\eta(w, e)$$

for every edge $e = \{v, w\}$. This bidirected graph is not unique. An incidence matrix of (Σ, γ) is a $|V| \times |E|$ -matrix $H(\Sigma)$ such that

$$H(\Sigma)_{v,e} = \begin{cases} \eta(v, e) & v \in e \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 3.13. $H(\Sigma)$ has rank $|V| - b(\Sigma)$.

Proof. See [23, Theorem IV.1]. □

We will apply this rank result to bipartite graphs.

Definition 3.14. Let Σ be a signed bipartite graph with vertex set $V(\Sigma) = V^0(\Sigma) \sqcup V^1(\Sigma)$. Suppose that every vertex in $V^0(\Sigma)$ is incident to exactly two edges. Define a new bidirected graph $(\hat{\Sigma}, \eta)$ with

- vertex set $V^0(\Sigma)$,
- edge set $V^1(\Sigma)$, and
- bidirection defined by

$$\eta(v, e) = \gamma(v, e)$$

where $v \in V^0(\Sigma)$ and $e \in V^1(\Sigma)$ are connected by an edge in Σ .

The vertices incident to an edge in $V^1(\Sigma)$ are those vertices in $V^0(\Sigma)$ connected to it by an edge in Σ . Effectively, to obtain $\hat{\Sigma}$ we merge any two edges incident to a vertex in $V^0(\Sigma)$ into a single edge. Then using Theorem 3.13 we have

$$\text{rank}(B(\Sigma)) = \text{rank}(H(\hat{\Sigma})) = |V^0(\Sigma)| - b(\hat{\Sigma}). \quad (17)$$

Note that the rank does not depend on the choice of η since a different choice would amount to multiplying the corresponding row with -1 . If $\Gamma_\beta^0(X, p)$ satisfies the assumption of Definition 3.14 we can apply the construction $\Sigma \mapsto \hat{\Sigma}$ to the graph $\Gamma_\beta^0(X, p)$. We define

$$b(X, p) = b(\hat{\Gamma}_\beta^0(X, p)). \quad (18)$$

Theorem 3.15. *Let X be a simplicial set generated by 2-simplices $\sigma_1, \sigma_2, \dots, \sigma_N$ such that each $\partial\sigma_i$ consists of either three distinct non-degenerate 1-simplices or two distinct non-degenerate 1-simplices and a remaining degenerate 1-simplex. Consider a twisted distribution $p \in \text{sDist}_\beta(X)$ satisfying the following conditions:*

- for each generating 2-simplex σ , $p_\sigma^{ab} = 0$ for at least one pair $(a, b) \in \mathbb{Z}_2^2$, and
- every non-degenerate 1-simplex of $\bar{X} = X/Z_p$ belongs precisely to two generating 2-simplices.

Then we have

$$\text{rank}(p) = |(Z_p)_1^\circ| + |\bar{X}_2^\circ| - b(\bar{X}, \bar{p}).$$

Proof. By Lemma 3.9, Proposition 3.11, and Equation (17) we have

$$\text{rank}(p) = |(Z_p)_1^\circ| + \text{rank}(\Gamma_\beta^0(\bar{X}, \bar{p})) = |(Z_p)_1^\circ| + |X_2^\circ| - b(X, p).$$

The assumption on p implies that the number of the vertices of $\hat{\Gamma}_\beta^0(\bar{X}, \bar{p})$ is given by $|X_2^\circ|$. \square

Note that if the condition $p_\sigma^{ab} = 0$ fails for at least one pair of outcomes for every non-degenerate simplex then to compute the rank we can restrict to the simplicial subset for which this condition holds.

4 Examples

The rank formula in Theorem 3.15 is very useful in finding the vertices of the polytope of twisted distributions. We can partition $\text{sDist}_\beta(X)$ by fixing a simplicial subset $Z \subset X$ and considering those twisted distributions p with the property that $Z_p = Z$. Our approach will be to combine this observation with the following result.

Lemma 4.1. *Let $p \in \text{sDist}_\beta(X)$ be a vertex. Then*

$$|(Z_p)_1^\circ| - |(Z_p)_2^\circ| \geq |X_1^\circ| - |X_2^\circ|.$$

Proof. By Corollary 3.7 p is a vertex if and only if $\text{rank}(p) = |X_1^\circ|$. We have $\text{rank}(\bar{p}) \leq |X_2^\circ| - |(Z_p)_2^\circ|$ since Z_p is 1-dimensional. Then the result follows from Lemma 3.9. \square

The rank formula contains the number of balanced components of the associated graph. For the computation of the signs of the circle graphs we will use the following formula. See Figure (8).

Lemma 4.2. *Let $p \in \text{sDist}_\beta(X)$ and consider γ_p defined in Equation (15). For a non-degenerate 2-simplex σ whose boundary $\partial\sigma = \{x, y, z\}$ contains a deterministic edge, say z , with $p_z = \delta^a$ we have*

$$\gamma_p(x, \sigma)\gamma_p(y, \sigma) = (-1)^{a+\beta(\sigma)+1}.$$

Proof. The distribution $p_\sigma \in D(\mathbb{Z}_2^2)$ is zero if and only if (c, d) satisfies $c + d = a + \beta(\sigma) + 1$. We have $\gamma(x, \sigma) = (-1)^c$ and $\gamma(y, \sigma) = (-1)^d$. Hence the formula follows. \square




Figure 8

4.1 N -cycle scenario

For $N \geq 2$, let \tilde{C}_N denote the measurement space of the N -cycle scenario defined as the following simplicial set:

- Generating 2-simplices: $\sigma_1, \dots, \sigma_N$.
- Identifying relations:

$$d_{i'_1} \sigma_1 = d_{i_2} \sigma_2, d_{i'_2} \sigma_2 = d_{i_3} \sigma_3 \cdots d_{i'_N} \sigma_N = d_{i_1} \sigma_1$$

where $i_k \neq i'_k \in \{0, 1\}$ for $1 \leq k \leq N$.

The case $N = 4$ where the boundary is oriented in the counter-clockwise direction is depicted in Figure (3a). See also [19, Definition 4.1]. The edges on the boundary will be denoted by x_1, x_2, \dots, x_N .



Figure 9: (a) Deterministic interior edge. (b) Collapsed scenario.

Let $p \in \text{sDist}_\beta(\tilde{C}_N)$ be a vertex. Let us consider the possibilities for Z_p .

1. Z_p has a deterministic edge connecting a boundary vertex to the central vertex: In this case the quotient \tilde{C}_N/Z_p can be identified with a wedge sum of D_{K_i} 's of Example 2.15 where $K_i \leq N$. We know that every twisted distribution on D_{K_i} is non-contextual. Then Corollary 2.21 implies that p is non-contextual, hence a deterministic distribution. The situation is depicted in Figure (9).
2. All deterministic edges of Z_p are on the boundary: Note that by the previous case we know that p would be a deterministic distribution if Z_p contained a deterministic triangle. Assuming that Z_p does not contain a deterministic triangle, Lemma 4.1 gives

$$|(Z_p)^\circ_1| \geq |X_1^\circ| - |X_2^\circ| = 2N - N = N.$$

This forces Z_p to be the boundary $\partial\tilde{C}_N$ of the disk. Assume that $p|_{\partial\tilde{C}_N} = \delta^s$. Then we have

$$\delta_{x_i}^s = \delta^{a_i}, \quad i = 1, 2, \dots, N$$

for some $a_i \in \mathbb{Z}_2$. The graph $\hat{\Gamma}_\beta^0(\bar{X}, \bar{p})$ is given by a circle C with edges e_1, e_2, \dots, e_N as in Figure (10a). The sign of the circle is given by

$$\gamma_{\bar{p}}(C) = \prod_{i=1}^N \gamma_{\bar{p}}(e_i) = (-1)^N \prod_{i=1}^N (-1)^{a_i + \beta(\sigma_i) + 1} = (-1)^N \prod_{i=1}^N (-1)^{\beta(\sigma_i) + 1} = (-1)^{[\beta]}$$

and by the rank formula we obtain

$$\text{rank}(p) = \begin{cases} 2N & [\bar{\beta}] = 1 \\ 2N - 1 & [\bar{\beta}] = 0. \end{cases} \quad (19)$$

Therefore p is a vertex if and only if $[\bar{\beta}] = 1$, that is, $[\beta] = 1 + \sum_{i=1}^N a_i$.

As a special case let us take $[\beta] = 0$. In this case p is a vertex if and only if $\sum_{i=1}^N a_i = 1$. When $N = 4$ this reproduces the PR boxes in Example 2.17.



Figure 10: (a) The graph $\hat{\Gamma}_{\bar{\beta}}^0(\tilde{C}_4/\partial\tilde{C}_4)$. (b) When the boundary is collapsed the resulting space is a sphere.

4.2 Boundary of tetrahedron

Let $\partial\Delta^3$ denote the simplicial set given by the boundary of a tetrahedron with generating 2-simplices $\sigma_1, \sigma_2, \sigma_3, \sigma_4$. Let $p \in \text{sDist}_{\beta}(\partial\Delta^3)$ be a vertex. Then Lemma 4.1 gives

$$|(Z_p)_1^{\circ}| - |(Z_p)_2^{\circ}| \geq 4 - 2 = 2$$

and we have $|(Z_p)_2^{\circ}| = 0, 1$ or 4 . The last case gives a deterministic distribution if $[\beta] = 0$. Let us consider the remaining cases.

1. $|(Z_p)_2^{\circ}| = 0$: In this case $\hat{\Gamma}_{\bar{\beta}}^0(\bar{X}, \bar{p})$ is a circle C with edges e_1, e_2, e_3, e_4 ; see Figure (11a). Its sign is given by

$$\gamma_{\bar{p}}(C) = \prod_{i=1}^4 \gamma_{\bar{p}}(e_i) = (-1)^{a+\beta(\sigma_1)+1} (-1)^{b+\beta(\sigma_2)+1} (-1)^{a+\beta(\sigma_3)+1} (-1)^{b+\beta(\sigma_4)+1} = (-1)^{[\beta]}$$

and by the rank formula

$$\text{rank}(p) = \begin{cases} 6 & [\beta] = 1 \\ 5 & [\beta] = 0. \end{cases} \quad (20)$$

Therefore p is a vertex if and only if $[\beta] = 1$.

2. $|(Z_p)_2^{\circ}| = 1$: This case (Figure (11b)) reduces to the cycle scenario \tilde{C}_3 where $p|_{\partial\tilde{C}_3}$ is deterministic with $a + b + c = 0$. Therefore by Equation (19), p is a vertex if and only if $[\beta] = 1$.

In the case $[\beta] = 0$ we observe that all the vertices are deterministic, which provides an alternative approach to the computation in [1, Proposition 4.12].

4.3 Mermin torus

Let T denote the Mermin torus introduced in [13]. Let $p \in \text{sDist}_{\beta}(T)$ be a vertex. The distribution p is deterministic if and only if $[\beta] = 0$. Let us assume p is not deterministic. Lemma 4.1 gives

$$|(Z_p)_1^{\circ}| - |(Z_p)_2^{\circ}| \geq 3$$



Figure 11

which can only be satisfied if Z_p consists of three generating 1-simplices as in Figure (12a) or two generating triangles as in Figure (12b). In both cases $\hat{\Gamma}_\beta^0(\bar{X}, \bar{p})$ is a circle C . We compute the sign of this circle for each case:

1. In Figure (12a) we have

$$\gamma_{\bar{p}}(C) = (-1)^{a+\beta(\sigma_1)+1}(-1)^{b+\beta(\sigma_2)+1}(-1)^{c+\beta(\sigma_3)+1}(-1)^{a+\beta(\sigma_4)+1}(-1)^{b+\beta(\sigma_5)+1}(-1)^{c+\beta(\sigma_6)+1} = (-1)^{[\beta]}.$$

Therefore p is a vertex if and only if $[\beta] = 1$.

2. In Figure (12b) we have

$$\gamma_{\bar{p}}(C) = (-1)^{a+\beta(\sigma_1)+1}(-1)^{b+\beta(\sigma_3)+1}(-1)^{a+b+c+\beta(\sigma_2)+\beta(\sigma_5)+\beta(\sigma_4)+1}(-1)^{c+\beta(\sigma_6)+1} = (-1)^{[\beta]}$$

and p is a vertex if and only if $[\beta] = 1$.

Combining these two observations gives the main result of [13]. $\text{sDist}_\beta(T)$ has only deterministic vertices if $[\beta] = 0$. On the other hand, if $[\beta] = 1$ then there are two kinds of vertices as given in Figure (12).



Figure 12

References

- [1] C. Okay, A. Kharoof, and S. Ipek, “Simplicial quantum contextuality,” *Quantum*, vol. 7, 2023. doi: 10.22331/q-2023-05-22-1009.
- [2] J. S. Bell, “On the Einstein Podolsky Rosen paradox,” *Physics Physique Fizika*, vol. 1, pp. 195–200, Nov 1964. doi: 10.1103/PhysicsPhysiqueFizika.1.195.

- [3] S. Kochen and E. P. Specker, “The problem of hidden variables in quantum mechanics,” *Journal of Mathematics and Mechanics*, vol. 17, pp. 59–87, 1967. url: <https://www.jstor.org/stable/24902153>.
- [4] P. G. Goerss and J. F. Jardine, *Simplicial homotopy theory*. Springer Science & Business Media, 2009.
- [5] S. Abramsky and A. Brandenburger, “The sheaf-theoretic structure of non-locality and contextuality,” *New Journal of Physics*, vol. 13, no. 11, p. 113036, 2011. doi: 10.1088/1367-2630/13/11/113036. arXiv: 1102.0264.
- [6] I. Pitowsky, *Quantum Probability Quantum Logic*. Springer, 1989.
- [7] S. Popescu and D. Rohrlich, “Quantum nonlocality as an axiom,” *Foundations of Physics*, vol. 24, no. 3, pp. 379–385, 1994.
- [8] J. Barrett, N. Linden, S. Massar, S. Pironio, S. Popescu, and D. Roberts, “Nonlocal correlations as an information-theoretic resource,” *Physical Review A*, vol. 71, p. 022101, Feb 2005. doi: 10.1103/PhysRevA.71.022101. arXiv: quant-ph/0404097.
- [9] N. S. Jones and L. Masanes, “Interconversion of nonlocal correlations,” *Physical Review A*, vol. 72, no. 5, p. 052312, 2005.
- [10] C. Horne, M. Horne, A. Shimony, and H. Richard, “Proposed experiment to test local hidden-variable theories,” *Physical Review Letters*, p. 880, 1969.
- [11] N. D. Mermin, “Hidden variables and the two theorems of John Bell,” *Reviews of Modern Physics*, vol. 65, no. 3, p. 803, 1993. doi: 10.1103/RevModPhys.65.803.
- [12] C. Okay, S. Roberts, S. D. Bartlett, and R. Raussendorf, “Topological proofs of contextuality in quantum mechanics,” *Quantum Information & Computation*, vol. 17, no. 13-14, pp. 1135–1166, 2017. doi: 10.26421/QIC17.13-14-5. arXiv: 1701.01888.
- [13] C. Okay, H. Y. Chung, and S. Ipek, “Mermin polytopes in quantum computation and foundations,” *Quantum Information & Computation*, vol. 23, no. 9/10, pp. 733–782, 2023. doi: 10.26421/QIC23.9-10-2. arXiv: 2210.10186.
- [14] M. Zurel, C. Okay, and R. Raussendorf, “Hidden variable model for universal quantum computation with magic states on qubits,” *Physical Review Letters*, vol. 125, no. 26, p. 260404, 2020.
- [15] S. Ipek and C. Okay, “The degenerate vertices of the 2-qubit A-polytope and their update rules,” *arXiv preprint arXiv:2312.10734*, 2023.
- [16] R. S. Barbosa, A. Kharoof, and C. Okay, “A bundle perspective on contextuality: Empirical models and simplicial distributions on bundle scenarios,” *arXiv preprint arXiv:2308.06336*, 2023. doi: 10.48550/arXiv.2308.06336.
- [17] G. Friedman, “An elementary illustrated introduction to simplicial sets,” *arXiv preprint arXiv:0809.4221*, 2008. doi: 10.48550/arXiv.0809.4221.
- [18] J. P. May, *Simplicial objects in algebraic topology*. Van Nostrand Mathematical Studies, No. 11, D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1967.
- [19] A. Kharoof, S. Ipek, and C. Okay, “Topological methods for studying contextuality: N-cycle scenarios and beyond,” *Entropy*, vol. 25, no. 8, 2023. doi: 10.3390/e25081127.
- [20] A. Kharoof and C. Okay, “Simplicial distributions, convex categories and contextuality,” *preprint arXiv:2211.00571*, 2022. doi: 10.48550/arXiv.2211.00571.
- [21] C. A. Weibel, *An introduction to homological algebra*. No. 38, Cambridge university press, 1995. doi: 10.1007/b98977.

- [22] V. Chvátal, *Linear programming*. A Series of Books in the Mathematical Sciences, W. H. Freeman and Company, New York, 1983.
- [23] T. Zaslavsky, “Matrices in the theory of signed simple graphs,” in *Advances in discrete mathematics and applications: Mysore, 2008*, vol. 13 of *Ramanujan Math. Soc. Lect. Notes Ser.*, pp. 207–229, Ramanujan Math. Soc., Mysore, 2010.