

Comment on "Shot noise in a strange metal"

V. R. Shaginyan,^{1,2,*} A. Z. Msezane,² and G. S. Japaridze²

¹*Petersburg Nuclear Physics Institute, Gatchina, 188300, Russia*

²*Clark Atlanta University, Atlanta, GA 30314, USA*

The recent paper [1] is devoted to measurements of shot noise to probe excitations in nanowires of the heavy fermion (HF) metal YbRh₂Si₂. The authors observed that shot noise is strongly suppressed, and claim that the suppression cannot be attributed to either electron-phonon or electron-electron interactions in a Fermi liquid. Their observation suggests that the current is not carried by well-defined quasiparticles in the YbRh₂Si₂, and calls for similar research into other strange metals.

In this comment, we show that it is unlikely that the affected carriers in bulk YbRh₂Si₂ would have undergone any fragmentation. Rather, it is the electron-phonon interaction that suppresses noise. Indeed, experimental observations unambiguously show that the Wiedemann-Franz law holds in YbRh₂Si₂, implying that no fundamental breakdown of quasiparticle behavior occurs in the archetypical HF metal [2]. Moreover, detailed study of the temperature evolution of quasiparticles in the HF metal Sr₂RuO₄ demonstrates that quasiparticles persist up to temperatures above 200 K, far beyond the Fermi liquid regime [3].

It was shown that HF metals can exhibit a quasiclassical behavior that remains applicable to the description of the resistivity of HF metals due to the presence of a transverse zero-sound collective mode, reminiscent of the phonon mode in solids. It is demonstrated that at temperatures T , in excess of an extremely low Debye temperature $T_D \lesssim 1$ K, the resistivity $\rho(T) \propto T$, since the mechanism, forming this dependence, is the same as the electron-phonon mechanism that prevails at high temperatures in ordinary metals [4, 5]. The same mechanism allows one to explain the optical conductivity in YbRh₂Si₂ [6], and suppressing the shot noise in HF metals. It is worth noting that the current-carrying excitations in

nanowires of YbRh₂Si₂ can be different from the excitation of the bulk YbRh₂Si₂, since the nanowires can be represented by one dimensional HF metal that possesses special properties, see e.g. [7].

We conclude that there is no reason to assume the fragmentation of charge carriers, and we can refer to William Shakespeare's immortal comedy that the paper [1] is "Much Ado About Nothing."

* Electronic address: vrshag@thd.pnpi.spb.ru

- [1] L. Chen, D. T. Lowder, E. Bakali, A. M. Andrews, W. Schrenk, *et al.*, Shot noise in a strange metal. *Science* **382**, 907 (2023).
- [2] J.-Ph. Reid, M. A. Tanatar, R. Daou, R. Hu, C. Petrovic, and L. Taillefer, Wiedemann-Franz law and nonvanishing temperature scale across the field-tuned quantum critical point of YbRh₂Si₂. *Phys. Rev. B* **89**, 045130 (2014).
- [3] A. Hunter, S. Beck, E. Cappelli, F. Margot, M. Straub, Y. Alexanian *et al.*, Fate of Quasiparticles at High Temperature in the Correlated Metal Sr₂RuO₄. *Phys. Rev. Lett.* **131**, 236502 (2023).
- [4] V. R. Shaginyan, K. G. Popov, and V. A. Khodel, Quasiclassical physics and T -linear resistivity in both strongly correlated and ordinary metals. *Phys. Rev. B* **88**, 115103 (2013).
- [5] V.R. Shaginyan, A.Z. Msezane, and G.S. Japaridze, Peculiar Physics of Heavy-Fermion Metals: Theory versus Experiment. *Atoms* **10**, 67 (2022).
- [6] V. R. Shaginyan, A. Z. Msezane, M. V. Zverev, Transport properties of strongly correlated Fermi systems. *Symmetry* **15**, 2055 (2023).
- [7] V.A. Khodel, V.R. Shaginyan, and V.V. Khodel, New approach in the microscopic Fermi system theory. *Phys. Rep.* **249**, 1 (1994).