THE ATIYAH CLASS OF DG MANIFOLDS OF AMPLITUDE +1

SEOKBONG SEOL

ABSTRACT. A dg manifold of amplitude +1 in C^{∞} -context can be thought of as a derived intersection of a section s and the zero section of a vector bundle E. In this paper, we compute the Atiyah class of dg manifolds of amplitude +1. In particular, we prove that the Atiyah class of a dg manifold of amplitude +1 vanishes if and only if the intersection of s and the zero section is a clean intersection.

As an example, given two submanifolds X and Y of W, we investigate the Atiyah class of a dg manifold which can be thought of as a model for the derived intersection of X and Y in the sense of Behrend–Liao–Xu.

CONTENTS

1. Introduction	1
Notations and conventions	3
2. Backgrounds on dg manifolds	4
3. The Atiyah class and clean intersection	5
3.1. Clean intersection	6
3.2. Locality of the Atiyah class in positive amplitude	8
3.3. The Atiyah class of dg manifolds of amplitude $+1$	11
4. Proof of Theorem 3.2	16
4.1. Proof of $(L1) \Rightarrow (L2)$: sample cases	16
4.2. Proof of $(L1) \Rightarrow (L2)$: general case	19
5. The case of derived intersections	22
Acknowledgement	23
References	23

1. INTRODUCTION

The present paper is devoted to studying the Atiyah class of dg manifolds of amplitude +1. The notion of dg manifolds (a.k.a. Q-manifolds) is a generalisation of the notion of smooth manifolds, in a way that the algebra of smooth functions is enriched to include graded commutative elements. Dg manifolds have appeared in mathematical physics literature, in relation to BRST quantisation and AKSZ formalism [1, 7]. They naturally arise in various fields of mathematics, such as Lie theory, differential geometry and homotopy theory. Moreover, they are closely related to the emerging field of derived differential geometry [3, 4, 6].

Recall that a dg manifold is a \mathbb{Z} -graded manifold \mathcal{M} , equipped with a homological vector field, i.e. a degree +1 derivation Q of $C^{\infty}(\mathcal{M})$ satisfying $[Q, Q] = 2Q^2 = 0$. An interesting example is a dg manifold of amplitude +1, which is essentially a vector bundle E equipped with a section $s \in \Gamma(E)$: the graded algebra

The author is supported by the KIAS Individual Grant MG090801 at Korea Institute for Advanced Study.

of functions is $\Gamma(\Lambda^{-\bullet}E^{\vee})$ and the homological vector field is the interior product ι_s . A dg manifold of amplitude +1 can be thought of as a model for a derived intersection of the section s and the zero section of E.

In studying dg manifolds, the Atiyah class takes an important role. Originally, Atiyah [2] introduced the Atiyah class of holomorphic vector bundles as an obstruction to the existence of holomorphic connections. Later, the notion of the Atiyah class was generalised to dg manifolds. The Atiyah class of dg manifolds was first introduced by Shoikhet [21] in terms of Lie algebra cohomology and 1-jets of tangent bundles. It also appeared in the work of Lyakhovich–Mosman–Sharapov [16] (in their notation, it is B_1) and was studied systemically by Mehta–Stiénon–Xu [18]. It is a key ingredient in the formality theorem and Duflo–Kontsevich type theorem for dg manifolds [15, 21, 13], an extension of the famous work of Kontsevich [11]. It also appears in the construction of Kapranov's L_{∞} algebra for dg manifolds [20], which generalises the celebrated work of Kapranov [10]. See also [23]. Following [18], we recall the definition of the Atiyah class of a dg manifold below.

Let (\mathcal{M}, Q) be a dg manifold. The space of (1, 2)-tensors $\Gamma(\mathcal{M}; T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))$ equipped with the Lie derivative \mathcal{Q} along the homological vector field Q forms a cochain complex. Given an affine connection ∇ on \mathcal{M} , the (1, 2)-tensor $\operatorname{At}^{\nabla}$ of degree +1 defined by

$$At^{\nabla}(X,Y) = [Q, \nabla_X Y] - \nabla_{[Q,X]} Y - (-1)^{|X|} \nabla_X [Q,Y]$$

for $X, Y \in \mathfrak{X}(\mathcal{M})$, is a cocycle in the cochain complex $(\Gamma(\mathcal{M}; T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M})), \mathcal{Q})$. The element $\operatorname{At}^{\nabla}$ is called the Atiyah cocycle associated with ∇ and its cohomology class

$$\alpha_{(\mathcal{M},Q)} := [\operatorname{At}^{\nabla}] \in H^1(\Gamma(\mathcal{M}; T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M})), \mathcal{Q})$$

is independent of choice of connection ∇ . The cohomology class $\alpha_{(\mathcal{M},Q)}$ is called the Atiyah class of the dg manifold (\mathcal{M},Q) .

The Atiyah class of a dg manifold (\mathcal{M}, Q) is an obstruction to the existence of an affine connection compatible with the homological vector field Q. In particular, it was shown by Chen–Xiang–Xu [8] (see also [12]) that the Atiyah class of the dg manifold arising from a complex manifold is the Atiyah class of the complex manifold, introduced by Atiyah [2].

Naturally, one can ask the following question.

Question A. What is the Atiyah class of dg manifolds of amplitude +1?

This paper attempts to answer this question. Before we present our main theorem, we need the notion of clean intersection [5].

Given two embedded submanifolds X and Y of W, we say X intersects with Y cleanly if the intersection $Z := X \cap Y$ is a smooth manifold and $T_pX \cap T_pY = T_pZ$ for each $p \in Z$. It can be checked (see for instance [9, Proposition C.3.1]) that the embedded submanifolds X and Y intersect cleanly in W if and only if for each intersection point $p \in X \cap Y$, there exists an open neighbourhood U of p and a chart $\phi : U \to \mathbb{R}^{\dim W}$ such that $\phi(X \cap U)$ and $\phi(Y \cap U)$ are vector spaces in $\mathbb{R}^{\dim W}$. Thus, around each intersection point $p \in X \cap Y$, there is a submanifold $W' \subset W$ near p such that X and Y intersects transversally in W'. Naively speaking, the space obtained by clean intersection is as good as the space obtained by transversal intersection.

An answer to Question A is our main theorem.

Theorem B (Theorem 3.2). Let *E* be a vector bundle and let *s* be a section on *E*. The Atiyah class of a dg manifold $(E[-1], \iota_s)$ vanishes if and only if the intersection of *s* and the zero section 0_E is clean.

The proof of Thoerem B is based on the following observation: the Atiyah class of a dg manifold of positive amplitude is a local invariant (Proposition 3.9). More precisely, let (\mathcal{M}, Q) be a dg manifold of positive amplitude whose base manifold is M. For each open subset $U \subset M$, let (\mathcal{M}_U, Q_U) denote the dg submanifold of \mathcal{M} obtained by restricting the base manifold M to open subset U. Under this notation, we proved that the

Atiyah class of (\mathcal{M}, Q) vanishes if and only if there exists an open cover $\{U_i\}_{i \in I}$ of M such that the Atiyah class of $(\mathcal{M}_{U_i}, Q_{U_i})$ vanishes for all $i \in I$.

In general, such locality of the Atiyah class is not expected. For instance, the Atiyah class of a dg manifold arising from a complex manifold is an obstruction to the existence of holomorphic connection, but holomorphic connections always exist locally. In other words, if (\mathcal{M}, Q) is the dg manifold arising from a complex manifold X, then for each $p \in X$, there exists a small neighbourhood $U \subset X$ of p such that the Atiyah class of (\mathcal{M}_U, Q_U) vanishes, whilst, for many complex manifolds X, the Atiyah class of (\mathcal{M}, Q) does not vanish. In fact, the property which enables such locality follows from the linearity of the homological vector field over smooth functions on the base manifold. For arbitrary dg manifold (\mathcal{M}, Q) with base manifold M, the homological vector field Q is not necessarily $C^{\infty}(M)$ -linear. However, if \mathcal{M} is of positive amplitude, then by degree reason, the homological vector field Q is always $C^{\infty}(M)$ -linear.

Going back to the dg manifold $(E[-1], \iota_s)$ of amplitude +1, it suffices to consider the Atiyah class of $(E|_U[-1], \iota_{s|_U})$ for a small neighbourhood of each point $p \in M$. Under this setting, we explicitly compute an Atiyah cocycle and the coboundary operator in a local coordinate. Finally, we prove that the Atiyah class of $(E|_U[-1], \iota_{s|_U})$ vanishes for some open neighbourhood U of p if and only if the section s intersects cleanly along the zero section 0_E in an open neighbourhood of p.

As an example, we investigate the Atiyah class of a dg manifold $(\mathcal{M}_{X,Y}, Q)$ of amplitude +1 arising from embedded submanifolds X and Y of W. In fact, this dg manifold $(\mathcal{M}_{X,Y}, Q)$ is a model for the derived intersection $X \cap^h Y$ of X and Y, in the sense of [3]. We proved that the Atiyah class of $(\mathcal{M}_{X,Y}, Q)$ vanishes if and only if X and Y intersect cleanly.

The work of Behrend–Liao–Xu [3] addresses the category of dg manifolds of positive amplitude, rather than dg manifolds of amplitude +1. Thus, it would be interesting to extend our work to the Atiyah class of dg manifolds of positive amplitude. Moreover, they introduced a notion of weak-equivalence between dg manifolds of amplitude +1 (in fact, weak-equivalence in the category of positive amplitude) so that a quasi-smooth derived manifold can be thought of as a dg manifold of amplitude +1 (and more generally, the Atiyah class of dg manifolds of positive amplitude) is invariant under weak-equivalence.

Notations and conventions. Throughout this paper, the base field is the field of real numbers \mathbb{R} : vector spaces, manifolds, vector bundles, functions in this paper are over \mathbb{R} .

For any smooth function $f : \mathbb{R}^n \to \mathbb{R}$, we write $f = f(x_1, \dots, x_n)$. The notation x^i denotes the coordinate function $x^i : \mathbb{R}^n \to \mathbb{R}$ defined by $x^i(x_1, \dots, x_n) = x_i$.

We reserve the symbol M for a manifold exclusively. By a manifold, we mean a smooth manifold without boundary with Hausdorff and second countability properties. The sheaf of smooth functions on M is denoted by \mathcal{O}_M . The algebra of globally defined smooth functions on M is $C^{\infty}(M) = \mathcal{O}_M(M)$.

All gradings in this paper are \mathbb{Z} -gradings and the symbols \mathcal{M} is always a finite-dimensional graded manifold. The abbreviation 'dg' stands for 'differential graded'.

Let R be a graded ring. Given any element v in a graded R-module $V = \bigoplus_{k \in \mathbb{Z}} V^k$, the symbol |v| = dmeans that v is a homogeneous element of degree d, or equivalently $v \in V^d$. Whenever the symbol |v|appears, we assume that v is a homogeneous element. The suspension of the graded R-module V is V[1]whose homogeneous component of degree k is $(V[1])^k = V^{k+1}$. Given any graded vector bundle \mathcal{E} , we use the symbol $S(\mathcal{E})$ to denote the bundle of graded symmetric tensors of \mathcal{E} .

Let $E \to M$ be a vector bundle. By abuse of notation, a section $s \in \Gamma(E)$ is often identified with $s \in \Gamma(E[-1])$. Moreover, a *TM*-connection $\nabla : \Gamma(TM) \times \Gamma(E) \to \Gamma(E)$ on *E* is often identified with a *TM*-connection $\nabla : \Gamma(TM) \times \Gamma(E[-1]) \to \Gamma(E[-1])$ on E[-1].

SEOKBONG SEOL

2. BACKGROUNDS ON DG MANIFOLDS

In this section, we present some background materials on the Atiyah class of a dg manifold. Readers are invited to [17, 18] for more detail.

Let M be a smooth manifold and let \mathcal{O}_M be its sheaf of ring of smooth functions. A **graded manifold** \mathcal{M} with base M is a sheaf of graded commutative \mathcal{O}_M -algebra \mathcal{A} on M such that there exists a \mathbb{Z} -graded vector space V and for each $p \in M$, there exists an open neighbourhood $U \subset M$ of p satisfying

$$\mathcal{A}(U) \cong \mathcal{O}_M(U) \otimes S(V^{\vee})$$

where $S(V^{\vee})$ is the graded symmetric tensor of V^{\vee} . The space of global sections of the sheaf \mathcal{A} will often be denoted by $C^{\infty}(\mathcal{M})$.

We say a graded manifold \mathcal{M} is of amplitude [n, m], for $n \leq m$, if the graded vector space V is of the form

$$V = \bigoplus_{i=n}^{m} V_i$$

where the graded vector space V_i consists of vectors of degree *i*. If $0 < n \leq m$, then we say that \mathcal{M} is of positive amplitude. The amplitude [n, n] will be simply denoted by amplitude *n*.

A graded manifold \mathcal{M} is called finite dimensional if both $\dim M < \infty$ and $\dim V < \infty$. All graded manifolds in this paper will be finite dimensional.

Remark 2.1. In literature, such as [18, 20, 23], the sheaf of \mathcal{O}_M -algebra \mathcal{A} is defined using formal power series on V rather than polynomial functions. However, all results in this paper are valid for both versions.

Example 2.2. Let M be a finite dimensional smooth manifold and for each $i \in \mathbb{Z}$, let $E_i \to M$ be a vector bundle of finite rank over M. By denoting $E_i[-i] \to M$, we mean a graded vector bundle such that the space of sections $\Gamma(E_i[-i]) = \Gamma(E_i)[-i]$ is a graded $C^{\infty}(M)$ -module concentrated in degree i. The graded vector bundle $E = \bigoplus_i E_i[-i]$ forms a graded manifold \mathcal{M} : the algebra of functions is $\mathcal{A}(U) = \Gamma(U; S(E^{\vee}))$ where $S(E^{\vee})$ is the graded symmetric tensor of E^{\vee} . If $\operatorname{rank}(E_i) = 0$ for all but $n \leq i \leq m$, then the amplitude of the graded manifold \mathcal{M} is [n, m]. If such $n, m \in \mathbb{Z}$ exists, then \mathcal{M} is finite dimensional.

A graded vector bundle $\pi : \mathcal{E} \to \mathcal{M}$ is a vector bundle object in the category of graded manifolds. In terms of sheaves, a graded vector bundle is a sheaf of locally free graded \mathcal{A} -module on M. A section $s : \mathcal{M} \to \mathcal{E}$ is a morphism of graded manifolds such that $\pi \circ s = id_{\mathcal{M}}$. The $C^{\infty}(\mathcal{M})$ -module of all sections of \mathcal{E} over \mathcal{M} is denoted by $\Gamma(\mathcal{M}; \mathcal{E}) = \Gamma(\mathcal{E})$.

An important example of a graded vector bundle over \mathcal{M} is the **tangent bundle** $T\mathcal{M}$. A section of $T\mathcal{M}$ is called a vector field, and the space of vector fields $\Gamma(T\mathcal{M})$, often denoted as $\mathfrak{X}(\mathcal{M})$, is identified with the space of graded derivations $Der(C^{\infty}(\mathcal{M}))$. The graded derivations $Der(C^{\infty}(\mathcal{M}))$ has a Lie algebra structure, defined as a graded commutator, hence so does $\Gamma(T\mathcal{M})$.

Given a graded manifold \mathcal{M} , a $T\mathcal{M}$ -connection on a graded vector bundle \mathcal{E} is a \mathbb{R} -bilinear map $\nabla : \Gamma(T\mathcal{M}) \times \Gamma(\mathcal{E}) \to \Gamma(\mathcal{E})$ of degree 0 satisfying

(1)
$$\nabla_{fX}s = f \cdot \nabla_X s,$$

(2) $\nabla_X(f \cdot s) = X(f) \cdot s + (-1)^{|f| \cdot |X|} f \cdot \nabla_X s,$

for $f \in C^{\infty}(\mathcal{M}), X \in \mathfrak{X}(\mathcal{M})$ and $s \in \Gamma(\mathcal{E})$.

When $\mathcal{E} = T\mathcal{M}$, the $T\mathcal{M}$ -connection ∇ on $T\mathcal{M}$ is called an affine connection. We say an affine connection ∇ is torsion-free if

$$\nabla_X Y - (-1)^{|X| \cdot |Y|} \nabla_Y X = [X, Y]$$

for $X, Y \in \mathfrak{X}(\mathcal{M})$. It is well known that affine torsion-free connections always exist [14].

A **dg manifold** is a graded manifold \mathcal{M} together with a homological vector field, i.e. a vector field $Q \in \mathfrak{X}(\mathcal{M})$ of degree +1 satisfying $[Q, Q] = Q \circ Q - (-1)^1 Q \circ Q = 0$.

A **dg vector bundle** $\pi : (\mathcal{E}, Q_{\mathcal{E}}) \to (\mathcal{M}, Q)$ is a vector bundle object in the category of dg manifolds. See [18] for precise definition. In fact, given a dg manifold (\mathcal{M}, Q) , a graded vector bundle $\mathcal{E} \to \mathcal{M}$ has a dg vector bundle structure if and only if there exists an operator $\mathcal{Q} : \Gamma(\mathcal{E}) \to \Gamma(\mathcal{E})$ of degree +1 such that $\mathcal{Q}^2 = 0$ and

$$\mathcal{Q}(f \cdot e) = Q(f) \cdot e + (-1)^{|f|} f \cdot \mathcal{Q}(e)$$

for $f \in C^{\infty}(\mathcal{M})$ and $e \in \Gamma(\mathcal{E})$.

In particular, the tangent bundle $T\mathcal{M}$ naturally forms a dg vector bundle over (\mathcal{M}, Q) by setting $\mathcal{Q} = \mathcal{L}_Q = [Q, -]$ to be the Lie derivative along the homological vector field Q. The corresponding homological vector field $Q_{T\mathcal{M}}$ on $T\mathcal{M}$ is the tangent lift of the homological vector field Q on \mathcal{M} .

Consider a graded vector bundle $\mathcal{E} = T^{\vee} \mathcal{M} \otimes \operatorname{End}(T\mathcal{M})$ over \mathcal{M} . We define an operator \mathcal{Q} of degree +1 on the graded $C^{\infty}(\mathcal{M})$ -module $\Gamma(T^{\vee} \mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))$:

$$\mathcal{Q}: \Gamma(T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))^{\bullet} \to \Gamma(T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))^{\bullet+1}$$
(1)

by the Lie derivative along the homological vector field Q:

$$(\mathcal{Q}F)(X,Y) = [Q,F(X,Y)] - (-1)^k F([Q,X],Y) - (-1)^{k+|X|} F(X,[Q,Y])$$

for any (1, 2)-tensor $F \in \Gamma(T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))^k$ of degree k and vector fields $X, Y \in \mathfrak{X}(\mathcal{M})$. One can easily check that \mathcal{Q} induces a dg vector bundle structure on $\mathcal{E} = T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M})$ over (\mathcal{M}, Q) .

Now given an affine connection ∇ on \mathcal{M} , consider the (1, 2)-tensor $\operatorname{At}_{(\mathcal{M}, Q)}^{\nabla} \in \Gamma(T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))$ of degree +1, defined by

$$\operatorname{At}_{(\mathcal{M},Q)}^{\nabla}(X,Y) = [Q,\nabla_X Y] - \nabla_{[Q,X]} Y - (-1)^{|X|} \nabla_X [Q,Y]$$

for $X, Y \in \mathfrak{X}(\mathcal{M})$.

Proposition 2.3 ([18]). In the above setting, the following statements hold.

(1) If the affine connection ∇ on \mathcal{M} is torsion-free, then $\operatorname{At}_{(\mathcal{M},Q)}^{\nabla} \in \Gamma(S^2(T^{\vee}\mathcal{M}) \otimes T\mathcal{M})$. In other words,

$$\operatorname{At}_{(\mathcal{M},Q)}^{\nabla}(X,Y) = (-1)^{|X| \cdot |Y|} \operatorname{At}_{(\mathcal{M},Q)}^{\nabla}(Y,X).$$

- (2) The degree 1 element $\operatorname{At}_{(\mathcal{M},Q)}^{\nabla} \in \Gamma(T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))^1$ is a 1-cocycle.
- (3) The cohomology class $[At^{\nabla}_{(\mathcal{M},Q)}]$ does not depend on the choice of connection.

The element $\operatorname{At}_{(\mathcal{M},Q)}^{\nabla}$ is called the **Atiyah cocycle** associated with the affine connection ∇ . The cohomology class $\alpha_{(\mathcal{M},Q)} := [\operatorname{At}_{(\mathcal{M},Q)}^{\nabla}] \in H^1(\Gamma(T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))^{\bullet}, \mathcal{Q})$ is called the **Atiyah class** of the dg manifold (\mathcal{M}, Q) [18]. See also [21] and [16].

The Atiyah class of a dg manifold (\mathcal{M}, Q) is an obstruction to the existence of an affine connection ∇ on \mathcal{M} compatible with the homological vector field Q.

3. The Atiyah class and clean intersection

As shown in [3], the category of dg manifolds of positive amplitude is equivalent to the category of L_{∞} bundles. In particular, any dg manifold of amplitude +1 is of the form $(E[-1], \iota_s)$ where $E \to M$ is a smooth vector bundle and $s \in \Gamma(E)$ is a section of E. Explicitly, the graded algebra of smooth functions $C^{\infty}(E[-1])$ on E[-1], equipped with a homological vector field ι_s , is formulated as a cochain complex of $C^{\infty}(M)$ -modules

$$\cdots \to \Gamma(\Lambda^2 E^{\vee}) \xrightarrow{\iota_s} \Gamma(\Lambda^1 E^{\vee}) \xrightarrow{\iota_s} C^{\infty}(M) \to 0$$

where ι_s is the interior product with s.

Remark 3.1. The way we count the degree of $C^{\infty}(E[-1]) \cong \Gamma(\Lambda^{-\bullet}E^{\vee})$ here is indicated by the integer \bullet . This is a commonly used notation, especially in cochain complexes: de Rham complex, Chevalley–Eilenberg cochain complex, etc. However, in some cases, such as the equation in Lemma 3.13 or Eq. (11) below, the degree coming from \bullet and the degree coming from the graded space are mixed. Thus, one needs to be careful with counting the degree. To avoid confusion, one should view the graded space $\Gamma(\Lambda^{-\bullet}E^{\vee})$ as a graded space $\Gamma(S(E[-1])^{\vee})^{\bullet}$ and the section s should be viewed as $s \in \Gamma(E[-1])$.

Given a dg manifold $(E[-1], \iota_s)$, there are two natural embeddings: one is the section $s : M \to E$ and the other is the zero section $0_E : M \to E$. Then, one can consider their intersection $Z := im(s) \cap im(0_E)$. It turns out that the Atiyah class of the dg manifold $(E[-1], \iota_s)$ measures how good the intersection Z of s and 0_E is. The following is our main theorem.

Theorem 3.2. Let M be a smooth manifold and let E be a vector bundle over M. Given a section $s \in \Gamma(E)$, the Atiyah class of the dg manifold $(E[-1], \iota_s)$ vanishes if and only if the intersection of s and the zero section 0_E is clean.

3.1. Clean intersection. Recall that given two embedded submanifolds X and Y of W, we say X and Y intersect cleanly if their intersection $Z := X \cap Y$ is a manifold and $T_p Z = T_p X \cap T_p Y$ for all $p \in Z$. Moreover, we say the embeddings $f : X \to W$ and $g : Y \to W$ intersect cleanly if their images intersect cleanly.

In particular, let E be a vector bundle over M and let $s: M \to E$ be a section. Consider the intersection of s and the zero section 0_E , or equivalently, the intersection of X := im(s) and $Y := im(0_E)$ in W := E. In this case, the intersection $Z := X \cap Y$ is naturally identified with the zero locus $s^{-1}(0) \subset M$. Moreover, for each $p \in s^{-1}(0)$, the intersection $T_{s(p)}X \cap T_{s(p)}Y \subset T_{s(p)}E$ of the tangent spaces can be identified with a subspace of T_pM . For this, we need a map $Ds_p: T_pM \to E_p$ for each $p \in s^{-1}(0)$.

Note that if $p \in M$ satisfies $s(p) = 0 \in E_p$, then the fibre $T_{s(p)}E$ of the bundle TE at $s(p) \in E$ has a canonical splitting $T_{s(p)}E \cong T_pM \oplus E_p$. For each $p \in s^{-1}(0)$, the map $Ds_p : T_pM \to E_p$ is defined by $Ds_p = \operatorname{pr}_{E_p} \circ Ts_p$ where Ts_p is the tangent map of s at p and $\operatorname{pr}_{E_p} : T_{s(p)}E \to E_p$ is the projection map:

 $T_pM \xrightarrow{T_{s_p}} T_{s(p)}E \cong T_pM \oplus E_p \xrightarrow{\operatorname{pr}_{E_p} \triangleleft} E_p .$ (2)

Lemma 3.3. For each $p \in s^{-1}(0)$, the vector space $T_{s(p)}X \cap T_{s(p)}Y$ in $T_{s(p)}E$ is isomorphic to ker Ds_p . Moreover, if $s^{-1}(0)$ is a manifold, then it satisfies $T_p(s^{-1}(0)) \subset \ker Ds_p$.

Proof. For each $p \in M$, it is clear by definition that $T_{s(p)}X = \operatorname{im} Ts_p$ and $T_{0_E(p)}Y = \operatorname{im}(T0_E)_p$ where $Ts_p: T_pM \to T_{s(p)}E$ and $(T0_E)_p: T_pM \to T_{0_E(p)}E$ are the tangent maps of s and 0_E at p, respectively. Suppose that $p \in s^{-1}(0)$. Under the splitting $T_{s(p)}E \cong T_pM \oplus E_p$, the image of each tangent maps at $v \in T_pM$ is

$$Ts_p(v) = (v, Ds_p(v)),$$
 $(T0_E)_p(v) = (v, 0).$

Thus,

$$T_{s(p)}X \cap T_{s(p)}Y \cong \{(v,0) \in T_pM \oplus E_p \cong T_{s(p)}E : Ds_p(v) = 0\} \cong \ker Ds_p$$

Since $T(X \cap Y) \subset TX \cap TY$, the last statement follows immediately.

Corollary 3.4. If the intersection of s and 0_E is clean, then $\dim T_p(s^{-1}(0)) + \operatorname{rank} Ds_p = \dim M$.

$$\Box$$

Example 3.5. Consider a vector bundle $E = \mathbb{R}^2 \times \mathbb{R}^3 \to \mathbb{R}^2 = M$ with a section $s : \mathbb{R}^2 \to \mathbb{R}^2 \times \mathbb{R}^3$ defined by $s(x, y) = (x, y; x, xy, x^2)$. Then $s^{-1}(0) = \{(0, y) \in \mathbb{R}^2\} \cong \mathbb{R}$ is a manifold. Moreover, simple computation shows that

$$Ds_{(0,y)} = \begin{bmatrix} 1 & 0 \\ y & 0 \\ 0 & 0 \end{bmatrix}$$

and thus, ker $Ds_{(0,y)} = T_{(0,y)}(s^{-1}(0))$. Therefore, the triple (E, M, s) gives a clean intersection.

Remark 3.6. Even in the case when the connected submanifolds X and Y intersect cleanly, their intersection does not have to be connected, nor constant dimensional. Consider a vector bundle $E = \mathbb{R}^2 \times \mathbb{R}^3$ over $M = \mathbb{R}^2$. Let $U_1 = \{(x, y) : x < 6\}$ and $U_2 = \{(x, y) : x > 4\}$ be open sets in M and let $\{\rho_1, \rho_2\}$ be a partition of unity subordinate to $\{U_1, U_2\}$. For sections $s_1(x, y) = (x, y; x, xy, x^2)$ and $s_2(x, y) = (x, y; 10 - x, y, 0)$ of E, we define $s = \rho_1 \cdot s_1 + \rho_2 \cdot s_2$. Then one can check that the triple (E, M, s) gives a clean intersection, but $s^{-1}(0) = \{(x, y) | x = 0\} \cup \{(10, 0)\}$ is a union of two connected manifolds in different dimensions.

The following lemma will be useful.

Lemma 3.7. Let E be a vector bundle of rank m over $M = \mathbb{R}^n$. Suppose that a section $s \in \Gamma(E)$ satisfies

- (1) s(0) = 0,
- (2) $\operatorname{rank}(Ds_0) = r$,
- (3) s(x) = 0 for all $x \in \{0\}^r \times \mathbb{R}^{n-r}$,

then there exist a local chart $\phi : U \to \mathbb{R}^n$ around $0 \in M$ and a local frame $\{e_1, \dots, e_m\}$ of E around $0 \in M$ such that $s \in \Gamma(E)$ has a local expression

$$s \circ \phi^{-1}|_{\phi(U)} = x^1 \cdot e_1 + \dots + x^r \cdot e_r$$

where x^1, \dots, x^n are coordinate functions on \mathbb{R}^n .

Proof. Given any frame $\{\tilde{e}_1, \dots, \tilde{e}_m\}$ of E over M, the section $s \in \Gamma(E)$ can be written as $s = \sum_{i=1}^m s^i \cdot \tilde{e}_i$ for some functions $s^1, \dots, s^m \in C^{\infty}(M)$. Thus, we may identify the section $s \in \Gamma(E)$ with a map $s = (s^1, \dots, s^m) : \mathbb{R}^n \to \mathbb{R}^m$.

The condition (1) and (2) implies that, by the inverse function theorem (or, by [22, Theorem 2.9]), there exists a chart $\phi : U \to \mathbb{R}^n$ around $0 \in M$ such that $s \circ \phi^{-1}|_{\phi(U)} = (x^1, \cdots, x^r, \tilde{s}^{r+1}, \cdots, \tilde{s}^m)$ for some functions $\tilde{s}^{r+1}, \cdots, \tilde{s}^m$, by rearranging the order of the frame $\{\tilde{e}_1, \cdots, \tilde{e}_m\}$.

Moreover, by the condition (3), the functions $\tilde{s}^{r+1}, \dots, \tilde{s}^m$ are in the ideal generated by the first r coordinate functions x^1, \dots, x^r , (c.f. [19, Lemma 2.1] or [22, Lemma 3.2]). Indeed, we may assume U is convex, and if we denote $\mathbf{x} = (x_1, \dots, x_r)$ and $\mathbf{y} = (x_{r+1}, \dots, x_n)$, then by the fundamental theorem of calculus, we have

$$\tilde{s}^{i}(\mathbf{x}, \mathbf{y}) = \int_{0}^{1} \frac{d}{dt} \tilde{s}^{i}(t \cdot \mathbf{x}, \mathbf{y}) \, \mathrm{d}t = \sum_{j=1}^{r} x_{j} \cdot \left(\int_{0}^{1} \frac{\partial \tilde{s}^{i}}{\partial x^{j}}(t \cdot \mathbf{x}, \mathbf{y}) \, \mathrm{d}t \right) = \left(\sum_{j=1}^{r} x^{j} \cdot g_{j}^{i} \right) (\mathbf{x}, \mathbf{y})$$

for all $i = r + 1, \dots, m$, by setting $g_j^i(\mathbf{x}, \mathbf{y}) = \int_0^1 \frac{\partial \tilde{s}^i}{\partial x^j} (t \cdot \mathbf{x}, \mathbf{y}) dt$.

Finally, we obtain our desired expression of $s \circ \phi^{-1}|_{\phi(U)}$ by considering a new local frame $\{e_1, \dots, e_m\}$ of E around $0 \in M$, defined by

$$e_k = \begin{cases} \tilde{e}_k + \sum_{j=r+1}^m g_k^j \cdot \tilde{e}_j, & \text{if } k \le r \\ \tilde{e}_k, & \text{if } k > r \end{cases}$$

This completes the proof.

We now characterise the clean intersection of a section $s: M \to E$ along the zero section $0_E: M \to E$.

Lemma 3.8. Let E be a rank m vector bundle over an n-dimensional manifold M. A section $s \in \Gamma(E)$ intersects along the zero section 0_E cleanly if and only if for each $p \in s^{-1}(0)$, there exists a local chart $\phi: U \to \mathbb{R}^n$ around p and a frame $\{e_1, \dots, e_m\}$ of $E|_U$ such that

$$s \circ \phi^{-1}|_{\phi(U)} = x^1 \cdot e_1 + \dots + x^r \cdot e_r \tag{3}$$

where r is the rank of $Ds_p: T_pM \to E_p$.

Proof. Suppose that the section s is characterised by the formula (3) in a neighbourhood $U = U_p$ of each $p \in s^{-1}(0)$. Then clearly, s and 0_E intersect cleanly.

Conversely, suppose s and 0_E intersect cleanly. Then $s^{-1}(0) \subset M$ is a smooth manifold. By the tubular neighbourhood theorem, there exists an open neighbourhood $V \subset M$ of $s^{-1}(0)$ such that V is diffeomorphic to the normal bundle of $s^{-1}(0)$. In particular, for each $p \in s^{-1}(0)$, there exists an open neighbourhood $U \subset M$ of p and a diffeomorphism $\alpha : U \to \mathbb{R}^d \times \mathbb{R}^{n-d}$ such that $\alpha(p) = 0$ and $\alpha(s^{-1}(0) \cap U) = \mathbb{R}^d \times \{0\}^{n-d}$. Also, by Corollary 3.4, the rank of $Ds_p = n - d$.

Observe that we have shown that the composition $s \circ \alpha^{-1} : \mathbb{R}^n \to E|_U \cong \mathbb{R}^n \times \mathbb{R}^m$ satisfies all the conditions in Lemma 3.7. Thus, by Lemma 3.7, the proof is complete.

3.2. Locality of the Atiyah class in positive amplitude. Consider a dg manifold (over \mathbb{C}) arising from arising from complex manifold X whose dg algebra of functions is the Dolbeault complex of X. It is shown [8] that the Atiyah class of this dg manifold is identical to the Atiyah class of the complex manifold X — it is an obstruction to the existence of holomorphic connections [2]. Observe that if the base manifold X is \mathbb{C}^n , then there always is a holomorphic connection. Thus, the Atiyah class of a dg manifold, in general, is not expected to be characterised by the local data.

However, suppose we are given a dg manifold of positive amplitude. Then its algebra of functions is concentrated in non-positive degrees and moreover, its degree 0 component consists of smooth functions on the base manifold. Since the homological vector field increase the degree by 1, it acts on the smooth functions on the base manifold trivially. In this case, the Atiyah class is completely determined by local data. In particular, the Atiyah class of a dg manifold of positive amplitude vanishes if and only if it vanishes locally.

To be more precise, consider a dg manifold (\mathcal{M}, Q) of positive amplitude whose base manifold is M. By [3], the graded manifold \mathcal{M} corresponds to a negatively graded vector bundle $E \to M$. Given an open subset $U \subset M$, denote the graded manifold with base U which corresponds to the negatively graded vector bundle $E|_U \to U$ by \mathcal{M}_U . Naturally, \mathcal{M}_U is a graded submanifold of \mathcal{M} . Also, denote the restriction of the homological vector field Q to \mathcal{M}_U by Q_U . Thus, each open subset $U \subset M$ gives rise to a dg manifold (\mathcal{M}_U, Q_U) of positive amplitude.

Proposition 3.9. Let (\mathcal{M}, Q) be a dg manifold of positive amplitude, and let $\{U_i\}_{i \in I}$ be an open cover of the base manifold \mathcal{M} . The Atiyah class α of (\mathcal{M}, Q) is completely determined by $\{\alpha_i\}_{i \in I}$ where α_i is the Atiyah class of $(\mathcal{M}_{U_i}, Q_{U_i})$. In particular, $\alpha = 0$ if and only if $\alpha_i = 0$ for all $i \in I$.

The positive amplitude condition on (\mathcal{M}, Q) which distinguishes from arbitrary dg manifold is that, by degree reason, the homological vector field Q acts on the space of smooth functions on M trivially, i.e. Q(f) = 0 for $\forall f \in \mathcal{O}_M$.

To prove Proposition 3.9, we use the notion of sheaves. The following lemma might be known but could not find a reference.

Lemma 3.10. Let \mathcal{O}_M be the sheaf of algebra of smooth functions on M and let $(\mathcal{E}, d) = (\mathcal{E}^{\bullet}, d)$ be a sheaf of dg module over \mathcal{O}_M . Then the cohomology $H^{\bullet}(\mathcal{E}, d)$ of the cochain complex $(\mathcal{E}^{\bullet}, d)$ is naturally a sheaf of graded \mathcal{O}_M -modules.

Proof. By the assumption, the differential $d : \mathcal{E}^{\bullet} \to \mathcal{E}^{\bullet+1}$ is a morphism of sheaves of \mathcal{O}_M -modules. In particular, the differential d is \mathcal{O}_M -linear and is compatible with the restriction map. Thus, the cohomology group $H^{\bullet}(\mathcal{E}, d)$ is naturally a presheaf of graded \mathcal{O}_M -module. For simplicity of notation, we use the symbol $\mathcal{H} = \mathcal{H}^{\bullet}$ to denote the presheaf of graded \mathcal{O}_M -module $H^{\bullet}(\mathcal{E}, d)$.

To show that \mathcal{H} is a sheaf, we need to show the locality axiom and gluing axiom.

Fix an open subset $U \subset M$ and an open cover $\{U_i\}_{i \in I}$ of U. For each $n \in \mathbb{Z}$, we have a map $F^n : \mathcal{E}^n(U) \to \prod_{i \in I} \mathcal{E}^n(U_i)$ induced by the restriction map $\mathcal{E}^n(U) \to \mathcal{E}^n(U_i)$ for each $i \in I$. Moreover, if we denote the induced differential on $\prod_{i \in I} \mathcal{E}^n$ by $\check{d} = \prod_{i \in I} d|_{U_i}$, it satisfies $\check{d} \circ F^n = F^{n+1} \circ d$.

Given a partition of unity $\{\rho_i\}_{i\in I}$ subordinate to $\{U_i\}_{i\in I}$, consider a map $G^n : \prod_{i\in I} \mathcal{E}^n(U_i) \to \mathcal{E}^n(U)$ defined by

$$G^n: \{s_i\}_{i\in I} \mapsto \sum_{i\in I} \rho_i \cdot s_i$$

where $s_i \in \mathcal{E}^n(U_i)$. Note that for each fixed $i \in I$, we have $\rho_i|_{U_j} \equiv 0$ for all but finitely many $j \in I$. Thus, for each $p \in M$, there exists an open neighbourhood U such that

$$\begin{aligned} d(s)|_{U} &= d|_{U}(s|_{U}) = d|_{U} \Big(\sum_{i \in I} (\rho_{i}|_{U} \cdot s_{i}|_{U_{i} \cap U}) \Big) \\ &= \sum_{i \in I} d|_{U} (\rho_{i}|_{U} \cdot s_{i}|_{U_{i} \cap U}) = \sum_{i \in I} \rho_{i}|_{U} \cdot d|_{U_{i} \cap U} (s_{i}|_{U_{i} \cap U}) = \Big(\sum_{i \in I} \rho_{i} \cdot d|_{U_{i}} (s_{i}) \Big) \Big|_{U} \end{aligned}$$

by writing $s = \sum_{i \in I} \rho_i \cdot s_i$. By the locality axiom, it satisfies

$$d\left(\sum_{i\in I}\rho_i\cdot s_i\right) = \sum_{i\in I}d(\rho_i\cdot s_i) = \sum_{i\in I}\rho_i\cdot d|_{U_i}(s_i) \tag{4}$$

and therefore we have $d \circ G^n = G^{n+1} \circ \check{d}$.

Observe that, by construction, both F^{\bullet} and G^{\bullet} are chain maps and they satisfy $G^n \circ F^n = id_{\mathcal{E}^n(U)}$. It implies that on the level of cohomology, they induce maps

$$F_*: H^n(\mathcal{E}, d)(U) \rightleftharpoons \prod_{i \in I} H^n(\mathcal{E}(U_i), d|_{U_i}) \cong H^n\left(\prod_{i \in I} \mathcal{E}(U_i), \check{d}\right): G$$

for each $n \in \mathbb{Z}$ such that $G_* \circ F_* = \mathrm{id}_{H^{\bullet}(\mathcal{E},d)(U)}$. In particular, if $\alpha \in \mathcal{H}(U)$ satisfies $F_*(\alpha) = 0$ (i.e. $0 = \alpha|_{U_i} \in \mathcal{H}(U_i)$ for all $i \in I$), then $\alpha = G_* \circ F_*(\alpha) = 0$. This proves the locality axiom.

Consider an element $\{\alpha_i\}_{i\in I} \in \prod_{i\in I} \mathcal{H}^n$ such that

$$\alpha_i|_{U_i \cap U_j} = \alpha_j|_{U_i \cap U_j} \tag{5}$$

for all $i, j \in I$. If we pick a representative $s_i \in \ker d \cap \mathcal{E}^n(U_i)$ of α_i for each *i*, then Eq. (5) is equivalent to

$$s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j} + dw_{ij}$$

for some $w_{ij} \in \mathcal{E}^{n-1}(U_i \cap U_j)$. Now, consider $s = \sum_{i \in I} \rho_i \cdot s_i \in \mathcal{E}^n$. By Eq. (4), the element s is in ker d. Moreover, it satisfies

$$s|_{U_j} = (\sum_{i \in I} \rho_i \cdot s_i)|_{U_j} = \sum_{i \in I} \rho_i|_{U_j} \cdot s_i|_{U_i \cap U_j} = \sum_{i \in I} \rho_i|_{U_j} \cdot (s_j|_{U_i \cap U_j} + dw_{ij})$$
$$= \sum_{i \in I} \rho_i|_{U_j} \cdot s_j|_{U_i \cap U_j} + \sum_{i \in I} \rho_i|_{U_j} \cdot dw_{ij} = s_j + d(\sum_{i \in I} \rho_i|_{U_j} \cdot w_{ij})$$

since $\rho_i|_{U_j} \equiv 0$ for all but finitely many $j \in I$. Therefore, the cohomology class α represented by s satisfies $F_*(\alpha) = \{\alpha_i\}_{i \in I}$. This proves the gluing axiom, whence proves that $\mathcal{H} = H^{\bullet}(\mathcal{E}, d)$ is a sheaf of graded \mathcal{O}_M -module.

Let (\mathcal{M}, Q) be a dg manifold of positive amplitude with base manifold M. If $(\mathcal{E}, Q_{\mathcal{E}})$ is a dg vector bundle over (\mathcal{M}, Q) , then its space of sections $\Gamma(\mathcal{M}; \mathcal{E})$, equipped with the differential $d : \Gamma(\mathcal{M}; \mathcal{E}) \to \Gamma(\mathcal{M}; \mathcal{E})$ induced by $Q_{\mathcal{E}}$, becomes a sheaf of dg \mathcal{O}_M -module

$$U \mapsto (\Gamma(\mathcal{M}_U; \mathcal{E}_U), d_U)$$

where \mathcal{E}_U is the restriction of the bundle \mathcal{E} to the graded submanifold $\mathcal{M}_U \subset \mathcal{M}$, and the operator $d_U : \Gamma(\mathcal{M}_U; \mathcal{E}_U) \to \Gamma(\mathcal{M}_U; \mathcal{E}_U)$ is the differential such that the diagram

is commutative. Here, vertical maps send a section $s : \mathcal{M} \to \mathcal{E}$ to a section $s|_{\mathcal{M}_U} : \mathcal{M}_U \to \mathcal{E}_U$.

Remark 3.11. To be more precise with the notations above, the graded algebra $C^{\infty}(\mathcal{M}_U)$ of smooth functions on \mathcal{M}_U has a natural identification

$$C^{\infty}(\mathcal{M}_U) \cong C^{\infty}(U) \otimes_{C^{\infty}(M)} C^{\infty}(\mathcal{M})$$

and the inclusion $\mathcal{M}_U \hookrightarrow \mathcal{M}$ corresponds to the map $C^{\infty}(\mathcal{M}) \to C^{\infty}(\mathcal{M}_U)$ defined by $f \mapsto 1 \otimes f$. Moreover, we have a natural identification

$$\Gamma(\mathcal{M}_U; \mathcal{E}_U) \cong C^{\infty}(\mathcal{M}_U) \otimes_{C^{\infty}(\mathcal{M})} \Gamma(\mathcal{M}; \mathcal{E})$$

and under this identification, we have $d_U = Q_U \otimes 1 + 1 \otimes d$.

We have proved the following.

Corollary 3.12. Suppose that the dg manifold (\mathcal{M}, Q) with base M is of positive amplitude. If $(\mathcal{E}, Q_{\mathcal{E}})$ is a dg vector bundle over (\mathcal{M}, Q) , then the assignment

$$U \mapsto H^{\bullet}(\Gamma(\mathcal{M}_U; \mathcal{E}_U), d_U)$$

is a sheaf of \mathcal{O}_M -module where $H^{\bullet}(\Gamma(\mathcal{E}), d)$ is the cohomology of the dg space of sections $(\Gamma(\mathcal{M}; \mathcal{E}), d)$ of $(\mathcal{E}, Q_{\mathcal{E}})$.

Now we are ready to prove Proposition 3.9.

Proof of Proposition 3.9. Consider the graded vector bundle $\mathcal{E} = T^{\vee}\mathcal{M} \otimes T^{\vee}\mathcal{M} \otimes T\mathcal{M}$, equipped with homological vector field $Q_{\mathcal{E}}$ induced by the complete lift of Q. By Corollary 3.12, it induces a sheaf \mathcal{H} of graded \mathcal{O}_M -module, defined by

$$U \mapsto \mathcal{H}(U) = H^{\bullet}(\Gamma(\mathcal{M}_U; T^{\vee} \mathcal{M}_U \otimes T^{\vee} \mathcal{M}_U \otimes T \mathcal{M}_U), \mathcal{Q}_U)$$

where Q is defined by (1). Note that the Atiyah class $\alpha_{(\mathcal{M},Q)}$ of (\mathcal{M},Q) belongs to its global section $\mathcal{H}(M)$.

Given open sets $V \subset U \subset M$, let $\rho_{U,V} : \mathcal{H}(U) \to \mathcal{H}(V)$ be the restriction map of the sheaf \mathcal{H} . Now, to prove Proposition 3.9, it suffices to prove

$$\rho_{M,U}(\alpha_{(\mathcal{M},Q)}) = \alpha_{(\mathcal{M}_U,Q_U)}$$

where $\alpha_{(\mathcal{M},Q)}$ and $\alpha_{(\mathcal{M}_U,Q_U)}$ are the Atiyah class of (\mathcal{M},Q) and (\mathcal{M}_U,Q_U) , respectively. Indeed, this means that for any two open subsets $U, V \subset M$, it satisfies $\rho_{U,U\cap V}(\alpha_{(\mathcal{M}_U,Q_U)}) = \rho_{V,U\cap V}(\alpha_{(\mathcal{M}_V,Q_V)})$, and thus using the gluing axiom, the Atiyah class computed locally completely determines the Atiyah class of (\mathcal{M},Q) .

Given an open subset U of M, denote the inclusion of graded manifold by $i_U : \mathcal{M}_U \hookrightarrow \mathcal{M}$. Observe that for any graded vector bundle \mathcal{E} over \mathcal{M} , there is a canonical identification $i_U^* \mathcal{E} \cong \mathcal{E}_U$ where $i_U^* \mathcal{E}$ is the pullback bundle of \mathcal{E} over i_U . In particular, when $\mathcal{E} = \mathfrak{X}(\mathcal{M})$, we have $\mathcal{E}_U = \mathfrak{X}(\mathcal{M}_U) \cong C^{\infty}(\mathcal{M}_U) \otimes_{C^{\infty}(\mathcal{M})} \mathfrak{X}(\mathcal{M})$ and one can check that the restriction map $\mathfrak{X}(\mathcal{M}) \to \mathfrak{X}(\mathcal{M}_U)$ defined by

$$X \mapsto X|_U = 1 \otimes X \in C^{\infty}(\mathcal{M}_U) \otimes_{C^{\infty}(\mathcal{M})} \mathfrak{X}(\mathcal{M})$$

is a morphism of Lie algebra.

Moreover, if $\nabla : \mathfrak{X}(\mathcal{M}) \times \mathfrak{X}(\mathcal{M}) \to \mathfrak{X}(\mathcal{M})$ is an affine connection on \mathcal{M} , then the pullback connection $\nabla^{i_U} : \mathfrak{X}(\mathcal{M}_U) \times \mathfrak{X}(\mathcal{M}_U) \to \mathfrak{X}(\mathcal{M}_U)$, characterised by

$$\nabla_{X|_U}^{i_U} Y|_U = (\nabla_X Y)|_U$$

for $X, Y \in \mathfrak{X}(\mathcal{M})$, is again an affine connection on \mathcal{M}_U .

For the Atiyah cocycle At^{∇} corresponds to an affine connection ∇ on \mathcal{M} , we have

$$\begin{aligned} \operatorname{At}^{\nabla}(X,Y)|_{U} &= [Q, \nabla_{X}Y]|_{U} - (\nabla_{[Q,X]}Y)|_{U} - (-1)^{|X|} (\nabla_{X}[Q,Y])|_{U} \\ &= [Q_{U}, (\nabla_{X}Y)|_{U}] - \nabla^{i_{U}}_{[Q,X]|_{U}}Y|_{U} - (-1)^{|X|_{U}|} \nabla^{i_{U}}_{X|_{U}}[Q,Y]|_{U} \\ &= [Q_{U}, \nabla^{i_{U}}_{X|_{U}}Y|_{U}] - \nabla^{i_{U}}_{[Q_{U},X|_{U}]}Y|_{U} - (-1)^{|X|_{U}|} \nabla^{i_{U}}_{X|_{U}}[Q_{U},Y|_{U}] \\ &= \operatorname{At}^{\nabla^{i_{U}}}. \end{aligned}$$
(6)

Therefore, the induced restriction map $\rho_{M,U}$ satisfies $\rho_{M,U}(\alpha_{(\mathcal{M},Q)}) = \alpha_{(\mathcal{M}_U,Q_U)}$. This completes the proof of Proposition 3.9.

3.3. The Atiyah class of dg manifolds of amplitude +1. We now focus on dg manifolds of amplitude +1. In this section, we will present the explicit description of the space of vector fields of dg manifolds of amplitude +1 and its Lie algebra structure. Also, we will characterise torsion-free affine connections on dg manifolds of amplitude +1. Then, we will finally describe the Atiyah class of dg manifolds of amplitude +1 in both global and local picture.

For the remainder of this section, unless otherwise stated, $\pi : E \to M$ is a (usual) vector bundle over M and E[-1] is always considered as a graded manifold of amplitude +1, rather than a graded vector bundle.

3.3.1. Vector fields and affine connections. Recall that given an ordinary smooth vector bundle $\pi : E \to M$, there exists a short exact sequence

$$0 \to \Gamma(E; \pi^*E) \to \mathfrak{X}(E) \xrightarrow{\pi_*} \Gamma(E; \pi^*TM) \to 0$$

of $C^{\infty}(E)$ -modules. Analogously, for the graded manifold $\mathcal{M} = E[-1]$ with base M, the algebra of smooth functions is $C^{\infty}(\mathcal{M}) \cong \Gamma(\Lambda^{-\bullet}E^{\vee})$ and one obtains a short exact sequence

$$0 \to \Gamma(\Lambda^{-\bullet} E^{\vee}) \otimes_{C^{\infty}(M)} \Gamma(E[-1]) \xrightarrow{\iota} \mathfrak{X}(E[-1]) \xrightarrow{\pi_*} \Gamma(\Lambda^{-\bullet} E^{\vee}) \otimes_{C^{\infty}(M)} \mathfrak{X}(M) \to 0$$
(7)

of graded $\Gamma(\Lambda^{-\bullet}E^{\vee})$ -modules. Here, the map ι is the $\Gamma(\Lambda^{-\bullet}E^{\vee})$ -linear extension of the interior product and the map π_* is induced by the morphism of dg manifolds $E[-1] \to E \xrightarrow{\pi} M$.

Note that a $\Gamma(\Lambda^{-\bullet}E^{\vee})\text{-linear}$ map

$$\tau: \Gamma(\Lambda^{-\bullet} E^{\vee}) \otimes_{C^{\infty}(M)} \mathfrak{X}(M) \to \mathfrak{X}(E[-1])$$

satisfying $\pi_* \circ \tau = \text{id}$ is called a horizontal lift. Let ∇^E be a *TM*-connection on *E*. Extending its dual connection by derivation law, one obtains a *TM*-connection on $\Lambda^{-\bullet}E^{\vee}$, again denoted by ∇^E , by abuse of notation. The horizontal lift associated with ∇^E is the $\Gamma(\Lambda^{-\bullet}E^{\vee})$ -linear map

$$\tau = \tau^{\nabla^E} : \Gamma(\Lambda^{-\bullet} E^{\vee}) \otimes_{C^{\infty}(M)} \mathfrak{X}(M) \to \mathfrak{X}(E[-1]) \cong \operatorname{Der}(\Gamma(\Lambda^{-\bullet} E^{\vee}))$$

characterised by

$$\tau(X): \xi \mapsto \nabla_X^E \xi$$

for $X \in \mathfrak{X}(M)$ and $\xi \in \Gamma(\Lambda^{-\bullet}E^{\vee})$. In particular, we have $\nabla_X^E f = X(f)$ for $X \in \mathfrak{X}(M)$ and $f \in C^{\infty}(M)$, thus satisfies $\pi_* \circ \tau(X) = X$ for each $X \in \mathfrak{X}(M)$.

Lemma 3.13. Upon a choice of a TM-connection ∇^E on E, there exists a $\Gamma(\Lambda^{-\bullet}E^{\vee})$ -module isomorphism $\Gamma(\Lambda^{-\bullet}E^{\vee}) \otimes_{C^{\infty}(M)} (\Gamma(E[-1]) \oplus \Gamma(TM)) \cong \mathfrak{X}(E[-1])$

characterised by

$$a \mapsto \iota_a, \quad and \quad X \mapsto \nabla^E_X$$

for $X \in \mathfrak{X}(M)$ and $a \in \Gamma(E[-1])$.

We note that the vector field $\nabla_X^E \in \mathfrak{X}(E[-1])$ on the graded manifold E[-1] is a horizontal lift of the vector field $X \in \mathfrak{X}(M)$ on M.

Lemma 3.13 indicates that any homogeneous vector fields on E[-1] has degree at most +1. Moreover, by restricting to homogeneous component of degree +1 and 0, we have

$$\mathfrak{X}(E[-1])^{+1} \cong \Gamma(E[-1]), \qquad \mathfrak{X}(E[-1])^0 \cong \Gamma(\Lambda^1 E^{\vee} \otimes E[-1]) \oplus \Gamma(TM)$$
(8)

where the symbol $\mathfrak{X}(E[-1])^n$ denotes homogeneous component of degree n in $\mathfrak{X}(E[-1])$.

Now, the Lie bracket [-, -] on $\mathfrak{X}(E[-1])$ can be computed easily. Fix a *TM*-connection ∇^E on *E* and denote the horizontal lift of a vector field *X* by $\hat{X} := \nabla^E_X$. Then, for $a, b \in \Gamma(E[-1])$ and $X, Y \in \mathfrak{X}(M)$, we have

$$[\iota_a, \iota_b] = 0, \quad [\hat{X}, \iota_a] = \iota_{\nabla_X^E a}, \quad [\hat{X}, \hat{Y}] = [X, Y]_M - \iota_{R^{\nabla^E}(X, Y)}$$
(9)

where $[-,-]_M$ denotes the Lie bracket on TM and R^{∇^E} is the curvature of the connection ∇^E . Here the symbol $\iota_{R^{\nabla^E}(X,Y)}$ denotes the image of the element $R^{\nabla^E}(X,Y) \in \Gamma(E^{\vee}) \otimes_{C^{\infty}(M)} \Gamma(E[-1])$ under the map ι in (7).

Finally, we investigate torsion-free affine connections on the graded manifold E[-1].

Lemma 3.14. A torsion-free affine connection $\nabla : \mathfrak{X}(E[-1]) \times \mathfrak{X}(E[-1]) \to \mathfrak{X}(E[-1])$ is equivalent to a triple $(\nabla^E, \nabla^M, \beta)$ where

- (1) $\nabla^E : \Gamma(TM) \times \Gamma(E) \to \Gamma(E)$ is a TM-connection on E;
- (2) $\nabla^M : \Gamma(TM) \times \Gamma(TM) \to \Gamma(TM)$ is a torsion-free affine connection on M;
- (3) an element $\beta \in \Gamma(T^{\vee}M \otimes T^{\vee}M \otimes \operatorname{End}(E))$ satisfying

$$\beta(X,Y) - \beta(Y,X) = -R^{\nabla^E}(X,Y).$$

In particular, denoting $\hat{X} = \nabla_X^E \in \mathfrak{X}(E[-1])$ for each $X \in \mathfrak{X}(M)$, we have an explicit formula:

$$\nabla_{\iota_a}\iota_b = 0, \quad \nabla_{\iota_a}\hat{X} = 0, \quad \nabla_{\hat{X}}\iota_a = \iota_{\nabla_X^E a}, \quad \nabla_{\hat{X}}\hat{Y} = \widehat{\nabla}_X^M \widehat{Y} + \iota_{\beta(X,Y)}, \tag{10}$$

where $\iota_{\beta(X,Y)}$ is the image of $\beta(X,Y) \in \Gamma(\operatorname{End} E) \cong \Gamma(E^{\vee}) \otimes_{C^{\infty}(M)} \Gamma(E[-1])$ under the map ι in (7).

Proof. Given a triple $(\nabla^E, \nabla^M, \beta)$, it is easy to show that the relation (10) completely determines a torsion-free affine connection on E[-1].

Conversely, suppose that a torsion-free affine connection $\nabla : \mathfrak{X}(E[-1]) \times \mathfrak{X}(E[-1]) \to \mathfrak{X}(E[-1])$ is given. If we denote an arbitrary horizontal lift of $X \in \mathfrak{X}(M)$ by \hat{X} , by Lemma 3.13 and by the torsion-freeness, the connection ∇ is completely determined by values of the form $\nabla_{\iota_a}\iota_b$, $\nabla_{\hat{X}}\iota_a$ and $\nabla_{\hat{X}}\hat{Y}$ for $a, b \in \Gamma(E[-1])$ and $X, Y \in \mathfrak{X}(M)$.

First, by degree reason, for all $a, b \in \Gamma(E[-1])$, we have $\nabla_{\iota_a} \iota_b = 0$.

Second, observe that the assignment

$$(X,a) \mapsto \nabla_{\hat{X}} \iota_a \in \mathfrak{X}(E[-1])^{+1} \cong \Gamma(E[-1])$$

defines a *TM*-connection $\nabla^E : \Gamma(TM) \times \Gamma(E[-1]) \to \Gamma(E[-1])$ on E[-1]. Indeed, since $\hat{X}(f) = X(f)$ for $f \in C^{\infty}(M)$, it satisfies $\nabla_{\hat{X}} \iota_{f \cdot a} = X(f) \cdot \iota_a + f \cdot \nabla_{\hat{X}} \iota_a$. Thus, we have $\nabla_{\hat{X}} \iota_a = \iota_{\nabla_X^E a}$ for some ∇^E .

Moreover, the connection ∇^E is chosen independently of the choice of horizontal lifts. Denote a different horizontal lift of $X \in \mathfrak{X}(M)$ by \overline{X} . By the short exact sequence (7), $\hat{X} - \overline{X} \in \operatorname{in} \iota$. Thus, because $\nabla_{\iota_b}\iota_a = 0$ for all $a, b \in \Gamma(E[-1])$, we have $\nabla_{\hat{X}-\overline{X}}\iota_a = 0$ for all $a \in \Gamma(E[-1])$. This implies that ∇ induces ∇^E independently of the choice of horizontal lifts.

Similarly, the assignment

$$(X,Y) \mapsto \pi_*(\nabla_{\hat{X}} \hat{Y}) \in \Gamma(TM)$$

defines an affine connection $\nabla^M : \Gamma(TM) \times \Gamma(TM) \to \Gamma(TM)$ on M. Here $\pi_* : \mathfrak{X}(E[-1])^0 \to \Gamma(TM)$ is induced by the map in the short exact sequence (7). Similar argument shows that ∇^M is chosen independently of the choice of horizontal lifts.

Now, given a torsion-free affine connection ∇ on E[-1], we have ∇^E and ∇^M , which does not depend on the choice of horizontal lift. Now, we choose the horizontal lift \hat{X} of $X \in \mathfrak{X}(M)$ to be the horizontal lift $\hat{X} = \nabla^E_X$ associated with ∇^E . Define $\iota_{\beta(X,Y)} := \nabla_{\hat{X}} \hat{Y} - \widehat{\nabla^M_X Y}$.

Finally, it is easy to show that the torsion-free condition implies that the connection ∇^M is torsion-free and that the relation $\beta(X, Y) - \beta(Y, X) = -R^{\nabla^E}(X, Y)$ holds. This completes the proof.

3.3.2. Atiyah class: global description. Given a dg manifold (\mathcal{M}, Q) , the Atiyah class is the 1st cohomology class of the cochain complex $(\Gamma(\mathcal{M}; T^{\vee}\mathcal{M} \otimes \operatorname{End}(T\mathcal{M}))^{\bullet}, Q)$. In fact, by Proposition 2.3, it suffices to consider the subcomplex $(\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M}))^{\bullet}, Q)$. When $(\mathcal{M}, Q) = (E[-1], \iota_s)$, by Lemma 3.13, a TM-connection ∇^E on E induces a graded $C^{\infty}(E[-1])$ -module isomorphism

$$\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M})) \cong \Gamma(M; \Lambda^{-\bullet} E^{\vee} \otimes \operatorname{Hom}(S^2(E[-1] \oplus TM), E[-1] \oplus TM)).$$
(11)

It can be easily checked that

- (1) $\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M}))$ is concentrated in degree ≤ 1 .
- (2) degree +1 component of $\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M}))$ is

$$\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M}))^1 \cong \Gamma(M; \operatorname{Hom}(S^2(TM), E[-1])).$$

(3) degree 0 component of $\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M}))$ is

$$\Gamma(\mathcal{M}; \operatorname{Hom}(S^{2}(T\mathcal{M}), T\mathcal{M}))^{0} \cong \Gamma(M; \Lambda^{1}E^{\vee} \otimes \operatorname{Hom}(S^{2}(TM), E[-1])) \\ \oplus \Gamma(M; \operatorname{Hom}(E[-1] \otimes TM, E[-1])) \\ \oplus \Gamma(M; \operatorname{Hom}(S^{2}(TM), TM)).$$

The differential

$$\mathcal{Q}: \Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M}))^0 \to \Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M}))^1$$
(12)

decomposes into $Q = d_1 + d_2 + d_3$ where

- (1) The first map $d_1 : \Gamma(M; E^{\vee} \otimes \operatorname{Hom}(S^2(TM), E[-1])) \to \Gamma(M; \operatorname{Hom}(S^2(TM), E[-1]))$ is obtained by applying $\iota_s : \Gamma(E^{\vee}) \to C^{\infty}(M)$ on the first component.
- (2) The second map $d_2 : \Gamma(M; \operatorname{Hom}(E[-1] \otimes TM, E[-1])) \to \Gamma(M; \operatorname{Hom}(S^2(TM), E[-1]))$ is obtained by the following commutative diagram

$$\Gamma(M; \operatorname{Hom}(E[-1] \otimes TM, E[-1])) \xrightarrow{d_2} \Gamma(M; \operatorname{Hom}(S^2(TM), E[-1]))$$

$$\uparrow^{\operatorname{sym}^*}$$

$$\Gamma(M; \operatorname{Hom}(TM \otimes TM, E[-1]))$$

where $(\nabla^E s)^*$ is the pre-composition of $\nabla^E s : TM \to E[-1]$ on the first component of the domain and sym^{*} is the pre-composition of sym $: S^2(TM) \to TM \otimes TM$ defined by $\operatorname{sym}(X \odot Y) = X \otimes Y + Y \otimes X$.

(3) The last map $d_3 : \Gamma(M; \operatorname{Hom}(S^2(TM), TM)) \to \Gamma(M; \operatorname{Hom}(S^2(TM), E[-1]))$ is obtained by post-composition of $\nabla^E s : TM \to E[-1]$.

Here, we view $\nabla^E s : TM \to E[-1]$ as a morphism of vector bundles defined by $X \mapsto \nabla^E_X s$. This describes the first cohomology group of $(\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M})), \mathcal{Q})$. In particular, if s is a nowhere vanishing section, then its first cohomology group vanishes.

Lemma 3.15. Let $(\mathcal{M}, Q) = (E[-1], \iota_s)$ be the dg manifold associated with a vector bundle E and its section $s \in \Gamma(E)$. If s is a nowhere vanishing section, then the first cohomology group $H^1(\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), T\mathcal{M})), Q)$ vanishes.

Proof. If s is nowhere vanishing, then there exists a dual section $\xi \in \Gamma(E^{\vee})$ such that $\iota_s(\xi) = 1 \in C^{\infty}(M)$. Then the aforementioned map d_1 is surjective and thus the map Q in (12) is surjective. In other words, all 1-cocycles are in the coboundary, and this proves the lemma.

Corollary 3.16. If $s \in \Gamma(E)$ is nowhere vanishing, then the Atiyah class of $(E[-1], \iota_s)$ vanishes.

Next, we compute an Atiyah cocycle of the dg manifold $(E[-1], \iota_s)$.

Proposition 3.17. Consider a dg manifold $(E[-1], \iota_s)$. Given a torsion-free affine connection ∇ on the graded manifold E[-1], the associated Atiyah cocycle $\operatorname{At}^{\nabla} \in \Gamma(\mathcal{M}; S^2(T^{\vee}\mathcal{M}) \otimes T\mathcal{M})$ is completely determined by

$$\operatorname{At}^{\nabla}(\hat{X}, \hat{Y}) = \iota_{\nabla_X^E \nabla_Y^E s - \nabla_{\nabla_X^M Y}^E s + \beta(X, Y)s}$$

for $X, Y \in \mathfrak{X}(M)$ where $\nabla = (\nabla^E, \nabla^M, \beta)$ as in Lemma 3.14, and $\hat{X} = \nabla^E_X \in \mathfrak{X}(E[-1])$ denotes the horizontal lift of $X \in \mathfrak{X}(M)$.

Proof. Since At^{∇} is $C^{\infty}(\mathcal{M})$ -linear in both argument, it suffices to compute

$$\operatorname{At}^{\nabla}(\iota_a,\iota_b), \quad \operatorname{At}^{\nabla}(\iota_a,\hat{X}), \quad \operatorname{At}^{\nabla}(\hat{X},\hat{Y})$$

for $a, b \in \Gamma(E[-1])$ and $X, Y \in \mathfrak{X}(M)$. By degree reason, $\operatorname{At}^{\nabla}(\iota_a, \iota_b) = 0$ and $\operatorname{At}^{\nabla}(\iota_a, \hat{X}) = 0$.

By Lemma 3.14, we have

$$\begin{aligned} \operatorname{At}^{\nabla}(\hat{X}, \hat{Y}) &= [Q, \nabla_{\hat{X}} \hat{Y}] - \nabla_{[Q, \hat{X}]} \hat{Y} - \nabla_{\hat{X}} [Q, \hat{Y}] \\ &= [\iota_s, \widehat{\nabla_X^M Y} + \iota_{\beta(X, Y)}] + \nabla_{\iota_{\nabla_X^E s}} \hat{Y} + \nabla_{\hat{X}} (\iota_{\nabla_Y^E s}) \\ &= \iota_{\beta(X, Y)s - \nabla_{\nabla_X^M Y}^E s} + 0 + \iota_{\nabla_X^E \nabla_Y^E s}. \end{aligned}$$

This completes the proof.

Under the identification (11), the Atiyah cocycle At^{∇} corresponds to a bundle map $\alpha : S^2(TM) \to E[-1]$ of degree +1 defined by

$$\alpha(X,Y) = \nabla_X^E \nabla_Y^E s - \nabla_{\nabla_X^M Y}^E s + \beta(X,Y)s.$$

3.3.3. Atiyah class: local description. By Proposition 3.9, in order to investigate the Atiyah class of a dg manifold of amplitude +1, it suffices to consider the local model. Thus, we will compute an Atiyah cocycle and the cohomology operator explicitly in a local coordinate.

For the local computation, we assume that $M = \mathbb{R}^n$ and $E = \mathbb{R}^n \times \mathbb{R}^m$ is a rank *m*-bundle over M. We denote the coordinate functions of M by $\{x^1, \dots, x^n\}$, the corresponding basis of vector fields by $\{\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}\}$ and the corresponding basis of 1-forms by $\{dx^1, \dots, dx^n\}$. Also, a frame of E[-1] is denoted by $\{e_1, \dots, e_m\}$ and its dual frame on $(E[-1])^{\vee}$ is denoted by $\{\xi^1, \dots, \xi^m\}$.

Observe that the graded algebra of smooth functions $C^{\infty}(E[-1]) = \Gamma(\Lambda^{-\bullet}E^{\vee})$ on the graded manifold $\mathcal{M} = E[-1]$ is generated by $\{x^1, \dots, x^n, \xi^1, \dots, \xi^m\}$. Then we may identify $\frac{\partial}{\partial x^i}$ and e_j as an element of $\operatorname{Der}(C^{\infty}(E[-1]))$ satisfying

$$\frac{\partial}{\partial x^i}(x^k) = \delta_i^k, \quad \frac{\partial}{\partial x^i}(\xi^k) = 0, \quad e_j(x^k) = 0, \quad e_j(\xi^k) = \delta_j^k$$

where δ_i^j is the Kronecker delta. Thus the set $\{e_1, \dots, e_m, \frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}\}$ can be identified with a frame of the tangent bundle TE[-1] over E[-1].

Remark 3.18. More precisely, under the isomorphism in Lemma 3.13, the element $\frac{\partial}{\partial x^i} \in \text{Der}(C^{\infty}(E[-1]))$ is, in fact, a horizontal lift of $\frac{\partial}{\partial x^i} \in \mathfrak{X}(M)$ with respect to the connection $\nabla^E : \Gamma(TM) \times \Gamma(E) \to \Gamma(E)$ satisfying $\nabla^E_{\frac{\partial}{\partial x^i}} e_j = 0$ for all i, j.

One can naturally construct an affine connection ∇^0 on E[-1] which is trivial with respect to this frame. That is, $\nabla^0 : \mathfrak{X}(E[-1]) \times \mathfrak{X}(E[-1]) \to \mathfrak{X}(E[-1])$ satisfies

$$\nabla^{0}_{e_{i}}e_{j} = 0, \quad \nabla^{0}_{\frac{\partial}{\partial x^{i}}}e_{j} = 0, \quad \nabla^{0}_{e_{i}}\frac{\partial}{\partial x^{j}} = 0, \quad \nabla^{0}_{\frac{\partial}{\partial x^{i}}}\frac{\partial}{\partial x^{j}} = 0$$

for all possible i, j. In fact, by Lemma 3.14 and Remark 3.18, one can show that ∇^0 corresponds to $(\nabla^E, \nabla^M, \beta)$ where ∇^E and ∇^M are trivial connections and $\beta = 0$. Thus, ∇^0 is a torsion-free affine connection on E[-1].

Suppose that a section $s = s^1 \cdot e_1 + \cdots + s^m \cdot e_m \in \Gamma(E[-1])$ is given. By Proposition 3.17, the Atiyah cocycle $\operatorname{At}^{\nabla^0}$ is completely determined by

$$\operatorname{At}^{\nabla^0}\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right) = \sum_k \frac{\partial^2 s^k}{\partial x^i \partial x^j} \cdot e_k$$

Under the identification $\Gamma(\mathcal{M}; \operatorname{Hom}(S^2(T\mathcal{M}), \mathcal{M})) \cong \Gamma(\mathcal{M}; S^2(T^{\vee}\mathcal{M}) \otimes \mathcal{M})$, we may write

$$\operatorname{At}^{\nabla^{0}} = \sum_{i \leq j} \sum_{k} \frac{\partial^{2} s^{k}}{\partial x^{i} \partial x^{j}} \cdot dx^{i} \odot dx^{j} \otimes e_{k} \in \Gamma(M; S^{2}(T^{\vee}M) \otimes (E[-1])^{\vee}).$$
(13)

Now, the differential Q restricted to degree 0 has a decomposition

$$\mathcal{Q} = d_1 + d_2 + d_3 : \Gamma(\mathcal{M}; S^2(T^{\vee}\mathcal{M}) \otimes \mathcal{M})^0 \to \Gamma(\mathcal{M}; S^2(T^{\vee}\mathcal{M}) \otimes \mathcal{M})^1$$

as in Section 3.3.2, and they are explicitly written as

$$d_1(\xi^l \otimes dx^i \odot dx^j \otimes e_k) = s^l \cdot dx^i \odot dx^j \otimes e_k \tag{14}$$

$$d_2(\xi^l \otimes dx^j \otimes e_k) = \sum_{r=1}^n \partial_r s^l \cdot dx^r \odot dx^j \otimes e_k \tag{15}$$

$$d_3(dx^i \odot dx^j \otimes \partial_l) = \sum_{r=1}^m \partial_l s^r \cdot dx^i \odot dx^j \otimes e_r$$
(16)

for each l, i, j, k.

4. PROOF OF THEOREM 3.2

The goal of this section is to prove Theorem 3.2, which states that the Atiyah class of a dg manifold $(E[-1], \iota_s)$ vanishes if and only if the section $s : M \to E$ intersects with the zero section cleanly.

By Proposition 3.9, it suffices to consider the Atiyah class on a neighbourhood of each $p \in M$. Moreover, by Corollary 3.16, it suffices to consider the neighbourhood of each $p \in s^{-1}(0)$. Finally, by Lemma 3.8, the clean intersection of s along the zero section is characterised by Eq. (3). Thus, Theorem 3.2 is an immediate consequence of the following lemma.

Lemma 4.1. Let M be a smooth manifold of dimension n and let E be a rank m vector bundle over M. Let $s \in \Gamma(E[-1])$ be a section. For the dg manifold $(E[-1], \iota_s)$, the following are equivalent.

- (L1) For each $p \in M$ with s(p) = 0, there exists a neighbourhood $U \subset M$ of p such that the Atiyah class of $(E|_U[-1], \iota_{s|_U})$ vanishes.
- (L2) For each $p \in M$ with s(p) = 0, there exists a neighbourhood $U \subset M$ of p and a local chart $\phi: U \to \mathbb{R}^n$ such that

$$s \circ \phi^{-1}|_{\phi(U)} = x^1 \cdot e_1 + \dots + x^r \cdot e_r$$

for some frame $\{e_1, \dots, e_m\}$ of $E|_U$ where r is the rank of Ds_p as in (2) and x^1, \dots, x^n are the coordinate functions on \mathbb{R}^n .

The rest of this paper is devoted to proving Lemma 4.1. Observe that the proof of $(L2) \Rightarrow (L1)$ is easy to prove. Indeed, first observe that the diffeomorphism $\psi := \phi^{-1}|_{\phi(U)}$ induces an isomorphism of dg manifolds $(E|_U[-1], \iota_s) \cong (\psi^* E|_{\phi(U)}, \iota_{so\psi})$. Thus, it suffices to prove that the Atiyah class of $(\psi^* E|_{\phi(U)}, \iota_{so\psi})$ vanishes. By the formula (13), the Atiyah cocycle associated with the trivial connection with respect to the local frame vanishes, and hence the Atiyah class vanishes.

Before we begin the proof of $(L1) \Rightarrow (L2)$, we fix some conventions and notations. For the proof, without loss of generality, we may assume $M = \mathbb{R}^n$ and $E = \mathbb{R}^n \times \mathbb{R}^m$ and the point $p \in M$ will always be identified with the origin $0 \in \mathbb{R}^n$. Thus, s(0) = 0.

In the proof, we will mostly use the notations from Section 3.3.3. The standard coordinate functions on M will be denoted by $\{x^1, \dots, x^n\}$, the corresponding basis of vector fields will be denoted by $\{\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}\}$ and that of differential forms will be denoted by $\{dx^1, \dots, dx^n\}$. For simplicity, we sometimes write ∂_i instead of $\frac{\partial}{\partial x^i}$. The standard frame of E[-1] will be denoted by $\{e_1, \dots, e_m\}$ and its dual frame on $(E[-1])^{\vee}$ will be denoted by $\{\xi^1, \dots, \xi^m\}$. For a section $s \in \Gamma(E[-1])$, we write $s = s^1 \cdot e_1 + \dots + s^m \cdot e_m$ where $s^i \in C^{\infty}(M)$ for each i.

Finally, the symbols d_1 , d_2 and d_3 are used to mean the maps defined in (14) - (16).

The proof of $(L1) \Rightarrow (L2)$ for arbitrary $n = \dim M$ and $m = \operatorname{rank} E$ is full of complicated and technical computations. To show the idea of the proof, we begin with two sample cases.

4.1. **Proof of** $(L1) \Rightarrow (L2)$: sample cases.

4.1.1. *Case 1*. dim M = 1 and rank E = 1.

For simplicity, we write $x = x^1$, $e = e_1$ and $\xi = \xi^1$. Also, we write $s = s(x) \cdot e$.

In this case, observe that

- (1) the Atiyah cocycle with respect to the trivial connection is $s''(x) \cdot dx \odot dx \otimes e$,
- (2) the image of d_1 is $C^{\infty}(\mathbb{R})$ -generated by $s(x) \cdot dx \odot dx \otimes e$,
- (3) and the image of d_2 and d_3 are both generated by $s'(x) \cdot dx \odot dx \otimes e$.

Thus, if there exists a neighbourhood U of $0 \in M$ such that the Atiyah class of $(E|_U[-1], \iota_{s|_U})$ vanishes then s''(x) is generated by s(x) and s'(x) in U. In other words, the condition (L1) implies that there exists an open neighbourhood U of $0 \in M$ such that for some $f, g \in C^{\infty}(U)$, we have an equation

$$s''(x) = f(x) \cdot s(x) + g(x) \cdot s'(x)$$
(17)

for all $x \in U$.

Suppose $s'(0) \neq 0$. Then by Lemma 3.7, the function $s : \mathbb{R} \to \mathbb{R}$ in a neighbourhood of 0 can be written as s(x) = x, up to a local diffeomorphism. Thus, we may assume that s(x) = x. Since s''(x) = 0, by setting f(x) = g(x) = 0, the Atiyah class vanishes.

Suppose s'(0) = 0. We claim that the Eq. (17) holds for some f, g in a neighbourhood U of 0 only when $s(x) \equiv 0$.

In fact, by setting y(t) = (s(t), s'(t)) and F(t, a, b) = (b, f(t)a + g(t)b), the uniqueness of ODE (a.k.a. Cauchy–Lipschitz, Picard–Lindelöf) states that the initial value problem

$$(s'(t), s''(t)) = y'(t) = F(t, y(t)) = (s'(t), f(t)s(t) + g(t)s'(t)), \quad y(0) = (s(0), s'(0)) = (0, 0)$$

has a unique solution near 0. Since $s(x) \equiv 0$ is one of the solution, by the uniqueness, this is the only possibility. This completes the proof for Case 1.

4.1.2. *Case 2.*
$$M = \mathbb{R}^2$$
, $E = \mathbb{R}^2 \times \mathbb{R}^2$ and rank $Ds_0 = 1$

Denote the section $s \in \Gamma(E[-1])$ with s(0) = 0 by $s = s^1 \cdot e_1 + s^2 \cdot e_2 = (s^1, s^2)$. The condition rank $Ds_0 = 1$ means that the 2×2 matrix

$$\begin{bmatrix} \frac{\partial s^i}{\partial x^j}(0) \end{bmatrix}_{ij} = \begin{bmatrix} \partial_1 s^1(0) & \partial_2 s^1(0) \\ \partial_1 s^2(0) & \partial_2 s^2(0) \end{bmatrix}$$

has rank 1. By the inverse function theorem (or, by [22, Theorem 2.9]), we may assume that $s^1 = x^1$ and $\partial_2 s^2(0) = 0$.

For simplicity, we use the notation

$$\frac{\partial^2 s^k}{\partial x^i \partial x^j} = \partial_{ij} s^k$$

By (13), the Atiyah cocycle with respect to the trivial connection is

$$At = \sum_{i \le j} \sum_{k} \partial_{ij} s^k \cdot dx^i \odot dx^j \otimes e_k$$

17

and it can be written as a vector

$$At = \begin{vmatrix} \partial_{11}s^{1} \\ \partial_{12}s^{1} \\ \partial_{22}s^{1} \\ \partial_{11}s^{2} \\ \partial_{12}s^{2} \\ \partial_{22}s^{2} \end{vmatrix}$$

with respect to the ordered basis

 $(dx^1 \odot dx^1 \otimes e_1, dx^1 \odot dx^2 \otimes e_1, dx^2 \odot dx^2 \otimes e_1, dx^1 \odot dx^1 \otimes e_2, dx^1 \odot dx^2 \otimes e_2, dx^2 \odot dx^2 \otimes e_2).$

With the same ordered basis, the Eqs. (14)-(16) shows that the coboundary elements are in the image of the linear transformations induced by the left multiplication of the following 3 matrices

$$d_{1} = \begin{bmatrix} s^{1} & 0 & 0 & 0 & 0 & 0 & s^{2} & 0 & 0 & 0 & 0 & 0 \\ 0 & s^{1} & 0 & 0 & 0 & 0 & 0 & s^{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & s^{1} & 0 & 0 & 0 & 0 & 0 & s^{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & s^{1} & 0 & 0 & 0 & 0 & 0 & s^{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & s^{1} & 0 & 0 & 0 & 0 & 0 & s^{2} & 0 \\ 0 & 0 & 0 & 0 & s^{1} & 0 & 0 & 0 & 0 & 0 & s^{2} \end{bmatrix}$$
$$d_{2} = \begin{bmatrix} \partial_{1}s^{1} & 0 & 0 & 0 & \partial_{1}s^{2} & 0 & 0 & 0 \\ \partial_{2}s^{1} & \partial_{1}s^{1} & 0 & 0 & \partial_{2}s^{2} & \partial_{1}s^{2} & 0 & 0 \\ 0 & \partial_{2}s^{1} & \partial_{1}s^{1} & 0 & 0 & 0 & \partial_{2}s^{2} & 0 & 0 \\ 0 & 0 & \partial_{1}s^{1} & 0 & 0 & 0 & \partial_{1}s^{2} & 0 \\ 0 & 0 & \partial_{2}s^{1} & \partial_{1}s^{1} & 0 & 0 & 0 & \partial_{2}s^{2} & \partial_{1}s^{2} \\ 0 & 0 & 0 & \partial_{2}s^{1} & 0 & 0 & 0 & \partial_{2}s^{2} \end{bmatrix}$$

and

$$d_{3} = \begin{bmatrix} \partial_{1}s^{1} & \partial_{2}s^{1} & 0 & 0 & 0 & 0 \\ 0 & 0 & \partial_{1}s^{1} & \partial_{2}s^{1} & 0 & 0 \\ 0 & 0 & 0 & 0 & \partial_{1}s^{1} & \partial_{2}s^{1} \\ \partial_{1}s^{2} & \partial_{2}s^{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & \partial_{1}s^{2} & \partial_{2}s^{2} & 0 & 0 \\ 0 & 0 & 0 & 0 & \partial_{1}s^{2} & \partial_{2}s^{2} \end{bmatrix}$$

Since $s^1 = x^1$, the Atiyah cocycle is

$$At = \begin{bmatrix} 0\\0\\0\\\partial_{11}s^2\\\partial_{12}s^2\\\partial_{22}s^2 \end{bmatrix}$$

and the matrix d_2 is

$$d_2 = \begin{bmatrix} 1 & 0 & 0 & 0 & \partial_1 s^2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & \partial_2 s^2 & \partial_1 s^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \partial_2 s^2 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & \partial_1 s^2 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & \partial_2 s^2 & \partial_1 s^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \partial_2 s^2 \end{bmatrix}$$

by applying $s^1 = x^1$. Because of the first 4 columns of d_2 , the 1st, 2nd, 4th and 5th rows of At is always in the image of $Q = d_1 + d_2 + d_3$. Thus, to check whether the Atiyah cocycle At is in the coboundary, it suffices to consider 3rd and 6th rows. In other words, it now simplifies to the question of whether $\begin{bmatrix} 0\\ \partial_{22}s^2 \end{bmatrix}$ is in the image of the linear transformation induced by the left multiplication of the matrix

$$\begin{bmatrix} x^1 & 0 & s^2 & 0 & \partial_2 s^2 & 0 & 1 & 0 \\ 0 & x^1 & 0 & s^2 & 0 & \partial_2 s^2 & \partial_1 s^2 & \partial_2 s^2 \end{bmatrix}$$

where first 4 columns are from d_1 , column 5 and 6 are from d_2 and the rest 2 columns are from d_3 .

Suppose that the Atiyah class vanishes. Then there exists a set of functions $\{f_1, \dots, f_8\}$ such that

$$\begin{bmatrix} 0\\ \partial_{22}s^2 \end{bmatrix} = f_1 \begin{bmatrix} x\\ 0 \end{bmatrix} + f_2 \begin{bmatrix} 0\\ x \end{bmatrix} + f_3 \begin{bmatrix} s^2\\ 0 \end{bmatrix} + f_4 \begin{bmatrix} 0\\ s^2 \end{bmatrix} + f_5 \begin{bmatrix} \partial_2s^2\\ 0 \end{bmatrix} + f_6 \begin{bmatrix} 0\\ \partial_2s^2 \end{bmatrix} + f_7 \begin{bmatrix} 1\\ \partial_1s^2 \end{bmatrix} + f_8 \begin{bmatrix} 0\\ \partial_2s^2 \end{bmatrix}$$
(18)

holds. The 1st row of the Eq. (18) is equivalent to

$$f_7 = -x^1 \cdot f_1 - s^2 \cdot f_3 - f_5 \cdot \partial_2 s^2.$$

This means that f_7 is a $C^{\infty}(M)$ -linear combination of x^1 , s^2 and $\partial_2 s^2$. Together with the 2nd row of (18), it implies that $\partial_{22}s^2$ is also a $C^{\infty}(M)$ -linear combination of x^1 , s^2 and $\partial_2 s^2$.

Recall that we started with the conditions s(0) = 0 and $\partial_2 s^2(0) = 0$. If the Atiyah class vanishes, then we have

$$\partial_{22}s^2(x_1, x_2) \in \langle x^1, s^2(x_1, x_2), \partial_2 s^2(x_1, x_2) \rangle$$

and in particular, when $x_1 = 0$, we have

$$\partial_{22}s^2(0,x_2) \in \langle s^2(0,x_2), \partial_2 s^2(0,x_2) \rangle$$

where the symbol $\langle a, b \rangle$ denotes the ideal generated by functions $a, b \in C^{\infty}(M)$. That is, there exist functions $f, g \in C^{\infty}(\mathbb{R})$ such that

$$\partial_{22}s^2(0, x_2) = f(x_2) \cdot s^2(0, x_2) + g(x_2) \cdot \partial_2 s^2(0, x_2)$$

with the initial condition $\partial_2 s^2(0,0) = 0$ and $s^2(0,0) = 0$. The uniqueness theorem of ODE implies that $s^2(0, x_2) \equiv 0$ is the only solution. Then, by Lemma 3.7, the proof for Case 2 is completed.

4.2. **Proof of** $(L1) \Rightarrow (L2)$: general case. $M = \mathbb{R}^n$, $E = \mathbb{R}^n \times \mathbb{R}^m$ and rank $Ds_0 = r$.

Finally we move on to the general case. Observe that the proof of Case 1 and Case 2 consists of 2 steps.

Step 1. We pick up the essential information from d_1 , d_2 and d_3 , and reduce to the problem of ODE. Step 2. Use the uniqueness of ODE to complete the proof.

Assume that $M = \mathbb{R}^n$, $E = \mathbb{R}^n \times \mathbb{R}^m$ and rank $Ds_0 = r$. Again, by the inverse function theorem (or, by [22, Theorem 2.9]), we may assume that the given section $s = s^1 \cdot e_1 + \cdots + s^m \cdot e_m$ satisfies $s^1 = x^1, \cdots, s^r = x^r$. By the condition on rank, we have $\partial_i s^j(0) = 0$ for all i, j > r.

By (13), the Atiyah cocycle associated with the trivial connection is

$$At = \sum_{i \le j} \sum_{k} \frac{\partial^2 s^k}{\partial x^i \partial x^j} \cdot dx^i \odot dx^j \otimes e_k = \sum_{i \le j} \sum_{k} \partial_{ij} s^k \cdot dx^i \odot dx^j \otimes e_k$$

and the formulas of $Q = d_1 + d_2 + d_3$ are described in Eqs. (14)-(16).

Claim 1. If the Atiyah class of the dg manifold $(E[-1], \iota_s)$ vanishes, then for each triple (i, j, k) with i, j, k > r, we have

$$\partial_{ij}s^k \in \langle s^1, \cdots, s^m, \partial_i s^{r+1}, \cdots, \partial_i s^m, \partial_j s^{r+1}, \cdots, \partial_j s^m, \partial_{r+1}s^k, \cdots, \partial_n s^k \rangle$$

where $\langle a, b \rangle$ denotes the ideal generated by functions $a, b \in C^{\infty}(M)$.

Proof of Claim 1. Similarly to Case 2, we first investigate the image of d_2 . Observe that if $i \leq r$, then we have

$$d_2(\xi^i \otimes dx^j \otimes e_k) = dx^i \odot dx^j \otimes e_k$$

and similarly, if $j \leq r$, then we have

$$d_2(\xi^j \otimes dx^i \otimes e_k) = dx^i \odot dx^j \otimes e_k.$$

This implies that the component $\partial_{ij}s^k \cdot dx^i \odot dx^j \otimes e_k$ is always in the coboundary if either $i \leq r$ or $j \leq r$. Thus, it suffices to consider when both i, j > r.

SEOKBONG SEOL

Consider an element $b = \sum_{\alpha \leq \beta} \sum_{\gamma} c_{\alpha\beta}^{\gamma} dx^{\alpha} \odot dx^{\beta} \otimes e_{\gamma}$. Then, for a fixed triple (i, j, k) with i, j > r, the following must satisfy.

- By (14), if b ∈ im d₁, then the function c^k_{ij} is generated by {s¹, ..., s^m}.
 By (15), if b ∈ im d₂, then the function c^k_{ij} is generated by {∂_is^{r+1}, ..., ∂_is^m, ∂_js^{r+1}, ..., ∂_js^m}.
 By (16), if b ∈ im d₃, then the function c^k_{ij} is generated by {∂₁s^k, ..., ∂_ns^k}.

Thus, if the Atiyah class vanishes, then for each i, j, k with i, j > r,

г

$$\partial_{ij}s^k \in \langle s^1, \cdots, s^m, \partial_i s^{r+1}, \cdots, \partial_i s^m, \partial_j s^{r+1}, \cdots, \partial_j s^m, \partial_1 s^k, \cdots, \partial_n s^k \rangle .$$

Fix a triple (i, j, k') with i, j > r and $k' \leq r$. Since $s^{k'} = x^{k'}$, we have $\partial_l s^{k'} = 0$ if $l \neq k'$. The only coboundary of d_3 which has non-zero coefficient of $dx^i \odot dx^j \otimes e_{k'}$ comes from

$$d_3(dx^i \odot dx^j \otimes \partial_{k'}) = dx^i \odot dx^j \otimes e_{k'} + \sum_{p>r} \partial_{k'} s^p \cdot dx^i \odot dx^j \otimes e_p.$$
(19)

Moreover, note that d_3 is the only map that changes the index k of e_k . Summing up, our situation can be pictured as a matrix

$$At = \begin{bmatrix} \vdots \\ \partial_{ij}s^{a} \\ \vdots \\ \partial_{ij}s^{b} \\ \vdots \end{bmatrix} \leftrightarrow \begin{bmatrix} \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \cdots & s^{l} & 0 & \cdots & \partial_{i}s^{l} & 0 & \cdots & \partial_{a}s^{a}(=1) & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \cdots & 0 & s^{l} & \cdots & 0 & \partial_{i}s^{l} & \cdots & \partial_{a}s^{b} & \partial_{a+1}s^{b} & \cdots \\ \vdots & \vdots \end{bmatrix}$$

where $a \leq r < b, i, j > r$ and $1 \leq l \leq m$. In particular, if the Atiyah class vanishes, then LHS is the image of the linear transformation induced by the left multiplication of the matrix on the RHS — compare with (18).

By the hypothesis, the Atiyah cocycle At must be in the image of $Q = d_1 + d_2 + d_3$. In particular, for each fixed (i, j, k') with i, j > r and $k' \le r$,

$$0 = \partial_{ij}s^{k'} = \sum_{p=1}^{m} f_{ij,p}^{k'} \cdot s^p + \sum_{q=r+1}^{m} (g_{ij,q}^{k'} \cdot \partial_i s^q + \tilde{g}_{ij,q}^{k'} \cdot \partial_j s^q) + h_{ij}^{k',k'} \cdot \partial_{k'}s^k$$

for some $f_{ij,p}^{k'}, g_{ij,q}^{k'}, \tilde{g}_{ij,q}^{k'}, h_{ij}^{k',k'} \in C^{\infty}(M)$. Since $\partial_{k'}s^{k'} = 1$ for $k' \leq r$, we may conclude that

$$h_{ij}^{k',k'} \in \langle s^1, \cdots, s^m, \partial_i s^{r+1}, \cdots, \partial_i s^m, \partial_j s^{r+1}, \cdots, \partial_j s^m \rangle$$
(20)

if $k' \leq r$. Similarly, for each fixed (i, j, k) with i, j > r and k > r,

$$\partial_{ij}s^k = \sum_{p=1}^m f^k_{ij,p} \cdot s^p + \sum_{q=r+1}^m (g^k_{ij,q} \cdot \partial_i s^q + \tilde{g}^k_{ij,q} \cdot \partial_j s^q) + \sum_{t=1}^n h^{k,t}_{ij} \cdot \partial_t s^k$$

for some $f_{ij,p}^k$, $g_{ij,q}^k$, $\tilde{g}_{ij,q}^k$, $h_{ij}^{k,t} \in C^{\infty}(M)$. Moreover, by the condition (19), it must satisfy $h_{ij}^{k,t} = h_{ij}^{t,t}$ for all $t \leq r$. Because of (20), we conclude that

$$\partial_{ij}s^k \in \langle s^1, \cdots, s^m, \partial_i s^{r+1}, \cdots, \partial_i s^m, \partial_j s^{r+1}, \cdots, \partial_j s^m, \partial_{r+1} s^k, \cdots, \partial_n s^k \rangle.$$

This proves the claim.

٦

The above claim is equivalent to the following: if the Atiyah class of $(E[-1], \iota_s)$ vanishes, then for each (i, j, k) with i, j, k > r, it satisfies

$$\partial_{ij}s^k = \sum_{p=1}^m F^k_{ij,p} \cdot s^p + \sum_{q=r+1}^m (G^k_{ij,q} \cdot \partial_i s^q + \tilde{G}^k_{ij,q} \cdot \partial_j s^q) + \sum_{t=r+1}^n H^{k,t}_{ij} \cdot \partial_t s^k$$
(21)

for some $F_{ij,p}^k, G_{ij,q}^k, \tilde{G}_{ij,q}^k, H_{ij}^{k,t} \in C^\infty(M).$

By assumption, we have $s^1 = x^1, \dots, s^r = x^r$. Let $X \in \{0\}^r \times \mathbb{R}^{n-r} \subset \mathbb{R}^n$. Then the condition (21) induces that for each (i, j, k) with i, j, k > r,

$$\partial_{ij}s^{k}(X) = \sum_{p=r+1}^{m} F_{ij,p}^{k}(X) \cdot s^{p}(X) + \sum_{q=r+1}^{m} (G_{ij,q}^{k}(X) \cdot \partial_{i}s^{q}(X) + \tilde{G}_{ij,q}^{k}(X) \cdot \partial_{j}s^{q}(X)) + \sum_{t=r+1}^{n} H_{ij}^{k,t}(X) \cdot \partial_{t}s^{k}(X) \quad (22)$$

for some $F_{ij,p}^k, G_{ij,q}^k, \tilde{G}_{ij,q}^k, H_{ijk}^{k,t} \in C^{\infty}(M)$. Moreover, by assumption, we have

$$s^{\alpha}(0) = 0, \qquad \partial_{\beta}s^{\alpha}(0) = 0 \tag{23}$$

for all $\alpha, \beta > r$.

Claim 2. Suppose there exist functions $F_{ij,p}^k, G_{ij,q}^k, \tilde{G}_{ij,q}^k, H_{ij}^{k,t} \in C^{\infty}(M)$ with i, j, k, p, q, t > r. If Eq. (22) and Eq. (23) holds for all $i, j, k, \alpha, \beta > r$, then $s^k|_{\{0\}^r \times \mathbb{R}^{n-r}} = 0$ for all k > r.

Proof of Claim 2. Note first that all indices are larger than r. Thus, it suffices to consider when r = 0.

The claim follows from the uniqueness theorem of ODE. Assume that there exist $\nu = (v^1, \dots, v^n) \in \mathbb{R}^n$ such that $s^k(\nu) \neq 0$ for some k. Then, consider $y_{\nu} : \mathbb{R} \to \mathbb{R}^{m+nm}$ defined by

$$y_{\nu}(t) = (s^{1}(t \cdot \nu), \cdots, s^{m}(t \cdot \nu), \partial_{1}s^{1}(t \cdot \nu), \cdots, \partial_{1}s^{m}(t \cdot \nu), \partial_{2}s^{1}(t \cdot \nu), \cdots, \partial_{n}s^{m}(t \cdot \nu))$$

and a smooth function $F_\nu:\mathbb{R}^{1+m+nm}\to\mathbb{R}^{m+nm}$ defined by

$$\mathcal{F}_{\nu}(t, a_1, \cdots, a_m, b_{11}, \cdots, b_{1m}, b_{21}, \cdots, b_{nm}) = (c_1, \cdots, c_m, d_{11}, \cdots, d_{1m}, d_{21}, \cdots, d_{nm})$$

where

$$c_{\alpha} = \sum_{i} v^{i} \cdot b_{i\alpha}$$

and

$$d_{\alpha\beta} = \sum_{i} v_i \cdot \left(\sum_{p} (F_{i\alpha,p}^{\beta} \cdot a_p) + \sum_{q} (G_{i\alpha,q}^{\beta} b_{iq} + \tilde{G}_{i\alpha,q}^{\beta} b_{\alpha q}) + \sum_{t} (H_{i\alpha}^{\beta,t} \cdot b_{t\beta}) \right).$$

Then $y'_{\nu}(t) = \mathcal{F}_{\nu}(t, y_{\nu}(t))$ is exactly Eq. (22) when $X = t \cdot \nu \in \mathbb{R}^n$, and Eq. (23) is the initial condition $y_{\nu}(0) = 0$. Thus, by the uniqueness of ODE, we have a unique solution $y_{\nu}(t) \equiv 0$. This contradicts to the assumption that $s^k(\nu) \neq 0$. This completes the proof.

Finally, by Lemma 3.7, the proof of the general case of $(L1) \Rightarrow (L2)$ is completed.

SEOKBONG SEOL

5. The case of derived intersections

Given two embedded submanifolds X and Y of a smooth manifold W, there exists a dg manifold of amplitude +1, which can be thought of as a model for the derived intersection $X \cap^h Y$ of X and Y — see [3] for detail. Below, we briefly recall its construction.

To construct a dg manifold $(\mathcal{M}, Q) = (E[-1], \iota_s)$ of amplitude +1, it suffices to construct a vector bundle $E \to M$ and a section $s : E \to M$. We begin with the construction of the base manifold M.

Upon a choice of an affine connection ∇ on W, consider the collection P_W of all geodesic paths $\gamma : [0,1] \to W$ whose domain is the unit interval [0,1] — we call such geodesics by short geodesics. By the existence and uniqueness of ODE, P_W is identified with an open subset of TW. Thus, P_W is a smooth manifold. Moreover, the evaluation maps $ev_0 : P_W \to W$ and $ev_1 : P_W \to W$, defined by $ev_0(\gamma) = \gamma(0)$ and $ev_1(\gamma) = \gamma(1)$ for $\gamma \in P_W$, are submersions. Therefore, the space $X \times_W P_W \times_W Y$ consisting of all short geodesics γ satisfying $\gamma(0) \in X$ and $\gamma(1) \in Y$ is a smooth manifold. We set the base manifold $M = X \times_W P_W \times_W Y$.

Next, the vector bundle E over M consists of all covariantly constant vector fields along short geodesics $\gamma \in M$. More precisely, as a space, a point in E is a path $\tilde{\gamma} : [0, 1] \to TW$ satisfying

(1)
$$\pi \circ \tilde{\gamma} = \gamma \in P_W$$
,

(2) $\nabla_{\dot{\gamma}(t)}\tilde{\gamma}(t) = 0$ for all $t \in [0, 1]$

where $\pi : TW \to W$ is the tangent bundle and $\dot{\gamma}(t)$ is the derivative of $\gamma : [0,1] \to W$ in the direction of t. Note that once $\gamma = \pi \circ \tilde{\gamma}$ and $\tilde{\gamma}(0) \in T_{\gamma(0)}W$ are chosen, then $\tilde{\gamma}$ is uniquely determined by the parallel transport along γ . Therefore E is identified with the pullback bundle $\operatorname{ev}_0^* TW$ of the tangent bundle TW over the evaluation map $\operatorname{ev}_0 : M \to W$ and thus, E carries the finite dimensional vector bundle structure over M.

Finally, we define the section $s: M \to E$ by $s(\gamma) = \dot{\gamma}$, hence obtain a dg manifold $(E[-1], \iota_s)$ arising from a pair of embedded submanifolds X and Y of W.

We note that, by [3, Theorem 3.4], the dg manifold $(E[-1], \iota_s)$ is independent of choice of affine connections ∇ on W, up to isomorphism of dg manifolds. Also, the zero locus $s^{-1}(0)$ of s consists of constant paths $\gamma: I \to W$ satisfying $\gamma(0) \in X$ and $\gamma(1) \in Y$; the zero locus $s^{-1}(0)$ is the space of intersection $X \cap Y$.

Now, it is natural to ask what the Atiyah class associated with the derived intersection measures. The following proposition provides an answer to this question.

Proposition 5.1. Given two embedded submanifolds X and Y, the Atiyah class of dg manifold $(E[-1], \iota_s)$ constructed above vanishes if and only if X and Y intersect cleanly.

Proof. By Proposition 3.9 and Corollary 3.16, it suffices to consider the Atiyah class of $(E|_U[-1], \iota_{s|_U})$ where U is a neighbourhood of $p \in M$ such that s(p) = 0. Thus, we may assume that $W = \mathbb{R}^n$ and X and Y are embedded image of $i : \mathbb{R}^l \to W$ and $j : \mathbb{R}^m \to W$, i.e. $X = i(\mathbb{R}^l)$ and $Y = j(\mathbb{R}^m)$. Moreover, since the construction of M is independent of choice of affine connection on W, we may choose a trivial connection ∇ on W.

Now, the base manifold $M = X \times_W P_W \times_W Y$ consists of elements of the form (i(x), i(x)+t(j(y)-i(x)), j(y))where $x \in \mathbb{R}^l$ and $y \in \mathbb{R}^m$. Since the maps $i : \mathbb{R}^l \to W$ and $j : \mathbb{R}^m \to W$ are embeddings, we have a diffeomorphism $\mathbb{R}^l \times \mathbb{R}^m \cong M$. The bundle E is the pullback bundle of TW over the evaluation map $ev_0 : M \to W$, thus, we have an isomorphism $E \cong (\mathbb{R}^l \times \mathbb{R}^m) \times \mathbb{R}^n$. Under the identifications, the section s is characterised by $s(x, y) = j(y) - i(x) \in \mathbb{R}^n$.

Suppose that X and Y intersect cleanly. Then, by [9, Proposition C.3.1], for each $p \in X \cap Y$, there exists a neighbourhood $U \subset W$ of p and a local chart $\phi : U \to \mathbb{R}^{\dim W}$ such that $\phi(X \cap U)$ and $\phi(Y \cap U)$ are vector spaces. Thus, we may assume that the embeddings $i : \mathbb{R}^l \to W$ and $j : \mathbb{R}^l \to W$ are linear maps. Thus

the section s is a linear section. By (13), an Atiyah cocycle is the 2nd derivatives of s. Therefore, the Atiyah class of $(E[-1], \iota_s)$ vanishes if X and Y intersect cleanly.

Conversely, suppose that the Atiyah class of $(E[-1], \iota_s)$ vanishes. Then, by Theorem 3.2, the section s and the zero section 0_E intersect cleanly. In particular, $s^{-1}(0) \cong X \cap Y$ is a manifold. Thus, it remains to show that $T_p(X \cap Y) = T_pX \cap T_pY$, for each $p \in X \cap Y$. In particular, since $T_p(X \cap Y) \subset T_pX \cap T_pY$, it suffices to show that dim $T_p(X \cap Y) \ge \dim(T_pX \cap T_pY)$.

On the one hand, by Corollary 3.4, we have the identity rank $Ds_p = l + m - \dim T_p(X \cap Y)$ for each $p \in X \cap Y$. On the other hand, the rank of Ds_p is bounded above by $l + m - \dim(T_pX \cap T_pY)$. Indeed, the domain of s has dimension l + m, and for each $v \in T_pX \cap T_pY \subset T_pW$, there exists $v_x \in \mathbb{R}^l$ and $v_y \in \mathbb{R}^m$ such that $Ds_p(v_x, v_y) = v - v = 0$. In fact, for $i(p_x) = j(p_y) = p \in X \cap Y$, the vectors v_x and v_y are the unique vectors satisfying $v = Ti_{p_x}(v_x) = Tj_{p_y}(v_y)$ where Ti_{p_x} is the tangent map of the embedding i at p_x and similarly for Tj_{p_y} . As a result, we have $\dim T_p(X \cap Y) \ge \dim(T_pX \cap T_pY)$, and this completes the proof.

Acknowledgement

The author would like to thank Zhuo Chen, Dongwook Choa, Dongnam Ko, Camille Laurant-Gengoux, Hsuan-Yi Liao, Mathieu Stiénon, Maosong Xiang and Ping Xu for their interest in this work and their helpful comments. Also, the author is grateful to National Center for Theoretical Sciences, Institut Henri Poincaré and Pennsylvania State University for their generous support and hospitality.

References

- [1] M. Alexandrov et al. "The geometry of the master equation and topological quantum field theory". In: *Internat. J. Modern Phys. A* 12.7 (1997), pp. 1405–1429.
- [2] M. F. Atiyah. "Complex analytic connections in fibre bundles". In: Trans. Amer. Math. Soc. 85 (1957), pp. 181–207.
- [3] Kai Behrend, Hsuan-Yi Liao, and Ping Xu. *Differential graded manifolds of finite positive amplitude*. 2023. arXiv: 2307.10242 [math.DG].
- [4] Kai Behrend, Hsuan-Yi Liao, and Ping Xu. On the structure of étale fibrations of L_{∞} -bundles. 2023. arXiv: 2307.08179 [math.DG].
- [5] Raoul Bott. "On the iteration of closed geodesics and the Sturm intersection theory". In: *Comm. Pure Appl. Math.* 9 (1956), pp. 171–206.
- [6] David Carchedi. Derived Manifolds as Differential Graded Manifolds. 2023. arXiv: 2303.11140 [math.DG].
- [7] Alberto S. Cattaneo and Florian Schätz. "Introduction to supergeometry". In: *Rev. Math. Phys.* 23.6 (2011), pp. 669–690.
- [8] Zhuo Chen, Maosong Xiang, and Ping Xu. "Atiyah and Todd classes arising from integrable distributions". In: *J. Geom. Phys.* 136 (2019), pp. 52–67.
- [9] Lars Hörmander. *The analysis of linear partial differential operators. III.* Classics in Mathematics. Pseudodifferential operators, Reprint of the 1994 edition. Springer, Berlin, 2007, pp. viii+525.
- [10] Mikhail Kapranov. "Rozansky-Witten invariants via Atiyah classes". In: Compositio Math. 115.1 (1999), pp. 71– 113.
- [11] Maxim Kontsevich. "Deformation quantization of Poisson manifolds". In: Lett. Math. Phys. 66.3 (2003), pp. 157–216.
- [12] Hsuan-Yi Liao. "Atiyah Classes and Todd Classes of Pullback dg Lie Algebroids Associated with Lie Pairs". In: *Comm. Math. Phys.* 404.2 (2023), pp. 701–734.
- [13] Hsuan-Yi Liao and Seokbong Seol. "Keller admissible triples and Duflo theorem". In: J. Math. Pures Appl. (9) 174 (2023), pp. 1–43.
- [14] Hsuan-Yi Liao and Mathieu Stiénon. "Formal exponential map for graded manifolds". In: Int. Math. Res. Not. IMRN 3 (2019), pp. 700–730.
- [15] Hsuan-Yi Liao, Mathieu Stiénon, and Ping Xu. "Formality theorem for differential graded manifolds". In: C. R. Math. Acad. Sci. Paris 356.1 (2018), pp. 27–43.

REFERENCES

- [16] S. L. Lyakhovich, E. A. Mosman, and A. A. Sharapov. "Characteristic classes of *Q*-manifolds: classification and applications". In: *J. Geom. Phys.* 60.5 (2010), pp. 729–759.
- [17] Rajan Amit Mehta. *Supergroupoids, double structures, and equivariant cohomology*. Thesis (Ph.D.)–University of California, Berkeley. ProQuest LLC, Ann Arbor, MI, 2006, p. 133.
- [18] Rajan Amit Mehta, Mathieu Stiénon, and Ping Xu. "The Atiyah class of a dg-vector bundle". In: C. R. Math. Acad. Sci. Paris 353.4 (2015), pp. 357–362.
- [19] J. Milnor. *Morse theory*. Annals of Mathematics Studies, No. 51. Based on lecture notes by M. Spivak and R. Wells. Princeton University Press, Princeton, N.J., 1963, pp. vi+153.
- [20] Seokbong Seol, Mathieu Stiénon, and Ping Xu. "Dg manifolds, formal exponential maps and homotopy Lie algebras". In: *Comm. Math. Phys.* 391.1 (2022), pp. 33–76.
- [21] Boris Shoikhet. On the Duflo formula for L_{∞} -algebras and Q-manifolds. 1998. arXiv: math/9812009 [math.QA].
- [22] Michael Spivak. A comprehensive introduction to differential geometry. Vol. I. Second. Publish or Perish, Inc., Wilmington, Del., 1979, pp. xiv+668.
- [23] Mathieu Stiénon and Ping Xu. "Atiyah classes and Kontsevich-Duflo type theorem for DG manifolds". In: *Ho-motopy algebras, deformation theory and quantization*. Vol. 123. Banach Center Publ. Polish Acad. Sci. Inst. Math., Warsaw, 2021, pp. 63–110.

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY

Email address: azuredream89@kias.re.kr