A bridge between the circular and linear normal distributions

Frédéric Ouimet^a

^aDepartment of Mathematics and Statistics, McGill University, Montréal (Québec) Canada H3A 0B9

Abstract

In this short note, we present a refined approximation for the log-ratio of the density of the von $\text{Mises}(\mu, \kappa)$ distribution (also called the circular normal distribution) to the standard (linear) normal distribution when the concentration parameter κ is large. Our work complements the one of Hill (1976), who obtained a very similar approximation along with quantile couplings, using earlier approximations by Hill & Davis (1968) of Cornish-Fisher type. One motivation for this note is to highlight the connection between the circular and linear normal distributions through their circular variance and (linear) variance.

Keywords: asymptotic distribution, circular data, circular variance, circular normal distribution, circular statistics, directional statistics, local approximation, local limit theorem, normal approximation, variance, von Mises distribution, wrapped normal distribution 2020 MSC: Primary: 62E20 Secondary: 62E10

1. Introduction

For any reals $\mu \in [0, 2\pi)$ and $\kappa \in (0, \infty)$, the density function of the von Mises distribution (also referred to as the circular normal distribution) with mean direction μ and concentration parameter κ , henceforth denoted by von Mises (μ, κ) , is defined by

$$f_{\mu,\kappa}(\theta) = \frac{\exp\{\kappa\cos(\theta - \mu)\}}{2\pi I_0(\kappa)}, \quad \theta \in [0, 2\pi), \tag{1}$$

where I_{ν} denotes the modified Bessel function of the first kind of order $\nu \in \mathbb{N}_0 = \{0, 1, \ldots\}$. The circular expectation and variance of $\Theta \sim \text{von Mises}(\mu, \kappa)$ are well known to be

$$\mathsf{CE}(\Theta) \equiv \arg\{\mathsf{E}(e^{\mathsf{i}\Theta})\} = \mu \text{ and } \mathsf{CVar}(\Theta) \equiv 1 - \mathsf{E}\{\cos(\Theta - \mu)\} = 1 - \frac{I_1(\kappa)}{I_0(\kappa)} =: \nu_{\kappa}^2$$

see, e.g., Section 3.5.4 of Mardia & Jupp (2000). As $\kappa \to \infty$, note that

$$I_{0}(\kappa) \sim \frac{e^{\kappa}}{\sqrt{2\pi\kappa}} \left\{ 1 + \frac{1}{8\kappa} + \frac{9}{128\kappa^{2}} + \frac{75}{1024\kappa^{3}} + \mathcal{O}(\kappa^{-4}) \right\},$$

$$I_{1}(\kappa) \sim \frac{e^{\kappa}}{\sqrt{2\pi\kappa}} \left\{ 1 - \frac{3}{8\kappa} - \frac{15}{128\kappa^{2}} - \frac{105}{1024\kappa^{3}} + \mathcal{O}(\kappa^{-4}) \right\},$$
(2)

see, e.g., (Abramowitz & Stegun, 1964, p.377), so that

$$\nu_{\kappa}^{2} = 1 - \frac{I_{1}(\kappa)}{I_{0}(\kappa)} = \frac{1}{2\kappa} + \frac{1}{8\kappa^{2}} + \frac{1}{8\kappa^{3}} + \mathcal{O}(\kappa^{-4}), \quad \kappa \to \infty.$$
(3)

In particular, one has $\nu_{\kappa} \to 0$ as $\kappa \to \infty$.

Email address: frederic.ouimet2@mcgill.ca (Frédéric Ouimet)

The main goal of the present paper (Theorem 1) is to establish a local asymptotic expansion, as $\kappa \to \infty$, for the log-ratio of the von Mises density (1) to the (linear) Normal($\mu, 2\nu_{\kappa}^2$) density, viz.,

$$\frac{1}{\sqrt{2}\nu_{\kappa}}\phi\left(\frac{\delta_x}{\sqrt{2}}\right), \quad \text{where } \phi(z) := \frac{\exp(-z^2/2)}{\sqrt{2\pi}} \quad \text{and} \quad \delta_x := \frac{x-\mu}{\sqrt{1-I_1(\kappa)/I_0(\kappa)}}.$$
 (4)

Notice the extra scaling factor $\sqrt{2}$ for the (linear) standard deviation in (4). This factor would not appear if the expected limiting distribution was the usual Normal (μ, ν_{κ}^2) distribution.

To explain the extra scaling factor, consider a wrapped normal distribution with mean direction $\mu \in [0, 2\pi)$ and scaling parameter $\sigma \in (0, \infty)$, henceforth denoted by WNormal (μ, σ^2) . Its density function is given by the infinite sum

$$g_{\mu,v^2}(\theta) = \sum_{k=-\infty}^{\infty} \frac{1}{\sigma\sqrt{2\pi}} \exp\left\{-\frac{1}{2\sigma^2}(\theta - \mu + 2\pi k)^2\right\}, \quad \theta \in [0, 2\pi),$$
(5)

see, e.g., p. 50-51 of Mardia & Jupp (2000) or Section 2.2.6 of Jammalamadaka & SenGupta (2001); it has the same support as the von Mises distribution. The WNormal(μ, σ^2) distribution can be visualized as a Normal(μ, σ^2) distribution "folded" onto the unit circle. When the scaling parameter σ of the underlying normal distribution is small, most of the distribution's mass will be located near the mean direction μ , minimizing the "wrapping" effect. Indeed, the expression (5) essentially sums the normal densities from each "wrap" around the circle. For a small σ , the majority of the distribution's weight is around the mean direction μ , and the contributions from terms with $k \neq 0$ (i.e., terms from subsequent wraps around the circle) will be minimal. It is well known to be a limiting distribution of the von Mises(μ, κ) distribution when the concentration parameter κ tends to infinity, see Kent (1978) or Eq. (3.5.24) of Mardia & Jupp (2000) for details. The circular expectation and variance of $\Theta \sim WNormal(\mu, \sigma^2)$ are

$$\mathsf{CE}(\Theta) \equiv \arg\{\mathsf{E}(e^{i\Theta})\} = \mu$$
 and $\mathsf{CVar}(\Theta) \equiv 1 - \mathsf{E}\{\cos(\Theta - \mu)\} = 1 - \exp(-\sigma^2/2).$

The circular variance provides a measure of the dispersion of the distribution on the circle and is derived using the so-called mean resultant length, see, e.g., Section 2.3.1 of Mardia & Jupp (2000). For small σ , the mean resultant length is close to 1. Specifically, when σ^2 is small, one has the first-order Taylor approximation

$$\operatorname{CVar}(\Theta) = 1 - \exp(-\sigma^2/2) \approx \frac{\sigma^2}{2} = \left(\frac{\sigma}{\sqrt{2}}\right)^2,$$

where the aforementioned extra scaling factor $\sqrt{2}$ appears again. The asymptotic relationship between the circular variance of the von Mises distribution and the corresponding limiting wrapped normal distributions is therefore

$$\nu_{\kappa} = \frac{\sigma}{\sqrt{2}}, \quad \text{or equivalently}, \quad \sigma = \sqrt{2}\nu_{\kappa}.$$

This explains the extra scaling factor $\sqrt{2}$ in (4); it serves as a pivotal scaling relationship between the unit circle and the real line. In essence, it accounts for the additional variance caused by the subsequent wraps ($k \neq 0$) around the unit circle. This connection is also mentioned for example in Eq. (3.4.16) of Mardia & Jupp (2000).

2. Main results

Below, we state and prove a local approximation for the ratio of the von Mises density to the linear normal density function with appropriate mean and variance. A very similar approximation was proved decades ago by Hill (1976) along with quantile couplings, using earlier approximations by Hill & Davis (1968) of Cornish-Fisher type. Regarding other univariate continuous distributions, local approximations analogous to Theorem 1 and Corollary 2 below can be found for the central/noncentral chi-square distribution in Ouimet (2022a) and for the Student distribution in Ouimet (2022b).

Throughout the paper, the notation $u = \mathcal{O}(v)$ means that

$$\limsup_{\kappa \to \infty} |u/v| \le C \quad \text{or} \quad \limsup_{y \to \infty} |u/v| \le C$$

for some universal positive constant $C \in (0, \infty)$, depending on the context. Whenever C might depend on some parameter, we add a subscript. For example, $u = \mathcal{O}_{\eta}(v)$.

Theorem 1 (Local approximation). Let $\mu \in \mathbb{R}$ and $\eta \in (0, 1)$ be given. For any $\kappa \in (0, \infty)$,

$$B_{\mu,\kappa}(\eta) := \left\{ x \in (0,\infty) : |\delta_x| \le \eta \, \kappa^{1/2} \right\},\,$$

denotes the bulk of the von Mises distribution. Then, uniformly for $x \in B_{\mu,\kappa}(\eta)$, one has

$$\log\left\{\frac{\sqrt{2}\nu_{\kappa}f_{\mu,\kappa}(x)}{\phi(\tilde{\delta}_x)}\right\} = \frac{1}{\kappa}\left(\frac{\tilde{\delta}_x^4}{24} - \frac{\tilde{\delta}_x^2}{8}\right) + \frac{1}{\kappa^2}\left(-\frac{\tilde{\delta}_x^6}{720} + \frac{\tilde{\delta}_x^4}{48} - \frac{\tilde{\delta}_x^2}{8} + \frac{3}{64}\right) + \mathcal{O}_\eta\left(\frac{1+|\tilde{\delta}_x|^8}{\kappa^3}\right), \quad (6)$$

as $\kappa \to \infty$, where $\tilde{\delta}_x = \delta_x/\sqrt{2}$. Moreover, uniformly for $x \in B_{\mu,\kappa}(\kappa^{-1/4}\tilde{\eta})$ given some $\tilde{\eta} \in (0,\infty)$, one has

$$\frac{\sqrt{2}\nu_{\kappa}f_{\mu,\kappa}(x)}{\phi(\widetilde{\delta}_{x})} = 1 + \frac{1}{\kappa} \left(\frac{\widetilde{\delta}_{x}^{4}}{24} - \frac{\widetilde{\delta}_{x}^{2}}{8}\right) + \frac{1}{\kappa^{2}} \left(\frac{\widetilde{\delta}_{x}^{8}}{1152} - \frac{19\widetilde{\delta}_{x}^{6}}{2880} + \frac{11\widetilde{\delta}_{x}^{4}}{384} - \frac{\widetilde{\delta}_{x}^{2}}{8} + \frac{3}{64}\right) + \mathcal{O}_{\widetilde{\eta}} \left(\frac{1 + |\widetilde{\delta}_{x}|^{12}}{\kappa^{3}}\right).$$

$$\tag{7}$$

Open problem. Extend the result of Theorem 1 to the more general (spherical) von Mises-Fisher distribution. For the definition, see, e.g., Section 9.3.2 of Mardia & Jupp (2000).

Next, as a corollary, we integrate the local density approximation of Theorem 1 to obtain a normal local approximation for the von Mises cumulative distribution function.

Corollary 2. Using the same notation as in Theorem 1, one has, uniformly for $x \in B_{\mu,\kappa}(\eta)$ and as $\kappa \to \infty$,

$$F_{\mu,\kappa}(x) := \int_{-\infty}^{x} f_{\mu,\kappa}(y) dy$$
$$= \Phi(\widetilde{\delta}_{x}) + \phi(\widetilde{\delta}_{x}) \left\{ \frac{1}{\kappa} \left(\frac{\widetilde{\delta}_{x}^{3}}{24} \right) + \frac{1}{\kappa^{2}} \left(\frac{\widetilde{\delta}_{x}^{7}}{1152} - \frac{\widetilde{\delta}_{x}^{5}}{1920} + \frac{5\widetilde{\delta}_{x}^{3}}{192} - \frac{3\widetilde{\delta}_{x}}{64} \right) \right\} + \mathcal{O}_{\widetilde{\eta}} \left(\frac{1 + |\widetilde{\delta}_{x}|^{11}}{\kappa^{3}} \right),$$

where

$$\Phi(z) := \int_{-\infty}^{z} \phi(y) \mathrm{d}y, \quad z \in \mathbb{R},$$

denotes the cumulative distribution function of the standard normal distribution.

3. Proofs

Proof of Theorem 1. By taking the logarithm in (1), one has

$$\log\left\{\frac{\sqrt{2}\nu_{\kappa}f_{\mu,\kappa}(x)}{\phi(\delta_{x}/\sqrt{2})}\right\} = \kappa \cos(x-\mu) - \frac{1}{2}\log(2\pi) - \log\{I_{0}(\kappa)\} + \frac{1}{2}\log 2 + \frac{1}{2}\log\nu_{\kappa}^{2} + \frac{1}{4}\delta_{x}^{2}.$$
(8)

Since

$$x - \mu = \sqrt{\delta_x^2 \nu_{\kappa}^2} = \sqrt{\delta_x^2 \left\{ \frac{1}{2\kappa} + \frac{1}{8\kappa^2} + \frac{1}{8\kappa^3} + \mathcal{O}(\kappa^{-4}) \right\}},$$

(recall (3)) and

$$\cos(\sqrt{y}) = 1 - \frac{y}{2} + \frac{y^2}{24} - \frac{y^3}{720} + \mathcal{O}(y^4), \quad |y| < 1,$$

we know that

$$\kappa \cos(x-\mu) = \kappa \left[1 - \frac{\delta_x^2}{2} \left\{ \frac{1}{2\kappa} + \frac{1}{8\kappa^2} + \frac{1}{8\kappa^3} + \mathcal{O}(\kappa^{-4}) \right\} + \frac{\delta_x^4}{24} \left\{ \frac{1}{2\kappa} + \frac{1}{8\kappa^2} + \frac{1}{8\kappa^3} + \mathcal{O}(\kappa^{-4}) \right\}^2 - \frac{\delta_x^6}{720} \left\{ \frac{1}{2\kappa} + \frac{1}{8\kappa^2} + \frac{1}{8\kappa^3} + \mathcal{O}(\kappa^{-4}) \right\}^3 + \mathcal{O}_\eta \left(\frac{1+|\delta_x|^8}{\kappa^4} \right) \right] = \kappa - \frac{1}{4} \delta_x^2 + \frac{1}{\kappa} \left(\frac{\delta_x^4}{96} - \frac{\delta_x^2}{16} \right) + \frac{1}{\kappa^2} \left(-\frac{\delta_x^6}{5760} + \frac{\delta_x^4}{192} - \frac{\delta_x^2}{16} \right) + \mathcal{O}_\eta \left(\frac{1+|\delta_x|^8}{\kappa^3} \right).$$
(9)

By (2), (3) and the Taylor expansion,

$$\log(1+y) = y - \frac{y^2}{2} + \frac{y^3}{3} + \mathcal{O}(y^4), \quad |y| < 1,$$

one has

$$-\log\{I_0(\kappa)\} = -\kappa + \frac{1}{2}\log(2\pi\kappa) - \frac{1}{8\kappa} - \frac{1}{16\kappa^2} - \frac{25}{384\kappa^3} + \mathcal{O}(\kappa^{-4}),$$
$$\frac{1}{2}\log\nu_{\kappa}^2 = -\frac{1}{2}\log(2\kappa) + \frac{1}{8\kappa} + \frac{7}{64\kappa^2} - \frac{11}{384\kappa^3} + \mathcal{O}(\kappa^{-4}),$$

so that

$$-\frac{1}{2}\log(2\pi) - \log\{I_0(\kappa)\} + \frac{1}{2}\log 2 + \frac{1}{2}\log\nu_{\kappa}^2 = -\kappa + \frac{3}{64\kappa^2} - \frac{3}{32\kappa^3} + \mathcal{O}(\kappa^{-4}).$$
(10)

By putting (9) and (10) together in (8), we obtain

$$\log\left\{\frac{\sqrt{2\nu_{\kappa}}f_{\mu,\kappa}(x)}{\phi(\delta_{x}/\sqrt{2})}\right\} = \frac{1}{\kappa}\left(\frac{\delta_{x}^{4}}{96} - \frac{\delta_{x}^{2}}{16}\right) + \frac{1}{\kappa^{2}}\left(-\frac{\delta_{x}^{6}}{5760} + \frac{\delta_{x}^{4}}{192} - \frac{\delta_{x}^{2}}{16} + \frac{3}{64}\right) + \mathcal{O}_{\eta}\left(\frac{1+|\delta_{x}|^{8}}{\kappa^{3}}\right),$$

which proves (6).

To obtain (7) and conclude the proof, we take the exponential on both sides of the last equation with $\eta = \kappa^{-1/4} \tilde{\eta}$, and we expand the right-hand side with

$$e^{y} = 1 + y + \frac{y^{2}}{2} + \mathcal{O}(e^{U}y^{3}), \text{ for } -\infty < y \le U < \infty.$$

We get

$$\begin{aligned} \frac{\sqrt{2}\nu_{\kappa}f_{\mu,\kappa}(x)}{\phi(\delta_{x}/\sqrt{2})} &= 1 + \frac{1}{\kappa} \left(\frac{\delta_{x}^{4}}{96} - \frac{\delta_{x}^{2}}{16}\right) + \frac{1}{\kappa^{2}} \left(-\frac{\delta_{x}^{6}}{5760} + \frac{\delta_{x}^{4}}{192} - \frac{\delta_{x}^{2}}{16} + \frac{3}{64}\right) \\ &+ \frac{1}{2\kappa^{2}} \left(\frac{\delta_{x}^{4}}{96} - \frac{\delta_{x}^{2}}{16}\right)^{2} + \mathcal{O}_{\tilde{\eta}} \left(\frac{1 + |\delta_{x}|^{12}}{\kappa^{3}}\right) \\ &= 1 + \frac{1}{\kappa} \left(\frac{\delta_{x}^{4}}{96} - \frac{\delta_{x}^{2}}{16}\right) + \frac{1}{\kappa^{2}} \left(\frac{\delta_{x}^{8}}{18432} - \frac{19\delta_{x}^{6}}{23040} + \frac{11\delta_{x}^{4}}{1536} - \frac{\delta_{x}^{2}}{16} + \frac{3}{64}\right) + \mathcal{O}_{\tilde{\eta}} \left(\frac{1 + |\delta_{x}|^{12}}{\kappa^{3}}\right),\end{aligned}$$

which yields the claim (7) after some rearrangements. This concludes the proof. *Proof of Corollary 2.* By integrating from x to ∞ the result of Theorem 1, one gets

$$\begin{split} \int_{x}^{\infty} f_{\mu,\kappa}(y) \mathrm{d}y &= \Psi(\widetilde{\delta}_{x}) + \frac{1}{\kappa} \left(\frac{\Psi_{4}(\widetilde{\delta}_{x})}{24} - \frac{\Psi_{2}(\widetilde{\delta}_{x})}{8} \right) \\ &+ \frac{1}{\kappa^{2}} \left(\frac{\Psi_{8}(\widetilde{\delta}_{x})}{1152} - \frac{19\Psi_{6}(\widetilde{\delta}_{x})}{2880} + \frac{11\Psi_{4}(\widetilde{\delta}_{x})}{384} - \frac{\Psi_{2}(\widetilde{\delta}_{x})}{8} + \frac{3}{64} \right) + \mathcal{O}_{\widetilde{\eta}} \left(\frac{1 + |\widetilde{\delta}_{x}|^{11}}{\kappa^{3}} \right), \end{split}$$

where

$$\Psi(z) = \int_{z}^{\infty} \phi(y) dy, \quad \Psi_{j}(z) = \int_{z}^{\infty} y^{j} \phi(y) dy, \quad z \in \mathbb{R}, \ j \in \mathbb{N}.$$

Since

$$\begin{split} \Psi_2(z) &= z\phi(z) + \Psi(z), \\ \Psi_4(z) &= (3z + z^3)\phi(z) + 3\Psi(z), \\ \Psi_6(z) &= (15z + 5z^3 + z^5)\phi(z) + 15\Psi(z), \\ \Psi_8(z) &= (105z + 35z^3 + 7z^5 + z^7)\phi(z) + 105\Psi(z), \end{split}$$

- ()

one deduces

$$F_{\mu,\kappa}(x) = \Phi(\widetilde{\delta}_x) + \phi(\widetilde{\delta}_x) \left\{ \frac{1}{\kappa} \left(\frac{\widetilde{\delta}_x^3}{24} \right) + \frac{1}{\kappa^2} \left(\frac{\widetilde{\delta}_x^7}{1152} - \frac{\widetilde{\delta}_x^5}{1920} + \frac{5\widetilde{\delta}_x^3}{192} - \frac{3\widetilde{\delta}_x}{64} \right) \right\} + \mathcal{O}_{\widetilde{\eta}} \left(\frac{1 + |\widetilde{\delta}_x|^{11}}{\kappa^3} \right).$$

his concludes the proof.

This concludes the proof.

- ()

. . .

References

- Abramowitz, M., & Stegun, I. A. 1964. Handbook of Mathematical Functions With Formulas, Graphs, and Mathematical Tables. National Bureau of Standards Applied Mathematics Series, vol. 55. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. MR0167642.
- Hill, G. W. 1976. New approximations to the von Mises distribution. Biometrika, 63(3), 673-676. doi:10.2307/2335751.
- Hill, G. W., & Davis, A. W. 1968. Generalized asymptotic expansions of Cornish-Fisher type. Ann. Math. Statist., **39**, 1264–1273. MR226762.
- Jammalamadaka, S. R., & SenGupta, A. 2001. Topics in Circular Statistics. Series on Multivariate Analysis, vol. 5. World Scientific Publishing Co., Inc., River Edge, NJ. With 1 IBM-PC floppy disk (3.5 inch; HD).
- Kent, J. 1978. Limiting behaviour of the von Mises-Fisher distribution. Math. Proc. Cambridge Philos. Soc., 84(3), 531-536. MR501259.
- Mardia, K. V., & Jupp, P. E. 2000. Directional Statistics. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd., Chichester. MR1828667.
- Ouimet, F. 2022a. Refined normal approximations for the central and noncentral chi-square distributions and some applications. *Statistics*, **56**(4), 935–956. MR4466042.
- Ouimet, F. 2022b. Refined normal approximations for the Student distribution. J. Class. Anal., 20(1), 23–33. MR4449387.