Fractional non-homogeneous counting process

Nick Laskin^{*}

TopQuark Inc. Toronto, ON, M6P 2P2 Canada

Abstract

A new fractional non-homogeneous counting process has been introduced and developed using the Kilbas and Saigo three-parameter generalization of the Mittag-Leffler function. The probability distribution function of this process reproduces for certain set of the fractality parameters the famous Poisson and fractional Poisson probability distributions as well as the probability distribution function of a counting process displaying stretched exponential interarrival times distribution.

Applications of the developed fractional non-homogeneous counting process cover fractional compound process, the generalization of combinatorial polynomials and numbers, statistics of the fractional non-homogeneous counting process and a new representation of the Kilbas and Saigo function.

PACS numbers: 05.10.Gg; 05.45.Df; 42.50.-p.

Keywords: Fractional non-homogeneous counting process, fractional Poisson process, generating functions, combinatorial polynomials and numbers, Mittag-Leffler function, Kilbas and Saigo function

1 Introduction

Recently, a discrete probability distribution function

^{*}*E-mail address*: nlaskin@rocketmail.com

$$P_{\sigma}(n,x) = \frac{x^{\sigma n}}{n!} \exp(-x^{\sigma}), \quad x \ge 0, \quad 0 < \sigma \le 1, \quad n = 0, 1, 2, ..., \quad (1)$$

has been implemented to introduce and study stretched coherent states [1]. Substitution $x \to \lambda_{\sigma}^{1/\sigma} t$ leads to the probability distribution function of the following counting process

$$P_{\sigma}(n,t) = \frac{(\lambda_{\sigma}t^{\sigma})^n}{n!} \exp(-\lambda_{\sigma}t^{\sigma}), \qquad (2)$$

where λ_{σ} has units of $[\lambda_{\sigma}] = time^{-\sigma}$. This expression is recognized as a non-homogeneous Poisson process whose "rate" r(t) changes with time,

$$r(t) = \sigma \lambda_{\sigma} t^{\sigma - 1}.$$
 (3)

Then $P_{\sigma}(n,t)$ can be represented as follows,

$$P_{\sigma}(n,t) = \frac{(\Lambda_{\sigma}(t))^n}{n!} \exp(-\Lambda_{\sigma}(t)), \qquad (4)$$

where

$$\Lambda_{\sigma}(t) = \int_{0}^{t} d\tau r(\tau) = \sigma \lambda_{\sigma} \int_{0}^{t} d\tau \tau^{\sigma-1} = \lambda_{\sigma} t^{\sigma}.$$
 (5)

Given the probability distribution $P_{\sigma}(n, t)$, it is easy to obtain the probability distribution function of interarrival times, i.e., the time intervals between two consecutive arrivals. It turns out, that for the counting process described by the probability distribution function specified by Eq.(2), interarrival times probability distribution function $\psi_{\sigma}(\tau)$ has the form

$$\psi_{\sigma}(\tau) = \sigma \lambda_{\sigma} \tau^{\sigma-1} \exp(-\lambda_{\sigma} \tau^{\sigma}), \qquad \tau > 0, \tag{6}$$

which is the well-known Weibull distribution with the shape parameter σ and the scale parameter λ_{σ} . The probability distribution function $\psi_{\sigma}(\tau)$ is widely used to fit "fat-tail" economic, social and natural data and is called the *stretched exponential distribution* (see, for example, [2]).

The function $P_{\sigma}(n,t)$ satisfies the following system of the equations

$$\frac{\partial P_{\sigma}(n,t)}{\partial t} = \sigma \lambda_{\sigma} t^{\sigma-1} (P_{\sigma}(n-1,t) - P_{\sigma}(n,t)), \qquad n \ge 1, \tag{7}$$

$$\frac{\partial P_{\sigma}(0,t)}{\partial t} = -\sigma \lambda_{\sigma} t^{\sigma-1} P_{\sigma}(0,t), \qquad (8)$$

with some initial condition at t = 0,

$$P_{\sigma}(n,t=0) = p_0(n).$$
(9)

Usually, it is supposed that at t = 0 nothing had arrived, that is the initial condition is specified as $P_{\sigma}(n,t)|_{t=0} = p_0(n)$, where $p_0(n) = 0$, if $n \neq 0$ and $p_0(n) = 1$, if n = 0.

Thus, Eq.(7) is linear differential equation with respect to t and difference equation with respect to n, and is called a differential-difference equation. When $\sigma = 1$, we get the famous Poisson probability distribution function $P_1(n,t) \equiv P_{\sigma}(n,t)|_{\sigma=1}$, which satisfies the system of equations [3]

$$\frac{\partial P_1(n,t)}{\partial t} = \lambda_1 (P_1(n-1,t) - P_1(n,t)), \qquad n \ge 1,$$
(10)

$$\frac{\partial P_{\sigma}(0,t)}{\partial t} = -\lambda_1 P_{\sigma}(0,t), \qquad (11)$$

and has the form

$$P_1(n,t) = \frac{(\lambda_1 t)^n}{n!} \exp(-\lambda_1 t), \qquad n = 0, 1, 2, ...,.$$
(12)

where λ_1 has units of $[\lambda_1] = time^{-1}$. As can be seen from Eq.(6), when $\sigma = 1, \psi_{\sigma}(\tau)|_{\sigma=1} = \psi_1(\tau)$ becomes an exponential distribution,

$$\psi_1(\tau) = \lambda_1 \exp(-\lambda_1 \tau), \qquad \tau > 0. \tag{13}$$

Therefore, $P_{\sigma}(n, t)$ defined by Eq.(2) is generalization of the Poisson probability distribution function $P_1(n, t)$ defined by Eq.(12) and $\psi_{\sigma}(\tau)$ is generalization of the exponential distribution defined by Eq.(13).

Another important and widely used generalization of the Poisson process is fractional Poisson process developed to model and study a long-memory impact on the counting processes [4] empirically observed in complex quantum and classical systems. The quantum system example is the fluorescence intermittency of single CdSe quantum dots, that is, the fluorescence emission of single nanocrystals exhibits intermittent behavior, namely, a sequence of "light on" and "light off" states departing from Poisson statistics. In fact, the waiting time distribution in both states displays non-exponential behavior [5]. As examples of classical systems let's mention the distribution of waiting times between two consecutive transactions in financial markets [6] and another, which comes from network communication systems, where the duration of network sessions or connections exhibits non-exponential powerlaw behavior [7]. The probability distribution function of the fractional Poisson process satisfies the following system of fractional differential-difference equations

$${}_{0}^{C}D_{t}^{\mu}P_{\mu}(n,t) = \lambda_{\mu} \left(P_{\mu}(n-1,t) - P_{\mu}(n,t) \right), \qquad n \ge 1,$$
(14)

and

$${}_{0}^{C}D_{t}^{\mu}P_{\mu}(0,t) = -\lambda_{\mu}P_{\mu}(0,t), \qquad 0 < \mu \le 1,$$
(15)

subject to normalization condition

$$\sum_{n=0}^{\infty} P_{\mu}(n,t) = 1,$$
(16)

and the initial condition at t = 0, $P_{\mu}(n, t = 0) = \delta_{n,0}$, where μ is fractality parameter, the operator ${}_{0}^{C}D_{t}^{\mu}$ is the Caputo fractional derivative¹ whose action on a function f(t) is defined as [11]

$${}_{0}^{C}D_{t}^{\mu}\mathbf{f}(t) = \frac{1}{\Gamma(1-\mu)} \int_{0}^{t} \frac{d\tau(d\mathbf{f}(\tau)/d\tau)}{(t-\tau)^{\mu}}, \qquad 0 < \mu \le 1,$$
(17)

with $\Gamma(\mu)$ being the Gamma function, which has the familiar representation $\Gamma(\mu) = \int_{0}^{\infty} dt e^{-t} t^{\mu-1}$, $\operatorname{Re}\mu > 0$, and the parameter λ_{μ} has units $[\lambda_{\mu}] = time^{-\mu}$.

The solution of the system of the equations (14) and (15) has the form

$$P_{\mu}(n,t) = \frac{(\lambda_{\mu}t^{\mu})^n}{n!} \sum_{k=0}^{\infty} \frac{(k+n)!}{k!} \frac{(-\lambda_{\mu}t^{\mu})^k}{\Gamma(\mu(k+n)+1))}, \quad n = 0, 1, 2, ..., \quad 0 < \mu \le 1$$
(18)

¹The basic formulas on fractional calculus can be found in Refs. [8] - [10].

which is probability distribution function of the *fractional Poisson process* [4]. The $P_{\mu}(n,t)$ gives us the probability that in the time interval [0,t] we observe *n* arrivals.

Equations (14) and (15) are fractional generalization of Eqs.(7) and (8) for probability distribution function of the Poisson process. When $\mu = 1$, Eq.(18) turns into Eq.(12).

The goal of this work is to develop a system of fractional non-homogeneous differential-difference equations the exact solution of which is new fractional probability distribution of non-homogeneous fractional counting process. It turns out that this system reproduces for certain set of the fractality parameters the systems of equations (7) and (8), (10) and (11), (14) and (15). This goal was achieved by implementing the Kilbas and Saigo function [12], which is a three-parameter generalization of the Mittag-Leffler function [13].

The paper is organized as follows.

In Section 1 a non-homogeneous fractional differential-difference equation for the counting probability distribution functions has been developed and discussed. To solve this equation the probability generating function was calculated in Section 2. The fractional counting probability distribution function has been obtained in Section 3 using the probability generating function. Section 4 is devoted to the calculation of the moment generating function. The mean and variance of the new fractional counting process were found using the moment generating function. Interarrival time distribution was obtained in Section 5. Section 6 presents applications of new fractional counting probability distribution function covering fractional compound process and the generalization of combinatorial polynomials and numbers.

The results obtained are summarized in the Conclusion.

The Appendix A lists special cases of Kilbas and Saigo three-parameter function.

The Appendix B presents analytical expressions for the first four central moments of the introduced fractional non-homogeneous probability distribution.

2 Fundamentals

We define a new fractional non-homogeneous counting process by introducing the following system of fractional differential-difference equations for probability distribution function $P_{\mu,\beta}(n,t)$ of this process

$${}_{0}D_{t}^{\mu}P_{\mu,\beta}(n,t) = \lambda_{\mu+\beta}t^{\beta} \left(P_{\mu,\beta}(n-1,t) - P_{\mu,\beta}(n,t)\right), \quad n = 1, 2, 3, ..., \quad t \ge 0,$$
(19)

$${}_{0}D^{\mu}_{t}P_{\mu,\beta}(0,t) = -\lambda_{\mu+\beta}t^{\beta}P_{\mu,\beta}(0,t), \qquad (20)$$

subject to normalization condition

$$\sum_{n=0}^{\infty} P_{\mu,\beta}(n,t) = 1,$$
(21)

and the initial condition

$$P_{\mu,\beta}(n,t=0) = \delta_{n,0}.$$
 (22)

Here, μ and β are dimensionless fractality parameters subject to the conditions,

$$0 < \mu \le 1, \qquad -\mu < \beta \le 1 - \mu,$$
 (23)

the rate of arrivals $\lambda_{\mu+\beta}$ has units $[\lambda_{\mu+\beta}] = time^{-(\mu+\beta)}$, ${}_{0}D_{t}^{\mu}$ is the operator of the Caputo fractional time derivative of order μ defined by Eq.(17). We will see later where the conditions for μ and β come from.

The system of Eqs.(19) and (22) introduces the counting process with probability distribution function $P_{\mu,\beta}(n,t)$ of arriving *n* items (n = 0, 1, 2, ...)by time *t*, when the rate of arrivals depends on time *t*. In other words, the above defined counting process is generalization to non-homogeneous case (due to the dependency of the arrival rate on time) of the fractional Poisson process originally introduced and developed in [4].

When $\mu = 1$ and $\beta = 0$, the system of Eqs.(19) and (22) turns into the system of equations (10) and (11), which define the Poisson process with rate of arrival λ_1 .

When $\beta = 0$, the system of Eqs.(19) and (22) turns into the system of equations (14) and (15), which define the fractional Poisson process with rate of arrival λ_{μ} where $0 < \mu \leq 1$.

When $\mu = 1$, the system of Eqs.(19) and (22) turns into the the system of equations (7) and (8), which define the probability distribution function $P_{\sigma}(n, t)$, with $\sigma = 1 + \beta$, $0 < \sigma \leq 1$.

3 Probability generating function

To solve the system of equations (19) - (22) it is convenient to use the method of the generating function. Introducing the probability generating function $G_{\mu,\beta}(s,t)$

$$G_{\mu,\beta}(s,t) = \sum_{n=0}^{\infty} s^n P_{\mu,\beta}(n,t), \qquad (24)$$

allows to calculate $P_{\mu,\beta}(n,t)$ as

$$P_{\mu,\beta}(n,t) = \frac{1}{n!} \frac{\partial^n G_{\mu,\beta}(s,t)}{\partial s^n}|_{s=0}.$$
 (25)

Then, by multiplying Eqs.(19) and (20) by s^n , summing over n, we obtain the following fractional differential-difference equation for the probability generating function $G_{\mu}(s,t)$

$${}_{0}D^{\mu}_{t}G_{\mu}(s,t) = \lambda_{\mu+\beta}t^{\beta} \left(\sum_{n=1}^{\infty} s^{n}P_{\mu}(n-1,t) - \sum_{n=0}^{\infty} s^{n}P_{\mu}(n,t)\right) = (26)$$
$$\lambda_{\mu+\beta}t^{\beta}(s-1)G_{\mu}(s,t).$$

Thus, we come to the following equation

$${}_{0}D^{\mu}_{t}G_{\mu,\beta}(s,t) = \lambda_{\mu+\beta}t^{\beta}(s-1)G_{\mu,\beta}(s,t), \qquad (27)$$

subject to the initial condition

$$G_{\mu,\beta}(s,t=0) = 1,$$
 (28)

due to the definition Eq.(24) and the initial condition for $P_{\mu,\beta}(n,t)$ given by Eq.(22). The solution to Eq.(27) can be obtained using the ansatz [12]

$$G_{\mu,\beta}(s,t) = \sum_{n=0}^{\infty} \frac{c_n(\mu,\beta) \{\lambda_{\mu+\beta} t^{\mu+\beta}(s-1)\}^n}{\Gamma(n(\mu+\beta)+1)},$$
(29)

where $c_n(\mu, \beta)$ are still unknown coefficients and $\Gamma(\mu)$ is the Gamma function. Substituting this ansatz into Eq.(27) and taking into account that due to the definition of Caputo fractional derivative Eq.(17)

$${}_{0}D_{t}^{\mu}t^{n(\mu+\beta)} = \frac{\Gamma(n(\mu+\beta)+1)}{\Gamma(n(\mu+\beta)+1-\mu)}t^{n(\mu+\beta)},$$
(30)

we find that Eq.(27) is satisfied if the following recurrence relation holds,

$$c_{n+1}(\mu,\beta) = \frac{\Gamma(n(\mu+\beta)+\beta+1)}{\Gamma(n(\mu+\beta)+1)}c_n(\mu,\beta).$$
(31)

This recurrence relation allows us to represent Eq.(29) as

$$G_{\mu,\beta}(s,t) = c_0(\mu,\beta) + \tag{32}$$

$$\sum_{n=1}^{\infty} \{\lambda_{\mu+\beta} t^{\mu+\beta} (s-1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma(k(\mu+\beta)+\beta+1)}{\Gamma(k(\mu+\beta)+\mu+\beta+1)} c_0(\mu,\beta).$$

Taking into account the initial condition given by Eq.(28), we come to the conclusion that $c_0(\mu, \beta) = 1$. Therefore, the solution of the initial problem defined by Eqs.(27) and (28) is

$$G_{\mu,\beta}(s,t) = 1 + \sum_{n=1}^{\infty} \{\lambda_{\mu+\beta} t^{\mu+\beta}(s-1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma(k(\mu+\beta)+\beta+1)}{\Gamma(k(\mu+\beta)+\mu+\beta+1)}.$$
 (33)

It turns out that the probability generating function $G_{\mu,\beta}(s,t)$ can be expressed in terms of Kilbas and Saigo three-parameter generalization of the Mittag-Leffler function [12], defined as

$$E_{\alpha,m,l}(z) = \sum_{n=0}^{\infty} c_n z^n, \qquad z \in \mathbb{C},$$
(34)

where

$$c_n = \prod_{j=0}^{n-1} \frac{\Gamma(\alpha(jm+l)+1)}{\Gamma(\alpha(jm+l)+1)+1)}, \qquad n = 1, 2, ..., \qquad c_0 = 1, \qquad (35)$$

and

$$\alpha, m, l \in \mathbb{R}, \quad \alpha > 0, \quad m > 0 \quad \text{and} \quad \alpha(jm+l) \neq -1, -2, -3, \dots$$
 (36)

To show how the Kilbas and Saigo function appears in Eq.(33) we transform the arguments of the Gamma functions under the product sign in Eq.(33) as follows

$$\prod_{k=0}^{n-1} \frac{\Gamma(k(\mu+\beta)+\beta+1)}{\Gamma(k(\mu+\beta)+\mu+\beta+1)} = \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$
 (37)

Hence, it can be seen from Eqs.(33) and (37) that the power series holds for $G_{\mu,\beta}(s,t)$

$$G_{\mu,\beta}(s,t) = 1 + \sum_{n=1}^{\infty} \{\lambda_{\mu+\beta} t^{\mu+\beta}(s-1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$
 (38)

On the other hand, using the definitions given by Eqs.(34) and (35), we see that the right side of Eq.(38) is the following Kilbas and Saigo function²,

$$E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) = 1+$$

$$\sum_{n=1}^{\infty} z^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]},$$
(39)

with $z = \lambda_{\mu+\beta} t^{\mu+\beta} (s-1)$. Therefore, the generating function $G_{\mu,\beta}(s,t)$ has the form

$$G_{\mu,\beta}(s,t) = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(\lambda_{\mu+\beta}t^{\mu+\beta}(s-1)), \qquad (40)$$

where μ and β are subject to the conditions given by Eq.(23). Thus, we expressed in terms of Kilbas and Saigo three-parameter function the probability generating function $G_{\mu,\beta}(s,t)$ of the counting process introduced by the system of fractional differential-difference equations (19) and (20).

When $\mu = 1$ and $\beta = 0$, then according to Eq.(134) the generating function $G_{1,0}(s,t)$ takes the form

$$G_{1,0}(s,t) = \exp(\lambda_1 t(s-1)),$$

²The special cases of the function $E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)$ are presented in Appendix A.

which is the generating function of the Poisson probability distribution [3].

When $0 < \mu \leq 1$ and $\beta = 0$, then due to Eq.(136) we obtain from Eq.(38)

$$G_{\mu,0}(s,t) = 1 + \sum_{n=1}^{\infty} \frac{\{\lambda_{\mu} t^{\mu}(s-1)\}^n}{\Gamma(\mu n+1)} = E_{\mu}(\lambda_{\mu} t^{\mu}(s-1)), \qquad (41)$$

where $E_{\mu}(z)$ is the Mittag-Leffler function defined as [13]

$$E_{\mu}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\mu n+1)}.$$
(42)

Thus, Eq.(41) gives us the probability generating function of the fractional Poisson process [4].

When $\mu = 1$ and $-1 < \beta \le 0$, then due to Eq.(138) and having introduced the notations,

$$\sigma = 1 + \beta$$
 and $\overline{\lambda}_{\sigma} = \frac{\lambda_{\sigma}}{\sigma}$, (43)

we present Eq.(38) in the form

$$G_{\sigma}(s,t) = \exp(\overline{\lambda}_{\sigma}t^{\sigma}(s-1)), \qquad (44)$$

where $0 < \sigma \leq 1$, if $-1 < \beta \leq 0$. Thus, Eq.(44) is probability generating function of the probability distribution function given by Eq.(2).

When $\mu + \beta = 1$, then taking into account Eq.(139) we can express Eq.(38) in the form

$$G_{\mu,1-\mu}(s,t) = 1 + \sum_{n=1}^{\infty} \{\lambda t(s-1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma(k+2-\mu)}{\Gamma(k+2)},$$

or

$$G_{\mu,1-\mu}(s,t) = E_{\mu,\frac{1}{\mu},\frac{1-\mu}{\mu}}(\lambda t(s-1)),$$

where the following notation was introduced

$$\lambda = \lambda_{\mu+\beta} \mid_{\mu+\beta=1}. \tag{45}$$

3.1 Probability distribution function

Using the definition given by Eq.(25) we obtain the probability distribution function $P_{\mu,\beta}(n,t)$

$$P_{\mu,\beta}(n,t) = \frac{1}{n!} \frac{\partial^{n} G_{\mu,\beta}(s,t)}{\partial s^{n}}|_{s=0} = \frac{(\lambda_{\mu+\beta} t^{\mu+\beta})^{n}}{n!} E^{(n)}_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(-\lambda_{\mu+\beta} t^{\mu+\beta}), \quad (46)$$

where the following notation is introduced

$$E^{(n)}_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) = \frac{d^n}{dz^n} E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z).$$
(47)

The equation (46) can be rewritten as

$$P_{\mu,\beta}(n,t) = \frac{(-z)^n}{n!} E^{(n)}_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) |_{z=-\lambda_{\mu+\beta}t^{\mu+\beta}}, \qquad (48)$$

and

$$P_{\mu,\beta}(n=0,t) = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(-\lambda_{\mu+\beta}t^{\mu+\beta}).$$
(49)

The function $P_{\mu,\beta}(n,t)$ is normalized

$$\sum_{n=0}^{\infty} P_{\mu,\beta}(n,t) = \frac{(-z)^n}{n!} E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}^{(n)}(z)|_{z=-\lambda_{\mu+\beta}t^{\mu+\beta}} = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}^{(n)}(0) = 1, \quad (50)$$

due to the definition given by Eqs.(34) and (35), and positive due to the complete monotonicity of the three-parameter Kilbas and Saigo function $E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(-x)$ when μ and β are subject to the conditions³ defined by Eq.(23), see, [14], [15],

 3 The conditions

$$0 < \mu \le 1, \qquad -\mu < \beta, \tag{51}$$

follow directly from the condition of complete monotonicity of the three-parameter Kilbas and Saigo function [14]. We have added an additional condition

$$\beta \leq 1-\mu$$

to ensure consistency with the parameter σ defined by Eq.(43) and included in Eq.(1).

$$(-1)^{n} \frac{d^{n} E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(-x)}{dx^{n}} \ge 0, \qquad x \ge 0,$$
(52)

Therefore, $P_{\mu,\beta}(n,t)$ is indeed a probability distribution function. The power series representation for $P_{\mu,\beta}(n,t)$ is

$$P_{\mu,\beta}(n,t) = \frac{(\lambda_{\mu+\beta}t^{\mu+\beta})^n}{n!} \times$$
(53)

$$\sum_{m=0}^{\infty} \frac{(m+n)!}{m!} (-\lambda_{\mu+\beta} t^{\mu+\beta})^m \prod_{k=0}^{m+n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$

Replacing $\lambda_{\mu+\beta}t^{\mu+\beta}$ with $x^{\mu+\beta}$ in Eqs.(48) and (53), we obtain a new family of fractional probability distributions $P_{\mu,\beta}(x)$,

$$P_{\mu,\beta}(n,x) = \frac{(-z)^n}{n!} E^{(n)}_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)|_{z=-\lambda_{\mu+\beta}t^{\mu+\beta}},$$
(54)

or in a series form

$$P_{\mu,\beta}(n,x) = \frac{x^{(\mu+\beta)n}}{n!} \sum_{m=0}^{\infty} \frac{(m+n)!}{m!} \times$$

$$(55)$$

$$(-x^{(\mu+\beta)})^m \prod_{k=0}^{m+n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]},$$

where μ and β are subject to the conditions given by Eq.(23). When $\mu = 1$ and $\beta = 0$, using Eq.(134) we get from Eq.(46)

$$P_{1,0}(n,t) = \frac{(\lambda_1 t)^n}{n!} \exp(-\lambda_1 t),$$

which is the celebrated Poisson probability distribution. When $0 < \mu \leq 1$ and $\beta = 0$, using Eq.(136) we get from Eq.(46)

$$P_{\mu,0}(n,t) = \frac{(\lambda_{\mu}t^{\mu})^n}{n!} \sum_{m=0}^{\infty} \frac{(m+n)!}{m!} \frac{(-\lambda_{\mu}t^{\mu})^m}{\Gamma(\mu m+1)},$$

which is the fractional Poisson probability distribution [4].

When $\mu = 1$ and $-1 < \beta \le 0$, then according to Eq.(138) we obtain from Eq.(46)

$$P_{1,\beta}(n,t) \equiv P_{1+\beta}(n,t) = \frac{(\frac{\lambda_{1+\beta}}{1+\beta}t^{1+\beta})^n}{n!} \exp(-\frac{\lambda_{1+\beta}t^{1+\beta}}{1+\beta}).$$
 (56)

Using the notations given by Eq.(43), we represent Eq.(56) in the form

$$P_{\sigma}(n,t) = \frac{(\overline{\lambda}_{\sigma}t^{\sigma})^n}{n!} \exp(-\overline{\lambda}_{\sigma}t^{\sigma}), \quad 0 < \sigma \le 1,$$

which is recognized as Eq.(1).

With $\mu + \beta = 1$ and using Eq.(139) we can present Eq.(46) as

$$P_{\mu,1-\mu}(n,t) = \frac{(\lambda t)^n}{n!} E_{\mu,\frac{1}{\mu},\frac{1-\mu}{\mu}}^{(n)}(-\lambda t),$$

where λ is defined by Eq.(45). The series representation for $P_{\mu,1-\mu}(n,t)$ has the form

$$P_{\mu,1-\mu}(n,t) = \frac{(\lambda t)^n}{n!} \sum_{m=0}^{\infty} \frac{(m+n)!}{m!} (-\lambda t)^m \prod_{k=0}^{m+n-1} \frac{\Gamma(k+2-\mu)}{\Gamma(k+2)}.$$
 (57)

4 Moment generating function

An equation for the moment of any integer order m of the probability distribution function $P_{\mu,\beta}(n,t)$ introduced by Eq. (48) can be easily found using the moment generating function $M_{\mu,\beta}(s,t)$ defined as

$$M_{\mu,\beta}(s,t) = \sum_{n=0}^{\infty} e^{-sn} P_{\mu,\beta}(n,t).$$
 (58)

Then for the moment of m^{th} order we have

$$< n_{\mu,\beta}^m > = \sum_{n=0}^{\infty} n^m P_{\mu,\beta}(n,t) = (-1)^m \frac{\partial^m}{\partial s^m} M_{\mu,\beta}(s,t) |_{s=0} .$$
 (59)

Multiplying Eq. (48) by e^{-sn} and summing over the *n* we get

$$M_{\mu,\beta}(s,t) = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(\lambda_{\mu+\beta}t^{\mu+\beta}(e^{-s}-1)),$$
(60)

or in a series form

$$M_{\mu,\beta}(s,t) = 1 + \sum_{n=1}^{\infty} \{\lambda_{\mu+\beta} t^{\mu+\beta} (e^{-s} - 1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu}) + \frac{\beta}{\mu}) + 1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu}) + \frac{\beta}{\mu} + 1) + 1]}.$$
(61)

Replacing $\lambda_{\mu+\beta}t^{\mu+\beta}$ with $x^{\mu+\beta}$ in Eqs.(60) and (61) we get the moment generating function for the fractional generalized probability distributions $P_{\mu,\beta}(x)$,

$$M_{\mu,\beta}(s,x) = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(x^{\mu+\beta}(e^{-s}-1)),$$
(62)

and in series form

$$M_{\mu,\beta}(s,x) = 1 + \sum_{n=1}^{\infty} \{x^{\mu+\beta}(e^{-s}-1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$
 (63)

When $\mu = 1$ and $\beta = 0$, using Eq.(134) we get from Eq.(63)

$$M_{1,0}(s,x) = 1 + \sum_{n=1}^{\infty} \frac{(\lambda_1 t(e^{-s} - 1))^n}{n!} = \exp(\lambda_1 t(e^{-s} - 1)),$$

which is the moment generating function of the Poisson process. With $0 < \mu \leq 1$ and $\beta = 0$, using Eq.(136) we get from Eq.(63)

$$M_{\mu,0}(s,x) = E_{\mu}(\lambda_{\mu}t^{\mu}(e^{-s}-1)),$$

which is the moment generating function of the fractional Poisson process [4].

For $\mu = 1$ and $-1 < \beta \leq 0$, Eq.(63) gives us the moment generating function of the counting process, the probability distribution function of which is determined by Eq.(2)

$$M_{\sigma}(s,x) \equiv M_{1+\beta}(s,t) = \exp(\overline{\lambda}_{\sigma}t^{\sigma}(e^{-s}-1)), \qquad (64)$$

where σ and $\overline{\lambda}_{\sigma}$ are defined by Eq.(43). When $\mu + \beta = 1$, Eq.(62) goes to

$$M_{\mu,1-\mu}(s,t) = E_{\mu,\frac{1}{\mu},\frac{1-\mu}{\mu}}(\lambda t(e^{-s}-1),$$

and Eq.(63) goes to

$$M_{\mu,1-\mu}(s,t) = 1 + \sum_{n=1}^{\infty} \{\lambda t(e^{-s} - 1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma(k+2-\mu)}{\Gamma(k+2)},$$

where λ is defined by Eq.(45).

4.1 Mean and variance

The mean $\langle n_{\mu,\beta} \rangle$ can be calculated using the definition (59),

$$\langle n_{\mu,\beta} \rangle = -\frac{\partial}{\partial s} M_{\mu,\beta}(s,t) |_{s=0} ,$$
 (65)

with $M_{\mu,\beta}(s,t)$ given either by Eq.(60) or Eq.(61). Thus, we have

$$\langle n_{\mu,\beta} \rangle = \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)} \lambda_{\mu+\beta} t^{\mu+\beta}.$$
 (66)

It is easy to see that the moment of the second order $< n_{\mu,\beta}^2 >$ is

$$\langle n_{\mu,\beta}^2 \rangle = \frac{\partial^2}{\partial s^2} M_{\mu,\beta}(s,t) \mid_{s=0} =$$
 (67)

$$2\frac{\Gamma(\beta+1)\Gamma(\mu+2\beta+1)}{\Gamma(\mu+\beta+1)\Gamma(2\mu+2\beta+1)}(\lambda_{\mu+\beta}t^{\mu+\beta})^2 + \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}\lambda_{\mu+\beta}t^{\mu+\beta}.$$

Then variance $Var_{\mu,\beta}$ of the probability distribution $P_{\mu,\beta}(n,t)$ is

$$Var_{\mu,\beta} = < n_{\mu,\beta}^2 > - < n_{\mu,\beta} >^2 =$$
(68)

$$\left\{\frac{2\Gamma(\beta+1)\Gamma(\mu+2\beta+1)}{\Gamma(\mu+\beta+1)\Gamma(2\mu+2\beta+1)} - \left(\frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}\right)^2\right\} (\lambda_{\mu+\beta}t^{\mu+\beta})^2 + \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}\lambda_{\mu+\beta}t^{\mu+\beta},$$

here μ and β are subject to the conditions given by Eq.(23).

5 Interarrival times distribution

An important characteristic of any counting process is the time between successive counts/arrivals - the interarrival time. Interarrival time is a random variable having the probability distribution function $\psi_{\mu,\beta}(\tau)$,

$$\psi_{\mu,\beta}(\tau) = -\frac{d}{d\tau} \mathcal{P}_{\mu,\beta}(\tau), \tag{69}$$

where $\mathcal{P}_{\mu,\beta}(\tau)$ is the probability that a given interarrival time is greater or equal to τ [4]

$$\mathcal{P}_{\mu,\beta}(\tau) = 1 - \sum_{n=1}^{\infty} P_{\mu,\beta}(n,\tau) = P_{\mu,\beta}(0,\tau),$$
(70)

here $P_{\mu,\beta}(n,\tau)$ is defined by Eq.(53).

It follows from Eqs.(69) and (70) that the probability distribution function of the interarrival times has the form

$$\psi_{\mu,\beta}(\tau) = \frac{\mu+\beta}{\tau} \sum_{l=0}^{\infty} (-1)^l (l+1) (\lambda_{\mu+\beta} \tau^{\mu+\beta})^{l+1} \prod_{k=0}^l \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$
(71)

The Laplace transform $\varphi_{\mu,\beta}(u)$ of the interarrival probability distribution function is defined as

$$\varphi_{\mu,\beta}(u) = \int_{0}^{\infty} d\tau e^{-u\tau} \psi_{\mu,\beta}(\tau), \qquad (72)$$

and has the form

$$\varphi_{\mu,\beta}(u) = (\mu + \beta) \sum_{l=0}^{\infty} (-1)^{l} (l+1) (\lambda_{\mu+\beta})^{l+1} \frac{\Gamma[(\mu + \beta)(l+1)]}{u^{(\mu+\beta)(l+1)}} \times$$
(73)
$$\prod_{k=0}^{l} \frac{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu}) + 1]}{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu} + 1) + 1]}.$$

It easy to see that when $\mu = 1$ and $\beta = 0$, Eq.(71) transforms into the exponential distribution defined by Eq.(13). From Eq.(73) we obtain the Laplace transform of the exponential distribution

$$\varphi_{1,0}(u) = \frac{\lambda_1}{\lambda_1 + u}.$$

When $0 < \mu \leq 1$ and $\beta = 0$, Eq.(71) becomes the probability distribution of interarrival times of the fractional Poisson process

$$\psi_{\mu,0}(\tau) = \lambda_{\mu} \tau^{\mu-1} E_{\mu,\mu}(-\lambda_{\mu} \tau^{\mu}), \qquad (74)$$

where $E_{\mu,\mu}(z)$ is the two parameter Mittag-Leffler function defined by the series [16]

$$E_{\mu,\mu}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\mu n + \mu)}.$$
(75)

From and Eq.(73) it follows that the Laplace transform has the form [4]

$$\varphi_{\mu,0}(u) = \frac{\lambda_{\mu}}{\lambda_{\mu} + u^{\mu}}.$$

When $\mu = 1$ and $-1 < \beta \leq 0$, Eq.(71) goes to stretched exponential distribution defined by Eq.(6) if we take into account the notation introduced by Eq.(43). From Eq.(73) we obtain the Laplace transform of the stretched exponential distribution

$$\varphi_{\sigma}(u) = \sigma \overline{\lambda}_{\sigma} \sum_{l=0}^{\infty} \frac{(-\overline{\lambda}_{\sigma})^{l}}{l!} \frac{\Gamma(\sigma(l+1))}{u^{\sigma(l+1)}},$$
(76)

here σ and $\overline{\lambda}_{\sigma}$ are defined by Eq.(43). With $\mu + \beta = 1$ Eq.(71) goes to

$$\psi_{\mu,1-\mu}(\tau) = \frac{1}{\tau} \sum_{l=0}^{\infty} (-1)^l (l+1) (\lambda \tau)^{l+1} \prod_{k=0}^l \frac{\Gamma(k+2-\mu)}{\Gamma(k+2)},$$
(77)

and Eq.(73) goes to

$$\varphi_{\mu,1-\mu}(u) = \frac{\lambda}{u} \sum_{l=0}^{\infty} (-\frac{\lambda}{u})^l (l+1)! \prod_{k=0}^l \frac{\Gamma(k+2-\mu)}{\Gamma(k+2)},$$
(78)

where λ is defined by Eq.(45).

6 Applications

6.1 Fractional compound process

We call stochastic process $\{X_{\mu,\beta}(t), t \ge 0\}$ a fractional compound process if it is represented by

$$X_{\mu,\beta}(t) = \sum_{i=1}^{N_{\mu,\beta}(t)} Y_i,$$
(79)

where $\{N_{\mu,\beta}(t), t \geq 0\}$ is a fractional counting process whose probability distribution function is introduced by Eq.(48) or Eq.(53), and $\{Y_i, i = 1, 2, ...\}$ are independent and identically distributed random variables with probability distribution function $p(Y_i)$ for each Y_i . It is assumed that the process $\{N_{\mu,\beta}(t), t \geq 0\}$ and the sequence of random variables $\{Y_i, i = 1, 2, ...\}$ are independent.

The moment generating function $J_{\mu,\beta}(s,t)$ of fractional generalized compound process is defined as follows

$$J_{\mu,\beta}(s,t) = <\exp\{sX_{\mu,\beta}(t)\} >_{Y,N_{\mu,\beta}(t)},\tag{80}$$

where $\langle ... \rangle_{Y_i, N_{\mu,\beta}(t)}$ stands for average which includes two statistically independent averaging procedures:

1. Averaging $\langle \dots \rangle_Y$ over independent random variables Y_i , defined by

$$< ... >_{Y} = \int dY_{1}...dY_{n}p(Y_{1})...p(Y_{n})... = \prod_{i=1}^{n} \int dY_{i}p(Y_{i})....$$
 (81)

2. Averaging over random number of counts defined by

$$< ... >_{N_{\mu,\beta}(t)} = \sum_{n=0}^{\infty} P_{\mu,\beta}(n,t)...,$$
 (82)

where $P_{\mu,\beta}(n,t)$ is given by Eq.(48).

One can see from Eq.(80) that k^{th} order moment of fractional compound process $X_{\mu,\beta}(t)$ is obtained by differentiating $J_{\mu,\beta}(s,t)$ k times with respect to s and then setting s = 0, that is

$$\langle (X_{\mu,\beta}(t))^k \rangle_{Y,N_{\mu,\beta}(t)} = \frac{\partial^k}{\partial s^k} J_{\mu,\beta}(s,t)_{|s=0}.$$
(83)

To obtain equation for the moment generating function $J_{\mu,\beta}(s,t)$ we apply Eqs.(81) and (82) to Eq.(80)

$$J_{\mu,\beta}(s,t) = \sum_{n=0}^{\infty} < \exp\{sX_{\mu,\beta}(t)|N_{\mu,\beta}(t) = n\} >_{Y} P_{\mu,\beta}(n,t) =$$
(84)

$$\sum_{n=0}^{\infty} < \exp\{s(Y_1 + \ldots + Y_n)\} >_Y \times \frac{(-z)^n}{n!} E_{\mu, 1 + \frac{\beta}{\mu}, \frac{\beta}{\mu}}^{(n)}(z) \left|_{z = -\lambda_{\mu+\beta} t^{\mu+\beta}}\right|_{z = -\lambda_{\mu+\beta} t^{\mu+\beta}},$$

where we used the independence of $\{Y_1, Y_2, ...\}$ and $\{N_{\mu,\beta}(t), t \ge 0\}$ and the independence of the Y_i 's between themselves. Hence, setting

$$g(s) = \int dY_i p(Y_i) \exp\{sY_i\},\tag{85}$$

for the moment generating function of random variables Y_i , we obtain from Eq.(84) the moment generating function $J_{\mu,\beta}(s,t)$ of the fractional compound process

$$J_{\mu,\beta}(s,t) = \sum_{n=0}^{\infty} g^n(s) \times \frac{(-z)^n}{n!} \frac{d^n}{dz^n} \left(E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) \right) |_{z=-\lambda_{\mu+\beta}t^{\mu+\beta}} = (86)$$
$$E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(\lambda_{\mu+\beta}t^{\mu+\beta}(g(s)-1)).$$

By differentiating the above with respect to s and setting s = 0, it is easy to obtain the mean of the fractional compound process,

$$\langle X_{\mu,\beta}(t) \rangle_{Y_{i},N_{\mu,\beta}(t)} = \frac{\partial}{\partial s} J_{\mu,\beta}(s,t)|_{s=0} = \langle Y \rangle_{Y} \lambda_{\mu+\beta} t^{\mu+\beta} \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)},$$
(87)

which is a manifestation of independency between fractional counting process $\{N_{\mu,\beta}(t), t \geq 0\}$ and random variables Y_i .

6.2 New polynomials and associated numbers

Based on the probability distribution function defined by Eq.(54) we introduce polynomials $B_{\mu,\beta}(x,m)$,

$$B_{\mu,\beta}(x,m) = \sum_{n=0}^{\infty} n^m P_{\mu,\beta}(n,x) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} E^{(n)}_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(-x), \qquad (88)$$
$$B_{\mu,\beta}(x,0) = 1,$$

where the parameters μ , and β satisfy the conditions given by Eq.(23). From Eq.(88) at x = 1 we come to fractional numbers $B_{\mu,\beta}(m)$,

$$B_{\mu,\beta}(m) = B_{\mu,\beta}(x,m)|_{x=1} = \sum_{n=0}^{\infty} \frac{n^m}{n!} E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}^{(n)}(-1).$$
(89)

As an example, here are the first two newly introduced polynomials

$$B_{\mu,\beta}(x,1) = \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}x,$$
(90)

and

$$B_{\mu,\beta}(x,2) = \frac{\Gamma(\beta+1)\Gamma(\mu+2\beta+1)}{\Gamma(\mu+\beta+1)\Gamma(2\mu+2\beta+1)}x^2 + \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}x.$$
 (91)

The first two fractional numbers associated with the polynomials $B_{\mu,\beta}(x,m)$ are

$$B_{\mu,\beta}(1) = \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)},\tag{92}$$

and

$$B_{\mu,\beta}(2) = \frac{\Gamma(\beta+1)\Gamma(\mu+2\beta+1)}{\Gamma(\mu+\beta+1)\Gamma(2\mu+2\beta+1)} + \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}.$$
 (93)

Series representations of the polynomials $B_{\mu,\beta}(x,m)$ and numbers $B_{\mu,\beta}(m)$ are

$$B_{\mu,\beta}(x,m) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} \sum_{r=0}^{\infty} \frac{(r+n)!}{r!} (-x)^r \prod_{k=0}^{r+n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]},$$
(94)

$$B_{\mu,\beta}(m) = B_{\mu,\beta}(x,m)|_{x=1} = \sum_{n=0}^{\infty} \frac{n^m}{n!} \times$$
(95)

$$\sum_{r=0}^{\infty} \frac{(r+n)!}{r!} \prod_{k=0}^{r+n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]},$$

where we used Eq.(55).

6.2.1 Generating function of polynomials $B_{\mu,\beta}(x,m)$

The generating function of the polynomials $B_{\mu,\beta}(x,m)$ is defined as

$$\mathcal{B}_{\mu,\beta}(s,x) = \sum_{m=0}^{\infty} \frac{s^m}{m!} B_{\mu,\beta}(x,m).$$
(96)

To get the polynomial $B_{\mu,\beta}(x,m)$ we differentiate $\mathcal{B}_{\mu,\beta}(s,x)$ m times with respect to s, and then let s = 0. That is,

$$B_{\mu,\beta}(x,m) = \frac{\partial^m}{\partial s^m} \mathcal{B}_{\mu,\beta}(s,x)|_{s=0}.$$
(97)

To find an explicit equation for $\mathcal{B}_{\mu,\beta}(s,x)$, let's substitute Eq.(88) into Eq.(96)

$$\mathcal{B}_{\mu,\beta}(s,x) = \sum_{m=0}^{\infty} \frac{s^m}{m!} \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}^{(n)}(-x).$$

Summation over m gives us

$$\sum_{m=0}^{\infty} \frac{s^m}{m!} n^m = e^{sn},$$

and we obtain

$$\mathcal{B}_{\mu,\beta}(s,x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} e^{sn} E^{(n)}_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(-x) = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(x(e^s-1)), \qquad (98)$$

where the function $E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)$ is defined by Eq.(39). It is easy to see that the representation of the generating function $\mathcal{B}_{\mu,\beta}(s,x)$ in a power series has the form

$$\mathcal{B}_{\mu,\beta}(s,x) = 1 + \sum_{n=1}^{\infty} \{x(e^s - 1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu}) + 1]}{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu} + 1) + 1]},$$
(99)

where μ and β are subject to the conditions given by Eq.(23).

If we put x = 1 in Eq.(98), then we immediately come to the generating function $\mathcal{B}_{\mu,\beta}(s)$ of the numbers $B_{\mu,\beta}(m)$ introduced by Eq.(89)

$$\mathcal{B}_{\mu,\beta}(s) = \mathcal{B}_{\mu,\beta}(s,x)|_{x=1} = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(e^s - 1).$$
(100)

To get the numbers $B_{\mu,\beta}(m)$ we differentiate $\mathcal{B}_{\mu,\beta}(s)$ m times with respect to s, and then set s = 0. That is,

$$B_{\mu,\beta}(m) = \frac{\partial^m}{\partial s^m} \mathcal{B}_{\mu,\beta}(s)|_{s=0}.$$
 (101)

When $\mu = 1$ and $\beta = 0$, using Eq.(134) we get from Eq.(88)

$$B_{1,0}(x,m) = e^{-x} \sum_{n=0}^{\infty} n^m \frac{x^n}{n!},$$

which are the Bell polynomials [17]. The new formula Eq.(89) is a generalization of the so-called Dobiński relation known since 1877 [18] for the Bell numbers⁴

$$B_{\rm 1,0}(m) = e^{-1} \sum_{n=0}^{\infty} \frac{n^m}{n!}$$

When $0 < \mu \leq 1$ and $\beta = 0$, using Eq.(136) we get from Eq.(88)

⁴The number of ways a set of m elements can be partitioned into non-empty subsets.

$$B_{\mu,0}(x,m) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} E_{\mu,1,0}^{(n)}(-x) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} E_{\mu}^{(n)}(-x),$$

or in the form of a power series

$$B_{\mu,0}(x,m) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} \sum_{k=0}^{\infty} \frac{(n+k)!}{k!} \frac{(-x)^k}{\Gamma(\mu(n+k)+1)}, \quad B_{\mu,0}(x,0) = 1,$$

which are the fractional Bell polynomials [19]. Fractional generalization of the Dobiński relation is [19]

$$B_{\mu,0}(m) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} E_{\mu,1,0}^{(n)}(-1) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} E_{\mu}^{(n)}(-1).$$

With $\mu + \beta = 1$ and using Eq.(139) we can present Eq.(88) as

$$B_{\mu,1-\mu}(x,m) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} E_{\mu,\frac{1}{\mu},\frac{1-\mu}{\mu}}^{(n)}(-x),$$

where λ is defined by Eq.(45). The series representation for polynomials $B_{\mu,1-\mu}(x,m)$ has the form

$$B_{\mu,1-\mu}(x,m) = \sum_{n=0}^{\infty} n^m \frac{x^n}{n!} \sum_{m=0}^{\infty} \frac{(m+n)!}{m!} (-\lambda t)^m \prod_{k=0}^{m+n-1} \frac{\Gamma(k+2-\mu)}{\Gamma(k+2)}.$$
 (102)

6.3 Fractional combinatorial numbers associated with polynomials $B_{\mu,\beta}(x,m)$

Let us introduce fractional combinatorial numbers $S_{\mu,\beta}(m,l)$ associated with the polynomials $B_{\mu,\beta}(x,m)$ using the following equations

$$B_{\mu,\beta}(x,m) = \sum_{l=0}^{m} S_{\mu,\beta}(m,l) x^{l},$$
(103)

$$S_{\mu,\beta}(m,0) = \delta_{m,0}, \qquad S_{\mu,\beta}(m,l) = 0, \ l = m+1, \ m+2, \ \dots$$

where $B_{\mu,\beta}(x,m)$ is the polynomials given by Eq.(88) and the parameters μ , and β satisfy the conditions (23).

At x = 1, when the polynomials $B_{\mu,\beta}(x,m)$ become the fractional numbers, $B_{\mu,\beta}(m) = B_{\mu,\beta}(x,m)|_{x=1}$, Eq.(103) goes into the expression for the fractional numbers $B_{\mu,\beta}(m)$ in terms of fractional combinatorial numbers $S_{\mu,\beta}(m,l)$

$$B_{\mu,\beta}(m) = \sum_{l=0}^{m} S_{\mu,\beta}(m,l).$$
 (104)

To find a generating function of the fractional combinatorial numbers $S_{\mu,\beta}(m,l)$, let's expand the generating function $\mathcal{B}_{\mu,\beta}(s,x)$ given by Eq.(96). Upon substituting $B_{\mu}(x,m)$ from Eq.(103) we have the following chain of transformations

$$\mathcal{B}_{\mu,\beta}(s,x) = \sum_{m=0}^{\infty} \frac{s^m}{m!} \left(\sum_{l=0}^m S_{\mu,\beta}(m,l) x^l \right) =$$

$$\sum_{m=0}^{\infty} \frac{s^m}{m!} \left(\sum_{l=0}^{\infty} \theta(m-l) S_{\mu,\beta}(m,l) x^l \right) = \sum_{l=0}^{\infty} \left(\sum_{m=l}^{\infty} S_{\mu,\beta}(m,l) \frac{s^m}{m!} \right) x^l, \quad (105)$$

here $\theta(l)$ is the Heaviside step function, $\theta(l) = 1$, if $l \ge 0$ and $\theta(l) = 0$, if l < 0.

On the other hand, from Eq.(98), we have for $\mathcal{B}_{\mu,\beta}(s,x)$

$$\mathcal{B}_{\mu,\beta}(s,x) = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(x(e^s-1)) = 1+$$
(106)
$$\sum_{n=1}^{\infty} \{x(e^s-1)\}^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$

Upon comparing Eq.(105) and Eq.(106), we conclude that

$$\sum_{m=l}^{\infty} S_{\mu,\beta}(m,l) \frac{s^m}{m!} = (e^s - 1)^l \prod_{k=0}^{l-1} \frac{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu}) + 1]}{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu} + 1) + 1]}, \quad l = 1, 2, \dots$$
(107)

Let us introduce two generating functions $\mathcal{G}_{\mu,\beta}(s,l)$ and $\mathcal{F}_{\mu,\beta}(s,t)$ of the fractional combinatorial numbers $S_{\mu,\beta}(m,l)$,

$$\mathcal{G}_{\mu,\beta}(s,l) = \sum_{m=l}^{\infty} S_{\mu,\beta}(m,l) \frac{s^m}{m!} = (e^s - 1)^l \prod_{k=0}^{l-1} \frac{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu}) + 1]}{\Gamma[\mu(k(1 + \frac{\beta}{\mu}) + \frac{\beta}{\mu} + 1) + 1]},$$
(108)

and

$$\mathcal{F}_{\mu,\beta}(s,t) = \sum_{l=0}^{m} \sum_{m=l}^{\infty} S_{\mu,\beta}(m,l) \frac{s^m t^l}{m!} = E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(t(e^s-1)).$$
(109)

Then the fractional combinatorial numbers are obtained as follows,

$$S_{\mu,\beta}(m,l) = \frac{\partial^m}{\partial s^m} \mathcal{G}_{\mu,\beta}(s,l) |_{s=0} =$$
(110)

$$\prod_{k=0}^{l-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]} \frac{\partial^m}{\partial s^m} (e^s-1)^l |_{s=0} ,$$

and

$$S_{\mu,\beta}(m,l) = \frac{1}{l!} \frac{\partial^{m+l}}{\partial s^m \partial t^l} \mathcal{F}_{\mu,\beta}(s,t) |_{s=0,t=0} =$$

$$\frac{1}{l!} \frac{\partial^{m+l}}{\partial s^m \partial t^l} E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(t(e^s-1))|_{s=0,t=0},$$
(111)

where $l \leq m$ and $S_{\mu,\beta}(m,0) = \delta_{m,0}$. When $\mu = 1$ and $\beta = 0$ it follows from Eq.(110) that

$$S_{1,0}(m,l) = \frac{1}{l!} \frac{\partial^m}{\partial s^m} (e^s - 1)^l |_{s=0} .$$
 (112)

The right side of this equation is generating function of the standard Stirling numbers of the second kind [20]. Hence, $S_{1,0}(m,l)$ is the standard Stirling number of the second kind⁵ $S(m,l) \equiv S_{1,0}(m,l)$, and we conclude that

$$S_{\mu,\beta}(m,l) = l! \prod_{k=0}^{l-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]} S(m,l).$$
(113)

⁵The number of ways to partition a set of m elements into l non-empty sets.

Thus, we have represented fractional combinatorial numbers $S_{\mu,\beta}(m,l)$ through the standard Stirling number of the second kind S(m,l). It is easy to see, for example, that

$$S_{\mu,\beta}(m,1) = \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)},$$

$$S_{\mu,\beta}(m,2) = 2(2^{m-1}-1)\prod_{k=0}^{1}\frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]},$$

$$S_{\mu,\beta}(m,m-1) = \frac{m!(m-1)}{2}\prod_{k=0}^{m-2}\frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]},$$

$$S_{\mu,\beta}(m,m) = m!\prod_{k=0}^{m-1}\frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$

In a particular case, for $\mu = 1$ and $\beta = 0$, equations (108) and (109) turn into the well-know generating functions for standard Stirling numbers of the second kind S(m, l),

$$\mathcal{G}_{1,0}(s,l) = \sum_{m=l}^{\infty} S(m,l) \frac{s^m}{m!} = \frac{(e^s - 1)^l}{l!}, \qquad l = 0, 1, 2, ...,$$
(114)

and

$$\mathcal{F}_{1,0}(s,t) = \sum_{m=0}^{\infty} \sum_{l=0}^{m} S(m,l) \frac{s^m t^l}{m!} = \exp(t(e^s - 1)).$$
(115)

(see, Eqs.(2.17) and (2.18) in [21]).

6.3.1 Statistics of fractional probability distribution

We now use the fractional combinatorial numbers introduced by Eq.(103) to obtain the moments of the fractional non-homogeneous counting process with the probability distribution function $P_{\mu,\beta}(n,t)$ given by Eqs.(48) and (49). By the definition of the m^{th} order moment, we have

$$< n_{\mu,\beta}^{m} >= \sum_{n=0}^{\infty} n^{m} P_{\mu,\beta}(n,t) = \sum_{n=0}^{\infty} n^{m} \frac{(-z)^{n}}{n!} \frac{d^{n}}{dz^{n}} \left(E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) \right) |_{z=-\lambda_{\mu+\beta}t^{\mu+\beta}}.$$
(116)

It is easy to see that $\langle n_{\mu,\beta}^m \rangle$ is the fractional polynomial $B_{\mu,\beta}(\lambda_{\mu+\beta}t^{\mu+\beta},m)$ defined by Eq.(88),

$$\langle n_{\mu,\beta}^m \rangle = B_{\mu,\beta}(\lambda_{\mu+\beta}t^{\mu+\beta},m).$$

Then using the definition Eq.(103) we obtain

$$< n_{\mu,\beta}^m > = \sum_{l=0}^m S_{\mu,\beta}(m,l) (\lambda_{\mu+\beta} t^{\mu+\beta})^l,$$
 (117)

where $S_{\mu,\beta}(m,l)$ is fractional combinatorial numbers defined by Eq.(113). Therefore, the fractional combinatorial numbers $S_{\mu,\beta}(m,l)$ naturally appear in the power series over $\lambda_{\mu+\beta}t^{\mu+\beta}$ for the m^{th} order moment of the fractional non-homogeneous counting process introduced by Eqs.(19) - (23).

Using the analytical expression given by Eq.(113), let's list first four moments of the fractional non-homogeneous probability distribution $P_{\mu,\beta}(n,t)$. The first and second moments are given by Eqs.(66) and (67) respectively. Third order moment $\langle n^3_{\mu,\beta} \rangle$ is

$$< n_{\mu,\beta}^{3} >= \sum_{n=0}^{\infty} n^{3} P_{\mu,\beta}(n,t) =$$

$$6 \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)} \frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)} \frac{\Gamma(2\mu+3\beta+1)}{\Gamma(3\mu+3\beta+1)} (\lambda_{\mu+\beta}t^{\mu+\beta})^{3} +$$

$$6 \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)} \frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)} (\lambda_{\mu+\beta}t^{\mu+\beta})^{2} + \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)} \lambda_{\mu+\beta}t^{\mu+\beta},$$
(118)

and fourth order moment $\langle n_{\mu,\beta}^4 \rangle$ is

$$< n_{\mu,\beta}^4 > = \sum_{n=0}^{\infty} n^4 P_{\mu,\beta}(n,t) =$$

 $24\frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}\frac{\Gamma(2\mu+3\beta+1)}{\Gamma(3\mu+3\beta+1)}\frac{\Gamma(3\mu+4\beta+1)}{\Gamma(4\mu+4\beta+1)}(\lambda_{\mu+\beta}t^{\mu+\beta})^4+$

$$36\frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}\frac{\Gamma(2\mu+3\beta+1)}{\Gamma(3\mu+3\beta+1)}(\lambda_{\mu+\beta}t^{\mu+\beta})^{3}+$$
(119)

$$14\frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}(\lambda_{\mu+\beta}t^{\mu+\beta})^2 + \frac{\Gamma(\beta+1)}{\Gamma(\mu+\beta+1)}\lambda_{\mu+\beta}t^{\mu+\beta}.$$

When $0 < \mu \leq 1$ and $\beta = 0$, Eqs.(66), (67), (118) and (119) turn into equations for the first four moments of the fractional Poisson process [19].

In the case when $\mu = 1$ and $\beta = 0$, Eqs.(66), (67), (118) and (119) become the well-know equations for moments of the standard Poisson probability distribution with the parameter $\lambda_1 = \lambda_{\mu+\beta}|_{\mu=1,\beta=0}$ (see, Eqs.(22) - (24) in Ref. [22]).

When $\mu = 1$ and $-1 < \beta \leq 0$ Eqs.(66), (67), (118) and (119) become the equations for the moments of the process, defined by Eq.(1). Taking into account Eq.(43) we come to the known equations for the moments of the standard Poisson probability distribution with the parameter $\overline{\lambda}_{\sigma}$.

6.4 Variance, skewness and kurtosis excess of the fractional non-homogeneous counting process

To find analytical expressions for the variance, skewness and kurtosis excess of a fractional non-homogeneous counting process determined by the probability distribution function $P_{\mu,\beta}(n,t)$ given by Eq.(55), let's introduce the central moment of the m^{th} order $M_{\mu,\beta}(m)$,

$$M_{\mu,\beta}(m) = <(n - < n_{\mu,\beta} >)^m > = \sum_{n=0}^{\infty} (n - < n_{\mu,\beta} >)^m P_{\mu,\beta}(n,t) =$$

$$\sum_{n=0}^{\infty} \sum_{r=0}^{m} (-1)^{m-r} {m \choose r} n^r (< n_{\mu,\beta} >)^{m-r} P_{\mu,\beta}(n,t) =$$

$$\sum_{r=0}^{m} (-1)^{m-r} {m \choose r} (< n_{\mu,\beta} >)^{m-r} \sum_{l=0}^{r} S_{\mu,\beta}(r,l) (\lambda_{\mu+\beta} t^{\mu+\beta})^l,$$
(120)

where $\langle n_{\mu,\beta} \rangle$ is the first order moment defined by Eq.(66) and $S_{\mu,\beta}(r,l)$ is given by Eq.(113). The first four central moments $M_{\mu,\beta}(m)$, m = 1, 2, 3, 4,

are presented in the Appendix B. In terms of the central moments $M_{\mu,\beta}(m)$, m = 1, 2, 3, 4, the variance $Var_{\mu,\beta}$, skewness $s_{\mu,\beta}$, and kurtosis excess $k_{\mu,\beta}$ of the fractional non-homogeneous counting process are

$$Var_{\mu,\beta} = \mathcal{M}_{\mu,\beta}(2), \tag{121}$$

$$s_{\mu,\beta} = \frac{M_{\mu,\beta}(3)}{M_{\mu,\beta}^{3/2}(2)},$$
(122)

$$k_{\mu,\beta} = \frac{\mathcal{M}_{\mu,\beta}(4)}{\mathcal{M}_{\mu,\beta}^2(2)} - 3.$$
(123)

When $0 < \mu < 1$ and $\beta = 0$, Eqs.(121) - (123) turn into equations for the variance, skewness and kurtosis excess of the fractional Poisson process [19]. In the case where $\mu = 1$ and $\beta = 0$, Eqs.(121) - (123) become equations for the variance, skewness and kurtosis excess of the standard Poisson probability distribution, see, for example, Eqs.(25) - (27) in Ref. [22]. When $\mu = 1$ and $-1 < \beta \leq 0$ the variance, skewness and kurtosis excess of the process defined by Eq.(1) are exactly the same as for the Poisson process if we take into account that for the process defined by Eq.(1) we have

$$< n_{1,\beta} > = < n_{1+\beta} > |_{\mu=1} = \frac{\lambda_{1+\beta} t^{1+\beta}}{1+\beta},$$

which can be expressed in *Poisson-like form* using Eq.(43)

$$< n_{1+\beta} > \equiv < n_{\sigma} > = \overline{\lambda}_{\sigma} t^{\sigma}.$$

6.4.1 New representation of the Kilbas and Saigo function

To obtain a new representation for the Kilbas and Saigo function $E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)$ defined by Eq.(39) we use the generating function of the Stirling numbers of the first kind⁶

$$\frac{x!}{(x-m)!} = x(x-1)...(x-m+1) = \sum_{l=0}^{m} s(m,l)x^{l}.$$
(124)

 $^{^6\}mathrm{The}$ Stirling numbers of the first kind are defined as the coefficients s(m,l) in the expansion

$$(1+z)^n = \sum_{m=0}^{\infty} \frac{z^m}{m!} \sum_{l=0}^m s(m,l)n^l,$$
(125)

where s(m, l) stands for the Stirling numbers of the first kind, see Sec. 24.1.3 in [20]. Then, we evaluate expectations of both sides of Eq.(125) using the probability distribution function of fractional non-homogeneous counting process $P_{\mu,\beta}(n,\rho)$ given by Eq.(54)

$$\sum_{n=0}^{\infty} P_{\mu,\beta}(n,1)(1+z)^n = \sum_{n=0}^{\infty} P_{\mu,\beta}(n,1) \sum_{m=0}^{\infty} \frac{z^m}{m!} \sum_{l=0}^m s(m,l)n^l.$$
(126)

It is easy to see that the calculations on the left side of Eq.(126) lead to $E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)$, and we

$$E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) = \sum_{n=0}^{\infty} P_{\mu}(n,1) \sum_{m=0}^{\infty} \frac{z^m}{m!} \sum_{l=0}^{m} s(m,l)n^l.$$

Then, using the definition of the fractional numbers $B_{\mu,\beta}(m)$ given by Eq.(89) we obtain new representations of the Kilbas and Saigo function

$$E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) = \sum_{m=0}^{\infty} \frac{z^m}{m!} \sum_{l=0}^m s(m,l) B_{\mu,\beta}(l)$$
(127)

through Stirling numbers of the first kind and the fractional numbers $B_{\mu,\beta}(l)$. Using Eq.(104) the Kilbas and Saigo function can be expressed through Stirling numbers of the first kind and fractional combinatorial numbers $S_{\mu,\beta}(m,l)$

$$E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z) = \sum_{m=0}^{\infty} \frac{z^m}{m!} \sum_{l=0}^m \sum_{r=0}^l s(m,l) S_{\mu,\beta}(l,r).$$
(128)

The two equations (127) and (128) are new representations of the Kilbas and Saigo function.

From Eq.(127) we see that the Kilbas and Saigo function can be considered as generating function of the sum $\sum_{l=0}^{m} s(m, l) B_{\mu,\beta}(l)$, that is

$$\sum_{l=0}^{m} s(m,l) B_{\mu,\beta}(l) = \frac{\partial^{m}}{\partial z^{m}} E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)|_{z=0}.$$
 (129)

Using Eq.(39), we obtain the identity

$$\sum_{l=0}^{m} s(m,l) B_{\mu,\beta}(l) = m! \prod_{k=0}^{m-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$
 (130)

From Eq.(128) we see that the Kilbas and Saigo function can be considered as generating function of the sum $\sum_{l=0}^{m} \sum_{r=0}^{l} s(m,l)S_{\mu,\beta}(l,r)$, that is

$$\sum_{l=0}^{m}\sum_{r=0}^{l}s(m,l)S_{\mu,\beta}(l,r) = \frac{\partial^{m}}{\partial z^{m}}E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)|_{z=0}.$$

Using Eq.(39), we obtain the identity

$$\sum_{l=0}^{m} \sum_{r=0}^{l} s(m,l) S_{\mu,\beta}(l,r) = m! \prod_{k=0}^{m-1} \frac{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu})+1]}{\Gamma[\mu(k(1+\frac{\beta}{\mu})+\frac{\beta}{\mu}+1)+1]}.$$
 (131)

7 Conclusions

A new fractional non-homogeneous counting process has been introduced and developed using the Kilbas and Saigo three-parameter generalization of the Mittag-Leffler function. The probability distribution function of this process reproduces for certain set of the fractality parameters the famous Poisson and fractional Poisson probability distributions as well as the probability distribution function of a counting process displaying stretched exponential interarrival times distribution.

Applications of the developed fractional non-homogeneous counting process cover fractional compound process, the generalization of combinatorial polynomials and numbers and their generating functions, variance, skewness and kurtosis excess of the fractional non-homogeneous counting process expressed through four central moments and a new representation of the Kilbas and Saigo function.

The presented results will stimulate further development of the fundamentals and various applications in a new field of probability theory and stochastic processes, which can be called fractional stochastic processes and their applications.

8 Appendix A

8.1 Special cases of the Kilbas and Saigo function $E_{\mu,1+\frac{\beta}{\mu},\frac{\beta}{\mu}}(z)$

Defined by Eq.(39) the Kilbas and Saigo function has the following special cases.

• When $\mu = 1$ and $\beta = 0$, Eq.(39) goes to

$$E_{1,1,0}(z) = 1 + \sum_{n=1}^{\infty} z^n \prod_{k=0}^{n-1} \frac{\Gamma[k+1]}{\Gamma[(k+1)+1]},$$

Then due to

$$\prod_{k=0}^{n-1} \frac{\Gamma[k+1]}{\Gamma[(k+1)+1]} = \prod_{k=0}^{n-1} \frac{1}{k+1} = \frac{1}{n!},$$
(132)

where the following property of the Gamma function was used

$$\Gamma(z+1) = z\Gamma(z), \tag{133}$$

we obtain

$$E_{1,1,0}(z) = 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} = \exp(z).$$
(134)

• When $0 < \mu \leq 1$ and $\beta = 0$, Eq.(39) goes to

$$E_{\mu,1,0}(z) = 1 + \sum_{n=1}^{\infty} z^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu k+1]}{\Gamma[\mu(k+1)+1]}.$$

Then due to

$$\prod_{k=0}^{n-1} \frac{\Gamma[\mu k+1]}{\Gamma[\mu(k+1)+1]} = \frac{1}{\Gamma(\mu n+1)},$$
(135)

we obtain

$$E_{\mu,1,0}(z) = 1 + \sum_{n=1}^{\infty} \frac{z^n}{\Gamma(\mu n + 1)} = E_{\mu}(z), \qquad (136)$$

here $E_{\mu}(z)$ is the Mittag-Leffler function defined by Eq.(42).

• When $\mu = 1$ and $-1 < \beta \le 0$, Eq.(39) goes to

$$E_{1,1+\beta,\beta}(z) = 1 + \sum_{n=1}^{\infty} z^n \prod_{k=0}^{n-1} \frac{\Gamma[\mu(k(1+\beta)+\beta)+1]}{\Gamma[\mu(k(1+\beta)+\beta+1)+1]}.$$

Then due to

$$\prod_{k=0}^{n-1} \frac{\Gamma[(k(1+\beta)+\beta)+1]}{\Gamma[(k(1+\beta)+\beta+1)+1]} = \prod_{k=0}^{n-1} \frac{1}{(k+1)(\beta+1)} = \frac{1}{(1+\beta)^n} \frac{1}{n!},$$
 (137)

we obtain

$$E_{1,1+\beta,\beta}(z) = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \frac{z^n}{(1+\beta)^n} = \exp(\frac{z}{1+\beta}).$$
 (138)

• When $\mu + \beta = 1$, Eq.(39) goes to

$$E_{\mu,\frac{1}{\mu},\frac{1-\mu}{\mu}}(z) = 1 + \sum_{n=1}^{\infty} z^n \prod_{k=0}^{n-1} \frac{\Gamma(k+2-\mu)}{\Gamma(k+2)}.$$
 (139)

9 Appendix B

9.1 Four first central moments

Using Eq.(120), the first four central moments can be calculated. We expressed the results of the calculation in the form of a power series over the first-order moment $\langle n_{\mu,\beta} \rangle$:

$$M_{\mu,\beta}(1) = 0,$$
 (140)

$$M_{\mu,\beta}(2) = \left(\frac{2\Gamma(\mu+\beta+1)\Gamma(\mu+2\beta+1)}{\Gamma(\beta+1)\Gamma(2\mu+2\beta+1)} - 1\right) < n_{\mu,\beta} >^2 + < n_{\mu,\beta} >,$$
(141)

$$M_{\mu,\beta}(3) = 2\left(\frac{3\Gamma(\mu+\beta+1)}{\Gamma(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}\frac{\Gamma(\mu+\beta+1)}{\Gamma(\beta+1)}\frac{\Gamma(2\mu+3\beta+1)}{\Gamma(3\mu+3\beta+1)}-\right)$$

$$\frac{3\Gamma(\mu+\beta+1)}{\Gamma(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)} + 1) < n_{\mu,\beta} >^3 +$$
(142)

$$3\left(\frac{2\Gamma(\mu+\beta+1)}{\Gamma(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}-1\right) < n_{\mu,\beta} >^{2} + < n_{\mu,\beta} >,$$

and

$$\begin{split} \mathbf{M}_{\mu,\beta}(4) &= 3(8\frac{\Gamma^{3}(\mu+\beta+1)}{\Gamma^{3}(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}\frac{\Gamma(2\mu+3\beta+1)}{\Gamma(3\mu+3\beta+1)}\frac{\Gamma(3\mu+4\beta+1)}{\Gamma(4\mu+4\beta+1)} - \\ &\qquad 8\frac{\Gamma^{2}(\mu+\beta+1)}{\Gamma^{2}(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}\frac{\Gamma(2\mu+3\beta+1)}{\Gamma(3\mu+3\beta+1)} + \\ &\qquad 4\frac{\Gamma(\mu+\beta+1)}{\Gamma(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)} - 1) < n_{\mu,\beta} >^{4} + \\ &\qquad (143) \\ &\qquad 6(\frac{6\Gamma^{2}(\mu+\beta+1)}{\Gamma^{2}(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}\frac{\Gamma(2\mu+3\beta+1)}{\Gamma(3\mu+3\beta+1)} - \\ &\qquad 4\frac{\Gamma(\mu+\beta+1)}{\Gamma(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)} + 1) < n_{\mu,\beta} >^{3} + \end{split}$$

$$2\left(7\frac{\Gamma(\mu+\beta+1)}{\Gamma(\beta+1)}\frac{\Gamma(\mu+2\beta+1)}{\Gamma(2\mu+2\beta+1)}-2\right) < n_{\mu,\beta} >^{2} + < n_{\mu,\beta} > .$$

References

- [1] N. Laskin, Stretched Coherent States, https://arxiv.org/abs/2302.07563
- [2] J. Laherrere and D. Sornette, Stretched exponential distributions in nature and economy: "Fat tails" with characteristic scales, https://arxiv.org/pdf/cond-mat/9801293.pdf
- [3] W. Feller, An introduction to probability theory and its applications, Vol. 1, John Wiley & Sons, Inc., New York, 1968.

- [4] N. Laskin, Fractional Poisson process, Communications in Nonlinear Science and Numerical Simulation, 8 (2003) 201-213.
- [5] M. Kuno, D.P. Fromm, H. F. Hamann, A. Gallagher, and D.J. Nesbitt, Nonexponential blinking kinetics of single CdSe quantum dots: A universal power law behavior, J. Chem. Phys. **112** (2000) 3117-3120.
- [6] L. Sabatelli et al, Waiting time distributions in financial markets, The European Physical Journal B - Condensed Matter and Complex Systems, 27 (2002) 273-275.
- [7] W. Willinger, and V. Paxson, Where Mathematics Meets the Internet, Notices of the American Mathematical Society, 45 (1998) 961-970.
- [8] K.B. Oldham and J. Spanier, The Fractional Calculus, Academic Press, New York, 1974.
- [9] S.G. Samko, A.A. Kilbas, and O.I. Marichev, Fractional Integrals and Derivatives and Their Applications, Gordon and Breach Science Publishers, Langhorne, PA, 1993.
- [10] K.S. Miller and B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations, J. Wiley & Sons, New York, 1993.
- [11] M. Caputo, Linear model of dissipation whose Q is almost frequency independent-II, Geophys., J. R. Astron. Soc., 13 (1967) 529–539.
- [12] A. Kilbasi and M. Saigo, On Mittag-Leffler type function, fractional calculus operators and solutions of integral equations, Integral Transforms and Special Functions, 4, (1996) 355-370.
- [13] G. Mittag-Leffler, Sur la repriésentation analytique d'une branche uniforme d'une fonction monogène, Acta Mathematics, 29 (1905) 101-182.
- [14] L. Boudabsa, T. Simon, and P. Valloist, Fractional extreme distributions, Electron. J. Probab., 25 (2020) 1-20 (2020), DOI: 10.1214/20-EJP520
- [15] L. Boudabsa, Τ. Simon, Some Properties of Kilbasthe Saigo Function, Mathematics, 9 (2021)217,24pages, https://doi.org/10.3390/math9030217

- [16] A. Erdélyi, Ed., Higher Transcendental Functions, Vol.3, pp. 206-227 (Chapter 18 Miscellaneous Functions) McGraw-Hill, New York, 1955.
- [17] E. T. Bell, Exponential Polynomials, The Annals of Mathematics, Second Series, 35 (1934), 258-277.
- [18] G. Dobiński, Summirung der Reihe $\sum n^m/n!$, für m = 1, 2, 3, 4, 5, ...,Grunert Archiv (Arch. für M. und Physik) **61** (1877) 333-336.
- [19] N. Laskin, Some applications of the fractional Poisson probability distribution, J Math Phys, 50 (2009) 113513, 12 pages.
- [20] M. Abramowitz and I.A. Stegun, (Eds.). Stirling Numbers of the Second Kind. §24.1.4 in Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing, pp. 824-825, New York, Dover, 1972.
- [21] C. A. Charalambides and J. Singh: A review of the Stirling numbers, their generalizations and statistical applications, Communications in Statistics - Theory and Methods, 17 (1988) 2533-2595.
- [22] E.W. Weisstein, Poisson Distribution, From *MathWorld* - A Wolfram Web Resource, http://mathworld.wolfram.com/PoissonDistribution.html