ARTIN L-FUNCTIONS AND NONCOMMUTATIVE TORI

IGOR V. NIKOLAEV¹

ABSTRACT. Using the ideas of Deninger, we prove that the Artin *L*-functions coincide with such of the noncommutative tori. This result can be viewed as the Langlands reciprocity for noncommutative tori.

1. INTRODUCTION

The aim of our note is a conjecture [6, Conjecture 1] that every Artin *L*-function [Artin 1924] [1] can be written as the Euler product of the local zeta functions associated to the 2*n*-dimensional noncommutative tori [Rieffel 1990] [10]. This conjecture is known to be true for n = 0 and n = 1 [6, Theorem 1]. Here we prove the general case $n \ge 1$ (Theorem 1.5). Our proof is based on the ideas of [Deninger 2001] [2].

An *m*-dimensional noncommutative torus is the universal C^* -algebra \mathscr{A}_{Θ}^m generated by the unitary operators u_1, \ldots, u_m satisfying the commutation relations $u_j u_i = e^{2\pi i \theta_{ij}} u_i u_j$, $1 \leq i, j \leq m$ for a skew-symmetric matrix $T = (\theta_{ij}) \in M_m(\mathbf{R})$ [Rieffel 1990] [10]. Recall that if m = 2n then every generic T can be brought by the orthogonal linear transformations to the normal form diag $\begin{pmatrix} 0 & \theta_1 \\ -\theta_1 & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & \theta_n \\ -\theta_n & 0 \end{pmatrix}$, where $\theta_i > 0$ are linearly independent over \mathbf{Q} . The noncommutative torus is said to have real multiplication, if all θ_i are algebraic numbers [8, Section 1.1.5.2]; we write \mathscr{A}_{BM}^{2n} in this case.

Given the \mathscr{A}_{RM}^{2n} , one can think of θ_i as components of a normalized eigenvector $(1, \theta_1, \ldots, \theta_n)$ corresponding to the Perron-Frobenius eigenvalue $\lambda_B > 1$ of a positive matrix $B \in GL_{n+1}(\mathbf{Z})$. For each prime p we consider the characteristic polynomial det $(xI - B^{\pi(p)}) = x^{n+1} - a_1 x^n - \cdots - a_n x - 1 \in \mathbf{Z}[x]$, where $\pi(p)$ is an integer-valued function of p [8, Section 6.6.1].

Definition 1.1. The local zeta of the noncommutative torus \mathscr{A}_{RM}^{2n} is a function:

$$\zeta_p(\mathscr{A}_{RM}^{2n}, z) := \frac{1}{1 - a_1 z + a_2 z^2 - \dots - a_n z^n + p z^{n+1}}, \quad z \in \mathbf{C}.$$
 (1.1)

Remark 1.2. A motivation for (1.1) is as follows. Let k be a totally real field and denote by k_{CM} (k_{RM} , resp.) its complex (real, resp.) multiplication field, i.e. a totally imaginary (totally real, resp.) quadratic extension of k. Then the minimal polynomial of k_{CM} is an alternating sum of monomials of the minimal polynomial of the field k_{RM} . Indeed, since both k_{RM} and k_{CM} are quadratic extensions, their minimal polynomials consist of the monomials of even degrees, i.e. $p(x) = x^{2n} + a_2x^{2n-2} + \cdots + a_{2n-2}x^2 + a_{2n}$. Moreover, if x is a real root of p(x) then ix is a complex

²⁰¹⁰ Mathematics Subject Classification. Primary 11M55; Secondary 46L85.

Key words and phrases. Artin L-functions, noncommutative tori.

root of the polynomial $q(x) = x^{2n} - a_2 x^{2n-2} + \cdots + (-1)^{n+1} a_{2n-2} x^2 + (-1)^n a_{2n}$, i.e. an alternating sum of the monomials of p(x). Thus (1.1) is an alternating sum of monomials of the minimal polynomial of an algebraic integer of the field k_{RM} of norm p, where $x = z^{-1}$. We refer the reader to [8, Remark 6.6.3] for the explicit matrix formulas.

Definition 1.3. An *L*-function of the noncommutative torus \mathscr{A}_{RM}^{2n} is the product of the local zeta functions taken over all but a finite number of primes *p*:

$$L(\mathscr{A}_{RM}^{2n}, s) := \prod_{p} \zeta_{p}(\mathscr{A}_{RM}^{2n}, p^{-s}), \quad s \in \mathbf{C}.$$
 (1.2)

Recall that the fundamental problem in algebraic number theory is to describe how an ordinary prime p factors into prime ideals \mathfrak{P} in the ring of integers of an arbitrary finite extensions \mathbf{K} of the rational field \mathbf{Q} . Let $O_{\mathbf{K}}$ be the ring of integers of the extension **K** and $pO_{\mathbf{K}}$ a principal ideal; it is known that $pO_{\mathbf{K}} = \prod \mathfrak{P}_i$, where \mathfrak{P}_i are prime ideals of $O_{\mathbf{K}}$. If **K** is the Galois extension of **Q** and Gal (**K**|**Q**) is the corresponding Galois group, then each automorphism $q \in Gal$ (**K**|**Q**) "moves around" the ideals \mathfrak{P}_i in the prime decomposition of p over K. An isotropy subgroup of $Gal(\mathbf{K}|\mathbf{Q})$ (for given p) consists of the elements of $Gal(\mathbf{K}|\mathbf{Q})$ which fix all the ideals \mathfrak{P}_i . For simplicity, we shall assume that p is unramified in **K**, i.e. all \mathfrak{P}_i are distinct; in this case the isotropy subgroup is cyclic. The (conjugacy class of) generator in the cyclic isotropy subgroup of Gal ($\mathbf{K}|\mathbf{Q}$) corresponding to p is called the Frobenius element and denoted by Fr_p . The element $Fr_p \in Gal(\mathbf{K}|\mathbf{Q})$ describes completely the factorization of p over \mathbf{K} and the major problem is to express Fr_p in terms of arithmetic of the ground field **Q**. To handle this problem, it was suggested in [Artin 1924] [1] to consider an *m*-dimensional irreducible representation $\sigma_m : Gal(\mathbf{K}|\mathbf{Q}) \longrightarrow GL_m(\mathbb{C})$. The idea is to use the characteristic polynomial Char $(\sigma_m(Fr_p)) := \det(I_m - \sigma_m(Fr_p)z)$ of the matrix $\sigma_m(Fr_p)$; the polynomial is independent of the similarity class of $\sigma_m(Fr_p)$ in the group $GL_m(\mathbb{C})$ and thus provides an intrinsic description of the Frobenius element Fr_p .

Definition 1.4. An Artin *L*-function is the product taken over all but a finite set of primes *p*:

$$L(\sigma_m, s) := \prod_p \zeta_p(\sigma_m, p^{-s}), \text{ where } \zeta_p(\sigma_m, z) = \mathbf{Char}(\sigma_m(Fr_p)), s \in \mathbf{C}.$$

Our main result can be formulated as follows.

Theorem 1.5. ([8, Conjecture 6.6.1]) For each finite Galois extension **K** of the field **Q** and each irreducible representation σ_{n+1} : Gal (**K**|**Q**) \rightarrow GL_{n+1}(\mathbb{C}), there exists a noncommutative torus \mathscr{A}_{RM}^{2n} , such that:

$$L(\sigma_{n+1}, s) \equiv L(\mathscr{A}_{RM}^{2n}, s). \tag{1.3}$$

Remark 1.6. Theorem 1.5 can be viewed as an analog of the Langlands reciprocity for the Artin *L*-functions. Namely, such a reciprocity predicts a standard *L*-function at the RHS of (1.3) coming from an automorphic cuspidal representation of the adelic algebraic group GL(n) [Gelbart 1984] [4, Conjecture 1]. It follows from Theorem 1.5 that the even dimensional noncommutative tori encode critical arithmetic data which can be used in non-abelian class field theory and higher reciprocity laws [6, Conjecture 1].

The paper is organized as follows. A brief review of the preliminary facts is given in Section 2. Theorem 1.5 proved in Section 3. Some remarks can be found in Section 4.

2. Preliminaries

We briefly review Deninger's foliated spaces, noncommutative tori and their relation to the Drinfeld modules. We refer the reader to [Deninger 2001] [2], [Rieffel 1990] [10], and [9] for a detailed exposition.

2.1. Deninger's foliated spaces.

2.1.1. Foliated spaces. Let X be a smooth n-dimensional manifold. Consider a ddimensional foliation \mathcal{F} on X, i.e. the partition of X into a disjoint union of the d-dimensional submanifolds (called leaves) immersed in X. The tangent bundle $T\mathcal{F}$ of a foliation \mathcal{F} is the union of the tangent spaces to the leaves. The $T\mathcal{F}$ is a sub-bundle of the tangent bundle TX of the manifold X. The differential forms of order n along the leaves of \mathcal{F} are the smooth sections $\mathcal{A}^n_{\mathcal{F}}(X) = \Gamma(X, \Lambda^n T^*\mathcal{F})$, where $\Lambda^n T^*\mathcal{F}$ is a real vector bundle of dimension n. The exterior derivative along the leaves of \mathcal{F} is written as $d^n_{\mathcal{F}} : \mathcal{A}^n_{\mathcal{F}}(X) \to \mathcal{A}^{n+1}_{\mathcal{F}}(X)$ and satisfies the relation $d^{n+1}_{\mathcal{F}} \circ d^n_{\mathcal{F}} = 0$.

Definition 2.1. The reduced leafwise cohomology of foliation \mathcal{F} is given by the formula:

$$\overline{H}_{\mathcal{F}}^{n} = Ker \ d_{\mathcal{F}}^{n} / \overline{Im \ d_{\mathcal{F}}^{n-1}}, \tag{2.1}$$

where $\overline{Im \ d_{\mathcal{F}}^{n-1}}$ is the topological closure of $Im \ d_{\mathcal{F}}^{n-1}$ in the Fréchet topology on the set $\mathcal{A}^n_{\mathcal{F}}(X)$.

Remark 2.2. The cohomology $\overline{H}_{\mathcal{F}}^n$ can (and will be) infinite dimensional even if the leaves of \mathcal{F} are dense in X.

Consider a smooth flow $\phi : \mathbf{R} \times X \to X$ on the foliated manifold X given by the formula $(t, x) \mapsto \phi^t(x)$ for all $t \in \mathbf{R}$ and $x \in X$. The ϕ^t is assumed leaf-preserving and transversal to the leaves of foliation \mathcal{F} . The action ϕ^t_* of ϕ^t on $\overline{H}^n_{\mathcal{F}}$ is linear for all n and the infinitesimal generator of ϕ^t_* is denoted by Θ .

Let \mathcal{F} be a foliation of codimension 1 on the manifold X. The flow ϕ defines a riemannian metric on X which turns $\overline{H}_{\mathcal{F}}^n$ into a Hilbert space $\hat{H}_{\mathcal{F}}^n$. The linear operator ϕ_*^t on $\overline{H}_{\mathcal{F}}^n$ extends to a bounded operator on $\hat{H}_{\mathcal{F}}^n$.

Theorem 2.3. [2, Thm 3.3] For every test function $\varphi \in C_0^{\infty}(\mathbf{R})$ the $A_{\varphi} := \int_{\mathbf{R}} \varphi(t) \phi_*^t dt$ is a trace class operator on the Hilbert space $\hat{H}_{\mathcal{F}}^n$. Setting $Tr(\phi_* \mid \overline{H}_{\mathcal{F}}^n(X))(\varphi) = Tr A_{\varphi}$ defines a distribution on \mathbf{R} , so that:

$$\sum_{n=0}^{\dim \mathcal{F}} (-1)^n Tr \ (\phi_* \mid \overline{H}^n_{\mathcal{F}}(X))(\varphi) = \sum_{\gamma} l(\gamma) \sum_{k \in \mathbf{Z} - 0} \epsilon_{\gamma}(k) \delta_{kl(\gamma)} + \chi_{C_0}(\mathcal{F}, \mu) \delta_0,$$
(2.2)

where $l(\gamma)$ is the length of closed geodesic γ , $\epsilon_{\gamma}(k) = \operatorname{sgn} \det(1 - T_x \phi^{kl(\gamma)} | T_x \mathcal{F})$, δ_x is the Dirac delta function and $\chi_{C_0}(\mathcal{F}, \mu)$ is the Connes' Euler characteristic of \mathcal{F} with respect to the transverse measure μ .

2.1.2. Dedekind zeta function. Let \mathbf{K} be a number field. Consider the Dedekind zeta function:

$$\zeta_{\mathbf{K}}(s) = \sum_{I \subseteq O_{\mathbf{K}}} \frac{1}{(N_{\mathbf{K}|\mathbf{Q}}(I))^s},\tag{2.3}$$

where I runs through through the non-zero ideals of the ring of integers $O_{\mathbf{K}}$ and $N_{\mathbf{K}|\mathbf{Q}}(I)$ is the norm of I. We shall be using the following explicit formula (due to A. Weil) for $\zeta_{\mathbf{K}}(s)$, i.e. a relation between sums over the non-trivial zeros ρ of $\zeta_{\mathbf{K}}(s)$ and sums over prime ideals of the field \mathbf{K} .

Theorem 2.4. For every test function $\varphi \in C_0^{\infty}(\mathbf{R})$ the following equality holds:

$$1 - \sum_{\rho} e^{t\rho} + e^t = \sum_{\mathfrak{p}<\infty} \log N\mathfrak{p} \sum_{k=1}^{\infty} \delta_{k \log N\mathfrak{p}} + \sum_{\mathfrak{p}=\infty} \frac{1}{1 - e^{\kappa_{\mathfrak{p}}t}},$$
(2.4)

where $\kappa_{\mathfrak{p}} \in \{-2, -1\}$ and δ_x is the Dirac delta function.

2.1.3. Analogy [2, Section 4]. Similar to the case of varieties over finite fields, Deninger suggested that the leafwise reduced cohomology (2.1) is underlying an analog of the Lefschetz trace formula for the varieties over number fields [Deninger 2001] [2]. Unlike the case of finite fields, such a cohomology is no longer finite dimensional, see Remark 2.2. Comparing formulas (2.2) and (2.4), the following beautiful analogy can be observed.

Conjecture 2.5. Finite places \mathfrak{p} in the explicit formula (2.4) are closed geodesics $\gamma_{\mathfrak{p}}$ in formula (2.2), such that:

$$l(\gamma_{\mathfrak{p}}) = \log N\mathfrak{p}. \tag{2.5}$$

Remark 2.6. ([2]) To understand number theory in geometric terms, it is important to construct a foliated space (X, ϕ^t, \mathcal{F}) which realizes formula (2.5).

2.2. Noncommutative tori.

2.2.1. C^* -algebras. The C^* -algebra is an algebra \mathscr{A} over \mathbb{C} with a norm $a \mapsto ||a||$ and an involution $\{a \mapsto a^* \mid a \in \mathscr{A}\}$ such that \mathscr{A} is complete with respect to the norm, and such that $||ab|| \leq ||a|| ||b||$ and $||a^*a|| = ||a||^2$ for every $a, b \in \mathscr{A}$. Each commutative C^* -algebra is isomorphic to the algebra $C_0(X)$ of continuous complex-valued functions on some locally compact Hausdorff space X. Any other algebra \mathscr{A} can be thought of as a noncommutative topological space.

2.2.2. K-theory of C^* -algebras. By $M_{\infty}(\mathscr{A})$ one understands the algebraic direct limit of the C^* -algebras $M_n(\mathscr{A})$ under the embeddings $a \mapsto \operatorname{diag}(a, 0)$. The direct limit $M_{\infty}(\mathscr{A})$ can be thought of as the C^* -algebra of infinite-dimensional matrices whose entries are all zero except for a finite number of the non-zero entries taken from the C^* -algebra \mathscr{A} . Two projections $p, q \in M_{\infty}(\mathscr{A})$ are equivalent, if there exists an element $v \in M_{\infty}(\mathscr{A})$, such that $p = v^*v$ and $q = vv^*$. The equivalence class of projection p is denoted by [p]. We write $V(\mathscr{A})$ to denote all equivalence classes of projections in the C^* -algebra $M_{\infty}(\mathscr{A})$, i.e. $V(\mathscr{A}) := \{[p] : p = p^* =$ $p^2 \in M_{\infty}(\mathscr{A})\}$. The set $V(\mathscr{A})$ has the natural structure of an abelian semi-group with the addition operation defined by the formula $[p] + [q] := \operatorname{diag}(p,q) = [p' \oplus q']$, where $p' \sim p$, $q' \sim q$ and $p' \perp q'$. The identity of the semi-group $V(\mathscr{A})$ is given by [0], where 0 is the zero projection. By the K_0 -group $K_0(\mathscr{A})$ of the unital C^* -algebra \mathscr{A} one understands the Grothendieck group of the abelian semi-group $V(\mathscr{A})$, i.e. a completion of $V(\mathscr{A})$ by the formal elements [p] - [q]. The image of $V(\mathscr{A})$ in $K_0(\mathscr{A})$ is a positive cone $K_0^+(\mathscr{A})$ defining the order structure \leq on the abelian group $K_0(\mathscr{A})$. The pair $(K_0(\mathscr{A}), K_0^+(\mathscr{A}))$ is known as a dimension group of the C^* -algebra \mathscr{A} .

2.2.3. Noncommutative tori. A noncommutative *m*-torus is the universal C^* -algebra generated by *k* unitary operators u_1, \ldots, u_m ; the operators do not commute with each other, but their commutators $u_i u_j u_i^{-1} u_j^{-1}$ are fixed scalar multiples $\{\exp(2\pi i\theta_{ij}), | \theta_{ij} \in \mathbf{R}\}$ of the identity operator. The *m*-dimensional noncommutative torus, \mathscr{A}_T^m , is defined by a skew symmetric real matrix $T = (\theta_{ij}), 1 \leq i, j \leq k$. It is known that $K_0(\mathscr{A}_T^m) \cong K_1(\mathscr{A}_T^m) \cong \mathbf{Z}^{2^{m-1}}$. The canonical trace τ on the C^* -algebra \mathscr{A}_T^m defines a homomorphism from $K_0(\mathscr{A}_T^m)$ to the real line \mathbf{R} ; under the homomorphism, the image of $K_0(\mathscr{A}_T^m)$ is a \mathbf{Z} -module, whose generators $\tau = (\tau_i)$ are polynomials in θ_{ij} . The noncommutative tori \mathscr{A}_T^m and $\mathscr{A}_{T'}^m$ are Morita equivalent, if the matrices T and T' belong to the same orbit of a subgroup $SO(m, m \mid \mathbf{Z})$ of the group $GL_{2m}(\mathbf{Z})$, which acts on T by the formula T' = (AT + B) / (CT + D), where $(A, B, C, D) \in GL_{2m}(\mathbf{Z})$ and the matrices $A, B, C, D \in GL_m(\mathbf{Z})$ satisfy the conditions:

$$A^{t}D + C^{t}B = I, \quad A^{t}C + C^{t}A = 0 = B^{t}D + D^{t}B,$$
(2.6)

where I is the unit matrix and t at the upper right of a matrix means a transpose of the matrix.) The group $SO(m, m | \mathbf{Z})$ can be equivalently defined as a subgroup of the group $SO(m, m | \mathbf{R})$ consisting of linear transformations of the space \mathbf{R}^{2m} , which preserve the quadratic form $x_1x_{m+1} + x_2x_{k+2} + \cdots + x_kx_{2m}$.

2.3. Drinfeld modules. Let $A = \mathbf{F}_q[t]$ be the ring of polynomials in one variable over a finite field \mathbf{F}_q and $k = \mathbf{F}_q(t)$ its field of rational functions [Rosen 2002] [11, Chapter 1]. Recall that a polynomial $f \in k[x]$ is called additive in the ring k[x, y] if f(x + y) = f(x) + f(y). When char k = p the polynomial $\tau_p(x) = x^p$ is additive and each additive polynomial has the form $a_0x + a_1x^p + \cdots + a_rx^{p^r}$. The set of all additive polynomials is closed under addition and composition operations thus generating a ring of the non-commutative polynomials $k\langle \tau_p \rangle$ defined by the commutation relation $\tau_p a = a^p \tau_p$ for all $a \in k$. The Drinfeld module $Drin^r_A(k)$ of rank $r \geq 1$ is a homomorphism ρ :

$$A \xrightarrow{r} k \langle \tau_p \rangle, \tag{2.7}$$

given by a polynomial $\rho_a = a + c_1 \tau_p + c_2 \tau_p^2 + \cdots + c_r \tau_p^r$ with $c_i \in k$ and $c_r \neq 0$, such that for all $a \in A$ the constant term of ρ_a is a and $\rho_a \notin k$ for at least one $a \in A$ [Rosen 2002] [11, p. 200]. For each non-zero $a \in A$ the function field $k (\Lambda_{\rho}[a])$ is a Galois extension of k, such that its Galois group is isomorphic to a subgroup G of the matrix group $GL_r(A/aA)$, where $\Lambda_{\rho}[a] = \{\lambda \in \bar{k} \mid \rho_a(\lambda) = 0\}$ is a torsion submodule of the non-trivial Drinfeld module $Drin_A^r(k)$ [Rosen 2002] [11, Proposition 12.5]. Clearly, the abelian extensions correspond to the case r = 1.

Let G be a left cancellative semigroup generated by τ_p and all $a_i \in k$ subject to the commutation relations $\tau_p a_i = a_i^p \tau_p$. Let $C^*(G)$ be the semigroup C^* -algebra [Li 2017] [5]. For a Drinfeld module $Drin_A^r(k)$ defined by (2.7) we consider a homomorphism of the semigroup C^* -algebras:

$$C^*(A) \xrightarrow{r} C^*(k\langle \tau_p \rangle).$$
 (2.8)

It follows from (2.8) that the Drinfeld modules $Drin_A^r(k)$ are classified by the noncommutative tori \mathscr{A}_{RM}^{2r} ; we refer the reader to [9] for the details. A correspondence F between the Drinfeld modules and noncommutative tori is written as $\mathscr{A}_{RM}^{2r} = F(Drin_A^r(k))$. If r = 1 then the Drinfeld module (2.7) plays the rôle of an elliptic curve with complex multiplication \mathcal{E}_{CM} [Drinfeld 1974] [3, p. 594]. The set $\Lambda_{\rho}[a]$ consists of coefficients of the curve \mathcal{E}_{CM} which are equal to the value of the Weierstrass \wp -function at the torsion points of lattice $\Lambda_{CM} \subset \mathbf{C}$, where $\mathcal{E}_{CM} \cong \mathbf{C}/\Lambda_{CM}$. Likewise, if $r \geq 1$ then the algebra \mathscr{A}_{RM}^{2r} plays the rôle of the curve \mathcal{E}_{CM} given by coefficients $e^{2\pi i \theta_k}$ of the equations for \mathscr{A}_{RM}^{2r} . We therefore get $F(\Lambda_{\rho}[a]) = \{\log (\varepsilon) \ e^{2\pi i \theta_k} \ | \ 1 \leq k \leq 2r - 1\};$ see [9] for the notation and details.

Theorem 2.7. ([9]) The following is true:

(i) the map $F : Drin_A^r(k) \mapsto \mathscr{A}_{RM}^{2r}$ is a functor from the category of Drinfeld modules \mathfrak{D} to a category of the noncommutative tori \mathfrak{A} , which maps any pair of isogenous (isomorphic, resp.) modules $Drin_A^r(k)$, $\widetilde{Drin}_A^r(k) \in \mathfrak{D}$ to a pair of the homomorphic (isomorphic, resp.) tori $\mathscr{A}_{RM}^{2r}, \widetilde{\mathscr{A}_{RM}^{2r}} \in \mathfrak{A}$;

(ii) $F(\Lambda_{\rho}[a]) = \{e^{2\pi i \theta_k + \log \log \varepsilon} \mid 1 \le k \le 2r - 1\}, \text{ where } \mathscr{A}_{RM}^{2r} = F(Drin_A^r(k)) \text{ and } \Lambda_{\rho}[a] \text{ is the torsion submodule of the Drinfeld module } Drin_A^r(k);$

(iii) the Galois group $Gal\left(\mathbf{k}(e^{2\pi i\theta_k + \log\log\varepsilon}) \mid \mathbf{k}\right) \subseteq GL_r(A/aA)$, where \mathbf{k} is a subfield of the number field $\mathbf{Q}(e^{2\pi i\theta_k + \log\log\varepsilon})$.

Remark 2.8. Theorem 2.7 says that the Galois extensions **K** of the field **Q** arise from the noncommutative tori \mathscr{A}_{RM}^{2r} via the formula $\mathbf{K} = \mathbf{Q}(e^{2\pi i\theta_k + \log \log \varepsilon})$.

3. Proof of theorem 1.5

For the sake of clarity, let us outline the main ideas. We construct a foliated space (X, ϕ^t, \mathcal{F}) satisfying equation (2.5), see Lemma 3.2. This proves Conjecture 2.5 of [Deninger 2001] [2, Section 4]. The construction is as follows. Recall that the infinitesimal generator Θ is a bounded linear operator on the Hilbert space $\hat{H}^n_{\mathcal{F}}$, see Section 2.1.1. We consider a C^* -algebra \mathscr{Z} generated by Θ and all operators $\{A_{\varphi} \mid \varphi \in C_0^{\infty}(\mathbf{R})\}$, see Theorem 2.3. We prove that \mathscr{Z} is the maximal abelian subalgebra (masa) of a noncommutative torus \mathscr{A}^{2n}_{RM} , such that $\mathscr{A}^{2n}_{RM} \subset \mathscr{Z} \rtimes \mathbf{Z}$, where the crossed product is taken by the Vershik automorphism of the algebra \mathscr{Z} . It is not hard to see, that our construction is a functor $(X, \phi^t, \mathcal{F}) \mapsto \mathbf{K}$ realizing formula (2.5). Theorem 1.5 follows from (2.5) by comparing the Euler factors of the functions $L(\sigma_{n+1}, s)$ and $L(\mathscr{A}^{2n}_{RM}, s)$. Let us pass to a detailed argument.

Definition 3.1. Let $T^n \cong \mathbf{R}^n / \mathbf{Z}^n$ be an *n*-dimensional torus. The foliation \mathcal{F}_A on T^n is called Anosov's if dim $\mathcal{F}_A = n - 1$ and there exists an automorphism of the lattice \mathbf{Z}^n given by a positive integer matrix $A \in GL_{2n}(\mathbf{Z})$, such that the eigenvector $v_A \in \mathbf{R}^{2n}$ corresponding to the Perron-Frobenius eigenvalue $\lambda_A > 1$ of A is normal to every leaf of \mathcal{F}_A on the covering space \mathbf{R}^n .

Lemma 3.2. (Main lemma) Given foliated space $(T^{2n}, \phi^t, \mathcal{F}_A)$ there exists a Galois number field **K**, such that

$$l(\gamma_{\mathfrak{p}}) = \log N\mathfrak{p},\tag{3.1}$$

where $l(\gamma_{\mathfrak{p}})$ the length of a closed geodesic $\gamma_{\mathfrak{p}}$ and $N\mathfrak{p}$ the norm of the prime ideal \mathfrak{p} of the field **K**.

Proof. (i) Let $X := T^{2n} \cong \mathbf{R}^{2n}/\mathbf{Z}^{2n}$ be a smooth torus of dimension 2n. Consider the Anosov foliation \mathcal{F}_A on T^{2n} of codimension 1. The \mathcal{F}_A is covered by a family $\{l_\alpha\}$ of the parallel hyperplanes $l_\alpha \cong \mathbf{R}^{2n-1} \subset \mathbf{R}^{2n}$. Since \mathcal{F}_A is Anosov's foliation, there exists an automorphism of the lattice \mathbf{Z}^{2n} given by a positive integer matrix $A \in GL_{2n}(\mathbf{Z})$, such that the (scaled) eigenvector $v_A \in \mathbf{R}^{2n}$ corresponding to the Perron-Frobenius eigenvalue $\lambda_A > 1$ of A is normal to every leaf in the family $\{l_\alpha\}$. It is easy to see, that the flow ϕ^t transversal to \mathcal{F}_A is covered by a flow $\tilde{\phi}^t$ on \mathbf{R}^{2n} given by the equation $d\tilde{\phi}^t/dt = v_A$.

(ii) Let Θ be the infinitesimal generator of the flow ϕ^t , see Section 2.1.1. As explained, such a generator gives rise to a bounded linear operator (also denoted by Θ) acting on the Hilbert space $\hat{H}^n_{\mathcal{F}}$, which is canonically associated to the foliated space $(T^{2n}, \phi^t, \mathcal{F}_A)$. The operator Θ is self-adjoint. Let $\varphi \in C_0^{\infty}(\mathbf{R})$ be a test function and consider a trace class operator $A_{\varphi} = \int_{\mathbf{R}} \varphi(t) \phi^t_* dt$ on $\hat{H}^n_{\mathcal{F}}$. It can be verified that A_{φ} commutes with Θ and with each other for all test functions φ . The commutative C^* -algebra generated by Θ and all $\{A_{\varphi} \mid \varphi \in C_0^{\infty}(\mathbf{R})\}$ is denoted by \mathscr{Z} . We identify $\mathscr{Z} \cong C(X)$, where C(X) is the C^* -algebra of complex valued functions on a Hausdorff space X (Gelfand Theorem).

(iii) Consider an AF C^* -algebra \mathbb{A}_{RM}^{2n} given by its dimension group [8, Section 3.5.2]:

$$\mathbb{Z}^{2n} \xrightarrow{A} \mathbb{Z}^{2n} \xrightarrow{A} \mathbb{Z}^{2n} \xrightarrow{A} \dots, \qquad (3.2)$$

where $A \in GL_{2n}(\mathbf{Z})$ is a positive integer matrix corresponding to the foliated space $(T^{2n}, \phi^t, \mathcal{F}_A)$ as explained in (i). Let $\theta = (1, \theta_1, \ldots, \theta_{2n-1})$ be the scaled eigenvector v_A of matrix A and consider the Bratteli compactum X_{θ} of \mathbb{A}_{RM}^{2n} . It follows from [7, Theorem 1 and Remark 1] that $\mathscr{Z} \cong C(X_{\theta})$ and

$$\mathbb{A}_{RM}^{2n} \cong \mathscr{Z} \rtimes \mathbf{Z},\tag{3.3}$$

where the crossed product is taken by the Vershik automorphism of $C(X_{\theta})$. It is well known that the AF C^* -algebra \mathbb{A}_{RM}^{2n} contains a copy of the noncommutative torus \mathscr{A}_{RM}^{2n} [8, Theorem 3.5.3]. We therefore get a map $(T^{2n}, \phi^t, \mathcal{F}_A) \mapsto \mathscr{A}_{RM}^{2n}$ which preserves morphisms in the corresponding categories of objects.

(iv) We apply Theorem 2.7 and Remark 2.8 which say that the noncommutative tori \mathscr{A}_{RM}^{2n} generate the Galois extensions **K** of the field **Q**. Thus one gets a composition of maps which preserves morphisms between the objects in the corresponding categories:

$$(T^{2n}, \phi^t, \mathcal{F}_A) \mapsto \mathscr{A}_{RM}^{2n} \mapsto \mathbf{K}.$$
 (3.4)

(v) The Selberg zeta function $\zeta_{\mathcal{F}_A}(s)$ is given by the formula:

$$\zeta_{\mathcal{F}_A}(s) = \prod_{\gamma_{\mathfrak{p}}} \frac{1}{1 - l(\gamma_{\mathfrak{p}})^{-s}},\tag{3.5}$$

where $l(\gamma_{\mathfrak{p}})$ are lengths of all simple closed geodesics $\gamma_{\mathfrak{p}}$ of the foliated space $(T^{2n}, \phi^t, \mathcal{F}_A)$. Likewise, the Dedekind zeta function of the number field **K** is

$$\zeta_{\mathbf{K}}(s) = \prod_{\mathfrak{p} \subseteq O_{\mathbf{K}}} \frac{1}{1 - N(\mathfrak{p})^{-s}},\tag{3.6}$$

where \mathfrak{p} runs through all prime ideals of the ring of integers $O_{\mathbf{K}}$ of the field \mathbf{K} . In view of (3.4) the zeta functions $\zeta_{\mathcal{F}_A}(s)$ and $\zeta_{\mathbf{K}}(s)$ must coincide as well as their explicit formulas (2.2) and (2.4). The necessary condition for the latter is given by the equation (3.1). Lemma 3.2 is proved.

Definition 3.3. By the zeta function $\zeta_{\mathscr{A}_{RM}^{2n}}(s)$ of the noncommutative torus \mathscr{A}_{RM}^{2n} one understands the Selberg zeta function (3.5) associated to the foliated space $(T^{2n}, \phi^t, \mathcal{F}_A) \mapsto \mathscr{A}_{RM}^{2n}$ according to the formula (3.4).

Lemma 3.4. $\zeta_{\mathscr{A}_{RM}^{2n}}(s) = \left(L(\mathscr{A}_{RM}^{2n},s)\right)^{n+1}.$

Proof. Since σ_{n+1} : Gal (**K**|**Q**) \rightarrow GL_{n+1}(**C**) is an irreducible representation, one gets [Artin 1924] [1, p. 96]:

$$\zeta_{\mathbf{K}}(s) = (L(\sigma_{n+1}, s))^{n+1}, \qquad (3.7)$$

where $L(\sigma_{n+1}, s)$ is the Artin *L*-function. In view of Lemma 3.2 and equivalence $\zeta_{\mathbf{K}}(s) \equiv \zeta_{\mathscr{A}_{PM}^{2n}}(s)$, we conclude from (3.7) that:

$$\zeta_{\mathscr{A}_{RM}^{2n}}(s) = \left(\widetilde{L}(\mathscr{A}_{RM}^{2n}, s)\right)^{n+1}, \qquad (3.8)$$

where $\tilde{L}(\mathscr{A}_{RM}^{2n}, s)$ is an *L*-function with the Euler product constructed from the noncommutative torus \mathscr{A}_{RM}^{2n} . But all such *L*-functions (1.1) must coincide except for a finite number of its Euler factors $\zeta_p(\mathscr{A}_{RM}^{2n}, z)$. Therefore $\tilde{L}(\mathscr{A}_{RM}^{2n}, s) \equiv L(\mathscr{A}_{RM}^{2n}, s)$ up to a finite number of the $\zeta_p(\mathscr{A}_{RM}^{2n}, z)$. From (3.8) one gets $\zeta_{\mathscr{A}_{RM}^{2n}}(s) = (L(\mathscr{A}_{RM}^{2n}, s))^{n+1}$. Lemma 3.4 follows.

Corollary 3.5. $L(\sigma_{n+1}, s) \equiv L(\mathscr{A}_{RM}^{2n}, s)$ except for a finite number of the Euler factors.

Proof. From lemma 3.2 and definition 3.3, one obtains $\zeta_{\mathscr{A}_{RM}^{2n}}(s) \equiv \zeta_{\mathbf{K}}(s) = (L(\sigma_{n+1},s))^{n+1}$. We apply lemma 3.4 and get the equality $L(\sigma_{n+1},s) \equiv L(\mathscr{A}_{RM}^{2n},s)$. Corollary 3.5 is proved.

Theorem 1.5 follows from lemmas 3.2, 3.4 and corollary 3.5.

4. Remarks

Remark 4.1. The maximal abelian subalgebra $\mathscr{Z} \subset \mathscr{A}_{RM}^{2n}$ is generated by the linear operator Θ and trace class operators A_{φ} for all test functions $\varphi \in C_0^{\infty}(\mathbf{R})$. The spectrum of Θ is related to the Riemann Hypothesis for the Dedekind zeta function $\zeta_{\mathbf{K}}(s)$ [Deninger 2001] [2, Corollary 3.5].

Remark 4.2. The reduced leafwise cohomology $\overline{H}^{n}_{\mathcal{F}}(X)$ is receptacle of the Lefschetz Trace Formula for the varieties over the number fields [Deninger 2001] [2, Theorem 3.3]. It will be interesting to relate the $\overline{H}^{n}_{\mathcal{F}}(X)$ with the K-theory of the noncommutative tori \mathscr{A}^{2n}_{RM} .

References

- E. Artin, Über eine neue Art von L-Reihen, Abhandlungen aus dem Mathematischen Seminar der Hamburgerischen Universität, 3 (1924), 89-108.
- C. Deninger, Number theory and dynamical systems on foliated spaces, Jahrsber. Deutsch. Math.-Verein. 103 (2001), 79-100.
- 3. V. G. Drinfeld, Эллиптические модули, Матем. Сборник 94 (1974), 594-627.
- S. Gelbart, An elementary introduction to the Langlands program, Bull. Amer. Math. Soc. 10 (1984), 177-219.
- X. Li, Semigroup C^{*}-algebras, Operator algebras and applications the Abel Symposium 2015, 191-202, Abel Symp., 12, Springer, 2017.
- I. V. Nikolaev, Langlands reciprocity for the even dimensional noncommutative tori, Proc. Amer. Math. Soc. 139 (2011), 4153-4162.
- I. V. Nikolaev, On a Poisson summation formula for noncommutative tori, Complex Anal. Oper. Theory 13 (2019), 3117-3125.
- I. V. Nikolaev, Noncommutative Geometry, Second Edition, De Gruyter Studies in Math. 66, Berlin, 2022.
- I. V. Nikolaev, Non-abelian class field theory and higher dimensional noncommutative tori, arXiv:2309.05779
- M. A. Rieffel, Non-commutative tori a case study of non-commutative differentiable manifolds, Contemp. Math. 105 (1990), 191-211.
- 11. M. Rosen, Number Theory in Function Flelds, GTM 210, Springer, 2002.

 1 Department of Mathematics and Computer Science, St. John's University, 8000 Utopia Parkway, New York, NY 11439, United States.

Email address: igor.v.nikolaev@gmail.com