RESONANT SOLUTIONS OF THE NON-LINEAR SCHRÖDINGER EQUATION WITH PERIODIC POTENTIAL

AREIN DUAIBES, YULIA KARPESHINA

ABSTRACT. The goal is a construction of stationary solutions close to non-trivial combinations of two plane waves at high energies for a periodic non-linear Schrödinger equation in dimension two. The corresponding isoenergetic surfaces are described for every sufficiently large energy.

1. INTRODUCTION

Non-linear Schrödinger equation

$$iu_t = -\Delta u + Vu + \sigma |u|^2 u \tag{1}$$

describes a variety of physical phenomena in optics, acoustics, quantum condensate (Gross-Pitaevskii equation), hydrodynamics, plasma physics, etc. The case of a periodic potential V is of great interest. The equation has been studied for a long time. However, majority of the studies is in physics literature and concerns with one-dimensional situation. Higher dimensions are investigated mostly numerically, (e.g. [1] – [5]) for dimension two). Theoretical papers for periodic multidimensional situations are more sparse. We definitely have to refer here to [6]-[13] devoted to periodic initial value problems with V being zero or an operator of multiplication in the Fourier dual space. We approach (1) somewhat differently. We are interested in stationary solutions of (1) in multiple dimensions. It was proven in [15], [16] that there are stationary solutions of (1) close to plane waves $e^{i(\vec{k},\vec{x})}$ for an extensive set of \vec{k} in dimensions two and three. In this paper we show that there exist stationary solutions close to non-trivial combinations of two plane waves at high energies for a periodic non-linear Schroedinger equation in dimension two. The corresponding isoenergetic surfaces are described.

We start with considering a nonlinear polyharmonic equation:

$$(-\Delta)^{l} u(\vec{x}) + V(\vec{x})u(\vec{x}) + \sigma |u(\vec{x})|^{2} u(\vec{x}) = \lambda u(\vec{x}), \ \vec{x} \in \mathbb{R}^{n},$$
(2)

Date: January 18, 2024.

and the quasi-periodic boundary conditions:

$$\begin{cases} u(x_1, \cdots, x_s + 2\pi, \cdots, x_n) = e^{2\pi i t_s} u(x_1, \cdots, x_s, \cdots, x_n), \\ \frac{\partial}{\partial x_s} u(x_1, \cdots, x_s + 2\pi, \cdots, x_n) = e^{2\pi i t_s} \frac{\partial}{\partial x_s} u(x_1, \cdots, x_s, \cdots, x_n), \\ \vdots \\ \frac{\partial^{2l-1}}{\partial x_s^{2l-1}} u(x_1, \cdots, x_s + 2\pi, \cdots, x_n) = e^{2\pi i t_s} \frac{\partial^{2l-1}}{\partial x_s^{2l-1}} u(x_1, \cdots, x_s, \cdots, x_n), \\ s = 1, \cdots, n. \end{cases}$$
(3)

where l is an integer such that 2l > n or l = 1, n = 2 and $\sigma \in \mathbb{R}$. We consider a periodic potential $V(\vec{x})$. We assume that the potential V is a trigonometric polynomial with the elementary cell of periods $Q = [0, 2\pi]^n$:

$$V(\vec{x}) = \sum_{q \neq 0, |q| \le R_0} v_q e^{i\langle q, \vec{x} \rangle}, \quad 0 < R_0 < \infty,$$

$$\int_Q V(\vec{x}) d\vec{x} = 0.$$
(4)

The last assumption can be done without loss of generality. We require that V is not identically zero. We consider the case l = 1, n = 2 for a sufficiently small V : $||V|| < \varepsilon_*$, here $\varepsilon_* \neq \varepsilon_*(\lambda)$.

We start with discussing the linear operator:

$$H = (-\Delta)^l + V(\vec{x}) \tag{5}$$

in $L^2(\mathbb{R}^n)$. The spectral analysis of the operator H can be reduced to the analysis of operators $H(\vec{t}), \vec{t} \in K$, with $K = [0, 1)^n$ being the elementary cell of dual lattice. The operators $H(\vec{t})$ are defined by (5) and the quasiperiodic conditions (3) in $L^2(Q)$.

It is well-known that $H(\vec{t})$ has a discrete semi-bounded from below spectrum $\bigcup_{n=1}^{\infty} \lambda_n(\vec{t})$. By Bloch's theorem [14], the following is true: 1. The spectrum of H has band structure:

$$\Delta = \bigcup_{n=1}^{\infty} \bigcup_{\vec{t} \in K} \lambda_n(\vec{t}).$$

2. A complete system of generalized eigenfunctions of H can be obtained by extending eigenfunctions of $H(\vec{t})$ quasiperiodically to the whole space \mathbb{R}^n .

We use the notation $H_0(\vec{t})$ for $H(\vec{t})$ with V = 0, its eigenvalues being given by

$$p_j^{2l}(\vec{t}) = |\vec{P}_j(\vec{t})|^{2l},\tag{6}$$

where

$$\vec{P}_j(\vec{t}) = \vec{P}_j(\vec{0}) + \vec{t} = 2\pi j + \vec{t}, \ j \in \mathbb{Z}^n, \ \vec{t} \in K.$$

The corresponding eigenfunction is a plane wave

$$e^{i\langle \vec{P}_j(\vec{t}), \vec{x} \rangle}$$
 (7)

and the corresponding spectral projection is E_i defined by

$$(E_j)_{rm} = \delta_{jr} \delta_{jm} \tag{8}$$

in the basis (7) in $L^2(Q)$. Obviously any $\vec{k} \in \mathbb{R}^n$ can be uniquely written in the form $\vec{k} = \vec{P}_j(\vec{t})$ for some $j \in \mathbb{Z}^n$ and $\vec{t} \in K$. An isoenergetic surface $S_0(k)$ of H_0 is a set of $t \in K$ such that $p_j(t) = k$ for some $j \in \mathbb{Z}^n$. It looks like a sphere "packed" into K. Namely,

$$S_0 = \mathcal{K}S(k),\tag{9}$$

where S(k) is the sphere in \mathbb{R}^n centered at the origin with radius k and

$$\mathcal{K}: \mathbb{R}^n \to K, \quad \mathcal{K}\vec{P}_q(\vec{t}) = \vec{t}.$$
 (10)

The process of obtaining $S_0(k)$ starts by dividing S(k) by the dual lattice $\{\vec{P}_m(0)\}_{m\in\mathbb{Z}^n}$ into pieces and all the pieces are then translated into K.

Perturbation series for eigenvalues and their spectral projections for H(t) with respect to the operator $H_0(\vec{t})$ are obtained in [17]. When 2l > n, perturbation formulas are valid for large k and a set $\chi_0(k, \delta) \subset S_0(k)$ of \vec{t} , such that $p_j(\vec{t}) = k$ for some j and

$$\min_{q \neq 0} |p_j^{2l}(\vec{t}) - p_{j+q}^{2l}(\vec{t})| > k^{2l-n-\delta},\tag{11}$$

 $0 < \delta < 2l - n$. For n = 2, l = 1, some additional inequalities are needed. The inequality (11) means that \vec{t} is sufficiently far from self-intersections of $S_0(k)$. The set $\chi_0(k, \delta)$ has an asymptotically full measure on the isoenergetic surface $S_0(k)$ as $k \to \infty$. It is proven that for every $\vec{t} \in \chi_0(k, \delta)$, the operator H(t) has an eigenvalue close to $p_j^{2l}(\vec{t}) = |\vec{P}_j(\vec{t})|^{2l}$ with an eigenfunction close to $e^{i\langle \vec{P}_j(\vec{t}), \vec{x} \rangle}$. The exact formulas of the eigenvalue and its spectral projection are given in Section 2, Theorem 3. In [15] we proved an analog of Theorem 3 for the non-linear case (2), (3); see Theorem 14. We call (11) the non-resonant case for t. Correspondingly, $\chi_0(k, \delta)$ is the non-resonant set.

The vector \vec{k} is said to satisfy the von Laue Diffraction condition if

$$|\vec{k}| = |\vec{k} - \vec{P}_q(\vec{0})|, \tag{12}$$

for some $q \in \mathbb{Z}^n \setminus \{0\}$. If $\vec{t} : \vec{P_j}(\vec{t}) = \vec{k}$, then, \vec{t} obviously belongs to a self-intersection of $S_0(k)$, Therefore, $\vec{t} \notin \chi_0(k, \delta)$. Perturbation formulas for eigenvalues and spectral projections of $H(\vec{t})$ with respect to $H_0(\vec{t})$ don't work in this situation. The situation when \vec{k} is in a vicinity of (12) is called the resonant case. It turns out that in a vicinity of (12) the eigenvalues and its spectral projections can be approximated by those of a model matrix

$$\begin{pmatrix} p_j^{2l}(\vec{t}) & v_q \\ v_{-q} & p_{j-q}^{2l}(\vec{t}) \end{pmatrix}.$$
 (13)

Further, we assume:

$$q: v_q \neq 0. \tag{14}$$

We denote its eigenvalues and eigenvectors by $\hat{\lambda}^{\pm}$ and \hat{e}^{\pm} , correspondingly. In [17] we described a set $\chi_q(k,\delta) \subset S_0(k) \setminus \chi_0(k,\delta)$ of quasimomenta for which formulas for eigenvalues and spectral projections of $H(\vec{t})$ were constructed for every sufficiently

large k. Indeed, let set $\mathcal{K}S_q(k, n-2+\delta)$ be the image of the spherical layer $S_q(k, n-2+\delta)$ under the map \mathcal{K} given by (10). Here,

$$S_q(k, n-2+\delta) = \{ \vec{x} \in S(k) : \left| |\vec{x}|^2 - |\vec{x} - \vec{P}_q(0)|^2 \right| < 4k^{-n+2-\delta} \}.$$
(15)

In other words, we consider $S_q(k, n-2+\delta)$ being a neighborhood of (12) in the sphere S(k) and then shift it into the cube K using (10). In Section 2 here, we explicitly describe the set $\chi_q(k,\delta)$ which has an asymptotically full measure on $\mathcal{K}S_q(k, n-2+\delta)$. For \vec{t} in a small vicinity of $\chi_q(k,\delta)$ we give formulas for eigenvalues and spectral projections of $H(\vec{t})$. The corresponding eigenfunctions are close to non-trivial combinations of two plane waves $e^{i\langle\vec{k},\vec{x}\rangle}$ and $e^{i\langle\vec{k}-\vec{P}_q(\vec{0}),\vec{x}\rangle}$. The coefficients of plane waves are described through eigenvectors \hat{e}^{\pm} of (13). Thus, we have a pair of eigenfunctions for every \vec{k} . The set $\chi_q(k,\delta)$ is called the resonant set. The exact statement of the result is given in Section 2, Theorem 9. It is proven that the isoenergetic surface corresponding to the pair of solutions is close to two pieces of isoenergetic surface of (13), see Corollary 13.

Here, we will construct formulas for solutions u^{\pm} , λ^{\pm} of (2), (3) when \vec{k} is close to (12). Essentially, we make use of the results proved for the linear case, and apply a method of successive approximation. Namely, we consider the part of equation (2) with the nonlinear term, i.e., $V + \sigma |u^{\pm}|^2$, as an unknown potential. For the method of successive approximation, we define two maps, $\hat{\mathcal{M}}^{\pm}$, and construct two sequences of potentials, W_m^{\pm} converging to the potentials $W^{\pm} = V + \sigma |u^{\pm}|^2$.

The map \mathcal{M}^{\pm} is defined in Section 3.1. Then, the sequence of potentials $\{W_m^{\pm}\}_{m=0}^{\infty}$ is constructed as follows:

$$W_0^{\pm} = V, \tag{16}$$

$$.W_m^{\pm} = \hat{\mathcal{M}}^{\pm} W_{m-1}^{\pm}.$$
 (17)

It turns out that it is a Cauchy sequence converging to a function W with respect to a norm $\|\cdot\|_*$,

$$||W||_{*} = \sum_{j \in \mathbb{Z}^{n}} |w_{j}|, \tag{18}$$

where w_i 's are the Fourier coefficients of W. Namely, we show that

$$\|W^{\pm} - W_m^{\pm}\|_* \le (ck^{-\gamma})^{m+1},\tag{19}$$

for some $\gamma > 0$. Then, we show that convergence of $\{W_m^{\pm}\}_{m=0}^{\infty}$ to W^{\pm} leads to convergence of the sequences of the spectral projections $\{\hat{E}_m^{\pm}(\vec{t})\}_{m=0}^{\infty}$ and their corresponding eigenvalues $\{\hat{\lambda}_m^{\pm}(\vec{t})\}_{m=0}^{\infty}$ to $\hat{E}_W^{\pm}(\vec{t})$ and $\hat{\lambda}_W^{\pm}(\vec{t})$, here $\hat{E}_m^{\pm}(\vec{t})$, $\hat{\lambda}_m^{\pm}(\vec{t})$ and $\hat{E}_W^{\pm}(\vec{t})$, $\hat{\lambda}_W^{\pm}(\vec{t})$ are spectral projections and their eigenvalues of $H_0 + W_m^{\pm}$ and $H_0 + W^{\pm}$. Corresponding to $\hat{E}_W^{\pm}(\vec{t})$ eigenfunction $u^{\pm} \equiv \hat{u}_W^{\pm}$, see (70), (71), solves (2), (3). It is shown that u^{\pm} is close to a combination of two plane waves $e^{i\langle\vec{k},\vec{x}\rangle}$ and $e^{i\langle\vec{k}-\vec{P}_q(\vec{0}),\vec{x}\rangle}$ under some restriction on its amplitude. The coefficients of plane waves are described by eigenvectors \hat{e}^{\pm} of (13) with a good accuracy. The exact statement of the result is given by RESONANT SOLUTIONS OF THE NON-LINEAR SCHRÖDINGER EQUATION WITH PERIODIC POTENTIAL

Theorem 24. The sequences $\{W_m^{\pm}\}_{m=0}^{\infty}$, $\{\hat{E}_m^{\pm}(\vec{t})\}_{m=0}^{\infty}$ and $\{\hat{\lambda}_m^{\pm}(\vec{t})\}_{m=0}^{\infty}$. can be differentiated with respect to \vec{t} and maintain their convergence. As a result the asymptotic formulas (139), (140) for $\nabla \hat{E}_W^{\pm}(\vec{t})$, $\nabla \hat{\lambda}_W^{\pm}(\vec{t})$ are proven, and $|\nabla \hat{\lambda}_W^{\pm}(\vec{t})| \approx 2lk^{2l-1}$. It follows that the surface $\lambda^{\pm}(\vec{t}) = \lambda_0$ is in the $C(V)\lambda_0^{-\hat{\gamma}}$ -neighborhood of $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$ for every sufficiently large λ_0 , here $\hat{\lambda}^{\pm}$ are eigenvalues of (13) and $\hat{\gamma} > (2l-1)(2l)^{-1}$, see Theorem 29 and Corollary 30.

In Section 4 we consider the physically interesting case l = 1, n = 2. All considerations of the previous case can be done for l = 1, n = 2 under the assumption that Vis sufficiently small $||V||_* < \epsilon^9$, $0 < \epsilon < \epsilon_0$, $\epsilon_0 \neq \epsilon_0(\lambda)$, for the waves with a sufficiently small amplitude A, the restriction on A being given in terms of ϵ , see (148).

The isoenergetic surface $\lambda^{\pm}(\vec{t}) = \lambda_0$ is in a small -neighborhood of $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$.

If V is not small, then there are solutions close to non-trivial combinations of more than two plane waves. In this case the model operator is described not by (13), but by a Hill operator. This result will be proven in a forthcoming paper.

The paper is organized as follows. Preliminary results are described in Section 2, Section 2.1 containing the results for a linear case [17] and Section 2.2 containing the results for the non-linear equation, non-resonant case [15]. Chapter 3 is devoted to proving the main result for the case 2l > n. We consider l = 1, n = 2 in Chapter 4.

Acknowledgement. The authors are thankful to Günter Stolz for usefull discussions.

2. Preliminary Results

In this chapter, we present a brief review of previous results needed for proof of the main result of this paper.

2.1. Linear Polyharmonic Equation with Periodic Potential. Let us consider an operator in $L_2(Q)$ given by the differential expression:

$$H_{\alpha}(\vec{t})u = (-\Delta)^{l}u + \alpha V u, \qquad (20)$$

with the quasi-periodic boundary conditions (3). Here l is an integer such that 2l > n, $-1 \le \alpha \le 1$.

Perturbation series for eigenvalues and its spectral projections for $H_{\alpha}(t)$ are constructed on a nonsingular set χ_0 [17]. We include here a construction of $\chi_0(k, \delta)$ and discuss the perturbation theory for $H_{\alpha}(\vec{t})$.

Lemma 1. For an arbitrarily small positive, δ , $2\delta < 2l - n$, and sufficiently large $k > k_0(\delta)$, there exists a non-resonant set $\chi_0(k, \delta)$, belonging to the isoenergetic surface $S_0(k)$ of the free operator $H_0(t)$, such that, for any \vec{t} in it,

1. there exists a unique $m \in \mathbb{Z}^n$ such that $p_m^{2l}(\vec{t}) = k^{2l}$, (21)

2.
$$\min_{j \neq m} |p_j^{2l}(\vec{t}) - p_m^{2l}(\vec{t})| > k^{2l-n-\delta}.$$
(22)

Moreover, the nonsingular set has an asymptotically full measure on $S_0(k)$:

$$\frac{s(S_0(k) \setminus \chi_0(k,\delta))}{s(S_0(k))} = O(k^{-\delta/2}), \text{ as } k \to \infty,$$

where s(.) denotes the Lebesgue measure.

Corollary 2. Suppose \vec{t} belongs to the $(k^{-n+1-2\delta})$ -neighborhood in K of the resonant set $\chi_0(k, \delta)$, $0 < 2\delta < 2l-n$. Then for all z lying on the circle $C_0 = \{z \in \mathbb{C} : |z-k^{2l}| = k^{2l-n-\delta}\}$ and any i of \mathbb{Z}^n , the inequality

$$2\mid p_i^{2l}(\vec{t})-z\mid >k^{2l-n-\delta}$$

holds.

Let us introduce the functions $g_r(k, \vec{t})$ and the operator-valued functions $G_r(k, \vec{t})$, r = 0, 1, ..., for $\vec{t} \in \chi_0(k, \delta)$:

$$g_r(k,\vec{t}) = \frac{(-1)^r}{2\pi i r} \operatorname{Tr} \oint_{C_0} \left((H_0(\vec{t}) - z)^{-1} V \right)^r dz,$$
(23)

$$G_r(k,\vec{t}) = \frac{(-1)^{r+1}}{2\pi i} \oint_{C_0} \left((H_0(\vec{t}) - z)^{-1} V \right)^r (H_0(\vec{t}) - z)^{-1} dz.$$
(24)

The following theorem presents the main result for (20).

Theorem 3. Suppose \vec{t} belongs to the $(k^{-n+1-2\delta})$ -neighborhood in K of the nonresonant set $\chi_0(k,\delta)$, $0 < 2\delta < 2l - n$. Then for sufficiently large k, $k > k_0(||V||, \delta)$, there exists a single eigenvalue of the operator $H(\vec{t})$ in the interval $\varepsilon(k,\delta) \equiv (k^{2l} - k^{2l-n-\delta}, k^{2l} + k^{2l-n-\delta})$. It is given by the series

$$\lambda_j(\alpha, \vec{t}) = p_j^{2l}(\vec{t}) + \sum_{r=2}^{\infty} \alpha^r g_r(k, \vec{t}), \qquad (25)$$

converging absolutely in the disk $|\alpha| \leq 1$, where the index j is uniquely determined from the relation $p_j^{2l}(\vec{t}) \in \varepsilon(k, \delta)$ and the spectral projection, corresponding to $\lambda_j(\alpha, \vec{t})$ is given by the series

$$E_j(\alpha, \vec{t}) = E_j + \sum_{r=1}^{\infty} \alpha^r G_r(k, \vec{t}), \qquad (26)$$

which converges in the trace class \mathbf{S}_1 uniformly with respect to α in the disk $|\alpha| \leq 1$. Moreover, for the coefficients $g_r(k, \vec{t})$ and $G_r(k, \vec{t})$, the following estimates hold.

$$|g_r(k,\vec{t})| < k^{2l-n-\delta} k^{-(2l-n-2\delta)r},$$
(27)

$$\|G_r(k,\vec{t})\|_1 \le k^{-(2l-n-2\delta)r}.$$
(28)

The series (25) and (26) are differentiable termwise with respect to \vec{t} in the $(k^{-n+1-2\delta})$ -neighborhood in \mathbb{C}^n of the set $\chi_0(k, \delta)$, see [17]. Indeed, let

$$T(m) \equiv \frac{\partial^{|m|}}{\partial t_1^{m_1} \partial t_2^{m_2} \dots \partial t_n^{m_n}}, \text{ where } |m| \equiv m_1 + m_2 + \dots + m_n,$$
(29)

RESONANT SOLUTIONS OF THE NON-LINEAR SCHRÖDINGER EQUATION WITH PERIODIC POTENTIAL

$$m! \equiv m_1! m_2! \dots m_n!, \quad 0 \le |m| < \infty, \quad T(0)f \equiv f.$$
 (30)

Theorem 4. Under the conditions of Theorem 3 the series (25) and (26) can be differentiated with respect to \vec{t} any number of times, and they retain their asymptotic character. The coefficients $g_r(k, \vec{t})$ and $G_r(k, \vec{t})$ satisfy the following estimates in the $(k^{-n+1-2\delta})$ -neighborhood in \mathbb{C}^n of the nonsingular set $\chi_0(k, \delta)$:

$$|T(m)g_r(k,\vec{t})| < m!k^{-(2l-n-2\delta)(r-1)}k^{|m|(n-1+2\delta)}$$
(31)

$$||T(m)G_r(k,\vec{t})||_1 < m!k^{-(2l-n-2\delta)r}k^{|m|(n-1+2\delta)}.$$
(32)

Corollary 5. The following estimates hold for the perturbed eigenvalue and its spectral projection.

$$|T(m)(\lambda_j(\alpha, \vec{t}) - p_j^{2l}(\vec{t}))| < cm! k^{-(2l-n-2\delta)} k^{|m|(n-1+2\delta)}$$
(33)

$$||T(m)(E_j(\alpha, \vec{t}) - E_j)||_1 < cm! k^{-(2l - n - 2\delta)} k^{|m|(n - 1 + 2\delta)}$$
(34)

In particular,

$$|\lambda_j(\alpha, \vec{t})) - p_j^{2l}(\vec{t})| < ck^{-(2l-n-2\delta)}$$

$$(35)$$

$$\|E_{j}(\alpha, \vec{t}) - E_{j})\|_{1} < ck^{-(2l-n-2\delta)}$$
(36)

$$|\nabla_{\vec{t}}\lambda_{j}(\alpha,\vec{t}) - 2l\vec{P}_{j}(\vec{t})p_{j}^{2l-2}(\vec{t})| < ck^{-(2l-n-2\delta)+n-1+2\delta}.$$
(37)

Corollary 6. The surface $\lambda_j(\alpha, \vec{t}) = \lambda_0$ is in the real $\lambda_0^{-(4l-n+1-2\delta)}$ -neighborhood of $\chi_0(k, \delta)$ for every sufficiently large λ_0 .

Next, we consider formulas for $\vec{t} \in S_0(k) \setminus \chi_0(k, \delta)$. This means that there is at least one $q \in \mathbb{Z}^n$ such that

$$|p_{j-q}^{2l}(\vec{t}) - p_{j}^{2l}(\vec{t})| < k^{2l-n-\delta}, \ \vec{P_{j}}(\vec{t}) \in S(k).$$

Below, we present the main results and the perturbation formulas for an eigenvalue and its spectral projection for \vec{t} belonging to a "resonant set", $\chi_q(k,\delta) \subset S_0(k) \setminus \chi_0(k,\delta)$. We set $\alpha = 1$ for simplicity.

Let P_q be the diagonal operator in $l^2(\mathbb{Z}^n)$ defined by the formula

$$(P_q)_{mm} = \delta_{jm} + \delta_{j-q,m}.$$
(38)

We define the operator $\hat{H}_q(\vec{t})$ as

$$\hat{H}_q(\vec{t}) = H_0(\vec{t}) + P_q V P_q.$$
 (39)

Note that the matrix corresponding to this operator has only two non-diagonal elements, namely,

$$\hat{H}_q(\vec{t})_{j,j-q} = \overline{\hat{H}_q(\vec{t})}_{j-q,j} = v_q.$$

Thus $\hat{H}_q(\vec{t})$ has only one block (13), here we assume (14). The eigenvalues of this block matrix we denote by $\hat{\lambda}^+(\vec{t})$ and $\hat{\lambda}^-(\vec{t})$. They are given as

$$\hat{\lambda}^+(\vec{t}) = a + b \text{ and } \hat{\lambda}^-(\vec{t}) = a - b, \tag{40}$$

where $2a = p_j^{2l}(\vec{t}) + p_{j-q}^{2l}(\vec{t})$ and $2b = (4|v_q|^2 + (p_j^{2l}(\vec{t}) - p_{j-q}^{2l}(\vec{t}))^2)^{1/2}$. Note that $\hat{\lambda}^+(\vec{t}) - \hat{\lambda}^-(\vec{t}) = 2b \ge 2|v_q|$.

This means that the spectrum of $\hat{H}_q(\vec{t})$ is $\{\{p_i^{2l}(\vec{t})\}_{i\neq j,j-q}, \hat{\lambda}^+(\vec{t}), \hat{\lambda}^-(\vec{t})\}$.

Definition 1. The spectral projections of $\hat{H}_q(\vec{t})$ corresponding to $\hat{\lambda}^{\pm}(\vec{t})$ we denote by $= \hat{E}^{\pm}$.

Recall that $\mathcal{K}S_q(k, n-2+\delta)$ was defined in Section 1, see (10), (15).

Lemma 7. For an arbitrarily small positive δ , $2\delta < 2l - n$, and sufficiently large k, there exists a subset $\chi_q(k, \delta)$ of $\mathcal{K}S_q(k, n - 2 + \delta)$, such that the following conditions hold

1. there exists
$$j \in \mathbb{Z}^n$$
 such that $P_j(\vec{t}) = \vec{k}$,
2. $|p_{j-q}^2(\vec{t}) - p_j^2(\vec{t})| < k^{-n+2-\delta}$,
3. $\min_{i \neq j, j-q} |p_j^2(\vec{t}) - p_i^2(\vec{t})| > 2k^{-n+2-\delta\delta}$.

for all $\vec{t} \in \chi_q(k, \delta)$. Moreover, for any \vec{t} in the k-neighborhood of $\chi_q(k, \delta)$ in \mathbb{C}^n , there exists a unique $j \in \mathbb{Z}^n$ such that $|p_j^2(\vec{t}) - k^2| < 5k^{-n+2-6\delta}$ and the second and third conditions above are satisfied. Also, the set $\chi_q(k, \delta)$ has an asymptotically full measure on $\mathcal{K}S_q(k, n-2+\delta)$, that is

$$\frac{s(\mathcal{K}S_q(k,n-2+\delta)\setminus\chi_q(k,\delta))}{s(\mathcal{K}S_q(k,n-2+\delta))} = O(k^{-\delta/2}) \quad as \ k \to \infty.$$
(41)

The previous lemma means that $p_j^2(\vec{t})$ and $p_{j-q}^2(\vec{t})$ are close to each other, but they are sufficiently far away from the remaining eigenvalues.

Corollary 8. If \vec{t} belongs to the $(2k^{-n+1-7\delta})$ - neighborhood of $\chi_q(k,\delta)$, then for all z on the circle $C_1^+ = \{z : |z - \hat{\lambda}^+(\vec{t})| = d\}, d = \frac{1}{10}|v_q|$, both of the following inequalities are true:

$$2|p_m^{2l}(\vec{t}) - z| \ge k^{2l - n - 6\delta}, m \ne j, j - q_0,$$
(42)

$$|\hat{\lambda}^{-}(\vec{t}) - z| \ge d. \tag{43}$$

Similar corollary holds for $\hat{\lambda}^-, C_1^-$. Further for definiteness we consider $\hat{\lambda}^+, C_1^+$.

Let V be as in (4) and functions $\hat{g}_r^+(k, \vec{t})$ and $\hat{G}_r^+(k, \vec{t})$, $r \in \mathbb{N}$, $t \in \chi_q(k, \delta)$ be defined as follows:

$$\hat{g}_{r}^{+}(k,\vec{t}) = \frac{(-1)^{r}}{2\pi i r} \operatorname{Tr} \oint_{C_{1}^{+}} \left((\hat{H}_{q}(\vec{t}) - z)^{-1} \hat{V} \right)^{r} dz,$$
(44)

$$\hat{G}_{r}^{+}(k,\vec{t}) = \frac{(-1)^{r+1}}{2\pi i} \oint_{C_{1}^{+}} \left(\left(\hat{H}_{q}(\vec{t}) - z \right)^{-1} \hat{V} \right)^{r} \left(\hat{H}_{q}(\vec{t}) - z \right)^{-1} dz, \tag{45}$$

here and below

$$\hat{V} = V - P_q V P_q.$$

The following result is proven in [17].

Theorem 9. Suppose \vec{t} belongs to the $(k^{-n+1-7\delta})$ -neighborhood in K of the set $\chi_q(k, \delta)$, $0 < 9\delta < 2l - n$. Then for sufficiently large k, $k > k_1(V, \delta)$, in the interval $\hat{\varepsilon}(k, \delta) \equiv (\hat{\lambda}^+(\vec{t}) - k^{-\delta}, \hat{\lambda}^+(\vec{t}) + k^{-\delta})$, there exists a single eigenvalue of the operator $H(\vec{t})$. It is given by the series

$$\hat{\lambda}^{q,+}(\vec{t}) = \hat{\lambda}^{+}(\vec{t}) + \sum_{r=2}^{\infty} \hat{g}_{r}^{+}(k,\vec{t}).$$
(46)

The spectral projection, corresponding to $\hat{\lambda}^{q}(\vec{t})$ is given by

$$\hat{E}^{q,+}(\vec{t}) = \hat{E}^{+} + \sum_{r=1}^{\infty} \hat{G}^{+}_{r}(k,\vec{t}), \qquad (47)$$

which converges in the class \mathbf{S}_1 . The following estimates hold for $\hat{g}_r^+(k, \vec{t})$, $\hat{G}_r^+(k, \vec{t})$:

$$|\hat{g}_r^+(k,\vec{t})| < k^{-\gamma_1 r - \delta},\tag{48}$$

$$\|\hat{G}_{r}^{+}(k,\vec{t})\|_{1} < k^{-\gamma_{1}r},\tag{49}$$

where

$$\gamma_1 = (2l - n)/2 - 4\delta > 0. \tag{50}$$

Theorem 10. Under the conditions of Theorem 9, the series (46) and (47) can be differentiated termwise with respect to \vec{t} any number of times, and they retain their asymptotic character. The coefficients $\tilde{g}_r(k, \vec{t})$ and $\tilde{G}_r(k, \vec{t})$ satisfy the following estimates in the $(k^{-n+1-7\delta})$ -neighborhood in \mathbb{C}^n of the singular set $\chi_q(k, \delta)$:

$$|T(m)\hat{g}_{r}^{+}(k,\vec{t})| < m!k^{-\gamma_{1}r-\delta+|m|(n-1+7\delta)},$$
(51)

$$\|T(m)\hat{G}_{r}^{+}(k,\vec{t})\|_{1} < m!k^{-\gamma_{1}r+|m|(n-1+7\delta)}.$$
(52)

Corollary 11. The following estimates for the perturbed eigenvalue and its spectral projection hold:

$$|T(m)(\hat{\lambda}^{q,+}(\vec{t}) - \hat{\lambda}^{+}(\vec{t}))| < 2m! k^{-2\gamma_1 - \delta + |m|(n-1+7\delta)},$$
(53)

$$\|T(m)(\hat{E}^{q,+}(\vec{t}) - \hat{E}^{+}(\vec{t}))\|_{1} < 2m!k^{-\gamma_{1}+|m|(n-1+7\delta)}.$$
(54)

In particular, the following estimates are valid:

$$\hat{\lambda}^{q,+}(\vec{t}) - \hat{\lambda}^{+}(\vec{t})| < 2k^{-2\gamma_1 - \delta},$$
(55)

$$\|\hat{E}^{q,+}(\vec{t}) - \hat{E}^{+}\|_{1} < 2k^{-\gamma_{1}},\tag{56}$$

$$|\nabla \hat{\lambda}^{q,+}(\vec{t}) - \nabla \hat{\lambda}^{+}(\vec{t})| < 2k^{-2\gamma_1 + n - 1 + 6\delta}.$$
(57)

Corollary 12. An eigenfunction $u_0^+(\vec{x}, \vec{t})$ corresponding to $\hat{E}^{q,+}(\vec{t})$ satisfies the estimate:

$$\left\| e^{-i < \vec{k}, \vec{x} >} u_0^+(\vec{x}, \vec{t}) - \left(a_1 + a_2 e^{-i < \vec{P}_q(0), \vec{x} >} \right) \right\|_* < ck^{-\gamma_1}, \tag{58}$$

where (a_1, a_2) is an eigenvector of (13) corresponding to $\lambda^+(t)$.

The analogous results hold for $\hat{\lambda}^{q,-}(\vec{t}), \hat{E}^{q,-}(\vec{t})$.

Corollary 13. The surface $\hat{\lambda}^{q,\pm}(\vec{t}) = \lambda_0$ is in the real $\lambda_0^{-2\gamma_1 - 2l + 1 - \delta}$ -neighborhood of $\chi_q(k, \delta)$ for every sufficiently large λ_0 .

The results above hold for $V : ||V||_* < \infty$.

2.2. Nonlinear Periodic Polyharmonic Equation. In this section we consider equation (2) with the quasi-periodic boundary conditions (3). In [15, 16] we proved existence of a quasi-periodic solution of equation (2) being close to a plane wave $Ae^{i\langle \vec{k},\vec{x}\rangle}$ for every \vec{k} belonging to a non-resonant set $\chi_0(k)$, where A is a complex number with sufficiently small |A|. The main idea of the papers is to look at equation (2) as a linear equation. This can be done by considering the sum $V(\vec{x}) + \sigma |u(\vec{x})|^2$ as an unknown potential. We use the facts proved for the linear case.

Now we describe the technique used to find a function u that solves the nonlinear equation. Let \hat{l}_1 be the space of functions in Q with Fourier coefficients in $l_1(\mathbb{Z}^n)$, see (18). First, a sequence of operators $\{W_m\}_{m=0}^{\infty}$ is constructed via the map $\mathcal{M} : \hat{l}_1 \to \hat{l}_1$ defined as

$$\mathcal{M}W(\vec{x}) = V(\vec{x}) + \sigma |u_{\widetilde{W}}(\vec{x})|^2.$$
(59)

Here, $u_{\widetilde{W}}$ is an eigenfunction of the linear operator $(-\Delta)^l + \widetilde{W}$ with the same boundary conditions (3), where \widetilde{W} is defined by the formula

$$\widetilde{W}(\vec{x}) = W(\vec{x}) - \frac{1}{(2\pi)^n} \int_Q W(\vec{x}) \, d\vec{x}.$$
(60)

More precisely, $u_{\widetilde{W}}$ is defined as

$$u_{\widetilde{W}}(\vec{x}) = \sum_{s \in \mathbb{Z}^n} (E_{\widetilde{W}}(\vec{t}))_{sj} e^{i\langle \vec{P}_s(\vec{t}), \vec{x} \rangle}$$

$$= e^{i\langle \vec{P}_j(\vec{t}), \vec{x} \rangle} + \sum_{r=1}^{\infty} \sum_{s \in \mathbb{Z}^n} (G_{\widetilde{W}}^r(k, \vec{t}))_{sj} e^{i\langle \vec{P}_s(\vec{t}), \vec{x} \rangle},$$

(61)

here $j : \vec{P}_j(\vec{t}) = \vec{k}$. Formula (61) is analogous to (26). Second, the operator W_m is defined by a recurrence procedure as follows. Let $W_0 = V + \sigma |A|^2$. Define

$$W_{m+1} = \mathcal{M}W_m, \ m = 0, 1, 2, \dots$$
 (62)

Then, W_m is proved to be a Cauchy sequence of periodic potentials converging to a periodic potential W with respect to the norm $\|\cdot\|_*$, see (18). Note that for any m, there is a solution u_m of the equation

$$(-\Delta)^l u_m + \overline{W}_m u_m = \lambda_m u_m,$$

given by the formula (61). Moreover, the sequences of the functions $\{u_m\}_{m=0}^{\infty}$ and the eigenvalues $\{\lambda_m\}_{m=0}^{\infty}$ converge to a function $u_{\widetilde{W}}$ and a real number λ , respectively. It is shown that $u = u_{\widetilde{W}}$ solves the nonlinear equation (2). The limits $E_{\widetilde{W}}(\vec{t})$ and $\lambda_{\widetilde{W}}(\vec{t})$ of the sequences $\{E_m(\vec{t})\}_{m=0}^{\infty}$ and $\{\lambda_m(\vec{t})\}_{m=0}^{\infty}$, respectively, are given by the two series (25), (26) written for the potential \widetilde{W} instead of V. The following theorem holds when 2l > n.

Theorem 14. Let $0 < 2\delta < 2l - n$. Suppose \vec{t} belongs to the $(k^{-n+1-2\delta})$ -neighborhood in K of the non-resonant set $\chi_0(k, \delta)$, $k > k_1(||V||_*, \delta)$, and $A \in \mathbb{C} : |\sigma||A|^2 < k^{\gamma_0 - \delta}$, where

$$\gamma_0 = 2l - n - 2\delta. \tag{63}$$

Then, there is a function u, depending on \vec{t} as a parameter, and a real value $\lambda(\vec{t})$, such that they solve the equation (2) and the quasi-periodic boundary condition (3). The following estimates hold:

$$\lambda(\vec{t}) = p_j^{2l}(\vec{t}) + \sigma |A|^2 + O\left(k^{-\gamma_0}\right),\tag{64}$$

$$u(\vec{x}) = A e^{i \langle \vec{P}_j(\vec{t}), \vec{x} \rangle} \left(1 + \tilde{u}(\vec{x}) \right), \tag{65}$$

where \tilde{u} is periodic and

$$\|\tilde{u}\|_* < k^{-\gamma_0}.$$
 (66)

Moreover, the following estimates hold:

$$\left|\nabla\left(\lambda(\vec{t}) - p_j^{2l}(\vec{t}) - \sigma |A|^2\right)\right| < k^{2l-1-\gamma_0}.$$
(67)

Corollary 15. The surface $\lambda(\vec{t}) = \lambda_0$, $\lambda_0 \equiv k_0^{2l} + \sigma |A|^2$, is in the $k_0^{-(4l-n+1-2\delta)}$ -neighborhood of $\chi_0(k, \delta)$ for every fixed $A : |\sigma A^2| < k_0^{\gamma_0 - \delta}$ and sufficiently large λ_0 .

If l = 1, n = 2 an analogous theorem holds with somewhat different non-resonant set set $\chi_0(k, \delta)$, and a constant γ_0 .

3. Perturbation Theory for Non-Linear Polyharmonic Equation with Periodic Potential for 2l > n. Resonant Case.

In Section 3.1, we introduce maps $\hat{\mathcal{M}}^{\pm}$ and construct two sequences of potentials $\{W_m^{\pm}\}_{m=0}^{\infty}$. We will prove that they are Cauchy sequences converging to some potentials W^{\pm} . In Section 3.2 we prove the existence of solutions $u^{\pm}(\vec{t}, \vec{x}), \lambda = \lambda^{\pm}(\vec{t})$ of (2), (3) and obtain their estimates at the high energy region. The functions $u^{\pm}(\vec{t}, \vec{x})$ are shown to be close to non-trivial combinations of two plane waves. Finally, in Section 3.3, we derive estimates for the derivatives of $\lambda^{\pm}(\vec{t})$ and $u^{\pm}(\vec{t}, \vec{x})$ with respect to \vec{t} . The isoenergetic surface $\{t \in K, \lambda^{\pm}(\vec{t}) = \lambda_0\}$ is constructed for the resonant case for every sufficiently large λ_0 .

3.1. Construction of a Cauchy Sequence. Recall that the operator $\hat{H}_q(\vec{t})$ defined by (39) has the spectrum $\{\{p_i^{2l}(\vec{t})\}_{i\neq j,j=q}, \hat{\lambda}^+(\vec{t}), \hat{\lambda}^-(\vec{t})\}$. It is easy to verify that eigenvectors of the block submatrix of $\hat{H}_q(\vec{t})$ corresponding to the eigenvalues $\hat{\lambda}^{\pm}(\vec{t})$, see (40), are:

$$\hat{e}^{+} = (p_{j-q}^{2l}(\vec{t}) - \hat{\lambda}^{+}(\vec{t}), \ -v_q)^T$$
(68)

$$\hat{e}^{-} = (-v_{-q}, \ p_{j}^{2l}(\vec{t}) - \hat{\lambda}^{-}(\vec{t}))^{T}.$$
(69)

By Definition 1 the corresponding spectral projections are denoted by by $\hat{E}^{\pm}(\vec{t})$. It is clear that $|\hat{\lambda}^{+}(\vec{t}) - p_{j}^{2l}(\vec{t})| < 2|v_{q}|$ and $|\hat{\lambda}^{-}(\vec{t}) - p_{j}^{2l}(\vec{t})| < 2|v_{q}|$. We assume that $|v_{q}| \neq 0$.

The cases + and - are analogous. Further, for definiteness we consider the case +.

Wee use a geometric Lemma 7 and Corollary 8.

Definition 2. Let W be such that $||W||_* < \infty$, $\vec{t} \in \chi_q(k, \delta)$ and

$$u_W^+(\vec{x}) = \psi_W^+(\vec{x})e^{i\langle P_j(\vec{t}),\vec{x}\rangle},\tag{70}$$

where ψ_W^+ is periodic and

$$\psi_W^+(\vec{x}) = A \sum_{b \in \mathbb{Z}^n} \hat{E}_W^{q,+}(\vec{t})_{j-b,j} e^{i \langle \vec{P}_b(0), \vec{x} \rangle}$$
(71)

 $\hat{E}_W^{q,+}$ being given by (45), (47) with $W - P_q V P_q$ in (45) instead of \hat{V} .

Obviously, $u_W^+(x)$ satisfies $(-\Delta)^l u_W^+ + W u_W^+ = \lambda_W^+(t) u_W^+$, where $\lambda_W^+(t)$ is given by (46) with $W - P_q V P_q$ in (45) instead of \hat{V} .

Definition 3. Let the map
$$\hat{\mathcal{M}}^+: \hat{l}_1 \to \hat{l}_1$$
 be defined by
 $\hat{\mathcal{M}}^+ W(\vec{x}) = V(\vec{x}) + \sigma |u_W^+(\vec{x})|^2,$ (72)

here \hat{l}_1 is the space of functions in Q with Fourier coefficients in $l_1(\mathbb{Z}^n)$, see (18).

Next, we define the sequence $\{W_m^+\}_{m=0}^\infty$ using the map $\hat{\mathcal{M}}^+$. Let

$$W_0^+ = V, \tag{73}$$

$$W_m^+ = \hat{\mathcal{M}}^+ W_{m-1}^+.$$
(74)

For every m there is an eigenfunction $u_m^+(\vec{x})$ corresponding W_m^+ , see (70), and the corresponding eigenvalue number λ_m^+ . Note that this sequence W_m^+ is quite different from that for a non-resonance case given by (62).

Definition 4. Let, $\hat{E}_{-1}^+ \equiv \hat{E}^+$ be the spectral projection of the model operator \hat{H}_q corresponding to $\lambda^+(\vec{t})$, and $\hat{E}_0^+ = \hat{E}^{q,+}$, see (47). By analogy, $\hat{E}_m^+ \equiv \hat{E}_{W_m^+}^{q,+}(\vec{t})$, the formula (45), (47) being used for $E_{W_m^+}^+(\vec{t})$ with $W_m^+ - P_q V P_q$ instead of \hat{V} .

Clearly, \hat{E}_{-1}^+ is a one-dimensional projection corresponding to (68).

Definition 5. Let

$$u_{-1}^{+}(\vec{x}) = \psi_{-1}^{+}(\vec{x})e^{i\langle \vec{P}_{j}(\vec{t}), \vec{x} \rangle}, \tag{75}$$

$$u_0^+(\vec{x}) = \psi_0^+(\vec{x})e^{i\langle \vec{P}_j(\vec{t}), \vec{x} \rangle},\tag{76}$$

where ψ_{-1}^+ and ψ_0^+ are the periodic parts of u_{-1}^+ and u_0^+ , written in terms of $\hat{E}^+(\vec{t})$ and $E^{q,+}(\vec{t})$ as follows. First,

$$\psi_{-1}^{+}(\vec{x}) = A\left(\hat{E}^{+}(\vec{t})_{j,j} + \hat{E}^{+}(\vec{t})_{j-q,j}e^{-i\langle \vec{P}_{q}(0), \vec{x} \rangle}\right)$$
(77)

The sum in (77) has only two terms, because $\hat{E}_{lm}^+ = 0$ if $l, m \neq 0, -q$. Note that $u_{-1}^+(\vec{x})$ is a linear combination of $e^{i\langle \vec{P}_j(\vec{t}), \vec{x} \rangle}$ and $e^{i\langle \vec{P}_{j-q}(\vec{t}), \vec{x} \rangle}$.

12

Second,

$$\psi_0^+(\vec{x}) = A \sum_{b \in \mathbb{Z}^n} \hat{E}^{q,+}(\vec{t})_{j-b,j} e^{i \langle \vec{P}_b(0), \vec{x} \rangle},\tag{78}$$

see (47) and Definition 4.

Next, by analogy to (75) and (77),

$$u_{s}^{+}(\vec{x}) = \psi_{s}^{+}(\vec{x})e^{i\langle P_{j}(\vec{t}),\vec{x}\rangle},$$
(79)

 ψ_s^+ being periodic and given by the formula:

$$\psi_s^+(\vec{x}) = A \sum_{b \in \mathbb{Z}^n} \hat{E}_s^+(\vec{t})_{j-b,j} e^{i\langle \vec{P}_b(0), \vec{x} \rangle}.$$
(80)

Definition 6. Let T be an operator in $l_2(\mathbb{Z}^n)$. Then $||T||_0$ is defined by

$$||T||_0 = \frac{1}{2} \max_r \sum_p \left(|T_{pr}| + |T_{rp}| \right).$$
(81)

It is obvious that $\|\mathcal{AB}\|_0 \leq 2\|\mathcal{A}\|_0\|\mathcal{B}\|_0$.

Remind that $2\gamma_1 = 2l - n - 8\delta$, see (50).

Lemma 16. Let $\gamma_2 > 0$, $0 < 8\delta < 2l - n$. The following estimates hold for every $m=1, 2, ..., and \forall A \in \mathbb{C} : |\sigma||A|^2 < k^{-\gamma_2}$ when $\vec{t} \in \chi_q(k, \delta)$:

$$\|W_m^+ - W_{m-1}^+\|_* < (\hat{c}k^{-\gamma})^m, \tag{82}$$

$$\|W_m^+ - V\|_* < \sum_{r=1}^m (\hat{c}k^{-\gamma})^r, \tag{83}$$

$$\|\hat{E}_{0}^{+}(\vec{t}) - \hat{E}_{-1}^{+}(\vec{t})\|_{0} < \hat{c}k^{-\gamma},$$
(84)

$$\|\hat{E}_{m-1}^{+}(\vec{t}) - \hat{E}_{m-2}^{+}(\vec{t})\|_{0} < \hat{c}(\hat{c}k^{-\gamma})^{m-1}, \ m \ge 2,$$
(85)

where

$$\gamma = \min\{\gamma_1, \gamma_2\},\tag{86}$$

 $\hat{c} = \hat{c}(V)$ and k is sufficiently large: $k > k_0(V, \gamma_2, \delta)$.

Corollary 17. The sequence $\{W_m^+\}_{m=0}^{\infty}$ converges to a continuous and periodic potential W^+ with respect to the norm $\|\cdot\|_*$. The following estimate holds:

$$\|W^{+} - W_{m}^{+}\|_{*} < 2(\hat{c}k^{-\gamma})^{m+1}.$$
(87)

Proof. We use an induction. For the first step m = 1, we need to show (84) and

$$\|W_1^+ - W_0^+\|_* \le \hat{c}k^{-\gamma}.$$
(88)

We start with (84). This is a perturbative formula for a linear operator. It is given by (54) for m = 0 up to the notations in Definition 4. To prove (88) we consider two functions $u_{-1}^+(\vec{x})$ and $u_0^+(\vec{x})$, see Definition 5. Next, by (72) – (74),

$$\|W_1^+ - W_0^+\|_* = \left\||\sigma||u_0^+|^2\right\|_* \tag{89}$$

A. DUAIBES, YU. KARPESHINA

$$\begin{split} &\leq |\sigma| \left(|||u_{0}^{+}|^{2} - |u_{-1}^{+}|^{2}||_{*} + ||u_{-1}^{+}||_{*}^{2} \right) \\ &= |\sigma| \left(|||\psi_{0}^{+}|^{2} - |\psi_{-1}^{+}|^{2}||_{*} + ||\psi_{-1}^{+}||_{*}^{2} \right) \\ &\leq |\sigma| \left(|||\psi_{0}^{+}|^{2} - |\psi_{-1}^{+}|^{2} + 2i\Im(\bar{\psi}_{-1}^{+}\psi_{0}^{+})||_{*} + ||\psi_{-1}^{+}||_{*}^{2} \right) \\ &= |\sigma| \left(||(\psi_{0}^{+} - \psi_{-1}^{+})(\bar{\psi}_{0}^{+} + \bar{\psi}_{-1}^{+})||_{*} + ||\psi_{-1}^{+}||_{*}^{2} \right) \\ &\leq |\sigma| \left(||\psi_{0}^{+} - \psi_{-1}^{+}||_{*} ||\bar{\psi}_{0}^{+} + \bar{\psi}_{-1}^{+}||_{*} + ||\psi_{-1}^{+}||_{*}^{2} \right). \end{split}$$

By (54) for m = 0,

$$\|\psi_0^+ - \psi_{-1}^+\|_* \le |A| \|\hat{E}^{+q}(\vec{t}) - \hat{E}^+(\vec{t})\|_0 \le 2|A|k^{-\gamma_1},\tag{90}$$

Next, we notice that $\|\bar{\psi}_{-1}^+\|_* = \|\psi_{-1}^+\|_*$ and $|\hat{E}_{jj}^+| \leq 1$ and $|\hat{E}_{j-q,j}^+| \leq 1$. This implies that $\|\psi_{-1}^+\|_* \leq 2|A|$ because the only two nonzero elements of $\hat{E}^+(\vec{t})$ are \hat{E}_{jj}^+ and $\hat{E}_{j-q,j}^+$. Hence, by (90),

$$\|\bar{\psi}_{0}^{+} + \bar{\psi}_{-1}^{+}\|_{*} \leq \|\bar{\psi}_{0}^{+} - \bar{\psi}_{-1}^{+}\|_{*} + 2\|\bar{\psi}_{-1}^{+}\|_{*} \leq 2|A|k^{-\gamma_{1}} + 4|A|.$$

Since k is chosen to be sufficiently large, we can write

$$|\bar{\psi}_0^+ + \bar{\psi}_{-1}^+||_* \le 5|A|. \tag{91}$$

Considering the last line of (89) and using (90), (91) and the condition $|\sigma A|^2 < k^{-\gamma_2}$, we obtain that

$$\| \|W_1^+ - W_0^+\|_* \le |\sigma| \left(\left(2|A|k^{-\gamma_1} \right) 5|A| + 2|A|^2 \right) \le 10|\sigma| |A|^2 k^{-\gamma_1} + 2|\sigma| |A|^2$$
(92)
$$\le 3k^{-2l\gamma_2}.$$
(93)

for sufficiently large $k > k_0(||V||_*, d, \gamma_1)$. This proves (82) and (83) for m = 1.

Now, let us assume that (82) -(85) are satisfied for all $s = 1, 2, ..., m - 1, m \ge 2$, i.e.,

$$||W_s^+ - W_{s-1}^+||_* \le (\hat{c}k^{-\gamma})^s, \tag{94}$$

$$||W_s^+ - V||_* \le \sum_{r=1}^s (\hat{c}k^{-\gamma})^r, \tag{95}$$

$$\|\hat{E}_{s-1}^{+}(\vec{t}) - \hat{E}_{s-2}^{+}(\vec{t})\|_{0} \le \hat{c}(\hat{c}k^{-\gamma})^{s-1}.$$
(96)

First, let us prove (85). Considering that $H_0 + W_{m-1}^+ = \hat{H}_q + (W_{m-1}^+ - P_q V P_q)$ we define: $\hat{P}^+_{m-1} = (\hat{H}_1(\vec{x}) - \hat{v})^{-\frac{1}{2}} (W_{m-1}^+ - P_q V P_q) (\hat{H}_1(\vec{t}) - \hat{v})^{-\frac{1}{2}}$ (07)

$$B_{m-1}^+(z) = (H_q(\bar{t}) - z)^{-\frac{1}{2}} \left(W_{m-1}^+ - P_q V P_q \right) (H_q(\bar{t}) - z)^{-\frac{1}{2}}.$$
 (97) is quite obvious that

$$\|(I - P_q)(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}\|_0 = \|(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}(I - P_q)\|_0 \le k^{-\gamma_1};$$
(98)

$$\|P_q(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}\|_0 = \|(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}P_q\|_0 \le \frac{1}{\sqrt{d}};$$
(99)

here $z \in C_1^+$, d is the radius of C_1^+ , $d = \frac{1}{10}|v_q|$. Considering two above estimates and $W_{m-1}^+ - P_q V P_q = \hat{W}_{m-1}^+ + P_q (W_{m-1}^+ - V) P_q$, here $\hat{W}_m^+ = W_m^+ - P_q W_m^+ P_q$, (100)

 It

we obtain:

$$\max_{z \in C_1} \|\hat{B}_{m-1}^+(z)\|_0 \le (1 + \frac{2}{\sqrt{d}}) \|\hat{W}_{m-1}^+\|_0 k^{-\gamma_1} + d^{-1} \|P_q(W_{m-1}^+ - V)P_q\|_0.$$
(101)

Hence, by induction assumption (95),

$$\max_{z \in C_1} \|\hat{B}_{m-1}^+(z)\|_0 \le \beta k^{-\gamma}.$$
(102)

where $\beta(V) = 3(1 + \frac{2}{\sqrt{d}}) ||V|| + 2\hat{c}d^{-1}$. This means that

$$\|\hat{G}_{m-1,r}^{+}(k,\vec{t})\|_{0} \le \left(c\beta k^{-\gamma}\right)^{r},\tag{103}$$

where $\hat{G}^+_{m-1,r}(k,\vec{t})$ is given by (45) with $W^+_{m-1} - P_q V P_q$ instead of \hat{V} and c is an absolute constant. Clearly,

$$\|\hat{E}_{m-1}^{+}(\vec{t})\|_{0} \leq |\hat{E}^{+}(\vec{t})_{jj}| + |\hat{E}^{+}(\vec{t})_{j-q,j}| + \sum_{r=1}^{\infty} \|\hat{G}_{m-1,r}^{+}(k,\vec{t})\|_{0}$$

Since $|\hat{E}^+(\vec{t})_{jj}|, |\hat{E}^+_{m-1}(\vec{t})| \le 1$ and by (103), we get $||\hat{E}^+_{m-1}(\vec{t})||_0 \le 3$. Now, we note that for $m \ge 2$:

$$\begin{aligned} \|\hat{E}_{m-1}^{+}(\vec{t}) - \hat{E}_{m-2}^{+}(\vec{t})\|_{0} &\leq \sum_{r=1}^{\infty} \|\hat{G}_{m-1,r}^{+}(k,\vec{t}) - \hat{G}_{m-2,r}^{+}(k,\vec{t})\|_{0} \\ &\leq \frac{1}{2\pi} \sum_{r=1}^{\infty} \|\oint_{C_{1}^{+}} \left(\hat{H}_{q}(\vec{t}) - z\right)^{-\frac{1}{2}} \left[\hat{B}_{m-1}^{+}(z)^{r} - \hat{B}_{m-2}^{+}(z)^{r}\right] \left(\hat{H}_{q}(\vec{t}) - z\right)^{-\frac{1}{2}} dz\|_{0}, \\ &\leq \left(\frac{1}{2\pi} \int_{C_{1}^{+}} \|\left(\hat{H}_{q}(\vec{t}) - z\right)^{-1/2} \|_{0}^{2} ds\right) \left(\sum_{r=1}^{\infty} \max_{z \in C_{1}^{+}} \|\hat{B}_{m-1}^{+}(z)^{r} - \hat{B}_{m-2}^{+}(z)^{r}\|_{0}\right) \\ &\leq c \sum_{r=1}^{\infty} \max_{z \in C_{1}^{+}} \|\left(\hat{B}_{m-1}^{+}(z)^{r} - \hat{B}_{m-2}^{+}(z)^{r}\|_{0}\right). \end{aligned}$$
(104)

Consider $\|\hat{B}_{m-1}^+(z) - \hat{B}_{m-2}^+(z)\|_0$ for any $z \in C_1^+$. Arguing as in the proof of the estimate for $\|\hat{B}_{m-1}^+(z)\|_0$, see (101), (102), we obtain:

$$\|\hat{B}_{m-1}^{+}(z) - \hat{B}_{m-2}^{+}(z)\|_{0} \leq \beta \|W_{m-1}^{+} - W_{m-2}^{+}\|_{0}, \quad m \geq 2.$$
(105)

On the other hand, the right-hand side of (104), which we denote here by R for simplicity, can be estimated by using (94), (102), and (105):

$$R \leq c \sum_{r=1}^{\infty} \max_{z \in C_{1}^{+}} \left[\| \hat{B}_{m-1}^{+}(z) - \hat{B}_{m-2}^{+}(z) \|_{0} \left(\| \hat{B}_{m-1}^{+}(z) \|_{0} + \| \hat{B}_{m-2}^{+}(z) \|_{0} \right)^{r-1} \right]$$

$$\leq c \sum_{r=1}^{\infty} \beta \| W_{m-1}^{+} - W_{m-2}^{+} \|_{0} \left(2\beta k^{-\gamma} \right)^{r-1} \leq c\beta \| W_{m-1}^{+} - W_{m-2}^{+} \|_{0}$$

$$\leq c\beta (\hat{c}k^{-\gamma})^{m-1}.$$
(106)

Thus, (85) holds. Next, we consider $||W_m^+ - W_{m-1}^+||_0$. Obviously,

$$\|\psi_s^+\|_* \le |A| \|\hat{E}_s^+(\vec{t})\|_0.$$
(107)

Using (72), we easily get

$$\|W_{m}^{+} - W_{m-1}^{+}\|_{0} = \|\sigma |u_{m-1}^{+}|^{2} - \sigma |u_{m-2}^{+}|^{2}\|_{*} \le |\sigma| \|\psi_{m-1}^{+} - \psi_{m-2}^{+}\|_{*} \|\bar{\psi}_{m-1}^{+} + \bar{\psi}_{m-2}^{+}\|_{*}.$$
(108)

Now, formula (108) can be rewritten using (80) and (107):

$$\begin{split} \|W_{m}^{+} - W_{m-1}^{+}\|_{0} \leq |\sigma||A|^{2} \|\hat{E}_{m-1}^{+}(\vec{t}) - \hat{E}_{m-2}^{+}(\vec{t})\|_{0} \left(\|\hat{E}_{m-1}^{+}(\vec{t})\|_{0} + \|\hat{E}_{m-2}^{+}(\vec{t})\|_{0}\right) \\ \leq (\hat{c}k^{-\gamma})^{m}, \quad m \geq 2. \end{split}$$
(109)

Therefore, (82) and (83) hold.

3.2. Solution to Nonlinear Polyharmonic Equation with Periodic Potential for 2l > n. In this section we show that convergence of $\{W_m^+\}_{m=0}^{\infty}$ leads to convergence of the sequence of the spectral projections $\{\hat{E}_m^+(\vec{t})\}_{m=0}^{\infty}$ to that of the operator $H_0(\vec{t}) + W^+$ (in the norm $\|\cdot\|_0$). The sequence of the corresponding eigenvalues $\{\hat{\lambda}_m^+(\vec{t})\}_{m=0}^{\infty}$ converges to the corresponding eigenvalue of $H_0(\vec{t}) + W^+$.

Lemma 18. Let $\gamma_2 > 0$, $0 < 9\delta < 2l - n$. Suppose that \vec{t} belong to the $(k^{-n+1-7\delta})$ -neighborhood in K of the set $\chi_q(k, \delta)$. Then, for every sufficiently large $k > k_0(V, \gamma_2, \delta)$ and every $A \in \mathbb{C}$: $|\sigma||A|^2 < k^{-\gamma_2}$, the sequence $\{\hat{E}_m^+(\vec{t})\}_{m=0}^{\infty}$ converges with respect to the norm $\|\cdot\|_0$ to a one-dimensional spectral projection $\hat{E}_W^+(\vec{t})$ of $H_0(\vec{t}) + W^+$,

$$\|\hat{E}_m^+(\vec{t}) - \hat{E}_W^+(\vec{t})\|_0 \le \hat{c}(\hat{c}k^{-\gamma})^{m+1}, \ m = 0, 1, \dots$$
(110)

The projection $\hat{E}^+_W(\vec{t})$ is given as

$$\hat{E}_{W}^{+}(\vec{t}) = \hat{E}^{+}(\vec{t}) + \sum_{r=1}^{\infty} \hat{G}_{W,r}^{+}(k,\vec{t}), \qquad (111)$$

where. $\hat{G}^+_{W,r}(k,\vec{t})$ is given by (45) with with $W^+ - P_q V P_q$ instead of \hat{V} . The following estimate holds:

$$\|\hat{G}_{W,r}^{+}(k,\vec{t})\|_{0} \leq \hat{c} \left(\hat{c}k^{-\gamma}\right)^{r}.$$
(112)

Proof. Let $B_W^+(z)$ be defined by (97) with W^+ instead of W_{m-1}^+ . Next, we estimate

$$\|\hat{B}_m^+(z) - \hat{B}_W^+(z)\|_0 = \|(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}(W_m^+ - W^+)(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}\|_0$$

By (98), (99)

$$\|\hat{B}_{m}^{+}(z) - \hat{B}_{W}^{+}(z)\|_{0} \le \beta \|W_{m}^{+} - W^{+}\|_{*}.$$
(113)

Using Corollary 17, we get

$$\|\hat{B}_{m}^{+}(z) - \hat{B}_{W}^{+}(z)\|_{0} \le \hat{c}(\hat{c}k^{-\gamma})^{m+1}.$$
(114)

RESONANT SOLUTIONS OF THE NON-LINEAR SCHRÖDINGER EQUATION WITH PERIODIC POTENTIAL By (102),

$$\max_{z \in C_1} \|\hat{B}^+_W(z)\|_0 \le \beta k^{-\gamma}.$$
(115)

Next, we prove (110). Let us write

$$\begin{aligned} \left\| \hat{E}_{m}^{+}(\vec{t}) - \hat{E}_{W}^{+}(\vec{t}) \right\|_{0} &\leq \sum_{r=1}^{\infty} \left\| \hat{G}_{m,r}^{+}(k,\vec{t}) - \hat{G}_{W,r}^{+}(k,\vec{t}) \right\|_{0} \tag{116} \\ &\leq c \sum_{r=1}^{\infty} \max_{z \in C_{1}^{+}} \left\| \hat{B}_{m}^{+}(z)^{r} - \hat{B}_{W}^{+}(z)^{r} \right\|_{0} \\ &\leq c \sum_{r=1}^{\infty} \max_{z \in C_{1}^{+}} \left\| \|\hat{B}_{m}^{+}(z) - B_{W}^{+}(z) \|_{0} \left(\|\hat{B}_{m}^{+}(z)\|_{0} + \|B_{W}^{+}(z)\|_{0} \right)^{r-1} \right]. \end{aligned}$$

By (102) and (113),

$$\begin{aligned} \|\hat{E}_{m}^{+}(\vec{t}) - \hat{E}_{W}^{+}(\vec{t})\|_{0} &\leq \sum_{r=1}^{\infty} \left[d^{-1} \|W_{m}^{+} - W^{+}\|_{*} \left(2\beta k^{-\gamma} \right)^{r-1} \right] \\ &\leq \hat{c} \|W_{m}^{+} - W^{+}\|_{*}. \end{aligned}$$

Hence, by (87) we get (110). Since $||W_m^+ - W^+||_* \to 0$ as $m \to \infty$, we have that $\hat{E}_W^+(\vec{t})$ is the limit of $\hat{E}_m^+(\vec{t})$. Formulas above also prove that $\hat{G}_{m,r}^+(k,\vec{t})$ converges to $\hat{G}_{W,r}^+(k,\vec{t})$ in the norm $\|\cdot\|_0$. Now, it remains to prove (112). Indeed,

$$\begin{aligned} \|\hat{G}_{W,r}^{+}\|_{0} &= \left\|\frac{(-1)^{r+1}}{2\pi i} \oint_{C_{1}^{+}} \left(\hat{H}_{q}(\vec{t}) - z\right)^{-\frac{1}{2}} \left(\hat{B}_{W}^{+}(z)\right)^{r} \left(\hat{H}_{q}(\vec{t}) - z\right)^{-\frac{1}{2}} dz \right\|_{0} \\ &\leq \frac{1}{2\pi} \max_{z \in C_{1}^{+}} \|\hat{B}_{W}^{+}(z)\|_{0}^{r} \int_{C_{1}^{+}} \left\|\left(\hat{H}_{q}(\vec{t}) - z\right)^{-1}\right\|_{0} ds \leq (\hat{c}k^{-\gamma})^{r}. \end{aligned}$$

Let u_W^+, ψ_W^+ be defined by Definition 2 for $W = W^+$.

Lemma 19. Under the assumptions of Lemma 18 and, for every sufficiently large $k > k_0(V, \gamma_2, \delta)$ and every $A \in \mathbb{C} : |\sigma||A|^2 < k^{-\gamma_2}$, the function $\psi_W^+(\vec{x})$ is the limit of the sequence $\psi_m^+(\vec{x})$ in the norm $\|\cdot\|_*$. Moreover,

$$\|\psi_m^+ - \psi_W^+\|_* \le (\hat{c}k^{-\gamma})^{m+1}, \quad m = 0, 1, \dots$$
(117)

Proof. Using (110), we get

$$\begin{aligned} \|\psi_m^+ - \psi_W^+\|_* &\leq |A| \|\hat{E}_m^+(\vec{t}) - \hat{E}_W^+(\vec{t})\|_0 \\ &\leq |A| (\hat{c}k^{-\gamma})^{m+1}. \end{aligned}$$
(118)

Corollary 20. The sequence u_m^+ converges to \hat{u}_W^+ in C(Q).

Corollary 21. The operator $\hat{\mathcal{M}}^+$ maps the operator W^+ into W^+ . In other words, $\hat{\mathcal{M}}^+W^+ = W^+$.

Proof. We know that $W_m^+ \to W^+$ with respect to the norm $\|\cdot\|_*$. We also know that $W_{m+1}^+ = \hat{\mathcal{M}} W_m^+$ by the equation (74). It follows that $\hat{\mathcal{M}}^+ W_m^+ \to W^+$ as $m \to \infty$. On the other hand,

$$\|\hat{\mathcal{M}}^{+}W_{m}^{+} - \hat{\mathcal{M}}^{+}W^{+}\|_{*} \leq |\sigma|\|\psi_{m}^{+} - \psi^{+}\|_{*}\|\bar{\psi}_{m}^{+} + \bar{\psi}^{+}\|_{*}.$$

This implies, by (117), that $\hat{\mathcal{M}}^+ W_m^+ \to \mathcal{M}^+ W^+$ in the norm $\|\cdot\|_*$. Therefore, $\hat{\mathcal{M}}^+ W^+ = W^+$.

Remark 22. Note that W^+ depends on V, σ , A, \vec{t} .

We use the notations $\hat{\lambda}_m^+(\vec{t})$ and $\hat{\lambda}_W^+(\vec{t})$ for the eigenvalues corresponding to $\hat{E}_m^+(\vec{t})$ and $\hat{E}_W^+(\vec{t})$, respectively.

Lemma 23. Under assumptions of Lemma 18 and, for every sufficiently large $k > k_0(V, \gamma_2, \delta)$ and every $A \in \mathbb{C} : |\sigma||A|^2 < k^{-\gamma_2}$, the sequence $\hat{\lambda}_m^+(\vec{t})$ converges to $\hat{\lambda}_W^+(\vec{t})$. The limit $\hat{\lambda}_W^+(\vec{t})$ is defined by

$$\hat{\lambda}_{W}^{+}(\vec{t}) = \hat{\lambda}^{+}(\vec{t}) + \sum_{r=1}^{\infty} \hat{g}_{W,r}^{+}(k,\vec{t}).$$
(119)

Moreover,

$$|\hat{g}^{+}_{W,r}(k,\vec{t})| < \hat{c}(\hat{c}k^{-\gamma})^{r}, \qquad (120)$$

where $g_{W,r}^+$ is given by (44) with $W^+ - P_q V P_q$ instead of \hat{V} :

Proof. First, we prove (120). For simplicity, we denote here $\hat{E}^+(\vec{t})$ by E_0 and $E_1 = I - E_0$. Note that, for any r,

$$\oint_{C_1^+} \left(E_1 \hat{B}_m^+(z) E_1 \right)^r dz = 0$$

since $E_1 \hat{B}_m^+(z) E_1$ is holomorphic inside C_1^+ . Consider

$$\mathcal{F}_{m,r} = \oint_{C_1^+} (\hat{B}_m^+(z))^r dz$$

Then,

$$\begin{aligned} \mathcal{F}_{m,r} &= \oint_{C_1^+} \left[(\hat{B}_m^+(z))^r - (E_1 \hat{B}_m^+(z) E_1)^r \right] (dz) \\ &= \oint_{C_1^+} \left[\left((E_0 + E_1) \hat{B}_m^+(z) (E_0 + E_1) \right)^r - (E_1 \hat{B}_m^+(z) E_1)^r \right] dz \\ &= \sum \oint_{C_1^+} E_{i_1} \hat{B}_m^+(z) E_{i_2} \dots E_{i_r} \hat{B}_m^+(z) E_{i_{r+1}} dz, \end{aligned}$$

where the sum is taken over $i_1, ..., i_{r+1} = 0, 1$, and $\exists s$ such that $E_{i_s} = E_0$. Let $E_{i_{r_o}}$ be the first one to be equal to E_0 . Since E_0 belongs to the trace class, it is obvious that $E_{i_1}\hat{B}_m^+(z)E_{i_2}...E_{i_r}\hat{B}_m^+(z)E_{i_{r+1}}$ is also in the trace class. Hence, we get by (102)

$$\|E_{i_1}\hat{B}_m^+(z)E_{i_2}\dots E_{i_r}\hat{B}_m^+(z)E_{i_{r+1}}\|_1 \le \|\hat{B}_m^+(z)\|_0^{r_0}\|\hat{B}_m^+(z)\|_0^{r-r_0} \le (\beta k^{-\gamma})^r.$$
(121)

Thus,

$$\|\mathcal{F}_{m,r}\|_{1} = \left\|\sum \oint_{C_{1}^{+}} E_{i_{1}}\hat{B}_{m}^{+}(z)E_{i_{2}}...E_{i_{r}}\hat{B}_{m}^{+}(z)E_{i_{r+1}}dz\right\|_{1}$$

$$\leq \max_{z \in C_{1}^{+}} \|E_{i_{1}}\hat{B}_{m}^{+}(z)E_{i_{2}}...E_{i_{r}}\hat{B}_{m}^{+}(z)E_{i_{r+1}}\|_{1}\int_{C_{1}^{+}} ds \leq 2\pi(\beta k^{-\gamma})^{r} \qquad (122)$$

and $\mathcal{F}_{m,r}$ converges to

$$\mathcal{F}_r = \oint_{C_1^+} B_W^+(z)^r dz \tag{123}$$

in the trace class. Let $\hat{g}_{m,r}^+$ be given by (44) with $W_m^+ - P_q V P_q$ instead of \hat{V} . Since $\hat{g}_{m,r}(k,\vec{t}) = \frac{(-1)^r}{2\pi i r} \operatorname{Tr} \mathcal{F}_{m,r}$, it follows that $\hat{g}_{m,r}(k,\vec{t})$ converges to $\hat{g}_{W,r}(k,\vec{t})$. Then,

$$\hat{g}_{W,r}(k,\vec{t})| = \left|\frac{(-1)^r}{2\pi i r} Tr \mathcal{F}_r\right| \le \frac{r^{-1}}{2\pi} \cdot 2\pi (\beta \|V\|_* k^{-\gamma})^r \le \hat{c}(\hat{c}\|V\|_* k^{-\gamma})^r.$$
(124)

This proves (120). Now, it is clear that $\lambda_W^+(\vec{t})$ is the limit of $\{\hat{\lambda}_m^+(\vec{t})\}_{m=0}^{\infty}$ since

$$\left|\hat{\lambda}_{m}^{+}(\vec{t}) - \hat{\lambda}_{W}^{+}(\vec{t})\right| \leq \sum_{r=1}^{\infty} \left|\hat{g}_{m,r}(k,\vec{t}) - g_{W,r}(k,\vec{t})\right|.$$

The following theorem is the main result of the paper for the case 2l > n.

Theorem 24. Let $\gamma_2 > 0$, $0 < 9\delta < 2l - n$. Then, for each sufficiently large k: $k > k_0(V, \gamma_2, \delta)$, the following holds. If \vec{t} belongs to the $(k^{-n+1-7\delta})$ -neighborhood in K of the resonant set $\chi_q(k, \delta)$ and $A \in \mathbb{C} : |\sigma||A|^2 < k^{-\gamma_2}$, then, there is a pair of functions $u^{\pm}(\vec{x}, \vec{t})$ and the corresponding real values $\lambda^{\pm}(\vec{t})$, solving the equation (2) with the boundary conditions (3). Moreover, the following is true as $k \to \infty$:

$$u^{\pm}(\vec{x}, \vec{t}) = A e^{i \langle k, \vec{x} \rangle} \left(\psi^{\pm}_{-1}(\vec{x}, \vec{t}) + \phi^{\pm}(\vec{x}, \vec{t}) \right), \tag{125}$$

$$\lambda^{\pm}(\vec{t}) = \hat{\lambda}^{\pm}(\vec{t}) + O\left(k^{-\gamma}\right), \qquad (126)$$

where ψ_{-1}^{\pm} is given by (77) and Definition 1 (see also (68), (69)) and $\phi(\vec{x}, \vec{t})$ is periodic in \vec{x} , and satisfies:

$$\|\phi\|_* \le \hat{c}k^{-\gamma},\tag{127}$$

 $\gamma = \min\{\gamma_1, \gamma_2\}, \ 2\gamma_1 = 2l - n - 8\delta.$

From now on we set

$$u^{\pm}(\vec{x}, \vec{t}) = u^{\pm}_{W}(\vec{x}, \vec{t}), \quad \lambda^{\pm}(\vec{t}) = \hat{\lambda}^{\pm}_{W}(\vec{t}), \quad E^{\pm} = \hat{E}^{\pm}_{W}.$$
 (128)

Proof. We note that the function u_W^{\pm} defined by (70) and the value $\lambda_W^{\pm}(\vec{t})$ given in Lemma 23 by formula (119) solve the equation:

$$(H_0(\vec{t}) + W^{\pm}) u_W^{\pm}(\vec{x}, \vec{t}) = \hat{\lambda}_W^{\pm}(\vec{t}) u_W^{\pm}(\vec{x}, \vec{t}), \ \vec{x} \in Q, \ W^{\pm} = W^{\pm}(\vec{x}, \vec{t}, A),$$
(129)

and satisfying the boundary conditions (3). Using Corollary 21, we can rewrite equation (129) as (2). Considering (128) we finish the proof.

3.3. The Differentiability of the Eigenvalue and Its Spectral Projection.

Lemma 25. Under conditions of Lemma 16

$$\|\nabla_{\vec{t}}(W_m^{\pm} - W_{m-1}^{\pm})\|_* \le k^{2l-1} (\hat{c}k^{-\gamma})^m, \tag{130}$$

$$\|\nabla_{\vec{t}}(W_m^{\pm} - V)\|_* \le k^{2l-1} \sum_{r=1}^m (\hat{c}k^{-\gamma})^r,$$
(131)

$$\|\nabla_{\vec{t}} \left(\hat{E}_0^{\pm}(\vec{t}) - \hat{E}_{-1}^{\pm}(\vec{t}) \right) \|_0 \le \hat{c} k^{n-1+7\delta-\gamma}, \tag{132}$$

$$\|\nabla_{\vec{t}} \left(\hat{E}_{m-1}^{\pm}(\vec{t}) - \hat{E}_{m-2}^{\pm}(\vec{t}) \right)\|_{0} \le k^{2l-1} (\hat{c}k^{-\gamma})^{m-1}, \ m \ge 2.$$
(133)

Proof. The inequality (132) follows from the linear case inequality (54). To obtain the estimates (130), (131) and (133), we use the obvious inequality:

$$\left\| \nabla_{\vec{t}} \left(\hat{H}_q(\vec{t}) - z \right)^{-1/2} \right\|_0 < ck^{2l-1} d^{-3/2}, \quad z \in C_1^{\pm}$$

Further considerations are analogous to that in Lemma 16.

Lemma 26. Under conditions of Lemma 18 the sequences $\{\nabla_{\vec{t}} \hat{E}_m^{\pm}(\vec{t})\}_{m=0}^{\infty}$ converge to $\nabla_{\vec{t}} E^{\pm}(\vec{t})$ in $\|\cdot\|_0$ and

$$\left\|\nabla_{\vec{t}} \left(\hat{E}_m^{\pm}(\vec{t}) - E^{\pm}(\vec{t})\right)\right\|_0 \le k^{2l-1} (\hat{c}k^{-\gamma})^{m+1}.$$
(134)

The operator $\nabla_{\vec{t}} E^{\pm}(\vec{t})$ is given as

$$\nabla_{\vec{t}} E^{\pm}(\vec{t}) = \nabla_{\vec{t}} \hat{E}^{\pm}(\vec{t}) + \sum_{r=1}^{\infty} \nabla_{\vec{t}} \hat{G}^{\pm}_{W,r}(k,\vec{t}),$$
(135)

where

$$\|\nabla_{\vec{t}} \hat{G}^{\pm}_{W,r}(k,\vec{t})\|_{0} \le k^{2l-1} \left(\hat{c}k^{-\gamma}\right)^{r}.$$
(136)

The sequences $\nabla_{\vec{t}} \hat{\lambda}_m^{\pm}(\vec{t})$ converge to $\nabla_{\vec{t}} \lambda^{\pm}(\vec{t})$. The limits $\nabla_{\vec{t}} \lambda^{\pm}(\vec{t})$ are defined by

$$\nabla_{\vec{t}}\lambda^{\pm}(\vec{t}) = \nabla_{\vec{t}}\hat{\lambda}^{\pm}(\vec{t}) + \sum_{r=1}^{\infty} \nabla_{\vec{t}}\hat{g}^{\pm}_{W,r}(k,\vec{t}), \qquad (137)$$

where

$$|\nabla_{\vec{t}} \hat{g}^{\pm}_{W,r}(k,\vec{t})| < k^{2l-1} (\hat{c}k^{-\gamma})^r.$$
(138)

Proof. The proof is analogous to those of Lemma 18 and 23 up to differentiation, Lemma 25 being used.

The next theorem easily follows from the previous lemma.

Theorem 27. Under conditions of Theorem 24:

$$\|\nabla_{\vec{t}} \left(E^{\pm}(\vec{t}) - \hat{E}^{\pm}(\vec{t}) \right) \|_{0} < C(V) k^{2l-1-\gamma}, \tag{139}$$

$$\left|\nabla_{\vec{t}} \left(\lambda^{\pm}(\vec{t}) - \hat{\lambda}^{\pm}(\vec{t})\right)\right| < C(V)k^{2l-1-\gamma}.$$
(140)

Corollary 28.

$$\left|\nabla_{\vec{t}}\lambda^{\pm}(\vec{t})\right| = 2lk^{2l-1} \left(1 + O(k^{-\gamma}) + O(k^{-1})\right).$$
(141)

The corollary follows from the theorem and the obvious relation $\nabla \hat{\lambda}^{\pm}(\vec{t}) = 2lk^{2l-1}(1+O(k^{-1})).$

Let us consider the surface $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$ for a fixed λ_0 , $\lambda_0 > k_0^{2l}(V, \gamma_2, \delta)$. Note that the parts $\hat{\lambda}^+(\vec{t}) = \lambda_0$ and $\hat{\lambda}^-(\vec{t}) = \lambda_0$ do not intersect, since $v_q \neq 0$. Thus the deviation of the surface $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$ from the unperturbed one (V=0) is essential. The next theorem follows.

Theorem 29. If $\lambda_0 > k_0^{2l}(V, \gamma_2, \delta)$, then the surface $\lambda^{\pm}(\vec{t}) = \lambda_0$ is in the $C(V)\lambda_0^{-\hat{\gamma}}$ -neighborhood of $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$ for every sufficiently large λ_0 , here $\hat{\gamma} = (2l - 1 + \gamma)(2l)^{-1}$.

Corollary 30. The surfaces $\lambda^+(\vec{t}) = \lambda_0$ and $\lambda^-(\vec{t}) = \lambda_0$ do not intersect and located at the distance greater than $|v_q| + O\left(\lambda_0^{-\hat{\gamma}}\right)$ from each other as $\lambda_0 \to \infty$.

4. Solutions of Nonlinear Scrödinger Equation in Dimension Two.

In this section, we present resonant solutions of (2), (3) for n = 2, l = 1. The equation is

$$-\Delta u + Vu + \sigma |u|^2 u = \lambda u. \tag{142}$$

The proof of the result is analogous to that for 2l > n. Indeed, let $S(k, \epsilon)$ be given by (15) up to replacing of $4k^{-n+2-\delta}$ by ϵ :

$$S_q(k,\epsilon) = \{ \vec{x} \in S(k) : \left| |\vec{x}|^2 - |\vec{x} - \vec{P}_q(0)|^2 \right| < \epsilon \}.$$
(143)

We first state the geometric lemma.

Lemma 31. Let $0 < \epsilon < \varepsilon_0$. Then, for sufficiently largek, $k > k_1(q, \varepsilon_0)$, there exists a resonant set $\chi_q(k, \epsilon) \subset \mathcal{K}S_q(k, \epsilon)$ such that, for any $\vec{t} \in \chi_q(k, \epsilon)$, the followings hold:

1. there exists a unique $j \in \mathbb{Z}^n$ such that $|\vec{P}_j(\vec{t})| = k$, (144)

2.
$$|p_j^2(\vec{t}) - p_{j-q}^2(\vec{t})| < \epsilon,$$
 (145)

3.
$$\min_{m \neq j, j-q} |p_j^2(\vec{t}) - p_m^2(\vec{t})| > 2\epsilon^6.$$
(146)

Moreover, for any \vec{t} in the $(k^{-1}\epsilon^7)$ -neighborhood of $\chi_q(k,\epsilon)$ in \mathbb{C}^2 , there exists a unique $j \in \mathbb{Z}^2$ such that $|p_j^2(\vec{t}) - k^2| < 5\epsilon^7$ and the second and third conditions above are satisfied.

The set $\chi_q(k,\epsilon)$ has an asymptotically full measure on $\mathcal{K}S_q(k,\epsilon)$ as $\epsilon \to 0$, that is

$$\frac{s(\mathcal{K}S_q(k,\epsilon) \setminus \chi_q(k,\epsilon))}{s(\mathcal{K}S_q(k,\epsilon))} < c\epsilon, \quad c \neq c(k).$$
(147)

For all of the followings we assume that

$$||V||_* < \epsilon^9, \quad |v_q| > \epsilon^{10}, \quad |\sigma||A^2| < \epsilon^{11}.$$
 (148)

Corollary 32. If \vec{t} belongs to the $(\frac{1}{8}k^{-1}\varepsilon^{10})$ - neighborhood of $\chi_q(k,\epsilon)$, then for all z on the circle

$$C_1^+ = \{z : |z - \hat{\lambda}^+(\vec{t})| = d\}, \quad d = \frac{1}{3}\epsilon^{10},$$

both of the following inequalities are true:

$$2|p_m^2(\vec{t}) - z| \ge \epsilon^6, \ m \ne j, j - q,$$
(149)

$$|\hat{\lambda}^{\pm}(\vec{t}) - z| \ge \frac{1}{12} \epsilon^{10},$$
 (150)

 $\hat{\lambda}^{\pm}$ being the eigenvalues of (13).

Now, consider the map $\hat{\mathcal{M}}$ defined by (72). The following lemma can be proved by analogy with Lemma 16.

Lemma 33. There is ϵ_0 , $0 < \epsilon_0 \neq \epsilon_0(\lambda)$, such that for any $0 < \epsilon < \epsilon_0$ under the conditions (148) and for any sufficiently large $k: k > k_1(V, \epsilon_0)$, the following holds. Let \vec{t} belong to the $(\frac{1}{8}k^{-1}\varepsilon^{10})$ -neighborhood in K of the set $\chi_q(k, \epsilon)$. Then, for any m = 1, 2, ...:

$$\|W_m^+ - W_{m-1}^+\|_* \le \epsilon^{11} (c\epsilon)^{m-1}, \tag{151}$$

$$\|W_m^+ - V\|_* \le \epsilon^{11} \sum_{r=1}^m (c\epsilon)^{r-1}, \tag{152}$$

$$\|\hat{E}_{0}^{+}(\vec{t}) - \hat{E}_{-1}^{+}(\vec{t})\|_{0} \le c\epsilon,$$
(153)

$$\|\hat{E}_{m-1}^{+}(\vec{t}) - \hat{E}_{m-2}^{+}(\vec{t})\|_{0} \le (c\epsilon)^{m-1},$$
(154)

where c is an absolute constant.

Corollary 34. The sequence $\{W_m^+\}_{m=0}^{\infty}$ converges to a continuous and periodic function W with respect to the norm $\|\cdot\|_*$. The following estimate holds:

$$\|W^{+} - W_{m}^{+}\|_{*} \le 2\epsilon^{11} (c\epsilon)^{m}.$$
(155)

Proof. The following facts can be easily checked:

$$1 \cdot \|(I - P_q)(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}\|_0 = \|(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}(I - P_q)\|_0 < 2\epsilon^{-3},$$
(156)

$$2.\|P_q(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}\|_0 = \|(\hat{H}_q(\vec{t}) - z)^{-\frac{1}{2}}P_q\|_0 < c\epsilon^{-5},$$
(157)

 $P_q,\,\hat{H}_q$ being defined by (38), (39). Further we use an induction. For the first step we need to show that

$$\|\hat{E}_{0}^{+}(\vec{t}) - \hat{E}_{-1}^{+}(\vec{t})\|_{0} \le c\epsilon,$$
(158)

22

$$\|W_1^+ - W_0^+\|_* \le c\epsilon^{11}.$$
(159)

We start with (158). This is a perturbative formula for a linear operator. We use the series (46), (47). Indeed, we take $\hat{B}_0(z)$ given by

$$\hat{B}_0(z) = \left(\hat{H}_q(\vec{t}) - z\right)^{-\frac{1}{2}} \hat{W}_0 \left(\hat{H}_q(\vec{t}) - z\right)^{-\frac{1}{2}}, \quad \hat{W}_0 = V - P_q V P_q.$$
(160)

Using $||V||_* < \epsilon^9$ and (156), (157), we obtain:

$$\max_{z \in C_1^+} \|\hat{B}_0(z)\|_0 < c\epsilon.$$
(161)

Now, we consider $\hat{G}^+_{0,r}(k,\vec{t})$, see (45). Applying (161) we get:

$$\|\hat{G}_{0,r}^{+}(k,\vec{t})\|_{0} \le (c\varepsilon)^{r}.$$
 (162)

Hence, (158) follows. Let us estimate $||W_1^+ - W_0^+||_*$. We use again Definition 5 and (89)–(91). Applying the condition $|\sigma A|^2 < \epsilon^{11}$ we obtain (159).

Now, let us assume that (151) -(154) are satisfied for all s = 1, 2, ..., m - 1, i.e.,

$$\|W_s^+ - W_{s-1}^+\|_* \le \epsilon^{11} (c\epsilon)^{s-1} \tag{163}$$

$$\|W_s^+ - V\|_* \le \epsilon^{11} \sum_{r=1}^s (c\epsilon)^{r-1}, \tag{164}$$

$$\|\hat{E}_{s-1}^{+}(\vec{t}) - \hat{E}_{s-2}^{+}(\vec{t})\|_{0} \le (c\epsilon)^{s-1}.$$
(165)

First, let us prove (154). We define $\hat{B}_{m-1}^+(z)$ by (97). Considering as in (100), (101), we get:

$$\max_{z \in C_1} \|\hat{B}_{m-1}^+(z)\|_0 \le \|\hat{W}_{m-1}^+\|_0 \epsilon^{-8} + d^{-1} \|P_q(W_{m-1}^+ - V)P_q\|_0 < c\epsilon.$$
(166)

It follows $\|\hat{E}_{m-1}^+(\vec{t})\|_0 \leq 3$. Now, we consider (104) and further. The analog of (105) is

$$\|\hat{B}_{m-1}^{+}(z) - \hat{B}_{m-2}^{+}(z)\|_{0} \le d^{-1} \|W_{m-1}^{+} - W_{m-2}^{+}\|_{0}, \quad m \ge 2.$$
(167)

The analog of (106) is

$$R \le c \sum_{r=1}^{\infty} d^{-1} \|W_{m-1}^{+} - W_{m-2}^{+}\|_{0} (c\epsilon)^{r-1} \le c d^{-1} \|W_{m-1}^{+} - W_{m-2}^{+}\|_{0} \le c\epsilon (c\epsilon)^{m-2},$$

 $m \ge 2$. Thus, (154) follows. Next, we consider $||W_m^+ - W_{m-1}^+||_0$. Using the first line of (109), we obtain:

$$\|W_m^+ - W_{m-1}^+\|_0 < \epsilon^{11} (c\epsilon)^{m-1}.$$
(168)

Therefore, (151) and (152) hold.

The convergence of the sequences of the eigenvalues and their spectral projections follow.

Lemma 35. There is ϵ_0 , $0 < \epsilon_0 \neq \epsilon_0(\lambda)$, such that for any $0 < \epsilon < \epsilon_0$ under the conditions (148) and for any sufficiently large $k, k > k_1(V, \epsilon_0)$, the following holds. Let \vec{t} belong to the $(\frac{1}{8}k^{-1}\varepsilon^{10})$ -neighborhood in K of the set $\chi_q(k, \epsilon)$. Then, the sequence $\{\hat{E}_m^+(\vec{t})\}_{m=0}^{\infty}$ converges with respect to the norm $\|\cdot\|_0$ to a one-dimensional spectral projection $\hat{E}_W^+(\vec{t})$ of $H_0(\vec{t}) + W^+$ and

$$\|\hat{E}_m^+(\vec{t}) - \hat{E}_W^+(\vec{t})\|_0 < (c\epsilon)^{m+1}, \quad m \ge 0,$$
(169)

$$\|\hat{E}_{-1}^{+}(\vec{t}) - \hat{E}_{W}^{+}(\vec{t})\|_{0} < c\epsilon.$$
(170)

The projection $\hat{E}_W^+(\vec{t})$ is given by (111) where \hat{E}^+ is the spectral projection of $\hat{H}_q(\vec{t})$ corresponding to the eigenvalue $\hat{\lambda}^+(\vec{t})$ and $\hat{G}_{W,r}^+(k,\vec{t})$ is defined by (45) with $W^+ - P_q V P_q$ instead of \hat{V} . Moreover, the following estimate is valid:

$$\|\hat{G}_{Wr}^{+}(k,\vec{t})\|_{0} \le (\hat{c}\epsilon)^{r}.$$
(171)

Proof. The proof of the lemma is analogous to that of Lemma 18.

Using Definition 2, we obtain the following results analogous to Lemma 19, Corollary 20, and Corollary 21.

Lemma 36. There is ϵ_0 , $0 < \epsilon_0 \neq \epsilon_0(\lambda)$, such that for any $0 < \epsilon < \epsilon_0$ under the conditions (148) and for any sufficiently large $k, k > k_1(V, \epsilon_0)$, the following holds. Let \vec{t} belong to the $(\frac{1}{8}k^{-1}\varepsilon^{10})$ -neighborhood in K of the set $\chi_q(k, \epsilon)$. Then, the sequence $\hat{\lambda}_m^+(\vec{t})$ converges to $\lambda_W^+(\vec{t})$ given by (119), and for all r we have

$$|\hat{g}^+_{Wr}(k,\vec{t})| < \epsilon^{10} (c\epsilon)^r, \tag{172}$$

where $\hat{g}_{W,r}$ is given by (44) with $W^+ - P_q V P_q$ instead of \hat{V} .

Corollary 37. The sequence u_m^+ converges to \hat{u}_W^+ in C(Q).

Corollary 38. The operator $\hat{\mathcal{M}}^+$ maps the operator W^+ into W^+ . In other words, $\hat{\mathcal{M}}^+W^+ = W^+$.

Next, we present a solution of non-linear Scrödinger equation in the dimension two.

Theorem 39. There is ϵ_0 , $0 < \epsilon_0 \neq \epsilon_0(\lambda)$, such that for any $0 < \epsilon < \epsilon_0$ under conditions (148) and for any sufficiently large k, $k > k_1(V, \epsilon_0)$, the following holds. Suppose \vec{t} belongs to the $(\frac{1}{8}k^{-1}\varepsilon^{10})$ -neighborhood in K of the resonant set $\chi_q(k, \epsilon)$. Then, there is a pair of functions $u^{\pm}(\vec{x}, \vec{t})$ and the corresponding real values $\lambda^{\pm}(\vec{t})$, satisfying the equation

$$-\Delta u^{\pm} + V(\vec{x})u^{\pm} + \sigma |u^{\pm}|^2 u^{\pm} = \lambda^{\pm} u^{\pm}, \ \vec{x} \in Q,$$
(173)

and the quasi-periodic boundary condition (3). The followings hold:

$$u^{\pm}(\vec{x}, \vec{t}) = A e^{i \langle k, \vec{x} \rangle} \left(\psi^{\pm}_{-1}(\vec{x}, \vec{t}) + \phi^{\pm}(\vec{x}, \vec{t}) \right), \tag{174}$$

$$\lambda^{\pm}(\vec{t}) = \hat{\lambda}^{\pm}(\vec{t}) + O\left(\epsilon^{11}\right), \tag{175}$$

where ψ_{-1}^{\pm} is as defined by (77) and Definition 1 (see also (68), (69)) and $\phi^{\pm}(\vec{x}, \vec{t})$ is periodic in \vec{x} , and satisfies:

$$\|\phi^{\pm}\|_* \le \epsilon. \tag{176}$$

The proof of the theorem is similar to that of Theorem 24, the notation (128) being used.

Theorem 40. Under the conditions of Theorem 39 the series (111) and (119) can be differentiated with respect to \vec{t} retaining their asymptotic character. Moreover, the coefficients $\hat{g}^{\pm}_{W,r}(k,\vec{t})$ and $\hat{G}^{\pm}_{W,r}(k,\vec{t})$ satisfy the following estimates in the $(\frac{1}{8}k^{-1}\varepsilon^{10})$ neighborhood in C^2 of the set $\chi_q(k,\delta)$:

$$|\nabla_{\vec{t}}\hat{g}_{W,r}^{\pm}(k,\vec{t})| < (\hat{c}\varepsilon)^r k \tag{177}$$

$$\|\nabla_{\vec{t}}\hat{G}^{\pm}_{W,r}(k,\vec{t})\|_{0} < (c\varepsilon)^{r} (k\varepsilon^{-10}).$$
(178)

Corollary 41. The followings hold for the perturbed eigenvalue and its spectral projection:

$$\left|\nabla_{\vec{t}} \left(\lambda^{\pm}(\vec{t}) - \hat{\lambda}^{\pm}(\vec{t})\right)\right| < C(V)k\varepsilon$$
(179)

$$\|\nabla_{\vec{t}}(E^{\pm}(\vec{t}) - \hat{E}^{\pm})\|_{0} < C(V)k\varepsilon^{-9}.$$
(180)

Corollary 42.

$$|\nabla_{\vec{t}}\lambda^{\pm}(\vec{t})| = 2k(1+O(\epsilon)).$$
(181)

Let us consider the surface $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$ for a fixed $\lambda_0, \lambda_0 > k_1^2(V, \epsilon_0)$. Note that the parts $\hat{\lambda}^+(\vec{t}) = \lambda_0$ and $\hat{\lambda}^-(\vec{t}) = \lambda_0$ do not intersect, since $v_q \neq 0$. Thus the deviation of the surface $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$ from the unperturbed one (V=0) is essential. The next theorem follows.

Theorem 43. If $0 < \varepsilon < \epsilon_0$, $\lambda_0 > k_1^{2l}(V, \epsilon_0)$, then the curves $\lambda^{\pm}(\vec{t}) = \lambda_0$ are in the $C(V)\varepsilon^{11}\lambda_0^{-1/2}$ -neighborhood of the curves $\hat{\lambda}^{\pm}(\vec{t}) = \lambda_0$.

Corollary 44. The curves $\lambda^+(\vec{t}) = \lambda_0$ and $\lambda^-(\vec{t}) = \lambda_0$ do not intersect and located at the distance greater than $|v_q| + O\left(\varepsilon^{11}\lambda_0^{-1/2}\right)$ from each other as $\varepsilon \lambda_0^{-1/2} \to 0$.

References

- V. V. Konotop and M. Salerno, Modulational instability in Bose-Einstein condensates in optical lattices, Phys. Rev. A 65, 021602, 2002.
- [2] Pearl J.Y. Louis, Elena A. Ostrovskaya, Craig M. Savage and Yuri S. Kivshar, Bose-Einstein Condensates in Optical Lattices: Band-Gap Structure and Solitons, Phys. Rev. A 67, 013602, 2003.
 [3] C. J. Pethick, H. Smith, Bose-Einstein Condensation in Dilute Gases, Cambridge, 2008.
- [4] A. V. Yulin, Yu. V. Bludov, V. V. Konotop, V. Kuzmiak, and M. Salerno, Superfluidity breakdown of periodic matter waves in quasi-one-dimensional annular traps via resonant scattering with moving defects, Phys. Rev. A 87, 2013.
- [5] Alexey V. Yulin and Dmitry V. Skryabin, Out-of-gap Bose-Einstein solitons in optical lattices, Phys. Rev. A 67, 023611, 2003.

- [6] J. Bourgain "Quasi-Periodic Solutions of Hamiltonian Perturbations of 2D Linear Schrödinger Equations", Annals of Mathematics, Second Series, Vol. 148, No. 2 (Sep., 1998), pp. 363-439 (77 pages)
- [7] Carlos E. Kenig, " On the work of Jean Bourgain in Non-linear Dispersive Equations", Bulletin (New Series) of the AMS, Volume 58, Number 2, April 2021, Pages 173–189.
- [8] J. Bourgain, Construction of quasi-periodic solutions for Hamiltonian perturbations of linear equations and applications to nonlinear PDE, Internat. Math. Res. Notices 11 (1994), 21 pp.
- [9] Bourgain, J. Scattering in the energy space and below for 3D NLS.J. Anal. Math.75(1998), 267–297.
- [10] N. Burq and M. Zworski, Instability for the semiclassical nonlinear Schrödinger equation, Commun. Math. Phys. 260(1) (2005) 45–48.
- [11] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, Transfer of energy to high frequencies in the cubic defocusing nonlinear Schrödinger equation, Invent. Math. 181 (2010), no. 1, 39–113
- [12] Colliander, J.; Keel, M.; Staffilani, G.; Takaoka, H.; Tao, T. "Resonant decompositions and the I-method for the cubic nonlinear Schrödinger equation on \mathbb{R}^2 ." Discrete Contin. Dyn. Syst.21(2008), no.3, 665–686.
- [13] Hongzi Cong, Jianjun Liu, Xiaoping Yuan, "Stability of KAM tori for nonlinear Schrödinger equation," Memoirs of the American Mathematical Society; **239**, 1134, 2016.
- [14] Reed, M., Simon, B., *Methods of Modern Mathematical Physics*, Vol IV, 3rd ed., Academic Press, New York, San Francisco, London, 1987.
- [15] Yu. Karpeshina, Seonguk Kim, R. Shterenberg, Solutions of Nonlinear Polyharmonic Equation with Periodic Potentials, Analysis as a Tool in Mathematical Physics: In Memory of Boris Pavlov (Operator Theory: Advances and Applications, 276), pp 401–416, 2020.
- [16] Yu. Karpeshina, Seonguk Kim, R.Shterenberg "Solutions of Gross-Pitaevskii Equation with Periodic Potential in Dimension Three", Saint-Petersburg Math. Journal (Algebra i Analiz), 2023, 35:1, 204–225.
- [17] Yulia E. Karpeshina, Perturbation Theory for the Schrödinger Operator with a Periodic Potential, Springer, 1997.

DEPARTMENT OF MATHEMATICS, COLLEGE OF SAINT BENEDICT AND SAINT JOHN'S UNIVERSITY, 2850 ABBEY PLAZA COLLEGEVILLE, MINNESOTA 56321, USA

 $Email \ address: \verb"aduaibes001@csbsju.edu"$

Department of Mathematics, University of Alabama at Birmingham, University Hall, Room 4005, 1402 10th Avenue South, Birmingham AL 35294-1241, USA

Email address: karpeshi@uab.edu