

## ON THE SOLUTION SET OF SEMI-INFINITE TENSOR COMPLEMENTARITY PROBLEM

R. Deb<sup>a,1</sup> and A. K. Das<sup>b,2</sup><sup>a</sup>*Jadavpur University, Kolkata , 700 032, India.*<sup>b</sup>*Indian Statistical Institute, 203 B. T. Road, Kolkata, 700 108, India.*<sup>1</sup>*Email: rony.knc.ju@gmail.com*<sup>2</sup>*Email: akdas@isical.ac.in***Abstract**

In this paper, we introduce semi-infinite tensor complementarity problem to provide an approach for considering a more realistic situation of the problem. We prove the necessary and sufficient conditions for the existence of the solution set. In this context, we study the error bounds of the solution set in terms of residual function.

**Keywords:** Semi-infinite tensor complementarity problem, error bound, residual function,  $R_0$ -tensor.

**AMS subject classifications:** 90C30, 90C33, 15A69.

# 1 Introduction

The tensor complementarity problem,  $\text{TCP}(q, \mathcal{A})$ , was introduced by Song and Qi [37] and [36]. By reformulating the multilinear game as a tensor complementarity problem, Huang and Qi [12] established a bridge between the multilinear game and tensor complementarity problem. They demonstrated that finding a Nash equilibrium point of the multilinear game is equivalent to finding a solution to the resulting TCP.

Let  $\mathcal{A}$  be a tensor of order  $m$  and dimension  $n$ , i.e.,  $\mathcal{A} \in T_{m,n}$  and a vector  $q \in \mathbb{R}^n$ , the tensor complementarity problem denoted by  $\text{TCP}(q, \mathcal{A})$  is to find  $x \in \mathbb{R}^n$  such that,

$$q + \mathcal{A}x^{m-1} \geq 0, \quad x \geq 0, \quad \text{and} \quad x^T(q + \mathcal{A}x^{m-1}) = 0. \quad (1.1)$$

When the order of the tensor  $m = 2$  then the problem reduces to a linear complementarity problem. Let  $A$  be an  $n \times n$  real matrix and a vector  $q \in \mathbb{R}^n$ , the linear complementarity problem, denoted by  $\text{LCP}(q, A)$  is finding  $x \in \mathbb{R}^n$  such that

$$q + Ax \geq 0, \quad x \geq 0, \quad \text{and} \quad x^T(q + Ax) = 0. \quad (1.2)$$

The idea of complementarity considers a large number of optimization problems. The problems which can be constituted as linear complementarity problem includes linear programming, linear fractional programming, convex quadratic programming and the bimatrix game problem. It is well considered in the literature on mathematical programming and occurs in a number of applications in operations research, control theory, mathematical

---

<sup>1</sup>Corresponding author

economics, geometry and engineering. For recent works on this problem and applications see [18], [19], [27], [25], [20], [6], [9], [14], [16], [33] and [23] references cited therein.

The concept of PPT is originally motivated by the well-known linear complementarity problem, and applied in many other settings. The PPT is basically a transformation of the matrix of a linear system for exchanging unknowns with the corresponding entries of the right hand side of the system. For details see [5], [21], [32], [3] and [26].

Due to their prominence in scientific computing, complexity theory, and the theoretical underpinnings of linear complementarity problems, a number of matrix classes and their subclasses have received substantial study. For recent work on this problem and applications see [13], [4], [10], [29], [31], [7], [15]. For multivariate analysis and game problem, See [21], [17], [15], [30], [28], [24] and references cited therein.

An implicit assumption shared by the nonlinear complementarity problem is that the information about the mapping  $F$  and the cone involved are all fixed and completely independent of other related parameters. However, this type of formulation is unable to model all realistic situations of the problem and fails to explain the complete reality. For example, in optimal control or engineering design fields [2], the data of the problem involves a time parameter; in non-cooperative games (e.g., generalized Nash equilibrium [11]), the strategy of each player is dependent on the strategy of the other players in case of the realistic model. Here we introduce semi-infinite tensor complementarity (SITCP), a generalized version of nonlinear complementarity to accommodate more number of realistic situations into the model.

The paper is organised as follows. Section 2 presents some basic notations and results which are used in the next section. In section 3 we prove the existence of the solution set for semi-infinite tensor complementarity problem with some assumptions. We establish a connection between the solution set of semi-infinite tensor complementarity problem and the solution sets of its equivalent tensor complementarity problems. We establish the necessary and sufficient conditions for the error bound of solution set to be level bounded in terms of residual function.

## 2 Preliminaries

We begin by outlining the fundamental concepts and the notation that will be applied throughout the text. Here we consider the vectors, matrices and tensors of real entries. For any positive integer  $n$ , the set  $\{1, \dots, n\}$  is denoted by  $[n]$ . Let  $\mathbb{R}^n$  denote the  $n$ -dimensional Euclidean space and  $\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x \geq 0\}$ . Any vector  $x \in \mathbb{R}^n$  is a column vector unless specified otherwise. The Euclidean norm of a vector  $x$  is defined as  $\|x\|_2 = \sqrt{|x_1^2| + \dots + |x_n^2|}$ . The distance of  $x$  from  $A \subseteq \mathbb{R}^n$  is denoted as  $\text{dist}(x; A)$  and is defined as  $\text{dist}(x; A) = \inf\{(x, a) : a \in A\}$ . The diameter of a set is denoted as  $\text{diam}A$  and is defined as  $\text{diam}A = \sup_{x, y \in A} \|x - y\|$ . The unit ball in  $\mathbb{R}^n$  is  $\mathbb{B} = \{x \in \mathbb{R}^n : \|x\| \leq 1\}$ . An  $m$ th order  $n$  dimensional real tensor  $\mathcal{A} = (a_{i_1 \dots i_m})$  is a multidimensional array of entries  $a_{i_1 \dots i_m} \in \mathbb{R}$  where  $i_j \in [n]$  with  $j \in [m]$ . The set of all  $m$ th order  $n$  dimensional real tensors are denoted by  $T_{m, n}$ . Shao [34] introduced a product of tensors. Let  $\mathcal{A}$  with order  $q \geq 2$  and  $\mathcal{B}$  with order  $k \geq 1$  be two  $n$ -dimensional tensors. The product of  $\mathcal{A}$  and  $\mathcal{B}$  is a tensor  $\mathcal{C}$  of order  $(q-1)(k-1)+1$  and dimension  $n$  with entries  $c_{j\alpha_1 \dots \alpha_{m-1}} = \sum_{j_2, \dots, j_m \in [n]} a_{jj_2 \dots j_m} b_{j_2\alpha_1} \dots b_{j_m\alpha_{m-1}}$ , where  $j \in [n]$ ,  $\alpha_1, \dots, \alpha_{m-1} \in [n]^{k-1}$ .

Then for a tensor  $\mathcal{A} \in T_{m,n}$  and  $x \in \mathbb{R}^n$ ,  $\mathcal{A}x^{m-1} \in \mathbb{R}^n$  is a vector defined by

$$(\mathcal{A}x^{m-1})_i = \sum_{i_2, \dots, i_m=1}^n a_{ii_2 \dots i_m} x_{i_2} \cdots x_{i_m}, \quad \forall i \in [n],$$

and  $\mathcal{A}x^m \in \mathbb{R}$  is a scalar defined by

$$x^T \mathcal{A}x^{m-1} = \mathcal{A}x^m = \sum_{i_1, \dots, i_m=1}^n a_{i_1 \dots i_m} x_{i_1} \cdots x_{i_m}.$$

Given a vector  $q \in \mathbb{R}^n$  and a tensor  $\mathcal{A} \in T_{m,n}$  the set of feasible solution of  $\text{TCP}(q, \mathcal{A})$  is defined as  $\text{FEA}(q, \mathcal{A}) = \{x \in \mathbb{R}_+^n : q + \mathcal{A}x^{m-1} \geq 0\}$  and the solution set of  $\text{TCP}(q, \mathcal{A})$  as  $S = \text{SOL}(q, \mathcal{A}) = \{x \in \text{FEA}(q, \mathcal{A}) : x^T(q + \mathcal{A}x^{m-1}) = 0\}$ . A residual function  $r(x)$  is a global (local) error bound for  $\text{TCP}$  if  $\exists$  some constant  $c > 0$  (and  $\epsilon > 0$ ) such that for each  $x \in \mathbb{R}^n$  (when  $r(x) \leq \epsilon$ )

$$\text{dist}(x, S) \leq cr(x). \quad (2.1)$$

We consider some definitions and results which are required for the next sections.

DEFINITION 2.1: [35], [8] The  $i$ th row subtensor of  $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$  is denoted by  $R_i(\mathcal{A})$  and its entries are given as  $(R_i(\mathcal{A}))_{i_2 \dots i_m} = (a_{ii_2 \dots i_m})$ , where  $i_j \in [n]$  and  $2 \leq j \leq m$ .

DEFINITION 2.2: [39] A function  $f : \mathbb{R}^n \mapsto \mathbb{R}^n$  is said to be level bounded if for every  $\alpha \geq 0$  the level set  $\{x \in \mathbb{R}^n : f(x) \leq \alpha\}$  is bounded.

DEFINITION 2.3: [36] A tensor  $\mathcal{A} \in T_{m,n}$  is said to be a  $S$ -tensor if the system

$$\mathcal{A}x^{m-1} > 0, \quad x > 0$$

has a solution.

DEFINITION 2.4: [36] A tensor  $\mathcal{A} \in T_{m,n}$  is said to be a  $P$ -tensor, if for each  $x \in \mathbb{R}^n \setminus \{0\}$ , there exists an index  $i \in [n]$  such that  $x_i \neq 0$  and  $x_i(\mathcal{A}x^{m-1})_i > 0$ .

DEFINITION 2.5: [38], [36] A tensor  $\mathcal{A} \in T_{m,n}$  is said to be a  $R_0$ -tensor if the  $\text{TCP}(0, \mathcal{A})$  has unique zero solution.

DEFINITION 2.6: [41] For a given  $\epsilon \geq 0$  a residual function  $r(x)$  is said to be an  $\epsilon$ -error bound for  $\text{TCP}$  if  $\exists c > 0$  such that  $\text{dist}(x, S) \leq cr(x) + \epsilon \quad \forall x \in \mathbb{R}^n$ . If  $\epsilon = 0$ , the definition reduces to the error bound.

THEOREM 2.1: [22] A non-empty family  $\mathcal{A}$  of subsets of  $\mathbb{R}^n$  is said to have the finite intersection property (FIP) if the intersection over any finite subcollection of  $\mathcal{A}$  is non-empty.

THEOREM 2.2: [1] For a  $P$ -tensor  $\mathcal{A} \in T_{m,n}$  and any  $q \in \mathbb{R}^n$  the solution set of  $\text{TCP}(q, \mathcal{A})$  is nonempty and compact.

THEOREM 2.3: [38] If  $\mathcal{A} \in T_{m,n}$  is a an  $R_0$ -tensor then for  $q \in \mathbb{R}^n$ , the solution set of the  $\text{TCP}(q, \mathcal{A})$  is bounded.

### 3 Main results

We begin by introducing semi-infinite tensor complementarity problem  $\text{SITCP}(q(\omega), \mathcal{A}(\omega), \Omega)$ .

Find a vector  $x \in \mathbb{R}^n$  such that

$$x \geq 0, \quad F(x, \omega) \geq 0, \quad x^T F(x, \omega) = 0, \quad \omega \in \Omega \quad (3.1)$$

where  $F : \mathbb{R}^n \times \Omega \mapsto \mathbb{R}^n$ ,  $F = \mathcal{A}(\omega)x^{m-1} + q(\omega)$  and  $\Omega$  is a set in  $\mathbb{R}^p$ . The solution set of  $\text{SITCP}(q(\omega), \mathcal{A}(\omega), \Omega)$  is denoted by  $S^* = \text{SOL}(q(\omega), \mathcal{A}(\omega), \Omega) = \{x \geq 0 : \mathcal{A}(\omega)x^{m-1} + q(\omega) \geq 0, x^T(\mathcal{A}(\omega)x^{m-1} + q(\omega)) = 0, \forall \omega \in \Omega\}$ .

Here we establish the necessary and sufficient conditions for  $S^*$  to be non-empty.

**THEOREM 3.1:** Consider the  $\text{SITCP}(q(\omega), \mathcal{A}(\omega), \Omega)$  and  $\mathcal{A}(\omega_0)$  is an  $R_0$ -tensor for some  $\omega_0 \in \Omega$ .  $S^* \neq \emptyset$  if and only if  $\bigcap_{i=1}^p \text{SOL}(q(\omega_i), \mathcal{A}(\omega_i)) \neq \emptyset$  for finitely many points  $\omega_1, \dots, \omega_p \in \Omega$ .

*Proof.* If part: Since  $S^* \subseteq \bigcap_{i=1}^p \text{SOL}(q(\omega_i), \mathcal{A}(\omega_i))$ ,  $S^* \neq \emptyset$  implies that  $\bigcap_{i=1}^p \text{SOL}(q(\omega_i), \mathcal{A}(\omega_i)) \neq \emptyset$ .

Only if part: Since  $\mathcal{A}(\omega_0)$  is an  $R_0$ -tensor, by Theorem 2.3, the set  $\text{SOL}(q(\omega_0), \mathcal{A}(\omega_0))$  is bounded. Which in turn implies the boundedness of  $S^*$ . On the other hand, since  $\text{SOL}(q(\omega), \mathcal{A}(\omega))$  is closed for each  $\omega$ , so is  $S^*$ . Thus  $S^*$  is compact. By Theorem 2.1 of finite intersection of compact sets, we obtain the result.  $\square$

In the next result we find the position of  $S^*$ . For this purpose we consider the followings:

- (i)  $(\mathcal{A}_{\max})_{i_1 \dots i_m} = (\bar{a}_{i_1 \dots i_m}) = \max_{\omega \in \Omega} a_{i_1 \dots i_m}(\omega)$
- (ii)  $(\mathcal{A}_{\min})_{i_1 \dots i_m} = (a'_{i_1 \dots i_m}) = \min_{\omega \in \Omega} a_{i_1 \dots i_m}(\omega)$
- (iii)  $(q_{\max})_i = \max_{\omega \in \Omega} q_i(\omega)$
- (iv)  $(q_{\min})_i = \min_{\omega \in \Omega} q_i(\omega)$

**THEOREM 3.2:** Consider  $\text{SITCP}(q(\omega), \mathcal{A}(\omega), \Omega)$ . If  $\Omega$  is compact and  $\mathcal{A}(\omega)$  and  $q(\omega)$  are continuous on  $\Omega$ , then  $S^* \subseteq \text{SOL}(q_{\max}, \mathcal{A}_{\max}) \cap \text{SOL}(q_{\min}, \mathcal{A}_{\min})$ . Furthermore, suppose in each row sub-tensor,  $R_i(\mathcal{A}(\omega))$  and in each row of  $q(\omega)$ ,  $q_i(\omega)$  the minimum (and maximum) is attained by a common  $\omega'$  (and  $\bar{\omega}$ ), i.e., for each  $i = 1, 2, \dots, n$ , there exist  $\omega'_i, \bar{\omega}_i \in \Omega$  such that  $R_i(\mathcal{A}_{\min}) = R_i(\mathcal{A}(\omega'_i))$ ,  $(q_{\min})_i = q(\omega'_i)_i$  and  $R_i(\mathcal{A}_{\max}) = R_i(\mathcal{A}(\bar{\omega}_i))$ ,  $(q_{\max})_i = q(\bar{\omega}_i)_i$ . Then

$$S^* = \text{SOL}(q_{\max}, \mathcal{A}_{\max}) \cap \text{SOL}(q_{\min}, \mathcal{A}_{\min}).$$

*Proof.* By the rules of maximization and minimization for the summation of functions in Exercise 1.36 of [39], we have,

$$\max_{\omega \in \Omega} (\mathcal{A}(\omega)x + q(\omega)) \geq \mathcal{A}_{\max}x + q_{\max} \quad (3.2)$$

$$\min_{\omega \in \Omega} (\mathcal{A}(\omega)x + q(\omega)) \leq \mathcal{A}_{\min}x + q_{\min} \quad (3.3)$$

for all  $x \geq 0$ . By using the inequalities (3.2) and (3.3) and following an argument similar to that for Theorem 2.2 of [41], we obtain

$$S^* \subseteq \text{SOL}(q_{\max}, \mathcal{A}_{\max}) \cap \text{SOL}(q_{\min}, \mathcal{A}_{\min}). \quad (3.4)$$

Now to prove the second part let us assume that the conditions of Theorem 3.2 hold. Then  $(\mathcal{A}_{\min}x^{m-1})_i = (\mathcal{A}(\omega')x^{m-1})_i$  and  $(\mathcal{A}_{\min}x^{m-1})_i + (q_{\min})_i = (\mathcal{A}(\omega'_i)x^{m-1} + q(\omega'_i))_i$ . Therefore  $\text{SOL}(q_{\min}, \mathcal{A}_{\min}) = \text{SOL}(q(\omega'_i), \mathcal{A}(\omega'_i))$ . Similarly we have  $\text{SOL}(q_{\max}, \mathcal{A}_{\max}) = \text{SOL}(q(\bar{\omega}_i), \mathcal{A}(\bar{\omega}_i))$ . Since  $S^* = \bigcap_{\omega \in \Omega} \text{SOL}(q(\omega), \mathcal{A}(\omega))$ , using (3.2) and (3.3) we conclude that

$$S^* = \text{SOL}(q_{\max}, \mathcal{A}_{\max}) \cap \text{SOL}(q_{\min}, \mathcal{A}_{\min}).$$

□

Here we provide an example to illustrate the result of Theorem 3.2.

EXAMPLE 3.1: Let  $\mathcal{A}(\omega) \in T_{3,2}$  and  $q(\omega) \in \mathbb{R}^2$  be such that  $a_{111} = (1 - 2\omega^3)$ ,  $a_{121} = 1 - \omega$ ,  $a_{112} = (1 - \omega)$ ,  $a_{122} = -1$ ,  $a_{211} = a_{212} = a_{221} = 0$ ,  $a_{222} = -\omega^2$  and  $q(\omega) = \begin{pmatrix} 1 \\ \omega^2 \end{pmatrix}$  and  $\omega \in \Omega = [0, 1]$ . Then  $\mathcal{A}(\omega)x^2 = \begin{pmatrix} (1 - 2\omega^3)x_1^2 + 2(1 - \omega)x_1x_2 - x_2^2 \\ -\omega^2x_2^2 \end{pmatrix}$ . Now consider the SITCP( $q(\omega), \mathcal{A}(\omega), \Omega$ ) which is to find  $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$  such that

$$x_1 \geq 0; \quad (\mathcal{A}(\omega)x^2)_1 \geq 0; \quad x_1[(1 - 2\omega^3)x_1^2 + 2(1 - \omega)x_1x_2 - x_2^2 + 1] = 0, \quad (3.5)$$

$$x_2 \geq 0; \quad (\mathcal{A}(\omega)x^2)_2 \geq 0; \quad x_2[-\omega^2x_2^2 + \omega^2] = 0. \quad (3.6)$$

Solving the equations (3.5) and (3.6) we obtain the solution set for  $\omega \in \Omega$ , which is  $\left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} \frac{2(1-\omega)}{1-2\omega^3} \\ 1 \end{pmatrix} \right\}$ . For  $\omega = 1$  we get  $\frac{2(1-\omega)}{1-2\omega^3} = 0$  so  $S^* = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}$ . Now for the given tensor  $\mathcal{A}(\omega)$ , let  $\mathcal{A}_{\max} = (\bar{a}_{ijk}) \in T_{3,2}$ . Then we obtain  $\bar{a}_{111} = 1$ ,  $\bar{a}_{121} = \bar{a}_{112} = 1$ ,  $\bar{a}_{122} = -1$  and  $\bar{a}_{211} = \bar{a}_{212} = \bar{a}_{221} = \bar{a}_{222} = 0$  and  $q_{\max} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ . Then the TCP( $q_{\max}, \mathcal{A}_{\max}$ ) which is finding  $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$  such that

$$x_1 \geq 0; \quad (\mathcal{A}_{\max}x^2)_1 \geq 0; \quad x_1[x_1^2 + 2x_1x_2 - x_2^2 + 1] = 0, \quad (3.7)$$

$$x_2 \geq 0; \quad (\mathcal{A}_{\max}x^2)_2 \geq 0; \quad x_2[0 + 1] = 0. \quad (3.8)$$

Solving (3.7) and (3.8) we get  $\text{SOL}(q_{\max}, \mathcal{A}_{\max}) = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ .

Again, let  $\mathcal{A}_{\min} = (a'_{ijk}) \in T_{3,2}$ . Then we obtain  $a'_{111} = -1$ ,  $a'_{121} = a'_{112} = 0$ ,  $a'_{122} = -1$  and  $a'_{211} = a'_{212} = a'_{221} = 0$ ,  $a'_{222} = -1$  and  $q_{\min} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ . Then TCP( $q_{\min}, \mathcal{A}_{\min}$ ) is finding  $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$  such that

$$x_1 \geq 0; \quad (\mathcal{A}_{\min}x^2)_1 \geq 0; \quad x_1[-x_1^2 - x_2^2 + 1] = 0, \quad (3.9)$$

$$x_2 \geq 0; \quad (\mathcal{A}_{\min}x^2)_2 \geq 0; \quad x_2[-x_2^2] = 0. \quad (3.10)$$

Solving (3.9) and (3.10) we have  $\text{SOL}(q_{\min}, \mathcal{A}_{\min}) = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}$ .

Thus  $S^* \supset \text{SOL}(q_{\max}, \mathcal{A}_{\max}) \cap \text{SOL}(q_{\min}, \mathcal{A}_{\min})$ , i.e., the inclusion is strict.

Now we replace  $q(\omega)$  by  $\bar{q} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ . Then the SITCP( $\bar{q}, \mathcal{A}(\omega), \Omega$ ) which is to find  $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$  such that

$$x_1 \geq 0; \quad (\mathcal{A}(\omega)x^2)_1 \geq 0; \quad x_1[(1 - 2\omega^3)x_1^2 + 2(1 - \omega)x_1x_2 - x_2^2 + 1] = 0, \quad (3.11)$$

$$x_2 \geq 0; \quad (\mathcal{A}(\omega)x^2)_2 \geq 0; \quad x_2[-\omega^2x_2^2 + 1] = 0. \quad (3.12)$$

Solving the equations (3.11) and (3.12) we obtain the solution set for  $\omega \in \Omega$ , which is  $\left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \alpha(\omega) \\ \frac{1}{\omega} \end{pmatrix} \right\}$ . Here  $\alpha(\omega)$  is the positive root of the equation

$$\omega^2(1 - 2\omega^3)x_1^2 - 2\omega(1 - \omega)x_1 - (1 - \omega^2) = 0.$$

For  $\omega = 1$  we get  $\alpha(1) = 0$ . Therefore  $S^* = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ .

Then  $\text{TCP}(q_{\max}, \mathcal{A}_{\max}) = \text{TCP}(\bar{q}, \mathcal{A}_{\max})$ .  $\text{SOL}(\bar{q}, \mathcal{A}_{\max}) = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ .

Now,  $\text{TCP}(\bar{q}, \mathcal{A}_{\min})$  is finding  $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$  such that

$$x_1 \geq 0; \quad (\mathcal{A}_{\min}x^2)_1 \geq 0; \quad x_1[-x_1^2 - x_2^2 + 1] = 0, \quad (3.13)$$

$$x_2 \geq 0; \quad (\mathcal{A}_{\min}x^2)_2 \geq 0; \quad x_2[-x_2^2 + 1] = 0. \quad (3.14)$$

Solving (3.13) and (3.14) we have  $\text{SOL}(\bar{q}, \mathcal{A}_{\min}) = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}$ . In this case we have  $S^* = \text{SOL}(q_{\max}, \mathcal{A}_{\max}) \cap \text{SOL}(q_{\min}, \mathcal{A}_{\min})$ .

For the next result we define semi-infinite  $S$ -tensor.

**DEFINITION 3.7:** A tensor  $\mathcal{A}(\omega) \in T_{m,n}$  is said to be a semi-infinite  $S$ -tensor with respect to the set  $\omega$  if  $\exists$  a vector  $x > 0$  such that  $\mathcal{A}(\omega)x^{m-1} > 0, \forall \omega \in \Omega$ .

Here we establish a connection between the semi-infinite  $S$ -tensor and the feasibility of SITCP( $q(\omega), \mathcal{A}(\omega), \Omega$ ).

**THEOREM 3.3:** Consider the SITCP( $q(\omega), \mathcal{A}(\omega), \Omega$ ) where  $\Omega$  is compact and all the elements of  $\mathcal{A}(\omega)$  are continuous on  $\Omega$ . Then  $\mathcal{A}(\omega)$  is a semi-infinite  $S$ -tensor relative to  $\Omega$  if and only if SITCP( $q(\omega), \mathcal{A}(\omega), \Omega$ ) is feasible for all  $q(\omega) \in C(\Omega)$  where  $C(\Omega)$  denotes all continuous mapping on  $\Omega$ .

*Proof.* If part: Since  $\mathcal{A}(\omega)$  is a semi-infinite  $S$ -tensor,  $\exists x > 0$  such that  $\mathcal{A}(\omega)x^{m-1} > 0$ . Then  $\exists$  a sufficiently small scalar  $\lambda > 0$  such that  $\mathcal{A}(\omega)x^{m-1} \geq \lambda e > 0$ , for all  $\omega \in \Omega$  where  $e = (1, \dots, 1)^T$ . Now choose  $\alpha > 0$  with  $\alpha e > -q_{\min}$ . Choosing  $\bar{\alpha} = \left(\frac{\alpha}{\lambda}\right)^{\frac{1}{m-1}} > 0$  we have  $\bar{\alpha}x > 0$ . Also,  $\mathcal{A}(\omega)(\bar{\alpha}x)^{m-1} = \frac{\alpha}{\lambda} \mathcal{A}(\omega)x^{m-1} \geq \alpha e > -q_{\min}$ . Thus for  $\bar{\alpha}x > 0$  we have  $\mathcal{A}(\omega)(\bar{\alpha}x)^{m-1} + q(\omega) \geq \mathcal{A}(\omega)(\bar{\alpha}x)^{m-1} + q_{\min} > 0$ . Therefore,  $\bar{\alpha}x$  is a feasible point of SITCP( $q(\omega), \mathcal{A}(\omega), \Omega$ ).

Only if part: Let  $q(\omega) := \tilde{q} < 0$  for all  $\omega \in \Omega$ . Let the SITCP( $\tilde{q}$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ) be feasible. Then  $\exists$  a vector  $x \geq 0$  such that  $\mathcal{A}(\omega)x^{m-1} + \tilde{q} \geq 0 \implies \mathcal{A}(\omega)x^{m-1} \geq -\tilde{q} > 0$  for all  $\omega \in \Omega$ . Since  $\mathcal{A}(\omega)x^{m-1}$  is continuous on  $\Omega$  there exists a sufficiently small  $\lambda > 0$  such that  $x + \lambda e > 0$ , and  $\mathcal{A}(\omega)(x + \lambda e)^{m-1} > 0$ .  $\square$

The following corollary provides a necessary condition for feasibility of the solution set of SITCP( $q(\omega)$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ).

**COROLLARY 3.1:** Consider the SITCP( $q(\omega)$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ). Suppose  $\Omega$  is compact and  $\mathcal{A}(\omega)$  is continuous on  $\Omega$ . If  $\mathcal{A}_{\min}$  is an  $S$ -tensor, then SITCP( $q(\omega)$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ) is feasible for all  $q(\omega) \in C(\Omega)$ .

*Proof.* Let  $\mathcal{A}_{\min}$  be an  $S$ -tensor. Then for some  $x > 0$  we have  $\mathcal{A}_{\min}x^{m-1} > 0$ . From the definition of  $\mathcal{A}_{\min}$  it follows that  $\mathcal{A}(\omega)x^{m-1} > 0$  for all  $\omega \in \Omega$ . This implies that  $\mathcal{A}(\omega)$  is a semi-infinite  $S$ -tensor. Hence by Theorem 3.3, we conclude that SITCP( $q(\omega)$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ) is feasible for all  $q(\omega) \in C(\Omega)$ .  $\square$

Now we establish an  $\epsilon$ -error bound for the solution set of semi-infinte tensor complementarity problem. Here  $\epsilon$  represents the degree of approximation of the set  $S^*$

**THEOREM 3.4:** Consider the SITCP( $q(\omega)$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ). Suppose the solution set  $S^*$  is nonempty. If  $\mathcal{A}(\omega_0)$  is an  $P$ -tensor for some  $\omega_0 \in \Omega$ , then there exist  $c > 0$  and  $\epsilon > 0$  with  $\epsilon \leq \text{diam}(\text{SOL}(q(\omega_0), \mathcal{A}(\omega_0)))$  such that

$$\text{dist}(x, S^*) \leq c r(x) + \epsilon$$

where residual function is  $r_y(x) = \max_{\omega \in \Omega} \|\min\{x, [\mathcal{A}(\omega)(x - y)]^{\frac{1}{m-1}} + [\mathcal{A}(\omega)y^{m-1}]^{\frac{1}{m-1}}\}\|$ , and  $y \in (\text{SOL}(q(\omega_0), \mathcal{A}(\omega_0)))$ .

*Proof.* Since  $\mathcal{A}(\omega_0)$  is an  $P$ -tensor,  $\text{SOL}(q(\omega_0), \mathcal{A}(\omega_0))$  is bounded. This implies  $\exists$  an  $\epsilon > 0$  such that  $\text{SOL}(q(\omega_0), \mathcal{A}(\omega_0)) \subseteq S^* + \epsilon\mathbb{B}$ . Consequently,

$$\text{dist}(x, S^*) \leq \text{dist}(x, \text{SOL}(q(\omega_0), \mathcal{A}(\omega_0))) + \epsilon, \forall x \in \mathbb{R}^n. \quad (3.15)$$

Notice that  $S^* \subseteq \text{SOL}(q(\omega_0), \mathcal{A}(\omega_0))$ . Therefore the diameter of the set  $\text{SOL}(q(\omega_0), \mathcal{A}(\omega_0))$  is an upper bound of  $\epsilon$ . By Theorem 3.2 of [40] for the tensor complementarity problem TCP( $q(\omega_0)$ ,  $\mathcal{A}(\omega_0)$ ), there exists  $c > 0$  such that  $\forall x \in \mathbb{R}^n$ ,

$$\text{dist}(x, \text{SOL}(q(\omega_0), \mathcal{A}(\omega_0))) \leq c \|\min\{x, [\mathcal{A}(\omega_0)(x - y)]^{\frac{1}{m-1}} + [\mathcal{A}(\omega_0)y^{m-1}]^{\frac{1}{m-1}}\}\|. \quad (3.16)$$

From (3.15) and (3.16) the desired result follows.  $\square$

For the next result we define semi-infinite  $R_0$ -tensor and establish a connection between  $R_0$ -tensor and semi-infinite  $R_0$ -tensor.

**DEFINITION 3.8:** The tensor  $\mathcal{A}(\omega)$  is said to be a semi-infinite  $R_0$ -tensor relative to a set  $\Omega$  if the SITCP( $0$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ) has zero as its unique solution, i.e.,

$$x \geq 0, \mathcal{A}(\omega)x \geq 0, x^T \mathcal{A}(\omega)x = 0, \quad \forall \omega \in \Omega \implies x = 0. \quad (3.17)$$

**THEOREM 3.5:** Consider the SITCP( $q(\omega)$ ,  $\mathcal{A}(\omega)$ ,  $\Omega$ ). If  $\mathcal{A}(\omega_0)$  is an  $R_0$ -tensor for some  $\omega_0 \in \Omega$ . Then  $\mathcal{A}(\omega)$  is a semi-infinite  $R_0$ -tensor relative to  $\Omega$ .

*Proof.* Since  $\mathcal{A}(\omega_0)$  is an  $R_0$ -tensor, we have  $\text{SOL}(0, \mathcal{A}(\omega_0)) = \{0\}$ . Also,  $S^* \subseteq \bigcap_{\omega \in \Omega} \text{SOL}(0, \mathcal{A}(\omega), \Omega)$ . Therefore  $S^* = \{0\}$ . Hence the result.  $\square$

Here we prove the necessary and sufficient conditions for the error bound of solution set to be level bounded in terms of residual function. To solve the semi-infinite tensor complementarity problem is equivalent to finding  $x \in \mathbb{R}^n$  such that  $x \in \text{SOL}(q(\omega), \mathcal{A}(\omega))$  for all  $\omega \in \Omega$ . However, in many cases, it is possible to obtain  $x \in \text{SOL}(q(\omega), \mathcal{A}(\omega))$  for some  $\omega$ . In this case, it is important to provide a quantitative measure of the closeness of each  $x \in \mathbb{R}^n$  to each individual set  $\text{SOL}(q(\omega), \mathcal{A}(\omega))$  in terms of some residual functions  $r(x)$ . In other words, we find  $c > 0$  such that

$$\text{dist}(x, \text{SOL}(q(\omega), \mathcal{A}(\omega))) \leq cr(x), \quad \forall \omega \in \Omega, \quad \forall x \in \mathbb{R}^n$$

which is equivalent to

$$\max_{\omega \in \Omega} \text{dist}(x, \text{SOL}(q(\omega), \mathcal{A}(\omega))) \leq cr(x), \quad \forall x \in \mathbb{R}^n.$$

Here  $c$  is said to be weak error bound. The importance of weak error bound is that the solution  $S^*$  is need not be nonempty as required in case of error bound.

**THEOREM 3.6:** Consider the  $\text{SITCP}(q(\omega), \mathcal{A}(\omega), \Omega)$ . Suppose  $\Omega$  is compact and  $\mathcal{A}(\omega)$  and  $q(\omega)$  are continuous. Then the residual function,  $r(x) = \max_{\omega \in \Omega} \|\min(x, \mathcal{A}(\omega)x + q(\omega))\|^2$  is level bounded if and only if the tensor  $\mathcal{A}(\omega)$  is a semi-infinite  $R_0$ -tensor relative to  $\Omega$ .

*Proof.* Only if part: We prove the first result by contrapositive method. By this approach we first assume that  $r(x)$  is not level bounded. Then  $\exists$  a sequence  $\{x_n\} \mapsto \infty$  as  $n \mapsto \infty$ ,  $\{r(x_n)\}$  is bounded. We assume that  $\frac{x_n}{\|x_n\|^{m-1}}$  converge to the limit  $x_0$  with  $\|x_0\| = 1$ . Taking into account the continuity of  $q(\omega)$  and  $\mathcal{A}(\omega)$  and the compactness of  $\Omega$ , we see that  $r(x)$  is continuous and  $q(\omega)$  is bounded on  $\Omega$ . Hence,  $\lim_{n \rightarrow \infty} \frac{r(x_n)}{\|x_n\|^{m-1}} = 0$  and  $\lim_{n \rightarrow \infty} \frac{q(\omega)}{\|x_n\|^{m-1}} = 0$  for all  $\omega \in \Omega$ . Now

$$\frac{r(x_n)}{\|x_n\|^{2(m-1)}} = \max_{\omega \in \Omega} \left\| \min \left\{ \frac{x_n}{\|x_n\|^{m-1}}, \frac{\mathcal{A}(\omega)x + q(\omega)}{\|x_n\|^{m-1}} \right\} \right\|^2. \quad (3.18)$$

Taking limit of both sides of the equation (3.18) as  $n \mapsto \infty$  we obtain,

$$\max_{\omega \in \Omega} \left\| \min \left\{ x_0, \mathcal{A}(\omega)x_0^{m-1} \right\} \right\|^2 = 0.$$

This means that a nonzero vector  $x_0$  is a solution of  $\text{SITCP}(0, \mathcal{A}(\omega), \Omega)$ . Hence  $\mathcal{A}(\omega)$  is not a semi-infinite  $R_0$ -tensor. This completes the proof.

If part: Suppose on the contrary that the  $\text{SITCP}(0, \mathcal{A}(\omega), \Omega)$  has a nonzero vector  $x$  as a solution. Let  $I(x) = \{i : x_i = 0\}$  and  $J(x) = \{i : x_i > 0\}$ . The compactness of  $\Omega$  and the continuity of  $q$  ensures that  $q(\omega)$  is bounded on  $\Omega$ . Thus there exists a scalar  $K > 0$  such that, for any  $k \geq K$ ,

$$kx_i \geq q_i(\omega) \text{ for all } \omega \in \Omega \text{ and } i \in J(x). \quad (3.19)$$

Given any  $k \geq K$ , we have

$$\begin{aligned} r(kx) &= \max_{\omega \in \Omega} \|\min(kx, k^{m-1}\mathcal{A}(\omega)x^{m-1} + q(\omega))\|^2 \\ &\leq \sum_{i=1}^n \max_{\omega \in \Omega} \{\min(kx_i, (k^{m-1}\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega))\}^2 \end{aligned} \quad (3.20)$$



Now we consider the following cases.

Case-1. We consider the case when  $i \in J(x)$ . Then  $((\mathcal{A}(\omega)x^{m-1})_i = 0$ . It follows from (3.19) that

$$\max_{\omega \in \Omega} \{\min\{kx_i, k^{m-1}(\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega)\}\}^2 = \max_{\omega \in \Omega} q_i(\omega)^2. \quad (3.21)$$

Case-2. We consider the case when  $i \in I(x)$ . If  $k^{m-1}(\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega) \geq 0$ , we have

$$\{\min(kx_i, k^{m-1}(\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega))\}^2 = 0. \quad (3.22)$$

If  $k^{m-1}(\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega) < 0$ , then by the fact  $q_i(\omega) \leq k^{m-1}(\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega) < 0$  we obtain

$$\{\min(kx_i, k^{m-1}(\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega))\}^2 \leq q_i(\omega)^2. \quad (3.23)$$

Thus combining (3.22) and (3.23) we have

$$\max_{\omega \in \Omega} [\min(kx_i, k^{m-1}(\mathcal{A}(\omega)x^{m-1})_i + q_i(\omega))]^2 \leq \max_{\omega \in \Omega} q_i(\omega)^2. \quad (3.24)$$

Putting the facts (3.20), (3.21) and (3.24) together, it follows that

$$r(kx) \leq \sum_{i=1}^n \max_{\omega \in \Omega} q_i(\omega)^2 < \infty$$

for all  $k \geq K$ . This contradicts the level boundedness of  $r(x)$ .  $\square$

## 4 Conclusion

In this paper we introduce semi-infinite tensor complementarity problem to accommodate more realistic situation of the problem. We show that the solution set of semi-infinite tensor complementarity problem exists with some assumption. An example is illustrated in detail to establish the result. We establish a connection between semi-infinite tensor and its equivalent tensors to obtain the solution of semi-infinite tensor complementarity problem. Finally, we show that the error bound of the solution set is level bounded in terms of residual function.

## 5 Acknowledgment

The author R. Deb is grateful to the Council of Scientific & Industrial Research (CSIR), India, for providing financial assistance through the Junior Research Fellowship program.

## References

- [1] Xue-Li Bai, Zheng-Hai Huang, and Yong Wang. Global uniqueness and solvability for tensor complementarity problems. *Journal of Optimization Theory and Applications*, 170(1):72–84, 2016.

- [2] Qihong Chen, Delin Chu, and Roger CE Tan. Optimal control of obstacle for quasi-linear elliptic variational bilateral problems. *SIAM journal on control and optimization*, 44(3):1067–1080, 2005.
- [3] A K Das. Properties of some matrix classes based on principal pivot transform. *Annals of Operations Research*, 243(1):375–382, 2016.
- [4] A K Das, R Jana, and Deepmala. On generalized positive subdefinite matrices and interior point algorithm. In *International Conference on Frontiers in Optimization: Theory and Applications*, pages 3–16. Springer, 2016.
- [5] A K Das, R Jana, and Deepmala. Finiteness of criss-cross method in complementarity problem. In *International Conference on Mathematics and Computing*, pages 170–180. Springer, 2017.
- [6] A K Das, R Jana, and Deepmala. Invex programming problems with equality and inequality constraints. *Transactions of A. Razmadze Mathematical Institute*, 172(3):361–371, 2018.
- [7] A K Das, R Jana, and Deepmala. Some aspects on solving transportation problem. *arXiv preprint arXiv:1808.05351*, 2018.
- [8] R Deb and A K Das. More on semipositive tensor and tensor complementarity problem. In *International Conference on Mathematics and Computing*, pages 147–156. Springer, 2023.
- [9] A Dutta and A K Das. On some properties of k-type block matrices in the context of complementarity problem. In *Mathematics and Computing: ICMC 2022, Vellore, India, January 6–8*, pages 143–154. Springer, 2023.
- [10] A Dutta, R Jana, and A K Das. On column competent matrices and linear complementarity problem. In *Proceedings of the Seventh International Conference on Mathematics and Computing*, pages 615–625. Springer Singapore, 2022.
- [11] Francisco Facchinei, Andreas Fischer, and Veronica Piccialli. Generalized nash equilibrium problems and newton methods. *Mathematical Programming*, 117(1-2):163–194, 2009.
- [12] Zheng-Hai Huang and Liqun Qi. Formulating an n-person noncooperative game as a tensor complementarity problem. *Computational Optimization and Applications*, 66(3):557–576, 2017.
- [13] R Jana, A K Das, and A Dutta. On hidden  $Z$ -matrix and interior point algorithm. *Opsearch*, 56(4):1108–1116, 2019.
- [14] R Jana, A K Das, and V N Mishra. Iterative descent method for generalized leontief model. *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences*, 91(2):237–244, 2021.
- [15] R Jana, A K Das, and S Sinha. On processability of Lemke’s algorithm. *Applications & Applied Mathematics*, 13(2), 2018.

- [16] R Jana, A K Das, and S Sinha. On semimonotone star matrices and linear complementarity problem. *arXiv preprint arXiv:1808.00281*, 2018.
- [17] R Jana, A Dutta, and A K Das. More on hidden  $Z$ -matrices and linear complementarity problem. *Linear and Multilinear Algebra*, 69(6):1151–1160, 2021.
- [18] S R Mohan, S K Neogy, and A K Das. More on positive subdefinite matrices and the linear complementarity problem. *Linear Algebra and its Applications*, 338(1-3):275–285, 2001.
- [19] S R Mohan, S K Neogy, and A K Das. On the classes of fully copositive and fully semimonotone matrices. *Linear Algebra and its Applications*, 323(1-3):87–97, 2001.
- [20] S R Mohan, S K Neogy, and A K Das. A note on linear complementarity problems and multiple objective programming. *Mathematical programming*, 100(2):339–344, 2004.
- [21] P Mondal, S Sinha, S K Neogy, and A K Das. On discounted ARAT semi-markov games and its complementarity formulations. *International Journal of Game Theory*, 45(3):567–583, 2016.
- [22] James Munkres. *Topology james munkres second edition*.
- [23] S K Neogy, R B Bapat, and A K Das. Optimization models with economic and game theoretic applications. *Annals of Operations Research*, 243(1):1–3, 2016.
- [24] S K Neogy and A K Das. Linear complementarity and two classes of structured stochastic games. *Operations Research with Economic and Industrial Applications: Emerging Trends*, eds: SR Mohan and SK Neogy, Anamaya Publishers, New Delhi, India, pages 156–180, 2005.
- [25] S K Neogy and A K Das. On almost type classes of matrices with  $Q$ -property. *Linear and Multilinear Algebra*, 53(4):243–257, 2005.
- [26] S K Neogy and A K Das. Principal pivot transforms of some classes of matrices. *Linear Algebra and its Applications*, 400:243–252, 2005.
- [27] S K Neogy and A K Das. Some properties of generalized positive subdefinite matrices. *SIAM Journal on Matrix Analysis and Applications*, 27(4):988–995, 2006.
- [28] S K Neogy and A K Das. *Mathematical programming and game theory for decision making*, volume 1. World Scientific, 2008.
- [29] S K Neogy and A K Das. On singular  $N_0$ -matrices and the class  $Q$ . *Linear Algebra and its Applications*, 434(3):813–819, 2011.
- [30] S K Neogy and A K Das. On weak generalized positive subdefinite matrices and the linear complementarity problem. *Linear and Multilinear Algebra*, 61(7):945–953, 2013.
- [31] S K Neogy, A K Das, and R B Bapat. *Modeling, computation and optimization*, volume 6. World Scientific, 2009.
- [32] S K Neogy, A K Das, and A Gupta. Generalized principal pivot transforms, complementarity theory and their applications in stochastic games. *Optimization Letters*, 6(2):339–356, 2012.

- [33] S K Neogy, A K Das, S Sinha, and A Gupta. On a mixture class of stochastic game with ordered field property. In *Mathematical programming and game theory for decision making*, pages 451–477. World Scientific, 2008.
- [34] Jia-Yu Shao. A general product of tensors with applications. *Linear Algebra and its Applications*, 439(8):2350–2366, 2013.
- [35] Jiayu Shao and Lihua You. On some properties of three different types of triangular blocked tensors. *Linear Algebra and its Applications*, 511:110–140, 2016.
- [36] Yisheng Song and Liqun Qi. Properties of some classes of structured tensors. *Journal of Optimization Theory and Applications*, 165(3):854–873, 2015.
- [37] Yisheng Song and Liqun Qi. Properties of tensor complementarity problem and some classes of structured tensors. *Annals of Applied Mathematics*, 2017.
- [38] Yisheng Song and Gaohang Yu. Properties of solution set of tensor complementarity problem. *Journal of Optimization Theory and Applications*, 170(1):85–96, 2016.
- [39] RT Rockafellar RJ-B Wets. *Variational analysis* spring-verlag berlin. 1998.
- [40] Mengmeng Zheng, Ying Zhang, and Zheng-Hai Huang. Global error bounds for the tensor complementarity problem with a p-tensor. *Journal of Industrial & Management Optimization*, 15(2):933, 2019.
- [41] JC Zhou, NH Xiu, and Jein-Shan Chen. Solution properties and error bounds for semi-infinite complementarity problems. *Journal of Industrial and Management Optimization*, 9(1):99–115, 2013.