




Comment on “Multiparty quantum mutual information: An alternative definition”

Jaehak Lee ^{1,2,*}, Gibeom Noh,³ Changsuk Noh ³ and Jiyong Park ^{4,†}

¹*Department of Physics and Astronomy, Seoul National University, Seoul 08826, Korea*

²*Center for Quantum Information, Korea Institute of Science and Technology, Seoul 02792, Korea*

³*Department of Physics, Kyungpook National University, Daegu 41566, Korea*

⁴*School of Basic Sciences, Hanbat National University, Daejeon 34158, Korea*

(Dated: January 2, 2024)

We show that, contrary to the claim by Kumar [Phys. Rev. A **96**, 012332 (2017)], the quantum dual total correlation of an n -partite quantum state cannot be represented as the quantum relative entropy between $n - 1$ copies of the quantum state and the product of n different reduced quantum states for $n \geq 3$. Specifically, we argue that the latter fails to yield a finite value for generalized n -partite Greenberger-Horne-Zeilinger states.

In [1], a quantum version of the dual total correlation [2] for an n -partite quantum state ρ was proposed as

$$I_n(\rho) \equiv \sum_{k=1}^n S(\rho_{\bar{k}}) - (n-1)S(\rho), \quad (1)$$

where $S(\tau) = -\text{tr}(\tau \log \tau)$ is the von Neumann entropy of τ and $\rho_{\bar{k}} = \text{tr}_k \rho$ denotes the $(n-1)$ -partite quantum state obtained by taking the partial trace on the k th party of ρ . In addition, it was claimed that Eq. (1) can be represented as

$$I_n(\rho) = J_n(\rho) \equiv S(\rho^{\otimes(n-1)} \| \bigotimes_{k=1}^n \rho_{\bar{k}}), \quad (2)$$

where $\rho^{\otimes j}$ represents j copies of ρ and $S(\tau \| \sigma)$ is the quantum relative entropy of τ with respect to σ [3],

$$S(\tau \| \sigma) = \begin{cases} \text{tr}(\tau \log \tau) - \text{tr}(\tau \log \sigma) & \text{if } \text{supp}(\tau) \subseteq \text{supp}(\sigma) \\ \infty & \text{otherwise,} \end{cases} \quad (3)$$

where the support of ω is the Hilbert space spanned by the eigenstates of ω with non-zero eigenvalue [4].

It is well known that $I_2(\rho)$ can be represented as the quantum relative entropy of a global quantum state ρ with respect to the product of two local quantum states ρ_1 and ρ_2 , i.e., $I_2(\rho) = S(\rho \| \rho_1 \otimes \rho_2)$ [5]. We emphasize that the parties between global and local quantum states must be properly matched to avoid the infinity in Eq. (3). While $\text{supp}(\rho) \subseteq \text{supp}(\rho_1 \otimes \rho_2)$ is always met, $\text{supp}(\rho) \subseteq \text{supp}(\rho_2 \otimes \rho_1)$ is not satisfied in general. It leads to a discrepancy between $I_2(\rho) = S(\rho \| \rho_1 \otimes \rho_2)$ and $J_2(\rho) = S(\rho \| \rho_2 \otimes \rho_1)$. For instance, we have $I_2(\rho) = 0$ and $J_2(\rho) = \infty$ for $\rho = |\psi_1\rangle\langle\psi_1| \otimes |\psi_2\rangle\langle\psi_2|$ satisfying $\langle\psi_1|\psi_2\rangle = 0$.

One can generalize this observation to the case of $n \geq 3$. Looking into the order of the parties in $\rho^{\otimes(n-1)}$ and $\bigotimes_{k=1}^n \rho_{\bar{k}}$, one immediately sees that they are mismatched

for all n because the former and the latter start with the first and second parties of ρ , respectively. Similar to the case of $n = 2$, if we look into a product state $\rho = \bigotimes_{k=1}^n |\psi_k\rangle\langle\psi_k|$ satisfying $\langle\psi_i|\psi_j\rangle = \delta_{i,j}$, we obtain $I_n(\rho) = 0$ and $J_n(\rho) = \infty$.

One may think that rearranging the parties of quantum states can resolve the support condition problem. It is possible for $n = 2$ but impossible for $n \geq 3$. We show this by investigating the case of $\rho = |\phi\rangle\langle\phi|$ with $|\phi\rangle = \sqrt{p}|0\rangle^{\otimes n} + \sqrt{1-p}|1\rangle^{\otimes n}$. The state of $|\phi\rangle^{\otimes(n-1)}$ is represented by the superposition of the basis states having the multiples of n , i.e., $\{0, n, 2n, \dots, n(n-1)\}$, copies of $|1\rangle$. On the other hand, the eigenstates of $\bigotimes_{k=1}^n \rho_{\bar{k}}$ have multiples of $n-1$, i.e., $\{0, (n-1), 2(n-1), \dots, n(n-1)\}$, copies of $|1\rangle$. As n and $n-1$ are coprimes, $|\phi\rangle^{\otimes(n-1)}$ is orthogonal to the eigenstates of $\bigotimes_{k=1}^n \rho_{\bar{k}}$ except for $|0\rangle^{\otimes n(n-1)}$ and $|1\rangle^{\otimes n(n-1)}$, which means $\text{supp}(\rho^{\otimes(n-1)}) \not\subseteq \text{supp}(\bigotimes_{k=1}^n \rho_{\bar{k}})$ for $n \geq 3$. Importantly, the observation just described remains valid even if we rearrange the parties. Therefore, we have $I_n(\rho) = -n[p \log p + (1-p) \log(1-p)]$ and $J_n(\rho) = \infty$ independent of the matching structure between $\rho^{\otimes(n-1)}$ and $\bigotimes_{k=1}^n \rho_{\bar{k}}$.

In [1], the equivalence between Eqs. (1) and (2), i.e., $I_n(\rho) = J_n(\rho)$, was invoked in order to prove the non-negativity and monotonicity of $I_n(\rho)$ under a partial trace and completely positive maps. In addition, the proof for non-negativity was reproduced in [6]. Our counterexample invalidates these proofs which rely on the facts that the relative entropy is non-negative and non-increasing under completely positive and trace-preserving (CPTP) maps. Interestingly, however, one can find alternative proofs for the non-negativity and monotonicity in [7]. In Ref. [7], $I_n(\rho)$ was proposed as a quantum secrecy monotone, and was shown to be non-negative and non-increasing under local CPTP maps, owing to strong subadditivity of the conditional quantum mutual entropy. Therefore, $I_n(\rho)$ is a suitable monotonic measure of multi-partite correlations, while $J_n(\rho)$ is not.

*Electronic address: jaehak.lee.201@gmail.com

†Electronic address: jiyong.park@hanbat.ac.kr

J.L. and J.P. acknowledge support from the National Research Foundation of Korea funded by the government

of Korea (MSIT) (Grants No. NRF-2022M3K4A1097117 and No. NRF-2019R1G1A1002337, respectively).

-
- [1] A. Kumar, Multiparty quantum mutual information: An alternative definition, *Phys. Rev. A* **96**, 012332 (2017).
 - [2] T. S. Han, Nonnegative entropy measures of multivariate symmetric correlations, *Inf. Control* **36**, 133 (1978).
 - [3] M. M. Wilde, *Quantum Information Theory* (Cambridge University Press, Cambridge, 2013).
 - [4] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2010).
 - [5] V. Vedral, The role of relative entropy in quantum information theory, *Rev. Mod. Phys.* **74**, 197 (2002).
 - [6] Y. Guo and L. Huang, Complete monogamy of the multipartite quantum mutual information, *Phys. Rev. A* **107**, 042409 (2023).
 - [7] N. J. Cerf, S. Massar, and S. Schneider, Multipartite classical and quantum secrecy monotones, *Phys. Rev. A* **66**, 042309 (2002).