Constant-roll inflation with tachyon field in the holographic braneworld

Marko Stojanovic^{*1}, Neven Bilić^{†2}, Dragoljub D. Dimitrijevic^{‡3}, Goran S. Djordjevic^{§3}, and Milan Milosevic^{¶3}

¹Faculty of Medicine, University of Niš, Serbia ²Division of Theoretical Physics, Rudjer Bošković Institute, Zagreb, Croatia ³Department of Physics, University of Niš, Serbia

January 2, 2024

Abstract

We study inflation driven by the tachyon field in the holographic braneworld by assuming the second slow-roll parameter η is constant. The parameter η can be either defined by the tachyon scalar field and the Hubble parameter or by the Hubble parameter only. By assuming a constant η , we derive and numerically solve a differential equation for the Hubble expansion rate. We calculate numerically the scalar spectral index and the tensor-to-scalar ratio. We confront the results with the observational data and find some constraints on the free model parameters. The swampland conjectures are discussed in the context of the constant-roll inflation, with some accent on the holographic model.

1 Introduction

Inflation is the most promising candidate for solving the horizon problem and other related problems of the standard Big Bang cosmology [1], by demanding an extremely short period in the early universe during which the universe rapidly expands. Moreover, the quantum fluctuations generated during inflation lead to the primordial density perturbation responsible for the large-scale structure formation.

An outstanding achievement has been made in the theoretical prediction of the inflation parameters, assuming that the inflation is governed by a single scalar field (the inflaton), whose origin is still unknown. One of the popular classes of inflationary models is based on the dynamics of the tachyon field originating in M or string theory [2]. It is known [3] that the decay of unstable D-branes in string theory is followed by a production of pressureless gas with non-zero energy density, which is required for a phase of inflationary expansion. This is one of main motivations for applications of the tachyon field in the theory of inflation.

The constant-roll inflation is based on the assumption that during inflation, the second slow-roll parameter η is an arbitrary constant and not necessarily small [4]. This formalism has been applied in many different scenarios of inflation [5–13]. In this work, for the first time, we apply the formalism of the constant-roll inflation to the holographic second Randall-Sundrum (RSII) model with a tachyon field [14, 15]. This paper is an extension of our previous work in which we studied the constant-roll tachyon inflation in the RSII cosmology [16]. We have shown that the observational parameters obtained in the constant-roll approach for the RSII model are in better agreement with observational data than those obtained in the slow-roll approximation [17].

Motivated by this result, we examine whether similar behavior will occur in the tachyon model in the holographic braneworld. The dynamics of the model considered here are based on a scenario in which the brane, with an effective tachyon field, is located at the boundary of a 5-dim asymptotically Anti-de Sitter (AdS_5) space. This type of model we refer to as the holographic braneworld [18]. The effective

^{*}marko.stojanovic@pmf.edu.rs

[†]bilic@irb.hr

[‡]ddrag@pmf.ni.ac.rs

[§]gorandj@junis.ni.ac.rs

[¶]milan.milosevic@pmf.edu.rs

four-dimensional Einstein equations on the holographic boundary of AdS_5 yield a modified Friedmann equation with a quartic term $\propto H^4$, where H is Hubble parameter [18, 19]. Due to this modification in the Friedmann equation, it is interesting to consider the constant-roll inflation in the (holographic) cosmology on the AdS_5 boundary. The constant-roll inflation has recently been analyzed, assuming that inflation is driven by (generalized) holographic dark energy [20, 21]. It is important to stress that these models are not equivalent to our model.

The paper is organized as follows. In Section 2, we give a brief review of the tachyon dynamics in the holographic braneworld. The scenario of the constant-roll inflation in the holographic RSII model is discussed in Section 3. In Section 4, we present the details of our numerical results. In Section 5, we discuss the swampland criteria for the holographic constant-roll RSII inflation model. A summary and conclusions are given in Section 6.

2 The tachyon dynamics in the holographic braneworld

In this section, we briefly introduce the basic property of the tachyon dynamics in the holographic braneworld. In a braneworld scenario, our universe is a brane moving in the higher dimensional bulk, with only gravity allowed to propagate in the bulk. In the RSII model, two 3+1 dimensional branes with opposite tensions are embedded in a 4+1 spacetime with AdS₅ geometry. The observers reside on the positive tension brane, and the negative tension brane is pushed off to infinity. If the brane is located at the boundary of an AdS₅ space, the effective Einstein equations on the brane can be obtained in the framework of anti-de Sitter/conformal field theory (AdS/CFT) correspondence and holographic renormalization [18]. The effective Einstein equations yield the holographic Friedmann equations. For a spatially flat FLRW geometry on the brane, the holographic Friedmann equations are [14]

$$h^2 - \frac{1}{4}h^4 = \frac{\kappa^2}{3}\ell^4\rho,$$
 (1)

$$\dot{h}\left(1-\frac{1}{2}h^2\right) = -\frac{\kappa^2}{2}\ell^3(p+\rho),$$
(2)

where p and ρ are the pressure and the density of the fluid associated with the field on the brane and $h \equiv \ell H$ is a dimensionless Hubble expansion rate. The overdot in (2) denotes a derivative with respect to time measured in units of ℓ . The fundamental coupling parameter κ is related to the AdS₅ curvature radius ℓ and the four-dimensional Newton constant $G_{\rm N}$

$$\kappa^2 = \frac{8\pi G_{\rm N}}{\ell^2}.\tag{3}$$

In (2), we omit the contribution of "dark radiation" [22]. The physically acceptable solution of the equation (1), as a quadratic equation for h^2 , is of the form

$$h^2 = 2\left(1 - \sqrt{\frac{\kappa^2}{3}\ell^4\rho}\right). \tag{4}$$

In the low-density limit ($\kappa^2 \ell^4 \rho \ll 1$), the equation (4) reduces to the standard Friedmann equation, ensures that the holographic cosmology reduces to the standard cosmology. According to (2), the expansion rate is a monotonously decreasing function of time. In this model, the evolution is constrained by the physical range of the expansion rate $0 \leq h^2 \leq 2$. The evolution of the universe starts with an initial $h_i < \sqrt{2}$ and with negative \dot{h}_i .

For a fluid on the brane described by the tachyon field, the tachyon Lagrangian of the Dirac-Born-Infeld (DBI) type is of the form

$$\mathcal{L} = -\ell^{-4} V(\theta/\ell) \sqrt{1 - g^{\mu\nu} \theta_{,\mu} \theta_{,\nu}}, \qquad (5)$$

where the tachyon field θ has units of ℓ . The tachyonic potential $V(\theta/\ell)$ satisfies the following properties

$$V(0) = \text{const}, \quad V_{,\theta}(\theta > 0) < 0, \quad V(|\theta| \to \infty) \to 0, \tag{6}$$

where the subscript , θ denotes a derivative with respect to θ . The pressure p and energy density ρ are the components of the energy-momentum tensor $T_{\mu\nu}$

$$T_{\mu\nu} = (\rho + p)u_{\mu}u_{\nu} - pg_{\mu\nu}, \tag{7}$$

where the components of the four velocity are $u_{\mu} \equiv \partial_{\mu}\theta / \sqrt{\partial_{\mu}\theta \partial^{\mu}\theta}$ and $u_{\alpha}u^{\alpha} = 1$. Assuming spatial isotropy and homogeneity at the background, $\theta = \theta(t)$, from

$$T_{\mu\nu} \equiv \frac{2}{\sqrt{-g}} \frac{\delta \sqrt{-g} \mathcal{L}}{\delta g^{\mu\nu}},\tag{8}$$

one finds

$$p = -\ell^{-4}V\sqrt{1-\dot{\theta}^2},\tag{9}$$

$$\rho = \frac{\ell^{-4}V}{\sqrt{1 - \dot{\theta}^2}}.$$
(10)

The model is presented in more detail in ref. [14]. In the following, we will analyze the constant-roll inflation based on this model, with the slow-roll parameter η being constant.

3 Constant-roll inflation

The dominant inflationary scenario nowadays, known as the slow-roll inflation, is based on the assumption that the inflaton field varies very slowly and inflation lasts long enough to overcome some problems in the standard Bing Bang cosmology. In the slow-roll inflation models, the slow-roll parameters

$$\epsilon = -\frac{\ell \dot{h}}{h^2},\tag{11}$$

$$\eta = \frac{\ell \ddot{\theta}}{h \dot{\theta}},\tag{12}$$

are small. Inflation ends when the parameter ϵ crosses unity. The modification of the slow-roll inflation $(\eta \approx 0)$ or ultra slow-roll inflation $(\eta = -3)$ leads to the constant-roll inflation defined by the condition [4]

$$\eta = \text{const.}$$
 (13)

In this work, we apply the condition (13) to the tachyon dynamics in the holographic braneworld. Combining Friedmann's equations, (1) and (2), with (9) and (10) we obtain

$$\dot{\theta} = -\frac{2\ell}{3} \frac{h_{,\theta}}{h^2} \frac{1 - \frac{1}{2}h^2}{1 - \frac{1}{4}h^2},\tag{14}$$

where according to the Hamilton-Jacobi formalism the expansion rate h is taken as a function of field θ , i.e. $h = h(\theta)$, yielding $\dot{h} = h_{,\theta}\dot{\theta}$. By taking the time derivative of (14) we obtain

$$\ddot{\theta} = -\frac{2\ell}{3} \frac{1 - \frac{1}{2}h^2}{1 - \frac{1}{4}h^2} \frac{1}{h^3} \left(hh_{,\theta\theta} - 2h_{,\theta}^2\right) \dot{\theta} + \frac{\ell}{3} \frac{h_{,\theta}^2}{h} \frac{1}{(1 - \frac{1}{4}h^2)^2} \dot{\theta}.$$
(15)

Then, using the condition (13), the equations (14) and (15), and replacing them in the equation (12) we derive the differential equation for the expansion rate

$$hh_{,\theta\theta} - 2h_{,\theta}^2 \left(1 + \frac{h^2}{4(1 - \frac{1}{2}h^2)(1 - \frac{1}{4}h^2)} \right) + \frac{3}{2\ell^2} \eta h^4 \frac{1 - \frac{1}{4}h^2}{1 - \frac{1}{2}h^2} = 0.$$
(16)

The equation (16) is more complicated than the corresponding equation in the tachyon constant-roll model in standard cosmology (derived from the Einstein-Hilbert action) and will also be solved numerically.

One of the main advantages of constant-roll inflation is that there is no need to specify the type of potential. The potential can be reconstructed when the dynamics of the model is known. Combining (1), (10) and (14), we obtain the potential as an implicit function of the tachyon field

$$V = \frac{3}{\kappa^2} h^2 (1 - \frac{1}{4}h^2) \sqrt{1 - \frac{4\ell^2}{9} \frac{h_{,\theta}^2}{h^4} \left(\frac{1 - \frac{1}{2}h^2}{1 - \frac{1}{4}h^2}\right)^2}.$$
(17)

In the limit when $h^2 \ll 1$, the equations (16) and (17) agree with those derived in tachyon constant-roll model in standard cosmology [13].

There are plenty of definitions of the slow-roll parameters. Besides the most common, given by (11) and (12), it proves advantageous to define the slow-roll parameters recursively

$$\varepsilon_0 \equiv h_*/h,\tag{18}$$

$$\varepsilon_{i+1} \equiv \frac{d\ln|\varepsilon_i|}{dN}, \quad i \ge 0, \tag{19}$$

where h_* is the expansion rate at an arbitrarily chosen time and N is the number of e-folds defined as

$$N \equiv \frac{1}{\ell} \int_{t_i}^{t_f} h dt, \tag{20}$$

where t_i and t_f denote the times of the beginning and the end of inflation, respectively. To solve the most important problems in the early universe, the number of e-folds during inflation should be $N \simeq 60$. From the definition for the parameters ε_i one obtains

$$\varepsilon_{i+1} = \frac{\ell \dot{\varepsilon}_i}{h \varepsilon_i}.$$
(21)

The relationships between $\varepsilon_1 \equiv \epsilon$, ε_2 and η depend on the form of the Friedmann equation. Using (21) together with (14), for the first two slow-roll parameters ε_i we obtain

$$\varepsilon_1 = -\ell \frac{\dot{h}}{h^2} = \frac{2\ell^2}{3} \frac{h_{,\theta}^2}{h^4} \frac{1 - \frac{1}{2}h^2}{1 - \frac{1}{4}h^2},\tag{22}$$

$$\varepsilon_2 = \ell \frac{\dot{\varepsilon}_1}{h\varepsilon_1} = \eta - \varepsilon_1 \left(\frac{h_{,\theta\theta}}{h_{,\theta}^2} h - 2 \right).$$
(23)

Substituting $h_{,\theta\theta}$ from (16) into (23), ε_2 takes the following simple form

$$\varepsilon_2 = 2\eta - \frac{h^2}{2(1 - \frac{1}{2}h^2)(1 - \frac{1}{4}h^2)}\varepsilon_1.$$
(24)

In our model, unlike the constant-roll inflation model in the standard cosmology [23], the expression for ε_2 contains the additional term proportional to h^2 , causing the change of ε_2 during the time. The parameter ε_3 is equal to

$$\varepsilon_3 = \frac{1}{\varepsilon_2} \left(\frac{h^2 (1 - \frac{1}{8}h^4)}{(1 - \frac{1}{2}h^2)^2 (1 - \frac{1}{4}h^2)^2} \varepsilon_1^2 - \frac{h^2}{2(1 - \frac{1}{2}h^2)(1 - \frac{1}{4}h^2)} \varepsilon_1 \varepsilon_2 \right).$$
(25)

In the limit when $h^2 \ll 1$, the contribution of the additional term can be neglected, i.e., $\varepsilon_2 \simeq 2\eta$, causes that the parameter ε_2 is almost constant and the parameter ε_3 tends to zero.

4 Numerical calculations

In this section, we present the results for the observational parameters and the number of e-folds, obtained by numerically solving the equation for the Hubble expansion rate, equation (16), for fixed η . This equation is a second-order differential equation for the tachyon field θ . To solve this equation numerically, we need to choose initial conditions, i.e., to fix the values of h and $h_{,\theta}$ for a chosen value θ . Let these be the values $h_i \equiv h(\theta_i)$ and $h_{,\theta_i} \equiv h_{,\theta}(\theta_i)$ that correspond to the value of the field θ_i at the time of the beginning of inflation ($\theta_i = \theta(t_i)$). For this specific choice, we will fix the initial conditions from the values of the observational parameters. As we will see, this approach allows us to estimate the range of parameter η in the model with the corresponding predictions that should be in good agreement with observational data. For numerical solving of (16), bearing in mind form of (14), without loss of generality, we can set $t_i = 0$ and $\theta_i = 0$.

The observational parameters, the scalar spectral index n_s , and the tensor-to-scalar ratio r have been calculated in Ref. [24] for the tachyon model in the holographic braneworld up to the second order in the slow-roll parameters

$$r = 16\varepsilon_{1i} \left[1 + C\varepsilon_{2i} - \frac{2(2-h^2)}{3(4-h^2)}\varepsilon_{1i} \right],$$
(26)

$$n_{\rm s} = 1 - 2\varepsilon_{1\rm i} - \varepsilon_{2\rm i} - \left(2 - \frac{8h^2}{3(4-h^2)^2}\right)\varepsilon_{1\rm i}^2 - \left(3 + 2C - \frac{2(2-h^2)}{3(4-h^2)}\right)\varepsilon_{1\rm i}\varepsilon_{2\rm i} - C\varepsilon_{2\rm i}\varepsilon_{3\rm i},\tag{27}$$

where $C \simeq -0.73$ and $\varepsilon_{ii} \equiv \varepsilon_i(\theta_i)$ are the values of the slow-roll parameters at the beginning of inflation . The expressions (26) and (27) at linear order in the slow-roll parameters ε_i agree with the expressions obtained in the standard scalar field inflation with a tachyon field (see e.g. ref. [23])

$$r \simeq 16\varepsilon_{1i},$$
 (28)

$$n_{\rm s} \simeq 1 - 2\varepsilon_{1\rm i} - \varepsilon_{2\rm i}.\tag{29}$$

The calculated parameters given in (26) and (27) may be confronted with the observed values from the Planck data [25]

$$n_{\rm s} = 0.9649 \pm 0.0042,\tag{30}$$

$$r < 0.056.$$
 (31)

Now, we calculate a functional relationship between the observational parameters, given by expressions at linear order in the slow-roll parameters, and the initial conditions h_i and $h_{,\theta_i}$. From (28) and (29), using (24), we obtain

$$\eta = \frac{1}{2} \left(1 - n_{\rm s} - \frac{r}{8} + \frac{h_{\rm i}^2}{(1 - \frac{1}{2}h_{\rm i}^2)(1 - \frac{1}{4}h_{\rm i}^2)} \frac{r}{32} \right).$$
(32)

The term proportional to h_i^2 , that results from (24), affects the value of the parameter η , which is another crucial difference in this model compared to the model in standard cosmology. Using (22), the expression (28) can be rewritten as

$$h_{,\theta i} = -\sqrt{\frac{3}{32}} r \frac{1 - \frac{1}{4} h_i^2}{1 - \frac{1}{2} h_i^2} h_i^4, \tag{33}$$

where we have chosen the minus sign, according to the demand that $\dot{h} < 0$. Clearly, for each choice of $n_{\rm s}$, r and $h_{\rm i}$ the parameter η and the value $h_{,\theta i}$ are fixed by (32) and (33).

4.1 Numerical results

Now, we present the results for some fixed values of the parameters. In the following, we treat h_i , besides η , as an additional free parameter of the model, the value of which lies in the interval $0 < h_i < \sqrt{2}$.

In Fig. 1 we plot η and $h_{,\theta i}$ as function of h_i for fixed and arbitrary $n_s = 0.966$ and r = 0.022. As we shall see, for the values of the observational parameters, which are well inside the observational region, the model can provide the inflation stage with $N \approx 60$. We will use these values to illustrate further results. It can be seen from Fig. 1 that the value of parameter η changes significantly only for those



Figure 1: η versus h_i (left panel) and h_{θ_i} versus h_i (right panel) for fixed $n_s = 0.966$ and r = 0.022.

values of h_i which are near to the maximum value $h_{i \max} = \sqrt{2}$. For the other allowed values for h_i , the parameter η is almost constant and very close to the value obtained in the standard tachyon constant-roll inflation ($\eta \approx 0.0153$) [13]. As shown in Fig. 1, $|h_{\theta i}|$ increases with h_i which indicates that there may be a deviation in the duration of inflation (e-folds number) in our model compared to the tachyon constant-roll model in standard cosmology [13]. We will discuss those results in the next section.

In Fig. 2, we plot η as a function of h_i varying the values of the observational parameters. Again, the most significant range of the values of h_i is near its maximum, and the value of the η is more affected by the change of r than by the change of n_s .



Figure 2: η versus h_i . Left panel: r = const = 0.01, $n_s = 0.96$ (solid red line), $n_s = 0.965$ (dashed blue line), $n_s = 0.97$ (dash-dotted green line). Right panel: $n_s = \text{const} = 0.965$, r = 0.01 (solid red line), r = 0.02 (dashed blue line), r = 0.03 (dash-dotted green line).

In Fig. 3 we present a numerical solution to the equation (16) and the reconstructed potential, obtained for $n_{\rm s} = 0.966$, r = 0.022 and $h_{\rm i} = 1.4$. For this choice of the parameters, using expressions (32) and (33), we find $\eta = 0.0817$ and $h_{\theta i} = -0.4495$. It is obvious from Fig. 3 that the reconstructed potential has all properties of a tachyon potential, defined by (6).



Figure 3: The expansion rate (left panel) and the reconstructed potential (right panel) versus tachyon field in the model with $\eta = 0.0817$, $h_i = 1.4$ and $h_{,\theta i} = -0.4495$.

The expression (14) may be integrated numerically, yielding the time t as a function of θ . With its help, the time evolution of the expansion rate and the tachyon field can be plotted (Fig. 4).



Figure 4: The time evolution of the expansion rate (left panel) and the tachyon field (right panel) in the model with $\eta = 0.0817$, $h_i = 1.4$ and $h_{,\theta i} = -0.4495$.

The slow-roll parameters ε_1 and ε_2 as functions of θ are shown in Fig. 5.



Figure 5: The slow-roll parameters ε_1 and ε_2 versus tachyon field calculated numerically in the model with $\eta = 0.0817$, $h_i = 1.4$ and $h_{,\theta i} = -0.4495$.

The parameter ε_1 increases and exceeds unity (for $\theta_f = 415.5$), defining the end of inflation. As already mentioned, for $h^2 \ll 1$ the value ε_2 is almost constant. Furthermore, it is obvious from the shape of the curve in Fig. 5 that the parameter ε_2 changes significantly only at the beginning of inflation.

Now, we calculate the number of e-folds associated with the observational parameters, the values of which we choose according to the restrictions from the observational data. Using (14) the expression (20) becomes

$$N = \frac{1}{\ell} \int_{\theta_{i}}^{\theta_{f}} \frac{h}{\dot{\theta}} d\theta = -\frac{3}{2\ell^{2}} \int_{\theta_{i}}^{\theta_{f}} \frac{h^{3}}{h_{,\theta}} \frac{(1 - \frac{1}{4}h^{2})}{(1 - \frac{1}{2}h^{2})} d\theta,$$
(34)

where $\theta_{\rm f} = \theta(t_{\rm f})$ is fixed by $\varepsilon_1(\theta_{\rm f}) = 1$. To calculate the number of e-folds, we proceed as follows. For some chosen $n_{\rm s}$, r and $h_{\rm i}$, using (32) and (33), we calculate the parameter η and the initial value $h_{,\theta_{\rm i}}$. Then, we obtain $h(\theta)$ solving (15), for given η , with the initial conditions $h_{\rm i}$ and $h_{,\theta_{\rm i}}$. From the condition for the end of inflation ($\varepsilon_1(\theta_{\rm f}) = 1$) we find $\theta_{\rm f}$. Finally, we use numerical integration in expression (34) to find N. For $n_{\rm s} = 0.966$, r = 0.022 and $h_{\rm i} = 1.4$ the number of e-folds is N = 60.2.

In Fig. 6, we plot N as a function of h_i and as a function of η . Note that the change in the number N due to the variation of the parameter n_s is significant only for small values of h_i . This result is expected, considering how the parameter η depends on the observable parameters n_s and r (see Fig. 2). As mentioned earlier, the number of e-folds is also affected by the value of h_i . Clearly, as the value of the parameter h_i increases, the number of e-folds decreases. Therefore, the interval of h_i in which the model gives inflation with $N \simeq 60$ is narrower than the model allows (see Fig. 1). In addition, Fig. 6 indicates that the value of the parameter η should be close to $\eta \approx 0.08$ to have inflation with $N \simeq 60$. In comparison, the tachyon constant-roll inflation in standard cosmology for the values $n_s = 0.9615$ and



Figure 6: Theoretical prediction for the functions N of h_i (left panel) and N of η (right panel) for several fixed observational parameters. On the left panel, r = 0.02 (red line), 0.025 (blue line), 0.03 (green line), where $n_s = 0.96$ is represented by solid line and $n_s = 0.965$ by dashed line. The right panel shows the function N of η for randomly chosen observational parameters $n_s = 0.965$ and r = 0.02 (solid red line) and $n_s = 0.965$ and r = 0.02 (solid red line)

r = 0.0197, with $\eta = 0.0153$, predicts inflation with N = 121 [13]. In our model, the chosen values of

the observational parameters inside the allowed range of the Planck data, with $h_i = 1.4$ (and $\eta = 0.077$), correspond to N = 64.4.

4.2 Scalar spectral index and tensor-to-scalar ratio

Note that previous results for the number of e-folds were obtained using the functional relationship between the observational parameters and the initial conditions h_i and $h_{,\theta i}$, equations (32) and (33). More accurate results can be obtained using (26) and (27) and applying the standard approach based on calculating observation parameters n_s and r for a given e-folds number N, described in details in refs. [14], [15] and [17]. For randomly chosen η , h_i , and $h_{,\theta i}$, the equation (16) and the differential equation for the number of e-folds $dN/d\theta = h/(\ell\dot{\theta})$, are solved simultaneously from $\theta_i = 0$ to some arbitrary value θ_f which is large enough, to provide the end of inflation ($\varepsilon_1(\theta_f) = 1$). Then, demanding $N = N(\theta_i) - N(\theta_f)$ the value θ_i is obtained for given N. The value θ_i is then used to find ε_{ii} and observational parameters (Fig. 7). According to the previous considerations, we choose the values for the free parameters in ranges $0.08 \le \eta \le 0.085$ and $1.39 \le h_i \le 1.41$. Initial values of $h_{,\theta i}$, which must be negative, are assumed to be in the interval $-0.01 \le h_{,\theta i} \le -0.9$ provided that $\varepsilon_1(\theta_i) \ll 1$.



Figure 7: r versus n_s diagrams, in the models with $\eta = \text{const}$ (left panel) and $\bar{\eta} = \text{const}$ (right panel), with observational constraints from Planck Collaboration 2018 [25]. The dots are obtained for randomly chosen η , h_i and $h_{,\theta i}$ in the intervals $0.08 \le \eta \le 0.085$, $1.39 \le h_i \le 1.41$ and $-0.01 \le h_{,\theta i} \le -0.9$ (left panel), and in the intervals $-0.001 \le \eta \le -0.04$, $1.38 \le h_i \le 1.414$ and $-0.1 \le h_{,\theta i} \le -0.9$ (right panel). The colors represent different N: N = 55 (red), N = 60 (blue), and N = 65 (green).

We can easily extend our consideration to other constant-roll models using the above procedure for calculating the observational parameters. The second slow-roll parameter can be defined in terms of the Hubble expansion rate only

$$\bar{\eta} = \frac{\ell h}{h \dot{h}}.$$
(35)

The parameter $\bar{\eta}$ is different from η except in the canonical single field inflation where these two parameters are equivalent [16]. Using the same procedure as in Section 3, we derive a differential equation for the Hubble expansion rate $\bar{\eta}$

$$hh_{,\theta\theta} - h_{,\theta}^2 \left(1 + \frac{h^2}{4(1 - \frac{1}{2}h^2)(1 - \frac{1}{4}h^2)} \right) - \frac{3}{2\ell^2} \bar{\eta} h^4 \frac{1 - \frac{1}{4}h^2}{1 - \frac{1}{2}h^2} = 0.$$
(36)

By applying the numerical procedure, we calculate the observational parameters from (26) and (27) in the model with $\bar{\eta} = \text{const}$, where ε_2 and ε_3 are given by

$$\varepsilon_2 = 2(\varepsilon_1 - \bar{\eta}),\tag{37}$$

$$\varepsilon_3 = 2\varepsilon_1. \tag{38}$$

As expected, the prediction for the observational parameters (Fig. 7) in the holographic braneworld agrees with the model-independent prediction in ref. [16], obtained analytically for any model that satisfies the condition $\bar{\eta} = \text{const.}$ As we will shortly see, this fact will be of crucial importance for consideration of the connection between our model and the swampland criteria in quantum gravity.

5 The swampland criteria and the braneworld inflation

It is known that the idea of swampland [26] appeared in the context of the investigation of low-energy effective field theories (non)compatible with quantum gravity. Much attention and effort have been paid to the investigation of criteria [27] related to "swampland" or to "string theory landscape" to distinguish between effective field theories that can or cannot be UV-completed to a quantum theory of gravity.

Investigation of swampland criteria in cosmology, more precisely the theory of inflation, is pretty recent and presented in very few research articles, especially in the case of braneworld and holographic inflation. While some inflationary models are not compatible with these criteria, inflationary models in the braneworld scenario, in principle, have the potential to evade the swampland constraints [5].

Let us initiate consideration of the swampland criteria in the context of our RSII holographic constantroll model. The swampland criteria read as follows [28]

$$\frac{|\Delta\theta|}{M_{\rm Pl}} < c,\tag{39}$$

$$M_{\rm Pl} \frac{|V_{,\theta}|}{V} > c',\tag{40}$$

where c and c' are constants of order one and $M_{\rm Pl} = (8\pi G_{\rm N})^{-1/2} \simeq 2.4 \cdot 10^{18}$ GeV. Further investigations have shown that the constant c' could be of order 0.1 [29]. From the swampland criteria, it may be seen that the range traversed by the field is bounded from above, and there is a lower bound on the gradient of the potential. As pointed out in Refs. [30] and [31], the swampland conjectures will be met if at least one of the conditions is satisfied. In the following, we will examine whether the second criterion (40) is satisfied in our model. To the best of our knowledge, there has been no attempt in the literature to check the swampland criteria in a holographic inflation model with dynamics similar to the one in our model.

First, consider the second criterion, using the parameter η as presented in (12). The second criterion is in strong tension with the single-field slow-roll inflation model because the slow-roll parameter

$$\varepsilon_1 = \frac{M_{\rm Pl}^2}{2} \frac{V_{,\theta}^2}{V^2} \tag{41}$$

should be smaller than unity to have accelerated expansion phase. The definition (41) is only valid in the slow-roll approximation, in a model with a canonical scalar field in the standard cosmology [32]. For tachyon field the therm $V_{,\theta}/V$ can be calculated using the equation for $\theta(t)$ [23]

$$\frac{\ell\ddot{\theta}}{1-\dot{\theta}^2} + 3h\dot{\theta} + \frac{\ell V_{,\theta}}{V} = 0, \tag{42}$$

yielding

$$\frac{\ell V_{,\theta}}{V} = -3h\dot{\theta} \left(1 + \frac{\eta}{3(1 - \dot{\theta}^2)}\right). \tag{43}$$

For the small and constant η it leads to

$$\frac{\ell V_{,\theta}}{V} \simeq -3h\dot{\theta}.\tag{44}$$

By combining the time derivative of $h = l\dot{a}/a$ with Eq. (2), we obtain

$$\frac{\ddot{a}}{a} = \frac{h^2}{\ell^2} \left(1 - \frac{3}{2} \dot{\theta}^2 \frac{1 - \frac{1}{2} h^2}{1 - \frac{1}{4} h^2} \right).$$
(45)

From this, it follows that the condition for accelerated expansion $\ddot{a} > 0$ will be met if

$$\dot{\theta}^2 < \frac{2}{3} \frac{1 - \frac{1}{4}h^2}{1 - \frac{1}{2}h^2},\tag{46}$$

and, when $h^2 \ll 1$, inflation ends for $\dot{\theta}_{\rm f} \simeq \sqrt{2/3}$. According to (44), the second swampland criteria requires $h_{\rm f} > 0.1/\sqrt{6} \approx 0.04$. As we can see (Fig. 3), the calculated Hubble expansion rate h tends to a very small value towards the end of inflation. One may conclude that the second criterion will not be satisfied for those values of h that provide agreement with Planck data.

Next, consider the second swampland criterion using the parameter $\bar{\eta}$ given by (35), which does not depend explicitly on the field θ ! Remember that η and $\bar{\eta}$ are the same only in canonical single-field inflation, while in our model, they are different, and, as we will see, it will lead us to a different scenario. In this case, using the derivative of (17) with respect to the tachyon field, we obtain

$$\frac{\ell V_{,\theta}}{V} = -\frac{\left(1 - \frac{h^2}{2}\right) \left(-36h^3 \left(1 - \frac{h^2}{4}\right) - 8hh_{,\theta}^2 + 8\left(1 - \frac{h^2}{2}\right)h_{,\theta\theta}\right) h_{,\theta}}{18h^4 \left(1 - \frac{h^2}{4}\right)^2 - 8\left(1 - \frac{h^2}{2}\right)^2 h_{,\theta}^2}.$$
(47)

From Fig. 8, we can conclude that our holographic model provides the second swampland criterion to be satisfied during inflation for a wide range of the relevant parameters. For instance, for $\bar{\eta} = \text{const.}$ and N = 65, the following set of observational parameters is obtained: $n_{\rm s} = 0.964$, r = 0.055, that are in a good agreement with the current observational data. We can roughly estimate that the second swampland criterion is satisfied for $n_{\rm s} > 0.96$ and r > 0.05. A similar behavior for $V_{,\theta}/V$ is obtained in Ref. [5].



Figure 8: The value of $\ell V_{,\theta}/V$ versus N in the model with $\eta = \text{const}$ (left panel) and $\bar{\eta} = \text{const}$ (right panel). The parameter values are $\eta = 0.0816$, $h_i = 1.4046$, $h_{,\theta i} = -0.4503$ (left panel) and $\bar{\eta} = -0.0082$, $h_i = 1.2170$, $h_{,\theta i} = -0.1964$ (right panel).

6 Summary and conclusions

We have studied constant-roll inflation driven by the tachyon-field inflaton in the the holographic braneworld. In particular, we have focused on two models with constant roll conditions $\eta = \text{const}$ and $\bar{\eta} = \text{const}$, respectively. As expected, the inflation dynamics is modified due to the differences in the holographic Friedmann equations compared with those in the standard approach. Owing to a complicated form of the equation for the Hubble rate, we have mainly used numerical calculations throughout this work.

We have compared the theoretical results for observational parameters with the Planck data. We have deduced optimal values of our free parameters for which the model predictions are in good agreement with the observational data. In comparison with the results of the tachyon constant-roll inflation in standard cosmology [13], our model gives significantly lower values for the number of e-folds, much closer to a widely accepted $N \simeq 60$.

We have found that the formulation in which the second slow-roll parameter $\bar{\eta}$ is constant (35) meets the swampland criteria. In this case, the obtained values for the observational parameters n_s and r are still in the range of the Planck data but slightly less favorable than the values obtained with constant second slow-roll parameter η (12).

It is worth noting that the swampland "problem" for holographic inflation may be closely related to another fundamental problem of braneworld inflation. Namely, according to our model, inflation on the holographic brane occurs under the influence of the scalar inflaton field defined on the brane. However, after the Big Bang, in the era of quantum gravity, inflation might be driven by another field, for instance, some moduli scalar field in the bulk, as a possible manifestation of supergravity or superstring theory. In that case, an effective low-energy theory consistent with the swampland conjectures may not match completely with our effective gravity on the brane.

As a point for more detailed consideration of the swampland criteria in brenworld inflation, note that the equations of the gravity field on the brane that lead to holographic cosmology are not derived strictly from some fundamental action analogous to the Einstein-Hilbert action. Given this point, the investigation of fulfilling the swampland criteria for the constant-roll inflation presented in Ref. [16] is a subject of ongoing research.

Acknowledgments

This work has been supported by the ICTP-SEENET-MTP project NT-03 Cosmology-Classical and Quantum Challenges and the COST Action CA18108 "Quantum gravity phenomenology in the multimessenger approach". M. Stojanovic acknowledges the support provided by The Ministry of Science, Technological Development and Innovation of the Republic of Serbia under contract 451-03-47/2023-01/2000113. D. D. Dimitrijevic, G. S. Djordjevic, and M. Milosevic acknowledge the support provided by the same Ministry 451-03-47/2023-01/2000124. In addition, G. S. Djordjevic acknowledges the support of the CEEPUS Program RS-1514-03-2223 "Gravitation and Cosmology" as well as the hospitalities of colleagues at the University of Banja Luka and CERN-TH. N. Bilić acknowledges the hospitality of the Department of Physics at the University of Niš, where part of his work was completed.

References

- [1] A. H. Guth, Phys. Rev. D 23 (1981), 347-356 doi:10.1103/PhysRevD.23.347
- [2] A. Sen, JHEP 04 (2002), 048 doi:10.1088/1126-6708/2002/04/048 [arXiv:hep-th/0203211 [hep-th]].
- [3] M. Fairbairn and M. H. G. Tytgat, Phys. Lett. B 546 (2002), 1-7 doi:10.1016/S0370-2693(02)02638-2 [arXiv:hep-th/0204070 [hep-th]].
- [4] H. Motohashi, A. A. Starobinsky and J. Yokoyama, JCAP 09, 018 (2015) doi:10.1088/1475-7516/2015/09/018 [arXiv:1411.5021 [astro-ph.CO]].
- [5] A. Mohammadi, T. Golanbari, S. Nasri and K. Saaidi, Phys. Rev. D 101 (2020) no.12, 123537 doi:10.1103/PhysRevD.101.123537 [arXiv:2004.12137 [gr-qc]].
- [6] S. Lahiri, [arXiv:1807.01329 [hep-th]].
- S. D. Odintsov and V. K. Oikonomou, Class. Quant. Grav. 37 (2020) no.2, 025003 doi:10.1088/1361-6382/ab5c9d [arXiv:1912.00475 [gr-qc]].
- [8] Z. Yi and Y. Gong, JCAP 03 (2018), 052 doi:10.1088/1475-7516/2018/03/052 [arXiv:1712.07478 [gr-qc]].
- [9] L. Anguelova, P. Suranyi and L. C. R. Wijewardhana, JCAP 02 (2018), 004 doi:10.1088/1475-7516/2018/02/004 [arXiv:1710.06989 [hep-th]].
- [10] Q. Gao, Y. Gong and Q. Fei, JCAP 05 (2018), 005 doi:10.1088/1475-7516/2018/05/005 [arXiv:1801.09208 [gr-qc]].
- [11] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, Class. Quant. Grav. 34 (2017) no.24, 245012 doi:10.1088/1361-6382/aa92a4 [arXiv:1704.05945 [gr-qc]].
- [12] D. H. Nguyen, T. M. Pham and T. Q. Do, Eur. Phys. J. C 81 (2021) no.9, 839 doi:10.1140/epjc/s10052-021-09652-1 [arXiv:2107.14115 [gr-qc]].
- [13] A. Mohammadi, K. Saaidi and T. Golanbari, Phys. Rev. D 97 (2018) no.8, 083006 doi:10.1103/PhysRevD.97.083006 [arXiv:1801.03487 [hep-ph]].
- [14] N. Bilic, D. D. Dimitrijevic, G. S. Djordjevic, M. Milosevic and M. Stojanovic, JCAP 08 (2019), 034 doi:10.1088/1475-7516/2019/08/034 [arXiv:1809.07216 [gr-qc]].
- [15] M. Milošević, N. Bilić, D. D. Dimitrijević, G. S. Djordjević and M. Stojanović, AIP Conf. Proc. 2075 (2019) no.1, 090009 doi:10.1063/1.5091223 [arXiv:1809.04939 [gr-qc]].
- [16] M. Stojanovic, N. Bilic, D. D. Dimitrijevic, G. S. Djordjevic and M. Milosevic, [arXiv:2306.02423 [gr-qc]].

- [17] N. Bilic, D. Dimitrijevic, G. Djordjevic and M. Milosevic, Int. J. Mod. Phys. A 32 (2017) no.05, 1750039 doi:10.1142/S0217751X17500397 [arXiv:1607.04524 [gr-qc]].
- [18] N. Bilic, Phys. Rev. D 93 (2016) no.6, 066010 doi:10.1103/PhysRevD.93.066010 [arXiv:1511.07323 [gr-qc]].
- [19] P. S. Apostolopoulos, G. Siopsis and N. Tetradis, Phys. Rev. Lett. 102, 151301 (2009) doi:10.1103/PhysRevLett.102.151301 [arXiv:0809.3505 [hep-th]].
- [20] A. Mohammadi, Phys. Dark Univ. 36, 101055 (2022) doi:10.1016/j.dark.2022.101055 [arXiv:2203.06643 [gr-qc]].
- [21] S. Nojiri, S. D. Odintsov and T. Paul, Phys. Lett. B 841, 137926 (2023) doi:10.1016/j.physletb.2023.137926 [arXiv:2304.09436 [gr-qc]].
- [22] N. Bilic, [arXiv:1802.02775 [hep-th]].
- [23] D. A. Steer and F. Vernizzi, Phys. Rev. D 70 (2004), 043527 doi:10.1103/PhysRevD.70.043527 [arXiv:hep-th/0310139 [hep-th]].
- [24] N. R. Bertini, N. Bilic and D. C. Rodrigues, Phys. Rev. D 102 (2020) no.12, 123505 [erratum: Phys. Rev. D 105 (2022) no.12, 129901] doi:10.1103/PhysRevD.102.123505 [arXiv:2007.02332 [gr-qc]].
- [25] Y. Akrami *et al.* [Planck], Astron. Astrophys. **641** (2020), A10 doi:10.1051/0004-6361/201833887 [arXiv:1807.06211 [astro-ph.CO]].
- [26] C. Vafa, [arXiv:hep-th/0509212 [hep-th]].
- [27] G. Obied, H. Ooguri, L. Spodyneiko and C. Vafa, [arXiv:1806.08362 [hep-th]].
- [28] A. Kehagias and A. Riotto, Fortsch. Phys. 66 (2018) no.10, 1800052 doi:10.1002/prop.201800052 [arXiv:1807.05445 [hep-th]].
- [29] A. Mohammadi, T. Golanbari, H. Sheikhahmadi, K. Sayar, L. Akhtari, M. A. Rasheed and K. Saaidi, Chin. Phys. C 44, no.9, 095101 (2020) doi:10.1088/1674-1137/44/9/095101 [arXiv:2001.10042 [gr-qc]].
- [30] V. K. Oikonomou, K. R. Revis, I. C. Papadimitriou and M. M. Pegioudi, Int. J. Mod. Phys. D 32 (2023), 06 doi:10.1142/S0218271823500347 [arXiv:2303.14724 [gr-qc]].
- [31] S. D. Odintsov, V. K. Oikonomou and F. P. Fronimos, Phys. Rev. D 107 (2023), 08 doi:10.1103/PhysRevD.107.084007 [arXiv:2303.14594 [gr-qc]].
- [32] Ø. Grøn, Universe 4 (2018) no.2, 15 doi:10.3390/universe4020015
- [33] C. M. Lin, K. W. Ng and K. Cheung, Phys. Rev. D 100 (2019) no.2, 023545 doi:10.1103/PhysRevD.100.023545 [arXiv:1810.01644 [hep-ph]].