Degenerate preconditioned backward-backward splitting for inclusion problem

Pankaj Gautam^a and V. Vetrivel^b

^aDepartment of Mathematical Sciences, NTNU Trondheim, Norway; ^bDepartment of Mathematics, IIT Madras, India

ARTICLE HISTORY

Compiled January 2, 2024

ABSTRACT

In this work, we introduce the notion of warped Yosida regularization and study the asymptotic behaviour of the orbit of dynamical systems generated by warped Yosida regularization, which includes Douglas-Rachford dynamical system. We analyze an algorithm where the inclusion problem is first approximated by a regularized one and then the preconditioned regularization parameter is reduced to converge to a solution of original problem. We propose and investigate backward-backward splitting using degenerate preconditioning for monotone inclusion problems. The applications provide a tool for finding a minima of a preconditioned regularization of the sum of two convex functions.

KEYWORDS

Monotone operator; Yosida regularization; Preconditioning; Backward-backward splitting.

1. Introduction

In this work, we aim to solve the inclusion problem:

find
$$(x,y) \in \mathcal{H}^2$$
 such that $(0,0) \in (I - R + \lambda(M^{-1}A \times M^{-1}B))(x,y),$ (1)

where $A, B : \mathcal{H} \to 2^{\mathcal{H}}$ are two set-valued operators, $R : \mathcal{H}^2 \to \mathcal{H}^2$ is an operator defined by $(x, y) \mapsto (y, x)$, and $M : \mathcal{H} \to \mathcal{H}$ is linear, bounded, self-adjoint and positivesemidefinite operator, and \mathcal{H} is a real Hilbert space. The dual inclusion problem of (1) is

find
$$(x^*, y^*) \in \mathcal{H}^2$$
 such that $(0,0) \in ((I-R)^{-1} + (M^{-1}A \times M^{-1}B)^{-1} \circ (I/\lambda))(x^*, y^*)$
(2)

The solution of problems (1) and (2) are related to warped Yosida regularization of structured monotone inclusion problem of the form:

find
$$x \in \mathcal{H}$$
 such that $0 \in (A+B)x$. (3)

CONTACT Pankaj Gautam. Email: pgautam908@gmail.com

To find the solution of problem (3), the Yosida regularization of problem (3) is firstly introduced by Mahey and Tao [1], and Yosida regularization of an operator is introduced in [2,3].

If M = I, (1) becomes the problem of finding

$$(x,y) \in \mathcal{H}^2$$
 such that $(0,0) \in I - R + \lambda(A \times B)(x,y),$ (4)

proposed in [4], in which the authors have shown the asymptotic behaviour of the sequences generated by composition of resolvents in the connection with the solution of (4) and its dual by using Yosida regularization of problem (3). The resolvent and Yosida regularization of a maximal monotone operator are very important tools in nonlinear analysis. Bui et al. [5] have proposed a generalized resolvent called warped resolvent by using an auxiliary operator. The main motivation to define warped resolvent is to construct and investigate splitting methods by different choices of kernels. Bredies et al. [6] have analyzed Chambolle-Pock, Forward Douglas-Rachford and Peaceman-Rachford splitting methods with the help of warped resolvent by using linear kernel.

In section 3, we introduce and investigate the properties of warped Yosida approximation. In section 4, we analyze the existence, uniqueness, weak convergence of the dynamical systems governed by the warped Yosida regularization and observed Douglas-Rachford dynamical system as a particular case. Section 5 describes the preconditioned Yosida regularization of the monotone inclusion problem and analyzes the preconditioned backward-backward splitting for problem (1). Section 6 provides backward-backward splitting methods for an optimization problem.

2. Preminilaries

This section is devoted to some important definitions and results from nonlinear analysis and operator theory. Throughout the paper, $\mathcal{G}(T)$ is used to denote the graph of the set-valued operator $T : \mathcal{H} \to 2^{\mathcal{H}}$, which is defined as $\mathcal{G}(T) = \{(x, Tx) : x \in D(T)\}$, where D(T) denotes the domain of T. Let Z(T) and $\operatorname{Fix}(T)$ denote the zero and set of fixed point of an operator T, respectively. Symbol \mathbb{N} and \mathbb{R} are used to denote the set of natural numbers and set of real numbers, respectively.

Definition 2.1. Let $T : \mathcal{H} \to 2^{\mathcal{H}}$ be a set-valued operator. Then T is said to be monotone if, $\forall x_1, x_2 \in \mathcal{H}$ and $u_1 \in T(x_1), u_2 \in T(x_2)$,

$$0 \le \langle x_1 - x_2, u_1 - u_2 \rangle.$$

T is said to be maximally monotone if there exists no monotone operator $S : \mathcal{H} \to 2^{\mathcal{H}}$ such that $\mathcal{G}(S)$ properly contains $\mathcal{G}(T)$, i.e., for every $(x_1, u_1) \in \mathcal{H} \times \mathcal{H}$,

$$(x_1, u_1) \in \mathcal{G}(T) \Leftrightarrow \langle x_1 - x_2, u_1 - u_2 \rangle \ge 0, \ \forall (x_2, u_2) \in \mathcal{G}(T).$$

Let M be a bounded linear operator on \mathcal{H} . M is said to be self-adjoint if $M^* = M$, where M^* denotes the transpose conjugate of operator M. A self-adjoint operator Mon \mathcal{H} is said to be positive definite if $\langle M(x), x \rangle > 0$ for every nonzero $x \in \mathcal{H}$ ([7]). Define the M-inner product $\langle \cdot, \cdot \rangle_M$ on \mathcal{H} by $\langle x, y \rangle_M = \langle x, M(y) \rangle$ for all $x, y \in \mathcal{H}$. The corresponding M-norm is defined by $||x||_M^2 = \langle x, Mx \rangle$ for all $x \in \mathcal{H}$. **Definition 2.2.** Let D be a nonempty subset of $H, T : D \to \mathcal{H}$ an operator and $M : \mathcal{H} \to \mathcal{H}$ a positive definite operator. Then T is said to be

(i) nonexpansive with respect to *M*-norm if

$$||Tx_1 - Tx_2||_M \le ||x_1 - x_2||_M \ \forall x_1, x_2 \in \mathcal{H};$$

(ii) *M*-cocoercive if

$$||Tx_1 - Tx_2||_{M^{-1}}^2 \leq \langle x_1 - x_2, Tx_1 - Tx_2 \rangle$$
, for all $x_1, x_2 \in \mathcal{H}$.

Lemma 2.3. [8] Let C be a nonempty subset of \mathcal{H} and $x : [0, \infty) \to \mathcal{H}$ be a map. Assume that

- (i) $\lim_{t \to \infty} ||x(t) x^*||$ exists, for every $x^* \in C$;
- (ii) $\stackrel{i\to\infty}{every}$ weak sequential cluster point of the map x is in C.

Then there exists $x_{\infty} \in C$ such that $x(t) \rightharpoonup x_{\infty}$ as $t \rightarrow \infty$.

Throughout the paper we consider operator $M : \mathcal{H} \to \mathcal{H}$ as a preconditioner, i.e., a linear, bounded, self-adjoint and positive semidefinite operator, and assume that \mathcal{H} is a real Hilbert space with norm $\|\cdot\|$ and inner product $\langle\cdot,\cdot\rangle$.

Definition 2.4. [6] An admissible preconditioner for the operator $T : \mathcal{H} \to 2^{\mathcal{H}}$ is a linear, bounded, self-adjoint and positive semi-definite operator $M : \mathcal{H} \to \mathcal{H}$ such that warped resolvent

$$J^M_{\gamma T} = (M + \gamma T)^{-1} \circ M \tag{5}$$

is single-valued and has full domain.

It is easy to check that $J_{\gamma T}^{M}$ is neither everywhere defined $(J_{\gamma T}^{M}0 = \emptyset)$ nor singlevalued. For this reason, instead of imposing the maximal monotonicity of T, we directly require (5), which, in the context of splitting methods, is a reasonable assumption.

Proposition 2.5. [9] Let $M : \mathcal{H} \to \mathcal{H}$ be an admissible preconditioning and $T : \mathcal{H} \to 2^{\mathcal{H}}$ be an operator such that $M^{-1}A$ is M-monotone. Then $J^{M}_{\gamma T}$ is M-firmly nonexpansive, i.e.,

$$\|J_{\gamma T}^{M} x - J_{\gamma T}^{M} y\|_{M}^{2} + \|(I - J_{\gamma T}^{M}) x - (I - J_{\gamma T}^{M}) y\|_{M}^{2} \le \|x - y\|_{M}^{2}, \ \forall x, y \in \mathcal{H}.$$

3. Warped Yosida regularization

In this section, we define warped Yosida regularization and provide its properties and some characterizations.

Definition 3.1. Let X be a reflexive Banach space with dual space X^* . Assume that $(\emptyset \neq)C \subseteq X, M: C \to X^*$ and $T: X \to 2^{X^*}$ are such that $\operatorname{ran}(M) \subset \operatorname{ran}(T + \gamma M)$ and $T + \gamma M$ is injective. For any $\gamma \in (0, \infty)$, warped Yosida regularization of T with kernel M is defined by $T_{\gamma}^M = \frac{1}{\gamma} \left(M - M \circ J_{\gamma T}^M \right)$, where $J_{\gamma T}^M$ is the warped resolvent of T [5].

Example 3.2. Let *C* be a non-empty subset of *X* and $\phi : X \to (-\infty, \infty]$ be a proper convex lower semicontinuous map. Let $\gamma > 0$. Assume that $M : C \to X^*$ is an operator with $\operatorname{ran}(M) \subset \operatorname{ran}(M + \gamma \partial \phi)$ and $M + \gamma \partial \phi$ is injective. Then warped Yosida regularization of $\partial \phi$ is $(\partial \phi)^M_{\gamma} = \frac{1}{\gamma}(M - M \circ \operatorname{prox}^M_{\gamma \partial \phi})$, where $\operatorname{prox}^M_{\gamma \partial \phi} = (M + \gamma \partial \phi)^{-1} \circ M$. Let *M* be an injective operator, then warped Yosida regularization of $\partial \phi$ is described by the following variational inequality:

$$z = (\partial \phi)^M_{\gamma} \Leftrightarrow (\forall y \in X) \ \langle y - x + \gamma M^{-1}z, \gamma z \rangle + \phi(x - \gamma M^{-1}z) \le \phi(z) \ \forall (x, z) \in X \times X$$

Example 3.3. Let $T: X \to 2^{X^*}$ be a maximal monotone operator such that $Z(T) \neq \emptyset$. Suppose that $f: X \to (-\infty, \infty]$ is an admissible function such that $D(T) \subset int \ D(f)$. Set $M = \nabla f$. Then $T_{\gamma}^{\nabla f}$ is a well defined warped Yosida regularization defined in [10].

Now, we provide an example of warped Yosida regularization with respect to different choices of the admissible function f.

Example 3.4. Let $A : (0, \infty) \to \mathbb{R}$ be a monotone mapping. Define an admissible function (Boltzmann-Shannon entropy) $\mathcal{BS} : (0, \infty) \to (0, \infty)$ as $x \mapsto x \log x - x$. Then warped resolvent of A is [11]

$$J_A^{\mathcal{BS}}x = (\log + A)^{-1} \circ Ax = xe^{Ax}$$

and the warped Yosida regularization is

$$A^{\mathcal{BS}}x = \nabla \mathcal{BS}x - \nabla \mathcal{BS} \circ J_A^{\mathcal{BS}}x$$
$$= \log x - \log(xe^{Ax})$$
$$= \log x - \log x - \log e^{Ax}$$
$$= -Ax.$$

Now, we explore some properties and characteristics of warped Yosida regularization.

Proposition 3.5. Let $T : \mathcal{H} \to 2^{\mathcal{H}}$ and $M : \mathcal{H} \to \mathcal{H}$ be an admissible preconditioner. Then we have the following:

 $\begin{array}{l} (i) \ \left(J_{\gamma T}^{M}(x), T_{\gamma}^{M}(x)\right) \in \mathcal{G}(T), \ x \in \mathcal{H}. \\ (ii) \ 0 \in T(x) \ if \ and \ only \ if \ 0 \in T_{\gamma}^{M}(x), \ x \in \mathcal{H}. \\ (iii) \ T_{\gamma}^{M} = \left(\gamma M^{-1} + T^{-1}\right)^{-1}. \\ (iv) \ T_{\gamma}^{M} = J_{\gamma^{-1}T^{-1}}^{M^{-1}} \circ \gamma^{-1}M. \\ (v) \ T_{\gamma+\lambda}^{M} = \left(T_{\gamma}^{M}\right)_{\lambda}^{M}. \\ (vi) \ Let \ x, y \in \mathcal{H}. \ Then \ x = T_{\gamma}^{M}y \Leftrightarrow (y - \gamma M^{-1}x, x) \in \mathcal{G}(T). \\ (vii) \ \overline{D(T)} \ is \ convex, \ D(J_{\gamma T}^{M}) = D(T_{\gamma}^{M}) \ and \ ran(J_{\gamma T}^{M}) = D(T). \end{array}$

Proof. (i) For $x \in \mathcal{H}$, we have

$$J_{\gamma T}^{M}(x) = (M + \gamma T)^{-1} \circ M(x) \Leftrightarrow M(x) \in (M + \gamma T) \circ J_{\gamma T}^{M}(x)$$

$$\Leftrightarrow \frac{1}{\gamma} \left(M - M \circ J_{\gamma T}^{M} \right)(x) \in T \left(J_{\gamma T}^{M}(x) \right) \Leftrightarrow T_{\gamma}^{M}(x) \in T \left(J_{\gamma T}^{M}(x) \right).$$

(ii) For $x \in \mathcal{H}$,

$$0 \in T(x) \Leftrightarrow 0 \in \gamma T(x) \Leftrightarrow M(x) \in (M + \gamma T)(x)$$

$$\Leftrightarrow x \in (M + \gamma T)^{-1} \circ M(x) \Leftrightarrow M(x) \in M(J^M_{\gamma T}(x))$$

$$\Leftrightarrow 0 \in (M - M \circ J^M_{\gamma T})(x) \Leftrightarrow 0 \in \gamma T^M_{\gamma} x \Leftrightarrow 0 \in T^M_{\gamma} x.$$

(iii) Let $x, y \in \mathcal{H}$. Indeed,

$$x \in T^{M}_{\gamma} y \Leftrightarrow x \in \frac{1}{\gamma} \left(M - M \circ J^{M}_{\gamma T} \right) (y)$$

$$\Leftrightarrow \gamma x \in \left(M - M \circ J^{M}_{\gamma T} \right) (y) \Leftrightarrow M(y) - \gamma x \in (M \circ J^{M}_{\gamma T})(y)$$

$$\Leftrightarrow M^{-1}(M(y) - \gamma x) \in J^{M}_{\gamma T}(y) \Leftrightarrow y - \gamma M^{-1}(x) \in (M + \gamma T)^{-1} \circ M(y)$$

$$\Leftrightarrow (M + \gamma T)(y - \gamma M^{-1}(x)) \in M(y) \Leftrightarrow M(y) + \gamma T(y) - \gamma x - \gamma^{2} T \circ M^{-1}(x) \in M(y)$$

$$\Leftrightarrow x \in T(y - \gamma M^{-1}(x)) \Leftrightarrow x \in (\gamma M^{-1} + T^{-1})^{-1}(y).$$
(6)

(iv) Let $x, y \in \mathcal{H}$. From part (i),

$$\begin{aligned} x \in T^M_{\gamma} y \Leftrightarrow x \in (\gamma M^{-1} + T^{-1})^{-1}(y) \\ \Leftrightarrow x \in (M^{-1} + \gamma^{-1}T^{-1})^{-1} \circ M^{-1}(\gamma^{-1}M(y)) \\ \Leftrightarrow x \in J^{M^{-1}}_{\gamma^{-1}T^{-1}} \circ \gamma^{-1}M(y). \end{aligned}$$

(v) From part(i), for $x, y \in \mathcal{H}$,

$$\begin{aligned} x \in T^M_{(\gamma+\lambda)}(y) \Leftrightarrow &x \in ((\gamma+\lambda)M^{-1}+T^{-1})^{-1}(y) \\ \Leftrightarrow &x \in T(y-(\gamma+\lambda)M^{-1}(x)) \\ \Leftrightarrow &x \in T(y-\gamma M^{-1}y-\lambda M^{-1}y) \\ \Leftrightarrow &x \in T^M_\gamma(y-\lambda M^{-1}(y)) \\ \Leftrightarrow &x \in (T^M_\gamma)^M_\lambda y. \end{aligned}$$

(vi) For $x, y \in \mathcal{H}$ and from (6), we have

$$y \in T^M_{\gamma} x \Leftrightarrow y \in T(x - \gamma M^{-1} y) \Leftrightarrow (x - \gamma M^{-1} y, y) \in \mathcal{G}(T).$$
(7)

Proposition 3.6. Let $T : \mathcal{H} \to 2^{\mathcal{H}}$ be a maximal monotone operator and $M : \mathcal{H} \to \mathcal{H}$ be an admissible preconditioner and $\gamma > 0$. Then, we have the following:

(i) T^M_{γ} is γ -cocoercive with respect to M, i.e.,

$$\gamma \|T_{\gamma}^{M} x_{1} - T_{\gamma}^{M} x_{2}\|_{M^{-1}}^{2} \leq \langle x_{1} - x_{2}, T_{\gamma}^{M} x_{1} - T_{\gamma}^{M} x_{2} \rangle.$$
(8)

and hence maximal monotone and L-Lipschitz continuous, for some L > 0.

- (ii) $J_{\gamma T}^{M}$ is *M*-nonexpansive. (iii) Let $x_{\gamma} = J_{\gamma T}^{M} x$. Then $x_{\gamma} \rightarrow P_{\overline{dom(T)}}^{M} x$ as $\lambda \downarrow 0$, where $P_{\overline{dom(T)}}^{M} x =$ $\arg\min_{y\in\overline{dom(T)}}\|x-y\|_M.$
- (iv) For x, y, and $p \in \mathcal{H}$

$$(y,p) = \left(J_{\gamma T}^{M} x, T_{\gamma}^{M} x\right) \Leftrightarrow \begin{cases} (y,p) \in \mathcal{G}(T), \\ x = y + \gamma M^{-1} p. \end{cases}$$

Proof. (i) Let $x_1, x_2 \in \mathcal{H}$. Then

$$\begin{split} \|T_{\gamma}^{M}x_{1} - T_{\gamma}^{M}x_{2}\|_{M^{-1}}^{2} &= \langle M^{-1}(T_{\gamma}^{M}x_{1} - T_{\gamma}^{M}x_{2}), T_{\gamma}^{M}x_{1} - T_{\gamma}^{M}x_{2} \rangle \\ &= \frac{1}{\gamma^{2}} \langle (I - J_{\gamma T}^{M})x_{1} - (I - J_{\gamma T}^{M})x_{2}, (I - J_{\gamma T}^{M})x_{1} - (I - J_{\gamma T}^{M})x_{2} \rangle_{M} \\ &= \frac{1}{\gamma^{2}} \| (I - J_{\gamma T}^{M})x_{1} - (I - J_{\gamma T}^{M})x_{2} \|_{M}^{2}. \end{split}$$

Since $I - J_{\gamma T}^{M}$ is *M*-firmly nonexpansive, hence

$$\begin{split} \|T_{\gamma}^{M}x_{1} - T_{\gamma}^{M}x_{2}\|_{M-1}^{2} &\leq \frac{1}{\gamma^{2}} \langle x_{1} - x_{2}, (I - J_{\gamma T}^{M})x_{1} - (I - J_{\gamma T}^{M})x_{2} \rangle_{M} \\ &= \frac{1}{\gamma^{2}} \langle x_{1} - x_{2}, M(I - J_{\gamma T}^{M})x_{1} - M(I - J_{\gamma T}^{M})x_{2} \rangle \\ &= \frac{1}{\gamma} \langle x_{1} - x_{2}, T_{\gamma}^{M}x_{1} - T_{\gamma}^{M}x_{2} \rangle. \end{split}$$

In the similar manner, we can also show that

$$\langle x_1 - x_2, T_{\gamma}^M x_1 - T_{\gamma}^M x_2 \rangle_M \ge \gamma \|T_{\gamma}^M x_1 - T_{\gamma}^M x_2\|^2.$$
 (9)

- (ii) It follows from [12, Proposition 3.1]
- (iii) Let $\gamma \in (0,1)$ and $(y,v) \in \mathcal{G}(T)$. Then by the monotonicity of T, we have

$$\begin{aligned} \langle x_{\gamma} - y, Mx - Mx_{\gamma} - \gamma v \rangle &\geq 0 \\ \Rightarrow \|x_{\gamma} - y\|_{M}^{2} &\leq \langle x_{\gamma} - y, x - y \rangle_{M} + \|x_{\gamma} - y\|_{M} \|M^{-1}v\|_{M} \\ \Rightarrow \|x_{\gamma} - y\|_{M} &\leq \|x - y\|_{M} + \gamma \|M^{-1}v\|_{M}, \end{aligned}$$

which yields $(x_{\gamma})_{\gamma \in (0,1)}$ is bounded. Rest part of the proof follows from the proof of [13, Theorem 23.48].

(iv) From the definitions of warped resolvent and warped Yosida regularization, we have

$$\begin{cases} y = J_{\gamma T}^M x, \\ p = T_{\gamma}^M x, \end{cases} \Leftrightarrow \begin{cases} (y, \gamma^{-1}(Mx - My) \in \mathcal{G}(T), \\ p = \frac{1}{\gamma}(Mx - My), \end{cases} \Leftrightarrow \begin{cases} (y, p) \in \mathcal{G}(T), \\ x = y + \gamma M^{-1}p, \end{cases}$$

Proposition 3.7. Let $T : \mathcal{H} \to 2^{\mathcal{H}}$ be a maximal monotone operator and $M : \mathcal{H} \to \mathcal{H}$ be an adimissable preconditioner. Then, for any $x \in \mathcal{H}$, we have the following:

(i)
$$J^{M}_{\mu T} x \left(\frac{\mu}{\gamma} x + (1 - \frac{\mu}{\gamma}) J^{M}_{\gamma T} x \right) = J^{M}_{\gamma T} x.$$

(ii) $\|J^{M}_{\gamma T} x - J^{M}_{\mu T} x\| \leq \frac{\beta}{\alpha} \left(1 - \frac{\mu}{\gamma} \right) \|M^{-1} T^{M}_{\gamma} x\|$

Proof. (i) For $x \in \mathcal{H}$ and $\mu = \lambda \gamma$, we have

$$\begin{split} x &\in ((M+\gamma T)^{-1} \circ M)^{-1} \circ (M+\gamma T)^{-1} \circ Mx \\ \Leftrightarrow x &\in M^{-1} \circ (M+\gamma T) \circ J_{\gamma T}^{M} x \\ \Leftrightarrow (Mx - M \circ J_{\gamma T}^{M} x) \in \gamma T(J_{\gamma T}^{M} x) \\ \Leftrightarrow \lambda Mx - (1-\lambda) M \circ J_{\gamma T}^{M} x \in (M+\mu T) J_{\gamma T}^{M} x \\ \Leftrightarrow (M+\mu T)^{-1} \circ M(\lambda x + (1-\lambda) J_{\gamma T}^{M} x) = J_{\gamma T}^{M} x \\ \Leftrightarrow J_{\mu T}^{M} (\lambda x + (1-\lambda) J_{\gamma T}^{M} x) = J_{\gamma T}^{M} x. \end{split}$$

Putting $\lambda = \frac{\mu}{\gamma}$, we have

$$J_{\mu T}^{M} x \left(\frac{\mu}{\gamma} x + (1 - \frac{\mu}{\gamma}) J_{\gamma T}^{M} x\right) = J_{\gamma T}^{M} x.$$

(ii) From (i), we get

$$\begin{split} \|J_{\gamma T}^{M} x - J_{\mu T}^{M} x\| &= \|J_{\mu T}^{M} x \left(\frac{\mu}{\gamma} x + (1 - \frac{\mu}{\gamma}) J_{\gamma T}^{M} x\right) - J_{\mu T}^{M} x\| \\ &\leq \frac{\beta}{\alpha} \left\| \left(1 - \frac{\mu}{\gamma}\right) (x - J_{\gamma T}^{M} x) \right\| \\ &= \frac{\beta}{\alpha} \left(1 - \frac{\mu}{\gamma}\right) \|M^{-1} T_{\gamma}^{M} x\|. \end{split}$$

-		

4. Motivation to define warped Yosida regularization

Consider the monotone inclusion problem:

find
$$u \in \mathcal{H}$$
 such that $0 \in Tu$, (10)

where $T: \mathcal{H} \to 2^{\mathcal{H}}$ is a maximal monotone operator defined on Hilbert space \mathcal{H} . The differential inclusion to solve problem (10) is

$$\begin{cases} \dot{u}(t) \in -Tu(t) \\ u(t_0) = u_0 \in \mathcal{H}. \end{cases}$$
(11)

In general, the differential inclusion (11) is not well-posed. For example, if we consider an operator $T : \mathbb{R}^2 \to \mathbb{R}^2$ defined by

$$T(x,y) = (-y,x),$$

then the orbit of (11) does not converge to zero of T. But we have $Z(T) = Z(T_{\gamma}^{M})$. In order to overcome this difficulty, we shall consider the following dynamical system

$$\begin{cases} \dot{u}(t) + T_{\gamma}^{M} u(t) = 0\\ u(t_0) = u_0 \in \mathcal{H}. \end{cases}$$
(12)

In the next result, we study the properties of the orbit u(t) generated by the dynamical system (12).

Proposition 4.1. Let $T : \mathcal{H} \to 2^{\mathcal{H}}$ be a maximal monotone operator with $T^{-1}(0) \neq \emptyset$ and M be an admissible preconditioner. Let $\gamma > 0$ and $u_0 \in D(T)$. Then we have the following identities for the unique solution u(t) of dynamical systems (12):

(i) $u \in C^1(\mathbb{R}, \mathcal{H})$ and $u(t) \in D(T), \forall t > 0.$

(ii) The orbit u is bounded and $\dot{u} \in L^2([t_0,\infty);\mathcal{H})$.

(iii) u(t) converges weakly to u^* , for some $u^* \in T^{-1}(0)$.

Proof. (i) It follows from the fact that T^M_{γ} is Lipschitz continuous. (ii) Let $u^* \in T^{-1}(0)$. Define an anchor

$$h(t) = \frac{1}{2} \|u(t) - u^*\|_M^2$$

From (12), we get

$$\dot{h}(t) + \langle u(t) - u^*, T_{\gamma}^M x(t) \rangle_M = 0.$$

Since $u^* \in T^{-1}(0) = (T^M_{\gamma})^{-1}(0)$ and by (9), we deduce

$$\dot{h}(t) + \gamma \|T_{\gamma}^{M}u(t)\|^{2} \leq 0$$

$$\Rightarrow \dot{h}(t) + \gamma \|\dot{u}(t)\|^{2} \leq 0.$$
(13)

Hence $t \mapsto h(t)$ is a monotonically decreasing function. Since $t \mapsto h(t)$ is a locally absolutely continuous function, there exists $N_1 \in \mathbb{R}$ such that

$$h(t) \leq N_1$$
 for all $t \in [t_0, \infty)$,

which concludes that h(t) is a bounded function and hence u(t) is also bounded function. Integrating (13), we get a real number N_2 such that

$$h(t) + \gamma \int_{t_0}^t \|\dot{u}(t)\|^2 \le N_2 \text{ for all } t \in [t_0, \infty).$$
 (14)

Using the boundedness of h, we conclude that $\dot{u} \in L^2([t_0, \infty); \mathcal{H})$.

(iii) By Lemma 2.3, and maximal monotonicity of T^M_{γ} , the orbit weakly converges to u^* for some $u^* \in T^{-1}(0)$.

Interestingly, from T_{γ}^{M} one can obtain different splitting based dynamical systems by different choices of operators T and M, e.g., Douglas-Rachford, forward-Douglas-Rachford dynamical systems.

Let $u = (x, y) \in \mathcal{H} := H^2$, where H is a real Hilbert space. Consider $T : H^2 \to 2^{H^2}$ and $M : H^2 \to H^2$ be the operators defined by

$$T = \begin{bmatrix} \alpha A & I \\ -I & (\alpha B)^{-1} \end{bmatrix}, \quad M = \begin{bmatrix} I & -I \\ -I & I \end{bmatrix}, \tag{15}$$

where $A, B: H \to 2^H$ are maximal monotone operators. With this choice of operator T, problem (10) is converted into

find
$$u \in H$$
 such that $0 \in (A+B)u$, (16)

equivalently, $0 \in (A + B)x$ if and only if there exists $y \in H$ such that $0 \in \alpha Ax + y$ and $0 \in \alpha Bx$. Now, for T and M defined by (15) we have

$$\begin{split} J_T^M \\ &= \begin{bmatrix} \alpha A + I & 0 \\ -2I & (\alpha B)^{-1} + I \end{bmatrix}^{-1} \circ \begin{bmatrix} I & -I \\ -I & I \end{bmatrix} \\ &= \begin{bmatrix} (\alpha A + I)^{-1} & -(\alpha A + I)^{-1} \\ 2((\alpha B)^{-1} + I)^{-1}(\alpha A + I)^{-1} - ((\alpha B)^{-1} + I)^{-1} & -2((\alpha B)^{-1} + I)^{-1}(\alpha A + I)^{-1} + ((\alpha B)^{-1} + I)^{-1} \end{bmatrix} \\ &= \begin{bmatrix} J_{\alpha A} & -J_{\alpha A} \\ 2J_{(\alpha B)^{-1}}J_{\alpha A} - J_{(\alpha B)^{-1}} & -2J_{(\alpha B)^{-1}}J_{\alpha A} + J_{(\alpha B)^{-1}} \end{bmatrix}. \end{split}$$

So,

$$\begin{split} T^{M} &= \begin{bmatrix} I & -I \\ -I & I \end{bmatrix} - \begin{bmatrix} J_{\alpha A} - 2J_{(\alpha B)^{-1}}J_{\alpha A} + J_{(\alpha B)^{-1}} & -J_{\alpha A} + 2J_{(\alpha B)^{-1}}J_{\alpha A} - J_{(\alpha B)^{-1}} \\ -J_{\alpha A} + 2J_{(\alpha B)^{-1}}J_{\alpha A} - J_{(\alpha B)^{-1}} & J_{\alpha A} - 2J_{(\alpha B)^{-1}}J_{\alpha A} + J_{(\alpha B)^{-1}} \end{bmatrix} \\ &= \begin{bmatrix} \frac{I - R_{\alpha A}R_{\alpha B}}{2} & \frac{R_{\alpha A}R_{\alpha B} - I}{2} \\ \frac{R_{\alpha A}R_{\alpha B} - I}{2} & \frac{I - R_{\alpha A}R_{\alpha B}}{2} \end{bmatrix}, \end{split}$$

where $R_{\alpha A}$ and $R_{\alpha B}$ are reflected resolvents of the operators A and B, respectively. Hence from dynamical system (12), we get

$$\begin{cases} \dot{x}(t) = -\frac{I - R_{\alpha A} R_{\alpha B}}{2} (x(t) - y(t)) \\ \dot{y}(t) = -\frac{I - R_{\alpha A} R_{\alpha B}}{2} (y(t) - x(t)) \\ x(0) = x_0, y(0) = y_0. \end{cases}$$

By substituting z(t) = x(t) - y(t), we obtain that

$$\begin{cases} \dot{z}(t) + z(t) = R_{\alpha A} R_{\alpha B} z(t) \\ z(0) = z_0, \end{cases}$$

which is a Douglas-Rachford (without preconditionig) dynamical system investigated in [14].

5. Preconditioned backward-backward splitting

This section generalizes the regularization of sum of two monotone operators [1] using preconditioner. A preconditioned regularization of monotone inclusion problem

find
$$x \in \mathcal{H}$$
 such that $0 \in Ax + Bx$ (17)

is

find
$$x \in \mathcal{H}$$
 such that $0 \in A_{\lambda}^{M} x + Bx$. (18)

Note that, if $0 \in A^M_{\lambda} x + Bx$,

$$M^{-1} \circ (M + \lambda B)x \in J^M_{\lambda A} x$$
$$x \in (M + \lambda B)^{-1} \circ M \circ J^M_{\lambda A} x$$
$$x \in J^M_{\lambda B} \circ J^M_{\lambda A} x.$$
(19)

We study the convergence of sequence $\{x_n\}$ defined by: $\{x_n\} \in \mathcal{H}$ and $\{x_n\}$ solves the problem (18). Assume that problem (18) has a solution for $\lambda > 0$. Then for each $n, \{x_n\}$ satisfies:

$$x_n = J^M_{\lambda B} \circ J^M_{\lambda A} x_n. \tag{20}$$

Let $y_n = A_{\lambda}^M x_n$. In the next result, we study the convergence analysis of sequences $\{x_n\}$ and $\{y_n\}$.

Theorem 5.1. Let $A, B : \mathcal{H} \to 2^{\mathcal{H}}$ be two maximal monotone operators. Then we have the following:

- (i) Assume that x is a solution of (17) and $y \in Ax \cap (-Bx)$. Then $||y_n|| \le ||y||$ for any $\lambda > 0$.
- (ii) Let x be a limit point of $\{x_n\}$ and the sequence $\{y_n\}$ be bounded. Then, x solves (17).
- (iii) If $\{x_n\}$ has a limit point, then problem (17) has a solution if and only if $\{y_n\}$ is bounded.
- (iv) If $\{x_n\}$ is bounded and problem (17) has a unique solution, then $x_n \to x^*$ and $y_n \to y^*$ which is the element of minimum norm in $Ax^* \cap (-Bx^*)$.

Proof. (i) As $y \in -Bx$ and $y_n \in -Bx_n$, so using the monotonicity of B, we have

$$\langle y_n - y, x_n - x \rangle \le 0. \tag{21}$$

Also

$$y_n = A_{\lambda}^M x_n = \frac{1}{\lambda} \left(M x_n - M \circ J_{\lambda A}^M x_n \right)$$

which implies that

$$Mx_n = \lambda y_n + M \circ J^M_{\lambda A} x_n \tag{22}$$

From (21) and (29), we have

$$\langle y_n - y, x_n - x \rangle = \langle y_n - y, \lambda M^{-1} y_n + J^M_{\gamma A} x_n - x \rangle \le 0,$$

i.e.,

$$\langle y_n - y, \lambda M^{-1} y_n \rangle \leq -\langle y_n - y, J_{\lambda A}^M x_n - x \rangle.$$

Since $y \in Ax$, $y_n = A_{\lambda}^M x_n \in A(J_{\lambda A}^M)$ and A is monotone, we get

 $||y_n||^2 \le ||y|| ||y_n||.$

(ii) By Proposition 3.5(vii) and (20), we have $x_n \in D(B)$. Again, by Proposition 3.5(vii) and (22), we obtain $x_n \in \lambda M^{-1}y_n + D(A)$. As operator M and sequence $\{y_n\}$ are bounded, the first term of the sequence $\{x_n\}$ tends to 0 as $\lambda \downarrow 0$ and limit point x must belong to $\overline{D(A)}$. Hence $x \in \overline{D(A)} \cap \overline{D(B)}$.

Since $\{y_n\}$ is bounded, we can consider a convergent subsequence $\{y_{n_k}\}$ of $\{y_n\}$. Let y be the limit of $\{y_{n_k}\}$. Consider $z_{n_k} = J^M_{\lambda A} x_{n_k}$. Then $y_{n_k} \in A z_{n_k}$ for any k. Now,

$$\begin{aligned} \|z_{n_{k}} - x\|_{M} &= \|z_{n_{k}} - J_{\lambda A}^{M} x + J_{\lambda A}^{M} x - x\|_{M} \\ &\leq \|J_{\lambda A}^{M} x_{n_{k}} - J_{\lambda A}^{M} x\|_{M} + \|J_{\lambda A}^{M} x - x\|_{M} \\ &\leq \|x_{n_{k}} - x\|_{M} + \|J_{\lambda A}^{M} x - x\|_{M}. \end{aligned}$$

Since $\|\cdot\|$ and $\|\cdot\|_M$ are equivalent norms, by Proposition 3.6(iii), the second term of the above inequality tends to 0 as $\lambda \downarrow 0$ (as $x \in \overline{D(A)}$), and hence $z_{n_k} \to x$.

As A is a closed map, $y_{n_k} \to y$ and $z_{n_k} \to x$, hence $y \in Ax$. Also $-y_{n_k} \in Bx_{n_k}$ and B is a closed map, $-y \in Bx$. Thus x solves (17).

- (iii) It follows from (i) and (ii).
- (iv) Let x^* is an unique solution of the problem (17). Since $\{x_n\}$ is bounded and from part (ii), we conclude that its limit point is a solution of (17). Hence x^* is the unique limit point of the sequence $\{x_n\}$ and $x_n \to x^*$. Suppose that y^* is an element of minimum norm in $Ax^* \cap (-Bx^*)$. Then from part (i), we get

$$||y_n|| \le ||y^*||.$$

Let $y_n \to y$. Then $||y|| \leq ||y^*||$. Also from part (2), we have $y \in Ax^* \cap (-Bx^*)$, and hence $y = y^*$. hence $y_n \to y^*$.

In the similar manner, we can show that the problem:

find
$$y \in \mathcal{H}$$
 such that $0 \in Ay + B^M_\lambda y$ (23)

is also a preconditioned regularization of problem (17). The dual of the problems (18) and (23) are

find
$$x^* \in H$$
 such that $0 \in \widetilde{A_{\lambda}^M} x^* + B^{-1} x^*$, (24)

and

find
$$y^* \in H$$
 such that $0 \in A^{-1}y^* + \widetilde{B^M_\lambda}y^*$, (25)

respectively, where $\widetilde{A} = (-I) \circ A^{-1} \circ (-I)$.

In the next result, we show the relation between the solution set of the problems (1), (2), (18) and (23), which are denoted by S, S^*, E and F, respectively.

Proposition 5.2. The following identities hold:

- $\begin{array}{ll} (i) \ E = \operatorname{Fix}(J_{\lambda B}^{M}J_{\gamma A}^{M}) = J_{\lambda B}^{M}(F), \ and \ F = \operatorname{Fix}(J_{\lambda A}^{M}J_{\lambda B}^{M}) = J_{\lambda A}^{M}(E).\\ (ii) \ S = (F \times E) \cap \mathcal{G}(J_{\lambda B}^{M})\\ (iii) \ S^{*} = \ \{(\lambda u^{*}, \lambda v^{*})\} \ such \ that \ v^{*} \ = \ -u^{*}, \ where \ u^{*} \ = \ J_{\frac{A^{-1} + \tilde{B}}{\lambda}}^{M^{-1}}(0) \ and \ v^{*} \ = \ J_{\frac{A^{+1} + \tilde{B}}{\lambda}}^{M^{-1}}(0).\\ (iv) \ S^{*} = -R \circ S. \end{array}$
- (v) $J^{M}_{\lambda B}|_{F}: F \to E: x \mapsto x + \lambda M^{-1}u^{*}$ is a bijective map with inverse $J^{M}_{\lambda A}|_{E}: E \to F: y \mapsto y + \lambda M^{-1}v^{*}.$

Proof. (i) Let $x \in F$. Then

$$0 \in Ax + B^{M}_{\lambda}x \Leftrightarrow 0 \in \lambda Ax + Mx - M \circ J^{M}_{\lambda B}x$$
$$\Leftrightarrow M^{-1} \circ (M + \lambda A)x \in J^{M}_{\lambda B}x$$
$$\Leftrightarrow x \in \operatorname{Fix}(J^{M}_{\lambda A}J^{M}_{\lambda B}).$$

In the similar manner, one can show that $E = \operatorname{Fix}(J_{\lambda B}^{M}J_{\lambda A}^{M})$. For $x \in \operatorname{Fix}(J_{\lambda B}^{M}J_{\lambda A}^{M})$, we have $J_{\lambda A}^{M}x \in \operatorname{Fix}(J_{\lambda A}^{M}J_{\lambda B}^{M})$. Hence $J_{\lambda A}^{M}(E) \subset F$. Similarly, $J_{\lambda B}^{M}(F) \subset E$. Again by these last two inclusions, we obtain that $F = J_{\lambda A}^{M}(E)$ and $E = J_{\lambda B}^{M}(F)$.

(ii) Let $(x, y) \in S$. Then

$$\begin{aligned} &-R(x,y) \in \lambda(A \times B)(x,y) \\ \Leftrightarrow &(My,Mx) \in (Mx,My) + \lambda(A \times B)(x,y) \\ \Leftrightarrow &My \in Mx + \lambda Ax \text{ and } Mx = My + \lambda By \\ \Leftrightarrow &x = J^M_{\lambda A}y \text{ and } y = J^M_{\lambda B}x \\ \Leftrightarrow &x \in \operatorname{Fix}(J^M_{\lambda A}J^M_{\lambda B}), \ y \in \operatorname{Fix}(J^M_{\lambda B}J^M_{\lambda A}) \text{ and } y = J^M_{\lambda B}x \\ \Leftrightarrow &(x,y) \in F \times E \text{ and } (x,y) = \mathcal{G}(J^M_{\lambda B}). \end{aligned}$$

(iii) For $(x^*, y^*) \in S^*$, we have

$$(0,0) \in R^{-1}(x^*, y^*) + \left(A^{-1}\left(\frac{x^*}{\lambda}\right) \times B^{-1}\left(\frac{x^*}{\lambda}\right)\right).$$

Then, there exists $(x, y) \in S$ such that

$$\begin{cases} (x,y) \in A^{-1}\left(\frac{x^*}{\lambda}\right) \times B^{-1}\left(\frac{x^*}{\lambda}\right), \\ (-x,-y) \in R^{-1}(x^*,y^*), \end{cases} \Rightarrow \begin{cases} (x^*,y^*) \in \lambda(A \times B)(x,y), \\ (-x^*,-y^*) = R(x,y), \end{cases}$$

which implies that

$$\begin{cases} x \in A^{-1}(x^*/\lambda), \\ -y \in \widetilde{B}(x^*/\lambda), \\ x^* = M(y-x) = -y^*. \end{cases}$$
(26)

On the other hand,

$$\begin{cases} -x \in \widetilde{A}(y^*/\lambda), \\ y \in B^{-1}(y^*/\lambda), \\ x^* = M(y-x) = -y^*. \end{cases}$$
(27)

From (26) and (27), we get

$$\begin{cases} -M^{-1}x^* = x - y \in (A^{-1} + \widetilde{B})(x^*/\lambda), \\ -M^{-1}y^* = y - x \in (\widetilde{A} + B^{-1})(y^*/\lambda), \end{cases}$$

i.e.,

$$\begin{cases} x^* = \lambda J_{\frac{A^{-1} + \tilde{B}}{\lambda}}^{M^{-1}}(0) = \lambda u^*, \\ y^* = \lambda J_{\frac{\tilde{A} + B^{-1}}{\lambda}}^{M^{-1}}(0) = \lambda v^*. \end{cases}$$

$$(28)$$

In the next results, we show the convergence of backward-backward splitting algorithm to problem (1).

Theorem 5.3. Let $M : \mathcal{H} \to \mathcal{H}$ be an admissible preconditioner and $A, B : \mathcal{H} \to 2^{\mathcal{H}}$ be two maximal monotone operators with $S \neq \emptyset$ and $\lambda > 0$. Fix $x_0 \in \mathcal{H}$ and for $n \ge 0$ set

$$y_n = J^M_{\lambda B} x_n, \ x_{n+1} = J^M_{\lambda A} y_n.$$
⁽²⁹⁾

Then we have the following:

- (i) The sequence $\{(x_n, y_n)\}$ converges weakly to a point in S.
- (ii) For every $(\bar{x}, \bar{y}) \in S$,

$$\sum_{n \in \mathbb{N}} \|(x_n - y_n) - (\bar{x} - \bar{y})\|_M^2 < \infty, \text{ and}$$
$$\sum_{n \in \mathbb{N}} \|(x_{n+1} - y_n) - (\bar{x} - \bar{y})\|_M^2 < \infty.$$

(iii) The sequence $\{(y_n - x_n, x_{n+1} - y_n)\}$ converges strongly to $\lambda(M^{-1}u^*, M^{-1}v^*)$. **Proof.** Let $(x^*, y^*) \in S$. By Proposition 2.5 and (29), we have

$$\begin{aligned} \|x_{n+1} - x^*\|_M^2 &= \left\|J_{\lambda A}^M J_{\lambda B}^M x_n - J_{\lambda A}^M J_{\lambda B}^M \bar{x}\right\|_M^2 \\ &\leq \left\|J_{\lambda B}^M x_n - J_{\lambda B}^M \bar{x}\right\|_M^2 - \left\|(I - J_{\lambda A}^M) J_{\lambda B}^M x_n - (I - J_{\lambda A}^M) J_{\lambda B}^M \bar{x}\right\|_M^2 \\ &\leq \|x_n - \bar{x}\|_M^2 - \left\|(I - J_{\lambda B}^M) x_n - (I - J_{\lambda B}^M) \bar{x}\right\|_M^2 \\ &- \left\|(I - J_{\lambda A}^M) J_{\lambda B}^M x_n - (I - J_{\lambda A}^M) J_{\lambda B}^M \bar{x}\right\|_M^2 \\ &= \|x_n - \bar{x}\|_M^2 - \|(x_n - y_n) - (\bar{x} - \bar{y})\|_M^2 - \|(y_n - x_{n+1}) - (\bar{y} - \bar{x})\|_M^2, \end{aligned}$$

which implies that

$$\sum_{n \in \mathbb{N}} \| (x_n - y_n) - (\bar{x} - \bar{y}) \|_M^2 + \| (x_{n+1} - y_n) - (\bar{x} - \bar{y}) \|_M^2 \| \le \| x_0 - \bar{x} \|_M^2,$$
(30)

which concludes (ii).

From Proposition 5.2(i) and (v), we get

$$\overline{y} = J^M_{\lambda B} \overline{x} = \overline{x} + \lambda M^{-1} u^* \text{ and } \overline{x} = J^M_{\lambda A} \overline{y} = \overline{y} + \lambda M^{-1} v^*.$$
(31)

From (30) and (31), we obtain $y_n - x_n \to \overline{y} - \overline{x} = \lambda M^{-1} u^*$ and $x_{n+1} - y_n \to \overline{x} - \overline{y} = \lambda M^{-1} v^*$, which concludes (iii) and also the fact that $x_{n+1} - x_n \to 0$.

Now, since $J_{\gamma A}^M J_{\gamma B}^M$ is nonexpansive, the sequence $\{x_n\}$ converges weakly to a fixed point x of $J_{\gamma A}^M J_{\gamma B}^M$. Let $y = J_{\gamma B}^M x$. From Proposition 5.2 (i) and (ii), we obtain that $(x, y) \in S$. Hence, sequence $\{y_n\}$ converges weakly to y, as $y_n - x_n \to y - x$. Therefore, $\{(x_n, y_n)\}$ converges weakly to a point in S.

Theorem 5.4. Let $M : \mathcal{H} \to \mathcal{H}$ be an admissible preconditioner and $A, B : \mathcal{H} \to 2^{\mathcal{H}}$ be two maximal monotone operators such that $S = \emptyset$ and $\gamma > 0$. Let the sequence $\{(x_n, y_n)\}$ be defined by (29). Then $||x_n|| \to \infty$ and $||y_n|| \to \infty$.

Proof. For every $n \in \mathbb{N}$, we have $x_n = T^n x_0$, where $T = J^M_{\lambda A} J^M_{\lambda B}$ is and by [4, Fact 2.2], T is strongly nonexpansive. So $||x_n|| \to \infty$ as $n \to \infty$. Similarly, we can show $||y_n|| \to \infty$.

6. An application

Let $\Gamma_0(\mathcal{H})$ be a collection of proper, convex and lower-semicontinuous from \mathcal{H} to $(-\infty, \infty]$ and $f, g \in \Gamma_0(\mathcal{H})$. Consider the function

$$\Phi: \mathcal{H} \times \mathcal{H} \to (-\infty, \infty]: (x, y) \mapsto f(x) + g(y) + \frac{1}{2\lambda} \|x - y\|_M^2,$$
(32)

where $M : \mathcal{H} \to \mathcal{H}$ is a linear, bounded, self-adjoint and positive semi-definite operator. By taking A and B to be the convex subdifferential of f and g, respectively, by Proposition 5.2 and Theorem 5.3, the following results hold for the problem: min $\Phi(\mathcal{H} \times \mathcal{H})$.

Corollary 6.1. (i) $S = \operatorname{argmin}(\Phi)$. (ii) $E = \operatorname{Fix}(\operatorname{prox}_{\lambda g}^{M} \operatorname{prox}_{\lambda f}^{M})$ and $F = \operatorname{Fix}(\operatorname{prox}_{\lambda f}^{M} \operatorname{prox}_{\lambda g}^{M})$. (iii) $S = (F \times E) \cap \mathcal{G}(\operatorname{prox}_{\lambda g}^{M})$.

Corollary 6.2. Let $M : \mathcal{H} \to \mathcal{H}$ be an admissible preconditioner, $f, g \in \Gamma_0(\mathcal{H})$ with $S \neq \emptyset$ and $\lambda > 0$. Fix $x_0 \in \mathcal{H}$ and for $n \ge 0$ set

$$y_n = \operatorname{prox}_{\lambda g}^M x_n, \ x_{n+1} = \operatorname{prox}_{\lambda f}^M y_n.$$

Then we have the following:

- (i) The sequence $\{(x_n, y_n)\}$ converges weakly to a point in S.
- (ii) For every $(\bar{x}, \bar{y}) \in S$,

$$\sum_{n \in \mathbb{N}} \|(x_n - y_n) - (\bar{x} - \bar{y})\|_M^2 < \infty, \text{ and}$$
$$\sum_{n \in \mathbb{N}} \|(x_{n+1} - y_n) - (\bar{x} - \bar{y})\|_M^2 < \infty.$$

(iii) The sequence $\{(y_n - x_n, x_{n+1} - y_n)\}$ converges strongly to $\lambda(M^{-1}u^*, M^{-1}v^*)$.

References

- [1] Mahey P, Tao PD. Partial regularization of the sum of two maximal monotone operators. ESAIM: Mathematical Modelling and Numerical Analysis. 1993;27(3):375–392.
- [2] Moreau JJ. Proximité et dualité dans un espace hilbertien. Bulletin de la Société mathématique de France. 1965;93:273-299.
- [3] Yosida K. Functional analysis. Springer Science & Business Media; 2012.
- [4] Bauschke HH, Combettes PL, Reich S. The asymptotic behavior of the composition of two resolvents. Nonlinear Analysis: Theory, Methods & Applications. 2005;60(2):283–301.
- [5] Bui MN, Combettes PL. Warped proximal iterations for monotone inclusions. Journal of Mathematical Analysis and Applications. 2020;491(1):124315.
- [6] Bredies K, Chenchene E, Lorenz DA, et al. Degenerate preconditioned proximal point algorithms. SIAM Journal on Optimization. 2022;32(3):2376–2401.
- [7] Limaye BV. Functional analysis. New Age International; 1996.
- [8] Bot RI, Csetnek ER. Second order forward-backward dynamical systems for monotone inclusion problems. SIAM Journal on Control and Optimization. 2016;54(3):1423–1443.
- Bredies K, Sun H. Preconditioned douglas-rachford splitting methods for convex-concave saddle-point problems. SIAM Journal on Numerical Analysis. 2015;53(1):421-444.
- [10] Reich S, Sabach S. Two strong convergence theorems for a proximal method in reflexive banach spaces. Numerical Functional Analysis and Optimization. 2010;31(1):22-44.
- [11] Sabach S. Iterative methods for solving optimization problems. Technion-Israel Institute of Technology, Faculty of Mathematics; 2012.
- [12] Dixit A, Sahu D, Gautam P, et al. An accelerated forward-backward splitting algorithm for solving inclusion problems with applications to regression and link prediction problems. choice. 2021;4(1):2.
- [13] Bauschke HH, Combettes PL. Correction to: Convex analysis and monotone operator theory in hilbert spaces. Springer; 2017.
- [14] Gautam P, Som K, Vetrivel V. Parameterized douglas-rachford dynamical system for monotone inclusion problems. Applied Set-Valued Analysis and Optimization. 2023; 5(1):19-29.