

Gravitational wave footprints from Higgs-portal scalegenesis with multiple dark chiral scalars

He-Xu Zhang* and Shinya Matsuzaki†

Center for Theoretical Physics and College of Physics, Jilin University, Changchun, 130012, China

Hiroyuki Ishida‡

Center for Liberal Arts and Sciences, Toyama Prefectural University, Toyama 939-0398, Japan

We discuss the gravitational wave (GW) spectra predicted from the electroweak scalegenesis of the Higgs portal type with a large number of dark chiral flavors, which many flavor QCD would underlie and give the dynamical explanation of the negative Higgs portal coupling required to trigger the electroweak symmetry breaking. We employ the linear-sigma model as the low-energy description of dark many flavor QCD and show that the model undergoes ultra-supercooling due to the produced strong first-order thermal phase transition along the (approximately realized) flat direction based on the Gildener-Weinberg mechanism. Passing through evaluation of the bubble nucleation/percolation, we address the reheating and relaxation processes, which are generically non-thermal and nonadiabatic. Parametrizing the reheating epoch in terms of the e-folding number, we propose proper formulae for the redshift effects on the GW frequencies and signal spectra. It then turns out that the ultra-supercooling predicted from the Higgs-portal scalegenesis generically yields none of GW signals with the frequencies as low as nano Hz, instead, prefers to give the higher frequency signals, which still keeps the future prospected detection sensitivity, like at LISA, BBO, and DECIGO, etc. We also find that with large flavors in the dark sector, the GW signals are made further smaller and the peak frequencies higher. Characteristic phenomenological consequences related to the multiple chiral scalars include the prediction of dark pions with the mass much less than TeV scale, which is also briefly addressed.

PACS numbers:

I. INTRODUCTION

The origin of mass and the electroweak symmetry breaking is not sufficiently accounted for in the standard model (SM), although the SM-like Higgs was discovered [1, 2]: in the SM, the sign of the Higgs mass parameter is necessarily assumed to be negative, to realize the electroweak symmetry breaking, which is given by hand. This is indeed the longstanding and unsolved issue still left at present, with which the gauge hierarchy problem or fine tuning problem is also associated.

One idea to tackle this issue is to consider the so-called classical scale invariance, which was originally inspired by Bardeen's argument [3]. The classical scale invariance sets the Higgs mass parameter to be zero at some scale in the renormalization group evolution, say, at the Planck scale, so that the Higgs mass will not be generated. It has been so far suggested that this classical scale invariance for the Higgs potential at the Planck scale can be realized as an infrared fixed point nonperturbatively generated by quantum gravitational effects [4–8].

It might be interesting to argue also that the observed SM-like Higgs is supposed to have the profile along a nearly scale-invariant direction, i.e., the flat direction in the electroweak-broken phase, which can be manifested by the small Higgs quartic coupling, $\lambda_H = (m_h^2/2v_{\text{EW}}^2) \simeq 1/8 \ll 1$ [9, 10]: taking the limit $\lambda_H \rightarrow 0$ thus leads to the flat Higgs potential keeping nonzero vacuum expectation value $v_{\text{EW}} \simeq 246$ GeV and the mass $m_h \simeq 125$ GeV.

Given the classical scale invariance, the scalegenesis has to be triggered by new physics, like a dark sector. The simplest idea along this conformal extension of the SM is to predict one SM-singlet scalar, S , allowing coupling to the Higgs doublet via forming the portal with a real scalar [11] or an extra $U(1)$ -charged scalar [12], or a generic complex scalar with or without CP violation [13–15], such as $|H|^2 S^2$. Those dark sector scalars together with the SM-like Higgs develop the flat direction and the classical scale invariance is spontaneously and explicitly broken by the dimensional transmutation at the quantum loop level, due to what is called Coleman-Weinberg mechanism [16]

*hxzhang18@163.com

†synya@jlu.edu.cn

‡ishidah@pu-toyama.ac.jp

and/or Gildener-Weinberg mechanism [17]. This is the scalegenesis of one kind, what we may call the Higgs portal scalegenesis.

In the simplest Higgs-portal scalegenesis where the singlet scalar is introduced, the portal coupling is necessarily assumed to be negative. There even including the radiative corrections, one needs to require the portal coupling to be negative by hand, otherwise any models can never realize the electroweak symmetry breaking (see, e.g., [18], and references therein). Actually, this is the same drawback as what the SM possesses in terms of the negative Higgs mass parameter. Therefore, the thus simple-minded Higgs portal scalegenesis still calls for some new physics.

One way out is to further predict an additional dark sector with a new gauge symmetry U_{B-L} or $U(1)_X$ under which the new scalar is charged (i.e., that is the $B-L$ Higgs), the negative portal coupling can be generated by the renormalization group evolution [19–21].

Another type of the dynamical origin of the negative portal coupling has been proposed in a unified way in [22]. It is mandatory to link with an underlying (almost) scale-invariant dark QCD with many flavors. In this scenario, the Higgs portal partner, a dilaton, arises as a composite-singlet scalar generated from the scale-invariant many flavor QCD. The scale anomaly induced via the Gildener-Weinberg/Coleman-Weinberg mechanism can also be interpreted, by the anomaly matching, as the nonperturbative scale anomaly coupled to the composite dilaton, which is generated by the dynamical chiral-scale breaking in many flavor QCD. Along this scenario, generically plenty of dark hadron spectra will be predicted due to the many flavor structure in the scale-invariant dark QCD sector, which could be testable at collider experiments and/or through footprints in cosmological observations. Such an almost-scale invariant feature has also been applied to inflationary scenarios with the small-field inflation of the Coleman-Weinberg (CW) type [23, 24].

In this paper, we focus on many flavor QCD scenario in a view of the underlying theory for the Higgs portal scalegenesis, and discuss the gravitational-wave (GW) footprints in cosmology arising from the cosmological phase transition along the flat direction. We in particular take the number of flavors (N_f) for scale-invariant many flavor QCD with the number of colors $N_c = 3$ to be 8, as a definite benchmark model, though we will keep arbitrary N_f when discussing analytic features. This setup has been definitely clarified, in lattice simulations, to be the scale-invariant QCD along with presence of the chiral broken phase [25–27] and the light dilaton [28–31] (when the eight fermions are in the fundamental representation of the gauge group).

We work on the scale-invariant linear sigma model as the low-energy description of underlying many flavor QCD, where the SM sector couples to through the Higgs portal coupling. With the currently available observables and constraints related to the Higgs sector at hand, we analyze the cosmological phase transition and show that in the case of the benchmark model with $N_f = 8$ the ultra-supercooling is generated and the nucleation/percolation of true-vacuum bubbles. The large flavor dependence on the cosmological phase transition is also discussed. Then we evaluate the GW signals sourced from the ultra-supercooling.

In the literature [32], GW spectra produced from the ultra-supercooling in many flavor QCD with $N_f = 8$ have been discussed based on the scale-invariant linear sigma model description as the low-energy effective theory. This is, however, not the Higgs portal scalegenesis, but what is called (many-flavor) walking technicolor [33], where the composite dilaton (called technidilaton [34]) plays the role of the SM-like Higgs itself.

Our particular claim is also on the evaluation of the redshift effect on the produced GWs. This redshift arises through the reheating epoch due to releasing the false vacuum energy (latent heat) into the SM thermal plasma via the Higgs portal coupling. Parametrizing the reheating epoch in terms of the e-folding number, we propose proper formulae for the redshift effects on the GW frequencies and signal spectra.

We find that the ultra-supercooling with large N_f generically yields none of GW signals with the frequencies as low as nano Hz (namely, no signal in NANO Grav 15yr [35] and also in other nano Hz signal prospects [36–38]), instead, prefers to give the higher frequency signals. The thus characteristically produced GW spectra, however, still keep having the future prospected detection sensitivity, like at LISA [39, 40], BBO [41, 42], and DECIGO [43, 44], etc. We also find that with large flavors in the dark sector, the GW signals are made further smaller and the peak frequencies higher.

Characteristic phenomenological consequences related to the multiple chiral scalars include the prediction of dark pions with the mass much less than TeV scale, which is also briefly addressed.

II. THE MODEL SET-UP

We begin by modeling the Lagrangian having the $U(N_f)_L \times U(N_f)_R$ chiral symmetry and the classical scale invariance at some ultraviolet scale (above TeV) as

$$\mathcal{L} = \mathcal{L}_{\overline{\text{SM}}} + \text{Tr} [\partial_\mu M^\dagger \partial^\mu M] - V(H, M), \quad (1)$$

where $\mathcal{L}_{\overline{\text{SM}}}$ is the SM Lagrangian without the Higgs potential term, and the scale-invariant potential is

$$V(H, M) = \lambda_1 (\text{Tr}[M^\dagger M])^2 + \lambda_2 \text{Tr} [(M^\dagger M)^2] + \lambda_{\text{mix}} |H|^2 \text{Tr}[M^\dagger M] + \lambda_h |H|^4. \quad (2)$$

The $N_f \times N_f$ matrix field M transforms under the chiral symmetry of $U(N_f)_L \times U(N_f)_R$ as

$$M \rightarrow g_L M g_R^\dagger, \quad g_L, g_R \in U(N_f). \quad (3)$$

The chiral $U(N_f) \times U(N_f)$ symmetry is assumed to be broken down to the diagonal subgroup $U(N_f)_V$, as well as the usual electroweak symmetry breaking by the vacuum expectation value (VEV) of the SM Higgs field H :

$$\langle M \rangle = \frac{\phi}{\sqrt{2N_f}} \mathbb{I}, \quad \langle H \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ h \end{pmatrix}. \quad (4)$$

The tree-level potential is thus read off from Eq.(2)

$$V_{\text{tree}} = \frac{1}{4} \left(\lambda_1 + \frac{\lambda_2}{N_f} \right) \phi^4 + \frac{\lambda_{\text{mix}}}{4} h^2 \phi^2 + \frac{\lambda_h}{4} h^4. \quad (5)$$

The vacuum stability at tree level imposes the constraints on couplings as

$$\lambda_h \geq 0, \quad \lambda_{\text{mix}}^2 \leq 4 \left(\lambda_1 + \frac{\lambda_2}{N_f} \right) \lambda_h. \quad (6)$$

Following the Gildener-Weinberg approach [17], in order to find the flat direction, we set

$$\begin{aligned} h &= \chi \sin \theta, \\ \phi &= \chi \cos \theta. \end{aligned} \quad (7)$$

Then the tree-level potential in Eq.(6) is rewritten as

$$V_{\text{tree}} = \frac{\chi^4}{4} \left[\left(\lambda_1 + \frac{\lambda_2}{N_f} \right) \cos^4 \theta + \lambda_{\text{mix}} \cos^2 \theta \sin^2 \theta + \lambda_h \sin^4 \theta \right]. \quad (8)$$

The flat direction condition, which requires V_{tree} to vanish and to be stationary along this direction, yields

$$\tan^2 \theta = \frac{-\lambda_{\text{mix}}}{2\lambda_h}, \quad \lambda_{\text{mix}}^2 = 4 \left(\lambda_1 + \frac{\lambda_2}{N_f} \right) \lambda_h \quad (9)$$

at a certain RG scale μ_{GW} .

Around the VEV in the flat direction, the scalar fields M and H can be expanded as

$$M = \frac{\phi + \sigma + i\eta}{\sqrt{2N_f}} \cdot \mathbb{I}_{N_f \times N_f} + \sum_{a=1}^{N_f^2-1} (\xi^a + i\pi^a) T^a, \quad H = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ h + \tilde{h} \end{pmatrix}, \quad (10)$$

where (σ, η) are the scalar fields and (ξ^a, π^a) are the pseudoscalar fields. T_a are the generators of $SU(N_f)$ group in the fundamental representation and normalized as

$$\text{Tr}(T^a T^b) = \delta^{ab}/2. \quad (11)$$

The field-dependent mass squares for those fields then read

$$\begin{aligned} m_\sigma^2(\chi) &= 3 \left(\lambda_1 + \frac{\lambda_2}{N_f} \right) \chi^2 \cos^2 \theta + \frac{\lambda_{\text{mix}}}{2} \chi^2 \sin^2 \theta = 2 \left(\lambda_1 + \frac{\lambda_2}{N_f} \right) \chi^2 \cos^2 \theta, \\ m_{\xi^a}^2(\chi) &= \left(\lambda_1 + \frac{3\lambda_2}{N_f} \right) \chi^2 \cos^2 \theta + \frac{\lambda_{\text{mix}}}{2} \chi^2 \sin^2 \theta = \frac{2\lambda_2}{N_f} \chi^2 \cos^2 \theta, \\ m_\eta(\chi) &= m_{\pi^a}^2(\chi) = 0, \quad m_{\tilde{h}}^2(\chi) = -\lambda_{\text{mix}} \chi^2 \cos^2 \theta. \end{aligned} \quad (12)$$

The angle θ defined in Eq.(7) simultaneously diagonalizes the $h - \chi$ mixing mass matrix,

$$\mathcal{M}^2 = \begin{pmatrix} 2\lambda_h v_h^2 & \lambda_{\text{mix}} v_h v_\phi \\ \lambda_{\text{mix}} v_h v_\phi & 2(\lambda_1 + \frac{\lambda_2}{N_f}) v_\phi^2 \end{pmatrix}, \quad (13)$$

in such a way that

$$\begin{pmatrix} h_1 \\ h_2 \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \tilde{h} \\ \sigma \end{pmatrix}. \quad (14)$$

This eigenvalue system gives the tree level mass eigenvalues for the mass eigenstates h_1 and h_2 a

$$m_{h_1}^2 = -\lambda_{\text{mix}} v_\chi^2, \quad m_{h_2}^2 = 0. \quad (15)$$

At this moment, h_2 thus becomes massless (called the scalon [17]) having its profile along the flat direction. At the one-loop level, this h_2 acquires a radiative mass as the flat direction is lifted by the quantum corrections, and becomes what is called the pseudo-dilaton due to the explicit-scale symmetry breaking. On the other hand, h_1 has the profile perpendicular to the flat direction, identified as the SM-like Higgs, observed at the LHC with $m_{h_1} = 125$ GeV, which does not develop its mass along the flat direction.

Current experimental limits on the mixing angle θ can be read off from the total signal strength of the Higgs coupling measurements at the LHC [45], which can conservatively be evaluated as

$$\sin^2 \theta = \frac{v_h^2}{v_\chi^2} \lesssim 0.10, \quad \text{i.e.,} \quad v_\chi \gtrsim 778 \text{ GeV}. \quad (16)$$

III. GILDENER-WEINBERG TYPE SCALEGENESIS AND THERMAL CORRECTIONS

Along the flat direction, the one-loop effective potential at zero temperature in the $\overline{\text{MS}}$ scheme and using Landau gauge takes the form ^{#1}

$$V_1(\chi) = A\chi^4 + B\chi^4 \ln \frac{\chi^2}{\mu_{\text{GW}}^2}, \quad (17)$$

with

$$\begin{aligned} A &= \frac{\cos^4 \theta}{16\pi^2} \left[(\lambda_1 + \frac{\lambda_2}{N_f})^2 \left(\ln \left(2(\lambda_1 + \frac{\lambda_2}{N_f}) \cos^2 \theta \right) - \frac{3}{2} \right) + (N_f^2 - 1) \frac{\lambda_2^2}{N_f^2} \left(\ln \left(\frac{2\lambda_2}{N_f} \cos^2 \theta \right) - \frac{3}{2} \right) \right. \\ &\quad \left. + \frac{\lambda_{\text{mix}}^2}{4} \left(\ln (|\lambda_{\text{mix}}| \cos^2 \theta) - \frac{3}{2} \right) \right] + \frac{1}{64\pi^2 v_\chi^4} \sum_{i=t,Z,W^\pm} (-1)^s n_i m_i^4 \left(\ln \frac{m_i^2}{v_\chi^2} - c_i \right), \\ B &= \frac{\cos^4 \theta}{16\pi^2} \left[(\lambda_1 + \frac{\lambda_2}{N_f})^2 + (N_f^2 - 1) \frac{\lambda_2^2}{N_f^2} + \frac{\lambda_{\text{mix}}^2}{4} \right] + \frac{1}{64\pi^2 v_\chi^4} \sum_{i=t,Z,W^\pm} (-1)^s n_i m_i^4, \end{aligned} \quad (18)$$

where $s = 1$ (0) for fermions (bosons), $c_i = \frac{1}{2}$ ($\frac{3}{2}$) for transverse (longitudinal) polarizations of gauge bosons and $c_i = \frac{3}{2}$ for the particles of other species. The numbers of degree of freedom (d.o.f.) n_i for $i = t, Z, W^\pm$ are 12, 3, 6, respectively, and their masses can be written as $m_i^2(\chi) = m_i^2 \frac{\chi^2}{v_\chi^2}$. The nonzero VEV of χ is associated with the renormalization scale μ_{GW} via the stationary condition:

$$\frac{\partial V_1(\chi)}{\partial \chi} = 0 \quad \Rightarrow \quad \mu_{\text{GW}} = v_\chi \exp\left(\frac{A}{2B} + \frac{1}{4}\right). \quad (19)$$

^{#1} We take the Landau gauge for the SM gauge loop contributions. In this case the Nambu-Goldstone boson loop contributions are field-independent at the leading order in the resummed perturbation theory, so that they are decoupled in the effective potential analysis.

Correspondingly, the effective potential can be rewritten as

$$V_1(\chi) = B\chi^4 \left(\ln \frac{\chi^2}{v_\chi^2} - \frac{1}{2} \right) + V_0, \quad V_0 = \frac{Bv_\chi^4}{2} \simeq \frac{\lambda_2^2 v_\chi^4}{32\pi^2} \frac{N_f^2 - 1}{N_f^2}, \quad (20)$$

where V_0 is the vacuum energy determined by $V_1(v_\chi) = 0$, and the last approximation has been made by taking into account the flat direction condition Eq.(9) and the inputs for the Higgs mass and the electroweak scale. The stability condition $B > 0$ gives the lower bound on the coupling λ_2 for any v_χ . The full mass of the dilaton is obtained by the second derivative of $V_1(\chi)$ estimated at the VEV

$$M_\chi^2 = \left. \frac{\partial^2 V_1(\chi)}{\partial \chi^2} \right|_{\chi=v_\chi} = 8Bv_\chi^2. \quad (21)$$

To be phenomenologically realistic, we need to introduce an explicit scale and chiral breaking term, otherwise there are plenty of massless Nambu-Goldstone bosons left in the universe. However, as long as the explicit breaking is small enough that the flat direction can still approximately work, the to-be-addressed characteristic features on the cosmological phase transition and the gravitational wave production will not substantially be altered. Later we will come back to this point in a view of the phenomenological consequences related to the predicted gravitational spectra (see Summary and Discussion).

To study the thermal phase transition of the chiral symmetry and gravitational waves, we turn now to calculating the effective potential at finite temperature. By following the standard procedure, the one-loop thermal corrections are evaluated as

$$V_{1,T}(\chi, T) = \frac{T^4}{2\pi^2} J_B \left(\frac{m_\sigma^2(\chi)}{T^2} \right) + \frac{(N_f^2 - 1) T^4}{2\pi^2} J_B \left(\frac{m_{\xi^i}^2(\chi)}{T^2} \right) + \frac{T^4}{2\pi^2} J_B \left(\frac{m_h^2(\chi)}{T^2} \right) + \frac{T^4}{2\pi^2} \left[\sum_{i=t,Z,W} (-1)^{2s} n_i J_{B/F} \left(\frac{m_i^2(\chi)}{T^2} \right) \right], \quad (22)$$

with the bosonic/fermionic thermal functions

$$J_{B/F}(y^2) = \int_0^\infty dt t^2 \ln \left(1 \mp e^{-\sqrt{t^2 + y^2}} \right). \quad (23)$$

It has been shown that the perturbative expansion will break down since in the high-temperature limit higher loop contributions can grow as large as the tree-level and one-loop terms [46, 47]. To improve the validity of the perturbation, we adopt the *truncated full dressing* resummation procedure [46], which is performed by the replacement $m_i^2(\chi) \rightarrow m_i(\chi)^2 + \Pi_i(T)$ in the full effective potential. The thermal masses $\Pi_i(T)$ are computed as follows:

$$\begin{aligned} \Pi_{\sigma/\xi^i}(T) &= \frac{T^2}{6} \left[(N_f^2 + 1)\lambda_1 + 2N_f\lambda_2 + \frac{\lambda_{\text{mix}}}{4} \right], \quad \Pi_h(T) = T^2 \left(\frac{\lambda_h}{4} + \frac{y_t^2}{4} + \frac{3g^2}{16} + \frac{g'^2}{16} + \frac{\lambda_{\text{mix}}}{24} + \frac{N_f^2}{12} \lambda_{\text{mix}} \right), \\ \Pi_W^L(T) &= \frac{11}{6} g^2 T^2, \quad \Pi_W^T(T) = 0, \quad \Pi_Z^L(T) = \frac{11}{6} (g^2 + g'^2) T^2, \quad \Pi_Z^T(T) = 0. \end{aligned} \quad (24)$$

Here, the corresponding SM Yukawa and gauge couplings are defined as $g = 2m_W/v_h$, $g' = 2\sqrt{m_Z^2 - m_W^2}/v_h$ and $y_t = \sqrt{2}m_t/v_h$. $\Pi_{W/Z}^L(T)$ denote the thermal masses of the longitudinal mode of the gauge bosons, while transverse modes $\Pi_{W/Z}^T(T)$ are protected not to be generated due to the gauge invariance and thus do not receive a mass at leading order in the perturbation theory. Note that while the top quark receives the thermal mass, there are no zero modes $\omega_n = (2n + 1)\pi T$, and as a result no IR divergences appear in the fermion propagator. In general, the contributions from the daisy resummation are less important due to the fact that the phase transition completes well below the critical temperature in the supercooling case. However, the thermal mass with such large number of degrees of freedom, $\mathcal{O}(N_f^2)$, will become ten times as big as the field-dependent mass around the barrier, so that it's necessary to include the daisy contributions.

Taking into account the flat direction condition in Eq.(9) together with the inputs for the Higgs mass m_h , the electroweak scale v_h , and the SM gauge and top quark masses, we see that the total one-loop effective potential, V_1 in Eq.(20) plus $V_{1,T}$ in Eq.(22), is evaluated as a function of λ_2 and v_χ . From the next section, we shall discuss the cosmological phase transition in this parameter space.

IV. COSMOLOGICAL PHASE TRANSITION: ULTRA-SUPERCOOLING AND NUCLEATION

Since it is the CW-type potential, the phase transition becomes of first order and the strength is found to be very strong $\langle\chi\rangle/T_c \simeq \gg 1$ in a wide range of the coupling parameter space, where T_c denotes the critical temperature at which the false and true vacua get degenerated. In the Hubble expanding universe the first order phase transition proceeds by the bubble nucleation. The nucleation rate per unit volume/time of the bubble $\Gamma(T)$ can be computed as

$$\Gamma(T) \simeq T^4 \left(-\frac{S_3(T)}{2\pi T} \right)^{3/2} \exp \left(-\frac{S_3(T)}{T} \right), \quad (25)$$

where the $S_3(T)$ is the $\mathcal{O}(3)$ symmetric bounce action at T :

$$S_3(T) = 4\pi \int_0^\infty d^3r r^2 \left(\frac{1}{2} \left(\frac{d\bar{\chi}}{dr} \right)^2 + V_{\text{eff}}(\bar{\chi}, T) \right). \quad (26)$$

The normalizable bubble profile $\bar{\chi}(r)$ can be obtained by numerically solving the equation of motion,

$$\frac{d^2\bar{\chi}}{dr^2} + \frac{2}{r} \frac{d\bar{\chi}}{dr} = \frac{dV_{\text{eff}}(\bar{\chi}, T)}{d\bar{\chi}}, \quad (27)$$

with the boundary conditions

$$\left. \frac{2}{r} \frac{d\bar{\chi}(r)}{dr} \right|_{r=0} = 0, \quad \bar{\chi}(r)|_{r=\infty} = 0. \quad (28)$$

The nucleation temperature T_n is defined when the bubble nucleation rate for the first time catches up with the Hubble expansion rate:

$$\frac{\Gamma(T_n)}{H(T_n)^4} \sim 1, \quad (29)$$

namely,

$$\frac{S_3(T_n)}{T_n} - \frac{3}{2} \log \left(\frac{S_3(T_n)}{2\pi T_n} \right) \sim 4 \log \frac{T_n}{H(T_n)}, \quad (30)$$

where $H^2(T) = [\Delta V(T) + \rho_{\text{rad}}(T)]/3M_{\text{pl}}^2$, which, for the supercooled phase transition, can be well approximated by the vacuum energy part $H_V = \Delta V(T_n)/3M_{\text{pl}}^2$. In Fig. 1 we display the contour plot of T_n in the parameter space on the (λ_2, v_χ) plane for a couple of reference values for T_n up to 10 GeV. The size of the dark chiral symmetry N_f has been taken to be 8 as a benchmark inspired by the large-flavor walking underlying theory as noted in the Introduction. In the plot we have discarded the case with $T_n < T_{\text{QCD}}$ because in that case instead of the Higgs portal, the QCD phase transition would trigger the EW phase transition, as addressed in the literature [48–51], which is off our current scope.

The contour plot shown in Fig. 1 is qualitatively identical to the one discussed in [52] except for the size of the relevant couplings. The discrepancy comes from the quite different number of the dark-sector particles contributing to the one-loop effective potential: the present case is, say, N_f^2 (see Eqs.(20) and (22)), while the model in the literature only includes one. In particular since a large number of thermal loop contributions are created in the present model, the smaller size of the coupling is sufficient to achieve the nucleation over the Hubble rate. The percolation will process qualitatively in a similar manner as well. In the literature, it has been shown that the null percolation regime, due to too small size of the coupling strength, is fully overlapped with the region with $T_n < T_{\text{QCD}}$, which starts when v_χ gets as large as $\sim 10^4$ GeV, and that the percolation temperature T_p is almost identical to T_n in a wide parameter space as in the contour plot, Fig. 1. These features follow also in the present model.

The GW spectrum resulting from the cosmological-first order phase transition can be parametrized by two parameters α and β . The former α measures the strength of the first order phase transition, which is given by the ratio of the latent heat released from the false vacuum to the radiation energy density:

$$\alpha \equiv \frac{1}{\rho_{\text{rad}}(T_n)} \left(-\Delta V(T_n) + T_n \left. \frac{d\Delta V}{dT} \right|_{T=T_n} \right) \simeq \frac{\Delta V(T_n)}{\rho_{\text{rad}}(T_n)}, \quad (31)$$




FIG. 1: The contour plot of the nucleation temperature T_n in the (λ_2, v_χ) plane with $N_f = 8$. The blue-shaded regime corresponding to the case $T_n < T_{\text{QCD}}$ is discarded in the present study, which will actually be covered with the null percolation regime due to too small coupling size of λ_2 .

where $\Delta V(T)$ is the difference of the effective potential at the true and false vacua. The value of α turns out to be extremely large, $\alpha \gg 1$, for the ultra-supercooled phase transition. The latter parameter β and its normalized one $\tilde{\beta}$ are defined as

$$\tilde{\beta} \equiv \frac{\beta}{H(T_n)} = T_n \left. \frac{d}{dT} \left(\frac{S_3}{T} \right) \right|_{T=T_n}, \quad (32)$$

which measures the duration of the phase transition and the characteristic frequency of the GW through the mean bubble radius at collisions.

Another remark should be made on the characteristic temperature directly related to the peak frequency of GW, that is the reheating temperature T_r . It is usually argued that the estimate of T_r depends on whether the rate of the χ decay to the SM sector (Γ_{dec}) becomes smaller or larger than the Hubble parameter, that we shall classify in more details below.

- (i) In the case with $\Gamma_{\text{dec}} \gg H(T_p)$, where the reheating is supposed to be processed instantaneously after the end of the supercooling, and the whole energy accumulated at the false vacuum is expected to be immediately converted into the radiation. The resulting reheating temperature is determined by assuming the full conversion of the vacuum energy into the radiation [50, 53]

$$\begin{aligned} \rho_{\text{rad}}(T_r) &\simeq \rho_{\text{rad}}(T_p) + \rho_{\text{vac}}(T_p) \simeq \rho_{\text{vac}}(T_p) \\ \Rightarrow T_r &\simeq (1 + \alpha)^{1/4} T_p \simeq \left(\frac{30\Delta V}{\pi^2 g_r} \right)^{1/4} \equiv T_{\text{vac}}, \end{aligned} \quad (33)$$

where in the last line we have taken into account $\alpha \gg 1$ for the ultra-supercooling case.

- (ii) In the case with $\Gamma_{\text{dec}} \ll H(T_p)$, the reheating process is supposed to work so slowly that χ is allowed to roll down and oscillate around the true vacuum until $\Gamma_{\text{dec}} \sim H(T_p)$, where the universe undergoes the matter-dominated period, In that case, the reheating temperature T_r reads [50, 54, 55]

$$T_r \simeq T_{\text{vac}} \sqrt{\frac{\Gamma_{\text{dec}}}{H(T_p)}}. \quad (34)$$

To summarize, the reheating temperature can be expressed as

$$T_r \simeq T_{\text{vac}} \min \left(1, \frac{\Gamma_{\text{dec}}}{H(T_p)} \right)^{1/2}, \quad T_{\text{vac}} = \left(\frac{30\Delta V}{\pi^2 g_r} \right)^{1/4} \simeq \left(\frac{30V_0}{\pi^2 g_r} \right)^{1/4}, \quad (35)$$

which indicates that T_r is controlled by the size of the χ decay rate to the SM. As will be discussed in more details, however, in the present study we do not refer to the size of the χ decay rate in addressing the reheating process as classified in way as above. More crucial to notice is that at any rate, T_r almost simply scales as (see also Eq.(20))

$$T_r \propto T_{\text{vac}} \propto \lambda_2^{1/2} v_\chi. \quad (36)$$

We now discuss the correlation between the cosmological phase transition parameters, α and β , and the nucleation temperature T_n or T_p .

First of all, see the two panels in Fig. 2, which (in the left panel) show the v_χ dependent on T_n varying λ_2 within the allowed regime as in Fig. 1 and (in the right panel) the v_χ dependence for β . In the left panel we observe that T_n linearly grows as v_χ increases for any λ_2 . This trend is closely tied with the scalegenesis feature #2: only one dimensional parameter v_χ is dominated in the theory after the dimensional transmutation, hence at finite temperature the dimensionless bubble action can be almost fully controlled by the dimensionless ratio v_χ/T_n once λ_2 is fixed, which means T_n linearly changes with the variation of v_χ . This can more quantitatively be viewed as follows: given that S_3/T_n is a function of (v_χ/T_n) , which is fixed to $\simeq 140$, then the stationary condition of S_3/T_n leads to $dv_\chi/dT_n = v_\chi/T_n$, hence $T_n \propto v_\chi$. Likewise, one can prove that $\tilde{\beta}$ is insensitive to increasing v_χ as plotted in the right panel of Fig. 2. This trend can be understood by noting that $\tilde{\beta} = T_n \partial(S_3/T_n)/\partial T_n = -v_\chi/T_n \partial(S_3/T_n)/\partial(v_\chi/T_n)$.

Second, we recall the scaling property of T_r in Eq.(36), $T_r \propto v_\chi$. Since both T_n and T_r linearly grow with v_χ , α defined as in Eq.(31) follows the same trend as what $\tilde{\beta}$ does. Thus we have

$$\alpha \sim \text{const.}, \quad \tilde{\beta} \sim \text{const.}, \quad \text{in } v_\chi, \quad (37)$$

Note, furthermore, that for a larger α as in Eq.(33), the slope of T_n with respect to v_χ is almost completely fixed as $\alpha^{-1/4}$. Those cosmological phase transition features are thus characteristic to the (almost) scale invariant setup.

In comparison, in the literature [52] with a similar scale-invariant setup, α and $\tilde{\beta}$ have been evaluated at not $T = T_n$, but at T_p , where the latter does not exhibit a simple scaling property with respect to v_χ unlike the former. Therefore, in the literature α and $\tilde{\beta}$ look sensitive to increase of v_χ . The discrepancy between T_n and T_p is thought to get significant when the GW production with the reheating process taken into account is addressed. A conventional estimate will be based on the instantaneous reheating with $T_r \simeq (1 + \alpha)^{1/4} T_p$ as in Eq.(33). Assuming the entropy conservation involving the reheating epoch one may then get the redshifted GW spectra and frequency at present day, which are scaled with T_p . However, as we will clarify more explicitly in the next section, it turns out that it is not T_p or T_n but T_r that sets the scale of the GW spectra and frequencies.




FIG. 2: Left: The plot of T_n vs. v_χ with λ_2 varied in the allowed range as in Fig. 1; Right: $\tilde{\beta}$ vs. v_χ with the same varied range of λ_2 .

#2 To be phenomenologically realistic, the scale invariance must be approximate even at the classical level. However, the general trends addressed here will not significantly be affected as long as small enough explicit scale breaking is taken into account, as noted also in the previous section. See Summary and Discussion, for more details.

V. GRAVITATIONAL WAVE PRODUCTION: PROSPECTS FOR NANO HZ AND HIGHER FREQUENCY SIGNALS

In this section, we explore the stochastic GW background produced sourced by the strong first order phase transitions produced in the present model setup. The resulting GW signals come from three processes: the collisions of bubble walls, the sound waves in the plasma, and the magnetohydrodynamics turbulence in the plasma, i.e.

$$h^2\Omega_{\text{GW}} \simeq h^2\Omega_{\text{coll}} + h^2\Omega_{\text{sw}} + h^2\Omega_{\text{turb}}. \quad (38)$$

In the strongly supercooled phase transitions, $\alpha \gg 1$, the transition temperatures are low enough that the plasma can be ignored and thus there is negligible friction to stop the bubble wall accelerating before it collides with other bubbles. Therefore, most of the released latent heat goes into the bubble walls and accelerates the bubbles without being bound and hence run away [56, 57] almost with the speed of light $v_w \sim c$. In this case, the bubble collisions give the dominant contribution to the GW spectrum. The efficiency factor κ_{coll} that characterizes the energy transfer between the vacuum energy and the kinetic energy of the bubble wall reads

$$\kappa_{\text{coll}} = 1 - \frac{\alpha_\infty}{\alpha}, \quad \alpha_\infty \simeq \frac{30}{24\pi^2} \frac{\sum_i c_i \Delta m_i^2}{g_p T_p^2}. \quad (39)$$

Here, the sum running over i considers all relativistic particles in the false vacuum and heavy and nonrelativistic ones in the true vacuum: Δm_i^2 is the difference of their (field-dependent) squared masses; g_p corresponds to the effective number of relativistic d.o.f. in the false vacuum; c_i is equal to n_i as in Eq.(18) for bosons and $\frac{1}{2}n_i$ for fermions, with n_i the number of the d.o.f. for species i . In the present model, which predicts the ultra-supercooling, we can safely take $\kappa_{\text{coll}} \sim 1$, which is due to the fact that

$$\frac{\alpha_\infty}{\alpha} \propto \frac{T_p^2}{v_\chi^2} \ll 1. \quad (40)$$

We also need to take into account the redshift factor ($\frac{a_p}{a_0}$) which describes the Hubble evolution acting on the GWs from when it is produced at the epoch corresponding to the scale factor a_p up until today at a_0 . We intercept $\frac{a_p}{a_0}$ by the epoch ($a_r = a(T_r)$), at which the latent heat released from the false vacuum starts to get efficient enough to be converted into the radiation, to be dominated over the universe (regarded as the end of the reheating): $\frac{a_p}{a_0} = \frac{a_p}{a_r} \cdot \frac{a_r}{a_0}$ #3. Since the reheating process is nonadiabatic and cannot simply be described by the thermodynamics, we instead of temperature monitor $\frac{a_p}{a_r}$ in terms of the e-folding number N_e , which is accumulated during the period from when one bubble is nucleated up to the end of the reheating #4. The latter part, $\frac{a_r}{a_0}$, is totally thermal, hence can simply be scaled by the entropy conservation per comoving volume: $s(T_r)a_r^3 = s(T_0)a_0^3$ with the thermal entropy density $s(T) = \frac{2\pi^2}{45} g_{*s}(T)T^3$.

One might think about constructing a couple of the Boltzmann equations with respect to the radiation energy density and the energy densities of χ and the SM Higgs, to which χ decays via the Higgs portal, and evaluate what is like ‘‘matter-radiation’’ equality at which the reheating temperature T_r can be defined. However, this approach cannot go beyond the level of the ensemble average approximation of the dynamics, i.e., sort of a classical level not incorporating the nonadiabatic and nonperturbative relaxation dynamics till the universe is fully radiated starting from the end of the supercooling in the de-Sitter expansion. Thus, there would be still lots of uncertainties involved if one addresses the reheating by naively referring to such Boltzmann equations with the size of the χ decay rate. Therefore, at this moment in our best reasonable way, we parametrize the epoch during the reheating process by the e-folding, as noted above, and simply assume the instantaneous reheating without referring to the size of the χ decay rate as classified in Eqs.(33) and (34).

Thus at this moment we write the redshift factor as

$$\frac{a_p}{a_0} = \frac{a_p}{a_r} \frac{a_r}{a_0} = e^{-N_e} \cdot \frac{g_0^{1/3} \cdot T_0}{g_r^{1/3} \cdot T_r}, \quad (41)$$

#3 For the exponential nucleation phase transitions as in the present model case, the percolation temperature T_p should not be far so much below the temperature at which bubbles collide. Therefore, it is appropriate to choose the temperature at which GWs are produced as the percolation temperature.

#4 Similar evaluation of the redshift factor in terms of the e-folding number has been made in [58], which is applied to the inflationary epoch, not the reheating process that, instead, the authors assumed to be matter dominated or kination dominated.

where $g_0 \simeq 2 + \frac{4}{11} \times \frac{7}{8} \times 2N_{\text{eff}}$ with $N_{\text{eff}} = 3.046$ [45] and g_r are the d.o.f. at the present-day temperature $T_0 = 2.725\text{K}$ and at the reheating temperature, respectively. The effective d.o.f. for the entropy density and of energy density has been assumed to be identical each other, i.e., assuming no extra entropy production other than the one created passing through the reheating epoch.

To make comparison with the conventional formula of the peak frequency, based on inclusion of the entropy conservation during the reheating epoch [57],

$$f_{\text{coll}} \Big|_{\text{conventional}} = 1.65 \times 10^{-5} \text{Hz} \times \left(\frac{0.62}{v_w^2 - 0.1v_w + 1.8} \right) \times \left(\frac{\beta}{H(T_p)} \right) \left(\frac{T_p}{100 \text{ GeV}} \right) \left(\frac{g_p}{100} \right)^{\frac{1}{6}}, \quad (42)$$

we rewrite Eq.(41) as follows:

$$\begin{aligned} \frac{a_p}{a_0} &= \frac{a_p a_r}{a_r a_0} = e^{-N_e} \frac{g_0^{1/3} T_0}{g_r^{1/3} T_r} H(T_r) \frac{H(T_p)}{H(T_r)} \frac{1}{H(T_p)} \\ &= e^{-N_e} \frac{g_0^{1/3} T_0}{g_r^{1/3} T_r} \frac{g_r^{1/2} \pi T_r^2}{3\sqrt{10} M_{\text{pl}}} \frac{H(T_p)}{H(T_r)} \frac{1}{H(T_p)} \\ &= e^{-N_e} \left(\frac{\rho(T_p)}{\rho(T_r)} \right)^{1/2} \frac{100^{7/6} g_0^{1/3} \pi T_0}{3\sqrt{10} M_{\text{pl}}} \left(\frac{g_r}{100} \right)^{1/6} \frac{T_r}{100 \text{ GeV}} \frac{1}{H(T_p)}, \end{aligned} \quad (43)$$

where we have used the Friedmann equations $3M_{\text{pl}}^2 H_r^2 = \rho(T_r) = \frac{\pi^2}{30} g_r T_r^4$ and $3M_{\text{pl}}^2 H_p^2 = \rho(T_p)$. The redshifted peak frequency is thus evaluated as

$$f_{\text{coll}} = e^{-N_e} \left(\frac{\rho_p}{\rho_r} \right)^{1/2} \times 1.65 \times 10^{-5} \text{Hz} \times \left(\frac{0.62}{v_w^2 - 0.1v_w + 1.8} \right) \times \left(\frac{\beta}{H(T_p)} \right) \left(\frac{T_r}{100 \text{ GeV}} \right) \left(\frac{g_p}{100} \right)^{\frac{1}{6}}, \quad (44)$$

which is compared to the conventional formula in Eq.(42):

$$f_{\text{coll}} = e^{-N_e} \left(\frac{\rho_p}{\rho_r} \right)^{1/2} \left(\frac{T_r}{T_p} \right) \times f_{\text{coll}} \Big|_{\text{conventional}}. \quad (45)$$

This implies that even when the GW is produced at the QCD scale or so, the nano Hz frequency is unlikely to be realized. One might still suspect that if an inflationary stage, after the tunneling for the flat enough CW-type potential, is present, it could suppress the peak frequency due to a huge amount of the accumulated e-folding N_e , so that the nano Hz signal could be generated. However, this would not be the case: the tunneling exit point is supposed to be within the inflation region, which requires that the coupling λ_2 is tiny enough that no percolation takes place and the stationary condition $B > 0$ in Eq.(18) is also violated, thus no bubble collision, nor GWs induced from the first-order phase transition.

Thus it is clarified that the peak frequency is shifted to higher by scaling with T_r .

GW spectra sourced from the bubble wall are evaluated based on the simulations of bubble wall, leading to the following fitting function with the conventional redshift incorporated [59]:

$$\Omega_{\text{coll}} h^2 \Big|_{\text{conventional}} = 1.67 \times 10^{-5} \left(\frac{H(T_p)}{\beta} \right)^2 \left(\frac{\kappa_{\text{coll}} \alpha}{1 + \alpha} \right)^2 \left(\frac{100}{g_p} \right)^{\frac{1}{3}} \times \left(\frac{0.11v_w^3}{0.42 + v_w^2} \right) S_{\text{coll}}(f), \quad (46)$$

where $S_{\text{coll}}(f)$ parametrizes the spectral shape, which is given also by the fitting procedure to be [59]

$$S_{\text{coll}}(f) = \frac{3.8 \left(\frac{f}{f_{\text{coll}}} \right)^{2.8}}{1 + 2.8 \left(\frac{f}{f_{\text{coll}}} \right)^{3.8}}. \quad (47)$$

These GWs also get redshifted similarly to the peak frequency as

$$\Omega_{\text{coll}} h^2 = e^{-4N_e} \left(\frac{\rho_p}{\rho_r} \right) \times \Omega_{\text{coll}} h^2 \Big|_{\text{conventional}}, \quad (48)$$




FIG. 3: Plots of GW energy spectra for several benchmarks of the present model with $N_f = 8$ in comparison with future prospected detector sensitivities [60, 61].

which generically tends to get suppressed by the e-folding N_e and (ρ_p/ρ_r) .

From the refined formulae Eqs.(45) and (48), we see that f_{coll} linearly grows as v_χ because T_r gets larger as v_χ gets larger as in Eq.(36), while $\Omega_{\text{GW}}h^2$ is insensitive to increase of v_χ .

In Fig. 3 we plot the GW spectra for several values of (λ_2, v_χ) for $N_f = 8$ with the instantaneous reheating ($\rho_p = \rho_r$ and $N_e = 0$) assumed in Eqs.(45) and (48). together with the prospected sensitivity curves [60, 61]. As evident from the newly proposed formula on the peak frequency in Eq.(45), the GW peaks are generically shifted toward higher due to the significant dependence of $T_r(\propto v_\chi)$, in comparison with a similar scalegenesis prediction in the literature [52]. In fact, the displayed three GW signals have been sourced from the ultra-supercooled first-order phase transitions at lower nucleation/percolation temperatures $T_p = 100$ MeV (for blue curve) and 10 GeV (for both black and red curves), which are typically thought to be low enough to realize the GW peak signals around nano Hz simply following the conventional formula in Eq.(42). Nevertheless, the produced signals following the proposed formula Eq.(45) peak at much higher frequencies, say, ranged from 10^{-4} Hz to 10^{-2} Hz, as seen from Fig. 3. This is manifested by the linear T_r dependence in the peak frequency formula, Eq.(45), in which currently we have $T_r \simeq 41$ GeV (for blue curve), 70 GeV (for black curve), and 5.2 TeV (for red curve), respectively.

Consequently, even the smaller v_χ (i.e. lower new physics scale ~ 1 TeV) can easily reach the LISA prospect and other higher frequency prospects (BBO and DECIGO, and so forth), though the GW signals would generically be as small as the lower bounds of the prospects.

On the other side of the same coin, we can conclude that nano (or less nano) Hz signals cannot be reached by the ultra-supercooled scalegenesis of this sort, because of the inevitable “blueshift” of the GW frequency: if the nano Hz signal is imposed to realize, i.e., simply $T_r \sim 100$ MeV, then Eq.(34) requires $v_\chi \sim 1$ TeV with $g_r = \mathcal{O}(100)$, which leads to $\alpha \gg 1$, hence extremely tiny T_n or T_p . Thus T_p would be required to be around \sim MeV or less, which is actually inside the excluded regime with no percolation (See Fig. 1).

This is the generic prediction of the Higgs portal scalegenesis with ultra-supercooling, not specialized to the large N_f dark chiral setup.

The large N_f models, e.g., with $N_f = 8$, as what we currently focus on, tends to make $\tilde{\beta}$ larger, while the N_f dependence in α gets almost insensitive in the GW signals sourced from collisions, because anyhow the ultra-supercooling merely provides huge α as noted around Eqs.(39) and (40) to give $\kappa_{\text{coll}} \sim 1$ irrespective to the precise large number of N_f . Thus, the large N_f case tends to further “blueshift” the peak frequency of the GW and make the GW signal strength smaller, due to the produced large $\tilde{\beta}$.

VI. SUMMARY AND DISCUSSION

In this paper, we have discussed GW spectra predicted from the electroweak scalegenesis of the Higgs portal type, with a particular interest in the dark-scalar sector having a large number of dark chiral flavors. We have modeled the dark-sector by the linear sigma model description, which has the chiral $U(N_f) \times U(N_f)$ symmetry and is coupled to the SM Higgs via the Higgs portal with keeping the classical scale invariance. Working on the Gildener-Weinberg mechanism, we have observed that models of this class undergoes a strong first-order thermal phase transition and ultra-supercooling. Evaluation of the bubble nucleation/percolation has clarified the possibility of generation of the GWs sourced from the supercooling, which is accessible in a wide parameter space of the model (see Fig 1). We also clarified the characteristic features for α and $\tilde{\beta}$ stemming from the consequence of the Higgs-portal scalegenesis irrespective to the case with or without large dark-sector flavors (Eq.(37)).

Our particular emphasis has been provided in evaluation of the reheating and relaxation processes, which necessarily and significantly arise from the ultra-supercooling predicted from the scale-invariant setup of the present model class. Being generically non-thermal and nonadiabatic, we proposed to parametrize such a reheating epoch in terms of the e-folding number, not assuming the conventional entropy conservation as in thermodynamic cosmology. This approach has derived refined formulae for the redshift effects on the peak frequencies of the produced GWs and the GW signal strengths: equations (45) and (48). Particularly, it has been clarified that the peak frequencies finally get “blueshifted”, in comparison with the conventional approach based on the thermal entropy conservation including the reheating epoch (see Eq.(45)).

We then observed that the ultra-supercooling predicted from the Higgs-portal scalegenesis generically yields none of nano Hz GW signal, instead, prefers to give the higher frequency signals (Fig. 3), which still keeps the future prospected detection sensitivity, like at LISA, BBO, and DECIGO, etc. We also found that with large flavors in the dark sector, the GW signals are made further smaller and the peak frequencies higher.

In closing, we give comments on the explicit chiral and scale breaking, which is, to be phenomenologically viable, necessary to be incorporated into the present model. The explicit breaking needs to be so small that the flat direction we have worked on is still approximately operative in searching for the true vacuum, as has been addressed in the literature [23, 24, 62, 63]. Given such a small enough breaking term, the model will predict light pseudo Nambu-Goldstone bosons (dark pions) with the number of N_f^2 or $N_f^2 - 1$ (with one decoupled due to the axial anomaly), depending on the underlying theory for the linear sigma model description.

The dark-pion mass term plays a role of the tadpole term for the χ potential does the job, which will make the false vacuum shifted depending on temperature, until the supercooling ends, as was clarified in [24]. The origin of such a chiral and scale breaking could be linked with presence of the gravitational dilaton which couples to the underlying dark QCD fermion bilinear $\bar{F}F$: $V_{\text{tadpole}} = \mathcal{C} \cdot e^{\varphi} \bar{F}F \approx -\mathcal{C} \cdot e^{\varphi} \langle -\bar{F}F \rangle \cdot \text{Tr}[M^\dagger + M] = -\mathcal{C} \cdot \langle -\bar{F}F \rangle \cdot \chi \cos \theta + \dots$ with φ and \mathcal{C} being the dilaton and a constant coupling, respectively. As long as the size of the potential coupling λ_2 is sizable enough as in the desired regime displayed in Fig. 1, both the percolation and nucleation can be realized to not substantially affect what we have addressed and clarified so far in the scale-invariant limit.

The precise size of the dark pion mass highly depends on the underlying theory. Generically the dark pions will be stable to be the dark matter candidate otherwise the dark isospin symmetry is violated. The dark pions ($\pi_D^A \equiv (\eta, \pi^a)$ in Eq.(10)) are expected to be light and can couple to the SM particles via the Higgs portal with λ_{mix} , which is $< 10^{-3}$ for $v_\chi \sim 1$ TeV (See Eq.(15)). Therefore the dark pions can be pair-produced at the Large Hadron Collider (LHC) experiment via the Higgs production processes like $pp \rightarrow h \rightarrow \chi^* \rightarrow \pi_D^A \pi_D^A$ with the final state identified as a large missing energy. The cross section is fixed by the size of v_χ , λ_2 , m_{π_D} , and N_f . Hunting the pseudodilaton χ is also accessible at the LHC-run 3 via the Higgs portal coupling. Those would be interesting study in light of the LHC-run 3 with high luminosity, hence would provide a complementary phenomenological consequence of the Higgs-portal scalegenesis with large flavors, to be explored elsewhere.

In the thermal history, the dark pions as dark matters can be produced via the annihilation into the lighter SM particles, presumably, diphoton, via the Higgs portal coupling including the χ exchange: $\pi_D^A \pi_D^A \rightarrow \chi^* - h^* \rightarrow \gamma\gamma$. This annihilation cross section roughly goes like $\langle \sigma_{\text{ann}} v \rangle \sim \frac{(\lambda_2 v_\chi)^2 \cdot (\lambda_{\text{mix}} v_\chi v_h)^2}{m_\chi^4 m_h^4} \cdot \text{Br}[h \rightarrow \gamma\gamma]$. Still, however, this process needs to assume the portal coupling to be thermalized with the SM thermal plasma, which can be evaluated by equating the conversion rate $\Gamma_{\chi \rightarrow hh}$ and the Hubble rate. The conversion rate is roughly $\Gamma_{\chi \rightarrow h} = n_\chi \langle \sigma v \rangle \sim \frac{\lambda_{\text{mix}}^2}{m_\chi^2} (m_\chi T)^{3/2} e^{-m_\chi/T}$ with $m_\chi \geq \mathcal{O}(\text{TeV})$ and $|\lambda_{\text{mix}}| = m_h^2/v_\chi^2 \lesssim 10^{-3}$ for $v_\chi \gtrsim 1$ TeV. Comparing this with $H \sim \sqrt{g_*(T)} \cdot T^2/M_{\text{pl}}$, we see that the conversion is thermally decoupled at $T < 50$ GeV for $g_*(T) \sim 100$. As v_χ gets larger, the decoupling temperature will be lower. If the conversion is still operative, the freeze-out of the dark pion annihilation is expected to happen when $m_{\pi_D}/T \sim 20$, i.e., $m_{\pi_D} \sim 1$ TeV for $v_\chi \sim 1$ TeV, which is too big to be consistent with the small-explicit breaking criterion. Since the SM Higgs is a nonrelativistic particle at $T < 50$ GeV, there would be no chance to make the freeze-in mechanism [64, 65] work for the dark pion as well. Thus the thermal relic abundance is unlikely

efficiently produced, hence is expected not to be the dark matter candidate.

The dark pions could still be produced non-thermally via the coherent oscillation mechanism, just like axionlike particles. Since the dark pions can develop the potential of the cosine form, $V(\pi_D) = m_\pi^2 f_{\pi_D}^2 (1 - \cos \frac{\pi_D}{f_{\pi_D}})$, where f_{π_D} denotes the decay constant, perhaps, $\ll v_\chi$ for many flavor QCD case [23, 24]. The size of the energy density per flavor accumulated by the coherent oscillation depends on f_{π_D} , m_{π_D} , and the initial phase of the dark pion (so-called the misalignment angle), which would be highly subject to the modeling of the underlying theory, to be pursued in another publication.

Acknowledgments

This work was supported in part by the National Science Foundation of China (NSFC) under Grant No.11747308, 11975108, 12047569, and the Seeds Funding of Jilin University (S.M.), and Toyama First Bank, Ltd (H.I.).

-
- [1] G. Aad *et al.* [ATLAS], Phys. Lett. B **716**, 1-29 (2012) doi:10.1016/j.physletb.2012.08.020 [arXiv:1207.7214 [hep-ex]].
 - [2] S. Chatrchyan *et al.* [CMS], Phys. Lett. B **716**, 30-61 (2012) doi:10.1016/j.physletb.2012.08.021 [arXiv:1207.7235 [hep-ex]].
 - [3] W. A. Bardeen, FERMILAB-CONF-95-391-T.
 - [4] M. Shaposhnikov and C. Wetterich, Phys. Lett. B **683**, 196-200 (2010) doi:10.1016/j.physletb.2009.12.022 [arXiv:0912.0208 [hep-th]].
 - [5] C. Wetterich and M. Yamada, Phys. Lett. B **770**, 268-271 (2017) doi:10.1016/j.physletb.2017.04.049 [arXiv:1612.03069 [hep-th]].
 - [6] A. Eichhorn, Y. Hamada, J. Lumma and M. Yamada, Phys. Rev. D **97**, no.8, 086004 (2018) doi:10.1103/PhysRevD.97.086004 [arXiv:1712.00319 [hep-th]].
 - [7] J. M. Pawłowski, M. Reichert, C. Wetterich and M. Yamada, Phys. Rev. D **99**, no.8, 086010 (2019) doi:10.1103/PhysRevD.99.086010 [arXiv:1811.11706 [hep-th]].
 - [8] C. Wetterich, [arXiv:1901.04741 [hep-th]].
 - [9] S. Matsuzaki, H. Ohki and K. Yamawaki, [arXiv:1608.03691 [hep-ph]].
 - [10] K. Yamawaki, [arXiv:1803.07271 [hep-ph]].
 - [11] R. Hempfling, Phys. Lett. B **379**, 153-158 (1996) doi:10.1016/0370-2693(96)00446-7 [arXiv:hep-ph/9604278 [hep-ph]].
 - [12] K. A. Meissner and H. Nicolai, Phys. Lett. B **648**, 312-317 (2007) doi:10.1016/j.physletb.2007.03.023 [arXiv:hep-th/0612165 [hep-th]].
 - [13] L. Alexander-Nunneley and A. Pilaftsis, JHEP **09**, 021 (2010) doi:10.1007/JHEP09(2010)021 [arXiv:1006.5916 [hep-ph]].
 - [14] A. Farzinia, H. J. He and J. Ren, Phys. Lett. B **727**, 141-150 (2013) doi:10.1016/j.physletb.2013.09.060 [arXiv:1308.0295 [hep-ph]].
 - [15] E. Gabrielli, M. Heikinheimo, K. Kannike, A. Racioppi, M. Raidal and C. Spethmann, Phys. Rev. D **89**, no.1, 015017 (2014) doi:10.1103/PhysRevD.89.015017 [arXiv:1309.6632 [hep-ph]].
 - [16] S. R. Coleman and E. J. Weinberg, Phys. Rev. D **7**, 1888-1910 (1973) doi:10.1103/PhysRevD.7.1888
 - [17] E. Gildener and S. Weinberg, Phys. Rev. D **13**, 3333 (1976) doi:10.1103/PhysRevD.13.3333
 - [18] F. Sannino and J. Virkajärvi, Phys. Rev. D **92**, no.4, 045015 (2015) doi:10.1103/PhysRevD.92.045015 [arXiv:1505.05872 [hep-ph]].
 - [19] S. Iso and Y. Orikasa, PTEP **2013**, 023B08 (2013) doi:10.1093/ptep/pts099 [arXiv:1210.2848 [hep-ph]].
 - [20] M. Hashimoto, S. Iso and Y. Orikasa, Phys. Rev. D **89**, no.5, 056010 (2014) doi:10.1103/PhysRevD.89.056010 [arXiv:1401.5944 [hep-ph]].
 - [21] Y. Hamada, K. Tsumura and M. Yamada, Eur. Phys. J. C **80**, no.5, 368 (2020) doi:10.1140/epjc/s10052-020-7929-3 [arXiv:2002.03666 [hep-ph]].
 - [22] H. Ishida, S. Matsuzaki and R. Ouyang, Chin. Phys. C **44**, no.11, 111002 (2020) doi:10.1088/1674-1137/abb07f [arXiv:1907.09176 [hep-ph]].
 - [23] H. Ishida and S. Matsuzaki, Phys. Lett. B **804**, 135390 (2020) doi:10.1016/j.physletb.2020.135390 [arXiv:1912.09740 [hep-ph]].
 - [24] H. X. Zhang, S. Matsuzaki and H. Ishida, Phys. Lett. B **846**, 138256 (2023) doi:10.1016/j.physletb.2023.138256 [arXiv:2306.15471 [hep-ph]].
 - [25] T. Appelquist *et al.* [LSD], Phys. Rev. D **90**, no.11, 114502 (2014) doi:10.1103/PhysRevD.90.114502 [arXiv:1405.4752 [hep-lat]].
 - [26] Y. Aoki *et al.* [LatKMI], Phys. Rev. D **89**, 111502 (2014) doi:10.1103/PhysRevD.89.111502 [arXiv:1403.5000 [hep-lat]].
 - [27] A. Hasenfratz, D. Schaich and A. Veernala, JHEP **06**, 143 (2015) doi:10.1007/JHEP06(2015)143 [arXiv:1410.5886 [hep-lat]].
 - [28] Y. Aoki *et al.* [LatKMI], Phys. Rev. D **96**, no.1, 014508 (2017) doi:10.1103/PhysRevD.96.014508 [arXiv:1610.07011 [hep-lat]].
 - [29] Y. Aoki *et al.* [LatKMI], Phys. Rev. D **96**, no.1, 014508 (2017) doi:10.1103/PhysRevD.96.014508 [arXiv:1610.07011 [hep-lat]].

- [30] T. Appelquist, R. C. Brower, G. T. Fleming, A. Hasenfratz, X. Y. Jin, J. Kiskis, E. T. Neil, J. C. Osborn, C. Rebbi and E. Rinaldi, *et al.* Phys. Rev. D **93**, no.11, 114514 (2016) doi:10.1103/PhysRevD.93.114514 [arXiv:1601.04027 [hep-lat]].
- [31] T. Appelquist *et al.* [Lattice Strong Dynamics], Phys. Rev. D **99**, no.1, 014509 (2019) doi:10.1103/PhysRevD.99.014509 [arXiv:1807.08411 [hep-lat]].
- [32] K. Miura, H. Ohki, S. Otani and K. Yamawaki, JHEP **10**, 194 (2019) doi:10.1007/JHEP10(2019)194 [arXiv:1811.05670 [hep-ph]].
- [33] K. Yamawaki, M. Bando and K. i. Matumoto, Phys. Rev. Lett. **56**, 1335 (1986) doi:10.1103/PhysRevLett.56.1335
- [34] M. Bando, K. i. Matumoto and K. Yamawaki, Phys. Lett. B **178**, 308-312 (1986) doi:10.1016/0370-2693(86)91516-9
- [35] G. Agazie *et al.* [NANOGrav], Astrophys. J. Lett. **951**, no.1, L8 (2023) doi:10.3847/2041-8213/acdac6 [arXiv:2306.16213 [astro-ph.HE]].
- [36] H. Xu, S. Chen, Y. Guo, J. Jiang, B. Wang, J. Xu, Z. Xue, R. N. Caballero, J. Yuan and Y. Xu, *et al.* Res. Astron. Astrophys. **23**, no.7, 075024 (2023) doi:10.1088/1674-4527/acdfa5 [arXiv:2306.16216 [astro-ph.HE]].
- [37] J. Antoniadis *et al.* [EPTA and InPTA:], Astron. Astrophys. **678**, A50 (2023) doi:10.1051/0004-6361/202346844 [arXiv:2306.16214 [astro-ph.HE]].
- [38] D. J. Reardon, A. Zic, R. M. Shannon, G. B. Hobbs, M. Bailes, V. Di Marco, A. Kapur, A. F. Rogers, E. Thrane and J. Askew, *et al.* Astrophys. J. Lett. **951**, no.1, L6 (2023) doi:10.3847/2041-8213/acdd02 [arXiv:2306.16215 [astro-ph.HE]].
- [39] P. Amaro-Seoane *et al.* [LISA], [arXiv:1702.00786 [astro-ph.IM]].
- [40] C. Caprini, M. Chala, G. C. Dorsch, M. Hindmarsh, S. J. Huber, T. Konstandin, J. Kozaczuk, G. Nardini, J. M. No and K. Rummukainen, *et al.* JCAP **03**, 024 (2020) doi:10.1088/1475-7516/2020/03/024 [arXiv:1910.13125 [astro-ph.CO]].
- [41] V. Corbin and N. J. Cornish, Class. Quant. Grav. **23**, 2435-2446 (2006) doi:10.1088/0264-9381/23/7/014 [arXiv:gr-qc/0512039 [gr-qc]].
- [42] G. M. Harry, P. Fritschel, D. A. Shaddock, W. Folkner and E. S. Phinney, Class. Quant. Grav. **23**, 4887-4894 (2006) [erratum: Class. Quant. Grav. **23**, 7361 (2006)] doi:10.1088/0264-9381/23/15/008
- [43] S. Kawamura, T. Nakamura, M. Ando, N. Seto, K. Tsubono, K. Numata, R. Takahashi, S. Nagano, T. Ishikawa and M. Musha, *et al.* Class. Quant. Grav. **23**, S125-S132 (2006) doi:10.1088/0264-9381/23/8/S17
- [44] K. Yagi and N. Seto, Phys. Rev. D **83**, 044011 (2011) [erratum: Phys. Rev. D **95**, no.10, 109901 (2017)] doi:10.1103/PhysRevD.83.044011 [arXiv:1101.3940 [astro-ph.CO]].
- [45] R. L. Workman *et al.* [Particle Data Group], PTEP **2022**, 083C01 (2022) doi:10.1093/ptep/ptac097
- [46] D. Curtin, P. Meade and H. Ramani, Eur. Phys. J. C **78**, no.9, 787 (2018) doi:10.1140/epjc/s10052-018-6268-0 [arXiv:1612.00466 [hep-ph]].
- [47] E. Senaha, Symmetry **12**, no.5, 733 (2020) doi:10.3390/sym12050733
- [48] E. Witten, Nucl. Phys. B **177**, 477-488 (1981) doi:10.1016/0550-3213(81)90182-6
- [49] S. Iso, P. D. Serpico and K. Shimada, Phys. Rev. Lett. **119**, no.14, 141301 (2017) doi:10.1103/PhysRevLett.119.141301 [arXiv:1704.04955 [hep-ph]].
- [50] T. Hambye, A. Strumia and D. Teresi, JHEP **08**, 188 (2018) doi:10.1007/JHEP08(2018)188 [arXiv:1805.01473 [hep-ph]].
- [51] L. Sagunski, P. Schicho and D. Schmitt, Phys. Rev. D **107**, no.12, 123512 (2023) doi:10.1103/PhysRevD.107.123512 [arXiv:2303.02450 [hep-ph]].
- [52] K. Kawana, Phys. Rev. D **105**, no.10, 103515 (2022) doi:10.1103/PhysRevD.105.103515 [arXiv:2201.00560 [hep-ph]].
- [53] J. Ellis, M. Lewicki and J. M. No, JCAP **04**, 003 (2019) doi:10.1088/1475-7516/2019/04/003 [arXiv:1809.08242 [hep-ph]].
- [54] J. Ellis, M. Lewicki, J. M. No and V. Vaskonen, JCAP **06**, 024 (2019) doi:10.1088/1475-7516/2019/06/024 [arXiv:1903.09642 [hep-ph]].
- [55] M. Kierkla, A. Karam and B. Swiezevska, JHEP **03**, 007 (2023) doi:10.1007/JHEP03(2023)007 [arXiv:2210.07075 [astro-ph.CO]].
- [56] D. Bodeker and G. D. Moore, JCAP **05**, 009 (2009) doi:10.1088/1475-7516/2009/05/009 [arXiv:0903.4099 [hep-ph]].
- [57] C. Caprini, M. Hindmarsh, S. Huber, T. Konstandin, J. Kozaczuk, G. Nardini, J. M. No, A. Petiteau, P. Schwaller and G. Servant, *et al.* JCAP **04**, 001 (2016) doi:10.1088/1475-7516/2016/04/001 [arXiv:1512.06239 [astro-ph.CO]].
- [58] H. An, X. Tong and S. Zhou, Phys. Rev. D **107**, no.2, 023522 (2023) doi:10.1103/PhysRevD.107.023522 [arXiv:2208.14857 [hep-ph]].
- [59] S. J. Huber and T. Konstandin, JCAP **09**, 022 (2008) doi:10.1088/1475-7516/2008/09/022 [arXiv:0806.1828 [hep-ph]].
- [60] C. J. Moore, R. H. Cole and C. P. L. Berry, Class. Quant. Grav. **32**, no.1, 015014 (2015) doi:10.1088/0264-9381/32/1/015014 [arXiv:1408.0740 [gr-qc]].
- [61] K. Schmitz, JHEP **01**, 097 (2021) doi:10.1007/JHEP01(2021)097 [arXiv:2002.04615 [hep-ph]].
- [62] S. Iso, K. Kohri and K. Shimada, Phys. Rev. D **91**, no.4, 044006 (2015) doi:10.1103/PhysRevD.91.044006 [arXiv:1408.2339 [hep-ph]].
- [63] G. Cacciapaglia, D. Y. Cheong, A. Deandrea, W. Isnard and S. C. Park, JCAP **10**, 063 (2023) doi:10.1088/1475-7516/2023/10/063 [arXiv:2307.01852 [hep-ph]].
- [64] M. Blennow, E. Fernandez-Martinez and B. Zaldivar, JCAP **01**, 003 (2014) doi:10.1088/1475-7516/2014/01/003 [arXiv:1309.7348 [hep-ph]].
- [65] L. J. Hall, K. Jedamzik, J. March-Russell and S. M. West, JHEP **03**, 080 (2010) doi:10.1007/JHEP03(2010)080 [arXiv:0911.1120 [hep-ph]].