# Factor Importance Ranking and Selection using Total Indices

Chaofan Huang and V. Roshan Joseph<sup>\*</sup>

H. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, GA, 30332

January 15, 2024

#### Abstract

Factor importance measures the impact of each feature on output prediction accuracy. Many existing works focus on the model-based importance, but an important feature in one learning algorithm may hold little significance in another model. Hence, a factor importance measure ought to characterize the feature's predictive potential without relying on a specific prediction algorithm. Such algorithm-agnostic importance is termed as *intrinsic importance* in Williamson et al. (2023), but their estimator again requires model fitting. To bypass the modeling step, we present the equivalence between predictiveness potential and total Sobol' indices from global sensitivity analysis, and introduce a novel consistent estimator that can be directly estimated from noisy data. Integrating with forward selection and backward elimination gives rise to FIRST, Factor Importance Ranking and Selection using Total (Sobol') indices. Extensive simulations are provided to demonstrate the effectiveness of FIRST on regression and binary classification problems, and a clear advantage over the state-of-the-art methods.

Keywords: Sensitivity analysis; Sobol' indices; Variable importance; Variable selection.

<sup>\*</sup>Corresponding author: roshan@gatech.edu.

### 1 Introduction

An important problem in statistics and machine learning is to understand which input variables affect the output variable and to rank them according to their importance based on the available input-output data. Traditionally, this is done by fitting a linear regression model and using the standardized coefficients (t-values or p-values) to assess the significance of the variables. More advanced techniques exist that decomposes the overall  $R^2$  into sum of contributions from each variable (Grömping, 2007). Nevertheless, the overly simplistic linear regression model would fail to capture non-linearities and higher-order interactions, resulting in misleading estimates for the factor importance, and more importantly, the chance of missing important variables.

With the advent of nonparameteric regression and machine learning models, e.g., generalized additive model (Hastie, 2017; Lin and Zhang, 2006), Gaussian process (Rasmussen et al., 2006), tree models (Breiman, 2001; Friedman, 2001), neural networks (LeCun et al., 2015), etc., more sophisticated learning algorithms are recommended over linear regression to improve the prediction performance. Many factor importance measures are also proposed to reflect how much the predictive accuracy of a given algorithm depends on a particular variable (Breiman, 2001; Gevrey et al., 2003; Grömping, 2009; Wei et al., 2015; Lundberg and Lee, 2017; Fisher et al., 2019; Murdoch et al., 2019; Wojtas and Chen, 2020), i.e., the decrease in predictiveness (e.g.  $R^2$  for regression and accuracy for classification) when a variable is dropped or permuted. While such importance measures provide some interpretability about the black-box model, there is a chance that these models would fail to capture the complex input-output relationship correctly, and thus potentially leading to biased importance measures. Sometimes it could also happen that several learning algorithms fit the data almost equally well, yet each provide different importance measures, complicating the determination of the correct importance. To reduce the risk of model dependence, an ensemble of models is also investigated (Fisher et al., 2019; Williamson et al., 2021, 2023), but at a higher computational cost, especially for big data.

Given the concern and limitation of the model-based approaches, Williamson et al. (2023) argue that the factor importance should be *algorithm-agnostic*. It should measure the population-level predictive potential of the factor, without being tied to any prediction algorithm. This is referred to as the *intrinsic importance* in Williamson et al. (2023), which is defined as the predictiveness gap between the *optimal* prediction function using all the variables versus that fitted on the subset after removing the factors of interest. Albeit being model-agnostic, the estimator from Williamson et al. (2023) again involves fitting predictive models twice, one on all variables and the other on the carefully selected subset. They claim that with flexible enough ensemble of learning techniques, the risk of systematic bias from model misspecification can be minimized. Further improvement to detect null importance requires sample-splitting and cross-fitting, which will result in larger computational expense.

On the other hand, factor importance is also of interest in sensitivity analysis (Da Veiga et al., 2021), where the objective is to identify inputs of computer code that have the most substantial influence on the output and fix the ones that the output is not sensitive to. Various techniques have been proposed, including Morris screening method (Morris, 1991), variance-based (total) Sobol' indices (Sobol', 1993), etc. Recently, Hart and Gremaud (2018) draw the equivalence between the total Sobol' index and the predictiveness potential of a factor, more specifically, the optimal approximation error introduced without using the factor of interest. This demonstrates the potential of estimating the intrinsic factor importance via total Sobol' indices. Though many efficient Monte Carlo based methods are available for the total Sobol' indices estimation, they all require having assess to the computer model. One solution is to construct a surrogate model (Gaussian process) and analytically derive the indices assumption rarely holds in real data applications, not to mention about the challenges associated with the model fitting.

To overcome the aforementioned limitations, we propose an efficient algorithm for estimating total (Sobol') indices directly from the randomly scattered noisy data. Our method is built upon the nearest-neighbor estimator proposed by Broto et al. (2020). We further adapt our estimation algorithm with forward selection and backward elimination to propose FIRST, Factor Importance Ranking and Selection using Total (Sobol') indices. To the best of our knowledge, FIRST is the first factor importance procedure that does not involve any model fitting, and hence safeguarding against the danger of model misspecification.

There is also another line of research aiming to assess the significance of the variables nonparametrically (Székely et al., 2007; Li et al., 2012; Chatterjee, 2021; Azadkia and Chatterjee, 2021; Huang et al., 2022). Although the measures from these techniques can be used for factor selection, the ranking of the importance is not reliable especially when the inputs are correlated. Other approaches consider the aggregation of local importance such as derivative (Bai et al., 2014; Ye and Xie, 2012; Yang et al., 2016), sharing similar spirit of the derivative-based sensitivity measure (Sobol' and Kucherenko, 2009). Again this kind of measure is valid for factor selection, but when there is strong non-linearity in the model, Sobol' and Kucherenko (2009) point out the derivative-based measure could yield misleading conclusion about the factor importance.

The article is organized as follows. Seciont 2 reviews the total Sobol' indices and presents its equivalence to the intrinsic importance measures. Section 3 discusses a novel procedure for estimating the total Sobol' indices directly from noisy data. Section 4 proposes FIRST for factor importance and selection. Extensive numerical comparison on both synthetic and real data are provided in Section 5 for regression and Section 6 for binary classification. We conclude the article with some remarks in Section 7.

### 2 Factor Importance and Total Sobol' Indices

This section we define the factor importance through its predictive potential, then review the total Sobol' indices (Sobol', 1993; Homma and Saltelli, 1996; Sobol', 2001) from the global sensitivity analysis literature, and last demonstrate the equivalence between the two.

Let  $\mathbf{X} = (X_1, \ldots, X_p) \in \mathcal{X} \subseteq \mathbb{R}^p$  be a *p*-dimensional random vectors with joint distribution  $\mathbb{P}_{\mathcal{X}}$ . Let the scalar output  $Y = f^*(\mathbf{X}) \in \mathcal{Y} \subseteq \mathbb{R}$  be some deterministic function of the input  $\mathbf{X}$ , where  $f^* \in \mathcal{F}$ , the class of functions from  $\mathcal{X} \to \mathcal{Y}$  that are square integrable over  $\mathbb{P}_{\mathcal{X}}$ . In this section, assume that the model  $f^*$  is known and there is no random error associated with Y. The randomness in Y is exclusively from the uncertainty in  $\mathbf{X}$ .

Let us first formally define the notations. For any non-empty  $u \subset 1: p$ , let  $\mathbf{X}_u$  be the

elements of  $\mathbf{X}$  with index u,  $\mathcal{X}_u$  be the sample space of  $\mathbf{X}_u$ , and  $\mathcal{F}_u := \{f \in \mathcal{F} : f(\mathbf{x}) = f(\mathbf{x}') \text{ for any } \mathbf{x}, \mathbf{x}' \in \mathcal{X} \text{ such that } \mathbf{x}_u = \mathbf{x}'_u\}$  be the class functions in  $\mathcal{F}$  that depend only on the elements of the input with index in u. Denote |u| be its cardinality,  $-u := 1 : p \setminus u$  be its complement, and  $\mathcal{P}(u)$  be its power set (the set of all subsets of u). For singleton  $\{i\}$ , abbreviate  $\mathbf{X}_{\{i\}}$  and  $\mathbf{X}_{-\{i\}}$  to  $X_i$  and  $\mathbf{X}_{-i}$  respectively.

#### 2.1 Factor Importance

The concept of quantifying the importance of an input through its predictive power is rooted in many *model-based* factor importance measures (Darlington, 1968; Breiman, 2001; Lundberg and Lee, 2017; Fisher et al., 2019). However, when the model is misspecified, the model-based measures cannot truly reflect the relevance between the inputs and the output, e.g., linear model often fails to capture the contribution from the nonlinear effect accurately. Hence, the factor importance should be measured by its population-level predictiveness potential, i.e., it should not be contingent on any specific prediction model. This idea is proposed as the *intrinsic* importance in Williamson et al. (2023).

Let  $V(f; f^*, \mathbb{P}_{\mathcal{X}})$  be a measure of the predictiveness for any function  $f \in \mathcal{F}$ , e.g., surrogate model, when the true model is  $f^* \in \mathcal{F}$  with input distribution  $\mathbb{P}_{\mathcal{X}}$ . The larger the  $V(f; f^*, \mathbb{P}_{\mathcal{X}})$ , the more predictive f is. Some popular predictiveness measures include  $R^2$  for regression, accuracy for classification, etc. It is obvious that the function in  $\mathcal{F}$  that maximizes  $V(f; f^*, \mathbb{P}_{\mathcal{X}})$  is the oracle model  $f^*$  itself, i.e.,  $f^* = \arg \max_{f \in \mathcal{F}} V(f; f^*, \mathbb{P}_{\mathcal{X}})$ . Similarly, define  $f^*_{-u}$  be the function in  $\mathcal{F}_{-u}$  that maximize  $V(f; f^*, \mathbb{P}_{\mathcal{X}})$ , which is the best possible predictive model without using variables  $\mathbf{X}_u$ . The population-level importance  $\psi_u$ of the factors  $\mathbf{X}_u$  relative to the all input  $\mathbf{X}$  is defined as

$$\psi_u := V(f^*; f^*, \mathbb{P}_{\mathcal{X}}) - V(f^*_{-u}; f^*, \mathbb{P}_{\mathcal{X}}), \tag{1}$$

which quantifies the oracle approximation error introduced by excluding variables  $\mathbf{X}_u$ .  $\psi_u$ can be viewed as the optimal lower bound for the approximation error to  $f^*(\mathbf{X})$  for any predictive model training only on  $\mathbf{X}_{-u}$ . By construction,  $\psi_u \ge 0$ . The larger value of  $\psi_u$ implies that  $\mathbf{X}_u$  is more important in terms of its predictive potential, and  $\psi_u = 0$  if and only if the true model  $f^*$  does not depend on  $\mathbf{X}_u$  at all. For the rest of this section we will introduce the total Sobol' indices, and then show it coincides with the intrinsic importance  $\psi_u$  (1) when  $R^2$  is the predictiveness measure.

### 2.2 Total Sobol' Indices

For any function  $f \in \mathcal{F}$  that is square integrable over the input distribution  $\mathbb{P}_{\mathcal{X}}$ , Sobol' (1993, 2001) show that it can be decomposed into

$$f(\mathbf{X}) = f_0 + \sum_{k=1}^p \sum_{|u|=k} f_u(\mathbf{X}_u),$$
(2)

where

$$f_0 = \mathbb{E}[f(\mathbf{X})],$$
  

$$f_i(X_i) = \mathbb{E}[f(\mathbf{X})|X_i] - f_0,$$
  

$$f_{i,j}(X_i, X_j) = \mathbb{E}[f(\mathbf{X})|X_i, X_j] - f_i(X_i) - f_j(X_j) - f_0,$$
  

$$\vdots$$
  

$$f_u(\mathbf{X}_u) = \mathbb{E}[f(\mathbf{X})|\mathbf{X}_u] - \sum_{v \in \mathcal{P}(u) \setminus u} f_v(\mathbf{X}_v).$$

When the input variables are *independent*, the decomposition is unique, and  $f_u(\mathbf{X}_u)$ 's have zero mean and are mutually orthogonal, yielding similar decomposition for output variance,

$$\operatorname{Var}[f(\mathbf{X})] = \sum_{k=1}^{p} \sum_{|u|=k} \operatorname{Var}[f_u(\mathbf{X}_u)],$$

which is also known as the ANOVA (analysis of variance) decomposition (Efron and Stein, 1981). The Sobol' index (Sobol', 1993) of  $\mathbf{X}_u$  under independent inputs is defined as

$$S_u := \frac{\operatorname{Var}[f_u(\mathbf{X}_u)]}{\operatorname{Var}[f(\mathbf{X})]}.$$
(3)

It is easy to see that  $0 \leq S_u \leq 1$  and  $\sum_{k=1}^p \sum_{|u|=k} S_u = 1$ . The *first-order* Sobol' indices (Homma and Saltelli, 1996) is proposed for capturing the output variance that is associated with the main effect, i.e., the first-order Sobol' index of  $X_i$  is defined as

$$S_i := \frac{\operatorname{Var}[f_i(X_i)]}{\operatorname{Var}[f(\mathbf{X})]} = \frac{\operatorname{Var}[\mathbb{E}\{f(\mathbf{X})|X_i\}]}{\operatorname{Var}[f(\mathbf{X})]} = \frac{\operatorname{Var}[f(\mathbf{X})] - \mathbb{E}[\operatorname{Var}\{f(\mathbf{X})|X_i\}]}{\operatorname{Var}[f(\mathbf{X})]}, \quad (4)$$

where the last equality follows from the variance decomposition formula.  $S_i$  measures the expected proportional reduction in  $\operatorname{Var}[f(\mathbf{X})]$  if  $X_i$  is fixed to some constant, so intuitively the higher the  $S_i$ , the more important  $X_i$  is. However,  $S_i$  fails to account for any effect from the interaction between  $X_i$  and other inputs. To overcome this limitation, Homma and Saltelli (1996) proposed the *total* Sobol' index  $S_i^{\text{tot}}$  that not only capture the main effect but also all the interaction effects involving  $X_i$ ,

$$S_i^{\text{tot}} := \sum_{v \in \mathcal{P}(1:p):\{i\} \cap v \neq \emptyset} S_v = 1 - \sum_{v \in \mathcal{P}(1:p \setminus \{i\})} S_v, \tag{5}$$

and thus  $S_i \leq S_i^{\text{tot}}$  and the difference reflects the output variance attributed by the interaction effects. However, when the inputs are *correlated*, the  $f_u(\mathbf{X}_u)$ 's in the decomposition (2) are no longer orthogonal. Li et al. (2010) propose the following generalization,

$$\tilde{S}_u := \frac{\operatorname{Cov}(f_u(\mathbf{X}_u), f(\mathbf{X}))}{\operatorname{Var}[f(\mathbf{X})]},\tag{6}$$

which reduces to (3) for independent inputs. The first-order Sobol' indices (4) and the total Sobol' indices (5) can be defined in the same way using  $\tilde{S}_u$ , and in what follows we focus on this generalized definition of the Sobol' indices. Li et al. (2010) shows that  $\sum_{k=1}^{p} \sum_{|u|=k} \tilde{S}_u = 1$ , but  $\tilde{S}_u$  could be negative depending on  $\text{Cov}(f_u(\mathbf{X}_u), f(\mathbf{X}))$ , and hence it is possible to see  $S_i > S_i^{\text{tot}}$  in the presence of dependent inputs. This negative contribution from the interaction effect creates ambiguity when trying to interpret Sobol' indices as a measure for factor importance (Chastaing et al., 2015; Song et al., 2016; Owen and Prieur, 2017; Hart and Gremaud, 2018). On the other hand, Kucherenko et al. (2012) show that

equivalently the total Sobol' index of  $X_i$  is

$$S_i^{\text{tot}} = \frac{\text{Var}[f(\mathbf{X})] - \text{Var}[\mathbb{E}\{f(\mathbf{X})|\mathbf{X}_{-i}\}]}{\text{Var}[f(\mathbf{X})]} = \frac{\mathbb{E}[\text{Var}\{f(\mathbf{X})|\mathbf{X}_{-i}\}]}{\text{Var}[f(\mathbf{X})]} \ge 0,$$

showing that  $S_i^{\text{tot}}$  captures the expected proportional residual variance if all other factors  $\mathbf{X}_{-i}$  are fixed. Hence, the smaller the  $S_i^{\text{tot}}$ , the less important  $X_i$  is, and  $S_i^{\text{tot}} = 0$  if and only if the model f does not include  $X_i$ . Moreover, the above definition of the total Sobol' indices generalizes to any subset  $u \subset 1:p$ , i.e.,

$$S_u^{\text{tot}} = \frac{\mathbb{E}[\operatorname{Var}\{f(\mathbf{X}) | \mathbf{X}_{-u}\}]}{\operatorname{Var}[f(\mathbf{X})]}.$$
(7)

To address the interpretation issue of the Sobol' indices in the presence of dependent inputs, there is also a growing interest on investigating other global sensitivity analysis methods, including the density-based measure (Borgonovo, 2007; Borgonovo et al., 2011; Plischke et al., 2013), derivative-based measure (Sobol' and Kucherenko, 2009), Shapley effect (Owen, 2014; Song et al., 2016; Broto et al., 2020), etc. For a comprehensive review, please see Borgonovo and Plischke (2016), Razavi et al. (2021), and Da Veiga et al. (2021).

#### 2.3 Equivalence under Regression

The  $\mathbb{R}^2$  predictiveness measure is the conventional choice for regression. It is defined as

$$V(f; f^*, \mathcal{P}_{\mathcal{X}}) = 1 - \frac{\mathbb{E}[\{f(\mathbf{X}) - f^*(\mathbf{X})\}^2]}{\operatorname{Var}[f^*(\mathbf{X})]}$$

which measures the proportion of variability in  $f^*(\mathbf{X})$  that can be explained by  $f(\mathbf{X})$  under the input distribution  $\mathbb{P}_{\mathcal{X}}$ .

**Theorem 1.** When  $R^2$  is the predictiveness measure,  $\psi_u = S_u^{tot}$  for  $u \subset 1: p$ .

*Proof.* When  $R^2$  is the predictiveness measure, it is obvious that  $V(f^*; f^*, \mathcal{P}_{\mathcal{X}}) = 1$  and

$$f_{-u}^{*}(\mathbf{X}_{-u}) = \arg \max_{f \in \mathcal{F}_{-u}} V(f; f^{*}, \mathcal{P}_{\mathcal{X}})$$
  
$$= \arg \max_{f \in \mathcal{F}_{-u}} \left( 1 - \frac{\mathbb{E}[\{f(\mathbf{X}_{-u}) - f^{*}(\mathbf{X})\}^{2}]}{\operatorname{Var}[f^{*}(\mathbf{X})]} \right)$$
  
$$= \arg \min_{f \in \mathcal{F}_{-u}} \mathbb{E}[\{f(\mathbf{X}_{-u}) - f^{*}(\mathbf{X})\}^{2}]$$
  
$$= \mathbb{E}[f^{*}(\mathbf{X})|\mathbf{X}_{-u}]$$

is the projection of  $f^*(\mathbf{X})$  onto  $\mathcal{F}_{-u}$ . It follows that the factor importance of  $\mathbf{X}_u$  is

$$\begin{split} \psi_u &= V(f^*; f^*, \mathbb{P}_{\mathcal{X}}) - V(f^*_{-u}; f^*, \mathbb{P}_{\mathcal{X}}) \\ &= \frac{\mathbb{E}[\{\mathbb{E}[f^*(\mathbf{X}) | \mathbf{X}_{-u}] - f^*(\mathbf{X})\}^2]}{\operatorname{Var}[f^*(\mathbf{X})]} \\ &= \frac{\mathbb{E}[\operatorname{Var}\{f^*(\mathbf{X}) | \mathbf{X}_{-u}\}]}{\operatorname{Var}[f^*(\mathbf{X})]} \\ &= S^{\operatorname{tot}}_u, \end{split}$$

which coincides with the total Sobol' index of  $\mathbf{X}_u$ . The third equality follows from the fact that  $\mathbb{E}[(\mathbb{E}[Y|X] - Y)^2] = \mathbb{E}[\operatorname{Var}(Y|X)]$  (Lemma 1 in Appendix A).

Hart and Gremaud (2018) provide an alternative proof for this approximation theoretic perspective of the total Sobol' indices. With this equivalence, computing the factor importance  $\psi_u$  reduces to the computation of the total Sobol' index  $S_u^{\text{tot}}$ . Hence, utilizing the procedure presented in the next section for estimating the total Sobol' indices directly from data, we could circumvent the model fitting step for approximating  $f^*$  and  $f_{-u}^*$  that are required by the existing estimator of  $\psi_u$  proposed by Williamson et al. (2023). This also avoid the risk of systematic bias from model misspecification.

## 3 Estimating Total Sobol' Index from Noisy Data

In the computer experiments literature, various Monte Carlo methods have been proposed for estimating the total Sobol' indices (7) efficiently, including Pick-and-Freeze estimator

(Homma and Saltelli, 1996), Jansen-Sobol estimator (Jansen, 1999; Sobol', 2001; Saltelli et al., 2010; Kucherenko et al., 2012), Double Monte Carlo estimator (Song et al., 2016), etc. These methods all require knowing about (i) the model  $f^*$ , or at the very least, the ability to evaluate it for any given input, and (ii) the input distribution and its conditional in closed-form. However, neither is available for our problem, where the goal is to assess the importance of the variables solely based on the data  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ . One possible way to overcome these obstacles is to fit (i) a surrogate model  $\hat{f}$  on  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$  to emulate the unknown true model  $f^*$  and (ii) an empirical distribution estimator on  $\{\mathbf{x}^{(n)}\}_{n=1}^N$  with the capability of simulating samples from  $X_i | \mathbf{X}_{-i}$ . As alluded to in the previous sections, we try to avoid any model fitting because (i) it can be an arduous task by itself, especially with big data, and (ii) it can lead to estimation bias if the model is misspecified. On the other hand, Monte Carlo methods require simulating synthetic inputs, but as criticized in Mase et al. (2022), the trustworthiness of the importance measures derived from these synthetic inputs is questionable because they can be "unlikely, physically impossible, or even logically impossible". Hence, in this section we propose an efficient algorithm for computing the total Sobol' indices (7) using only the collected noisy data  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .

#### 3.1 Design-based Double Monte Carlo Estimator

Let us start by reviewing the Double Monte Carlo estimator proposed in Song et al. (2016) that lays the crucial groundwork for the total Sobol' indices estimator using only data. Recall that the total Sobol' index  $S_i^{\text{tot}}$  of  $X_i$  with respect to function  $f^*$  is defined as

$$S_i^{\text{tot}} = \frac{\mathbb{E}[\operatorname{Var}\{f^*(\mathbf{X}) | \mathbf{X}_{-i}\}]}{\operatorname{Var}[f^*(\mathbf{X})]}.$$

We focus on the estimation for a single index, i.e.,  $u = \{i\}$  for some  $i \in 1: p$ , because it is often the interest to assess the importance of individual factor rather than a group of them, but the estimation procedure generalizes for any subset  $u \subset 1: p$ .

The denominator  $\operatorname{Var}[f^*(\mathbf{X})]$  can be estimated by simple Monte Carlo using samples

from  $\mathbb{P}_{\mathcal{X}}$ . The main difficulty comes from estimating the numerator,

$$T_i = \mathbb{E}[\operatorname{Var}\{f^*(\mathbf{X})|\mathbf{X}_{-i}\}].$$
(8)

For simplicity we refer  $T_i$  as the total Sobol' effect of  $X_i$ , the un-normalized version of the total Sobol' index. Song et al. (2016) proposed to use a double loop Monte Carlo: the inner loop for the conditional variance using  $N_I$  designs and the outer loop for the expectation using  $N_O$  designs, i.e.,

$$\hat{T}_{i}^{\rm mc} = \frac{1}{N_O} \sum_{m=1}^{N_O} \left\{ \frac{1}{N_I - 1} \sum_{j=1}^{N_I} \left( f^*(\mathbf{x}_{-i}^{(m)}, x_i^{(m,j)}) - \frac{1}{N_I} \sum_{l=1}^{N_I} f^*(\mathbf{x}_{-i}^{(m)}, x_i^{(m,l)}) \right)^2 \right\}, \qquad (9)$$

where  $\{\mathbf{x}_{-i}^{(m)}\}_{m=1}^{N_O}$  are i.i.d. samples from the distribution of  $\mathbf{X}_{-i}$  and  $\{x_i^{(m,j)}\}_{j=1}^{N_I}$  conditionally to  $\mathbf{x}_{-i}^{(m)}$  is i.i.d. with respect to the distribution of  $X_i | \mathbf{X}_{-i} = \mathbf{x}_{-i}^{(m)}$ . The Double Monte Carlo estimator  $\hat{T}_i^{\text{mc}}$  (9) is unbiased for  $T_i$  (8), and it is suggested that  $N_I = 3$  is sufficient (Song et al., 2016). When  $N_I = 2$ , the Double Monte Carlo estimator (9) reduces to the generalized Jansen-Sobol estimator in Kucherenko et al. (2012),

$$\hat{T}_{i}^{js} = \frac{1}{N_{O}} \sum_{m=1}^{N_{O}} \frac{1}{2} \left( f^{*}(\mathbf{x}_{-i}^{(m)}, x_{i}^{(m,1)}) - f^{*}(\mathbf{x}_{-i}^{(m)}, x_{i}^{(m,2)}) \right)^{2}.$$

#### **3.2** Estimation from Clean Data

Before presenting the algorithm for estimating the total Sobol' indices from the noisy data, we first discuss the estimation using clean i.i.d. samples  $\{(\mathbf{x}^{(n)}, y^{(n)} = f^*(\mathbf{x}^{(n)}))\}_{n=1}^N$ , i.e., there is no random noise associated with the output. It is worth emphasizing that here  $f^*$  is neither known analytically nor assessable for evaluation, so the Double Monte Carlo estimator (9) is not applicable. To overcome this limitation, a consistent estimator based on the nearest-neighbor method is proposed in Broto et al. (2020). The key is to replace the inner loop designs  $\{(\mathbf{x}_{-i}^{(m)}, x_i^{(m,j)})\}_{j=1}^{N_I}$  by the  $N_I$  samples in  $\{\mathbf{x}^{(n)}\}_{n=1}^N$  with the values of their -i entries closest to  $\mathbf{x}_{-i}^{(m)}$ . More formally, let  $k_{-i}^{(m)}(j)$  be the index such that  $\mathbf{x}_{-i}^{(k_{-i}^{(m)}(j))}$ is the *j*-th closest element to  $\mathbf{x}_{-i}^{(m)}$  in  $\{\mathbf{x}_{-i}^{(n)}\}_{n=1}^N$ , with ties broken by random selection. The nearest-neighbor version of the Double Monte Carlo estimator (9) for  $T_i$  is

$$\hat{T}_{i}^{\mathrm{nn}} = \frac{1}{N_{O}} \sum_{m=1}^{N_{O}} \left\{ \frac{1}{N_{I} - 1} \sum_{j=1}^{N_{I}} \left( y^{(k_{-i}^{(m)}(j))} - \frac{1}{N_{I}} \sum_{l=1}^{N_{I}} y^{(k_{-i}^{(m)}(l))} \right)^{2} \right\},\tag{10}$$

where  $\{\mathbf{x}_{-i}^{(m)}\}_{m=1}^{N_O}$  for the outer loop Monte Carlo can either be all the samples  $\{\mathbf{x}_{-i}^{(n)}\}_{n=1}^N$ or a random subsample (with or without replacement) of  $\{\mathbf{x}_{-i}^{(n)}\}_{n=1}^N$  when N is large. For convenience, in what follows we refer to (10) as the nearest-neighbor estimator. Under some mild assumptions on the smoothness of the input distribution  $\mathbb{P}_{\mathcal{X}}$ , Broto et al. (2020) proved that  $\hat{T}_i^{nn}$  converges to  $T_i$  in probability as  $N, N_O \to \infty$  for any model function f that is bounded. The key is to recognize that as  $N \to \infty$ , any nearest neighbor  $\mathbf{x}_{-i}^{(k_{-i}^{(m)}(j))}$ converges almost surely to  $\mathbf{x}_{-i}^{(m)}$ . It follows that  $X_i | \mathbf{X}_{-i} = \mathbf{x}_{-i}^{(k_{-i}^{(m)}(j))}$  converges weakly to  $X_i | \mathbf{X}_{-i} = \mathbf{x}_{-i}^{(m)}$ , and hence  $\{\mathbf{x}^{(k_{-i}^{(m)}(j))}\}_{j=1}^{N_I}$  can be viewed as i.i.d. samples simulated from  $X|\mathbf{X}_{-i} = \mathbf{x}_{-i}^{(m)}$ . Moreover, Broto et al. (2020) further show that the rate of convergence for  $\hat{T}_i^{nn}$  (10) is almost  $o_p(N^{-1/2})$  by imposing a stronger assumption that (i) the first derivative of model function f exist and is continuous and (ii) the input variables  $\mathbf{X}$  are continuous, real-valued, restricted to a compact set where the density is lower-bounded (e.g., uniform or truncated Gaussian). The consistency and the rate of convergence is independent of (i) the problem dimension p and (ii) the number of samples  $N_I$  for estimating the conditional variance. For the denominator  $\operatorname{Var}[f^*(\mathbf{X})]$  of the total Sobol' indices, one can use the consistent estimator,  $\frac{1}{N-1}\sum_{n=1}^{N}(y^{(n)}-\frac{1}{N}\sum_{l=1}^{N}y^{(l)})^2$ .

### **3.3** Extension to Noisy Data

Suppose now that the output is contaminated by random measurement error, i.e.,

$$Y = \tilde{f}(\mathbf{X}, \epsilon) = f^*(\mathbf{X}) + \epsilon, \tag{11}$$

where  $\epsilon$  is the random noise with zero mean and finite second moment and is *independent* of the input **X**. We will develop a consistent estimator for both  $T_i = \mathbb{E}[\operatorname{Var}\{f^*(\mathbf{X})|\mathbf{X}_{-i}\}],$ the total Sobol' effect of  $X_i$  with respect to the true model  $f^*$ , and  $\operatorname{Var}[f^*(\mathbf{X})]$  given the i.i.d. noisy samples  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$  of (11), where  $y^{(n)} = f^*(\mathbf{x}^{(n)}) + \epsilon^{(n)}$ .

Consider under the framework of analysis of computer experiment that the random noise  $\epsilon$  in (11) can be treated as another stochastic variable, similar to **X**, supplied as an input to the deterministic function  $\tilde{f}$ . The variance of  $\epsilon$  is equivalent to  $\tilde{T}_{\epsilon}$ , the total Sobol' effect of  $\epsilon$  with respect to the noisy model  $\tilde{f}$ , i.e.,

$$\tilde{T}_{\epsilon} = \mathbb{E}[\operatorname{Var}\{\tilde{f}(\mathbf{X},\epsilon)|\mathbf{X}\}] = \mathbb{E}[\operatorname{Var}\{f^*(\mathbf{X})+\epsilon|\mathbf{X}\}] = \mathbb{E}[\operatorname{Var}(\epsilon)] = \operatorname{Var}[\epsilon].$$

The nearest-neighbor estimator (10) for  $\tilde{T}_{\epsilon}$  using noisy data  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^{N}$  is

$$\hat{\tilde{T}}_{\epsilon}^{nn} = \frac{1}{N_O} \sum_{m=1}^{N_O} \left\{ \frac{1}{N_I - 1} \sum_{j=1}^{N_I} \left( y^{(k_{1:p}^{(m)}(j))} - \frac{1}{N_I} \sum_{l=1}^{N_I} y^{(k_{1:p}^{(m)}(l))} \right)^2 \right\}.$$
(12)

From the discussion in Subsection 3.2,  $\hat{T}_{\epsilon}^{nn}$  is a consistent estimator for  $\tilde{T}_{\epsilon}$ , and hence also a consistent estimator for  $\operatorname{Var}[\epsilon]$ . Here  $k_{1:p}^{(m)}(j)$  is the index for the *j*-th nearest-neighbor to  $\mathbf{x}^{(m)}$  in  $\{\mathbf{x}^{(n)}\}_{n=1}^{N}$ , where  $\{\mathbf{x}^{(m)}\}_{m=1}^{N_0}$  for the outer loop Monte Carlo can either be all the samples  $\{\mathbf{x}^{(n)}\}_{n=1}^{N}$  or a random subsample. Similarly, for  $\tilde{T}_{i,\epsilon}$ , the total Sobol' effect of  $X_i$ and  $\epsilon$  jointly with respect to the noisy model  $\tilde{f}$ , we have

$$\tilde{T}_{i,\epsilon} = \mathbb{E}[\operatorname{Var}\{f^*(\mathbf{X}) + \epsilon | \mathbf{X}_{-i}\}] = \mathbb{E}[\operatorname{Var}\{f^*(\mathbf{X}) | \mathbf{X}_{-i}\} + \operatorname{Var}(\epsilon)] = T_i + \operatorname{Var}[\epsilon],$$

because  $\mathbf{X}_{-i}$  and  $\epsilon$  are *independent*. The corresponding nearest-neighbor estimator is

$$\hat{\tilde{T}}_{i,\epsilon}^{nn} = \frac{1}{N_O} \sum_{m=1}^{N_O} \left\{ \frac{1}{N_I - 1} \sum_{j=1}^{N_I} \left( y^{(k_{-i}^{(m)}(j))} - \frac{1}{N_I} \sum_{l=1}^{N_I} y^{(k_{-i}^{(m)}(l))} \right)^2 \right\}.$$
(13)

By the Slutzky's theorem,  $\hat{T}_i = \hat{T}_{i,\epsilon}^{nn} - \hat{T}_{\epsilon}^{nn}$  is a consistent estimator for  $T_i$ . Following the same line of thought,  $\operatorname{Var}[Y] = \operatorname{Var}[\tilde{f}(\mathbf{X},\epsilon)] = \operatorname{Var}[f^*(\mathbf{X})] + \operatorname{Var}[\epsilon]$ , and a consistent



Figure 1: Comparison of NANNE estimator with  $N_I = 2$  (NANNE-2) and  $N_I = 3$  (NANNE-3) to the nearest-neighbor estimator (10) with  $N_I = 3$  (NN-3) on the noisy data simulated from the Ishigami function. The analytical total Sobol' indices are marked by the red dashed lines, and the boxplots summarize the estimations from 100 independent runs.

estimator for  $\operatorname{Var}[Y]$  is

$$\widehat{\operatorname{Var}[Y]} = \frac{1}{N-1} \sum_{n=1}^{N} \left( y^{(n)} - \frac{1}{N} \sum_{l=1}^{N} y^{(l)} \right)^2.$$
(14)

Again by the Slutzky's theorem,  $\operatorname{Var}[f^*(\mathbf{X})] = \widehat{\operatorname{Var}[Y]} - \widehat{T}_{\epsilon}^{\operatorname{nn}}$  is a consistent estimator for  $\operatorname{Var}[f^*(\mathbf{X})]$ . Numerically this estimator is not always positive given the noisy data, we clip its value to 0 in the actual implementation. Moreover,  $\operatorname{Var}[f^*(\mathbf{X})] = 0$  indicates that the output variance is exclusively from the random noise, and hence  $\widehat{S}_i^{\operatorname{tot}} = 0$  for all  $i \in 1:p$ .

When  $\operatorname{Var}(\widehat{f^*(\mathbf{X})}) > 0$ , a consistent estimator for the total Sobol' index  $S_i^{\text{tot}}$  of  $X_i$  is

$$\hat{S}_i^{\text{tot}} = \frac{\hat{T}_i}{\widehat{\text{Var}[f^*(\mathbf{X})]}} = \frac{\hat{\tilde{T}}_{i,\epsilon}^{\text{nn}} - \hat{\tilde{T}}_{\epsilon}^{\text{nn}}}{\widehat{\text{Var}[Y]} - \hat{\tilde{T}}_{\epsilon}^{\text{nn}}}$$

Numerically the numerator can again be negative, and hence we also clip its value to 0. Let us refer to this as the Noise-Adjusted Nearest-NEighbor (NANNE) estimator, and the steps involved in its computation are outlined in Algorithm 1. K-d tree (Bentley, 1975) can be employed to accelerate the nearest neighbor search in practice.

Figure 1 compares NANNE and the nearest-neighbor (10) estimators on the noisy data

#### Algorithm 1 Noise-Adjusted Nearest-NEighbor (NANNE) estimator.

- Input: (i) noisy samples {(**x**<sup>(n)</sup> ∈ ℝ<sup>p</sup>, y<sup>(n)</sup> ∈ ℝ)}<sup>N</sup><sub>n=1</sub>, (ii) number of samples for inner loop Monte Carlo N<sub>I</sub>, and (iii) number of samples for outer loop Monte Carlo N<sub>O</sub>.
   Compute Var[Y] via (14).
   Compute Var[f<sup>\*</sup>(**X**)] = max (Var[Y] Î<sup>\*</sup><sub>ϵ</sub>, 0).
   if Var[f<sup>\*</sup>(**X**)] > 0 then
   for i = 1,..., p do
   Compute Î<sup>\*</sup><sub>i,ϵ</sub> via (13).
- 8: Compute  $\hat{T}_i = \max\left(\hat{T}_{i,\epsilon}^{nn} \hat{T}_{\epsilon}^{nn}, 0\right)$ . 9: Compute  $\hat{S}_i^{\text{tot}} = \hat{T}_i / \operatorname{Var}[f^*(\mathbf{X})]$ . 10: end for
- 11: else 12: Let  $\hat{S}_{i}^{\text{tot}} = 0$  for  $i = 1, \dots, p$ . 13: end if
- 14: **Output:** the total Sobol' indices  $\{\hat{S}_i^{\text{tot}}\}_{i=1}^p$ .

 $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$  simulated from the Ishigami function (Ishigami and Homma, 1990),

$$f^*(\mathbf{X}) = \sin(X_1) + 7\sin^2(X_2) + 0.1X_3^4\sin(X_1), \quad X \sim \text{Uniform}[-\pi,\pi]^3.$$

The random noise  $\epsilon^{(n)} \sim \mathcal{N}(0, 1)$ . We run the simulation with N = 10,000 samples and use all of them for the outer loop Monte Carlo designs  $\{\mathbf{x}^{(m)}\}_{m=1}^{N_O}$ , i.e.,  $N_O = N = 10,000$ . From Figure 1 it is evident that in the presence of noise, the nearest-neighbor estimator consistently overestimates the total Sobol' indices, whereas our proposed NANNE estimator is able to address this overestimation issue and provide an accurate estimate. For the nearest-neighbor estimator, Broto et al. (2020) suggested using  $N_I = 3$  for the conditional variance estimation. However, from the simulation result in Figure 1 we find that for the NANNE estimator,  $N_I = 2$  yields a more robust estimation when the data is noisy. Hence, in practice for the regression problem we suggest using  $N_I = 2$  for the NANNE estimator.

### 4 Rethinking Factor Importance

Recall from Subsection 2.1, the factor importance  $\psi_u$  of  $\mathbf{X}_u$  is defined relative to all the input  $\mathbf{X}$ . The natural question that follows is *what should be this full input*  $\mathbf{X}$ ? For analysis of computer experiments where the model  $f^*$  is known, the answer is straightforward, i.e.  $\mathbf{X}$  should be all the inputs required by  $f^*$ . However, for the usual statistical modeling problem where we only have noisy data  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ , the answer becomes ambiguous: should  $\mathbf{X}$  be (i) all the variables that are collected or (ii) only the variables that the *unknown* underlying model  $f^*$  depends on? To answer this question, let us first consider the following linear Gaussian example,

$$f^*(\mathbf{X}) = X_1 + X_2 + 0X_3,$$

where  $\mathbf{X} \sim \mathcal{N}(0, \Sigma)$  with  $\operatorname{Var}(X_i) = 1$  for all *i*,  $\operatorname{Cov}(X_1, X_2) = \operatorname{Cov}(X_1, X_3) = 0$ , and  $\operatorname{Cov}(X_2, X_3) = 0.9$ , i.e.,  $X_2$  and  $X_3$  are strongly correlated while both are independent of  $X_1$ . Analytically solving for the factor importance relative to all input  $\mathbf{X} = [X_1, X_2, X_3]$ , we have  $\psi_1^{(1)} = 0.500$  and  $\psi_2^{(1)} = 0.095$ . On the other hand, if the factor importance is computed relative to only the factors  $[X_1, X_2]$  that are influential in the model, we have  $\psi_1^{(2)} = 0.500$  and  $\psi_2^{(2)} = 0.500$ . Clearly, the latter importance measure is evidently more in line with our expectation. The reason that  $\psi_2^{(1)}$  is significantly smaller than  $\psi_2^{(2)}$  is that  $X_3$  serves as a good proxy for  $X_2$  due to their strong correlation, though  $X_3$  itself does not play a role in the true model. Nevertheless, if  $X_2$  is collected as one of the predictors, we want neither  $X_3$  show up as important nor the importance of  $X_2$  depend on  $X_3$ . Hence, given data  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ , factor selection precedes factor importance, and the importance should be measured relative to only factors that are relevant to the true model. By factor selection, we imply filtering out variables that are not part of the true model, including those that could be correlated with the output but only through their dependency with the true model variables, e.g.,  $X_3$  in the linear Gaussian example.

From the equivalence of the total Sobol' indices and the intrinsic factor importance, a factor is not important if its total Sobol' index is zero, and it should be removed from the list

#### Algorithm 2 NANNE-BE: NANNE after Backward Elimination.

 Input: (i) noisy samples {(x<sup>(n)</sup> ∈ ℝ<sup>p</sup>, y<sup>(n)</sup> ∈ ℝ)}<sup>N</sup><sub>n=1</sub>, (ii) number of samples for inner loop Monte Carlo N<sub>I</sub>, and (iii) number of samples for outer loop Monte Carlo N<sub>O</sub>.
 Initialization: A ← 1:p, Ŝ<sup>tot</sup><sub>i</sub> ← 1/p for i ∈ A;
 do
 A ← {i ∈ A : Ŝ<sup>tot</sup><sub>i</sub> > 0}; {Ŝ<sup>tot</sup><sub>i∈A</sub> ← NANNE({(x<sup>(n)</sup><sub>A</sub>, y<sup>(n)</sup>)}<sup>N</sup><sub>n=1</sub>, N<sub>I</sub>, N<sub>O</sub>);
 while |A| > 0 and min<sub>i∈A</sub> Ŝ<sup>tot</sup><sub>i</sub> = 0;
 Ŝ<sup>tot</sup><sub>i</sub> ← 0 for i ∈ 1:p\A.
 Output: the total Sobol' indices {Ŝ<sup>tot</sup><sub>i</sub>}<sup>p</sup><sub>i=1</sub> after factor selection.

of inputs for the factor importance computation. Hence, a natural extension of the NANNE estimator to accommodate factor selection and importance simultaneously is by backward elimination. The key idea is to start with all the available factors, next compute the total Sobol' indices via NANNE and filter out factors with the total Sobol' index value of zero, and then re-estimate the total Sobol' indices using only the remaining variables. The above procedure is repeated until the total Sobol' indices are all positive for the surviving factors. We refer this as NANNE-BE, NANNE with Backward Elimination that provides the factor importance estimation after discarding the irrelevant variables. Please see Algorithm 2 for the detailed procedure.

However, since NANNE estimator is developed upon the nearest-neighbor estimator, it also suffers from the *curse of dimensionality*: all the points are nearly in equal distance from one another when the dimension *p* is large. This is especially problematic when many of the factors are irrelevant for the target prediction, causing very noisy estimation and hence undermining the robustness of NANNE and NANNE-BE. To address the curse of dimensionality issue, we propose to first choose a promising set of candidates by greedy forward selection to reduce the dimensionality, and then compute the importance exclusively on the candidate factors. For the factor that is not in the candidate set, zero importance is assigned. Similar to the standard forward selection procedures (Efroymson, 1960), we start from the empty set and add one factor at a time. The selection criterion we consider is to maximize the oracle predictive power, i.e., the maximal possible variance that can be explained from the selected factors. In the context of Sobol' effect, this is the sum of the main effects and all the possible interaction effects of the selected factors. Let u denote the set of indices for the selected factors, the variance of  $f^*(\mathbf{X})$  that can be explained is

$$V_u = \sum_{v \in \mathcal{P}(u)} \operatorname{Cov}(f_v^*(X_v), f^*(\mathbf{X})) = \operatorname{Var}[\mathbb{E}\{f^*(\mathbf{X}) | \mathbf{X}_u\}].$$

Moreover, let  $\tilde{V}_u$  be the corresponding effect of  $\mathbf{X}_u$  with respect to the noisy model (11)  $Y = \tilde{f}(\mathbf{X}, \epsilon) = f^*(\mathbf{X}) + \epsilon$ . Since  $\epsilon$  is independent of  $\mathbf{X}$ , we have

$$\tilde{V}_u = \operatorname{Var}[\mathbb{E}\{\tilde{f}(\mathbf{X},\epsilon)|\mathbf{X}_u\}] = \operatorname{Var}[\mathbb{E}\{f^*(\mathbf{X})+\epsilon|\mathbf{X}_u\}] = \operatorname{Var}[\mathbb{E}\{f^*(\mathbf{X})|\mathbf{X}_u\}] = V_u.$$

By the variance decomposition formula,

$$V_u = \tilde{V}_u = \operatorname{Var}[Y] - \mathbb{E}[\operatorname{Var}\{\tilde{f}(\mathbf{X})|\mathbf{X}_u\}] = \operatorname{Var}[Y] - \tilde{T}_{-u},$$

where  $\tilde{T}_{-u}$  is the total Sobol' effect (8) of  $\mathbf{X}_{-u}$  with respect to the noisy model  $\tilde{f}$ . Given noisy data  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ , the nearest-neighbor estimator of  $\tilde{T}_{-u}$  is

$$\hat{\tilde{T}}_{-u} = \frac{1}{N_O} \sum_{m=1}^{N_O} \left\{ \frac{1}{N_I - 1} \sum_{j=1}^{N_I} \left( y^{(k_u^{(m)}(j))} - \frac{1}{N_I} \sum_{l=1}^{N_I} y^{(k_u^{(m)}(l))} \right)^2 \right\},\$$

where  $k_u^{(m)}(j)$  is the index for the *j*-th nearest-neighbor to  $\mathbf{x}_u^{(m)}$  in  $\{\mathbf{x}^{(n)}\}_{n=1}^N$ , with  $\{\mathbf{x}_u^{(m)}\}_{m=1}^{N_O}$ be either all the samples or a random subsample of  $\{\mathbf{x}_u^{(n)}\}_{n=1}^N$ . It follows that a consistent estimator for  $V_u$  is

$$\hat{V}_u = \widehat{\operatorname{Var}[Y]} - \hat{\tilde{T}}_{-u},\tag{15}$$

where  $\widehat{\operatorname{Var}[Y]}$  is a consistent estimator of  $\operatorname{Var}[Y]$  from (14). The forward selection terminates when no variable can further improve the variance that can be explained. Given that the greedy forward selection could sometimes include a few irrelevant factors, we apply NANNE-BE (backward elimination) to filter them before computing the importance measure  $\{\hat{\psi}_i\}_{i=1}^p$ . The detail procedure is presented in Algorithm 3, and we name it FIRST which

#### Algorithm 3 FIRST: forward selection followed by NANNE-BE.

- 1: Input: (i) noisy samples  $\{(\mathbf{x}^{(n)} \in \mathbb{R}^p, y^{(n)} \in \mathbb{R})\}_{n=1}^N$ , (ii) number of samples for inner loop Monte Carlo  $N_I$ , and (iii) number of samples for outer loop Monte Carlo  $N_O$ . 2: Initialization:  $k = -1, A \leftarrow \emptyset, \{i_0\} = \emptyset, \hat{V}_{\emptyset} = 0;$
- 3: do
- 4:  $k \leftarrow k+1;$
- $A \leftarrow A \cup \{i_k\};$ 5:

Choose  $i_{k+1} \in 1: p \setminus A$  such that  $\hat{V}_{A \cup \{i_{k+1}\}}$  (15) is maximized, i.e., 6:

$$i_{k+1} = \arg \max_{i \in 1: p \setminus A} \hat{V}_{A \cup \{i\}};$$

7: while |A| < p and  $\hat{V}_{A \cup \{i_{k+1}\}} > \hat{V}_A$ ; 8:  $\{\hat{S}_i^{\text{tot}}\}_{i \in A} \leftarrow \text{NANNE-BE}(\{(\mathbf{x}_A^{(n)}, y^{(n)})\}_{n=1}^N, N_I, N_O);$ 9:  $\hat{\psi}_i \leftarrow \hat{S}_i^{\text{tot}}$  for  $i \in A$  and  $\hat{\psi}_i \leftarrow 0$  for  $i \in 1: p \setminus A$ ; 10: **Output:** the factor importance  $\{\hat{\psi}_i\}_{i=1}^p$ .

stands for Factor Importance Ranking and Selection using Total indices. Compared to the existing stepwise procedures, FIRST is independent of learning algorithms, i.e., all the estimation are done directly from data, and hence not only avoiding the computational burden of model fitting but more importantly the risk of model misspecification.

#### 5 Simulations on Regression Problems

We now compare the performance of our proposed  $FIRST^1$  algorithm (Algorithm 3) to a comprehensive list of factor importance/selection procedures in various regression settings. The following models are considered:

• Rescaled version of the Ishigami function (Ishigami and Homma, 1990):

$$f^*(\mathbf{X}) = \sin(2\pi X_1 - \pi) + 7\sin^2(2\pi X_2 - \pi) + 0.1(2\pi X_3 - \pi)^4 \sin(2\pi X_1 - \pi).$$
(16)

• Modified version of the heavy-tailed nonlinear function (Huang et al., 2022):

$$f^*(\mathbf{X}) = \frac{2\log(X_1^2 + X_2^4)}{\cos(X_1) + \sin(X_3)} + \frac{X_2^2 \exp(X_3)}{\sqrt{1.1 - X_6}}.$$
(17)

<sup>&</sup>lt;sup>1</sup>The implementation of **FIRST** is available in the R package **first**.

• Modified version of the Friedman function (Friedman, 1991):

$$f^*(\mathbf{X}) = 10\sin(\pi X_1 X_7) + 20(X_8 - 0.5)^2 + 10X_9 + 5X_{10} - 20X_9 X_{10} - 10.$$
(18)

**Data Generation Mechanism** To account for the correlated input scenario, we use Gaussian copula (Nelsen, 2006; Kucherenko et al., 2012) to model this dependency,

$$\mathbf{X} \sim C(G_1(X_1), \cdots, G_p(X_p); \mathbf{\Sigma}_{\mathbf{X}}) = \Phi_p(\Phi^{-1}(G_1(X_1)), \cdots, \Phi^{-1}(G_p(X_p)); \mathbf{\Sigma}))$$

where  $G_i$ 's are the univariate marginal cumulative distributions,  $C(\cdot; \Sigma_{\mathbf{X}})$  is the Gaussian copula with correlation matrix  $\Sigma_{\mathbf{X}}$ ,  $\Phi_p$  is the *p*-variate cumulative Gaussian distribution function with covariance matrix  $\Sigma$ , and  $\Phi^{-1}$  is the inverse univariate cumulative Gaussian distribution function. Note that there is a mapping from  $\Sigma_{\mathcal{X}}$  to  $\Sigma$ , but for our simulation study purpose, we focus on modeling  $\Sigma$  by letting  $\Sigma_{ij} = \rho^{|i-j|}$  for some  $0 \le \rho \le 1$  to control the level of correlation between the inputs. To sample  $\mathbf{X}$ , we first sample  $\mathbf{Z} \sim \mathcal{N}(0, \Sigma)$ then apply transformation  $X_i = G_i^{-1}(\Phi(Z_i))$  for each dimension. Uniform distribution over [0, 1] is used for all the marginal  $G_i$ 's, e.g.,  $\mathbf{X}$  follows the uniform distribution over the unit hypercube  $[0, 1]^p$  when  $\Sigma = \mathbf{I}_p$ . To obtain the noisy output, we add a random Gaussian noise  $\epsilon$  that is independent of  $\mathbf{X}$ , i.e.,

$$Y = f^*(\mathbf{X}) + \epsilon, \quad \epsilon \sim \mathcal{N}(0, 1).$$

For all simulations in this section, we generate N = 1,000 noisy samples.

**Groundtruth Factor Importance** Given its equivalence to the total Sobol' indices, the groundtruth factor importance measures  $\{\psi_i^*\}_{i=1}^p$  can be computed using the Double Monte Carlo estimator (9) with  $N_I = 2$  and  $N_O = 100,000$ . The Double Monte Carlo estimator requires sampling from the conditional distribution  $X_i | \mathbf{X}_{-i} = \mathbf{x}_{-i}$ , which can be done by first computing  $z_j = \Phi^{-1}(G_j(x_j))$  for  $j \in -i$ , then sampling  $Z_i$  conditional on  $\mathbf{Z}_{-i} = \mathbf{z}_{-i}$ , which is a conditional Gaussian distribution, and last applying transformation  $X_i = G_i^{-1}(\Phi(Z_i))$ . Recall from the discussion in Section 4, the groundtruth factor importance is

measured relative only to those variables that are presented in the true model.

#### 5.1 Low Dimensional Setting

For factor importance we compare FIRST (Algorithm 3) to the following competitors:

- VIMP-R2: sample-splitted, cross-fitted estimator (Williamson et al., 2023) of the R<sup>2</sup>-based intrinsic factor importance ψ<sub>i</sub> (1) implemented in the R package vimp (Williamson, 2023). For estimating the oracle functions f\* and f\*<sub>-i</sub> in (1), we use the random forest implemented in the R package ranger (Wright and Ziegler, 2017). For all other configurations, we follow the default values recommended in the package.
- FILTER: "filterVarImp" implemented in the R package caret (Kuhn et al., 2020), which assess the importance of each factor individually by the  $R^2$  of any univariate nonparametric model such as loess smoother fitted between the output and the factor.
- LASSO: Lasso (Tibshirani, 1996) implemented in the R package glmnet (Friedman et al., 2010). The regularization parameters are chosen by "lambda.1se". The factor importance is evaluated by the square of the coefficients.
- COSSO: COSSO (Lin and Zhang, 2006) implemented in the R package cosso (Zhang and Lin, 2023). The regularization parameters are chosen by the default tuning procedure provided in the package. The factor importance is evaluated by the variance of each component, i.e., suppose  $\hat{f}(\mathbf{X}) = \sum_{i=1}^{p} \hat{f}_i(X_i)$  is the additive model learned by COSSO, the importance of  $X_i$  is characterized by  $\operatorname{Var}[\hat{f}_i(X_i)]$ .
- RF: random forest (Breiman, 2001) implemented in the R package ranger (Wright and Ziegler, 2017). The factor importance is assessed by the mean decrease in the impurity, which is mean squared error in the regression setting.
- RF-SHAP: SHAP (SHapley Additive exPlanations) factor importance (Lundberg and Lee, 2017) based on the random forest. The Shapley value of each data instance is computed using the R package fastshap (Greenwell, 2021). The factor importance is evaluated by the mean absolute Shapley value across the data (Molnar, 2020).



Figure 2: Comparison of FIRST to other factor importance procedures on the Friedman function (18) with dimension p = 10 and correlation  $\rho = 0.8$ . For a better visual comparison, the importance measures are normalized to have a sum of one. For each procedure, 100 independent runs are performed, and the results are summarized by the boxplots, with the groundtruth marked by the red crosses. The average Kendall rank correlation coefficients (19), the higher the better, are shown in the top right corner.

• RF-VS: random forest with variable selection (Genuer et al., 2010) implemented in the R package VSURF (Genuer et al., 2015). Random forest are fitted only on the selected variables obtained from the "interpretation step". The importance is defined by the mean decrease in the impurity for the selected variables, and 0 otherwise.

For FIRST, we utilize  $N_I = 2$  nearest-neighbor for estimating the conditional variance. Figure 2 compares their performance on the p = 10 dimensional Friedman function (18) with input correlation  $\rho = 0.8$ . Our proposed FIRST procedure visually aligns the best with the groundtruth importance (red crosses), while some competitors, e.g. FILTER, RF, and RF-SHAP, struggle from the correlation in the input. The simple linear model LASSO overlooks the subtle nonlinear effect. COSSO and RF-VS are able to provide reasonable estimations, but at the computational costs that are about 6 and 30 times that of FIRST, respectively. Similar to FIRST, VIMP-R2 also aims to directly estimate intrinsic factor importance  $\psi_i$  (1) but through building estimators for the oracle functions  $f^*$  and  $f^*_{-i}$ . From



Figure 3: Comparison of FIRST to other factor importance procedures on the Ishigami, Heavy-tailed, and Friedman function across different correlation ( $\rho$ ) levels. The Kendall rank correlation coefficients (19) is averaged over 100 independent runs.

Figure 2 we can see that VIMP-R2 (i) is much more computationally expensive and (ii) has difficulty in filtering out variables that are deemed unimportant, but it does provide good importance estimation for the last four variables. To quantify the performance comparison, we consider the Kendall rank correlation coefficient  $\tau$  between the groundtruth  $\{\psi_i\}_{i=1}^p$  and the estimation  $\{\hat{\psi}_i\}_{i=1}^p$ ,

$$\tau = \frac{2}{p(p-1)} \sum_{i < j} \operatorname{sign}(\psi_i - \psi_j) \times \operatorname{sign}(\hat{\psi}_i - \hat{\psi}_j),$$
(19)

with value of 1 indicated perfect agreement between the two rankings, and -1 for perfect disagreement. This metric is chosen because for factor importance, the emphasis is on the ranking rather than the exact value. This also offers a fair comparison among different methods, since some captures the importance in squared error while other captures in absolute error. Again from Figure 2, FIRST stands out as the clear winner that best correlates in ranking with the groundtruth.

A more comprehensive comparison is provided in Figure 3 for different correlation level on (i) Ishigami function (16) with p = 6, (ii) heavy-tailed nonlinear function (17) with p = 8, and (iii) Friedman function (18) with p = 10. We can see that most of the time FIRST outperforms the other methods, and it is robust to the various input dependency level. COSSO and RF-VS occasionally take the lead, but their performances vary across



Figure 4: Comparison of FIRST to other factor selection procedures on the Ishigami, Heavytailed, and Friedman function across different correlation levels. For each procedure, 100 independent runs are performed, and the percentage of times that the model variables are exactly selected is reported.

different correlation level, especially on the Ishigami function. The other methods are evidently inferior and generally suffer from strong correlation presented in the input.

Now let us also examine the factor selection performance. Given that FILTER, RF, and RF-SHAP do not have a good criterion that automatically determines the number of variables to select, we omit them in this comparison. In addition to LASSO, COSSO, VIMP-R2, and RF-VS, we also compare FIRST to KFOCI (kernel feature ordering by conditional independence), a model-free forward stepwise variable selection procedure based on the kernel partial correlation (Huang et al., 2022) implemented in the R package KPC (Huang, 2021). The default configurations suggested in the package are used for KFOCI. Figure 4 reports the proportion of times that the true model variables are exactly selected, e.g., for Ishigami function (16) they are  $\{X_1, X_2, X_3\}$ . FIRST achieves the best possible performance on the Ishigami function, in particular, it selects  $\{X_1, X_2, X_3\}$  exactly 100% of the times for all correlation levels. For the Friedman function (18), FIRST remains the top performing method, especially when the inputs are strongly correlated ( $\rho = 0.9$ ): all the competitors fail completely while FIRST can still identify the true model variables exactly in about half of the times. For the Heavy-tailed nonlinear function (17) though FIRST slightly falls short of RF-VS and KFOCI, its performance is still competitive, selecting the correct model variables at least 85% of the cases. Overall FIRST stands out as the robust option against



Figure 5: Comparison of FIRST to other factor importance / selection procedures on the p = 100 dimensional Friedman (18) function across different correlation levels. For each procedure, 100 independent runs are performed, and the following metrics are reported: average Kendall rank correlation coefficient (19) / the percentage of times that the model variables are exactly selected / the average runtime.

	Ishigami (16)			Friedman (18)		
	$\rho = 0.0$	$\rho = 0.5$	$\rho = 0.9$	$\rho = 0.0$	$\rho = 0.5$	$\rho = 0.9$
p = 50	1.00/1.00/6s	0.99/1.00/6s	1.00/1.00/6s	0.99/1.00/10s	1.00/1.00/9s	0.96/0.67/5s
p = 100	1.00/1.00/7s	1.00/1.00/7s	1.00/1.00/7s	1.00/1.00/13s	1.00/1.00/9s	0.96/0.58/8s
p = 200	1.00/1.00/24s	1.00/1.00/24s	1.00/1.00/24s	1.00/1.00/22s	1.00/1.00/25s	0.96/0.62/21s
p = 500	1.00/1.00/59s	1.00/1.00/58s	1.00/1.00/58s	1.00/0.99/55s	1.00/1.00/103s	0.95/0.60/84s
p = 1000	1.00/1.00/117s	1.00/1.00/117s	1.00/1.00/115s	1.00/0.99/168s	$1.00/1.00/207 \mathrm{s}$	0.95/0.58/361s

Table 1: Performance of FIRST on the Ishigami and Friedman function with dimension p = 50, 100, 200, 500, 1000. For each setting, 100 independent runs are performed, and the following metrics are reported: average Kendall rank correlation coefficient (19) / the percentage of times that the model variables are exactly selected / the average runtime.

various levels of input correlation, and moreover it is computationally more efficient than its key competitors COSSO, RF-VS and KFOCI<sup>2</sup>.

#### 5.2 High Dimensional Setting

Figure 5 compares the performance of FIRST to its competitors on the 100-dimensional Friedman function (18). Again only  $X_1, X_7, X_8, X_9, X_{10}$  are important, and the rest 95 variables do not influence the output at all. We use N = 1,000 noisy samples for the simulations. We can see that FIRST not only outperforms the other procedures on both

<sup>&</sup>lt;sup>2</sup>For the Friedman function example with dimension p = 10 and correlation  $\rho = 0.8$  in Figure 2, the average runtime of KFOCI is 70.15 seconds, which is 40 times greater than the runtime of FIRST.

factor importance and selection, but more importantly it is orders of magnitude faster over its key competitors COSSO, RF-VS, and KFOCI. Moreover, Table 1 reports the simulation results of FIRST on p = 50, 100, 200, 500, 1000. For the Ishigami (16) function, FIRST is able to (i) identify the true model parameters exactly and (ii) learn the importance ranking accurately on all dimensions. Similar findings are also observed for the Friedman (18) function except on the extremely difficult strongly correlated case ( $\rho = 0.9$ ), yet the performance of FIRST does not degrade as the dimension increases. On the other hand, though the computational time of FIRST is reasonable for moderate (p = 200) dimensional problem, it is getting computationally more expensive as the dimension further increases. To address this computational challenge, in Appendix B we provide a fast version of FIRST that borrows idea from the *effect sparsity* principle (Wu and Hamada, 2011) to more efficiently filter out variables that are not important. Unfortunately it also comes with the trade-off of sacrificing some accuracy.

### 5.3 Real Data Example

In this subsection we study the performance of FIRST on some real world datasets. Let us start with the Abalone dataset (Nash et al., 1994) that is available on the UCI Machine Learning repository (Asuncion and Newman, 2007). This dataset consists of N = 4,177instances, and the objective is to predict the age of the abalone using the other eight physical measurements, including sex (male/female/infant), length (the longest shell measurement), diameter (perpendicular to length), height, whole weight, shucked weight (weight of the meat), viscera weight (weight of the gut after bleeding), and shell weight. Among these eight predictors, FIRST filters out height from the set of relevant variables for the age prediction. By fitting a random forest on the remaining seven variables, the out-of-bag prediction mean squared errors (OOB-MSE) is 4.59, achieving better results than the 4.61 OOB-MSE from the random forest fit on all eight predictors. Moreover, among the 256 possible subsets of the eight factors, the seven factors selected by FIRST yields the *best random forest model* with the smallest OOB-MSE. Table 2 reports the factor importance estimated by FIRST, and we can see that the ranking of the importance measures is in good

Factor	FIRST Importance	Increment in OOB-MSE if Dropped
Sex (categorical)	0.016~(6)	0.098(4)
Length (continuous)	0.012(7)	0.010(7)
Diameter (continuous)	0.022~(4)	0.020(6)
Height (continuous)	0.000(8)	0.000(8)
Whole weight (continuous)	0.040(2)	0.237(2)
Shucked weight (continuous)	0.094(1)	0.597(1)
Viscera weight (continuous)	0.019~(5)	0.059~(5)
Shell weight (continuous)	0.031~(3)	0.222(3)

Table 2: The factor importance estimated by **FIRST** on the Abalone dataset. The increment in out-of-bag prediction mean squared errors (OOB-MSE) if the factor is dropped from the *best random forest model* is also reported. The ranking is provided inside the parentheses.

agreement with the predictive power proxy by the increment in OOB-MSE if the variable is dropped from the *best random forest model*. This again demonstrate that the factor importance measure by **FIRST** nicely captures the predictive capability of the variables.

Now we further study its factor selection performance on four other popular real data examples: Auto MPG data (Asuncion and Newman, 2007), Boston Housing data (Harrison Jr and Rubinfeld, 1978), concrete compressive strength data (Yeh, 1998), and meat spectroscopy data (Thodberg, 1993). For each dataset, we perform 80%/20% train/test split after filtering out the instances with missing values, next apply FIRST to select important factors from the train set, then fit a random forest on the train set using FIRST selected variables, and last evaluate it on the test set to obtain the test mean squared errors (MSE). We repeat this process for 100 times and record the 20%, 50%, and 80% quantiles for (i) the number of selected factors and (ii) the ratio of test MSE to that from random forest fitting on the entire set of variables. From Table 3 we can see that the test MSE ratio is generally around one, showing that the predictive power does not degrade much from using the variables selected by FIRST. Moreover, on the meat spectroscopy dataset, FIRST is able to reduce the problem dimension from 100 to around 4, while still maintaining a comparable prediction performance to the random forest fit with all 100 factors.

Dataset	# Selected Factors	Test MSE Ratio
Auto MPG $(N = 392, p = 7)$	4[4,5]	1.04[0.98, 1.11]
Boston Housing $(N = 506, p = 13)$	8[6,9]	1.02[0.93, 1.14]
Concrete Compressive Strength $(N = 1030, p = 8)$	6[6,7]	$0.93[0.88,\!0.97]$
Meat Spectroscopy $(N = 215, p = 100)$	4[2,6]	0.93[0.77, 1.07]

Table 3: The number of selected factors and the ratio between the test MSE obtained from the random forest fitted on the selected covariates and the entire set of covariates. The median [20% quantile, 80% quantile] are reported from 100 independent replications.

### 6 Extension to Binary Classification Problems

In Section 5 we have demonstrated the excellent performance of FIRST on the regression problems. A natural question is whether this can be extended for binary classification problems. To answer this question, we first present simulation results of FIRST (Algorithm 3) on both synthetic and real data binary classification problems, and then provide intuition on why FIRST can be applied directly without any modification. Different from regression that  $N_I = 2$  nearest-neighbor are sufficient for estimating the conditional variance, from our simulation we find that  $N_I = 3$  is more robust for the binary classification problem. For the results presented in this section,  $N_I = 3$  is used for FIRST.

### 6.1 Synthetic Data Example

Similar to the regression synthetic example, we model the input X using Gaussian copula with marginals being uniform distributions and covariance matrix  $\Sigma$  defined by  $\Sigma_{ij} = \rho^{|i-j|}$ , where  $0 \leq \rho \leq 1$  controls the level of correlation. To obtain the binary outcome, we first convert the function output to a probability by the inverse-probit transformation, and then perform Bernoulli sampling using this probability, i.e.,

$$Y = \text{Bernoulli}(\Phi(f^*(\mathbf{X}))),$$

where  $\Phi$  is the standard Gaussian cumulative distribution function. N = 1,000 noisy binary output samples are generated for the simulations.

For the binary classification problem, Williamson et al. (2023) suggest that (i) clas-



Figure 6: Comparison of FIRST to other factor selection procedures on the Ishigami and the Friedman binary classification problem for different correlation ( $\rho$ ) levels. For each procedure, 100 independent runs are performed, and the following metrics are reported: the percentage of times that the model variables are exactly selected / average true positive rate of the selected variables / average false positive rate of the selected variables.

sification accuracy and (ii) area under the receiver operating characteristic curve (AUC) as the measures of predictiveness V for computing the factor importance  $\psi_u$  (1). The equivalence between the total Sobol' indices and the importance using  $R^2$  predictiveness measure in the regression setting no longer holds here. Hence, there is no straightforward way to compute the groundtruth factor importance. We focus on the factor selection perspective for the performance evaluation. Three metrics are reported: (i) the proportion of times that the model variables are exactly selected, (ii) the true positive rate (TPR), the probability of success detection, of the selected variables, and (iii) the false positive rate (FPR), the probability of false alarm, of the selected variables. We compare **FIRST** to the

Dataset	# Selected Factors	Test Accuracy Ratio
A vs. S Letter Recognition $(N = 1537, p = 16)$	8[7,9]	0.99[0.99, 1.00]
Australian Credit Approval $(N = 690, p = 14)$	5[4,6]	0.98[0.97, 1.01]
Breast Cancer $(N = 683, p = 9)$	5[4,6]	0.99[0.98, 1.00]
Pima Indians Diabetes $(N = 392, p = 8)$	3[2,4]	0.98[0.94, 1.03]
Titanic $(N = 714, p = 7)$	4[3,5]	0.99[0.97, 1.01]

Table 4: The number of selected factors and the ratio between the test accuracy obtained from the random forest fitted on the selected covariates and the entire set of covariates. The median [20% quantile, 80% quantile] are reported from 100 independent replications.

following competitors: LASSO, COSSO, RF-VS, VIMP-ACC, and KFOCI. VIMP-ACC is the classification adaptation of the VIMP-R2 considered in the regression comparison, which replaces the  $R^2$  predictiveness measure by the classification accuracy<sup>3</sup>. From Figure 6 we can see that FIRST is among one of the top performers: FIRST is marginally less effective than COSSO on the Ishigami function (16), and only outperformed by RF-VS on the Friedman function (18). FIRST achieves excellent false positive rate, demonstrating the reliability of the variables selected by FIRST. For the true positive rate, FIRST is not doing as good on the Friedman function, occasionally missing some true model variables. However, overall FIRST maintains a good balance between the true positive rate and the false positive rate compared to its competitors.

### 6.2 Real Data Example

Let us study the performance of FIRST on five real data applications: A vs. S Letter Recognition(Asuncion and Newman, 2007), Australian credit approval (Asuncion and Newman, 2007), Breast Cancer (Wolberg and Mangasarian, 1990), Pima Indians Diabetes (Ripley, 2007), and Titanic (Cukierski, 2012). Similar to the real data regression examples in Subsection 5.3, we first filter out the instances with missing values. Next, we split the dataset into 80% training and 20% testing and perform factor selection via FIRST on the train set. Last, we fit a random forest model on the train set using only the variables identified by FIRST and then evaluate the accuracy on the test set. Table 4 reports the ratio of the afore-

<sup>&</sup>lt;sup>3</sup>The simulations in Williamson et al. (2023) shows that using accuracy and AUC yield similar results.

mentioned test accuracy to that obtained by fitting random forest on all the factors. We can see that with FIRST, using only about half of the factors is able to achieve a prediction performance that is comparable to the random forest fit with all the covariates.

#### 6.3 Connection to the Impurity Measure

We now provide some intuition on why FIRST works for the binary classification problem. Recall that the total Sobol' index of  $X_i$  is

$$S_i^{\text{tot}} = \frac{\mathbb{E}[\operatorname{Var}\{Y | \mathbf{X}_{-i}\}]}{\operatorname{Var}[Y]},$$

where  $Y = \mathbb{1}(f^*(\mathbf{X}) > 0)$  is now a binary random variable. We have  $\operatorname{Var}[Y|\mathbf{X}_{-i}] = p_{-i}(1-p_{-i})$  where  $p_{-i} = \mathbb{E}[\mathbb{1}(f^*(\mathbf{X}) > 0)|\mathbf{X}_{-i}]$  is the probability of Y = 1 conditional on  $\mathbf{X}_{-i}$ . In the binary classification problem, p(1-p) is half of the Gini impurity measure, with maximum at p = 1/2 and minimum at p = 0 or p = 1. When the full model  $f^*(\mathbf{X})$  is known, the Gini impurity is 0 everywhere since we know whether Y = 1 with probability one. The total Sobol' index  $S_i^{\text{tot}}$  can then be interpreted as measuring the expected increment in the Gini impurity if  $X_i$  is dropped from the model, and the larger the increment the more important  $X_i$  is. This closely aligns with the importance definition of the tree models via the average reduction in the Gini impurity (Breiman, 2001). It is also straightforward to see that  $X_i$  is not part of the model if and only if  $S_i^{\text{tot}} = 0$ , and hence demonstrating that FIRST is appropriate for factor selection.

### 7 Conclusions

In this article we propose FIRST (Algorithm 3), a novel procedure for estimating factor importance directly from noisy data. FIRST is motivated from the equivalence between the total Sobol' indices and the intrinsic population-level variable importance, circumventing the model fitting step that is required by the existing factor importance procedures in the literature. Extensive simulations are provided in Section 5 and Section 6 to demonstrate that FIRST not only provides trustworthy factor importance estimation and selection across problems with varying correlation levels and dimensions, but more importantly it outshines its main competitors by completing the task in only a fraction of the computational time.

### Acknowledgments

This research is supported by U.S. National Science Foundation grants DMS-2310637 and DMREF-1921873.

## References

- Asuncion, A. and Newman, D. (2007), "UCI machine learning repository," URL https://archive.ics.uci.edu/.
- Azadkia, M. and Chatterjee, S. (2021), "A simple measure of conditional dependence," The Annals of Statistics, 49, 3070–3102.
- Bai, E.-w., Li, K., Zhao, W., and Xu, W. (2014), "Kernel based approaches to local nonlinear non-parametric variable selection," Automatica, 50, 100–113.
- Bentley, J. L. (1975), "Multidimensional binary search trees used for associative searching," Communications of the ACM, 18, 509–517.
- Borgonovo, E. (2007), "A new uncertainty importance measure," Reliability Engineering & System Safety, 92, 771–784.
- Borgonovo, E., Castaings, W., and Tarantola, S. (2011), "Moment independent importance measures: New results and analytical test cases," *Risk Analysis: An International Journal*, 31, 404–428.
- Borgonovo, E. and Plischke, E. (2016), "Sensitivity analysis: A review of recent advances," *European Journal of Operational Research*, 248, 869–887.

Breiman, L. (2001), "Random forests," Machine learning, 45, 5–32.

- Broto, B., Bachoc, F., and Depecker, M. (2020), "Variance reduction for estimation of Shapley effects and adaptation to unknown input distribution," SIAM/ASA Journal on Uncertainty Quantification, 8, 693–716.
- Chastaing, G., Gamboa, F., and Prieur, C. (2015), "Generalized Sobol sensitivity indices for dependent variables: numerical methods," *Journal of statistical computation and simulation*, 85, 1306–1333.
- Chatterjee, S. (2021), "A new coefficient of correlation," *Journal of the American Statistical* Association, 116, 2009–2022.
- Chen, W., Jin, R., and Sudjianto, A. (2005), "Analytical variance-based global sensitivity analysis in simulation-based design under uncertainty," *Journal of mechanical design*, 127, 875–886.
- Cukierski, W. (2012), "Titanic Machine Learning from Disaster," Retrieved November 30, 2023 from https://kaggle.com/competitions/titanic.
- Da Veiga, S., Gamboa, F., Iooss, B., and Prieur, C. (2021), Basics and Trends in Sensitivity Analysis: Theory and Practice in R, SIAM.
- Darlington, R. B. (1968), "Multiple regression in psychological research and practice." *Psychological bulletin*, 69, 161.
- Efron, B. and Stein, C. (1981), "The jackknife estimate of variance," *The Annals of Statistics*, 586–596.
- Efroymson, M. A. (1960), "Multiple regression analysis," Mathematical methods for digital computers, 191–203.
- Fisher, A., Rudin, C., and Dominici, F. (2019), "All Models are Wrong, but Many are Useful: Learning a Variable's Importance by Studying an Entire Class of Prediction Models Simultaneously." J. Mach. Learn. Res., 20, 1–81.

- Friedman, J., Hastie, T., and Tibshirani, R. (2010), "Regularization Paths for Generalized Linear Models via Coordinate Descent," *Journal of Statistical Software*, 33, 1–22, URL https://www.jstatsoft.org/v33/i01/.
- Friedman, J. H. (1991), "Multivariate adaptive regression splines," The annals of statistics, 19, 1–67.
- (2001), "Greedy function approximation: a gradient boosting machine," Annals of statistics, 1189–1232.
- Genuer, R., Poggi, J.-M., and Tuleau-Malot, C. (2010), "Variable selection using random forests," *Pattern recognition letters*, 31, 2225–2236.
- (2015), "VSURF: an R package for variable selection using random forests," The R Journal, 7, 19–33.
- Gevrey, M., Dimopoulos, I., and Lek, S. (2003), "Review and comparison of methods to study the contribution of variables in artificial neural network models," *Ecological modelling*, 160, 249–264.
- Greenwell, B. (2021), fastshap: Fast Approximate Shapley Values, URL https://CRAN. R-project.org/package=fastshap. R package version 0.0.7.
- Grömping, U. (2007), "Estimators of relative importance in linear regression based on variance decomposition," *The American Statistician*, 61, 139–147.
- (2009), "Variable importance assessment in regression: linear regression versus random forest," The American Statistician, 63, 308–319.
- Harrison Jr, D. and Rubinfeld, D. L. (1978), "Hedonic housing prices and the demand for clean air," Journal of environmental economics and management, 5, 81–102.
- Hart, J. and Gremaud, P. A. (2018), "An approximation theoretic perspective of Sobol'indices with dependent variables," *International Journal for Uncertainty Quan*tification, 8.

- Hastie, T. J. (2017), "Generalized additive models," in *Statistical models in S*, Routledge, 249–307.
- Homma, T. and Saltelli, A. (1996), "Importance measures in global sensitivity analysis of nonlinear models," *Reliability Engineering & System Safety*, 52, 1–17.
- Huang, Z. (2021), KPC: Kernel Partial Correlation Coefficient. R package version 0.1.0.
- Huang, Z., Deb, N., and Sen, B. (2022), "Kernel partial correlation coefficient—a measure of conditional dependence," *The Journal of Machine Learning Research*, 23, 9699–9756.
- Ishigami, T. and Homma, T. (1990), "An importance quantification technique in uncertainty analysis for computer models," in [1990] Proceedings. First international symposium on uncertainty modeling and analysis, IEEE.
- Jansen, M. J. (1999), "Analysis of variance designs for model output," Computer Physics Communications, 117, 35–43.
- Kucherenko, S., Tarantola, S., and Annoni, P. (2012), "Estimation of global sensitivity indices for models with dependent variables," *Computer physics communications*, 183, 937–946.
- Kuhn, M., Wing, J., Weston, S., Williams, A., Keefer, C., Engelhardt, A., Cooper, T., Mayer, Z., Kenkel, B., Team, R. C., et al. (2020), "Package 'caret'," *The R Journal*, 223.
- LeCun, Y., Bengio, Y., and Hinton, G. (2015), "Deep learning," nature, 521, 436–444.
- Li, G., Rabitz, H., Yelvington, P. E., Oluwole, O. O., Bacon, F., Kolb, C. E., and Schoendorf, J. (2010), "Global sensitivity analysis for systems with independent and/or correlated inputs," *The journal of physical chemistry A*, 114, 6022–6032.
- Li, R., Zhong, W., and Zhu, L. (2012), "Feature screening via distance correlation learning," Journal of the American Statistical Association, 107, 1129–1139.
- Lin, Y. and Zhang, H. H. (2006), "Component selection and smoothing in multivariate nonparametric regression," *The Annals of Statistics*, 34, 2272–2297.

- Lundberg, S. M. and Lee, S.-I. (2017), "A unified approach to interpreting model predictions," Advances in neural information processing systems, 30.
- Mase, M., Owen, A. B., and Seiler, B. B. (2022), "Variable importance without impossible data," arXiv preprint arXiv:2205.15750.
- Molnar, C. (2020), Interpretable machine learning, URL https://christophm.github. io/interpretable-ml-book/.
- Morris, M. D. (1991), "Factorial sampling plans for preliminary computational experiments," *Technometrics*, 33, 161–174.
- Murdoch, W. J., Singh, C., Kumbier, K., Abbasi-Asl, R., and Yu, B. (2019), "Interpretable machine learning: definitions, methods, and applications," *arXiv preprint arXiv:1901.04592*.
- Nash, W. J., Sellers, T. L., Talbot, S. R., Cawthorn, A. J., and Ford, W. B. (1994), "The population biology of abalone (haliotis species) in tasmania. i. blacklip abalone (h. rubra) from the north coast and islands of bass strait," *Sea Fisheries Division, Technical Report*, 48, p411.
- Nelsen, R. B. (2006), An introduction to copulas, Springer.
- Oakley, J. E. and O'Hagan, A. (2004), "Probabilistic sensitivity analysis of complex models: a Bayesian approach," Journal of the Royal Statistical Society: Series B (Statistical Methodology), 66, 751–769.
- Owen, A. B. (2014), "Sobol'indices and Shapley value," SIAM/ASA Journal on Uncertainty Quantification, 2, 245–251.
- Owen, A. B. and Prieur, C. (2017), "On Shapley value for measuring importance of dependent inputs," SIAM/ASA Journal on Uncertainty Quantification, 5, 986–1002.
- Plischke, E., Borgonovo, E., and Smith, C. L. (2013), "Global sensitivity measures from given data," *European Journal of Operational Research*, 226, 536–550.

- Rasmussen, C. E., Williams, C. K., et al. (2006), Gaussian processes for machine learning, volume 1, Springer.
- Razavi, S., Jakeman, A., Saltelli, A., Prieur, C., Iooss, B., Borgonovo, E., Plischke, E., Piano, S. L., Iwanaga, T., Becker, W., et al. (2021), "The future of sensitivity analysis: An essential discipline for systems modeling and policy support," *Environmental Modelling* & Software, 137, 104954.
- Ripley, B. D. (2007), Pattern recognition and neural networks, Cambridge university press.
- Saltelli, A., Annoni, P., Azzini, I., Campolongo, F., Ratto, M., and Tarantola, S. (2010), "Variance based sensitivity analysis of model output. Design and estimator for the total sensitivity index," *Computer physics communications*, 181, 259–270.
- Sobol', I. M. (1993), "On sensitivity estimation for nonlinear mathematical models," *Mathematical Modeling and Computational Experiments*, 1, 407–414.
- (2001), "Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates," *Mathematics and computers in simulation*, 55, 271–280.
- Sobol', I. M. and Kucherenko, S. (2009), "Derivative based global sensitivity measures and their link with global sensitivity indices," *Mathematics and Computers in Simulation*, 79, 3009–3017.
- Song, E., Nelson, B. L., and Staum, J. (2016), "Shapley effects for global sensitivity analysis: Theory and computation," SIAM/ASA Journal on Uncertainty Quantification, 4, 1060–1083.
- Székely, G. J., Rizzo, M. L., and Bakirov, N. K. (2007), "Measuring and testing dependence by correlation of distances," *The Annals of Statistics*, 35, 2769–2794.
- Thodberg, H. H. (1993), "Ace of bayes: Application of neural networks with pruning," *Technical report*.
- Tibshirani, R. (1996), "Regression shrinkage and selection via the lasso," Journal of the Royal Statistical Society Series B: Statistical Methodology, 58, 267–288.

- Wei, P., Lu, Z., and Song, J. (2015), "Variable importance analysis: A comprehensive review," *Reliability Engineering & System Safety*, 142, 399–432.
- Williamson, B. D. (2023), vimp: Perform Inference on Algorithm-Agnostic Variable Importance, URL https://CRAN.R-project.org/package=vimp. R package version 2.3.3.
- Williamson, B. D., Gilbert, P. B., Carone, M., and Simon, N. (2021), "Nonparametric variable importance assessment using machine learning techniques," *Biometrics*, 77, 9– 22.
- Williamson, B. D., Gilbert, P. B., Simon, N. R., and Carone, M. (2023), "A general framework for inference on algorithm-agnostic variable importance," *Journal of the American Statistical Association*, 118, 1645–1658.
- Wojtas, M. and Chen, K. (2020), "Feature importance ranking for deep learning," Advances in Neural Information Processing Systems, 33, 5105–5114.
- Wolberg, W. H. and Mangasarian, O. L. (1990), "Multisurface method of pattern separation for medical diagnosis applied to breast cytology." *Proceedings of the national academy* of sciences, 87, 9193–9196.
- Wright, M. N. and Ziegler, A. (2017), "ranger: A Fast Implementation of Random Forests for High Dimensional Data in C++ and R," *Journal of Statistical Software*, 77, 1–17.
- Wu, C. J. and Hamada, M. S. (2011), Experiments: planning, analysis, and optimization, John Wiley & Sons.
- Yang, L., Lv, S., and Wang, J. (2016), "Model-free variable selection in reproducing kernel Hilbert space," *The Journal of Machine Learning Research*, 17, 2885–2908.
- Ye, G.-B. and Xie, X. (2012), "Learning sparse gradients for variable selection and dimension reduction," *Machine learning*, 87, 303–355.
- Yeh, I.-C. (1998), "Modeling of strength of high-performance concrete using artificial neural networks," *Cement and Concrete research*, 28, 1797–1808.

Zhang, H. H. and Lin, C.-Y. (2023), cosso: Fit Regularized Nonparametric Regression Models Using COSSO Penalty, URL https://CRAN.R-project.org/package=cosso. R package version 2.1-2.

# Appendices

## A Proof of Lemma 1

Lemma 1.  $\mathbb{E}[(\mathbb{E}[Y|X] - Y)^2] = \mathbb{E}[\operatorname{Var}(Y|X)].$ 

Proof.

$$\mathbb{E}[(\mathbb{E}(Y|X) - Y)^2] = \mathbb{E}[\mathbb{E}(Y|X)^2] + \mathbb{E}[Y^2] - 2\mathbb{E}[\mathbb{E}(Y|X)Y]$$

$$= \mathbb{E}[\mathbb{E}(Y|X)^2] + \mathbb{E}[Y^2] - 2\mathbb{E}[\mathbb{E}\{\mathbb{E}(Y|X)Y|X\}]$$

$$= \mathbb{E}[\mathbb{E}(Y|X)^2] + \mathbb{E}[Y^2] - 2\mathbb{E}[\mathbb{E}(Y|X)^2]$$

$$= \mathbb{E}[Y^2] - \mathbb{E}[\mathbb{E}(Y|X)^2]$$

$$= (\mathbb{E}[Y^2] - \mathbb{E}[Y]^2) - (\mathbb{E}[\mathbb{E}(Y|X)^2] - \mathbb{E}[Y]^2)$$

$$= (\mathbb{E}[Y^2] - \mathbb{E}[Y]^2) - (\mathbb{E}[\mathbb{E}(Y|X)^2] - \mathbb{E}[\mathbb{E}(Y|X)]^2)$$

$$= \operatorname{Var}[Y] - \operatorname{Var}[\mathbb{E}(Y|X)]$$

$$= \mathbb{E}[\operatorname{Var}(Y|X)].$$

## **B** Additional Simulation Results on Regression

Figure S1 compares the performance of FIRST to its competitors on the 100-dimensional Ishigami function (16) with N = 1,000 noisy samples. Similar to the comparison on the Friedman function in Figure 5, FIRST outperforms the other methods for factor importance ranking in terms of the Kendall rank correlation coefficient. Moreover, FIRST is able to identify the true variables  $\{X_1, X_2, X_3\}$  exactly 100% of the times for different correlation levels. Compared to its key competitors RF-VS (for factor importance ranking) and KFOCI (for factor selection), FIRST only uses a fraction of their runtimes to achieve a better and more robust performance on this 100 dimensional example.

Though the computational time is fast for the 100 dimensional examples (Figure 5 and



Figure S1: Comparison of FIRST to other factor importance / selection procedures on the p = 100 dimensional Ishigami (16) function across different correlation levels. For each procedure, 100 independent runs are performed, and the following metrics are reported: average Kendall rank correlation coefficient (19) / the percentage of times that the model variables are exactly selected / the average runtime.

S1), from Table 1 we can see that FIRST is getting computationally more expensive as the dimension grows up. For the strongly correlated ( $\rho = 0.9$ ) 1,000-dimensional input Friedman function example, on average FIRST takes 361 seconds to complete the computation. To further speed up FIRST, we incorporate the *effect sparsity* principle(Wu and Hamada, 2011). The key is to filter out irrelevant factors more efficiently to obtain a parsimonious model. Instead of considering all factors except for the selected ones as candidate in each forward selection step, we only keep the variables that are likely to improve the output variance that can be explained in the candidate set, i.e., in each forward selection step we remove variables that would diminish the explainable variance from the candidate set. Algorithm S1 details this fast version of FIRST, and the difference from FIRST (Algorithm 3) is highlighted in red. Table S1 reports the comparison of FIRST and FIRST-FAST on the p = 50, 100, 200, 500, 1000 dimensional Friedman function. Clearly we can see that FIRST-FAST is able to reduce the runtime of FIRST by more than 3 times, but in the sacrifice of worse true positive rate, i.e., FIRST-FAST is more likely to leave out some crucial variables. If the computational time is not a major bottleneck, we would recommend FIRST for its better accuracy.

#### Algorithm S1 FIRST-FAST.

- 1: Input: (i) noisy samples  $\{(\mathbf{x}^{(n)} \in \mathbb{R}^p, y^{(n)} \in \mathbb{R})\}_{n=1}^N$ , (ii) number of samples for inner loop Monte Carlo  $N_I$ , and (iii) number of samples for outer loop Monte Carlo  $N_O$ .
- 2: Initialization:  $k = -1, A \leftarrow \emptyset, \{i_0\} = \emptyset, \hat{V}_{\emptyset} = 0, C \leftarrow \{1:p\};$
- 3: **do**
- $4: \qquad k \leftarrow k+1;$
- 5:  $A \leftarrow A \cup \{i_k\};$
- $6: \qquad C \leftarrow C \setminus \{i_k\};$
- 7: Choose  $i_{k+1} \in C$  such that  $\hat{V}_{A \cup \{i_{k+1}\}}$  (15) is maximized, i.e.,

$$i_{k+1} = \arg \max_{i \in 1: p \setminus A} \hat{V}_{A \cup \{i\}},$$

8: and remove  $i \in C$  from C such that  $\hat{V}_{A \cup \{i\}} < \hat{V}_A$ , i.e.,

$$C \leftarrow C \setminus \{i \in C : V_{A \cup \{i\}} < V_A\};\$$

- 9: while |C| > 0;
- 10:  ${\hat{S}_i^{\text{tot}}}_{i \in A} \leftarrow \text{NANNE-BE}({(x_A^{(n)}, y^{(n)})}_{n=1}^N, N_I, N_O);$ 11:  $\hat{\psi}_i \leftarrow \hat{S}_i^{\text{tot}}$  for  $i \in A$  and  $\hat{\psi}_i \leftarrow 0$  for  $i \in 1: p \setminus A;$
- 12: **Output:** the factor importance  $\{\hat{\psi}_i\}_{i=1}^p$ .

	$\rho = 0.0$		$\rho = 0.5$		$\rho = 0.9$	
	FIRST	FIRST-FAST	FIRST	FIRST-FAST	FIRST	FIRST-FAST
p = 50	0.99/1.00/0.00/10s	0.88/0.80/0.00/3s	1.00/1.00/0.00/9s	0.93/0.87/0.00/3s	0.96/0.94/0.00/5s	0.91/0.84/0.00/2s
p = 100	1.00/1.00/0.00/13s	0.89/0.81/0.00/9s	1.00/1.00/0.00/9s	0.94/0.89/0.00/3s	0.96/0.93/0.00/8s	0.92/0.85/0.00/3s
p = 200	1.00/1.00/0.00/22s	0.89/0.81/0.00/6s	1.00/1.00/0.00/25s	0.94/0.89/0.00/7s	0.96/0.93/0.00/21s	0.92/0.85/0.00/6s
p = 500	1.00/1.00/0.00/55s	0.89/0.81/0.00/14s	1.00/1.00/0.00/103s	0.94/0.88/0.00/28s	0.95/0.92/0.00/84s	0.91/0.84/0.00/24s
p = 1000	$1.00/1.00/0.00/168 \mathrm{s}$	$0.88/0.81/0.00/42 \mathrm{s}$	$1.00/1.00/0.00/207 \mathrm{s}$	$0.94/0.88/0.00/55 \mathrm{s}$	$0.95/0.92/0.00/361 \mathrm{s}$	$0.92/0.85/0.00/105 \mathrm{s}$

Table S1: Comparison of FIRST and FIRST-FAST on the Friedman function with dimension p = 50, 100, 200, 500, 1000. For each setting, 100 independent runs are performed, and the following metrics are reported: average Kendall rank correlation coefficient (19) / true positive rate / false positive rate / the average runtime.

### C Minor Implementation Details

We want to point out a minor implementation detail of the NANNE estimator (Algorithm 1) that is different from the nearest-neighbor estimator of Broto et al. (2020). For the nearest-neighbor estimator, when there are multiple nearest neighbors that are at the same distance, Broto et al. (2020) suggests to break the tie by random selection and obtain exactly  $N_I$  nearest neighbors for the conditional variance computation. In actual implementation,

we notice that the random selection introduces noise to the estimation, especially when majority of the variables are categorical. Hence, instead of breaking tie by random selection, our proposed NANNE estimator computes the conditional variance using all the samples that are within the distance of the  $N_I$ -th nearest neighbor. This minor modification exhibits a more robust performance on the real world datasets.