WEAK COMPACTONS OF NONLINEARLY DISPERSIVE KdV AND KP EQUATIONS

S.C. ANCO¹ AND M.L. GANDARIAS²

¹Department of Mathematics and Statistics Brock University St. Catharines, ON L2S3A1, Canada

²DEPARTMENT OF MATHEMATICS FACULTY OF SCIENCES, UNIVERSITY OF CÁDIZ PUERTO REAL, CÁDIZ, SPAIN, 11510

ABSTRACT. A weak formulation is devised for the K(m, n) equation which is a nonlinearly dispersive generalization of the gKdV equation having compacton solutions. With this formulation, explicit weak compacton solutions are derived, including ones that do not exist as classical (strong) solutions. Similar results are obtained for a nonlinearly dispersive generalization of the gKP equation in two dimensions, which possesses line compacton solutions.

keywords: compacton; weak solution; travelling wave; cut off; nonlinear dispersion; KdV equation; K(m, n) equation; line compacton; KP equation

1. INTRODUCTION

In the study of dispersive nonlinear wave equations, solutions of basic importance are travelling waves

$$u = U(\xi), \quad \xi = x - ct \tag{1}$$

where c is the wave speed. Stable travelling waves for which the profile U goes to zero for large $|\xi|$ describe solitary waves. Physical and mathematical features of such solutions have been extensively for a large number of dispersive nonlinear wave equations.

Another kind of travelling wave which has attracted much interest are compactons that describe waves with compact spatial support — see the review article [1] and references therein. For a given wave equation, a compacton is a strong (classical) solution (1) whose profile is, without loss of generality via translation, supported on $-L \leq \xi \leq L$ and vanishing for $|\xi| > L$, with L > 0. Such solutions can be constructed by starting with a travelling wave such that $U(\xi)$ and derivatives of $U(\xi)$ that appear in the equation each vanish at $\xi = \pm L$, which allows extending $U(\xi)$ to be 0 outside of $-L \leq \xi \leq L$.

Compactons were first found for a nonlinearly-dispersive generalization of the KdV equation, known as the K(m, n) equation [2],

$$u_t + a(u^m)_x + b(u^n)_{xxx} = 0 (2)$$

where m, n > 0 are the nonlinearity powers, and $a, b \neq 0$ are the coefficients. For n = 1, this is the gKdV equation, which reduces to the KdV equation when m = 2. Nonlinear dispersion arises for $n \neq 1$. The simplest examples of a compacton have a profile given by a

cosine function in the cases $1 < m = n \leq 3$ [2] and $n = 1, \frac{1}{2} < m < 1$ [3, 4], with U being C^3 . The latter case has linear dispersion but sub-linear convection.

However, many papers in the literature focus on compact wave profiles that are only C^0 or C^1 ; it is crucial to emphasize that, for any given dispersive nonlinear wave equation, such profiles are not actual solutions because they lack sufficient differentiability to satisfy the equation. These formal piecewise "solutions" are not very meaningful and cannot arise in the context of evolution of initial data.

Nevertheless, as explained in the present work, compactons can be defined with less differentiability than strong solutions. In particular, there is a well-known weak formulation of nonlinear wave equations which can be adapted to obtain compactons as weak solutions satisfying a dispersive nonlinear wave equation in a pointwise distributional sense. This makes precise a notion of distributional compactons introduced originally in Ref. [5]. Importantly, weakening the differentiability allows for additional solutions, which do not exist as strong solutions.

The weak formulation of compactons will be considered here for the K(m, n) equation (2) as well as an analogous nonlinearly-dispersive generalization of the KP equation given by

$$(u_t + a(u^m)_x + b(u^n)_{xxx})_x + \sigma u_{yy} = 0$$
(3)

with nonlinearity powers m, n > 0 $(m, n \neq 1)$, and coefficients $a, b \neq 0$, $\sigma^2 = 1$. This is called the KP(m, n) equation. It reduces to the KP equation when n = 1 and m = 2. The KP equation possesses line solitary waves [6], which are 2 dimensional counterparts of KdV solitary waves. Correspondingly, a *line compacton* is a travelling wave of the form

$$u = U(\xi), \quad \xi = x + \mu y - \nu t \tag{4}$$

such that its profile U has compact support. The parameters (μ, ν) determine both the direction and the speed of the wave. Specifically, $c = \nu/\sqrt{1+\mu^2}$ is the speed, and $\theta = \arctan(\mu)$ is the angle with respect to the positive x axis.

A complete classification of compacton strong solutions for the KP(m, n) equation has been obtained recently in Ref. [7]. This yielded many types of explicit compactons whose profiles are given by powers of several different functions — cosine and sine, Jacobi cn and sn, and quadratics. The main results of the present work consist of finding:

- additional explicit KP(m, n) compactons with different profiles
- less restrictive existence conditions for KP(m, n) compactons as weak solutions
- K(m, n) compactons have two-dimensional counterparts given by KP(m, n) line compactons

2. TRAVELLING WAVE ODE AND SOLUTION PROFILES

Travelling waves (1) for the K(m, n) equation (2) are given by the nonlinear third-order ODE

$$(-kU + aU^m + b(U^n)'')' = 0 (5)$$

where

$$k = c \tag{6}$$

A similar fourth-order ODE

$$(-kU + aU^m + b(U^n)'')'' = 0 (7)$$

arises for line travelling waves (4) for the KP(m, n) equation (3), with

$$k = \nu - \sigma \mu^2 = c |\sec \theta| - \sigma \tan \theta \tag{8}$$

Note that every solution of the third-order ODE (5) will satisfy the fourth-order ODE (7).

As shown in Ref. [7], it is convenient to change variables

$$U = V^{1/n} \tag{9}$$

Integration of the ODE (5) with respect to ξ followed by use of the integrating factor V' yields a first-order separable ODE for $V(\xi)$

$$V^{\prime 2} = E + CV + BV^{1+1/n} - AV^{1+m/n}$$
(10)

where E and C are constants of integration, and where

$$B = \frac{2nk}{(n+1)b}, \quad A = \frac{2na}{(m+n)b} \tag{11}$$

This ODE has the form of the energy equation of a nonlinear oscillator. It possess a wide variety of solutions that vanish at the endpoints of an interval $-L \leq \xi \leq L$. Under suitable conditions on n and m, these solutions can be cutoff at $\xi = \pm L$ to get a compacton strong solution whose profile is centered at $\xi = 0$ and vanishes for $|\xi| \geq L$.

2.1. Compacton profiles. Corresponding compactons of the K(m, n) equation (2) and the KP(m, n) equation (3) have the form

$$u = U_{c}(\xi) = U(\xi)H(L - |\xi|)$$
(12)

where $H(\xi)$ is the step function. A complete classification of compactons (12), which exist as strong solutions, is presented in Ref. [7] for the KP(m, n) equation. Solutions were obtained with explicit profiles $U(\xi)$ of the following types:

algebraic:
$$\alpha (1 - \beta \xi^2)^p$$
 (13a)

cosine:
$$\alpha \cos(\beta \xi)^p$$
 (13b)

sine: $\alpha \sin(\beta \xi)^p$ (13c)

Jacobi cn:
$$\alpha \operatorname{cn}(\beta\xi, l)^p$$
, $\alpha \operatorname{cn}(\beta(\xi + \frac{1}{2}L), l)^p$ (13d)

Jacobi sn:
$$\alpha \operatorname{sn}(\beta \xi, l)^p$$
, $\alpha \operatorname{sn}(\beta (\xi + L), l)^p$, (13e)

where α , β are real constants, l is a real or imaginary constant, and p is a positive power. Profiles (13a), (13b), along with the first of the two profiles (13d) and the second of the two profiles (13e), are symmetric in ξ , whereas the other profiles are antisymmetric when p is an odd integer or an odd fraction.

It is worth noting that the Jacobi cn and sn profiles (13d)-(13e) can be written in several different ways by means of various identities (see e.g. Ref. [8]):

$$cn(z,l)^{2} + sn(z,l)^{2} = 1, \quad dn(z,l)^{2} = 1 - l^{2}sn(z,l)^{2} = 1 - l^{2} + l^{2}cn(z,l)^{2}$$

$$sn(z,1/l) = lsn(z/l,l), \quad cn(z,1/l) = dn(z/l,l)$$

$$sn(z,li) = \frac{sn(\sqrt{1+l^{2}}z,l/\sqrt{1+l^{2}})^{2}}{\sqrt{1+l^{2}}dn(\sqrt{1+l^{2}}z,l/\sqrt{1+l^{2}})^{2}}, \quad cn(z,li) = \frac{cn(\sqrt{1+l^{2}}z,l/\sqrt{1+l^{2}})^{2}}{dn(\sqrt{1+l^{2}}z,l/\sqrt{1+l^{2}})^{2}}$$
(14)

Note that here, as in Ref. [7], the Maple convention is used for the parameter in the Jacobi cn and sn functions, which is the square root of the standard parameter for these functions.

The following type of symmetric profile

rational Jacobi cn:
$$\alpha \left(\frac{\operatorname{cn}(\beta\xi, l) + \kappa}{\operatorname{cn}(\beta\xi, l) + \gamma}\right)^q, \quad \kappa, \gamma \neq 0$$
 (15)

cannot be reduced to any of the previous profiles (13d) and (13e). It will be considered in section 4.

3. Weak formulation for compacton solutions

A standard weak formulation [9] for nonlinear PDEs will be adapted here for the K(m, n) equation (2) and the KP(m, n) equation (3).

In general, a weak formulation of a PDE with a dependent variable u is an integral equation that is obtained by first assuming that u is a strong solution, then multiplying the PDE by a test function and integrating by parts to leave as few as derivatives on u as necessary so that u has the lowest regularity possible. (Note that this does not require explicitly introducing a notion of weak derivatives. Instead u will simply be assumed to belong to a suitable function space.)

3.1. Weak form of K(m, n) equation. Let $\psi(t, x)$ be a test function, namely smooth with compact support. Then the weak formulation of the K(m, n) equation (2) is given by

$$\iint_{\mathbb{R}\times\mathbb{R}} (u\psi_t + au^m\psi_x + bu^n\psi_{xxx}) \, dt \, dx = 0 \tag{16}$$

with $u(t,x) \in C^0(\mathbb{R} \times \mathbb{R})$ being a weak solution iff this integral equation holds for all test functions.

In case of travelling waves (1), the integral equation (16) can be simplified via two steps. First, change variables $(t, x) \to (\tau, \xi)$ where $\tau = t$, yielding $\iint_{\mathbb{R} \times \mathbb{R}} (U(\psi_{\tau} - c\psi_{\xi}) + aU^m\psi_{\xi} + bU^n\psi_{\xi\xi\xi}) d\tau d\xi = 0$. Next, use $\int_{\mathbb{R}} U\psi_{\tau} d\tau = -\int_{\mathbb{R}} U_{\tau}\psi d\tau$ due to $U_{\tau} = 0$. This gives

$$\iint_{\mathbb{R}\times\mathbb{R}} \left((-cU + aU^m)\psi_{\xi} + bU^n\psi_{\xi\xi\xi} \right) d\tau \, d\xi = 0 \tag{17}$$

whereby $u = U(\xi)$ is weak travelling wave solution of the K(m, n) equation (2) if this integral equation holds for all test functions $\psi(\xi, \tau)$.

Similarly, the travelling wave ODE (5)-(6) has the weak formulation

$$\int_{\mathbb{R}} \left((-kU + aU^m)\phi' + bU^n\phi''' \right) d\xi = 0$$
(18)

with $U(\xi) \in C^0(\mathbb{R})$ being a weak solution if this integral equation holds for all test functions $\phi(\xi)$.

Comparison of the integral equations (18) and (17) shows that $U(\xi)$ is a weak solution of the travelling wave ODE (5)–(6) when (and only when) $u = U(\xi)$ is a weak solution of the K(m, n) equation (2).

3.2. Weak form of KP(m, n) equation. The preceding weak formulations extend straightforwardly to the KP(m, n) equation (3): $u(t, x, y) \in C^0(\mathbb{R} \times \mathbb{R})$ is weak solution if the integral equation

$$\iiint_{\mathbb{R}\times\mathbb{R}^2} (u\psi_{tx} + \sigma u\psi_{yy} + au^m\psi_{xx} + bu^n\psi_{xxxx}) \,dt\,dx\,dy = 0 \tag{19}$$

holds for all test functions $\psi(t, x, y)$. In case of line travelling waves (4), this integral equation simplifies to the form

$$\iiint_{\mathbb{R}\times\mathbb{R}^2} ((-kU+aU^m)\psi_{\xi\xi}+bU^n\psi_{\xi\xi\xi\xi}) \,d\tau \,d\zeta \,d\xi = 0$$
(20)

using variables $\tau = t, \zeta = y$, where k is expression (8).

The travelling wave ODE (7)-(8) has the similar weak formulation

$$\int_{\mathbb{R}} ((-kU + aU^m)\phi'' + bU^n\phi'''') d\xi = 0$$
(21)

with $U(\xi) \in C^0(\mathbb{R})$ being a weak solution if this integral equation holds for all test functions $\phi(\xi)$. Thus, $U(\xi)$ is a weak solution of the travelling wave ODE (7)–(8) when (and only when) $u = U(\xi)$ is a weak solution of the K(m, n) equation (2).

3.3. Derivation of existence conditions for weak solutions. Conditions for a travelling wave $U(\xi)$ to yield a compacton weak solution $U_{\rm c}(\xi) = U(\xi)H(L-|\xi|)$ will now be derived from the respective integral equations (18) and (21). The following preliminary result is needed.

Lemma 3.1. For any function $f(\xi)$ that is at least C^1 , and any test function $\phi(\xi)$,

$$\int_{\mathbb{R}} \phi'(\xi) f(\xi) H(L - |\xi|) \, d\xi = -\phi(L) f(L) - \phi(-L) f(-L) - \int_{\mathbb{R}} \phi(\xi) f'(\xi) H(L - |\xi|) \, d\xi \quad (22)$$

is an identity.

This can be proved with standard methods (see e.g. Ref. [10]) by means of integration by parts and properties of the Dirac delta distribution $\delta(L - |\xi|) = H'(L - |\xi|)$.

To proceed using this Lemma, substitute the expression $U_{\rm c}(\xi) = U(\xi)H(L-|\xi|)$ into the integral equations (18) and (21) and use $H^m = H^n = H$ to get

$$0 = \int_{\mathbb{R}} \left((-kU_{\rm c} + aU_{\rm c}^m)\phi' + bU_{\rm c}^n\phi''' \right) d\xi = \int_{\mathbb{R}} \left((-kU + aU^m)H\phi' + bU^nH\phi''' \right) d\xi \qquad (23)$$

and

$$0 = \int_{\mathbb{R}} \left((-kU_{\rm c} + aU_{\rm c}^m)\phi'' + bU_{\rm c}^n\phi'''' \right) d\xi = \int_{\mathbb{R}} \left((-kU + aU^m)H\phi'' + bU^nH\phi'''' \right) d\xi = 0 \quad (24)$$

Next, a repeated application of the relation (22) yields

$$0 = \int_{\mathbb{R}} \left((-kU + aU^m)' + b(U^n)''' \right) H\phi \, d\xi - \left(\phi'' A_1 - \phi' A_2 + \phi A_3 \right) \Big|_{\xi = \pm L}$$
(25)

and

$$0 = \int_{\mathbb{R}} \left((-kU + aU^m)'' + b(U^n)'''' \right) H\phi \, d\xi - \left(\phi'''A_1 - \phi''A_2 + \phi'A_3 + \phi A_4 \right) \Big|_{\xi = \pm L}$$
(26)

where

$$A_1 = bU^n, \quad A_2 = b(U^n)', \quad A_3 = -kU + aU^m + b(U^n)''$$
 (27)

and

$$A_4 = (-kU + aU^m)' + b(U^n)'''$$
(28)

with $a \neq 0$ and $b \neq 0$.

In equations (25) and (26), the integral term vanishes when $U(\xi)$ satisfies the respective travelling wave ODEs. This leaves the boundary terms. Since $\phi|_{\xi=\pm L}$, $\phi'|_{\xi=\pm L}$, $\phi''|_{\xi=\pm L}$, $\phi''|_{\xi$

$$A_i|_{\xi=\pm L} = 0, \quad i = 1, 2, 3 \tag{29}$$

are satisfied, as well as the condition

$$A_4|_{\xi=\pm L} = 0 \tag{30}$$

in the case of equation (26).

Remark 3.1. Existence conditions (29) and (30) can be derived in an alternative way directly from the K(m,n) and KP(m,n) equations through substitution of the cutoff profile expression (12) and use of the following steps: firstly, simplify $H(L - |\xi|)^q = H(L - |\xi|)$ for any q > 0; secondly, integrate by parts and use $H(\xi \pm L)' = \delta(\xi \pm L)$; thirdly, collect coefficients of separate singular terms and set each coefficient to zero. This corresponds to the PDEs being formally satisfied in pointwise distributional sense.

These conditions (29) and (30) will now be investigated in detail. Suppose that $U(\xi)$ has behaviour

$$U(\xi) \sim U_0 \left(L \mp \xi \right)^p \text{ as } \xi \sim \pm L \tag{31}$$

for some power $p \neq 0$, where $U_0 \neq 0$ is a constant.

The first and second conditions $A_1 = A_2 = 0$ require that, near $\xi = \pm L$, both expressions

$$A_1 \sim bU_0^n (L \mp \xi)^{pn} \quad \text{and} \quad A_2 \sim bpn U_0^n (L \mp \xi)^{pn-1}$$
(32)

must go to zero. This implies p > 0 and pn > 1, since n > 0. The third condition similarly requires that

$$A_3 \sim -kU_0^p (L \mp \xi)^p + aU_0^m (L \mp \xi)^{pm} + bpn(pn-1)U_0^n (L \mp \xi)^{pn-2}$$
(33)

goes to zero. This reduces to $(L \mp \xi)^{pn-2} \sim 0$ due to the previous two conditions combined with m > 0, which implies pn > 2.

Thus, the three conditions (29) jointly hold iff pn > 2.

The fourth condition (30) is more complicated to analyze. Near the cut off, A_4 is given by the sum of the terms

$$-kpU_0^p(L\mp\xi)^{p-1}, \quad apmU_0^m(L\mp\xi)^{pm-1}, \quad bpn(pn-1)(pn-2)U_0^n(L\mp\xi)^{pn-3}$$
(34)

with $a \neq 0$, $b \neq 0$, and p > 0. It will be convenient to consider separately the cases k = 0 and $k \neq 0$.

Case k = 0: The second and third terms either each go to zero, which gives pm > 1 and pn > 3, or they mutually cancel, which holds when $pm - 1 = pn - 3 \le 0$ and $apmU_0^m + bpn(pn - 1)(pn - 2)U_0^n = 0$. The first case is equivalent to $p > \max(\frac{1}{m}, \frac{3}{n})$, while the second case gives $p = \frac{2}{n-m}$ and $n \ge 3m$, and also $apmU_0^m + bpn(pn - 1)(pn - 2)U_0^n = 0$.

Case $k \neq 0$: The first term either goes to zero or has to cancel the third term, since $m \neq 1$. In the latter case, the middle term must go to zero, which gives pm - 1 > 0 and $p - 1 = pm - 3 \leq 0$ as well as $kU_0^p = bn(pn - 1)(pn - 2)U_0^n$. The two inequalities imply $1 \geq p = \frac{2}{n-1} > \frac{1}{m}$, and hence $n \geq 3$. In the former case, p must be greater than 1, and the sum of the second and third terms must vanish. This can happen in several different ways depending on whether m > 1 or m < 1.

If m > 1, then p > 1 implies pm > 1, and so the second term goes to term. This leaves the third term which then must also go to zero. Hence pn - 3 > 0, and all of these inequalities together then require $p > \max(1, \frac{3}{n})$.

If m < 1, then either the second term and the third term mutually cancel or they separately go to zero. The latter case has pm - 1 > 0 and pn - 3 > 0. All of the inequalities together thereby require $p > \max(\frac{1}{m}, \frac{3}{n})$. The former case gives $pm - 1 = pn - 3 \le 0$ and also $apmU_0^m + bpn(pn - 1)(pn - 2)U_0^n = 0$. This requires $\frac{1}{m} \ge p = \frac{2}{n-m} > 1$ using m < 1, which then implies $2 + m > n \ge 3m$.

In each of the previous cases, the inequality pn > 2 can be checked to hold. This completes the analysis of the conditions (29) and (30), which establishes the following key result necessary for existence of weak compactons.

Theorem 3.1. Suppose a compacton profile (12) has the behaviour (31) at the end points of its support.

(i) It is a weak solution of the K(m,n) equation, with m, n > 0 and $m \neq 1$, iff $U(\xi) \in C^3(\mathbb{R})$ satisfies the travelling wave ODE (5)–(6) and p obeys

$$p > 2/n. \tag{35}$$

(ii) It is a weak solution of the KP(m, n) equation, with m, n > 0 and $m \neq 1$, iff $U(\xi) \in C^4(\mathbb{R})$ satisfies the travelling wave ODE (7)-(8) and n, m, p, U₀ obey at least one of

$$p > \max(1/m, 3/n), \quad k = 0;$$
 (36a)

$$p > \max(1/m, 3/n), \quad m < 1, \quad k \neq 0;$$
 (36b)

$$p > \max(1, 3/n), \quad m > 1, \quad k \neq 0;$$
 (36c)

$$1 \ge p = 2/(n-1), \quad 2m+1 > n \ge 3, \quad k \ne 0, \quad U_0 = \left(\frac{k}{2b} \frac{(n-1)^2}{(n-2)(n+1)}\right)^{\frac{n-1}{(n-2)(n+1)}};$$
 (36d)

$$p = 2/(n-m), \quad n \ge 3m, \quad k = 0, \quad U_0 = \left(\frac{-a}{2b}\frac{(n-m)^2}{n(n+m)}\right)^{\frac{1}{n-m}};$$
 (36e)

$$p = 2/(n-m), \quad m+2 > n \ge 3m, \quad k \ne 0, \quad U_0 = \left(\frac{-a}{2b}\frac{(n-m)^2}{n(n+m)}\right)^{\frac{1}{n-m}}.$$
 (36f)

In all six cases (36), the inequality (35) is satisfied.

3.4. Existence conditions for strong solutions. In comparison, the conditions for existence of compacton strong solutions (12) for the K(m, n) and KP(m, n) equations are simply that $U(\xi)$ satisfies the respective travelling wave ODE pointwise and that U, U', U'', U''' vanish at $\xi = \pm L$ and, additionally, that U'''' vanishes at $\xi = \pm L$ in the case of the KP(m, n) equation.

Proposition 3.1. Suppose a compacton profile (12) is a weak solution of the K(m, n) equation or the KP(m, n) equation, with the behaviour (31) at the end points of its support. (i) It is a strong solution of the K(m, n) equation, with m, n > 0 and $m \neq 1$, iff p obeys the condition

$$p > 3. \tag{37}$$

(ii) It is a strong solution of the KP(m,n) equation, with m, n > 0 and $m \neq 1$, iff p obeys the condition

$$p > 4. \tag{38}$$

In Ref. [7], less stringent conditions were considered, which arise when the travelling wave ODE is multiplied by powers of U to remove all negative powers. Such solutions need not be classical solutions of the unmodified ODE. The modified ODE, however, does not possess a weak formulation similar to the integral equations (18) and (21). Specifically, there will be additional terms that contain U'^2 and U'^3 and lead to products of Dirac delta distributions.

4. Explicit weak compactons

All weak compactons (12) obtained from the cut off of travelling wave solutions with the various types of profiles (13) will now be summarized for the K(m, n) and KP(m, n) equations. The steps consist of:

(1) substitute the specific form for the travelling wave into the ODE (5);

(2) determine the solution parameters α , β , k, q by solving an overdetermined system of algebraic equations;

(3) extract the cutoff L and the power p in the behaviour (31) near the cutoff;

(4) impose the existence conditions in Theorem 3.1.

4.1. Solution profiles. The preceding steps have been carried out in Maple and yield the following weak compacton solutions, which have a symmetric profile. Hereafter, K denotes the complete elliptic integral (see e.g. Ref. [8]).

Algebraic compactons:

$$u = \left(\frac{k(3n+1)}{2a(n+1)}\right)^{\frac{2}{n-1}} \left(1 - \frac{a^2(n+1)(n-1)^2}{2kbn(3n+1)^2}\xi^2\right)^{\frac{2}{n-1}} H(L - |\xi|), \quad L = \frac{3n+1}{(n-1)|a|} \sqrt{\frac{2n|kb|}{n+1}}, \quad (39)$$
$$m = \frac{1}{2}(n+1), \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = \operatorname{sgn}(b)$$

$$u = \left(\frac{a(n+1)}{2k}\right)^{\frac{1}{n-1}} \left(1 + \frac{k^2(n-1)^2}{abn(n+1)^2}\xi^2\right)^{\frac{1}{n-1}} H(L - |\xi|), \quad L = \frac{(n+1)\sqrt{n|ab|}}{(n-1)|k|}, \quad (40)$$
$$m = 2 - n, \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = -\operatorname{sgn}(b)$$

Cosine compactons:

$$u = \left(\frac{2nk}{(n+1)a}\right)^{\frac{1}{n-1}} \cos\left(\frac{n-1}{2n}\sqrt{\frac{a}{b}}\xi\right)^{\frac{2}{n-1}} H(L-|\xi|), \quad L = \frac{n}{n-1}\sqrt{\frac{b}{a}}\pi,$$

$$m = n, \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = \operatorname{sgn}(b)$$
(41)

$$u = \left(\frac{2a}{k(m+1)}\right)^{\frac{1}{1-m}} \cos\left(\sqrt{\frac{-k}{4b}}(1-m)\xi\right)^{\frac{2}{1-m}} H(L-|\xi|), \quad L = \sqrt{\frac{|b|}{|k|}}(1-m)\pi, \quad (42)$$

$$n = 1, \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = -\operatorname{sgn}(b)$$

Jacobi cn compactons:

$$u = \left(\frac{a(n+1)}{k(3n-1)}\right)^{\frac{1}{2-2n}} \operatorname{cn}\left((1-n)\sqrt{\frac{-a}{nb}}\sqrt[4]{\frac{k}{a(3n-1)(n+1)}}\xi, \frac{1}{\sqrt{2}}\right)^{\frac{2}{1-n}} H(L-|\xi|),$$

$$L = \frac{\sqrt{n|b|}}{1-n}\sqrt[4]{\frac{(3n-1)(n+1)}{ak}} K(\frac{1}{\sqrt{2}}),$$

$$m = 2n - 1, \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = -\operatorname{sgn}(b)$$

$$u = \left(\frac{k(3n-1)}{a(n+1)}\right)^{\frac{1}{2n-2}} \operatorname{cn}\left((n-1)\sqrt{\frac{a}{bn}}\sqrt[4]{\frac{k}{a(3n-1)(n+1)}}\xi, \frac{1}{\sqrt{2}}\right)^{\frac{2}{n-1}} H(L-|\xi|),$$

$$L = \frac{\sqrt{n|b|}}{n-1}\sqrt[4]{\frac{(3n-1)(n+1)}{ak}} K(\frac{1}{\sqrt{2}}),$$

$$m = 2n - 1, \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = \operatorname{sgn}(b)$$

$$(43)$$

Jacobi sn compactons:

$$u = \left(\frac{a(n+1)}{k(3n-1)}\right)^{\frac{1}{2-2n}} \operatorname{sn}\left((1-n)\sqrt{\frac{-a}{2bn}}\sqrt[4]{\frac{k}{a(3n-1)(n+1)}}(\xi+L), i\right)^{\frac{2}{1-n}} H(L-|\xi|),$$

$$L = \frac{\sqrt{2n|b|}}{1-n}\sqrt[4]{\frac{(3n-1)(n+1)}{ak}} K(i),$$

$$m = 2n - 1, \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = -\operatorname{sgn}(b)$$

$$u = \left(\frac{k(3n-1)}{a(n+1)}\right)^{\frac{1}{2n-2}} \operatorname{sn}\left((n-1)\sqrt{\frac{a}{2bn}}\sqrt[4]{\frac{k}{a(3n-1)(n+1)}}(\xi+L), i\right)^{\frac{2}{n-1}} H(L-|\xi|),$$

$$L = \frac{\sqrt{2n|b|}}{n-1}\sqrt[4]{\frac{(3n-1)(n+1)}{ak}} K(i),$$

$$m = 2n - 1, \quad \operatorname{sgn}(k) = \operatorname{sgn}(a) = \operatorname{sgn}(b)$$
(45)

The same steps can be applied to the new type of profiles (15). This yields the following six new compacton solutions. (In particular, for each solution, m differs compared to the previous Jacobi cn and sn solutions.)

rational Jacobi cn compactons:

$$u = \left(\frac{(5+3\sqrt{3})k(2n-1)}{-2a(n+1)}\right)^{\frac{1}{3n-3}} \left(\frac{\operatorname{cn}\left((n-1)\sqrt[6]{\frac{12\sqrt{3}ak^2}{b^3n^3(n+1)^2(2n-1)}}(\xi+L), \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) - 1}{\operatorname{cn}\left((n-1)\sqrt[6]{\frac{12\sqrt{3}ak^2}{b^3n^3(n+1)^2(2n-1)}}(\xi+L), \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) + \sqrt{3} + 2}\right)^{\frac{1}{n-1}} H(L-|\xi|),$$

$$L = \frac{2}{n-1}\sqrt[6]{\frac{b^3n^3(n+1)^2(2n-1)}{12\sqrt{3}ak^2}}} \operatorname{K}\left(\frac{\sqrt{2}(\sqrt{3}-1)}{4}\right),$$

$$m = 3n-2, \quad \operatorname{sgn}(a) = \operatorname{sgn}(b) = -\operatorname{sgn}(k) \tag{47}$$

$$u = \left(\frac{(5+3\sqrt{3})k(2n-1)}{-2a(n+1)}\right)^{\frac{1}{3n-3}} \left(\frac{\operatorname{cn}\left((n-1)\sqrt[6]{\frac{12\sqrt{3}ak^2}{b^3n^3(n+1)^2(2n-1)}}\xi, \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) + 1}{\operatorname{cn}\left((n-1)\sqrt[6]{\frac{12\sqrt{3}ak^2}{b^3n^3(n+1)^2(2n-1)}}\xi, \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) - \sqrt{3} - 2}\right)^{\frac{1}{n-1}} H(L-|\xi|),$$

$$L = \frac{2}{n-1}\sqrt[6]{\frac{b^3n^3(n+1)^2(2n-1)}{12\sqrt{3}ak^2}}} \operatorname{K}\left(\frac{\sqrt{2}(\sqrt{3}-1)}{4}\right),$$

$$m = 3n-2, \quad \operatorname{sgn}(a) = \operatorname{sgn}(b) = -\operatorname{sgn}(k)$$

$$(48)$$

$$u = \left(\frac{(5+3\sqrt{3})a(n+1)}{k(2n-1)}\right)^{\frac{1}{3-3n}} \left(\frac{\operatorname{cn}\left((1-n)\sqrt[6]{\frac{-12\sqrt{3}ak^2}{b^3n^3(n+1)^2(2n-1)}}\xi, \frac{\sqrt{2}(\sqrt{3}+1)}{4}\right) + \sqrt{3} - 2}{\operatorname{cn}\left((1-n)\sqrt[6]{\frac{-12\sqrt{3}ak^2}{b^3n^3(n+1)^2(2n-1)}}\xi, \frac{\sqrt{2}(\sqrt{3}+1)}{4}\right) + 1}\right)^{\frac{1}{1-n}} H(L-|\xi|),$$

$$L = \frac{1}{1-n}\sqrt[6]{\frac{|b|^3n^3(n+1)^2(2n-1)}{12\sqrt{3}|a|k^2}}\operatorname{arccn}\left(\sqrt{3} - 2, \frac{\sqrt{2}(\sqrt{3}+1)}{4}\right),$$

$$m = 3n - 2, \quad \operatorname{sgn}(a) = -\operatorname{sgn}(b) = \operatorname{sgn}(k)$$
(49)

$$u = \left(\frac{(5+3\sqrt{3})a(n+1)}{-2k(5n-1)}\right)^{\frac{2}{3-3n}} \left(\frac{\operatorname{cn}\left((1-n)\sqrt[6]{\frac{-3\sqrt{3}a^2k}{2b^3n^3(n+1)(5n-1)^2}}(\xi+L), \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) - 1}{\operatorname{cn}\left((1-n)\sqrt[6]{\frac{-3\sqrt{3}a^2k}{2b^3n^3(n+1)(5n-1)^2}}(\xi+L), \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) + \sqrt{3} + 2}\right)^{\frac{2}{1-n}} H(L-|\xi|)$$

$$L = \frac{2}{1-n}\sqrt[6]{\frac{2|b|^3n^3(n+1)(5n-1)^2}{3\sqrt{3}a^2|k|}}} \operatorname{K}\left(\frac{\sqrt{2}(\sqrt{3}-1)}{4}\right),$$

$$m = \frac{1}{2}(3n-1), \quad \operatorname{sgn}(a) = -\operatorname{sgn}(b) = \operatorname{sgn}(k) \tag{50}$$

$$u = \left(\frac{(5+3\sqrt{3})a(n+1)}{-2k(5n-1)}\right)^{\frac{2}{3-3n}} \left(\frac{\operatorname{cn}\left((1-n)\sqrt[6]{\frac{-3\sqrt{3}a^2k}{2b^3n^3(n+1)(5n-1)^2}}\xi, \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) + 1}{\operatorname{cn}\left((1-n)\sqrt[6]{\frac{-3\sqrt{3}a^2k}{2b^3n^3(n+1)(5n-1)^2}}\xi, \frac{\sqrt{2}(\sqrt{3}-1)}{4}\right) - \sqrt{3} - 2}\right)^{\frac{2}{1-n}} H(L-|\xi|),$$

$$L = \frac{2}{1-n}\sqrt[6]{\frac{2|b|^3n^3(n+1)(5n-1)^2}{3\sqrt{3}a^2|k|}}} \operatorname{K}\left(\frac{\sqrt{2}(\sqrt{3}-1)}{4}\right),$$

$$m = \frac{1}{2}(3n-1), \quad \operatorname{sgn}(a) = -\operatorname{sgn}(b) = \operatorname{sgn}(k) \tag{51}$$

$$u = \left(\frac{(5+3\sqrt{3})k(5n-1)}{a(n+1)}\right)^{\frac{2}{3n-3}} \left(\frac{\operatorname{cn}\left((n-1)\sqrt[6]{\frac{3\sqrt{3a^2k}}{2b^3n^3(n+1)(5n-1)^2}}\xi, \frac{\sqrt{2}(\sqrt{3}+1)}{4}\right) + \sqrt{3} - 2}{\operatorname{cn}\left((n-1)\sqrt[6]{\frac{3\sqrt{3a^2k}}{2b^3n^3(n+1)(5n-1)^2}}\xi, \frac{\sqrt{2}(\sqrt{3}+1)}{4}\right) + 1}\right)^{\frac{2}{1-n}} H(L-|\xi|),$$

$$L = \frac{1}{n-1}\sqrt[6]{\frac{2|b|^3n^3(n+1)(5n-1)^2}{3\sqrt{3a^2}|k|}} \operatorname{arccn}\left(\sqrt{3} - 2, \frac{\sqrt{2}(\sqrt{3}+1)}{4}\right),$$

$$m = \frac{1}{2}(3n-1), \quad \operatorname{sgn}(a) = \operatorname{sgn}(b) = \operatorname{sgn}(k) \tag{52}$$

There is an sn counterpart for each compacton (47)–(52). 8888

All of these solutions (41)-(46) exist both as strong and weak solutions. As seen in Table 1, the weak existence conditions are less restrictive than the strong existence conditions for the following solutions: (39)-(42), (44), (46), (47), (48), (52); for the other solutions, the weak conditions imply the strong conditions.

Similar results hold for compactons that have an antisymmetric profile: (13c) and the second of (13d) as well as the first of (13e), when p is an odd integer or an odd fraction.

type	profile	p	weak $K(m, n)$	strong $K(m, n)$	weak $KP(m, n)$	strong $KP(m, n)$
algebraic	(39)	$\frac{2}{n-1}$	1 < n	1 < n < 5/3	1 < n < 3	1 < n < 3/2
	(40)	$\frac{1}{n-1}$	1 < n < 2	1 < n < 4/3	1 < n < 2	1 < n < 5/4
cosine	(41)	$\frac{2}{n-1}$	1 < n	1 < n < 5/3	1 < n < 3	1 < n < 3/2
	(42)	$\frac{2}{1-m}$	0 < m < 1	1/3 < m < 1	0 < m < 1	1/2 < m < 1
cn	(43)	$\frac{2}{1-n}$	1/2 < n < 1			
	(44)	$\frac{2}{n-1}$	1 < n	1 < n < 5/3	1 < n < 3	1 < n < 3/2
sn	(45)	$\frac{\frac{2}{1-n}}{\frac{2}{n-1}}$	1/2 < n < 1			
	(46)	$\frac{2}{n-1}$	1 < n	1 < n < 5/3	1 < n < 3	1 < n < 3/2
rational cn	(47), (48), (52)	$\frac{2}{n-1}$	1 < n	$1 < n < \frac{5}{3}$	1 < n	$1 < n < \frac{3}{2}$
	(49)	$\frac{1}{1-n}$	$\frac{2}{3} < n < 1$	$\frac{2}{3} < n < 1$	$\frac{2}{3} < n < 1$	$\frac{3}{4} < n < 1$
	(50), (51)	$\frac{4}{1-n}$	$\frac{1}{3} < n < 1$			

TABLE 1. Existence conditions of the symmetric compactons

4.2. Examples when compactons exist only as weak solutions. The preceding results show that there are specific K(m, n) and KP(m, n) equations such the only explicit compactons of the form (13) are weak solutions.

For example: algebraic type (39), cosine type (41), Jacobi cn and sn types (44) and (46), and rational Jacobi cn type (52) are admitted as weak solutions by both equations in the case n = 2, $\operatorname{sgn}(a) = \operatorname{sgn}(b)$, with m = 3/2, 2, 3, 3, 5/2 respectively, but fail to exist as strong solutions. Moreover, in each of these cases, all of the other types (40), (42), (43) and (45), (47)–(51) are not solutions since they have either $\operatorname{sgn}(a) = -\operatorname{sgn}(b)$ or an opposite relation between $\operatorname{sgn}(a)$ and $\operatorname{sgn}(k)$.

Another example is the second cosine type (42), which has n = 1. It is admitted only as a weak solution by both equations in the case m = 1/4, with sgn(a) = -sgn(b).

4.3. Examples of numerical weak compactons. The existence conditions (29) and (30) for weak compactons enable finding numerical solutions of the respective integral equations (17) and (20). It is most convenient to work in terms of the variable $V(\xi) = U(\xi)^n$, which satisfies

$$V'' = 2(C + (1 + 1/n)BV^{1/n} - (1 + m/n)AV^{m/n})$$
(53)

obtained from differentiation of the first-order ODE (10).

In the case of the K(m, n) equation, existence condition (29) is equivalent to the cutoff conditions

$$V(\pm L) = 0, \quad V'(\pm L) = 0, \quad V''(\pm L) = 0 \tag{54}$$

on solutions $V(\xi)$ of ODE (53) in an interval $-L \leq \xi \leq L$. These conditions together require that C = 0 and hence E = 0, as seen from the ODEs (53) and (10), whereby the first condition then implies that the second and third conditions hold automatically. To obtain a weak compacton solution that has a symmetric profile, it is sufficient to take an initial condition $V(0) = V_0 \neq 0$, V'(0) = 0, along with $\operatorname{sgn}(V''(0)) = -\operatorname{sgn}(V_0)$, in the ODE (53). The initial value V_0 here is determined by the first-order ODE (10), which yields the algebraic relation $BV_0^{1/n} - AV_0^{m/n} = 0$. Thus,

$$V_0 = (B/A)^{n/(m-1)} = \left(\frac{(n+m)k}{(n+1)a}\right)^{n/(m-1)}$$
(55)

from expressions (11) for B and A. The sign condition thereby becomes

$$(1-m)(B^{m-n}/A^{1-n})^{1/(m-1)} = \frac{1-m}{b} \left(\frac{k^{m-n}}{a^{1-n}}\right)^{1/(m-1)} < 0$$
(56)

In addition, the endpoints $\pm L$ are determined by the integral

$$L = \int_{0}^{V_0} \frac{dV}{\sqrt{B(1 - (V/V_0)^{(m-1)/n})V^{1+1/n}}} = \int_{0}^{V_0} \frac{dV}{\sqrt{A((V_0/V)^{m/n} - 1)V^{1+m/n}}}$$
(57)

through integration of ODE (10).

The case of the KP(m, n) equation is similar, with the additional existence condition (30) which is equivalent to the cutoff condition $V'''(\pm L) = 0$. This condition holds as a consequence of the previous conditions (54), as seen from differentiation of the second order ODE (53). Thus, a symmetric weak compaction solution is given by solving the ODE (53) with the same initial condition $V(0) = V_0 \neq 0$, V'(0) = 0, using relations (55), (56), (57).

It is straightforward to obtain and plot the resulting numerical weak compactons expressed in terms of the original variable $U(\xi) = V(\xi)^{1/n}$. Two examples are shown in Fig. 1. They have $m = \frac{1}{4}(5n-1)$ and m = 5n - 4, respectively.




FIGURE 1. Profile of symmetric weak compactons: (left) n = 2, m = 9/4, a = b = k = 1; (right) n = 9/10, m = 1/2, a = -b = k = 1

Explicit symmetric compactons cannot be found for these particular equations, K(9/4, 2) and K(1/2, 9/10), as well as KP(9/4, 2) and KP(1/2, 9/10).

A similar analysis can be applied to obtain numerical weak compactons having an antisymmetric profile.

5. Concluding remarks

Weak solutions provide a general setting for formulating and understanding compactons, such that classical (strong) solutions are a special case. In this setting, there exist weak compactons that have insufficient differentiability to be classical solutions. Additionally, it enables finding numerical compacton solutions with minimal differentiability. New types of explicit compacton profiles have been derived for both the K(m, n) equation (2), which is a generalization of the gKdV equation having nonlinear dispersion, and for its two-dimensional counterpart called the KP(m, n) equation (3), which is an analogous generalization of the gKP equation. These new compactons comprise weak solutions as well as classical solutions. In particular, examples of compactons that exist only as weak solutions are given.

The same method is applicable more generally to other nonlinearly dispersive equations, such as generalizations of the KdV and KP equations with different forms of nonlinearities [11, 12].

Acknowledgments

SCA is supported by an NSERC Discovery grant. MLG gratefully acknowledges support of Junta de Andalucia FQM-201 group.

References

- [1] P. Rosenau, A. Zilburg, Compactons, J. Phys. A: Math. Theor. 51 (2018), 343001.
- [2] P. Rosenau, J.M. Hyman, Compactons: solitons with finite wavelength, Phys. Rev. Lett. 70 (1993), 564-567.
- [3] P. Rosenau, Compactification of patterns by a singular convection or stress, Phys. Rev. Lett. (2007), 234102.
- [4] D.E. Pelinovsky, A.V. Slunyaev, A.V. Kokorina, E.N. Pelinovsky, Stability and interaction of compactons in the sublinear KdV equation, Commun. Nonlin. Sci. Numer. Simulat. 101 (2021), 105855.
- [5] P. Rosenau, Nonlinear dispersion and compacton structure, Phys. Rev. Lett. 73 (1994), 1737–1741.
- [6] B.B. Kadomstev and V.I. Petviashvili, On the stability of waves in weakly dispersive media, Sov. Phys. Dokl. 15 (1970), 539–541.
- [7] S.C. Anco, M.L. Gandarias, Nonlinearly dispersive KP equations with new compacton solutions, Nonlin. Analysis: Real World Appl. 75 (2024), 103964.
- [8] M. Abramowitz, I. Stegun, Handbook of Mathematical Functions, Applied Mathematics Series 55, National Bureau of Standards, 1964.
- [9] L.C. Evans, Partial Differential Equations, Amer. Math. Soc., Providence, 1998.
- [10] I. Gel'fand, G. Shilov, *Generalized functions*, Academic Press, New York, 1964.
- [11] P. Rosenau, A. Oron, On compactons induced by a non convex convection, Commun. Nonlin. Sci. Numer. Sim. 19 (2014), 1329.
- [12] Y.-X. Yu, New compacton solutions and solitary pattern solutions for modified nonlinearly dispersive mK(m,n,a,b) equation, Commun. Theor. Phys. 52 (2009) 637–640.