Euler-Poincaré Reduction with Broken Symmetries on Riemannian Manifolds and Optimal Control Applications*

Jacob R. Goodman † and Leonardo J. Colombo ‡

Abstract. This paper studies the reduction by symmetry of variational problems on Lie groups and Riemannian homogeneous spaces. We derive the reduced equations of motion in the case of Lie groups endowed with a left-invariant metric, and on Lie groups that admits a bi-invariant metric. We repeated this analysis for Riemannian homogeneous spaces, where we derive the reduced equations by considering an alternative variational problem written in terms of a connection on the horizontal bundle of the underlying Lie group. We study also the case that the underlying Lie group admits a bi-invariant metric space. These ideas are applied to geodesics for a rigid body on SO(3) to derive geodesic equations of motion on the unit sphere S^2 , geodesics on S^2 as a Riemannian symmetric space endowed with a bi-invariant metric and optimal control problems for applications to robotic manipulators.

Key words. Euler-Poincaré reduction, Symmetry breaking, Optimal control, Variational methods on Riemannian manifolds.

AMS subject classifications. 68Q25, 68R10, 68U05

1. Introduction. The major drawback of utilizing variationally defined curves in the task of path planning is that they are frequently impossible to compute explicitly, and often computationally expensive to approximate numerically. However, in many applications, the configuration manifold is a Lie group or a Riemannian homogeneous space, which possess special structures that allow to express the equations of motion of a mechanical system as some lower-order ODE evolving on the Lie algebra of the underlying Lie group (together with a reconstruction equation). The Lie algebra, in turn, carries a natural vector space structure, and therefore is well-suited for classical numerical integration schemes.

Dimensionality reduction for large scale systems has become an active problem of interest within the automatic control and robotics communities. In large robotic swarms, guidance and trajectory planning algorithms for coordination while optimizing qualitative features for the swarm of multiple robots are determined by solutions of nonlinear equations which demand a high-computational costs along its integration [6]. The construction of methods for reduction of dimensionality permits fast computations for the generation of optimal trajectories [8].

The Euler-Poincaré equations are a paradigmatic example of reduction of Lagrangian mechanical systems where the configuration space is a Lie group G and the Lagrangian is

^{*}Submitted to the SIAM Journal on Applied Dynamical Systems on December 28, 2023.

Funding: The authors acknowledge financial support from Grant PID2022-137909NB-C21 funded by MCIN/AEI/ 10.13039/501100011033. The project that gave rise to these results received the support of a fellowship from "Ia Caixa" Foundation (ID 100010434). The fellowship code is LCF/BQ/DI19/11730028.

[†]Antonio de Nebrija University, Departamento de Informática, Escuela Politécnica Superior, C. de Sta. Cruz de Marcenado, 27, 28015, Madrid, Spain (jacob.goodman@nebrija.es).

[‡]Centre for Automation and Robotics (CSIC-UPM), Ctra. M300 Campo Real, Km 0,200, Arganda del Rey - 28500 Madrid, Spain (leonardo.colombo@car.upm-csic.es).

G-invariant (see [14, 15, 20]). In this case, it induces a reduced Lagrangian $\ell : \mathfrak{g} \to \mathbb{R}$, where \mathfrak{g} is the Lie algebra of a Lie group *G*. Applying adapted variational techniques, it is possible to derive the equations of motion for ℓ , the Euler-Poincaré equations. This procedure is called Euler-Poincaré reduction (see [15, 17] and references therein). The Euler-Poincaré equations are explicitly written as

(1.1)
$$\frac{d}{dt}\frac{\delta\ell}{\delta\xi} - ad_{\xi}^*\frac{\delta\ell}{\delta\xi} = 0$$

where ad_{ξ}^* is the dual operator of the adjoint endomorphism $ad_{\xi} : \mathfrak{g} \to \mathfrak{g}$ defined via the Lie bracket on \mathfrak{g} by $ad_{\xi}\tilde{\xi} = [\xi, \tilde{\xi}]$ where $\xi, \tilde{\xi} \in \mathfrak{g}$. Typically, it is used the functional derivative notation $\delta \ell / \delta \xi$ since the equations are also valid for infinite dimensional Lie algebras. As particular examples, these equations include the equations for rigid bodies and fluids, but in the latter case, one must use infinite dimensional Lie algebras. In the variational deduction of the Equations (1.1), it is necessary to develop a careful analysis of the admissible infinitesimal variations to deduce the equations. More precisely, the variations are obtained by a reduction procedure of the admissible variations of the unreduced Lagrangian $L : TG \to \mathbb{R}$. Of course, using the corresponding Legendre transformation, it is possible to rewrite Equations (1.1) as the Lie-Poisson equations on \mathfrak{g}^* [17]. Observe that, as an essential feature, Equations (1.1) initially involve half of the degrees of freedom as compared with the usual Euler-Lagrange equations for $L : TG \to \mathbb{R}$ although after solving Equations (1.1) it is necessary to use the reconstruction equations $\dot{g} = g\xi$ from the trivialization of the Lie group to recover the solutions on the Lie group. Moreover, Equations (1.1) are first-order differential equations while the standard Euler-Lagrange equations are second-order ones.

The inclusion of potentials functions in the Lagrangian system under study often complicates the reduction process. The potential typically must admit certain symmetries to allow for reduction. Generally, the potential will not share all the symmetries of the kinetic component of the Lagrangian but instead will be invariant under the action of some isotropy subgroup of the underlying Lie group. Such problems are typically referred to as reduction with symmetry breaking. The problem of reduction with symmetry breaking has been studied in [14, 11] for variational principles of arbitrary order and where the parameter space is an arbitrary smooth manifold, for contact Lagrangian systems in [2], in the context of optimization problems on manifolds in [10], and for optimal control applications in [3], [13], [7], [21]. For homogeneous spaces, reduction with symmetry breaking has only been studied in [22].

The main goal of this paper is to establish a general theory for reduction with symmetry breaking on Lie groups endowed with a left-invariant (respectively, bi-invariant) metric on a Riemannian manifold, and on Riemannian homogeneous spaces; and to apply the construction to optimal control problems of affine-connection control systems on Lie groups with partial symmetry-breaking cost functions. For the optimal control problem, the generated optimal trajectories may be found in a cost-effective manner. This is motivated by the desire to apply these results to the path-planning of robotic systems (rigid body motions and robotic manipulators, for example), where such spaces frequently appear, and cannot be addressed using the methods applied in the literature.

The structure of the paper is as follows. In Section 2 we introduce the basic background on Riemannian manifolds and Momentum Maps. Section 3 describes the geometry of systems

3

on Lie-groups endowed with a left-invariant Riemannian metric. In Section 4, we derive Euler-Poincaré equations for systems endowed with a left-invariant metric and in Section 5 we study the problem of reduction by symmetries for systems with symmetry breaking. In Section 6 we study Euler-Poincaré Reduction with broken symmetries on Riemannian homogeneous space. Finally, in Section 7, we consider an application to optimal control of left-invariant homogeneous systems inspired by path planning for robotic manipulators. Multiple examples are provided throughout the papers.

2. Riemannian Manifolds. Let $(Q, \langle \cdot, \cdot \rangle)$ be an *n*-dimensional Riemannian manifold, where Q is an *n*-dimensional smooth manifold and $\langle \cdot, \cdot \rangle$ is a positive-definite symmetric covariant 2-tensor field called the Riemannian metric. That is, to each point $q \in Q$ we assign a positive-definite inner product $\langle \cdot, \cdot \rangle_q : T_q Q \times T_q Q \to \mathbb{R}$, where $T_q Q$ is the tangent space of Q at q and $\langle \cdot, \cdot \rangle_q$ varies smoothly with respect to q. The length of a tangent vector is determined by its norm, defined by $||v_q|| = \langle v_q, v_q \rangle^{1/2}$ with $v_q \in T_q Q$. For any $p \in Q$, the Riemannian metric induces an invertible map $\cdot^{\flat} : T_p Q \to T_p^* Q$, called the flat map, defined by $X^{\flat}(Y) = \langle X, Y \rangle$ for all $X, Y \in T_p Q$. The inverse map $\cdot^{\sharp} : T_p^* Q \to T_p Q$, called the sharp map, is similarly defined implicitly by the relation $\langle \alpha^{\sharp}, Y \rangle = \alpha(Y)$ for all $\alpha \in T_p^* Q$. Let $C^{\infty}(Q)$ and $\Gamma(TQ)$ denote the spaces of smooth scalar fields and smooth vector fields on Q, respectively. The sharp map provides a map from $C^{\infty}(Q) \to \Gamma(TQ)$ via $\operatorname{grad} f(p) = df_p^{\sharp}$ for all $p \in Q$, where $\operatorname{grad} f$ is called the gradient vector field of $f \in C^{\infty}(Q)$. More generally, given a map $V : Q \times \cdots \times Q \to \mathbb{R}$ (with m copies of Q), we may consider the gradient vector field of V with respect to i^{th} component as $\operatorname{grad}_i V(q_1, \ldots, q_m) = \operatorname{grad} U(q_i)$, where $U(q) = V(q_1, \ldots, q_{i-1}, q, q_{i+1}, \ldots, q_m)$ for all $q, q_1, \ldots, q_m \in Q$ (and similarly for the differential with respect to the i^{th} component).

Vector fields are a special case of smooth sections of vector bundles. In particular, given a vector bundle (E, Q, π) with total space E, base space Q, and projection $\pi : E \to Q$, where E and Q are smooth manifolds, a *smooth section* is a smooth map $X : Q \to E$ such that $\pi \circ X = \mathrm{id}_Q$, the identity function on Q. We similarly denote the space of smooth sections on (E, Q, π) by $\Gamma(E)$. A connection on (E, Q, π) is a map $\nabla : \Gamma(TQ) \times \Gamma(E) \to \Gamma(TQ)$ which is $C^{\infty}(Q)$ -linear in the first argument, \mathbb{R} -linear in the second argument, and satisfies the product rule $\nabla_X(fY) = X(f)Y + f\nabla_X Y$ for all $f \in C^{\infty}(Q)$, $X \in \Gamma(TQ)$, $Y \in \Gamma(E)$. The connection plays a role similar to that of the directional derivative in classical real analysis. The operator ∇_X which assigns to every smooth section Y the vector field $\nabla_X Y$ is called the *covariant derivative* (of Y) with respect to X.

We now specialize our attention to affine connections, which are connections on TQ. Let $q: I \to Q$ be a smooth curve parameterized by $t \in I \subset \mathbb{R}$, and denote the set of smooth vector fields along q by $\Gamma(q)$. Then for any affine connection ∇ on Q, there exists a unique operator $D_t: \Gamma(q) \to \Gamma(q)$ (called the *covariant derivative along* q) which agrees with the covariant derivative $\nabla_{\dot{q}} \tilde{W}$ for any extension \tilde{W} of W to Q. A vector field $X \in \Gamma(q)$ is said to be parallel along q if $D_t X \equiv 0$.

The covariant derivative makes it possible to define a particularly important family of smooth curves on Q called *geodesics*, which are defined as the smooth curves γ satisfying $D_t \dot{\gamma} =$ 0. Moreover, geodesics induce a map $\exp_q : T_q Q \to Q$ called the *exponential map* defined by $\exp_q(v) = \gamma(1)$, where γ is the unique geodesic verifying $\gamma(0) = q$ and $\dot{\gamma}(0) = v$. In particular, \exp_q is a diffeomorphism from some star-shaped neighborhood of $0 \in T_q Q$ to a convex open

This manuscript is for review purposes only.

neighborhood \mathcal{B} (called a goedesically convex neighborhood) of $q \in Q$. It is well-known that the Riemannian metric induces a unique torsion-free and metric compatible connection called the Riemannian connection, or the Levi-Civita connection. Along the remainder of this paper, we will assume that ∇ is the Riemannian connection, unless explicitly said otherwise. For additional information on connections, we refer the reader to [4, 18]. When the covariant derivative D_t corresponds to the Levi-Civita connection, geodesics can also be characterized as the critical points of the length functional $L(\gamma) = \int_0^1 ||\dot{\gamma}|| dt$ among all unit-speed piecewise regular curves $\gamma : [a, b] \to Q$ (that is, where there exists a subdivision of [a, b] such that γ is smooth and satisfies $\dot{\gamma} \neq 0$ on each subdivision). Equivalently, we may characterize geodesics by the critical points of the energy functional $\mathcal{E} = \frac{1}{2} \int_a^b ||\dot{\gamma}||^2 dt$ among all C^1 piecewise smooth curves $\gamma : [a, b] \to Q$ parameterized by arc-length. The length functional induces a metric $d : Q \times Q \to \mathbb{R}$ called the Riemannian distance via $d(p,q) = \inf\{L(\gamma) : \gamma \text{ regular}, \gamma(a) = p, \gamma(b) = q\}$.

If we assume that Q is *complete* (that is, (Q, d) is a complete metric space), then by the Hopf-Rinow theorem, any two points x and y in Q can be connected by a (not necessarily unique) minimal-length geodesic $\gamma_{x,y}$. In this case, the Riemannian distance between x and y can be defined by $d(x,y) = \int_0^1 \left\| \frac{d\gamma_{x,y}}{ds}(s) \right\| ds$. Moreover, if y is contained in a geodesically convex neighborhood of x, we can write the Riemannian distance by means of the Riemannian exponential as $d(x,y) = \|\exp_x^{-1}y\|$.

Given $q_a, q_b \in Q$ and $a < b \in \mathbb{R}$, we will say that a curve $\gamma : [a, b] \to Q$ is admissible if it is of Sobolev class $H^1([a, b], Q)$ (that is, the chart representation of γ is of Sobolev class $H^2([a, b], \mathbb{R}^n)$ for any smooth chart on Q), and satisfies $\gamma(a) = q_a, \gamma(b) = q_b$. The space of admissible curves will be denoted by $\Omega_{q_a,q_b}^{a,b}(Q)$ (though we will frequently drop subscripts, superscripts, and the dependency on the underlying manifold when there is no confusion), and it has the structure of a smooth infinite-dimensional manifold. Its tangent space $T_x\Omega$ consists of all Sobolev class H^1 vector fields along x satisfying X(a) = X(b) = 0. It should be noted that restricting \mathcal{E} to Ω instead of the space of C^1 piecewise smooth curves satisfying the boundary conditions results in the same set of geodesics.

An admissible variation of $\gamma \in \Omega$ is a family of curves $\Gamma : (-\epsilon, \epsilon) \times [a, b] \to Q$ such that:

- 1. $\Gamma(s, \cdot) \in \Omega$ for all $s \in (-\epsilon, \epsilon)$,
- 2. $\Gamma(\cdot, t)$ is smooth for all $t \in [a, b]$.
- 3. $\Gamma(0,t) = \gamma(t)$ for all $t \in [a,b]$.

The variational vector field corresponding to Γ is defined by $\partial_s \Gamma(0, t)$. It can be seen that any $X \in T_{\gamma}\Omega$ is the variational vector field of some admissible variation, and conversely, the variational vector field of any admissible variation belongs to $T_{\gamma}\Omega$. Following the literature, we frequently denote an admissible variation of γ by γ_s , and its corresponding variational vector field by $\delta\gamma$.

Finally, we need an important lemma regarding the commutativity of derivatives of an admissible variation, often referred to as the *Symmetry Lemma*:

5

Lemma 2.1 (Symmetry Lemma). If Γ is an admissible variation of $\gamma \in \Omega$, then

$$(2.1) D_s \partial_t \Gamma(s,t) = D_t \partial_s \Gamma(s,t)$$

for all $(s,t) \in (-\epsilon,\epsilon) \times [a,b]$.

2.1. The momentum map. Let G be a Lie group with Lie algebra \mathfrak{g} , and suppose that G acts on a smooth manifold Q on the left. For any $\xi \in \mathfrak{g}$, we define the *infinitesimal generator* associated to ξ , denoted by ξ_Q , as

(2.2)
$$\xi_Q(q) = \frac{d}{dt}\Big|_{t=0} (\operatorname{Exp}(t\xi)q),$$

for all $q \in Q$, where $\text{Exp} : \mathfrak{g} \to G$ is the Lie exponential map on G.

Informally, we think of the infinitesimal generator as a map $\mathfrak{g} \times Q \to TQ$ which describes the *infinitesimal action* of \mathfrak{g} on Q, despite not being a group action in the usual sense. Moreover, the map $\mathfrak{g} \to \Gamma(TQ)$ which maps $\xi \mapsto \xi_Q$ is linear and satisfies $[\xi, \eta]_{\mathfrak{g}} = -[\xi_Q, \eta_Q]$ for all $\xi, \eta \in \mathfrak{g}$, where $[\cdot, \cdot]_{\mathfrak{g}}$ is the Lie bracket on \mathfrak{g} and $[\cdot, \cdot]$ is the Lie bracket of vector fields. The infinitesimal generator induces another important map, called the *momentum* map $J: T^*Q \to \mathfrak{g}^*$, which is defined implicitly by the relation

(2.3)
$$\langle J(\omega), \xi \rangle_* = \langle \omega, \xi_Q(q) \rangle_*,$$

for all $q \in Q, \omega \in T_q^*Q, \xi \in \mathfrak{g}$. Momentum maps turn out to be a vital concept in the process of reduction by symmetry, as we will see in Section 4. Moreover, they serve as the connection between the symmetry of a manifold (invariance under group actions) and conserved quantities, a result which is often known as *Noether's Theorem* [15].

3. Riemannian Geometry on Lie Groups. Let G be a Lie group with Lie algebra $\mathfrak{g} := T_e G$, where e is the identity element of G. The left-translation map $L : G \times G \to G$ provides a group action of G on itself under the relation $L_g h := gh$ for all $g, h \in G$. Given any inner-product $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$ on \mathfrak{g} , left-translation provides us with a Riemannian metric $\langle \cdot, \cdot \rangle$ on G via the relation:

$$\langle X_g, Y_g \rangle := \left\langle g^{-1} X_g, g^{-1} Y_g \right\rangle_{\mathfrak{q}}$$

for all $g \in G, X_g, Y_g \in T_gG$. Such a Riemannian metric is called *left-invariant*, and it follows immediately that there is a one-to-one correspondence between left-invariant Riemannian metrics on G and inner products on \mathfrak{g} , and that $L_g : G \to G$ is an isometry for all $g \in G$ by construction. Any Lie group equipped with a left-invariant metric is complete as a Riemannian manifold. In the remainder of the section, we assume that G is equipped with a left-invariant Riemannian metric.

In the following L_{g^*} stands for the push-forward of L_g , which is well-defined because $L_g: G \to G$ is a diffeomorphism for all $g \in G$. We call a vector field X on G left-invariant if $L_{g*}X = X$ for all $g \in G$, and we denote the set of all left-invariant vector fields on G by $\mathfrak{X}_L(G)$. It is well-known that the map $\phi: \mathfrak{g} \to \mathfrak{X}_L(G)$ defined by $\phi(\xi)(g) = L_{g*}\xi$ for all $\xi \in \mathfrak{g}, g \in G$ is an isomorphism between vector spaces. This isomorphism allows us to construct an operator $\nabla^{\mathfrak{g}}: \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ defined by:

(3.1)
$$\nabla^{\mathfrak{g}}_{\xi}\eta := \nabla_{\phi(\xi)}\phi(\eta)(e),$$

for all $\xi, \eta \in \mathfrak{g}$, where ∇ is the Levi-Civita connection on G corresponding to the left-invariant Riemannian metric $\langle \cdot, \cdot \rangle$. Although $\nabla^{\mathfrak{g}}$ is not a connection, we shall refer to it as the *Rieman*nian g-connection corresponding to ∇ because of the similar properties that it satisfies (see [13]):

Lemma 3.1. $\nabla^{\mathfrak{g}} : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ is \mathbb{R} -bilinear, and for all $\xi, \eta, \sigma \in \mathfrak{g}$, the following relations hold:

- 1. $\nabla^{\mathfrak{g}}_{\xi}\eta \nabla^{\mathfrak{g}}_{\eta}\xi = [\xi,\eta]_{\mathfrak{g}},$ 2. $\langle \nabla^{\mathfrak{g}}_{\sigma}\xi,\eta \rangle + \langle \xi, \nabla^{\mathfrak{g}}_{\sigma}\eta \rangle = 0.$

Remark 3.2. We may consider the Riemannian \mathfrak{g} -connection as an operator

 $\nabla^{\mathfrak{g}}: C^{\infty}([a,b],\mathfrak{g}) \times C^{\infty}([a,b],\mathfrak{g}) \to C^{\infty}([a,b],\mathfrak{g})$ in a natural way, namely, if $\xi, \eta \in C^{\infty}([a,b],\mathfrak{g})$, we can write $(\nabla_{\xi}^{\mathfrak{g}}\eta)(t) := \nabla_{\xi(t)}^{\mathfrak{g}}\eta(t)$ for all $t \in [a, b]$. With this notation, Lemma 3.1 works identically if we replace $\xi, \eta, \sigma \in \mathfrak{g}$ with $\xi, \eta, \sigma \in C^{\infty}([a, b], \mathfrak{g})$.

We define an operator $\mathrm{ad}^{\dagger} : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ by $\mathrm{ad}_{\mathcal{E}}^{\dagger} \eta = (\mathrm{ad}_{\mathcal{E}}^* \eta^{\flat})^{\sharp}$ for all $\xi, \eta \in \mathfrak{g}$. It follows that ad^* is bilinear, and satisfies $\left\langle \mathrm{ad}_{\xi}^{\dagger}\eta, \sigma \right\rangle = \left\langle \mathrm{ad}_{\xi}^*\eta^{\flat}, \sigma \right\rangle_* = \left\langle \eta^{\flat}, \mathrm{ad}_{\xi}\sigma \right\rangle_* = \left\langle \eta, \mathrm{ad}_{\xi}\sigma \right\rangle$ for all $\xi, \eta, \sigma \in \mathfrak{g}$. Hence, ad^{*} is nothing more than the adjoint of the adjoint action with respect to the Riemannian metric on G. This leads to the following decomposition of $\nabla^{\mathfrak{g}}$ (see Theorem 5.40 of [5], for instance

Lemma 3.3. For all $\xi, \eta \in \mathfrak{g}$ the Riemannian \mathfrak{g} -connection can be expressed as

$$abla_{\xi}^{\mathfrak{g}}\eta = rac{1}{2}\left([\xi,\eta]_{\mathfrak{g}} - \mathrm{ad}_{\xi}^{\dagger}\eta - \mathrm{ad}_{\eta}^{\dagger}\xi
ight).$$

Given a basis $\{A_i\}$ of \mathfrak{g} , we may write any vector field X on G as $X = X^i \phi(A_i)$, where $X^i: G \to \mathbb{R}$, where we have adopted the Einstein sum convention. If X is a vector field along some smooth curve $g: [a, b] \to G$, then we may equivalently write $X = X^i g A_i$, where now $X^i: [a,b] \to \mathbb{R}$ and $gA_i =: L_qA_i$. We denote $\dot{X} = \dot{X}^iA_i$, which may be written in a coordinate-free fashion via $\dot{X}(t) = \frac{d}{dt} \left(L_{g(t)^{-1*}} X(t) \right)$. We now wish to understand how the Levi-Civita connection ∇ along a curve is related to the Riemannian g-connection $\nabla^{\mathfrak{g}}$. This relation is summarized in the following result (see [13]):

Lemma 3.4. Consider a Lie group G with Lie algebra g and left-invariant Levi-Civita connection ∇ . Let $g: [a,b] \to G$ be a smooth curve and X a smooth vector field along g. Then the following relation holds for all $t \in [a, b]$:

(3.2)
$$D_t X(t) = g(t) \left(\dot{X}(t) + \nabla^{\mathfrak{g}}_{\xi} \eta(t) \right).$$

Lemma 3.5. Suppose that $\mathcal{Y}, \mathcal{Z} : [a, b] \to \mathfrak{g}$. Then:

(3.3)
$$\frac{d}{dt} \langle \mathcal{Y}, \mathcal{Z} \rangle = \left\langle \dot{\mathcal{Y}}, \mathcal{Z} \right\rangle + \left\langle \mathcal{Y}, \dot{\mathcal{Z}} \right\rangle$$

Proof. Let $\mathcal{T} := \gamma^{-1} \dot{\gamma}$. From Lemma 3.4, the metric compatibility of ∇ , and the left-invariance of the metric, we have that:

$$\begin{split} \frac{d}{dt} \langle \mathcal{Y}, \mathcal{Z} \rangle &= \frac{d}{dt} \langle \gamma \mathcal{Y}, \gamma \mathcal{Z} \rangle \\ &= \langle \nabla_T \gamma \mathcal{Y}, \gamma \mathcal{Z} \rangle + \langle \gamma \mathcal{Y}, \nabla_T \gamma \mathcal{Z} \rangle \,, \\ &= \left\langle \gamma \left(\dot{\mathcal{Y}} + \nabla^{\mathfrak{g}}_{\mathcal{T}} \mathcal{Y} \right), \gamma \mathcal{Z} \right\rangle + \left\langle \gamma \mathcal{Y}, \gamma \left(\dot{\mathcal{Z}} + \nabla^{\mathfrak{g}}_{\mathcal{T}} \mathcal{Z} \right) \right\rangle , \\ &= \left\langle \dot{\mathcal{Y}} + \nabla^{\mathfrak{g}}_{\mathcal{T}} \mathcal{Y}, \mathcal{Z} \right\rangle + \left\langle \mathcal{Y}, \dot{\mathcal{Z}} + \nabla^{\mathfrak{g}}_{\mathcal{T}} \mathcal{Z} \right\rangle , \\ &= \left\langle \dot{\mathcal{Y}}, \mathcal{Z} \right\rangle + \left\langle \mathcal{Y}, \dot{\mathcal{Z}} \right\rangle + \left\langle \nabla^{\mathfrak{g}}_{\mathcal{T}} \mathcal{Y}, \mathcal{Z} \right\rangle + \left\langle \mathcal{Y}, \nabla^{\mathfrak{g}}_{\mathcal{T}} \mathcal{Z} \right\rangle , \\ &= \left\langle \dot{\mathcal{Y}}, \mathcal{Z} \right\rangle + \left\langle \mathcal{Y}, \dot{\mathcal{Z}} \right\rangle , \\ &= \left\langle \dot{\mathcal{Y}}, \mathcal{Z} \right\rangle + \left\langle \mathcal{Y}, \dot{\mathcal{Z}} \right\rangle , \end{split}$$

where in the last equality we have applied Lemma 3.1 in the context of Remark 3.2.

Remark 3.6. Here we see a stark divergence from [1]. Namely, it is claimed in Proposition 5.3 that $\langle \dot{\mathcal{Y}}, \mathcal{Z} \rangle + \langle \mathcal{Y}, \dot{\mathcal{Z}} \rangle = 0$, implying that $\frac{d}{dt} \langle \mathcal{Y}, \mathcal{Z} \rangle = 0$. Considering the case where $\mathcal{Z} = \mathcal{Y}$, this would then imply that every curve in the Lie algebra has constant length, which cannot be true. Consider for instance a curve $\xi(t)$ of constant length c, and define $\eta(t) = f(t)\xi(t)$ for some smooth non-constant real-valued function f. Then $\langle \eta(t), \eta(t) \rangle = f(t)^2 \langle \xi(t), \xi(t) \rangle = (cf(t))^2$, which is non-constant by assumption.

In anticipation of working with variational principles, we momentarily switch our attention to variations. Consider a curve $g \in \Omega$ and an admissible proper variation $\Gamma : (-\epsilon, \epsilon) \times [a, b] \to G$ of g with variational vector field $\frac{\partial}{\partial s}\Big|_{s=0} \Gamma(s,t) = \delta g(t)$. Denote the vector fields along the main and transverse curves of Γ as $T(s,t) := \frac{\partial}{\partial t} \Gamma(s,t)$ and $S(s,t) = \frac{\partial}{\partial s} \Gamma(s,t)$, respectively. Further denote their pullback (via left-translations) to the Lie algebra \mathfrak{g} by $\mathcal{T}(s,t) := \Gamma(s,t)^{-1}T(s,t)$ and $\mathcal{S}(s,t) := \Gamma(s,t)^{-1}S(s,t)$ for all $(s,t) \in (-\epsilon,\epsilon) \times [a,b]$. Observe the following relations:

$$\begin{split} T(0,t) &= \dot{g}(t), \\ S(0,t) &= \delta g(t), \end{split} \qquad \qquad \mathcal{T}(0,t) &= g(t)^{-1} \dot{g}(t), \\ \mathcal{S}(0,t) &= g(t)^{-1} \delta g(t), \end{split}$$

Consider a basis $\{A_i\}_{i=1}^{\dim(G)}$ of \mathfrak{g} , and write $\mathcal{T} = a^i(s,t)A_i$ and $\mathcal{S} = b^i(s,t)A_i$ for some Sobolev class H^1 coordinate functions a^i, b^i on $(-\epsilon, \epsilon) \times [a, b]$. We denote $\dot{\mathcal{S}} = \frac{\partial b^i}{\partial t}(s,t)A_i$ and $\mathcal{T}' = \frac{\partial a^i}{\partial s}(s,t)A_i$. Then Lemma 3.4 gives the following corollary:

Corollary 3.7. The relationships

(3.4)
$$\nabla_T S(s,t) = \Gamma(s,t) \left(\dot{\mathcal{S}}(s,t) + \nabla^{\mathfrak{g}}_{\mathcal{T}} \mathcal{S}(s,t) \right)$$

(3.5)
$$\nabla_S T(s,t) = \Gamma(s,t) \left(\mathcal{T}'(s,t) + \nabla_S^{\mathfrak{g}} \mathcal{T}(s,t) \right)$$

hold for all $(s,t) \in (-\epsilon,\epsilon) \times [a,b]$ among all proper variations Γ of g, with S,T,S,T as defined above. Moreover, we have that

(3.6)
$$\mathcal{T}' = \dot{\mathcal{S}} + [\mathcal{T}, \mathcal{S}]_{\mathfrak{g}}.$$

Proof. For equation (3.4), we need to only apply Lemma 3.4 with $g(t) = \Gamma(s, t)$ and X(t) = S(s,t) for fixed $s \in (-\epsilon, \epsilon)$. Similarly, equation (3.5) follows with $g(t) = \Gamma(s,t)$ and X(t) = T(s,t) for fixed $t \in [a, b]$. Finally, equation (3.6) follows from (3.4) and (3.5) together with the Symmetry Lemma 2.1.

Remark 3.8. It is stated in lemma 5.2 of [1] that $\mathcal{T}' = \dot{S}$ (so that, in particular, $[\mathcal{T}, \mathcal{S}] \equiv 0$). We provide a counterexample to this assertion now. Consider $G = \operatorname{GL}(2, \mathbb{R})$ with Lie algebra $\mathfrak{gl}(2, \mathbb{R})$. Let $g : [0, 1] \to \operatorname{GL}(2, \mathbb{R})$ be defined by $g(t) = \begin{bmatrix} t & -1 \\ 1 & 0 \end{bmatrix}$, and consider the admissible proper variation $\Gamma(s, t) = \begin{bmatrix} t + f(s, t) & -1 \\ 1 + f(s, t) & 0 \end{bmatrix}$, where f is a smooth function such that f(0, t) = f(s, 0) = f(s, 1) = 0 and 1 + f(s, t) > c > 0 for all $(s, t) \in (-\epsilon, \epsilon) \times [0, 1]$ and some $c \in \mathbb{R}$. It follows that $\Gamma(s, t)^{-1} = \frac{1}{1+f(s, t)} \begin{bmatrix} 0 & 1 \\ -(1+f(s, t)) & t+f(s, t) \end{bmatrix}$ and so:

$$\begin{split} T &= \frac{\partial}{\partial t} \Gamma = \begin{bmatrix} 1+f_t & 0\\ f_t & 0 \end{bmatrix}, \qquad \qquad S &= \frac{\partial}{\partial s} \Gamma = \begin{bmatrix} f_s & 0\\ f_s & 0 \end{bmatrix}, \\ \mathcal{T} &= \frac{1}{1+f} \begin{bmatrix} f_t & 0\\ tf_t - f - (t+1) & 0 \end{bmatrix}, \qquad \qquad \mathcal{S} &= \frac{1}{1+f} \begin{bmatrix} f_s & 0\\ f_s(t-1) & 0 \end{bmatrix}, \end{split}$$

where f_t, f_s denote the partial derivatives of f with respect to t and s, respectively. We show that $[\mathcal{T}, \mathcal{S}] \neq 0$ for appropriate choice in f. The Lie bracket is simply the matrix commutator, so in particular we need to show that $\mathcal{TS} \neq \mathcal{ST}$. This reduces to the condition

(3.7)
$$f_s(tf_t - f - (t+1)) = f_s f_t(t-1)$$

for all $(s,t) \in (\epsilon,\epsilon) \times [0,1]$. We choose f(s,t) = st(t-1) for $s \in (-\frac{1}{2}, \frac{1}{2})$, which clearly satisfies all the constraints imposed on f. It is easy to see that $f(s, \frac{1}{2}) = \frac{1}{4}s$, $f_t(s, \frac{1}{2}) = 0$, and $f_s(s, \frac{1}{2}) = -\frac{1}{4}$. By direct computation, we see that (3.7) is not satisfied at $t = \frac{1}{2}$ for any $s \in (-\frac{1}{2}, \frac{1}{2})$.

3.1. Bi-invariant metrics. Now we wish to discuss another important class of Riemannian metrics on a Lie group, the so-called *bi-invariant* (or Ad-*invariant*) metrics. These are the Riemannian metrics $\langle \cdot, \cdot \rangle$ on G which are both left- and right-invariant. Unlike left- and right-invariant metrics, not every Lie group G admits a bi-invariant metric. The following result from [9] provides necessary and sufficient conditions for the existence of a bi-invariant metric.

Lemma 3.9. A connected Lie group admits a bi-invariant metric if and only if it is isomorphic to the Cartesian product of a compact Lie group and a finite-dimensional vector space. Moreover, such a metric is unique up to scalar multiplication.

Despite this limitation, many important examples of Lie groups satisfy the conditions: In particular, two of our most important model spaces SO(3) and \mathbb{R}^3 . This is because SO(3) is a compact Lie group, and \mathbb{R}^3 is a finite-dimensional vector space. Bi-invariant metrics have many nice properties that greatly simplify calculations in practice. First, it is clear that for all $g \in G$, $X, Y \in T_g G$, we have $\langle X, Y \rangle = \langle gXg^{-1}, gYg^{-1} \rangle = \langle \mathrm{Ad}_g X, \mathrm{Ad}_g Y \rangle$ where $\mathrm{Ad} : G \times \mathfrak{g} \to \mathfrak{g}$ is the adjoint operator (this is why such metrics are also called Ad-invariant). Let $\xi, \eta, \sigma \in \mathfrak{g}$.

9

Then, $\langle \eta, \sigma \rangle = \langle \operatorname{Ad}_{\operatorname{Exp}(t\xi)} \eta, \operatorname{Ad}_{\operatorname{Exp}(t\xi)} \sigma \rangle$. Differentiating at t = 0 and making use of Lemma 3.5, we see that $0 = \langle \mathrm{ad}_{\xi}\eta, \sigma \rangle + \langle \eta, \mathrm{ad}_{\xi}\sigma \rangle$, which implies that $\langle \mathrm{ad}_{\xi}^{\dagger}\eta, \sigma \rangle = \langle -\mathrm{ad}_{\xi}\eta, \sigma \rangle$. Hence, $\mathrm{ad}_{\xi}^{\dagger}\eta = -\mathrm{ad}_{\xi}\eta = [\eta, \xi] \text{ for all } \xi, \eta \in \mathfrak{g}.$

Lemma 3.10. Consider a Lie group G equipped with a bi-invariant metric. Let ∇ be the Levi-Civita connection and $\nabla^{\mathfrak{g}}$ be the corresponding Riemannian g-connection. Then:

$$\begin{split} 1. \ \nabla^{\mathfrak{g}}_{\xi} \eta &= \frac{1}{2} [\xi, \eta], \\ 2. \ R(\xi, \eta) \sigma &= \frac{1}{4} [[\xi, \eta], \sigma], \end{split}$$

for all $\xi, \eta, \sigma \in \mathfrak{a}$.

The Lie exponential map $\operatorname{Exp}: \mathfrak{g} \to G$ is defined for all $\xi \in \mathfrak{g}$ by $\operatorname{Exp}(\xi) = \gamma(1)$, where γ is the unique solution to $\dot{\gamma} = \gamma \xi$ with $\gamma(0) = e$. The Lie exponential map Exp : $\mathfrak{g} \to G$ agrees with the restriction of the Riemannian exponential map $\exp : TG \to G$ to the Lie algebra, where exp is taken with respect to the Levi-Civita connection of a bi-invariant metric. In particular, this implies that geodesics through the identity are just one-parameter subgroups (and vice verse).

Theorem 3.11. Consider a Lie group G equipped with a bi-invariant metric. For all $\xi \in \mathfrak{g}$, we have $\operatorname{Exp}(\xi) = \exp_e(\xi)$, where Exp is the Lie exponential map, and $\exp_e: \mathfrak{g} \to G$ is the Riemannian exponential map at the identity $e \in G$ with respect to the Levi-Civita connection.

Theorem 3.11 turns out to be very useful in many applications on compact matrix Lie groups, due to the simple expression for the Lie exponential map. In cases where we are able to calculate the Logarithmic map $\text{Log}: G \to \mathfrak{g}$ directly (on some neighborhood of the identity element $e \in G$), we may also calculate the Riemannian distance explicitly. In particular, we find that d(e, g) = ||Log(g)|| wherever it is defined.

4. Euler-Poincaré Reduction by a Lie Group Symmetry. Let G be a Lie group equipped with a left-invariant metric $\langle \cdot, \cdot \rangle$. We saw in Section 2 that the critical points of the energy functional $\mathcal{E}: \Omega \to \mathbb{R}$ are geodesics. That is, they are the smooth curves $g \in \Omega$ such that $D_t \dot{g} =$ 0 on the full interval [a, b], where D_t is the covariant derivative along g induced by the Levi-Civita connection of $\langle \cdot, \cdot \rangle$. Observe that from Lemma 3.4, this implies that $g\left(\dot{\xi} + \nabla_{\dot{\xi}}^{\mathfrak{g}}\dot{\xi}\right) \equiv 0$, where $\xi := g^{-1}\dot{g}$. Since left-translation is a diffeomorphism, we obtain the following reduction of geodesics:

Theorem 4.1. Let G be a Lie group equipped with a left-invariant metric, and suppose that $q:[a,b] \rightarrow G$ is a geodesic with respect to the Levi-Civita connection induced by this metric if and only if $\xi := g^{-1}\dot{g}$ satisfies the Euler-Poincaré equations

(4.1)
$$\dot{\xi} + \nabla^{\mathfrak{g}}_{\varepsilon} \xi = 0$$

on [a, b].

Theorem 4.1 tells us that solving $D_t \dot{g} = 0$ on a Lie group with a left-invariant metric is equivalent to solving (4.1) (often called the *reduced geodesic equation*, or referred as the Euler-Poincaré equation for a geodesic) together with the reconstruction equation $\dot{g} = g\xi$. Hence, we have replaced a second-order equation with two first-order equations. In mechanical applications, these equations represent the kinematics (the reconstruction equation) and the

This manuscript is for review purposes only.

dynamics (the reduced geodesic equation) of some mechanical system. Moreover, (4.1) evolves on the Lie algebra \mathfrak{g} , so that we have removed the non-linearity of the manifold from our dynamics. This allows for the use of classical numerical integration schemes in approximating solutions, which can only be done for equations evolving on non-linear spaces by using local coordinate systems (which often lead to poor behavior, especially when dealing with chart transitions). In all, the process of converting some ODE/variational principle evolving on a Lie group G into an equation evolving on \mathfrak{g} (together with a reconstruction equation) is known as reduction by symmetry.

Remark 4.2. Reduction by symmetry may be framed more generally in the language of Lagrangian mechanics. That is, given a function $L: TG \to \mathbb{R}$ called the Lagrangian, we seek to minimize the action $\mathcal{A}[g] = \int_{a}^{b} L(g, \dot{g}) dt$ on Ω . It is well-known that the critical points of \mathcal{A} satisfy the Euler-Lagrange equations $\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{g}}\right) + \frac{\partial L}{\partial g} = 0$. If the Lagrangian is left-invariant, meaning that $L(hg, h\dot{g}) = L(g, \dot{g})$ for all $g, h \in G$, then we may consider the reduced Lagrangian $l : \mathfrak{g} \to \mathbb{R}$ given by $l(\xi) := L(e, \xi) = L(g, \dot{g})$, where $\xi := g^{-1}\dot{g}$. Then, the critical points of the reduced action $\mathcal{A}_{red}[\xi] = \int_a^b l(\xi)dt$ among variations of the form $\delta\xi = \dot{\eta} + [\xi, \eta]$ (note that this is just (3.6)), where η is arbitrary admissible proper variation, satisfy the Euler-Poincaré equations $\frac{d}{dt}\left(\frac{\partial l}{\partial \xi}\right) = \mathrm{ad}_{\xi}^* \frac{\partial l}{\partial \xi}$. Moreover, it can be seen that g satisfies the Euler-Lagrange equations corresponding to L if and only if $\xi := g^{-1}\dot{g}$ satisfies the Euler-Poincaré equations corresponding to l [14]. The equation $\dot{g} = g\xi$ is known as the reconstruction equation. The reduced variational principle is an example of a *constrained* variational principle, since we are constraining the set of admissible variations.

The energy function \mathcal{E} is precisely the action corresponding to the Lagrangian $L(g, \dot{g}) =$ $\frac{1}{2}\|\dot{g}\|^2$. Since the Riemannian metric is left-invariant, it follows that the Lagrangian too is left-invariant, and so the reduced Lagrangian takes the form $l(\xi) = \frac{1}{2} \|\xi\|^2$. It is straightforward to show that $\frac{\partial l}{\partial \xi} = \xi^{\flat}$, so that the Euler-Poincaré equations associated to l are given by $\dot{\xi}^{\flat} = \mathrm{ad}_{\epsilon}^* \xi^{\flat}$, which is equivalent to (4.1) by Lemma 3.3. Notice that the Euler-Poincaré equations corresponding to l naturally live on the dual of the Lie algebra \mathfrak{g}^* . It is only through the metric that we are able to convert them into equations on \mathfrak{g} .

4.1. Example: Geodesics for Rigid Body on SO(3). It is well-known that the motion of a rigid body can be modeled on G = SO(3) equipped with the left-invariant Riemannian metric $\langle \dot{R}_1, \dot{R}_2 \rangle = \operatorname{tr}(\dot{R}_1 \mathbb{M} \dot{R}_2^T)$ for all $R \in \operatorname{SO}(3), \dot{R}_1, \dot{R}_2 \in T_R \operatorname{SO}(3)$, where \mathbb{M} is a symmetric positive-definite 3×3 matrix called the *coefficient of inertia matrix*. On the Lie algebra $\mathfrak{so}(3)$, the metric takes the form of the inner-product $\langle \hat{\Omega}_1, \hat{\Omega}_2 \rangle = \Omega_1^T \mathbb{J}\Omega_2$, where $\hat{\cdot} : \mathbb{R}^3 \to \mathfrak{so}(3)$ is the hat isomorphism defined by $(x_1, x_2, x_3)^T \mapsto \begin{bmatrix} 0 & -x_3 & x_2 \\ x_3 & 0 & -x_1 \\ -x_2 & x_1 & 0 \end{bmatrix}$, and \mathbb{J} is a symmetric

positive-definite 3×3 matrix called the moment of inertia

Suppose that G is equipped with the Levi-Civita connection induced by the above metric, and suppose that $R : [a,b] \to \mathrm{SO}(3)$ is a geodesic. Define $\hat{\Omega} := R^{-1}\dot{R} \in \mathfrak{so}(3)$. Then, from Theorem 4.1, we have $\hat{\Omega} = \mathrm{ad}_{\hat{\Omega}}^{\dagger}\hat{\Omega}$. First, observe that $\langle \hat{\Omega}, \hat{\eta} \rangle = \Omega^T \mathbb{J}\eta = (\mathbb{J}\Omega) \cdot \eta$ for all $\hat{\Omega}, \hat{\eta} \in \mathfrak{so}(3)$. Hence, we may identify $\hat{\Omega} \in \mathfrak{so}(3)$ with $\mathbb{J}\Omega \in \mathbb{R}^3$ under the hat isomorphism. Moreover,

$$\begin{split} \left\langle \mathrm{ad}_{\hat{\Omega}}^{\dagger}\hat{\sigma},\hat{\eta}\right\rangle &= \left\langle \hat{\sigma},\mathrm{ad}_{\hat{\Omega}}\hat{\eta}\right\rangle = \left\langle \hat{\sigma},\widehat{\Omega\times\eta}\right\rangle \\ &= \sigma^{T}\mathbb{J}(\Omega\times\eta) = (\mathbb{J}\sigma)\cdot(\Omega\times\eta) = (\mathbb{J}\sigma\times\Omega)\cdot\eta, \end{split}$$

where "×" refers to the vector cross product of $\Omega \in \mathbb{R}^3$ with $\eta \in \mathbb{R}^3$. Hence, we may identify $\operatorname{ad}_{\hat{\Omega}}^{\dagger} \hat{\sigma} \in \mathfrak{so}(3)$ with $\mathbb{J}\sigma \times \Omega \in \mathbb{R}^3$. It then follows that (4.1) is equivalent to $\mathbb{J}\dot{\Omega} = \mathbb{J}\Omega \times \Omega$, which is recognized as Euler's equation for a rigid body.

In the case that $\mathbb{M} = \mathbb{J} = I$, the metric is in fact bi-invariant (and is the unique biinvariant metric on SO(3) up to scalar multiplication by Lemma 3.9). In this case, we have $\operatorname{Exp} : \mathfrak{so}(3) \to \operatorname{SO}(3)$ is diffeomorphism between $\{\hat{\Omega} : \Omega \in \mathbb{R}^3, \ \Omega^T \Omega \leq \pi^2\}$ and $\{R \in \operatorname{SO}(3) : \operatorname{tr}(R) \neq -1\}$, and the logarithmic map is given by

(4.2)
$$\operatorname{Log}(R) = \begin{cases} 0, & R = I\\ \frac{\phi(R)}{\sin(\phi(R))}(R - R^T), & R \neq I \end{cases}$$

where $\phi(R) := \arccos(\frac{1}{2}(\operatorname{tr}(R)-1))$. Moreover, we have $\|\operatorname{Log}(R)\| = \phi(R)$ (see [5], Proposition 5.7 for more details).

5. Euler-Poincaré Reduction with Broken Symmetry. Consider the action functional $\mathcal{A}: \Omega \to \mathbb{R}$ given by

(5.1)
$$\mathcal{A}[g] = \int_{a}^{b} \left(\frac{1}{2} \|\dot{g}(t)\| - V(g(t))\right) dt,$$

where $V: G \to \mathbb{R}$ is a smooth scalar field called the *potential*. Note that the potential may be a potential energy corresponding to a physical system, such as the heavy top (see Example 5.2) or an *artificial potential* used to modify the behavior of the resulting minimizers, such as in the case of the obstacle avoidance problem (see Example 5.3). It is easy to see from Remark 4.2 that the critical points of (5.1) are the smooth solutions to the Euler-Lagrange equations

(5.2)
$$D_t \dot{g}(t) = -\text{grad}V(g(t)),$$

for all $t \in [a, b]$. In general, the process of reduction by symmetry used in Theorem 4.1 for the energy functional (that is, \mathcal{A} with $V \equiv 0$) will only apply to \mathcal{A} when V is invariant under left-translations. However, this is seldom the case in applications, at which point the potential is said to "break the symmetry" of the energy functional. Rather, it is more common that V possesses a *partial symmetry*.

Definition 5.1. We say that the potential $V : G \to \mathbb{R}$ admits a partial symmetry over some smooth manifold M if there exists a left-action $\Psi : G \times M \to M$ and a function $V_{\text{ext}} : G \times M \to \mathbb{R}$ called the extended potential such that:

- 1. For some $\alpha_0 \in M$ and for all $g \in G$, we have $V_{\text{ext}}(g, \alpha_0) = V(g)$.
- 2. V_{ext} is invariant under the left-action $\chi : G \times (G \times M) \to G \times M$ given by $\chi_h(g, \alpha) = (hg, \Psi_h(\alpha))$ for all $g, h \in G, \alpha \in M$.

Before stating the main theorem, we provide a useful identity in applications.

Lemma 5.2. Let G be a Lie group equipped with a left-invariant metric, M a smooth manifold, and $\Psi: G \times M \to M$ a left-action. If $f: G \times M \to \mathbb{R}$ is a smooth function such that $f(hg, \Psi_h(p)) = f(g, p)$ for all $h, g \in G, p \in M$, then:

(5.3)
$$d_1 f(e, p) = -J(d_2 f(e, p))$$

for all $p \in M$, where $J: T^*M \to \mathfrak{g}^*$ is the momentum map associated to the Ψ .

Proof. Let $\xi \in \mathfrak{g}$. Then, on one hand,

$$\frac{d}{dt}\Big|_{t=0} f(\operatorname{Exp}(t\xi), p) = \langle d_1 f(e, p), \xi \rangle_*$$

On the other hand, $f(\operatorname{Exp}(t\xi), p) = f(e, \Psi_{\operatorname{Exp}(-t\xi)}(p))$, so that

$$\frac{d}{dt}\Big|_{t=0} f(\operatorname{Exp}(t\xi), p) = \langle d_2 f(e, p), \frac{d}{dt}\Big|_{t=0} \Psi_{\operatorname{Exp}(-t\xi)}(p) \rangle_*$$
$$= \langle d_2 f(e, p), -\xi_M(p) \rangle_*$$
$$= \langle -J(d_2 f(e, p)), \xi \rangle_*$$

Note that the previous lemma also allows relating the gradients of such a function f. In particular,

(5.4)
$$\operatorname{grad}_1 f(e,p) = -J(\operatorname{grad}_2 f(e,p)^{\flat})^{\sharp}$$

Theorem 5.3. Let G be a Lie group equipped with a left-invariant metric, and suppose that $g:[a,b] \to G$ is a critical point of the action function defined in (5.1). Then g is smooth, and if the potential admits a partial symmetry over some smooth manifold M, then $\xi := g^{-1}\dot{g}$ and $\alpha := \Psi_{g^{-1}}(\alpha_0)$ satisfy

(5.5)
$$\dot{\xi} + \nabla^{\mathfrak{g}}_{\xi} \xi = -\operatorname{grad}_{1} V_{\operatorname{ext}}(e, \alpha),$$

(5.6)
$$\dot{\alpha} = -\xi_M(\alpha),$$

Proof. Suppose that g_s is an admissible variation of g with $\eta := g^{-1} \delta g = g^{-1} \frac{\partial g_s}{\partial s}|_{s=0}$. Then, it must be the case that

$$0 = \frac{\partial}{\partial s}\Big|_{s=0} \int_{a}^{b} \left(\frac{1}{2} \|\dot{g}_{s}\|^{2} - V(g_{s})\right) dt = \int_{a}^{b} \left(\langle D_{s}\dot{g}_{s}, \dot{g}_{s}\rangle - \frac{\partial}{\partial s}V(g_{s})\right)\Big|_{s=0} dt.$$

From the proof of Theorem 4.1, it follows that

$$\int_{a}^{b} \left\langle D_{s} \dot{g}_{s}, \dot{g}_{s} \right\rangle \Big|_{s=0} dt = \int_{a}^{b} \left\langle D_{t} \delta g, \dot{g} \right\rangle dt.$$

This manuscript is for review purposes only.

We now handle the term involving the potential.

$$\begin{split} \int_{a}^{b} \frac{\partial}{\partial s} V(g_{s}) \Big|_{s=0} dt &= \int_{a}^{b} \frac{\partial}{\partial s} V_{\text{ext}}(g_{s}, \alpha_{0}) \Big|_{s=0} dt \\ &= \int_{a}^{b} \frac{\partial}{\partial s} V_{\text{ext}}(g^{-1}g_{s}, \alpha) \Big|_{s=0} dt \\ &= \int_{a}^{b} \langle \text{grad}_{1} V_{\text{ext}}(e, \alpha), \eta \rangle \, dt, \end{split}$$

Hence,

$$\delta \mathcal{A}[g] = \int_{a}^{b} \langle D_t \delta g, \dot{g} \rangle - \langle L_{g^*} \operatorname{grad}_1 V_{\operatorname{ext}}(e, \alpha), \delta g \rangle \, dt = 0.$$

Observe that g([a, b]) is bounded by the continuity of g. Since G is complete, there exists some $r \in \mathbb{R}$ and $g_0 \in G$ such that g([a, b]) is contained within the closed geodesic ball of radius r centered at g_0 , denoted by $\bar{B}_r(g_0)$. It is known that the space $C_0^{\infty}([a, b], \bar{B}_r(g_0))$ of smooth functions with compact support in $\bar{B}_r(g_0)$ is dense in $H^1([a, b], B_r(g_0))$. Hence, there exists a sequence $(g_n) \subset C_0^{\infty}([a, b], \bar{B}_r(g_0))$ of smooth curves such that $g_n \to g$ uniformly with respect to the H^1 norm. Repeating the above analysis for the sequence g_n , and integrating by parts, we then find that

$$\delta \mathcal{A}[g_n] = -\int_a^b \left\langle D_t \dot{g}_n + L_{g_n^*} \operatorname{grad}_1 V_{\operatorname{ext}}(e, \alpha_n), \delta g_n \right\rangle dt \to 0$$

Taking $\delta g_n(t) = f(t)D_t \dot{g}_n$ for some non-zero $f:[a,b] \to \mathbb{R}$ such that $f(t) \ge 0$ for all $t \in [a,b]$ and f(a) = f(b) = 0, it is easily seen by the fundamental lemma of the calculus of variations that $D_t \dot{g}_n \to 0$. Therefore, $g_n \to g$ with respect to the H^2 -norm by uniqueness of limits. Hence, g is of H^2 regularity, from which equation (5.5) follows immediately (as does smoothness upon expanding in local coordinates and applying Gronwall's inequality). Equation (5.6) is a direct consequence of the definition $\alpha = \Psi_{q^{-1}}(\alpha_0)$.

We now discuss some important special cases of Theorem 5.3.

5.1. Euler-Poincaré reduction with advected parameters. Consider Theorem 5.3 in the case where the parameter manifold M = W is a finite-dimensional real vector space on which G acts on the left by linear transformations (that is, $\Psi_g : W \to W$ is linear for all $g \in G$). We implicitly define a left-action $\Theta : G \times W^* \to W^*$ on the dual space via $\langle \Theta_g(w), x \rangle_* = \langle w, \Psi_{g^{-1}}(x) \rangle_*$ for all $w \in W^*, x \in W$. Consider the map $\rho_x : \mathfrak{g} \to W$ given by $\rho_x(\xi) = \xi_W(x)$, where ξ_W is the infinitesimal generator associated to Ψ . Then the dual map $\rho_x^* : W^* \to \mathfrak{g}^*$ satisfies $\langle \rho_x^*(w), \xi \rangle_* = \langle w, \rho_x(\xi) \rangle_*$ for all $x \in W, w \in W^*, \xi \in \mathfrak{g}$. Hence, through the identification $T^*W = W \times W^*$, we see that the momentum map $J : W \times W^* \to \mathfrak{g}^*$ is given by $J(x, w) = \rho_x^*(w)$. Another common notation used here is the diamond operator $\diamond : W \times W^* \to \mathfrak{g}^*$ defined by $x \diamond w = \rho_x^*(w)$ for all $x \in W, w \in W^*$. In all, in the case that V

admits a partial symmetry over W, and making use of Lemma 5.2, equations (5.5)-(5.6) take the form

(5.7)
$$\dot{\xi} + \nabla^{\mathfrak{g}}_{\mathcal{E}} \xi = \rho^*_a (d_2 V_{\text{ext}}(e,\alpha))^{\sharp},$$

(5.8)
$$\dot{\alpha} = -\rho_{\alpha}(\xi)$$

5.2. Example: The heavy top. Here we consider the classical example of a heavy top as an application of our formalism. The heavy top is a rigid body with coefficient of inertia matrix I which rotates about some fixed point (called the pivot), and which is under a constant gravitational field (see [17, 15]). We consider two frames: a fixed body frame which is attached to the heavy top at some fixed point (in such a frame, the heavy top is seen as a stationary object), and an inertial frame taken with respect to some fixed point of the ambient space. We let $v_0 \in \mathbb{R}^3$ denote the vector pointing from the pivot to the center of mass of the heavy top in the body frame. We may now model the attitude of the heavy top at any point in time as $Rv_0 \in \mathbb{R}^3$, where $R \in SO(3)$ is the rotation matrix which moves between the body and inertial frames.

It can be seen that the rotational kinetic energy of the system is then given by $K = \frac{1}{2} \operatorname{tr}(\dot{R} \mathbb{I} \dot{R}^T)$, which we recognize a scalar multiple of the left-invariant Riemannian metric on SO(3) described in Example 4.1. The potential energy is given by $V(R) = mgRv_0 \cdot e_3$, where $m, g \in \mathbb{R}$ denote the mass of the heavy top and acceleration due to gravity, respectively, $e_3 \in \mathbb{R}^3$ is the unit vector pointing vertically upwards in the inertial frame, and $Rv_0 \cdot e_3$ is just the standard 'dot product' of vectors in \mathbb{R}^3 .

The dynamics for the system can then be found by minimizing the action functional

$$\mathcal{A}[g] = \int_{a}^{b} L(g, \dot{g}) dt$$

over some appropriate space of admissible curves, where $L(g, \dot{g}) = K(g, \dot{g}) - V(g)$ is the Lagrangian of the system. Choosing the admissible curves to be Ω (as defined in Section 2), we then see that this minimization problem corresponds to that given in (5.1), where the parameter manifold is $M = \mathbb{R}^3$. We consider the left-action Ψ : SO(3) × $\mathbb{R}^3 \to \mathbb{R}^3$ defined by $\Psi(R, \alpha) = R\alpha$ for all $R \in SO(3), \alpha \in \mathbb{R}^3$, from which we derive the left-action $\chi : SO(3) \times (SO(3) \times \mathbb{R}^3) \to SO(3) \times \mathbb{R}^3$ given by $\chi_{R'}(R, \alpha) = (R'R, R'\alpha)$. It is clear that the extended potential

$$V_{\text{ext}}(R,\alpha) = mgRv_0 \cdot \alpha$$

is then SO(3)-invariant with respect to χ , and satisfies $V_{\text{ext}}(R, e_3) = V(R)$ for all $R \in \text{SO}(3)$. In particular, this is exactly the case studied in Example 5.1, so that the reduced dynamics are given by (5.7)-(5.8) with our choice in V_{ext} .

It is easily seen that $d_2V_{\text{ext}}(I,\alpha) = mgv_0$ for all $\alpha \in \mathbb{R}^3$. Now, by definition,

$$\rho_{\alpha}(\hat{\Omega}) = \frac{d}{dt} \Big|_{t=0} \Psi_{\text{Exp}(t\hat{\Omega})}(\alpha) = \frac{d}{dt} \Big|_{t=0} \text{Exp}(t\hat{\Omega})\alpha = \hat{\Omega}\alpha = \Omega \times \alpha$$

for all $\Omega, \alpha \in \mathbb{R}^3$. Moreover, for all $\Pi \in \mathbb{R}^3$,

$$\langle \rho_{\alpha}^{*}(\Pi), \hat{\Omega} \rangle = \langle \Pi, \rho_{\alpha}(\hat{\Omega}) \rangle = \Pi \cdot (\Omega \times \alpha) = (\alpha \times \Pi) \cdot \Omega = \langle \alpha \times \Pi, \hat{\Omega} \rangle,$$

from which we find that $\rho_{\alpha}^{*}(\Pi) = \alpha \times \Pi$. In particular, $\rho_{\alpha}^{*}(d_2 V_{\text{ext}}(I, \alpha)) = mg \alpha \times v_0$. For any $\Pi, \Omega \in \mathbb{R}^3$,

$$\Pi \cdot \Omega = \langle \breve{\Pi}, \hat{\Omega} \rangle_* = \langle \breve{\Pi}^{\sharp}, \hat{\Omega} \rangle = \mathbb{J} \left(\breve{\Pi}^{\sharp} \right)^{\vee} \cdot \Omega,$$

from which it follows that $\check{\Pi}^{\sharp} = \widehat{\mathbb{J}^{-1}\Pi}$. Hence,

$$\rho_{\alpha}^{*}(d_{2}V_{\text{ext}}(I,\alpha))^{\sharp} = mg\mathbb{J}^{-1}(\alpha \times v_{0}).$$

Together with the analysis done in Example 4.1, it follows that the reduced dynamics for the heavy top are given by

$$\begin{split} \mathbb{J}\hat{\Omega} &= \mathbb{J}\Omega \times \Omega + mg(\alpha \times v_0), \\ \dot{\alpha} &= \alpha \times \Omega. \end{split}$$

5.3. Example: Obstacle avoidance on Lie groups. Suppose G is a Lie group equipped with a left-invariant metric, $g_0 \in G$ is some point-obstacle, and that we wish to find near-geodesic trajectories $g : [a,b] \to G$ satisfying the boundary conditions $g(a) = g_a$ and $g(b) = g_b$ while avoiding g_0 (or rather, some ball $B_r(g_0)$ centered at g_0). We choose an avoidance family of artificial potentials of the form:

(5.9)
$$V(g) = f(d^2(g, g_0); \tau, k, D)$$

where $d: G \times G \to \mathbb{R}$ is the Riemannian distance function on G, τ, k, D are some non-negative parameters, and $f: \mathbb{R} \to \mathbb{R}$ satisfies the following properties:

1. f is smooth and non-negative.

- 2. $f(0; \tau, k, D) = \tau$ is the unique global maximum of f.
- 3. $\lim_{k\to\infty} f(x;\tau,k,D) = 0$ for all |x| > D.
- 4. $\lim_{k \to \infty} f(x; \tau, D) = \tau$ for all |x| < D.

Following the results of [12], it can be shown that for any D > R, and τ, k sufficiently large, the corresponding minimizers of \mathcal{A} avoid $B_R(g_0)$ provided that $g_a \notin B_D(g_0)$ and $g_b \notin B_D(g_0)$. One example of such an avoidance family of artificial potentials is given by

(5.10)
$$V(g) = \begin{cases} e\tau \exp\left(-\frac{1}{1 - (d(g,g_0)/D)^{2k}}\right), & d(g,g_0) < D\\ 0, & \text{otherwise} \end{cases}$$

From which we see that the potential only "engages" when the corresponding minimizer enters the ball of radius D around g_0 . In other words, the minimizers behave as geodesics when far away from the obstacles ("far away" being a concept controlled by the parameter D), and prioritize the obstacle-avoidance task when sufficiently close to g_0 .

Since the Riemannian distance corresponding to a left-invariant Riemannian metric is itself left-invariant (that is, left-translation is an isometry), it follows that the extended artificial potential given by

(5.11)
$$V_{\text{ext}}(g,h) = f(d^2(g,h);\tau,k,D)$$

is left-invariant under left-translation on $G \times G$, and satisfies $V_{\text{ext}}(g, g_0) = V(g)$ for all $g \in G$. In other words, V admits a partial symmetry over M = G. If we suppose that g is contained within a geodesically convex neighborhood of g_0 , then it can be shown that

$$\operatorname{grad}_1 V_{\operatorname{ext}}(g,h) = -2f'(\|\exp_g^{-1}(h)\|^2) \exp_g^{-1}(h).$$

Moreover, the infinitesimal generator corresponding left-translation is simply the push-forward of right-translation. Hence, equations (5.5)-(5.6) take the form

(5.12)
$$\dot{\xi} + \nabla^{\mathfrak{g}}_{\xi} \xi = 2f'(\|\exp_e^{-1}(\alpha)\|^2) \exp_e^{-1}(\alpha),$$

(5.13)
$$\dot{\alpha} = -\xi\alpha.$$

where $\xi = g^{-1}\dot{g}$, and $\alpha = g^{-1}g_0$.

Observe that, in the case where G is equipped with a bi-invariant metric, (5.12) takes the form:

$$\dot{\xi} + \nabla^{\mathfrak{g}}_{\xi} \xi = 2f'(\|\operatorname{Log}(\alpha)\|^2)\operatorname{Log}(\alpha)$$

Using (4.2), in the special case that G = SO(3) is equipped with the bi-invariant metric described in Example 4.1, we find that the critical points of \mathcal{A} are given by

(5.14)
$$\dot{\Omega} = \mathbb{J}\Omega \times \Omega + \frac{2f'(\phi(H)^2)\phi(H)}{\sin(\phi(H))}(H - H^T),$$

(5.15)
$$H = -\hat{\Omega}H.$$

where $H := R^T R_0$ for the point obstacle $R_0 \in SO(3)$, and $\Omega := R^T \dot{R}$.

6. Euler-Poincaré Reduction with Broken Symmetries on Riemannian Homogeneous Space. We now discuss spaces which are not necessarily Lie groups themselves, but nonetheless possess certain symmetries and invariances that allow for the similar results to be obtained: *Homogeneous spaces*. Let G be a connected Lie group. A homogeneous space H of G is a smooth manifold on which G acts transitively. Any Lie group is itself a homogeneous space, where the transitive action is given by left-translation (or right-translation).

Suppose that $\Psi: G \times H \to H$ is a transitive left-action, which we denote by $gx := \Psi_g(x)$. It can be shown that for any $x \in H$, we have $G/\operatorname{Stab}(x) \cong H$ as differentiable manifolds, where $\operatorname{Stab}(x) := \{g \in G \mid gx = x\}$ denotes the *stabilizer subgroup* (also called the *isotropy subgroup*) of x, and $G/\operatorname{Stab}(x)$ denotes the space of equivalence classes determined by the equivalence relation $g \sim h$ if and only if $gh^{-1} \in \operatorname{Stab}(x)$ (or, equivalently, the set of left-cosets of $\operatorname{Stab}(x)$). Moreover, for any closed Lie subgroup $K \subset G$, the left-action $\Phi: G \times G/K \to G/K$ satisfying $\Phi_g([h]) = [gh]$ for all $g, h \in G$ is transitive, and so G/K is a homogeneous space. Hence, we may assume without loss of generality that H := G/K is a homogeneous space of G for some closed Lie subgroup K.

Let $\pi : G \to H$ be the canonical projection map. We define the vertical subspace at $g \in G$ by $\operatorname{Ver}_g := \operatorname{ker}(\pi_*|_g)$, from which we may construct the vertical bundle as $VG := \bigcup_{g \in G} \{g\} \times \operatorname{Ver}_g$. Given a Riemannian metric $\langle \cdot, \cdot \rangle_G$ on G, we define the horizontal subspace

at any point $g \in G$ (with respect to $\langle \cdot, \cdot \rangle_G$) as the orthogonal complement of Ver_g . That is, Hor $_g := \operatorname{Ver}_g^{\perp}$. Similarly, we define the horizontal bundle as $HG := \bigsqcup_{g \in G} \{g\} \times \operatorname{Hor}_g$. Both the vertical and horizontal bundles are vector bundles, and are in fact subbundles of the tangent bundle TG. It is clear that $T_gG = \operatorname{Ver}_g \oplus \operatorname{Hor}_g$ for all $g \in G$, so that the Lie algebra \mathfrak{g} of Gadmits the decomposition $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{h}$, where \mathfrak{s} is the Lie algebra of K and $\mathfrak{h} \cong T_{\pi(e)}H$. We denote the orthogonal projections onto the vertical and horizontal subspaces by \mathcal{V} and \mathcal{H} .

A section $Z \in \Gamma(HG)$ is called a *horizontal vector field*. That is, $Z \in \Gamma(TG)$ and $Z(g) \in$ Hor_g for all $g \in G$. A vector field $Y \in \Gamma(TG)$ is said to be π -related to some $X \in \Gamma(TH)$ if $\pi_*Y_g = X_{\pi(g)}$ for all $g \in G$. If in addition $Y \in \Gamma(HG)$, we say that Y is a *horizontal lift* of X. We further define a horizontal lift of a smooth curve $q : [a, b] \to H$ as a smooth curve $\tilde{q} : [a, b] \to G$ such that $\pi \circ \tilde{q} = q$ and $\dot{\tilde{q}}(t)$ is horizontal for all $t \in [a, b]$. We have the following result from [16]:

Lemma 6.1. Let H be a homogeneous space of G and $X \in \Gamma(TH)$. Then:

- 1. For all $X \in \Gamma(TH)$, there exists a unique horizontal lift \tilde{X} of X. That is, the map $\tilde{\cdot} : \Gamma(TH) \to \Gamma(HG)$ sending $X \mapsto \tilde{X}$ is \mathbb{R} -linear and injective.
- 2. For all smooth curves $q : [a,b] \to H$ and $q_0 \in \pi^{-1}(\{q(a)\})$, there exists a unique horizontal lift $\tilde{q} : [a,b] \to H$ of q satisfying $\tilde{q}(a) = q_0$, called the horizontal lift of X which is π -related to X.

In general, the horizontal lift map will not be surjective, as it need not be the case that $Z_g = Z_h$ whenever $\pi(g) = \pi(h)$ for $Z \in \Gamma(HG)$. The image of the horizontal lift map will be denoted by $\mathcal{B}(G) \subset \Gamma(HG)$, and its elements will be called *basic vector fields*. In particular, $\mathcal{B}(G) \cong \Gamma(TH)$ as an \mathbb{R} -vector space, and a vector field $Z \in \Gamma(TG)$ is basic if and only if it is the horizontal lift of some vector field in $\Gamma(TH)$. Basic vector fields are precisely those which can be pushed-forward to a smooth non-zero vector field on H under π . We may similarly call a curve $g : [a,b] \to G$ basic when it is a horizontal lift of some curve $q : [a,b] \to H$, and a vector field $\tilde{X} \in \Gamma(g)$ basic when it is horizontal and π -related to some $X \in \Gamma(q)$. That is, $(\pi_*)_{g(t)}\tilde{X}(t) = X(t)$ for all $t \in [a,b]$. However, it turns out that these distinctions are redundant, as any smooth curve $g : [a,b] \to G$ satisfying $\dot{g}(t) \in \operatorname{Hor}_{g(t)}$ for all $t \in [a,b]$ is necessarily a horizontal lift of a unique $q : [a,b] \to H$ (namely, $q := \pi \circ g$). This is due to the fact that the following diagram commutes:

From which we find the following lemma (see [13] for more details):

Lemma 6.2. Suppose that $q : [a, b] \to H$ and $\tilde{q} : [a, b] \to G$ is a horizontal lift of q. Then, for any $\tilde{\eta} : [a, b] \to \mathfrak{h}$, there exists a unique $X \in \Gamma(q)$ such that its horizontal lift \tilde{X} along \tilde{q} satisfies $L_{\tilde{q}(t)^{-1}*}\tilde{X}(t) = \tilde{\eta}(t)$ for all $t \in [a, b]$.

6.1. Riemannian Homogeneous Spaces. Consider a connected Lie group G and a homogeneous space H = G/K of G. Since H is a smooth manifold, it can be equipped with a

Riemannian metric. As when discussing Riemannian metrics on Lie groups in Section 3, we are interested in those metrics $\langle \cdot, \cdot \rangle_H$ which in some sense preserve the structure of the homogeneous space. In this case, we wish to choose $\langle \cdot, \cdot \rangle_H$ so that the canonical projection map $\pi: G \to H$ is a *Riemannian submersion*. That is, so that $\pi_*|_g$ is a linear isometry between Hor_g and $T_{\pi(g)}H$ for all $g \in G$. In such a case, we call H a *Riemannian homogeneous space*. It is clear that if H is a Riemannian homogeneous space, then $\langle \mathcal{H}(X), \mathcal{H}(Y) \rangle_G = \langle \pi_* X, \pi_* Y \rangle_H$ for all $X, Y \in T_g G, g \in G$. In particular, $\langle \tilde{X}, \tilde{Y} \rangle_G = \langle X, Y \rangle_H$ for all $X, Y \in T_g G, g \in G$. The metric $\langle \cdot, \cdot \rangle_H$ is said to be *G-invariant* if it is invariant under the left-action Φ_g for all $g \in G$. It can be shown that every homogeneous space H = G/K that admits a *G*-invariant metric is *reductive*. That is, the Lie algebra admits a decomposition $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{h}$, where \mathfrak{s} is the Lie algebra of K, and $\mathfrak{h} \subset \mathfrak{g}$ satisfies $[\mathfrak{s}, \mathfrak{h}] \subset \mathfrak{h}$. In particular, $\mathfrak{h} \cong T_{\pi(e)}(G/K)$ as vector spaces (note that \mathfrak{h} need not be a Lie subalgebra of \mathfrak{g}).

Lemma 6.3. Suppose that $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{h}$ is a reductive Lie algebra. Then $[\mathfrak{s}, \mathfrak{h}] \subset \mathfrak{h}$.

Proof. Since \mathfrak{s} is Lie subalgebra of \mathfrak{g} , we have by definition that $[\mathfrak{s},\mathfrak{s}] \subset \mathfrak{s}$. Let $\xi, \eta \in \mathfrak{s}$ and $\sigma \in \mathfrak{h}$. Then,

$$0 = \langle [\xi, \eta], \sigma \rangle = \left\langle \frac{d}{dt} \Big|_{t=0} \operatorname{Ad}_{\operatorname{Exp}(t\xi)}(\eta), \sigma \right\rangle,$$
$$= \frac{d}{dt} \Big|_{t=0} \left\langle \operatorname{Ad}_{\operatorname{Exp}(t\xi)}(\eta), \sigma \right\rangle.$$

It follows that the map $t \mapsto \langle \operatorname{Ad}_{\operatorname{Exp}(t\xi)}(\eta), \sigma \rangle$ is constant. In particular, since

$$\left\langle \operatorname{Ad}_{\operatorname{Exp}(t\xi)}(\eta), \sigma \right\rangle \Big|_{t=0} = \langle \eta, \sigma \rangle = 0,$$

we have $\langle \operatorname{Ad}_{\operatorname{Exp}(t\xi)}(\eta), \sigma \rangle = 0$ for all $\xi, \eta \in \mathfrak{s}, \sigma \in \mathfrak{m}, t \in \mathbb{R}$. Since the exponential map is a local diffeomorphism, and $\mathfrak{s} = \{X \in \mathfrak{g} \mid \operatorname{Exp}(tX) \in H \text{ for all } t \in \mathbb{R}\}$, it follows that there exists an open subset $V \subset G$ containing e such that $\operatorname{Ad}_g(\mathfrak{s}) \subset \mathfrak{s}$ for all $g \in V \cap H$. Since H is connected by assumption, there exists some finite collection of vectors $\xi_1, \ldots, \xi_n \in \mathfrak{h}$ such that $\operatorname{Exp}(\xi_1) \cdots \operatorname{Exp}(\xi_n) = h$. Hence, for all $\xi \in \mathfrak{h}$, $\operatorname{Ad}_h(\xi) = \operatorname{Ad}_{\operatorname{Exp}(\xi_1)} \circ \cdots \circ \operatorname{Ad}_{\operatorname{Exp}(\xi_n)}(\xi) \in \mathfrak{h}$.

The following result establishes an equivalency between the existence of a G-invariant metrics on H, and the existence of a left-invariant metric on G for which H is a Riemannian homogeneous space.

Lemma 6.4. If G is equipped with a left-invariant metric $\langle \cdot, \cdot \rangle_G$, and $\pi : G \to H$ is a Riemannian submersion, then $\langle \cdot, \cdot \rangle_H$ is G-invariant. Moreover, if $\langle \cdot, \cdot \rangle_H$ is G-invariant, then there exists a left-invariant metric $\langle \cdot, \cdot \rangle_G$ on G such that $\pi : G \to H$ is a Riemannian submersion.

Proof. First suppose G is equipped with a left-invariant metric $\langle \cdot, \cdot \rangle_G$, and that $\pi : G \to H$ is a Riemannian submersion. Let $q \in H, X, Y \in T_qH, g \in G$, and let \tilde{X}, \tilde{Y} be horizontal lifts of X, Y. Then,

$$\begin{split} \left\langle \Phi_{g*}X, \Phi_{g*}Y \right\rangle_{H} &= \left\langle \pi_{*}^{-1} \circ \Phi_{g*}X, \pi_{*}^{-1} \circ \Phi_{g*}Y \right\rangle_{G} = \left\langle \pi_{*}^{-1} \circ \Phi_{g*} \circ \pi_{*}\tilde{X}, \pi_{*}^{-1} \circ \Phi_{g*} \circ \pi_{*}\tilde{Y} \right\rangle_{G} \\ &= \left\langle L_{g*}\tilde{X}, L_{g*}\tilde{Y} \right\rangle_{G} \end{split}$$

where the last equality follows from (6.1). By the left-invariance of $\langle \cdot, \cdot \rangle_G$, we then have

$$\left\langle L_{g*}\tilde{X}, L_{g*}\tilde{Y}\right\rangle_G = \left\langle \tilde{X}, \tilde{Y}\right\rangle_G = \left\langle X, Y\right\rangle_H.$$

In particular, $\langle \cdot, \cdot \rangle_H$ is *G*-invariant.

Now suppose that H is equipped with a G-invariant metric $\langle \cdot, \cdot \rangle_H$. Since H is reductive, we may define the orthogonal projections $\mathcal{V} : \mathfrak{g} \to \mathfrak{s}$ and $\mathcal{H} : \mathfrak{g} \to \mathfrak{h}$, where $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{h}$. Define an inner product on \mathfrak{h} via $\langle \xi, \eta \rangle_{\mathfrak{h}} := \langle \pi_* \xi, \pi_* \eta \rangle_H$ for all $\xi, \eta \in \mathfrak{h}$. Let $\langle \cdot, \cdot \rangle_{\mathfrak{s}}$ be any inner product on \mathfrak{s} , and define an inner product on \mathfrak{g} by $\langle \xi, \eta \rangle_{\mathfrak{g}} := \langle \mathcal{V}(\xi), \mathcal{V}(\eta) \rangle_{\mathfrak{s}} + \langle \mathcal{H}(\xi), \mathcal{H}(\eta) \rangle_{\mathfrak{h}}$. We extend $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$ to a left-invariant Riemannian metric $\langle \cdot, \cdot \rangle_G$ on G via left-translations. Now, for all $X, Y \in \operatorname{Hor}_g$,

$$\begin{split} \left\langle \pi_{*} \big|_{g} X, \pi_{*} \big|_{g} Y \right\rangle_{H} &= \left\langle \Phi_{g^{-1*}} \circ \pi_{*} \big|_{g} X, \Phi_{g^{-1*}} \circ \pi_{*} \big|_{g} Y \right\rangle_{H} \\ &= \left\langle \pi_{*}^{-1} \big|_{e} \circ \Phi_{g^{-1*}} \circ \pi_{*} \big|_{g} X, \pi_{*}^{-1} \big|_{e} \circ \Phi_{g^{-1*}} \circ \pi_{*} \big|_{g} Y \right\rangle_{\mathfrak{h}} \\ &= \left\langle L_{g^{*}} \circ \pi_{*}^{-1} \big|_{e} \circ \Phi_{g^{-1*}} \circ \pi_{*} \big|_{g} X, L_{g^{*}} \circ \pi_{*}^{-1} \big|_{e} \circ \Phi_{g^{-1*}} \circ \pi_{*} \big|_{g} Y \right\rangle_{G} \end{split}$$

By (6.1), we have that $L_{g^*} \circ \pi_*^{-1} |_e \circ \Phi_{g^{-1*}} \circ \pi_* |_g$ is the identity map on Hor_g, so that $\langle X, Y \rangle_G = \langle \pi_* |_g X, \pi_* |_g Y \rangle_H$ for all $X, Y \in \text{Hor}_g$. Hence π is a Riemannian submersion, and so $(H, \langle \cdot, \cdot \rangle_H)$ is a Riemannian homogeneous space with respect to $(G, \langle \cdot, \cdot \rangle_G)$.

In general, there need not exist G-invariant metrics on a homogeneous space. The following theorem from [16] describes a sufficient condition that is commonly fulfilled in an application.

Theorem 6.5. Suppose that G and H := G/K is a homogeneous space of G. If Stab(x) is compact for all $x \in H$, then there exists a G-invariant metric on H.

Hence, if $\operatorname{Stab}(x)$ is compact for all $x \in H$ (which happens for example if H is compact, or equivalently, that G is compact), then by Lemma 6.4, there exists a left-invariant metric on G and a G-invariant metric on H for which H is a Riemannian homogeneous space. For the remainder of the paper, we suppose that G is a simply connected Lie group equipped with a left-invariant metric, and that H is a Riemannian homogeneous space of G. Denote the Levi-Civita connections on H and G with respect to these metrics by ∇ and $\tilde{\nabla}$, respectively. The following lemma from [13] outlines some useful properties of $\tilde{\nabla}$.

Lemma 6.6. Let $X, Y \in \Gamma(TH)$, and $\tilde{X}, \tilde{Y} \in \mathcal{B}(G)$ be the horizontal lifts of X and Y, respectively. Further suppose that $V \in \Gamma(VG)$. Then, the following identities hold:

(6.2)
$$\tilde{\nabla}_{\tilde{X}}\tilde{Y} = \widetilde{\nabla_X Y} + \frac{1}{2}\mathcal{V}([\tilde{X}, \tilde{Y}]),$$

(6.3)
$$\tilde{\nabla}_{\tilde{X}} V = \tilde{\nabla}_V \tilde{X}$$

Apart from the Levi-Civita connection, there is another important connection that we will utilize throughout this section when working on (Riemannian) homogeneous spaces. We define the *horizontal connection* $\tilde{\nabla}^{\mathcal{H}} : \Gamma(TG) \times \Gamma(HG) \to \Gamma(HG)$ on the horizontal bundle by

(6.4)
$$\tilde{\nabla}_W^{\mathcal{H}} Z = \mathcal{H}(\tilde{\nabla}_W Z),$$

for all $W \in \Gamma(TG)$, $Z \in \Gamma(HG)$. It is easy to see that $\tilde{\nabla}^{\mathcal{H}}$ satisfies all the conditions required to be a connection. Moreover, $\tilde{\nabla}^{\mathcal{H}}$ is simply the projection of the Levi-Civita connection on G onto the horizontal bundle.

Remark 6.7. Observe that while $\tilde{\nabla}^{\mathcal{H}}$ is only a connection in the strict sense on the domain $\Gamma(TG) \times \Gamma(HG)$, it is still a well-defined operation on the larger domain $\Gamma(TG) \times \Gamma(TG)$. In fact, since K is a closed Lie subgroup of G, it is an isometrically embedded submanifold when equipped with the induced metric under the inclusion $i: K \hookrightarrow G$. In this case, it is easy to see that $\tilde{\nabla}^{\mathcal{H}}_X Y = \mathbb{I}(X,Y)$, where \mathbb{I} is the second fundamental form. We choose to separate the notations, however, as the contexts in which these objects are discussed are quite different. In particular, the second fundamental form is typically viewed as a 2-tensor field defined over the space of vertical vector fields.

We now outline some additional properties satisfied by $\tilde{\nabla}^{\mathcal{H}}$ and basic vector fields from [13], which follow immediately from Lemma 6.6 and the fact that $\tilde{\nabla}$ is metric compatible and torsion-free:

Lemma 6.8. Let $X, Y \in \Gamma(TH)$, and $\tilde{X}, \tilde{Y} \in \mathcal{B}(G)$ be the horizontal lifts of X and Y, respectively. Further suppose that $W, Z \in \Gamma(HG)$, and $P \in \Gamma(TG)$. Then, the following identities hold:

(6.5)
$$\tilde{\nabla}_{\tilde{X}}^{\mathcal{H}}\tilde{Y} = \tilde{\nabla}_{X}\tilde{Y}$$

(6.6)
$$P \langle W, Z \rangle_G = \left\langle \tilde{\nabla}_P^{\mathcal{H}} W, Z \right\rangle_G + \left\langle W, \tilde{\nabla}_P^{\mathcal{H}} Z \right\rangle_G$$

(6.7)
$$\nabla_W^{\mathcal{H}} Z - \nabla_Z^{\mathcal{H}} W = \mathcal{H}([W, Z])$$

As in Section 3, we may define an operator $\tilde{\nabla}^{\mathfrak{h}} : \mathfrak{h} \times \mathfrak{h} \to \mathfrak{h}$, called the *Riemannian* \mathfrak{h} -connection, via

(6.8)
$$\tilde{\nabla}^{\mathfrak{h}}_{\xi}\eta = \left(\tilde{\nabla}^{\mathcal{H}}_{\phi(\xi)}\phi(\eta)\right)(e).$$

It is clear that $\tilde{\nabla}^{\mathfrak{h}}_{\xi}\eta = \mathcal{H}(\tilde{\nabla}^{\mathfrak{g}}_{\xi}\eta)$, where $\tilde{\nabla}^{\mathfrak{g}}$ is the Riemannian \mathfrak{g} -connection corresponding to the Levi-Civita connection $\tilde{\nabla}$ on G. Therefore, we obtain the explicit expression

(6.9)
$$\tilde{\nabla}^{\mathfrak{h}}_{\xi}\eta = \frac{1}{2}\mathcal{H}([\xi,\eta]_{\mathfrak{g}} - \mathrm{ad}^{\dagger}_{\xi}\eta - \mathrm{ad}^{\dagger}_{\eta}\xi).$$

Given any C^1 curve g, we may further define the operator $\tilde{D}_t^{\mathcal{H}}X := \tilde{\nabla}_g^{\mathcal{H}}X$ for all C^1 horizontal vector fields X along g. It is clear from Lemma 6.8 and Lemma 6.1 that if g is a basic curve and X is a C^1 horizontal vector field along $\pi \circ g$, then $\tilde{D}_t^{\mathcal{H}}\tilde{X} = \widetilde{D_tX}$. We now express Lemma 3.4 in terms of the Riemannian \mathfrak{h} -connection and horizontal connection:

Lemma 6.9. Let $g : [a,b] \to G$ be a basic curve and X a smooth horizontal vector field along g. Suppose that $\xi(t) = g(t)^{-1}\dot{g}(t)$ and $\eta(t) = g(t)^{-1}X(t)$. Then the following relation holds for all $t \in [a,b]$:

(6.10)
$$\tilde{D}_t^{\mathcal{H}} X(t) = g(t) \left(\dot{X}(t) + \tilde{\nabla}_{\xi(t)}^{\mathfrak{h}} \eta(t) \right).$$

EULER-POINCARÉ REDUCTION WITH BROKEN SYMMETRIES ON RIEMANNIAN MANIFOLDS 21

6.2. Euler-Poincaré reduction on Riemannian homogeneous spaces. Throughout the section, we assume that G is a Lie group with left-invariant metric, and H := G/K is a Riemannian homogeneous space of G. For some smooth scalar field $V: H \to \mathbb{R}$, we consider the action function $\mathcal{A}_H : \Omega(H) \to \mathbb{R}$ on H as in (5.1). As usual, our aim is to find the critical points of \mathcal{A}_H over Ω . Since H is a Riemannian manifold, this may be done via variational analysis. However, in many cases, it will be more convenient to work on G—especially in cases where we can apply reduction by symmetry.

To that end, we seek variational principles on G which are equivalent to the variational principle $\delta \mathcal{A}_H = 0$ on H (whose solutions are given by (5.2)). Ultimately, this equivalency will come from the relationship between the metrics on G and H. First, suppose that the boundary conditions on the spaces of admissible curves $\Omega(G) = \Omega^{a,b}_{(g_a,w_a),(g_b,w_b)}$ and $\Omega(H) = \Omega^{a,b}_{(q_a,v_a),(q_b,v_b)}$ are chosen such that:

$$\pi(g_a) = q_a, \qquad \pi(g_b) = q_b, \qquad \pi_* w_a = v_a, \qquad \pi_* w_b = v_b$$

In particular, suppose that $q \in \Omega(G)$ and let $q := \pi \circ q$. Then, it is clear that $q \in \Omega(H)$, and $\dot{q} = \pi_* \dot{g}$. Hence, $\|\dot{q}\|_H = \|\pi_* \dot{g}\|_H = \|\mathcal{H}(\dot{g})\|_G$. On the other hand, suppose that $q \in \Omega(H)$ and let g be a horizontal lift of q. Then for some g_a, g_b , we have $g \in \Omega_{g_a, g_b}(G)$, and $\|\mathcal{H}(\dot{g})\|_G =$ $\|\dot{g}\|_G = \|\dot{q}\|_H$. This motivates the action functional $\mathcal{A}_G : \Omega(G) \to \mathbb{R}$ given by

(6.11)
$$\tilde{\mathcal{A}}_G = \int_a^b \left(\frac{1}{2} \|\mathcal{H}(\dot{g}(t))\|_G^2 - \tilde{V}(g(t))\right) dt,$$

where $\tilde{V} := V \circ \pi$. We have the following Proposition:

Proposition 6.10. Let $q \in \Omega(H)$ and $g \in \Omega(G)$ satisfy $\pi \circ g = q$ on [a, b]. Then q is a critical point of \mathcal{A}_H if and only if g is a critical point of $\tilde{\mathcal{A}}_G$.

Proof. Suppose that q is a critical point of \mathcal{A}_H , and consider an admissible variation g_s of g. It follows that $q_s := \pi \circ g_s$ is an admissible variation of q, and $\dot{q}_s = \pi_* \dot{g}$. Hence,

$$\delta \tilde{\mathcal{A}}_G[g] = \frac{\partial}{\partial s} \Big|_{s=0} \int_a^b \left(\frac{1}{2} \|\mathcal{H}(\dot{g}_s)\|_G^2 - \tilde{V}(g_s) \right) dt$$
$$= \frac{\partial}{\partial s} \Big|_{s=0} \int_a^b \left(\frac{1}{2} \|\dot{q}_s\|_H^2 - V(q_s) \right) dt,$$
$$= \mathcal{A}_H[q] = 0.$$

Therefore, g is a critical point of $\tilde{\mathcal{A}}_G$. Now suppose that g is a critical point of $\tilde{\mathcal{A}}_G$, and consider an admissible variation q_s of q. Let \tilde{q}_s be the horizontal lift of q_s at g(a), and consider the family of curves defined by $g_s(t) = \tilde{q}_s(t)\tilde{q}_{\frac{s(t-a)}{b-a}}(b)^{-1}g(b)$ for all $(s,t) \in (-\epsilon,\epsilon) \times [a,b]$. Observe

that

1.
$$g_0(t) = \tilde{q}_0(t)\tilde{q}_0(b)^{-1}g(b) = g(t)$$
 for all $t \in [a, b]$,

2.
$$g_s(a) = \tilde{q}_s(a)\tilde{q}_0(b)^{-1}g(b) = g(a) \ s \in (-\epsilon, \epsilon)$$

- 2. $g_s(a) = q_s(a)q_0(c) g(a) + c \in (-, c),$ 3. $g_s(b) = \tilde{q}_s(b)\tilde{q}_s^{-1}(b)g(b) = g(b)$, for all $s \in (-\epsilon, \epsilon)$. 4. $\pi \circ g_s \equiv \pi \circ \tilde{q}_s \equiv q_s$, since $\tilde{q}_{\frac{s(t-a)}{b-a}}(b)^{-1}$ and g(b) belong to the same equivalency class for all $(s,t) \in (\epsilon,\epsilon) \times [a,b]$.

It follows that g_s is indeed an admissible variation of g, so that

$$\delta \mathcal{A}_{H}[q] = \frac{\partial}{\partial s} \Big|_{s=0} \int_{a}^{b} \left(\frac{1}{2} \| \dot{q}_{s} \|_{H}^{2} - V(q_{s}) \right) dt$$
$$= \frac{\partial}{\partial s} \Big|_{s=0} \int_{a}^{b} \left(\frac{1}{2} \| \mathcal{H}(\dot{g}_{s}) \|_{G}^{2} - \tilde{V}(q_{s}) \right) dt$$
$$= \tilde{\mathcal{A}}_{G}[g] = 0.$$

From Proposition 6.10, we can generate critical points of \mathcal{A}_H by finding critical points of $\tilde{\mathcal{A}}_G$ on then projecting them onto H. Assuming that $V \circ \pi$ admits a partial symmetry, we could then, in principle, apply symmetry reduction to $\tilde{\mathcal{A}}_G$. However, the inclusion of the horizontal projection \mathcal{H} in $\tilde{\mathcal{A}}_G$ makes the analysis difficult. Our goal now is to find a *constrained* variational principle for $\mathcal{A}_G : \Omega \to \mathbb{R}$ defined by (5.1) on G with the potential \tilde{V} .

Definition 6.11. We call an admissible variation g_s of a basic curve $g \in \Omega(G)$ for which both $\dot{g}_s(t)$ and $\frac{\partial}{\partial s}g_s(t)$ are horizontal vector fields a basic variation of g.

Remark 6.12. Note that the variation defined by $g_s(t) = g(t)$ for all s is trivially a basic variation, so that the definition is well-defined.

Proposition 6.13. Suppose that $g \in \Omega(G)$ is a critical point of $\tilde{\mathcal{A}}_G$. If g is basic, then $\delta \mathcal{A}_G[g] = 0$ among all basic variations.

Proof. Let g_s be a basic variation of g. Then,

$$\begin{split} \delta \mathcal{A}[g] &= \frac{\partial}{\partial s} \Big|_{s=0} \int_{a}^{b} \left(\frac{1}{2} \| \dot{g}_{s}(t) \|_{G}^{2} - \tilde{V}(g_{s}(t)) \right) dt \\ &= \frac{\partial}{\partial s} \Big|_{s=0} \int_{a}^{b} \left(\frac{1}{2} \| \mathcal{H}(\dot{g}_{s}(t)) \|_{G}^{2} - \tilde{V}(g_{s}(t)) \right) dt \\ &= \delta \tilde{\mathcal{A}}[g] = 0. \end{split}$$

We now derive necessary conditions on g to solve the constrained variational principle on G.

Proposition 6.14. A basic curve $g \in \Omega$ satisfies the variational principle $\delta \mathcal{A}_G[g] = 0$ among all basic variations of g if and only if g is smooth and satisfies

(6.12)
$$\tilde{D}_t^{\mathcal{H}} \dot{g}(t) = -\operatorname{grad} \tilde{V}(g(t)),$$

for all $t \in [a, b]$.

Proof. Let g_s be a basic variation of G with variational vector field δg . Then, from Lemma 6.8 and the Symmetry Lemma,

$$\begin{split} \delta \mathcal{A}_G[g] &= \frac{\partial}{\partial s} \Big|_{s=0} \int_a^b \left(\frac{1}{2} \| \dot{g}_s(t) \|_G^2 - \tilde{V}(g_s(t)) \right) dt \\ &= \int_a^b \left(\left\langle \tilde{D}_s^{\mathcal{H}} \dot{g}_s, \dot{g}_s \right\rangle_G - \left\langle \operatorname{grad} \tilde{V}(g_s), \frac{\partial}{\partial s} \dot{g}_s \right\rangle_G \right) \Big|_{s=0} dt \\ &= \int_a^b \left(\left\langle \tilde{D}_t^{\mathcal{H}} \delta g, \dot{g} \right\rangle_G - \left\langle \operatorname{grad} \tilde{V}(g), \delta g \right\rangle_G \right) dt. \end{split}$$

This manuscript is for review purposes only.

As in the proof of Theorem 5.3, the smoothness of g can be shown via a bootstrapping method. Integrating once by parts, we then obtain

$$\delta \mathcal{A}_G[g] = \int_a^b \left\langle \tilde{D}_t^{\mathcal{H}} \dot{g} + \operatorname{grad} \tilde{V}(g), \delta g \right\rangle_G dt = 0$$

for all basic variational vector fields δg . Now let $h \in G$, $X \in \operatorname{Ver}_h$, and observe that $\left\langle \operatorname{grad} \tilde{V}(h), X \right\rangle_G = d\tilde{V}(X) = dV \circ \pi_*(X) = 0$. Hence $\operatorname{grad} \tilde{V} \in \Gamma(HG)$. Moreover, if we consider any $\tilde{Y} \in \operatorname{Hor}_h$, then:

$$\left\langle \operatorname{grad} \tilde{V}(h), \tilde{Y} \right\rangle_G = d\tilde{V}(\tilde{Y}) = dV(Y) = \left\langle \operatorname{grad} V(\pi(h)), Y \right\rangle_H = \left\langle \operatorname{grad} \widetilde{V(\pi(h))}, \tilde{Y} \right\rangle_G,$$

where $Y := \pi_* \tilde{Y}$. It follows that $\operatorname{grad} \tilde{V}$ is precisely the horizontal lift of $\operatorname{grad} V$, so that $\operatorname{grad} \tilde{V}$ is basic. Hence,

$$\int_{a}^{b} \left\langle \tilde{D}_{t}^{\mathcal{H}} \dot{g} + \operatorname{grad} \tilde{V}(g), \delta g \right\rangle_{G} dt = \int_{a}^{b} \left\langle \tilde{D}_{t}^{\mathcal{H}} \dot{g} + \operatorname{grad} \widetilde{V(\pi(g))}, \delta g \right\rangle_{G} dt.$$

It follows from 6.1 and the fact that G is simply connected that every horizontal variational vector field is the variational vector field of some basic variation, from which (6.12) follows.

Corollary 6.15. If $q \in \Omega(H)$ is a critical point of \mathcal{A}_H , then its horizontal lift \tilde{q} satisfies equation (6.12). Moreover, if $g \in \Omega(G)$ is smooth, basic, and satisfies (6.12), then $\pi \circ g$ is a critical point of \mathcal{A}_H .

Proof. The first direction is an immediate consequence of Propositions 6.13, 6.10, and 6.14. Now suppose that $g \in \Omega(G)$ is basic and satisfies (6.12). Since g is smooth and basic, the curve $q := \pi \circ g$ is smooth and satisfies $\tilde{D}_t^{\mathcal{H}} \dot{g} = \widetilde{D_t} \dot{q}$. Moreover, from the proof of Proposition 6.14, we have $\operatorname{grad} \tilde{V}(q) = \operatorname{grad} \tilde{V}(q)$. Hence,

$$0 = \tilde{D}_t^{\mathcal{H}} \dot{g} + \operatorname{grad} \tilde{V}(g) = D_t \dot{q} + \operatorname{grad} V(q),$$

so that $D_t \dot{q} + \text{grad}V(q) = 0$ on [a, b], which are precisely the Euler-Lagrange equations for \mathcal{A}_H .

Now that we have found a (constrained) variational principle on G which is equivalent to $\delta A_H = 0$ on H, we seek to apply reduction by symmetry under the assumption that V admits a partial symmetry over some smooth manifold M.

Theorem 6.16. Suppose that \tilde{V} admits a partial symmetry over some smooth manifold M. Then, $g \in \Omega(G)$ satisfies (6.12) if and only if $\xi := g^{-1}\dot{g}$ and $\alpha := \Psi_{g^{-1}}(\alpha_0)$ satisfy the Euler-Poincaré equations

(6.13)
$$\dot{\xi} + \tilde{\nabla}^{\mathfrak{h}}_{\xi} \xi = -\operatorname{grad}_{1} \tilde{V}_{\mathrm{ext}}(e, \alpha),$$

(6.14)
$$\dot{\alpha} = -\xi_M(\alpha),$$

on [a, b]. Moreover, g is basic if and only if $\xi(a) = g(a)^{-1}\dot{g}(a) \in \mathfrak{h}$.

Proof. From Lemma 6.9, we have $\tilde{D}_t^{\mathcal{H}}\dot{g} = g(\dot{\xi} + \tilde{\nabla}_{\xi}^{\mathfrak{h}}\xi)$. Hence, if g_s is a basic variation of g, then from the proofs of Propositions 6.14 and Theorem 5.3,

$$\begin{split} \frac{\partial}{\partial s}\Big|_{s=0} \int_{a}^{b} \left(\frac{1}{2} \|\dot{g}_{s}(t)\|_{G}^{2} - \tilde{V}_{\text{ext}}(e, g_{s}(t))\right) dt &= \int_{a}^{b} \left(\left\langle \tilde{D}_{t}^{\mathcal{H}} \dot{g}, \delta g \right\rangle_{G} - \frac{\partial}{\partial s}\Big|_{s=0} \tilde{V}(g_{s}(t))\right) dt \\ &= \int_{a}^{b} \left\langle \dot{\xi} + \tilde{\nabla}_{\xi}^{\mathfrak{h}} \xi + \text{grad}_{1} \tilde{V}_{\text{ext}}(e, \alpha), \eta \right\rangle_{G} dt, \end{split}$$

where $\eta := g^{-1} \delta g \in \mathfrak{h}$. Equations (6.13)-(6.14) follow immediately.

To see that g is basic if and only if $\xi(a) \in \mathfrak{h}$, observe that (6.13) can be written in the form

$$\xi = X(\xi, \alpha),$$

where $X : \mathfrak{g} \times M \to \mathfrak{h}$ is everywhere horizontal. Hence, $\dot{\xi}(t) \in \mathfrak{h}$ for all $t \in [a, b]$. If $\mathcal{B} = \{\alpha_1, \ldots, \alpha_n\}$ is an ordered basis for $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{s}$ such that $\{\alpha_1, \ldots, \alpha_k\}$ is an ordered basis for \mathfrak{h} , and $\{\alpha_{k+1}, \ldots, \alpha_n\}$ is an ordered basis for \mathfrak{s} , it then follows that the coordinates of $\dot{\xi}$ with respect to \mathcal{B} take the form

$$[\xi(t)]_{\mathcal{B}} = \begin{bmatrix} \dot{\xi}_1(t) & \cdots & \dot{\xi}_k(t) & 0 & \cdots & 0 \end{bmatrix}^T$$

from which we find, upon integration, that the coordinates of ξ are of the form

$$[\xi]_{\mathcal{B}} = \begin{bmatrix} \xi_1(t) & \cdots & \xi_k(t) & \xi_{k+1}(a) & \cdots & \xi_n(a) \end{bmatrix}^T,$$

from which it is clear that $\xi(t) \in \mathfrak{h}$ for all $t \in [a, b]$ if and only if $\xi(a) \in \mathfrak{h}$ (in other words, that $\xi_{k+1}(a) = \cdots = \xi_n(a) = 0$).

Remark 6.17. Theorem 6.16 tells us that we may search for solutions $\delta \mathcal{A}_H = 0$ over $\Omega(H)$ by integrating the Euler-Poincaré equations (6.13)-(6.14), solving the reconstruction equation $\dot{g} = g\xi$, and then projecting these solutions onto H. Numerically, our BVP can be solved by applying a shooting method over horizontal initial velocities. Moreover, note that solutions are guaranteed to exist by the existence of solutions to $\delta \mathcal{A}_H = 0$.

6.3. Bi-invariant metrics and Riemannian Symmetric Spaces. Suppose G is a Lie group equipped with a left-invariant metric, and H := G/K is a Riemannian homogeneous space. It is easily seen that the metric $\langle \cdot, \cdot \rangle_G$ is right K-invariant. That is, $\langle Xk, Yk \rangle_G = \langle X, Y \rangle_G$ for all $k \in K$. In the case that the metric $\langle \cdot, \cdot \rangle_G$ is bi-invariant, it follows from Lemma 3.10 that

(6.15)
$$\tilde{\nabla}^{\mathfrak{h}}_{\xi}\eta = \frac{1}{2}\mathcal{H}\big([\xi,\eta]\big).$$

Another important special case of spaces that appears frequently in applications are *Rie-mannian symmetric spaces*, which are Riemannian homogeneous spaces such that there exists an involutive automorphism $\sigma : G \to G$ with $K = \{g \in G \ \sigma(g) = g\}$. It can be seen that every Riemannian symmetric space of the form G/K satisfies the *Cartan Decomposition*:

$$(6.16) \qquad [\mathfrak{s},\mathfrak{s}] \subset \mathfrak{s}, \quad [\mathfrak{s},\mathfrak{h}] \subset \mathfrak{h}, \quad [\mathfrak{h},\mathfrak{h}] \subset \mathfrak{s}.$$

The condition $[\mathfrak{s},\mathfrak{s}] \subset \mathfrak{s}$ follows from the fact that \mathfrak{s} is a Lie subalgebra of \mathfrak{g} , and $[\mathfrak{s},\mathfrak{h}] \subset \mathfrak{h}$ follows from the fact that every Riemannian homogeneous space is reductive. The novel condition, then, is $[\mathfrak{h},\mathfrak{h}] \subset \mathfrak{s}$, which is not in general true for Riemannian homogeneous spaces. Moreover, if G is simply connected (an assumption that we always make), then every Riemannian homogeneous space satisfying (6.16) is a Riemannian symmetric space. It is neither necessary nor sufficient that $\langle \cdot, \cdot \rangle_G$ be bi-invariant in order for G/K to be Riemannian symmetric, though it is the case in many important applications. In the case that H is a Riemannian symmetric space, it is easily seen from equation (6.9) and the Cartan decomposition (6.16) that

(6.17)
$$\tilde{\nabla}^{\mathfrak{h}}_{\varepsilon}\eta = 0$$

6.4. Example: Obstacle avoidance on Riemannian Homogeneous Spaces. Suppose that G is a Lie group equipped with a left-invariant metric, and H is a Riemannian homogeneous space of G. As in Example 5.3, we consider an obstacle avoidance problem on H, with point-obstacle $q_0 \in H$. Towards that end, we design the artificial potential $V : H \to \mathbb{R}$ as given by (5.9). The lifted potential then takes the form:

$$\tilde{V}(g) = f(d_H^2(\pi(g), q_0))$$

where $d_H : H \times H \to \mathbb{R}$ is the Riemannian distance function on H. Since the Riemannian metric on H is G-invariant with respect to the transitive group action Φ of G on H, so too is d_H . Hence, the extended potential $V_{\text{ext}} : G \times G \to \mathbb{R}$ given by

$$\widetilde{V}_{\mathrm{ext}}(g,h) = f(d_H^2(\pi(g),\pi(h)))$$

is invariant under left-translation, and satisfies $\tilde{V}_{\text{ext}}(g, g_0) = \tilde{V}(g)$ for all $g_0 \in \pi^{-1}(\{q_0\})$. In other words, \tilde{V} admits a partial symmetry over G. In principle, we may now calculate the Euler-Poincaré equations given by (6.13)-(6.14), from which critical points of \mathcal{A}_H can be calculated through the reconstruction equation together with $q = \pi(g)$.

However, there is seemingly a major problem with this approach: the advantages obtained by considering the Euler-Poincaré equations on G instead of the Euler-Lagrange equations on H are somewhat superficial, as we still must be able to calculate the gradient vector field of the Riemannian distance on H at all points along the minimizers. In particular, it is not generally the case that $d_G(g,h) = d_H(\pi(g), \pi(h))$, so that we cannot directly take advantage of the additional symmetries afforded by the Lie group (particularly in the case where the metric is bi-invariant, which allows identifying the Lie logarithmic map with the inverse Riemannian exponential map at the identity as we did in Example 5.3). To address this issue, we take the following Lemma from [13]:

Lemma 6.18. Suppose that G is a Lie-group equipped with a left-invariant Riemannian metric, and H is a Riemannian homogeneous space of G. Then for all $g \in G$, and for all $h \in G$ contained within a geodesically convex neighborhood of g, there exists a unique $h^* \in \pi^{-1}(\{\pi(h)\})$ such that $d_G(g, h^*) = d_H(\pi(g), \pi(h))$. Moreover, the function $\theta : G \times G \to G$ defined implicitly by $\theta(g, h) = h^*$ is smooth and left-invariant.

In fact, we are able to calculate the map θ explicitly, which was not previously done in the literature:

Lemma 6.19. Suppose that $g, h \in G$ are contained within a goedesically convex neighborhood on which the exponential map on G is a diffeomorphism, then

(6.18)
$$\theta(g,h) = \exp_g(\mathcal{H}(\exp_g^{-1}(h))).$$

Proof. Since g and h are contained within a geodesically convex neighborhood, it follows that, for $X := \exp_g^{-1}(h)$, the curve $c(t) = \exp_g(tX)$ is the unique geodesic satisfying c(0) = g and $\dot{c}(0) = X$, and we have c(1) = h. Let \tilde{c} be the curve defined by $\tilde{c}(t) = \pi \circ \overline{\exp_g(tX)}$ with $\tilde{c}(0) = g$. It follows from Proposition 6.14 is a geodesic, and $\dot{\tilde{c}}(0) = \mathcal{H}(X)$. Moreover, the geodesic c^* defined by $c^*(t) = \exp_g(t\mathcal{H}(X))$ satisfies $c^*(0) = g$ and $\dot{c}^*(0) = \mathcal{H}(X)$. Hence $c^* = \tilde{c}$. It follows that $\pi \circ \exp_g(tX) = \pi \circ \exp_g(t\mathcal{H}(X))$, and so $\exp_g(\mathcal{H}(X)) := \exp_g(\mathcal{H}(\exp_g^{-1}(h))) \in \pi^{-1}(\{\pi(h)\})$.

Hence, as long as $g, h \in G$ can be connected by a geodesic, our extended artificial potential takes the form:

(6.19)
$$\tilde{V}_{\text{ext}}(g,h) = f(d_G^2(g,\theta(g,h)))$$

where θ is given by (6.18). In particular, if G is equipped with a bi-invariant metric, then

 $\operatorname{grad}_{1} \tilde{V}_{\operatorname{ext}}(e, \alpha) = -2f'(\|\operatorname{Log}(\theta(e, \alpha))\|_{G}^{2})\operatorname{Log}(\theta(e, \alpha)).$

Using (6.18), it is easily seen that

$$\theta(e, \alpha) = \operatorname{Exp}(\mathcal{H}(\operatorname{Log}(\alpha))),$$

so that

(6.20)
$$\operatorname{grad}_{1} \tilde{V}_{\text{ext}}(e, \alpha) = -2f'(\|\mathcal{H}(\operatorname{Log}(\alpha))\|_{G}^{2})\operatorname{Log}(\mathcal{H}(\operatorname{Log}(\alpha))),$$

after which the Euler-Poincaré equations can be calculated by Proposition 6.16.

Remark 6.20. Note that the projected curve $q = \pi \circ g$ avoids q_0 if and only if g avoids the entire fiber $\pi^{-1}(\{q_0\})$. However, this is handled by our methodology. Indeed, suppose that f is designed so that g avoids the point $\theta(g, g_0)$ (which is always possible by correctly choosing the underlying parameters τ, k, D [12]). That is, $d_G(g(t), \theta(g(t), g_0)) > R$ for all $t \in [a, b]$ and some $R \in \mathbb{R}$. Assume towards contradiction that there exists some point $g^* \in \pi^{-1}(\{q_0\})$ and $T \in [a, b]$ such that $d_G(g(T), g^*) \leq R$. Then,

$$R \ge d_G(g(T), g^*) \ge d_H(\pi(g(T)), \pi(g^*)) = d_G(g(T), \theta(g(T), g^*)) = d_G(g(T), \theta(g(T), g_0)),$$

since g_0 and g^* belong to the same fiber.

6.5. Example: Euler-Poincaré Equations on S^2 . Consider the Lie group G = SO(3) equipped with the bi-invariant metric defined by $\langle \hat{\Omega}_1, \hat{\Omega}_2 \rangle_{SO(3)} = \Omega_1^T \Omega_2$ for all $\Omega_1, \Omega_2 \in \mathbb{R}^3$, where we've made use of the hat map defined in Example 4.1 to identify $\mathfrak{so}(3) \cong \mathbb{R}^3$. We let SO(3) act on \mathbb{R}^3 from the matrix multiplication (i.e. $\Phi : SO(3) \times \mathbb{R}^3 \to \mathbb{R}^3$ given by

 $\begin{array}{l} (R,x) \mapsto Rx). \text{ We let } K \subset \mathrm{SO}(3) \text{ be the Lie subgroup given by the elements of the form} \\ k = \begin{bmatrix} \cos(\theta) & \sin(\theta) & 0 \\ -\sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ with } \theta \in S^1. \end{array}$

It is clear that $K \cong SO(2)$ and $K = Stab(e_3)$, where $e_3 = \begin{bmatrix} 0 & 0 & 1 \end{bmatrix}^T$ and $Stab(e_3)$ denotes the stabilizer subgroup of the left-action Φ . Moreover, the restriction of Φ to the unit sphere S^2 (embedded in \mathbb{R}^3 via inclusion) is transitive. It follows that the quotient space $H = SO(3)/SO(2) \cong S^2$ is a homogeneous space of SO(3), and the projection map $\pi : SO(3) \to S^2$ is given by $\pi(R) = Re_3$ for all $R \in SO(3)$.

We consider the orthonormal basis $\{\hat{\Omega}_1, \hat{\Omega}_2, \hat{\Omega}_3\}$ of $\mathfrak{g} = \mathfrak{so}(3)$ given by

(6.21)
$$\hat{\Omega}_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \hat{\Omega}_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \quad \hat{\Omega}_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

It is clear from the definition of the hat map that $\hat{\Omega}_1 = \hat{e}_1, \hat{\Omega}_2 = \hat{e}_2$, and $\hat{\Omega}_3 = \hat{e}_3$. Moreover, we have $\pi_*\hat{\Omega}_1 = -e_2, \ \pi_*\hat{\Omega}_2 = e_1$, and $\pi_*\hat{\Omega}_3 = 0$, so that, in particular, $\mathfrak{s} = \ker(\pi_*|_I) = \operatorname{span}\{\hat{\Omega}_3\}$ and $\mathfrak{h} = \mathfrak{s}^{\perp}$. Define an inner product on $T_{e_3}S^2$ via the relation $\langle X, Y \rangle_{T_{e_3}S^2} := \langle \pi_*^{-1}X, \pi_*^{-1}Y \rangle_{\mathrm{SO}(3)}$ for all $X, Y \in T_{e_3}S^2$. Since $\{e_1, e_2\}$ forms a basis for $T_{e_3}S^2$, it is clear that—letting $X = X^1e_1 + X^2e_2$ and $Y = Y^1e_1 + Y^2e_2$ —

$$\begin{split} \langle X, Y \rangle_{T_{e_3}S^2} &= X^1 Y^1 \langle e_1, e_1 \rangle_{T_{e_3}S^2} + X^1 Y^2 \langle e_1, e_2 \rangle_{T_{e_3}S^2} + X^2 Y^1 \langle e_2, e_1 \rangle_{T_{e_3}S^2} \\ &+ X^2 Y^2 \langle e_2, e_2 \rangle_{T_{e_3}S^2} , \\ &= X^1 Y^1 \left\langle \hat{\Omega}_1, \hat{\Omega}_1 \right\rangle_{\mathrm{SO}(3)} + X^1 Y^2 \left\langle \hat{\Omega}_1, \hat{\Omega}_2 \right\rangle_{\mathrm{SO}(3)} + X^2 Y^1 \left\langle \hat{\Omega}_2, \hat{\Omega}_1 \right\rangle_{\mathrm{SO}(3)} \\ &+ X^2 Y^2 \left\langle \hat{\Omega}_2, \hat{\Omega}_2 \right\rangle_{\mathrm{SO}(3)} , \\ &= X^1 Y^1 + X^2 Y^2 , \end{split}$$

so that $\langle \cdot, \cdot \rangle_{T_{e_3}S^2}$ is just the standard Euclidean metric with respect to the basis $\{e_1, e_2\}$. We extend this inner product to an SO(3)-invariant Riemannian metric on S^2 by left-action. That is,

(6.22)
$$\langle X, Y \rangle_{S^2} = \left\langle R^T X, R^T Y \right\rangle_{T_{e_3}S^2} = R^T X \cdot R^T Y = X \cdot Y$$

for all $X, Y \in T_q S^2$ and for any $R \in SO(3)$ such that $\pi(R) = q$. It is clear that with this metric, S^2 becomes a Riemannian homogeneous space, and from the matrix commutator relations,

$$[\hat{\Omega}_1, \hat{\Omega}_2] = \hat{\Omega}_3, \qquad \qquad [\hat{\Omega}_1, \hat{\Omega}_3] = -\hat{\Omega}_2, \qquad \qquad [\hat{\Omega}_2, \hat{\Omega}_3] = \hat{\Omega}_1,$$

it is immediately clear that the decomposition $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{h}$ satisfies Cartan's Decomposition (6.16), so that S^2 is in fact a Riemannian symmetric space of SO(3). We see also that the horizontal projection can be calcualted by

(6.23)
$$\mathcal{H}(\hat{\Omega}) = \widehat{\Omega \times e_3}$$

for all $\hat{\Omega} \in \mathfrak{so}(3)$, so that its norm is given by

(6.24)
$$\|\mathcal{H}(\hat{\Omega})\|_G = \|\hat{\Omega}\|_G^2 \sin(\theta),$$

where $\theta = \arccos\left(\frac{\Omega \cdot e_3}{\|\Omega\|}\right)$, so that $\sin(\theta) = \sqrt{1 - \left(\frac{\Omega \cdot e_3}{\|\Omega\|}\right)^2}$.

Consider a curve $q : [a, b] \to S^2$, and let $R : [a, b] \to SO(3)$ be a horizontal lift of q. Hence, $q(t) = R(t)e_3$ for all $t \in [a, b]$. Let $X \in \Gamma(q)$, and denote its horizontal lift by \tilde{X} . It follows that $\tilde{X} = R\hat{\Omega}$ for some curve $\hat{\Omega} : [a, b] \to \mathfrak{h}$. Then,

$$D_t X = \pi_* (\tilde{D}_t^{\mathcal{H}} \tilde{X}) = (\tilde{D}_t^{\mathcal{H}} \tilde{X}) e_3 = R(\dot{\hat{\Omega}} + \tilde{\nabla}_{R^T \dot{R}}^{\mathfrak{h}} \hat{\Omega}) e_3.$$

Since S^2 is a Riemannian symmetric space, we have from (6.17) that $\tilde{\nabla}^{\mathfrak{h}}_{B^T\dot{B}}\hat{\Omega} = 0$. Hence,

$$D_t X = R\dot{\Omega}e_3 = R(\dot{\Omega} \times e_3) = R\dot{\Omega} \times Re_3 = R\dot{\Omega} \times q.$$

Let $\omega := R\Omega$. Then, on one hand, $\dot{\omega} = \dot{R}\Omega + R\dot{\Omega} = R(\Omega \times \Omega) + R\dot{\Omega} = R\dot{\Omega}$, so that $D_t X = \dot{\omega} \times q$. On the other hand, we have $X = \tilde{X}e_3 = R\Omega = R\Omega \times q = \omega \times q$. Since X is everywhere orthogonal to q, this further implies that $\omega = q \times X$, so that $\dot{\omega} = \dot{q} \times X + q \times \dot{X}$. Therefore, $D_t X = (\dot{q} \times X) \times q - q \times (q \times \dot{X})$. It is easy to see from the vector triple product rule that $(\dot{q} \times X) \times q = 0$, so that

$$(6.25) D_t X = -q \times (q \times X),$$

which is precisely the orthogonal projection of \dot{X} onto the tangent bundle. It follows that the critical points of \mathcal{A} on S^2 satisfy

(6.26)
$$q \times (q \times \ddot{q}) = -\operatorname{grad} V(q)$$

Suppose now that $\tilde{V}(R) := V(Re_3)$ admits a partial symmetry. Then, from Proposition 6.16, together with equation (6.17), the Euler-Poincaré equations (6.13)-(6.14) take the form

(6.27)
$$\dot{\Omega} = -\operatorname{grad}_1 \tilde{V}(I, \alpha),$$

(6.28)
$$\dot{\alpha} = -\xi_M(\alpha),$$

together with the reconstruction equations $\dot{R} = R\hat{\Omega}$ and $q = Re_3$.

7. Optimal Control of left-invariant homogeneous systems. Let H be a Riemannian homogeneous space of some connected Lie group G equipped with a left-invariant Riemannian metric $\langle \cdot, \cdot \rangle_G$. As in Section 6.2, we suppose that the Lie algebra decomposes as $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{s}$, where \mathfrak{s} is the Lie algebra of the closed Lie subgroup $K \subset G$ such that $H \cong G/K$, and $\mathfrak{h} \cong T_{\pi(e)}H$, where $\pi: G \to H$ is the canonical projection, and e is the identity element of G. We further suppose that $f: TH \to TH$ is a diffeomorphism for which there exists positive constants $c, C \in \mathbb{R}$ such that:

(7.1)
$$c\|u\|_{H} \le \|f(q,u)\|_{H} \le C\|u\|_{H}$$

for all $(q, u) \in TH$.

Remark 7.1. In many applications, f will in fact be linear in the second argument. In such a case, for every $q \in H$, there exists positive constants c_q, C_q such that $c_q ||u||_H \leq ||f(q, u)||_H \leq C_q ||u||_H$ for all $u \in T_q H$. (7.1) will automatically follow, for instance, in the case that H is compact.

We wish to solve the following optimal control problem:

(7.2)
$$\min_{q,u} \int_{a}^{b} \left(\frac{1}{2} \|u\|_{H}^{2} - V(q(t))\right) dt$$

subject to $q \in \Omega$, and $\dot{q} = f(q, u)$, where $V : H \to \mathbb{R}$ is a smooth scalar field.

The fact that f is a diffeomorphism makes it possible to view the optimal control problem (7.2) as a variational problem over curves $q \in \Omega$. That is due to the fact that, for any such q, there will exist a unique control input u such that $\dot{q} = f(q, u)$. Moreover, the inequality given in the second assumption guarantees that

$$\frac{1}{C^2} \int_a^b \left(\frac{1}{2} \| f(q, u) \|_H^2 - V(q(t)) \right) dt \le \int_a^b \left(\frac{1}{2} \| u \|_H^2 - V(q(t)) \right) dt \le \frac{1}{c^2} \int_a^b \left(\frac{1}{2} \| f(q, u) \|_H^2 - V(q(t)) \right) dt.$$

It follows that the optimal control problem (7.2) is equivalent to minimizing the actionfunctional $\mathcal{A} : \Omega \to \mathbb{R}$ defined in equation (5.1), from which the optimal control can be recovered via the inverse kinematics $u = f^{-1}(q, \dot{q})$. We summarize these results with the following proposition:

Proposition 7.2. The pair (q, u) solve the optimal control problem (7.2) if and only if $q \in \Omega$ is smooth, and

$$D_t \dot{q} = -\text{grad}V(q)$$
$$u = f^{-1}(q, \dot{q})$$

We may now utilize all the techniques developed in Section 6.2 to calculate the critical points of \mathcal{A} , and reduce them by symmetry under the assumption that $\tilde{V} := V \circ \pi$ admits a partial symmetry. In particular, we have the following corollary:

Corollary 7.3. Suppose that \tilde{V} admits a partial symmetry over some smooth manifold Q. Then, the pair (q, u) solve the optimal control problem (7.2) if and only if $q = \pi \circ g$ and $u = f^{-1}(\pi \circ g, \pi_* \dot{g})$, where g is a basic curve such that $\xi := g^{-1}\dot{g}$ and $\alpha := \Psi_{g^{-1}}(\alpha_0)$ satisfy (6.13) - (6.14).

7.1. Optimal Control of Robotic Manipulators. Robotic manipulators are one of the most fundamental robots in engineering, with an expansive list of applications from assembly lines, to medicine, to prosthetic exoskeletons [19]. They are characterized by programmable mechanical limbs connected by various types of joints, the most common of which are:

- **Revolute joints**, which allow for twisting about a fixed axis $(S^1 \cong SO(2))$.
- **Prismatic joints**, which allow for linear translation along a fixed axis (\mathbb{R}) .
- Cylindrical joints, which allow for rotation and translation about a fixed axis $(S^1 \times \mathbb{R} \cong SO(2) \times \mathbb{R})$.
- Ball and socket joints, which allow for rotations about all three axes (S^2) .

An infinite variety of robotic manipulators can be constructed by pairing these joints (together with some mechanical limbs) appropriately. A typical control problem is to design some kinematic controls such that the end effector (the end point of the manipulator, at which some sort of tool is attached to allow it to interact with its environment) track a given trajectory. This is a classical kinematic control problem, which has been exhaustively studied in the literature [19].

An associated optimal control problem that may arise (for example, in an assembly line) is to find a control-trajectory pair such that the end effector interpolates some set of knot points while minimizing energy consumption, and potentially with the additional task of avoiding static obstacles in the environment, or avoiding certain configurations. This energy consumption, in turn, is directly related to the magnitude of the control.

As described above, the typical joint types are all homogeneous spaces (or Lie groups), so that the configuration space H for such a robotic manipulator will be the product of Lie groups and homogeneous spaces, which is itself a homogeneous space. The position of the end effector will thus be described by some map $\varphi : H \to \mathbb{R}^3$. That is, we wish to solve the optimal control problem (7.2) for some function $V : H \to \mathbb{R}$ which depends on the particular task and environment at hand.

7.1.1. Example. Consider a robotic manipulator with N revolute joints actuating mechanical limbs of lengths l_i , i = 1, ..., N. At the end, we place a single spherical joint to which the end effector is attached (for example, a robotic hand). We suppose that we track a point on the end effector of distance L from the spherical joint. Thus, the configuration space is given by $H = SO(2)^N \times S^2$.

Fix some intertial frame \mathcal{O} in \mathbb{R}^3 , and consider the family of frames \mathcal{O}_i given by translating \mathcal{O} to the center of the i^{th} revolute joint. At the origin of \mathcal{O}_i , we may choose a unit vector $v_i \in \mathbb{R}^3$ pointing in the direction of the axis fixed by the i^{th} revolute joint with respect to \mathcal{O}_i . We then embed each copy of SO(2) into SO(3) in such a way that v_i is fixed. In particular, this generates a family of closed Lie subgroups $K_i \subset$ SO(3) such that SO(2) $\cong K_i$, so that $H \cong \left(\prod_{i=1}^N K_i\right) \times S^2$. Coordinates can be chosen so that there exists a vector $x_0 \in \mathbb{R}^3$ such that the position $x \in \mathbb{R}^3$ of the end effectors is given by:

$$x = \varphi(R_1, \dots, R_N, p) = \sum_{i=1}^n l_i R_i x_0 + Lp$$

where $R_i \in K_i$ and $p \in S^2$. In general, φ is not invertible, and numerical coordinate-based approximations must be used to calculate the initial and final values for the configuration variables R_1, \ldots, R_N, p such that $\varphi(R_1, \ldots, R_N, p) = x_a, x_b$ for some given desired initial and final positions $x_a, x_b \in \mathbb{R}^3$ for the end effector. This is often referred to as the inverse kinematic problem. We do not devote any time to such an analysis, and refer the reader to [19] for more information.

Given the desired boundary data, we now seek to solve the corresponding optimal control problem. The differential kinematics for the system are given by

- (7.3) $\dot{R}_i = R_i u_i, \quad \text{for } i = 1, \dots, N$
- (7.4) $\dot{p} = p \times \omega,$

where $u_i \in TK_i$ and $\omega \in TS^2$ are the controls. For $q = (R_1, \ldots, R_N, p) \in H$ and $u = (u_1, \ldots, u_N, \omega)$, it is clear that we can write the above dynamical system in the form $\dot{q} = f(q, u)$. Moreover, it is clear that f is a diffeomorphism with inverse

(7.5)
$$f^{-1}(q, \dot{q}) = \left(R_1^T \dot{R}_1, \dots, R_N^T \dot{R}_N, \dot{p} \times p \right)$$

and satisfies (7.1).

We equip each K_i with the bi-invariant metric on SO(3) described in Example 4.1, and equip S^2 with the round metric describe in Example 6.5. The configuration space H is subsequently equipped with the product metric. The optimal control problem (7.2) then takes the form

(7.6)
$$\min_{q,u} \int_{a}^{b} \left(\sum_{i=1}^{N} \frac{1}{2} \|u_i\|_{\mathrm{SO}(3)}^{2} + \frac{1}{2} \|\omega\|_{S^2}^{2} - V(q) \right) dt$$

subject to $q \in \Omega$ and (7.3)-(7.4), where $V : H \to \mathbb{R}$. Since (7.3)-(7.4) is a decoupled system, we may separate choose a potential of the form

$$V(q) = \sum_{i=1}^{N} V_i(R_i) + V_{N+1}(p)$$

and subsequently separate (7.6) into a series of N + 1 optimal control problems for each variable. That is, with the above choice for V, (7.6) is equivalent to:

(7.7)
$$\min_{u_i,R_i} \int_a^b \left(\frac{1}{2} \|u_i\|_{SO(3)}^2 - V_i(R_i)\right) dt$$

subject to (7.3) and the relevant boundary conditions for each R_i , together with

(7.8)
$$\min_{\omega,p} \int_{a}^{b} \left(\frac{1}{2} \|\omega\|_{S^{2}}^{2} - V_{N+1}(p)\right) dt$$

subject to (7.4). Each of the above optimal control problems can then be solved according to Proposition 7.2. We now seek to design the potentials towards the end of avoiding certain configurations that are not physically realizable. For example, the end effector has limited range of motion due to mechanical constraints, and the fact that it must not collide with the mechanical limbs to which it is connected. Mathematically, this can be interpreted as an obstacle avoidance problem, where the obstacle in question is a section of the unit sphere.

In particular, suppose that $\bar{R}_i \in K_i$ and $\bar{p} \in S^2$ are some point-obstacles. As in Example 5.3, we choose the avoidance family of potentials given by:

(7.9)
$$V_i(R_i) = f_i(d_{SO(3)}(R_i, R_i); \tau_i, k_i, D_i),$$

(7.10)
$$V_{N+1}(p) = f_{N+1}(d_{S^2}(p,\bar{p});\tau_{N+1},k_{N+1},D_{N+1}),$$

where $d_{SO(3)}$ and d_{S^2} are the Riemannian distance functions on SO(3) and S^2 , respectively, τ_i, k_i, D_i are some non-negative parameters for i = 1, ..., N + 1, and for each $1 \le i \le N + 1$, $f_i : \mathbb{R} \to \mathbb{R}$ satisfy the properties outlined in Example 5.3. We may now extend the potentials and consider their gradient vector fields, as in the previous examples. In particular, we find that

$$grad_{1}(V_{i})_{ext}(I,H) = \frac{-2f_{i}'(\phi(H)^{2})\phi(H)}{\sin(\phi(H))}(H - H^{T}),$$

$$grad_{1}(\tilde{V}_{N+1})_{ext}(I,Q) = \frac{-2f_{N+1}'(\phi(Q)^{2}\sin^{2}(\theta))\phi(Q)}{\sin(\phi(Q))}\left((Q - Q^{T})^{\vee} \times e_{3}\right)^{\wedge}$$

In total, the Euler-Poincaré equations corresponding to (7.7) are given by (5.14)-(5.15) for $f = f_i$, whereas the Euler-Poincaré equations corresponding to (7.8) are given by

(7.11)
$$\dot{\Omega}_{N+1} = \frac{2f'_{N+1}\left(\phi(Q)^2 \sin^2(\theta)\right)\phi(Q)}{\sin(\phi(H))}(Q - Q^T)^{\vee} \times e_3.$$

(7.12)
$$\dot{Q} = -\dot{\Omega}Q,$$

together with the reconstruction equation $p = \bar{R}Q^T e_3$, where $\bar{R}e_3 = \bar{p}$.

REFERENCES

- C. ALTAFINI, Reduction by group symmetry of second order variational problems on a semidirect product of lie groups with positive definite riemannian metric, ESAIM: Control, Optimisation and Calculus of Variations, 10 (2004), pp. 526–548.
- [2] A. ANAHORY SIMOES, L. COLOMBO, M. DE LEÓN, J. C. MARRERO, D. M. DE DIEGO, AND E. PADRÓN, *Reduction by symmetries of contact mechanical systems on lie groups*, arXiv preprint arXiv:2306.07028, (2023).
- [3] A. M. BLOCH, L. J. COLOMBO, R. GUPTA, AND T. OHSAWA, Optimal control problems with symmetry breaking cost functions, SIAM Journal on Applied Algebra and Geometry, 1 (2017), pp. 626–646.
- W. M. BOOTHBY, An introduction to differentiable manifolds and Riemannian geometry, Revised, vol. 120, Gulf Professional Publishing, 2003.
- [5] F. BULLO AND A. D. LEWIS, Geometric control of mechanical systems: modeling, analysis, and design for simple mechanical control systems, vol. 49, Springer, 2019.
- [6] L. COLOMBO, M. G. FERNÁNDEZ, AND D. M. DE DIEGO, Variational integrators for non-autonomous lagrangian systems, Journal of Computational and Applied Mathematics, 424 (2023), p. 114966.
- [7] L. COLOMBO AND E. STRATOGLOU, Lie-poisson reduction for optimal control of left-invariant control systems with subgroup symmetry, Reports on Mathematical Physics, 91 (2023), pp. 131–141.
- [8] L. J. COLOMBO AND D. V. DIMAROGONAS, Symmetry reduction in optimal control of multiagent systems on lie groups, IEEE Transactions on Automatic Control, 65 (2020), pp. 4973–4980.
- [9] J. GALLIER AND J. QUAINTANCE, Differential Geometry and Lie Groups, vol. 13, Springer, 2020.
- [10] F. GAY-BALMAZ, D. D. HOLM, D. M. MEIER, T. S. RATIU, AND F. X. VIALARD, Invariant higher-order variational problems, Communications in Mathematical Physics, 309 (2012), pp. 413–458.
- [11] F. GAY-BALMAZ AND C. TRONCI, Reduction theory for symmetry breaking with applications to nematic systems, Physica D: Nonlinear Phenomena, 239 (2010), pp. 1929–1947.
- [12] J. R. GOODMAN AND L. J. COLOMBO, Variational obstacle avoidance with applications to interpolation problems in hybrid systems, IFAC-PapersOnLine, 54 (2021), pp. 82–87.
- [13] J. R. GOODMAN AND L. J. COLOMBO, Reduction by symmetry in obstacle avoidance problems on riemannian manifolds, arXiv preprint arXiv:2207.13574. To appear in SIAM Journal of Applied Algebra and Geometry, (2022).
- [14] D. HOLM, J. E. MARSDEN, AND T. S. RATIU, The euler-poincaré equations and semidirect products with applications to continuum theories, Advances in Mathematics, 137 (1998), pp. 1–81.

EULER-POINCARÉ REDUCTION WITH BROKEN SYMMETRIES ON RIEMANNIAN MANIFOLDS 33

- [15] D. D. HOLM, T. SCHMAH, AND C. STOICA, Geometric mechanics and symmetry: from finite to infinite dimensions, vol. 12, Oxford University Press, 2009.
- [16] J. M. LEE, Introduction to Smooth Manifolds, Springer Science + Business Media, second ed., 2013.
- [17] J. E. MARSDEN AND T. S. RATIU, Introduction to mechanics and symmetry: a basic exposition of classical mechanical systems, vol. 17, Springer Science & Business Media, 2013.
- [18] J. MILNOR, Morse Theory. (AM-51), Volume 51, Princeton university press, 2016.
- [19] R. M. MURRAY, Z. LI, AND S. S. SASTRY, A mathematical introduction to robotic manipulation, CRC press, 2017.
- [20] H. POINCARÉ, Sur une forme nouvelle des équations de la mécanique, CR Acad. Sci, 132 (1901), pp. 369– 371.
- [21] E. STRATOGLOU, A. A. SIMOES, AND L. J. COLOMBO, Reduction in optimal control with broken symmetry for collision and obstacle avoidance of multi-agent system on lie groups., Communications in Analysis & Mechanics (CAM), 15 (2023).
- [22] C. VIZMAN, Lagrangian reduction on homogeneous spaces with advected parameters, Symmetry, Integrability and Geometry: Methods and Applications, 11 (2015).