

Copulas.jl: A fully Distributions.jl-compliant copula package

Oskar Laverny ¹¶ and Santiago Jimenez ²

¹*Aix Marseille Univ, Inserm, IRD, SESSTIM, Sciences Economiques & Sociales de la Santé & Traitement de l'Information Médicale, ISSPAM, Marseille, France.*

²*Federal University of Pernambuco*

¶ *Corresponding author*

16 November 2023

Summary

Copulas are functions that describe dependence structures of random vectors, without describing their univariate marginals. In statistics, the separation is sometimes useful, the quality and/or quantity of available information on these two objects might differ. This separation can be formally stated through Sklar's theorem:

Theorem: existence and uniqueness of the copula (Sklar, 1959): For a given d -variate absolutely continuous random vector \mathbf{X} with marginals X_1, \dots, X_d , there exists a unique function C , the copula, such that

$$F(\mathbf{x}) = C(F_1(x_1), \dots, F_d(x_d)),$$

where F, F_1, \dots, F_d are respectively the distributions functions of $\mathbf{X}, X_1, \dots, X_d$.

Copulas are standard tools in probability and statistics, with a wide range of applications from biostatistics, finance or medicine, to fuzzy logic, global sensitivity and broader analysis. A few standard theoretical references on the matter are (Joe, 1997), (Nelsen, 2006), (Joe, 2014), and (Durante & Sempi, 2015).

The Julia package `Copulas.jl` brings most standard copula-related features into native Julia: random number generation, density and distribution function evaluations, fitting, construction of multivariate models through Sklar's theorem, and many more related functionalities. Copulas being fundamentally distributions of random vectors, we fully comply with the `Distributions.jl` API (Besançon et al., 2021; Lin et al., 2019), the Julian standard for implementation of random variables and random vectors. This compliance allows inter-

operability with other packages based on this API such as, e.g., [Turing.jl](#) (Ge et al., 2018) and several others.

Statement of need

The R package `copula` (Hofert et al., 2020; Ivan Kojadinovic & Jun Yan, 2010; Jun Yan, 2007; Marius Hofert & Martin Mächler, 2011) is the gold standard when it comes to sampling, estimating, or simply working around dependence structures. However, in other languages, the available tools are not as developed and/or not as recognised. We bridge the gap in the Julian ecosystem with this Julia-native implementation. Due to the very flexible type system in Julia, our code expressiveness and tidyness will increase its usability and maintainability in the long-run. Type-stability allows sampling in arbitrary precision without requiring more code, and Julia's multiple dispatch yields most of the below-described applications.

There are competing packages in Julia, such as [BivariateCopulas.jl](#) which only deals with a few models in bivariate settings but has very nice graphs, or [DatagenCopulaBased.jl](#), which only provides sampling and does not have exactly the same models as `Copulas.jl`. While not fully covering out both of these package's functionality (mostly because the three projects chose different copulas to implement), `Copulas.jl` is clearly the most fully featured, and brings, as a key feature, the compliance with the broader ecosystem.

Examples

SklarDist: sampling and fitting examples

The `Distributions.jl`'s API provides a `fit` function. You may use it to simply fit a compound model to some dataset as follows:

```
using Copulas, Distributions, Random
```

```
# Define the marginals and the copula, then use Sklar's theorem:
```

```
X1 = Gamma(2,3)
X2 = Pareto(0.5)
X3 = Binomial(10,0.8)
C = ClaytonCopula(3,0.7)
X = SklarDist(C, (X1,X2,X3))
```

```
# Sample from the model:
```

```
x = rand(D,1000)
```

```
# You may estimate the model as follows:
```

```
Dhat = fit(SklarDist{FrankCopula,Tuple{Gamma,Normal,Binomial}}, x)
```

```
# Although you'll probably get a bad fit !
```

The API does not fix the fitting procedure, and only loosely specify it, thus the implemented default might vary on the copula. If you want more control, you may turn to bayesian estimation using `Turing.jl` (Ge et al., 2018):

```
using Turing
@model function model(dataset)
  # Priors
  theta ~ TruncatedNormal(1.0, 1.0, theta, Inf)
  gamma ~ TruncatedNormal(1.0, 1.0, 0.25, Inf)
  eta ~ Beta(1,1)
  delta ~ Exponential(1)

  # Define the model through Sklar's theorem:
  X1 = Gamma(2,theta)
  X2 = Pareto(gamma)
  X3 = Binomial(10,eta)
  C = ClaytonCopula(3,delta)
  X = SklarDist(C, (X1,X2,X3))

  # Add the loglikelihood to the model :
  Turing.Turing.@addlogprob! loglikelihood(D, dataset)
end
```

The Archimedean API

Archimedean copulas are a huge family of copulas that has seen a lot of theoretical work. Among others, you may take a look at (McNeil & Nešlehová, 2009). We use `WilliamsonTransformations.jl`'s implementation of the Williamson d -transform to sample from any archimedean copula, including for example the `ClaytonCopula` with negative dependence parameter in any dimension, which is a first to our knowledge.

The API is consisting of the following functions:

```
phi(C::MyArchimedean, t) # Generator
williamson_dist(C::MyArchimedean) # Williamson d-transform
```

So that implementing your own archimedean copula only requires to subset the `ArchimedeanCopula` type and provide your generator as follows:

```
struct MyUnknownArchimedean{d,T} <: ArchimedeanCopula{d}
  theta::T
end
phi(C::MyUnknownArchimedean,t) = exp(-t*C.theta)
```

The obtained model can be used as follows:

```
C = MyUnknownCopula{2,Float64}(3.0)
spl = rand(C,1000) # sampling
```

```

cdf(C, spl)           # cdf
pdf(C, spl)           # pdf
loglikelihood(C, spl) # llh

```

The following functions have defaults but can be overridden for performance:

```

phi_inv(C::MyArchimedean, t) # Inverse of phi
phi_1(C::MyArchimedean, t) # first derivative of phi
phi_d(C::MyArchimedean, t) # dth derivative of phi
tau(C::MyArchimedean) # Kendall tau
tau_inv(::Type{MyArchimedean}, tau) = # Inverse kendall tau
fit(::Type{MyArchimedean}, data) # fitting.

```

Broader ecosystem

The package is starting to get used in several other places of the ecosystem. Among others, we noted:

- The package `GlobalSensitivity.jl` exploit `Copulas.jl` to provide Shapley effects implementation, see [this documentation](#).
- `EconomicScenarioGenerators.jl` uses dependence structures between financial assets.

Acknowledgments

References

- Besançon, M., Papamarkou, T., Anthoff, D., Arslan, A., Byrne, S., Lin, D., & Pearson, J. (2021). Distributions.jl: Definition and modeling of probability distributions in the JuliaStats ecosystem. *Journal of Statistical Software*, 98(16), 1–30. <https://doi.org/10.18637/jss.v098.i16>
- Durante, F., & Sempi, C. (2015). *Principles of copula theory*. Chapman and Hall/CRC. <https://doi.org/10.1201/b18674>
- Ge, H., Xu, K., & Ghahramani, Z. (2018). Turing: A language for flexible probabilistic inference. *International Conference on Artificial Intelligence and Statistics, AISTATS 2018, 9-11 April 2018, Playa Blanca, Lanzarote, Canary Islands, Spain*, 1682–1690. <http://proceedings.mlr.press/v84/ge18b.html>
- Hofert, M., Kojadinovic, I., Maechler, M., & Yan, J. (2020). *Copula: Multivariate dependence with copulas*. <https://CRAN.R-project.org/package=copula>
- Ivan Kojadinovic, & Jun Yan. (2010). Modeling multivariate distributions with continuous margins using the copula R package. *Journal of Statistical Software*, 34(9), 1–20. <https://doi.org/10.18637/jss.v034.i09>
- Joe, H. (1997). *Multivariate models and multivariate dependence concepts*. CRC press. <https://doi.org/10.1201/9780367803896>
- Joe, H. (2014). *Dependence modeling with copulas*. CRC press.
- Jun Yan. (2007). Enjoy the joy of copulas: With a package copula. *Journal of Statistical Software*, 21(4), 1–21. <https://doi.org/10.18637/jss.v021.i04>

- Lin, D., White, J. M., Byrne, S., Bates, D., Noack, A., Pearson, J., Arslan, A., Squire, K., Anthoff, D., Papamarkou, T., Besançon, M., Drugowitsch, J., Schauer, M., & contributors, other. (2019). *JuliaStats/Distributions.jl: a Julia package for probability distributions and associated functions*. <https://doi.org/10.5281/zenodo.2647458>
- Marius Hofert, & Martin Mächler. (2011). Nested archimedean copulas meet R: The nacopula package. *Journal of Statistical Software*, 39(9), 1–20. <https://doi.org/10.18637/jss.v039.i09>
- McNeil, A. J., & Nešlehová, J. (2009). Multivariate Archimedean copulas, d-monotone functions and L1-norm symmetric distributions. *The Annals of Statistics*, 37(5B), 3059–3097. <https://doi.org/10.1214/07-AOS556>
- Nelsen, R. B. (2006). *An introduction to copulas* (2nd ed). Springer. ISBN: 978-0-387-28659-4
- Sklar, A. (1959). Fonction de répartition dont les marges sont données. *Inst. Stat. Univ. Paris*, 8, 229–231.