

# ON INTERNAL CATEGORIES AND CROSSED OBJECTS IN THE CATEGORY OF MONOIDS

ILIA PIRASHVILI

ABSTRACT. It is a well-known fact that the category  $\mathbf{Cat}(\mathbf{C})$  of internal categories in a category  $\mathbf{C}$  has a description in terms of crossed modules, when  $\mathbf{C} = \mathbf{Gr}$  is the category of groups. The proof of this result heavily uses the fact that any split epimorphism decomposes as a semi-direct product. An equivalent statement does not hold in the category  $\mathbf{Mon}$  of monoids. In a previous work on quadratic algebras, [3] I constructed an internal category in the category of monoids, see Section 6. Based on this construction, this paper will introduce the notion of a crossed semi-bimodule and show that it gives rise to an object in  $\mathbf{Cat}(\mathbf{Mon})$ . I will also relate this new notion to the crossed semi-modules introduced earlier by A. Patchkoria [2].

## 1. INTRODUCTION

The notion of a crossed module was introduced by Whitehead in his study of homotopy theory [4]. A crossed module is a group homomorphism  $\partial : K \rightarrow A$ , together with an action of  $A$  on the group  $K$ , satisfying some conditions. Starting with such data, one can construct the category  $\mathbf{Cat}(\partial : K \rightarrow A)$ . Objects are elements of  $A$ , while morphisms from  $b$  to  $a$  are essentially elements  $x$  of  $K$ , such that  $b = \partial(x)a$  and the composition is induced by the multiplication in  $K$ .

One can readily check that  $\mathbf{Cat}(\partial : K \rightarrow A)$  is an internal category in the category of groups, meaning that all structural maps are group homomorphisms. It is well-known that in this way, one obtains an isomorphism of the category of internal categories in the category of groups, and the category of crossed modules [1].

This paper will introduce the notion of a crossed semi-bimodule, which is a pair of monoids  $A, K$ , and a triple of maps  $A \times K \xrightarrow{\circ} A$ ,  $A \times K \xrightarrow{\lambda} K$ ,  $K \times A \xrightarrow{\rho} K$ , satisfying certain conditions. We will show that one can construct a category  $\mathbf{Cat}(A, K, \circ)$  from such a data, which is internal in the category of monoids.

We will show that crossed semi-bimodules for which  $\lambda$  is a trivial homomorphism, is equivalent to the category of crossed semi-modules

introduced by A. Patchkoria [2]. We will also show that in the cases when both monoids  $A$  and  $K$  are groups, our notion is essentially equivalent to a crossed module.

In the last section, we will describe a particular crossed module, which arose during my work on quadratic étale algebras [3]. It is an example which does not come from crossed semi-modules.

## 2. CROSSED (SEMI-)MODULES AND CROSSED SEMI-BIMODULES

**2.1. Internal categories.** Let  $\mathbf{A}$  be a category with finite limits. Recall that an internal category  $\mathbf{C}$  in  $\mathbf{A}$  is the following data:

- i) A pair of objects  $C_0$  and  $C_1$  of  $\mathbf{A}$ , called, respectively, the *object of objects* and *object of morphisms* of  $\mathbf{C}$ .
- ii) Four morphisms  $d_0^1, d_1^1 : C_1 \rightarrow C_0$ ,  $s_0^0 : C_0 \rightarrow C_1$  and  $d_1^2 : C_2 \rightarrow C_1$ , called, respectively, the *target*, *source*, *unit* and *composite*. Here,  $C_2$  is the following pullback:

$$\begin{array}{ccc} C_2 & \xrightarrow{d_2^2} & C_1 \\ d_0^2 \downarrow & & \downarrow d_0^1 \\ C_1 & \xrightarrow{d_1^1} & C_0. \end{array}$$

The morphisms  $d_0^1, d_1^1, d_1^2$  and  $s_0^0$  satisfy the standard simplicial identities.

**2.2. Crossed modules and internal categories in  $\mathbf{Gr}$ .** Let  $\mathbf{Gr}$  be the category of groups and group homomorphisms. It was known since the 60's that the internal categories in  $\mathbf{Gr}$  are equivalent to crossed modules. Crossed modules were introduced by Whitehead [4] in the study of homotopy theory.

**Definition 2.1.** A crossed module is a group homomorphism  $\partial : K \rightarrow A$ , together with a right action of the group  $A$  on the group  $K$ , denoted by  $x^a$ , where  $x \in K$ ,  $a \in A$ . It must satisfy the following equalities

- (1)  $\partial(x^a) = a^{-1}\partial(x)a$ ,
- (2)  $y^{\partial x} = x^{-1}yx$

for all  $x, y \in K$  and  $a \in A$ .

Denote by  $\mathbf{Xmod}$  the category of crossed modules. From this, we can construct an internal category in  $\mathbf{Gr}$  in the following way:

Let  $(K, A, \partial)$  be a crossed module. Define  $C_0 = A$  and  $C_1$  to be  $A \times K$  as a set. The group structure is defined by

$$(a, x)(b, y) = (ab, x^b y).$$

This construction is compare with [1].

The group  $C_2$  is  $A \times K \times K$  as a set with the following operation

$$(a, x, y)(b, u, v) = (ab, x^b u, y^{b\partial u} v).$$

Next, we put

$$s_0^0(a) = (a, 1), \quad d_i^1(a, x) = \begin{cases} a, & \text{if } i = 0 \\ a\partial(x), & \text{if } i = 1 \end{cases}$$

$$s_i^1(a, x) = \begin{cases} (a, x, 1) & \text{if } i = 0 \\ (a, 1, x) & \text{if } i = 1 \end{cases}$$

$$d_i^2(a, x, y) = \begin{cases} (a, x) & \text{if } i = 0 \\ (a, xy) & \text{if } i = 1 \\ (a\partial(x), y) & \text{if } i = 2 \end{cases}$$

$$s_i^1(a, x) = \begin{cases} (a, x, 1) & \text{if } i = 0 \\ (a, 1, x) & \text{if } i = 1. \end{cases}$$

One checks that all maps are group homomorphisms and in this way, obtains an internal category in **Gr**. This is denoted by  $\text{Cat}(K \xrightarrow{\partial} A)$ . It is well-known that any internal category in **Gr** is of this form, and the functor

$$\mathbf{Xmod} \rightarrow \text{Cat}(\mathbf{Gr}), \quad (\mathbf{K}, \mathbf{A}, \partial) \mapsto \text{Cat}(\mathbf{K} \xrightarrow{\partial} \mathbf{A})$$

is an equivalence of categories [1].

**2.3. Crossed semi-modules.** A. Patchkoria [2] introduced the notion of a crossed semi-module and proved that the category of crossed semi-modules is equivalent to the category of Schreier categorical objects in the category of monoids.

**Definition 2.2.** *A crossed semi-module [2] is a monoid homomorphism  $\partial : K \rightarrow A$ , together with a right action of the monoid  $A$  on the monoid  $K$ , denoted by  $x^a$ ,  $x \in K$ ,  $a \in A$ . It must satisfy the following equalities*

- i)  $a\partial(x^a) = \partial(x)a$ ,
- ii)  $yx^{\partial y} = xy$

for all  $a \in A$ ,  $x, y \in K$ .

By defining morphisms in the analogous way to groups, we can form a category, which we will denote by **Xsmod**.

Starting from a crossed semi-module, one can construct an internal category in **Mon**, the category of monoids, in the same exact way as the crossed modules.

### 3. CROSSED SEMI-BIMODULES AND INTERNAL CATEGORIES IN THE CATEGORY OF MONOIDS

**3.1. Definition and first consequences.** The following definitions are probably more appropriate for monoids than crossed semi-modules.

**Definition 3.1.** *A crossed semi-bimodule consists of monoids  $A$  and  $K$ , together with maps*

$$A \times K \xrightarrow{\circ} A; \quad (a, x) \mapsto a \circ x,$$

$$A \times K \xrightarrow{\lambda} K; \quad (a, x) \mapsto {}^a x,$$

$$K \times A \xrightarrow{\rho} K; \quad (x, a) \mapsto x^a.$$

The first map makes  $A$  into a right  $K$ -set, the second map makes  $K$  into a left  $A$ -monoid and the third map makes  $K$  into a right  $A$ -monoid. Moreover, the following conditions need to hold for every  $x, y \in K$  and  $a, b \in A$ :

- (1)  $({}^a x)^b = {}^a(x^b)$ ,
- (2)  $(ab) \circ ({}^a x) = a(b \circ x)$ ,
- (3)  $(ab) \circ (x^b) = (a \circ x)b$ ,
- (4)  $({}^a y)x^{b \circ y} = x^b({}^{a \circ x} y)$ .

For simplicity, instead of saying that  $(K, A, \circ, \lambda, \rho)$  is a crossed semi-bimodule, we will say that  $(K, A, \circ)$  is a crossed semi-bimodule

**Definition 3.2.** *A morphism from a crossed semi-bimodule  $(K, A, \circ)$  to a crossed semi-bimodule  $(K', A', \circ)$  is given by the homomorphisms  $\kappa : K \rightarrow K'$  and  $\alpha : A \rightarrow A'$ , for which the following conditions hold*

- (1)  $\alpha(a \circ x) = \alpha(a) \circ \kappa(x)$ ,
- (2)  $\kappa({}^a x) = \alpha(a)\kappa(x)$ ,
- (3)  $\kappa(x^a) = \kappa(x)^{\alpha(a)}$ .

Denote the category of crossed semi-bimodules by  $\mathbf{Xbsmod}$ . We will need also a weaker version.

**Definition 3.3.** *A weak morphism from a crossed semi-bimodule  $(K, A, \circ)$  to a crossed semi-bimodule  $(K', A', \circ)$  is given by a monoid homomorphism  $\kappa : A \rightarrow A'$  and by the map  $\gamma : A \times K \rightarrow K'$ , such that the following conditions hold*

- (1)  $\gamma(a, xy) = \gamma(a, x)\gamma(a, y)$ ,
- (2)  $\kappa(a) \circ \gamma(a, x) = \kappa(a \circ x)$ ,
- (3)  $\kappa(a)\gamma(b, y)\gamma(a, x)^{\kappa(a) \circ \gamma(b, y)} = \gamma(ab, {}^a y x^{b \circ y})$ .

Any morphism  $(\kappa, \alpha)$  gives rise to a weak morphism  $(\kappa, \gamma)$ , where  $\gamma(a, x) = \alpha(x)$ . Weak morphisms can be composed as follows

$$(\kappa', \gamma') \circ (\kappa, \gamma) = (\kappa' \circ \kappa, \gamma''),$$

where  $\gamma''(a, x) = \gamma'(\kappa(a), \gamma(a, x))$ . Thus, we obtain the category  $\mathbf{Xbsmod}_w$ . Isomorphisms in  $\mathbf{Xbsmod}_w$  are called weak isomorphisms.

Recall that a *semi-bimodule* over  $A$  is a commutative monoid  $K$ , together with left and right monoid actions of  $A$  on  $K$ , satisfying Equality i) of Definition 3.1.

**Lemma 3.4.** *The category of semi-bimodule over a monoid  $A$  is equivalent to the full subcategory of crossed semi-bimodules over  $A$ , for which*

$$a \circ x = a$$

*holds for all  $a \in A$  and all  $x \in K$ .*

The proof is just a straightforward verification.

**Lemma 3.5.** *Let  $(K, A, \circ)$  be a crossed semi-bimodule. Define*

$$\partial : K \rightarrow A$$

*by  $\partial(y) = 1 \circ y$ . For any  $x, y \in K$ , one has*

$$yx^{\partial(y)} = x^{\partial(x)}y.$$

*Proof.* Substitute  $a = 1 = b$  in Identity iv) of the definition of a crossed semi-bimodule 3.1 to obtain  $yx^{1 \circ y} = x^{1 \circ x}y$ , and the result follows.  $\square$

### 3.2. Crossed semi-bimodules and internal categories in $\mathbf{Mon}$ .

**Proposition 3.6.** *Let  $(K, A, \circ)$  be a crossed semi-bimodule.*

i) *Define the multiplication  $\bullet$  on  $A \times K$  by*

$$(a, x) \bullet (b, y) = (ab, {}^a y x^{b \circ y}).$$

*This defines a monoid structure on the set  $A \times K$ . This monoid is denoted by  $A \bowtie K$ .*

ii) *Consider the set  $A \times K \times K$  with the multiplication*

$$(a, x, y) \bullet (b, u, v) = (ab, {}^a u x^{b \circ u}, {}^{a \circ u} v y^{b \circ u \circ v}).$$

*This induces a monoid structure denoted by  $A \bowtie K \bowtie K$ .*

iii) *Define the maps*

$$A \bowtie K \bowtie K \begin{array}{c} \xrightarrow{d_0^2} \\ \xrightarrow{d_0^2} \\ \xrightarrow{d_0^2} \end{array} A \bowtie K \begin{array}{c} \xrightarrow{d_0^1} \\ \xrightarrow{d_0^1} \\ \xrightarrow{d_0^1} \end{array} A; \quad A \xrightarrow{s_0^0} A \bowtie K \begin{array}{c} \xrightarrow{s_0^1} \\ \xrightarrow{s_1^1} \\ \xrightarrow{s_1^1} \end{array} A \bowtie K \bowtie K$$

*by*

$$s_0^0(a) = (a, 1), \quad s_0^1(a, x) = (a, x, 1), \quad s_1^1(a, x) = (a, 1, x).$$

$$d_0^1(a, x) = a, \quad d_1^1(a, x) = a \circ x,$$

$$d_0^2(a, x, y) = (a, x), \quad d_1^2(a, x, y) = (a, xy), \quad d_2^2(a, x, y) = (a \circ x, y).$$

These maps satisfy all the simplicial identities and are homomorphisms.

iv) The diagram

$$\begin{array}{ccc} A \bowtie K \bowtie K & \xrightarrow{d_2^2} & A \bowtie K \\ d_0^2 \downarrow & & \downarrow d_0^1 \\ A \bowtie K & \xrightarrow{d_1^1} & A \end{array}$$

is a pullback diagram.

v) The above defines an internal category, denoted  $\mathbf{Cat}(A, K, \circ)$ , in the category  $\mathbf{Mon}$  of monoids.

*Proof.* i) It is clear that  $(1, 1)$  serves as the unit of  $\bullet$ . For associativity, we have

$$\begin{aligned} ((a, x)(b, y))(c, z) &= (ab, {}^a y x^{b \circ y})(c, z) \\ &= (abc, {}^{ab} z ({}^a y x^{b \circ y})^{c \circ z}) \\ &= (abc, {}^{ab} z ({}^a y)^{c \circ z} x^{(b \circ y)(c \circ z)}). \end{aligned}$$

On the other hand

$$\begin{aligned} (a, x)((b, y)(c, z)) &= (a, x)(bc, {}^b z y^{c \circ z}) \\ &= (abc, {}^{ab} z ({}^a y)^{c \circ z} x^{(bc) \circ ({}^b z y^{c \circ z})}). \end{aligned}$$

Thus, we need to check the identity

$$(1) \quad (b \circ y)(c \circ z) = (bc) \circ ({}^b z y^{c \circ z}).$$

Equalities (2) and (3) of Definition 3.1  $a = b \circ y, b = c, x = z$  yield

$$\begin{aligned} (b \circ y)(c \circ z) &= ((b \circ y)c) \circ ({}^{b \circ y} z) \\ &= (bc) \circ y^c \circ ({}^{b \circ y} z). \end{aligned}$$

Equality (4) of the same Definition now gives Identity (1).

ii) This follows from parts iii) and iv).

iii) We will only check that the maps  $d_1^1, d_1^2, d_2^2$  are homomorphisms. The rest is clear.

**First claim:**  $d_1^1$  is a homomorphism. Thanks to Identity (1), we have

$$\begin{aligned} d_1^1(a, x)d_1^1(b, y) &= (a \circ x)(b \circ y) \\ &= (ab) \circ ({}^a y x^{b \circ y}) \\ &= d_1^1((a, x) \bullet (b, y)). \end{aligned}$$

**Second claim:**  $d_1^2$  is a homomorphism. We have

$$\begin{aligned} d_1^2((a, x, y) \bullet (b, u, v)) &= d_1^2(ab, {}^a u x^{bou}, {}^{aou} v y^{bouov}) \\ &= (ab, {}^a u x^{bou} {}^{aou} v y^{bouov}). \end{aligned}$$

On the other hand

$$\begin{aligned} d_1^2(a, x, y) \bullet d_1^2(b, u, v) &= (a, xy) \bullet (b, uv) \\ &= (ab, {}^a (uv)(xy)^{bouv}) \\ &= (ab, {}^a u {}^a v x^{bou} y^{bou}). \end{aligned}$$

Since  $\circ$  is an action, we have  $b \circ (uv) = b \circ u \circ v$ , and based on Equality (4) of the definition of a crossed semi-bimodule 3.1, we conclude that these expressions are equal.

**Third claim:**  $d_2^2$  is a homomorphism. As for the first claim, we have

$$\begin{aligned} d_2^2((a, x, y) \bullet (b, u, v)) &= d_2^2(ab, {}^a u x^{bou}, {}^{aou} v y^{bouov}) \\ &= ((a \circ x)(b \circ u, {}^{aou} v y^{bouov}) \\ &= (a \circ x, y) \bullet (b \circ u, v) \\ &= d_2^2(a, x, y) \bullet d_2^2(b, u, v). \end{aligned}$$

iv) Take two elements  $(a, x)$  and  $(b, y)$  in  $A \bowtie K$  such that  $d_1^1(a, x) = d_1^1(b, y)$ . Equivalently  $a \circ x = b$ . Thus, there exists a unique element in  $A \bowtie K \bowtie K$ , namely  $(a, x, y)$ , such that  $d_0^2(a, x, y) = (a, x)$  and  $d_2^2(a, x, y) = (b, y)$ . The result follows.

v) This is a consequence of iii) and iv).  $\square$

It is clear that if  $(\kappa, \alpha)$  is a morphism of crossed semi-bimodules, it defines an internal functor  $\mathbf{Cat}(K, A, \circ) \rightarrow \mathbf{Cat}(K', A', \circ)$ . In this way, we obtain a functor  $\mathbf{Xbsmod} \rightarrow \mathbf{Cat}(\mathbf{Mon})$ , from the category of crossed semi-bimodules to the category of internal categories in the category of monoids.

It is less obvious, but readily checked, that any weak morphism  $(\kappa, \gamma)$  from  $(K, A, \circ) \rightarrow (K', A', \circ)$  also defines an internal functor  $\mathbf{Cat}(K, A, \circ) \rightarrow \mathbf{Cat}(K', A', \circ)$ . On object objects, it is given by  $\kappa$ . On morphism objects, it is defined by  $(a, x) \mapsto (\kappa(a), \gamma(a, x))$ . On the object of composable morphisms, the internal functor is given by  $(a, x, y) \mapsto (\kappa(a), \gamma(a, x), \gamma(a, y))$ . This gives us a functor

$$\mathbf{Xbsmod}_w \rightarrow \mathbf{Cat}(\mathbf{Mon}).$$

**Lemma 3.7.** *Let  $(K, A, \circ)$  be a crossed semi-bimodule. Define a new operation  $\diamond$  on  $K$  by*

$$x \diamond y := yx^{\partial(y)}.$$

Let  $K^{tw}$  be the set  $K$ , equipped with the operation  $\diamond$ . Then  $K^{tw}$  is a monoid.

*Proof.* By comparing  $\diamond$  to the multiplication on  $A \bowtie K$ , we see that

$$(1, x) \bullet (1, y) = (1, x \diamond y).$$

It follows that  $\diamond$  is associative and  $1 \diamond x = x = x \diamond 1$ .  $\square$

#### 4. RELATIONSHIP BETWEEN CROSSED SEMI-MODULES AND CROSSED SEMI-BIMODULES

The aim of this section is to show that crossed semi-modules are exactly the crossed semi-bimodules for which  ${}^a x = x$ , for every  $a \in A$  and every  $x \in K$ .

**Proposition 4.1.** i) Let  $\partial : K \rightarrow A$  be a crossed semi-module. For elements  $a \in M$  and  $x \in G$ , define

$${}^a x = x, \quad a \circ x = a\partial(x).$$

This yields obtains a crossed semi-bimodule structure, denoted by  $\Phi(K, A, \partial)$ .

ii) Conversely, assume we have a crossed semi-bimodule for which  ${}^a x = a$ , for all  $a \in A$  and  $x \in K$ . Define  $\partial : K \rightarrow A$  by

$$\partial(x) = 1 \circ x.$$

We obtain a crossed semi-module.

iii) The constriction in i) defines a full and faithful functor  $\Phi : \mathbf{Xsmod} \rightarrow \mathbf{Xbsmod}$  from the category of crossed semi-modules to the category of crossed semi-bimodules. Moreover, it induces an isomorphism of  $\mathbf{Xsmod}$  to the full subcategory of crossed semi-bimodules for which  ${}^a x = x$ , for all  $a \in A$  and  $x \in K$ .

*Proof.* i) Equality (1) of Definition 3.1 is obvious. We have

$$(ab) \circ ({}^a x) = (ab) \circ (x) = ab\partial(x) = a(b \circ x),$$

which proves Equality (2) of Definition 3.1. We also have

$$(ab) \circ (x^b) = ab\partial(x^b) = a\partial(x)b = (a \circ x)b,$$

which gives us Equality (3). Finally, we have

$$({}^a y)x^{b \circ y} = yx^{b\partial(y)} = x^b y = x^b ({}^{a \circ x} y),$$

and part i) follows.

ii) Assume  ${}^a x = x$ . By putting  $b = 1$  in Identity (2) of the definition of a crossed semi-bimodule Definition 3.1, we obtain

$$a \circ x = a\partial(x).$$

By the definition of  $\partial$  and setting  $a = \partial(x)$ , we get

$$\partial(xy) = 1 \circ (xy) = (1 \circ x) \circ y = \partial(x) \circ y = \partial(x)\partial(y).$$

Hence,  $\partial$  is a monoid homomorphism. It remains to show that the defining relations of crossed semi-modules are satisfied. To this end, we put  $m = 1$  in Identity (3) of Definition 3.1 to obtain

$$b\partial(x^b) = b \circ x^b = (1 \circ x)b = \partial(x)b.$$

Quite similarly, taking  $b = 1$  in Identity (4) of Definition 3.1 gives us

$$yx^{\partial(y)} = xy,$$

and part ii) proved.

iii) By our construction, in any crossed semi-bimodule coming from  $\mathbf{Xsmod}$ , we have  $\partial(x) = 1 \circ x$ . This implies that  $\Phi$  is a full and faithful functor. The rest follows from part ii).  $\square$

## 5. THE CASE WHEN $K$ AND $A$ ARE GROUPS

Recall that if  $(K, A, \circ)$  is a crossed semi-bimodule, we have a map

$$\partial : K \rightarrow A$$

defined by  $\partial(x) = 1 \circ x$ .

In this section, we will investigate crossed semi-bimodules  $(K, A, \circ)$ , such that  $K$  and  $A$  are groups. Our first observation is that the action  $A \times K \xrightarrow{\circ} A$  can be reconstructed from the map  $\partial$ .

**Lemma 5.1.** *Let  $(K, A, \circ)$  be a crossed semi-bimodule such that  $K$  and  $A$  are groups. For all  $x, z \in K$  and  $a \in A$ , one has*

- i)  $a \circ x = a\partial(a^{-1}x)$ ,
- ii)  $\partial(a^{-1}z^a) = a^{-1}\partial(z)a$ ,
- iii)  $\partial(xz^{\partial(x)}) = \partial(z)\partial(x)$ .

*Proof.* i) Putting  $x = a^{-1}t$  and  $b = 1$  in Equality ii) of Definition 3.1, we obtain  $a \circ t = a\partial(a^{-1}t)$ .

ii) Having in mind i), Identity 3 of Definition 3.1 can be rewritten as

$$ab\partial((ab)^{-1}x^b) = a\partial(a^{-1}x)b.$$

If we cancel both sides by  $a$  and replace  $x$  by  $a^{-1}y$ , we obtain

$$b\partial(b^{-1}y^b) = \partial(y)b,$$

and the result follows.

iii) Since  $\circ$  is an action of  $K$  on  $A$ , we have

$$a \circ (xy) = (a \circ x) \circ y.$$

Take  $a = 1$  and use Identity i) to rewrite the last equality as

$$(2) \quad \partial(xy) = \partial(x)\partial((\partial(x))^{-1}y).$$

Take  $y = z^{\partial(x)}$  and use part ii) to obtain

$$\begin{aligned} \partial(xz^{\partial(x)}) &= \partial(x)\partial(\partial(x)^{-1}z^{\partial(x)}) \\ &= \partial(x)(\partial(x))^{-1}\partial(z)\partial(x) \\ &= \partial(z)\partial(x). \end{aligned}$$

□

**Proposition 5.2.** *Let  $K$  and  $A$  be groups. Assume there are given maps*

$$A \times K \xrightarrow{\lambda} K; \quad (a, x) \mapsto {}^a x,$$

$$K \times A \xrightarrow{\rho} K; \quad (x, a) \mapsto x^a$$

*such that the first map makes  $K$  into a left  $A$ -monoid, the second map makes  $K$  into a right  $A$ -monoid and these actions are compatible. That is to say*

$$({}^a x)^b = {}^a(x^b), \quad a, b \in A, x \in K.$$

*Additionally, assume there is given a map  $\partial : K \rightarrow A$ , such that the following identities hold:*

- i)  $\partial(xy) = \partial(x)\partial(\partial(x)^{-1}y)$
- ii)  $\partial({}^{b^{-1}}z^b) = b^{-1}\partial(z)b$
- iii)  $yx^{\partial(y)} = x^{\partial(x)}y.$

*Define the map  $A \times K \xrightarrow{\circ} A$  by  $a \circ x = a\partial(a^{-1}x)$ . This gives us a crossed semi-bimodule. Moreover, any crossed semi-bimodule for which  $K$  and  $A$  are groups can be obtained in this manner.*

*Proof.* By putting  $x = 1 = y$  in i), we see that  $\partial(1) = 1$ . Next, we show that  $\circ$  defines a right action of  $K$  on  $A$ . Indeed, we have  $a \circ 1 = a\partial(a^{-1}1) = a$ . We also have

$$a \circ (xy) = a\partial(a^{-1}x^{\partial(x)}y).$$

Putting  $u = a^{-1}x$  and  $v = a^{-1}y$ , the previous equality takes on the form  $a \circ (xy) = a\partial(uv)$ . Using Identity i) yields

$$a \circ (xy) = a\partial(u)\partial\left(\partial(u)^{-1}v\right).$$

On the other hand

$$\begin{aligned}
(a \circ x) \circ y &= (a\partial^{(a^{-1}x)}) \circ y \\
&= (a\partial(u)) \circ y \\
&= a\partial(u)\partial\left(\partial^{(u)^{-1}a^{-1}}y\right) \\
&= a\partial(u)\partial\left(\partial^{(u)^{-1}}v\right).
\end{aligned}$$

Comparing these expressions we see that

$$a \circ (xy) = (a \circ x) \circ y.$$

Thus,  $\circ$  defines the action of  $K$  on  $A$ . It remains to check the identities (ii)-(iv) of Definition 3.1. We have

$$\begin{aligned}
(ab) \circ ({}^ax) &= ab\partial^{(b^{-1}a^{-1}ax)} \\
&= ab\partial^{(b^{-1}x)} \\
&= a(b \circ x).
\end{aligned}$$

This proves Identity (2) of the Definition of a corssed semi-bimodule 3.1. We also have

$$\begin{aligned}
(ab) \circ x^b &= ab\partial^{(b^{-1}a^{-1}x^b)} \\
&= abb^{-1}\partial^{(a^{-1}x)}b \\
&= a\partial^{(a^{-1}x)}b \\
&= (a \circ x)b.
\end{aligned}$$

and Identity (3) of Definition 3.1 follows. Here, we used Identity ii). To prove Identity (4) of Definition 3.1, we first show it in the particular case when  $a = 1$ . Setting  $z = y^{b^{-1}}$  gives us

$$\begin{aligned}
yx^{b \circ y} &= yx^{b\partial^{(b^{-1}y)}} \\
&= z^b x^{b\partial^{(b^{-1}z^b)}} \\
&= z^b x^{\partial(z)b} \\
&= (zx^{\partial(z)})^b.
\end{aligned}$$

Using Identity iii) now gives us

$$\begin{aligned}
yx^{b \circ y} &= (x^{(\partial x z)})^b \\
&= x^b \partial(x) z^b \\
&= x^b \partial(x) y.
\end{aligned}$$

For the general case, put  $t = a^{-1}x$ . Thus,  $x = {}^a t$  and we have

$$\begin{aligned} {}^a y x^{b \circ y} &= {}^a y ({}^a t^{b \circ y}) \\ &= {}^a (y t^{b \circ y}). \end{aligned}$$

The case  $a = 1$  now gives us

$$\begin{aligned} {}^a y x^{b \circ y} &= {}^a (t^{b \partial(t)} y) \\ &= x^b {}^{a \partial(t)} y \\ &= x^b {}^{a \partial(a^{-1}x)} y \\ &= x^b ({}^{a \circ x} y). \end{aligned}$$

This proves Identity (3) of Definition 3.1.  $\square$

**Proposition 5.3.** *Let  $(K, A, \circ)$  be a crossed semi-bimodule such that  $K$  and  $A$  are groups. Recall that we defined a monoid  $K^{st}$  in Lemma 3.7, which, as a set, is  $K$ . The monoid structure  $\diamond$ , however, is given by*

$$x \diamond y := y x^{\partial(y)} = x^{\partial(x)} y,$$

where  $\partial(y) = 1 \circ y$ , as usual.

- i) The map  $\partial : K^{tw} \rightarrow A$  is a homomorphism.
- ii) Define a new action of  $A$  on the  $K^{st}$  by

$$x^{*a} = a^{-1} x^a,$$

This gives us

$$\partial(x^{*a}) = a^{-1} \partial(x) a.$$

- iii) The monoid  $K^{tw}$  is, indeed, a group.
- iv) We have

$$y \diamond x^{*\partial y} = x \diamond y.$$

*Proof.* i) We have

$$\begin{aligned} \partial(x \diamond y) &= \partial(y x^{\partial(y)}) \\ &= \partial(x) \partial(y), \end{aligned}$$

according to part iii) of Lemma 5.1.

- ii) This is a restatement of part ii) of Lemma 5.1.
- iii) Put

$$x^{\flat} = \partial(x)^{-1} (x^{-1})$$

We have

$$\begin{aligned} x \diamond x^{\flat} &= x(\partial(x)(x^{\flat})) \\ &= x(x^{-1}) \\ &= 1. \end{aligned}$$

Thus, the  $\diamond$ -inverse of  $x$  is  $x^b$ .

iv) We have

$$\begin{aligned} y \diamond x^{*\partial y} &= y \diamond (\partial y^{-1} x^{\partial(y)}) \\ &= y^{\partial(y)} (\partial y^{-1} x^{\partial(y)}) \\ &= y x^{\partial(y)}. \end{aligned}$$

□

**Corollary 5.4.** *Let  $(K, A, \circ)$  be a crossed semi-bimodule such that  $K$  and  $A$  are groups. Then  $(K^{tw}, A, \partial)$  is a crossed module.*

*Proof.* We already proved that  $K^{tw}$  is a group and  $\partial$  is a homomorphism, see Proposition 5.3. The group  $A$  acts on  $K$  via  $x^{*a} = {}^{a^{-1}}x^a$ . This action respects the  $\diamond$  operation. Indeed, we have

$$\begin{aligned} (x \diamond y)^{*a} &= (yx^{\partial(y)})^{*a} \\ &= ({}^{a^{-1}}y^a) ({}^{a^{-1}}(x^{\partial y})^a). \end{aligned}$$

We also have

$$\begin{aligned} (x^{*a}) \diamond (y^{*a}) &= ({}^{a^{-1}}x^a) \diamond ({}^{a^{-1}}y^a) \\ &= ({}^{a^{-1}}y^a) ({}^{a^{-1}}x^a)^{\partial({}^{a^{-1}}y^a)} \\ &= ({}^{a^{-1}}y^a) ({}^{a^{-1}}x^{\partial(y)a}). \end{aligned}$$

Comparing these expressions, we see that  $(x \diamond y)^{*a} = (x^{*a}) \diamond (y^{*a})$ . Finally Identities i) and ii) of Definition 2.1 hold, thanks to parts ii) and iv) of Proposition 5.3. □

We constructed a functor  $\mathbf{Xmod} \rightarrow \mathbf{Xbsmod}_{\mathbf{Gr}}$  in Lemma 3.4, whose image lies in the subcategory  $\mathbf{Xbsmod}_{\mathbf{Gr}}$  whose objects are such crossed semi-bimodules, for which both  $K$  and  $A$  are groups. Moreover, in Proposition 5.3, we constructed the functor  $\mathbf{Xbsmod}_{\mathbf{Gr}} \rightarrow \mathbf{Xmod}$  in the reverse direction. It is easy to check that the composite

$$\mathbf{Xmod} \rightarrow \mathbf{Xbsmod}_{\mathbf{Gr}} \rightarrow \mathbf{Xmod}$$

is isomorphic to the identity functor. The next Lemma will show that, though the composite

$$\mathbf{Xbsmod}_{\mathbf{Gr}} \rightarrow \mathbf{Xmod} \rightarrow \mathbf{Xbsmod}_{\mathbf{Gr}}$$

is in general not isomorphic to the identity, it is weakly isomorphic to it.

**Proposition 5.5.** *Let  $(K, A, \circ)$  be a crossed semi-bimodule such that  $K$  and  $A$  are groups. Denote by  $(K^{tw}, A, \partial)$  the crossed semi-bimodule*

corresponding to the crossed module  $(K^{tw}, A, \circ)$ , constructed in Proposition 5.3.

The maps  $\text{ld} : A \rightarrow A$  and  $\gamma : A \times K^{tw} \rightarrow K$  define a weak isomorphism  $(K^{tw}, A, \partial) \rightarrow (K, A, \partial)$ , where  $\gamma(a, x) = {}^a x$ .

*Proof.* The fact that these maps define a weak morphism is a straightforward computation. It is an isomorphism as the inverse morphism is given by  $\text{ld} : A \rightarrow A$  and  $\gamma^{[-1]} : A \times K \rightarrow K^{tw}$ , where the latter is defined by  $\gamma^{[-1]}(a, x) = a^{-1}x$ .  $\square$

## 6. A CONSTRUCTION RELATED TO RANK TWO COMMUTATIVE ALGEBRAS

The notion of crossed semi-bimodules arose during my work on rank two commutative algebras [3]. In more detail, let  $R$  be a commutative ring and  $p, q \in R$ , such that

$$pq + 2 = 0.$$

We define a crossed semi-bimodule, denoted by  $\text{Qu}(R) := (\mathbf{K}(R), \mathbf{A}(R), \circ)$ . Here,  $\mathbf{K}(R)$  is the multiplicative monoid of  $2 \times 2$  matrices of the form

$$\begin{pmatrix} 1 & r \\ 0 & s \end{pmatrix},$$

where  $r, s \in R$ . The monoid  $\mathbf{A}(R)$ , as a set, is simply  $R^2$ . Elements of  $\mathbf{A}(R)$  are denoted by  $[a, b]$ . The monoid operation is defined by

$$[a, b][c, d] = [ac, a^2d + bc^2 + p^2bd].$$

One easily sees that in this way, we obtain a commutative monoid with  $[1, 0]$  as the unit.

**Lemma 6.1.** *For elements*

$$k = \begin{pmatrix} 1 & r \\ 0 & s \end{pmatrix} \in \mathbf{K}(R) \quad \text{and} \quad m = [a, b] \in \mathbf{A}(R),$$

we set

$$m \circ k = [as - pr, s^2b - qr sa - r^2].$$

This defines an action of the monoid  $\mathbf{K}(R)$  on the set  $\mathbf{A}(R)$ .

*Proof.* Taking  $r = 0$  and  $s = 1$ , we see that  $m \circ 1 = m$ . Next, take

$$k_1 = \begin{pmatrix} 1 & r_1 \\ 0 & s_1 \end{pmatrix}. \quad \text{We have}$$

$$\begin{aligned} (m \circ k) \circ k_1 &= [as - pr, s^2b - qr sa - r^2] \circ \begin{pmatrix} 1 & r_1 \\ 0 & s_1 \end{pmatrix} \\ &= [(as - pr)s_1 - pr_1, s_1^2(s^2b - qr sa - r^2) - qr_1s_1(as - pr) - r_1^2]. \end{aligned}$$

On the other, hand we have

$$\begin{aligned} m \circ (kk_1) &= [a, b] \circ \begin{pmatrix} 1 & r_1 + rs_1 \\ 0 & ss_1 \end{pmatrix} \\ &= [ass_1 - p(r_1 + rs_1), s^2s_1^2b - q(r_1 + rs_1)ss_1a - (r_1 + rs_1)^2]. \end{aligned}$$

Comparing these expression and using the fact that  $pq = -2$ , we see that

$$(m \circ k) \circ k_1 = m \circ (kk_1),$$

implying the lemma.  $\square$

**Proposition 6.2.** *For elements*

$$k = \begin{pmatrix} 1 & r \\ 0 & s \end{pmatrix} \in \mathbf{K}(R) \quad \text{and} \quad m = [a, b] \in \mathbf{A}(R)$$

we set

$$k^m = \begin{pmatrix} 1 & ra \\ 0 & s \end{pmatrix} = {}^m k.$$

In this way, we obtain a crossed semi-bimodule  $(\mathbf{K}(R), \mathbf{A}(R), \circ)$ .

*Proof.* For  $n = [c, d]$ , we have

$$\begin{aligned} (mn) \circ ({}^m k) &= [ac, a^2d + bc^2 + p^2bd] \circ \begin{pmatrix} 1 & ra \\ 0 & s \end{pmatrix} \\ &= [acs - pra, s^2(a^2d + bc^2 + p^2bd) - qarsac - r^2a^2] \end{aligned}$$

On the other hand

$$\begin{aligned} m(n \circ k) &= [a, b][cs - pr, s^2d - qrsc - r^2] \\ &= [acs - par, a^2(s^2d - qrsc - r^2) + b(cs - pr)^2 + p^2b(s^2d - qrsc - r^2)]. \end{aligned}$$

Since  $pq = -2$ , we obtain

$$(mn) \circ ({}^m s) = m(n \circ s).$$

For

$$h = \begin{pmatrix} 1 & u \\ 0 & v \end{pmatrix} \in K(R),$$

we have

$$\begin{aligned} ({}^m h)k^{n \circ h} &= \begin{pmatrix} 1 & ua \\ 0 & v \end{pmatrix} \begin{pmatrix} 1 & crv - pur \\ 0 & s \end{pmatrix} \\ &= \begin{pmatrix} 1 & crv - pur + asu \\ 0 & sv \end{pmatrix} \\ &= \begin{pmatrix} 1 & rc \\ 0 & s \end{pmatrix} \begin{pmatrix} 1 & asu - pru \\ 0 & v \end{pmatrix} \\ &= k^n ({}^{m \circ k} h) \end{aligned}$$

□

The internal category in **Mon**, corresponding to  $(\mathbf{K}(R), \mathbf{A}(R), \circ)$ , is intimately related to the Galois algebras over the Hopf  $R$ -algebra  $J^{pq}$ , thanks to [3]. The latter is freely generated by 1 and  $x$  as an  $R$ -module. The multiplication and comultiplication in  $J^{pq}$  is given by

$$x^2 = qx, \quad \Delta(x) = 1 \otimes x + x \otimes 1 + p(x \otimes x).$$

#### REFERENCES

- [1] S. MACLANE. Categories for working mathematician. Springer (2nd edition), 1998.
- [2] A. PATCHKORIA. Crossed semi-modules and Schreier internal categories in the category of monoids. Georgian Mathematical Journal. 5 (1998) 575-581.
- [3] I. PIRASHVILI. On the group of quadratic algebras and stacks. Glasgow Mathematical Journal, 61(2018) 151-167.
- [4] J.H.C. WHITEHEAD. Combinatorial homotopy II. Bull. AMS, 55(1949), 453–496