

# Nonrigid Surface Registration: Gradient Calculations

Dániel Unyi

Department of Telecommunications and Media Informatics  
 Budapest University of Technology and Economics  
 Műegyetem rkp. 3., H-1111 Budapest, Hungary

## 1 Rigid registration

The goal of rigid registration is to align a source surface  $\mathcal{X}$  to a target surface  $\mathcal{Y}$ . The alignment process involves iteratively transforming  $\mathcal{X}$  closer and closer to  $\mathcal{Y}$ , such that  $\mathcal{X} = \mathcal{Z}^0 \rightarrow \mathcal{Z}^1 \rightarrow \mathcal{Z}^2 \rightarrow \dots \rightarrow \mathcal{Z}^T = \mathcal{Y}$ . In other words, in each iteration we seek a rotation matrix  $\tilde{\mathbf{R}}$  and a translation vector  $\tilde{\mathbf{t}}$  such that  $\mathcal{X}^{t+1} = \tilde{\mathbf{R}}(\mathbf{R}^t \mathcal{X} + \mathbf{t}^t) + \tilde{\mathbf{t}} = \tilde{\mathbf{R}}\mathcal{X}^t + \tilde{\mathbf{t}}$ , which corresponds to minimizing the error function

$$w_1 \sum_{i=1}^N \|\Pi_{\mathcal{Y}}(\mathbf{z}_i^{t+1}) - \mathbf{z}_i^{t+1}\|_2^2 + w_2 \sum_{i=1}^N \|\tilde{\mathbf{R}}\mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2$$

$\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$  and  $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_n$  are sets of points sampled from the surfaces  $\mathcal{X}$  and  $\mathcal{Z}$ , and  $\Pi_{\mathcal{Y}}(\mathbf{z}_i)$  denotes the closest point on  $\mathcal{Y}$  to the point  $\mathbf{z}_i$ .  $w_1$  and  $w_2$  are simple weighting factors. By applying two simplifications, we can trace the minimization problem back to solving a system of linear equations. First, there is nonlinearity in  $\Pi_{\mathcal{Y}}(\mathbf{z}_i^{t+1})$ , so we use the previous estimate  $\Pi_{\mathcal{Y}}(\mathbf{z}_i^t)$  which is constant<sup>1</sup>. Second, there is also nonlinearity in  $\tilde{\mathbf{R}}$ . Assuming small rotations, we can linearize the rotation matrices using  $\cos(\cdot) \approx 1$  and  $\sin(\cdot) \approx 0$ :

$$\begin{aligned} \tilde{\mathbf{R}} &= \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \gamma & -\sin \gamma \\ 0 & \sin \gamma & \cos \gamma \end{bmatrix} \\ \tilde{\mathbf{R}} &\approx \begin{bmatrix} 1 & -\gamma & \beta \\ \gamma & 1 & -\alpha \\ -\beta & \alpha & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & -\gamma & \beta \\ \gamma & 0 & -\alpha \\ -\beta & \alpha & 0 \end{bmatrix} \end{aligned}$$

By introducing  $\tilde{\mathbf{r}} = [\alpha \ \beta \ \gamma]$ , we can substitute  $\tilde{\mathbf{R}}$  with  $\mathbf{I} + \tilde{\mathbf{r}} \times$ . Adding Tikhonov regularization might be beneficial in ensuring that the rotations remain small.

---

<sup>1</sup>For efficient closest point lookup, a k-d tree has to be built in advance.

## 1.1 Gradient calculations

We have to minimize the following function with respect to  $\tilde{\mathbf{r}}$ ,  $\tilde{\mathbf{t}}$ , and  $\mathbf{z}_j^{t+1}$ :

$$\arg \min_{\tilde{\mathbf{r}}, \tilde{\mathbf{t}}, \mathbf{z}_j^{t+1}} w_1 \sum_{i=1}^N \|\Pi_{\mathcal{Y}}(\mathbf{z}_i^t) - \mathbf{z}_i^{t+1}\|_2^2 + w_2 \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2$$

- Minimizing w.r.t.  $\tilde{\mathbf{r}}$ :

$$\begin{aligned} \frac{\partial}{\partial \tilde{\mathbf{r}}} \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2 &= 0 \\ \sum_{i=1}^N \mathbf{x}_i^t \times (\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}) &= \mathbf{0} \\ - \sum_{i=1}^N \mathbf{X}_i^t \mathbf{X}_i^t \tilde{\mathbf{r}} + \sum_{i=1}^N \mathbf{X}_i^t \tilde{\mathbf{t}} - \sum_{i=1}^N \mathbf{X}_i^t \mathbf{z}_i^{t+1} &= \mathbf{0} \end{aligned}$$

- Minimizing w.r.t.  $\tilde{\mathbf{t}}$ :

$$\begin{aligned} \frac{\partial}{\partial \tilde{\mathbf{t}}} \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2 &= 0 \\ \sum_{i=1}^N \mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1} &= \mathbf{0} \\ - \sum_{i=1}^N \mathbf{X}_i^t \tilde{\mathbf{r}} + N \tilde{\mathbf{t}} - \sum_{i=1}^N \mathbf{z}_i^{t+1} &= - \sum_{i=1}^N \mathbf{x}_i^t \end{aligned}$$

- Minimizing w.r.t.  $\mathbf{z}_j^{t+1}$ :

$$\begin{aligned} \frac{\partial}{\partial \mathbf{z}_j^{t+1}} (w_1 \sum_{i=1}^N \|\Pi_{\mathcal{Y}}(\mathbf{z}_i^t) - \mathbf{z}_i^{t+1}\|_2^2 + w_2 \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2) &= 0 \\ \frac{w_1}{w_2} (\mathbf{z}_j^{t+1} - \Pi_{\mathcal{Y}}(\mathbf{z}_j^t)) + \mathbf{z}_j^{t+1} - \mathbf{x}_j^t - \tilde{\mathbf{r}} \times \mathbf{x}_j^t - \tilde{\mathbf{t}} &= \mathbf{0} \\ \mathbf{X}_j^t \tilde{\mathbf{r}} - \tilde{\mathbf{t}} + \left(1 + \frac{w_1}{w_2}\right) \mathbf{z}_j^{t+1} &= \frac{w_1}{w_2} \Pi_{\mathcal{Y}}(\mathbf{z}_j^t) + \mathbf{x}_j^t \end{aligned}$$

The result is a system of  $6 + 3N$  linear equations with the same number of unknowns.

$$\mathbf{x} = \begin{bmatrix} \tilde{\mathbf{r}} \\ \tilde{\mathbf{t}} \\ \mathbf{z}_1^{t+1} \\ \mathbf{z}_2^{t+1} \\ \vdots \\ \mathbf{z}_N^{t+1} \end{bmatrix}$$

$$\mathbf{b} = \begin{bmatrix} \mathbf{0} \\ -\sum_{i=1}^N \mathbf{x}_i^t \\ \frac{w_1}{w_2} \Pi_{\mathcal{Y}}(\mathbf{z}_1^t) + \mathbf{x}_1^t \\ \frac{w_1}{w_2} \Pi_{\mathcal{Y}}(\mathbf{z}_2^t) + \mathbf{x}_2^t \\ \vdots \\ \frac{w_1}{w_2} \Pi_{\mathcal{Y}}(\mathbf{z}_N^t) + \mathbf{x}_N^t \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_{rr} & \mathbf{A}_{tr} & \mathbf{A}_{z_1r} & \mathbf{A}_{z_2r} & \dots & \mathbf{A}_{z_Nr} \\ \mathbf{A}_{rt} & \mathbf{A}_{tt} & \mathbf{A}_{z_1t} & \mathbf{A}_{z_2t} & \dots & \mathbf{A}_{z_Nt} \\ \mathbf{A}_{rz_1} & \mathbf{A}_{tz_1} & \mathbf{A}_{z_1z_1} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{A}_{rz_2} & \mathbf{A}_{tz_2} & \mathbf{0} & \mathbf{A}_{z_2z_2} & \dots & \mathbf{0} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{A}_{rz_N} & \mathbf{A}_{tz_N} & \mathbf{0} & \mathbf{0} & \dots & \mathbf{A}_{z_Nz_N} \end{bmatrix}$$

where

$$\mathbf{A}_{rr} = - \sum_{i=1}^N \mathbf{X}_i^t \mathbf{X}_i^t$$

$$\mathbf{A}_{tr} = \sum_{i=1}^N \mathbf{X}_i^t$$

$$\mathbf{A}_{z_i r} = - [\mathbf{X}_1^t \quad \mathbf{X}_2^t \quad \dots \quad \mathbf{X}_N^t]$$

$$\mathbf{A}_{rt} = - \sum_{i=1}^N \mathbf{X}_i^t$$

$$\mathbf{A}_{tt} = N\mathbf{I}$$

$$\mathbf{A}_{z_i t} = - [\mathbf{I} \quad \mathbf{I} \quad \dots \quad \mathbf{I}]$$

$$\mathbf{A}_{rz_j} = \begin{bmatrix} \mathbf{X}_1^t \\ \mathbf{X}_2^t \\ \vdots \\ \mathbf{X}_N^t \end{bmatrix}$$

$$\mathbf{A}_{tz_j} = - \begin{bmatrix} \mathbf{I} \\ \mathbf{I} \\ \vdots \\ \mathbf{I} \end{bmatrix}$$

$$\mathbf{A}_{z_i z_j} = \left(1 + \frac{w_1}{w_2}\right) \begin{bmatrix} \mathbf{I} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{I} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{I} \end{bmatrix}$$

## 2 As-rigid-as-possible registration

In each iteration we also compute the local rotation matrices  $\tilde{\mathbf{R}}_i$ :

$$w_1 \sum_{i=1}^N \|\Pi_{\mathcal{Y}}(\mathbf{z}_i^{t+1}) - \mathbf{z}_i^{t+1}\|_2^2 + w_2 \sum_{i=1}^N \|\tilde{\mathbf{R}} \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2 + w_3 \sum_{i=1}^N \sum_{k \in \mathcal{N}(i)} \|\tilde{\mathbf{R}}_i(\mathbf{x}_k^t - \mathbf{x}_i^t) - (\mathbf{z}_k^{t+1} - \mathbf{z}_i^{t+1})\|_2^2$$

We apply the same linearization to the local rotation matrices as we do to the global rotation matrix. Connectivity has to be defined among the points of the source surface, resulting a graph (or a mesh specifically) with graph Laplacian  $\mathbf{L}$ . The neighbours of point  $i$  are denoted by  $\mathcal{N}(i)$ .

## 2.1 Gradient calculations

We have to minimize the following function with respect to  $\tilde{\mathbf{r}}$ ,  $\tilde{\mathbf{t}}$ ,  $\tilde{\mathbf{r}}_j$ , and  $\mathbf{z}_j^{t+1}$ :

$$\begin{aligned} \arg \min_{\tilde{\mathbf{r}}, \tilde{\mathbf{t}}, \tilde{\mathbf{r}}_j, \mathbf{z}_j^{t+1}} & w_1 \sum_{i=1}^N \|\Pi_{\mathcal{Y}}(\mathbf{z}_i^{t+1}) - \mathbf{z}_i^{t+1}\|_2^2 + w_2 \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2 + \\ & + w_3 \sum_{i=1}^N \sum_{k \in \mathcal{N}(i)} \|(\mathbf{x}_k^t - \mathbf{x}_i^t) + \tilde{\mathbf{r}}_i \times (\mathbf{x}_k^t - \mathbf{x}_i^t) - (\mathbf{z}_k^{t+1} - \mathbf{z}_i^{t+1})\|_2^2 \end{aligned}$$

- Minimizing w.r.t.  $\tilde{\mathbf{r}}$ :

$$\begin{aligned} \frac{\partial}{\partial \tilde{\mathbf{r}}} \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2 &= 0 \\ \sum_{i=1}^N \mathbf{x}_i^t \times (\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}) &= \mathbf{0} \\ - \sum_{i=1}^N \mathbf{X}_i^t \mathbf{X}_i^t \tilde{\mathbf{r}} + \sum_{i=1}^N \mathbf{X}_i^t \tilde{\mathbf{t}} - \sum_{i=1}^N \mathbf{X}_i^t \mathbf{z}_i^{t+1} &= \mathbf{0} \end{aligned}$$

- Minimizing w.r.t.  $\tilde{\mathbf{t}}$ :

$$\begin{aligned} \frac{\partial}{\partial \tilde{\mathbf{t}}} \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2 &= 0 \\ \sum_{i=1}^N \mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1} &= \mathbf{0} \\ - \sum_{i=1}^N \mathbf{X}_i^t \tilde{\mathbf{r}} + N \tilde{\mathbf{t}} - \sum_{i=1}^N \mathbf{z}_i^{t+1} &= - \sum_{i=1}^N \mathbf{x}_i^t \end{aligned}$$

- Minimizing w.r.t.  $\tilde{\mathbf{r}}_j$ :

$$\begin{aligned} \frac{\partial}{\partial \tilde{\mathbf{r}}_j} \sum_{i=1}^N \sum_{k \in \mathcal{N}(i)} \|(\mathbf{x}_k^t - \mathbf{x}_i^t) + \tilde{\mathbf{r}}_i \times (\mathbf{x}_k^t - \mathbf{x}_i^t) - (\mathbf{z}_k^{t+1} - \mathbf{z}_i^{t+1})\|_2^2 &= 0 \\ \sum_{k \in \mathcal{N}(j)} (\mathbf{x}_k^t - \mathbf{x}_j^t) \times ((\mathbf{x}_k^t - \mathbf{x}_j^t) + \tilde{\mathbf{r}}_j \times (\mathbf{x}_k^t - \mathbf{x}_j^t) - (\mathbf{z}_k^{t+1} - \mathbf{z}_j^{t+1})) &= \mathbf{0} \\ - \sum_{k \in \mathcal{N}(j)} (\mathbf{X}_k^t - \mathbf{X}_j^t)(\mathbf{X}_k^t - \mathbf{X}_j^t) \tilde{\mathbf{r}}_j + \sum_{k \in \mathcal{N}(j)} (\mathbf{X}_k^t - \mathbf{X}_j^t)(\mathbf{z}_j^{t+1} - \mathbf{z}_k^{t+1}) &= \mathbf{0} \end{aligned}$$

- Minimizing w.r.t.  $\mathbf{z}_j^{t+1}$ :

$$\begin{aligned} & \frac{\partial}{\partial \mathbf{z}_j^{t+1}} \left( w_1 \sum_{i=1}^N \|\Pi_{\mathcal{Y}}(\mathbf{z}_i^t) - \mathbf{z}_i^{t+1}\|_2^2 + w_2 \sum_{i=1}^N \|\mathbf{x}_i^t + \tilde{\mathbf{r}} \times \mathbf{x}_i^t + \tilde{\mathbf{t}} - \mathbf{z}_i^{t+1}\|_2^2 + \right. \\ & \quad \left. + w_3 \sum_{i=1}^N \sum_{k \in \mathcal{N}(i)} \|(\mathbf{x}_k^t - \mathbf{x}_j^t) + \tilde{\mathbf{r}}_i \times (\mathbf{x}_k^t - \mathbf{x}_j^t) - (\mathbf{z}_k^{t+1} - \mathbf{z}_j^{t+1})\|_2^2 \right) = 0 \end{aligned}$$

$$\begin{aligned} & w_1(\mathbf{z}_j^{t+1} - \Pi_{\mathcal{Y}}(\mathbf{z}_j^t)) + w_2(\mathbf{z}_j^{t+1} - \mathbf{x}_j^t - \tilde{\mathbf{r}} \times \mathbf{x}_j^t - \tilde{\mathbf{t}}) + \\ & + w_3 \sum_{k \in \mathcal{N}(j)} 2(\mathbf{x}_k^t - \mathbf{x}_j^t) + (\tilde{\mathbf{r}}_k + \tilde{\mathbf{r}}_j) \times (\mathbf{x}_k^t - \mathbf{x}_j^t) - 2(\mathbf{z}_k^{t+1} - \mathbf{z}_j^{t+1}) = \mathbf{0} \end{aligned}$$

$$\begin{aligned} & (w_1 + w_2)\mathbf{z}_j^{t+1} - 2w_3 \sum_{k \in \mathcal{N}(j)} (\mathbf{z}_k^{t+1} - \mathbf{z}_j^{t+1}) + w_2 \mathbf{X}_j^t \tilde{\mathbf{r}} - w_2 \tilde{\mathbf{t}} - w_3 \sum_{k \in \mathcal{N}(j)} (\mathbf{X}_k^t - \mathbf{X}_j^t)(\tilde{\mathbf{r}}_j + \tilde{\mathbf{r}}_k) = \\ & = w_1 \Pi_{\mathcal{Y}}(\mathbf{z}_j^t) + w_2 \mathbf{x}_j^t - 2w_3 \sum_{k \in \mathcal{N}(j)} (\mathbf{x}_k^t - \mathbf{x}_j^t) \end{aligned}$$

The result is a system of  $6 + 6N$  linear equations with the same number of unknowns.

$$\begin{aligned} \mathbf{x} &= \begin{bmatrix} \tilde{\mathbf{r}} \\ \tilde{\mathbf{t}} \\ \tilde{\mathbf{r}}_1^{t+1} \\ \tilde{\mathbf{r}}_2^{t+1} \\ \vdots \\ \tilde{\mathbf{r}}_N^{t+1} \\ \mathbf{z}_1^{t+1} \\ \mathbf{z}_2^{t+1} \\ \vdots \\ \mathbf{z}_N^{t+1} \end{bmatrix} \\ \mathbf{b} &= \begin{bmatrix} \mathbf{0} \\ -\sum_{i=1}^N \mathbf{x}_i^t \\ \mathbf{0} \\ \mathbf{0} \\ \vdots \\ \mathbf{0} \\ w_1 \Pi_{\mathcal{Y}}(\mathbf{z}_1^t) + w_2 \mathbf{x}_1^t + 2w_3 \sum_{k=1}^N \mathbf{L}_{1k} \mathbf{x}_k^t \\ w_1 \Pi_{\mathcal{Y}}(\mathbf{z}_2^t) + w_2 \mathbf{x}_2^t + 2w_3 \sum_{k=1}^N \mathbf{L}_{2k} \mathbf{x}_k^t \\ \vdots \\ w_1 \Pi_{\mathcal{Y}}(\mathbf{z}_N^t) + w_2 \mathbf{x}_N^t + 2w_3 \sum_{k=1}^N \mathbf{L}_{Nk} \mathbf{x}_k^t \end{bmatrix} \end{aligned}$$

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_{rr} & \mathbf{A}_{tr} & \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{A}_{z_1r} & \mathbf{A}_{z_2r} & \dots & \mathbf{A}_{z_Nr} \\ \mathbf{A}_{rt} & \mathbf{A}_{tt} & \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{A}_{z_1t} & \mathbf{A}_{z_2t} & \dots & \mathbf{A}_{z_Nt} \\ \mathbf{0} & \mathbf{0} & \mathbf{A}_{r_1r_1} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{A}_{z_1r_1} & \mathbf{A}_{z_2r_1} & \dots & \mathbf{A}_{z_Nr_1} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{A}_{r_2r_2} & \dots & \mathbf{0} & \mathbf{A}_{z_1r_2} & \mathbf{A}_{z_2r_2} & \dots & \mathbf{A}_{z_Nr_2} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \dots & \mathbf{A}_{r_Nr_N} & \mathbf{A}_{z_1r_N} & \mathbf{A}_{z_2r_N} & \dots & \mathbf{A}_{z_Nr_N} \\ \mathbf{A}_{rz_1} & \mathbf{A}_{tz_1} & \mathbf{A}_{r_1z_1} & \mathbf{A}_{r_2z_1} & \dots & \mathbf{A}_{r_Nz_1} & \mathbf{A}_{z_1z_1} & \mathbf{A}_{z_2z_1} & \dots & \mathbf{A}_{z_Nz_1} \\ \mathbf{A}_{rz_2} & \mathbf{A}_{tz_2} & \mathbf{A}_{r_1z_2} & \mathbf{A}_{r_2z_2} & \dots & \mathbf{A}_{r_Nz_2} & \mathbf{A}_{z_1z_2} & \mathbf{A}_{z_2z_2} & \dots & \mathbf{A}_{z_Nz_2} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{A}_{rz_N} & \mathbf{A}_{tz_N} & \mathbf{A}_{r_1z_N} & \mathbf{A}_{r_2z_N} & \dots & \mathbf{A}_{r_Nz_N} & \mathbf{A}_{z_1z_N} & \mathbf{A}_{z_2z_N} & \dots & \mathbf{A}_{z_Nz_N} \end{bmatrix}$$

where

$$\mathbf{A}_{rr} = - \sum_{i=1}^N \mathbf{X}_i^t \mathbf{X}_i^t$$

$$\mathbf{A}_{tr} = \sum_{i=1}^N \mathbf{X}_i^t$$

$$\mathbf{A}_{z_i r} = - [\mathbf{X}_1^t \quad \mathbf{X}_2^t \quad \dots \quad \mathbf{X}_N^t]$$

$$\mathbf{A}_{rt} = - \sum_{i=1}^N \mathbf{X}_i^t$$

$$\mathbf{A}_{tt} = N\mathbf{I}$$

$$\mathbf{A}_{z_i t} = - [\mathbf{I} \quad \mathbf{I} \quad \dots \quad \mathbf{I}]$$

$$\mathbf{A}_{r_i r_j} = - \begin{bmatrix} \sum_{k=1}^N \mathbf{L}_{1k} (\mathbf{X}_1^t - \mathbf{X}_k^t)^2 & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \sum_{k=1}^N \mathbf{L}_{2k} (\mathbf{X}_2^t - \mathbf{X}_k^t)^2 & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \sum_{k=1}^N \mathbf{L}_{Nk} (\mathbf{X}_N^t - \mathbf{X}_k^t)^2 \end{bmatrix}$$

$$\mathbf{A}_{z_i r_j} = - \begin{bmatrix} - \sum_{k=1}^N \mathbf{L}_{1k} (\mathbf{X}_1^t - \mathbf{X}_k^t) & \mathbf{L}_{12} (\mathbf{X}_1 - \mathbf{X}_2) & \dots & \mathbf{L}_{1N} (\mathbf{X}_1 - \mathbf{X}_N) \\ \mathbf{L}_{21} (\mathbf{X}_2 - \mathbf{X}_1) & - \sum_{k=1}^N \mathbf{L}_{2k} (\mathbf{X}_2^t - \mathbf{X}_k^t) & \dots & \mathbf{L}_{2N} (\mathbf{X}_2 - \mathbf{X}_N) \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{L}_{N1} (\mathbf{X}_N - \mathbf{X}_1) & \mathbf{L}_{N2} (\mathbf{X}_N - \mathbf{X}_2) & \dots & - \sum_{k=1}^N \mathbf{L}_{Nk} (\mathbf{X}_N^t - \mathbf{X}_k^t) \end{bmatrix}$$

$$\mathbf{A}_{rz_j} = w_2 \begin{bmatrix} \mathbf{X}_1^t \\ \mathbf{X}_2^t \\ \vdots \\ \mathbf{X}_N^t \end{bmatrix}$$

$$\mathbf{A}_{tz_j} = -w_2 \begin{bmatrix} \mathbf{I} \\ \mathbf{I} \\ \vdots \\ \mathbf{I} \end{bmatrix}$$

$$\mathbf{A}_{r_i z_j} = -w_3 \begin{bmatrix} \sum_{k=1}^N \mathbf{L}_{1k} (\mathbf{X}_1^t - \mathbf{X}_k^t) & \mathbf{L}_{12} (\mathbf{X}_1 - \mathbf{X}_2) & \dots & \mathbf{L}_{1N} (\mathbf{X}_1 - \mathbf{X}_N) \\ \mathbf{L}_{21} (\mathbf{X}_2 - \mathbf{X}_1) & \sum_{k=1}^N \mathbf{L}_{2k} (\mathbf{X}_2^t - \mathbf{X}_k^t) & \dots & \mathbf{L}_{2N} (\mathbf{X}_2 - \mathbf{X}_N) \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{L}_{N1} (\mathbf{X}_N - \mathbf{X}_1) & \mathbf{L}_{N2} (\mathbf{X}_N - \mathbf{X}_2) & \dots & \sum_{k=1}^N \mathbf{L}_{Nk} (\mathbf{X}_N^t - \mathbf{X}_k^t) \end{bmatrix}$$

$$\mathbf{A}_{z_i z_j} = (w_1 + w_2) \begin{bmatrix} \mathbf{I} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{I} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{I} \end{bmatrix} + 2w_3 \begin{bmatrix} \mathbf{L}_{11}\mathbf{I} & \mathbf{L}_{12}\mathbf{I} & \dots & \mathbf{L}_{1N}\mathbf{I} \\ \mathbf{L}_{21}\mathbf{I} & \mathbf{L}_{22}\mathbf{I} & \dots & \mathbf{L}_{2N}\mathbf{I} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{L}_{N1}\mathbf{I} & \mathbf{L}_{N2}\mathbf{I} & \dots & \mathbf{L}_{NN}\mathbf{I} \end{bmatrix}$$

### 3 Accelerating the convergence: point-to-plane error

In this case, we also measure the distance from the tangent plane belonging to the closest point, adding an extra term with weighting factor  $w_4$ :

$$w_4(\mathbf{n}_i^T(\Pi_{\mathcal{Y}}(\mathbf{z}_i^t) - \mathbf{z}_i^{t+1}))^2$$

where  $\mathbf{n}_i$  is the surface normal at point  $i$ . The gradient of this term w.r.t  $\mathbf{z}_j^{t+1}$  is

$$2w_4(\mathbf{n}_i \circ \mathbf{n}_i)(\mathbf{z}_i^{t+1} - \Pi_{\mathcal{Y}}(\mathbf{z}_i^t))^2$$

### References

1. Bouaziz, S., Tagliasacchi, A., Li, H., & Pauly, M. (2016). Modern techniques and applications for real-time non-rigid registration. In SIGGRAPH ASIA 2016 Courses (pp. 1-25).
2. Sorkine, O., & Alexa, M. (2007, July). As-rigid-as-possible surface modeling. In Symposium on Geometry processing (Vol. 4, pp. 109-116).
3. Low, K. L. (2004). Linear least-squares optimization for point-to-plane icp surface registration. Chapel Hill, University of North Carolina, 4(10), 1-3.