## Investigation of hidden symmetry in  $AdS_5$  black hole with Heun equation

Jafar Sadeghi $*1$  $*1$ , Mohammad Ali S. Afshar  $*^2$  $*^2$ , , Mohammad Reza Alipour  $*3$  $*3$ , Saeed Noori Gashti\*, $\frac{1}{4}$  $\frac{1}{4}$  $\frac{1}{4}$ 

> <sup>⋆</sup>Department of Physics, Faculty of Basic Sciences, University of Mazandaran P. O. Box 47416-95447, Babolsar, Iran

†School of Physics, Damghan University, P. O. Box 3671641167, Damghan, Iran

#### Abstract

Investigating the existence of algebra and finding hidden symmetries in physical systems is one of the most important aspects for understanding their behavior and predicting their future. Expanding this unique method of study to cosmic structures and combining past knowledge with new data can be very interesting and lead to discovering new ways to analyze these systems. However, studying black hole symmetries always presents many complications and sometimes requires computational approximations. For example, checking the existence of Killing vectors and then calculating them is not always an easy task. It becomes much more difficult as the structure and geometry of the system become more complex. In this work, we will show that if the wave equations with a black hole background can be converted in the form of general Heun equation, based on its structure and coefficients, the algebra of the system can be easily studied, and computational and geometrical complications can be omitted. For this purpose, we selected two  $AdS_5$  black holes: Reissner-Nordström (R-N) and Kerr, and analyzed the Klein-Gordon equation with the background of these black holes. Based on this concept, we observed that the radial part of the R-N black hole and both the radial and angular parts of the Kerr black hole could be transformed into the general form of the Heun equation. As a result, according to the algebraic structure that governs the Heun equation and its coefficients, one can easily achieve generalized  $sl(2)$  algebra.

Keywords: $AdS_5$  Black Holes; Klein–Gordon Particle; Generalized  $sl(2)$  Algebra.

<sup>&</sup>lt;sup>1</sup>Email: pouriya@ipm.ir<br><sup>2</sup>Email: m.a.s.afshar@gr

<span id="page-0-0"></span><sup>2</sup>Email: m.a.s.afshar@gmail.com

<span id="page-0-1"></span><sup>3</sup>Email: mr.alipour@stu.umz.ac.ir

<span id="page-0-3"></span><span id="page-0-2"></span><sup>4</sup>Email: saeed.noorigashti@stu.umz.ac.ir

## Contents



## <span id="page-1-0"></span>1 Introduction

It must be admitted that throughout the history of physics, conservation laws are the most basic tool of a physicist to study a dynamic and playful universe. Undoubtedly, the first step in studying such a system is to check for the existence of constants and stability within it. In this way, the existence of "symmetries" and "periodicity" is the most important data that these systems provide, hidden from the effects of time and without any changes. When symmetry and periodicity are raised, the forced relationship of system elements together (algebra) and the geometric rules of this repetition (topology) play a more fundamental role.

Sometimes this role even becomes so powerful that it dominates the basic cognitive tools of a physicist and, as a result, makes decisions about the physics of the system based only on geometry, topology, and algebra, regardless of conventional methods. The meaning of these words is more tangible for particle physicists and cosmologists because we are not dealing with a crystal where everything can be tested and verified.

This is a fact! All we have is a black hole in infinity, and we are so far away that all our seemingly advanced measuring tools seem practically unusable or useless. This is where the value of long-range measurement tools such as algebra, which can reproduce system generators based on inductive reasoning, geometry, and comparison of symmetric behaviors, is determined. Then, with the help of the golden key of Noter's theorem, the relevant constants and currents will be easily available to us so that we can safely make decisions about the behaviors and results of the system.

But there are two ways to use the algebraic method: first of all, like a mathematician, regardless of whether the results are meaningful or meaningless from a physical point of view, consider the black hole as a mathematical entity and be satisfied only with obtaining logical results. Or be ourselves, a physicist that prioritizes conceptualization and subjectivity and use algebra only as a study tool. We have chosen the second approach in this article.

Also the study of hidden symmetries in black holes is not a new concept and is often based on analyzing the wave field equation with a black hole background and then identifying the Killing vectors and the Killing-Yano tensor. This method has been used in the basic work of Castro, Mallory, and Strominger for the Kerr-Newman black hole [\[1\]](#page-15-0), as well as in subsequent works such as [\[2–](#page-15-1)[5\]](#page-15-2), which were based on Castro and Maloney's approach for R-N black holes and 5D black holes. It is well-known that constructing conformal coordinates and generating the Killing vectors can be a challenging task when dealing with complex black hole geometries. We propose that if it is possible to convert the obtained differential equations into general form of Heun's equation, perhaps using this method to identify the algebra of hidden symmetries in field-particle equations with a black hole background is more appropriate, since it can provide a direct and faster way to reveal the algebra and hidden symmetries of black holes, which we will show that for the several different examples.

Prior to concluding this section, it is imperative to underscore that the transformation of various physical systems' differential equations into recognized forms of Heun's equations has elevated the significance of these equations in propelling the exploration of physical systems [\[10–](#page-16-0)[19\]](#page-16-1). Various topics, such as the Schrödinger equation with anharmonic potential, the Teukolsky linear perturbation theory for the Schwarzschild and Kerr metrics, traversable wormholes, quantum Rabi models, the perturbations of rotating and non-rotating black holes, the gravitational lensing by black holes and wormholes, the quantum tunneling, the Hawking radiation, the quasinormal modes, Klein-Gordon, Dirac, Schrödinger, Maxwell, and Einstein equations in various space-times, etc., have motivated many researchers in recent decades to solve, interpret and justify differential equations arising from physical systems that turn into different forms of Heun's equations. [\[28](#page-17-0)[–43\]](#page-18-0).

Now, based on all the explanations above, we will organize the article in the following manner.In sections 2 and 3, we provide an explanation of algebra, Heun's equation, and work motivation. In section 4, we will give a general overview of Heun equations. In section 5, we investigate the scalar field equation in the background of the  $RN - AdS_5$  black hole and the Heun equation of its radial part by introducing the Klein–Gordon particle around it. We identify its operators and the properties of displacement and consider the generalized  $sl(2)$  algebra for the  $RN - AdS_5$ black hole. In section 6, we will follow the same process for the  $Kerr-AdS_5$  black hole. Finally, we describe the results in section 7.

## <span id="page-3-0"></span>2 Warming the mind

Maloney, Castro, and Strominger [\[1\]](#page-15-0) showed at the beginning of the last decade that the physical structure of a Kerr-Newman black hole can be strongly influenced by the algebra of conformal field theory. They had their reasons for this concept: for example, the wave equation around a black hole in a certain limit ("near region") can be calculated as a quadratic Casimir of an  $sl(2, R) \times sl(2, R)$  algebra. They were able to calculate the entropy of the black hole using the concept of central charge near the extreme limit points and showed that the absorption cross section for scalars in the near-region can be reinterpreted as a standard finite temperature CFT absorption cross section [\[3\]](#page-15-3). This new path, and the checking of its validity, prompted others to move to other black holes as R-N to study the hidden symmetries of this black hole [\[2\]](#page-15-1). But the main point in all of these works was that they all used the conventional method during this path. It means Maloney's method! The basis of Castro and Maloney's work was that they considered the Klein-Gordon wave equation with the background of the Kerr black hole. Then, they obtained the radial and angular equations by separating the variables. Although they pointed out that these equations are solvable according to the Heun form, they went on to construct conformal coordinates and consider generators for the vector field on these coordinates. Then, they showed that these generators follow the  $sl(2, R)$  algebra. In other words, they rely on the conventional method of using Killing vectors to identify symmetry. However, in this work, we demonstrate that it is not necessary to construct a new vector field to obtain the hidden algebras of a black hole. Instead, it can be achieved as soon as the wave equation appears in the form of Heun. Depending on the type of algebraic structure of the equation and its coefficients, we can directly proceed to generate and reconstruct the algebra, and then verify its validity by checking the Lie bracket.

### <span id="page-3-1"></span>3 Why Heun?

Bender and Dunne [\[6\]](#page-15-4) showed that there is a logical correspondence between orthogonal polynomials and quasi-exactly solvable (QES) models. In other words, they demonstrated that the wave function in quantum mechanics can act as a generator for orthogonal polynomials expressed in terms of energy. Typically, the algebraic approach is used to analyze QES models, where the Hamiltonian is represented as a non-linear combination of Lie algebra generators. Based on this concept, Turbiner claimed in [\[7\]](#page-16-2) that most QES models are influenced by the  $sl(2, Q)$  algebra (although it was later discovered that some QES models do not necessarily follow the  $sl(2)$  algebra). He was able to show that for different models of specific potentials of this class of Schrödinger equations, these equations can be written using the structure of polynomials and Hamiltonian reconstruction in terms of generators of the  $sl(2)$  algebra. Therefore, it can be said that due to the basic and structural similarity of the Schrödinger and Klein-Gordon equations, differential equations resulting from physical systems whose solutions are somehow the result of orthogonal polynomials may lead us to the existence of a hidden algebra in the physical system. Heun's equation is one such equation. For this reason, in this article, this feature can be used as a practical, easy, and short way to get rid of the construction of Killing vectors, which usually result from long paths with complex computational problems. The important point is that Castro and Maloney's method confirms the result obtained based on this method.

### <span id="page-4-0"></span>4 Overview of Heun Equation

In 1889, Karl M. W. L. Heun introduced a new equation [\[8\]](#page-16-3), a linear ordinary differential equation of second-order type ,which has four regular singular points at zero, one, an arbitrary point between zero and one, and infinity on the complex plane.In fact, this equation is a generalization of the hypergeometric equation, which undoubtedly has found many applications in various branches of science, including in physics, especially in general relativity and astrophysics. To study about this equation and its properties, we can refer to the book of Ronveaux [\[9\]](#page-16-4), which is considered to be comprehensive in this field.

<span id="page-4-1"></span>Based on this, the general form of Heun's equation will be as follows

$$
y''(x) + \left[\frac{\eta}{x} + \frac{\delta}{x-1} + \sum_{i=1}^k \frac{\epsilon_i}{x-a_i}\right] y'(x) + \frac{\mu \nu x^k + \sum_{i=1}^k p_i x^{i-1}}{x(x-1) \Pi_{i=1}^k (x-a_i)} y(x) = 0,\tag{1}
$$

where each singularity  $a_i$  has the characteristic  $(0, 1 - \epsilon_i), (0, 1 - \eta), (0, 1 - \delta)$  as well as  $(\mu, \nu)$ at  $x = 0, x = 1$  and  $x = \infty$ . Therefore, considering the Fuchsian relation, we will have,

$$
\alpha + \beta + 1 = \eta + \delta + \sum_{i=1}^{k} \epsilon_i.
$$
 (2)

<span id="page-4-5"></span><span id="page-4-2"></span>So, the Heun equation [\(1\)](#page-4-1) in the case of  $k = 1$  is as follows,

$$
y''(x) + \left[\frac{\eta}{x} + \frac{\delta}{x - 1} + \frac{\epsilon_1}{x - a}\right]y'(x) + \frac{\alpha\beta x + p_1}{x(x - 1)(x - a)}y(x) = 0,
$$
\n(3)

furthermore, we can rewrite Heun's differential equation [\(3\)](#page-4-2) in a general form by using the functions  $g_1(x)$ ,  $g_2(x)$ , and  $g_3(x)$ ,

<span id="page-4-3"></span>
$$
g_1(x)y''(x) + g_2(x)y'(x) + g_3(x)y(x) = 0,
$$
\n(4)

<span id="page-4-4"></span>where

$$
g_1(x) = a_0 x^3 + a_1 x^2 + a_2 x + a_3,
$$
  
\n
$$
g_2(x) = a_4 x^2 + a_5 x + a_6,
$$
  
\n
$$
g_3(x) = a_7 x + a_8.
$$
\n(5)

With  $a_{i(i=0,1,...,8)} \in R$ , we can express the coefficients of  $a_i$  by comparing equations [\(3\)](#page-4-2) and [\(4\)](#page-4-3) in terms of the parameters of Heun's equation. Additionally, as it shown in [\[20](#page-16-5)[–22\]](#page-17-1), we can rewrite the Heun equation using equations [\(5\)](#page-4-4) and [\(4\)](#page-4-3) in terms of operators  $P^0$ ,  $P^-$ ,  $P^+$ , and  $F(x\frac{d}{dx})$ , such that they follow the generalized  $sl(2)$  algebra and the following commutation relation holds for them,

$$
[P^0, P^{\pm}] = \pm \alpha P^{\pm} \qquad [P^+, P^-] = FP^0,
$$
\n(6)

<span id="page-5-2"></span><span id="page-5-1"></span>where

$$
P^{0} = \alpha x \frac{d}{dx} + \beta
$$
  
\n
$$
P^{+} = a_{0} x^{3} \frac{d^{2}}{dx^{2}} + a_{4} x^{2} \frac{d}{dx} + a_{7} x
$$
  
\n
$$
P^{-} = a_{2} x \frac{d^{2}}{dx^{2}} + a_{6} \frac{d}{dx}
$$
  
\n
$$
F(x \frac{d}{dx}) = a_{1} x^{2} \frac{d^{2}}{dx^{2}} + a_{5} x \frac{d}{dx} + a_{8}.
$$
\n(7)

Also note that we have the following conditions to establish equations  $(6)$  and  $(7)$ ,

$$
\alpha = -\frac{4a_0a_2}{a_1}, \qquad \beta = -\frac{a_6a_7}{a_8}
$$
  
\n
$$
2\alpha a_1 + \alpha a_5 + a_1 \beta = -(6a_0a_2 + 3a_4a_2 + 3a_0a_6)
$$
  
\n
$$
\alpha a_5 + \alpha a_8 + a_5 \beta = -(2a_2a_4 + 2a_6a_4 + 2a_2a_7).
$$
\n(8)

<span id="page-5-3"></span>The calculations related to equation  $(8)$  are in Appendix A. Note that the  $P<sup>0</sup>$  form is significant because it exhibits the symmetry algebra of the mode. By considering the  $P^0$  form in relation [\(7\)](#page-5-2), we can establish the generalized  $sl(2)$  algebra. Additionally,  $\alpha$  and  $\beta$  are special parameters that establish the relation [\(6\)](#page-5-1). In the following, we will discuss the existence of scalar field symmetry in the background of  $RN - AdS_5$  and  $Kerr - AdS_5$  black holes.

## <span id="page-5-0"></span>5  $RN - AdS_5$  black hole

First, we introduce the  $RN - AdS_5$  black hole. The line element of the  $RN - AdS_5$  black hole is as follows [\[23\]](#page-17-2),

$$
ds^{2} = -f(r)dt + \frac{1}{f(r)}dr^{2} + r^{2}d\Omega_{3}^{2},
$$
\n(9)

where  $d\Omega_3^2$  represents the metric of the unit three-sphere. Additionally, the function  $f(r)$  is defined as follows,

$$
f(r) = 1 - \frac{M}{r^2} + \frac{Q^2}{r^4} + r^2 = \frac{\Delta_r}{r^4} = \frac{(r^2 - r_+^2)(r^2 - r_-^2)(r^2 - r_0^2)}{r^4},\tag{10}
$$

where M and Q are the black hole's mass and charge, respectively, and the AdS radius is  $L = 1$ . There is a correspondence between the roots  $\Delta_r$  and the black hole's Killing horizons, whose radial positions will be parameterized by the largest real root of  $r_{+}$ , corresponding to the outer horizon. The roots are as follows,

$$
r_{-}^{2} = \frac{1}{2} \bigg[ -1 - r_{+}^{2} + \sqrt{(1 + r_{+}^{2})^{2} + \frac{4Q^{2}}{r_{+}^{2}}} \bigg],
$$
  
\n
$$
r_{0}^{2} = \frac{1}{2} \bigg[ -1 - r_{+}^{2} - \sqrt{(1 + r_{+}^{2})^{2} + \frac{4Q^{2}}{r_{+}^{2}}} \bigg].
$$
\n(11)

One can use the Klein-Gordon equation in curved space to derive the field equation near the black hole [\[23\]](#page-17-2),

$$
\frac{1}{\sqrt{-g}}D_{\mu}(\sqrt{-g}g^{\mu\nu}D_{\nu})\Psi - m^2\Psi = 0,
$$
\n(12)

<span id="page-6-0"></span>where  $D_{\mu} = \nabla_{\mu} - iqA_{\mu}$ ,  $A_{\mu} = (-\sqrt{3}Q/2r^2, 0, 0, 0, 0)$ , and m and q are the electromagnetic potential, mass, and charge of the scalar field, respectively. Also, we can relate the mass of the scalar field in equation [\(12\)](#page-6-0) to the CFT dimension, such that  $m^2 = \Delta(\Delta - 4)$ . In [\[23\]](#page-17-2), the differential equation for the radial part of equation [\(12\)](#page-6-0) for the scalar field near the  $RN - AdS_5$ black hole was obtained by defining  $z = \frac{r^2 - r^2}{r^2 - r^2}$  $\frac{r^2-r_0^2}{r^2-r_0^2},$ 

<span id="page-6-1"></span>
$$
F''(z) + \left[\frac{1-\theta_{-}}{z} + \frac{1-\theta_{+}}{z-z_{0}} + \frac{\Delta-1}{z-1}\right]F'(z) + \left[\frac{k_{1}k_{2}}{z(z-1)} - \frac{z_{0}(z_{0}-1)K_{0}}{z(z-z_{0})(z-1)}\right]F(z) = 0, \quad (13)
$$

where

$$
k_1 = \frac{1}{2}(\theta_+ + \theta_- - \theta_0 - \Delta)
$$
  
\n
$$
k_2 = \frac{1}{2}(\theta_+ + \theta_- + \theta_0 - \Delta)
$$
  
\n
$$
z_0 = \frac{r_+^2 - r_-^2}{r_+^2 - r_0^2}
$$
  
\n
$$
K_0 = -\frac{[(\theta_+ + \theta_- + 1)^2 - 1 - \theta_0^2]}{4z_0} - \frac{[(2 - \Delta)^2 + 2(\theta_+ - 1)(1 - \Delta) - 2]}{4(z_0 - 1)}
$$
  
\n
$$
-\frac{\ell(\ell + 2) + \Delta(\Delta - 4)r_-^2 - \omega^2}{4z_0(z_0 - 1)(r_+^2 - r_0^2)},
$$
\n(14)

where  $\ell$  and  $\omega$  are the angular momentum quantum number and the quasinormal frequency, respectively. Also, parameters  $\{\theta_n, n = +, -, 0\}$  are  $\theta_n = \frac{i}{2\pi}$ .  $\frac{i}{2\pi T_n}$   $(\omega - \sqrt{3}qQ/2r_n^2)$ . We rewrite equation  $(13)$  as equation  $(4)$ ,

$$
(z3 - z2(z0 + 1) + zz0)F''(z)
$$
  
+ [(1 -  $\theta$  -  $\theta$ <sub>+</sub> +  $\Delta$ )z<sup>2</sup> + ( $\theta$ <sub>+</sub> +  $\theta$  - + z<sub>0</sub> $\theta$  - z<sub>0</sub> $\Delta$  - 2)z + z<sub>0</sub>(1 -  $\theta$  -)]F'(z) (15)  
+ [k<sub>1</sub>k<sub>2</sub>z - k<sub>1</sub>k<sub>2</sub>z<sub>0</sub> - z<sub>0</sub>k<sub>0</sub>(z<sub>0</sub> - 1)]F(z) = 0.

<span id="page-7-1"></span><span id="page-7-0"></span>Now, by comparing relations [\(15\)](#page-7-0), [\(5\)](#page-4-4), [\(4\)](#page-4-3), and [\(3\)](#page-4-2), we can obtain values for  $a_i$ ,  $\alpha$ , and  $\beta$ ,

$$
a_0 = 1, \quad a_1 = -(1 + z_0), \quad a_2 = z_0, \quad a_3 = 0
$$
  
\n
$$
a_4 = 1 + \Delta - \theta_+ - \theta_-, \quad a_5 = -2 + \theta_+ - z_0 \Delta + \theta_- + z_0 \theta_-, \quad a_6 = z_0 (1 - \theta_-) \tag{16}
$$
  
\n
$$
a_7 = k_1 k_2, \quad a_8 = -z_0 [k_1 k_2 + (z_0 - 1) K_0], \quad \alpha = -k_1, \quad \beta = -k_2.
$$

Using relations [\(16\)](#page-7-1) and [\(7\)](#page-5-2), we obtain the operators  $P^+$ ,  $P^-$ ,  $P^0$ , and  $F(z_{\frac{d}{dz}})$ ,

<span id="page-7-2"></span>
$$
P^{0} = -k_{1}z\frac{d}{dz} - k_{2}
$$
  
\n
$$
P^{+} = z^{3}\frac{d^{2}}{dz^{2}} + (1 + \Delta - \theta_{+} - \theta_{-})z^{2}\frac{d}{dz} + k_{1}k_{2}z
$$
  
\n
$$
P^{-} = z_{0}x\frac{d^{2}}{dz^{2}} + z_{0}(1 - \theta_{-})\frac{d}{dz}
$$
  
\n
$$
F(z\frac{d}{dz}) = -(1 + z_{0})z^{2}\frac{d^{2}}{dz^{2}} + (-2 + \theta_{+} - z_{0}\Delta + \theta_{-} + z_{0}\theta_{-})z\frac{d}{dz} + -z_{0}[k_{1}k_{2} + (z_{0} - 1)K_{0}].
$$
\n(17)

According to equation [\(8\)](#page-5-3), we obtain the conditions for establishing algebra for equation [\(17\)](#page-7-2) as follows,

$$
\alpha = \frac{4z_0}{z_0 + 1}, \qquad \beta = \frac{(1 - \theta_-)k_1k_2}{k_1k_2 + K_0(z_0 - 1)}
$$
  
\n
$$
\alpha\theta_+ - 4\alpha - \beta - z_0(\alpha(\Delta + 2) + \beta - 3(\Delta + 4) + 3\theta_+) + \theta_-(\alpha + (\alpha - 6)z_0) = 0
$$
  
\n
$$
(\theta_- + \theta_+ - 2)(\alpha + \beta) - \alpha K_0 z_0^2
$$
  
\n
$$
+ z_0(\theta_-(\alpha + \beta - 2\Delta + 2\theta_+ - 6) - \Delta(\alpha + \beta - 4) + 2\theta_-^2 - 4\theta_+ - (\alpha - 2)k_1k_2 + \alpha K_0 + 4) = 0.
$$
  
\n(18)

Since the operators mentioned above satisfy the relation [\(6\)](#page-5-1), the radial part of the scalar field near the  $RN - AdS_5$  black hole can exhibit generalized  $sl(2)$  symmetry. Furthermore, the quadratic Casimir operator  $C$ , which is associated with all current operators of the model, can be obtained using the formula  $C = \frac{1}{2}$  $\frac{1}{2}(P^-P^+ + P^+P^-) + P^0P^0.$ 

## <span id="page-8-0"></span>6  $Kerr - AdS_5$  black hole

The metric of the  $Kerr - AdS_5$  spacetime is given by [\[24](#page-17-3)[–26\]](#page-17-4),

$$
ds^{2} = -\frac{\Delta}{\rho^{2}} \left( dt - \frac{a_{1} \sin^{2} \theta}{\Xi_{1}} d\phi_{1} - \frac{a_{2} \cos^{2} \theta}{\Xi_{2}} d\phi_{2} \right)^{2} + \frac{\rho^{2}}{\Delta} dr^{2} + \frac{\rho^{2}}{\Delta_{\theta}} d\theta^{2} + \frac{\Delta_{\theta} \sin^{2} \theta}{\rho^{2}} \left( a_{1} dt + \frac{r^{2} + a_{1}^{2}}{\Xi_{1}^{2}} d\phi_{1} \right)^{2} + \frac{\Delta_{\theta} \cos^{2} \theta}{\rho^{2}} \left( a_{2} dt + \frac{r^{2} + a_{2}^{2}}{\Xi_{2}^{2}} d\phi_{2} \right)^{2} + \frac{1 + r^{2} / L^{2}}{r^{2} \rho^{2}} \left( a_{1} a_{2} dt - \frac{a_{2} (r^{2} + a_{1}^{2}) \sin^{2} \theta}{\Xi_{1}} d\phi_{1} - \frac{a_{1} (r^{2} + a_{2}^{2}) \cos^{2} \theta}{\Xi_{2}} d\phi_{2} \right)^{2},
$$
(19)

where,

$$
\Delta := \frac{1}{r^2} (r^2 + a_1^2)(r^2 + a_2^2)(1 + r^2/L^2) - 2M = \frac{1}{r^2 L^2} (r^2 - r_0^2)(r^2 - r_+^2)(r^2 - r_-^2),
$$
  
\n
$$
\Delta_{\theta} := 1 - \frac{a_1^2}{L^2} \cos^2 \theta - \frac{a_2^2}{L^2} \sin^2 \theta,
$$
  
\n
$$
\rho^2 := r^2 + a_1^2 \cos^2 \theta + a_2^2 \sin^2 \theta,
$$
  
\n
$$
\Xi_1 := 1 - \frac{a_1^2}{L^2},
$$
  
\n
$$
\Xi_2 := 1 - \frac{a_2^2}{L^2},
$$
  
\n(20)

where  $0 \le \phi_1, \phi_2 \le 2\pi, 0 \le \theta \le \pi/2$ . The rotational parameters a and b are constrained such that  $a^2, b^2 \leq \ell^{-2}$ . Also, M is the mass parameter,  $a_1$  and  $a_2$  are spin parameters, and  $r_k$  (where  $k = +, -, 0$ ) describe the horizon radii. We assume that they are distinct and that  $r_0^2 < r_-^2 < r_+^2$ . Based on the above explanations,  $r_0$  is purely imaginary, while  $r_-$  and  $r_+$  are real and satisfy  $0 < r_{-} < r_{+}$ . Therefore, r<sub>-</sub> represents the radius of the inner horizon, and r<sub>+</sub> represents the radius of the outer horizon. As shown in [\[27\]](#page-17-5) we will have,

$$
r_0^2 + r_-^2 + r_+^2 + a_1^2 + a_2^2 + L^2 = 0,
$$
  
\n
$$
r_0^2 r_-^2 + r_0^2 r_+^2 + r_-^2 r_+^2 - a_1^2 a_2^2 + (2M - a_1^2 - a_2^2) L^2 = 0,
$$
  
\n
$$
r_0^2 r_-^2 r_+^2 + a_1^2 a_2^2 L^2 = 0.
$$
\n(21)

The angular velocities and Hawking temperatures at each event horizon, denoted by  $r_n$  where  $n = +, -, 0$ , are obtained according to [\[27\]](#page-17-5),

$$
\Omega_{n,1} \equiv \frac{a_1 \Xi_1}{r_n^2 + a_1^2}, \quad \Omega_{n,2} \equiv \frac{a_2 \Xi_1}{r_n^2 + a_2^2}, \quad T_n = \frac{r_n (r_n^2 - r_i^2)(r_n^2 - r_j^2)}{2\pi L^2 (r_n^2 + a_1^2)(r_n^2 + a_2^2)}, \quad i, j \neq n. \tag{22}
$$

We solve the Klein-Gordon equation [\(12\)](#page-6-0) in the background of the  $Kerr - AdS_5$  black hole. Also, considering  $\Psi = \psi(r)\psi(\theta)e^{-i\omega t + im_1\phi_1 + im_2\phi_2}$  (where  $m_1$  and  $m_2$  are integers) and using the separation of variables [\[27\]](#page-17-5), we convert the differential equation into two radial and angular parts.

#### <span id="page-9-0"></span>6.1 Angular section

In the angular section, when  $|a_1| \neq |a_2|$ , and by changing the variables, as shown in [\[27\]](#page-17-5) we will have,

$$
x = \sin^2 \theta, \quad \psi(\theta) = (x - 1)^{m_2/2} x^{m_1/2} (x - x_0)^{\tau/2} S(x)
$$

$$
x_0 = \frac{\Xi_1}{\Xi_1 - \Xi_2}, \quad \tau = \frac{m_1 a_1}{L} + \frac{m_2 a_2}{L} + \omega L.
$$
(23)

<span id="page-9-1"></span>Then, as shown in [\[27\]](#page-17-5) for the Heun equation we will have,

$$
\frac{d^2S(x)}{dx^2} + \left[\frac{\gamma_a}{x} + \frac{\delta_a}{x-1} + \frac{\epsilon_a}{x-x_0} + \right] \frac{dS(x)}{dx} + \frac{\alpha_a\beta_a x + p_a}{x(x-1)(x-x_0)}S(x) = 0,\tag{24}
$$

where

$$
p_a = -\frac{1}{2} [(m_1 + m_2 + m_1 m_2)x_0 + (m_1 + 1)(\tau + m_1)]
$$
  
\n
$$
-\frac{x_0}{4\Xi_1} [\omega^2 L^2 + m^2 a_1^2 - \lambda + \Xi_2 (m_2^2 + m_1^2 - \tau^2)],
$$
  
\n
$$
\alpha_a = \frac{1}{2} (\tau + m_1 + m_2 + \sqrt{m^2 L^2 + 4} + 2), \quad \beta_a = \frac{1}{2} (\tau + m_1 + m_2 - \sqrt{m^2 L^2 + 4} + 2),
$$
  
\n
$$
\epsilon_a = 1 + \tau, \quad \delta_a = 1 + m_2, \quad \gamma_a = 1 + m_1.
$$
\n(25)

Now, we can rewrite equation [\(24\)](#page-9-1) as follows,

$$
(x3 - x2(x0 + 1) + xx0)S''(x)
$$
  
+ 
$$
[(\gammaa + \deltaa + \epsilona)x2 - (\gammaa + x0\gammaa + x0\deltaa + \epsilona)x + x0\gammaa]S'(x)
$$
  
+ 
$$
[\alphaa\betaa + pa]S(x) = 0.
$$
 (26)

<span id="page-9-2"></span>Therefore, by comparing relations [\(24\)](#page-9-1), [\(5\)](#page-4-4), [\(4\)](#page-4-3), and [\(3\)](#page-4-2), we obtain values for  $\alpha$ ,  $\beta$ , and  $a_i$ .

$$
a_0 = 1, \quad a_1 = -(1 + x_0), \quad a_2 = x_0, \quad a_3 = 0
$$
  
\n
$$
a_4 = \gamma_a + \delta_a + \epsilon_a, \quad a_5 = -(\gamma_a + x_0\gamma_a + x_0\delta_a + \epsilon_a), \quad a_6 = \gamma_a x_0
$$
  
\n
$$
a_7 = \alpha_a \beta_a, \quad a_8 = p_a, \quad \alpha = \alpha_a, \quad \beta = \beta_a.
$$
\n(27)

Also, by using relations [\(27\)](#page-9-2) and [\(7\)](#page-5-2), we obtain the operators  $P^+$ ,  $P^-$ ,  $P^0$ , and  $F(x\frac{d}{dx})$ 

$$
P^{0} = \alpha_{a}x\frac{d}{dx} + \beta_{a}
$$
  
\n
$$
P^{+} = x^{3}\frac{d^{2}}{dx^{2}} + (\gamma_{a} + \delta_{a} + \epsilon_{a})x^{2}\frac{d}{dx} + \alpha_{a}\beta_{a}x
$$
  
\n
$$
P^{-} = x_{0}x\frac{d^{2}}{dx^{2}} + \gamma_{a}x_{0}\frac{d}{dx}
$$
  
\n
$$
F(x\frac{d}{dx}) = -(1 + x_{0})x^{2}\frac{d^{2}}{dx^{2}} - (\gamma_{a} + x_{0}\gamma_{a} + x_{0}\delta_{a} + \epsilon_{a})x\frac{d}{dx} + p_{a}.
$$
\n(28)

So, we will have according to equation [\(8\)](#page-5-3),

$$
\alpha = \frac{4x_0}{x_0 + 1}, \qquad \beta = -\frac{x_0 \alpha_a \beta_a \gamma_a}{p_a}
$$
  
\n
$$
- \alpha \gamma_a - \alpha \epsilon_a - x_0 \left( (\alpha - 6) \gamma_a + (\alpha - 3) \delta_a - 3 \epsilon_a + 2 \alpha + \beta - 6 \right) - 2 \alpha - \beta = 0
$$
  
\n
$$
\alpha p_a - (\alpha + \beta) (\gamma_a + \epsilon_a)
$$
  
\n
$$
+ x_0 \left( -\gamma_a \left( -2 \delta_a - 2 \epsilon_a + \alpha + \beta - 2 \right) + 2 \alpha_a \beta_a - \alpha \delta_a - \beta \delta_a + 2 \gamma_a^2 + 2 \delta_a + 2 \epsilon_a \right) = 0
$$
  
\n
$$
.
$$
\n(29)

Since the above operators obey relation  $(6)$ , we can say that there is generalized  $sl(2)$  symmetry for the angular part of the scalar field near the  $ker - AdS_5$  black hole.

#### <span id="page-10-0"></span>6.2 Radial section

In the radial section, the following transformation is applied [\[27\]](#page-17-5),

$$
z = \frac{r^2 - r_+^2}{r^2 - r_0^2}, \quad \psi(r) = (z - 1)^{\sigma/2} (z - z_0)^{\theta - 2} z^{\theta + 2} R(z)
$$
  
\n
$$
z_0 = \frac{r_-^2 - r_+^2}{r_-^2 - r_0^2}, \quad \theta_n = \frac{i}{2\pi T_n} (\omega - m_1 \Omega_{k,1} - m_2 \Omega_{k,2}), \quad \sigma = 2 + \sqrt{4 + m^2 L^2},
$$
\n(30)

<span id="page-10-1"></span>using the above transformations, the radial part of equation will become into the form of Heun equation [\[27\]](#page-17-5),

$$
\frac{d^2R(z)}{dz^2} + \left[\frac{\gamma_r}{z} + \frac{\delta_r}{z-1} + \frac{\epsilon_r}{z-z_0} + \right] \frac{dR(z)}{dz} + \frac{\alpha_r \beta_r x + p_r}{z(z-1)(z-z_0)} R(z) = 0, \tag{31}
$$

where

$$
p_r = \frac{1}{4} \left[ \frac{(\lambda + m^2 r_+^2 - \omega^2 L^2) L^2}{r_-^2 - r_0^2} + 2z_0 (\sigma + \sigma \theta_+ - \theta_+) + (\theta_- + \theta_+ + 2)(\theta_- + \theta_+) - \theta_0^2 \right]
$$
  
\n
$$
\alpha_r = \frac{1}{2} (\sigma + \theta_0 + \theta_- + \theta_+), \quad \beta_r = \frac{1}{2} (\sigma - \theta_0 + \theta_- + \theta_+),
$$
  
\n
$$
\epsilon_r = \theta_- + 1, \quad \delta_r = \sigma - 1, \quad \gamma_r = \theta_+ + 1.
$$
\n(32)

<span id="page-11-0"></span>Note that equation [\(2\)](#page-4-5) holds. We rewrite equation [\(31\)](#page-10-1) as equation [\(2\)](#page-4-5)

$$
(z3 - z2(z0 + 1) + zz0)R''(z)+ [(\gammar + \deltar + \epsilonr)z2 - (\gammar + z0\gammar + z0\deltar + \epsilonr)z + z0\gammar]R'(z)+ [\alphar\betar + pr]R(r) = 0.
$$
\n(33)

<span id="page-11-1"></span>Now, by comparing relations  $(33)$ ,  $(5)$ ,  $(4)$  and  $(3)$  we get

$$
a_0 = 1, \quad a_1 = -(1 + z_0), \quad a_2 = z_0, \quad a_3 = 0
$$
  
\n
$$
a_4 = \gamma_r + \delta_r + \epsilon_r, \quad a_5 = -(\gamma_r + z_0\gamma_r + z_0\delta_r + \epsilon_r), \quad a_6 = \gamma_r z_0
$$
  
\n
$$
a_7 = \alpha_r \beta_r, \quad a_8 = p_r, \quad \alpha = \alpha_r, \quad \beta = \beta_r.
$$
\n(34)

Therefore, using relations [\(34\)](#page-11-1) and [\(7\)](#page-5-2), we obtain the following operators,

$$
P^{0} = \alpha_{r} z \frac{d}{dz} + \beta_{r}
$$
  
\n
$$
P^{+} = z^{3} \frac{d^{2}}{dz^{2}} + (\gamma_{r} + \delta_{r} + \epsilon_{r}) z^{2} \frac{d}{dz} + \alpha_{r} \beta_{r} z
$$
  
\n
$$
P^{-} = z_{0} z \frac{d^{2}}{dz^{2}} + \gamma_{r} z_{0} \frac{d}{dz}
$$
  
\n
$$
F(z \frac{d}{dz}) = -(1 + z_{0}) z^{2} \frac{d^{2}}{dz^{2}} - (\gamma_{r} + z_{0} \gamma_{r} + z_{0} \delta_{r} + \epsilon_{r}) z \frac{d}{dz} + p_{r}.
$$
\n(35)

Here, we can obtain,

$$
\alpha = \frac{4z_0}{z_0 + 1}, \qquad \beta = -\frac{z_0 \alpha_r \beta_r \gamma_r}{p_r}
$$
  
\n
$$
-2\alpha - \beta - \gamma_r \alpha - \alpha \epsilon_r - z_0 (2\alpha + \beta + (\alpha - 6)\gamma_r + (\alpha - 3)\delta_r - 3\epsilon_r - 6) = 0
$$
  
\n
$$
\alpha p_r - (\alpha + \beta) (\gamma_r + \epsilon_r)
$$
  
\n
$$
+ z_0 (-\gamma_r (\alpha + \beta - 2\delta_r - 2\epsilon_r - 2) + 2\alpha_r \beta_r - \alpha \delta_r - \beta \delta_r + 2\gamma_r^2 + 2\delta_r + 2\epsilon_r) = 0.
$$
\n(36)

Note that the operators mentioned above are valid with respect to relation [\(6\)](#page-5-1). Therefore, we can conclude that the radial part of the scalar field near the Kerr black hole follows the generalized sl(2) algebra. Our research reveals that the scalar field near the  $RN - AdS_5$ black hole only adheres to the generalized  $sl(2)$  algebra in the radial part, whereas near the  $Kerr - AdS_5$  black hole, it follows the generalized  $sl(2)$  algebra in both the radial and angular parts.

### <span id="page-12-0"></span>7 Concluding remarks

In the investigation of hidden symmetries in black holes, conventional methods are either based on a purely algebraic and geometric approach or on the method of finding symmetries based on the Killing vector, which has a more dominant physical approach [\[1](#page-15-0)[–5\]](#page-15-2). However, each of these methods has its complexities. As the metric function and geometric structure of the system become more complex, these methods may lose their effectiveness due to the complexity or difficulty of the problem. By studying the method of Castro and Maloney in the search for hidden symmetries of Kerr's black hole and other works based on this method, as well as articles that did not seek to find symmetry in cosmic structures, we realized that Klein Gordon's equations with a black hole background often become differential equations that can be solved with orthogonal polynomials. We also know that studies of Schrödinger equations in quasi-exact solvable models have shown a logical relationship between orthogonal polynomials and the method of solving these models. The similarity between the Klein-Gordon and Schrödinger equations on one hand, and the properties of orthogonal polynomials on the other, led us to consider whether the differential equation resulting from the system, which has been transformed into a special and well-known form (Heun's equation) according to the orthogonal polynomials, can be a good candidate for directly finding the hidden algebra of the system. To this end, we selected two black holes in  $AdS_5$ , namely R-N and Kerr, and analyzed the Klein-Gordon equation with the background of these black holes. As we have seen that the Klein-Gordon differential equations in the background of Reissner-Nordström and Kerr black holes can be transformed into the general form of Heun's equation by using a suitable change of variable, both in the radial and angular parts. As a result, we can easily achieve the algebraic structure based on the algebraic structure that governs the Heun equation and its coefficients. We selected the general form of Heun and demonstrated that the same results can be obtained more simply and efficiently based on the structure of Heun's coefficients, which is a shorter method than the alternative. Our work is accurate as it matches the previous works precisely. Another interesting point is that in the work of Castro and Maloney, they were forced to impose restrictions on the omega function to remove powers higher than 2 and divide the space into two parts - near and far [\[1\]](#page-15-0). Unlike any other application of this approximation for the Castro and Maloney method, it seems that no preconditions need to be applied on the way to the algebra in our method. This method, in addition to being more concise and practical, is also more general and comprehensive across the entire space.

Finally, it may be worthwhile to ask what the purpose and benefit of finding hidden symmetries, such as  $sl(2)$ , for the studied systems are and what new perspectives they offer for further research? To answer this question briefly, we can say that: The  $sl(2)$  symmetry implies that the wave equations with a black hole background can be related to a two-dimensional conformal field theory (CFT) via the AdS/CFT correspondence. This means that the black hole physics can be described by a dual CFT that lives on the boundary of the AdS space. The  $sl(2)$  generators act as the Virasoro operators of the CFT and the Casimir operator corresponds to the central charge. The  $sl(2)$  symmetry also enables us to find the eigenvalues and eigenfunctions of the Heun equation using the representation theory of the algebra [\[44,](#page-18-1) [45\]](#page-19-0). However, In this article, our goal is not to explore these properties, but to demonstrate that we can use the wave equations with a black hole background to reveal the hidden symmetries in a system, without resorting to the common methods that may sometimes be complicated.

In addition to all the mentioned concepts,we also suggest some questions that can serve as motivations for further research.

Could renowned variants of Heun's equation, such as the Trigonometric, Jacobi's Elliptic,Confluent, or Weierstrass's Form [\[46\]](#page-19-1), be employed by reconfiguring the coefficients to unveil concealed algebras and symmetries inherent in the system?

Finally, this fundamental and historical question: if it is possible to solve the differential equations of a physical system using orthogonal polynomials, is it possible to point out the existence of a hidden algebra in the system based on the algebraic structure of these polynomials?

# <span id="page-14-0"></span>8 Appendix A: Calculations related to generator's communication

In this section, we want to calculate the equation [\(7\)](#page-5-2), for a general function  $\psi(x)$ , to obtain the necessary conditions for establishing this relationship [\[20](#page-16-5)[–22\]](#page-17-1).

$$
[P^+, P^-] \psi(x) = FP^0 \psi(x). \tag{37}
$$

<span id="page-14-1"></span>First from the left side we will have

$$
p^{+}p^{-}\psi(x) = a_{7}a_{2}x^{2}\psi''(x) + a_{7}a_{6}x\psi'(x)
$$
  
+  $a_{4}a_{2}x^{3}\psi^{(3)}(x) + a_{4}a_{2}x^{2}\psi''(x) + a_{4}a_{6}x^{2}\psi''(x)$   
+  $a_{0}a_{2}x^{4}\psi^{(4)}(x) + 2a_{0}a_{2}x^{3}\psi^{(3)}(x) + a_{0}a_{6}x^{3}\psi^{(3)}(x),$  (38)

and

<span id="page-14-2"></span>
$$
p^-p^+\psi(x) = 2a_2a_7x\psi'(x) + a_6a_7x\psi'(x) + 2a_2a_4x\psi'(x) + 2a_4a_6x\psi'(x) + a_6a_7\psi(x) + 6a_0a_2x^2\psi''(x) + 4a_2a_4x^2\psi''(x) + 3a_0a_6x^2\psi''(x) + a_4a_6x^2\psi''(x) + a_2a_7x^2\psi''(x) + a_0a_2x^4\psi^{(4)}(x) + 6a_0a_2x^3\psi^{(3)}(x) + a_2a_4x^3\psi^{(3)}(x) + a_0a_6x^3\psi^{(3)}(x).
$$
\n(39)

<span id="page-14-3"></span>Using the equations [\(38\)](#page-14-1) and [\(39\)](#page-14-2), we have

$$
(p^+p^- - p^-p^+) \psi(x) = (-4a_0a_2x^3\psi^{(3)}(x) - (6a_0a_2 + 3a_2a_4 + 3a_0a_6)x^2\psi''(x)
$$
  
-(2a\_2a\_4 + 2a\_6a\_4 + 2a\_2a\_7)x\psi'(x) - a\_6a\_7\psi(x). (40)

<span id="page-14-4"></span>Now, we rewrite the relation [\(40\)](#page-14-3) as below

$$
(p^+p^- - p^-p^+) \psi(x) = \left(-4a_0a_2x^3 \frac{d^3}{dx^3} - (6a_0a_2 + 3a_2a_4 + 3a_0a_6)x^2 \frac{d^2}{dx^2} - (2a_2a_4 + 2a_6a_4 + 2a_2a_7)x \frac{d}{dx}
$$
\n
$$
- a_6a_7)\psi(x).
$$
\n(41)

<span id="page-14-5"></span>From the righthand side of equation (37) we will also have

$$
Fp^{0}\psi(x) = \left(\alpha a_{1}x^{3}\frac{d^{3}}{dx^{3}} + \left(2\alpha a_{1} + \alpha a_{5} + a_{1}\beta\right)x^{2}\frac{d^{2}}{dx^{2}} + \left(\alpha a_{5} + \alpha a_{8} + a_{5}\beta\right)x\frac{d}{dx} + a_{8}\beta\right)\psi(x),
$$
\n(42)

with the comparison of equations[\(41\)](#page-14-4) and [\(42\)](#page-14-5) it will be concluded that

$$
\alpha = -\frac{4a_0a_2}{a_1}, \qquad \beta = -\frac{a_6a_7}{a_8}
$$
  
\n
$$
2\alpha a_1 + \alpha a_5 + a_1 \beta = -(6a_0a_2 + 3a_4a_2 + 3a_0a_6)
$$
  
\n
$$
\alpha a_5 + \alpha a_8 + a_5 \beta = -(2a_2a_4 + 2a_6a_4 + 2a_2a_7).
$$
\n(43)

Now, by using the above relations , the necessary conditions for establishing the equation [\(7\)](#page-5-2) will be obtained.

## Acknowledgments

The authors would like to express their appreciation to Professor Adel Rezaei-Aghdam for his valuable feedback and advice, which has improved the quality and academic accuracy of this manuscript.

## <span id="page-15-0"></span>References

- <span id="page-15-1"></span>[1] Castro, Alejandra, Alexander Maloney, and Andrew Strominger. "Hidden conformal symmetry of the Kerr black hole." Physical Review D 82.2 (2010): 024008.
- [2] Chen, Chiang-Mei, and Jia-Rui Sun. "Hidden conformal symmetry of the Reissner-Nordstrøm black holes." Journal of High Energy Physics 2010.8 (2010): 1-11.
- <span id="page-15-3"></span>[3] Krishnan, Chethan. "Hidden conformal symmetries of five-dimensional black holes." Journal of High Energy Physics 2010.7 (2010): 1-13.
- [4] Franzin, Edgardo, and Ivica Smolić. "A new look at hidden conformal symmetries of black holes." Journal of High Energy Physics 2011.9 (2011): 1-11.
- <span id="page-15-2"></span>[5] Rasmussen, Jørgen. "On hidden symmetries of extremal Kerr–NUT–AdS–dS black holes." Journal of Geometry and Physics 61.5 (2011): 922-926.
- <span id="page-15-4"></span>[6] Bender, Carl M., and Gerald V. Dunne. "Quasi-exactly solvable systems and orthogonal polynomials." Journal of Mathematical Physics 37.1 (1996): 6-11.
- <span id="page-16-3"></span><span id="page-16-2"></span>[7] Turbiner, A. V. "Quasi-exactly-solvable problems and sl (2) algebra." Communications in Mathematical Physics 118 (1988): 467-474.
- <span id="page-16-4"></span>[8] Heun, Karl. "Zur Theorie der Riemann'schen Functionen zweiter Ordnung mit vier Verzweigungspunkten." Mathematische Annalen 33 (1888): 161-179.
- <span id="page-16-0"></span>[9] Ronveaux, A., ed. (1995), Heun's differential equations, Oxford Science Publications, The Clarendon Press Oxford University Press, ISBN 978-0-19-859695-0, MR 1392976
- [10] Decarreau, A., et al. "FORMES CANONIQUES DES EQUATIONS CONFLUENTES DE L'EQUATION DE HEUN." Annales de la Societe Scientifique de Bruxelles 92. (1978): 53.
- [11] Decarreau, Alain, Pascal Maroni, and A. Robert. "Sur les ´equations confluentes de l'équation de Heun." Ann. Soc. Sc. Bruxelles 92. (1978): 151–189.
- [12] Figueiredo, Bartolomeu DB. "On some solutions to generalized spheroidal wave equations and applications." Journal of Physics A: Mathematical and General 35.12 (2002): 2877.
- [13] Figueiredo, B. D. "Ince's limits for confluent and double-confluent Heun equations." Journal of mathematical physics 46.11 (2005).
- [14] Figueiredo, B. D. "Generalized spheroidal wave equation and limiting cases." Journal of Mathematical Physics 48.1 (2007).
- [15] Kimura, Tosihusa. "by Linear Transformations." Funkcialaj Ekvacioj 13 (1970): 213-232.
- [16] Leaver, Edward W. "Solutions to a generalized spheroidal wave equation: Teukolsky's equations in general relativity, and the two-center problem in molecular quantum mechanics." Journal of mathematical physics 27.5 (1986): 1238-1265.
- [17] Wilson, Alan Herries. "A generalised spheroidal wave equation." Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character 118.780 (1928): 617-635.
- [18] Teitelboim, Claudio. "Gravitation and hamiltonian structure in two spacetime dimensions." Physics Letters B 126.1-2 (1983): 41-45.
- <span id="page-16-1"></span>[19] Maldacena, Juan, Douglas Stanford, and Zhenbin Yang. "Conformal symmetry and its breaking in two-dimensional nearly anti-de Sitter space." Progress of Theoretical and Experimental Physics 2016.12 (2016): 12C104.
- <span id="page-16-5"></span>[20] Debergh, N. "The relation between polynomial deformations of sl (2, R) and quasi-exact solvability." Journal of Physics A: Mathematical and General 33.40 (2000): 7109.
- <span id="page-17-1"></span>[21] Shokri, Mehdi, Jafar Sadeghi, and Mohammad Reza Setare. "The generalized sl (2, R) and su (1, 1) in non-minimal constant-roll inflation." Annals of Physics 429 (2021): 168487..
- [22] Sadeghi, J., and M. R. Alipour. "Klein–Gordon particle near RN black hole, generalized sl (2) algebra and harmonic oscillator energy." International Journal of Modern Physics A 34.31 (2019): 1950196.
- <span id="page-17-2"></span>[23] Amado, Julián Barragán, Bruno Carneiro Da Cunha, and Elisabetta Pallante. "Quasinormal modes of scalar fields on small Reissner-Nordström-AdS 5 black holes." Physical Review D 105.4 (2022): 044028.
- <span id="page-17-3"></span>[24] Myers, Robert C., and Michael J. Perry. "Black holes in higher dimensional space-times." Annals of Physics 172.2 (1986): 304-347.
- [25] Hawking, Stephen W., Christian J. Hunter, and Marika M. Taylor-Robinson. "Rotation and the AdS-CFT correspondence." Physical Review D 59.6 (1999): 064005.
- <span id="page-17-4"></span>[26] Gibbons, Gary W., Malcolm J. Perry, and Christopher N. Pope. "The first law of thermodynamics for Kerr–anti-de Sitter black holes." Classical and Quantum Gravity 22.9 (2005): 1503.
- <span id="page-17-5"></span>[27] Noda, Sousuke, and Hayato Motohashi. "Spectroscopy of Kerr-Ads5 spacetime with the Heun function: Quasinormal modes, greybody factor, and evaporation." PRD 106.6 (2022): 064025.
- <span id="page-17-0"></span>[28] Fiziev, Plamen P. "Novel relations and new properties of confluent Heun's functions and their derivatives of arbitrary order." Journal of Physics A: Mathematical and Theoretical 43.3 (2009): 035203.
- [29] Fiziev, Plamen, and Denitsa Staicova. "Application of the confluent Heun functions for finding the quasinormal modes of nonrotating black holes." Physical Review D 84.12 (2011): 127502.
- [30] Fiziev, Plamen, and Denitsa Staicova. "Solving systems of transcendental equations involving the Heun functions." AIP Conference Proceedings. Vol. 1458. No. 1. American Institute of Physics, 2012.
- [31] Fiziev, Plamen P. "Novel representation of the general Heun's functions. Back to the beginning." Proceedings of the International Conference Days on Diffraction 2014. IEEE, 2014.
- [32] Fiziev, Plamen, and Denitsa Staicova. "Solving systems of transcendental equations involving the Heun functions." AIP Conference Proceedings. Vol. 1458. No. 1. American Institute of Physics, 2012.
- [33] Fiziev, Plamen P. "Teukolsky-Starobinsky identities: A novel derivation and generalizations." Physical Review D 80.12 (2009): 124001.
- [34] Fiziev, Plamen P. "Classes of exact solutions to the Teukolsky master equation." Classical and Quantum Gravity 27.13 (2010): 135001.
- [35] Fiziev, Plamen P. "Exact solutions of Regge–Wheeler equation and quasi-normal modes of compact objects." Classical and Quantum Gravity 23.7 (2006): 2447.
- [36] Staicova, Denitsa, and Plamen Fiziev. "New results for electromagnetic quasinormal and quasibound modes of Kerr black holes." Astrophysics and Space Science 358 (2015): 1-12.
- [37] Momeni, D., Koblandy Yerzhanov, and Ratbay Myrzakulov. "Quantized black hole and Heun function." Canadian Journal of Physics 90.9 (2012): 877-881.
- [38] Birkandan, Tolga, and M. Hortaçsu. "Heun-type solutions for Schwarzschild metric with electromagnetic fields." Europhysics Letters 119.2 (2017): 20002.
- [39] Dariescu, Ciprian, Marina-Aura Dariescu, and Cristian Stelea. "Dirac equation on the Kerr–Newman spacetime and heun functions." Advances in High Energy Physics 2021 (2021): 1-10.
- [40] Sakallı, İzzet, Kimet Jusufi, and Ali Övgün. "Analytical solutions in a cosmic string Born–Infeld-dilaton black hole geometry: quasinormal modes and quantization." General Relativity and Gravitation 50 (2018): 1-24.
- [41] Ovgün, Ali, Izzet Sakallı, and Joel Saavedra. "Quasinormal modes of a Schwarzschild black hole immersed in an electromagnetic universe." Chinese Physics C 42.10 (2018): 105102.
- [42] Vieira, H. S., and V. B. Bezerra. "Confluent Heun functions and the physics of black holes: Resonant frequencies, Hawking radiation and scattering of scalar waves." Annals of Physics 373 (2016): 28-42.
- <span id="page-18-0"></span>[43] Suzuki, Hisao, Eiichi Takasugi, and Hiroshi Umetsu. "Perturbations of Kerr-de Sitter black holes and Heun's equations." Progress of theoretical physics 100.3 (1998): 491-505.
- <span id="page-18-1"></span>[44] Yee, Ho-Ung. "A note on AdS/CFT dual of SL (2, Z) action on 3D conformal field theories with U (1) symmetry." Physics Letters B 598.1-2 (2004): 139-148.
- <span id="page-19-0"></span>[45] Klebanov, Igor R., and Edward Witten. "AdS/CFT correspondence and symmetry breaking." Nuclear Physics B 556.1-2 (1999): 89-114.
- <span id="page-19-1"></span> $[46]$ https://dlmf.nist.gov/31.2#iii