

# Calibration of the distribution of the pair-rule proteins Even-skipped and Fushi-tarazu in the *Drosophila* embryo

Catarina Dias, Rui Dilão

University of Lisbon, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisbon, Portugal

## Abstract

From a mass action law approach, we modelled and calibrated the distributions of the seven-stripe patterns of Even-skipped (*Eve*) and Fushi-tarazu (*Ftz*) proteins in the *Drosophila* embryo. The regulators of *Eve* are stripe-specific enhancers from the gap family of proteins. We achieved remarkable data matching of the *Eve* stripe pattern. We have identified the putative repressive combinations for three *Ftz* enhancers, and we have explored the relationship between *Eve* and *Ftz* for complementary patterns. Extended work to infer the Wingless (*Wg*) 14 stripe pattern from *Eve* and *Ftz* enhancers have been proposed, given clues for the hierarchical structure of *Drosophila*'s genetic expression network during early development prior to cellularisation.

**Keywords:** *Drosophila melanogaster* early development, Genetic segmentation network, Stripe enhancers of *Eve* and *Ftz*, Segment-polarity protein enhancers

## 1. Introduction

The emergence of periodic band structures along the antero-posterior axis of the *Drosophila* embryo during early development characterises *Drosophila* embryogenesis, one of the most thoroughly studied gene regulatory networks (Nusslein-Volhard (2006)). Broad domains of maternal proteins regulate the expression of a set of zygotic genes – the gap genes, along the anteroposterior axis of the embryo (Alves *et al.* (2006)). At a later stage, a new set of proteins form seven-stripe complementary patterns – pair-rule proteins, which the most studied pair is the *Even-skipped* (*Eve*) and *Fushi-tarazu* (*Ftz*). This cascade of gene-expressed proteins (maternal → gap → pair-rule) terminates with the segment-polarity gene-expressed proteins that double the pattern to fourteen stripes, after which cellularisation begins, and cells allocate to one of the stripes, defining a segmented larvae body plan.

It is presumed that each *Eve* stripe, and possibly each *Ftz* stripe, is formed and controlled individually by a specific set of transcription factors that regulate individual enhancers (Nusslein-Volhard (2006)). Over the past decades, *eve*'s transcriptional control was studied via mutation experiments and mRNA sequence analyses.

The most recent depiction of the *eve* locus includes five separate enhancers: two dual stripe elements 3+7 and 4+6, and three single stripe elements 1, 2 and 5 (Lim *et al.* (2018)). The classical view of the stripe 2 enhancer includes two transcription repressors, *Krüppel* (*Kr*) and *Giant* (*Gt*), and two transcription activators, *Bicoid* (*Bcd*) and *Hunchback* (*Hb*), who work synergistically (Small *et al.* (1992), Simpson-Brose *et al.* (1994)).

However, Vincent *et al.* (2018) proposed an alternative hypothesis that *Hb* represses the stripe 2 enhancer, and that its repression can be counteracted by nearby sequences, possibly by *Caudal* (*Cad*). As for stripe 5 little information is provided, but it is proposed to be regulated by two transcription repressors, *Kr* and *Gt* (Fujioka *et al.* (1999)). Finally, the two dual stripe enhancers are believed to be regulated by the same transcription repressors, *Hb* and *Knirps* (*Kni*) (Small *et al.* (1996), Fujioka *et al.* (1999)). In the latter case, it is suggested that the enhancers are sensitive to repression by the number and affinity to repressor-binding sites, implying that the 3+7 and 4+6 enhancers would respond autonomously to different amounts of *Hb* and *Kni* (Clyde *et al.* (2003), Crocker *et al.* (2016)). It is also suggested that although stripe 7 is not separable from stripe 3, its activation requires sequences outside of the minimal stripe 3 element. This led to the suggestion of an extra enhancer, a 2+7 dual element, which would be an extended version of the minimal *eve* stripe 2 enhancer (Vincent *et al.* (2018)). To date, practically nothing is known about the formation of stripe 1, with only a small mention that it may be activated by *Hb* in Fujioka *et al.* (1999).

Despite the many experiments for *Eve*, there is still no consensus on the minimum set of its regulators or any successful mathematical modelling, calibration, or spatial distribution of the full seven-stripe pattern of *Eve*. It is even less clear how the formation of the *Ftz* pattern occurs. The latest hypothesis (Schroeder *et al.* (2011)) suggests three enhancers initiating stripe pairs 1+5, 2+7, and 3+6, and the *Zebra* enhancer driving the expression of stripe 4, but, as far as we know, no transcription factors are known or reported in the literature.

In this paper, we develop a simple kinetic framework that accurately predicts and calibrates the formation of *Eve* stripe pattern, providing a minimum set of gap regulators for each *eve* enhancer. We use the mass-action law to describe every

Email addresses: catarina.reis.dias@tecnico.ulisboa.pt (Catarina Dias), ruidilao@tecnico.ulisboa.pt (Rui Dilão)


biological process describing the expression of the *Eve* along the anteroposterior axis of the *Drosophila* embryo, and we anchor this work on the data provided by the FlyEx database (Kozlov *et al.* (2000), Myasnikova *et al.* (2001), Poustelnikova *et al.* (2004), Pisarev *et al.* (2008)). Based on the success of this modelling approach for *Eve*, we are able to predict the position of six of the seven stripes of *Ftz*.

## 2. Methods

To describe all the genetic regulatory interactions that occur during the embryonic segmentation of *Drosophila*, we adopted the Alves *et al.* (2005) approach. We considered that the kinetic mechanism describing the non-regulated production of a protein P is



where G is the template gene for P. We are assuming that DNA splicing has occurred, the transcription and translation from DNA to protein and the concentration of polymerases are represented by the rate constant  $k_1$ , and that  $d$  is the intrinsic degradation rate of protein P. The generic kinetic mechanism that describes gene repression is



where R is a repressor transcription factor,  $G_R$  is a gene complex and  $k_i$  are reaction rates. In mechanism (2), the regulation of the transcription factor is not considered since we are only interested in the repressed translation states of gene G.

The regulation of the *Eve* stripes as depicted in Figure 1 is obtained by several combinations of the basic mechanisms (1)-(2). The choice of wild-type *eve* locus has been adapted from Lim *et al.* (2018), and we have added two more repressor locus sites for *Tll* and *Hkb*, defining the anterior and posterior regions of the *Drosophila*'s embryo.

The complete list of kinetic mechanisms specific to each enhancer locus for *Eve* and *Ftz* is presented in the Appendices. We applied the mass-action law to each kinetic diagram and obtained the differential equations describing the expression of the corresponding *Eve* and *Ftz* stripes along the anteroposterior axis of the *Drosophila*'s embryo. The differential equation models to be calibrated were derived using the software package *kinetics* (Dilão *et al.* (2010)).



Figure 1: Schematic of the wild-type *eve* locus with two dual stripe elements 3+7 and 4+6 and three single stripe enhancers 1, 2 and 5. Adapted from Lim *et al.* (2018). Two repressor locus sites for *Tll* and *Hkb* were added, defining the anterior and posterior regions of the *Drosophila*'s embryo where *Eve* expression is absent at steady state.

In this modelling approach for the calibration of data, we have not considered diffusion mechanisms for proteins or mRNA. In general, reaction-diffusion-type models are used to describe the formation of maternal and gap gene families of proteins (Alves *et al.* (2006), Dilão *et al.* (2010b), Dilão (2014)). We have assumed that stripes of the pair-rule proteins are formed only due to the action of repressing transcription factors distributed along the anteroposterior axis of the *Drosophila*'s embryo, with a clear steady-state expression at the fourteenth mitotic cycle of the *Drosophila* zygote.

As the distributions of the concentrations of the different proteins along the *Drosophila*'s embryo are not measured in well-defined units, the equations were modified so that the variables of interest in the model incorporate a calibration parameter  $\alpha_P$ , specific to each protein P. For each protein P we introduce a new variable

$$\bar{P} = \alpha_P P \quad (3)$$

where  $\alpha_P$  is the weight of a constituent in each reaction. Then, the differential equations in time were solved numerically, with each function taking the experimental spatial distribution of the modelled proteins as input, symbolised by  $\bar{P}$ . Model equations with the respective weights and remaining parameters are listed in the Appendices.

To determine the set of parameters that produce a calibrated steady-state solution that best explains the spatial distribution of *Eve*, we implemented a Monte Carlo algorithm where, at each iteration, every weight parameter  $\alpha_i$ , reaction rates  $k_i$  and  $d$  were generated randomly in certain domains. A fitness function evaluates the quality of the steady-state solutions of the model equations when compared with the data. For each set of  $m$  parameters  $\vec{p} = (p_1, \dots, p_m)$ ,  $n$  experimental data points along the embryo  $\{(x_i, B_{exp}(x_i))\}_{i=1}^n$ , and a function  $B(x_i; \vec{p})$  which approximates the data set, the fitness function is defined as the average of the squared residuals

$$\chi^2(\vec{p}) = \frac{1}{n} \sum_{i=1}^n [B(x_i; \vec{p}) - B_{exp}(x_i)]^2. \quad (4)$$

After  $M$  iterations of the Monte Carlo algorithm, the set of parameters providing the lower  $\chi^2$  value was chosen and used to provide the best fit to the experimental data of each model.

All model results correspond to the best fit out of half a million trials with Monte Carlo optimisation ( $M = 0.5 \times 10^6$ ).

Data was taken from the FlyEx database (Kozlov *et al.* (2000), Myasnikova *et al.* (2001), Poustelnikova *et al.* (2004), Pisarev *et al.* (2008)), except for *Hunkabein* (*Hkb*), which was taken from the SuperFly database (Cicin-Sain *et al.* (2014)). Following common data processing practices (Surkova *et al.* (2008)), we used a Laplacian filter to remove background noise and levelled all distributions to eliminate residual fluorescent concentrations (Alves-Pires *et al.* (1998)). The size of the embryos was rescaled to  $L=1$ . The final gap protein profiles along the normalised anteroposterior position in the embryo of *Drosophila* and the corresponding data are shown in Figure 2.

To obtain the transcriptional regulators for *Eve*'s stripes, we analysed the distribution of gap proteins along *Drosophila*'s




Figure 2: Distribution of gap protein (experimental – dots – and filtered) along the anteroposterior axis of the *Drosophila*'s embryo from cycle 14A4. Ordinate coordinates for each protein are arbitrary and do not correlate. FlyEx datasets: pe3 (*Kr*, *Gt*), rb9 (*Hb*, *Kni*) and th4 (*Tll*). Data for *Hkb* is from the SuperFly database: batch 220310, embryo 016. The embryo lengths were rescaled to  $L = 1$ .

embryo (Figure 3). The observed distributions suggested that the gap transcription factors act as repressors of *eve* and work together in a combinatorial manner. Therefore, we adopted the concept of individual *eve* enhancers, which only allow *Eve*'s production when none of its repressive regulators binds to it. Assuming a kinetic system where the protein *Eve* is freely produced and has  $n$  repressive regulators  $R$ , each repressor contributes to *Eve*'s production and its anteroposterior differentiated distribution. Higher production of *Eve* will occur in regions with lower sums of repressors' concentration, resulting in a localised stripe formation. We tested all the possible combinations of two, three, and four gap proteins, assigning each protein a weight factor to mimic the calibration parameters  $\alpha_P$ , and obtained minimal sets of repressive transcriptional regulators.




Figure 3: Distribution of *Eve*'s stripes and gap proteins along the *Drosophila*'s embryo. The gap data are from stage 14A4 and *Eve*'s data is from stage 14A8. Ordinate coordinates for each protein are arbitrary and do not correlate.

In the modelling, we made several assumptions. We presumed that the concentration of *Eve*'s gene template,  $\bar{G}_e$  is constant, (3), as mitosis does not occur at the stage of embryogenesis considered, and that  $\bar{G}_e$  has a fixed availability in the embryo. Therefore, we have arbitrarily set  $\bar{G}_e(0) = 250$  and  $\alpha_{G_e} = 1$ . To the reference protein *Eve*, has been assigned the weight  $\alpha_E = 1$ , against which each regulator's contribution was measured. All repressor-gene complexes  $G_R(0)$  were set to zero since they should not be formed at the initial stage. For the other initial conditions, we used the distributions of the corresponding gap regulators in cycle 14A4 shown in Figure 3.

### 3. Results and Discussion

In the *Drosophila* wild-type embryo, the pair-rule stripes are limited to a specific area, and *Eve* is not present at the anterior and posterior poles. We determined that the *Eve* stripe region is well-defined by the proteins *Tailless* (*Tll*) and *Huckebein* (*Hkb*), only produced at the embryo ends. These proteins restrict the expression of *Eve* to the desired region by repressing its transcription at the poles. Therefore, the identified terminal regulators integrated every model developed to describe the formation of *Eve*'s stripes.

#### 3.1. *Eve*'s stripes definition via a gap repressive system

$k_{eve-s1}$	$k_{eve-s2}$	$k_{eve-s3}$	$k_{eve-s4}$	$k_{eve-s5}$	$k_{eve-s6}$	$k_{eve-s7}$
0.8584	0.8584	0.9500	3.1100	0.9150	0.8584	1.2900

Table 1: *Eve* production rate values for stripe 1-7 that best fit the experimental data for *Eve* at time class 8, cycle 14 of *Drosophila* embryogenesis shown in Figure 4.


	$k_{kni}$	$k_{-kni}$	$\alpha_{kni}$	$k_{hkb}$	$k_{-hkb}$	$\alpha_{hkb}$	d
<b>1</b>	2.8995	0.0002	0.5667	-	-	-	0.7906
<b>2</b>	5.8021	0.0304	0.9269	-	-	-	0.7906
<b>3+7</b>	4.0241	0.1620	0.1153	1.4049	0.0078	0.1523	0.7906
<b>4+6</b>	7.4764	0.0647	0.6720	7.1382	0.0424	0.0200	0.7906
<b>5</b>	-	-	-	3.6212	0.0657	0.0103	0.7906
	$k_{kr}$	$k_{-kr}$	$\alpha_{kr}$	$k_{gt}$	$k_{-gt}$	$\alpha_{gt}$	
<b>1</b>	3.3110	0.0001	0.0065	0.1294	0.0098	0.5990	
<b>2</b>	0.7502	0.0048	0.0419	0.1091	0.0705	0.0819	
<b>3+7</b>	-	-	-	-	-	-	
<b>4+6</b>	-	-	-	1.5716	0.0428	0.7840	
<b>5</b>	7.4532	0.0356	0.5312	6.5580	0.0374	0.0894	
	$\chi_1^2$	$\chi_2^2$	$\chi_3^2$	$\chi_4^2$	$\chi_5^2$	$\chi_6^2$	$\chi_7^2$
	16.59	10.48	173.40	1301.54	18.55	317.45	386.02
	$\varepsilon_1$	$\varepsilon_2$	$\varepsilon_3$	$\varepsilon_4$	$\varepsilon_5$	$\varepsilon_6$	$\varepsilon_7$
	0.07	0.05	0.11	0.33	0.07	0.27	0.33

Table 2: Parameter values that best fit the experimental data of *Eve*'s stripes at time class 8 of mitotic cycle 14 of *Drosophila* embryogenesis shown in Figure 4. Terminal gap repression parameters were fixed with the 3+7 simulation:  $k_{tll} = 2.2956$ ,  $k_{-tll} = 0.0747$ ,  $\alpha_{tll} = 0.7925$ ,  $k_{hkb} = 2.4229$ ,  $k_{-hkb} = 10^{-5}$ ,  $\alpha_{hkb} = 0.7988$ . Model results after 500,000 trials with Monte Carlo, with calibration parameters  $\alpha$  determined within the interval  $[0,1]$ . The parameters  $\chi_i^2$  are the averages for the squares residuals for each strip, and  $\varepsilon_i = \sqrt{\chi_i^2/B_{i,max}^2}$  is an estimate of the relative error of each stripe fit.

Initially, we believed that the production rate of *Eve* was the same across all cells. However, we found that adjusting *Eve*'s production spatially was necessary to improve the accuracy of the fits. For this reason, we considered the production rate as a function of space and adjusted it stripe by stripe to best match the data (Table 1). This approach is supported by recent studies demonstrating mRNA translation's deep heterogeneity, including protein synthesis rates (Sonneveld *et al.* (2020)).

### 3.1.1. Single stripe 1 enhancer of *Eve*

To explain the formation of stripe 1 of *Eve* protein, we propose a novel enhancer model regulated by the repressive proteins *Tll*, *Hkb*, *Kr*, *Kni* and *Gt*, with kinetic mechanisms



where  $G_e$  is the gene template for the protein *Eve*. The corresponding evolution equations derived from the mass action law are shown in Appendix A. The parameters that best fit the data for stripe 1 are shown in Tables 1 and 2. The parameters in the kinetic mechanisms (5) correspond to the following parameters in Tables 1 and 2: (Table 1)  $k_1 = k_{eve-s1}$ , (Table 2)  $k_3 = k_{kr}$ ,

$k_{-3} = k_{-kr}$ ,  $k_4 = k_{kni}$ ,  $k_{-4} = k_{-kni}$ ,  $k_5 = k_{gt}$ ,  $k_{-5} = k_{-gt}$ ,  $k_6 = k_{tll}$ ,  $k_{-6} = k_{-tll}$ ,  $k_7 = k_{hkb}$  and  $k_{-7} = k_{-hkb}$ .

In Figure 4, we show the position of stripe 1 along the antero-posterior axis of the *Drosophila* embryo, showing a remarkable agreement with the experimental data, reflected in a relative error of  $\varepsilon_1 = 7\%$  (Table 2).


### 3.1.2. Single stripe 2 enhancer of *Eve*

When testing the joint formation of stripes 2 and 5 of *Eve* with only *Gt* and *Kr*, we realised that the only option was to separate their regulation with two single stripe enhancers. Thus, we propose *Kr*, *Kni*, and *Gt* to be the minimal gap set required to explain stripe 2, just as for stripe 1, with the same kinetic mechanisms (5), but different parameters. In this case, (Table 1)  $k_1 = k_{eve-s2}$  and the other parameters are defined as in the stripe 1 case.

With this assumption, we obtained a very precise prediction of stripe 2 – Figure 4, with a relative error lower than  $\varepsilon_2 = 5\%$  (Table 2). Whereas for stripe 1 *Gt* and *Kni* had more or less the same level of repression, now we find a greater affinity to both *Gt* and *Kr* compared to *Kni* (Table 2).

### 3.1.3. Dual stripe enhancer 3+7 of *Eve*

We propose a dual stripe enhancer model for stripes 3 and 7 of *Eve* with the repressive regulators *Tll*, *Hkb*, *Kni* and *Hb*:



The equations associated with these kinetic mechanisms are shown in Appendix A.


The parameters that best fit the data for stripes 3 and 7 are shown in Tables 1 and 2. The parameters in the kinetic mechanisms (6) correspond to the following parameters in Tables 1 and 2: (Table 1)  $k_1 = k_{eve-s3}$ , or  $k_1 = k_{eve-s7}$ , or  $k_1 = 0.8584$  away from the stripes 3 and 7 regions; (Table 2)  $k_3 = k_{kni}$ ,  $k_{-3} = k_{-kni}$ ,  $k_4 = k_{hkb}$ ,  $k_{-4} = k_{-hkb}$ ,  $k_5 = k_{tll}$ ,  $k_{-5} = k_{-tll}$ ,  $k_6 = k_{hkb}$  and  $k_{-6} = k_{-hkb}$ .

The steady-state solution predicted – Figure 4 – shows the location, width, and amplitude of stripe 3 practically match the data, reflected in a low relative error of  $\varepsilon_3 = 11\%$ . Concerning stripe 7, we see that the anterior border forms correctly, but its location and width are not precise, with a high relative error of  $\varepsilon_7 = 33\%$ . This may be due to an excessive *Hkb* repression in the posterior embryo pole. The *Hkb* profile overlaps with stripe 7, suggesting that our model will always underestimate this stripe with the initial conditions data we had access to.

We showed that the proper functioning of this enhancer requires *Kni* to have a slightly higher repression weight, and at a faster rate, than *Hb* (Table 2).

### 3.1.4. Dual stripe enhancer 4+6 of *Eve*

For stripes 4 and 6 of the *Eve* pattern, we propose a model with the repressive regulators *Tll*, *Hkb*, *Kni* and *Gt* and *Hb*:



The equations associated with these kinetic mechanisms are shown in Appendix A. The parameters that best fit the data for stripes 4 and 6 are shown in Tables 1 and 2. The parameters in the kinetic mechanisms (7) correspond to the following parameters in Tables 1 and 2: (Table 1)  $k_1 = k_{eve-s4}$ , or  $k_1 = k_{eve-s6}$ , or  $k_1 = 0.8584$  away from the stripes 4 and 6 regions; (Table 2)  $k_3 = k_{kni}$ ,  $k_{-3} = k_{-kni}$ ,  $k_4 = k_{gt}$ ,  $k_{-4} = k_{-gt}$ ,  $k_5 = k_{hkb}$ ,  $k_{-5} = k_{-hkb}$ ,  $k_6 = k_{tll}$ ,  $k_{-6} = k_{-tll}$ ,  $k_7 = k_{hkb}$  and  $k_{-7} = k_{-hkb}$ .


Regarding stripes 4 and 6, our studies showed that an extra repressor to *Hb* and *Kni*, *Gt*, is necessary for a more accurate description of their development. We achieved an overall correct prediction of both stripe's localisation (Figure 4). There are, however, localisation deviations of both stripes that increase

significantly their relative errors to around  $\varepsilon_4 = 33\%$  and  $\varepsilon_6 = 27\%$ .

In this case, we found it necessary for *Hb* to have a much higher repressive contribution than *Kni* and *Gt*, contrary to the 3+7 case. This sensitivity difference was proposed in the literature (Clyde *et al.* (2003), Crocker *et al.* (2016)) and is now confirmed with our simple enhancer models. Furthermore, our results prove to be consistent with the computationally predicted number of binding sites for each enhancer: Clyde *et al.* (2003) identify 12 *Hb* and 16 *Kni* binding sites in the 3+7 enhancer, and 11 *Hb* and 4 *Kni* binding sites in the 4+6 enhancer. The proportions between these numbers are very similar to those between the calibration  $\alpha$ 's we computed:  $\alpha_{Kni} = 0.1152$  and  $\alpha_{Hb} = 0.1523$  for the 3+7 enhancer, and  $\alpha_{Kni} = 0.6720$  and  $\alpha_{Hb} = 0.0200$  for the 4+6 enhancer.

### 3.1.5. Single stripe 5 enhancer of *Eve*

We found that *Tll*, *Hkb*, *Kr* and *Gt* are sufficient to generate stripe 5 of *Eve*. However, when we added *Hb* as a repressive regulator, we improved the fit to the data, lowering the relative error from around  $\varepsilon_5 \simeq 13\%$  to less than  $\varepsilon_5 = 8\%$  (Figure 4, Table 2). This latter enhancer has *Hb* as the main repressive regulator, followed by *Gt*, and both have much higher weighting factors than *Kr*. Therefore to regulate stripe 5, we have chosen the kinetic mechanism



The equations associated with these kinetic mechanisms are shown in Appendix A. The parameters that best fit the data for stripe 5 are shown in Tables 1 and 2. The parameters in the kinetic mechanisms (8) correspond to the following parameters in Tables 1 and 2: (Table 1)  $k_1 = k_{eve-s5}$ , (Table 2)  $k_3 = k_{kr}$ ,  $k_{-3} = k_{-kr}$ ,  $k_4 = k_{gt}$ ,  $k_{-4} = k_{-gt}$ ,  $k_5 = k_{hkb}$ ,  $k_{-5} = k_{-hkb}$ ,  $k_6 = k_{tll}$ ,  $k_{-6} = k_{-tll}$ ,  $k_7 = k_{hkb}$  and  $k_{-7} = k_{-hkb}$ .

Overall, we accurately predicted all the stripes in the *Eve* pattern with a purely repressive gap regulating system. We found no justification or need for any gap transcription activator role. Moreover, our model does not require any maternal input. We propose that any effect on the *Eve* pattern observed in maternal gene mutation experiments comes from collateral damage to gap regulation and consequent disturbance in the formation of the gap domains.

## 3.2. Simulating gap mutations of *Eve*

An important step in validating our model was the introduction of gap mutations. We removed one regulator at a time in each model and observed the effects on the *Eve* pattern of time class 8 of mitotic cycle 14 (Figure 5).

For a *kr* mutation introduced in stripe 2 (Figure 5, S2,  $kr^-$ ) and stripe 5 enhancers (Figure 5, S5,  $kr^-$ ), we observe the fusion




Figure 4: Predictions for *eve*'s stripe-specific enhancers 1, 2, 5, 3+7 and 4+6 and full *Eve* pattern prediction obtained by combining the results of each enhancer model. The figure shows the best fits of each model to the experimental data of *Eve* stripes (dotted lines) at time class 8 of mitotic cycle 14 (14A8) of *Drosophila* embryogenesis. We obtained the parameter values of  $k_i$  and  $\alpha_i$  by a Monte Carlo optimisation after 500,000 trials which are summarised in Table 2. The adjusted *Eve* production rates  $k_i$  are shown in Table 1.

of stripes 2 and 3 and the fusion of stripes 4 and 5. Such a result coincides with the experimental results of Surkova *et al.* (2013) and with the anterior expansion of stripe 5 reported in Fujioka *et al.* (1999). However, we do not observe the weakening of stripe 4 and the posterior expansion of stripe 6, reported in that same article, or even a more pronounced expression of stripe 2. We point out that we are not considering the collateral effects of each mutation in other gap gene's expression, and thus, we

are disregarding any indirect alteration of *Eve*'s pattern. For instance, a *kr* mutation can influence all other gap genes *gt*, *kni* and *hb* expression domains (Figure 2), involved in forming stripes 4 and 6.

Next, we analysed the introduction of a *gt* mutation (Figure 5, S2 and S5, *gt*<sup>-</sup>), in which stripes 1 and 2 merge as well as stripes 5 and 6. These results agree with the experiments carried out by Frash *et al.* (1987), and also with Small *et al.* (1991) and




Figure 5: Effects on *Eve*'s pattern of the introduction of gap gene mutations  $kr^-$ ,  $gt^-$ ,  $kni^-$  and  $hb^-$  in each enhancer model, cycle 14A8 of *Drosophila* embryogenesis. A  $kr^-$  embryo shows two fused domains, 2-3 and 4-5, coincident with the experimental results of Surkova *et al.* (2008) and Fujioka *et al.* (1999). A  $gt^-$  embryo presents a merging of stripes 1 and 2 as of stripes 5 and 6, in agreement with Frash *et al.* (1987), Small *et al.* (1991) and Stanojevic *et al.* (1991). A  $kni^-$  embryo presents a merging of stripes in the domain 3 to 7, just as observed in Surkova *et al.* (2008). A  $hb^-$  embryo shows an anterior fused domain up to stripe 3 or 4, a posterior expansion of stripe 7, and a possible overlap with stripe 6. Once *sgsin*, this is in line with the mutation experiments studied in Frash *et al.* (1987), Small *et al.* (1991) and Fujioka *et al.* (1999).

Stanojevic *et al.* (1991) that report the expansion of the anterior border of stripe 2. Although an intensity reduction for stripe 7 is shown in Frash *et al.* (1987), we do not verify that. On the contrary, we observe the appearance of a seventh stripe that did not exist before on this model.

As for the  $kni$  mutation, it was introduced in both enhancers 3+7 and 4+6 (Figure 5, S3+7,  $kni^-$  and S4+6,  $kni^-$ ). They show a fusion of stripes 3 to 7 and a fusion of stripes 4 to 6, the behaviour expected according to Surkova *et al.* (2013) experiments.

Finally, we studied the introduction of an  $hb$  mutation in the 3+7 and 4+6 enhancers (Figure 5, S3+7,  $hb^-$  and S4+6,  $hb^-$ ). For the 3+7 case we observe a broad fusion domain starting at the anterior end of the embryo and encompassing stripes 1, 2 and 3, and a posterior expansion of stripe 7. As for the 4+6 enhancer, we see a fused expression domain in the anterior region up to stripe 4 and an expansion of the posterior border of stripe 6, overlapping with stripe 7. Both cases are in line with the mutation experiences made by Frash *et al.* (1987), Small *et al.* (1996) and Fujioka *et al.* (1999) as they report a total or partial deletion of stripes 1 to 4, with a broad anterior band of expression instead, the expansion of stripes 3 and 7, and the expansion of stripes 4 and 6.

Both  $tll^-$  and  $hkb^-$  mutations were tested, but they did not cause significant changes to *Eve*'s pattern aside from a lack of definition in the embryo poles. Contrarily to what was expected (Small *et al.* (1996), Frash *et al.* (1987), Fujioka *et al.* (1999), Janssens *et al.* (2013), Ilsley *et al.* (2013)), a  $tll$  mutation did not abolish stripe 7. Once more, we believe that the abolition

of this stripe is somehow related to the direct interference with the regulation of other gap genes, such as  $kni$  and  $gt$ , that is caused by a  $tll$  mutation. Furthermore, we hypothesise that the experimentally recorded effects of a mutation in maternal genes  $nos$  and  $tor$  are once again the effect of the mis-expression of gap genes. For example,  $tor$  regulates the terminal gap genes, so a  $tor$  mutation is expected to alter *Eve* in a similar way to a mutation in  $tll$  or  $hkb$ .

### 3.3. *Ftz*'s stripes can be partially defined via a gap repressive system




Figure 6: Schematic of the hypothesised regulation of the wild-type *ftz* locus with three dual stripe elements 1+5, 2+7 and 3+6 stripe enhancers. Enhancers for stripe 4 are unknown.

Following a reasoning in every way similar to the formation of *Eve* stripes, we tested the viability of stripe-specific enhancers driving *Ftz* stripes. The gap protein combinations we identified as possibly regulating *Ftz* correspond to those found to regulate *eve*'s transcription: our thermodynamics approach is compatible with two dual stripe elements 2+7 and 3+6 regulated by *Hb* and *Kni* and another dual stripe enhancer 1+5 regulated by *Kr* and *Gt*. We also assume that *Tll* and *Hkb* repress the production of *Ftz* in the anterior and posterior regions of the *Drosophila*'s embryo (Figure 6).




Figure 7: Predictions for *ftz*'s stripe-specific enhancers 1+5, 2+7, 3+6 and full *Ftz* pattern prediction obtained by combining the results of each enhancer model. The figure shows the best fits of each model to the experimental data of *Ftz* stripes (dotted lines) at time class 8 of mitotic cycle 14 (14A8) of *Drosophila*'s embryogenesis. We obtained the parameter values of  $k_i$  and  $\alpha_i$  by a Monte Carlo optimisation after 500,000 trials which are summarised in Table 4. The adjusted *Ftz* production rates  $k_1$  are listed in Table 3.

### 3.3.1. Dual stripe enhancer 1+5 of *Ftz*

We initially thought that *Gt* and *Kr* were enough to regulate the 1+5 enhancer, but we discovered that adding a third regulator, *Kni*, is essential. In this case, the kinetic regulatory mechanism is similar to the one for stripes 1 and 2 of *Eve*, (5), with  $G_e \rightarrow G_f$ , where  $G_f$  is the gene template for the production of *Ftz*. The corresponding evolution equations derived from the mass action law are shown in Appendix A. The parameters that best fit the data for stripes 1 and 5 are shown in Tables 3 and 4: (Table 3)  $k_1 = k_{ftz-s1}$ , or  $k_1 = k_{ftz-s5}$ , or  $k_1 = 0.5823$  away from the stripes 1 and 5 regions; (Table 4)  $k_3 = k_{kr}$ ,  $k_{-3} = k_{-kr}$ ,  $k_4 = k_{kni}$ ,  $k_{-4} = k_{-kni}$ ,  $k_5 = k_{gt}$ ,  $k_{-5} = k_{-gt}$ ,  $k_6 = k_{tll}$  and  $k_{-6} = k_{-tll}$ ,  $k_7 = k_{hkb}$  and  $k_{-7} = k_{-hkb}$ .

In Figure 7, we show the position of stripes 1 and 5 along the anteroposterior axis of the *Drosophila*'s embryo, showing a remarkable agreement with the experimental data, reflected in relative errors of  $\varepsilon_1 = 20\%$  and  $\varepsilon_5 = 18\%$ , respectively (Table 2). The mismatches to the experimental data are the stripe widths and a very slight deviation of the stripe 1 peak.

We found this enhancer to have a higher affinity to *Kr* and *Kni* than to *Gt* (Table 4).

$k_{ftz-s1}$	$k_{ftz-s2}$	$k_{ftz-s3}$	$k_{ftz-s4}$	$k_{ftz-s5}$	$k_{ftz-s6}$	$k_{ftz-s7}$
1.3100	0.6800	1.8910	-	1.500	0.5630	3.3400

Table 3: *Ftz* production rate values that best fit the experimental data at time class 8 of mitotic cycle 14 of *Drosophila* embryogenesis shown in Figure 7.

### 3.3.2. Dual stripe enhancer 2+7 of *Ftz*

The position of stripes 2 and 7 are predicted with an enhancer model with repressive regulators *Hb*, *Kni*, *Tll* and *Hkb*. In this case, the kinetic regulatory mechanism is similar to the one for dual stripes 3+7 of *Eve*, (6), with  $G_e \rightarrow G_f$ . The corresponding evolution equations derived from the mass action law are shown in Appendix A. The parameters that best fit the data for stripes 2 and 7 are shown in Tables 3 and 4: (Table 3)  $k_1 = k_{ftz-s2}$ , or  $k_1 = k_{ftz-s7}$ , or  $k_1 = 0.5823$  away from the stripes 2 and 7 regions; (Table 4)  $k_3 = k_{kni}$ ,  $k_{-3} = k_{-kni}$ ,  $k_4 = k_{hb}$ ,  $k_{-4} = k_{-hb}$ ,  $k_5 = k_{tll}$  and  $k_{-5} = k_{-tll}$ ,  $k_6 = k_{hkb}$  and  $k_{-6} = k_{-hkb}$ .

An enhancer model with repressive regulators *Hb* and *Kni* predicted stripe 7 with a reasonable agreement with the data and a low relative error of about  $\varepsilon_7 = 18\%$ . Its location was

	$k_{kni}$	$k_{-kni}$	$\alpha_{kni}$	$k_{hb}$	$k_{-hb}$	$\alpha_{hb}$	d
<b>1+5</b>	5.6887	0.0205	0.9909	-	-	-	0.7104
<b>2+7</b>	7.3807	0.0034	0.0434	4.5349	0.0375	0.5605	0.7104
<b>3+6</b>	3.1028	0.0877	0.3160	7.7090	0.0265	0.0440	0.7104
	$k_{kr}$	$k_{-kr}$	$\alpha_{kr}$	$k_{gt}$	$k_{-gt}$	$\alpha_{gt}$	
<b>1+5</b>	3.6834	0.0748	0.2489	0.3609	0.0001	0.3461	
<b>2+7</b>	-	-	-	-	-	-	
<b>3+6</b>	-	-	-	-	-	-	
	$\chi_1^2$	$\chi_2^2$	$\chi_3^2$	$\chi_4^2$	$\chi_5^2$	$\chi_6^2$	$\chi_7^2$
	201.46	1051.69	106.03	-	198.03	253.16	325.30
	$\varepsilon_1$	$\varepsilon_2$	$\varepsilon_3$	$\varepsilon_4$	$\varepsilon_5$	$\varepsilon_6$	$\varepsilon_7$
	0.20	0.36	0.09	-	0.18	0.19	0.18

Table 4: Parameter values that best fit the experimental data of *Ftz*'s stripes at time class 8 of mitotic cycle 14 of *Drosophila*'s embryogenesis shown in Figure 4. Terminal gap repression parameters fixed with the 2+7 simulation for *Tll* and with the 1+5 simulation for *Hkb*:  $k_{tll} = 5.2564$ ,  $k_{-tll} = 0.0937$ ,  $\alpha_{tll} = 0.4560$ ,  $k_{hkb} = 8.3229$ ,  $k_{-hkb} = 0.0500$ ,  $\alpha_{hkb} = 0.2900$ . Model results after 500,000 trials with Monte Carlo, with calibration parameters  $\alpha$  determined within the interval  $[0,1]$ .  $\varepsilon_i = \sqrt{\chi_i^2/B_{i,max}^2}$  is an estimate of the relative error of each stripe fit.

precise, but its width was underestimated. As for stripe 2, it has a well-described anterior border but an extended posterior border. Such can be an indication that the *Hb* repression weight ( $\alpha_{Hb} = 0.5605$ ) determined is not enough. Together with the width mismatch, this leads to a high relative error of  $\varepsilon_2 = 36\%$ .

The proper functioning of this enhancer requires a higher repression weight from *Kni* versus *Hb* (Table 4).

### 3.3.3. Dual stripe enhancer 3+6 of *Ftz*

Finally, we studied the control mechanisms of the dual stripe enhancer 3+6, which we propose to be regulated by the same proteins as the 2+7 enhancer of *Ftz* and the 3+7 enhancer of *Eve*: *Hb*, *Kni*, *Tll* and *Hkb*. The predicted stripes precisely match the experimental data, reflected in the low relative errors for stripes 3 and 6 of  $\varepsilon_3 = 9\%$  and  $\varepsilon_6 = 19\%$ .

Contrarily to what is observed for the dual stripe enhancer 2+7, the correct functioning of the 3+6 enhancer relies on a higher concentration of *Hb* versus *Kni* ( $\alpha_{Hb} = 0.0440$  and  $\alpha_2 = 0.3160$ ). Thus, despite depending on the same regulators as the 2+7, enhancer 3+6 has a greater affinity or a greater number of binding sites for *Hb* than for *Kni*. Note that this relationship between the *ftz*'s 2+7 and 3+6 enhancers is exactly the same as between *eve*'s 3+7 and 4+6 enhancers. If these two dual stripe enhancers are indeed involved in *Ftz*'s pattern formation, it would be interesting to determine experimentally the number of binding sites in each and compare them to the calibration  $\alpha$ 's we computationally predicted.




Figure 8: Best fits of the experimental data of the full *Eve* and *Ftz* patterns at time class 8 of mitotic cycle 14 (14A8) of *Drosophila* embryogenesis. These predictions were obtained following a stripe-specific enhancer theory. No regulators were identified for *Ftz*'s stripe 4.

We proved that stripe-specific elements can partially define the *Ftz* pattern along the anteroposterior axis of the *Drosophila*'s embryo, with stripe 4 being the only one that can't be explained with gap transcription factors.

### 3.4. Final predictions for *Eve* and *Ftz* patterns

Modelling *Eve* and *Ftz* pattern development with a purely repressive gap regulating system was successful, with the two proteins demonstrating the desired complementarity. In Figure 8, we combine the two distributions obtained in Figures 4 and 7. We provided a new mechanism for the regulation of stripe 1, with a repressive combination of five gap proteins, making us reconsider the *Hb* activation proposal by Fujioka *et al.* (1999).





Figure 9: Simulation of direct repression between *Eve* and *Ftz* as the main mechanism to develop the *Ftz* complementary pattern. Three scenarios were tested. **Left:** *Eve*'s pattern is primarily formed by its five enhancers and *Ftz*, by some unknown process, starts developing with the **(left)** same or **(middle)** similar pattern to *Eve*'s. Epistasis quickly overrides this mechanism and becomes the main process creating the two complementary patterns. **(right)** *Ftz* starts to be produced from its stripe-specific enhancers, and then epistasis overrides and becomes the main mechanism. We assume the *ftz* enhancers react to *eve*'s enhancers parameters. For example, we could have as an initial condition the 1+5 enhancer giving rise to a stripe overlapping *Eve*'s stripe 1, and the 2+7 and 3+6 enhancers originating stripes coinciding with *Eve* stripe pairs 3+7 and 4+6. Parameter values: **(left)**  $k_1 = 3.54, k_2 = 3.54, k_3 = 0.1, k_4 = 0.1, k_5 = 2.22, k_{-5} = 5.00 \times 10^{-4}, k_6 = 3.66, k_{-6} = 5.00 \times 10^{-4}, \alpha_{eve} = 0.370, \alpha_{ftz} = 0.478$ ; **(middle)**  $k_1 = 3.54, k_2 = 3.54, k_3 = 0.1, k_4 = 0.1, k_5 = 2.34, k_{-5} = 5.00 \times 10^{-4}, k_6 = 3.66, k_{-6} = 5.00 \times 10^{-4}, \alpha_{eve} = 0.370, \alpha_{ftz} = 0.498$ ; **(right)**  $k_1 = 3.54, k_2 = 3.54, k_3 = 0.1, k_4 = 0.1, k_5 = 1.40, k_{-5} = 5.00 \times 10^{-4}, k_6 = 3.00, k_{-6} = 5.00 \times 10^{-4}, \alpha_{eve} = 0.394, \alpha_{ftz} = 0.796$ . The effects of the distribution of terminal proteins *Tll* and *Hkb* were not considered.

The final *Eve* and *Ftz* patterns have  $\chi^2$  values of  $\chi^2_{Eve} = 2224$  and  $\chi^2_{Ftz} = 2136$ , which are dependent on the system of units of experimental data. On the other hand, this depends on the variability of embryo lengths and the experimental conditions used to obtain each protein profile. These variabilities, although small, can introduce imprecisions in the stripe location predictions. Ideally, to be objective, we would have to use profiles belonging to the same embryo.

### 3.5. An alternative *Ftz* pattern formation via *eve-ftz* epistasis

In the models developed, stripe 4 expressions cannot be explained by gap transcriptional repression. Additionally, the two patterns we obtained (Figure 8) are not perfectly complementary. This suggests yet another level of complexity as, for example, the introduction of cross-repression between *Eve* and *Ftz* to adjust the precise position of the stripes and develop *Ftz*'s stripe 4. This hypothesis advanced by Frash *et al.* (1987), together with studies by Lim *et al.* (2018) on the temporal dynamics of the two proteins, led us to consider the possibility of another, simpler mechanism that can override the stripe specific enhancers action where *Eve* and *Ftz* directly repress each other's transcription.

To test this self-repression hypothesis, we introduced the simplified kinetic mechanism




and we mimicked the striped patterns using sums of Gaussian functions, each centred in individual stripes. The evolution equations of the kinetic mechanism (9) are shown in Appendix B.

In Figure 9, we show the steady states obtained with the evolution equations associated with kinetic mechanism (9) for three

different biological scenarios. In every case, this epistasis mechanism proved sufficient to produce two perfectly complementary stripe patterns. This epistasis concept, although successful in theory, implies that the *Ftz* pattern formation is dependent on *Eve*. At first glance, this does not agree with the mutation experiments (Harding *et al.* (1986)). However, we point out that there are few experiments of this type and little information on the regulation of *ftz* for a conclusive study. Moreover, we must not forget that *eve* and *ftz* are part of a complex gene expression network. We can conjecture a cross-repression network between the pair-rule such that other genes similar to *Eve* assume its role in providing periodic input in the case of a gene mutation.

### 3.6. Modeling segment-polarity pattern formation

Our modelling strategy can be extended to infer the behaviour of other genes in the *Drosophila* embryogenesis, just as the segment-polarity gene *Wg*. *Wg* has 14 stripes perfectly developed in the narrow stripes of cells that separate *Eve* and *Ftz* (Ingham *et al.* (1988)). We built a simple model



where  $G_{wg}$  is the gene template for the free production of the protein *Wg*, and *Eve* and *Ftz* are repressive regulators. The corresponding evolution equations derived from the mass action law are shown in Appendix C. The patterns of *Eve* and *Ftz* were simulated, summing localised Gaussian functions, each centred in an individual stripe. The model results in Figure 10 show the full development of 11 of the 14 stripes. However, there is an indication that two other stripes would form, one between *Eve* and *Ftz*'s first stripes and another between their last stripes. A fourteenth stripe is expected just before the beginning of

*Eve*'s pattern. This pattern would be fixed with terminal protein repression, as in the pair-rule case. We do not know, however, which proteins could play this role for the segment-polarity family.




Figure 10: Simulation of a segment-polarity pattern (solid black line, SP), through the repressive action of *Eve* and *Ftz* in *Drosophila* embryogenesis. We observe the clear formation of 11 of the expected 14 stripes. Steady-state solution evaluated at time  $t = 100$ . Parameter values:  $k_1 = 13.65$ ,  $k_2 = 1.00$ ,  $k_3 = 5.00$ ,  $k_4 = 5.00$ ,  $k_{-3/-4} = 10^{-2}$ ,  $\alpha_1 = 0.570$ ,  $\alpha_2 = 0.600$ . The effects of the distribution of putative terminal repressing proteins were not considered.

This demonstrates that with access to experimental data for segment-polarity expression patterns, it would be easy to analyse the possible pair-rule combinations, determine transcriptional activators and repressors for this last hierarchical level of gene regulation, and calibrate the respective patterns, regardless of the embryo in question.

#### 4. Conclusions

We studied the gene regulatory network of *Drosophila* embryogenesis, focusing on modelling and calibrating the seven-stripe patterns of the proteins *Eve* and *Ftz*. With the kinetic mass-action law describing transcriptional regulation, we modelled five stripe-specific enhancers for *eve*. We proposed a purely repressive gap system with novel minimal regulator combinations as depicted in Figures 1 and 6. Our model predictions matched the data remarkably well and provided a complete description of *Eve*'s development, including the definition of the anterior and posterior *Drosophila* embryo poles and the calibration of the *Eve* production rate across the embryo. Then, we identified three gap repressive combinations driving three *ftz* enhancers and successfully achieved *Ftz*'s pattern development calibration, except stripe 4. We explored an alternative mechanism of epistasis between *eve* and *ftz* that proved to be sufficient to originate two perfectly complementary patterns, having thus identified two different mechanisms for *Ftz* formation. Lastly, we extended our work to infer the behaviour of the segment polarity protein *Wg* from *Eve* and *Ftz* repressive interactions.


#### References

- F. Alves and R. Dilão, A simple framework to describe the regulation of gene expression in prokaryotes, *Comptes Rendus Biologies*, 328 (2005) 429-444, doi: 10.1016/j.crv.2005.01.009.
- F. Alves and R. Dilão, Modeling segmental patterning in *Drosophila*: Maternal and gap genes, *J. Theo. Biol.* 241 (2006) 342-359, doi: 10.1016/j.jtbi.2005.11.034.
- R. Alves-Pires, R. Dilão, H. Neves, L. Parreira and J. Sainhas, Anisotropy-free Laplacian filters, contour detection, and 3D image reconstruction for confocal microscopy imaging, In F. Muge, M. Piedade and J. C. Pinto (ed.), "Proceedings of the RecPad98, 10th Portuguese Conference on Pattern Recognition", CVRM/IST, pp. 247-251, 1998, ISBN: 972-97711-0-3.
- A. Berrocal, N. Lammers, H. Garcia, G. Hernan and M. B. Eisen, Kinetic sculpting of the seven stripes of the *Drosophila even-skipped* gene, *eLife*, 9 (2020) e61635, doi: 10.7554/eLife.61635.
- V. C. Calhoun and M. Levine, Long-range enhancer-promoter interactions in the *ScrAntp* interval of the *Drosophila* Antennapedia complex, *PNAS*, 100 (2003) 9878-9883, doi: 10.1073/pnas.1233791100.
- D. Cicin-Sain, A. H. Pulido, A. Crombach, K. R. Wotton, E. Jiménez-Guri, J.-F. Taly, G. Roma and J. Jaeger, SuperFly: a comparative database for quantified spatio-temporal gene expression patterns in early dipteran embryos, *Nucleic Acids Research*, 43 (2014) D751-D755, doi: 10.1093/nar/gku1142.
- D. Clyde, M. Corado and X. Wu, A self-organizing system of repressor gradients establishes segmental complexity in *Drosophila*, *Nature*, 426 (2003) 849-853, doi: 10.1038/nature02189.
- J. Crocker, G. R. Iisley and D. L. Stern, Quantitatively predictable control of *Drosophila* transcriptional enhancers in vivo with engineered transcription factors, *Nature Genetics*, 48 (2016) 292-298.
- C. R. Dearolf, J. Topol and C. S. Parker, Transcriptional control of *Drosophila* fushi tarazu zebra stripe expression, *Genes Dev*, 3 (1989) 384-398, doi: 10.1101/gad.3.3.384.
- R. Dilão and D. Muraro, A Software Tool to Model Genetic Regulatory Networks. Applications to the Modeling of Threshold Phenomena and of Spatial Patterning in *Drosophila*, *PLOS ONE*, 5 (2010) 1-10, doi: 10.1371/journal.pone.0010743.
- R. Dilão and D. Muraro, Calibration and validation of a genetic regulatory network model describing the production of the protein Hunchback in *Drosophila* early development, *Comptes Rendus Biologies*, 333 (2010) 779-788, DOI:10.1016/j.crv.2010.09.003.
- R. Dilão, Bicoid mRNA diffusion as a mechanism of morphogenesis in *Drosophila* early development, *Comptes Rendus Biologies*, 337 (2014) 679-682, DOI: 10.1016/j.crv.2014.09.004.
- M. Frash and M. Levine, Complementary patterns of *even-skipped* and *fushi-tarazu* expression involve their differential regulation by a common set of segmentation genes in *Drosophila*, *Genes & Development*, 1(9) (1987) 981-995, DOI: 10.1101/gad.1.9.981.
- M. Fujioka, Y. Emi-Sarker, G. L. Yusibova, T. Goto and J. B. Jaynes, Analysis of an even-skipped rescue transgene reveals both composite and discrete neuronal and early blastoderm enhancers, and multi-stripe positioning by gap gene repressor gradients, *Development*, 126 (1999) 2527-2538.
- K. Harding, C. Rushlow, H. J. Doyle, T. Hoey and M. Levine, Cross-Regulatory Interactions Among Pair-Rule Genes in *Drosophila*, *Science*, 233 (1986) 953-959, doi: 10.1126/science.3755551.
- Y. Hiromi and W. J. Gehring, Regulation and function of the *Drosophila* segmentation gene fushi-tarazu, *Cell*, 50 (1987) 963-974, doi: 10.1016/0092-8674(87)90523-X.
- G. R. Iisley, J. Fisher, R. Apweiler, A. H. DePace and N. M. Luscombe, Cellular resolution models for even skipped regulation in the entire *Drosophila* embryo, *eLife*, 2 (2013).
- P. Ingham, N. Baker, A. Martinez-Arias, Regulation of segment polarity genes in the *Drosophila* blastoderm by *fushi tarazu* and *even skipped*, *Nature*, 331 (1988) 73-75, doi: 10.1038/331073a0.
- H. Janssens, Lack of tailless leads to an increase in expression variability in *Drosophila* embryos, *Developmental Biology*, vol. 377 (2013) 305-317.
- K. Kozlov, E. Myasnikova, J. Reinitz and D. Kosman, Method for spatial registration of the expression patterns of *Drosophila* segmentation genes using wavelets, *Computational Technologies*, 5 (2000) 112-119.
- B. Lim, T. Fukaya, T. Heist and M. Levine, Temporal dynamics of pair-rule stripes in living *Drosophila* embryos, *PNAS*, 115 (2018) 8376-8381, doi: 10.1073/pnas.1810430115.

- E. Myasnikova, A. Samsonova, K. Kozlov, M. Samsonova and J. Reinitz, Registration of the expression patterns of *Drosophila* segmentation genes by two independent methods, *Bioinformatics*, 17 (2001) 3-12, doi: 10.1038/nrg1724.
- C. Nusslein-Volhard, *Coming to life: how genes drive development*, Yale University Press, 2006.
- S. M. Parkhurst and D. Ish-Horowicz, Mis-regulating segmentation gene expression in *Drosophila*, *Development*, 111 (1991) 1121-1135.
- A. D. Peel, A. D. Chipman and M. Akam, Arthropod segmentation: Beyond the *Drosophila* paradigm, *Nature Reviews Genetics*, 12 (2005) 905-916, doi: 10.1038/nrg1724.
- A. Pisarev, E. Poustelnikova, M. G. Samsonova and J. Reinitz, FlyEx, the quantitative atlas on segmentation gene expression at cellular resolution, *Nucleic Acids Research*, 37 (2008) D560 - D566.
- E. Poustelnikova, A. Pisarev, M. Blagov, M. Samsonova and J. Reinitz, A database for management of gene expression data in situ, *Bioinformatics*, 20 (2004) 2212-2221.
- M. D. Schroeder, C. Greer and U. Gaul, How to make stripes: deciphering the transition from non-periodic to periodic patterns in *Drosophila* segmentation, *Development*, 138 (2011) 3067-3078, doi: 10.1242/dev.062141.
- M. Simpson-Brose, J. Treisman and C. Desplan, Synergy between the hunchback and bicoid morphogens is required for anterior patterning in *Drosophila*, *Cell*, 78 (1994) 855-865, doi: 10.1016/S0092-8674(94)90622-X.
- S. Small, R. Kraut, T. Hoey, R. Warrior and M. Levine, Transcriptional regulation of a pair-rule stripe in *Drosophila*, *Genes and Development*, 5 (1991) 827-839, doi: 10.1101/gad.5.5.827.
- S. Small and A. Blair and M. Levine, Regulation of even-skipped stripe 2 in the *Drosophila* embryo, *The EMBO Journal*, 1 (1992) 4047-4057.
- S. Small, A. Blair and M. Levine, Regulation of Two Pair-Rule Stripes by a Single Enhancer in the *Drosophila* Embryo, *Developmental Biology*, 175 (1996) 314-324, doi: 10.1006/dbio.1996.0117.
- S. Sonneveld, B. M. P. Verhagen and M. E. Tanenbaum, Heterogeneity in mRNA translation, *Trends in Cell Biology*, 30 (2020) 606-618, doi: 10.1016/j.tcb.2020.04.008.
- D. Stanojevic, S. Small and M. Levine, Regulation of a Segmentation Stripe by Overlapping Activators and Repressors in the *Drosophila* Embryo, *Science*, 254 (1991) 1385-1387, doi: 10.1126/science.1683715.
- S. Surkova, D. Kosman, K. Kozlov, Manu, E. Myasnikova, A. A. Samsonova, A. Spirov, C. E. Vanario-Alonso, M. Samsonova and J. Reinitz, Characterization of the *Drosophila* segment determination morphome, *Developmental Biology*, 313 (2008) 844-862, doi: 10.1016/j.ydbio.2007.10.037.
- S. Surkova, E. Golubkova, Manu, L. Panok, L. Mamon, J. Reinitz and M. Samsonova, Quantitative dynamics and increased variability of segmentation gene expression in the *Drosophila* Krüppel and knirps mutants, *Developmental Biology*, 376 (2013) 99-112, doi: 10.1016/j.ydbio.2013.01.008.
- B. J. Vincent, M. V. Staller, F. López-Rivera, M. D. J. Bragdon, E.C. G. Pym, K. M. Biette, Z. Wunderlich, T. T. Harden, J. Estrada and A. H. DePace, Hunchback is counter-repressed to regulate even-skipped stripe 2 expression in *Drosophila* embryos, *PLoS Genetics*, 14 (2018).

## Appendix A. Stripe-specific *eve* enhancers

### Appendix A.1. Transcriptional regulation model for stripes 1 and 2 of *Eve*, and dual stripe 1+5 of *Ftz*



In the case of *Eve* we considered the gene template  $G \rightarrow G_e$  and, in the case of *Ftz*,  $G \rightarrow G_f$ . The stripes 1 and 2 production models of *Eve* and the production model for dual stripe 1+5 have different rates.

The production of proteins *Eve* or *Ftz* described by the kinetic equations (A.1) are

$$\begin{aligned}
\overline{\text{Eve}}'(t) &= -d\overline{\text{Eve}}(t) + \alpha_1 k_1 F(\cdot) \\
\overline{Kr}'(t) &= k_{-3} \left[ \alpha_2 G_{R1}(0) - \overline{Kr}(t) + \overline{Kr}(0) \right] - k_3 \overline{Kr}(t) F(\cdot) \\
\overline{Kni}'(t) &= k_{-4} \left[ \alpha_3 G_{R2}(0) - \overline{Kni}(t) + \overline{Kni}(0) \right] - k_4 \overline{Kni}(t) F(\cdot) \\
\overline{Gt}'(t) &= k_{-5} \left[ \alpha_4 G_{R3}(0) - \overline{Gt}(t) + \overline{Gt}(0) \right] - k_5 \overline{Gt}(t) F(\cdot) \\
\overline{Tll}'(t) &= k_{-6} \left[ \alpha_6 G_{R4}(0) - \overline{Tll}(t) + \overline{Tll}(0) \right] - k_6 \overline{Tll}(t) F(\cdot) \\
\overline{Hkb}'(t) &= k_{-7} \left[ \alpha_7 G_{R5}(0) - \overline{Hkb}(t) + \overline{Hkb}(0) \right] - k_7 \overline{Hkb}(t) F(\cdot),
\end{aligned}$$


where

$$F(\cdot) = \left( \frac{\overline{G}(0)}{\alpha_0} + \frac{\overline{Kr}(t) - \overline{Kr}(0)}{\alpha_2} + \frac{\overline{Kni}(t) - \overline{Kni}(0)}{\alpha_3} + \frac{\overline{Gt}(t) - \overline{Gt}(0)}{\alpha_4} + \frac{\overline{Tll}(t) - \overline{Tll}(0)}{\alpha_6} + \frac{\overline{Hkb}(t) - \overline{Hkb}(0)}{\alpha_7} \right),$$

$P$  or  $\overline{P} = \alpha_p P$  represent a specific protein. At each anteroposterior  $x$ -coordinate of the embryo, the repressive protein data are introduced at  $\overline{P}(0)$ . The initial conditions for each repressor complex are  $G_{R1}(0) = \dots = G_{R5}(0) = 0$  and  $\overline{G}(0) = 250$ . The initial conditions for *Eve* and *Ftz* models are  $\overline{\text{Eve}}(0) = 0$  and  $\overline{\text{Ftz}}(0) = 0$ .

Concerning weight parameter we have:  $\alpha_0 = \alpha_{G_e/G_f}$ ,  $\alpha_1 = \alpha_{eve/ftz}$ ,  $\alpha_2 = \alpha_{kr}$ ,  $\alpha_3 = \alpha_{kni}$ ,  $\alpha_4 = \alpha_{gt}$ ,  $\alpha_5 = \alpha_{hb}$ ,  $\alpha_6 = \alpha_{tll}$  and  $\alpha_7 = \alpha_{hkb}$ . In all the simulations, we have assumed  $\alpha_0 = 1$ . These conditions are fixed for all the models analysed.

### Appendix A.2. Transcriptional regulation model for dual stripes 3+7 of *Eve*, and dual stripe 2+7 and 3+6 of *Ftz*




The production of protein *Eve* or *Ftz* described by the kinetic equations (A.2) are

$$\begin{aligned}
\overline{\text{Eve}}'(t) &= -d\overline{\text{Eve}}(t) + \alpha_1 k_1 F(\cdot) \\
\overline{Kni}'(t) &= k_{-3} \left[ \alpha_3 G_{R1}(0) - \overline{Kni}(t) + \overline{Kni}(0) \right] - k_3 \overline{Kni}(t) F(\cdot) \\
\overline{Hb}'(t) &= k_{-4} \left[ \alpha_5 G_{R2}(0) - \overline{Hb}(t) + \overline{Hb}(0) \right] - k_4 \overline{Hb}(t) F(\cdot) \\
\overline{Tll}'(t) &= k_{-5} \left[ \alpha_6 G_{R3}(0) - \overline{Tll}(t) + \overline{Tll}(0) \right] - k_5 \overline{Tll}(t) F(\cdot) \\
\overline{Hkb}'(t) &= k_{-6} \left[ \alpha_7 G_{R4}(0) - \overline{Hkb}(t) + \overline{Hkb}(0) \right] - k_6 \overline{Hkb}(t) F(\cdot),
\end{aligned}$$

where

$$F(\cdot) = \left( \frac{\overline{G}(0)}{\alpha_0} + \frac{\overline{Kni}(t) - \overline{Kni}(0)}{\alpha_3} + \frac{\overline{Hb}(t) - \overline{Hb}(0)}{\alpha_5} + \frac{\overline{Tll}(t) - \overline{Tll}(0)}{\alpha_6} + \frac{\overline{Hkb}(t) - \overline{Hkb}(0)}{\alpha_7} \right).$$

Appendix A.3. Transcriptional regulation model for dual stripes 4+6 of *Eve*




The production of protein *Eve* described by the kinetic equations (A.3) is

$$\begin{aligned}
 \overline{\text{Eve}}'(t) &= -d\overline{\text{Eve}}(t) + \alpha_1 k_1 F(\cdot) \\
 \overline{\text{Kni}}'(t) &= k_{-3} \left[ \alpha_3 G_{R1}(0) - \overline{\text{Kni}}(t) + \overline{\text{Kni}}(0) \right] - k_3 \overline{\text{Kni}}(t) F(\cdot) \\
 \overline{\text{Gt}}'(t) &= k_{-4} \left[ \alpha_4 G_{R2}(0) - \overline{\text{Gt}}(t) + \overline{\text{Gt}}(0) \right] - k_4 \overline{\text{Gt}}(t) F(\cdot) \\
 \overline{\text{Hb}}'(t) &= k_{-5} \left[ \alpha_5 G_{R3}(0) - \overline{\text{Hb}}(t) + \overline{\text{Hb}}(0) \right] - k_5 \overline{\text{Hb}}(t) F(\cdot) \\
 \overline{\text{Tll}}'(t) &= k_{-6} \left[ \alpha_6 G_{R4}(0) - \overline{\text{Tll}}(t) + \overline{\text{Tll}}(0) \right] - k_6 \overline{\text{Tll}}(t) F(\cdot) \\
 \overline{\text{Hkb}}'(t) &= k_{-7} \left[ \alpha_7 G_{R5}(0) - \overline{\text{Hkb}}(t) + \overline{\text{Hkb}}(0) \right] - k_7 \overline{\text{Hkb}}(t) F(\cdot),
 \end{aligned}$$

where

$$F(\cdot) = \left( \frac{\overline{G_e}(0)}{\alpha_0} + \frac{\overline{\text{Kni}}(t) - \overline{\text{Kni}}(0)}{\alpha_3} + \frac{\overline{\text{Gt}}(t) - \overline{\text{Gt}}(0)}{\alpha_4} + \frac{\overline{\text{Hb}}(t) - \overline{\text{Hb}}(0)}{\alpha_5} + \frac{\overline{\text{Tll}}(t) - \overline{\text{Tll}}(0)}{\alpha_6} + \frac{\overline{\text{Hkb}}(t) - \overline{\text{Hkb}}(0)}{\alpha_7} \right).$$

Appendix A.4. Transcriptional regulation model for stripe 5 of *Eve*




The production of protein *Eve* described by the kinetic equations (A.4) is

$$\begin{aligned}
 \overline{\text{Eve}}'(t) &= -d\overline{\text{Eve}}(t) + \alpha_1 k_1 F(\cdot) \\
 \overline{\text{Kr}}'(t) &= k_{-3} \left[ \alpha_2 G_{R1}(0) - \overline{\text{Kr}}(t) + \overline{\text{Kr}}(0) \right] - k_3 \overline{\text{Kr}}(t) F(\cdot) \\
 \overline{\text{Gt}}'(t) &= k_{-4} \left[ \alpha_4 G_{R2}(0) - \overline{\text{Gt}}(t) + \overline{\text{Gt}}(0) \right] - k_4 \overline{\text{Gt}}(t) F(\cdot) \\
 \overline{\text{Hb}}'(t) &= k_{-5} \left[ \alpha_5 G_{R3}(0) - \overline{\text{Hb}}(t) + \overline{\text{Hb}}(0) \right] - k_5 \overline{\text{Hb}}(t) F(\cdot) \\
 \overline{\text{Tll}}'(t) &= k_{-6} \left[ \alpha_6 G_{R4}(0) - \overline{\text{Tll}}(t) + \overline{\text{Tll}}(0) \right] - k_6 \overline{\text{Tll}}(t) F(\cdot) \\
 \overline{\text{Hkb}}'(t) &= k_{-7} \left[ \alpha_7 G_{R5}(0) - \overline{\text{Hkb}}(t) + \overline{\text{Hkb}}(0) \right] - k_7 \overline{\text{Hkb}}(t) F(\cdot),
 \end{aligned}$$

where

$$F(\cdot) = \left( \frac{\overline{G_e}(0)}{\alpha_0} + \frac{\overline{\text{Kr}}(t) - \overline{\text{Kr}}(0)}{\alpha_2} + \frac{\overline{\text{Gt}}(t) - \overline{\text{Gt}}(0)}{\alpha_4} + \frac{\overline{\text{Hb}}(t) - \overline{\text{Hb}}(0)}{\alpha_5} + \frac{\overline{\text{Tll}}(t) - \overline{\text{Tll}}(0)}{\alpha_6} + \frac{\overline{\text{Hkb}}(t) - \overline{\text{Hkb}}(0)}{\alpha_7} \right).$$

## Appendix B. Alternative pattern formation: *eve-ftz* epistasis




The production of proteins *Eve* and *Ftz* described by the kinetic equations (B.1) is

$$\begin{aligned}
\overline{\text{Eve}}'(t) &= \frac{\alpha_2 k_{-4} (\alpha_1 G_{R2}(0) - \overline{G}_f(t) + \overline{G}_f(0))}{\alpha_1} - \frac{k_4 \overline{\text{Eve}}(t) \overline{G}_f(t)}{\alpha_1} - k_5 \overline{\text{Eve}}(t) + \frac{\alpha_2 k_1 \overline{G}_e(t)}{\alpha_0} \\
\overline{\text{Ftz}}'(t) &= \frac{\alpha_3 k_{-3} (\alpha_0 G_{R1}(0) - \overline{G}_e(t) + \overline{G}_e(0))}{\alpha_0} - \frac{k_3 \overline{\text{Ftz}}(t) \overline{G}_e(t)}{\alpha_0} - k_6 \overline{\text{Ftz}}(t) + \frac{\alpha_3 k_2 \overline{G}_f(t)}{\alpha_1} \\
\overline{G}_e'(t) &= k_{-3} (\alpha_0 G_{R1}(0) - \overline{G}_e(t) + \overline{G}_e(0)) - \frac{k_3 \overline{\text{Ftz}}(t) \overline{G}_e(t)}{\alpha_3} \\
\overline{G}_f'(t) &= k_{-4} (\alpha_1 G_{R2}(0) - \overline{G}_f(t) + \overline{G}_f(0)) - \frac{k_4 \overline{\text{Eve}}(t) \overline{G}_f(t)}{\alpha_2}.
\end{aligned}$$

Concerning weight parameter we have:  $\alpha_0 = \alpha_{G_e}$ ,  $\alpha_1 = \alpha_{G_f}$ ,  $\alpha_2 = \alpha_{eve}$ ,  $\alpha_3 = \alpha_{ftz}$ . In all the simulations, we have assumed  $\alpha_0 = 1$  and  $\alpha_1 = 1$ .

## Appendix C. Segment-Polarity pattern formation



The production of proteins *Eve*, *Ftz* *Wg* and described by the kinetic equations (C.1) is

$$\begin{aligned}
\overline{\text{Eve}}'(t) &= k_{-3} \left[ \alpha_1 G_{R1}(0) - \overline{\text{Eve}}(t) + \overline{\text{Eve}}(0) \right] + k_3 \overline{\text{Eve}}(t) \left( \frac{\overline{\text{Eve}}(0) - \overline{\text{Eve}}(t)}{\alpha_1} + \frac{\overline{\text{Ftz}}(0) - \overline{\text{Ftz}}(t)}{\alpha_2} - \frac{\overline{G}_{wg}(0)}{\alpha_0} \right) \\
\overline{\text{Ftz}}'(t) &= k_{-4} \left[ \alpha_2 G_{R2}(0) - \overline{\text{Ftz}}(t) + \overline{\text{Ftz}}(0) \right] + k_4 \overline{\text{Ftz}}(t) \left( \frac{\overline{\text{Eve}}(0) - \overline{\text{Eve}}(t)}{\alpha_1} + \frac{\overline{\text{Ftz}}(0) - \overline{\text{Ftz}}(t)}{\alpha_2} - \frac{\overline{G}_{wg}(0)}{\alpha_0} \right) \\
\overline{\text{Wg}}'(t) &= \alpha_3 k_1 \left( \frac{\overline{\text{Eve}}(t) - \overline{\text{Eve}}(0)}{\alpha_1} + \frac{\overline{\text{Ftz}}(t) - \overline{\text{Ftz}}(0)}{\alpha_2} + \frac{\overline{G}_{wg}(0)}{\alpha_0} \right) - k_2 \overline{\text{Wg}}(t).
\end{aligned}$$

Concerning weight parameter we have:  $\alpha_0 = \alpha_{G_{wg}}$ ,  $\alpha_1 = \alpha_{eve}$ ,  $\alpha_2 = \alpha_{ftz}$ ,  $\alpha_3 = \alpha_{wg}$ . In all the simulations, we have assumed  $\alpha_0 = 1$  and  $\alpha_3 = 1$ .