THE PITCH-CLASS INTEGER THEOREM

CALEB SCOTT ALONS

ABSTRACT. Mathematical music theory has assumed without proof that musical notes can be associated with the equivalence classes of \mathbb{Z}_n . We contest the triviality of this assertion, which we call the Pitch-class Integer Theorem (PCIT). Since the existing literature assumes the PCIT without proof, the mathematics to rigorously treat the PCIT does not yet exist. Thus, we construct an axiomatic proof of the PCIT to support the existing mathematical models of music theory.

1. INTRODUCTION

Although musical harmony is well-understood and standardized from the musical perspective, this subject is comparatively undeveloped on the mathematical side. There does not currently exist a well-established and universally agreed upon canon of definitions and concepts, and most papers in this subject provide their own internal definitions for music theory concepts. Moreover, such a canon is far from achieving standardization in mathematical music theory. One particular concept, however, has gained traction and is universally accepted: the association of musical notes with the equivalence classes of finite groups of integers.

Fiore proposed an integer model of pitch, stating that we could translate pitch classes into numbers as follows [1]:

C = 0,	$C\sharp = D\flat = 1,$
D=2,	$D\sharp = E\flat = 3,$
E = 4,	F = 5,
$F\sharp = G\flat = 6,$	G=7,
$G\sharp = A\flat = 8,$	A=9,
$A\sharp = B\flat = 10,$	B = 11.

Fiore assumed that these pitch classes were the elements of the group \mathbb{Z}_{12} . Fiore's model was echoed later by Zhang, who postulated that pitches that are separated by an integer multiple of an octave or are enharmonically equivalent belong to the same pitch class [5]. Zhang's definition of an ordered pitch-class interval assumes that pitch classes correspond to \mathbb{Z}_{12} . Table 1 gives Zhang's pitch-class integer model, which was proposed independently from Fiore [5].

Around the same time, Wright wrote the book *Mathematics and Music* to show how mathematics models music more broadly. In his chapter on harmony, he assumes that intervals are measured in semitones and that the set of intervals is identified with the set \mathbb{Z} of integers, with an integer k corresponding to the interval of k semitones [4]. Under this

Date: March 8, 2024.

	Integer	Possible Notation
	0	$B\sharp, C, D\flat$
	1	$C\sharp, D\flat$
	2	C×, D, Eb
	3	D♯, E♭
	4	$D \times, E, F \flat$
	5	E♯, F, G♭
	6	F#, Gb
	7	F×, G, Ab
	8	G♯, A♭
	9	G×, A, B♭
	10	A♯, B♭
	11	A×, B, Cb
LE 1	. Zhang'	s Pitch-Class Integer Mo

TABLE 1. Zhang's Pitch-Class Integer Model.

assertion, Wright claims that equivalence modulo 12 is octave identification and proposes the following pitch-class integer model: two intervals, k semitones and l semitones, are equivalent modulo octave if and only if $k \equiv l \mod 12$. Kennedy wrote an undergraduate senior thesis that sought to use group theory to qualify *n*-note tonal systems. Her base assertion is to let the group \mathbb{Z}_n be the model for *n*-note tonal systems [2]. That is, Kennedy expanded Wright's model to accommodate a tonal system that includes an arbitrary number of notes, rather than strictly 12. We note that Zhang, Wright, and Kennedy do not cite Fiore's initial paper, nor do they reference each other's research.

Most recently, Tsok, Jelten, and Gamson published an article that cites the research done by Fiore and Zhang [3]. They reinforce the pitch-class integer model and assert that the musical notes form a group of integers modulo 12 such that

$$\mathbb{Z}_{12} = \{ C, C\sharp, D, D\sharp, E, F, F\sharp, G, G\sharp, A, A\sharp, B \}.$$

We observe that pitch-class integer models have been independently developed by several authors and are currently assumed without proof. This paper seeks to formalize the association of musical notes with the equivalence classes of \mathbb{Z}_n , which we call the Pitch-class Integer Theorem (PCIT).

2. PITCHES, OCTAVES, AND TUNING SETS

We first address our underlying assumption in this paper: we assume without proof that sound waves exist and express this assumption by defining pitch.

Definition 2.1 (Pitch, Lower Pitch, Higher Pitch, Same Pitch). A pitch f is a positive real number that corresponds to the frequency in hertz of a sound wave. Let $f_1 < f_2$ be pitches. We say that f_1 is a lower pitch than f_2 and that f_2 is a higher pitch than f_1 . If $f_1 = f_2$, we say that f_1, f_2 are the same pitch.

Definition 2.2 (*kth* Octave of f, Base Octave). Let f be a pitch. The *kth* octave of fis the interval $o_k(f) = [2^k f, 2^{k+1} f], k \in \mathbb{Z}$. The **base octave** of f is $o_0(f) = [f, 2f]$.

Proposition 2.3. Let f, f' be pitches. Then $\forall f' \in \mathbb{R}^+, \exists k \in \mathbb{Z} \text{ such that } f' \in o_k(f)$.

Proof. If $f' \in \mathbb{R}^+$, then $\exists k \in \mathbb{Z}$ such that $f' \in o_k(f)$ because $\bigcup_{k \in \mathbb{Z}} o_k(f) = \mathbb{R}^+$.

Definition 2.4 (Tuning Set). Let $o_k(f)$ be the *k*th octave of f and let $\tau_k = \{\nu_{k,i}\}_{i=0}^n$ be a partition of $o_k(f)$. The **tuning set** $\mathcal{T}_n(f)$ is the set

$$\mathcal{T}_n(f) = \bigcup_{k \in \mathbb{Z}} \tau_k,$$

where $\nu_{k,n} = \nu_{k+1,0}$.

Example 2.5. Let $\mathcal{T}_3(f)$ be a tuning set. Figure 1 shows the octaves $o_k(f)$, $o_{k+1}(f)$ and their respective partitions τ_k , τ_{k+1} .




FIGURE 1. The octaves $o_k(f), o_{k+1}(f)$ with partitions τ_k, τ_{k+1} .

3. STEPS AND STEP SETS

Definition 3.1 (Step, Step Set: Uniform, Non-uniform). Suppose $\mathcal{T}_n(f)$ is a tuning set. A step is a rule $\delta_i : \mathcal{T}_n(f) \to \mathcal{T}_n(f)$ defined by

$$\nu_{k,i+1} = \nu_{k,i} * \delta_i \qquad \forall \nu_{k,i} \in \mathcal{T}_n(f).$$

A step set is the collection of steps $\Delta_n = \{\delta_i\}_{i=0}^{n-1}$. A step set Δ_n is uniform if $\delta_i = \delta$ $\forall i = 0, \ldots, n-1$; otherwise, Δ_n is non-uniform.

The equation $\nu_{k,i+1} = \nu_{k,i} * \delta_i$ says that the pitch that results from the rule δ_i applied to $\nu_{k,i}$ is $\nu_{k,i+1}$. If δ belongs to a uniform step set Δ_n , we denote

$$u_{k,i} * \delta^j = \nu_{k,i} * \underbrace{\delta * \cdots * \delta}_{j}.$$

We describe the relationship between two pitches in a tuning set using steps. We say that the pitches $\nu_{k,p}$, $\nu_{k,q}$ are |p-q| steps apart. If p > q, we say that $\nu_{k,p}$ is p-q steps above $\nu_{k,q}$, and if p < q, we say that $\nu_{k,p}$ is q-p steps below $\nu_{k,q}$.

Example 3.2. Let $\mathcal{T}_4(f)$ be a tuning set. We define a non-uniform step set Δ_4 , where each step $\delta_i \in \Delta_4$ is defined by $\nu_{k,i+1} = \nu_{k,i} * \delta_i$, $\forall \nu_{k,i} \in \mathcal{T}_4(f)$. The partition of $o_3(f)$ together with Δ_4 is shown in Figure 2.




FIGURE 2. The partition of $o_3(f)$ together with Δ_4 .

We can use steps to express each $\nu_{3,i} \in o_3(f)$ with respect to $\nu_{3,0}$. See Figure 3.




FIGURE 3. The partition of $o_3(f)$ together with Δ_4 with respect to $\nu_{3,0}$.


Example 3.3. Let $\mathcal{T}_5(f)$ be a tuning set; let Δ_5 be a uniform step set, where each step $\delta \in \Delta_5$ is defined by

$$\nu_{k,i+1} = \nu_{k,i} + \frac{\nu_{k,5} - \nu_{k,0}}{5} \qquad \forall \nu_{k,i} \in \mathcal{T}_5(f).$$

We see that each pitch $\nu_{k,j}$ can be found with the equation

$$\nu_{k,j} = \nu_{k,0} + \left(\frac{\nu_{k,5} - \nu_{k,0}}{5}\right) j.$$

Suppose we choose $\nu_{k,0} = 100$ Hz; we use the above formula to compute pitches in $\mathcal{T}_5(100)$.



Example 3.4. Let $\mathcal{T}_4(f)$ be a tuning set, and let $\Delta_4 = \{\delta_i\}_{i=0}^3$ be a non-uniform step set, where each step $\delta_i \in \Delta_4$ is defined by

$$\nu_{k,i+1} = \begin{cases} 5\nu_{k,0}/4, & i = 0\\ \nu_{k,1} + \nu_{k,0}/5, & i = 1\\ \nu_{k,2} + \nu_{k,0}/20, & i = 2\\ \nu_{k,3} + \nu_{k,0}/2, & i = 3 \end{cases} \quad \forall \nu_{k,i} \in \mathcal{T}_4(f).$$

Suppose $\nu_{k,0} = 500$ Hz; then $\mathcal{T}_4(500)$ together with Δ_4 yields the set of pitches (in Hz)

$$\left\{\cdots, \frac{725}{2}, 375, 500, 625, 725, \cdots\right\}.$$




FIGURE 5. The tuning set $\mathcal{T}_4(500)$ together with Δ_4 defined in Example 3.4.

4. TUNING SPACES

Definition 4.1 (Tuning Space, Standard Pitch). A **tuning space** $\langle \mathcal{T}_n(\varsigma), \Delta_n \rangle$ is a tuning set $\mathcal{T}_n(\varsigma)$ together with a step set Δ_n , where $\varsigma \in \mathbb{R}^+$ is an arbitrarily selected pitch such that $\forall \nu_{k,i} \in \mathcal{T}_n(\varsigma), \nu_{k,i} = 2^k \varsigma * \delta_0 * \cdots * \delta_{i-1}$. The pitch ς is the **standard pitch** of the tuning space. We denote tuning spaces with \mathcal{T}_n , reserving the full notation for explicitly-defined tuning spaces.

Definition 4.2 (Chromatic, Non-chromatic). A tuning space is **chromatic** if its step set is uniform. A **non-chromatic** tuning space is a tuning space that is not chromatic.

Definition 4.3 (*n*-tone Equal Temperament (*n*-TET)). Let $\mathcal{T}_n(\varsigma)$ be a tuning set and Δ_n^{ET} be the step set containing steps δ_{ET} defined by $\nu_{k,j+1} = \sqrt[n]{2} \nu_{k,j}, \forall \nu_{k,j} \in \mathcal{T}_n(\varsigma)$. The tuning space $\langle \mathcal{T}_n(\varsigma), \Delta_n^{\text{ET}} \rangle$ is *n*-tone equal temperament (*n*-TET) tuned to ς . The pitch equation for *n*-TET tuned to ς is

$$\nu_{k,j} = 2^k \varsigma \cdot \sqrt[n]{2^j} = \sqrt[n]{2^{j+kn}} \varsigma.$$

Example 4.4. Let $\langle \mathcal{T}_{12}(440), \Delta_{12}^{\text{ET}} \rangle$ be 12-TET tuned to 440 Hz. We have the pitch equation $\nu_{k,j} = \sqrt[n]{2^{j+kn}} \varsigma = 440 \sqrt[12]{2^{j+12k}} \text{ Hz.}$

Although tedious, we show the full list of base octave computations (in Hz) to honor their historical significance in music:

$$\begin{split} \nu_{0,0} &= 440 \sqrt[12]{2^0} = 440, & \nu_{0,7} = 440 \sqrt[12]{2^7} = 440 \sqrt[12]{128} \approx 659.255, \\ \nu_{0,1} &= 400 \sqrt[12]{2^1} = 440 \sqrt[12]{2} \approx 466.164, & \nu_{0,8} = 440 \sqrt[12]{2^8} = 440 \sqrt[3]{4} \approx 698.456, \\ \nu_{0,2} &= 440 \sqrt[12]{2^2} = 440 \sqrt[6]{2} \approx 493.883, & \nu_{0,9} = 440 \sqrt[12]{2^9} = 440 \sqrt[4]{8} \approx 739.989, \\ \nu_{0,3} &= 440 \sqrt[12]{2^3} = 440 \sqrt[4]{2} \approx 523.251, & \nu_{0,10} = 440 \sqrt[12]{2^{10}} = 440 \sqrt[6]{32} \approx 783.989, \\ \nu_{0,4} &= 440 \sqrt[12]{2^4} = 440 \sqrt[3]{2} \approx 554.365, & \nu_{0,11} = 440 \sqrt[12]{2^{11}} = 440 \sqrt[12]{2^{048}} \approx 830.609, \\ \nu_{0,5} &= 440 \sqrt[12]{2^5} = 440 \sqrt[12]{32} \approx 587.330, & \nu_{0,12} = 440 \sqrt[12]{2^{12}} = 880. \\ \nu_{0,6} &= 440 \sqrt[12]{2^6} = 440\sqrt{2} \approx 622.254, \end{split}$$

Definition 4.5 (*n*-equal Divisions of the Octave (*n*-EDO)). Let $\mathcal{T}_n(\varsigma)$ be a tuning set and Δ_n^{EDO} be the step set containing steps δ_{EDO} defined by $\nu_{k,j+1} = \nu_{k,j} + \nu_{k,0}/n, \forall \nu_{k,j} \in \mathcal{T}_n(\varsigma)$. The tuning space $\langle \mathcal{T}_n(\varsigma), \Delta_n^{\text{EDO}} \rangle$ is *n*-equal divisions of the octave (*n*-EDO) tuned to ς . The pitch equation for *n*-EDO tuned to ς is

$$\nu_{k,j} = 2^k \varsigma + \frac{2^k j \varsigma}{n} = 2^k \varsigma \left(1 + \frac{j}{n}\right).$$

5. OCTAVE EQUIVALENCE AND NOTES

Theorem 5.1. Let $\nu_{a,p}, \nu_{b,q} \in \mathcal{T}_n$ a tuning space; let \sim be a relation such that $\nu_{a,p} \sim \nu_{b,q} \iff p = q$. Then \sim is an equivalence relation on \mathcal{T}_n .

Proof. Let $\nu_{a,p}, \nu_{b,q}, \nu_{c,r} \in \mathcal{T}_n$ a tuning space; let \sim be a relation such that $\nu_{a,p} \sim \nu_{b,q} \iff p = q$. Since $\nu_{a,p} \sim \nu_{a,p} \Rightarrow p = p$, then \sim is reflexive. Since $\nu_{a,p} \sim \nu_{b,q} \Rightarrow p = q \Rightarrow q = p \Rightarrow \nu_{b,q} \sim \nu_{a,p}$, then \sim is symmetric. Since $\nu_{a,p} \sim \nu_{b,q}, \nu_{b,q} \sim \nu_{c,r} \Rightarrow p = q, q = r \Rightarrow p = r \Rightarrow \nu_{a,p} \sim \nu_{c,r}$, then \sim is transitive.

Definition 5.2 (Octave Equivalence Relation). The equivalence relation defined in Theorem 5.1 is the octave equivalence relation.

Proposition 5.3. Let $\nu_{x,i} \leq \nu_{y,j} \in \mathcal{T}_n$ a tuning space. Then $\nu_{x,i}, \nu_{y,j}$ are octave equivalent $\iff \nu_{y,j} = 2^{y-x} \nu_{x,i}$.

Proof. Let $\nu_{x,i} \leq \nu_{y,j} \in \mathcal{T}_n$ a tuning space; let \sim be octave equivalence; let $\nu_{x,i} \sim \nu_{y,j}$. Then $\nu_{y,j} = \nu_{y,i}$, and since $\nu_{x,i} \leq \nu_{y,j}$, we have that $x \leq y$. If x = y, then

$$\nu_{y,j} = 2^{y-x} \nu_{x,i} \Rightarrow \nu_{y,i} = 2^{y-y} \nu_{y,i}$$
$$\Rightarrow \nu_{y,i} = \nu_{y,i}.$$

If x < y, we show that

$$2^{y-x}\nu_{x,i} = 2^{x+(y-x)}\varsigma * \delta_0 * \cdots * \delta_{i-1}$$
$$= 2^y\varsigma * \delta_0 * \cdots * \delta_{i-1}$$
$$= \nu_{y,i}$$
$$= \nu_{u,j}.$$

We now assume $\nu_{y,j} = 2^{y-x} \nu_{x,i}$. We show that

$$2^{y}\varsigma * \delta_0 * \cdots * \delta_{j-1} = 2^{y-x} \cdot 2^x\varsigma * \delta_0 * \cdots * \delta_{i-1} = 2^y\varsigma * \delta_0 * \cdots * \delta_{i-1}.$$

Then i = j, so $\nu_{x,i} \sim \nu_{y,j}$.

Definition 5.4 (Note, Chromatic Note). Let ~ be the octave equivalence relation on a tuning space \mathcal{T}_n . A **note** is an equivalence class $[\nu_{k,i}] = \{\nu_{k,x} \in \mathcal{T}_n \mid \nu_{k,x} \sim \nu_{k,i}\}$. The notes of a chromatic tuning space are called **chromatic notes**.

Lemma 5.5. Let \mathcal{T}_n be a tuning space with notes $[\nu_{k,i}]$, $i = 0, \ldots, n-1$. Then $[\nu_{k,n}] = [\nu_{k,0}]$.

Proof. Let \mathcal{T}_n be a tuning space with notes $[\nu_{k,i}]$, $i = 0, \ldots, n-1$. Definition 2.4 gives that $[\nu_{k,n}] = [\nu_{k+1,0}]$, so we show that

$$\begin{aligned} [\nu_{k,n}] &= [\nu_{k+1,0}] \\ &= \{\nu_{k+1,x} \in \mathcal{T}_n \mid \nu_{k+1,x} \sim \nu_{k+1,0}\} \\ &= \{\dots, \nu_{-1,0}, \nu_{0,0}, \nu_{1,0}, \nu_{2,0}, \nu_{3,0}, \dots\} \\ &= \{\nu_{k,x} \in \mathcal{T}_n \mid \nu_{k,x} \sim \nu_{k,0}\} \\ &= [\nu_{k,0}]. \end{aligned}$$

Example 5.6. Let 12-TET be tuned to 440 Hz. The 12 notes of 12-TET have been given designated musical notations which are shown in Table 2. We formalize two significant notes in music.

Definition 5.7 (Concert A, Middle C). Let 12-TET be tuned to 440 Hz with notes given by Table 2. Concert A is the pitch $\nu_{0,0} \in A$, and middle C is the pitch $\nu_{-1,3} \in C$.

Definition 5.8 (Tuning Note). The **tuning note** of a tuning space \mathcal{T}_n is the note $[\nu_{k,0}]$. Moreover, the tuning note contains the standard pitch of \mathcal{T}_n .

$[u_{k,i}]$	Musical Notations	
$[u_{k,0}]$	G×, A, B♭	
$[u_{k,1}]$	$A\sharp, B\flat$	
$[u_{k,2}]$	$A \times, B, C \flat$	
$[u_{k,3}]$	$B\sharp, C, D\flat$	
$[u_{k,4}]$	C \sharp , D \flat	
$[u_{k,5}]$	C×, D, E♭	
$[u_{k,6}]$	$D\sharp, E\flat$	
$[u_{k,7}]$	$D \times, E, F \flat$	
$[u_{k,8}]$	E♯, F, G♭	
$[u_{k,9}]$	$F\sharp, G\flat$	
$[u_{k,10}]$	F×, G, A♭	
$[\nu_{k,11}]$	$G\sharp, A\flat$	
a chromatic notes of 19 TET t		

TABLE 2. The chromatic notes of 12-TET tuned to 440 Hz.

6. HARMONY GROUPS

Lemma 6.1. Let $[\nu_{k,i}]$ be a note of \mathcal{T}_n a tuning space. Then $[\nu_{k,i}] * \delta_i = [\nu_{k,i+1}]$. Proof. Let $[\nu_{k,i}] = \{\nu_{k,x} \in \mathcal{T}_n \mid \nu_{k,x} \sim \nu_{k,i}\}$ be a note of \mathcal{T}_n a tuning space. Then $[\nu_{k,i}] * \delta_i = \{\nu_{k,x} \in \mathcal{T}_n \mid \nu_{k,x} \sim (\nu_{k,i} * \delta_i)\} = \{\nu_{k,x} \in \mathcal{T}_n \mid \nu_{k,x} \sim \nu_{k,i+1}\} = [\nu_{k,i+1}].$

Lemma 6.2. Let $[\nu_{k,x}]$ be a note of \mathcal{T}_n a tuning space. Then $[\nu_{k,x}] = [\nu_{k,0}] * \delta_0 * \cdots * \delta_{x-1}$. *Proof.* Let $[\nu_{k,x}]$ be a note of \mathcal{T}_n a tuning space. If x = 0, then trivially we have that $[\nu_{k,x}] = [\nu_{k,0}]$. If 0 < x < n-1, then

$$[\nu_{k,x}] = [\nu_{k,x-x}] * \delta_{x-x} * \dots * \delta_{x-1} = [\nu_{k,0}] * \delta_0 * \dots * \delta_{x-1}$$

Definition 6.3 (Note Set). Let ~ be the octave equivalence relation on a tuning space \mathcal{T}_n . The **note set** \mathcal{N}_n of \mathcal{T}_n is the set containing the notes of \mathcal{T}_n .

Example 6.4. The note set \mathcal{N}_{12} of 12-TET tuned to 440 Hz is

 $\{A, A\sharp, B, C, C\sharp, D, D\sharp, E, F, F\sharp, G, G\sharp\}.$

Table 2 shows that the elements of \mathcal{N}_{12} have 2 or 3 equivalent musical notations.

Proposition 6.5. The order of note set \mathcal{N}_n is n.

Proof. Definition 6.3 gives that $|\mathcal{N}_n| = |\{[\nu_{k,i}]\}_{i=0}^{n-1}| = n.$

Definition 6.6 (Harmonic Addition). The binary operation $+ : (\mathcal{N}_n \times \mathcal{N}_n) \to \mathcal{N}_n$ defined by $[\nu_{k,x}] + [\nu_{k,y}] = [\nu_{k,x+y}]$ is harmonic addition.

Lemma 6.7. Note sets are closed under harmonic addition.

Proof. Let \mathcal{N}_n be a note set; let + be harmonic addition. Suppose $[\nu_{k,x}], [\nu_{k,y}] \in \mathcal{N}_n$. If $0 \le x + y < n$, then $[\nu_{k,x+y}] \in \mathcal{N}_n$. If $n \le x + y \le 2n - 2$, then $0 \le x + y - n < n - 1$ and $[\mu_{k,x+y}] = [\mu_{k,x+y}] =$

$$[\nu_{k,x+y}] = [\nu_{k,n+(x+y-n)}] = [\nu_{k,n}] + [\nu_{k,x+y-n}] = [\nu_{k,0}] + [\nu_{k,x+y-n}] = [\nu_{k,x+y-n}] \in \mathcal{N}_n$$

CALEB SCOTT ALONS

Theorem 6.8. A note set \mathcal{N}_n together with harmonic addition is a group.

Proof. Let $[\nu_{k,a}], [\nu_{k,b}], [\nu_{k,c}] \in \mathcal{N}_n$ a note set; let + be harmonic addition. We show that + is associative:

$$([\nu_{k,a}] + [\nu_{k,b}]) + [\nu_{k,c}] = [\nu_{k,a+b}] + [\nu_{k,c}]$$

= $[\nu_{k,(a+b)+c}]$
= $[\nu_{k,a+(b+c)}]$
= $[\nu_{k,a}] + [\nu_{k,b+c}]$
= $[\nu_{k,a}] + ([\nu_{k,b}] + [\nu_{k,c}])$

Since $\forall [\nu_{k,x}] \in \mathcal{N}_n$, $[\nu_{k,0}] + [\nu_{k,x}] = [\nu_{k,x}] + [\nu_{k,0}] = [\nu_{k,x}]$, we see that + has identity $[\nu_{k,0}] \in \mathcal{N}_n$. We lastly show that $\forall [\nu_{k,a}] \in \mathcal{N}_n$, $\exists [\nu_{k,n-a}] \in \mathcal{N}_n$ such that $[\nu_{k,a}] + [\nu_{k,n-a}] = [\nu_{k,n-a}] + [\nu_{k,a}] = [\nu_{k,n}] = [\nu_{k,n}]$.

Definition 6.9 (Harmony Group). A harmony group of n notes \mathcal{H}_n is a note set \mathcal{N}_n together with harmonic addition.

7. THE PITCH-CLASS INTEGER THEOREM

Theorem 7.1 (Pitch-class Integer Theorem). Let \mathcal{H}_n be a harmony group of n notes. Then $\mathcal{H}_n \simeq \mathbb{Z}_n$.

Proof. Let $\varphi : \mathcal{H}_n \to \mathbb{Z}_n$ be a map defined by $\varphi([\nu_{k,x}]) = x$. Since $\varphi([\nu_{k,x}]) = \varphi([\nu_{k,y}]) \Rightarrow x = y$, then φ is injective. We show that $\forall z \in \mathbb{Z}_n, \exists [\nu_{k,z}] \in \mathcal{H}_n$ such that $\varphi([\nu_{k,z}]) = z$, so φ is surjective. Since $\varphi([\nu_{k,x}] + [\nu_{k,y}]) = \varphi([\nu_{k,x+y}]) = x + y = \varphi([\nu_{k,x}]) + \varphi([\nu_{k,y}])$, we conclude that φ is a bijective homomorphism, so $\mathcal{H}_n \simeq \mathbb{Z}_n$.

ACKNOWLEDGMENTS

I first thank the LORD for his guidance, patience, grace, and mercy. I would not have this paper without his help and mentoring. I would like to thank both Dr. Andrew S.I.D. Lang and Dr. Elizabeth Carlson for the many discussions involving this paper. I would also like to thank Daniel T. Hanson for helpful discussions. This work was partially supported by the Stanley White (SW) Foundation through the Oral Roberts University Computing & Mathematics Department.

References

- [1] T. M. Fiore. Music and mathematics.
- [2] K. Kennedy. Mathematics and music: Using group theory to qualify n-note tonal systems. Undergraduate Thesis, May 2015.
- [3] S. H. Tsok, N. B. Jelten, and D. G. Gamson. Application of group theory to musical notes. African Journal of Mathematics and Statistics Studies, 1(1):10–20, May 2018.
- [4] D. Wright. Mathematics and Music. Washington University, St. Louis, MO 63130, April 2009.
- [5] A. Zhang. The framework of music theory as represented with groups. Technical report, Carnegie Mellon University, January 2009.

(Caleb Scott Alons) Computing & Mathematics Department, Oral Roberts University, Tulsa, OK 74136, USA

Email address, Caleb Scott Alons: calebalons@oru.edu