## JONATHAN ROSENBERG

ABSTRACT. We discuss twisted cohomology, not just for ordinary cohomology but also for K-theory and other exceptional cohomology theories, and discuss several of the applications of these in mathematical physics. Our list of applications is by no means exhaustive, but we are hoping that it is extensive enough to give the reader a feel for the possible applications of twisted theories in many different contexts. We also give many suggestions for further reading, but this subject has now expanded to the point where the bibliography is necessarily very incomplete.

# Contents

1. Twisted ordinary cohomology	1
1.1. Cohomology with local coefficients	1
1.2. Twisted de Rham cohomology	2
2. Twisted generalized cohomology theories	3
2.1. "Classical" twisted K-theory	3
2.2. "Exotic" twisted K-theory	5
2.3. Twisted theories in general	6
3. Some physical applications	7
3.1. D-brane charges in Type II string theory	7
3.2. Topological T-duality	8
3.3. The Verlinde formula and the work of Freed-Hopkins-Teleman	9
3.4. Brane charges in the WZW model	9
3.5. Orientifold brane charges	10
3.6. Applications in condensed matter theory	10
3.7. Applications of other twisted cohomology theories	10
References	11

## 1. TWISTED ORDINARY COHOMOLOGY

1.1. Cohomology with local coefficients. The story of twisted cohomology begins with the classical subject of *cohomology with local coefficients* [101]. One way to motivate this is to think about the Leray-Serre spectral sequence of a fibration  $F \to E \xrightarrow{p} B$ , where F, E, and B are nice topological spaces, say CW-complexes. When B is simply connected, the Leray-Serre spectral sequence

<sup>2020</sup> Mathematics Subject Classification. Primary 55N25; Secondary 19L50 81T50.

 $Key\ words\ and\ phrases.$  local coefficients, twisted cohomology, anomaly, K-theory, generalized cohomology.

 $H^p(B, H^q(F; G)) \Rightarrow H^{p+q}(E; G)$  enables one in many cases to compute the cohomology of E in terms of the cohomologies of B and F. However, when B is path connected but not simply connected, the fundamental group of B can act nontrivially on the cohomology  $H^q(F; G)$  of the fiber, and  $x \mapsto H^q(p^{-1}(x); G)$  can become a nontrivial locally constant sheaf on the base B. The cohomology of this sheaf is called cohomology with local coefficients. In the case where F, E, and Bare the classifying spaces of groups K, L, and M, and the fibration  $F \to E \xrightarrow{p} B$ comes from a group extension  $1 \to K \to L \to M \to 1$ , the Leray-Serre spectral sequence specializes to the Hochschild-Serre spectral sequence of the group extension  $H^p(M, H^q(K; G)) \Rightarrow H^{p+q}(L; G)$ , where one has to take into account the action of the quotient group M on the cohomology of the normal subgroup K.

Another important place where cohomology with local coefficients is needed is in expressing Poincaré duality for non-orientable manifolds. Suppose  $M^n$  is a connected compact manifold. If M is orientable, Poincaré duality says that cap product with the fundamental homology class  $[M] \in H_n(M; \mathbb{Z})$  (for some choice of orientation) gives an isomorphism from  $H^k(M; \mathbb{Z})$  to  $H_{n-k}(M; \mathbb{Z})$ . When M is not orientable, this can no longer work since there is no fundamental homology class  $[M] \in H_n(M; \mathbb{Z})$ ; in fact  $H_n(M; \mathbb{Z}) = 0$ . In this case, the first Stiefel-Whitney class  $w_1 \in H^1(M; \mathbb{Z}/2)$  is non-zero. We can repair Poincaré duality by observing that there is a twisted fundamental class [M] in  $H_n(M)$  with local coefficients given by  $w_1$ , and that cap product with [M] implements an isomorphism from  $H^k(M; \mathbb{Z})$  to  $H_{n-k}(M; \mathbb{Z})$ , where  $\mathbb{Z}$  denotes the locally constant sheaf locally isomorphic to  $\mathbb{Z}$ on which  $\pi_1(M)$  acts via  $w_1(M)$ . (Think of  $w_1$  as taking values in  $\{\pm 1\}$  and let it act on  $\mathbb{Z}$  by multiplication.) This was first observed by Steenrod in [101] and has played a big role in manifold theory ever since.

1.2. Twisted de Rham cohomology. Cohomology with local coefficients is the first example of twisted cohomology. But there are other twists of ordinary cohomology, even of de Rham cohomology on manifolds (which is naturally isomorphic to ordinary cohomology with coefficients in  $\mathbb{R}$  or  $\mathbb{C}$ ). The following construction seems to have originally been "folklore," but a hint of it appears in [9], with a longer discussion in [18, §9.3], and a full exposition in [104].

Let M be a manifold and let  $\alpha$  be a closed k-form on M, where k is odd. Then  $d'_{\alpha} = d + \alpha \wedge$  maps even forms to odd forms and odd forms to even forms. (However, it increases degree by 1 only when  $\alpha$  is a 1-form, i.e., k = 1.) And  $d'_{\alpha}$  is a differential, i.e.,

$$(d'_{\alpha})^{2}\omega = (d + \alpha \wedge)(d\omega + \alpha \wedge \omega) = d^{2}\omega - \alpha \wedge d\omega + \alpha \wedge d\omega + (\alpha \wedge \alpha) \wedge \omega = 0$$

for any form  $\omega$ . Thus the cohomology  $H^{\bullet}_{\alpha}(M)$  of the 2-periodic complex

$$\cdot \xrightarrow{d'_{\alpha}} \Omega^{\operatorname{even}}(M) \xrightarrow{d'_{\alpha}} \Omega^{\operatorname{odd}}(M) \xrightarrow{d'_{\alpha}} \Omega^{\operatorname{even}}(M) \xrightarrow{d'_{\alpha}} \cdots$$

can be called the *twisted cohomology* of M with twist  $\alpha$ . A simple calculation [104, Theorem 5] shows that  $d'_{\alpha}$  and  $d'_{\alpha'}$  are conjugate under  $e^{-\eta}$  if  $\alpha' - \alpha = d\eta$ , so  $H^{\bullet}_{\alpha}(M)$  only depends on the de Rham class of  $\alpha$ . Note that (unless  $\alpha$  is a 1-form) we will only think of it as being  $\mathbb{Z}/2$ -graded rather than  $\mathbb{Z}$ -graded, but when  $\alpha = 0$ , it coincides with the usual de Rham cohomology of M,  $\mathbb{Z}/2$ -graded via even and odd degree forms. Also note (see [104, §3.3]) that  $d'_{\alpha}$  is the total differential of a double complex, where the horizontal differential is the usual differential d and the vertical differential is  $\alpha \wedge$ . In the spectral sequence of the double complex, the first

differential  $d_1$  is d and so  $E_2$  is just the usual de Rham cohomology  $H^{\bullet}_{deR}(M)$ , and the next differential  $d_k$  is cup product with the de Rham class  $[\alpha]$  of  $\alpha$ . So the next stage  $E_{k+1}$  in the spectral sequence for  $H^{\bullet}_{\alpha}(M)$  is the cohomology of  $[\alpha] \cup \_$  acting on  $H^{\bullet}_{deR}(M)$ . However, in general there can be still higher differentials. Atiyah and Segal [11, Proposition 6.1] have shown that these higher differentials all come from iterated Massey products with  $\alpha$ .

There is one case where twisted cohomology reduces to cohomology with local coefficients. Assume k = 1 so that  $\alpha$  is a 1-form. Since  $H^1_{deR}(M) \cong H^1(M; \mathbb{R}) \cong$ Hom $(\pi_1(M), \mathbb{R}), [\alpha]$  can be identified with a group homomorphism  $\pi_1(M) \to \mathbb{R}$ . There is an associated locally constant sheaf  $\mathcal{L}$  and one can easily check that  $H^{\bullet}_{\alpha}(M) \cong H^{\bullet}(M, \mathcal{L})$ , which is cohomology with local coefficients [104, §3.1].

The kind of twisted cohomology we have just been discussing is based on de Rham cohomology, so it ignores torsion phenomena. In [37] is a proposal for a kind of integral twisted cohomology based on homotopy theory. Namely, given a cohomology class  $\alpha \in H^k(X; \mathbb{Z})$  (the main case of interest is when k = 3), one can identify  $\alpha$  with the classifying invariant for a principal fibration  $K(\mathbb{Z}, k-1) \to E \to X$ . The Serre spectral sequence for this fibration will have  $E_2^{p,0} = H^p(X; \mathbb{Z})$  and first differential  $d_k$  sending the generator  $\iota$  of  $H^{k-1}(K(\mathbb{Z}, k-1); \mathbb{Z}) \cong \mathbb{Z}$  to  $\alpha \in H^k(X; \mathbb{Z})$ , and sending  $x \otimes \iota$  to  $\pm \alpha \cup x$ , just as for the spectral sequence in twisted de Rham cohomology. It is proposed in [37] to take the  $\alpha$ -twisted cohomology of X to be the cohomology of E, though this has the drawbacks that if  $\alpha = 0$ , it computes  $H^{\bullet}(X \times K(\mathbb{Z}, k-1))$ and not  $H^{\bullet}(X)$ , and even if X is finite-dimensional, this will usually have cohomology in arbitrarily large degree. So it seems a more reasonable definition would be the image  $E_{\infty}^{p,0}$  of the edge homomorphism  $H^p(X; \mathbb{Z}) \to H^p(E; \mathbb{Z})$ , which agrees with  $H^p(X; \mathbb{Z})$  when  $\alpha = 0$  and always vanishes beyond the dimension of X.

# 2. Twisted generalized cohomology theories

2.1. "Classical" twisted K-theory. When it comes to recent impact on mathematical physics, one can easily argue that twists of K-theory and other generalized cohomology theories (not satisfying the Eilenberg-Mac Lane dimension axiom) have played a much bigger role than twisted ordinary cohomology. We shall begin with K-theory, the most familiar and important generalized cohomology theory, and then move on later to other theories.

The story of twisted K-theory begins with the work of Karoubi and Donovan [42, 43]. They actually worked with  $\mathbb{Z}/2$ -graded algebras, but for simplicity we'll ignore this here. Then they observed that if one looks at Azumaya algebras over a compact Hausdorff space X (algebras of the form  $A = \Gamma(X, \mathcal{A})$ , where  $\mathcal{A}$  is a locally trivial bundle over X with fibers of the form  $M_n(D)$ , the  $n \times n$  matrices over a finite-dimensional division algebra D over  $\mathbb{R}$  or  $\mathbb{C}$ , so that  $D \cong \mathbb{R}$ ,  $\mathbb{C}$  or  $\mathbb{H}$ ), the K-theory of the algebra A looks like a version of  $KO^{\bullet}(X)$ ,  $KU^{\bullet}(X)$ , or  $KSp^{\bullet}(X)$  (depending on which division algebra one is using) with local coefficients, where the local coefficients come from the twisting of the bundle  $\mathcal{A}$ . In, say, the complex case where  $\mathcal{A}$  has fibers  $M_n(\mathbb{C})$ ,  $\mathcal{A}$  is classified by the homotopy class of a map from X to the classifying space  $BPGL_n(\mathbb{C}) \simeq BPU(n)$ , and because of the Morita invariance of K-theory, the K-theory of A only depends on the Brauer group class of the Azumaya algebra, which is a torsion class in  $H^3(X;\mathbb{Z})$ . (In fact, by a theorem of Serre, every torsion class in  $H^3(X;\mathbb{Z})$  arises this way.)

The theory of continuous-trace algebras, for which a good reference is [89], provides a substitute for Azumaya algebras when the characteristic class in  $H^3(X;\mathbb{Z})$ , now called the Dixmier-Douady class, is no longer torsion. A "stable" continuoustrace algebra, say over a connected second-countable locally compact Hausdorff space X, is a C<sup>\*</sup>-algebra of the form  $A = \Gamma_0(X, \mathcal{A})$ , where  $\mathcal{A}$  is a locally trivial bundle over X with fibers isomorphic to  $\mathcal{K}$ , the compact operators on a separable infinite-dimensional Hilbert space  $\mathcal{H}$  (over  $\mathbb{R}$ ,  $\mathbb{C}$  or  $\mathbb{H}$ ), and structure group the  $C^*$ -algebra automorphisms of  $\mathcal{K}$ . Such an algebra is classified by the homotopy class of a map from X to the classifying space  $BPU(\mathcal{H})$ . Since the unitary group  $U(\mathcal{H})$  is contractible (in the natural topology, the strong operator topology), this classifying space has the homotopy type of the Eilenberg-Mac Lane space  $K(\mathbb{Z},3)$ in the complex case (since the center of  $U(\mathcal{H})$  is the circle group  $\mathbb{T}$  and thus  $PU(\mathcal{H})$ is a  $K(\mathbb{Z},2)$ , or  $K(\mathbb{Z}/2,2)$  in the real or quaternionic cases (since then the center of  $U(\mathcal{H})$  consists just of  $\{\pm 1\}$ , and thus  $PU(\mathcal{H})$  is a  $K(\mathbb{Z}/2,1)$ ). For details on all of this see [91]. Thus given  $\delta \in H^3(X;\mathbb{Z})$ , one has twisted K-groups  $K^{\bullet}_{\delta}(X)$  in the complex case, and given  $\delta \in H^2(X; \mathbb{Z}/2)$ , one has twisted K-groups  $KO^{\bullet}_{\delta}(X)$  and  $KSp^{\bullet}_{\bullet}(X)$  in the real and quaternionic cases, obtained by taking the topological K-theory of the appropriate continuous-trace algebras.

There is a slight technical complication in that to get a functorial map  $K^{\bullet}_{\delta'}(Y) \to K^{\bullet}_{\delta}(X)$ , one needs a map  $f: X \to Y$  that induces a \*-homomorphism of the appropriate  $C^*$ -algebras, and thus a map of bundles. This involves a choice of Čech cocycles representing  $\delta$  and  $\delta'$  such that f carries one cocycle to the other. More formal definitions of twisted K-theory and discussions of this and other fine points may be found in [10, 11] and [68].

As one would expect, there is a compatibility between twisted K-theory and twisted ordinary cohomology, given by the Chern character. If M is a compact manifold (the compactness is not so essential, but is assumed here to avoid complications if M is not of finite type) and  $\delta \in H^3(M; \mathbb{Z})$ , then the (twisted) Chern character gives a natural transformation from  $K^{\bullet}_{\delta}(M)$  to  $H^{\bullet}_{\delta}(M)$  which is an isomorphism after tensoring the K-theory with  $\mathbb{R}$ . (Here one should take a closed 3-form representing the integral cohomology class  $\delta$ .) For references, see [18], [79], and [11].

One immediate use for twisted K-theory, pointed out in the original work by Karoubi and Donovan, involves the Thom isomorphism. Suppose X is a compact Hausdorff space and V is a real vector bundle of rank k over X. If V is orientable, a choice of orientation on V gives rise to a Thom isomorphism  $\tau: H^{\bullet}(X) \xrightarrow{\cong} H_c^{k+\bullet}(V)$ , where the cohomology on the right is taken with compact support. (Alternatively, one could take the relative cohomology of the disk bundle of V relative to its boundary the sphere bundle, or take the reduced cohomology of the Thom space.) When V is not orientable, one gets a Thom isomorphism only if the cohomology of V (with compact support) is replaced by twisted cohomology with twist  $w_1, w_1$  the first Stiefel-Whitney class of the bundle.

This suggests a similar picture for K-theory. (Here we only deal with complex K-theory for simplicity, but the same would work for KO with spin<sup>c</sup> replaced by spin.) Suppose V is equipped with a spin<sup>c</sup> structure. Then again one gets a Thom isomorphism  $\tau \colon K^{\bullet}(X) \xrightarrow{\cong} K^{k+\bullet}(V)$ , where the K-theory is taken with compact support. (A proof of this when V is a complex vector bundle is in [8, Theorem 4.7].) If V does *not* admit a spin<sup>c</sup> structure, then this fails, but it can be repaired

4

by replacing  $K^{k+\bullet}(V)$  with twisted K-theory. For example, if V is oriented but not spin<sup>c</sup>, one should use  $K^{k+\bullet}_{W_3(V)}(V)$ , the twisted K-theory for the obstruction  $W_3(V) \in \text{Tors}\,H^3(V;\mathbb{Z})$  to a spin<sup>c</sup> structure on V (see [43, §6] or [68, §4]).

In general it is not so obvious how to compute twisted K-theory, but there are two useful techniques in this direction. One is the (twisted) Atiyah-Hirzebruch spectral sequence  $H^p(X, K^q) \Rightarrow K_{\delta}^{p+q}(X)$ . This has the same  $E_2$  term as the Atiyah-Hirzebruch spectral sequence for untwisted K-theory, but the difference is that the twist  $\delta$  appears in the first nonzero differential  $d_3$ , which is now Sq<sup>3</sup> +  $\delta \cup$ \_ [90, 10]. A second quite different technique is to use a theorem of Khorami [70], which shows that for  $\delta \in H^3(X;\mathbb{Z})$  and P the associated  $PU(\mathcal{H})$ -bundle over X, one has  $K_{\bullet}^{\delta}(X) \cong K_{\bullet}(P) \otimes_{K_0(\mathbb{CP}^{\infty})} \mathbb{Z}$ , where the map  $K_0(\mathbb{CP}^{\infty}) \to \mathbb{Z}$  is rather subtle and is not flat as a map of rings. (The ring structure on  $K_0(\mathbb{CP}^{\infty})$  comes from the Pontryagin product.) Note that this result is formulated in K-twisted homology, but one can dualize via the universal coefficient theorem (which applies to continuous-trace algebras [28]) to compute twisted K-cohomology.

2.2. "Exotic" twisted K-theory. What we have described so far is what one could call "classical" twisted K-theory, known since the work of Karoubi and Donovan in 1969–1970. At this point quite a number of generalizations are known. First of all, there is an equivariant theory for spaces equipped with an action of a compact Lie group G. This is discussed, for example, in [10, §6] or [68, §6]. In the complex (ungraded) case, the twisting is by a class in equivariant cohomology  $H^3_G(X;\mathbb{Z})$ .

A second variant is *twisted* KR-theory, developed in the work of Moutuou [84, 85] and in [44]. To explain this let's start with a review of Atiyah's KR-theory [7]. This is a cohomology theory based on vector bundles on "Real spaces," (locally compact) spaces X equipped with an involutive homeomorphism  $\iota$ . The vector bundles are required to have a conjugate-linear involution compatible with the involution on the base space X. The motivating example of a Real space is the set of complex points of an algebraic variety defined over  $\mathbb{R}$ , with  $\iota$  given by the action of  $\operatorname{Gal}(\mathbb{C}/\mathbb{R})$ . Like KO, KR-theory is 8-periodic, and when  $\iota = \operatorname{id}_X, KR^{\bullet}(X, \iota) = KO^{\bullet}(X)$ . On the other hand, when  $X = Y \sqcup Y$  and  $\iota$  interchanges the two copies of Y, a Real vector bundle on X is uniquely determined by a complex vector bundle on Y, so that  $KR^{\bullet}(X) \cong K^{\bullet}(Y)$ .

However, as we have already seen in the discussion of twisted real K-theory, at some level the division algebras  $\mathbb{R}$  and  $\mathbb{H}$  have to be treated on an equal footing. This insight is reinforced by the fact, due to Donald Anderson, that  $KO^{\bullet}$  and  $KSp^{\bullet}$  are naturally duals of each other, whereas  $K^{\bullet}$  (complex K-theory) is selfdual. This motivates the consideration [44] of a twisted version of KR theory, in which for  $(X, \iota)$  a Real space, each connected component of the fixed set  $X^{\iota}$  of  $\iota$ is assigned a sign  $\pm$ . When all the signs are +, one recovers the usual  $KR^{\bullet}$ , but for each - sign, the restriction of  $KR^{\bullet}_{\pm}(X)$  to the associated component of  $X^{\iota}$  is  $KSp^{\bullet}$  rather than  $KO^{\bullet}$ . Still another version of twisted KR-theory, with some applications to Grothendieck-Witt groups of real algebraic varieties, may be found in [69].

There is another way of twisting K-theory which, even in the case of nonequivariant complex K-theory, produces something new. This approach was first introduced in [2] with special attention to twisted K-theory, and was also studied in [64]. In the next subsection we will see how this generalizes to other cohomology theories.

For simplicity we just deal with complex K-theory. As is well known, the classifying space for stable complex vector bundles, which is also the classifying space for K-theory, is  $\mathbb{Z} \times BU$ , where the  $\mathbb{Z}$  factor is used to record the rank of the bundle and BU is the classifying space of the infinite unitary group. Twisting by a class in  $H^3$ comes, as we have seen before, from the identification of  $BPU(\mathcal{H})$ , the classifying space for the projective unitary group of a separable infinite-dimensional Hilbert space, with a  $K(\mathbb{Z}, 3)$  space, and the fact that one can replace U by unitaries of the form 1 + (compact), which are preserved under conjugation by  $PU(\mathcal{H})$ . However,  $PU(\mathcal{H})$  does not account for the full set of homotopy automorphisms of K-theory. To study these one should build a topological monoid  $GL_1(K)$ , which in other language was computed in [72] — see [1, Chapter 6] and [6] for an explanation. It turns out that

$$GL_1(K) \simeq K(\mathbb{Z}/2, 0) \times K(\mathbb{Z}, 2) \times BSU^{\otimes}$$

and thus

 $BGL_1(K) \simeq K(\mathbb{Z}/2, 1) \times K(\mathbb{Z}, 3) \times BBSU^{\otimes}.$ 

Here the first two factors are "classical": the  $K(\mathbb{Z}/2,0)$  comes from complex conjugation on U and the  $K(\mathbb{Z},2)$  comes from  $PU(\mathcal{H})$ . But the last factor is more mysterious, and leads to new twists of K-theory.  $BSU^{\otimes}$  is the infinite loop space classifying virtual complex vector bundles of rank and determinant one, equipped with the tensor product structure. After localization at any prime p, it is equivalent to the usual BSU, so it has lots of nonzero homotopy groups. Then a homotopically nontrivial map  $X \to BBSU^{\otimes}$  can be used to construct an exotic twist of K-theory, different from the twists that come from elements of  $H^3(X; \mathbb{Z})$ .

2.3. Twisted theories in general. Now that we have examined a large number of examples of twisted cohomology or cohomology with local coefficients, it is time to discuss a general framework for such theories, and to explain how this framework applies to cohomology theories other than ordinary cohomology and K-theory. This theory was developed by Ando, Blumberg, and Gepner *et al.* in [4, 5, 3]. Alternative approaches due to May and Sigurdsson and to Hebestreit, Sagave, and Schlichtkrull may be found in [81, 65, 64]. To avoid very complicated homotopy theoretic problems, the treatment here will be a bit sloppy, but should convey the general idea. For details of how to make everything rigorous, please see the original references.

Let's start with a homotopy-theoretic interpretation of cohomology with local coefficients. Recall that if G is an abelian group, then  $H^k(X; G)$  can be identified with the set of homotopy classes of maps  $X \to K(G, k)$  into the Eilenberg-Mac Lane space K(G, k). For cohomology with local coefficients  $H^k(X, \underline{G})$ ,  $\underline{G}$  a local coefficient system coming from an action of  $\pi_1(X)$  on G, one can similarly identify the local cohomology with  $\pi_1(X)$ -equivariant homotopy classes of equivariant maps from the universal cover  $\widetilde{X}$  of X to K(G, k). This approach won't generalize to more general twisted theories, but we can instead create a fiber bundle

$$E = X \times_{\pi_1(X)} K(G, k) \to X \times_{\pi_1(X)} * = X$$

over X with fibers that are copies of K(G, k). Then  $H^k(X, \underline{G})$  can be identified with homotopy classes of sections of this bundle.

A similar approach will work with more general cohomology theories. A generalized cohomology theory E is always represented in the homotopy category by a spectrum, which is a sequence  $\{E_k\}_{k\geq 0}$  of (based) spaces with connecting maps  $f_k: \Sigma E_k \to E_{k+1}$ . Then we define

$$E^{n}(X) = \varinjlim_{k} [\Sigma^{k-n} X, E_{k}],$$

where  $[\_,\_]$  denotes homotopy classes of (based) maps and we map  $f: \Sigma^{k-n}X \to E_k$  to  $\Sigma f: \Sigma^{k-n+1}X \to \Sigma E_k$  and then compose with  $f_k$  to get a map  $\Sigma^{k-n}X \to E_{k+1}$ . The definition is rigged so that  $E^{n+1}(\Sigma X) \cong E^n(X)$ , which is one of the key properties of a cohomology theory.

For present purposes we will mostly be interested in generalized cohomology theories E that come with a suitable notion of cup products. In this case, we need some extra structure on E, summarized by saying that it is a *ring spectrum*, i.e., is equipped with a suitable multiplication map  $E \wedge E \xrightarrow{\mu_E} E$  that is homotopy associative. There are various axioms to satisfy, but these hold in most of the standard examples, such as ordinary cohomology with coefficients in an abelian group G (where E is what's called an *Eilenberg-Mac Lane spectrum HG*, where  $E_k = K(k, G)$  and  $\Sigma K(k, G) \to K(k + 1, G)$  is adjoint to the equivalence  $K(k, G) \xrightarrow{\simeq} \Omega K(k + 1, G)$ ), real or complex K-theory, and oriented, unoriented, spin, and complex cobordism. Given a ring spectrum E, we also have the notion of an E-module spectrum, which is a spectrum F with a multiplication map  $\mu_F : E \wedge F \to F$  such that

$$E \wedge E \wedge F \xrightarrow{\mu_E \wedge \operatorname{id}_F} E \wedge F$$

$$\downarrow_{\operatorname{id}_E \wedge \mu_F} \qquad \qquad \downarrow_{\mu_F}$$

$$E \wedge F \xrightarrow{\mu_F} F$$

is homotopy commutative.

Twisted *E*-cohomology of *X* comes from taking a bundle of *E*-module spectra over *X*, with fibers that are equivalent to *E*, and taking homotopy classes of sections. In the case of a trivial bundle  $X \times E \xrightarrow{\text{pr}_1} X$ , a section is just a map  $f: X \to E$ , so we recover the homotopy group [X, E], which is  $E^0(X)$ , and similarly after suspending for other values of *n*. The sort of bundle needed is what is called a "parameterized spectrum" in the literature. The set of these bundles is parameterized by homotopy classes of maps  $X \to BGL_1(E)$ , so this definition recovers our original definition of cohomology with local coefficients or twisted *K*-theory. An essentially equivalent definition involves giving a more general notion of Thom space (a "Thom spectrum") and taking its homotopy groups.

### 3. Some physical applications

Let's now outline a few of the major applications of twisted cohomology in mathematical physics.

3.1. **D-brane charges in Type II string theory.** Our first example comes from type II string theory. In this theory, spacetime is a 10-dimensional manifold X (usually spin) and it contains submanifolds  $N \subseteq X$  called *D-branes*, of even dimension in type IIB and of odd dimension in type IIA, on which "open" strings can begin and end. (The letter D stands for "Dirichlet," from Dirichlet boundary conditions.) These D-branes carry *Chan-Paton bundles* and Ramond-Ramond charges. It is now generally believed that D-brane charges should take values in K-theory (of even degree in type IIB, of odd degree in type IIA) rather than in

ordinary cohomology [82, 35, 105]. However, this statement is really only true when the *B*-field of string theory is topologically trivial (i.e., given by a closed 2-form). If this is not the case, then there is an "*H*-flux"  $H \in H^3(X; \mathbb{Z})$ , and D-brane charges should take their values in *H*-twisted *K*-theory [105, 67, 26].

Let's give a quick review of the physical argument for the need for twisted Ktheory. The key fact is that the Freed-Witten anomaly formula on open strings [57] forces, for N a D-brane in spacetime  $X, W_3(N)$  to agree with the pullback of H from X to N. To phrase things another way (see [83, §4] or [67]), neither the gauge field associated to the Chan-Paton bundle nor the B-field as a 2-form are completely well-defined on the D-brane, but the indeterminacies have to cancel so that the path integral is well defined. When H = 0, that means that N is a spin<sup>c</sup> manifold; in fact, in classical IIB string theory, one often thinks of the D-branes as being complex submanifolds of X (which are of course automatically spin<sup>c</sup>). But in general, the D-branes are not spin<sup>c</sup>, so the Poincaré dual of the Chan-Paton class in K-homology is not well-defined. However, the fact that N is "twisted spin<sup>c</sup>" means that one does have a Poincaré dual in H-twisted K-homology, which one can push forward to a class in  $K_{\bullet}^{H}(X)$  or to its dual  $K_{H}^{\bullet}(X)$ .

A slightly different use of twisted K-theory in string theory comes from an attempt to relate type IIB string theory to the  $E_8$  gauge theory in eleven dimensions that underlies M-theory [78]. In this picture the twisting of K-theory in  $H^3(X;\mathbb{Z})$  can be identified with the homotopy class of a map  $X \to E_8$  (since  $E_8$  is a good approximation to a  $K(\mathbb{Z},3)$  space) and since  $[X, E_8] = [X, \Omega B E_8] \cong [\Sigma X, B E_8]$  we obtain an  $E_8$  bundle over a circle bundle Y over X.

3.2. Topological T-duality. The concept discussed above in subsection 3.1, that D-brane charges naturally live in twisted K-theory, has an interesting consequence which can be used to test various physical hypotheses. Namely, physicists predict many "dualities" in which seemingly different mathematical models give rise to the same physical phenomena. An example of such dualities is T-duality (or target-space duality or torus duality) in type II string theory, in which two different type II string theories on possibly different spacetimes X and X', obtained by replacing certain circles in X by their duals, are actually physically equivalent. It was first observed in [20, 21] that under T-duality, in general both the topology of X and the H-flux in  $H^3(X;\mathbb{Z})$  have to change. Twisted K-theory imposes a constraint on this change of topology; since the D-brane charge group is given by twisted K-theory, under a T-duality between (X, H) and (X', H'), one needs to have  $K^{\bullet}_{\mathbf{H}}(X) \cong K^{\bullet+k}_{\mathbf{H}'}(X')$ , where k is the number of T-dualities involved.

The Bouwknegt-Evslin-Mathai observation led to a fairly large industry to studying "topological T-duality," trying to understand the algebraic topology underlying the physics of T-duality. This began with twisted cohomology [25] and moved on to a full understanding of how twisted K-theory is related to the H-flux on a principal torus bundle [74, 31, 30, 75, 34, 38, 92]. Interesting phenomena that occur in the case of higher-dimensional torus bundles are that sometimes T-duals don't exist, are non-unique, or are noncommutative. The interested reader will find a very rich literature on these topics, of which we have only scratched the surface.

Still another variant of this situation comes from studying topological T-duality for *non-principal* circle or torus bundles. For this variant of the theory see [13, 14, 15, 76].

Or one can replace bundles by "bundles with degeneration" or stacks. This generalization of topological T-duality is studied in [32, 86, 33, 80, 87].

Another generalization of topological T-duality is what has been called "spherical T-duality," in which circle bundles are replaced by pricipal SU(2)-bundles. (Recall that SU(2) is topologically just  $S^3$ .) Such bundles are classified by homotopy classes of maps into  $BSU(2) \cong \mathbb{HP}^{\infty}$ . In the simplest cases where one considers SU(2)-bundles over a closed oriented 4-manifold M, these bundles are parametrized by  $H^4(M;\mathbb{Z})\cong\mathbb{Z}$  and the total space of each bundle is a closed oriented 7-manifold P. In [22, 23, 24] it is shown that one gets isomorphisms of twisted cohomology between  $(P, H), H \in H^7(P;\mathbb{Z})$ , and a dual bundle  $(\hat{P}, \hat{H}), \hat{H} \in H^7(\hat{P};\mathbb{Z})$ . There is also an isomorphism of twisted K-theory with twisting H in the more general sense of [2]. When the base of the bundle has dimension bigger than 4, or one considers non-principal SU(2)-bundles, then the situation is more complicated and spherical T-duals may not exist or may be non-unique.

Finally, there are some extensions of topological T-duality that interact with other subjects such as Fourier-Mukai and Langlands duality. For a few results in this direction see [17, 39, 29, 77].

3.3. The Verlinde formula and the work of Freed-Hopkins-Teleman. One of the most striking applications of twisted (equivariant) K-theory to mathematical physics is the work of Freed, Hopkins, and Teleman [52, 53, 55, 54] on using twisted K-theory to explain and understand the "Verlinde formula" [103]. The latter, which on the face of it has nothing to do with twisted K-theory, computes the structure constants for the "fusion rule" for multiplication of the primary fields in the conformal field theory on  $\mathbb{CP}^1$  with three punctures coming from a positive energy (projective) representation of the loop group LG of a compact Lie group G. Roughly speaking, their result is that a positive energy representation determines a twist  $\tau$  in  $H^3_G(G; \mathbb{Z})$  (where G acts on itself by conjugation), the sum of the level of the representation and the dual Coxeter number, and the fusion ring can be identified naturally with  $\tau K^{\bullet}_G(G)$ , the  $\tau$ -twisted equivariant K-theory, with ring operation coming from Pontryagin product. The actual details are more complicated and include many auxiliary results of independent interest.

3.4. Brane charges in the WZW model. Related to the Freed-Hopkins-Teleman theorem is the computation of the non-equivariant twisted K-theory of a compact Lie group G, which should classify the D-branes in the WZW theory of a given level. A first attack on the problem of computing these twisted K-groups was made in [50, 73, 19, 58, 59], followed by a more comprehensive approach by Braun [27], still partially conjectural since it relied on a conjecture about the commutative algebra of Verlinde rings (which is known in some cases but may still be open in others). Finally, using rather complicated techniques from algebraic topology, Douglas proved Braun's result unconditionally in [46], though in a different form. The result of [27] is that for G a simply connected simple compact Lie group of rank n and twisting h > 0 in  $H^3(G;\mathbb{Z}) \cong \mathbb{Z}, K^h_{\bullet}(X)$  is (as a ring) the tensor product of an exterior algebra over  $\mathbb{Z}$  on n-1 odd-degree generators with a finite cyclic group of order c(G,h) = h/gcd(h, y(G)), where the constant y(G) can be described easily in terms of n for the classical groups (for example it is  $lcm(1, 2, \dots, n)$  for  $A_n = SU(n+1)$ ) and where it takes the value 60 for  $G_2$ , 27720 for  $F_4$  and  $E_6$ , lcm $(1, 2, \dots, 17)$  for  $E_7$ , and lcm $(1, 2, \dots, 29)$  for  $E_8$ . Other proofs of many of these results were given

in [77] and [94]. Some of these calculations illuminate certain conjectured physics dualities such as "level-rank duality" or "strange duality."

3.5. Orientifold brane charges. Orientifold string theories are a variant of string theory where one assumes that spacetime X is equipped with an involution  $\iota$ , and the map  $\Sigma \to X$  of a string worldsheet into X is assumed to be equivariant for the worldsheet parity operator  $\Omega$  on  $\Sigma$  and  $\iota$  on X. Such theories include type I string theory, for example, obtained from type IIB string theory on X by taking  $\iota = \mathrm{id}_X$ . Components of the fixed set  $X^{\iota}$  are usually called O-planes ("O" for orientifold). As explained in [105, §5.2] and in [66, 62], D-brane charges in orientifold theories should take their values in KR-theory. However, the KR-theory can be twisted because the gauge field on an O-plane can be of either orthogonal or symplectic type, leading to a need for sign-twisted KR-theory as discussed above in subsection 2.2. Examples of calculations with this theory in physically relevant situations were given in [44]. Then in [45], a complete analysis was made of all the possible twisted orientifold string theories on an elliptic curve (with holomorphic or antiholomorphic involution), and the twisted KR-theory was computed for each one. Isomorphisms between the various groups (further studied in [93]) agree with physics-based predictions of dualities between the various theories. In these calculations, one special case of twisted KR-theory that occurs is actually twisted KO-theory in the sense of Karoubi and Donovan. This case comes up in the treatment of Witten's theory of "type I string theory without vector structure," introduced in [106].

3.6. Applications in condensed matter theory. Twisted K-theory appears not only in string theory but also in condensed matter physics. The story begins with the "tenfold way" approach to classification of topological states of matter via K-theory [71, 12]. Most of the physical properties of solids come from the quantum mechanics of the electrons that are not tightly bound (those with energies above the Fermi level). Depending on the dimension and the symmetry class (whether or not one has time-reversal symmetry, parity symmetry, etc.), the wave functions of these electrons can be viewed as sections of a vector bundle with certain symmetry properties, and thus live in a certain K-group that classifies the states. It was shown in [56] that a more refined analysis (for example, taking into account the various crystallographic groups), leads sometimes to twisted K-theory, which can help explain phenomena such as topological insulators. By now there is quite a large mathematical physics literature involving twisted cohomology or twisted Ktheory in the study of topological states of matter. A few representative papers are [16, 36, 63, 88, 41, 60, 51, 61, 102, 99, 98, 40].

3.7. Applications of other twisted cohomology theories. Up to this point, most of the physics applications we have discussed have concerned twisted K-theory of one form or another. In this last subsection we mention some applications of twists of other cohomology theories. One example that seems to have physical applications is twisted Morava K-theory K(2) at the prime 2, or 2-local *tmf* (topological modular forms), which is closely related. At the prime 2, K(2) is periodic of period 6 and has a twisting given by a class in  $H^4(X, \mathbb{Z}_{(2)})$ , which the authors of [95, 96, 100] relate to the notion of "twisted String structures," which bear the same relation to M-theory that that twisted spin<sup>c</sup>-structures do in relation to string theory. (Just as anomaly cancellation for D-branes in the presence of an H-flux

requires D-branes to take their charges in twisted K-theory, a similar anomaly cancellation in M-theory suggests that branes should have charges in twisted K(2).)

Another example of a twisted cohomology theory that comes up in M-theory is twisted cohomotopy — see [47, 49, 48, 97]. Note that stable cohomotopy is in some sense the universal cohomology theory. (Unstable) cohomotopy is not exactly a cohomology theory; in fact cohomotopy sets are not always groups, but it reduces to stable cohomotopy in some circumstances. And calculations suggest that anomaly cancellation in M-theory should force charges to live in twisted cohomotopy groups.

# References

- John Frank Adams, Infinite loop spaces, Annals of Mathematics Studies, No. 90, Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1978. MR 505692
- Matthew Ando, Andrew J. Blumberg, and David Gepner, Twists of K-theory and TMF, Superstrings, geometry, topology, and C\*-algebras, Proc. Sympos. Pure Math., vol. 81, Amer. Math. Soc., Providence, RI, 2010, pp. 27–63. MR 2681757
- \_\_\_\_\_, Parametrized spectra, multiplicative Thom spectra and the twisted Umkehr map, Geom. Topol. 22 (2018), no. 7, 3761–3825. MR 3890766
- Matthew Ando, Andrew J. Blumberg, David Gepner, Michael J. Hopkins, and Charles Rezk, An ∞-categorical approach to R-line bundles, R-module Thom spectra, and twisted R-homology, J. Topol. 7 (2014), no. 3, 869–893. MR 3252967
- 5. \_\_\_\_, Units of ring spectra, orientations and Thom spectra via rigid infinite loop space theory, J. Topol. 7 (2014), no. 4, 1077–1117. MR 3286898
- 6. Benjamin Antieau, David Gepner, and José Manuel Gómez, Actions of  $K(\pi, n)$  spaces on K-theory and uniqueness of twisted K-theory, Trans. Amer. Math. Soc. **366** (2014), no. 7, 3631–3648. MR 3192610
- M. F. Atiyah, K-theory and reality, Quart. J. Math. Oxford Ser. (2) 17 (1966), 367–386. MR 206940
- 8. \_\_\_\_\_, Bott periodicity and the index of elliptic operators, Quart. J. Math. Oxford Ser. (2) 19 (1968), 113–140. MR 228000
- Michael Atiyah, K-theory past and present, Sitzungsberichte der Berliner Mathematischen Gesellschaft, Berliner Math. Ges., Berlin, 2001, arXiv:math/0012213, pp. 411–417. MR 2091892
- Michael Atiyah and Graeme Segal, *Twisted K-theory*, Ukr. Mat. Visn. 1 (2004), no. 3, 287– 330, arXiv:math/0407054. MR 2172633
- Twisted K-theory and cohomology, Inspired by S. S. Chern, Nankai Tracts Math., vol. 11, World Sci. Publ., Hackensack, NJ, 2006, pp. 5–43. MR 2307274
- John C. Baez, The tenfold way, Notices Amer. Math. Soc. 67 (2020), no. 10, 1599–1601. MR 4575939
- David Baraglia, Topological T-duality with monodromy, String-Math 2011, Proc. Sympos. Pure Math., vol. 85, Amer. Math. Soc., Providence, RI, 2012, pp. 293–302. MR 2985336
- <u>—</u>, Topological T-duality for general circle bundles, Pure Appl. Math. Q. 10 (2014), no. 3, 367–438. MR 3282986
- <u>Topological T-duality for torus bundles with monodromy</u>, Rev. Math. Phys. 27 (2015), no. 3, 1550008, 55. MR 3342758
- Maissam Barkeshli, Parsa Bonderson, Meng Cheng, and Zhenghan Wang, Symmetry Fractionalization, Defects, and Gauging of Topological Phases, Phys. Rev. B 100 (2019), no. 11, 115147.
- O. Ben-Bassat, J. Block, and T. Pantev, Non-commutative tori and Fourier-Mukai duality, Compos. Math. 143 (2007), no. 2, 423–475. MR 2309993
- Peter Bouwknegt, Alan L. Carey, Varghese Mathai, Michael K. Murray, and Danny Stevenson, Twisted K-theory and K-theory of bundle gerbes, Comm. Math. Phys. 228 (2002), no. 1, 17–45. MR 1911247
- Peter Bouwknegt, Peter Dawson, and David Ridout, D-branes on group manifolds and fusion rings, J. High Energy Phys. (2002), no. 12, 065, 22. MR 1960468
- Peter Bouwknegt, Jarah Evslin, and Varghese Mathai, *T-duality: topology change from H-flux*, Comm. Math. Phys. **249** (2004), no. 2, 383–415. MR 2080959

- <u>\_\_\_\_\_</u>, Topology and H-flux of T-dual manifolds, Phys. Rev. Lett. **92** (2004), no. 18, 181601,
   3. MR 2116165
- 22. \_\_\_\_\_, Spherical T-duality, Comm. Math. Phys. 337 (2015), no. 2, 909–954. MR 3339166
- Spherical T-duality II: an infinity of spherical T-duals for non-principal SU(2)bundles, J. Geom. Phys. 92 (2015), 46–54. MR 3328056
- <u>Spherical T-duality and the spherical Fourier-Mukai transform</u>, J. Geom. Phys. 133 (2018), 303–314. MR 3850273
- Peter Bouwknegt, Keith Hannabuss, and Varghese Mathai, *T-duality for principal torus bundles*, J. High Energy Phys. (2004), no. 3, 018, 10. MR 2062361
- Peter Bouwknegt and Varghese Mathai, D-branes, B-fields and twisted K-theory, J. High Energy Phys. (2000), no. 3, Paper 7, 11. MR 1756434
- Volker Braun, Twisted K-theory of Lie groups, J. High Energy Phys. (2004), no. 3, 029, 15. MR 2061550
- L. G. Brown, The universal coefficient theorem for Ext and quasidiagonality, Operator algebras and group representations, Vol. I (Neptun, 1980), Monogr. Stud. Math., vol. 17, Pitman, Boston, MA, 1984, pp. 60–64. MR 731763
- Ulrich Bunke and Thomas Nikolaus, *T-duality via gerby geometry and reductions*, Rev. Math. Phys. 27 (2015), no. 5, 1550013, 46. MR 3361543
- Ulrich Bunke, Philipp Rumpf, and Thomas Schick, The topology of T-duality for T<sup>n</sup>-bundles, Rev. Math. Phys. 18 (2006), no. 10, 1103–1154. MR 2287642
- Ulrich Bunke and Thomas Schick, On the topology of T-duality, Rev. Math. Phys. 17 (2005), no. 1, 77–112. MR 2130624
- , T-duality for non-free circle actions, Analysis, geometry and topology of elliptic operators, World Sci. Publ., Hackensack, NJ, 2006, pp. 429–466. MR 2246781
- Ulrich Bunke, Thomas Schick, and Markus Spitzweck, Periodic twisted cohomology and Tduality, Astérisque (2011), no. 337, vi+134. MR 2797285
- Ulrich Bunke, Thomas Schick, Markus Spitzweck, and Andreas Thom, *Duality for topological abelian group stacks and T-duality*, K-theory and noncommutative geometry, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008, pp. 227–347. MR 2482327
- Yeuk-Kwan E. Cheung and Zheng Yin, Anomalies, branes, and currents, Nuclear Phys. B 517 (1998), no. 1-3, 69–91. MR 1621204
- Ching-Kai Chiu, Jeffrey C. Y. Teo, Andreas P. Schnyder, and Shinsei Ryu, Classification of topological quantum matter with symmetries, Rev. Mod. Phys. 88 (2016), no. 3, 035005.
- Andrés Collinucci and Jarah Evslin, Twisted homology, J. High Energy Phys. (2007), no. 3, 058, 24. MR 2313935
- Calder Daenzer, A groupoid approach to noncommutative T-duality, Comm. Math. Phys. 288 (2009), no. 1, 55–96. MR 2491618
- Calder Daenzer and Erik van Erp, T-duality for Langlands dual groups, Adv. Theor. Math. Phys. 18 (2014), no. 6, 1267–1285. MR 3285609
- Giuseppe De Nittis and Kiyonori Gomi, Differential geometric invariants for time-reversal symmetric Bloch bundles, II: The low-dimensional "quaternionic" case, Algebr. Geom. Topol. 23 (2023), no. 7, 2925–2974. MR 4647667
- Giuseppe De Nittis and Kyonori Gomi, The cohomology invariant for class DIII topological insulators, Ann. Henri Poincaré 23 (2022), no. 10, 3587–3632. MR 4483417
- Peter Donovan and Max Karoubi, Groupe de Brauer et coefficients locaux en K-théorie, C. R. Acad. Sci. Paris Sér. A-B 269 (1969), A387–A389. MR 256386
- <u>—</u>, Graded Brauer groups and K-theory with local coefficients, Inst. Hautes Études Sci. Publ. Math. (1970), no. 38, 5–25. MR 282363
- 44. Charles Doran, Stefan Méndez-Diez, and Jonathan Rosenberg, *T-duality for orientifolds and twisted KR-theory*, Lett. Math. Phys. **104** (2014), no. 11, 1333–1364. MR 3267662
- <u>String theory on elliptic curve orientifolds and KR-theory</u>, Comm. Math. Phys. 335 (2015), no. 2, 955–1001. MR 3316647
- Christopher L. Douglas, On the twisted K-homology of simple Lie groups, Topology 45 (2006), no. 6, 955–988. MR 2263220
- Domenico Fiorenza, Hisham Sati, and Urs Schreiber, Twisted cohomotopy implies M-theory anomaly cancellation on 8-manifolds, Comm. Math. Phys. 377 (2020), no. 3, 1961–2025. MR 4121615

12

- Twisted cohomotopy implies level quantization of the full 6d Wess-Zumino term of the M5-brane, Comm. Math. Phys. 384 (2021), no. 1, 403–432. MR 4252880
- Twisted cohomotopy implies twisted string structure on M5-branes, J. Math. Phys. 62 (2021), no. 4, Paper No. 042301, 16. MR 4242295
- Stefan Fredenhagen and Volker Schomerus, Branes on group manifolds, gluon condensates, and twisted K-theory, J. High Energy Phys. (2001), no. 4, Paper 7, 30. MR 1834409
- Daniel S. Freed and Michael J. Hopkins, *Reflection positivity and invertible topological phases*, Geom. Topol. 25 (2021), no. 3, 1165–1330. MR 4268163
- Daniel S. Freed, Michael J. Hopkins, and Constantin Teleman, Twisted equivariant K-theory with complex coefficients, J. Topol. 1 (2008), no. 1, 16–44. MR 2365650
- 53. \_\_\_\_\_, Loop groups and twisted K-theory I, J. Topol. 4 (2011), no. 4, 737–798. MR 2860342
- 54. \_\_\_\_\_, Loop groups and twisted K-theory III, Ann. of Math. (2) 174 (2011), no. 2, 947– 1007. MR 2831111
- 55. \_\_\_\_\_, Loop groups and twisted K-theory II, J. Amer. Math. Soc. 26 (2013), no. 3, 595–644. MR 3037783
- Daniel S. Freed and Gregory W. Moore, *Twisted equivariant matter*, Ann. Henri Poincaré 14 (2013), no. 8, 1927–2023. MR 3119923
- Daniel S. Freed and Edward Witten, Anomalies in string theory with D-branes, Asian J. Math. 3 (1999), no. 4, 819–851. MR 1797580
- Matthias R. Gaberdiel and Terry Gannon, The charges of a twisted brane, J. High Energy Phys. (2004), no. 1, 018, 24. MR 2045888
- 59. \_\_\_\_\_, D-brane charges on non-simply connected groups, J. High Energy Phys. (2004), no. 4, 030, 27. MR 2080884
- Kiyonori Gomi, Yosuke Kubota, and Guo Chuan Thiang, Twisted crystallographic T-duality via the Baum-Connes isomorphism, Internat. J. Math. 32 (2021), no. 10, Paper No. 2150078, 65. MR 4311720
- Kiyonori Gomi and Guo Chuan Thiang, 'Real' gerbes and Dirac cones of topological insulators, Comm. Math. Phys. 388 (2021), no. 3, 1507–1555. MR 4340936
- Sergei Gukov, K-theory, reality, and orientifolds, Comm. Math. Phys. 210 (2000), no. 3, 621–639. MR 1777343
- Keith C. Hannabuss, Varghese Mathai, and Guo Chuan Thiang, *T-duality simplifies bulk-boundary correspondence: the parametrised case*, Adv. Theor. Math. Phys. **20** (2016), no. 5, 1193–1226. MR 3604557
- Fabian Hebestreit and Steffen Sagave, Homotopical and operator algebraic twisted K-theory, Math. Ann. 378 (2020), no. 3-4, 1021–1059. MR 4163521
- Fabian Hebestreit, Steffen Sagave, and Christian Schlichtkrull, Multiplicative parametrized homotopy theory via symmetric spectra in retractive spaces, Forum Math. Sigma 8 (2020), Paper No. e16, 84. MR 4079640
- Kentaro Hori, D-branes, T-duality, and index theory, Adv. Theor. Math. Phys. 3 (1999), no. 2, 281–342. MR 1736794
- Anton Kapustin, D-branes in a topologically nontrivial B-field, Adv. Theor. Math. Phys. 4 (2000), no. 1, 127–154. MR 1807598
- Max Karoubi, Twisted K-theory—old and new, K-theory and noncommutative geometry, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008, pp. 117–149. MR 2513335
- Max Karoubi and Charles Weibel, Twisted K-theory, real A-bundles and Grothendieck-Witt groups, J. Pure Appl. Algebra 221 (2017), no. 7, 1629–1640. MR 3614970
- Mehdi Khorami, A universal coefficient theorem for twisted K-theory, J. Topol. 4 (2011), no. 3, 535–542. MR 2832567
- Alexei Kitaev, Periodic table for topological insulators and superconductors, Advances in Theoretical Physics: Landau Memorial Conference, Chernogolovka, 2008, AIP Conference Proceedings, vol. 1134, American Institute of Physics, 2009, arXiv:0901.2686, pp. 22–34.
- I. Madsen, V. Snaith, and J. Tornehave, *Infinite loop maps in geometric topology*, Math. Proc. Cambridge Philos. Soc. 81 (1977), no. 3, 399–430. MR 494076
- Juan Maldacena, Gregory Moore, and Nathan Seiberg, *D-brane instantons and K-theory charges*, J. High Energy Phys. (2001), no. 11, Paper 62, 42. MR 1877986
- Varghese Mathai and Jonathan Rosenberg, T-duality for torus bundles with H-fluxes via noncommutative topology, Comm. Math. Phys. 253 (2005), no. 3, 705–721. MR 2116734

- T-duality for torus bundles with H-fluxes via noncommutative topology. II. The high-dimensional case and the T-duality group, Adv. Theor. Math. Phys. 10 (2006), no. 1, 123–158. MR 2222224
- T-duality for circle bundles via noncommutative geometry, Adv. Theor. Math. Phys. 18 (2014), no. 6, 1437–1462. MR 3285613
- 77. \_\_\_\_\_, Group dualities, T-dualities, and twisted K-theory, J. Lond. Math. Soc. (2) 97 (2018), no. 1, 1–23. MR 3764065
- Varghese Mathai and Hisham Sati, Some relations between twisted K-theory and E<sub>8</sub> gauge theory, J. High Energy Phys. (2004), no. 3, 016, 22. MR 2062363
- Varghese Mathai and Danny Stevenson, Chern character in twisted K-theory: equivariant and holomorphic cases, Comm. Math. Phys. 236 (2003), no. 1, 161–186. MR 1977885
- Varghese Mathai and Siye Wu, Topology and flux of T-dual manifolds with circle actions, Comm. Math. Phys. 316 (2012), no. 1, 279–286. MR 2989461
- J. P. May and J. Sigurdsson, *Parametrized homotopy theory*, Mathematical Surveys and Monographs, vol. 132, American Mathematical Society, Providence, RI, 2006. MR 2271789
- Ruben Minasian and Gregory Moore, K-theory and Ramond-Ramond charge, J. High Energy Phys. (1997), no. 11, Paper 2, 7. MR 1606278
- Gregory Moore, K-theory from a physical perspective, Topology, geometry and quantum field theory, London Math. Soc. Lecture Note Ser., vol. 308, Cambridge Univ. Press, Cambridge, 2004, pp. 194–234. MR 2079376
- El-kaïoum M. Moutuou, Twisted groupoid KR-theory, doctoral thesis, Université de Lorraine and Universität Paderborn, 2012, available at https://hal.univ-lorraine.fr/.
- Equivariant KK-theory for generalised actions and Thom isomorphism in groupoid twisted K-theory, J. K-Theory 13 (2014), no. 1, 83–113. MR 3177819
- Ashwin S. Pande, Topological T-duality and Kaluza-Klein monopoles, Adv. Theor. Math. Phys. 12 (2008), no. 1, 185–215. MR 2369414
- <u>Topological T-duality for stacks using a Gysin sequence</u>, Adv. Theor. Math. Phys. **22** (2018), no. 6, 1535–1591. MR 3952356
- Emil Prodan and Yitzchak Shmalo, The K-theoretic bulk-boundary principle for dynamically patterned resonators, J. Geom. Phys. 135 (2019), 135–171. MR 3872629
- Iain Raeburn and Dana P. Williams, Morita equivalence and continuous-trace C<sup>\*</sup>-algebras, Mathematical Surveys and Monographs, vol. 60, American Mathematical Society, Providence, RI, 1998. MR 1634408
- Jonathan Rosenberg, Homological invariants of extensions of C<sup>\*</sup>-algebras, Operator algebras and applications, Part 1 (Kingston, Ont., 1980), Proc. Sympos. Pure Math., vol. 38, Amer. Math. Soc., Providence, RI, 1982, pp. 35–75. MR 679694
- <u>\_\_\_\_\_</u>, Continuous-trace algebras from the bundle theoretic point of view, J. Austral. Math. Soc. Ser. A 47 (1989), no. 3, 368–381. MR 1018964
- 92. \_\_\_\_\_, Topology, C\*-algebras, and string duality, CBMS Regional Conference Series in Mathematics, vol. 111, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2009. MR 2560910
- <u>main and the set of the set of</u>
- <u>A new approach to twisted K-theory of compact Lie groups</u>, Algebr. Geom. Topol. 20 (2020), no. 1, 135–167. MR 4071368
- Hisham Sati, Geometric and topological structures related to M-branes, Superstrings, geometry, topology, and C\*-algebras, Proc. Sympos. Pure Math., vol. 81, Amer. Math. Soc., Providence, RI, 2010, pp. 181–236. MR 2681765
- 96. \_\_\_\_\_, Geometric and topological structures related to M-branes II: Twisted string and string<sup>c</sup> structures, J. Aust. Math. Soc. **90** (2011), no. 1, 93–108. MR 2810946
- Hisham Sati and Urs Schreiber, Twisted cohomotopy implies M5-brane anomaly cancellation, Lett. Math. Phys. 111 (2021), no. 5, Paper No. 120, 25. MR 4318438
- 98. \_\_\_\_\_, Anyonic defect branes and conformal blocks in twisted equivariant differential (TED) K-theory, Rev. Math. Phys. **35** (2023), no. 6, Paper No. 2350009, 65. MR 4612787
- 99. \_\_\_\_\_, Anyonic topological order in twisted equivariant differential (TED) K-theory, Rev. Math. Phys. 35 (2023), no. 3, Paper No. 2350001, 72. MR 4568912
- 100. Hisham Sati and Craig Westerland, Twisted Morava K-theory and E-theory, J. Topol. 8 (2015), no. 4, 887–916. MR 3431663

14

- 101. N. E. Steenrod, Homology with local coefficients, Ann. of Math. (2) 44 (1943), 610–627. MR 9114
- 102. Luuk Stehouwer, Jan de Boer, Jorrit Kruthoff, and Hessel Posthuma, Classification of crystalline topological insulators through K-theory, Adv. Theor. Math. Phys. 25 (2021), no. 3, 723–775. MR 4403305
- Erik Verlinde, Fusion rules and modular transformations in 2D conformal field theory, Nuclear Phys. B 300 (1988), no. 3, 360–376. MR 954762
- 104. Toni Aliza Watson, *Twisted cohomology groups*, M.A. thesis, University of Maryland, 2006, available at ProQuest Dissertations & Theses Global.
- 105. Edward Witten, D-branes and K-theory, J. High Energy Phys. (1998), no. 12, Paper 19, 41. MR 1674715
- 106. \_\_\_\_\_, Toroidal compactification without vector structure, J. High Energy Phys. (1998), no. 2, Paper 6, 43. MR 1615617

Department of Mathematics, University of Maryland, College Park, MD 20742-4015, USA

Email address: jmr@umd.edu

URL: http://www.math.umd.edu/~jmr/