

Many-body non-Hermitian Skin Effect for Multipoles

Jacopo Gliozzi,¹ Giuseppe De Tomasi,¹ and Taylor L. Hughes¹

¹*Department of Physics and Institute for Condensed Matter Theory, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801-3080, USA*

In this work, we investigate the fate of the non-Hermitian skin effect in one-dimensional systems that conserve the dipole moment and higher moments of an associated global U(1) charge. Motivated by field theoretical arguments and lattice model calculations, we demonstrate that the key feature of the non-Hermitian skin effect for m -pole conserving systems is the generation of an $(m+1)$ th multipole moment. For example, in contrast to the conventional skin effect where charges are anomalously localized at one boundary, the dipole-conserving skin effect results in charges localized at both boundaries, in a configuration that generates an extremal *quadrupole* moment. In addition, we explore the dynamical consequences of the m -pole skin effect, focusing on charge and entanglement propagation. Both numerically and analytically, we provide evidence that long-time steady-states have Fock-space localization and an area-law scaling of entanglement entropy, which serve as quantum indicators of the skin effect.

Introduction.— A central postulate of quantum mechanics is the unitarity of time evolution, which implies the conservation of probability during dynamics. However, interactions between a quantum system and its environment can induce dissipation, decoherence, or wavefunction collapse, destroying unitarity. Recent advances in controlled experimental techniques that allow for the manipulation of single qubits, along with the need to overcome decoherence in quantum devices, have led to increasing interest in such non-unitary dynamics as a paradigm to describe open and monitored quantum systems [1–5]. Relaxing unitarity allows for a rich phenomenology that ranges from novel topological phases [6–13] to phase transitions in the entanglement structure of non-equilibrium steady-states [14–20].

One approach to non-unitary dynamics is through non-Hermitian Hamiltonians, which model quantum systems subject to dissipation [21, 22] or continuous measurements and postselection [23–27]. Remarkably, non-Hermitian couplings can lead to unique phenomena with no Hermitian counterpart, such as the non-Hermitian skin effect, an anomalous localization of extensively many eigenstates at the boundary of a system [28–31], which has been observed experimentally [32–35]. While the boundary localization of particles is an essentially classical effect due to non-reciprocal hopping, it also has implications for the spread of quantum correlations during time evolution. For example, the skin effect has been shown to hinder entanglement growth and can trigger a volume-to-area law transition in the entanglement entropy of long-time steady-states [36]. In addition, the skin effect challenges the traditional understanding of a bulk-boundary correspondence in topological systems [37, 38].

Previous studies of the non-Hermitian skin effect have been mostly restricted to non-interacting Hamiltonians with U(1) particle-number symmetry. Moreover, the few extensions to interacting systems have relied on the existence of a non-interacting limit [39–42]. In this work, we




FIG. 1. (a-b) Schematic of many-body non-Hermitian charge and dipole skin effect, which generate a dipole and quadrupole moment, respectively. (c-d) Time evolution of the local charge density ρ_{charge} from a random uniform state for (c) charge-conserving and (d) dipole-conserving Hamiltonians with a non-Hermitian imbalanced kinetic term.

investigate the fate of the non-Hermitian skin effect in intrinsically interacting systems that conserve multipole moments of a U(1) charge. Such conservation laws constrain particle motion and can dramatically affect the out-of-equilibrium dynamics [43–53]. For instance, in systems where the total dipole moment is conserved, isolated charges cannot move, leading to anomalously slow charge transport and even ergodicity breaking.

The interplay between multipole conservation laws, which restrict particle motion, and non-reciprocal hoppings, which pump particles to a boundary through the non-Hermitian skin effect, creates a complex scenario for charge transport. For example, charge flows to a single boundary with only U(1) symmetry, but in dipole-conserving systems charges must flow symmetrically, e.g., outwards to both boundaries (see Fig. 1). Motivated by

the hierarchical structure of multipoles, we are able to make significant progress by defining the m -pole skin effect in terms of an $(m + 1)$ -pole moment in the charge distribution of eigenstates. We also investigate the dynamical consequences of this generalized skin effect by examining charge and entanglement propagation following a global quench. In particular, we provide analytical and numerical evidence that the m -pole skin effect leads to an area-law scaling of entanglement entropy in the long-time steady-state. Importantly, our definition also encompasses the standard “charge” skin effect, and helps identify experimentally accessible probes in systems with strong interactions.

Charge skin effect.— Before discussing higher multipole moments, let us first revisit the non-Hermitian skin effect in the context of U(1) charge conservation. We focus on one-dimensional fermionic chains of length L . The paradigmatic non-Hermitian model having the skin effect is the Hatano-Nelson Hamiltonian [54, 55]:

$$H_{\text{HN}}(g) = -\sum_j (e^g c_j^\dagger c_{j+1} + e^{-g} c_{j+1}^\dagger c_j), \quad (1)$$

where c_j^\dagger is the fermionic creation operator at site j , and g parameterizes the left-right imbalance in charge hopping. With periodic boundary conditions (PBCs), the single-particle spectrum of (1) winds around $E = 0$ in the complex plane, and the eigenstates are extended plane waves. In contrast, for open boundary conditions (OBCs) the spectrum is real-valued, and all single-particle eigenstates are exponentially localized at one boundary. This stark difference between PBC and OBC spectra is the hallmark of the non-Hermitian skin effect for non-interacting systems [28, 29].

To extend this definition to interacting systems, we first need a many-body diagnostic of the skin effect. Unlike the single-particle case, an extensive number of particles cannot be exponentially localized at a boundary in a many-body state because of the Pauli exclusion principle, which forbids fermions from clustering at the same site [56, 57]. Nevertheless, features of the single-particle phenomenology persist. For instance, the N -particle spectrum of (1) winds in the complex plane when the PBCs are twisted by a phase. In OBCs, the spectrum still becomes real-valued, and the eigenstates exhibit a left-right imbalance of charge. This asymmetry in the charge density of open eigenstates has been interpreted as a many-body extension of the skin effect [58].

While such diagnostics have been used for the Hatano-Nelson model with Hubbard interactions [41, 42], we aim for a definition that applies even in the absence of a non-interacting limit. Independently of particle number or interactions, the skin effect arises from the imbalanced kinetic term, which preferentially moves electrons to one edge and holes to the other. The resulting charge distribution, which is depicted in Fig. 1(a), exhibits an electric

dipole moment, which is defined by

$$P^{(1)} = \sum_{j=1}^L j \rho_j, \quad (2)$$

where ρ_j is the deviation of the local density at site j , $n_j = c_j^\dagger c_j$, from the average density per site, $\bar{n} = N/L$. We then define the non-Hermitian skin effect as the presence of an extensive dipole moment in the N -particle eigenstates with OBCs, formalizing the previous notion of a left-right asymmetry in density. Importantly, such a dipole moment arises both in the distribution of eigenstates and dynamically from time evolution, as shown in Fig. 1(c). We will see below that this definition allows for a direct generalization to systems that conserve arbitrary U(1) multipole moments.

For the Hatano-Nelson model, the relation between imbalanced charge-hopping and dipole moment can be understood by applying an “imaginary gauge transformation” [54, 55]:

$$c_j^\dagger \mapsto R^{-1} c_j^\dagger R = e^{gj} c_j^\dagger, \quad R = e^{-gP^{(1)}}, \quad (3)$$

where $P^{(1)}$ is the dipole moment in Eq. (2). In a chain with OBCs, this is a similarity transformation that maps $H_{\text{HN}}(g)$ to $H_{\text{HN}}(0)$, the latter being a *Hermitian* Hamiltonian without imbalance. An immediate consequence of this map is that the OBC spectrum of $H_{\text{HN}}(g)$ must be real. Furthermore, single-particle eigenstates of the Hermitian $H_{\text{HN}}(0)$ are standing wave-like, so applying the map in reverse yields eigenstates of the original Hamiltonian that are exponentially localized at an edge. Indeed, a single-particle state $|\psi_{0\alpha}\rangle = \sum_j \alpha_j c_j^\dagger |0\rangle$ is mapped back to $|\psi_{g\alpha}\rangle = \sum_j e^{gj} \alpha_j c_j^\dagger |0\rangle$, which is localized within $\sim |g|^{-1}$ of the boundary as long as α_j is not fine-tuned to counteract the exponential dependence. Regardless of particle number, the operator R tends to project onto states that maximize or minimize the dipole operator (depending on the sign of g). Hence, while a generic many-body eigenstate $|\psi_0\rangle$ of $H_{\text{HN}}(0)$ does not exhibit charge imbalance, the corresponding eigenstate of $H_{\text{HN}}(g)$, $|\psi_g\rangle = R^{-1} |\psi_0\rangle$, necessarily has an extremal dipole moment [59].

Although the map in (3) is applicable only to Hamiltonians having a real-valued spectrum, we anticipate that imbalanced charge hopping also induces a dipole moment in more general models. We can argue this heuristically using the effective field theory approach for non-Hermitian systems developed in Ref. 60. There, the authors couple the kinetic term of a non-Hermitian Hamiltonian to a background U(1) gauge field through a Peierls factor $c_j^\dagger e^{iA_{j,j+1}} c_{j+1}$, and derive an effective response action for the vector potential A_x in the continuum limit,

$$S_E = W(E) \int dx A_x, \quad (4)$$




FIG. 2. (a) Many body energy spectrum of H_{dip} at half-filling and with $P^{(1)} = 0$, for system size $L = 12$ and $g_1 = 2g_2/3 \equiv g = -1, t = 0.6$. For OBCs (red dots), the spectrum is real, while for PBCs (empty dots) the spectrum is complex and winds as the boundary conditions are twisted (blue lines). (b) Quadrupole moment, $P^{(2)}$ in Eq. (6), of the right eigenstates of the open chain with (red) and without (grey) imbalance. All eigenstates with $g \neq 0$ exhibit close to maximal quadrupole moment (dashed line), a signature of the dipole skin effect. (c) Local charge density and (d) local dipole density of 30 mid-spectrum states for $L = 20$ and OBCs. The charge distribution exhibits a quadrupole moment as opposite dipoles are piled up at opposite edges. The distribution of some eigenstates of the analogous Hermitian model is shown in black dashed lines for comparison.

where E is a reference energy, and $W(E)$ is the winding of the PBC spectrum around E . This effective action encodes the response of quasiparticle modes of energy E , and though $W(E)$ depends on the underlying Hamiltonian, the form of the action is fixed by $U(1)$ symmetry. The main observable associated with Eq. (4) is a persistent unidirectional current: $j^x(E) \equiv \delta S_E[A_x]/\delta A_x = W(E)$ [61]. Expressing the current in terms of the total dipole moment, $j^x = \partial_t P^{(1)}$, it follows that the physical response of non-Hermitian $U(1)$ -conserving system is the generation of a dipole moment when $W(E)$ is non-vanishing.

Multipole skin effect.— Having defined the charge skin effect in terms of the dipole operator, we now turn to the higher multipole case. In particular, we will show that the dipole-conserving skin effect is associated with a *quadrupole moment* in the charge distribution. This generalizes to a hierarchical structure in which the m -pole skin effect is defined through the $(m + 1)$ -pole moment. To demonstrate, we begin by considering an analytically tractable model that conserves both $U(1)$ charge $N =$

$\sum_j n_j$ and dipole moment $P^{(1)}$ [62],

$$H_4(g) = - \sum_j \left(e^g c_j^\dagger c_{j+1} c_{j+2} c_{j+3}^\dagger + e^{-g} c_j c_{j+1}^\dagger c_{j+2}^\dagger c_{j+3} \right). \quad (5)$$

This Hamiltonian generalizes the Hatano-Nelson model to imbalanced dipole-hopping terms, where $d_j^\dagger \equiv c_j^\dagger c_{j+1}$ creates a local dipole at site j (see Fig. 1(b)). Like the Hatano-Nelson model, the many-body spectrum of $H_4(g)$ is complex for PBCs and real for OBCs.

Taking L to be divisible by four for simplicity, we consider the model at half-filling $N = L/2$ and fixed dipole moment $P^{(1)} = 0$, i.e., the largest symmetry sector. Because H_4 exhibits strong Hilbert space fragmentation, this symmetry sector itself splits into exponentially many disconnected Krylov sub-sectors [47, 63]. The largest of these sub-sectors, which is still exponentially small in L , is integrable and can be mapped to a free-fermion chain of length $L/2$ [63]. The mapping proceeds by grouping sites $2j - 1$ and $2j$ of the original chain into a new composite site, which hosts four degrees of freedom associated with the occupations of the original sites: $|00\rangle, |01\rangle, |10\rangle, |11\rangle$. Composite sites are exactly singly-occupied in the largest sub-sector, and the allowed states $|10\rangle$ and $|01\rangle$ can be interpreted as a local dipole and anti-dipole, respectively. As a result, the dipole creation and annihilation operators on odd sites, d_{2j-1}^\dagger and d_{2j-1} , act as genuine fermionic operators on this subspace.

Only terms in Eq. (5) having odd j act nontrivially in this sector, and the resulting effective Hamiltonian can be written as $H_4^{\text{sector}} = \sum_k (e^g d_k^\dagger d_{k+1} + e^{-g} d_k d_{k+1}^\dagger)$, where k labels the composite sites [63]. This is just the Hatano-Nelson model, though with the fermions interpreted as local dipoles instead of charges. As a result, at least in this sector, the non-Hermitian skin effect is manifest as a dipole moment of *dipoles*, and therefore a quadrupole moment of charges. The buildup of dipoles (anti-dipoles) at the right (left) end of an open chain is depicted in Fig. 1(b), and generates an extensive quadrupole moment:

$$P^{(2)} = \sum_{j=1}^L j^2 \rho_j. \quad (6)$$

In contrast to the ordinary skin effect, there is no asymmetry in the charge distribution for eigenstates with OBCs, and particles are localized at both boundaries.

A few comments about the generality of this model are in order. Although this quadrupolar response is particularly clear in the example above, the mapping between $H_4(g)$ and a free-fermion chain holds only in a few integrable subsectors. Furthermore, the strong Hilbert space fragmentation precludes ergodicity, and thus makes it difficult to observe the skin effect dynamically. Finally, because the dipole operator $P^{(1)}$ does not commute with translations [64], local dipoles themselves are not always

well-defined. For example, a sequence of alternating positive and negative charges can be viewed as a chain of either local dipoles *or* anti-dipoles.

To circumvent these restrictions, we consider a more general dipole-conserving Hamiltonian:

$$H_{\text{dip}}(g_1, g_2) = - \sum_j \left(h_j^{(2)}(g_1) + t h_j^{(3)}(g_2) \right), \quad (7)$$

$$h^{(k)}(g) = e^g c_j^\dagger c_{j+1} c_{j+k} c_{j+k+1}^\dagger + e^{-g} \times \text{h.c.},$$

which reduces to Eq. (5) for $t = 0$, but otherwise contains longer range dipole-hopping terms that prevent strong Hilbert space fragmentation [47]. Once again, we fix $N = L/2$ and $P^{(1)} = 0$. The resulting PBC spectrum winds around the complex plane when the boundary conditions are twisted, as shown via exact diagonalization in Fig. 2(a).

Many-body spectral windings generically diverge in the thermodynamic limit [41], so we focus on OBCs. The model in Eq. (7) is not integrable and cannot be mapped to a free-fermion Hamiltonian. Nevertheless, for the particular choice of parameters $g_2 = 3g_1/2$, there exists a similarity transformation analogous to Eq. (3) that maps $H_{\text{dip}}(g, 3g/2)$ to the Hermitian Hamiltonian $H_{\text{dip}}(0, 0)$:

$$c_j^\dagger \mapsto R^{-1} c_j^\dagger R = e^{gj^2/4} c_j^\dagger, \quad R = e^{-gP^{(2)}/4}. \quad (8)$$

The spectrum of $H_{\text{dip}}(g, 3g/2)$ is therefore real, and the eigenstates of the non-Hermitian model are related to the Hermitian eigenstates by R^{-1} , which manifestly projects onto states with large positive or negative quadrupole moment $P^{(2)}$. We claim that this is a fundamental signature of the ‘‘dipole skin effect’’, which, as we have seen, arises in generic dipole-conserving Hamiltonians with non-reciprocal hoppings. Motivated by the charge- and dipole-conserving cases, we can generalize the definition to systems that conserve up to m th multipole moments: the non-Hermitian m -pole skin effect is marked by the generation of an extensive $(m + 1)$ -pole moment in the open-boundary eigenstates [65].

We confirm the dipole skin effect through exact diagonalization of $H_{\text{dip}}(g_1, g_2)$: the real OBC spectrum is shown in Fig. 2(a) for $L = 12$, while the quadrupole moments of the eigenstates plotted in Fig. 2(b) are near-maximal. The charge density profiles of a few mid-spectrum eigenstates for $L = 20$ are shown in Fig. 2(c): here the quadrupole moment is manifest as a buildup of positive charge at the system boundaries and negative charge in the middle. Following Refs. [66, 67], we can also define a bounded local dipole density $\rho_j^{\text{dip}} = - \sum_{k \leq j} \rho_k$ [68], which we plot for the same eigenstates in Fig. 2(d). The imbalance in dipole hopping yields a dipole moment of dipoles, i.e., a quadrupole moment of the charge, which should be contrasted with the Hermitian case, where ρ_j and ρ_j^{dip} are uniformly distributed across the sample (dashed lines).




FIG. 3. Dynamical probes of the late-time steady-state after evolution with a charge-conserving (inset: dipole-conserving) non-Hermitian Hamiltonian with non-reciprocal hopping imbalance g . (a) The half-chain von Neumann entanglement entropy exhibits area-law scaling. (b) The inverse participation ratio converges with system size, indicating Fock space localization. Both quantities are averaged over random initial product states.

The similarity transformation in (8) helps prove the existence of the skin effect, but generic non-reciprocal Hamiltonians with complex OBC spectra exhibit the same phenomenology. Indeed, we have checked numerically that our results are qualitatively unchanged for arbitrary values of g_1 and g_2 . This stability can be justified by an effective field theory that we propose for dipolar response. The background gauge field for dipole symmetry, A_{xx} , is a (one-component) rank-2 tensor and gives the Peierls phase associated with a dipole-hopping process [69]. By analogy with (4) we expect a response action of $S_E = \int dx j^{xx} A_{xx}$ for generic dipole-conserving non-reciprocal systems, whose primary observable is a dipole current $j^{xx} = \partial_t P^{(2)}$ that generates a quadrupole moment in open boundaries [70].

Dynamical signatures.—Having defined and examined the skin effect in systems that conserve multipole moments, we now focus on its implications for dynamics. Dynamical probes are especially important because they can be implemented in experimental platforms to characterize the skin effect. Concretely, we will study the state of a system at time t undergoing dynamics modeled by a non-Hermitian Hamiltonian H :

$$|\psi(t)\rangle = \frac{e^{-itH} |\psi(0)\rangle}{\|e^{-itH} |\psi(0)\rangle\|}. \quad (9)$$

We have already discussed the dynamics of charge: the m -pole skin effect causes charged particles to flow and generate an $(m + 1)$ -pole moment. Now, we turn to the evolution of quantum correlations, focusing on the entanglement scaling. Entanglement is quantified by the von Neumann entropy $S(t) = -\text{Tr}[\rho_{L/2} \log \rho_{L/2}]$, where $\rho_{L/2}$ is the bipartite reduced density matrix of the evolved state $|\psi(t)\rangle$ for a spatial cut that divides the system in half. Generic closed quantum systems, such as $H_{\text{dip}}(0, 0)$, thermalize under their own time evolution. It follows

that generic states evolve to have volume-law entanglement, $S \sim O(L)$. This behavior should be distinguished from area-law entanglement, $S \sim O(1)$, which is typically found only for ground states.

In addition to entanglement, we also investigate the spread of $|\psi(t)\rangle$ in Fock space through the inverse participation ratio, $IPR_2(t) = \sum_{\underline{n}} |\langle \underline{n} | \psi(t) \rangle|^4$, where $\{|\underline{n}\rangle\}$ is the many-body computational basis. The evolved state spans the entire Hilbert space if $IPR_2 \sim 2^{-D_2 L}$ with fractal dimension $D_2 = 1$, and is localized if $D_2 = 0$. Let us consider a generic m -pole conserving Hamiltonian having asymmetric hopping such that a similarity transformation $R^{-1} H^m(g) R = H^m(0)$ exists, with $R = e^{-\beta(g)P^{(m+1)}} = e^{-\beta(g)\sum_j j^{m+1}\rho_j}$. Starting from a random product state in the computational basis, $|\psi(0)\rangle = \prod_{\{s_j\}} c_{s_j}^\dagger |0\rangle$, we can use the transformation to write

$$IPR_2(t) = \frac{\sum_{\underline{n}} e^{-4\beta\sum_j j^{m+1}n_j} |\langle \underline{n} | e^{-itH^{(m)}(0)} |\psi(0)\rangle|^4}{\left(\sum_{\underline{n}} e^{-2\beta\sum_j j^{m+1}n_j} |\langle \underline{n} | e^{-itH^{(m)}(0)} |\psi(0)\rangle|^2\right)^2}. \quad (10)$$

Assuming that the corresponding Hermitian evolution reaches a generic state, e.g., via thermalization, such that $|\langle \underline{n} | e^{-itH^{(m)}(0)} |\psi(0)\rangle|^2 \sim 2^{-L}$ for large t , we obtain

$$|IPR_2(t \rightarrow \infty)| \approx \prod_{j=1}^L \frac{(1 + e^{-4\beta j^{m+1}})}{(1 + e^{-2\beta j^{m+1}})^2}. \quad (11)$$

This expression is bounded by $|\log IPR_2| \leq \sum_{j=1}^L \left| \log \left(1 - \frac{2e^{-2\beta j^{m+1}}}{(1 + e^{-2\beta j^{m+1}})^2} \right) \right|$, which converges as $L \rightarrow \infty$, implying localization in Fock space ($D_2 \rightarrow 0$). This should be viewed as a generalization of the edge localization that results from the single-particle skin effect. Although fermion repulsion prevents many-body states from localizing at the physical boundary, the m -pole skin effect extremizes the $(m+1)$ -pole moment of any initial state, driving time evolution to a highly restricted region of the Fock space. Finally, localization in Fock space implies that the entanglement entropy of the steady-state has area-law scaling [71].

We test our conjecture by performing exact numerical simulations of charge- and dipole-conserving time evolution. Fig. 3 (a-b) shows $S(t \rightarrow \infty)$ and $IPR_2(t \rightarrow \infty)$ for the $U(1)$ charge skin effect using $H_{\text{HN}}(g)$ in Eq. 1 (for various system sizes as a function of the non-reciprocity g). In the insets of Fig. 3 (a-b), we consider the case of the dipole skin effect with $H_4(g)$ in Eq. 5. In agreement with our analytical analysis, we find that the final states exhibit area-law scaling for the entanglement entropy and are localized in Fock space.

Conclusions— Nonreciprocal hopping leads to the localization of charge at one boundary of a 1D chain. On the other hand, m -pole conservation imposes strong constraints on charge transport. The competition between these two effects gives rise to a new class of non-

Hermitian skin effects in intrinsically interacting systems. In this work, we studied these generalized skin effects using lattice models, field theory arguments, and numerics. Our central result, that eigenstates exhibiting the m -pole skin effect have an $(m+1)$ -pole moment, naturally implies a definition of the ordinary skin effect beyond the non-interacting regime.

Furthermore, this definition provides an experimentally accessible signature of the m -pole skin effect, which may then be observed in ultra-cold atom platforms. For example, non-reciprocal charge hopping can be engineered by coupling an optical lattice to a collective one-body loss term induced by dissipation [72]. Placing this lattice in a strongly tilted external field then effectively imposes dipole-conservation [73–75], and the ensuing dynamics is governed by the dipole skin effect. Looking beyond local densities of conserved charges, we have also provided a truly quantum characterization of the skin effect, showing that it results in area-law scaling of the entanglement entropy in long-time steady-states.

We have seen an intimate connection between the symmetries of a system and its non-Hermitian dynamics. To this end, it would be interesting to consider models with other types of symmetries, like higher-form or subsystem symmetries, as well as patterns of symmetry-breaking. It may also be possible to extend the m -pole skin effect to higher dimensions and connect it to many-body topological invariants, both fruitful directions for future research.

G. D. T. acknowledges the support from the EPiQS Program of the Gordon and Betty Moore Foundation. T.L.H. and J. G. acknowledge support from the U.S. Office of Naval Research (ONR) Multidisciplinary University Research Initiative (MURI) under Grant No. N00014-20-1-2325 on Robust Photonic Materials with Higher-Order Topological Protection.

-
- [1] M. P. Fisher, V. Khemani, A. Nahum, and S. Vijay, *Annual Review of Condensed Matter Physics* **14**, 335 (2023), <https://doi.org/10.1146/annurev-conmatphys-031720-030658>.
 - [2] A. C. Potter and R. Vasseur, Entanglement dynamics in hybrid quantum circuits, in *Entanglement in Spin Chains: From Theory to Quantum Technology Applications*, edited by A. Bayat, S. Bose, and H. Johanneson (Springer International Publishing, Cham, 2022) pp. 211–249.
 - [3] E. Altman, K. R. Brown, G. Carleo, L. D. Carr, E. Demler, C. Chin, B. DeMarco, S. E. Economou, M. A. Eriks-son, K.-M. C. Fu, M. Greiner, K. R. Hazzard, R. G. Hulet, A. J. Kollár, B. L. Lev, M. D. Lukin, R. Ma, X. Mi, S. Misra, C. Monroe, K. Murch, Z. Nazario, K.-K. Ni, A. C. Potter, P. Roushan, M. Saffman, M. Schleier-Smith, I. Siddiqi, R. Simmonds, M. Singh, I. Spielman, K. Temme, D. S. Weiss, J. Vučković, V. Vuletić, J. Ye,

- and M. Zwierlein, *PRX Quantum* **2**, 017003 (2021).
- [4] E. Chertkov, Z. Cheng, A. C. Potter, S. Gopalakrishnan, T. M. Gatterman, J. A. Gerber, K. Gilmore, D. Gresh, A. Hall, A. Hankin, M. Matheny, T. Mengle, D. Hayes, B. Neyenhuis, R. Stutz, and M. Foss-Feig, *Nature Physics* **19**, 1799 (2023).
- [5] Z. G. Yuto, Ashida, and M. Ueda, *Advances in Physics* **69**, 249 (2020), <https://doi.org/10.1080/00018732.2021.1876991>.
- [6] D. Leykam, K. Y. Bliokh, C. Huang, Y. D. Chong, and F. Nori, *Phys. Rev. Lett.* **118**, 040401 (2017).
- [7] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, *Physical Review X* **9**, 041015 (2019).
- [8] K. Kawabata, M. Sato, and K. Shiozaki, *Physical Review B* **102**, 205118 (2020), arxiv:2008.07237 [cond-mat, physics:math-ph, physics:physics, physics:quant-ph].
- [9] D. S. Borgia, A. J. Kruchkov, and R.-J. Slager, *Phys. Rev. Lett.* **124**, 056802 (2020).
- [10] Y. Ma and T. L. Hughes, *Physical Review B* **108**, L100301 (2023).
- [11] N. Okuma and M. Sato, *Annual Review of Condensed Matter Physics* **14**, 83 (2023).
- [12] A. Lavasani, Y. Alavirad, and M. Barkeshli, *Nat. Phys.* **17**, 342 (2021).
- [13] A. Lavasani, Y. Alavirad, and M. Barkeshli, *Phys. Rev. Lett.* **127**, 235701 (2021).
- [14] Y. Li, X. Chen, and M. P. A. Fisher, *Phys. Rev. B* **98**, 205136 (2018).
- [15] B. Skinner, J. Ruhman, and A. Nahum, *Phys. Rev. X* **9**, 031009 (2019).
- [16] A. Chan, R. M. Nandkishore, M. Pretko, and G. Smith, *Phys. Rev. B* **99**, 224307 (2019).
- [17] S. Choi, Y. Bao, X.-L. Qi, and E. Altman, *Phys. Rev. Lett.* **125**, 030505 (2020).
- [18] Y. Bao, S. Choi, and E. Altman, *Phys. Rev. B* **101**, 104301 (2020).
- [19] M. J. Gullans and D. A. Huse, *Phys. Rev. X* **10**, 041020 (2020).
- [20] Z. Weinstein, Y. Bao, and E. Altman, *Phys. Rev. Lett.* **129**, 080501 (2022).
- [21] H. Feshbach, C. E. Porter, and V. F. Weisskopf, *Phys. Rev.* **96**, 448 (1954).
- [22] I. Rotter, *J. Phys. A: Math. Theor.* **42**, 153001 (2009).
- [23] M. Ueda, *Phys. Rev. A* **41**, 3875 (1990), 9903562.
- [24] J. Dalibard, Y. Castin, and K. Mølmer, *Phys. Rev. Lett.* **68**, 580 (1992).
- [25] R. Dum, P. Zoller, and H. Ritsch, *Phys. Rev. A* **45**, 4879 (1992).
- [26] M. B. Plenio and P. L. Knight, *Rev. Mod. Phys.* **70**, 101 (1998).
- [27] A. J. Daley, *Adv. Phys.* (2014).
- [28] T. E. Lee, *Phys. Rev. Lett.* **116**, 133903 (2016).
- [29] S. Yao and Z. Wang, *Phys. Rev. Lett.* **121**, 086803 (2018).
- [30] M.-H. L. Xiujuan Zhang, Tian Zhang and Y.-F. Chen, *Advances in Physics: X* **7**, 2109431 (2022), <https://doi.org/10.1080/23746149.2022.2109431>.
- [31] R. Lin, T. Tai, L. Li, and C. H. Lee, *Frontiers of Physics* **18**, 53605 (2023).
- [32] A. Ghatak, M. Brandenbourger, J. van Wezel, and C. Coulais, *Proceedings of the National Academy of Sciences* **117**, 29561 (2020), <https://www.pnas.org/doi/pdf/10.1073/pnas.2010580117>.
- [33] T. Helbig, T. Hofmann, S. Imhof, M. Abdelghany, T. Kiessling, L. W. Molenkamp, C. H. Lee, A. Szameit, M. Greiter, and R. Thomale, *Nature Physics* **16**, 747 (2020).
- [34] X. Zhang, Y. Tian, J.-H. Jiang, M.-H. Lu, and Y.-F. Chen, *Nature Communications* **12**, 5377 (2021).
- [35] L. Zhang, Y. Yang, Y. Ge, Y.-J. Guan, Q. Chen, Q. Yan, F. Chen, R. Xi, Y. Li, D. Jia, S.-Q. Yuan, H.-X. Sun, H. Chen, and B. Zhang, *Nature Communications* **12**, 6297 (2021).
- [36] K. Kawabata, T. Numasawa, and S. Ryu, *Physical Review X* **13**, 021007 (2023).
- [37] L. Xiao, T. Deng, K. Wang, G. Zhu, Z. Wang, W. Yi, and P. Xue, *Nature Physics* **16**, 761 (2020).
- [38] K. Zhang, Z. Yang, and C. Fang, *Nature Communications* **13**, 2496 (2022).
- [39] D.-W. Zhang, Y.-L. Chen, G.-Q. Zhang, L.-J. Lang, Z. Li, and S.-L. Zhu, *Phys. Rev. B* **101**, 235150 (2020).
- [40] T. Liu, J. J. He, T. Yoshida, Z.-L. Xiang, and F. Nori, *Phys. Rev. B* **102**, 235151 (2020).
- [41] K. Kawabata, K. Shiozaki, and S. Ryu, *Physical Review B* **105**, 165137 (2022).
- [42] S.-B. Zhang, M. M. Denner, T. Bzdusek, M. A. Sentef, and T. Neupert, *Physical Review B* **106**, L121102 (2022), arxiv:2201.12653 [cond-mat].
- [43] S. Pai, M. Pretko, and R. M. Nandkishore, *Phys. Rev. X* **9**, 021003 (2019).
- [44] V. Khemani, M. Hermele, and R. Nandkishore, *Phys. Rev. B* **101**, 174204 (2020).
- [45] J. Feldmeier, P. Sala, G. De Tomasi, F. Pollmann, and M. Knap, *Phys. Rev. Lett.* **125**, 245303 (2020).
- [46] T. Rakovszky, P. Sala, R. Verresen, M. Knap, and F. Pollmann, *Physical Review B* **101**, 125126 (2020).
- [47] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, *Physical Review X* **10**, 011047 (2020).
- [48] J. Feldmeier and M. Knap, *Phys. Rev. Lett.* **127**, 235301 (2021).
- [49] A. G. Burchards, J. Feldmeier, A. Schuckert, and M. Knap, *Phys. Rev. B* **105**, 205127 (2022).
- [50] J. Feldmeier, W. Witczak-Krempa, and M. Knap, *Phys. Rev. B* **106**, 094303 (2022).
- [51] J. Gliozzi, J. May-Mann, T. L. Hughes, and G. De Tomasi, *Phys. Rev. B* **108**, 195106 (2023).
- [52] P. Zechmann, J. Boesl, J. Feldmeier, and M. Knap, *arXiv e-prints*, arXiv:2310.16084 (2023), arXiv:2310.16084 [cond-mat.quant-gas].
- [53] O. Ogunnaike, J. Feldmeier, and J. Y. Lee, *Phys. Rev. Lett.* **131**, 220403 (2023), arXiv:2304.13028 [cond-mat.str-el].
- [54] N. Hatano and D. R. Nelson, *Physical Review Letters* **77**, 570 (1996).
- [55] N. Hatano and D. R. Nelson, *Phys. Rev. B* **56**, 8651 (1997).
- [56] E. Lee, H. Lee, and B.-J. Yang, *Phys. Rev. B* **101**, 121109 (2020).
- [57] T. Liu, J. J. He, T. Yoshida, Z.-L. Xiang, and F. Nori, *Phys. Rev. B* **102**, 235151 (2020).
- [58] F. Alsallom, L. Herviou, O. V. Yazyev, and M. Brzezińska, *Physical Review Research* **4**, 033122 (2022).
- [59] The many-body eigenstates of $H_{\text{HN}}(g)$ are Slater determinants of the single-particle states $|E(g)\rangle = R^{-1}|E(0)\rangle$.
- [60] K. Kawabata, K. Shiozaki, and S. Ryu, *Physical Review Letters* **126**, 216405 (2021).
- [61] With open boundaries, the effective action must include boundary degrees of freedom to preserve gauge invari-

- ance. Physically, these are external sources and sinks of charge that allow a steady-state current to exist.
- [62] The dipole moment is only conserved modulo the system size L in periodic boundary conditions.
- [63] S. Moudgalya, A. Prem, R. Nandkishore, N. Regnault, and B. A. Bernevig, [Thermalization and its absence within krylov subspaces of a constrained hamiltonian](#) (2021), [arXiv:1910.14048](#).
- [64] A. Gromov, A. Lucas, and R. M. Nandkishore, [Physical Review Research](#) **2**, 033124 (2020).
- [65] Multipole moments are origin-dependent when lower, conserved moments do not vanish, but for any choice of origin, the m -pole skin effect always tends to extremize the $(m + 1)$ -pole moment.
- [66] S. Moudgalya, A. Prem, D. A. Huse, and A. Chan, [Phys. Rev. Res.](#) **3**, 023176 (2021).
- [67] P. Zechmann, E. Altman, M. Knap, and J. Feldmeier, [Physical Review B](#) **107**, 195131 (2023).
- [68] This comes from expressing the dipole operator as $P^{(1)} = -\sum_{j=1}^L (L-j)\rho_j$ (since $\sum_j \rho_j = 0$ by definition) and then rewriting the sum as $-\sum_{j=1}^{L-1} \sum_{k=1}^j \rho_j$.
- [69] M. Pretko, [Phys. Rev. B](#) **95**, 115139 (2017).
- [70] Y. You, F. J. Burnell, and T. L. Hughes, [Phys. Rev. B](#) **103**, 245128 (2021).
- [71] G. De Tomasi and I. M. Khaymovich, [Phys. Rev. Lett.](#) **124**, 200602 (2020).
- [72] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, [Phys. Rev. X](#) **8**, 031079 (2018).
- [73] E. Guardado-Sanchez, A. Morningstar, B. M. Spar, P. T. Brown, D. A. Huse, and W. S. Bakr, [Phys. Rev. X](#) **10**, 011042 (2020).
- [74] S. Scherg, T. Kohlert, P. Sala, F. Pollmann, B. Hebbe Madhusudhana, I. Bloch, and M. Aidelsburger, [Nat. Commun.](#) **12**, 1 (2021).
- [75] T. Kohlert, S. Scherg, P. Sala, F. Pollmann, B. Hebbe Madhusudhana, I. Bloch, and M. Aidelsburger, [Phys. Rev. Lett.](#) **130**, 010201 (2023).