

THE ODD FERMION

DANIEL S. FREED, MICHAEL J. HOPKINS, AND CONSTANTIN TELEMAN

ABSTRACT. In this short note we use the geometric approach to (topological) field theory to address the question: Does an odd number of quantum mechanical fermions make sense?

Old questions: Does an odd number of quantum mechanical fermions make sense? And, if so, what is the dimension of the Hilbert space of states?

Answers: Yes, it makes sense as an anomalous theory. The standard definition of ‘dimension’ gives 0, and the “twisted dimension” for a single fermion is $\sqrt{2}$.

Key points:

- The theory is anomalous: it is a 1-dimensional field theory relative to the 2-dimensional *Arf theory*.
- The theory is topological, so computations are easily carried out.
- The state space is a module over the Clifford algebra $Cl_1^{\mathbb{C}}$; ‘dimension’ and ‘twisted dimension’ are relative concepts—relative to the Clifford algebra action.
- The tensor product construction of a theory of $N_1 + N_2$ fermions works as it should.

This question has garnered some interest over the years, and the theory of an odd number of fermions has sometimes been deemed inconsistent. See [DGG], [W, §2.1], and [SS, footnote 1] for recent comments; the latter contains several additional references to the literature. That the state space is to be regarded as a module over $Cl_1^{\mathbb{C}}$ already appears in [F1, Remark 5.10].

We thank Nati Seiberg for his perspective. His lecture at the November, 2023 annual meeting of the Simons Collaboration on Global Categorical Symmetries inspired us to write this note.

Quantum mechanics of N fermions

Let N be a positive integer, which we presently assume is odd. Consider the quantum mechanical theory of N free massless spinor fields. It is a 1-dimensional theory which, after Wick rotation, has background fields a Riemannian metric and spin structure. The fluctuating field is a spinor field with values in a real vector space of dimension N . For N even, one can work out this theory as a standard exercise; see [F2, FP16] for example.

Remark 1. We assume from the beginning that the spinor field is fermionic, or odd (anticommuting). Therefore, its quantization generates fermionic states. Furthermore, Bose-Fermi statistics are encoded in a $\mathbb{Z}/2\mathbb{Z}$ -grading on the state space, and algebras of observables are also $\mathbb{Z}/2\mathbb{Z}$ -graded;

Date: November 28, 2023.

This material is based upon work supported by the National Science Foundation under Grant Number DMS-2005286 and by the Simons Foundation Award 888988 as part of the Simons Collaboration on Global Categorical Symmetries. This work was performed in part at Aspen Center for Physics, which is supported by National Science Foundation grant PHY-2210452.

they are *superalgebras*. Alternatively, one can consider the system in which the spinor field is bosonic and there are no fermionic states; see the discussion of a lattice version in [SS, §1.1].

Let W be an N -dimensional real inner product space. The spin structure on a spin Riemannian 1-manifold X is given as a double cover $\tilde{X} \rightarrow X$. A spinor field ψ on X is a function $\tilde{\psi}: \tilde{X} \rightarrow \Pi W$ that is odd under the deck transformation of $\tilde{X} \rightarrow X$. Here ΠW is the parity-reversed vector space, i.e., W as an odd vector space.¹ The lagrangian density is

$$(2) \quad L = \frac{1}{2} \langle \psi, D_X \psi \rangle d\mu_X,$$

where μ_X is the Riemannian measure on X and D_X is the Dirac operator on X . The latter is $D_X \psi = \partial \tilde{\psi}$ for ∂ the positively oriented unit norm vector field on \tilde{X} .

Compute canonical quantization on $Y = \text{pt}$ as follows. First, solve the classical equations of motion on $\mathbb{R} \times Y$ to obtain a symplectic vector space, which in this case is odd. Here the classical equation asserts that ψ is locally constant, so the vector space of solutions is canonically isomorphic to ΠW . After complexification, classical observables form the algebra $\text{Sym}^\bullet \Pi W_{\mathbb{C}}^*$. This is the finite dimensional exterior algebra of $W_{\mathbb{C}}^*$ with the $\mathbb{Z}/2\mathbb{Z}$ -grading that makes $\wedge^1 W_{\mathbb{C}}^* = W_{\mathbb{C}}^*$ odd. Second, form the *deformation quantization* using the Poisson bracket derived from the lagrangian (2). The result is the Clifford algebra $\text{Cliff}(W_{\mathbb{C}}^*)$ for the complexified dual inner product; the Clifford algebra carries its usual $\mathbb{Z}/2\mathbb{Z}$ -grading. Third, quantum states are defined by producing an irreducible $\mathbb{Z}/2\mathbb{Z}$ -graded module M over $\text{Cliff}(W_{\mathbb{C}}^*)$. Here is where the parity of N shows up. If N is even, then the complex algebra $\text{Cliff}(W_{\mathbb{C}}^*)$ of observables is the full endomorphism algebra of M . In this case M is a $\mathbb{Z}/2\mathbb{Z}$ -graded vector space of dimension

$$(3) \quad \dim M = 2^{N/2-1} \mid 2^{N/2-1};$$

the notation ‘ $b \mid f$ ’ indicates the dimensions of the even (Bose) and odd (Fermi) homogeneous subspaces of a $\mathbb{Z}/2\mathbb{Z}$ -graded vector space. For N odd the action of $\text{Cliff}(W_{\mathbb{C}}^*)$ on M has a nontrivial *commutant*:² the subalgebra of $\text{End}(M)$ consisting of operators that commute with every element of $\text{Cliff}(W_{\mathbb{C}}^*)$ is isomorphic to $\text{Cl}_1^{\mathbb{C}}$, the complex Clifford algebra on a single odd generator δ .

Example 4 ($N = 1$). Let $W = \mathbb{R}$ with its standard inner product, so $\text{Cliff}(W_{\mathbb{C}}^*) = \text{Cl}_1^{\mathbb{C}}$ is the complex superalgebra generated by an odd element γ . An irreducible module is $M = \mathbb{C}^{1 \mid 1}$ with $\gamma = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. The commutant is generated by $\delta = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$.

Key point: One must consider the state space as a module over the commutant. Explanation: In this case the algebra of complex observables A is the commutant of the commutant B inside the full endomorphism algebra $\text{End}(M)$, so is identified as $\text{End}_B(M)$, the algebra of endomorphisms of M as a B -module. Typically, one must complete the module M to obtain a Hilbert space of states, but in this case M is finite dimensional and no completion is necessary.

¹More precisely, $\Pi W = \Pi \otimes W$, where Π is the canonical odd line.

²As always in super algebra, the commutant is taken in the $\mathbb{Z}/2\mathbb{Z}$ -graded sense: odd elements a_1, a_2 commute if $a_2 a_1 = -a_1 a_2$.

Remark 5. An extra “spectator fermion” has appeared in many previous works, for example [tH, LW, K]. It plays a similar role to the commutant, but the construction in the previous paragraph applies generally and leads to the commutant as part of the anomaly.

Next, we compute (heuristically, using a regularization) the path integral on $X = S^1$. For simplicity take $N = 1$, as in [Example 4](#); the general case is the N^{th} power of the computation here. There are two spin structures: *bounding* (antiperiodic, Neveu-Schwarz) and *non-bounding* (periodic, Ramond). The relevant one for us is the bounding spin structure. Let X be a Riemannian circle of length L . The path integral of the Lagrangian (2) is the pfaffian $\text{pfaff } D_X$ of the Dirac operator. It is naturally an element of the real Pfaffian line $\text{Pfaff } D_X$. That line carries a natural inner product, and the norm square of $\text{pfaff } D_X$ is the determinant of D_X . The set of eigenvalues of D_X is

$$(6) \quad \left\{ \frac{2\pi i n}{L} : n \in \mathbb{Z} + \frac{1}{2} \right\}.$$

The computation of the product of eigenvalues is

$$(7) \quad \prod_{n \in \mathbb{Z} + 1/2} \left(\frac{2\pi i}{L} n \right) = \prod_{n \in \mathbb{Z} + 1/2} n = \prod_{n \text{ odd}} n = \frac{\prod_{n \neq 0} n}{\prod_{\substack{n \text{ even} \\ n \neq 0}} n} = \frac{\prod_{n \neq 0} n}{\prod_{n \neq 0} 2n} = 2.$$


We use ζ -functions to regulate the infinite products. In the first and second steps we use that the ζ -regularized cardinality of \mathbb{Z} , and so too of $\mathbb{Z} + 1/2$, is zero. In the last step we use the corollary that the ζ -regularized cardinality of $\mathbb{Z}^{\neq 0}$ is -1 . The pfaffian is the square root of (7), determined here up to sign.

Remark 8.

- (1) The path integral is independent of the length L : the theory is *topological*.
- (2) That the state space M is a module over $C\ell_1^{\mathbb{C}}$ signals that the theory is *anomalous*, as confirmed by the fact that the path integral is an element of a line rather than a number.
- (3) In the next section we use a canonical element of $\text{Pfaff } D_X$ to identify the pfaffian $\text{pfaff } D_X$ as a number: $\sqrt{2}$.
- (4) For the nonbounding spin structure the Dirac operator has a 1-dimensional kernel, and hence the $\mathbb{Z}/2\mathbb{Z}$ -graded Pfaffian line is odd in the sense of super vector spaces. The nonzero kernel also implies that the pfaffian element is zero. In fact, this is a consequence of the oddness of the Pfaffian line: in the category of super vector spaces every homomorphism $\mathbb{C} \rightarrow \Pi$ from an even line to an odd line is zero.

N fermions as a boundary topological field theory


In the Wick-rotated setting an anomaly is a once-categorified invertible field theory, and the anomalous theory is defined relative to it. Both theories are defined on the same bordism category, so in particular have the same dimension. It often happens that the anomaly extends to a full theory in one higher dimension. See [F3] for discussion and motivation.

FIGURE 1. The $\text{Cl}_1^{\mathbb{C}}$ -module M

Fix N to be odd. We work with topological field theories defined on spin manifolds and taking values in the Morita 2-category of complex superalgebras. The preceding implies that the anomaly theory of N fermions assigns the algebra³ $\text{Cl}_1^{\mathbb{C}}$ to a point and the Pfaffian line of Dirac to a circle. The resulting once-categorified 1-dimensional invertible theory α is the truncation of the 2-dimensional invertible ‘‘Arf theory’’; see [Gu, DeGu] for example. The partition function on a closed spin 2-manifold W is $\alpha(W) = (-1)^{\text{Arf } W}$, where $\text{Arf } W$ is the Arf invariant of the spin structure: 0 for even spin structures and 1 for odd spin structures. The quantum theory of N fermions is a boundary theory of α ; the combined theory ($\alpha + \text{boundary}$) assigns the left $\text{Cl}_1^{\mathbb{C}}$ -module M to the bordism depicted in Figure 1. Green depicts components of the boundary colored by the module theory, and the red arrow tells if an uncolored boundary component is incoming or outgoing.

In a spin theory the usual dimension of the value of the theory on a codimension one closed manifold Y is computed by taking Cartesian product of Y with the *nonbounding* S^1 : the spin double cover of the nonbounding S^1 is the product double cover, so it computes the trace of the identity map in whatever category we are in. Furthermore, this trace lives in the Hochschild *homology* of the value of Y . In our case, for $Y = \text{pt}$ the Arf theory on the Cartesian product evaluates to $\alpha(S_{\text{nonbounding}}^1) = \Pi$, where Π is the odd line. As mentioned in Remark 8(4), an odd line has no nonzero elements. Hence the usual dimension is 0.

Remark 9. This is the ‘usual dimension’, or categorical dimension, in the category of $\text{Cl}_1^{\mathbb{C}}$ -modules. By comparison, the ‘usual dimension’ in the category of super vector spaces is $\dim \mathbb{C}^{b|f} = b - f$. This is oft referred to as ‘trace $(-1)^F$ ’. The ‘twisted dimension’ we are after is the categorical trace of the grading operator, which for the super vector space $\mathbb{C}^{b|f}$ equals $b + f$. We now compute ‘twisted dimension’ in the category of $\text{Cl}_1^{\mathbb{C}}$ -modules.

FIGURE 2. The twisted dimension of M

Consider the Cartesian product of the bordism in Figure 1 with the *bounding* circle S_{bounding}^1 . The theory ($\alpha + \text{boundary}$) computes the supertrace of the grading operator on M —what we are calling the *twisted dimension* of M —which lives in the Hochschild *cohomology* $\alpha(S_{\text{bounding}}^1)$ of the algebra $A = \text{Cl}_1^{\mathbb{C}}$. The Hochschild cohomology is an algebra, so has a canonical element: the unit for multiplication. Since the Arf theory α is invertible, the Hochschild cohomology is a line, and so the unit is a basis element. We use this basis element to define the twisted dimension as a

³The Clifford algebra is invertible in the Morita 2-category of superalgebras. This is equivalent to its being *central simple*. See [Wa, De].

number, the value of the theory on **Figure 2**. Compute the square of this number by evaluating the squared theory $\alpha^{\otimes 2}$ on **Figure 2**. Observe that $\alpha^{\otimes 2}(\text{pt}) = \text{Cl}_1^{\mathbb{C}} \otimes \text{Cl}_1^{\mathbb{C}} \cong \text{Cl}_2^{\mathbb{C}}$, and the square of $(\alpha + \text{boundary})$ assigns the left $\text{Cl}_2^{\mathbb{C}}$ -module $M \otimes M$ to the bordism in **Figure 1**. Observe that every finite dimensional $A = \text{Cl}_1^{\mathbb{C}}$ -module is free, so has the form $M = A \otimes V$ for an even vector space V . For N fermions (N odd) we have $\dim V = 2^{(N-1)/2}$. The Morita equivalence $\text{Cl}_2^{\mathbb{C}} \simeq \mathbb{C}$ identifies $\text{Cl}_2^{\mathbb{C}}$ -modules with super vector spaces; under this equivalence $M \otimes M$ corresponds to the super vector space $V \otimes V \otimes A$. By **Remark 9**, the twisted dimension of $M \otimes M$ is $2(\dim V)^2$. This is the square of the twisted dimension of M , hence the twisted dimension of M equals

$$(10) \quad \sqrt{2} \dim V = 2^{(N-1)/2} \sqrt{2}.$$

For a single fermion $N = 1$, this twisted dimension equals $\sqrt{2}$.

Remark 11.

- (1) Tensor products work as they should. For example, the tensor product of the state spaces in two theories with N odd is the tensor product of two $\text{Cl}_1^{\mathbb{C}}$ -modules, which is a $\text{Cl}_2^{\mathbb{C}}$ -module. As above, apply the Morita equivalence $\text{Cl}_2^{\mathbb{C}} \simeq \mathbb{C}$ to identify $\text{Cl}_2^{\mathbb{C}}$ -modules with super vector spaces. The twisted dimension of the resulting super vector space is $1/2$ the product of the twisted dimensions of the $\text{Cl}_1^{\mathbb{C}}$ -modules.
- (2) There are variants of this story on pin manifolds. In that case the anomaly theory is more involved; see [DeGu].

References

- [De] Pierre Deligne, *Notes on spinors*, Quantum Fields and Strings: a course for mathematicians, Vol. 1, 2 (Princeton, NJ, 1996/1997), Amer. Math. Soc., Providence, RI, 1999, pp. 99–135. [4](#)
- [DeGu] Arun Debray and Sam Gunningham, *The Arf-Brown TQFT of pin⁻ surfaces*, Topology and quantum theory in interaction, Contemp. Math., vol. 718, Amer. Math. Soc., Providence, RI, 2018, pp. 49–87. [arXiv:1803.11183](#). [4](#), [5](#)
- [DGG] Diego Delmastro, Davide Gaiotto, and Jaume Gomis, *Global anomalies on the Hilbert space*, **JHEP** **11** (2021), 142, [arXiv:2101.02218](#) [[hep-th](#)]. [1](#)
- [F1] Daniel S Freed, *Anomalies and invertible field theories*, Proc. Symp. Pure Math, Proc. Sympos. Pure Math., vol. 88, Amer. Math. Soc., Providence, RI, 2014, pp. 25–45. [arXiv:1404.7224](#). [1](#)
- [F2] Daniel S. Freed *et. al.*, *Homework*, pp. 609–717, American Mathematical Society, Providence, RI, 1999. in Quantum Fields and Strings: A Course for Mathematicians. Vol. 1. eds P. Deligne, P. Etingof, D. S. Freed, L. C. Jeffrey, D. Kazhdan, J. W. Morgan, D. R. Morrison, E. Witten. [1](#)
- [F3] Daniel S. Freed, *What is an anomaly?*, [arXiv:2307.08147](#). [3](#)
- [Gu] Sam Gunningham, *Spin Hurwitz numbers and topological quantum field theory*, **Geom. Topol.** **20** (2016), no. 4, 1859–1907, [arXiv:1201.1273](#). [4](#)
- [K] Alexei Kitaev, *Unpaired Majorana fermions in quantum wires*, Physics-Uspekhi **44** (2001), no. 10S, 131, [arXiv:cond-mat/0010440](#). [3](#)
- [LW] Jaehoon Lee and Frank Wilczek, *Algebra of Majorana Doubling*, **Phys. Rev. Lett.** **111** (2013), no. 22, 226402, [arXiv:1307.3245](#) [[cond-mat.supr-con](#)]. [3](#)
- [SS] Nathan Seiberg and Shu-Heng Shao, *Majorana chain and Ising model – (non-invertible) translations, anomalies, and emanant symmetries*, [arXiv:2307.02534](#) [[cond-mat.str-el](#)]. [1](#), [2](#)

- [tH] Gerard 't Hooft, *Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking*, *NATO Sci. Ser. B* **59** (1980), 135–157. [3](#)
- [W] Edward Witten, *Anomalies and Nonsupersymmetric D-Branes*, [arXiv:2305.01012 \[hep-th\]](#). [1](#)
- [Wa] C. T. C. Wall, *Graded Brauer groups*, *J. Reine Angew. Math.* **213** (1963/1964), 187–199. [4](#)

HARVARD UNIVERSITY, DEPARTMENT OF MATHEMATICS, SCIENCE CENTER ROOM 325, 1 OXFORD STREET,
CAMBRIDGE, MA 02138

Email address: dafr@math.harvard.edu

HARVARD UNIVERSITY, DEPARTMENT OF MATHEMATICS, SCIENCE CENTER ROOM 325, 1 OXFORD STREET,
CAMBRIDGE, MA 02138

Email address: mjh@math.harvard.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, 970 EVANS HALL #3840, BERKELEY, CA 94720-
3840

Email address: teleman@berkeley.edu