arXiv:2401.04281v2 [nucl-th] 6 Mar 2024

Comment on the article "Detailed study of the astrophysical direct capture reaction ${}^{6}\text{Li}(p,\gamma)^{7}$ Be in a potential model approach" by E. M. Tursunov, S. A. Turakulov, and K. I. Tursunmakhatov

S. B. Dubovichenko,¹ A.S. Tkachenko,¹ and R. Ya. Kezerashvili^{2,3}

¹Fesenkov Astrophysical Institute, 050020 Almaty, Kazakhstan

²New York City College of Technology, City University of New York, Brooklyn, 11201 New York, USA

³Graduate School and University Center, City University of New York, 10016 New York, USA

We explicitly present the comparison of the results for the astrophysical *S*-factor and reaction rate for the ${}^{6}\text{Li}(p,\gamma)^{7}\text{Be}$ capture process at astrophysical energies, presented in Phys. Rev. Phys. Rev. C **105**, 065806 (2022) and Phys. Rev. C **108**. 065801 (2023) obtained within the famework of potential models. We demonstrate that both potential model approaches describe successfully the astrophysical *S*-factor and reaction rate similtaneusly and reproduce the LUNA Collaboration [Phys. Rev. C **102**, 052802(R) (2020)] results.

Keywords: low and astrophysical energies, p⁶Li system, thermonuclear reaction rate, potential cluster model

In the recent paper [1] there is a statement: "Very recently, a detailed study of the above ${}^{6}\text{Li}(p,\gamma)^{7}$ Be direct capture process at astrophysical energies was performed within the potential model [28]. Various versions of the potential model have been suggested; however, none of them describe the astrophysical *S* –factor and the reaction rates simultaneously. More precisely, the temperature dependence of the reaction rates of the LUNA Collaboration [25] was not reproduced within that model. Thus, the question of whether a potential model can simultaneously describe the astrophysical *S* –factor and the reaction rates remains open."

Reference [28] is our article [2], and [25] is the LUNA Collaboration [3]. This statement is written without making any comparison between authors' results and results [2] obtained within a potential model approach. The statement motivated us to compare the astrophysical S –factor and reaction rate obtained in Refs. [2] and [1]. To shed light on the contradiction between this statement and our results [2], we present a comparison of calculations for astrophysical factor and reaction rate reported in [2] and [1] along with reported in literature experimental data.

In Fig. 1*a* we present the results of calculations for the astrophysical *S*-factor in the framework of different models and all reported experimental data, including the most recent Luna Collaboration [3] measurements. Our astrophysical *S*-factor is given with a solid red curve, and results [1] are presented with the pink curve. The *R*-matrix fit of the data from LUNA Collaboration [3] is represented with the solid blue curve. A solid green curve was obtained by Kiss et al. [11] using the weighted means of the ANCs from the analysis of the ⁶Li(³He,d)⁷Be transfer reaction within the modified two-body potential method (MTBPM). In addition, [11] contains the results for the *S*-factor of the ⁶Li(p,γ)⁷Be reaction calculated within the MTBPM, using the values of ANCs obtained from the analysis of the experimental astrophysical *S*-factors of the ⁶Li(p,γ)⁷Be reaction [3]. This



FIG. 1. (Color online) (a) Comparison of ${}^{6}\text{Li}(p,\gamma)^{7}\text{Be}$ reaction astrophysical *S*-factors. Experimental data are from $\blacktriangle - [5], \blacklozenge - [8], \blacksquare - [3], \diamondsuit - [7], \square - [4], \bigtriangledown - [6], \bigstar - [9]$. Results of calculations: red curve – our work [2]; pink curve – Ref. [1]; blue curve – Ref. [3]; black and green curves – Ref. [11]. (b) Comparison of the astrophysical reaction rates from [1–3, 12] in the range of 0.001 to 1 *T*₉, normalized to the NACRE rate [13]. The shaded area represents the uncertainties from LUNA [3] (pale red band).

result is given in Fig. 1*a* with the solid black curve. One can see that both [2] and [1] potential model approaches provide results within the error bar of the LUNA data [3]. There is a slight difference between them, which is not surprising, since in [1] slight modifications of the potentials from [2] were used. The uncertainty of experimental data allows to select other options for the potentials to describe the LUNA data [3]. Thus, the potential model [2] reproduces LUNA Collaboration [3] data for the astrophysical S – factor.

Next, we compare the reaction rates from Refs. [2] and [1]. A comparison of the reaction rates from [2] and [1] with the experimental data of the LUNA Collaboration [3] and the results of the NACRE II compilation [12], normalized to the NACRE rate [13] in the temperature range of $T_9 = 0.001 - 1$ is shown in Fig. 1*b*. The shaded area represent the uncertainties from the LUNA Collaboration [3]. The deviation between the adopted reaction rate obtained in [3] and our calculation [2] in the range of $T_9 = 0.001 - 1$ does not exceed 5%, while the deviation of the results [1] in a temperature range of $T_9 = 0.001 - 0.015$ is larger. The analytical approximations of the reaction rates reported in [3], [2], and [1] are different, but the corresponding reaction rates are within the uncertainties given by the LUNA Collaboration [3].

Thus, both [2] and [1] single-channel potential model approaches reproduce not only the absolute values of the reaction rates of Ref. [3] but also the temperature dependence of the reaction rates. Consequently, both [2] and [1] potential model approaches can describe simultaneously the *S* –factor and reaction rate reported by the LUNA Collaboration [3] for the ⁶Li(p,γ)⁷Be reaction. Therefore, it's not clear how the results of [1] differ from ours [2] and what is the novelty of [1]. In addition, it is important to mention that the advantage of the potential model approach in [2] relates to a realistic description of the p^6 Li interaction in scattering and bound-state channels and it does not need the use of any additional fitting parameters for reproducing experimental data in contrast to [1].

- [1] E. M. Tursunov, S. A. Turakulov, K. I. Tursunmakhatov, Detailed study of the astrophysical direct capture reaction ${}^{6}\text{Li}(p,\gamma)^{7}\text{Be}$ in a potential model approach. Phys. Rev. C **108**. 065801 (2023). DOI: 10.1103/PhysRevC.108.065801
- [2] S. B. Dubovichenko, A. S. Tkachenko, R. Ya. Kezerashvili, N. A. Burkova, and A. V. Dzhazairov-Kakhramanov ${}^{6}Li(p,\gamma)^{7}Be$ reaction rate in the light of the new data of the Laboratory for Underground Nuclear Astrophysics. Phys. Rev. C **105**, 065806 (2022). DOI: 10.1103/PhysRevC.105.065806
- [3] D. Piatti, T. Chillery, R. Depalo, M. Aliotta, D. Bemmerer, A. Best, et al., Underground experimental study finds no evidence of lowenergy resonance in the ${}^{6}\text{Li}(p,\gamma)^{7}\text{Be}$ reaction. Phys. Rev. C **102**, 052802(R) (2020).
- [4] S. Bashkin and R. R. Carlson, Phys. Rev. 97, 1245 (1955).
- [5] Z. E. Switkowski, J. C. Heggie, D. L. Kennedy, D. G. Sargood, F. C. Barker, and R. H. Spear, Nucl. Phys. A 331, 50 (1979).
- [6] R. Ostojić, K. Subotić, and B. Stepančić, Nuovo Cimento A 76, 73 (1983).
- [7] T. Paradellis, data taken from K. Arai, D. Baye, and P. Descouvemont, Nucl. Phys. A 699, 963 (2002).
- [8] J. J. He, S. Z. Chen, C. E. Rolfs, S. W. Xu, J.Hu, X. W. Ma,M. Wiescher, et al., Phys. Lett. B 725, 287 (2013).
- [9] A. Amar and N. Burtebayev, J. Nucl. Sci. 1, 15 (2014).
- [10] [15] K. Arai, D. Baye, and P. Descouvemont, Nucl. Phys. A 699, 963 (2002).
- [11] G. G. Kiss, M. La Cognata, R. Yarmukhamedov, et al., Phys. Rev. C 104, 015807 (2021).
- [12] Y. Xu, K. Takahashi, S. Goriely, M. Arnould, M. Ohta, and H. Utsunomiya, Nucl. Phys. A 918, 61 (2013).
- [13] C. Angulo, M. Arnould, M. Rayet, et al., Nucl. Phys. A 656, 3 (1999).