

OPTIMAL CONVERGENCE RATE FOR MIRROR DESCENT METHODS WITH SPECIAL TIME-VARYING STEP SIZES RULES

MOHAMMAD S. ALKOUSA, FEDOR S. STONYAKIN, ASMAA M. ABDO,
AND MOHAMMAD M. ALCHEIKH

ABSTRACT. In this paper, the optimal convergence rate $O(N^{-1/2})$ (where N is the total number of iterations performed by the algorithm), without the presence of a logarithmic factor, is proved for mirror descent algorithms with special time-varying step sizes, for solving classical constrained non-smooth problems, problems with the composite model and problems with non-smooth functional (inequality types) constraints. The proven result is an improvement on the well-known rate $O(\log(N)N^{-1/2})$ for the mirror descent algorithms with the time-varying step sizes under consideration. It was studied a new weighting scheme assigns smaller weights to the initial points and larger weights to the most recent points. This scheme improves the convergence rate of the considered mirror descent methods, which in the conducted numerical experiments outperform the other methods providing a better solution in all the considered test problems.

1. INTRODUCTION

The first method for unconstrained minimization of non-smooth convex function was proposed with fixed step size in [31]. In the next years, there were developed several strategies for choosing the step sizes [29]. Among them, are fixed step length, non-summable diminishing step, square summable but not summable step [11], Polyak step [29], and many others (see Table 1, below). The optimal convergence rate of the subgradient method is known as $O(N^{-1/2})$, where N is the total number of iterations performed by the algorithm [4, 12, 20, 25]. This rate was proved by using many rules of the step size. One of them is fixed (constant), namely $\gamma_k = O(1)/\sqrt{N}$ for $k = 1, \dots, N$ (with special chosen of the constant $O(1)$), which minimizes the upper bound of the difference between the best value of the objective function attained (i.e., the best value observed during the minimization process) and the optimal value. But most of the time it is preferable to use the time-varying step size since the constant step size rule requires the a priori

Key words and phrases. Convex optimization, Non-smooth problem, Composite problem, Problem with functional constraints, Mirror descent, Mirror-C descent, Optimal convergence rate, Sub-optimal convergence rate, Time-varying step size.

Preprint under update.

knowledge of the total number of iterations employed by the method. From the mentioned fixed steps, the inspiration was to take $\gamma_k = O(1)/\sqrt{k}$ for $k = 1, \dots, N$, but the convergence rate, with this rule, became sub-optimal $O(\log(N)N^{-1/2})$. Recently in [39], for the projected subgradient method it was proved the optimal convergence rate using the previously mentioned time-varying step size with a new weighting scheme for the generated points each iteration of the algorithm.

The next important advancement in this area is related to the development of the mirror descent method which originated in [23, 24] and was later analyzed in [5]. It is considered as the non-Euclidean extension of standard subgradient methods. This method is used in many applications, see [8, 18, 26, 27, 35] and references therein. The standard subgradient methods employ the Euclidean distance function with a suitable step size in the projection step. Mirror descent extends the standard projected subgradient methods by employing a nonlinear distance function with an optimal step size in the nonlinear projection step [22]. The mirror descent method is also applicable to optimization problems in Banach spaces where gradient descent is not [14]. An extension of the mirror descent method for constrained problems was proposed in [6, 24]. Usually, the step size and stopping criterion for the mirror descent method require knowing the Lipschitz constant of the objective function and constraints, if any. Adaptive step sizes, which do not require this information, are considered for unconstrained problems in [7], and for constrained problems in [6].

In addition to the previously mentioned problems, there is a more general class of problems, which is non-smooth optimization problems with non-smooth functional (inequality types) constraints (see problem (5.1) or its equivalence (5.2)). This class of problems arises and attracts widespread interest in many areas of modern large-scale optimization and its applications [9, 30]. On continuous optimization with functional constraints, there is a long history of studies. The monographs in this area include [10, 28]. Some of the works on first-order methods for convex optimization problems with convex functional constraints include [3, 17, 21, 32, 33, 34, 37] for the deterministic setting and [1, 2, 19, 38] for the stochastic setting.

As we pointed out the mirror descent algorithm was proposed and analyzed a long time ago for different step size rules. However, the optimal convergence rate $O(N^{-1/2})$ of the algorithm was first proved, as for the subgradient method, with the fixed step size $\gamma_k = O(1)/\sqrt{N}$ for $k = 1, \dots, N$. But, when the step sizes are time-varying, namely $\gamma_k = O(1)/\sqrt{k}$, it was proved the sub-optimal convergence rate $O(\log(N)N^{-1/2})$, see [4, 12, 20, 25] for details. Although the constant step size rule is relatively easy to analyze but has the disadvantage of requiring the a priori knowledge of the total number of iterations employed by the method. In practical situations, the number of iterations is not fixed a priori, and stopping criteria different than merely fixing the total number of iterations is usually imposed. This is why

dynamic, or time-varying (i.e., non-constant) step size rules are important [12].

In this paper, we extend the results of [39] to the mirror descent methods. We proved the optimal convergence rate, without the presence of a logarithmic factor, for mirror descent methods with special time-varying step sizes, for solving classical constrained non-smooth problems (see Sect. 3), constrained composite problems (see Sect. 4), and problems with functional constraints (see Sect. 5). We demonstrate the efficiency of the theoretical studied results by some conducted experiments for some constrained (with and without functional constraints) optimization problems, such as best approximation problem, Fermat-Torricelli-Steiner problem, smallest covering ball problem and the maximum of a finite collection of linear functions.

The paper consists of an introduction and 5 main sections. In Sect. 2 we mentioned the basic facts and tools and mirror descent basics. Sect. 3 devoted to the proof of the optimal convergence rate of the mirror descent algorithm with time-varying step size rules for solving the classical constrained non-smooth optimization problem. In Sect. 4 the same analysis, that in Sect. 3, performed for more general optimization problems, namely the constrained composite optimization problem. Section 5 is devoted to the problem with functional constraints, which is a more general class of non-smooth optimization problems. In Sect. 6 we present some numerical experiments which demonstrate the effectiveness of the proposed algorithms and analysis.

2. FUNDAMENTALS

Let $(\mathbf{E}, \|\cdot\|)$ be a normed finite-dimensional vector space, with an arbitrary norm $\|\cdot\|$, and \mathbf{E}^* be the conjugate space of \mathbf{E} with the following norm

$$\|y\|_* = \max_{x \in \mathbf{E}} \{\langle y, x \rangle : \|x\| \leq 1\},$$

where $\langle y, x \rangle$ is the value of the continuous linear functional $y \in \mathbf{E}^*$ at $x \in \mathbf{E}$.

Let $Q \subset \mathbf{E}^n$ be a compact convex set, and $\psi : Q \rightarrow \mathbb{R}$ be a proper closed differentiable and σ -strongly convex (called prox-function or distance generating function) with $\sigma > 0$. The corresponding Bregman divergence is defined as

$$V_\psi(x, y) = \psi(x) - \psi(y) - \langle \nabla \psi(x), y - x \rangle, \quad \forall x, y \in Q.$$

For the Bregmann divergence, it holds the following inequality

$$(2.1) \quad V_\psi(x, y) \geq \frac{\sigma}{2} \|y - x\|^2, \quad \forall x, y \in Q.$$

The following identity, known as the three points identity, is essential in the analysis of the mirror descent method.

Lemma 2.1. (*three points identity*) [13] *Suppose that $\psi : \mathbf{E} \rightarrow (-\infty, \infty]$ is proper closed, convex, and differentiable over $\text{dom}(\partial(\psi))$. Let $a, b \in$*

$\text{dom}(\partial(\psi))$ and $c \in \text{dom}(\psi)$. Then it holds

$$(2.2) \quad \langle \nabla \psi(b) - \nabla \psi(a), c - a \rangle = V_\psi(c, a) + V_\psi(a, b) - V_\psi(c, b).$$

Fenchel-Young inequality. For any $a \in \mathbf{E}, b \in \mathbf{E}^*$, it holds the following inequality

$$(2.3) \quad |\langle a, b \rangle| \leq \frac{\|a\|^2}{2\lambda} + \frac{\lambda \|b\|_*}{2}, \quad \forall \lambda > 0.$$

In what follows, we denote the subdifferential of f at x by $\partial f(x)$, and the subgradient of f at any point x by $\nabla f(x) \in \partial f(x)$.

3. MIRROR DESCENT METHOD FOR CONSTRAINED PROBLEM

In this section, we consider the following constrained optimization problem

$$(3.1) \quad \min_{x \in Q} f(x),$$

where f is a convex Lipschitz continuous function (non-smooth) with Lipschitz constant $M_f > 0$, i.e.,

$$(3.2) \quad |f(x) - f(y)| \leq M_f \|x - y\|, \quad \forall x, y \in Q.$$

For problem (3.1), we use Algorithm 1, under consideration

$$(3.3) \quad \max_{x \in Q} V_\psi(x^*, x) \leq V_\psi(x^*, x^1) < \infty,$$

where $x^1 \in Q$ is a chosen (dependently on Q) initial point and x^* is an optimal solution of (3.1) nearest to x^1 .

Algorithm 1 Mirror descent method.

Require: step sizes $\{\gamma_k\}_{k \geq 1}$, initial point $x^1 \in Q$ s.t. (3.3) holds, number of iterations N .

- 1: **for** $k = 1, 2, \dots, N$ **do**
 - 2: Calculate $\nabla f(x^k) \in \partial f(x^k)$,
 - 3: $x^{k+1} = \arg \min_{x \in Q} \left\{ \langle x, \nabla f(x^k) \rangle + \frac{1}{\gamma_k} V_\psi(x, x^k) \right\}$.
 - 4: **end for**
-

Theorem 3.1. Let f be an M_f -Lipschitz convex function, and assume $\|\nabla f(x)\|_* \leq M_f, \forall x \in Q$. Then for problem (3.1), by Algorithm 1, with a positive non-increasing sequence of step sizes $\{\gamma_k\}_{k \geq 1}$, for any fixed $m \geq -1$, it satisfies the following inequality

$$(3.4) \quad f(\hat{x}) - f(x^*) \leq \frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \left(\frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \right),$$

where $\hat{x} = \frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$, and x^* is an optimal solution of (3.1).

Proof. Let $\tilde{f}(x) := \langle x, \nabla f(x^k) \rangle + \frac{1}{\gamma_k} V_\psi(x, x^k)$, then from Algorithm 1 we have $x^{k+1} = \arg \min_{x \in Q} \tilde{f}(x)$. By the optimality condition, we get

$$\langle \nabla \tilde{f}(x^{k+1}), x - x^{k+1} \rangle \geq 0, \quad \forall x \in Q.$$

Thus,

$$\langle \gamma_k \nabla f(x^k) + \nabla \psi(x^{k+1}) - \nabla \psi(x^k), x - x^{k+1} \rangle \geq 0, \quad \forall x \in Q.$$

In particular for $x = x^*$, we have

$$(3.5) \quad \langle \nabla \psi(x^k) - \nabla \psi(x^{k+1}) - \gamma_k \nabla f(x^k), x^* - x^{k+1} \rangle \leq 0.$$

By using the definition of a subgradient of f at x^* , we get

$$\begin{aligned} \gamma_k \left(f(x^k) - f(x^*) \right) &\leq \gamma_k \langle \nabla f(x^k), x^k - x^* \rangle \\ &= \underbrace{\langle \nabla \psi(x^k) - \nabla \psi(x^{k+1}) - \gamma_k \nabla f(x^k), x^* - x^{k+1} \rangle}_{:=J_1} \\ &\quad + \underbrace{\langle \nabla \psi(x^{k+1}) - \nabla \psi(x^k), x^* - x^{k+1} \rangle}_{:=J_2} \\ &\quad + \underbrace{\langle \gamma_k \nabla f(x^k), x^k - x^{k+1} \rangle}_{:=J_3}. \end{aligned}$$

From (3.5), we find $J_1 \leq 0$. By Lemma 2.1, we have

$$\begin{aligned} J_2 &= -\langle \nabla \psi(x^k) - \nabla \psi(x^{k+1}), x^* - x^{k+1} \rangle \\ &= -\left(V_\psi(x^*, x^{k+1}) + V_\psi(x^{k+1}, x^k) - V_\psi(x^*, x^k) \right) \\ &= V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1}) - V_\psi(x^{k+1}, x^k). \end{aligned}$$

By the Fenchel-Young inequality (2.3), with $\lambda = \sigma > 0$, we find

$$J_3 \leq \frac{\gamma_k^2}{2\sigma} \|\nabla f(x^k)\|_*^2 + \frac{\sigma}{2} \|x^k - x^{k+1}\|^2.$$

Therefore, we get

$$\begin{aligned} \gamma_k \left(f(x^k) - f(x^*) \right) &\leq V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1}) - V_\psi(x^{k+1}, x^k) \\ &\quad + \frac{\gamma_k^2}{2\sigma} \|\nabla f(x^k)\|_*^2 + \frac{\sigma}{2} \|x^k - x^{k+1}\|^2. \end{aligned}$$

From (2.1), we have

$$V_\psi(x^{k+1}, x^k) \geq \frac{\sigma}{2} \|x^{k+1} - x^k\|^2.$$

Thus, we get the following inequality

$$(3.6) \quad f(x^k) - f(x^*) \leq \frac{1}{\gamma_k} \left(V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1}) \right) + \frac{\gamma_k}{2\sigma} \|\nabla f(x^k)\|_*^2.$$

By multiplying both sides of (3.6) by $\frac{1}{\gamma_k^m}$, and taking the sum from 1 to N , we get

$$\begin{aligned} \sum_{k=1}^N \frac{1}{\gamma_k^m} \left(f(x^k) - f(x^*) \right) &\leq \sum_{k=1}^N \frac{1}{\gamma_k^{m+1}} \left(V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1}) \right) \\ &\quad + \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{2\sigma\gamma_k^{m-1}}. \end{aligned}$$

Since f is a convex function, we have

$$\begin{aligned} \left(\sum_{k=1}^N \frac{1}{\gamma_k^m} \right) \left[f \left(\frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k \right) - f(x^*) \right] &\leq \\ &\leq \sum_{k=1}^N \frac{1}{\gamma_k^m} \left(f(x^k) - f(x^*) \right). \end{aligned}$$

Therefore, by setting $\hat{x} := \frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$, we get

$$\begin{aligned} \left(\sum_{k=1}^N \gamma_k^{-m} \right) (f(\hat{x}) - f(x^*)) &\leq \frac{1}{\gamma_1^{m+1}} (V_\psi(x^*, x^1) - V_\psi(x^*, x^2)) \\ &\quad + \sum_{k=2}^{N-1} \frac{1}{\gamma_k^{m+1}} (V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1})) \\ &\quad + \frac{1}{\gamma_N^{m+1}} (V_\psi(x^*, x^N) - V_\psi(x^*, x^{N+1})) + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \\ &\leq \frac{1}{\gamma_1^{m+1}} V_\psi(x^*, x^1) + \frac{1}{\gamma_N^{m+1}} V_\psi(x^*, x^N) + \sum_{k=2}^{N-1} \frac{1}{\gamma_k^{m+1}} V_\psi(x^*, x^k) \\ &\quad - \frac{1}{\gamma_1^{m+1}} V_\psi(x^*, x^2) - \sum_{k=2}^{N-1} \frac{1}{\gamma_k^{m+1}} V_\psi(x^*, x^{k+1}) + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \\ &= \frac{1}{\gamma_1^{m+1}} V_\psi(x^*, x^1) + \sum_{k=2}^N \frac{1}{\gamma_k^{m+1}} V_\psi(x^*, x^k) - \frac{1}{\gamma_1^{m+1}} V_\psi(x^*, x^2) \\ &\quad - \sum_{k=3}^N \frac{1}{\gamma_{k-1}^{m+1}} V_\psi(x^*, x^k) + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \end{aligned}$$

Due to (3.3), we find that $V_\psi(x^*, x^k) \leq V_\psi(x^*, x^1)$, $\forall k = 2, \dots, N$. Hence we get

$$\begin{aligned}
& \left(\sum_{k=1}^N \gamma_k^{-m} \right) (f(\hat{x}) - f(x^*)) \leq \frac{1}{\gamma_1^{m+1}} V_\psi(x^*, x^1) + \sum_{k=2}^N \frac{1}{\gamma_k^{m+1}} V_\psi(x^*, x^k) \\
& \quad - \sum_{k=2}^N \frac{1}{\gamma_{k-1}^{m+1}} V_\psi(x^*, x^k) + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \\
& \leq \frac{1}{\gamma_1^{m+1}} V_\psi(x^*, x^1) + V_\psi(x^*, x^1) \sum_{k=2}^N \left(\frac{1}{\gamma_k^{m+1}} - \frac{1}{\gamma_{k-1}^{m+1}} \right) + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \\
& = \frac{1}{\gamma_1^{m+1}} V_\psi(x^*, x^1) + V_\psi(x^*, x^1) \left(-\frac{1}{\gamma_1^{m+1}} + \frac{1}{\gamma_N^{m+1}} \right) + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \\
& = \frac{1}{\gamma_N^{m+1}} V_\psi(x^*, x^1) + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}.
\end{aligned}$$

Finally, by dividing by $\sum_{k=1}^N \gamma_k^{-m}$, we get the desired inequality

$$f(\hat{x}) - f(x^*) \leq \frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \left(\frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \right).$$

□

As a special case of Theorem 3.1, with a special value of the parameter m , we can deduce the well-known sub-optimal convergence rate of mirror descent algorithm $O\left(\frac{\log(N)}{\sqrt{N}}\right)$, see [4, 12, 20, 25], with the following (adaptive and non-adaptive) time-varying step size rules

$$(3.7) \quad \gamma_k = \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}, \quad \text{or} \quad \gamma_k = \frac{\sqrt{2\sigma}}{\|\nabla f(x^k)\|_* \sqrt{k}}, \quad k = 1, 2, \dots, N.$$

Corollary 3.2. *Let f be an M_f -Lipschitz convex function, and assume $\|\nabla f(x)\|_* \leq M_f, \forall x \in Q$. Then for problem (3.1), by Algorithm 1, with $m = -1$, and the time-varying step sizes given in (3.7), it satisfies the following sub-optimal convergence rate*

$$(3.8) \quad f(\tilde{x}) - f(x^*) \leq \frac{M_f (V_\psi(x^*, x^1) + 1 + \log(N))}{\sqrt{\sigma}} \frac{1}{\sqrt{N}} = O\left(\frac{\log(N)}{\sqrt{N}}\right),$$

where $\tilde{x} = \frac{1}{\sum_{k=1}^N \gamma_k} \sum_{k=1}^N \gamma_k x^k$.

Proof. By setting $m = -1$ in (3.4), we get the following inequality

$$(3.9) \quad f(\tilde{x}) - f(x^*) \leq \frac{V_\psi(x^*, x^1) + \frac{1}{2\sigma} \sum_{k=1}^N \gamma_k^2 \|\nabla f(x^k)\|_*^2}{\sum_{k=1}^N \gamma_k},$$

where $\tilde{x} = \frac{1}{\sum_{k=1}^N \gamma_k} \sum_{k=1}^N \gamma_k x^k$.

Case 1 (non-adaptive rule). When $\gamma_k = \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (3.9) we find

$$f(\tilde{x}) - f(x^*) \leq \frac{M_f}{\sqrt{2\sigma}} \frac{V_\psi(x^*, x^1) + \sum_{k=1}^N \frac{1}{k}}{\sum_{k=1}^N \frac{1}{\sqrt{k}}},$$

where $\tilde{x} = \frac{1}{\sum_{k=1}^N \frac{1}{\sqrt{k}}} \sum_{k=1}^N \frac{1}{\sqrt{k}} x^k$. But

$$\sum_{k=1}^N \frac{1}{k} \leq 1 + \log(N), \quad \text{and} \quad \sum_{k=1}^N \frac{1}{\sqrt{k}} \geq 2\sqrt{N+1} - 2, \quad \forall N \geq 1.$$

Therefore,

$$\begin{aligned} f(\tilde{x}) - f(x^*) &\leq \frac{M_f}{\sqrt{2\sigma}} \frac{V_\psi(x^*, x^1) + 1 + \log(N)}{2\sqrt{N+1} - 2} \\ &\leq \frac{M_f}{\sqrt{\sigma}} \frac{V_\psi(x^*, x^1) + 1 + \log(N)}{\sqrt{N}}, \quad \forall N \geq 1. \end{aligned}$$

Where in the last inequality, we used the fact $2\sqrt{2}(\sqrt{N+1}-1) \geq \sqrt{N}$, $\forall N \geq 1$.

Case 2 (adaptive rule). When $\gamma_k = \frac{\sqrt{2\sigma}}{\|\nabla f(x^k)\|_* \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (3.9) we find

$$\begin{aligned} f(\tilde{x}) - f(x^*) &\leq \frac{V_\psi(x^*, x^1) + \sum_{k=1}^N \frac{1}{k}}{\sum_{k=1}^N \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}} = \frac{M_f}{\sqrt{2\sigma}} \frac{V_\psi(x^*, x^1) + \sum_{k=1}^N \frac{1}{k}}{\sum_{k=1}^N \frac{1}{\sqrt{k}}} \\ &\leq \frac{M_f}{\sqrt{\sigma}} \frac{V_\psi(x^*, x^1) + 1 + \log(N)}{\sqrt{N}}, \end{aligned}$$

where $\tilde{x} = \frac{1}{\sum_{k=1}^N (\|\nabla f(x^k)\|_* \sqrt{k})^{-1}} \sum_{k=1}^N \left(\|\nabla f(x^k)\|_* \sqrt{k} \right)^{-1} x^k$. □

Also, from Theorem 3.1, with a special value of the parameter m , we can obtain the optimal convergence rate of mirror descent algorithm $O\left(\frac{1}{\sqrt{N}}\right)$, with the time-varying step sizes given in (3.7).

Corollary 3.3. *Let f be an M_f -Lipschitz convex function, and assume $\|\nabla f(x)\|_* \leq M_f, \forall x \in Q$. Then for problem (3.1), by Algorithm 1, with $m = 0$, and the time-varying step sizes given in (3.7), it satisfies the following optimal convergence rate*

$$(3.10) \quad f(\bar{x}) - f(x^*) \leq \frac{M_f (2 + V_\psi(x^*, x^1))}{\sqrt{2\sigma}} \frac{1}{\sqrt{N}} = O\left(\frac{1}{\sqrt{N}}\right),$$

where $\bar{x} = \frac{1}{N} \sum_{k=1}^N x^k$.

Proof. By setting $m = 0$ in (3.4), we get the following inequality

$$(3.11) \quad f(\bar{x}) - f(x^*) \leq \frac{1}{N} \left(\frac{V_\psi(x^*, x^1)}{\gamma_N} + \frac{1}{2\sigma} \sum_{k=1}^N \gamma_k \|\nabla f(x^k)\|_*^2 \right),$$

where $\bar{x} = \frac{1}{N} \sum_{k=1}^N x^k$.

Case 1 (non-adaptive rule). When $\gamma_k = \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (3.11) we find

$$f(\bar{x}) - f(x^*) \leq \frac{1}{N} \left(\frac{M_f \sqrt{N} V_\psi(x^*, x^1)}{\sqrt{2\sigma}} + \frac{M_f}{\sqrt{2\sigma}} \sum_{k=1}^N \frac{1}{\sqrt{k}} \right).$$

But

$$\sum_{k=1}^N \frac{1}{\sqrt{k}} \leq 2\sqrt{N}, \quad \forall N \geq 1.$$

Therefore,

$$f(\bar{x}) - f(x^*) \leq \frac{1}{N} \frac{M_f}{\sqrt{2\sigma}} \left(\sqrt{N} V_\psi(x^*, x^1) + 2\sqrt{N} \right) = \frac{M_f(2 + V_\psi(x^*, x^1))}{\sqrt{2\sigma}} \frac{1}{\sqrt{N}}.$$

Case 2 (adaptive rule). When $\gamma_k = \frac{\sqrt{2\sigma}}{\|\nabla f(x^k)\|_* \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (3.11) we find

$$\begin{aligned} f(\bar{x}) - f(x^*) &\leq \frac{1}{N} \left(\frac{\sqrt{N} \|\nabla f(x^N)\|_* V_\psi(x^*, x^1)}{\sqrt{2\sigma}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\sqrt{2\sigma}}{\sqrt{k}} \|\nabla f(x^k)\|_* \right) \\ &\leq \frac{1}{N} \left(\frac{M_f \sqrt{N} V_\psi(x^*, x^1)}{\sqrt{2\sigma}} + \frac{M_f}{\sqrt{2\sigma}} \sum_{k=1}^N \frac{1}{\sqrt{k}} \right) \\ &\leq \frac{1}{N} \frac{M_f}{\sqrt{2\sigma}} \left(\sqrt{N} V_\psi(x^*, x^1) + 2\sqrt{N} \right) \\ &= \frac{M_f(2 + V_\psi(x^*, x^1))}{\sqrt{2\sigma}} \frac{1}{\sqrt{N}}. \end{aligned}$$

□

Remark 3.4. By setting any $m > -1$ in (3.4) (in Theorem 3.1), the convergence rate of Algorithm 1 with the time-varying step sizes given in (3.7), will be optimal $O\left(\frac{1}{\sqrt{N}}\right)$ without presence of a $\log(N)$ factor. In comparison with the sub-optimal convergence rate (3.8), when $m > 0$, the weighting scheme $\frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$ assigns smaller weights to the initial points and larger weights to the most recent points that generated by Algorithm 1. This fact will be shown in numerical experiments.

4. MIRROR DESCENT METHOD FOR CONSTRAINED COMPOSITE PROBLEM

In this section, we will consider a more general problem than problem (3.1), namely

$$(4.1) \quad \min_{x \in Q} \{F(x) = f(x) + h(x)\},$$

where $Q = \text{dom}(h) \cap \text{dom}(\partial\psi)$, f and h satisfy the following assumption

Assumption 4.1. (properties of f and h)

- (1) f and h are proper closed and convex;
- (2) $\text{dom}(h) \subseteq \text{int}(\text{dom}(f))$;
- (3) f is M_f -Lipschitz continuous on $\text{dom}(h)$ for any $M_f > 0$.

We also, assume that the optimal set of the problem (4.1) is nonempty.

For the problem (4.1), with an additional assumption that h is non-negative, we mention the following well-defined algorithm [4].

Algorithm 2 Mirror-C descent method.

Require: step sizes $\{\gamma_k\}_{k \geq 1}$, initial point $x^1 \in Q$ s.t. (3.3) holds, number of iterations N .

- 1: **for** $k = 1, 2, \dots, N$ **do**
 - 2: Calculate $\nabla f(x^k) \in \partial f(x^k)$,
 - 3: $x^{k+1} = \arg \min_{x \in Q} \{\gamma_k \langle x, \nabla f(x^k) \rangle + \gamma_k h(x) + V_\psi(x, x^k)\}$.
 - 4: **end for**
-

Let us mention the following key lemma for the analysis of the Algorithm 2.

Lemma 4.2. [4] *Let*

- (1) $\psi : \mathbf{E} \rightarrow (-\infty, \infty]$ *be a proper closed and convex function differentiable over* $\text{dom}(\partial\psi)$,
- (2) $\varphi : \mathbf{E} \rightarrow (-\infty, \infty]$ *be a proper closed and convex function satisfying* $\text{dom}(\partial\varphi) \subseteq \text{dom}(\partial\psi)$,
- (3) $\psi + \mathbb{I}_{\text{dom}(\varphi)}$ *be a σ -strongly convex, where \mathbb{I}_A is the indicator function of the set A .*

Assume that $b \in \text{dom}(\partial(\psi))$, and let a be defined by

$$a = \arg \min_{x \in \mathbf{E}} \{\varphi(x) + V_\psi(x, b)\}.$$

Then we have

$$(4.2) \quad \langle \nabla\psi(b) - \nabla\psi(a), u - a \rangle \leq \varphi(u) - \varphi(a), \quad \forall u \in \text{dom}(\varphi).$$

Theorem 4.3. *Suppose that Assumption 4.1 holds and h is a non-negative function. Then for problem (4.1), by Algorithm 2, with a positive non-increasing sequence of step sizes $\{\gamma_k\}_{k \geq 1}$, for any fixed $m \geq -1$, it satisfies*

the following inequality

(4.3)

$$F(\hat{x}) - F(x^*) \leq \frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \left(\frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}} \right),$$

where $\hat{x} = \frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$, and x^* is an optimal solution of (4.1).

Proof. From (4.2), with $b = x^k, a = x^{k+1}$ and $\varphi(x) = \gamma_k \langle x, \nabla f(x^k) \rangle + \gamma_k h(x)$, we get the following inequality

(4.4)

$$\langle \nabla \psi(x^k) - \nabla \psi(x^{k+1}), u - x^{k+1} \rangle \leq \gamma_k \langle u - x^{k+1}, \nabla f(x^k) \rangle + \gamma_k h(u) - \gamma_k h(x^{k+1}).$$

Now, from three points identity (2.2), with $a = x^{k+1}, b = x^k$ and $c = u$, we get

(4.5)

$$\langle \nabla \psi(x^k) - \nabla \psi(x^{k+1}), u - x^{k+1} \rangle = V_\psi(u, x^{k+1}) + V_\psi(x^{k+1}, x^k) - V_\psi(u, x^k).$$

By combining (4.4) and (4.5), we get the following inequality

$$(4.6) \quad \begin{aligned} V_\psi(u, x^{k+1}) + V_\psi(x^{k+1}, x^k) - V_\psi(u, x^k) &\leq \gamma_k \langle u - x^{k+1}, \nabla f(x^k) \rangle \\ &\quad + \gamma_k h(u) - \gamma_k h(x^{k+1}). \end{aligned}$$

From (4.6), we find

$$(4.7) \quad \begin{aligned} \gamma_k \langle \nabla f(x^k), x^k - u \rangle + \gamma_k h(x^{k+1}) - \gamma_k h(u) &\leq V_\psi(u, x^k) - V_\psi(u, x^{k+1}) \\ &\quad - V_\psi(x^{k+1}, x^k) + \langle \gamma_k \nabla f(x^k), x^k - x^{k+1} \rangle. \end{aligned}$$

Therefore, by (2.1) and Fenchel-Young inequality (2.3) with $\lambda = \sigma$, from (4.7) we get the following

(4.8)

$$\begin{aligned} \gamma_k \langle \nabla f(x^k), x^k - u \rangle + \gamma_k h(x^{k+1}) - \gamma_k h(u) &\leq V_\psi(u, x^k) - V_\psi(u, x^{k+1}) \\ &\quad - \frac{\sigma}{2} \|x^{k+1} - x^k\|^2 + \frac{\gamma_k^2}{2\sigma} \|\nabla f(x^k)\|_*^2 + \frac{\sigma}{2} \|x^{k+1} - x^k\|^2. \\ &= V_\psi(u, x^k) - V_\psi(u, x^{k+1}) + \frac{\gamma_k^2}{2\sigma} \|\nabla f(x^k)\|_*^2. \end{aligned}$$

Since f is a convex function, it holds

$$f(u) - f(x^k) \geq \langle \nabla f(x^k), u - x^k \rangle \implies f(x^k) - f(u) \leq \langle \nabla f(x^k), x^k - u \rangle,$$

and by setting $u = x^*$ in (4.8), we get the following inequality

$$\begin{aligned} \gamma_k \left(f(x^k) - f(x^*) \right) + \gamma_k h(x^{k+1}) - \gamma_k h(x^*) &\leq V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1}) \\ &\quad + \frac{\gamma_k^2}{2\sigma} \|\nabla f(x^k)\|_*^2, \end{aligned}$$

which means

(4.9)

$$f(x^k) + h(x^{k+1}) - F(x^*) \leq \frac{1}{\gamma_k} \left(V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1}) \right) + \frac{\gamma_k}{2\sigma} \|\nabla f(x^k)\|_*^2.$$

By multiplying both sides of (4.9) by $\frac{1}{\gamma_k^m}$, and taking the sum from 1 to N , we get

$$\begin{aligned} \sum_{k=1}^N \frac{1}{\gamma_k^m} \left(f(x^k) + h(x^{k+1}) - F(x^*) \right) &\leq \sum_{k=1}^N \frac{1}{\gamma_k^{m+1}} \left(V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1}) \right) \\ &\quad + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}. \end{aligned}$$

Now, in a similar way as in the proof of Theorem 3.1, we conclude

$$(4.10) \quad \sum_{k=1}^N \frac{1}{\gamma_k^m} \left(f(x^k) + h(x^{k+1}) - F(x^*) \right) \leq \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}.$$

By adding the term $\frac{h(x^1)}{\gamma_1^m} - \frac{h(x^{N+1})}{\gamma_N^m}$ to both sides of (4.10), we get the following

$$\begin{aligned} \sum_{k=2}^N \frac{1}{\gamma_k^m} \left(f(x^k) + h(x^{k+1}) - F(x^*) \right) &+ \frac{f(x^1)}{\gamma_1^m} + \frac{h(x^2)}{\gamma_1^m} - \frac{F(x^*)}{\gamma_1^m} + \frac{h(x^1)}{\gamma_1^m} \\ &- \frac{h(x^{N+1})}{\gamma_N^m} \leq \frac{h(x^1)}{\gamma_1^m} - \frac{h(x^{N+1})}{\gamma_N^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}, \end{aligned}$$

Because of h is a non-negative, we have $\frac{h(x^{N+1})}{\gamma_N^m} > 0$, and then we get the following

$$\begin{aligned} \sum_{k=2}^N \left(\frac{1}{\gamma_k^m} f(x^k) + \frac{1}{\gamma_k^m} h(x^{k+1}) - \frac{1}{\gamma_k^m} F(x^*) \right) &+ \frac{1}{\gamma_1^m} (F(x^1) - F(x^*)) + \frac{h(x^2)}{\gamma_1^m} \\ &- \frac{h(x^{N+1})}{\gamma_N^m} \leq \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}, \end{aligned}$$

which means

$$\begin{aligned} \sum_{k=2}^N \left(\frac{1}{\gamma_k^m} f(x^k) - \frac{1}{\gamma_k^m} F(x^*) \right) &+ \sum_{k=2}^N \frac{1}{\gamma_k^m} h(x^{k+1}) + \frac{1}{\gamma_1^m} (F(x^1) - F(x^*)) \\ &+ \frac{h(x^2)}{\gamma_1^m} - \frac{h(x^{N+1})}{\gamma_N^m} \leq \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}. \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{k=2}^N \left(\frac{1}{\gamma_k^m} f(x^k) - \frac{1}{\gamma_k^m} F(x^*) \right) &+ \sum_{k=3}^{N+1} \frac{1}{\gamma_{k-1}^m} h(x^k) + \frac{1}{\gamma_1^m} (F(x^1) - F(x^*)) \\ &+ \frac{h(x^2)}{\gamma_1^m} - \frac{h(x^{N+1})}{\gamma_N^m} \leq \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \sum_{k=2}^N \left(\frac{1}{\gamma_k^m} f(x^k) - \frac{1}{\gamma_k^m} F(x^*) \right) + \frac{1}{\gamma_1^m} (F(x^1) - F(x^*)) + \sum_{k=2}^N \frac{1}{\gamma_{k-1}^m} h(x^k) \leq \\ & \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}. \end{aligned}$$

Thus, we get

$$\begin{aligned} & \sum_{k=2}^N \left(\frac{1}{\gamma_k^m} f(x^k) + \frac{1}{\gamma_{k-1}^m} h(x^k) - \frac{1}{\gamma_k^m} F(x^*) \right) + \frac{1}{\gamma_1^m} (F(x^1) - F(x^*)) \leq \\ & \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}. \end{aligned}$$

Because of the sequence $\{\gamma_k\}_{k \geq 1}$ non-increasing, we have $\frac{1}{\gamma_{k-1}^m} \geq \frac{1}{\gamma_k^m}$, and then we conclude

$$\begin{aligned} & \sum_{k=2}^N \left(\frac{1}{\gamma_k^m} f(x^k) + \frac{1}{\gamma_k^m} h(x^k) - \frac{1}{\gamma_k^m} F(x^*) \right) + \frac{1}{\gamma_1^m} (F(x^1) - F(x^*)) \leq \\ & \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}, \end{aligned}$$

which means

$$\sum_{k=1}^N \frac{1}{\gamma_k^m} (F(x^k) - F(x^*)) \leq \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}.$$

From the convexity of the function F , we have

$$\begin{aligned} & \left(\sum_{k=1}^N \frac{1}{\gamma_k^m} \right) \left[F \left(\frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k \right) - F(x^*) \right] \leq \\ & \leq \sum_{k=1}^N \frac{1}{\gamma_k^m} (F(x^k) - F(x^*)). \end{aligned}$$

Therefore, by setting $\hat{x} = \frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$, we get the following inequality

$$(4.11) \quad \left(\sum_{k=1}^N \frac{1}{\gamma_k^m} \right) (F(\hat{x}) - F(x^*)) \leq \frac{h(x^1)}{\gamma_1^m} + \frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla f(x^k)\|_*^2}{\gamma_k^{m-1}}.$$

By dividing both sides of (4.11) by $\sum_{k=1}^N \gamma_k^{-m}$, we get the desired inequality (4.3). □

As a special case of Theorem 4.3, with a special value of the parameter m , we can deduce the well-known sub-optimal convergence rate of the mirror-C descent method (Algorithm 2), namely $O\left(\frac{\log(N)}{N}\right)$, with the time-varying step sizes given in (3.7) (see [4, 12, 20, 25]).

Corollary 4.4. *Suppose that Assumption 4.1 holds and h is a non-negative function. Then for problem (4.1), by Algorithm 2, with $m = -1$, and the time-varying step sizes given in (3.7), it satisfies the following sub-optimal convergence rate*

$$(4.12) \quad \begin{aligned} F(\tilde{x}) - F(x^*) &\leq \frac{M_f}{\sqrt{\sigma}} \left(\frac{\sqrt{2\sigma}}{M_f} h(x^1) + V_\psi(x^*, x^1) + 1 + \log(N) \right) \frac{1}{\sqrt{N}} \\ &= O\left(\frac{\log(N)}{\sqrt{N}}\right), \end{aligned}$$

where $\tilde{x} = \frac{1}{\sum_{k=1}^N \gamma_k} \sum_{k=1}^N \gamma_k x^k$.

Proof. By setting $m = -1$ in (4.3), we get the following inequality

$$(4.13) \quad F(\tilde{x}) - F(x^*) \leq \frac{\gamma_1 h(x^1) + V_\psi(x^*, x^1) + \frac{1}{2\sigma} \sum_{k=1}^N \gamma_k^2 \|\nabla f(x^k)\|_*^2}{\sum_{k=1}^N \gamma_k},$$

where $\tilde{x} = \frac{1}{\sum_{k=1}^N \gamma_k} \sum_{k=1}^N \gamma_k x^k$.

Case 1 (non-adaptive rule).

When $\gamma_k = \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (4.13), with $\tilde{x} = \frac{1}{\sum_{k=1}^N \frac{1}{\sqrt{k}}} \sum_{k=1}^N \frac{1}{\sqrt{k}} x^k$, we find

$$\begin{aligned} F(\tilde{x}) - F(x^*) &\leq \frac{M_f}{\sqrt{2\sigma}} \frac{\frac{\sqrt{2\sigma}}{M_f} h(x^1) + V_\psi(x^*, x^1) + \sum_{k=1}^N \frac{1}{k}}{\sum_{k=1}^N \frac{1}{\sqrt{k}}} \\ &\leq \frac{M_f}{\sqrt{2\sigma}} \frac{\frac{\sqrt{2\sigma}}{M_f} h(x^1) + V_\psi(x^*, x^1) + 1 + \log(N)}{2\sqrt{N+1} - 2} \\ &\leq \frac{M_f}{\sqrt{\sigma}} \frac{\frac{\sqrt{2\sigma}}{M_f} h(x^1) + V_\psi(x^*, x^1) + 1 + \log(N)}{\sqrt{N}}. \end{aligned}$$

Where in the last inequality, we used the fact $2\sqrt{2}(\sqrt{N+1}-1) \geq \sqrt{N}$, $\forall N \geq 1$.

Case 2 (adaptive rule). When $\gamma_k = \frac{\sqrt{2\sigma}}{\|\nabla f(x^k)\|_* \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (4.13), we find

$$\begin{aligned} F(\tilde{x}) - F(x^*) &\leq \frac{\frac{\sqrt{2\sigma}}{M_f} h(x^1) + V_\psi(x^*, x^1) + \sum_{k=1}^N \frac{1}{k}}{\sum_{k=1}^N \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}} \\ &\leq \frac{M_f \frac{\sqrt{2\sigma}}{M_f} h(x^1) + V_\psi(x^*, x^1) + 1 + \log(N)}{\sqrt{\sigma} \sqrt{N}}, \end{aligned}$$

where $\tilde{x} = \frac{1}{\sum_{k=1}^N (\|\nabla f(x^k)\|_* \sqrt{k})^{-1}} \sum_{k=1}^N \left(\|\nabla f(x^k)\|_* \sqrt{k} \right)^{-1} x^k$. \square

Also, from Theorem 4.3, with a special value of the parameter m , we can obtain the optimal convergence rate $O\left(\frac{1}{\sqrt{N}}\right)$ of Algorithm 2, with the time-varying step sizes given in (3.7).

Corollary 4.5. *Suppose that Assumption 4.1 hold and h is a non-negative function, with $h(x^1) = 0$. Then for problem (4.1), by Algorithm 2, with $m = 0$, and the time-varying step sizes given in (3.7), it satisfies the following optimal convergence rate*

$$F(\bar{x}) - F(x^*) \leq \left(\frac{M_f V_\psi(x^*, x^1) + 2M_f}{\sqrt{2\sigma}} \right) \frac{1}{\sqrt{N}} = O\left(\frac{1}{\sqrt{N}}\right),$$

where $\bar{x} = \frac{1}{N} \sum_{k=1}^N x^k$.

Proof. By setting $m = 0$ in (4.3), and since $h(x^1) = 0$, we get the following inequality

$$(4.14) \quad F(\bar{x}) - F(x^*) \leq \frac{1}{N} \left(\frac{V_\psi(x^*, x^1)}{\gamma_N} + \frac{1}{2\sigma} \sum_{k=1}^N \gamma_k \|\nabla f(x^k)\|_*^2 \right),$$

where $\bar{x} = \frac{1}{N} \sum_{k=1}^N x^k$.

Case 1 (non-adaptive rule). When $\gamma_k = \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (4.14) we find

$$\begin{aligned} F(\bar{x}) - F(x^*) &\leq \frac{1}{N} \left(\frac{M_f \sqrt{N} V_\psi(x^*, x^1)}{\sqrt{2\sigma}} + \frac{M_f}{\sqrt{2\sigma}} \sum_{k=1}^N \frac{1}{\sqrt{k}} \right) \\ &\leq \frac{1}{N} \frac{M_f}{\sqrt{2\sigma}} \left(\sqrt{N} V_\psi(x^*, x^1) + 2\sqrt{N} \right) \\ &= \frac{M_f (2 + V_\psi(x^*, x^1))}{\sqrt{2\sigma}} \frac{1}{\sqrt{N}}. \end{aligned}$$

Case 2 (adaptive rule). When $\gamma_k = \frac{\sqrt{2\sigma}}{\|\nabla f(x^k)\|_* \sqrt{k}}$, $k = 1, 2, \dots, N$, and since $\|\nabla f(x^k)\|_* \leq M_f$, then by substitution in (4.14) we find

$$\begin{aligned} F(\bar{x}) - F(x^*) &\leq \frac{1}{N} \left(\frac{M_f \sqrt{N} V_\psi(x^*, x^1)}{\sqrt{2\sigma}} + \frac{M_f}{\sqrt{2\sigma}} \sum_{k=1}^N \frac{1}{\sqrt{k}} \right) \\ &\leq \frac{M_f(2 + V_\psi(x^*, x^1))}{\sqrt{2\sigma}} \frac{1}{\sqrt{N}}. \end{aligned}$$

□

Remark 4.6. By setting any $m > -1$ in (4.3) with the assumption that $h(x^1) = 0$, the convergence rate of Algorithm 2 with the time-varying step sizes given in (3.7), will be optimal without the presence of a $\log(N)$ factor, namely $O\left(\frac{1}{\sqrt{N}}\right)$. In comparison with the sub-optimal convergence rate (4.12), when $m > 0$, the weighting scheme $\frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$ assigns smaller weights to the initial points and larger weights to the most recent points that generated by Algorithm 2.

5. MIRROR DESCENT METHOD FOR PROBLEMS WITH FUNCTIONAL CONSTRAINTS

In this section, we focus on a more general class of problems, which is a class of optimization problems with functional constraints (inequality constraints)

$$(5.1) \quad \min \{f(x) : x \in Q \text{ and } g_i(x) \leq 0 \text{ for all } i = 1, 2, \dots, p\},$$

where f and g_i (for all $i = 1, \dots, p$) are convex and non-smooth functions given on a closed convex set Q .

It is clear that instead of a set of functionals $\{g_i(\cdot)\}_{i=1}^p$ we can see one functional constraint $g : Q \rightarrow \mathbb{R}$, such that $g(x) = \max_{1 \leq i \leq p} \{g_i(x)\}$. Therefore, by this setting, the problem (5.1) will be equivalent to the following constrained minimization problem

$$(5.2) \quad \min_{x \in Q, g(x) \leq 0} f(x).$$

Let x^* be a solution of (5.2), we say that $\hat{x} \in Q$ is an ε -solution to (5.2) if $f(\hat{x}) - f(x^*) \leq \varepsilon$ and $g(\hat{x}) \leq \varepsilon$.

To solve problem (5.2), we propose Algorithm 3 which represents a generalization of Algorithm 1 to solve the general problem (5.2).

The needed point (output of Algorithm 3) will be selected among the points x^k for which $g(x^k) \leq \varepsilon$. Therefore, we will call step k productive if $g(x^k) \leq \varepsilon$. If the reverse inequality $g(x^k) \geq \varepsilon$ holds then step k will be called non-productive. Let I and J denote the set of productive and non-productive steps, respectively. $|I|$ and $|J|$ denote the number of productive

and non-productive steps, respectively. Let us also set $\gamma_k := \gamma_k^f$ if $k \in I$ and $\gamma_k := \gamma_k^g$ if $k \in J$.

Algorithm 3 Mirror descent method for problems with functional constraints.

Require: $\varepsilon > 0$, initial point $x^1 \in Q$, step sizes $\{\gamma_k^f\}_{k \geq 1}, \{\gamma_k^g\}_{k \geq 1}$, number of iterations N .

```

1:  $I \rightarrow \emptyset, J \rightarrow \emptyset$ .
2: for  $k = 1, 2, \dots, N$  do
3:   if  $g(x^k) \leq \varepsilon$  then
4:     Calculate  $\nabla f(x^k) \in \partial f(x^k)$ ,
5:      $x^{k+1} = \arg \min_{x \in Q} \left\{ \langle x, \nabla f(x^k) \rangle + \frac{1}{\gamma_k^f} V_\psi(x, x^k) \right\}$ .
6:      $k \rightarrow I$       "productive step"
7:   else
8:     Calculate  $\nabla g(x^k) \in \partial g(x^k)$ ,
9:      $x^{k+1} = \arg \min_{x \in Q} \left\{ \langle x, \nabla g(x^k) \rangle + \frac{1}{\gamma_k^g} V_\psi(x, x^k) \right\}$ .
10:     $k \rightarrow J$       "non-productive step"
11:   end if
12: end for
```

For Algorithm 3, we have the following result.

Theorem 5.1. *Let f and g are Lipschitz convex functions with constants $M_f > 0$ and $M_g > 0$, respectively. Assume that $\|\nabla f(x)\|_* \leq M_f$ and $\|\nabla g(x)\|_* \leq M_g$, for all $x \in Q$. Then for problem (5.2), with positive non-increasing sequences of step sizes $\{\gamma_k^f\}_{k \geq 1}, \{\gamma_k^g\}_{k \geq 1}$, for any fixed $m \geq -1$, and $\varepsilon > 0$, after N iterations of Algorithm 3, it satisfies the following inequality*

$$(5.3) \quad f(\hat{x}) - f(x^*) < \frac{1}{\sum_{k \in I} (\gamma_k^f)^{-m}} \left(\frac{V_\psi(x^*, x^1)}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k \in I} \frac{\|\nabla f(x^k)\|_*^2}{(\gamma_k^f)^{m-1}} \right. \\ \left. + \frac{1}{2\sigma} \sum_{k \in J} \frac{\|\nabla g(x^k)\|_*^2}{(\gamma_k^g)^{m-1}} - \varepsilon \sum_{k \in J} (\gamma_k^g)^{-m} \right),$$

where $\hat{x} = \frac{1}{\sum_{k \in I} (\gamma_k^f)^{-m}} \sum_{k \in I} (\gamma_k^f)^{-m} x^k$ and x^* is an optimal solution of (5.2).

Proof. Similar to what was done in proving Theorem 3.1, we find that for all $k \in I$,

$$(5.4) \quad \frac{f(x^k) - f(x^*)}{(\gamma_k^f)^m} \leq \frac{V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1})}{(\gamma_k^f)^{m+1}} + \frac{\|\nabla f(x^k)\|_*^2}{2\sigma (\gamma_k^f)^{m-1}},$$

and for all $k \in J$,

$$(5.5) \quad \frac{g(x^k) - g(x^*)}{(\gamma_k^g)^m} \leq \frac{V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1})}{(\gamma_k^g)^{m+1}} + \frac{\|\nabla g(x^k)\|_*^2}{2\sigma (\gamma_k^g)^{m-1}},$$

By taking the summation, in each side of (5.4) and (5.5), over productive and non-productive steps, with $\gamma_k = \gamma_k^f$ if $k \in I$ and $\gamma_k = \gamma_k^g$ if $k \in J$, we get

$$\begin{aligned} & \sum_{k \in I} (\gamma_k^f)^{-m} (f(x^k) - f(x^*)) + \sum_{k \in J} (\gamma_k^g)^{-m} (g(x^k) - g(x^*)) \\ & \leq \sum_{k=1}^N \frac{1}{\gamma_k^{m+1}} (V_\psi(x^*, x^k) - V_\psi(x^*, x^{k+1})) + \frac{1}{2\sigma} \sum_{k \in I} \frac{\|\nabla f(x^k)\|_*^2}{(\gamma_k^f)^{m-1}} \\ & \quad + \frac{1}{2\sigma} \sum_{k \in J} \frac{\|\nabla g(x^k)\|_*^2}{(\gamma_k^g)^{m-1}}. \end{aligned}$$

Also, in a similar way as in the proof of Theorem 3.1, and since for any $k \in J$, we have

$$(5.6) \quad g(x^k) - g(x^*) \geq g(x^k) > \varepsilon > 0.$$

Therefore we get

$$\begin{aligned} \sum_{k \in I} (\gamma_k^f)^{-m} (f(x^k) - f(x^*)) & < \frac{1}{\gamma_N^{m+1}} V_\psi(x^*, x^1) + \frac{1}{2\sigma} \sum_{k \in I} \frac{\|\nabla f(x^k)\|_*^2}{(\gamma_k^f)^{m-1}} \\ & \quad + \frac{1}{2\sigma} \sum_{k \in J} \frac{\|\nabla g(x^k)\|_*^2}{(\gamma_k^g)^{m-1}} - \varepsilon \sum_{k \in J} (\gamma_k^g)^{-m}. \end{aligned}$$

Since f is a convex function, we have

$$\begin{aligned} \left(\sum_{k \in I} (\gamma_k^f)^{-m} \right) \left[f \left(\frac{1}{\sum_{k \in I} (\gamma_k^f)^{-m}} \sum_{k \in I} (\gamma_k^f)^{-m} x^k \right) - f(x^*) \right] & \leq \\ & \leq \sum_{k \in I} (\gamma_k^f)^{-m} (f(x^k) - f(x^*)). \end{aligned}$$

Therefore, by setting $\hat{x} := \frac{1}{\sum_{k \in I} (\gamma_k^f)^{-m}} \sum_{k \in I} (\gamma_k^f)^{-m} x^k$, we get

$$\begin{aligned} (5.7) \quad \left(\sum_{k \in I} (\gamma_k^f)^{-m} \right) (f(\hat{x}) - f(x^*)) & < \frac{1}{\gamma_N^{m+1}} V_\psi(x^*, x^1) + \frac{1}{2\sigma} \sum_{k \in I} \frac{\|\nabla f(x^k)\|_*^2}{(\gamma_k^f)^{m-1}} \\ & \quad + \frac{1}{2\sigma} \sum_{k \in J} \frac{\|\nabla g(x^k)\|_*^2}{(\gamma_k^g)^{m-1}} - \varepsilon \sum_{k \in J} (\gamma_k^g)^{-m}. \end{aligned}$$

By dividing both sides of the previous inequality by $\sum_{k \in I} (\gamma_k^f)^{-m}$, we get the desired inequality (5.3).

Finally, let us show that $|I| \neq 0$. For this let us assume that $|I| = 0$, therefore $|J| = N$, i.e., all steps are non-productive. From (5.6), we get

$$\sum_{k=1}^N \frac{g(x^k) - g(x^*)}{\gamma_k^m} > \sum_{k=1}^N \frac{\varepsilon}{\gamma_k^m} = \frac{\varepsilon M^m}{(\sqrt{2}\sigma)^m} \sum_{k=1}^N (\sqrt{k})^m,$$

and for all $k \in J = \{1, \dots, N\}$, with the assumption that $V_\psi(x^*, x^1) \leq \theta$ for some $\theta > 0$, we get

$$\begin{aligned} \sum_{k=1}^N \frac{g(x^k) - g(x^*)}{\gamma_k^m} &\leq \frac{\theta}{\gamma_N^{m+1}} + \frac{1}{2\sigma} \sum_{k=1}^N \frac{\|\nabla g(x^k)\|_*^2}{\gamma_k^{m-1}} \\ &\leq \frac{M^{m+1}}{(\sqrt{2}\sigma)^{m+1}} \left(\theta (\sqrt{N})^{m+1} + \sum_{k=1}^N (\sqrt{k})^{m-1} \right). \end{aligned}$$

But, one can numerically verify that for a sufficiently big number of iterations N (dependently on suitable values of the parameters $\theta > 0, m \geq -1, M > 0, \varepsilon > 0, \sigma > 0$), the following inequality holds

$$(5.8) \quad \frac{M^{m+1}}{(\sqrt{2}\sigma)^{m+1}} \left(\theta (\sqrt{N})^{m+1} + \sum_{k=1}^N (\sqrt{k})^{m-1} \right) < \frac{\varepsilon M^m}{(\sqrt{2}\sigma)^m} \sum_{k=1}^N (\sqrt{k})^m.$$

So, we have a contradiction. It means that $|I| \neq 0$. \square

Remark 5.2. Note that the reverse inequality of (5.8), i.e.,

$$\sum_{k=1}^N (\sqrt{k})^m \leq \frac{M}{\varepsilon \sqrt{2}\sigma} \left(\theta (\sqrt{N})^{m+1} + \sum_{k=1}^N (\sqrt{k})^{m-1} \right),$$

for any $m \geq -1, M > 0, \theta > 0, \sigma > 0$ and $\varepsilon \leq \frac{M}{\sqrt{2}\sigma}$, holds for at least $N = 1$. This means that by choosing $\varepsilon \leq \frac{M}{\sqrt{2}\sigma}$ ($\forall M > 0, \sigma > 0$), we have at least one productive step for any $m \geq -1$ and $\theta > 0$.

Remark 5.3. From inequality (5.7) (which is equivalent to the inequality (5.3)), with $\gamma_q = \gamma_q^f$ if $q \in I$ and $\gamma_q = \gamma_q^g$ if $q \in J$, with

$$(5.9) \quad \hat{x} = \frac{1}{\sum_{i \in I} (\gamma_i^f)^{-m}} \sum_{i \in I} (\gamma_i^f)^{-m} x^i,$$

for any $k \geq 1$, we find

$$\begin{aligned}
\left(\sum_{i \in I} (\gamma_i^f)^{-m} \right) (f(\hat{x}) - f(x^*)) &< \frac{1}{\gamma_k^{m+1}} V_\psi(x^*, x^1) + \frac{1}{2\sigma} \sum_{i \in I} \frac{\|\nabla f(x^i)\|_*^2}{(\gamma_i^f)^{m-1}} \\
&+ \frac{1}{2\sigma} \sum_{j \in J} \frac{\|\nabla g(x^j)\|_*^2}{(\gamma_j^g)^{m-1}} - \varepsilon \sum_{i=1}^k (\gamma_i)^{-m} + \varepsilon \sum_{i \in I} (\gamma_i^f)^{-m} \\
&= \varepsilon \sum_{i \in I} (\gamma_i^f)^{-m} - \left(\varepsilon \sum_{i=1}^k (\gamma_i)^{-m} - \frac{V_\psi(x^*, x^1)}{\gamma_k^{m+1}} - \frac{1}{2\sigma} \sum_{i \in I} \frac{\|\nabla f(x^i)\|_*^2}{(\gamma_i^f)^{m-1}} \right. \\
&\quad \left. - \frac{1}{2\sigma} \sum_{j \in J} \frac{\|\nabla g(x^j)\|_*^2}{(\gamma_j^g)^{m-1}} \right).
\end{aligned}$$

From this, without relying on prior knowledge of the number of iterations that the algorithm performs, we can set for any $k \geq 1$,

$$(5.10) \quad \varepsilon \sum_{i=1}^k (\gamma_i)^{-m} \geq \frac{V_\psi(x^*, x^1)}{\gamma_k^{m+1}} + \frac{1}{2\sigma} \sum_{i \in I} \frac{\|\nabla f(x^i)\|_*^2}{(\gamma_i^f)^{m-1}} + \frac{1}{2\sigma} \sum_{j \in J} \frac{\|\nabla g(x^j)\|_*^2}{(\gamma_j^g)^{m-1}},$$

as a stopping criterion of Algorithm 3 and as a result we conclude

$$\left(\sum_{i \in I} (\gamma_i^f)^{-m} \right) (f(\hat{x}) - f(x^*)) < \varepsilon \sum_{i \in I} (\gamma_i^f)^{-m},$$

i.e., $f(\hat{x}) - f(x^*) \leq \varepsilon$.

For all $i \in I$ it holds that $g(x^i) \leq \varepsilon$, and since g is convex, then we have

$$g(\hat{x}) \leq \frac{1}{\sum_{i \in I} (\gamma_i^f)^{-m}} \sum_{i \in I} (\gamma_i^f)^{-m} g(x^i) \leq \varepsilon.$$

Thus after the stop criterion (5.10) is met we find that \hat{x} given in (5.9) represents an ε -solution to problem (5.2).

Now let us analyze the convergence of Algorithm 3, by taking the following time-varying step size rules

$$(5.11) \quad \gamma_k = \begin{cases} \gamma_k^f := \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}, & \text{or } \gamma_k^f := \frac{\sqrt{2\sigma}}{\|\nabla f(x^k)\|_* \sqrt{k}}; & \text{if } k \in I, \\ \gamma_k^g := \frac{\sqrt{2\sigma}}{M_g \sqrt{k}}, & \text{or } \gamma_k^g := \frac{\sqrt{2\sigma}}{\|\nabla g(x^k)\|_* \sqrt{k}}; & \text{if } k \in J. \end{cases}$$

Let us assume that $V_\psi(x^*, x^1) \leq \theta$, for some $\theta > 0$, $M := \max\{M_f, M_g\}$. By using the non-adaptive rules from (5.11) (we also can conclude the same results if we take the adaptive step size rules), and since $\|\nabla f(x^k)\|_* \leq M_f \leq M$ and $\|\nabla g(x^k)\|_* \leq M_g \leq M$, then for any $m \geq 1$, from Theorem 5.1 we

have

$$\begin{aligned}
f(\hat{x}) - f(x^*) &< \frac{(\sqrt{2}\sigma)^m}{M^m \sum_{k \in I} (\sqrt{k})^m} \left(\frac{\theta M^{m+1} (\sqrt{N})^{m+1}}{(\sqrt{2}\sigma)^{m+1}} + \right. \\
&\quad \left. + \frac{1}{2\sigma} \sum_{k=1}^N \frac{M^{m+1} (\sqrt{k})^{m-1}}{(\sqrt{2}\sigma)^{m-1}} \right) \\
&= \frac{M}{\sqrt{2}\sigma} \cdot \frac{1}{\sum_{k \in I} (\sqrt{k})^m} \cdot \left(\theta (\sqrt{N})^{m+1} + \sum_{k=1}^N (\sqrt{k})^{m-1} \right) \\
&\leq \frac{M}{\sqrt{2}\sigma} \cdot \frac{1}{\sum_{k \in I} (\sqrt{k})^m} \cdot \left(\theta (\sqrt{N})^{m+1} + N (\sqrt{N})^{m-1} \right) \\
&= \frac{M(1+\theta) (\sqrt{N})^{m+1}}{\sqrt{2}\sigma} \cdot \frac{1}{\sum_{k \in I} (\sqrt{k})^m}.
\end{aligned}$$

Now, by setting $\frac{M(1+\theta)(\sqrt{N})^{m+1}}{\sqrt{2}\sigma} \cdot \frac{1}{\sum_{k \in I} (\sqrt{k})^m} \leq \varepsilon$ and since $|I| \leq N$, we get

$$\frac{M(1+\theta) (\sqrt{N})^{m+1}}{\sqrt{2}\sigma N (\sqrt{N})^m} \leq \frac{M(1+\theta) (\sqrt{N})^{m+1}}{\sqrt{2}\sigma} \cdot \frac{1}{\sum_{k \in I} (\sqrt{k})^m} \leq \varepsilon.$$

Thus,

$$\frac{M(1+\theta)}{\sqrt{N}\sqrt{2}\sigma} \leq \varepsilon \implies N \geq \frac{M^2(1+\theta)^2}{2\sigma\varepsilon^2}.$$

Hence, we can formulate the following result.

Corollary 5.4. *Let f and g are Lipschitz convex functions with constants $M_f > 0$ and $M_g > 0$, respectively. Assume that $\|\nabla f(x)\|_* \leq M_f$ and $\|\nabla g(x)\|_* \leq M_g$, for all $x \in Q$, and $V_\psi(x^*, x^1) \leq \theta$, for some $\theta > 0$, and let $\varepsilon > 0$, $M := \max\{M_f, M_g\}$. Then after*

$$(5.12) \quad N = \left\lceil \frac{M^2(1+\theta)^2}{2\sigma\varepsilon^2} \right\rceil = O\left(\frac{1}{\varepsilon^2}\right),$$

iterations of Algorithm 3, for any fixed $m \geq 1$, with step size rules given in (5.11), it satisfies

$$f(\hat{x}) - f(x^*) < \varepsilon, \quad \text{and} \quad g(\hat{x}) \leq \varepsilon,$$

where $\hat{x} = \frac{1}{\sum_{k \in I} (\gamma_k^f)^{-m}} \sum_{k \in I} (\gamma_k^f)^{-m} x^k$.

The estimate (5.12), is optimal for the class of non-smooth optimization problems under consideration.

Now, by setting $m = 0$ in (5.3), with $\bar{x} = \frac{1}{|I|} \sum_{k \in I} x^k$, we get

$$\begin{aligned} f(\bar{x}) - f(x^*) &< \frac{1}{|I|} \left(\frac{V_\psi(x^*, x^1)}{\gamma_N} + \sum_{k \in I} \frac{\|\nabla f(x^k)\|_*^2 \gamma_k^f}{2\sigma} + \sum_{k \in J} \frac{\|\nabla g(x^k)\|_*^2 \gamma_k^g}{2\sigma} \right), \\ &\leq \frac{1}{|I|} \left(\frac{M\theta\sqrt{N}}{\sqrt{2\sigma}} + \frac{M}{\sqrt{2\sigma}} \sum_{k \in I} \frac{1}{\sqrt{k}} + \frac{M}{\sqrt{2\sigma}} \sum_{k \in J} \frac{1}{\sqrt{k}} \right) \\ &= \frac{M}{|I|\sqrt{2\sigma}} \left(\theta\sqrt{N} + \sum_{k=1}^N \frac{1}{\sqrt{k}} \right) \leq \frac{M}{|I|\sqrt{2\sigma}} (\theta\sqrt{N} + 2\sqrt{N}) \\ &= \frac{M\sqrt{N}(2+\theta)}{|I|\sqrt{2\sigma}}. \end{aligned}$$

Now, by setting $\frac{M\sqrt{N}(2+\theta)}{|I|\sqrt{2\sigma}} \leq \varepsilon$ and since $|I| \leq N$, we get

$$\frac{M(2+\theta)}{\sqrt{N}\sqrt{2\sigma}} \leq \varepsilon \implies N \geq \frac{M^2(2+\theta)^2}{2\sigma\varepsilon^2}.$$

Hence, for $m = 0$, we can formulate the following result.

Corollary 5.5. *Let f and g are Lipschitz convex functions with constants $M_f > 0$ and $M_g > 0$, respectively. Assume that $\|\nabla f(x)\|_* \leq M_f$ and $\|\nabla g(x)\|_* \leq M_g$, for all $x \in Q$, and $V_\psi(x^*, x^1) \leq \theta$, for some $\theta > 0$, and let $\varepsilon > 0, M := \max\{M_f, M_g\}$. Then after*

$$N = \left\lceil \frac{M^2(2+\theta)^2}{2\sigma\varepsilon^2} \right\rceil = O\left(\frac{1}{\varepsilon^2}\right),$$

iterations of Algorithm 3, with $m = 0$, and step size rules given in (5.11), it satisfies

$$f(\bar{x}) - f(x^*) < \varepsilon, \quad \text{and} \quad g(\bar{x}) \leq \varepsilon,$$

where $\bar{x} = \frac{1}{|I|} \sum_{k \in I} x^k$.

Remark 5.6. By setting $m = -1$ in (5.3), we deduce the sub-optimal convergence rate for Algorithm 3. In this case, with $\tilde{x} = \frac{1}{\sum_{k \in I} \frac{1}{\sqrt{k}}} \sum_{k \in I} \frac{1}{\sqrt{k}} x^k$, we have

$$\begin{aligned} f(\tilde{x}) - f(x^*) &\leq \frac{M}{\sqrt{2\sigma}} \cdot \frac{1}{\sum_{k \in I} \sqrt{k}} \cdot \left(\theta + \sum_{k=1}^N \frac{1}{k} \right) \\ &\leq \frac{M}{\sqrt{2\sigma}} \cdot \frac{1}{2\sqrt{|I|+1}-2} \cdot (\theta + 1 + \log(N)) \\ &\leq \frac{M(\theta + 1 + \log(N))}{\sqrt{\sigma}\sqrt{|I|}}. \end{aligned}$$

The presence of the logarithm in the last inequalities is what causes the sub-optimality in the convergence of algorithm 3 for $m = -1$.

Remark 5.7. For solving problem (5.1) (or its equivalence (5.2)), when we have a very big number of functional constraints, we can modify Algorithm 3 as in [32]. When we have a non-productive step k , i.e., $g(x^k) > \varepsilon$, then instead of calculating a (sub)gradient of the functional constraint with max-type $g(x) = \max_{1 \leq i \leq p} \{g_i(x^k)\}$, we calculate a (sub)gradient of one functional g_i , for which we have $g_i(x^k) > \varepsilon$. This idea of the proposed modification of Algorithms 3 allows saving the running time of the algorithm due to consideration of not all functional constraints on non-productive steps.

6. NUMERICAL EXPERIMENTS

To show the advantages of the proposed algorithms and effects of the weighting scheme for generated points \hat{x} in Theorem 3.1 (and the same points in Theorem 4.3), in the convergence of the studied algorithms, a series of numerical experiments were performed for some problems with a geometrical nature in addition to the problem of minimizing the maximum of a finite collection of linear functions.

We compare the performance of the studied algorithms with the projected subgradient method using different famous step size rules that are listed in Table 1 (which is inspired from [16]). Therefore, in our experiments, we take the standard Euclidean prox-structure, namely $\psi(x) = \frac{1}{2}\|x\|_2^2$ which is 1-strongly functions (i.e., $\sigma = 1$) and the corresponding Bergman divergence is $V_\psi(x, y) = \frac{1}{2}\|x - y\|_2^2$. In all experiments, we take the set Q as a unit ball in \mathbb{R}^n with the center at $0 \in \mathbb{R}^n$. All compared methods start from the same initial point $x^1 = \left(\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}\right) \in Q \subset \mathbb{R}^n$. In the AdaGrad algorithm, there is an assumption that $\|x^1 - x^*\|_2^2 \leq 2\theta_0^2$, thus for the taken feasible set in our experiments, we can take $\theta_0 = \sqrt{2}$.

The comparison of the methods is done in terms of the difference $\hat{f}_k - f_{\min}$, where \hat{f}_k denotes the value of the objective function f at the averaged points (namely at $\hat{x}_k = \frac{1}{k} \sum_{j=1}^k x^j$, for all cases in Table 1, except to the case of "quad grad", where in this case we have $\hat{x}_k = \left(\sum_{j=1}^k \gamma_j\right)^{-1} \sum_{j=1}^k \gamma_j x^j$) and f_{\min} denotes the minimal value of the objective function computed by SciPy, a package for solving many different classes of optimization problems (when the dimension of the space \mathbb{R}^n is not big (for example when $n \leq 200$)). Also, we did a comparison of the methods when \hat{f}_k denotes the best value of $f(x^k)$ attained, i.e., $\hat{f}_k = \min\{f(x^1), \dots, f(x^k)\}$, see Remarks 2 and 5.

6.1. Best approximation problem. The considered problem in this subsection is connected with the problem of the best approximation of the distance between a point and a given set Q . For this problem, let $A \notin Q$ be

Abbreviation	Step sizes	formula of γ_k
constant step	constant step size [11]	$\gamma_k = 0.1$
fixed length	fixed step length [11]	$\gamma_k = \frac{0.2}{\ \nabla f(x^k)\ _*}$
nonsum	non-summable diminishing step [11]	$\gamma_k = \frac{0.1}{\sqrt{k}}$
sqrsum nonsum	square summable but not summable step [11]	$\gamma_k = \frac{0.5}{k}$
quad grad	quadratic of the norm of the gradient [36]	$\gamma_k = \frac{0.2}{\ \nabla f(x^k)\ _*^2}$
AdaGrad	AdaGrad algorithm [15]	$\gamma_k = \frac{\theta_0}{\sqrt{\sum_{j=1}^k \ \nabla f(x^j)\ _*^2 + \alpha}}$
Polyak step	Polyak step size [29]	$\gamma_k = \frac{f(x^k) - f^*}{\ \nabla f(x^k)\ _*^2}$
new non-adaptive	this paper (3.7)	$\gamma_k = \frac{\sqrt{2\sigma}}{M_f \sqrt{k}}$
new adaptive	this paper (3.7)	$\gamma_k = \frac{\sqrt{2\sigma}}{\ \nabla f(x^k)\ _* \sqrt{k}}$

Table 1. The used step sizes in the compared methods.

a given point, we need to solve the following optimization problem

$$(6.1) \quad \min_{x \in Q} \{f(x) := \|x - A\|_2\}.$$

The point A is randomly generated from a uniform distribution over $[0, 1)$, such that $\|A\|_2 = 10$, therefore the distance between the point A and the considered unit ball Q is equal to 9, i.e., $f^* = 9$. Here, we mention that this problem is constructed so that it can be used the Polyak step size, which requires knowing the optimal value f^* . The results of the comparison, for problem (6.1) with $n = 1000$ are presented in Fig. 1. In this figure \hat{f}^k denotes the value of the objective function f at the averaged points in each iteration of all compared algorithms. From this figure, we can see that Algorithm 1, with non-adaptive and adaptive step size rules (3.7), works better than other algorithms, where the difference between the performance of Algorithm 1 and the rest of algorithms with steps in Table 1 is clear and significant. We also note that there is no significant difference in the performance of the Algorithm 1 when using the adaptive or non-adaptive step size rules steps

Remark 6.1. If we take $\hat{f}_k = \min\{f(x^1), \dots, f(x^k)\}$ for all algorithms with steps in Table 1, instead of the value of the objective function at the averaged points in each iteration, then we also can see that that Algorithm 1 still works better. Although the work of other algorithms will improve and be better than the results in Figure 1.

Remark 6.2. As we mentioned in Remark 3.4, in comparison with the sub-optimal convergence rate (3.8), when $m > 0$, the weighting scheme




Figure 1. Results of Algorithm 1 for problem (6.1) with $n = 1000$.




Figure 2. Results of Algorithm 1 for problem (6.1) with $n = 1000$. Here $\hat{f}_k = \min\{f(x^1), \dots, f(x^k)\}$.

$\frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$ (see Theorem 3.1) assigns smaller weights to the initial points and larger weights to the most recent points that generated by Algorithm 1. In Fig. 3, we can see this fact, where when we increase the value of the parameter m , we can see that convergence of Algorithm 1 and its performance becomes significantly better.




Figure 3. Results of Algorithm 1 for problem (6.1) with $n = 1000$ with different values of the parameter m .

6.2. Fermat–Torricelli–Steiner problem. Let $A_j \in \mathbb{R}^n, j = 1, \dots, T$ be a given set of T points, and let us consider an analogue of the well-known Fermat–Torricelli–Steiner problem. For this we need to solve the following optimization problem

$$(6.2) \quad \min_{x \in Q} \left\{ f(x) := \frac{1}{T} \sum_{j=1}^T \|x - A_j\|_2 \right\}.$$

The points $A_j, j = 1, 2, \dots, T$ are randomly generated from a uniform distribution over $[0, 1]$. We run all algorithms (except the algorithm with Polyak step size since we cannot know the optimal value f^* for problem (6.2)), with the same initial point $x^1 = \left(\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}} \right) \in Q$.

The results of the comparison, for problem (6.2) with $n = 200$ and $T = 25$, are presented in Fig. 4. The reason here for taking $n = 200, T = 25$ is that we calculated the value f_{\min} using SciPy which does not well work for large values. In Fig. 4, \hat{f}^k denotes the value of the objective function f at the

averaged points in each iteration of all compared algorithms. From this figure, we can see that Algorithm 1, with non-adaptive and adaptive step size rules (3.7), outperforms the other methods providing a better solution.




Figure 4. Results of Algorithm 1 for problem (6.2) with $n = 200, T = 25$.

Remark 6.3. As in Remark 6.1. If we take $\hat{f}_k = \min\{f(x^1), \dots, f(x^k)\}$ for all algorithms with steps in Table 1, instead of the value of the objective function at the averaged points in each iteration, then we also can see that that Algorithm 1 still works better at the first iterations and after that, it works the same as other algorithms (except algorithm with "nonsum" and "sqsum nonsum").

Remark 6.4. As in Remark 6.2. We can see that when we increase the value of the parameter m in the weighting scheme $\frac{1}{\sum_{k=1}^N \gamma_k^{-m}} \sum_{k=1}^N \gamma_k^{-m} x^k$ (see Theorem 3.1) the convergence of Algorithm 1 and its performance becomes better.

Remark 6.5. Because of SciPy does not work well for large values of n and T , we conduct experiments without calculating f_{\min} . We run algorithms with $n = 1000$ and $T = 100$. The results are presented in Fig. 7. In this figure, we show the dynamics of \hat{f}_k (the value of the objective function at the averaged points in each iteration) as a function of k . From Fig. 7, as in the previous results, we also see that Algorithm 1, with non-adaptive and adaptive step size rules (3.7), outperforms the other methods providing a




Figure 5. Results of Algorithm 1 for problem (6.2) with $n = 200, T = 25$. $\hat{f}_k = \min\{f(x^1), \dots, f(x^k)\}$.




Figure 6. Results of Algorithm 1 for problem (6.2) with $n = 200, T = 25$ and different values of the parameter m .

better solution. Note that in the performance of the Algorithm 1, there is a difference if we choose adaptive or non-adaptive step sizes.




Figure 7. Results of Algorithm 1 for problem (6.2) with $n = 1000, T = 100$.

6.3. Smallest covering ball problem. Let $A_j \in \mathbb{R}^n, j = 1, \dots, T$ be a given set of T points, and let us consider an analogue of the well-known smallest covering ball problem. For this we need to solve the following optimization problem

$$(6.3) \quad \min_{x \in Q} \left\{ f(x) := \max_{1 \leq j \leq T} \|x - A_j\|_2 \right\}.$$

The points $A_j, j = 1, 2, \dots, T$ are randomly generated from a uniform distribution over $[0, 1)$. We run all algorithms (except the algorithm with Polyak step size since we cannot know the optimal value f^* for problem (6.2)), with the same initial point $x^1 = \left(\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}} \right) \in Q$.

The results of the comparison, for problem (6.3) with $n = 200$ and $T = 25$, are presented in Fig. 4 (in the left, where we show the dynamics of $\hat{f}_k - f_{\min}$ as a function of k , and f_{\min} calculated by SciPy), and with $n = 1000$ and $T = 100$ (in the right, where we show the dynamics of \hat{f}_k as a function of k).

From Fig. 8, we also see that Algorithm 1, with non-adaptive and adaptive step size rules (3.7), outperforms the other methods providing a better solution, without any difference if we choose adaptive or non-adaptive step sizes.




Figure 8. Results of Algorithm 1 for problem (6.3) with $n = 200, T = 25$ (in the left). And with $n = 1000, T = 100$ (in the right).

6.4. Maximum of a finite collection of linear functions. In this subsection, we consider the minimization problem of a function that represents a point-wise maximum of a finite collection of linear functions. This problem has the following form:

$$(6.4) \quad \min_Q f(x) := \max \{f_i(x) = \langle a_i, x \rangle + b_i, i = 1, 2, \dots, T\},$$

where $a_i \in \mathbb{R}^n$, and $b_i \in \mathbb{R}$ ($\forall i = 1, \dots, T$).

The vectors a_j and constants b_j , for $j = 1, 2, \dots, T$, are randomly generated from a uniform distribution over $[0, 1]$. We run all algorithms (except the algorithm with Polyak step size since we cannot know the optimal value f^* for problem (6.2)), with the same initial point $x^1 = \left(\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}\right) \in Q$.

The results of the comparison, for problem (6.3) with $n = 200$ and $T = 25$, are presented in Fig. 9 (in the left, where we show the dynamics of $\hat{f}_k - f_{\min}$ as a function of k , and f_{\min} calculated by SciPy), and with $n = 1000$ and $T = 100$ (in the right, where we show the dynamics of \hat{f}_k as a function of k).

From Fig. 9, we also see that Algorithm 1, with non-adaptive and adaptive step size rules (3.7), outperforms the other methods providing a better solution, without any difference if we choose adaptive or non-adaptive step sizes.

6.5. Problems with functional constraints. In this subsection, we will show the advantages of the proposed Algorithm 3 with adaptive step size rules given in (5.11), and compare its work with algorithms 1 and 2 from [3], which were proposed to solve the problem (5.1). We will consider the problems (6.1), (6.2), (6.3) and (6.4), with the following functional constraints

$$(6.5) \quad g(x) = \max \{g_i(x) = \langle \alpha_i, x \rangle - \beta_i, \quad i = 1, \dots, p\},$$

where $\alpha_i \in \mathbb{R}^n$ and $\beta_i \in \mathbb{R}$ are randomly generated from a uniform distribution over $[0, 1]$.




Figure 9. Results of Algorithm 1 for problem (6.4) with $n = 200, T = 25$ (in the left). And with $n = 1000, T = 100$ (in the right).

As in the previous, we take Q as a unit ball in \mathbb{R}^n with the center at $0 \in \mathbb{R}^n$. All compared methods start from the same initial point $x^1 = 0 \in Q$. We take $n = 1000, p = 100$ functional constraints in (6.5) and $\varepsilon = 10^{-3}$.

The results of the comparison are presented in Fig. 10 and Fig. 11. In these figures, we show the values of the objective function in each problem at the points that correspond to the set of productive steps generated by the compared algorithms.

From Figures 10 and 11, we can see how the proposed Algorithm 3 is the best, and we note that the work of this algorithm will be better if we increase the value of the parameter m .




Figure 10. Results of Algorithm 3 and Algorithms 1, 2 in [3], for problem (6.1) (in the left) and problem (6.2) (in the right), with functional constraints (6.5), and $n = 1000, p = 100, T = 100$.

REFERENCES

- [1] M. S. Alkousa: On Modification of an Adaptive Stochastic Mirror Descent Algorithm for Convex Optimization Problems with Functional Constraints. Forum for Interdisciplinary Mathematics. Springer, Singapore. pp. 47–63.
- [2] M. S. Alkousa: On Some Stochastic Mirror Descent Methods for Constrained Online Optimization Problems. Computer Research and Modeling, 11(2), pp. 205–217, 2019.




Figure 11. Results of Algorithm 3 and Algorithms 1, 2 in [3], for problem (6.3) (in the left) and problem (6.4) (in the right), with functional constraints (6.5), and $n = 1000, p = 100, T = 100$.

- [3] A. Bayandina, P. Dvurechensky, A. Gasnikov, F. Stonyakin, A. Titov: Mirror descent and convex optimization problems with non-smooth inequality constraints. In: Large-Scale and Distributed Optimization, Springer, Cham pp. 181–213, 2018.
- [4] A. Beck: First-order methods in optimization. Society for Industrial and Applied Mathematics, 2017.
- [5] A. Beck, M. Teboulle: Mirror descent and nonlinear projected subgradient methods for convex optimization. *Oper. Res. Lett.*, 31(3), pp. 167–175, 2003.
- [6] A. Beck, A. Ben-Tal, N. Guttman-Beck, L. Tetruashvili: The comirror algorithm for solving nonsmooth constrained convex problems. *Operations Research Letters*, **38**(6), pp. 493–498, 2010.
- [7] A. Ben-Tal, A. Nemirovski: Lectures on Modern Convex Optimization. Society for Industrial and Applied Mathematics, Philadelphia 2001.
- [8] A. Ben-Tal, T. Margalit, A. Nemirovski: The Ordered Subsets Mirror Descent Optimization Method with Applications to Tomography. *SIAM Journal on Optimization*, **12**(1), pp. 79–108, 2001.
- [9] A. Ben-Tal, A. Nemirovski: Robust truss topology design via semidefinite programming. *SIAM J. Optim.* 7(4), 991–1016 (1997)
- [10] D. P. Bertsekas: Constrained optimization and Lagrange multiplier methods. Academic press, 2014.
- [11] S. Boyd, L. Xiao, A. Mutapcic: Subgradient methods. lecture notes of EE392o, Stanford University, Autumn Quarter, 2004(01).
- [12] S. Bubeck: Convex optimization: algorithms and complexity. *Foundations and Trends in Machine Learning*, **8**(3–4), pp. 231–357, 2015. <https://arxiv.org/pdf/1405.4980.pdf>
- [13] G. Chen, M. Teboulle: Convergence analysis of a proximal like minimization algorithm using Bregman functions, *SIAM J. Optim.* 3 (1993) 538–543.
- [14] T. T. Doan, S. Bose, D. H. Nguyen, C. L. Beck: Convergence of the Iterates in Mirror Descent Methods. *IEEE Control Systems Letters*, **3**(1), pp. 114–119, 2019.
- [15] J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12(Jul.):2121–2159, 2011.
- [16] O. P. Ferreira, G. N. Grapiglia, E. M. Santos, J. C. O. Souza: A subgradient method with non-monotone line search. *Computational Optimization and Applications*, 84(2), 397–420 (2023).

- [17] O. Fercoq, A. Alacaoglu, I. Necoara, V. Cevher: Almost surely constrained convex optimization. Proceedings of the 36th International Conference on Machine Learning, PMLR 97, pp. 1910–1919, 2019.
- [18] A. B. Juditsky, A. V. Nazin, A. B. Tsybakov, N. Vayatis: Recursive Aggregation of Estimators by the Mirror Descent Algorithm with Averaging. Problems of Information Transmission, **41**(4), pp. 368–384, 2005.
- [19] G. Lan, Z. Zhou: Algorithms for stochastic optimization with functional or expectation constraints. Comput Optim Appl 76, 461–498 (2020).
- [20] G. Lan: First-order and Stochastic Optimization Methods for Machine Learning. Springer Nature Switzerland AG 2020
- [21] Q. Lin, R. Ma, T. Yang: Level-set methods for finite-sum constrained convex optimization. In International Conference on Machine Learning, pp. 3118–3127, 2018.
- [22] D. V. N. Luong, P. Parpas, D. Rueckert, B. Rustem: A Weighted Mirror Descent Algorithm for Nonsmooth Convex Optimization Problem. J Optim Theory Appl **170**(3), pp. 900–915, 2016.
- [23] A. Nemirovskii: Efficient methods for large-scale convex optimization problems. Ekonomika i Matematicheskie Metody, 1979. (in Russian)
- [24] A. Nemirovsky, D. Yudin: Problem Complexity and Method Efficiency in Optimization. J. Wiley & Sons, New York 1983.
- [25] Yu. Nesterov: Lectures on convex optimization. Switzerland: Springer Optimization and Its Applications, 2018.
- [26] A. V. Nazin, B. M. Miller: Mirror Descent Algorithm for Homogeneous Finite Controlled Markov Chains with Unknown Mean Losses. Proceedings of the 18th World Congress The International Federation of Automatic Control Milano (Italy) August 28 - September 2, 2011.
- [27] A. Nazin, S. Anulova, A. Tremba: Application of the Mirror Descent Method to Minimize Average Losses Coming by a Poisson Flow. European Control Conference (ECC) June 24–27, 2014.
- [28] J. Nocedal, S. J. Wright: Numerical Optimization. Springer, New York, 2006.
- [29] B. Polyak. Introduction to optimization. Optimization Software, New York (1987).
- [30] S. Shpirko, Yu. Nesterov: Primal-dual subgradient methods for huge-scale linear conic problem. SIAM J. Optim. **24**(3), 1444–1457 (2014)
- [31] N. Shor: On the structure of algorithms for the numerical solution of optimal planning and design problems (in Russian) Ph. D. dissertation, Cybernetics Institute, Academy of Sciences of the Ukrainian SSR, Kiev (1964).
- [32] F. S. Stonyakin, M. S. Alkousa, A. N. Stepanov, M. A. Barinov: Adaptive mirror descent algorithms in convex programming problems with Lipschitz constraints. Trudy Instituta Matematiki i Mekhaniki UrO RAN, 2018, Volume 24, Number 2, Pages 266–279.
- [33] F. S. Stonyakin, M. Alkousa, A. N. Stepanov, A. A. Titov: Adaptive Mirror Descent Algorithms for Convex and Strongly Convex Optimization Problems with Functional Constraints. J. Appl. Ind. Math. **13**, 557–574 (2019).
- [34] A. A. Titov, F. S. Stonyakin, A. V. Gasnikov, M. S. Alkousa: Mirror Descent and Constrained Online Optimization Problems. Optimization and Applications. 9th International Conference OPTIMA-2018 (Petrovac, Montenegro, October 1–5, 2018). Revised Selected Papers. Communications in Computer and Information Science, 974, pp. 64–78, 2019.
- [35] A. Tremba, A. Nazin: Extension of a saddle point mirror descent algorithm with application to robust PageRank. 52nd IEEE Conference on Decision and Control December 10–13, 2013.
- [36] E. Vorontsova, R. Hildebrand, A. Gasnikov, F. Stonyakin: Convex optimization. arXiv:2106.01946 <https://arxiv.org/pdf/2106.01946.pdf> (in Russian)

- [37] Y. Xu: Iteration complexity of inexact augmented lagrangian methods for constrained convex programming. Mathematical Programming, Series A, pp. 1–46, 2019.
- [38] Y. Xu: Primal-dual stochastic gradient method for convex programs with many functional constraints. SIAM Journal on Optimization Vol. 30, Iss. 2 (2020)
- [39] Z. Zhu, Y. Zhang, Y. Xia: Convergence Rate of Projected Subgradient Method with Time-varying Step-sizes. arxiv preprint 2023. <https://optimization-online.org/wp-content/uploads/2023/12/Subgradient-Method-2.pdf>

(M. S. Alkousa) MOSCOW INSTITUTE OF PHYSICS AND TECHNOLOGY, RUSSIA.
Email address: `mohammad.alkousa@phystech.edu`

(F. S. Stonyakin) MOSCOW INSTITUTE OF PHYSICS AND TECHNOLOGY AND V. I. VERNADSKY CRIMEAN FEDERAL UNIVERSITY, RUSSIA.
Email address: `fedyor@mail.ru`

(A. M. Abdo) MATHEMATICS DEPARTMENT, FACULTY OF SCIENCE, DAMASCUS UNIVERSITY, DAMASCUS, SYRIA.
Email address: `asmaa.abdo@damascusuniversity.edu.sy`

(M. M. Alcheikh) MATHEMATICS DEPARTMENT, FACULTY OF SCIENCE, DAMASCUS UNIVERSITY, DAMASCUS, SYRIA.
Email address: `mohammad.alcheikh@damascusuniversity.edu.sy`