### Information scrambling and butterfly velocity in quantum spin glass chains

Venkata Lokesh K. Y<sup>1</sup>, Surajit Bera<sup>2</sup> and Sumilan Banerjee<sup>2</sup>

<sup>1</sup> Centre of High Energy Physics, Indian Institute of Science, Bangalore 560012, India

<sup>2</sup> Centre for Condensed Matter Theory, Department of Physics, Indian Institute of Science, Bangalore 560012, India\*

(Dated: January 11, 2024)

We make lattice generalization of two well-known zero dimensional models of quantum spin glass, Sachdev-Ye (SY) and spherical quantum p-spin glass model, to one dimension for studying crossovers in non-local scrambling dynamics due to glass transition, complex dynamics, and quantum and thermal fluctuations in paramagnetic (PM) and spin glass (SG) phases. In the SY chain of quantum dots, each described by infinite-range random Heisenberg model of N spin-SSU(M) spins, we obtain the quantum Lyapunov exponent  $\lambda_{\rm L}$  and butterfly velocity  $v_B$  as a function of temperature T and the quantum parameter S across the PM-SG phase boundary using a bosonic spinon representation in the large-N, M limit. In particular, we extract asymptotic T and S dependence, e.g., power laws, for  $\lambda_{\rm L}$  and  $v_B$  in different regions deep inside the phases and near the replica symmetry breaking SG transition. We find the chaos to be non-maximal almost over the entire phase diagram. Very similar results for chaos indicators are found for the p-spin glass chain as a function of temperature and a suitable quantum parameter  $\Gamma$ , with some important qualitative differences. In particular,  $\lambda_{\rm L}$  and  $v_B$  exhibit a maximum, coinciding with onset of complex glassy relaxation, above the glass transition as a function of T and  $\Gamma$  in the PM phase of the p-spin glass model. In contrast, the maximum is only observed as a function of S, but not with temperature, in the PM phase of SY model. The maximum originates from enhanced chaos due to maximal complexity in the glassy landscape. Thus, the results in the SY model indicate very different evolution of glassy complexity with quantum and thermal fluctuations.


#### 1. INTRODUCTION

In a generic quantum many-body system, quantum information encoded in local degrees of freedom in the initial state scrambles to spread globally over the entire system due to its time evolution. The dynamics of quantum information, e.g., how rapidly local information scrambles, characterized by a rate or the quantum Lyapunov exponent  $\lambda_{\rm L}$ , and how fast the information propagates from one spatial region to other, quantified by a speed or the butterfly velocity  $v_B$ , can provide crucial insights into the non-trivial dynamics of plethora of quantum manybody systems<sup>1-4</sup>. These insights are complementary to that from transport and usual dynamical correlations. As a result, information scrambling has become an important theme of research for understanding holographic duality<sup>5</sup>, black hole information scrambling<sup>1,2</sup>, non-Fermi liquids<sup>3,6–9</sup> and quantum thermalization<sup>4,10,11</sup>. For systems close to a semiclassical limit, information scrambling can often be connected to chaos 12-16, namely extreme sensitivity of the classical trajectories to initial conditions.

For systems with many metastable configurations, information scrambling via quantum dynamics may become enhanced when the dynamics samples these multitude of configurations, leading to extreme sensitivity to initial conditions in the classical limit. Glasses<sup>17–19</sup>, which possess extensive number of metastable states<sup>18–22</sup>, are among the most prominent examples of such systems. At low temperatures, the system remains trapped near one of the metastable configurations for long times and explore the phase space extremely slowly. As a result, glasses exhibit many fas-

cinating dynamical properties ranging from complex relaxation with anomalously large time scales 17,23, aging and memory effects<sup>24</sup>, unusual transport properties such as the violation of Stokes-Einstein relations<sup>25</sup>, dynamical heterogeneity<sup>19,26</sup> etc. In particular, spin glasses<sup>23,27,28</sup> with quenched randomness undergo ergodicity breaking thermodynamic phase transition at low temperature. e.g., for mean-field or infinite-range models, to replica symmetry broken<sup>29,30</sup> spin glass phase. Recent work<sup>31</sup> on a well-known solvable zero-dimensional or infinite-range model of quantum spin glass, namely the quantum p-spin glass model<sup>32–34</sup>, have shown that chaos, as characterized by the Lyapunov exponent  $\lambda_{\rm L}$ , becomes stronger over a region above the glass transition where the dynamics exhibits typical signatures<sup>17</sup> of complex glassy relaxation. In this region of enhanced chaos, the dynamics samples exponentially large number of saddle points of the classical potential energy landscape and becomes maximally complex<sup>35,36</sup>, before eventually getting trapped within the basin of attraction of one of the glassy minima below the glass transition. In this work, we study scrambling dynamics in two solvable, albeit spatially extended models, of quantum spin glasses and show how the measures of information scrambling,  $v_B$  and  $\lambda_L$ , are influenced by glass transition, complex dynamics, and quantum and thermal fluctuations in various paramagnetic and spin glass phases.

In line with the growing interests  $^{31,35,37}$  in understanding the relation between scrambling and the complex dynamics of spin glass (SG) systems, we consider two well-known large-N zero dimensional (0+1 D) quantum spin glass models, namely (a) Sachdev-Ye (SY) model, a random Heisenberg spin glass  $^{38-40}$ , and (b) quantum spherical p-spin glass model  $^{32-34}$ , and make lattice gen-






FIG. 1.1. An one-dimensional (1D) generalization of (a) SY model, (b) p-spin model for p=3. Each lattice site contains a quantum dot described by zero-dimensional SY model in (a) and quantum spherical p-spin glass model in (b) with  $N\gg 1$  spins (blue circles). In (a), the coupling between nearest neighbour sites are random two-spin interaction with one from each lattice site for the SY model. In the case of p-spin model in (b), the random nearest neighbour interaction is between two spins from one site and one form the other and vice versa.

eralizations of these models to one dimension (1 + 1 D), as shown schematically in Fig. 1.1. These generalizations allow us to meaningfully define a butterfly velocity  $v_B$ , along with  $\lambda_L$ , from an out-of-time-order correlation (OTOC) function<sup>3,41</sup>. The SY model is parent to the celebrated fermionic Sachdev-Ye-Kitaev (SYK) model<sup>3,6,42</sup>, which has emerged as a paradigm to study quantum many-body chaos, information scrambling<sup>8,9,41</sup>, as well as strongly correlated metals<sup>43</sup>. The SYK model is maximally chaotic, namely it saturates the remarkable upper bound<sup>2,3,6</sup>  $2\pi k_{\rm B}T/\hbar$  for  $\lambda_{\rm L}$ . There have been numerous lattice and higher dimensional generalizations 41,44-51 of the zero dimensional SYK model, with a view of studying transport and information propagation in extended interacting systems. Our lattice generalizations of the spin glass models pave the way to access non-local dynamical correlations and spatio-temporal information scrambling in spatially extended strongly disordered systems that undergo ergodicity-breaking glass transitions.

Earlier works on quantum chaos in spin glass systems<sup>31,35,37</sup> have only looked at the Lyapunov exponent, e.g., its dependence on temperature T and quantum fluctuation parameter in the paramagnetic and spin glass phases, in the zero dimensional quantum p-spin glass model, i.e., for a single *p-spin glass quantum dot* (see Fig. 1.1). On the other hand, studies on spatially extended large-N systems, such as lattice generalizations of the SYK model<sup>41,44–51</sup>, have been limited to non (replica) symmetry broken phases, such as Fermi and non-Fermi liquids metals. Our lattice generalizations of the spin glass models retain the same large-N saddle points as the zero dimensional models, and capture replica symmetry broken spin glass (SG) and unbroken paramagnetic (PM) phases, such as a spin liquid for SY quantum dot. Thus, the spatial locality in these extended models permits us to explore information scrambling across

replica-symmetry breaking spin glass phase transition. In particular, we obtain the variations of the Lyapunov exponent and butterfly velocity with thermal and quantum fluctuations across the PM-SG phase transition and various crossovers between dynamically different paramagnetic states, e.g., spin liquid and local moment states in the SY model. We also find the asymptotic dependence of  $\lambda_{\rm L}$  and  $v_B$  on temperature T and quantum/classical parameter in different regions deep inside the phases as well as near the replica symmetry breaking SG transitions.

Temperature dependence of chaos indicators, especially that of the butterfly velocity, has been studied in several earlier works in various contexts, for examples, for solvable large-N models<sup>41,52–54</sup>, perturbatively in interaction for weakly interacting systems<sup>55</sup> and in 1/Nexpansion in strongly interacting systems<sup>7</sup>, via numerical methods<sup>54</sup> as well as in the semiclassical limit of spin models<sup>56–58</sup>. For a maximally chaotic  $[\lambda_L = 2\pi T]$  $(\hbar, k_{\rm B} = 1)$ ] interacting diffusive metal built from SYK  $dots^{41}$ ,  $v_B$  has been found to follow a  $\sqrt{T}$  dependence. Also, in this case, the quantity,  $v_B^2/\lambda_L$ , which has the dimension of the diffusion constant, turns out to be exactly equal to energy diffusion constant, as expected from holographic theories<sup>59</sup>. Similar relation,  $D=v_B^2/4\lambda_L$  has been found to be satisfied for electron diffusion constant D in weakly interacting diffusive metal<sup>55,60</sup>, where  $\lambda_{\rm L} \sim T^{d/2}, \ v_B \sim T^{d/4}$  in d dimension, and a temperature independent D is determined by elastic scattering. A power-law T dependence of  $v_B$  is similar to that at a critical point<sup>57,61</sup>, i.e.,  $v_B \sim T^{1-1/z}$  for a dynamical exponent z > 1. In a strongly interacting system with critical Fermi surface of N-species fermions coupled with U(1) gauge field, power-law behaviours,  $\lambda_{\rm L} \sim T$ ,  $v_B \sim T^{1/3}$ , with energy diffusion constant  $\sim v_B^2/\lambda_{\rm L}$ , have been obtained<sup>7</sup> at leading order in 1/N-expansion. The Lyapunov exponent and butterfly velocity also have been computed in (2+1 d) O(N) non-linear sigma model, where, in the quantum critical region<sup>53</sup>,  $\lambda_{\rm L} \sim T/N$ , and  $v_B \sim c$ , a constant close to the speed of light in the theory. Similar T independent  $v_B$  is found in the O(N)symmetry broken ordered phase<sup>53</sup> with  $\lambda_{\rm L} \sim T^3$ , whereas  $\lambda_{\rm L} \sim e^{-\tilde{E}_g/T}$ , due to the gap  $E_g$  in the disordered phase for  $T \ll E_g$ , and  $v_B$  obeys a  $\sqrt{T}$  behaviour<sup>53,54</sup>. The latter is consistent with numerical calculations in 1D quantum Ising model<sup>54</sup>. Power-law temperature dependence of  $\lambda_{\rm L}$  and  $v_B$  is also observed<sup>56,57</sup> in the ordered and disordered phases of spin models with semiclassical dynamics.

Our results for the dependence of  $\lambda_{\rm L}$  and  $v_B$  on temperature T and the quantum/classical parameter S in the chain of SY quantum dots are summarized on the schematic phase diagram of Fig. 1.2. In this model, the dots are described by infinite-range random Heisenberg model of spin-S SU(M) spins on N sites, where  $N, M \to \infty$ . Like in the lattice generalization<sup>41</sup> of the SYK model, the SY chain model, in a bosonic spinon representation, is described by the same saddle point of




FIG. 1.2. Schematic phase diagram for chaos in SY **model.** Dependence of Lyapunov exponent  $\lambda_{L}$  and butterfly velocity  $v_B$  on temperature T and quantum parameter S are shown in the paramagnetic (PM) and spin glass (SG) phases.  $\lambda_{\rm L}$  and  $v_B$  increases as a power law in S for small S in the PM phase, and as a power law in T for low temperature in the SG phase. Both the chaos indicators increases slowly, as  $\sim \ln T$ , above the glass transition  $T_{SG}$ , before saturating to a constant at high temperature. In the SG phase,  $\lambda_{\rm L}$  decreases as  $\sim 1/S^2$ , whereas  $v_B \sim e^{aS}$  increases exponentially in S. There is also maximum (not shown) in both  $\lambda_{\rm L}$  and  $v_B$  as a function of S in the PM phase, at intermediate S, but there are no maxima as a function of T (see main text). The region bounded by  $T_{SG}(S)$  and  $T/\tilde{J}=1$  ( $\tilde{J}$  coupling constant) line (dotted line) demarcates the quantum critical region within the PM phase. Arrows indicate the direction of increase of the parameter.

the zero-dimensional quantum dots with the strength of the coupling J renormalized by the inter-dot interaction. As a result,  $\lambda_{\rm L}$ , which is extracted from a local OTOC, has the same S and T dependence of a single dot, or the original SY model<sup>38</sup>. On the contrary, the butterfly velocity can only be defined in a lattice model like the SY chain of Fig. 1.1(a). To compare with the chaotic properties of the SY chain, we also calculate the butterfly velocity from the non-local OTOC as a function of T and quantum parameter  $\Gamma$  (defined later) in a similar lattice model [Fig. 1.1(b)] of quantum spherical p-spin spin glass model<sup>32–34</sup>, as summarized in Fig. 1.3 (schematic). Again,  $\lambda_{\rm L}(\Gamma,T)$  of the p-spin glass chain is identical to that of the corresponding zero-dimensional model, and was studied in ref.31. Our results are summarized below.

(1) As shown in Fig. 1.2, we find a power-law increase with S in the SY model, i.e.,  $\lambda_{\rm L}(S,T) \simeq \lambda_{\rm L}(0,T) + aS^{\alpha_{\lambda}}$  and  $v_B(S,T) \simeq v_B(0,T) + aS^{\alpha_{\nu}}$  for small S in the PM phase with  $\alpha_{\lambda(v)} \approx 0.5$  at a fixed temperature, like in the quantum critical region for T < J, where two types of PM behaviours, spin liquid and local moment, compete. In contrast to the power-law or gap-like temperature dependence seen in the disordered phase of many other models  $^{53,54,56,57}$ , a ln T variation of both  $\lambda_{\rm L}$  and  $v_B$  are seen in the PM phase pf the SY model close to the




FIG. 1.3. Schematic phase diagram for chaos in p-spin glass model. Dependence of  $v_B$  on temperature T and quantum parameter  $\Gamma$  are shown in the PM and SG phases. Similar dependence has been seen for  $\lambda_L$  in Ref.31. The continuous PM-SG phase transition  $T_d(\Gamma)$  [or  $T_{SG}(\Gamma)$ ] (solid line) terminates at tricritical point and thereafter the transition is first order (hatched line). There is a maximum in  $v_B$  at  $T_m(\Gamma)$  (dotted line), both as a function of T and  $\Gamma$ .  $v_B$  increases as power law with T and  $\Gamma$  at low temperature and small  $\Gamma$ , respectively, in the SG phase. The butterfly velocity increases as  $\sim \ln T$  and as a power law in  $\Gamma$  above the glass transition in the PM phase, before the maximum. Arrows indicate the direction of increase of the parameter.

transition temperature  $T_{SG}$ . Similar  $\ln T$  dependence is seen for the PM phase of the *p*-spin glass model, where  $v_B \sim \Gamma^{\alpha_v}$  with  $\alpha_v \approx 1$ .

- (2) On the contrary, in the SG phase of SY model, power-law temperature dependence,  $\lambda_{\rm L}(S,T) \sim T^{\beta_{\lambda}}$ , and a power-law increment  $v_B(S,T) \simeq v_B(S,0) + aT^{\beta_v}$ , with  $\beta_{\lambda} \approx 1 - 1.5$  and  $\beta_{v} \approx 1.5$ , are observed. The velocity  $v_B(S,0)$  extrapolated to zero temperature is finite, indicating information spreading in the quantum SG ground state of the SY chain. Very similar behaviours of  $\lambda_{\rm L}(\Gamma,T)$  and  $v_B(\Gamma,T)$  occurs in the SG phase of p-spin glass. In the SG phase of SY model at a fixed T, the Lyapunov exponent and butterfly velocity exhibit contrasting trends with increasing S or decreasing quantum fluctuations;  $\lambda_{\rm S}$  decreases as  $\sim 1/S^2$ , whereas  $v_B$ increases exponentially with S. These behaviours are different from the p-spin glass model where both  $\lambda_{\rm L}^{31}$ and  $v_B$  decreases with decreasing quantum fluctuation, e.g.,  $v_B(\Gamma, T) \simeq v_B(0, T) + a\Gamma^{\alpha_v}$  with  $\alpha_v \approx 0.5$ .
- (3) The Lyapunov exponent and butterfly velocity have smooth crossovers across the PM-SG transitions in both SY and p-spin glass models. However,  $v_B$  exhibits a dip as a function of S at the SG transition for the SY chain, indicating slowing down of information spread at the transition. This is unlike the  $v_B(\Gamma)$  which monotonically changes across the SG transition in the p-spin glass.
- (4) As mentioned earlier,  $\lambda_{\rm L}(\Gamma, T)$  exhibits a maximum<sup>31</sup> [ $T_m(\Gamma)$  in Fig. 1.3] in the PM phase of the p-spin glass model as function of both  $\Gamma$  and T. The maximum

corresponds to enhanced chaos due to the sampling of maximum complexity of the glassy landscape at  $T_m^{35,36}$ . The same maximum is found in  $v_B$  for p-spin glass. However, the maximum in  $\lambda_{\rm L}$  and  $v_B$  is seen in the PM phase of the SY model only as a function of S, but not as a function of T. This implies very different evolution of complexity with quantum and thermal fluctuations in SY model, unlike that in the p-spin glass model.

(5) For computing the butterfly velocity, we implement a numerical approach alternative to the more standard semi-analytical  $single-mode\ ansatz^{52,62}$ . The results from the numerical method agree very well with those from the single-mode ansatz over the entire phase diagrams of SY and p-spin glass chains, thus, independently verifying the applicability of the single-mode ansatz even for the replica symmetry broken spin glass phase. Using the single-mode ansatz, we find that chaos is non-maximal almost over the entire phase diagram for SY and p-spin glass models.

We also connect our results for the chaos parameters with spectral characteristics of various regions of the phase diagram of the two models, like spin liquid, local moment, classical and quantum paramagnets, and spin glass. The rest of the paper is organized as follows. In Sec. 2, we discuss the one dimensional lattice generalizations of the zero-dimensional SY and quantum p-spin glass model. The large-N saddle-point equations for the PM and SG phased for both the models are described in Sec. 3, followed by review of the phase diagrams known from earlier works. Sec. 4 describes the formalism for calculating local OTOC and the Lyapunov exponent using ladder approximation for the PM and SG phases for the SY model. This section reports the results for  $\lambda_L(S,T)$ in the SY model and compare them with those in the p-spin glass model. The formalism for computing the butterfly velocity from non-local OTOC using two different methods, a numerical method and the semi-analytical single-mode ansatz<sup>62</sup>, is discussed in Sec. 5 for both SY and p-spin glass models. The results for  $v_B$  as a function of T and quantum parameter for the two models are described in this section. The appendices (Appendix A, Appendix B) give additional details of the derivations of the saddle-point equations and methods for their numerical self-consistent solutions. Appendix A 2 discusses the results for the spectral properties for the SY model. Appendix C provides details of the numerical computation of  $v_B$ .

# 2. LATTICE GENERALIZATIONS OF SOLVABLE QUANTUM SPIN GLASS MODELS

As mentioned earlier, we consider one-dimensional (1D) chains of zero-dimensional random quantum Heisenberg and spherical *p*-spin glass quantum dots, as shown in Fig. 1.1 and discussed below.

#### A. 1D chain of Sachdev-Ye (SY) spin-glass quantum dots

We make a lattice generalization of the well-known zero-dimensional SY model of random quantum Heisenberg model by arranging  $x=1,\ldots,L$  quantum dots of SY model, where each dot contains  $i=1,\ldots,N$  sites. The model is described by the Hamiltonian

$$\mathcal{H} = \frac{1}{2\sqrt{MN}} \sum_{x,i\neq j} J_{ij,x} S_{i\alpha\beta x} S_{j\beta\alpha x} + \frac{1}{\sqrt{MN}} \sum_{x,i,j} J'_{ij,x} S_{i\alpha\beta x} S_{j\beta\alpha,x+1}.$$
 (2.1)

Here the first term represents all-to-all infinite-range interactions within a dot, and the second term the nearestneighbour inter-dot coupling. The couplings  $J_{ij,x}$  and  $J'_{ij,x}$  are drawn from Gaussian distribution, independent dently at each site with zero mean and variances  $\overline{J_{ij}^2} = J^2$ and  $\overline{J_{ij}^{\prime 2}} = J^{\prime 2}$ .  $S_{i\alpha\beta x}$  denote the spin degrees of freedom, the generators of the SU(M) group for spin S, at each i and dot x, where  $\alpha, \beta = 1, \dots, M$ . The model is exactly solvable in the large N limit  $(N \to \infty)$  followed by the large M limit  $(M \to \infty)$ , i.e., thermodynamics, as well as equilibrium and non-equilibrium properties can be exactly obtained in this limit through large-N saddlepoint equations, as known from previous works<sup>38–40,63</sup> on the zero-dimensional model. The effect of quantum fluctuations in the model can be tuned by varying the spin S and the classical and semiclassical regimes can be probed by taking the large  $S(S \to \infty)$  limit while appropriately scaling the energy /temperature scales<sup>40</sup>. The M=2 limit of the model is the usual SU(2) Heisenberg model, which has been studied via quantum Mote Carlo  $(QMC)^{64}$  after taking the  $N \to \infty$  limit. The large M limit was found to capture the properties of the SU(2)model quite well.

In Sec. 3 A, we discuss the equilibrium phase diagram obtained from the large-N saddle-point equations as a function of temperature T and spin S. The large-N saddle point is obtained via the standard bosonic representation of the SU(M) spin<sup>38</sup>. Subsequently, our main results on the dynamical phase diagram for chaos, characterized via the Lyapunov exponent  $\lambda_{\rm L}$  and butterfly velocity  $v_B$ , are discussed in Secs. 4, 5.

#### B. 1D chain of spherical p-spin glass quantum dots

To compare and contrast the results for  $\lambda_{\rm L}$  and  $v_B$  in the SY chain, we consider the lattice generalization of another paradigmatic model of quantum glasses, the quantum spherical p-spin glass model<sup>32–34,65</sup>. The 1D chain of the p-spin glass quantum dots  $(x=1,\ldots,L)$  is

described by the Hamiltonian

$$\mathcal{H} = \sum_{x,i} \frac{\pi_{i,x}^2}{2\mathcal{M}} + \frac{1}{3!} \sum_{x,ijk} J_{ijk,x} s_{i,x} s_{j,x} s_{k,x}$$

$$+ \frac{1}{2!} \sum_{x,ijk} s_{i,x} (J_{ijk,x}^+ s_{j,x+1} s_{k,x+1} + J_{ijk,x}^- s_{j,x-1} s_{k,x-1}),$$

$$(2.2)$$

for p = 3. The intra- and inter-dot all-to-all couplings among sites i = 1, ..., N in the dots,  $J_{ijk,x}$  and  $J_{ijk,x}^{\pm}$ , are Gaussian random numbers with zero mean and variances  $3J^2/(2N^2)$  and  $3J'^2/(2N^2)$ , respectively. The commutation relation of the spin  $s_{i,x}$  with the momentum  $\pi_{i,x}$ ,  $[s_{i,x}, \pi_{j,x'}] = i\hbar \delta_{ij}\delta_{xx'}$  induces the quantum dynamics. The spherical constraint  $\sum_i s_i^2 = N$  makes the model non trivial. Essentially, when the quantum dots are uncoupled (J'=0), the Hamiltonian describes quantum particles with mass  $\mathcal{M}$  moving on the surface of an Ndimensional hypersphere. The inter-dot interaction couples particles in neighbouring dots. The advantage of this model lies in the fact that one can take a classical limit of the model by continuously tuning a quantum fluctuation parameter  $\Gamma = \hbar^2/\mathcal{M}J$  to zero<sup>31,37</sup>. The classical model for  $\hbar \to 0$  limit (fixed  $\mathcal{M}$ ) for a single dot gives rise dynamics identical to the mode coupling theory (MCT) dynamics in the super-cooled liquid regime of structural glasses<sup>17,19,31,66,67</sup>. The limit  $\mathcal{M} \to \infty$  while keeping  $\hbar$ fixed lead to a different classical limit of infinitely heavy mass<sup>31,37</sup>. In this work we vary the dimensionless quantum parameter  $\Gamma$ .

In Sec. 3B, we discuss the equilibrium phase diagram of the model as a function of T and  $\Gamma$ . The results for  $\lambda_{\rm L}$  and  $v_B$  are discussed in Secs. 4 and 5 and compared with those from the lattice SY model.

#### 3. EQUILIBRIUM PHASE DIAGRAMS OF THE SPIN GLASS CHAINS

We first discuss the equilibrium phase diagram of the quantum spin glass chains [Eq.(2.1) and Eq.(2.2)] as a function of T and quantum fluctuation parameters, S or  $\Gamma$  [Figs. 3.1, 3.2]. The equilibrium phase diagrams are obtained by solving large-N saddle-point equations. As we discuss below, due to the particular choice of inter-dot couplings in Eqs.(2.1),(2.2), the large-N saddle-point equations of the lattice models are identical to the zero-dimensional saddle-point equations<sup>38,65</sup> of the single dot with renormalized intra-dot coupling.

#### A. Sachdev-Ye chain

To capture both (replica) symmetry broken ordered SG and disordered paramagnetic (PM) phases we use bosonic spinons to represent spin-S SU(M) generators as  $S_{i\alpha\beta x} = b^{\dagger}_{i\alpha x}b_{i\beta x} - S\delta_{\alpha\beta}$ , where  $b_{i\alpha x}$ ,  $b^{\dagger}_{i\alpha x}$  ( $\alpha = 1, ..., M$ )




FIG. 3.1. Mean-field phase diagram of the bosonic spinon model howing the paramagnetic and spin glass phases. Spin glass phase below  $T_{SG}$  (solid line) is determined by the marginality criterion (see main text).  $T_{eq}$  (thick dashed line) is determined by equilibrium criterion. The regimes within spin glass phase separated by blue lines are identified via a large S expansion of saddle point equations  $^{40}$  I, II and III are classical, semi-classical and quantum spin glass regions respectively. IV is the quantum critical regime in PM phase. We perform the calculations of Lyapunov exponent and butterfly velocity across the SG-PM transition for several S as a function of S (along horizontal dashed lines S0, and S1 as a function of S2 (along horizontal dashed lines S3.

are bosonic annihilation and creation operators with the constraint  $\sum_{\alpha} b_{i\alpha x}^{\dagger} b_{i\alpha x} = SM \ (S>0)$  on the number of bosons at each i,x to fix the spin of the representation to S. The saddle-point equations, their solutions and resulting phase diagram for the bosonic representation have been discussed in detail for the zero-dimensional model in the pioneering works<sup>38–40</sup>. We briefly discuss the derivation of saddle point equations for the SY chain model [Eq.(2.1)] in Appendix A.

The disorder averaged Green's function  $G_x^{ab}(\tau,\tau')=G^{ab}(\tau,\tau')=-\frac{1}{M}\sum_{\alpha}\langle T_{\tau}b_{i\alpha x}^a(\tau)\bar{b}_{i\alpha x}^b(\tau')\rangle$  is independent of i,x and determines the equilibrium dynamics of the model. The Green's function is obtained from the saddle point equations in the limit  $N\to\infty$  followed by  $M\to\infty$  as

$$(G^{-1})^{ab}(i\omega_k) = (i\omega_k + \lambda)\,\delta_{ab} - \Sigma^{ab}(i\omega_k) \qquad (3.1a)$$

$$\Sigma^{ab}(\tau) = \tilde{J}^2 [G^{ab}(\tau)]^2 G^{ba}(-\tau)$$
 (3.1b)

where  $\omega_k=2\pi kT$  is bosonic Matsubara frequency with k an integer  $(k_B=1), T$  the temperature in units of  $\tilde{J}$ . We have introduced the replicas  $a=1,\ldots,n$  to carry out the disorder average.  $G(\mathrm{i}\omega_k)=\int_0^\beta d\tau e^{\mathrm{i}\omega_k\tau}G(\tau)$  and  $\beta$  is the inverse temperature. Lagrange multiplier,  $\lambda$  is introduced to impose the constraint,  $G^{aa}(\tau=0^-)=-S$ . The above saddle-point equations are same as the zero-dimensional ones<sup>38,39</sup> with renormalized coupling  $\tilde{J}^2=J^2+J'^2$  due to the nearest-neighbour inter-dot coupling in Eq.(2.1). In the limit  $n\to 0$ , the replica structure of the Green's function determines the PM and SG phases

as we discuss below.

#### 1. Paramagnetic phase

In the PM phase one employs a replica diagonal and symmetric ansatz,  $G^{ab}(\tau) = G(\tau)\delta_{ab}$ . Thus, the saddle-point equations reduce to,

$$[G(i\omega_k)]^{-1} = (i\omega_k + \lambda) - \Sigma(i\omega_k)$$
 (3.2a)

$$\Sigma(\tau) = \tilde{J}^2[G(\tau)]^2G(-\tau) \tag{3.2b}$$

Analytic continuation from Matsubara to real frequency  $\mathrm{i}\omega_k\to\omega+\mathrm{i}0^+$  gives the saddle point equation for retarded Green's function

$$[G_R(\omega)]^{-1} = \omega + \lambda - \Sigma_R(\omega) \tag{3.3}$$

where  $\Sigma_R(\omega) = \Sigma(\mathrm{i}\omega_k \to \omega + \mathrm{i}0^+)$  and  $\Sigma(\mathrm{i}\omega_k) = \int_0^\beta d\tau e^{\mathrm{i}\omega_k\tau}\Sigma(\tau)$ .  $\lambda$  is determined by the constraint on  $G(\tau=0^-)$ . The procedure to numerically solve for the retarded Green's function is discussed in Appendix A.

### 2. Spin glass phase

Following earlier works<sup>40</sup>, we describe the spin glass phase via the following one-step replica symmetry breaking (1RSB) ansatz for the bosonic Green's function that appears in Eq.(3.1).

$$G^{ab}(\tau) = \tilde{G}(\tau)\delta_{ab} - g\epsilon_{ab} \tag{3.4}$$

or,  $G^{ab}(i\omega_k) = \tilde{G}(i\omega_k)\delta_{ab} - \beta g\epsilon_{ab}\delta_{\omega_k,0}$ , and

$$\Sigma^{ab}(\tau) = \tilde{\Sigma}(\tau)\delta_{ab} - \tilde{J}^2 g^3 \epsilon_{ab}, \qquad (3.5)$$

where  $\epsilon_{ab}=1$  in the diagonal block of size  $m\times m$ , and 0 otherwise, for the  $n\times n$  matrices. Here  $\tilde{G}(\tau)$  is the regular part of the Green's function such that  $\tilde{G}(\tau\to\infty)\to 0$  at T=0. The Edward-Anderson (EA) spin-glass order parameter is obtained from  $q_{\rm EA}=\lim_{\tau\to\infty}(1/M^2)\sum_{\alpha\beta}\langle S_{i\alpha\beta x}(\tau)S_{i\beta\alpha x}(0)\rangle=\lim_{\tau\to\infty}G^{aa}(\tau)G^{aa}(-\tau)=g^2$  at zero temperature. Using  $[G^{-1}]^{ab}(\mathrm{i}\omega_k)=A(\mathrm{i}\omega_k)\delta_{ab}+B(\mathrm{i}\omega_k)\epsilon_{ab}$  and the standard replica matrix inversion, we get

$$A(i\omega_k) = \frac{1}{\tilde{G}(i\omega_k)},\tag{3.6a}$$

$$B(i\omega_k) = \frac{\beta g}{[\tilde{G}(i\omega_k) - m\beta g]\tilde{G}(i\omega_k)}.$$
 (3.6b)

The saddle point equation for  $\tilde{G}$  is therefore,

$$[\tilde{G}(i\omega_k)]^{-1} = i\omega_k - \frac{\tilde{J}g}{\Theta} - [\tilde{\Sigma}(i\omega_k) - \tilde{\Sigma}(i\omega_k = 0)] \quad (3.7a)$$

where

$$\tilde{\Sigma}(\tau) = \tilde{J}^2 \left[ \tilde{G}^2(\tau) \tilde{G}(-\tau) - 2g\tilde{G}(\tau) \tilde{G}(-\tau) - g\tilde{G}^2(\tau) + 2g^2 \tilde{G}(\tau) + g^2 \tilde{G}(-\tau) \right]$$
(3.7b)

In the above, we have eliminated  $\lambda$  using the equation for  $i\omega_k = 0$  and the parameterization  $\tilde{G}(i\omega_k = 0) = -\Theta/\tilde{J}g$ . The saddle point equation for g is obtained from Eq.(3.6b) as,

$$m\beta = \frac{1}{\tilde{J}g^2} \left( \frac{1}{\Theta} - \Theta \right) \tag{3.8}$$

The equations Eq. 3.7a, Eq. 3.7b and Eq. 3.8 along with the constraint,  $\tilde{G}(\tau=0^-)=g-S$  provide a closed set of self-consistent saddle-point equations for the SG phase. The characteristic of 1RSB symmetry breaking solution is that the saddle point equations form a one-parameter family, parameterized here by  $\Theta$  or equivalently by the breaking point m. To obtain this parameter we impose the usual marginality criterion<sup>40</sup>, namely we fix  $\Theta = \Theta_R$ , and hence m, such that the lowest eigenvalue of the fluctuation matrix in the replica space vanishes.

The marginality criterion leads to a gapless  $\chi''(\omega) \propto \omega$  with a delta function peak at  $\omega = 0$ , where  $\chi^{aa}(\tau) = G^{aa}(\tau)G^{aa}(-\tau) = (1/M^2)\sum_{\alpha\beta}\langle S_{i\alpha\beta x}(\tau)S_{i\beta\alpha x}(0)\rangle$  is the local spin susceptibility<sup>40</sup>. The equilibrium criterion, which minimizes free energy with respect to the breaking point m, leads to a gapped spectrum<sup>40</sup>. Analytic continuation of Eq.(3.7a) gives the equation for retarded Green's function  $G_B^{ab}(\omega) = G_R(\omega)\delta_{ab}$  with

$$[G_R(\omega)]^{-1} = \omega - \frac{\tilde{J}g}{\Theta_R} - [\Sigma_R(\omega) - \Sigma_R(\omega = 0)] \quad (3.9)$$

where  $\Sigma_R(\omega) = \tilde{\Sigma}(i\omega_k \rightarrow \omega + i0^+)$  and  $\tilde{\Sigma}(i\omega_k) =$  $\int_0^\beta d\tau e^{i\omega_k \tau} \tilde{\Sigma}(\tau)$ .  $\lambda$  is fixed by the constraint  $\tilde{G}(\tau=0^-)=$ g-S, g is calculated from the relation Eq. 3.8. The procedure for numerically calculating the retarded Green's function  $G_R(\omega)$  is discussed in Appendix A. For a fixed S, the temperature  $T_{SG}$ , below which the spins freeze into a SG order is identified by  $m(T_{SG}) = 1$ , where the SG order parameter  $q_{\rm EA}$  jumps discontinuously to zero in the PM phase<sup>40</sup>. A similar procedure is followed to obtain  $S_{SG}$  for a given temperature T. Unlike earlier  $works^{40,68}$ , in this work we mostly use the real-frequency saddle-point solutions to find the Lagrange multiplier  $\lambda$ , EA order parameter  $q_{\rm EA}$  in the SG phase, the PM-SG phase boundary, as well as the retarded Green's function  $G_R(\omega)$ , and chaos diagnostics  $\lambda_L$  and  $v_B$ . The numerical methods employed for obtaining the real-frequency saddle-point solutions are discussed in Appendix A1a. The imaginary-time saddle-point equations are only used for deriving the real-frequency saddle-point equations via analytical continuation, and occasionally, to check the consistency of the thermodynamic quantities like  $\lambda$ ,  $q_{EA}$ and  $T_{SG}(S)$  computed from the real-frequency solution. For the sake of completeness, we discuss the numerical method used for the imaginary-time solution in Appendix A 1 b. The same procedure is followed for the *p*-spin glass model.

#### 3. Phase transition and crossovers

The equilibrium phase diagram of the SY chain model is identical to the zero-dimensional case, which has been studied in detail in the  $N, M \to \infty$  limit in both bosonic and fermionic representation of the SU(M) spins for general  $S^{38-40}$ . The phase diagram has been also studied for finite M in the  $N \to \infty$  limit, using 1/M expansion<sup>69</sup> and for small S, e.g., S = 1/2, through QMC<sup>64</sup> in the  $N \to \infty$  limit, as well as via exact diagonalization for finite  $N^{70}$ . In Fig. 3.1, we reproduce the schematic large-N, M phase diagram obtained via bosonic representation in ref.40. This is consistent with QMC result<sup>64</sup> for S=1/2 for  $N\to\infty$ . The system undergoes a PM to SG phase transition at a temperature  $T_{SG}\sim (2/3\sqrt{3})\tilde{J}S^2$ , where the transition temperature,  $T_{SG}^{eq}$  obtained from the equilibrium criterion is slightly smaller than the temperature calculated via the marginal stability criterion. We refer to this latter temperature as  $T_{SG}$ . This corresponds to a dynamical transition, similar to that in the p-spin glass model<sup>33,65</sup>, where the spin relaxation time diverges and glassy aging behaviour commences in the non-equilibrium dynamics<sup>63</sup>, approaching from the PM side.

The point S = 0, T = 0 corresponds to a quantum critical point with a spin liquid ground state, akin to SYK non-Fermi liquid<sup>3,6</sup>. The spin liquid state exhibits so-called 'marginal spectrum' i.e.,  $\chi''(\omega) \sim$  $\operatorname{sgn}(\omega)$  at T=0, or  $\chi''(\omega,T)\sim \tanh(\omega/2T)$  at finite temperature, and diverging spin susceptibility  $\chi(T)$  ~  $\ln(T/J)$ , whereas the spectral density of the bosons  $\rho(\omega) = -(1/\pi) \text{Im} G_R(\omega) \sim -\text{sgn}(\omega)/\sqrt{|\omega|}$  has a characteristic power-law divergence at T=0. At any finite S the ground state is a spin glass and the spin liquid state becomes metastable up to a small maximum  $S = S_{max} \simeq 0.052^{40}$  at T = 0. However, as shown in Fig. 3.1, a quantum critical regime with marginal spectrum persists at finite temperature,  $T_{SG} < T \lesssim \tilde{J}$ , up to  $S \lesssim 2$ . Apart from the spin liquid, another paramagnetic solution to the Eqs. (3.2), the so-called 'local moment' solution<sup>40</sup> coexists in the phase diagram. This leads to a high-temperature  $(T \gg \tilde{J})$  Curie susceptibility  $\chi(T) \simeq S(S+1)/T$  which gradually decreases to  $\chi(T) \simeq S^2/T$  as  $T \to 0$ , i.e., below  $T_{SG}$ , due to partial screening of the spins due to spin-spin interaction. Of course, for  $T < T_{SG}$ , the local moment solution is metastable. In appendix Appendix A2, we discuss the numerically obtained bosonic spectral function  $\rho(\omega)$ across the PM-SG transition as a function of S for fixed T, and as a function of T for fixed S, in the regime of interest for our chaos calculations (Secs. Appendix A1b, **5**).

Based on the large-S expansions in ref.40, the SG phase in Fig. 3.1 can be divided into three regimes – I. a classical regime for  $\tilde{J}S \lesssim T \lesssim \tilde{J}S^2$  where all non-zero Matsubara frequency components of the Green's function in eq.(3.1) can be neglected and spins become commuting vectors of length S, II. a semiclassical regime for  $\tilde{J}\sqrt{S} \lesssim T \lesssim \tilde{J}S$ , where  $\rho(\omega) \sim \omega$  for  $\omega \to 0$  is obtained by 1/S expansion of the saddle-point equations in the SG phase (Sec. 3A2) after scaling  $\omega$  and T by  $\tilde{J}S$ , and III. a quantum regime for  $0 \le T \lesssim \tilde{J}\sqrt{S}$ , deduced from the 1/S expansion of the internal energy and linear in T specific heat. In this work, we do not study the chaos across these various quantum-classical crossovers in detail. As shown in Fig. 3.1, we focus on changes in the chaotic properties, like  $\lambda_{\rm L}$  and  $v_{\rm B}$ , from different regions of the PM phase to the SG phase across the SG

Here, we only study the chaos in the SY chain model [Eq.(2.1)] using bosonic representation. The model can also be studied in the large N,M limit via fermionic representation  $^{38,39,69}$ . In this case, the ground state is a spin liquid for all  $0 \le S \le 1$  accessible in the fermionic representation. Thus system is maximally chaotic with  $\lambda_{\rm L} = 2\pi T$  ( $\hbar = 1$ ) for  $T \to 0$ . However, for finite M the SG order can be recovered below  $T_{SG} \sim \tilde{J} \exp(-\sqrt{M\pi})^{40,69}$ . In the SG ground state at T = 0, one obtains the marginal spectrum  $\chi''(\omega) \sim \text{sgn}(\omega)$  above a characteristic frequency  $\omega^* = \tilde{J}q_{\rm EA}$ , i.e., over a range  $\omega^* < |\omega| \ll \tilde{J}$ , and the marginal SG behaviour for  $\chi''(\omega)$  for smaller frequencies  $0 < |\omega| < \omega^*$ . The EA order parameter in the fermionic case is  $q_{\rm EA} \sim 1/M$ .

#### B. p-spin glass chain

As in the case of SY model, the equilibrium dynamics of the *p*-spin glass chain [Eq.(2.2)] is described by the disorder averaged Green's function  $Q_{ab}(\tau, \tau') = (1/N) \sum_i \overline{\langle \mathcal{T}_{\tau} s_i^a(\tau) s_i^b(\tau') \rangle}$ . This model is solvable in the large N limit and is described by the Dyson equation,

$$(Q^{-1})^{ab}(i\omega_k) = \left(\omega_k^2/\Gamma + z\right)\delta_{ab} - \Sigma^{ab}(i\omega_k) \qquad (3.10a)$$
$$\Sigma^{ab}(\tau) = \frac{3\tilde{J}^2}{2}[Q_{ab}(\tau)]^2, \qquad (3.10b)$$

where  $\tilde{J}^2 = J^2 + J'^2$ . The derivation of the above saddle-point equations, which are identical to that of the zero-dimensional p-spin glass model<sup>65</sup>, is discussed in Appendix B. The spherical constraint  $Q_{aa}(\tau=0)=1$  is imposed by the Lagrange multiplier z. Similar to the case of SY model, in the limit  $n \to 0$ , we have the replica diagonal PM and 1RSB SG phases. In the PM phase, the saddle point equations, after analytical continuation, simplify to

$$[Q_R(\omega)]^{-1} = -\omega^2/\Gamma + z - \Sigma_R(\omega),$$
  
$$\Sigma_R(\omega) = \Sigma_R(i\omega_k \to \omega + i0^+)$$
 (3.11a)

In the SG phase, we consider the 1RSB solution, which is exact<sup>65</sup>. Using this ansatz, we can write the Dyson equation as,

$$Q^{ab}(\tau) = (q_d(\tau) - q_{\rm EA})\delta_{ab} + q_{\rm EA}\epsilon_{ab}$$
 (3.12)

which after Fourier transformation becomes,

$$Q^{ab}(i\omega_k) = (q_d(i\omega_k) - \tilde{q}_{EA})\delta_{ab} + \tilde{q}_{EA}\epsilon_{ab}, \qquad (3.13)$$

where  $\tilde{q}_{\rm EA} = \beta q_{\rm EA} \delta_{\omega_k,0}$ . Further, it is convenient to write  $q_d(\tau) = \tilde{Q}(\tau) + q_{\rm EA}$ , i.e., in terms of a regular part, such that  $\tilde{Q}(\tau \to \infty) \to 0$  at T = 0, and EA order parameter  $q_{\rm EA}$ . This can be inverted in the replica space as,

$$[Q^{-1}]^{ab}(i\omega_k) = A(i\omega_k)\delta_{ab} + B(i\omega_k)\epsilon_{ab}$$
 (3.14)

with

$$A(i\omega_k) = \frac{1}{q_d(i\omega_k) - \tilde{q}_{EA}}, \qquad (3.15a)$$

$$B(i\omega_k) = \frac{-\tilde{q}_{\rm EA}}{q_d(i\omega_k)^2 + (m-2)\tilde{q}_{EA}q_d(i\omega_k) - (m-1)\tilde{q}_{\rm EA}^2}$$
(3.15b)

where  $q_{\rm EA}$  is Edward-Anderson order parameter and  $\epsilon_{ab}=1$  for the diagonal blocks and zero otherwise.

This leads to the saddle point equations,

$$\omega_k^2/\Gamma + z = \frac{1}{\tilde{Q}(i\omega_k)} + \tilde{\Sigma}(i\omega_k), \qquad (3.16)$$

and

$$\frac{3(\beta m)^2}{2}q_{\rm EA}^3 = \frac{x_p^2}{1+x_p},\tag{3.17a}$$

$$y = \beta q_{\rm EA}/q_d(0)$$
 and  $x_p = my/(1-y)$ . (3.17b)

where  $\tilde{\Sigma}(\mathrm{i}\omega_k)=\frac{3\tilde{J}^2}{2}\int_0^\beta d\tau e^{\mathrm{i}\omega_k\tau}(q_d^2(\tau)-q_{\mathrm{EA}}^2)$ . The marginality criterion leads to  $x_p=1^{65}$ , which determines the breaking point, m. The saddle-point equations along with the constraint  $\tilde{Q}(\tau=0)=1-q_{\mathrm{EA}}$ , can be analytically continued to real frequency and solved numerically. The details are discussed in the Appendix B. We show the phase diagram determined by the solutions of the saddle point equation in Fig. 3.2.

## 4. OUT-OF-TIME-ORDERED CORRELATOR (OTOC) AND LYAPUNOV EXPONENT

To characterize the many-body quantum chaos in the SY and p-pin glass chain models [Eqs. (2.1), (2.2)], we first extract Lyapunov exponent from *on-site* OTOC. The calculation of Lyapunov exponent in the chain models are identical, with the effective coupling  $\tilde{J} = \sqrt{J^2 + J'^2}$ , to that in the corresponding zero-dimensional models, as have been done already for the p-spin glass in Ref.31. Thus, we focus on the effective zero-dimensional




FIG. 3.2. Mean-field phase diagram of quantum spherical p-spin glass model<sup>65</sup> as a function of temperature T and dimensionless quantum fluctuation parameter  $\Gamma$  (see main text). The solid line denotes second order phase transition between SG and PM phases. The second order line terminates at a tricritical point and the transition becomes first order for low temperature and larger  $\Gamma$ . Spin glass phase is determined by the marginality criterion. We calculate the Lyapunov exponent and butterfly velocity across the SG-PM transition for two values of T as a function of  $\Gamma$  (along vertical dashed lines x and y) and for two values of  $\Gamma$  as a function of T (along horizontal dashed lines u and v).




FIG. 4.1. Schwinger-Keldysh contour with four real-time branches separated by  $\beta/4$  in imaginary time, used for calculating OTOC.

SY model, where the dot indices x in Eq.(2.1) drop out, and compare the results for  $\lambda_{\rm L}(T)$  across the SG transition with that in the zero-dimensional p-spin glass model<sup>31</sup>.

To diagnose the quantum chaos in the SY model, we consider two disorder-averaged four-point 'regularized' OTOCs<sup>3,6,12</sup> for the bosonic operators on a Schwinger Keldysh (SK) contour Fig. 4.1, with two forward and two backward branches, equally spaced in imaginary time

with separation  $\beta/4$ , namely

$$\begin{split} F_{1}(t_{1},t_{2}) &= \frac{1}{N^{2}M^{2}} \sum_{ij,\alpha\beta} \overline{\text{Tr}[yb_{i\alpha}^{\dagger}(t_{1})yb_{j\beta}^{\dagger}(0)yb_{i\alpha}(t_{2})yb_{j\beta}(0)]} \\ &\qquad \qquad (4.1a) \\ F_{2}(t_{1},t_{2}) &= \frac{1}{N^{2}M^{2}} \sum_{ij,\alpha\beta} \overline{\text{Tr}[yb_{i\alpha}(t_{1})yb_{j\beta}^{\dagger}(0)yb_{i\alpha}^{\dagger}(t_{2})yb_{j\beta}(0)]}, \\ &\qquad \qquad (4.1b) \end{split}$$

where  $y^4 = \exp(-\beta \mathcal{H})/\text{Tr}[\exp(-\beta \mathcal{H})]$ . In this work, we do not consider the OTOC of the physical spin operators  $S_{i\alpha\beta x}$ . Calculations of such an OTOC would involve 8-point bosonic correlation functions. We assume the maximum Lyapunov exponent in the large-N, M SY model to be the same for OTOCs for bosonic and spin operators.

The calculation of the OTOCs in Eq.(4.1) on the SK contour (Fig. 4.1) can be done in the replica diagonal PM phase following a procedure very similar to SYK and related models<sup>2,3,8</sup>. However below the dynamical transition temperature  $T_{SG}$ , in the SG phase, calculating any dynamic correlation function is more involved since the system always stays out of equilibrium<sup>63,71</sup>. Alternatively, as discussed in Ref.31 for p-spin glass, the OTOCs can be calculated in the marginal SG phase by replicated generating function<sup>31</sup>  $\mathbb{Z}^n$  on the four-branch contour [Fig. 4.1], i.e.,

$$\mathcal{Z}^{n} = \frac{1}{Z^{n}} \int \mathcal{D}(\bar{b}_{i\alpha}^{a}, b_{i\alpha}^{a}) \mathcal{D}\lambda_{i}^{a} \exp(iS)$$
 (4.2)

where a = 1, ..., n are the replica indices and the action

$$\begin{split} \mathrm{i}S = & \left[ \mathrm{i} \int_{\mathcal{C}} dz \sum_{\mathrm{i}\alpha a} \left\{ \bar{b}^a_{i\alpha} \mathrm{i} \partial_z b^a_{i\alpha} - \lambda^a_i \left( \bar{b}^a_{i\alpha}(z) b^a_{i\alpha}(z) - SM \right) \right\} \\ & - \frac{1}{\sqrt{MN}} \frac{1}{2!} \mathrm{i} \int_{\mathcal{C}} dz \sum_{ij,\alpha\beta,a} J_{ij} \bar{b}^a_{i\alpha}(z) b^a_{i\beta}(z) \bar{b}^a_{j\beta}(z) b^a_{j\alpha}(z) \right], \end{split}$$

where z is the complex time variable along the SK contour  $\mathcal{C}$  [Fig. 4.1]. Here Z is the equilibrium partition function which appears due to the initial thermal density matrix<sup>31</sup>. The initial density matrix corresponds to the replica symmetric saddle point above  $T_{SG}$ , and 1RSB marginal SG saddle point below  $T_{SG}$ , as discussed in Sec. 3 A. After performing the disorder averaging, we can take the  $n \to 0$  limit. The resulting time-translation invariant dynamical correlations and responses are identical to those obtained from the 1RSB saddle point solutions in the marginal SG phase. Using this replicated generating

function, the OTOC can now be obtained as,

$$F_{1}^{ab}(t_{1}, t_{2}) = \frac{1}{N^{2}M^{2}} \sum_{ij,\alpha\beta} \langle yb_{i\alpha4}^{\dagger a}(t_{1})yb_{j\beta3}^{\dagger a}(0)yb_{i\alpha2}^{b}(t_{2})yb_{j\beta1}^{b}(0) \rangle \quad (4.4a)$$

$$F_{2}^{ab}(t_{1}, t_{2}) = \frac{1}{N^{2}M^{2}} \sum_{ij,\alpha\beta} \langle yb_{i\alpha4}^{a}(t_{1})yb_{j\beta3}^{\dagger a}(0)yb_{i\alpha2}^{\dagger b}(t_{2})yb_{j\beta1}^{b}(0) \rangle \quad (4.4b)$$

where the superscripts 1, 2, 3, 4 denote the branch on the SK contour and the average  $\langle \dots \rangle$  is with respect to replicated generating function [Eq.(4.2)].

FIG. 4.2. Diagrammatic representation of the kernel equation in Eq.(4.6b) for  $\mathcal{O}(1/(NM))$  term,  $(\mathcal{F}_{1,2}^a)$  in the OTOC  $F_{1,2}^a(t_1,t_2)$ . The solid horizontal lines denote dressed retarded and advanced propagators  $G_R$ ,  $G_A$  and the vertical rung denotes the Wightmann correlations  $G_{lr}^+$  and  $G_{lr}^-$ . The dashed line represents disorder averaging. We suppress replica indices of the vertices in the ladder diagram to avoid cluttering.

The diagrams that contribute to the OTOC can be arranged in powers of 1/(NM).  $\mathcal{O}(1)$  diagrams are disconnected and do not contribute to the growth of OTOC. The contribution,  $\mathcal{F}_{\mu}^{aabb}$  at  $\mathcal{O}(1/(NM))$  form the ladder diagrams (Fig. 4.2) grow exponentially, i.e., for  $F_{\mu}^{aabb}(t_1,t_2)=F_{\mu}^{0aabb}(t_1,t_2)+(1/(NM))\mathcal{F}_{\mu}^{aabb}(t_1,t_2)+\mathcal{O}(1/(N^2M^2))$  ( $\mu=1,2$ ),  $\mathcal{F}_{\mu}\sim e^{\lambda_{\rm L}t}$ . The ladder diagrams can be written in the form of a Bethe-Salpeter-like equation, similar to SYK-type models<sup>6,8,72</sup>,

$$\begin{split} \mathcal{F}_{1}^{aabb}(t_{1},t_{2}) &= \sum_{c} \int_{t_{3},t_{4}} \left[ K_{11}^{aacc}(t_{1},t_{2},t_{3},t_{4}) \mathcal{F}_{1}^{ccbb}(t_{3},t_{4}) \right. \\ &+ K_{12}^{aacc}(t_{1},t_{2},t_{3},t_{4}) \mathcal{F}_{2}^{ccbb}(t_{3},t_{4}) \right] \qquad (4.5a) \\ \mathcal{F}_{2}^{aabb}(t_{1},t_{2}) &= \sum_{c} \int_{t_{3},t_{4}} \left[ K_{21}^{aacc}(t_{1},t_{2},t_{3},t_{4}) \mathcal{F}_{1}^{ccbb}(t_{3},t_{4}) \right. \\ &+ K_{22}^{aacc}(t_{1},t_{2},t_{3},t_{4}) \mathcal{F}_{2}^{ccbb}(t_{3},t_{4}) \right]. \qquad (4.5b) \end{split}$$

Here  $\int_{t_3,t_4} = \int dt_3 dt_4$ . In the chaotic growth regime,  $\lambda_{\rm L}^{-1} \lesssim t < \lambda_{\rm L}^{-1} \ln N$ , the propagators along the horizontal lines, from  $t_1$  to  $t_3$  and from  $t_2$  to  $t_4$  in Fig. 4.2 can be approximated by retarded and advanced propagators  $^{3,6,62}$ . Since the retarded and advanced propagators are replicadiagonal in both PM and SG phases, the kernel equations

become,

$$\mathcal{F}_{1}^{a}(t_{1}, t_{2}) = \int dt_{3}dt_{4}K_{11}^{a}(t_{1}, t_{2}, t_{3}, t_{4})\mathcal{F}_{1}^{a}(t_{3}, t_{4})$$

$$+ \int dt_{3}dt_{4}K_{12}^{a}(t_{1}, t_{2}, t_{3}, t_{4})\mathcal{F}_{2}^{a}(t_{3}, t_{4}) \qquad (4.6a)$$

$$\mathcal{F}_{2}^{a}(t_{1}, t_{2}) = \int dt_{3}dt_{4}K_{21}^{a}(t_{1}, t_{2}, t_{3}, t_{4})\mathcal{F}_{1}^{a}(t_{3}, t_{4})$$

$$+ \int dt_{3}dt_{4}K_{22}^{a}(t_{1}, t_{2}, t_{3}, t_{4})\mathcal{F}_{2}^{a}(t_{3}, t_{4}), \qquad (4.6b)$$

where  $\mathcal{F}_{\mu}^{a}(t_1, t_2) \equiv \mathcal{F}_{\mu}^{aaaa}(t_1, t_2)$  and the kernels are given by

$$\begin{split} K_{11}^{a}(t_{1},t_{2},t_{3},t_{4}) = & 2\tilde{J}^{2}G_{A}(t_{31})G_{R}(t_{24})G_{lr}^{+}(t_{43})G_{lr}^{-}(t_{34})\\ K_{12}^{a}(t_{1},t_{2},t_{3},t_{4}) = & \tilde{J}^{2}G_{A}(t_{31})G_{R}(t_{24})G_{lr}^{+}(t_{43})G_{lr}^{+}(t_{43})\\ K_{21}^{a}(t_{1},t_{2},t_{3},t_{4}) = & \tilde{J}^{2}G_{R}(t_{13})G_{A}(t_{42})G_{lr}^{-}(t_{34})G_{lr}^{-}(t_{34})\\ K_{22}^{a}(t_{1},t_{2},t_{3},t_{4}) = & 2\tilde{J}^{2}G_{R}(t_{13})G_{A}(t_{42})G_{lr}^{-}(t_{34})G_{lr}^{+}(t_{43}). \end{split}$$

Here all the Green's functions are replica diagonal and  $G_{lr}^{\pm}(t)$  refers to Wightmann correlations functions (Appendix A 2).

Using the exponential ansatz in the growth regime,  $\mathcal{F}_{\mu}^{a}(t_{1},t_{2})=e^{\lambda_{L}(t_{1}+t_{2})/2}f_{\mu}^{a}(t_{1}-t_{2})$  and taking the Fourier transform of the kernel equations [Eq.(4.6)], leads to an eigenvalue equations in the frequency space with eigenvalue,  $\lambda_{e}=1$ . For the PM phase, these eigenvalue equa-

tions are

$$\tilde{J}^{2}G_{A}\left(-\omega - i\frac{\lambda_{L}}{2}\right)G_{R}\left(-\omega + i\frac{\lambda_{L}}{2}\right)\left[2\int d\omega' g_{1}(\omega - \omega')\right]$$

$$f_{1}^{a}(\omega') + \int d\omega' g_{2}(-(\omega - \omega'))f_{2}^{a}(\omega')\right] = f_{1}^{a}(\omega) \quad (4.8a)$$

$$\tilde{J}^{2}G_{R}\left(\omega + i\frac{\lambda_{L}}{2}\right)G_{A}\left(\omega - i\frac{\lambda_{L}}{2}\right)\left[\int d\omega' g_{2}(\omega - \omega')\right]$$

$$f_{1}^{a}(\omega') + 2\int d\omega' g_{1}(\omega - \omega')f_{2}^{a}(\omega')\right] = f_{2}^{a}(\omega) \quad (4.8b)$$

where

$$g_1(\omega) = \frac{1}{4\pi^2} \int d\omega_1 G_{lr}^+(\omega_1) G_{lr}^-(\omega + \omega_1)$$
$$= \frac{1}{4} \int d\omega_1 \frac{\rho(\omega_1)\rho(\omega + \omega_1)}{\sinh\frac{\beta\omega_1}{2}\sinh\frac{\beta(\omega + \omega_1)}{2}}$$
(4.9a)

$$g_2(\omega) = \frac{1}{4\pi^2} \int d\omega_1 G_{lr}^+(\omega_1) G_{lr}^+(\omega - \omega_1)$$
$$= \frac{1}{4} \int d\omega_1 \frac{\rho(\omega_1)\rho(\omega - \omega_1)}{\sinh\frac{\beta\omega_1}{2}\sinh\frac{\beta(\omega - \omega_1)}{2}}. \tag{4.9b}$$

Here  $\rho(\omega) = -(1/\pi) \text{Im} G_R(\omega)$  is the bosonic spectral function (Appendix A). We discretize the kernel equations [Eqs. (4.8)] in frequency and diagonalize the resulting matrix kernel for different trial values of  $\lambda_L$ . Eventually, the Lyapunov exponent  $\lambda_L$  is obtained from the one for which the kernel has at least one eigenvalue  $\lambda_e = 1$ .

Similarly, for the 1RSB SG phase, the kernel equations are

$$\begin{split} \tilde{J}^2 G_A \left( -\omega - \mathrm{i} \frac{\lambda_\mathrm{L}}{2} \right) G_R \left( -\omega + \mathrm{i} \frac{\lambda_\mathrm{L}}{2} \right) \left[ 2 \left( g^2 f_1^a(\omega) + g \int \frac{d\omega'}{2\pi} \tilde{g}_{lr}(\omega - \omega') f_1^a(\omega') + g \int \frac{d\omega'}{2\pi} \tilde{g}_{lr}(\omega' - \omega) f_1^a(\omega') \right. \\ \left. + \int d\omega' \tilde{g}_1(\omega - \omega') f_1^a(\omega') \right) + g^2 f_2^a(\omega) + 2g \int \frac{d\omega'}{2\pi} \tilde{g}_{lr}(-\omega + \omega') f_2^a(\omega') + \int d\omega' \tilde{g}_2(-\omega + \omega') f_2^a(\omega') \right] = f_1^a(\omega) \quad (4.10a) \\ \tilde{J}^2 G_A \left( \omega - \mathrm{i} \frac{\lambda_\mathrm{L}}{2} \right) G_R \left( \omega + \mathrm{i} \frac{\lambda_\mathrm{L}}{2} \right) \left[ 2 \left( g^2 f_2^a(\omega) + g \int \frac{d\omega'}{2\pi} \tilde{g}_{lr}(\omega - \omega') f_2^a(\omega') + g \int \frac{d\omega'}{2\pi} \tilde{g}_{lr}(\omega' - \omega) f_2^a(\omega') \right. \\ \left. + \int d\omega' \tilde{g}_1(\omega - \omega') f_2^a(\omega') \right) + g^2 f_1^a(\omega) + 2g \int \frac{d\omega'}{2\pi} \tilde{g}_{lr}(\omega - \omega') f_1^a(\omega') + \int d\omega' \tilde{g}_2(\omega - \omega') f_1^a(\omega') \right] = f_2^a(\omega) \quad (4.10b) \end{split}$$

In the above equations,  $\tilde{g}_1(\omega)$  and  $\tilde{g}_2(\omega)$  are the same as in Eqs. (4.9a), (4.9b) in terms of the SG spectral function  $\rho(\omega)$ . We obtain  $\lambda_{\rm L}$  as a function of S and T by separately solving Eq. (4.8) for the PM phase, and Eq. (4.10)] for the SG phase, with the retarded, advanced and Wightmann functions obtained from the large-N saddle point equations (Appendix A).

The crucial difference between the PM and SG phases is encoded in the ladder kernerl [Eq.(4.7)] through the

Wightmann correlators. In the SG phase,

$$G_{lr}^{+}(\omega) = G_{lr}^{-}(\omega) = 2\pi g \delta(\omega) \epsilon_{ab} + \pi \rho(\omega) / (\sinh(\beta \omega/2)) \delta_{ab}$$
(4.11)

In contrast, the delta function term is absent in the PM phase, where  $g = \sqrt{q_{\rm EA}} = 0$ .

We compare the results for  $\lambda_{L}(S,T)$  obtained using the above formalism for the zero-dimensional SY model with the  $\lambda_{L}(\Gamma,T)$  for the effective zero-dimensional p-




FIG. 4.3. Variation of Lyapunov exponent with S in the SY model.  $\lambda_{\rm L}(S)$  for  $T=0.1,\,T=0.8$  and T=2.0 (in units of  $\tilde{J}$ ), across the PM-SG phase transition marked by the vertical lines at  $S_{SG}\simeq 0.48,\,1.29,\,2.1$ , respectively.  $\lambda_{\rm L}$  has a maximum in the PM phase and changes non-monotonically.  $\lambda_{\rm L}\sim 1/S^{1.7}$  in the SG phase.

spin glass model. The OTOC formalism for the latter is very similar to that of SY model, but somewhat simpler. To extract  $\lambda_{\rm L}(\Gamma, T)$ , we compute the following regularized OTOC<sup>31</sup> for the *p*-spin glass model,

$$F(t_1, t_2) = \frac{1}{N^2} \sum_{i,j} \overline{\text{Tr} \Big[ y s_i(t_1) y s_j(0) y s_i(t_2) y s_j(0) \Big]},$$
(4.12)

We skip the details of the formalism here since those have been already described in Ref.31. We briefly discuss a few additional details for the non-local OTOC for p-spin glass chain model [Eq.(2.2)] in Sec. 5 along with those for the SY chain [Eq.(2.1)].

#### A. Lyapunov exponent across spin glass transition in the SY model

We obtain  $\lambda_{\rm L}(S,T)$  in the PM and SG phases by separately solving the corresponding large-N saddle-point equations (Sec. 3) for the two phases. Specifically, as shown in Fig. 3.1, we take cuts along several S and T, namely, from small S (low-T) quantum critical region a. S=0.5 (d. T=0.1), from intermediate-S (intermediate-T) crossover region between quantum critical and local moment behaviour b. S=1 (e. T=0.8), and from large-S (high-T) local moment region c. S=2 (f. T=2), as a function of T (S). We set  $\tilde{J}=1$  as the unit of temperature (energy). The results for  $\lambda_{\rm L}(S,T)$  are qualitatively similar for all the cuts, as we discuss below.

S dependence of Lyapunov exponent.—We first show the Lyapunov exponent as a function of S in Fig. 4.3

from PM to spin glass phase for three temperatures, T =0.1, 0.8, 2.0. In all the cases, the Lyapunov exponent  $\lambda_{\rm L}$ has a non-monotonic dependence on S within the PM phase;  $\lambda_{\rm L}$  initially increases with S till a value  $S_m(T)$ where  $\lambda_{\rm L}$  has a broad peak, and then it decreases with further increase of S approaching the critical value of S,  $S_{SG}(T)$  for the spin glass transition, which is obtained from the breakpoint criterion  $m(T, S_{SG}) = 1$ . The nonmonotonic  $\lambda_{L}(S)$  in the PM phase can be heuristically understood from the behaviour of the spectral function. As discussed in Appendix A 2, in our numerical solution of the real-frequency saddle-point equation [Eq. (3.3)], e.g., at T = 0.1, 0.8, for small  $S \simeq 0.05 - 0.2$ , the spectral function has a local moment like form [see Figs.A.1(a), A.2(a)], with a peak at  $\omega \sim \omega_0 \sim T \ln[(S+1)/S] > T$  and very little spectral weight around  $\omega = 0$ . The peak moves closer to zero energy with increasing S and the spectral density increases around  $\omega = 0$  as the spectral function resembles the spin-liquid solution [Eq.(A.14)] more. As a result,  $\lambda_L$  is small at small S and increases with S till a maximum value in the PM phase. However, beyond the maximum, closer to the SG transition, there is a loss of spectral weight at intermediate energies or pseudogap<sup>68</sup> like feature, in between a narrower low-energy peak and a broad peak at high energies. This presumably leads to decreases of  $\lambda_{\rm L}$  approaching the SG transition. Thus,  $\lambda_{\rm L}$  exhibits a non-monotonic dependence on S in the PM region.

In the PM phase, especially in the quantum critical region IV in Fig. 3.1, for  $S \to 0$ , we expect dominantly spin liquid behaviour and  $\lambda_{\rm L}(S \to 0, T) \to \lambda_{\rm L}^{\rm SYK}(T)$ , i.e.,  $\lambda_{\rm L}$  should asymptotically approach the T-dependent Lyapunov exponent<sup>8</sup>,  $\lambda_{\rm L}^{\rm SYK}(T)$ , of the SYK spin liquid<sup>40</sup> at S = 0. Based on this expectation we fit our data with  $\lambda_{\rm L}(S,T) \simeq \lambda_{\rm L}(0,T) + aS^{\alpha_{\lambda}}$  in the PM phase for small values of S, before reaching the peak [Fig. 4.3]. As shown in Fig. 4.4(a) for T = 0.8, 1.0, we find reasonable fit with  $\alpha_{\lambda} \simeq 0.5 - 0.6$ . However, due to this  $\sqrt{S}$  dependence, there is a rapid decrease of  $\lambda_{\rm L}(S)$  for  $S \lesssim 0.2$ , as seen in Fig. 4.4(a). As a result, the extrapolated  $\lambda_{\rm L}(0,T)$  for  $S\to 0$  from the fit consistently turns out to be very low, and much smaller than  $\lambda_{L}^{SYK}(T)^{8}$ , even at low temperature  $T \simeq 0.1$ . Nevertheless, our real frequency saddle-point solutions, e.g., the computed bosonic spectral function, for  $S \leq 0.05$  at low T are consistent with the conformal spin liquid solutions  $^{38,40,42}$  for  $|\omega|, T \ll J$  as well as the numerical spin liquid solutions obtained in Ref. 68, as discussed in Appendix A 2. Thus, our results indicate strong S (and T) dependent corrections to the conformal limit of the spectral function, and, consequently,  $\lambda_{\rm L}=2\pi T$ , in the spin liquid region, even for small  $S \lesssim 0.05$ . We also note that our numerical saddle-point solutions at small S smoothly evolves to the solutions at higher S, as evident from the smooth evolution of  $\lambda_L(S)$  in Figs. 4.3, 4.4(a) for all T.

The Lyapunov exponent appears to have weak discontinuity/change of slope at  $S_{SG}$  for  $T=0.1,\ 2.0$ , which, however, could be a numerical artifact due to problem in




FIG. 4.4. S dependence of  $\lambda_{\rm L}$  in PM and SG phases of SY model.  $\lambda_{\rm L}$  is shown as a function of S. (a) In the PM phase, before reaching the maximum [Fig. 4.3],  $\lambda_{\rm L}$  varies as,  $\lambda_{\rm L}(S,T) \simeq \lambda_{\rm L}(0,T) + aS^{\alpha_{\lambda}}$ , shown as line fit to the data (circles) for T=0.8,2.0 with  $\alpha_{\lambda}\approx 0.5,0.6$ , respectively. (b)  $\lambda_{\rm L}$  decays with S as a power law in the SG phase, shown as straight line fit to the data (circles) on log-log scale.  $\lambda_{\rm L}\sim 1/S^{1.7}$  for T=0.8,2.0.

obtaining a converged solution of the saddle-point equations very close to the transition, as well as, due to difficulty in matching the PM and 1RSB SG solutions approaching from two different sides of the SG transition. We find  $\lambda_{L}(S)$  to be continuous for T = 0.8 across the SG transition. In the SG phase,  $\lambda_{L}(S)$  decreases monotonically as  $\sim 1/S^{1.7}$ , as shown in Fig. 4.4(b). The reduction of  $\lambda_{\rm L}$  with increasing S or its increase with decreasing S (increasing quantum fluctuations) can be understood qualitatively by drawing analogy with the similar dependence of  $\lambda_{\rm L}$  on  $\Gamma$  in the p-spin glass model. In the latter case, the kernel equations for  $\lambda_{\rm L}$  can be mapped to a solvable one-dimensional Schrödinger equation<sup>31</sup>. As a result, it can be shown by that  $\lambda_{\rm L} \sim T^2/q_{\rm EA}^{3/2}$ .  $\lambda_{\rm L}$  increases with increasing quantum fluctuations in the SG phase due to the reduction of EA order parameter. Similar relation between  $\lambda_{\rm L}$  and  $q_{\rm EA}$  can be expected for the SY model, however, the kernel equations [Eq.(4.10)]are much more complicated in this case, compared to the p-spin glass model<sup>31</sup>, and we could not map them into a solvable quantum mechanical problem.

Temperature dependence of  $\lambda_{\rm L}$ .— We show the temperature dependence of  $\lambda_{\rm L}$  in Fig. 4.5 for three values of S, S=0.5, 1.0, 2.0. In all these case, in contrast to non-monotonic dependence of  $\lambda_{\rm L}$  on  $S, \lambda_{\rm L}$  monotonically increases with temperature throughout SG and PM phases, with possibly a weak slope change at the transition [Fig. 4.5 (inset)]. The dependence of  $\lambda_{\rm L}$  on temperature, in SG and PM phases is summarized below. Similar to the case of p-spin glass<sup>31</sup>,  $\lambda_{\rm L}$  follows a power-law T dependence in the SG phase for SY model. For large  $S, \lambda_{\rm L} \sim T^{\beta_{\lambda}}$ , where the exponent  $\beta_{\lambda} \approx 1.5$  weakly depends on S, as




FIG. 4.5. Temperature dependence of Lyapunov exponent in SY model.  $\lambda_{\rm L}$  as function of T for S=0.5, S=1.0 and S=2.0 across the SG to PM phase transitions, marked by the vertical lines at  $T_{SG}\simeq 0.12,\ 0.49,\ 1.9$ , respectively. The inset shows zoomed in view near the transitions for smaller S values.

shown in Fig. 4.6. For small S,  $\lambda_{\rm L} \sim T$ , i.e., has a linear T dependence [Fig. 4.7(a)], albeit in a narrow range of temperature accessible for  $T \lesssim T_{SG}$  for small S.

In the PM phase for  $T > T_{SG}$ , for all values of S,  $\lambda_{\rm L}$  increases near the transition logarithmically in T as,  $a \ln T + c$ , with constants a and c that depend on S [Fig. 4.7(b)].  $\lambda_{\rm L}(T)$  eventually tends to saturate at high temperature, as shown in Fig. 4.7. The logarithmic dependence of  $\lambda_{L}(T)$ , albeit over a small temperature window near  $T_{SG}$  in the PM phase, even for relatively large S=2, is somewhat surprising. For classical spin models<sup>56,57</sup>,  $\lambda_{\rm L} \sim \sqrt{T}$  is found at high temperatures, e.g., above thermal phase transitions. The  $\sqrt{T}$  behaviour is expected for classical models from fairly generally considerations<sup>73</sup>. In our case, the initial increase of  $\lambda_{\rm L}(T)$  for  $T \gtrsim T_{SG}$  cannot be fitted with a power law like  $\sqrt{T}$ . Nevertheless, qualitatively, the increase of  $\lambda_{\rm L}$  is expected with increasing T due to gradual unfreezing of spins going away from  $T_{SG}$ . As discussed in Appendix A2, near  $T_{SG}$  in the PM phase, the spectral function has a sharp peak close to  $\omega = 0$  separated from high-frequency spectral weight with a pseudogap-like dip at intermediate frequencies. The sharp peak can be understood as a relic of the delta function peak  $\sim \delta(\omega)$  that appears in correlation functions (see Appendix A2) in the SG phase, e.g., in the Wightmann correlation function [Eq.(4.11)]. The delta function peak indicates static order and spin freezing. With increasing T, the low- and high-frequency peaks move closer filling up the pseudogap at intermediate frequencies due to unfreezing of spins. This spectral weight transfer correlates with increase of  $\lambda_{\rm L}$  with T. Like in the SYK model<sup>8</sup>,  $\lambda_{\rm L}$  is expected to saturate eventually at high temperature since the model has a bounded spectrum. This is unlike the p-spin glass model which has a unbounded spectrum due to unbounded kinetic energy. The latter renders interaction effect negligible<sup>31</sup> at high temperature in the *p*-spin glass model leading to decrease of  $\lambda_{\rm L} \sim 1/T^2$  for  $T \gg \tilde{J}$ .




FIG. 4.6. Power-law temperature dependence of  $\lambda_{\rm L}$  in the SG phase for large S in SY model. The straight lines show the power-law fit,  $\lambda_{\rm L} \sim T^{\beta_{\lambda}}$  to the data (circles), on log-log plot with  $\beta_{\lambda} = 1.6, 1.5$  for S = 2.0, 3.0, respectively.




FIG. 4.7. Temperature dependence of  $\lambda_{\rm L}$  in PM and SG phases of SY model. (a)  $\lambda_{\rm L}$  increases linearly with temperature as  $\lambda_{\rm L} \sim T$  in the SG phase for small S, shown for S=0.5,1.0 as lines fitted to the data (circles) on log-log scale. (b) In the PM phase,  $\lambda_{\rm L}$  shows logarithmic dependence on temperature,  $a \ln T + c$  (line) initially before it saturates to a constant value for large T (Fig. 4.5), as shown for S=0.5,1.0,2.0 in semi-log plot ( $\lambda_{\rm L}$  vs.  $\ln T$ ).

The non-monotonic behaviour of  $\lambda_{\rm L}(S)$  [Fig. 4.5] in the PM phase of the SY model is similar to the non-monotonic behaviour seen<sup>31</sup> for  $\lambda_{\rm L}(\Gamma)$  or  $\lambda_{\rm L}(\hbar)$ , i.e., as a function of quantum parameter, in the *p*-spin glass model [Eq.(2.2)] as shown in Fig. 4.8. However, in the latter model, in contrast to the  $\lambda_{\rm L}(T)$  [Fig. 4.3] in the SY model,  $\lambda_{\rm L}$  also shows non-monotonic dependence on T




FIG. 4.8. Variation of Lyapunov exponent with quantum parameter  $\Gamma$  in the *p*-spin glass model.  $\lambda_{\rm L}$  as function of  $\Gamma$  for T=0.2 and T=0.4 across the PM-SG phase transition marked by the vertical lines at  $\Gamma_{SG}\simeq 2.65,\ 1.35$  respectively.  $\lambda_{\rm L}$  changes non-monotonically and has a maximum above the the glass transition in the PM phase<sup>31</sup>.




FIG. 4.9. Temperature dependence of  $\lambda_{\rm L}$  in the p-spin glass model.  $\lambda_{\rm L}$  as function of T for  $\Gamma=0.5$  and  $\Gamma=2.0$  across the PM-SG phase transition marked by the vertical lines at  $T_{SG}\sim0.3,0.51$  respectively.  $\lambda_{\rm L}$  changes non-monotonically and has a maximum above the glass transition in the PM phase<sup>31</sup>

in the PM phase, as in Fig. 4.9. The broad peak related to the non-monotonic  $\lambda_{\rm L}(\Gamma,T)$  in the PM phase of the p-spin glass model correlates with the onset of complex glassy relaxation<sup>31</sup>, where the dynamics starts getting dominated by saddle points of the glassy energy landscape<sup>35</sup>. In the classical limit ( $\Gamma \to 0$ ) of the p-spin glass model, the origin of the broad peak was shown to be related with the interplay of the rapid increase of relaxation time approaching the glass transition, and weakening of interaction effects due to unbounded kinetic energy growth with increasing temperature. The interplay leads

to a crossover from strong to weak chaos with increasing T, manifested as a broad peak in  $\lambda_{\rm L}$  coinciding with the onset of two-step glassy relaxation regime<sup>31</sup>. Due to bounded spectrum of the SY model, such crossover from strong to weak chaos with temperature is absent in the SY model, and the  $\lambda_{\rm L}(T)$  monotonically increases with T in the PM phase till it saturates at high temperature [Fig. 4.3]. However, as already discussed, a non-monotonic  $\lambda_{\rm L}(S)$  is still observed in the PM phase due to change of spectral properties as a function of S.




FIG. 4.10. Onset of glassy relaxation with S in SY model. Correlation function C(t) in the PM phase near the PM-SG phase transition for a fixed value T=0.8 and different values of S. As we approach the transition from the PM phase, near the transition at  $S_{SG} \simeq 1.3$ , the two step glassy relaxation is more pronounced.

The two-step glassy relaxation, seen for the p-spin glass model<sup>31</sup>, can also be observed in the PM phase of the SY model above the glass transition. To this end, we calculate the correlation function using the Fluctuation dissipation theorem,  $C(t) = -(\mathbf{i}/M) \sum_{\alpha} \langle [b_{i\alpha x}(t), b_{i\alpha x}^{\dagger}(0)] \rangle = 2 \int (d\omega/2\pi) e^{-\mathbf{i}\omega t} \coth(\beta\omega/2) \mathrm{Im} G_R(\omega)$ . We show the correlation function for several S in Fig. 4.10 at T=0.8, and for several T in Fig. 4.11 at S=1, above the SG transition. The gradual onset of the two-step relaxation is evident. This suggests that the onset of two-step glassy relaxation is not a sufficient condition to realize a peak in the Lyapunov exponent.

#### 5. BUTTERFLY VELOCITY IN QUANTUM SPIN GLASS CHAINS

In this section we study the spatial spreading of chaos, first in the SY model and then in the *p*-spin glass model.




FIG. 4.11. Onset of glassy relaxation with temperature in SY model. Correlation function C(t) in the PM phase near the PM-SG phase transition for a fixed value of S=1.0 and different values of T. The two step glassy relaxation is more pronounced as we approach  $T_{SG}=0.5$  from the PM phase.

#### A. Butterfly velocity across spin glass transition in the SY chain

In the previous section, we calculated  $\lambda_{\rm L}$  which determines the temporal growth of the four-point OTOC. Due to the spatial locality of our generalized 1D SY chain model [Eq.(2.1)], we can study the spatio-temporal dynamics of growth and spread of chaos. The quantity which measures the propensity of spatial spreading of chaos is the butterfly velocity,  $v_B$ . The operators that form large commutators with an initial operator lie inside a region of space-time that forms a light  $cone^{41,74}$ , signifying a ballistic spread of chaos. The speed or velocity that defines the boundary of the light cone is defined as  $v_B$ . Like the Lyapunov exponent  $\lambda_L$ , the butterfly speed  $v_B$  provides a state-dependent measure of operator growth and spreading, and can be related<sup>75</sup> to the well-known Lieb-Robinson velocity bound<sup>76</sup>, that gives state-independent measure in terms of operator norm. To this end, we calculate the following disorder-averaged regularized spatio-temporal OTOCs for the SY model,

$$F_{1,x}(t_1, t_2) = \frac{1}{N^2 M^2} \sum_{ij\alpha\beta} \overline{\text{Tr}[y b_{i\alpha,x}^{\dagger}(t_1) y b_{j\beta,0}^{\dagger}(0) y b_{i\alpha,x}(t_2) y b_{j\beta,0}(0)]}$$
(5.1a)

$$F_{2,x}(t_1, t_2) = \frac{1}{N^2 M^2} \sum_{ij\alpha\beta} \overline{\text{Tr}[yb_{i\alpha,x}(t_1)yb_{j\beta,0}^{\dagger}(0)yb_{i\alpha,x}^{\dagger}(t_2)yb_{j\beta,0}(0)]}$$

$$(5.1b)$$

As discussed in Sec. 4 for the case of the onsite OTOCs, which is equivalent to OTOCs in zero-dimensional SY

FIG. 5.1. Diagrammatic representation of the Kernel equation in Eq. 5.2 for  $\mathcal{O}(1/(NM))$  term,  $\mathcal{F}_{\mu,x}^a(t_1,t_2)$  in the OTOC  $F_{\mu,x}^a(t_1,t_2)$  ( $\mu=1,2$ ). The solid horizontal lines denote dressed retarded and advanced propagators  $G_R$ ,  $G_A$  and the vertical rung denotes the Wightmann correlations  $G_{lr}^+$  and  $G_{lr}^-$ . The dashed and dotted lines denote disorder averaging over J and J' couplings, respectively in Eq.(2.1). Here  $x'=x\pm 1$ . We suppress replica indices for the vertices to avoid cluttering.

model, the above non-local OTOCs are computed via a replicated Keldysh field theory, where the OTOCs become replica diagonal up to  $\mathcal{O}(1/(NM))$  diagrams. The latter are arranged as ladder series that can be written in the form of Bethe-Salpeter-like equations, as shown in Fig. 5.1,

$$\begin{split} \mathcal{F}^{a}_{1,x}(t_{1},t_{2}) &= \int dx' \int dt_{3}dt_{4} \bigg[ K^{a}_{11,xx'}(t_{1},t_{2},t_{3},t_{4}) \\ \mathcal{F}^{a}_{1,x'}(t_{3},t_{4}) &+ K^{a}_{12,xx'}(t_{1},t_{2},t_{3},t_{4}) \mathcal{F}^{a}_{2,x'}(t_{3},t_{4}) \bigg], \\ & \qquad \qquad (5.2a) \\ \mathcal{F}^{a}_{2,x}(t_{1},t_{2}) &= \int dx' \int dt_{3}dt_{4} \bigg[ K^{a}_{21,xx'}(t_{1},t_{2},t_{3},t_{4}) \\ \mathcal{F}^{a}_{1,x'}(t_{3},t_{4}) &+ K^{a}_{22,xx'}(t_{1},t_{2},t_{3},t_{4}) \mathcal{F}^{a}_{2,x'}(t_{3},t_{4}) \bigg]. \\ & \qquad \qquad (5.2b) \end{split}$$

In the above, the kernels, which are diagrammatically shown in Fig. 5.1, are given in Eqs. (5.3). Although we have not indicated explicitly, all the Green's functions appearing in the kernel elements are replica diagonal. Since the saddle-point Green's function (Sec. 3A) of the extended 1D SY chain model are local and independent of x, the kernel elements are translation invariant, as in Eqs. (5.3). Thus, after Fourier transforming Eqs. (5.2) to momentum space, we obtain equations for the Fourier transform of  $\mathcal{F}^a_{\mu,x}(t_1,t_2)$  ( $\mu=1,2$ ),  $\mathcal{F}^a_{\mu,p}(t_1,t_2)$ . The kernel elements for a given momentum p are obtained by replacing  $\tilde{J}^2$  in the local kernel elements of Eqs. (4.7) by momentum dependent prefactors, as shown in Eqs. (5.4), where  $\alpha = J'^2/\tilde{J}^2$ . In this section, as discussed below, we calculate butterfly velocity  $v_B$  using two methods -(1)a fully numerical method, which uses a straightforward generalization of the exponential growth ansatz for the local OTOC (Sec. 4) for each momentum mode; we refer to this as real p method, and (2) a semi-analytic method. called the *single mode ansatz*, following ref.41. Here we refer to this as imaginary p method.

$$K_{11,xx'}^{a}(t_{1},t_{2};t_{3},t_{4}) = G_{A}(t_{31})G_{R}(t_{24})G_{lr}^{+}(t_{43})G_{lr}^{-}(t_{34})\left(2J^{2}\delta_{x,x'} + J'^{2}(\delta_{x,x'} + \delta_{x\pm 1,x'})\right)$$
(5.3a)

$$K_{12,xx'}^{a}(t_1, t_2; t_3, t_4) = G_A(t_{31})G_R(t_{24})G_{lr}^{+}(t_{43})G_{lr}^{+}(t_{43})\left(J^2\delta_{x,x'} + J'^2(\delta_{x\pm 1,x'})\right)$$
(5.3b)

$$K_{21,xx'}^{a}(t_1,t_2;t_3,t_4) = G_R(t_{13})G_A(t_{42})G_{lr}^{-}(t_{34})G_{lr}^{-}(t_{34})\left(J^2\delta_{x,x'} + J'^2(\delta_{x\pm 1,x'})\right)$$
(5.3c)

$$K_{22,xx'}^{a}(t_1,t_2;t_3,t_4) = G_R(t_{13})G_A(t_{42})G_{lr}^{-}(t_{34})G_{lr}^{+}(t_{43})\left(2J^2\delta_{x,x'} + J'^2(\delta_{x,x'} + \delta_{x\pm 1,x'})\right)$$
(5.3d)

$$K_{11,p}^{a}(t_{1},t_{2};t_{3},t_{4}) = \tilde{J}^{2}\left(2 - \alpha(1-2\cos p)\right)G_{A}(t_{31})G_{R}(t_{24})G_{lr}^{+}(t_{43})G_{lr}^{-}(t_{34})$$
(5.4a)

$$K_{12,p}^{a}(t_1, t_2; t_3, t_4) = \tilde{J}^2 \Big( 1 - \alpha (1 - 2\cos p) \Big) G_A(t_{31}) G_R(t_{24}) G_{lr}^{+}(t_{43}) G_{lr}^{+}(t_{43})$$
 (5.4b)

$$K_{21,p}^{a}(t_{1},t_{2};t_{3},t_{4}) = \tilde{J}^{2} \Big( 1 - \alpha(1-2\cos p) \Big) G_{R}(t_{13}) G_{A}(t_{42}) G_{lr}^{-}(t_{34}) G_{lr}^{-}(t_{34})$$
 (5.4c)

$$K_{22,p}^{a}(t_1, t_2; t_3, t_4) = \tilde{J}^2 \Big( 2 - \alpha (1 - 2 \cos p) \Big) G_R(t_{13}) G_A(t_{42}) G_{lr}^{-}(t_{34}) G_{lr}^{+}(t_{43})$$
 (5.4d)




FIG. 5.2. The growing part of OTOC,  $\mathcal{F}_{1,x}(t)$ , is plotted on the x-t plane in logarithmic scale, as indicated in the color bar at T=0.8 and S=1.0 using the real p method.

Real p method.— As in Sec. 4, we take an exponential growth ansatz,  $\mathcal{F}_{\mu,p}(t_1, t_2) = e^{\lambda_L(p)(t_1 + t_2)/2} f_{\mu,p}^a(t_1 - t_2),$ and extract the momentum-dependent Lyapunov exponent  $\lambda_{\rm L}(p)$  by demanding that the matrix kernel has at least one eigenvalue equal to 1 (typically degenerate). We compute the eigenvector corresponding to the eigenvalue 1 to obtain  $f_{\mu,p}^a(\omega)$ . To extract  $v_B$ , we calculate  $\mathcal{F}_{\mu,p}^a(t,t) = e^{\lambda_L(p)t} f_{\mu,p}^a(0)$ , where  $f_{\mu,p}(0) =$  $\int (d\omega/2\pi) f_{\mu,p}(\omega)$ . We Fourier transform to the position space,  $\mathcal{F}_{\mu,x}(t) \equiv \mathcal{F}_{\mu,x}^a(t,t) = (1/L) \sum_p e^{ixp} \mathcal{F}_{\mu,p}(t,t)$  where L is the number of sites or the number of discrete p values taken within the Brillouin zone  $[-\pi, \pi]$  for unit lattice spacing. We show in Fig. 5.2 that  $\mathcal{F}_{1,x}(t)$  exhibits quite distinctive light cone. The magnitude of  $\mathcal{F}_{\mu,p}(t,t)$ cannot be fixed, i.e., the eigenvector can be multiplied by an arbitrary factor, for the kernel equations. Thus, we compute the light-cones using the locus of (x,t) points corresponding to  $\mathcal{F}_{1,x}(t) = \mathcal{F}_{th}$ , for various threshold values  $\mathcal{F}_{th}$ , for which  $t \propto x$ . We find that this ballistic feature is violated for too small or too large values of  $\mathcal{F}_{th}$ . We extract  $v_B(\mathcal{F}_{th})$  from the resulting light cones and find that  $v_B(\mathcal{F}_{th})$  converges approximately to the

same value  $v_B$  over a wide range of  $\mathcal{F}_{th}$ , as shown in Fig. 5.3. We discuss the results for  $v_B(S,T)$  obtained from the real p method below along with that from the imaginary p method.




FIG. 5.3. Light cones at T = 0.8 and S = 1.0 for different values of  $\mathcal{F}_{th}$  obtained using the real p method. The same butterfly speed  $v_B$  is obtained for a wide range of  $\mathcal{F}_{th}$ .

Imaginary p method.— To calculate butterfly velocity through this method, we use the single-mode ansatz for OTOC and the results obtained by Gu and Kitaev<sup>62</sup>. Using the single mode ansatz<sup>77</sup> for early time OTOC, they proved a ladder identity which implies the existence of a pole and a saddle at two imaginary values of momentum that contribute to the growth of OTOC. We refer to ref.62 for the proof of the ladder identity and the details of the method. Using this ansatz we can write the solution of Eqs. (5.2) for the 1D SY chain model [Eq. (2.1)],

$$\mathcal{F}_{\mu,p}^{a}(t_{1},t_{2}) \approx \frac{e^{\lambda_{L}(p)(t_{1}+t_{2})/2}}{C(p)} \Upsilon_{\mu,p}^{R}(t_{12}) \Upsilon_{\mu,p}^{A}(0) \quad (5.5)$$

where  $\Upsilon^R_{\mu,p}(t)$  and  $\Upsilon^A_{\mu,p}(t)$  are the retarded and advanced vertex functions<sup>62</sup> which modify the OTOC by an overall magnitude. The dominant p dependence of the above

comes from a pole  $\lambda_{\rm L}=2\pi T~(k_{\rm B}=1,~\hbar=1)$  in C(p), since  $C(p)\propto\cos[\lambda_{\rm L}(p)/4T]^{62}$ . This pole leads to the maximal chaos for non-Fermi liquids in SYK and related models<sup>41,52,62</sup>. The asymptotic form of Eq.(5.5) ignores the initial correlations and any non-linear effects<sup>62,77</sup> and is valid only at the butterfly or chaos front, i.e.,  $1/N\ll \mathcal{F}_{\mu,p}(t_1,t_2)/N\ll 1$ . Fourier transforming Eq.(5.5) back to real space, we obtain

$$\mathcal{F}_{\mu,x}(t) \equiv \mathcal{F}_{\mu,x}^{a}(t,t) \approx \int_{-\infty}^{\infty} \frac{dp}{2\pi} \frac{e^{\lambda_{L}(p)t + ipx}}{\cos(\lambda_{L}(p)/4T)} f(p),$$
(5.6)

where f(p), which depends on the kernel [Eqs. (5.4)] and vertex functions in Eq. (5.5), is assumed to be analytical function of p and non vanishing in the momentum regime of interest<sup>62</sup>.

As in the real p method discussed above, we choose a fixed, albeit arbitrary, value of  $\mathcal{F}_{\mu,x}(t,t) \approx 1$ , which is at the lower end of the butterfly front  $^{62}$ , to obtain the light cone from the resulting locus of (x,t) points and extract  $v_B$ . For large x and t, the integral in Eq.(5.6) can be estimated by saddle-point method. The saddle-point  $p_s$  for the integrand is obtained from  $\lambda'_{\rm L}(p_s)t+{\rm i}x=0$ , where  $\lambda'_{\rm L}(p)$  is the derivative. This together with the condition  $\mathcal{F}_{\mu,x}(t,t)\approx 1$ , implying  $\lambda_{\rm L}(p)t+{\rm i}px=0$ , leads to the following equation for  $p_s$ 

$$\lambda_{\rm L}'(p_s) = \frac{\lambda_{\rm L}(p_s)}{p_s},\tag{5.7}$$

and a corresponding velocity

$$v_s = \left(\frac{x}{t}\right)_s = i\lambda'_{\rm L}(p_s).$$
 (5.8)

The saddle point leads to a contribution  $\mathcal{F}^s_{\mu,x}(t) \sim \exp\left[\lambda_{\rm L}(p_s)t + {\rm i}p_sx\right]$ . Similarly, the pole  $p_1$  of the integrand in Eq.(5.6), such that  $\lambda_{\rm L}(p_1) = 2\pi T$ , gives a contribution  $\mathcal{F}^p_{\mu,x}(t) \sim \exp\left[2\pi T t + {\rm i}p_1x\right]$  with an associated velocity

$$v_m = i \frac{2\pi T}{p_1}. (5.9)$$

To give rise to a well defined light cone,  $\mathcal{F}_{\mu,x}^s(t)$  ( $\mathcal{F}_{\mu,x}^p(t)$ ) should be non-oscillatory in space and time, implying that  $p_s$  ( $p_1$ ) is purely imaginary, i.e.,  $p_s = \mathfrak{i}|p_s|$  ( $p_1 = \mathfrak{i}|p_1|$ ), and  $\lambda_L(\mathfrak{i}|p_s|)$  is purely real. In Appendix C, we show that this is indeed the case by solving for  $\lambda_L(p)$  for purely imaginary  $p = \mathfrak{i}|p|$  from Eqs.(5.2) using the exponential growth ansatz, as discussed earlier for the real p method.  $\lambda_L(\mathfrak{i}|p|)$  is found to be a convex function of |p| as expected<sup>62</sup>. The light cone and the associated  $v_B$  are determined by either the saddle-point or the pole contribution to  $\mathcal{F}_{\mu,x}(t)$  depending on whether  $|p_s| < |p_1|$  or  $|p_s| > |p_1|$ , respectively. The conditions also translate to  $v_s(\equiv v(p_s)) < v_1(\equiv v(p_1))$  or  $v_s > v_1$ , respectively, where the velocity  $v(p) = \mathfrak{i}\lambda'_L(p)$ , since  $\lambda_L(\mathfrak{i}|p|)$  is a convex (up) function of p (Appendix C).

For the condition  $|p_s| < |p_1|$ , the deformed contour, that goes through the saddle point  $\mathfrak{i}|p_s|$  on the imaginary axis in the evaluation of the integral in Eq.(5.6), does not enclose the pole and hence the latter does not contribute. As discussed in refs.52 and 62, this condition is realized when  $\lambda_L(0) \ll 2\pi T$ , i.e., when the chaos is far away from maximal. In this case  $\mathcal{F}_{\mu,x}(t) \approx \mathcal{F}^s_{\mu,x}(t)$  and  $v_B = v_s$ . By making a small p approximation,  $\lambda_L(p) \simeq \lambda_L(0) - \tilde{D}p^2$  (with  $\tilde{D} > 0$ ), we obtain  $v_B = v_s = [4\tilde{D}\lambda_L(0)]^{1/2}$  and

$$\mathcal{F}_{\mu,x}(t) \sim e^{\lambda_{\rm L}(0)t(1-x^2/v_B^2t^2)}$$
 (5.10)

Here the quantity  $\tilde{D} = v_B^2/4\lambda_{\rm L}(0)$  has the dimension of diffusion constant. For certain holographic theories<sup>59</sup> and strongly interacting diffusive metals built from SYK-type quantum dots<sup>41,52</sup>,  $4\tilde{D}$  has been found to be exactly equal to energy diffusion constant at low temperature.  $\tilde{D}$  has also been found to be same as spin diffusion constant at high temperature in a classical spin liquid<sup>56</sup>, and even for classical XY model<sup>57</sup>. Assuming that such relation between  $\tilde{D}$  and actual diffusion constant hold, at least, over a limited range of (S,T) in the phased diagram of SY chain, we use  $\tilde{D}$  as a proxy for diffusion constant and plot it as a function of S and T below.

For the other condition  $|p_s| > |p_1|$ , realized for  $\lambda_L(0) \approx 2\pi T$ , the pole contributes and dominates over the saddle-point contribution, such that

$$\mathcal{F}_{\mu,x}(t) \approx \mathcal{F}_{\mu,x}^{p}(t) \sim e^{2\pi T t(1-|x|/v_B t)}$$
 (5.11)

and  $v_B = v_m$ , as can be deduced from  $\mathcal{F}_{\mu,x}(t) \approx 1$ . The above corresponds to maximal chaos, realized for SYK-type non-Fermi liquids<sup>41,52,62</sup>.

For most part of the phase diagram [Fig. 3.1] we have studied, we see that saddle point,  $|p_s|$  remains well below the pole,  $|p_1|$  and so the OTOC receives dominant contribution from the saddle point. See Appendix C for more details. Therefore the chaos front travels with a speed  $v_B = \mathrm{i}\lambda_L'(\mathrm{i}|p_s|)$  and the Lyapunov exponent  $\lambda_L$  is non-maximal, as was verified through direct calculation of  $\lambda_L(S,T)$  in Sec. 4. The SY chain can be analytically shown to be maximally chaotic in the limit  $S \to 0$ ,  $T \to 0$  following the same methods<sup>3,6,8</sup> used for calculating  $\lambda_L(T)$  in the SYK model.

We calculate  $v_B$  as a function of T and S, across the SG-PM phase transition, for S=0.5,1.0 and T=0.1,0.8 along the a, b and d, e cuts, respectively, shown in Fig. 3.1, using both real and imaginary p methods. As discussed below, we find that the results from the two methods match closely, validating the particular form of single-mode ansatz in Eq.(5.5) even in the replicasymmetry broken spin glass phase.

S dependence of butterfly velocity.— Fig. 5.4 shows  $v_B$  as a function of S for T=0.1 and T=0.8, going from spin liquid and spin liquid-local moment crossover region to the SG phase. As in the case of  $\lambda_L(S)$  [Fig. 4.3],  $v_B(S)$  exhibits a broad maximum in the PM phase. However, unlike  $\lambda_L(S)$  it has a dip at the phase transition  $S_{SG}$  and then it increases monotonically with S,




FIG. 5.4. Variation of butterfly velocity with S in the SY model.  $v_B$  across the SG phase transition ( $S_{SG}$ , vertical dashed lines) as a function of S at T=0.1,0.8 (in units of  $\tilde{J}=1$ ) and  $\alpha=0.25$ , calculated using both real and imaginary p methods, as indicated in the legends. For the real p method, we have used  $\mathcal{F}_{th}=0.1$  for T=0.1 and  $\mathcal{F}_{th}=0.5$  for T=0.8, with ranges (error bars)  $\mathcal{F}_{th}=0.5-2.0$  and  $\mathcal{F}_{th}=0.1-1.0$ , respectively, for a lattice of L=50 sites.




FIG. 5.5. S dependence of  $v_B$  in the PM and SG phases of SY model. (a) The The data (circles) for  $v_B(S,T)$  in the PM phase, before it reaches the maximum [Fig. 5.4], is fitted with  $v_B(0,T)+aS^{\alpha_v}$  (line) for T=0.8 with  $\alpha_v\approx 0.5$ . (b) Exponential dependence of  $v_B$  on S in the SG phase is shown as straight line on a semi-log plot ( $\ln v_B$  vs. S) for T=0.8.

as the system becomes more classical. Such dip in  $v_B$  at the phase transition has been also seen for classical spin models<sup>57,58</sup>, implying that the speed of information scrambling becomes slower at the transition in all these cases.

We find  $v_B$  to have power-law dependence,  $v_B(S,T) - v_B(0,T) \sim \sqrt{S}$  for small S, similar to  $\lambda_L$  [Fig. 4.4(a)]




FIG. 5.6.  $\tilde{D} \sim v_B^2/\lambda_{\rm L}$  across the phase transition  $(S_{SG},$  vertical dashed lines) as a function of S calculated for T=0.1 and T=0.8.




FIG. 5.7. Temperature dependence of butterfly velocity in SY model.  $v_B$  across the phase transition ( $T_{SG}$ , vertical dashed lines) as a function of temperature at S=0.5, 1.0,  $\alpha=0.25$ , using both real and imaginary p methods (see main text). The parameters and the conventions are same as those described in the caption of Fig. 5.4.

with a very small extrapolated  $v_B(0,T)\approx 0$  in the limit  $S\to 0$ , as shown in Fig. 5.5(a) for T=0.8. Again, this is related to the local moment like spectra at small  $S\lesssim 0.2$  [Figs.A.1(a), A.2(a)]. On the other hand,  $v_B$  increases exponentially with S in the SG phase as shown in Fig. 5.5(b), in contrast to power-law decay of  $\lambda_{\rm L}(S)$  [ig. 4.4(b)]. We also plot the proxy for the diffusion constant  $\tilde{D}\sim v_B^2/\lambda_{\rm L}$  as a function of S in Fig. 5.6.  $\tilde{D}$  increases monotonically with S from the PM to SG phase across the phase transition.

Temperature dependence of butterfly velocity.—  $v_B$  is shown as a function of T for S=1.0 across the PM-SG transition. We show  $v_B(T)$  for S=0.5 only in the PM phase, as we did compute  $v_B$  for the narrow temperature range  $T < T_{SG}$  in the SG phase for S=0.5. The butterfly velocity monotonically increases with temperature through  $T_{SG}$ , saturating at high temperature, similar to




FIG. 5.8. Temperature dependence of  $v_B$  in the SG and PM phases of SY model. (a)  $v_B(S,T)$  (circles) in the SG phase is fitted with  $v_B(S,T) \simeq v_B(S,0) + aT^{\beta_v}$  (line) for S=1.0 where  $v_B(S,0) \simeq 0.46$  and  $\beta_v \simeq 1.6$ . (b) Logarithmic fit (line) to  $v_B(T)$  in the PM phase,  $v_B \sim a \ln T + c$  is shown on a semi-log plot  $(v_B \text{ vs. } \ln T)$  for S=1.0.




FIG. 5.9.  $\tilde{D} \sim v_B^2/\lambda_{\rm L}$  as a function of temperature for S=0.5 and S=1.0 in SY model. The SG transition temperatures are shown by vertical dashed lines.

 $\lambda_{\rm L}(T)$  [Fig. 4.5]. As seen in Fig. 5.7,  $v_B$  approaches a non-zero value for  $T \to 0$  in the SG phase. This implies a finite speed  $v_B(S,0)$  of information scrambling at T=0 in the SG phase of the SY chain. As shown in Fig. 5.8(a),  $v_B(T) - v_B(S, 0) \sim T^{1.6}$ , i.e.,  $v_B$  increases with temperature as a power law from the T=0 value. Like the  $\lambda_{\rm L}(T)$  [Fig. 5.8(b)], the T dependence of  $v_B$  is logarithmic,  $\sim a \ln T + c$ , for  $T \gtrsim T_{SG}$  in the PM phase [Fig. 5.8(b)]. The proxy diffusion constant  $\tilde{D} \sim v_B^2/\lambda_L$ is plotted in Fig. 5.9. For  $S=1.0,\,\tilde{D}$  approaches a large value at low temperature  $T \ll T_{SG}$  in the SG phase, since  $v(S,T) \to v_B(S,0) \neq 0$  and  $\lambda_L \sim T$  for  $T \to 0$ . In the SG phase one naively expects the transport to slow down and D to decrease with temperature. Thus, it appears that a system made out of spin glass quantum dots, like in the SY chain [Eq. (2.1)], can lead to unusual enhanced

diffusion at low temperature. However, this needs to be verified by calculating actual diffusion coefficient D, instead of  $\tilde{D}$ . The latter also has dip at the transition and saturates at high temperature in the PM phase.

FIG. 5.10. Diagrammatic representation of the Kernel equation in Eq. 5.13 for  $\mathcal{O}(1/N)$  term,  $\mathcal{F}_x^a$  in the OTOC  $F_x^a(t_1,t_2)$ . The solid horizontal lines denote dressed retarded and advanced propagators  $G_R$ ,  $G_A$  and the vertical rung denotes the Wightmann correlation  $G_{lr}$  (Appendix B). The dashed and dotted lines denote disorder averaging over J and  $J^+$  couplings in Eq.( 2.2). Here  $x' = x \pm 1$ .

#### B. Butterfly velocity across spin glass transition in the p-spin glass chain

To compute the  $v_B(\Gamma, T)$  in the *p*-spin glass chain [Eq.(2.2)], we use methods similar to that for the SY chain discussed above. We calculate the following regularized OTOC defined as,

$$F_x(t_1, t_2) = \frac{1}{N^2} \sum_{i,j} \overline{\text{Tr} \left[ y s_{i,x}(t_1) y s_{j,0}(0) y s_{i,x}(t_2) y s_{j,0}(0) \right]}$$
(5.12)

Following the calculations discussed earlier for the SY chain, the diagrams that contribute to the OTOC can be grouped in powers of 1/N. The leading order  $\mathcal{O}(1)$  diagrams are disconnected and do not contribute to the growth. At the next leading order  $\mathcal{O}(1/N)$  we have "ladder" diagrams which contribute to the growth of OTOC in the scrambling regime,  $\lambda_{\rm L}^{-1} \lesssim t \lesssim \lambda_{\rm L}^{-1} \ln N$ . As earlier, these ladder diagrams can be written in the form of a Bethe-Salpeter equation shown in Fig. 5.10, i.e.

$$\mathcal{F}_{x}^{a}(t_{1}, t_{2}) = \int dx' \int dt_{3} dt_{4} K_{xx'}^{a}(t_{1}, t_{2}, t_{3}, t_{4}) \mathcal{F}_{x'}^{a}(t_{3}, t_{4}),$$
(5.13)

where the kernel is given by

$$K_{xx'}^{a}(t_1, t_2, t_3, t_4) = \left[ (3J^2 + J'^2)\delta_{x'x} + 2J'^2\delta_{x', x\pm 1} \right]$$
$$G_R(t_{13})G_R(t_{24})G_W(t_{34}) \quad (5.14)$$

After Fourier transforming  $\mathcal{F}_x^a(t_1, t_2)$  to momentum space, we obtain  $\mathcal{F}_p^a(t_1, t_2)$  and the Kernel

$$K_p^a(t_1, t_2, t_3, t_4) = \tilde{J}^2[3 + 2\alpha(2\cos p - 1)]$$
 (5.15)

where we have used,  $J^2 + {J'}^2 = \tilde{J}^2$  and  $\alpha = {J'}^2/\tilde{J}^2$ .

To calculate the butterfly velocity we again use the real and imaginary p methods discussed for the SY chain above [Sec.  $5\,\mathrm{A}$ ]. In the imaginary p method, we employ the single mode ansatz

$$F_p^a(t_1, t_2) \approx \frac{e^{\lambda_L(p)(t_1 + t_2)/2}}{C(p)} \Upsilon_p^R(t_{12}) \Upsilon_p^A(0)$$
 (5.16)

We skip the details of the implementations of the two methods here, since they are similar to those for the SY chain.  $v_B$  is calculated across the spin glass phase transition line as a function of T and  $\Gamma$ .




FIG. 5.11. Variation of butterfly velocity with quantum parameter  $\Gamma$  in the p-spin glass model.  $v_B$  across the p-spin glass phase transition as a function of  $\Gamma$  at T=0.4,  $\alpha=0.25$ . In the real p calculation  $v_B$  is calculated from the slope of light cone for  $\mathcal{F}_{th}=0.5$ , for a lattice of L=50 sites (see Sec. 5 A). The error bars in the real p results indicate variation of  $v_B$  over a range of threshold values  $\mathcal{F}_{th}=0.1-1.0$ .

 $\Gamma$  dependence of butterfly velocity.— Fig. 5.11 shows the butterfly velocity as a function of the quantum fluctuation parameter  $\Gamma$  for T=0.4 (in units of  $\tilde{J}=1$ ) (y cut in Fig. 3.2). Unlike the SY model [Fig. 5.4],  $v_B$ does not have a dip at the transition as function of the quantum parameter. In the PM phase,  $v_B$  exhibits nonmonotonic dependence on  $\Gamma$  with the broad maximum above the glass transition at  $\Gamma = \Gamma_{SG}$ , as in the case of  $\lambda_{\rm L}(\Gamma)$  [Fig. 4.8]. The butterfly velocity follows a power law,  $v_B(\Gamma, T) - v_B(0, T) \sim \sqrt{\Gamma}$ , with the quantum fluctuation parameter  $\Gamma$  for small  $\Gamma < \Gamma_{SG}$  with extrapolated  $v_B(0,T) \approx 0$  in the  $\Gamma \to 0$  limit, deep in the SG phase [Fig. 5.12(a)]. This is unlike the exponential dependence of  $v_B$  on quantum/classical parameter S for the SY chain [Fig. 4.4(b)].  $v_B$  varies almost linearly with  $\Gamma$ [Fig. 5.12(b)] in the PM phase close to the transition  $(\Gamma \gtrsim \Gamma_{SG})$  before reaching the maximum in Fig. 5.11. We also plot the proxy for diffusion constant  $\tilde{D} \sim v_B^2/\lambda_L$ in Fig. 5.13 for T=0.4.  $\tilde{D}$  is very small for small  $\Gamma$ and then rises to a peak around the transition  $\Gamma_{SG}$ , with




FIG. 5.12.  $\Gamma$  dependence of butterfly velocity in the SG and PM phases of p-spin glass model.  $v_B(\Gamma,T)$  (circles) is fitted (a) in the SG phase with  $v_B \simeq v_B(0,T) + a\Gamma^{\alpha_v}$  (line), where  $\alpha_v \simeq 0.5$  and  $v_B(0,T) \simeq 0.005$  for T=0.4, and (b) in the PM phase with  $v_B \sim \Gamma^{0.9}$  (line), shown on a log-log plot for T=0.4.




FIG. 5.13.  $\tilde{D} \sim v_B^2/\lambda_{\rm L}$  across the SG phase transition as a function of  $\Gamma$  calculated for T=0.4

further non-monotonic dependence on  $\Gamma$  in the PM phase for large  $\Gamma > \Gamma_{SG}$ .

Temperature dependence of butterfly velocity.— We demonstrate the overall temperature dependence of the butterfly velocity going from the SG to the PM phase through the transition at  $T_{SG}$  for  $\Gamma=2.0$  in Fig. 5.14.  $v_B(T)$  exhibits a broad peak in the PM phase, just like  $\lambda_{\rm L}(T)$  in Fig. 4.9. Figs. 5.15(a),(b) show that the butterfly velocity follows  $v_B(\Gamma,T)-v_B(\Gamma,0)\sim T^{1.5}$ , with substantial extrapolated zero-temperature value  $v_B(\Gamma,0)$  in the SG phase, and logarithmic T dependence in the PM phase for  $T\gtrsim T_{SG}$ , as in the SY model [Fig. 5.8]. The variation of resultant  $\tilde{D}\sim v_B^2/\lambda_{\rm L}$  with temperature [Fig. 5.16] is also very similar to that for the SY model, except, for the former there is no dip of  $\tilde{D}$  at the transition, unlike the dip in  $\tilde{D}$  for SY model, shown in Fig. 5.9.




FIG. 5.14. Temperature dependence of butterfly velocity in p-spin model.  $v_B$  across the phase transition as a function of temperature at  $\Gamma=2.0$ ,  $\alpha=0.25$ . The imaginary momentum calculation uses the single mode ansatz. In the real momentum calculation  $v_B$  is calculated from the edge of the light cone for values of  $\mathcal{F}_{th}=0.5$ , in a lattice of L=50 sites. Variation in  $v_B$  from nearby values of  $\mathcal{F}_{th}=0.1$  and  $\mathcal{F}_{th}=1.0$  is shown as error bar.




FIG. 5.15. Temperature dependence of  $v_B$  in PM and SG phases of p-spin glass model.. (a)  $v_B$  vs. T (circles) in SG phase is fitted as  $v_B \simeq v_B(\Gamma,0) + aT^{\beta_v}$  (line) for  $\Gamma = 2.0$ , where  $\beta_v \simeq 1.45$  and  $v_B(\Gamma,0) \simeq 0.43$ . (b) Logarithmic dependence of  $v_B$  on T (circles) in PM phase, shown as a straight line fit on a semi-log plot for  $\Gamma = 2.0$ .

#### 6. DISCUSSION

In this work, we have studied how quantum and thermal fluctuations influence propagation of quantum information, as diagnosed by butterfly velocity, along with chaotic growth rate characterized by a Lyapunov exponent, through one-dimensional generalizations of two paradigmatic zero-dimensional quantum spin glass mod-




FIG. 5.16.  $\tilde{D} \sim v_B^2/\lambda_{\rm L}$  across the phase transition as a function of T calculated for  $\Gamma = 2.0$ . in p-spin glass model

els, Sachdev-Ye and p-spin glass models. These lattice models allow us to map out variations of  $\lambda_{\rm L}$  and  $v_B$  with temperature and quantum parameter deep inside the phases and near the non-trivial replica symmetry breaking spin glass transitions, as well as through crossovers between competing paramagnetic phases, e.g., spin liquid and local moment behaviours, and across the SG transition. We find distinct evolution of glassy complexity, indicated by maximum, or lack thereof, in the Lyapunov exponent and butterfly velocity in the PM phase, as a function quantum and thermal fluctuations in the SY model. The maximum is seen only as a function of quantum parameter S, around the onset of two-step glassy relaxation, away from the glass transition from the PM side. The maximum is absent when the glass transition is approached by decreasing T. In contrast, the maximum occurs as function of both T and the quantum parameter  $\Gamma$  in the PM phase of p-spin glass. The maximum corresponds to intensified chaos due to maximal complexity arising from the sampling of exponentially large number of saddle points of the underlying glassy landscape. In the semiclassical limit, the maximal number of saddle points leads to extreme sensitivity to the initial condition<sup>35</sup> or chaos.

Our works needs to be extended in future in several directions. To better understand connection between glassy complexity  $^{35,37}$  and quantum many-body chaos, it will be worthwhile to compute the complexity for the SY model using both semiclassical methods and quantum Thouless-Anderson-Palmer (TAP) equations and quantum Thouless-Anderson-Palmer (TAP) equations as have been done for quantum p-spin glass. The asymptotic temperature dependence of  $\lambda_{\rm L}$  and  $v_B$  we obtain deep in the SG phase of SY and p-spin glass implies a finite  $v_B$  for the T=0 SG ground state, and, consequently, lead to rapid increase of the quantity  $\tilde{D}\sim v_B^2/\lambda_{\rm L}$  for  $T\to 0$ . As  $\tilde{D}$  often acts as a proxy to diffusion constant, it will be interesting to probe the possibility of such anomalously large diffusion through actual transport calculations in a chain of spin glass quantum dots.

Our real-frequency saddle-point calculations of  $\lambda_{\rm L}$  and

 $v_B$ , and corresponding bosonic spectral functions, in the SY model indicate subtle competition between spin liquid and local moment solutions, as well as strong S and T dependent corrections to the conformal solution for the SYK-like spin liquid maximal chaotic behaviour, even at small values of  $S \lesssim 0.05$  and low temperature  $T \sim 0.1$ . This implies intriguing crossover in the small S region towards the expected maximally chaotic SYK spin liquid for  $S \to 0$ . We have not studied this crossover and the relative stability and intricate competition between spin liquid and local moment solutions in the small Sregion in much detail in this work. It will be important to study the chaos and spectral properties in the small-Sregime in future work. Finally, it will be an exciting research direction to compute entanglement properties in various crossover regions and across the replica symmetry breaking spin glass transition in these large-N models following methods<sup>79,80</sup> similar to those applied for calculating entanglement entropy in lattice of SYK dots and in Hubbard model within dynamical mean field theory.

#### ACKNOWLDEGEMENTS

VL acknowledges support from CSIR, India and is grateful to the workstation provided by CHEP on which most of calculations in this work are done. SB acknowledges support from SERB (Grant No. CRG/2022/001062), DST, India, and QuST project of the DST, Govt. of India.

- \* venkatakumar@iisc.ac.in; surajit@iisc.ac.in; sumi lan@iisc.ac.in
- Yasuhiro Sekino and L. Susskind, "Fast scramblers," Journal of High Energy Physics 2008, 065 (2008), arXiv:0808.2096 [hep-th].
- <sup>2</sup> Juan Maldacena, Stephen H. Shenker, and Douglas Stanford, "A bound on chaos," Journal of High Energy Physics 2016, 1–17 (2016).
- <sup>3</sup> A. Kitaev, "A simple model of quantum holography," Talks at KITP, April 7, 2015 and May 27, 2015.
- <sup>4</sup> Subhayan Sahu, Shenglong Xu, and Brian Swingle, "Scrambling dynamics across a thermalization-localization quantum phase transition," Phys. Rev. Lett. **123**, 165902 (2019).
- <sup>5</sup> Kristan Jensen, "Chaos in ads<sub>2</sub> holography," Phys. Rev. Lett. **117**, 111601 (2016).
- <sup>6</sup> Juan Maldacena and Douglas Stanford, "Remarks on the sachdev-ye-kitaev model," Phys. Rev. D 94, 106002 (2016).
- Aavishkar A. Patel and Subir Sachdev, "Quantum chaos on a critical fermi surface," Proceedings of the National Academy of Sciences 114, 1844–1849 (2017), https://www.pnas.org/doi/pdf/10.1073/pnas.1618185114.
- Sumilan Banerjee and Ehud Altman, "Solvable model for a dynamical quantum phase transition from fast to slow scrambling," Phys. Rev. B 95, 134302 (2017).
- <sup>9</sup> Debanjan Chowdhury, Antoine Georges, Olivier Parcollet, and Subir Sachdev, "Sachdev-ye-kitaev models and beyond: Window into non-fermi liquids," Rev. Mod. Phys. 94, 035004 (2022).
- Mark Srednicki, "Chaos and quantum thermalization," Phys. Rev. E 50, 888–901 (1994).
- Chaitanya Murthy and Mark Srednicki, "Bounds on chaos from the eigenstate thermalization hypothesis," Phys. Rev. Lett. 123, 230606 (2019).
- <sup>12</sup> Igor L. Aleiner, Lara Faoro, and Lev B. Ioffe, "Microscopic model of quantum butterfly effect: Out-of-time-order correlators and traveling combustion waves," Annals of Physics 375, 378–406 (2016).
- Douglas Stanford, "Many-body chaos at weak coupling," Journal of High Energy Physics 2016, 9 (2016).
- <sup>14</sup> Efim B. Rozenbaum, Sriram Ganeshan, and Victor Galitski, "Lyapunov exponent and out-of-time-ordered corre-

- lator's growth rate in a chaotic system," Phys. Rev. Lett. 118, 086801 (2017).
- Thomas Scaffidi and Ehud Altman, "Chaos in a classical limit of the Sachdev-Ye-Kitaev model," Phys. Rev. B 100, 155128 (2019).
- Tianrui Xu, Thomas Scaffidi, and Xiangyu Cao, "Does scrambling equal chaos?" Phys. Rev. Lett. 124, 140602 (2020).
- W. Kob, "The Mode-Coupling Theory of the Glass Transition," arXiv e-prints, cond-mat/9702073 (1997), arXiv:cond-mat/9702073 [cond-mat.stat-mech].
- Pablo G. Debenedetti and Frank H. Stillinger, "Supercooled liquids and the glass transition," Nature 410, 259–267 (2001).
- Ludovic Berthier and Giulio Biroli, "Theoretical perspective on the glass transition and amorphous materials," Rev. Mod. Phys. 83, 587–645 (2011).
- Stephan Büchner and Andreas Heuer, "Potential energy landscape of a model glass former: Thermodynamics, anharmonicities, and finite size effects," Phys. Rev. E 60, 6507-6518 (1999).
- <sup>21</sup> Srikanth Sastry, "The relationship between fragility, configurational entropy and the potential energy landscape of glass-forming liquids," Nature (London) **409**, 164–167 (2001), arXiv:cond-mat/0011317 [cond-mat.stat-mech].
- Francesco Sciortino, "Potential energy landscape description of supercooled liquids and glasses," Journal of Statistical Mechanics: Theory and Experiment 2005, P05015 (2005).
- <sup>23</sup> K. Binder and A. P. Young, "Spin glasses: Experimental facts, theoretical concepts, and open questions," Rev. Mod. Phys. 58, 801–976 (1986).
- <sup>24</sup> E. Vincent, "Ageing, rejuvenation and memory: The example of spin-glasses," in *Ageing and the Glass Transition*, edited by Malte Henkel, Michel Pleimling, and Roland Sanctuary (Springer Berlin Heidelberg, Berlin, Heidelberg, 2007) pp. 7–60.
- <sup>25</sup> Jennifer A. Hodgdon and Frank H. Stillinger, "Stokeseinstein violation in glass-forming liquids," Phys. Rev. E 48, 207–213 (1993).
- <sup>26</sup> Smarajit Karmakar, Chandan Dasgupta, and Srikanth Sastry, "Analysis of dynamic heterogeneity in a glass for-

- mer from the spatial correlations of mobility," Phys. Rev. Lett. 105, 015701 (2010).
- <sup>27</sup> S. F. Edwards and P. W. Anderson, "Theory of spin glasses," Journal of Physics F Metal Physics 5, 965–974 (1975).
- <sup>28</sup> Marc Mezard, Giorgio Parisi, and Miguel Angel Virasoro, Spin Glass Theory and Beyond: AN Introduction to the Replica Method and its Applications (1987).
- <sup>29</sup> G. Parisi, "Infinite number of order parameters for spinglasses," Phys. Rev. Lett. 43, 1754–1756 (1979).
- Tommaso Castellani and Andrea Cavagna, "Spin-glass theory for pedestrians," Journal of Statistical Mechanics: Theory and Experiment 2005, P05012 (2005).
- <sup>31</sup> Surajit Bera, K. Y. Venkata Lokesh, and Sumilan Banerjee, "Quantum-to-classical crossover in many-body chaos and scrambling from relaxation in a glass," Phys. Rev. Lett. 128, 115302 (2022).
- <sup>32</sup> Th. M. Nieuwenhuizen, "Exactly solvable model of a quantum spin glass," Phys. Rev. Lett. **74**, 4289–4292 (1995).
- Theo M. Nieuwenhuizen and Felix Ritort, "Quantum phase transition in spin glasses with multi-spin interactions," Physica A Statistical Mechanics and its Applications 250, 8–45 (1998), arXiv:cond-mat/9706244 [cond-mat.dis-nn].
- <sup>34</sup> Leticia F. Cugliandolo, Daniel R. Grempel, and Constantino A. da Silva Santos, "From second to first order transitions in a disordered quantum magnet," Phys. Rev. Lett. 85, 2589–2592 (2000).
- <sup>35</sup> Lorenzo Correale, Anatoli Polkovnikov, Marco Schirò, and Alessandro Silva, "Probing chaos in the spherical p-spin glass model," arXiv e-prints, arXiv:2303.15393 (2023), arXiv:2303.15393 [cond-mat.dis-nn].
- <sup>36</sup> Antonio Auffinger and Gerard Ben Arous, "Complexity of random smooth functions on the high-dimensional sphere," arXiv e-prints, arXiv:1110.5872 (2011), arXiv:1110.5872 [math.PR].
- <sup>37</sup> Tarek Anous and Felix M Haehl, "The quantum p-spin glass model: a user manual for holographers," Journal of Statistical Mechanics: Theory and Experiment 2021, 113101 (2021).
- <sup>38</sup> Subir Sachdev and Jinwu Ye, "Gapless spin-fluid ground state in a random quantum heisenberg magnet," Phys. Rev. Lett. **70**, 3339–3342 (1993).
- <sup>39</sup> Antoine Georges, Olivier Parcollet, and Subir Sachdev, "Mean field theory of a quantum heisenberg spin glass," Phys. Rev. Lett. 85, 840–843 (2000).
- <sup>40</sup> A. Georges, O. Parcollet, and S. Sachdev, "Quantum fluctuations of a nearly critical heisenberg spin glass," Phys. Rev. B 63, 134406 (2001).
- <sup>41</sup> Yingfei Gu, Xiao-Liang Qi, and Douglas Stanford, "Local criticality, diffusion and chaos in generalized sachdev-yekitaev models," Journal of High Energy Physics 2017, 125 (2017).
- <sup>42</sup> Subir Sachdev, "Bekenstein-hawking entropy and strange metals," Phys. Rev. X 5, 041025 (2015).
- <sup>43</sup> Xue-Yang Song, Chao-Ming Jian, and Leon Balents, "Strongly correlated metal built from sachdev-ye-kitaev models," Phys. Rev. Lett. 119, 216601 (2017).
- <sup>44</sup> Xue-Yang Song, Chao-Ming Jian, and Leon Balents, "Strongly correlated metal built from sachdev-ye-kitaev models," Phys. Rev. Lett. 119, 216601 (2017).
- <sup>45</sup> Richard A. Davison, Wenbo Fu, Antoine Georges, Yingfei Gu, Kristan Jensen, and Subir Sachdev, "Thermoelectric transport in disordered metals without quasiparticles: The sachdev-ye-kitaev models and holography," Phys. Rev. B

- **95**, 155131 (2017).
- <sup>46</sup> Chao-Ming Jian, Zhen Bi, and Cenke Xu, "Model for continuous thermal metal to insulator transition," Phys. Rev. B 96, 115122 (2017).
- <sup>47</sup> Shao-Kai Jian and Hong Yao, "Solvable sachdev-ye-kitaev models in higher dimensions: From diffusion to many-body localization," Phys. Rev. Lett. 119, 206602 (2017).
- <sup>48</sup> Micha Berkooz, Prithvi Narayan, Moshe Rozali, and Joan Simón, "Higher dimensional generalizations of the SYK model," Journal of High Energy Physics 2017, 138 (2017), arXiv:1610.02422 [hep-th].
- <sup>49</sup> Pengfei Zhang, "Dispersive sachdev-ye-kitaev model: Band structure and quantum chaos," Phys. Rev. B 96, 205138 (2017).
- Debanjan Chowdhury, Yochai Werman, Erez Berg, and T. Senthil, "Translationally invariant non-fermi-liquid metals with critical fermi surfaces: Solvable models," Phys. Rev. X 8, 031024 (2018).
- Arijit Haldar, Sumilan Banerjee, and Vijay B. Shenoy, "Higher-dimensional sachdev-ye-kitaev non-fermi liquids at lifshitz transitions," Phys. Rev. B 97, 241106 (2018).
- <sup>52</sup> Haoyu Guo, Yingfei Gu, and Subir Sachdev, "Transport and chaos in lattice sachdev-ye-kitaev models," Phys. Rev. B 100, 045140 (2019).
- <sup>53</sup> Debanjan Chowdhury and Brian Swingle, "Onset of many-body chaos in the o(n) model," Phys. Rev. D **96**, 065005 (2017).
- <sup>54</sup> Subhayan Sahu and Brian Swingle, "Information scrambling at finite temperature in local quantum systems," Phys. Rev. B **102**, 184303 (2020).
- Aavishkar A. Patel, Debanjan Chowdhury, Subir Sachdev, and Brian Swingle, "Quantum butterfly effect in weakly interacting diffusive metals," Phys. Rev. X 7, 031047 (2017).
- Thomas Bilitewski, Subhro Bhattacharjee, and Roderich Moessner, "Temperature dependence of the butterfly effect in a classical many-body system," Phys. Rev. Lett. 121, 250602 (2018)
- <sup>57</sup> Sibaram Ruidas and Sumilan Banerjee, "Many-body chaos and anomalous diffusion across thermal phase transitions in two dimensions," SciPost Physics 11, 087 (2021), arXiv:2007.12708 [cond-mat.stat-mech].
- <sup>58</sup> Thomas Bilitewski, Subhro Bhattacharjee, and Roderich Moessner, "Classical many-body chaos with and without quasiparticles," Phys. Rev. B 103, 174302 (2021).
- Mike Blake, "Universal diffusion in incoherent black holes," Phys. Rev. D 94, 086014 (2016).
- <sup>60</sup> Brian Swingle and Debanjan Chowdhury, "Slow scrambling in disordered quantum systems," Phys. Rev. B 95, 060201 (2017).
- <sup>61</sup> Bo-Bo Wei, Gaoyong Sun, and Myung-Joong Hwang, "Dynamical scaling laws of out-of-time-ordered correlators," Phys. Rev. B 100, 195107 (2019).
- <sup>62</sup> Yingfei Gu and Alexei Kitaev, "On the relation between the magnitude and exponent of OTOCs," JHEP **02**, 075 (2019), arXiv:1812.00120 [hep-th].
- <sup>63</sup> Giulio Biroli and Olivier Parcollet, "Out-of-equilibrium dynamics of a quantum heisenberg spin glass," Phys. Rev. B 65, 094414 (2002).
- <sup>64</sup> D. R. Grempel and M. J. Rozenberg, "Fluctuations in a quantum random heisenberg paramagnet," Phys. Rev. Lett. 80, 389–392 (1998).
- <sup>65</sup> Leticia F. Cugliandolo, D. R. Grempel, and Constantino A. da Silva Santos, "Imaginary-time replica formalism study of a quantum spherical p-spin-glass model,"

Phys. Rev. B 64, 014403 (2001).

- <sup>66</sup> W. Götze, Complex Dynamics of Glass-Forming Liquids (Oxford University Press, Oxford, 2008).
- <sup>67</sup> David R. Reichman and Patrick Charbonneau, "Modecoupling theory," Journal of Statistical Mechanics: Theory and Experiment 2005, 05013 (2005), arXiv:condmat/0511407 [cond-mat.soft].
- Alberto Camjayi and Marcelo J. Rozenberg, "Quantum and thermal fluctuations in the su(n) heisenberg spin-glass model near the quantum critical point," Phys. Rev. Lett. 90, 217202 (2003).
- Maine Christos, Felix M. Haehl, and Subir Sachdev, "Spin liquid to spin glass crossover in the random quantum heisenberg magnet," Phys. Rev. B 105, 085120 (2022).
- Tiliana Arrachea and Marcelo J. Rozenberg, "Infinite-range quantum random heisenberg magnet," Phys. Rev. B 65, 224430 (2002).
- <sup>71</sup> Leticia F. Cugliandolo and Gustavo Lozano, "Real-time nonequilibrium dynamics of quantum glassy systems," Phys. Rev. B **59**, 915–942 (1999).
- Yingfei Gu, Alexei Kitaev, Subir Sachdev, and Grigory Tarnopolsky, "Notes on the complex Sachdev-Ye-Kitaev model," JHEP 02, 157 (2020), arXiv:1910.14099 [hep-th].
- Jorge Kurchan, "Quantum bound to chaos and the semiclassical limit," arXiv e-prints, arXiv:1612.01278 (2016), arXiv:1612.01278 [cond-mat.stat-mech].
- <sup>74</sup> Vedika Khemani, David A. Huse, and Adam Nahum, "Velocity-dependent lyapunov exponents in many-body quantum, semiclassical, and classical chaos," Phys. Rev. B 98, 144304 (2018).
- Daniel A. Roberts and Brian Swingle, "Lieb-robinson bound and the butterfly effect in quantum field theories," Phys. Rev. Lett. 117, 091602 (2016).
- <sup>76</sup> Elliott H. Lieb and Derek W. Robinson, "The finite group velocity of quantum spin systems," Communications in Mathematical Physics 28, 251–257 (1972).
- Alexei Kitaev and S. Josephine Suh, "The soft mode in the Sachdev-Ye-Kitaev model and its gravity dual," JHEP 05, 183 (2018), arXiv:1711.08467 [hep-th].
- <sup>78</sup> Giulio Biroli and Leticia F. Cugliandolo, "Quantum thouless-anderson-palmer equations for glassy systems," Phys. Rev. B 64, 014206 (2001).
- Arijit Haldar, Surajit Bera, and Sumilan Banerjee, "Rényi entanglement entropy of fermi and non-fermi liquids: Sachdev-ye-kitaev model and dynamical mean field theories," Phys. Rev. Res. 2, 033505 (2020).
- Surajit Bera, Arijit Haldar, and Sumilan Banerjee, "Dynamical mean-field theory for Rényi entanglement entropy and mutual Information in Hubbard Model," arXiv e-prints, arXiv:2302.10940 (2023), arXiv:2302.10940 [cond-mat.str-el].
- 81 Olivier Parcollet, Antoine Georges, Gabriel Kotliar, and Anirvan Sengupta, "Overscreened multichannel SU(n) kondo model: Large-n solution and conformal field theory," Phys. Rev. B 58, 3794–3813 (1998).

# Appendix A: Imaginary-time path integral and saddle-point equations for SY chain

We write the partition function  $\mathcal{Z}(\{J_{ij,x},J'_{ij,x}\})= \operatorname{Tr}[\exp(-\beta\mathcal{H})]$   $(\beta=1/T)$  of the disordered model of Eq.(2.1) as an imaginary-time path integral using the bosonic representation  $S_{i\alpha\beta x}=b^{\dagger}_{i\alpha x}b_{i\beta x}-S\delta_{\alpha\beta}$  with the constraint  $\sum_{\alpha}b^{\dagger}_{i\alpha x}b_{i\alpha x}=SM$   $(S\geq0)$  on the number of bosons at each i,x, to fix the spin of the bosonic representation to S. Here the (real) random couplings  $\{J_{ij,x},J'_{ij,x}\}$  are drawn from Gaussian distributions with zero mean and variances  $J^2$  and  $J'^2$ , respectively. We calculate disorder-averaged quantities by introducing replicas  $a=1,\ldots,n$  to obtain  $\overline{\mathcal{Z}^n}=\int \mathcal{D}(\bar{b},b)\mathcal{D}\lambda \exp[-S_{eff}],$  where

$$S_{eff} = \int_{0}^{\beta} d\tau \sum_{ix} \left[ \bar{b}_{i\alpha x}^{a} \partial_{\tau} b_{i\alpha x}^{a} - \lambda_{ix}^{a}(\tau) (\bar{b}_{i\alpha x}^{a} b_{i\alpha x}^{a} - SM) \right]$$

$$-\frac{1}{2MN} \int_{0}^{\beta} d\tau d\tau' \left[ \frac{J^{2}}{2} \sum_{i < j, x} S_{i\alpha \beta x}^{a}(\tau) S_{i\gamma \delta x}^{b}(\tau') S_{j\beta \alpha x}^{a}(\tau) \right]$$

$$S_{j\delta \gamma x}^{b}(\tau') + J'^{2} \sum_{i,j,x} S_{i\alpha \beta x}^{a}(\tau) S_{i\gamma \delta x}^{b}(\tau') S_{j\beta \alpha, x+1}^{a}(\tau) S_{j\delta \gamma, x+1}^{b}(\tau')$$

$$(A.1)$$

Here the field  $\lambda^a_{ix}(\tau)$  imposes the constraint on boson number and sum over repeated  $\alpha,\beta,\ldots$  and a,b indices are assumed. In the  $N\to\infty$  limit, introducing the Hubbard-Stratonovich field  $Q^{ab}_{\alpha\beta\gamma\delta,x}(\tau,\tau')$ , and its conjugate  $\Pi^{ab}_{\alpha\beta\gamma\delta,x}(\tau,\tau')$  to impose  $Q^{ab}_{\alpha\beta\gamma\delta,x}(\tau,\tau')=(1/N)\sum_i S^a_{i\alpha\beta x}(\tau)S^b_{i\gamma\delta x}(\tau')$ , we decouple the quartic terms above. Moreover, assuming  $\mathrm{SU}(M)$  symmetry, i.e.,  $Q^{ab}_{\alpha\beta\gamma\delta,x}(\tau,\tau')=Q^{ab}_{x}(\tau,\tau')\delta_{\alpha\gamma}\delta_{\beta\delta}$ , and similarly for  $\Pi$ , we obtain  $\overline{\mathcal{Z}^n}=\int \mathcal{D}Q\mathcal{D}\Pi\mathcal{D}(\bar{b},b)\mathcal{D}\lambda \exp[-S_{eff}]$  and the action

$$S_{eff} = \int_0^\beta d\tau \sum_{ix} \left[ \bar{b}_{i\alpha x}^a \partial_\tau b_{i\alpha x}^a - \lambda_{ix}^a(\tau) (\bar{b}_{i\alpha x}^a b_{i\alpha x}^a - SM) \right]$$

$$- MN \int_0^\beta d\tau d\tau' Q_x^{ab}(\tau, \tau') \left[ \frac{J^2}{4} Q_x^{ab}(\tau, \tau') + \frac{J'^2}{2} Q_{x+1}^{ab}(\tau, \tau') \right]$$

$$+ MN \int_0^\beta d\tau d\tau' \Pi_x^{ab}(\tau', \tau) \left[ Q_x^{ab}(\tau, \tau') - \frac{1}{NM^2} \sum_i \bar{b}_{i\alpha x}^a(\tau) b_{i\alpha x}^b(\tau') b_{i\beta x}^a(\tau) \bar{b}_{i\beta x}^b(\tau') \right]$$

$$(A.2)$$

Introducing  $G_x^{ab}(\tau,\tau') = -(1/M) \sum_{\alpha} b_{i\alpha x}^a(\tau) \overline{b}_{i\alpha x}^b(\tau')$  and the conjugate field  $\Sigma_x^{ab}(\tau,\tau')$  we get  $\overline{Z}^n = \int \mathcal{D}(Q,\Pi,G,\Sigma,\lambda) \exp[-MNS_{eff}]$  with

$$S_{eff} = \sum_{x} \left[ \operatorname{Tr} \ln \left[ (\partial_{\tau} - \lambda_{x}^{a}(\tau)) \delta_{ab} \delta(\tau - \tau') + \Sigma_{x}^{ab}(\tau, \tau') \right] \right.$$

$$\left. - \int_{0}^{\beta} d\tau d\tau' Q_{x}^{ab}(\tau, \tau') \left( \frac{J^{2}}{4} Q_{x}^{ab}(\tau, \tau') + \frac{J'^{2}}{2} Q_{x+1}^{ab}(\tau, \tau') \right) \right.$$

$$\left. + \int_{0}^{\beta} d\tau d\tau' \Pi_{x}^{ab}(\tau', \tau) \left( Q_{x}^{ab}(\tau, \tau') - G_{x}^{ab}(\tau, \tau') G_{x}^{ba}(\tau', \tau) \right) \right.$$

$$\left. + \int_{0}^{\beta} d\tau d\tau' \Sigma_{x}^{ab}(\tau, \tau') G_{x}^{ba}(\tau', \tau) + S \sum_{a} \int_{0}^{\beta} d\tau \lambda_{x}^{a}(\tau) \right]$$

$$\left. - \left( A.3 \right) \right.$$

$$\left. \left. \left( A.3 \right) \right.$$

where we have imposed  $\lambda^a_{ix}(\tau) = \lambda^a_x(\tau)$ . In the large N,M limit, minimization of the above action with respect to  $Q,\Pi,G,\Sigma$  and  $\lambda$ , with the assumption of a time-translation invariant (equilibrium) saddle point, e.g.,  $Q^{ab}_x(\tau,\tau') = Q^{ab}_x(\tau-\tau')$  and  $\lambda^a_x(\tau) = \lambda^a_x$ , leads to the following self-consistency equations for the bosonic Green's function

$$\begin{split} [G_x^{-1}]^{ab}(\mathrm{i}\omega_k) &= (\mathrm{i}\omega_k + \lambda_x^a)\delta^{ab} - \Sigma_x^{ab}(\mathrm{i}\omega_k) \\ \Sigma_x^{ab}(\tau) &= J^2 G_x^{ab}(\tau) G_x^{ab}(\tau) G_x^{ab}(-\tau) \\ &+ J'^2 G_x^{ab}(\tau) G_{x+1}^{ab}(\tau) G_{x+1}^{ab}(-\tau) \\ G_x^{aa}(\tau = 0^-) &= -S. \end{split} \tag{A.4}$$

Here  $\omega_k = 2k\pi\beta$  is the bosonic Matsubara frequency with integer k. Assuming lattice translation invariance,  $G_x^{ab}(\tau) = G^{ab}(\tau)$  and  $\lambda_x^a = \lambda^a$ , the above saddle point is the same as that [Eq.(3.1)] for the zero dimensional model<sup>38,39</sup> with the renormalized coupling  $\tilde{J}^2 = J^2 + J'^2$ .

Saddle point equation in PM phase— In the PM phase the Green's function is replica diagonal and symmetric. Thus, the saddle point equations in Eq.( 3.1), after analytic continuation to real frequency, i.e.,  $i\omega_k \to \omega + i0^+$ , becomes

$$[G_R(\omega)]^{-1} = \omega - \lambda - \Sigma_R(\omega)$$

$$\Sigma(i\omega_k) = \int_0^\beta d\tau e^{i\omega_k t} \Sigma(\tau) = \frac{\tilde{J}^2}{\beta^2} \sum_{k_1, k_2} G(i\omega_{k_1}) G(i\omega_{k_2})$$

$$G(i\omega_{k_1} + i\omega_{k_2} - i\omega_k)$$
(A.5b)

Using the spectral representation  $G(i\omega_k) = \int d\omega \rho(\omega)/(i\omega_k - \omega)$  with  $\rho(\omega) = -(1/\pi) \text{Im} G_R(\omega)$  in Eq.A.5b gives,

$$\Sigma(\mathbf{i}\omega_k) = \tilde{J}^2 \int d\omega_1 d\omega_2 d\omega_3 \frac{\rho(\omega_1)\rho(\omega_2)\rho(\omega_3)}{\mathbf{i}\omega_k - \omega_1 - \omega_2 + \omega_3}$$

$$[n_B(\omega_1)n_B(\omega_2)n_B(-\omega_3) + n_B(-\omega_1)n_B(-\omega_2)n_B(\omega_3)], \tag{A.6}$$

where  $\lambda^a = \lambda$ .

Using the identity  $1/(\omega - \omega_1 - \omega_2 + \omega_3 + i0^+) = -i \int_0^\infty dt e^{i(\omega - \omega_1 - \omega_2 + \omega_3 + i0^+)}$  and analytic continuation

gives the retarded self-energy

$$\Sigma_R(\omega) = -i\tilde{J}^2 \int_0^\infty dt e^{i\omega t} \left[ n_1(t) n_1(t) n_2(t) + n_3(t) n_3(t) n_4(t) \right]$$
(A.7)

where  $n_1(t) = \int d\omega e^{-i\omega t} \rho(\omega) n_B(-\omega)$ ,  $n_2(t) = \int d\omega e^{i\omega t} \rho(\omega) n_B(\omega)$ ,  $n_3(t) = \int d\omega e^{-i\omega t} \rho(\omega) n_B(\omega)$ , and  $n_4(t) = \int d\omega e^{i\omega t} \rho(\omega) n_B(-\omega)$ .

Saddle point equation in SG phase— Using the 1-RSB ansatz in Eq. 3.4, the analytically continued saddle point equation for the regular part of the green's function is given by

$$[G_R(\omega)]^{-1} = \omega - \frac{\tilde{J}g}{\Theta_R} - [\Sigma_R(\omega) - \Sigma_R(\omega = 0)],$$
  
$$m\beta = \frac{1}{\tilde{J}g^2} \left(\frac{1}{\Theta_R} - \Theta_R\right)$$
 (A.8)

Using the similar steps for analytic continuation as in the PM phase, we get the retarded self energy from Eq.(3.7b),

$$\Sigma_R(\omega) = -i\tilde{J}^2 \int_0^\infty dt e^{i\omega t} \Big[ \Big( n_1^2(t) n_2(t) + n_3^2(t) n_4(t) \Big) - 2g \Big( n_1(t) n_2(t) - n_3(t) n_4(t) \Big) - g \Big( n_3^2(t) - n_1^2(t) \Big) \Big] + 2g^2 \tilde{J}^2 G_R(\omega) + g^2 \tilde{J}^2 G_R^*(-\omega)$$
(A.9)

where  $n_i$ , i = 1, 2, 3, 4 are same as those defined in the PM phase. The two equations above can be solved numerically similar to PM phase calculation, to obtain the retarded Green's function,  $G_R(\omega)$ .

#### 1. Numerical solution of the saddle point equations

Here we provide the details of the numerical implementation of the self-consistent equations in imaginary as well as in real frequency.

#### a. Real-frequency solution

PM phase – The retarded Green function in PM phase can be obtained in the frequency domain by self-consistently solving Eq. 3.3. We write the self-energy  $\Sigma_R(\omega)$  in a form convenient for numerical calculation using FFT, as in Eq.A.7. We start with an initial ansatz for  $G_R(\omega)$  on a discrete range  $\omega \in [-\omega_{max}, \omega_{max}]$  and an initial value of  $\lambda$  and calculate  $\Sigma_R(\omega)$  from it. At each iteration,  $[G_R(\omega)]^{-1}$  is updated as,

$$G_{R,j+1}^{-1}(\omega) = (1-y)\Big(\omega - \lambda - \Sigma_{R,j}(\omega)\Big) + yG_{R,j}^{-1}(\omega)$$
(A.10)

At each iteration, we monitor the error  $|[G_{R,j+1}(\omega)]^{-1}| - |[G_{R,j}(\omega)]^{-1}|$  and repeat the

iteration with the updated  $[G_{R,j+1}(\omega)]^{-1}$  until it converges. We then check the constraint,  $G(\tau=0^-)=-\int_{-\infty}^{\infty}d\omega\rho(\omega)/(e^{\beta\omega}-1)=-S$ . If it' is not met, we repeat the procedure with a different  $\lambda$ . The procedure is repeated until the constraint is met.

SG phase – The retarded Green function,  $G_R(\omega)$  in the SG phase can be obtained by solving Eq. 3 A 2. Similar to PM phase, the retarded self-energy  $\Sigma_R(\omega)$  is written in a form suitable for implementation using FFT. Starting with an initial ansatz and an inital value of m, we calculate  $\Sigma_R(\omega)$ .  $G_R(\omega)$  is updated at each iteration as

$$G_{R,j+1}^{-1}(\omega) = (1-y) \left( \omega - \frac{\tilde{J}g}{\Theta_R} - \left[ \Sigma_{R,j}(\omega) - \Sigma_{R,j}(\omega = 0) \right] \right) + yG_{R,j}^{-1}(\omega)$$
(A.11)

We repeat the iteration with the updated  $[G_{R,j+1}(\omega)]^{-1}$  until it converges and check the constraint  $-\int_{-\infty}^{\infty} d\omega \rho(\omega)/(e^{\beta\omega}-1)-g=-S$ . m is varied to satisfy the constraint.

#### b. Imaginary-time solution

PM phase –We solve the saddle point Eq. 3.2 iteratively to obtain  $G(\mathrm{i}\omega_k)$  for  $k\in[0,k_{max}]$ . We start with an initial guess solution which could be the non-interacting part of the Green function and an initial guess value of the Lagrange multiplier  $\lambda$  and calculate  $\Sigma(\tau)$  using Eq. 3.2b. We then Fourier transform  $G(\tau)$  using fast-Fourier transform(FFT) and obtain  $G(\mathrm{i}\omega_k)$  and  $\Sigma(\mathrm{i}\omega_k)$ . After jth iteration,  $G_i(\mathrm{i}\omega_k)$  is updated as,

$$G_{j+1}(i\omega_k) = \frac{1-y}{i\omega_k - \lambda - \Sigma_j(i\omega_k)} + yG_j(i\omega_k) \quad (A.12)$$

where  $y \in (0,1)$  is used to ensure smooth convergence, which we choose typically between 0.2 and 0.3. After each iteration we monitor the error  $e(i\omega_k) = |G_{i+1}(i\omega_k) - G_i(i\omega_k)|$  for each k by comparing with  $\epsilon = 10^{-9}$ , the tolerance. The procedure is repeated with updated  $G_{i+1}(i\omega_k)$  after Fourier transforming back to time domain until  $e(i\omega_k) < \epsilon$  for each k. After convergence, we check if the constraint  $G(\tau = 0^-) = -S$  is met for the current value of  $\lambda$ . If not we repeat the procedure again after varying  $\lambda$ . The iterations stop only after the constraint is met within a tolerance.

 $SG\ phase\ -$  In the SG phase we solve for the regular part of the Green function,  $\tilde{G}(i\omega_k)$  in Eq. 3.7a using the modified self-energy  $\tilde{\Sigma}(\tau)$  in Eq. 3.7b. The procedure to solve the saddle point equation in SG phase is the similar to PM phase with the modified constraint,  $\tilde{G}(\tau=0^-)=g-S$ .

Starting with an initial ansatz and an initial value of the breaking point, m we calculate the  $\tilde{\Sigma}(\tau)$  and g from

Eq. 3.8. We then Fourier transform  $\tilde{G}(\tau)$  using fast-Fourier transform(FFT) and obtain  $\tilde{G}(i\omega_k)$  and  $\tilde{\Sigma}(i\omega_k)$ . At each iteration,  $\tilde{G}_i(i\omega_k)$  is updated as

$$\tilde{G}_{j+1}(i\omega_k) = \frac{1 - y}{i\omega_k - \tilde{J}g/\Theta - [\tilde{\Sigma}_j(i\omega_k) - \tilde{\Sigma}(i\omega_k = 0)]} + y\tilde{G}_j(i\omega_k)$$
(A.13)

The procedure is repeated until convergence and then we check if the constraint is met with current value of m. If not, we vary the value of m and repeat the iteration procedure until the constraint is met.

## 2. Spectral and correlation function in the SY model

In the SY model, we look at the nature of spectrum as a function of thermal and quantum fluctuations. As summarized in Sec. 3A3, following earlier works<sup>40,68</sup>, there are three types of solutions of the saddle point equations of Sec. 3A, (1) spin liquid, (2) local moment, and (3) marginal spin glass. These solutions can be distinguished via the bosonic spectral function  $\rho(\omega) = -(1/\pi) \text{Im} G_R(\omega)$ . In the PM phase, the spin liquid solution resides in the quantum critical region IV, as shown in Fig. 3.1, extending all the way down to T = 0 for S = 0. For small  $S < S_{max} \approx 0.052$ , the zero-temperature bosonic Green's function for complex frequency z is given by  $G(z) \sim e^{-i(\theta+\pi/4)}/\sqrt{z}$ , with Imz > 0 and  $\pi/4 < \theta < 3\pi/4$ , where the spectral asymmetry  $\theta$  is related to spin S via  $\theta/4 + \sin(2\theta)/4 =$  $1/2 + S^{38,40}$ . This leads to a T = 0 spectral function,  $\rho(\omega) \sim -\operatorname{sgn}(\omega)/\sqrt{|\omega|}$ , with characteristic square root divergence at  $\omega \to 0$ , and a finite-T conformal spectral function<sup>81</sup> in region IV of Fig. 3.1,

$$\rho(\omega) \sim \frac{\sinh{(\omega/2T)}}{\sqrt{T}} \Gamma\left(\frac{1}{4} + \mathrm{i}\frac{\omega - \alpha T}{2\pi T}\right) \Gamma\left(\frac{1}{4} - \mathrm{i}\frac{\omega - \alpha T}{2\pi T}\right), \tag{A.14}$$

where  $\sin(\pi/4 + \theta) = 1/\sqrt{1 + e^{-2\alpha}}$ . At finite temperature, the above spectral function,  $\rho(\omega) \sim \omega$  for  $|\omega| \lesssim T$ , has a peak at  $|\omega| \sim \mathcal{O}(T)$ , and thereafter decay as  $1/\sqrt{|\omega|}$  for  $\mathcal{O}(T) \lesssim |\omega| \lesssim J$ .

The other PM solution, the local moment solution co-exist, and has intricate competition  $^{40}$  with the spin liquid solution in the quantum critical region IV (Fig. 3.1). As a result, it is not always easy to distinguish the spin liquid solution and the local moment solution for general (S,T) in the quantum critical region. Ref.68 has shown the existence of spin liquid solution for small S<0.5 using analytical continuation of the numerical solution of the imaginary-time saddle-point equations [Eqs.( 3.2)]. Our results are consistent with that of Ref.68. The local moment becomes the dominant solution above the quantum critical region  $T\gtrsim J,$  in the PM phase. At




FIG. A.1. Spectral function as a function of S at T=0.1 in (a) PM phase and (b) SG phase.

high temperature, the local moment solution has a characteristic peak at  $\omega \sim \omega_0 \sim T \ln{[(S+1)/S]^{40}}$ . At very low temperature, below  $T_{SG}$  (Fig. 3.1), where the local moment solution is metastable, its spectral function has two peaks at  $\sim \pm 1/T$  and a delta function peak with  $\rho(\omega) \sim S\omega\delta(\omega)^{40}$ . The low-temperature local moment solution is not relevant for us.

For the marginal spin glass solution, on the other hand, one again obtains a linear-in- $\omega$  behaviour of the spectral function, termed as pseudogap<sup>68</sup>, where  $\rho(\omega) \sim \omega/S$  for low frequency and low temperature  $T \ll T_{SG}$ , with a high-frequency broad peak. As T increases towards glass transition, a narrow low-frequency peak around  $|\omega| \sim T$ appears on top of the linear- $\omega$  pseudogap background. which fills up with increasing temperature. The narrow peak around  $\sim T$  appears from the loss of spectral weight from the  $\delta$ -function part in the correlation function, e.g., the Wightmann function of Eq. (4.11), in the SG phase. The  $\delta(\omega)$  part in the correlation functions corresponds to static order and its weight is transferred to the narrow peak in the spectral function  $\rho(\omega)$  due to the gradual unfreezing of spins approaching the SG transition. The broad high-frequency peak at  $\omega \sim \mathcal{O}(JS)$  originates from precession of frozen spin around their local fields<sup>68</sup>.

We have studied the evolution of the spectral function along the fixed-T and fixed-S cuts (Fig. 3.1) employed for our chaos calculations in Secs. 4, 5. We find crossover between the PM solutions and transition from the PM to SG solution. Figs.A.1(a) and (b) show the evolution of  $\rho(\omega)$  at low temperature T=0.1 with S(d) cut, Fig. 3.1) for the PM and SG phases, respectively. The result for  $S=0.04 < S_{max}$  in Fig.A.1(a) is consistent with a spin liquid solution of Eq.(A.14). Similarly,  $\rho(\omega)$  for  $S=0.2 > S_{max}$  also resembles a spin liquid solution, though no analytical form is known in this case. The spectral function acquires low-energy and




FIG. A.2. Spectral function as a function of S at T=0.8 in (a) PM phase and (b) SG phase.

high-energy peaks, with a dip in spectral weight in between, with further increase of S towards  $S_{SG} \simeq 0.48$ , as shown in Fig.A.1(a). In the SG phase [Fig.A.1(a)],  $\rho(\omega)$  follows the expected marginal SG behaviour discussed above. Qualitatively almost the same progression of  $\rho(\omega)$  with S is seen in Fig.A.2 for the higher T=0.8 (e cut in Fig. 3.1) with  $S_{SG} \simeq 1.29$ .

Similarly, for fixed-S cuts a (S=0.5) and b (S=1.0) in Fig. 3.1, the spectral function evolves from that of the SG solutions for  $T < T_{SG} \simeq 0.12$ , 0.49 in Figs.A.3(a),A.4(a) to the spin liquid and local moment spectral functions in Figs.A.3(b),A.4(b). The narrow peak close to the transition  $T \gtrsim T_{SG}$  at  $\omega \sim \mathcal{O}(T)$  results from the melting of static order in the SG phase, as discussed earlier.




FIG. A.3. Spectral function as a function of T at S=0.5 in (a) SG phase and (b) PM phase.




FIG. A.4. Spectral function as a function of T at S=1.0 in (a) SG phase and (b) PM phase.

Wightmann correlation functions— The correlation functions needed in the calculation of chaos can be obtained from the knowledge of the retarded Green's function or the spectral function discussed above. We calculate  $G(\tau)$  using the spectral function from

$$G(\tau) = \begin{cases} \int_{-\infty}^{\infty} d\omega \frac{e^{-\omega\tau}\rho(\omega)}{e^{-\beta\omega} - 1} & (\tau > 0) \\ -\int_{-\infty}^{\infty} d\omega \frac{e^{-\omega\tau}\rho(\omega)}{e^{\beta\omega} - 1} & (\tau < 0) \end{cases}$$
(A.15)

Analytically continuing  $G(\tau)$  to real time gives Wightmann correlation functions, in the PM phase,

$$G_{lr}^{+}(t) = -G(it + \beta/2) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{-i\omega t} \frac{\pi \rho(\omega)}{\sinh\frac{\beta\omega}{2}}$$
 (A.16)

$$G_{lr}^{-}(t) = -G(it - \beta/2) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{-i\omega t} \frac{\pi \rho(\omega)}{\sinh\frac{\beta\omega}{2}} \quad (A.17)$$

We notice that  $G_{lr}^+(t) = G_{lr}^-(t)$  and hence  $G_{lr}^+(\omega) = G_{lr}^-(\omega) = \pi \rho(\omega)/\sinh(\beta \omega/2)$ . Similarly, in the SG phase,

$$G_{lr}^{+,ab}(t) = G_{lr}^{-,ab}(t) = g\epsilon_{ab} + \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{-i\omega t} \frac{\pi\rho(\omega)}{\sinh\frac{\beta\omega}{2}} \delta_{ab},$$
(A.18)

that allows to define  $G_{lr}^{+,ab}(\omega)=G_{lr}^{-,ab}(\omega)=2\pi g\delta(\omega)\epsilon_{ab}+\pi\rho(\omega)/(\sinh(\beta\omega/2))\delta_{ab}$ . The delta function term proportional to the EA order parameter g at  $\omega=0$  in the Wightmann correlation function indicates the presence of static spin glass order. Similar delta function term appears in the Fourier transform of other correlation functions, e.g.,  $G_>^{aa}(t)=-(\mathrm{i}/M)\sum_{\alpha}\langle b_{i\alpha x}^a\bar{b}_{i\alpha x}^a\rangle$ , and similarly  $G_<^{aa}(t)$ . We note that, since retarded Green's function,  $G_R^{aa}(t)=\theta(t)[G_>^{aa}(t)-G_<^{aa}(t)]$   $[\theta(t)$  is the Heaviside function] and and  $G_R^{aa}(\omega)$  is obtained by analytically continuing  $\mathrm{i}\omega_k\to\omega+\mathrm{i}0^+$  in Eq.(3.4),  $G_R^{aa}(t)$  is free of any static part even in the SG phase.

## Appendix B: Imaginary-time path integral and saddle point equations in *p*-spin glass chain

As in the case of SY chain [Appendix A], we write the partition function  $\mathcal{Z}(\{J_{ijk,x},J_{ijk,x}^{\pm}\})=\operatorname{Tr}[\exp(-\beta\mathcal{H})]$  ( $\beta=1/T$ ) of the disordered model of Eq.( 2.2), where  $J_{ijk,x}$  and  $J_{ijk,x}^{\pm}$  are Gaussian random number with zero mean and variances  $3J^2/(2N^2)$  and  $3J'^2/2N^2$ , as an imaginary-time path integral. We calculate disorder-averaged quantities by introducing replicas  $a=1,\ldots,n$ , and a Lagrangian multiplier  $z(\tau)$  to impose the spherical constraint, to obtain  $\overline{\mathcal{Z}^n}=\int \mathcal{D} s\exp[-S_{eff}]$ , where

$$S_{eff} = \sum_{x} \left\{ \int_{\tau} \left[ \frac{1}{2\Gamma} \left( \frac{\partial s_{ix}^{a}}{\partial \tau} \right)^{2} + \frac{z^{a}(\tau)}{2} \left( (s_{ix}^{a})^{2} - N \right) \right] - \frac{1}{4} \int_{\tau,\tau'} \sum_{ab} \left[ \frac{J^{2}}{N^{2}} \left( \sum_{i} s_{ix}^{a}(\tau) s_{ix}^{b}(\tau') \right)^{3} + \frac{J'^{2}}{N^{2}} \left( \sum_{i} s_{ix}^{a}(\tau) s_{ix}^{b}(\tau') \right) \left( \sum_{i} s_{ix+1}^{a}(\tau) s_{ix+1}^{b}(\tau') \right)^{2} + \frac{J'^{2}}{N^{2}} \left( \sum_{i} s_{ix}^{a}(\tau) s_{ix}^{b}(\tau') \right)^{2} \left( \sum_{i} s_{ix+1}^{a}(\tau) s_{ix+1}^{b}(\tau') \right) \right] \right\}$$
(B.1)

Here  $\int_{\tau} = \int_{0}^{\beta} d\tau$ . We now introduce the large-N field Q and its conjugate  $\Sigma$ , where  $Q_{x}^{ab}(\tau, \tau') = \frac{1}{N} \sum_{i} s_{ix}^{a}(\tau) s_{ix}^{b}(\tau')$ , through the identity

$$1 = \int DQ_x^{ab}(\tau, \tau') \delta\left(NQ_x^{ab}(\tau, \tau') - \sum_i s_{ix}^a(\tau) s_{ix}^b(\tau')\right)$$

$$= \int DQ_x^{ab}(\tau, \tau') D\Sigma_x^{ab}(\tau, \tau') \exp\left\{-\frac{1}{2} \int_{\tau, \tau'} \Sigma_x^{ab}(\tau, \tau')\right\}$$

$$\left(NQ_x^{ab}(\tau, \tau') - \sum_i s_{ix}^a(\tau) s_{ix}^b(\tau')\right)$$
(B.2)

Inserting the identity and integrating over  $s_{ix}(\tau)$  in  $\mathbb{Z}^n$  leads to the effective action

$$S_{eff} = \sum_{x} \left\{ \frac{N}{2} \operatorname{Tr} \ln \left[ \left( -\frac{1}{\Gamma} \partial_{\tau}^{2} + z^{a}(\tau) \right) \delta(\tau - \tau') \delta_{ab} \right. \right.$$

$$\left. - \Sigma_{x}^{ab}(\tau, \tau') \right] - \frac{N}{2} \sum_{a} \int_{\tau} z^{a}(\tau)$$

$$\left. + \frac{N}{2} \sum_{ab} \int_{\tau, \tau'} \left[ \Sigma_{x}^{ab}(\tau, \tau') Q_{x}^{ab}(\tau, \tau') - \frac{J^{2}}{4} Q_{x}^{ab}(\tau, \tau')^{3} \right.$$

$$\left. - \frac{J'^{2}}{4} \left( Q_{x}^{ab}(\tau, \tau') Q_{x+1}^{ab}(\tau, \tau')^{2} + Q_{x}^{ab}(\tau, \tau')^{2} Q_{x+1}^{ab}(\tau, \tau') \right) \right] \right\}$$
(B.3)

In the limit  $N \to \infty$ , we can take the saddle point with respect to  $\Sigma_x^{ab}(\tau,\tau')$ . This gives  $\boldsymbol{Q}_x^{-1} = \boldsymbol{Q}_0^{-1} - \boldsymbol{\Sigma}$ , where the matrix  $\boldsymbol{Q}_0^{-1}$  is given by  $(Q_0^{-1})^{ab}(\tau,\tau') = \left(-\frac{1}{\Gamma}\partial_{\tau}^2 + \frac{1}{\Gamma}\partial_{\tau}^2 + \frac{1}{\Gamma$ 

(B.9b)

 $z^a(\tau)$ ) $\delta(\tau - \tau')\delta_{ab}$ . Using this we eliminate  $\Sigma_x^{ab}(\tau, \tau')$  from the action and obtain effective action in terms of Q,

$$\begin{split} S_{eff} &= -\sum_{x} \left\{ \frac{N}{2} \text{Tr} \ln Q_{x} - \frac{N}{2} \sum_{a} \int_{\tau} z^{a}(\tau) \Big( Q_{x}^{aa}(\tau, \tau) - 1 \Big) \right. \\ &+ N \sum_{a,b} \int_{0}^{\beta} d\tau d\tau' \Big( \frac{1}{2\Gamma} \partial_{\tau}^{2} Q_{x}^{ab}(\tau, \tau') \delta(\tau - \tau') \delta_{ab} \Big) \\ &+ \frac{J^{2}}{4} Q_{x}^{ab}(\tau, \tau')^{3} + \frac{J'^{2}}{4} Q_{x}^{ab}(\tau, \tau') Q_{x+1}^{ab}(\tau, \tau')^{2} \\ &+ \frac{J'^{2}}{4} Q_{x}^{ab}(\tau, \tau')^{2} Q_{x+1}^{ab}(\tau, \tau') \Big\} \end{split} \tag{B.4}$$

Taking the saddle point with respect to Q gives the Dyson equation,

$$(Q_x^{-1})^{ab}(\tau,\tau') = \left(-\frac{1}{\Gamma}\partial_{\tau}^2 + z_a(\tau)\right)\delta(\tau - \tau')\delta_{ab}$$
$$-\frac{3J^2}{2}Q_x^{ab}(\tau,\tau')^2 - \frac{J'^2}{2}Q_{x+1}^{ab}(\tau,\tau')^2$$
$$-J'^2Q_x^{ab}(\tau,\tau')Q_{x+1}^{ab}(\tau,\tau') \tag{B.5}$$

Considering space and time translation invariant solutions for the Dyson equation, of the form  $Q_x^{ab}(\tau, \tau') = Q^{ab}(\tau - \tau')$  and  $z^a(\tau) = z^a$ , the above equation reduces to,

$$(Q^{-1})^{ab}(i\omega_k) = (\frac{1}{\Gamma}\omega_k^2 + z^a)\delta_{ab} - \Sigma_{ab}(\omega_k)$$
 (B.6a)  
$$\Sigma^{ab}(\tau) = \frac{3\tilde{J}^2}{2}Q^{ab}(\tau)^2$$
 (B.6b)

where  $\tilde{J}^2 = J^2 + J'^2$ . The above large-N saddle-point equations are same as that of the zero dimensional model<sup>65</sup>.

Saddle point equation in PM phase— Using the replica diagonal ansatz and analytically continuing to real frequency, we obtain the saddle-point equation for the retarded Green's function,

$$[Q_R(\omega)]^{-1} = -\frac{1}{\Gamma}\omega^2 + z - \Sigma_R(\omega)$$
 (B.7a)

$$\Sigma_R(\omega) = \Sigma(i\omega_k \to \omega + i0^+)$$
 (B.7b)

Following procedure similar to that in the case of SY model (Appendix A), the self-energy can be written as,

$$\Sigma_{R}(\omega) = i \frac{3J^{2}}{2} \int_{0}^{\infty} e^{i\omega t} \left( n_{1}^{2}(t) - n_{2}^{2}(t) \right), \qquad (B.8a)$$

$$n_{1}(t) = \int d\omega e^{-i\omega t} \rho(\omega) n_{B}(-\omega),$$

$$n_{2}(t) = \int d\omega e^{-i\omega t} \rho(\omega) n_{B}(\omega) \qquad (B.8b)$$

where  $\rho(\omega) = -(1/\pi) \text{Im} Q_R(\omega)$ . The saddle-point equations can be solved numerically following steps similar to that discussed in the case of SY model.

Saddle point equation in SG phase— Using 1RSB ansatz, after analytical continuation to real frequency  $\mathrm{i}\omega_k \to \omega + \mathrm{i}\eta$  only the regular part (see Sec. 3B) contributes to the retarded Green's function  $Q_R^{ab}(\omega) = Q_R(\omega)\delta_{ab} = \tilde{Q}(\mathrm{i}\omega_k \to \omega + \mathrm{i}\eta)\delta_{ab}$  such that

$$\begin{split} &[Q_{R}(\omega)]^{-1} = -\frac{1}{\Gamma}\omega^{2} + z - \Sigma_{R}(\omega), \\ &\Sigma_{R}(\omega) = \frac{3J^{2}}{2} \int_{0}^{\beta} d\tau e^{i\omega_{k}\tau} [\tilde{Q}(\tau)]^{2} + 3J^{2} q_{\text{EA}} Q_{R}(\omega) \\ &= \mathrm{i} \frac{3J^{2}}{2} \int_{0}^{\infty} e^{i\omega t} \Big( n_{1}^{2}(t) - n_{2}^{2}(t) \Big) + 3J^{2} q_{\text{EA}} Q_{R}(\omega), \end{split}$$

where

$$n_1(t) = \int d\omega e^{-i\omega t} \rho(\omega) n_B(-\omega),$$
 (B.9c)

$$n_2(t) = \int d\omega e^{-i\omega t} \rho(\omega) n_B(\omega)$$
 (B.9d)

The spherical constraint now takes the form,

$$-\int_{-\infty}^{\infty} d\omega \rho(\omega) n_B(\omega) = 1 - q_{\rm EA}$$
 (B.10)

Wightmann correlation functions— The Wightmann functions in the PM phase is given by,

$$Q_{lr}(t) = Q(it + \beta/2) = -\int \frac{d\omega}{2\pi} e^{-i\omega t} \frac{\pi \rho(\omega)}{\sinh\frac{\beta\omega}{2}}, \quad (B.11)$$

and

$$Q_{lr}^{ab}(t) = \epsilon_{ab} q_{\rm EA} - \delta_{ab} \int \frac{d\omega}{2\pi} e^{-i\omega t} \frac{\pi \rho(\omega)}{\sinh\frac{\beta\omega}{2}}$$
 (B.12)

in the SG phase.

### Appendix C: Calculation of butterfly velocity from the single mode ansatz

Here we briefly discuss the computation of  $v_B$  in SY model using the imaginary p method or the single mode ansatz, as described in the main text (Sec. 5A). By calculating  $\lambda_{\rm L}(p)$  for imaginary values of momentum, p=i|p|, using the exponential growth ansatz  $\mathcal{F}_{\mu,p}(t_1,t_2)=e^{\lambda_{\rm L}(p)(t_1+t_2)/2}f_{\mu,p}^a(t_1-t_2)$  in Eq.(5.2), we find  $p_1$  and  $p_s$ , as discussed in Sec. 5A. At T=0.8 and S=0.8, for example, we have  $p_1=2.91$  and  $p_s=1.9$ , as shown in Fig.C.1. Also, in Fig.C.2, we show that  $p_s$  is always less than  $p_1$ , e.g., at S=1.0.




FIG. C.1.  $\lambda_{\rm L}(p=i|p|)$  for T=0.8 and S=0.8 where  $p_1$  and  $p_s$  are marked.




FIG. C.2.  $p_1, p_s$  for a range of temperatures at S=1.0