

Viscosity Solutions for HJB Equations on the Process Space: Application to Mean Field Control with Common Noise

Jianjun Zhou* Nizar Touzi[†] and Jianfeng Zhang[‡]

Abstract

In this paper we investigate a path dependent optimal control problem on the process space with both drift and volatility controls, with possibly degenerate volatility. The dynamic value function is characterized by a fully nonlinear second order path dependent HJB equation on the process space, which is by nature infinite dimensional. In particular, our model covers mean field control problems with common noise as a special case. We shall introduce a new notion of viscosity solutions and establish both existence and comparison principle, under merely Lipschitz continuity assumptions. The main feature of our notion is that, besides the standard smooth part, the test function consists of an extra singular component which allows us to handle the second order derivatives of the smooth test functions without invoking the Ishii's lemma. We shall use the doubling variable arguments, combined with the Ekeland-Borwein-Preiss Variational Principle in order to overcome the noncompactness of the state space. A smooth gauge-type function on the path space is crucial for our estimates.

Keywords: Mean field control, common noise, viscosity solutions, HJB equations, Wasserstein space, path dependence, comparison principle, variational principle

2020 AMS Subject Classification: 49L25, 49N80, 35R15, 60H30, 93E20.

*College of Science, Northwest A&F University, Yangling 712100, Shaanxi, P. R. China, zhoujianjun@nwsuaf.edu.cn.

[†]New York University, Tandon School of Engineering, United States, nizar.touzi@nyu.edu.

[‡]Department of Mathematics, University of Southern California, United States, jianfenz@usc.edu. This author is partially supported by NSF grant DMS-2205972.

1 Introduction

Strongly motivated by mean field control problems, we consider a stochastic control problem whose data rely on the whole underlying state process. Consequently, its dynamic value function satisfies an infinite dimensional HJB equation on the process space. This framework is very general and has the following features:

- Our model covers the mean field control with common noise as a special case. In the state dependent case, the idea of lifting probability measures to the Hilbert space of random variables is due to Lions [47], and in the contexts of mean field games, Ahuja-Ren-Yang [1] employed this idea to study common noise.
- We allow for path dependence and thus our HJB equation is path dependent. For finite dimensional path dependent PDEs, we refer to a series of works by the authors and their collaborators: [28, 29, 30, 49, 56], and the references therein. We also refer to Wu-Zhang [54] and Cosso-Gozzi-Kharroubi-Pham-Rosestolato [18] for path dependent mean field control problems.
- We consider both the drift and volatility controls, with possibly degenerate volatilities. Consequently, besides infinite dimensionality, our HJB equation is a fully nonlinear degenerate second order PDE.
- We allow the problem to depend on the joint law of the state process and the control process. This is in the spirit of the mean field game of controls (which was called extended mean field game in the early stage), see e.g. Gomes-Voskanyan [38].

Initiated independently by Caines-Huang-Malhame [10] and Lasry-Lions [42], the theory of mean field games and mean field controls has received extremely strong attention in the literature. We refer to Lions's lecture [47] and the book Carmona-Delarue [13, 14] and the references therein for a general exposition of the theory. One popular approach in the literature is to consider PDEs on the Wasserstein space of probability measures. We should note that the master equations for mean field games have quite different nature than the HJB equations for mean field controls. In this paper we address the latter equations and our main focus is the comparison principle, while for mean field game master equations even classical solutions typically violate the comparison principle.

Partially due to its infinite dimensionality, such an HJB equation can rarely have a classical solution. In recent years there have been serious efforts on viscosity solutions for HJB equations for mean field control problems. We shall provide a literature review in the end

of this introduction. Our goal of this paper is to propose an appropriate notion of viscosity solution and show that the dynamic value function of our control problem is the unique viscosity solution of the HJB equation. In particular, we shall establish the comparison principle of the viscosity solution under merely Lipschitz continuity assumptions, by using the doubling variable arguments. Our results also imply the wellposedness of the second order HJB equation on the Wasserstein space of probability measures induced from mean field control problems with common noise. To the best of our knowledge, even in the mean field control framework, our model is most general and our conditions are the weakest.

Notice that a notion of viscosity solution is essentially determined by the set of test functions one chooses. In particular, the proof of the comparison principle relies heavily on this choice of test functions. Inspired by the viscosity solution theory for PDEs in infinite dimensions, see, e.g., Crandall-Lions [23, Definition 2.1], Li-Yong [43, Chapter 6 Definition 3.1], and Fabbri-Gozzi-Świąch [33, Definitions 3.32 & 3.35], our test functions take the form

$$\varphi + \phi, \tag{1.1}$$

where the first part φ is smooth (in appropriate sense) and thus is standard; the extra part ϕ is not smooth, but is absolutely continuous in time with respect to the Lebesgue measure. Our construction of ϕ is motivated from an observation in the constant volatility case. In this case, by a simple transformation one can convert the state process from a controlled SDE to a controlled ODE with random coefficients. Consequently, the resulting HJB equation becomes first order, whose viscosity solution is a lot easier to study. By applying the inverse transformation on the test functions for the latter equation, we obtain a candidate test function ϕ for our original dynamic value function, which turns out to be absolutely continuous in time but is in general nonsmooth. See Subsection 4.1 for details. For the general case with volatility controls, we tailor the construction of ϕ which in essence cancels the diffusion term. Indeed, technically ϕ is used to cancel some terms appearing in the doubling variable arguments, which involve the second order derivatives of φ and are otherwise hard to estimate. To our best knowledge, this type of test functions is new in the literature of viscosity solutions for mean field control problems.

Another important consequence of introducing the singular component ϕ of test functions is that we can establish the comparison principle without using the Ishii's lemma, even though we are using the doubling variable argument for a second order equation. Indeed, as we just explained, in the constant volatility case, the introduction of ϕ allows us to convert the HJB equation into a first order one, which does not require Ishii's lemma. For standard second order HJB equations, Ishii's lemma is used exactly to handle the second

order derivatives of the test functions. So our approach provides an alternative solution to this important issue. Moreover, our general framework covers the standard finite dimensional (path dependent) HJB equations, as well as HJB equations on Wasserstein space of probability measures, so our results imply the comparison principle for viscosity solutions of those equations as well, without using the Ishii’s lemma. However, we should point out that our notion of viscosity solutions is not equivalent to the “standard” ones for those equations. So our results do not imply directly the wellposedness results in the literature.

Unfortunately, even in the mean field framework, this function ϕ is typically not law invariant, which prohibits us from defining the viscosity solution intrinsically on the Wasserstein space. So, besides the advantage of covering the common noise case, technically we are required to consider functions on the process space. Another advantage for working directly on the process space is that, in the state dependent case, the square distance of two random variables is a smooth functional, but the square of 2-Wasserstein distance of two probability measures is not differentiable. While being more involved, our path dependent case will benefit from this feature as well. We should note that Soner-Yan [50, 51] introduced a nice norm on the Wasserstein space by using Fourier transformation, whose square is smooth. However, it is not clear how to extend this norm to serve for our purpose in the path dependent case.

One drawback of the process space is its lack of local compactness, which is crucial for the proof of the comparison principle. To circumvent this difficulty, we shall use the Ekeland-Borwein-Preiss variational principle, see Aubin-Ekeland [2] and Borwein-Preiss [8]. Roughly speaking, to optimize a continuous function on a non-compact space, one may construct an approximate function by using the so called gauge type functions such that the approximate function has a strict optimal argument. To serve as test functions for our purpose, we require these gauge type functions on processes to be smooth with desired estimates for their derivatives. This is achieved by utilizing the smooth gauge type function on continuous paths constructed by Zhou [56] for path dependent PDEs.

To prove the comparison principle, we shall first double the spatial variable and then double the temporal variable. This kind of two step approximations is standard for parabolic equations, see, e.g., Crandall-Ishii [21, Lemma 8] and Crandall-Ishii-Lions [22, Theorem 8.3]. However, due to a subtle adaptedness requirement of ϕ , we need a third step of approximation to derive the desired contradiction.

The rest of the paper is organized as follows. First, we conclude this introduction by providing a brief literature review on viscosity solutions for mean field control problems. In Section 2 we introduce our control problem and establish some basic properties of the

dynamic value function. In Section 3 we introduce smooth functions φ on process space, which leads to the target HJB equation. In Section 4 we construct the singular component ϕ and derive some crucial estimates. In Section 5 we propose our notion of viscosity solutions and present the main results. In Section 6 we illustrate how our general model covers the mean field control problem with common noise as a special case. Sections 7, 8 and 9 are devoted to the proof of the comparison principle for viscosity solutions. Finally in Appendix we complete some technical proofs.

Some literature review on viscosity solutions for mean field control problems. We first remark that these equations are by nature infinite dimensional, and thus the related works are intrinsically connected to the viscosity solution theory for infinite dimensional HJB equations, for which we refer to Lions [44, 45, 46] and the books Li-Yong [43] and Fabbri-Gozzi-Świąch [33]. For first order HJB equations on Wasserstein space, arising from mean field control problems with deterministic controls or other related problems, we refer to the works Bertucci [7], Cardaliaguet-Quincampoix [11], Conforti-Kraaij-Tonon [16], Feng-Katsoulakis [31], Gangbo-Nguyen-Tudorascu [36], Gangbo-Tudorascu [37], Jimenez-Marigonda-Quincampoix [39, 40]. These equations involve the Lions derivative $\partial_\mu V(t, \mu, x)$, but not the higher order derivative $\partial_{x\mu} V(t, \mu, x)$, and in the finite dimensional case they correspond to the standard first order HJB equations.

We are mainly interested in mean field control problems where the state process is a controlled diffusion, with or without volatility controls. These equations involve $\partial_{x\mu} V(t, \mu, x)$ (and $\partial_{\mu\mu} V$ when there is common noise) and in the finite dimensional case are equivalent to the standard second order HJB equations. In the path dependent setting, Wu-Zhang [54] proposed a notion of viscosity solutions by restricting the viscosity neighborhood of some point (t, μ) to certain compact set, and established the partial comparison principle. However, the full comparison principle requires certain perturbed equations to have a classical solution, which is in general not easy to verify, especially in the case with volatility control. Following the similar approach, Talbi-Touzi-Zhang [52, 53] established the complete wellposedness for a mean field optimal stopping problem. The work Cosso-Gozzi-Kharroubi-Pham-Rosestolato [19] removed the compactness requirement by applying the Ekeland-Borwein-Preiss variational principle. Based on a so called Gaussian-smoothed 2-Wasserstein distance, [19] constructed a finite dimensional smooth approximation of the value function and established the comparison principle by comparing viscosity semi-solutions with the value function. The works Burzoni-Ignazio-Reppen-Soner [9] and Soner-Yan [50] derived the comparison result by means of the doubling variable argument.

They do not invoke Ishii's lemma either, but due to a completely different nature than ours, see Remark 6.9 below for the detailed explanation. In particular, they introduced a smooth metric on the Wasserstein space by using the Fourier transformation. However, these works require certain uniform Lipschitz continuity on the controls. When the volatility is a positive constant and hence the equation is semilinear, this serious constraint was removed in the recent work Soner-Yan [51] and the comparison principle is established. More recently, by also using the Fourier-Wasserstein metric, Bayraktar-Ekren-Zhang [3] established an Ishii's lemma for functions on the Wasserstein space of probability measures, and considered a general mean field control problems with common noise. In the case with common noise with constant coefficient but without idiosyncratic noises, Gangbo-Mayorga-Swiech [34] and Mayorga-Swiech [48] studied the so called L -viscosity solution by lifting the equation to the Hilbert space. The work Daudin-Seeger [24] studied semilinear HJB equations, by applying the doubling variable argument with a further entropy penalization. Another recent work Daudin-Jackson-Seeger [25] studied semilinear equations with common noise and non-convex Hamiltonian, which allows to consider zero-sum game problem in the mean field setting. They established the comparison principle by exploiting the idea of [3] together with some delicate regularity estimates.

We should also mention the following works concerning potential mean field games, where the mean field control problem is involved automatically: Bensoussan-Graber-Yam [4, 5], Bensoussan-Tai-Yam [6], Carmona-Cormier-Soner [12], Cecchin-Delarue [15], and Gangbo-Meszaros [35]. Moreover, the works Cox-Kallblad-Larsson-Svaluto-Ferro [20] on controlled measure-valued martingales and Feng-Swiech [32] where the controlled dynamics involves a mixture of a Hamiltonian flow and a gradient flow are also closely related.

Some notations. We shall denote $x \cdot x' := \sum_{i=1}^n x_i x'_i$, for $x, x' \in \mathbb{R}^n$, and $M : M' := \text{tr}(M^\top M')$ with M^\top the transpose of M , for $M, M' \in \mathbb{R}^{m \times n}$. The corresponding subset of $d \times d$ -symmetric matrices is denoted by \mathbb{S}^d . Moreover, $|x|^2 := x \cdot x$, $|M|^2 := M : M$.

For a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, an Euclidian space E , and $p \geq 1$, we denote by $\mathbb{L}^p(\mathcal{F}_t; E)$ the space of \mathcal{F}_t -measurable E -valued random variables ξ_* such that $\|\xi_*\|_p^p := \mathbb{E}[|\xi_*|^p] < \infty$. Similarly, given a finite horizon $T > 0$, we denote by $\mathbb{L}^p(\mathbb{F}_{[t, T]}; E)$ the space of \mathbb{F} -progressively measurable E -valued processes ξ on $[t, T]$ such that $\mathbb{E}[\int_t^T |\xi_s|^p ds] < \infty$.

For the filtration $\mathbb{F} = \{\mathcal{F}_t\}_{0 \leq t \leq T}$, we assume \mathcal{F}_0 is rich enough to support any probability measure on \mathbb{R}^d , and $\mathcal{F}_t = \mathcal{F}_0 \vee \mathcal{F}_t^B$, where B is a d -dimensional Brownian motion on $(\Omega, \mathcal{F}, \mathbb{P})$. Moreover, we fix a sub-filtration $\mathbb{F}^0 = \{\mathcal{F}_t^0\}_{0 \leq t \leq T} \subset \mathbb{F}$, and denote $\mathbb{E}_t^0 := \mathbb{E}[\cdot | \mathcal{F}_t^0]$.

Denote $\mathbb{X} := C([0, T]; \mathbb{R}^d)$, equipped with the uniform norm $|\cdot|_\infty$. For any $p \geq 1$, let \mathcal{X}_p

denote the set of \mathbb{F} -progressively measurable continuous processes ξ with $\|\xi\|_p^p := \mathbb{E}[|\xi|_\infty^p] < \infty$, equipped with the norm $\|\cdot\|_p$. Note that $\xi(\omega) \in \mathbb{X}$ for all $\omega \in \Omega$. Let $\overline{\mathcal{X}}_{[t,T]}^p := [t, T] \times \mathcal{X}_p$ denote the time and state space.

In order to distinguish the dependence on the whole process, or more precisely on the (deterministic) mapping on $[0, T] \times \Omega$, from that on the realized paths of the process, we introduce the notation $\underline{\xi} = \xi$ to emphasize the dependence on the whole process. That is, for a function φ on $[0, T] \times \mathbb{X} \times \mathcal{X}_p$, we shall write $\varphi_t(\mathbf{x}, \underline{\xi})$ instead of $\varphi_t(\mathbf{x}, \xi)$. In particular, this allows us to express the following without confusion:

$$\varphi_t(\xi, \underline{\xi})(\omega) = \varphi_t(\xi(\omega), \underline{\xi}),$$

and, when not involving \mathbf{x} , the value $\varphi_t(\underline{\xi})$ is always deterministic. When it is more convenient, especially when the functions are state dependent, we may also use the notation $\varphi(t, \mathbf{x}, \underline{\xi}) = \varphi_t(\mathbf{x}, \underline{\xi})$. Moreover, throughout the paper we shall always assume all involved path dependent functions φ are adapted in the sense:

$$\varphi_t(\mathbf{x}, \underline{\xi}) = \varphi_t(\mathbf{x}_{\cdot \wedge t}, \underline{\xi}_{\cdot \wedge t}), \quad \text{for all } (t, \mathbf{x}, \xi) \in [0, T] \times \mathbb{X} \times \mathcal{X}_p. \quad (1.2)$$

2 Formulation of the process dependent control problem

We will consider open loop controls taking values in A , a domain in a Euclidian space.¹ Denote $\mathcal{A}_{[t,T]} := \mathbb{L}^2(\mathbb{F}_{[t,T]}; A)$ and $\mathcal{A}_t := \mathbb{L}^2(\mathcal{F}_t; A)$. Let:

$$\begin{aligned} (b, \sigma) &: [0, T] \times \Omega \times \mathbb{X} \times A \times \mathcal{X}_2 \times \mathcal{A}_T \longrightarrow (\mathbb{R}^d, \mathbb{R}^{d \times d}), \\ f &: [0, T] \times \mathcal{X}_2 \times \mathcal{A}_T \longrightarrow \mathbb{R}, \quad \text{and } g : \mathcal{X}_2 \longrightarrow \mathbb{R}. \end{aligned} \quad (2.1)$$

As usual, we omit the variable $\omega \in \Omega$ inside b and σ , and we recall the convention (1.2). For any $(t, \xi) \in \overline{\mathcal{X}}_{[0,T]}^2$ and $\alpha \in \mathcal{A}_{[t,T]}$, we consider the following path dependent SDE on $[t, T]$:

$$\begin{aligned} X_s^{t,\xi,\alpha} &= \xi_s, \quad s \in [0, t]; \\ dX_s^{t,\xi,\alpha} &= b_s(X^{t,\xi,\alpha}, \alpha_s, \underline{X}^{t,\xi,\alpha}, \underline{\alpha}_s) ds + \sigma_s(X^{t,\xi,\alpha}, \alpha_s, \underline{X}^{t,\xi,\alpha}, \underline{\alpha}_s) dB_s, \quad s \in [t, T]. \end{aligned} \quad (2.2)$$

Here X is path dependent, while α involves only the current state, and the notation $\underline{\alpha}_s$ refers to the whole random variable α_s . The value function of our control problem is:

$$\begin{aligned} V_t(\underline{\xi}) &:= \inf_{\alpha \in \mathcal{A}_{[t,T]}} J_t(\underline{\xi}, \underline{\alpha}), \quad (t, \xi) \in \overline{\mathcal{X}}_{[0,T]}^2, \\ \text{where } J_t(\underline{\xi}, \underline{\alpha}) &:= g(\underline{X}^{t,\xi,\alpha}) + \int_t^T f_s(\underline{X}^{t,\xi,\alpha}, \underline{\alpha}_s) ds. \end{aligned} \quad (2.3)$$

¹Although closed loop controls are expected to induce the same value function under appropriate regularity conditions, we refrain from considering this case for technical simplicity.

Throughout the paper, the following assumptions will be in force.

Assumption 2.1. (i) *The coefficients b, σ, f are progressively measurable in all variables and adapted in the sense of (1.2); in particular, b, σ are \mathbb{F} -progressively measurable; and $h(\cdot, \mathbf{0}, \cdot, \cdot, \underline{\mathbf{0}}, \cdot)$, for $h = b, \sigma$, $f(\cdot, \underline{\mathbf{0}}, \cdot)$ and $g(\underline{\mathbf{0}})$ are bounded² by a constant C_0 .*

(ii) *b, σ, f, g are uniformly Lipschitz continuous in (\mathbf{x}, ξ) with a Lipschitz constant L :*

$$\begin{aligned} |h_t(\mathbf{x}, a, \underline{\xi}, \underline{\alpha}_t) - h_t(\mathbf{x}', a, \underline{\xi}, \underline{\alpha}_t)| &\leq L|\mathbf{x}_{\cdot \wedge t} - \mathbf{x}'_{\cdot \wedge t}|_\infty, \quad h = b, \sigma; \\ |h_t(\mathbf{x}, a, \underline{\xi}, \underline{\alpha}_t) - h_t(\mathbf{x}, a, \underline{\xi}', \underline{\alpha}_t)| &\leq L\left(\mathbb{E}_t^0[|\xi_{\cdot \wedge t} - \xi'_{\cdot \wedge t}|_\infty^2]\right)^{\frac{1}{2}}, \quad \mathbb{P}\text{-a.s.}, \quad h = b, \sigma; \\ |f_t(\underline{\xi}, \underline{\alpha}_t) - f_t(\underline{\xi}', \underline{\alpha}_t)| &\leq L\|\xi_{\cdot \wedge t} - \xi'_{\cdot \wedge t}\|_2, \quad |g(\underline{\xi}) - g(\underline{\xi}')| \leq L\|\xi - \xi'\|_2, \end{aligned}$$

for all $(t, \mathbf{x}, \xi) \in [0, T] \times \mathbb{X} \times \mathcal{X}_p$ and $\alpha \in \mathcal{A}_{[0, T]}$.

Here $\mathbf{0}$ denotes the zero path and $\underline{\mathbf{0}} \in \mathcal{X}_2$ the zero stochastic process. Notice that the volatility σ is possibly degenerate, so there is no loss of generality in taking it as a square matrix and considering X and B with same dimension.

Lemma 2.2. *Let Assumption 2.1 hold true.*

(i) *For any $(t, \xi) \in \mathcal{X}_{[0, T]}^2$ and $\alpha \in \mathcal{A}_{[t, T]}$, the SDE (2.2) admits a unique strong solution $X^{t, \xi, \alpha}$. Moreover, for any $p \geq 2$, there exists a constant C_p , depending only on p, T, d , and the constants L, C_0 in Assumption 2.1, such that, for any $\xi' \in \mathcal{X}_p$,*

$$\begin{aligned} \|X^{t, \xi, \alpha}\|_p &\leq C_p(1 + \|\xi_{\cdot \wedge t}\|_p); \quad \|X^{t, \xi, \alpha} - X^{t, \xi', \alpha}\|_p \leq C_p\|\xi_{\cdot \wedge t} - \xi'_{\cdot \wedge t}\|_p; \\ \|X_{\cdot \wedge s}^{t, \xi, \alpha} - X_{\cdot \wedge s'}^{t, \xi, \alpha}\|_p &\leq C_p(1 + \|\xi_{\cdot \wedge t}\|_p)|s - s'|^{\frac{1}{2}}, \quad s, s' \in [t, T]. \end{aligned} \tag{2.4}$$

(ii) *The functions J and V are adapted in ξ in the sense of (1.2), and V satisfies the dynamic programming principle:*

$$V_t(\underline{\xi}) = \inf_{\alpha \in \mathcal{A}_{[t, T]}} \left\{ V_{t+\delta}(X^{t, \xi, \alpha}) + \int_t^{t+\delta} f_s(X^{t, \xi, \alpha}, \underline{\alpha}_s) ds \right\}, \text{ for all } (t, \xi) \in \overline{\mathcal{X}}_{[0, T]}^2, \delta \leq T - t. \tag{2.5}$$

Moreover, for the constant $C = C_2$ of (i), denoting $\Delta t := t - t'$, $\Delta \xi := \xi - \xi'$, we have:

$$\begin{aligned} |J_t(\underline{\xi}, \underline{\alpha}_t)| &\leq C(1 + \|\xi_{\cdot \wedge t}\|_2); \quad |J_t(\underline{\xi}, \underline{\alpha}_t) - J_t(\underline{\xi}', \underline{\alpha}_t)| \leq C\|\Delta \xi_{\cdot \wedge t}\|_2; \\ |V_t(\underline{\xi})| &\leq C(1 + \|\xi_{\cdot \wedge t}\|_2); \quad |V_t(\underline{\xi}) - V_{t'}(\underline{\xi}')| \leq C\left[\|\xi_{\cdot \wedge t} - \xi'_{\cdot \wedge t}\|_2 + (1 + \|\xi_{\cdot \wedge t}\|_2)|t - t'|^{\frac{1}{2}}\right]. \end{aligned} \tag{2.6}$$

The DPP (2.5) follows from similar arguments as in [18, Theorem 3.4]. All the involved estimates are rather standard, in particular the conditional \mathbb{L}^2 -type regularity of b, σ with respect to \underline{X} in Assumption 2.1 (ii) does not induce any difficulty. We thus omit the proof.

²This boundedness requirement is just for technical convenience. In particular, it can be relaxed so as to cover the linear quadratic case.

3 The HJB equation and classical solutions

Following the standard control theory, the DPP (2.5) induces an HJB equation for V . For this purpose, we first introduce derivatives for functions on $\overline{\mathcal{X}}_{[0,T]}^p$.

3.1 Smooth functions on $\overline{\mathcal{X}}_{[0,T]}^p$

For a generic metric space E and for $t \in [0, T)$, $p \geq 1$, let $C^0(\mathcal{X}_{[t,T]}^p; E)$ denote the space of adapted and continuous functions $\varphi : \mathcal{X}_{[t,T]}^p \rightarrow E$. In particular, $C^0(\mathcal{X}_{[t,T]}^p) := C^0(\mathcal{X}_{[t,T]}^p; \mathbb{R})$. Throughout this paper, $\mathcal{M}_{[t,T]}^p$ denotes the space of Itô processes $X \in \mathcal{X}_p$ such that

$$dX_s = \beta_s ds + \gamma_s dB_s, \quad s \in [t, T], \quad \text{for some } (\beta, \gamma) \in \mathbb{L}^p(\mathbb{F}_{[t,T]}; \mathbb{R}^d \times \mathbb{R}^{d \times d}). \quad (3.1)$$

Definition 3.1. For $\hat{t} < T$, $p \geq 2$, we say $\varphi \in C^{1,2}(\overline{\mathcal{X}}_{[\hat{t},T]}^p)$ if $\varphi \in C^0(\overline{\mathcal{X}}_{[\hat{t},T]}^p)$ and there exist $\partial_t \varphi \in C^0(\overline{\mathcal{X}}_{[\hat{t},T]}^p)$, $\partial_X \varphi \in C^0(\overline{\mathcal{X}}_{[\hat{t},T]}^p; \mathbb{L}^{\frac{p}{p-1}}(\mathcal{F}; \mathbb{R}^d))$, and $\partial_{\mathbf{x}X} \varphi \in C^0(\overline{\mathcal{X}}_{[\hat{t},T]}^p; \mathbb{L}^{\frac{p}{p-2}}(\mathcal{F}; \mathbb{S}^d))$, such that $\partial_t \varphi_t(\underline{\xi})$, $\partial_X \varphi_t(\underline{\xi})$ and $\partial_{\mathbf{x}X} \varphi_t(\underline{\xi})$ are \mathcal{F}_t -measurable for all $(t, \underline{\xi}) \in \overline{\mathcal{X}}_{[\hat{t},T]}^p$, and for any $X \in \mathcal{M}_{[\hat{t},T]}^p$, the following functional Itô formula holds:

$$d\varphi_t(\underline{X}) = \partial_t \varphi_t(\underline{X}) dt + \mathbb{E} \left[\partial_X \varphi_t(\underline{X}) \cdot \beta_t dt + \frac{1}{2} \partial_{\mathbf{x}X} \varphi_t(\underline{X}) : \gamma_t \gamma_t^\top dt \right], \quad t \in [\hat{t}, T]. \quad (3.2)$$

The following simple lemma is crucial for the above definition. The proof is similar to the justification of [29, Definition 2.8], and is postponed to Appendix.

Lemma 3.2. For any $\varphi \in C^{1,2}(\overline{\mathcal{X}}_{[\hat{t},T]}^p)$, the derivatives $\partial_t \varphi$, $\partial_X \varphi$, $\partial_{\mathbf{x}X} \varphi$ are unique.

Remark 3.3. (i) *The state dependent case:* Let $\varphi_t(\underline{\xi}) = \psi(t, \underline{\xi}_t)$ for some ψ defined on $[0, T] \times \mathbb{L}^2(\mathcal{F}_T)$. Then, $\partial_t \varphi = \partial_t \psi$ is standard and $\partial_X \varphi = \partial_X \psi$ is the Fréchet derivative:

$$\begin{aligned} \partial_t \psi(t, \underline{Y}) &= \lim_{\delta \downarrow 0} \frac{1}{\delta} [\psi(t + \delta, \underline{Y}) - \psi(t, \underline{Y})], \quad Y \in \mathbb{L}^2(\mathcal{F}_T); \\ \psi(t, \underline{Y} + \underline{Z}) &= \psi(t, \underline{Y}) + \mathbb{E} \left[\partial_X \psi(t, \underline{Y} + \underline{Z}) Z \right] + o(\|Z\|_2), \quad Y, Z \in \mathbb{L}^2(\mathcal{F}_T). \end{aligned}$$

Moreover, $\partial_{\mathbf{x}X} \varphi = \partial_{\mathbf{x}X} \psi$ can be determined by:

$$\partial_{\mathbf{x}X} \psi(t, \underline{X}_t) = \frac{d}{ds} \left\langle \partial_X \psi(t, \underline{X}_t + \underline{B}_s - \underline{B}_t), B_s \right\rangle \Big|_{s=t} \quad \text{for all } X \in \mathcal{M}_{[t,T]}^p.$$

(ii) *The law invariant case:* Let $\varphi_t(\underline{\xi}) = \check{\varphi}_t(\mathbb{P}_\xi)$ for some $\check{\varphi}$ smooth as in [54, Theorem 2.7], then we can easily see that $\varphi \in C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^2)$ with

$$\partial_t \varphi_t(\underline{\xi}) = \partial_t \check{\varphi}_t(\mathbb{P}_\xi), \quad \partial_X \varphi_t(\underline{\xi}) = \partial_\mu \check{\varphi}_t(\mathbb{P}_\xi, \xi), \quad \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}) = \partial_{\omega\mu} \check{\varphi}_t(\mathbb{P}_\xi, \xi).$$

(iii) For the general case, one may define the path derivatives first and then prove the Itô formula (3.2), as in [26, 17] for functions on $[0, T] \times \mathbb{X}$ and in [54, 18] for functions on $[0, T] \times \mathcal{P}_2(\mathbb{X})$. However, for the viscosity theory later, we will need only the Itô formula, so we follow the approach of [29] by using (3.2) directly to define the path derivatives.

Example 3.4. All smooth functions involved in this paper are of the form

$$\varphi_t(\underline{\xi}) := \mathbb{E}[\bar{\varphi}_t(\xi - \hat{\xi}_{\cdot, \wedge \hat{t}})], \quad (t, \xi) \in \bar{\mathcal{X}}_{[\hat{t}, T]}^p, \quad \text{for some fixed } (\hat{t}, \hat{\xi}) \in \bar{\mathcal{X}}_{[0, T]}^p,$$

where $p \geq 2$ and $\bar{\varphi} : [0, T] \times \mathbb{X} \rightarrow \mathbb{R}$ is continuous, adapted, and admits continuous pathwise Dupire's derivatives $(\partial_t \bar{\varphi}, \partial_{\mathbf{x}} \bar{\varphi}, \partial_{\mathbf{xx}} \bar{\varphi}) : [0, T] \times \mathbb{X} \rightarrow (\mathbb{R}, \mathbb{R}^d, \mathbb{S}^d)$ such that

$$|\partial_t \bar{\varphi}_t(\mathbf{x})| \leq C[1 + |\mathbf{x}_{\cdot, \wedge t}|_\infty^p], \quad |\partial_{\mathbf{x}} \bar{\varphi}_t(\mathbf{x})| \leq C[1 + |\mathbf{x}_{\cdot, \wedge t}|_\infty^{p-1}], \quad |\partial_{\mathbf{xx}} \bar{\varphi}_t(\mathbf{x})| \leq C[1 + |\mathbf{x}_{\cdot, \wedge t}|_\infty^{p-2}]. \quad (3.3)$$

Then $\varphi \in C^{1,2}(\bar{\mathcal{X}}_{[\hat{t}, T]}^p)$, and denoting $\xi' := \xi - \hat{\xi}_{\cdot, \wedge \hat{t}}$, we have

$$\partial_t \varphi_t(\underline{\xi}) := \mathbb{E}[\partial_t \bar{\varphi}_t(\xi')], \quad \partial_X \varphi_t(\underline{\xi}) := \partial_{\mathbf{x}} \bar{\varphi}_t(\xi'), \quad \partial_{\mathbf{xx}} \varphi_t(\underline{\xi}) := \partial_{\mathbf{xx}} \bar{\varphi}_t(\xi'). \quad (3.4)$$

Proof. For any $X \in \mathcal{M}_{[\hat{t}, T]}^p$, by the standard functional Itô formula (cf. [26, 17]) we have

$$d\bar{\varphi}_t(X') = \partial_t \bar{\varphi}_t(X') dt + \partial_{\mathbf{x}} \bar{\varphi}_t(X') \cdot dX_t + \frac{1}{2} \partial_{\mathbf{xx}} \bar{\varphi}_t(X') : d\langle X \rangle_t, \quad t \geq \hat{t}, \quad \text{with } X' := X - \hat{\xi}_{\cdot, \wedge \hat{t}}.$$

Take expectation on both sides and compare it with (3.2), we obtain (3.4) immediately. In particular, the integrability of $\partial_X \varphi$ and $\partial_{\mathbf{xx}} \varphi$ follows directly from (3.3) and the \mathbb{L}^p -integrability of X and $\hat{\xi}$, and it follows from the Burkholder-Davis-Gundy inequality that $\partial_{\mathbf{x}} \bar{\varphi}_t(X') \cdot \gamma_t dB_t$ is a true martingale. \blacksquare

3.2 HJB equation on the space of processes $\bar{\mathcal{X}}_{[0, T]}^2$

We now consider the following HJB equation on $\bar{\mathcal{X}}_{[0, T]}^2$:

$$\begin{aligned} \mathcal{L}U_t(\underline{\xi}) &:= \partial_t U_t(\underline{\xi}) + \inf_{\alpha_t \in \mathcal{A}_t} H_t(\underline{\xi}, \partial_X U_t(\underline{\xi}), \partial_{\mathbf{xx}} U_t(\underline{\xi}), \underline{\alpha}_t) = 0, \quad (t, \xi) \in \bar{\mathcal{X}}_{[0, T]}^2; \\ \text{where } H_t(\underline{\xi}, \underline{Z}, \underline{\Gamma}, \underline{\alpha}_*) &:= \mathbb{E} \left[(b_t(\cdot) \cdot \underline{Z} + \frac{1}{2} \sigma \sigma^\top(\cdot) : \underline{\Gamma}) (\xi, \alpha_*, \underline{\xi}, \underline{\alpha}_*) + f_t(\underline{\xi}, \underline{\alpha}_*) \right], \quad (3.5) \\ (t, \xi) &\in \bar{\mathcal{X}}_{[0, T]}^2, \quad \underline{Z} \in \mathbb{L}^2(\mathcal{F}_t, \mathbb{R}^d), \quad \underline{\Gamma} \in \mathbb{L}^\infty(\mathcal{F}_t, \mathbb{S}^d), \quad \alpha_* \in \mathcal{A}_T. \end{aligned}$$

Remark 3.5. When b, σ do not depend on $(\underline{X}, \underline{\alpha})$ and $g(\underline{X}) = \mathbb{E}[\bar{g}(X)]$, $f_t(\underline{X}, \underline{\alpha}_t) = \mathbb{E}[\bar{f}_t(X, \alpha_t)]$, for some deterministic function \bar{f}, \bar{g} , one can easily show that

$$V_t(\underline{\xi}) = \mathbb{E}[v_t(\xi)], \quad (3.6)$$

where v is associated with the standard path dependent HJB equation on $[0, T] \times \mathbb{X}$:

$$\partial_t v_t(\mathbf{x}) + \inf_{a \in A} \left[b_t(\mathbf{x}, a) \cdot \partial_{\mathbf{x}} v_t(\mathbf{x}) + \frac{1}{2} \sigma \sigma_t^\top(\mathbf{x}, a) : \partial_{\mathbf{x}\mathbf{x}} v_t(\mathbf{x}) + \bar{f}_t(\mathbf{x}, a) \right] = 0, \quad (3.7)$$

and $\partial_t v, \partial_{\mathbf{x}} v, \partial_{\mathbf{x}\mathbf{x}} v$ are Dupire's path derivatives. Moreover, in the state dependent case, (3.7) reduces to the standard HJB equation:

$$\partial_t v(t, x) + \inf_{a \in A} \left[b(t, x, a) \cdot \partial_x v(t, x) + \frac{1}{2} \sigma \sigma^\top(t, x, a) : \partial_{xx} v(t, x) + \bar{f}(t, x, a) \right] = 0. \quad (3.8)$$

As usual, we start with classical solutions.

Definition 3.6. (Classical solution) A functional $U \in C^{1,2}(\bar{\mathcal{X}}_{[0,T]}^2)$ is called a classical solution (resp. subsolution, supersolution) to the equation (3.5) if

$$-\mathcal{L}U_t(\underline{\xi}) = (\text{resp. } \leq, \geq) 0, \quad \forall (t, \xi) \in \bar{\mathcal{X}}_{[0,T]}^2.$$

In the context of classical solutions, we also need the regularity in t which is not included in our general Assumption 2.1.

Assumption 3.7. The functions $h = b, \sigma, f$ are locally uniformly continuous in $(t, a, \alpha_*) \in [0, T] \times A \times \mathcal{A}_T$ in the following sense: for any $R > 0$, there exists a modulus of continuity function ρ_R such that, for all $\mathbf{x} \in \mathbb{X}$, $\|\xi_{\cdot \wedge t}\|_2 \leq R$, and all $t \leq s$, $a, a' \in A$, $\alpha_*, \alpha'_* \in \mathcal{A}_T$,

$$|h_t(\mathbf{x}, a, \underline{\xi}, \underline{\alpha}_*) - h_s(\mathbf{x}_{\cdot \wedge t}, a', \underline{\xi}_{\cdot \wedge t}, \underline{\alpha}'_*)| \leq \rho_R(|s - t| + |a - a'| + \|\alpha_* - \alpha'_*\|_2).$$

Theorem 3.8. Let Assumptions 2.1 and 3.7 hold and assume the value function $V \in C^{1,2}(\bar{\mathcal{X}}_{[0,T]}^2)$. Then V is the unique classical solution of the HJB equation (3.5).

Combining the DPP (2.5) and the Itô formula (3.2), it is rather standard to verify that V satisfies (3.5). The uniqueness is not hard either, however, since it will be a consequence of the uniqueness of the viscosity solution later, we omit the proof.

4 The singular component of test functions

In general one can hardly expect (3.5) to have a classical solution. Our goal of this paper is to build an appropriate notion of viscosity solution which allows for a complete characterization of the value function through the corresponding dynamic programming equation. The main feature of our approach is that, besides the standard smooth functions $\varphi \in C^{1,2}(\bar{\mathcal{X}}_{[0,T]}^2)$, our test functions have an additional component which is singular in certain sense. To motivate this, we first consider a simple constant diffusion coefficient setting.

4.1 A motivating case

In this subsection we study heuristically the state dependent case with constant diffusion $\sigma \equiv I_{d \times d}$. In this case, it is natural to consider the following change of variables to convert the SDE into a random ODE:

$$\begin{aligned} \tilde{b}_s(x, \omega, a, \underline{X}_s, \underline{\alpha}_s) &:= b_s(x + B_s(\omega), \omega, a, \underline{X}_s + \underline{B}_s, \underline{\alpha}_s), \\ \tilde{f}_s(\underline{X}_s, \underline{\alpha}_s) &:= f_s(\underline{X}_s + \underline{B}_s, \underline{\alpha}_s), \quad \tilde{g}(\underline{X}_T) := g(\underline{X}_T + \underline{B}_T). \end{aligned} \quad (4.1)$$

Then, recalling (2.2) and (2.3), we have

$$\begin{aligned} V_t(\underline{\xi}_t) &= \tilde{V}_t(\underline{\xi}_t - \underline{B}_t), \quad \text{where} \quad \tilde{V}_t(\underline{\xi}_t) := \inf_{\alpha \in \mathcal{A}_{[t, T]}} \tilde{g}(\tilde{X}_T^{t, \underline{\xi}_t, \alpha}) + \int_t^T \tilde{f}_s(\tilde{X}_s^{t, \underline{\xi}_t, \alpha}, \underline{\alpha}_s) ds, \\ \text{and} \quad \tilde{X}_s^{t, \underline{\xi}_t, \alpha} &= \underline{\xi}_t + \int_t^s \tilde{b}_r(\tilde{X}_r^{t, \underline{\xi}_t, \alpha}, \alpha_r, \underline{X}_r^{t, \underline{\xi}_t, \alpha}, \underline{\alpha}_s) ds, \quad s \in [t, T]. \end{aligned} \quad (4.2)$$

This is a deterministic control problem, and thus (3.5) becomes a first order equation:

$$\tilde{\mathcal{L}}\tilde{U}_t(\underline{\xi}_t) := \partial_t \tilde{U}_t(\underline{\xi}_t) + \inf_{\alpha_t \in \mathcal{A}_t} \mathbb{E} \left[\tilde{b}_t(\xi_t, \alpha_t, \underline{\xi}_t, \underline{\alpha}_t) \cdot \partial_X \tilde{U}_t(\underline{\xi}_t) + \tilde{f}_t(\underline{\xi}_t, \underline{\alpha}_t) \right] = 0. \quad (4.3)$$

Since (4.3) does not involve $\partial_{xX} \tilde{U}$, it suffices to consider $\tilde{U} \in C^{1,1}([0, T] \times \mathbb{L}^2(\mathcal{F}_T))$. That is, there exist appropriate $\partial_t \tilde{U}$ and $\partial_X \tilde{U}$ satisfying the chain rule:

$$d\tilde{U}_t(\underline{X}_t) = \partial_t \varphi_t(\underline{X}_t) dt + \mathbb{E} \left[\partial_X \tilde{U}_t(\underline{X}_t) \cdot \beta_t dt \right], \quad \text{where } dX_t = \beta_t dt, \quad \text{and } \beta \in \mathbb{L}^p(\mathbb{F}, \mathbb{R}^d). \quad (4.4)$$

We now investigate the comparison principle for appropriately defined viscosity solution for the first order equation (4.3). Let \tilde{U}_1, \tilde{U}_2 be a viscosity sub and supersolution, respectively. As standard for first order equations, we consider the doubling variable approach:

$$\tilde{\psi}(t, \underline{\xi}_t, s, \underline{\zeta}_s) := \tilde{U}_1(t, \underline{\xi}_t) - \tilde{U}_2(s, \underline{\zeta}_s) - n \left[|t - s|^2 + \mathbb{E}[|\xi_t - \zeta_s|^2] \right] - \dots, \quad (4.5)$$

where “ \dots ” indicates appropriate further penalty functions so as to guarantee the existence of a maximizer, denoted as $(\hat{t}, \hat{s}, \hat{\xi}_{\hat{t}}, \hat{\zeta}_{\hat{s}})$. Then we shall consider a test function of \tilde{U}_1 in the form (omitting possibly additional terms):

$$\tilde{\varphi}_1(t, \underline{\xi}_t) := n \left[|t - \hat{s}|^2 + \mathbb{E}[|\xi_t - \hat{\zeta}_{\hat{s}}|^2] \right]. \quad (4.6)$$

Indeed, by this argument one can easily prove rigorously the comparison principle for (4.3).

We now turn back to (3.5). Recall (4.2), it is natural to consider $U_1(t, \underline{\xi}_t) := \tilde{U}_1(t, \underline{\xi}_t - \underline{B}_t)$ as a viscosity subsolution of (3.5), with test function induced by (4.6):

$$\varphi_1(t, \underline{\xi}_t) = \tilde{\varphi}_1(t, \underline{\xi}_t - \underline{B}_t) = n \left[|t - \hat{s}|^2 + \mathbb{E}[|\xi_t - B_t - \hat{\zeta}_{\hat{s}}|^2] \right]. \quad (4.7)$$

The above φ_1 has two features:

- (i) φ_1 is not law invariant, namely $\mathbb{P}_{\xi_t} = \mathbb{P}_{\xi'_t}$ does not imply $\varphi_1(t, \underline{\xi}_t) = \varphi_1(t, \underline{\xi}'_t)$. This is another motivation for us to consider functions on the process space $\overline{\mathcal{X}}_{[0,T]}^2$ directly, instead of the Wasserstein space of probability measures.
- (ii) φ_1 is not in $C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^2)$ in the sense of Definition 3.1. Indeed, assume $t \geq \hat{t} \vee \hat{s}$ and $dX_s = \beta_s ds + \gamma_s dB_s$, $s \geq t$, then

$$\frac{d}{ds} \varphi_1(s, \underline{X}_s) = 2n(s - \hat{s}) + 2n \mathbb{E}[(X_s - B_s - \hat{\zeta}_s) \cdot \beta_s + \frac{1}{2}(\gamma_s - I_{d \times d}) : (\gamma_s - I_{d \times d})^\top]. \quad (4.8)$$

Comparing this with (3.2), we see that $\partial_{\mathbf{x}X} \varphi_1$ does not exist and thus $\varphi_1 \notin C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^2)$. However, from (4.8) it is clear that $s \mapsto \varphi_1(s, \underline{X}_s)$ is absolutely continuous. This motivates us to consider test functions which are absolutely continuous in t but are not in $C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^2)$.

4.2 The singular component of test functions in the general case

We now introduce the singular component of test functions, denoted as ϕ , which is absolutely continuous in t but is not in $C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^2)$ in general. This part ϕ is new in the literature and is the main feature of our notion of viscosity solutions due to its crucial role in our proof of comparison principle.

For $\tilde{t} \in [0, T]$ and $p \geq 1$, let $\Xi_{\tilde{t}}^p$ denote the set of maps $(\tilde{b}, \tilde{\sigma}, \tilde{f})$ where

$$\begin{aligned} (\tilde{b}, \tilde{\sigma}) : [\tilde{t}, T] \times A \times \mathcal{A}_T &\longrightarrow \mathbb{L}^p(\mathbb{F}_{[\tilde{t}, T]}, (\mathbb{R}^d, \mathbb{R}^{d \times d})), \quad \tilde{f} : [\tilde{t}, T] \times \mathcal{A}_T \longrightarrow \mathbb{R} \text{ are continuous} \\ \text{with } \|(\tilde{b}, \tilde{\sigma}, \tilde{f})\|_p^p &:= \mathbb{E} \left[\sup_{[\tilde{t}, T] \times A \times \mathcal{A}_T} [|\tilde{b}|^p + |\tilde{\sigma}|^p] \right] + \sup_{[\tilde{t}, T] \times \mathcal{A}_T} |\tilde{f}|^p < \infty. \end{aligned} \quad (4.9)$$

Given $(\tilde{t}, \tilde{\xi}) \in \overline{\mathcal{X}}_{[0,T]}^p$ and $(\tilde{b}, \tilde{\sigma}, \tilde{f}) \in \Xi_{\tilde{t}}^p$, we introduce the maps defined for all $(t, \xi) \in \overline{\mathcal{X}}_{[\tilde{t}, T]}^p$:

$$\mathcal{I}_t^\alpha(\xi) := \mathcal{I}_t^{\tilde{b}, \tilde{\sigma}, \tilde{t}, \tilde{\xi}, \alpha}(\xi) := \xi_t - \tilde{\xi}_{\tilde{t}} - \int_{\tilde{t}}^t \tilde{b}_s^\alpha ds - \int_{\tilde{t}}^t \tilde{\sigma}_s^\alpha dB_s, \quad F_t^\alpha := F_t^{\tilde{f}, \tilde{t}, \alpha} := \int_{\tilde{t}}^t \tilde{f}_s^\alpha ds, \quad (4.10)$$

$$\text{where } \tilde{b}_s^\alpha := \tilde{b}_s(\alpha_s, \underline{\alpha}_s), \quad \tilde{\sigma}_s^\alpha := \tilde{\sigma}_s(\alpha_s, \underline{\alpha}_s), \quad \tilde{f}_s^\alpha := \tilde{f}_s(\underline{\alpha}_s).$$

We now introduce the class of singular test functions inspired from (4.6) and which allows us to handle the general setting of varying and even controlled diffusion coefficient.

Definition 4.1. For $\tilde{t} \in [0, T]$ and $p \geq 4$, we denote $C^+(\overline{\mathcal{X}}_{[\tilde{t}, T]}^p)$ the set of maps of the form:

$$\phi_t(\underline{\xi}) := \inf_{\alpha \in \mathcal{A}_{[\tilde{t}, T]}} \left\{ k \mathbb{E} \left[|\mathcal{I}_t^\alpha(\xi)|^p + |\mathcal{I}_t^\alpha(\xi')|^p \right] + \int_{\tilde{t}}^t \tilde{f}_s^\alpha ds \right\}, \text{ for all } (t, \xi) \in \overline{\mathcal{X}}_{[\tilde{t}, T]}^p, \quad (4.11)$$

for some $\tilde{\xi} \in \mathcal{X}_p$, $(t', \xi') \in \overline{\mathcal{X}}_{[\tilde{t}, T]}^p$, $(\tilde{b}, \tilde{\sigma}, \tilde{f}) \in \Xi_{\tilde{t}}^p$, and some constant $k \geq 0$.

Moreover, let $C^-(\overline{\mathcal{X}}_{[\tilde{t}, T]}^p)$ denote the set of ϕ such that $-\phi \in C^+(\overline{\mathcal{X}}_{[\tilde{t}, T]}^p)$.

Remark 4.2. (i) Due to the involvement of $\tilde{\xi}, \xi'$ in (4.11), ϕ is not law invariant. This explains partially that we need to work on \mathcal{X}_p instead of the Wasserstein space $\mathcal{P}_p(\mathbb{X})$.

(ii) All the results in this paper will remain true, after obvious modifications, if we restrict the $(\tilde{b}, \tilde{\sigma}, \tilde{f})$ in (4.11) to $(b, \sigma)(\hat{\xi}_{\cdot \wedge \tilde{t}}, \cdot, \hat{\xi}_{\cdot \wedge \tilde{t}}, \cdot)$ and $f(\hat{\xi}_{\cdot \wedge \tilde{t}}, \cdot)$ for some $\hat{\xi} \in \mathcal{X}_p$. We allow for the flexibility on $(\tilde{b}, \tilde{\sigma}, \tilde{f})$ so that our definition of viscosity solution is model independent.

(iii) In the uncontrolled diffusion case: $\sigma = \sigma_t(\xi, \underline{\xi})$, we may consider the simpler set $C^+(\overline{\mathcal{X}}_{[\tilde{t}, T]}^p)$ consisting of functions $\phi_t(\xi) := \mathbb{E}[|\xi_t - \tilde{\xi}_{\tilde{t}} - \tilde{\sigma}_{\tilde{t}}(B_t - B_{\tilde{t}})|^p]$ for some $\tilde{\xi}_{\tilde{t}} \in \mathbb{L}^p(\mathcal{F}_{\tilde{t}}; \mathbb{R}^d)$ and $\tilde{\sigma}_{\tilde{t}} \in \mathbb{L}^p(\mathcal{F}_{\tilde{t}}; \mathbb{S}^d)$. In this case the related arguments can be simplified significantly.

(iii) For the estimates later, we will often use the following equivalent formulation of ϕ :

$$\phi_t(\underline{\xi}) := \inf_{\alpha \in \mathcal{A}_{[\tilde{t}, T]}} \left[k\mathbb{E}[|\mathcal{I}_t^\alpha(\underline{\xi})|^p] + F_t^\alpha + \kappa(\alpha) \right], \text{ where } \kappa(\alpha) := k\mathbb{E}[|\mathcal{I}_{t'}^\alpha(\xi')|^p] - F_{t'}^\alpha. \quad (4.12)$$

The next result states that the function $t \mapsto \phi_t$ is absolutely continuous with respect to the Lebesgue measure, and provides some crucial estimates for the comparison principle of viscosity solutions. We observe that the density $\dot{\phi}$ of ϕ is not continuous in general.

Proposition 4.3. Let $\phi \in C^+(\overline{\mathcal{X}}_{[\tilde{t}, T]}^p)$ as in (4.11) with corresponding (t', ξ') . Then, for all $X \in \mathcal{M}_{[\tilde{t}, T]}^p$, the mapping $t \mapsto \phi_t(\underline{X})$ is absolutely continuous, with density $\dot{\phi}$ satisfying:

(i) when $t \geq t'$,

$$\int_t^{t+\delta} \dot{\phi}_s(\underline{X}) ds \leq \inf_{\alpha \in \mathcal{A}_{[t, T]}} \int_t^{t+\delta} \left[\frac{p(p-1)}{2} kI_s^p(\alpha) + \tilde{f}_s^\alpha \right] ds, \quad (4.13)$$

$$\text{where } I_s^p(\alpha) := \|\beta_s - \tilde{b}_s^\alpha\|_p \sup_{\alpha' \in \mathcal{A}_{[\tilde{t}, T]}} \|\mathcal{I}_s^{\alpha'}(X)\|_p^{p-1} + \|\gamma_s - \tilde{\sigma}_s^{\tilde{\alpha}}\|_p^2 \sup_{\alpha' \in \mathcal{A}_{[\tilde{t}, T]}} \|\mathcal{I}_s^{\alpha'}(X)\|_p^{p-2};$$

(ii) alternatively, when $t < t'$, for any $\delta > 0$, there exists $\alpha^\delta \in \mathcal{A}_{[\tilde{t}, T]} \subset \mathcal{A}_{[t, T]}$, which may depend on $X_{\cdot \wedge t}$ but is independent of $(\beta_s, \gamma_s)_{s \geq t}$, such that

$$\int_t^{t+\delta} \dot{\phi}_s(\underline{X}) ds \leq \int_t^{t+\delta} \left(\frac{p(p-1)}{2} kI_s^p(\alpha^\delta) + \tilde{f}_s^{\alpha^\delta} \right) ds + \delta^2. \quad (4.14)$$

The proof is postponed to Appendix.

5 Viscosity solutions of the HJB equation

In this section we propose a notion of viscosity solutions for (3.5). We shall set $p = 6$ in Definition 4.1. For any $U : \overline{\mathcal{X}}_{[0, T]}^2 \rightarrow \mathbb{R}$ and $(t, \xi) \in \overline{\mathcal{X}}_{[0, T]}^6$, denote

$$\mathfrak{F}_6^+ U_t(\underline{\xi}) := \left\{ (\varphi, \phi) \in C^{1,2}(\overline{\mathcal{X}}_{[t, T]}^6) \times C^+(\overline{\mathcal{X}}_{[\tilde{t}, T]}^6) \text{ for some } \tilde{t} \in [0, t] : \right.$$

$$[U - (\varphi + \phi)]_t(\underline{\xi}) = \sup_{(s, \zeta) \in \bar{\mathcal{X}}_{[t, T]}^6} [U - (\varphi + \phi)]_s(\underline{\zeta}); \quad (5.1)$$

$$\begin{aligned} \mathfrak{F}_6^- U_t(\underline{\xi}) &:= \left\{ (\varphi, \phi) \in C^{1,2}(\bar{\mathcal{X}}_{[t, T]}^6) \times C^-(\bar{\mathcal{X}}_{[t, T]}^6) \text{ for some } \tilde{t} \in [0, t] : \right. \\ &\quad \left. [U - (\varphi + \phi)]_t(\underline{\xi}) = \inf_{(s, \zeta) \in \bar{\mathcal{X}}_{[t, T]}^6} [U - (\varphi + \phi)]_s(\underline{\zeta}) \right\}. \end{aligned} \quad (5.2)$$

Due to the use of the singular test functions ϕ , we need to introduce the frozen state process defined for all $(t, \xi) \in \bar{\mathcal{X}}_{[0, T]}^2$ and $\alpha \in \mathcal{A}_{[t, T]}$ by:

$$\bar{X}_s^{t, \xi, \alpha} := \xi_s, \quad s \in [0, t]; \quad \bar{X}_s^{t, \xi, \alpha} := \xi_t + \int_t^s b_r^{t, \xi, \alpha} dr + \int_t^s \sigma_r^{t, \xi, \alpha} dB_r, \quad s \in [t, T], \quad (5.3)$$

where $h_s^{t, \xi, \alpha} := h_s(\xi_{\cdot \wedge t}, \alpha_s, \underline{\xi}_{\cdot \wedge t}, \underline{\alpha}_s)$, for $h = b, \sigma$, and $f_s^{t, \xi, \alpha} := f_s(\underline{\xi}_{\cdot \wedge t}, \underline{\alpha}_s)$, $s \geq t$.

We notice the slight difference between \bar{X} and the X in (2.2). In particular, in $h_s^{t, \xi, \alpha}$ the state process is frozen while the control part α_s is evolving in s .

Definition 5.1. (i) $U \in USC(\bar{\mathcal{X}}_{[0, T]}^2)$ is a viscosity subsolution of HJB equation (3.5) if, for all $(t, \xi) \in \bar{\mathcal{X}}_{[0, T]}^6$ and $(\varphi, \phi) \in \mathfrak{F}_6^+ U_t(\underline{\xi})$,

$$\partial_t \varphi_t(\underline{\xi}) + \liminf_{\delta \rightarrow 0} \inf_{\alpha \in \mathcal{A}_{[t, T]}} \frac{1}{\delta} \int_t^{t+\delta} [H_s(\underline{\xi}_{\cdot \wedge t}, \partial_X \varphi_t(\underline{\xi}), \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}), \underline{\alpha}_s) + \dot{\phi}_s(\bar{X}^{t, \xi, \alpha})] ds \geq 0. \quad (5.4)$$

(ii) $U \in LSC(\bar{\mathcal{X}}_{[0, T]}^2)$ is a viscosity supersolution of HJB equation (3.5) if, for all $(t, \xi) \in \bar{\mathcal{X}}_{[0, T]}^6$ and $(\varphi, \phi) \in \mathfrak{F}_6^- U_t(\underline{\xi})$,

$$\partial_t \varphi_t(\underline{\xi}) + \limsup_{\delta \rightarrow 0} \inf_{\alpha \in \mathcal{A}_{[t, T]}} \frac{1}{\delta} \int_t^{t+\delta} [H_s(\underline{\xi}_{\cdot \wedge t}, \partial_X \varphi_t(\underline{\xi}), \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}), \underline{\alpha}_s) + \dot{\phi}_s(\bar{X}^{t, \xi, \alpha})] ds \leq 0. \quad (5.5)$$

(iii) $U \in C^0(\bar{\mathcal{X}}_{[0, T]}^2)$ is a viscosity solution of HJB equation (3.5) if it is both a viscosity subsolution and a viscosity supersolution of (3.5).

Remark 5.2. (i) While in quite different forms, our idea of introducing the singular component ϕ is also inspired by works on viscosity solutions for PDEs in infinite dimensions, see, e.g., Crandall-Lions [23, Definition 2.1].

(ii) The standard definition of viscosity solutions amounts to setting $\phi = 0$. So a viscosity solution in our sense is always a viscosity solution in the standard sense. Consequently, by introducing the component ϕ , the existence of viscosity solutions becomes slightly harder, but as we will see it significantly helps for the comparison principle of viscosity solutions.

(iii) We take the integral form in the left side of (5.4) and (5.5) because $\dot{\phi}$ is discontinuous, in general. If it were continuous, then these expressions would reduce to a simpler and more standard form under Assumption 3.7:

$$\partial_t \varphi_t(\underline{\xi}) + \inf_{\alpha \in \mathcal{A}_{[t, T]}} [H_t(\underline{\xi}, \partial_X \varphi_t(\underline{\xi}), \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}), \underline{\alpha}_t) + \dot{\phi}_t(\bar{X}^{t, \xi, \alpha})].$$

Remark 5.3. (i) In light of (7.2) below, it is convenient to choose p as an even integer. Some estimates later will fail for $p = 4$, so we choose $p = 6$. Any larger even integer will serve for our purpose as well.

(ii) We shall prove the comparison principle only on $\overline{\mathcal{X}}_{[0,T]}^6$. Since \mathcal{X}_6 is dense in \mathcal{X}_2 , then by the continuity of U under $\|\cdot\|_2$ we obtain the comparison principle on $\overline{\mathcal{X}}_{[0,T]}^2$.

(iii) We should note that \mathcal{X}_6 is not compact under $\|\cdot\|_6$. We shall circumvent this difficulty by using the Ekeland-Borwein-Preiss variational principle.

We first state that our notion of viscosity solution is consistent with the classical solution.

Proposition 5.4. *Let Assumptions 2.1 and 3.7 hold true and $U \in C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^2)$. Then U is a viscosity subsolution (resp. supersolution) of (3.5) if and only if it is a classical subsolution (resp. supersolution) of (3.5).*

The proof of this result is rather standard, and thus is postponed to Appendix. Our main result of the paper is the following characterization of the value function by means of the corresponding HJB equation (3.5).

Theorem 5.5. *Under Assumption 2.1, the value function V is the unique viscosity solution of the equation (3.5) with terminal condition $V_T = g$ in the class of functions satisfying (2.6).*

The viscosity property of the value function V follows standard arguments and will be reported below. The uniqueness is as usual more challenging, and is a consequence of the following comparison result.

Theorem 5.6. *Let Assumption 2.1 hold true, and $U_1 \in USC(\overline{\mathcal{X}}_{[0,T]}^2)$, $U_2 \in LSC(\overline{\mathcal{X}}_{[0,T]}^2)$ be a viscosity subsolution and supersolution of HJB equation (3.5), respectively. Assume further that there exists a modulus of continuity function ρ_R for each $R > 0$ such that*

- (i) *one of U_1, U_2 satisfies the estimates in (2.6),*
- (ii) *and the other one satisfies for $h = U_1$ or $-U_2$, and for any $R > 0$:*

$$|h_t(\underline{\xi})| \leq C(1 + \|\underline{\xi}_{\cdot \wedge t}\|_2), \quad h_t(\underline{\xi}) - h_s(\underline{\xi}_{\cdot \wedge t}) \leq \rho_R(s - t), \quad \text{for all } t < s, \|\underline{\xi}_{\cdot \wedge t}\|_2 \leq R. \quad (5.6)$$

Then $U_1(T, \cdot) \leq U_2(T, \cdot)$, on \mathcal{X}_2 , implies that $U_1 \leq U_2$ on $\overline{\mathcal{X}}_{[0,T]}^2$.

We defer the proof of the last theorem to Sections 7, 8, and 9. This theorem also implies the following comparison result immediately.

Theorem 5.7. *Let Assumption 2.1 hold true, and $U_1 \in USC(\overline{\mathcal{X}}_{[0,T]}^2)$, $U_2 \in LSC(\overline{\mathcal{X}}_{[0,T]}^2)$ be a viscosity subsolution and supersolution of (3.5), respectively. Assume U_1, U_2 satisfy the estimates (5.6), and $U_1(T, \cdot) \leq g \leq U_2(T, \cdot)$ on \mathcal{X}_2 . Then $U_1 \leq U_2$ on $\overline{\mathcal{X}}_{[0,T]}^2$.*

Proof. Note that the value function V is a viscosity solution of (3.5) with terminal condition $V_T = g$ and satisfies (2.6). Then, by applying Theorem 5.6 on U_1 and V we have $U_1 \leq V$, and by applying Theorem 5.6 on V and U_2 we have $V \leq U_2$. Thus $U_1 \leq U_2$. ■

Remark 5.8. (i) *We remark that we will not use the Ishii's lemma in the proof of Theorem 5.6, despite that our control problem involves the diffusion term. This is not completely surprising because, as we explained in Subsection 4.1, the singular part of the test function ϕ is essentially to cancel the diffusion term.*

(ii) *In the setting of Remark 3.5, our HJB equation (3.5) reduces to standard equations (3.7) or (3.8). Defining v as a viscosity solution of (3.7) or (3.8) if V defined in (3.6) is a viscosity solution of the equation (3.5), then we obtain the comparison principle for the fully nonlinear second order (path dependent) PDE without using the Ishii's lemma.*

(iii) *However, we shall emphasize that the above definition of viscosity solution is not equivalent to the standard notion of viscosity solutions for standard HJB equations, e.g. in [22]. So we are not claiming that we can avoid the Ishii's lemma for the standard viscosity solutions for fully nonlinear second order HJB equations.*

(iv) *We also refer to Remark 6.9 below for a highly related comment.*

Proof of Theorem 5.5. As the uniqueness is implied by the comparison result of Theorem 5.6, we only focus on the existence part.

(i) We first prove the viscosity subsolution property. Fix $(t, \xi) \in \bar{\mathcal{X}}_{[0, T]}^6$ and $(\varphi, \phi) \in \mathfrak{F}_6^+ V_t(\underline{\xi})$. For any $\alpha \in \mathcal{A}_{[t, T]}$, recall (2.2) and denote $X^\alpha := X^{t, \xi, \alpha}$. By DPP (2.5) and (5.1) we have for any $\delta > 0$,

$$\begin{aligned} 0 &\leq V_{t+\delta}(\underline{X}^\alpha) - V_t(\underline{\xi}) + \int_t^{t+\delta} f_s(\underline{X}^\alpha, \underline{\alpha}_s) ds \\ &\leq [\varphi + \phi]_{t+\delta}(\underline{X}^\alpha) - [\varphi + \phi]_t(\underline{\xi}) + \int_t^{t+\delta} f_s(\underline{X}^\alpha, \underline{\alpha}_s) ds. \end{aligned}$$

Applying the Itô formula (3.2) on $\varphi_s(\underline{X}^\alpha)$ we have

$$\begin{aligned} 0 &\leq \frac{1}{\delta} \int_t^{t+\delta} \mathbb{E} \left[\partial_t \varphi_s(\underline{X}^\alpha) + \partial_X \varphi_s(\underline{X}^\alpha) \cdot b_s(X^\alpha, \alpha_s, \underline{X}^\alpha, \underline{\alpha}_s) \right. \\ &\quad \left. + \frac{1}{2} \partial_{\mathbf{X}\mathbf{X}} \varphi_s(\underline{X}^\alpha) : \sigma \sigma_s^\top(X^\alpha, \alpha_s, \underline{X}^\alpha, \underline{\alpha}_s) + f_s(\underline{X}^\alpha, \underline{\alpha}_s) + \dot{\phi}_s(\underline{X}^\alpha) \right] ds. \end{aligned}$$

Recall (5.3) and denote $\bar{X}^\alpha := \bar{X}^{t, \xi, \alpha}$. Note that $\phi_t(\underline{X}^\alpha) = \phi_t(\bar{X}^\alpha)$. We claim that

$$\sup_{\alpha \in \mathcal{A}_{[t, T]}} \Delta_\delta^\alpha = o(\delta), \quad \text{where} \quad \Delta_\delta^\alpha := \left| \int_t^{t+\delta} [\dot{\phi}_s(\underline{X}^\alpha) - \dot{\phi}_s(\bar{X}^\alpha)] ds \right|. \quad (5.7)$$

Together with the regularity of φ and b, σ, f , and recalling the $\psi^{t, \xi, \alpha}$ in (5.3), this implies:

$$0 \leq \inf_{\alpha \in \mathcal{A}_{[t, T]}} \frac{1}{\delta} \int_t^{t+\delta} \mathbb{E} \left[\partial_t \varphi_t(\underline{\xi}) + \partial_X \varphi_t(\underline{\xi}) \cdot b_s^{t, \xi, \alpha} + \frac{1}{2} \partial_{\mathbf{X}\mathbf{X}} \varphi_t(\underline{\xi}) : (\sigma \sigma^\top)_s^{t, \xi, \alpha} + f_s^{t, \xi, \alpha} + \dot{\phi}_s(\underline{X}^\alpha) \right] ds + o(1).$$

Send $\delta \rightarrow 0$, we obtain (5.4), namely the viscosity subsolution property of V .

We now prove (5.7). Recall (4.10) and (4.12) with corresponding $\tilde{t}, \tilde{\xi}, t', \xi', \tilde{b}, \tilde{\sigma}, \tilde{f}$ and $p = 6$. Note that $X_{\cdot \wedge t}^\alpha = \xi_{\cdot \wedge t} = \bar{X}_{\cdot \wedge t}^\alpha$, then $\phi_t(\underline{X}^\alpha) = \phi_t(\bar{X}^\alpha)$. Thus, by (4.12),

$$\begin{aligned} \Delta_\delta^\alpha &= \left| \phi_{t+\delta}(\underline{X}^\alpha) - \phi_{t+\delta}(\bar{X}^\alpha) \right| \leq k \sup_{\tilde{\alpha} \in \mathcal{A}_{[t, T]}} \Delta_\delta^{\alpha, \tilde{\alpha}}, \\ \text{where } \Delta_\delta^{\alpha, \tilde{\alpha}} &:= \left| \mathbb{E} \left[|\mathcal{I}_{t+\delta}^{\tilde{\alpha}}(X^\alpha)|^6 - |\mathcal{I}_{t+\delta}^{\tilde{\alpha}}(\bar{X}^\alpha)|^6 \right] \right|. \end{aligned} \quad (5.8)$$

Here, to distinguish with the α in (5.7), we use $\tilde{\alpha}$ in (4.12). Fix $\alpha, \tilde{\alpha}$ and denote

$$h_s^{\alpha, \tilde{\alpha}} := h_s(X^\alpha, \alpha_s, \underline{X}^\alpha, \underline{\alpha}_s) - \tilde{h}_s^{\tilde{\alpha}}, \quad \bar{h}_s^{\tilde{\alpha}} := h_s^{t, \xi, \tilde{\alpha}} - \tilde{h}_s^{\tilde{\alpha}}, \quad h = b, \sigma.$$

Recall (4.9). Then, by (2.2) we have

$$\begin{aligned} \mathbb{E} \left[|\mathcal{I}_{t+\delta}^{\tilde{\alpha}}(X^\alpha)|^6 \right] &= \mathbb{E} \left[\left| \mathcal{I}_t^{\tilde{\alpha}}(\xi) + \int_t^{t+\delta} b_s^{\alpha, \tilde{\alpha}} ds + \int_t^{t+\delta} \sigma_s^{\alpha, \tilde{\alpha}} dB_s \right|^6 \right] \\ &= \mathbb{E} \left[|\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^6 + \int_t^{t+\delta} \left[6 |\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^4 \mathcal{I}_t^{\tilde{\alpha}}(\xi) \cdot b_s^{\alpha, \tilde{\alpha}} + 3 |\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^4 |\sigma_s^{\alpha, \tilde{\alpha}}|^2 \right. \right. \\ &\quad \left. \left. + 12 |\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^2 |\mathcal{I}_t^{\tilde{\alpha}}(\xi) \sigma_s^{\alpha, \tilde{\alpha}}|^2 \right] ds \right] + o(\delta), \end{aligned}$$

where $o(\delta)$ is uniform in $\alpha, \tilde{\alpha}$. Similarly, by (5.3) we have

$$\begin{aligned} \mathbb{E} \left[|\mathcal{I}_{t+\delta}^{\tilde{\alpha}}(\bar{X}^\alpha)|^6 \right] &= \mathbb{E} \left[|\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^6 + \int_t^{t+\delta} \left[6 |\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^4 \mathcal{I}_t^{\tilde{\alpha}}(\xi) \cdot \bar{b}_s^{\tilde{\alpha}} \right. \right. \\ &\quad \left. \left. + 3 |\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^4 |\bar{\sigma}_s^{\tilde{\alpha}}|^2 + 12 |\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^2 |\mathcal{I}_t^{\tilde{\alpha}}(\xi) \bar{\sigma}_s^{\tilde{\alpha}}|^2 \right] ds \right] + o(\delta). \end{aligned}$$

Then

$$\begin{aligned} \Delta_\delta^{\alpha, \tilde{\alpha}} &\leq C \mathbb{E} \left[\int_t^{t+\delta} \left[|\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^5 |b_s^{\alpha, \tilde{\alpha}} - \bar{b}_s^{\tilde{\alpha}}| + |\mathcal{I}_t^{\tilde{\alpha}}(\xi)|^4 |\sigma_s^{\alpha, \tilde{\alpha}} + \bar{\sigma}_s^{\tilde{\alpha}}| |\sigma_s^{\alpha, \tilde{\alpha}} - \bar{\sigma}_s^{\tilde{\alpha}}| \right] ds \right] + o(\delta) \\ &\leq C \int_t^{t+\delta} \left[\|\mathcal{I}_t^{\tilde{\alpha}}(\xi)\|_6^5 \|b_s^{\alpha, \tilde{\alpha}} - \bar{b}_s^{\tilde{\alpha}}\|_6 + \|\mathcal{I}_t^{\tilde{\alpha}}(\xi)\|_6^4 \|\sigma_s^{\alpha, \tilde{\alpha}} + \bar{\sigma}_s^{\tilde{\alpha}}\|_6 \|\sigma_s^{\alpha, \tilde{\alpha}} - \bar{\sigma}_s^{\tilde{\alpha}}\|_6 \right] ds + o(\delta). \end{aligned}$$

Note that, for $s \in [t, t + \delta]$, by Assumption 2.1 and standard SDE estimates,

$$\begin{aligned} \|\mathcal{I}_t^{\tilde{\alpha}}(\xi)\|_6 &\leq C \left(\|\xi_t\|_6 + \|\tilde{\xi}_t\|_6 + \|(\tilde{b}, \tilde{\sigma}, \tilde{f})\|_6 \right); \\ \|\sigma_s^{\alpha, \tilde{\alpha}} + \bar{\sigma}_s^{\tilde{\alpha}}\|_6 &\leq C \left(1 + \|X_{\cdot \wedge s}^\alpha\|_6 + \|\bar{X}_{\cdot \wedge s}^\alpha\|_6 + \|(\tilde{b}, \tilde{\sigma}, \tilde{f})\|_6 \right) \leq C \left(1 + \|\xi_{\cdot \wedge t}\|_6 + \|(\tilde{b}, \tilde{\sigma}, \tilde{f})\|_6 \right); \\ \|b_s^{\alpha, \tilde{\alpha}} - \bar{b}_s^{\tilde{\alpha}}\|_6^6 &= \mathbb{E} \left[\left| b_s(X^\alpha, \alpha_s, \underline{X}^\alpha, \underline{\alpha}_s) - b_s(X_{\cdot \wedge t}^\alpha, \alpha_s, \underline{X}_{\cdot \wedge t}^\alpha, \underline{\alpha}_s) \right|^6 \right] \leq C \mathbb{E} \left[|X_{\cdot \wedge s}^\alpha - X_{\cdot \wedge t}^\alpha|_\infty^6 \right] = o(1). \end{aligned}$$

Similarly, $\|\sigma_s^{\alpha, \tilde{\alpha}} - \bar{\sigma}_s^{\tilde{\alpha}}\|_6 \leq o(1)$. Then, we derive (5.7) from (5.8):

$$\Delta_\delta^\alpha \leq Ck \left(1 + \|\xi_{\cdot, \wedge t}\|_6 + \|\tilde{\xi}_t\|_6 + \|(\tilde{b}, \tilde{\sigma}, \tilde{f})\|_6\right)^5 o(\delta) + o(\delta).$$

(ii) We next prove the viscosity supersolution property. Fix $(t, \xi) \in \bar{\mathcal{X}}_{[0, T]}^6$ and $(\varphi, \phi) \in \mathfrak{F}_6^- V_t(\underline{\xi})$. For any $\delta > 0$, by DPP (2.5) there exists $\alpha^\delta \in \mathcal{A}_{[t, T]}$ such that

$$\begin{aligned} 0 &\geq V_{t+\delta}(\underline{X}^{\alpha^\delta}) - V_t(\underline{\xi}) + \int_t^{t+\delta} f_s(\underline{X}^{\alpha^\delta}, \underline{\alpha}_s^\delta) ds - \delta^2 \\ &\geq [\varphi + \phi]_{t+\delta}(\underline{X}^{\alpha^\delta}) - [\varphi + \phi]_t(\underline{\xi}) + \int_t^{t+\delta} f_s(\underline{X}^{\alpha^\delta}, \underline{\alpha}_s^\delta) ds - \delta^2. \end{aligned}$$

Now following similar arguments as in (i) we can show

$$\partial_t \varphi_t(\underline{\xi}) + \overline{\lim}_{\delta \rightarrow 0} \frac{1}{\delta} \int_t^{t+\delta} \left[H_s(\underline{\xi}_{\cdot, \wedge t}, \partial_X \varphi_t(\underline{\xi}), \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}), \underline{\alpha}_s^\delta) + \dot{\phi}_s(\bar{X}^{\alpha^\delta}) \right] ds \leq 0.$$

This implies (5.5), and hence V is a viscosity supersolution. \blacksquare

6 Connection with the HJB equation on Wasserstein space

6.1 The mean field control problem with common noise

In this section, we explain that our general setting in this paper covers the mean field control problem with common noise as a special case. For a generic Polish space (E, d_E) , we denote by $\mathcal{P}_p(E)$ the set of probability measures μ on the Borel field $\mathcal{B}(E)$ with $\int_E d_E^p(\mathbf{x}, \mathbf{x}_0) \mu(d\mathbf{x}) < \infty$, for some (and hence for all) $\mathbf{x}_0 \in E$, equipped with the p -Wasserstein distance W_p . In this section, we decompose $B^\top = ((B^1)^\top, (B^0)^\top)$, where B^0 denotes the common noise, and set $\mathbb{F}^0 = \mathbb{F}^{B^0}$. Given $\xi \in \mathcal{X}_2$, let \mathbb{P}_ξ denote the law of ξ under \mathbb{P} , and $\mathbb{P}_{\xi|\mathcal{F}_t^0}$ the conditional law of ξ , conditional on \mathcal{F}_t^0 . Moreover, for the canonical process \mathbf{X} on \mathbb{X} , define $\mu_{\cdot, \wedge t} := \mu \circ \mathbf{X}_{\cdot, \wedge t}^{-1}$, the law of the stopped process. Let $\mathcal{X}_p^\perp(\mathbb{F}_t^0)$ denote the set of $\xi \in \mathcal{X}_p$ such that $\xi_{\cdot, \wedge t}$ is independent of \mathcal{F}_t^0 , and for given $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{X})$,

$$\mathcal{X}_p^\perp(\mathbb{F}_t^0; t, \mu) := \left\{ \xi \in \mathcal{X}_p^\perp(\mathbb{F}_t^0) : \mathbb{P}_{\xi_{\cdot, \wedge t}} = \mu_{\cdot, \wedge t} \right\}.$$

Our mean field control problem with common noise involves the following data:

$$(\tilde{b}, \tilde{\sigma}, \tilde{f}) : [0, T] \times \mathbb{X} \times A \times \mathcal{P}_2(\mathbb{X} \times A) \longrightarrow (\mathbb{R}^d, \mathbb{R}^{d \times d}, \mathbb{R}), \quad \check{g} : \mathbb{X} \times \mathcal{P}_2(\mathbb{X}) \longrightarrow \mathbb{R}. \quad (6.1)$$

We emphasize that $\tilde{b}, \tilde{\sigma}$ are deterministic here. Given $(t, \xi) \in \bar{\mathcal{X}}_{[0, T]}^2$, define

$$\check{V}_t(\underline{\xi}) := \inf_{\alpha \in \mathcal{A}_{[t, T]}} \check{J}_t(\underline{\xi}, \underline{\alpha}), \quad \check{J}_t(\underline{\xi}, \underline{\alpha}) := \mathbb{E} \left[\check{g}(X, \mathbb{P}_{X|\mathcal{F}_T^0}) + \int_t^T \check{f}_s(X, \alpha_s, \mathbb{P}_{(X, \alpha_s)|\mathcal{F}_s^0}) ds \right], \quad (6.2)$$

where $X := X^{t,\xi,\alpha}$ is the controlled state process defined by the SDE:

$$X_{\cdot\wedge t} = \xi_{\cdot\wedge t}, \text{ and } dX_s = \check{b}_s(X, \alpha_s, \mathbb{P}_{(X, \alpha_s)|\mathcal{F}_s^0}) ds + \check{\sigma}_s(X, \alpha_s, \mathbb{P}_{(X, \alpha_s)|\mathcal{F}_s^0}) dB_s, \quad s \in [t, T].$$

We note that here, by writing $\check{\sigma} = (\check{\sigma}^1, \check{\sigma}^0)$, we also allow the control to act on the diffusion coefficient $\check{\sigma}^0$ of the common noise.

Assumption 6.1. (i) *The data $\check{h} = \check{b}, \check{\sigma}, \check{f}$ are progressively measurable in all variables and adapted in the spirit of (1.2): $\check{h}_t(\mathbf{x}, a, \mathbb{P}_{(\xi, \alpha_*)}) = \check{h}_t(\mathbf{x}_{\cdot\wedge t}, a, \mathbb{P}_{(\xi_{\cdot\wedge t}, \alpha_*)})$; and $\check{h}(\mathbf{0}, \cdot, \mathbb{P}_{(\mathbf{0}, \cdot)})$ and $\check{g}(\mathbf{0}, \delta_0)$ are bounded by a constant C_0 .*

(ii) *$\check{h} = \check{b}, \check{\sigma}, \check{f}, \check{g}$ are uniformly Lipschitz continuous in $(\mathbf{x}, \mu) \in \mathbb{X} \times \mathcal{P}_2(\mathbb{X})$:*

$$|\check{h}_t(\mathbf{x}, a, \mathbb{P}_{(\xi, \alpha_*)}) - \check{h}_t(\mathbf{x}', a, \mathbb{P}_{(\xi', \alpha_*)})| \leq L \left[|\mathbf{x}_{\cdot\wedge t} - \mathbf{x}'_{\cdot\wedge t}|_\infty + W_2(\mathbb{P}_{\xi_{\cdot\wedge t}}, \mathbb{P}_{\xi'_{\cdot\wedge t}}) \right].$$

(iii) *For any $R > 0$, there exists a modulus of continuity function ρ_R such that, for $\check{h} = \check{b}, \check{\sigma}, \check{f}$, for all $t \in [0, T]$, $\mathbf{x} \in \mathbb{X}$, $a, a' \in A$, $\alpha_*, \alpha'_* \in \mathcal{A}_T$, and all $\|\xi_{\cdot\wedge t}\|_2 \leq R$,*

$$|\check{h}_t(\mathbf{x}, a, \mathbb{P}_{(\xi, \alpha_*)}) - \check{h}_t(\mathbf{x}, a', \mathbb{P}_{(\xi, \alpha'_*)})| \leq \rho_R(|a - a'| + \|\alpha_* - \alpha'_*\|_2).$$

Now given $\check{b}, \check{\sigma}, \check{f}, \check{g}$, define

$$\begin{aligned} h_t(\omega, \mathbf{x}, a, \underline{\xi}, \underline{\alpha}_*) &:= \check{h}_t(\mathbf{x}, a, \mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^0}(\omega)), \quad h = b, \sigma; \\ f_t(\underline{\xi}, \underline{\alpha}_*) &:= \mathbb{E}[\check{f}_t(\xi, \alpha_*, \mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^0})], \quad g(\underline{\xi}) := \mathbb{E}[\check{g}(\xi, \mathbb{P}_{\xi|\mathcal{F}_T^0})]. \end{aligned} \tag{6.3}$$

We remark that, given $(\xi, \alpha_*) \in \mathcal{X}_2 \times \mathcal{A}_T$, we know the joint law $\mathbb{P}_{(\xi, \alpha_*, B^0)}$ and hence the conditional law $\mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^0} = \mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^{B^0}}$, so the above functions are well defined. The following result verifies that (6.2) is a special case of the problem (2.3).

Proposition 6.2. *Let $\check{b}, \check{\sigma}, \check{f}, \check{g}$ satisfy Assumption 6.1 (i), (ii), then b, σ, f, g defined by (6.3) satisfy Assumption 2.1. Moreover, $V(t, \underline{\xi}) = \check{V}(t, \underline{\xi})$ for all $(t, \underline{\xi}) \in \overline{\mathcal{X}}_{[0, T]}^2$.*

Proof. We shall only verify the properties of b . Those for σ are similar and those for f and g are slightly easier. First, the properties in Assumption 2.1 (i) are obvious. Next, by omitting the dependence on (ω, a) , we have

$$\begin{aligned} |b_t(\mathbf{x}, \underline{\xi}, \underline{\alpha}_*) - b_t(\mathbf{x}', \underline{\xi}, \underline{\alpha}_*)| &= \left| \check{b}_t(\mathbf{x}, \mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^0}) - \check{b}_t(\mathbf{x}', \mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^0}) \right| \leq L |\mathbf{x}_{\cdot\wedge t} - \mathbf{x}'_{\cdot\wedge t}|_\infty; \\ |b_t(\mathbf{x}, \underline{\xi}, \underline{\alpha}_*) - b_t(\mathbf{x}, \underline{\xi}', \underline{\alpha}_*)| &= \left| \check{b}_t(\mathbf{x}, \mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^0}) - \check{b}_t(\mathbf{x}, \mathbb{P}_{(\xi', \alpha_*)|\mathcal{F}_t^0}) \right| \\ &\leq LW_2(\mathbb{P}_{(\xi, \alpha_*)|\mathcal{F}_t^0}, \mathbb{P}_{(\xi', \alpha_*)|\mathcal{F}_t^0}) \leq L \left(\mathbb{E}_t^0[|\xi_{\cdot\wedge t} - \xi'_{\cdot\wedge t}|_\infty^2] \right)^{\frac{1}{2}}. \end{aligned}$$

Moreover, by (6.3) it is clear that $X^{t,\xi,\alpha} = \check{X}^{t,\xi,\alpha}$. Then one can easily get $J_t(\underline{\xi}, \underline{\alpha}_t) = \check{J}_t(\underline{\xi}, \underline{\alpha}_t)$, and hence $V_t(\xi) = \check{V}_t(\xi)$. \blacksquare

Proposition 6.3. *Let Assumption 6.1 hold.*

(i) \check{V} is conditional-law invariant in the sense that $\check{V}_t(\underline{\xi}) = \check{V}_t(\underline{\xi}')$ for all $\xi, \xi' \in \overline{\mathcal{X}}_{[0,T]}^2$ satisfying $\mathbb{P}_{\xi, \wedge t | \mathcal{F}_t^0} = \mathbb{P}_{\xi', \wedge t | \mathcal{F}_t^0}$, a.s. Consequently, by abusing the notation \check{V} , we may define

$$\check{V}_t(\mu) := \check{V}_t(\underline{\xi}), \quad \text{for all } (t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{X}), \quad \text{where } \xi \in \mathcal{X}_2^\perp(\mathbb{F}^0; t, \mu). \quad (6.4)$$

(ii) For any $(t, \xi) \in \overline{\mathcal{X}}_{[0,T]}^2$, we have $V_t(\underline{\xi}) = \mathbb{E}[\check{V}_t(\mathbb{P}_{\xi | \mathcal{F}_t^0})]$.

(iii) For any $0 \leq t < t + \delta \leq T$, $\xi \in \mathcal{X}_2^\perp(\mathbb{F}^0; t, \mu)$, we have

$$\check{V}_t(\mu) = \inf_{\alpha \in \mathcal{A}_{[t,T]}} \mathbb{E} \left[\check{V}_{t+\delta}(\mathbb{P}_{X^{t,\xi,\alpha} | \mathcal{F}_{t+\delta}^0}) + \int_t^{t+\delta} \check{f}_s(X^{t,\xi,\alpha}, \alpha_s, \mathbb{P}_{(X^{t,\xi,\alpha}, \alpha_s) | \mathcal{F}_s^0}) ds \right]. \quad (6.5)$$

(iv) For the constant $C = C_2$ in Lemma 2.2 (ii),

$$\begin{aligned} |\check{V}_t(\mu)| &\leq C \left(1 + W_2(\mu, \wedge t, \delta_0) \right); \\ |\check{V}_t(\mu) - \check{V}_{t'}(\mu')| &\leq C \left(W_2(\mu, \wedge t, \delta_0) + (1 + W_2(\mu, \wedge t, \delta_0)) \sqrt{|t - t'|} \right). \end{aligned} \quad (6.6)$$

This result is not surprising, for example, [18, Theorem 3.6] proved the law invariance property when there is no common noise. However, due to its generality, we believe the precise form here is new in the literature. For completeness we sketch a proof in Appendix.

To derive the PDE from the DPP (6.5), as usual we need the appropriate Itô formula. The following result can be proved by combining [14, Theorem 4.17] and [54, Theorem 2.7], see also [27]. The precise meaning of the space $C_b^{1,2,2}([0, T] \times \mathbb{X} \times \mathcal{P}_2(\mathbb{X}))$ and that of the derivatives, as well as the proof are again postponed to Appendix.

Proposition 6.4. *Assume $U \in C_b^{1,2,2}([0, T] \times \mathbb{X} \times \mathcal{P}_2(\mathbb{X}))$. For $i = 1, 2$, consider*

$$dX_t^i := b_t^i dt + \sigma_t^{i,1} dB_t^1 + \sigma_t^{i,0} dB_t^0, \quad \text{and introduce the conditional law } \mu_{\cdot, \wedge t} := \mathbb{P}_{X_{\cdot, \wedge t}^2 | \mathcal{F}_t^0},$$

where $b^i \in \mathbb{L}^2(\mathbb{F}; \mathbb{R}^d)$ and $\sigma^i = (\sigma^{i,1}, \sigma^{i,0}) \in \mathbb{L}^2(\mathbb{F}; \mathbb{R}^{d \times d})$. Then

$$\begin{aligned} dU_t(X^1, \mu) &= \left[\partial_t U_t + \partial_x U_t \cdot b_t^1 + \frac{1}{2} \partial_{xx} U_t : \sigma_t^1 (\sigma_t^1)^\top \right] (X^1, \mu) dt + \partial_x U_t(X^1, \mu) \cdot \sigma_t^1 dB_t^1 \\ &\quad + \tilde{\mathbb{E}}_{\mathcal{F}_t} [\partial_\mu U_t(X^1, \mu, \tilde{X}_t^2) \cdot \tilde{\sigma}_t^{2,0}] dB_t^0 \\ &\quad + \mathbb{E}_{\mathcal{F}_t} \left[\partial_\mu U_t(\cdot, \tilde{X}_t^2) \cdot \tilde{b}_t^2 + \frac{1}{2} \partial_{\tilde{x}} \partial_\mu U_t(\cdot, \tilde{X}_t^2) : \tilde{\sigma}_t^2 (\tilde{\sigma}_t^2)^\top \right. \\ &\quad \left. + \partial_x \partial_\mu U(\cdot, \tilde{X}_t^2) : \sigma_t^{1,0} (\tilde{\sigma}_t^{2,0})^\top + \frac{1}{2} \partial_{\mu\mu} U_t(\cdot, \tilde{X}_t^2, \tilde{X}_t^2) : \tilde{\sigma}_t^{2,0} (\tilde{\sigma}_t^{2,0})^\top \right] (X^1, \mu) dt, \end{aligned} \quad (6.7)$$

where $(\tilde{X}^2, \tilde{b}^2, \tilde{\sigma}^2)$ and $(\bar{X}^2, \bar{b}^2, \bar{\sigma}^2)$ are conditionally independent copies of (X^2, b^2, σ^2) , conditional on \mathbb{F}^0 .

Now apply Proposition 6.4 on the DPP (6.5), we obtain the following second order path dependent HJB equation on the Wasserstein space: for $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{X})$,

$$\check{\mathcal{L}}\check{U}_t(\mu) := \partial_t \check{U}_t(\mu) + \inf_{\alpha_t \in \mathcal{A}_t^\perp(\mathcal{F}_t^0)} \check{H}_t(\mu, \partial_\mu \check{U}_t(\mu, \cdot), \partial_{\tilde{x}\mu} \check{U}_t(\mu, \cdot), \partial_{\mu\mu} \check{U}_t(\mu, \cdot, \cdot), \underline{\alpha}_t) = 0, \quad (6.8)$$

where, for $\alpha_t \in \mathcal{A}_t^\perp(\mathcal{F}_t^0)$ independent of \mathcal{F}_t^0 , $\xi \in \mathcal{X}_2^\perp(\mathbb{F}^0; t, \mu)$, and letting $(\tilde{\xi}, \tilde{\alpha}_t)$, $(\bar{\xi}, \bar{\alpha}_t)$ be independent copies of (ξ, α_t) ,

$$\begin{aligned} \check{H} := \mathbb{E} & \left[\partial_\mu \check{U}_t(\mu, \tilde{\xi}_t) \cdot \check{b}_t(\tilde{\xi}, \tilde{\alpha}_t, \mathbb{P}_{(\xi, \alpha_t)}) + \frac{1}{2} \partial_{\tilde{x}\mu} \check{U}_t(\mu, \tilde{\xi}_t) : \check{\sigma}_t \check{\sigma}_t^\top(\tilde{\xi}, \tilde{\alpha}_t, \mathbb{P}_{(\xi, \alpha_t)}) \right. \\ & \left. + \frac{1}{2} \partial_{\mu\mu} \check{U}_t(\mu, \tilde{\xi}_t, \bar{\xi}_t) : \check{\sigma}_t^0(\tilde{\xi}, \tilde{\alpha}_t, \mathbb{P}_{(\xi, \alpha_t)}) (\check{\sigma}_t^0)^\top(\bar{\xi}, \bar{\alpha}_t, \mathbb{P}_{(\xi, \alpha_t)}) + \check{f}_t(\xi, \alpha_t, \mathbb{P}_{(\xi, \alpha_t)}) \right]. \end{aligned} \quad (6.9)$$

Definition 6.5. We say $\check{U} \in [0, T] \times \mathcal{P}_2(\mathbb{X}) \rightarrow \mathbb{R}$ is a viscosity solution (resp. subsolution, supersolution) of the equation (6.8)-(6.9) if $U_t(\underline{\xi}) := \mathbb{E}[\check{U}_t(\mathbb{P}_{\xi|\mathcal{F}_t^0})]$ is a viscosity solution (resp. subsolution, supersolution) of the equation (3.5), where b, σ, f, g are defined by (6.3).

The following result is a direct consequence of Theorems 5.5 and 5.6.

Theorem 6.6. Let Assumption 6.1 hold.

- (i) The \check{V} defined by (6.4) is the unique viscosity solution, in the sense of Definition 6.5, to the equation (6.8)-(6.9) with terminal condition \check{g} .
- (ii) Let \check{U}_1, \check{U}_2 be a viscosity subsolution and viscosity supersolution, respectively, to the equation (6.8)-(6.9), in the sense of Definition 6.5. Assume, for $\check{h} = \check{U}_1$ and $-\check{U}_2$, for any $R > 0$, there exists a modulus of continuity function ρ_R such that

$$|\check{h}_t(\mu)| \leq C(1 + W_2(\mu_{\cdot \wedge t}, \delta_0)); \quad \check{h}_t(\mu) - \check{h}_s(\mu_{\cdot \wedge t}) \leq \rho_R(s - t), \quad \text{for all } t < s, W_2(\mu_{\cdot \wedge t}, \delta_0) \leq R.$$

If $\check{U}_1(T, \mu) \leq \int_{\mathbb{X}} \check{g}(\mathbf{x}, \mu) \mu(d\mathbf{x}) \leq \check{U}_2(T, \mu)$ for all $\mu \in \mathcal{P}_2(\mathbb{X})$, then $\check{U}_1 \leq \check{U}_2$ on $[0, T] \times \mathcal{P}_2(\mathbb{X})$.

We shall remark that the above notion of viscosity solution is not equivalent to the definitions in the literature as mentioned in Introduction. Therefore, our results do not imply the uniqueness or comparison principle for the viscosity solutions in those publications.

Remark 6.7. Given $\check{U} \in C_b^{1,2}([0, T] \times \mathcal{P}_2(\mathbb{X}))$, similarly to the φ_1 in (4.7), in general $U_t(\underline{\xi}) := \mathbb{E}[\check{U}_t(\mathbb{P}_{\xi|\mathcal{F}_t^0})]$ may not be in $C^{1,2}(\mathcal{X}_{[0,T]}^2)$ in the sense of Definition 3.1. So, while we are identifying the viscosity solutions for (6.8)-(6.9) and (3.5), in general we cannot identify their classical solutions. The situation is different when there is no common noise, see Proposition 6.8 below.

6.2 Mean field control without common noise

When there is no common noise, recall Remark 3.3 (ii) and define the classical semisolutions of (6.8)-(6.9) in the obvious manner. The following result is obvious.

Proposition 6.8. *Let Assumption 6.1 hold. Assume there is no common noise, and $s \in [t, T] \mapsto \check{h}_s(\mathbf{x}_{\cdot \wedge t}, a, \mathbb{P}_{(X_{\cdot \wedge t}, \alpha_*)})$ is uniformly continuous, uniformly in $(\mathbf{x}, a, X, \alpha_*)$. Let $\check{U} \in C^{1,2}([0, T] \times \mathcal{P}_2(\mathbb{X}))$ and denote $U_t(\underline{\xi}) := \check{U}(t, \mathbb{P}_{\underline{\xi}})$. Then $U \in C^{1,2}(\overline{\mathcal{X}}_{[0, T]}^2)$, and U is a classical solution (resp. sub-solution, supersolution) of (3.5)-(6.3) if and only if \check{U} is a classical solution (resp. sub-solution, supersolution) of (6.8)-(6.9).*

Remark 6.9. *When there is no common noise and in the state dependent setting, [9, 50, 51] proposed different notions of viscosity solutions for HJB equations on the Wasserstein space and established the comparison principle by doubling variable argument, also without invoking the Ishii's lemma. However, their mechanism is completely different from ours.*

(i) *Our HJB equation (3.5) is a second order equation. As in (4.5) we consider the following penalization in the doubling variable argument:*

$$\psi(t, \underline{\xi}, s, \underline{\zeta}) := U_1(t, \underline{\xi}) - U_2(s, \underline{\eta}) - n\mathbb{E}[|\xi - \eta|^2] - \dots$$

Here we consider the state dependent case for simplicity and ξ, ζ denote random variables instead of processes. Assume the above has optimal arguments $(t_n, \xi_n, s_n, \zeta_n)$. By standard arguments one can easily show $\lim_{n \rightarrow \infty} n\mathbb{E}[|\xi_n - \eta_n|^2] = 0$. Then naturally we would construct test functions (omitting the other terms):

$$\varphi_1(\underline{\xi}) := n\mathbb{E}[|\xi - \zeta_n|^2], \quad \varphi_2(\underline{\zeta}) := n\mathbb{E}[|\zeta - \xi_n|^2].$$

Ignoring the possible adaptedness issue, by (3.2) we have

$$\partial_X \varphi_1(\underline{\xi}) = 2n(\xi - \zeta_n), \quad \partial_{xX} \varphi_1(\underline{\xi}) = 2n; \quad \partial_X \varphi_2(\underline{\zeta}) = 2n(\zeta - \xi_n), \quad \partial_{xX} \varphi_2(\underline{\zeta}) = 2n.$$

This implies that,

$$\partial_X \varphi_1(\underline{\xi}_n) + \partial_X \varphi_2(\underline{\zeta}_n) = 0, \quad \text{but} \quad \partial_{xX} \varphi_1(\underline{\xi}_n) = \partial_{xX} \varphi_2(\underline{\zeta}_n) \quad (6.10)$$

Then we can have the desired cancellation for the first order derivatives $\partial_X \varphi$. For example,

$$\begin{aligned} \sup_{\alpha} \mathbb{E} \left[b(\xi_n, \alpha) \partial_X \varphi_1(\underline{\xi}_n) + b(\zeta_n, \alpha) \partial_X \varphi_2(\underline{\zeta}_n) \right] &= \sup_{\alpha} \mathbb{E} \left[(b(\xi_n, \alpha) - b(\zeta_n, \alpha)) 2n(\xi_n - \zeta_n) \right] \\ &\leq Cn\mathbb{E}[|\xi_n - \eta_n|^2] \rightarrow 0. \end{aligned}$$

However, there won't be such type of cancellation for the second order derivatives $\partial_{xX}\varphi$. This is the same issue the second order derivatives $\partial_{xx}\varphi$ face for standard HJB equations, and in that case Ishii's lemma is used exactly to overcome this difficulty. We alternatively introduced the singular component ϕ , instead of Ishii's lemma, to get around of this difficulty.

(ii) The HJB equation (6.8)-(6.9) on the Wasserstein space, however, should be viewed as a first order equation (again, when there is no common noise), as [9, 50, 51] observed, even though we are talking about the same value function: $V(t, \underline{\xi}) = \check{V}(t, \mathbb{P}_{\underline{\xi}})$ by Proposition 6.3 (ii). To see this, we use the setting in [51]. Introduce a smooth distance function on $\mathcal{P}_2(\mathbb{R})$ by using Fourier transformation: assuming state dependent and 1-dimensional, for simplicity, and for some constant k and $i = \sqrt{-1}$:

$$\rho^2(\mu, \nu) := \int_{\mathbb{R}} \frac{|\mathcal{F}_{\mu-\nu}(z)|^2}{1 + |z|^k} dz, \quad \text{where } \mathcal{F}_{\mu-\nu}(z) := \int_{\mathbb{R}} e^{-iz\lambda}(\mu - \nu)(d\lambda).$$

Consider the following penalization in the doubling variable argument:

$$\check{\psi}(t, \mu, s, \nu) := \check{U}_1(t, \mu) - \check{U}_2(s, \nu) - n\rho^2(\mu, \nu) - \dots$$

and assume it has optimal arguments (t_n, μ_n, s_n, ν_n) . Then we have $\lim_{n \rightarrow \infty} n\rho^2(\mu_n, \nu_n) = 0$, and we shall similarly construct test functions (again omitting the other terms):

$$\check{\varphi}_1(\mu) := n\rho^2(\mu, \nu_n), \quad \check{\varphi}_2(\nu) := n\rho^2(\mu_n, \nu).$$

By direct calculation, we have: denoting by Re the real part,

$$\begin{aligned} \partial_{\mu}\check{\varphi}_1(\mu, x) &= 2n \int_{\mathbb{R}} \frac{\text{Re}\left(ize^{izx}\mathcal{F}_{\mu-\nu_n}(z)\right)}{1 + |z|^k} dz, & \partial_{x\mu}\check{\varphi}_1(\mu, x) &= -2n \int_{\mathbb{R}} \frac{\text{Re}\left(|z|^2e^{izx}\mathcal{F}_{\mu-\nu_n}(z)\right)}{1 + |z|^k} dz; \\ \partial_{\nu}\check{\varphi}_2(\nu, x) &= 2n \int_{\mathbb{R}} \frac{\text{Re}\left(ize^{izx}\mathcal{F}_{\nu-\mu_n}(z)\right)}{1 + |z|^k} dz, & \partial_{x\nu}\check{\varphi}_2(\nu, x) &= -2n \int_{\mathbb{R}} \frac{\text{Re}\left(|z|^2e^{izx}\mathcal{F}_{\nu-\mu_n}(z)\right)}{1 + |z|^k} dz. \end{aligned}$$

This implies that, unlike (6.10) for the second equality,

$$\partial_{\mu}\check{\varphi}_1(\mu_n, x) + \partial_{\nu}\check{\varphi}_2(\nu_n, x) = 0, \quad \text{and} \quad \partial_{x\mu}\check{\varphi}_1(\mu_n, x) + \partial_{x\nu}\check{\varphi}_2(\nu_n, x) = 0. \quad (6.11)$$

So one may expect the desired cancellation for the terms involving $\partial_{x\mu}\check{\varphi}$, and in this sense the HJB equation should be viewed as a first order equation. However, we should note that the infinite dimensionality, in the sense that $\partial_{x\mu}\check{\varphi}(\mu, \cdot)$ is a function of x , will cause additional difficulty than the standard finite dimensional HJB equations.

(iii) When there is common noise, the HJB equation (6.8)-(6.9) involves $\partial_{\mu\mu}\check{U}$. Note that

$$\partial_{\mu\mu}\check{\varphi}_1(\mu, x, \tilde{x}) = 2n \int_{\mathbb{R}} \frac{\text{Re}\left(|z|^2e^{iz(x-\tilde{x})}\right)}{1 + |z|^k} dz = \partial_{\mu\mu}\check{\varphi}_2(\nu, x, \tilde{x}).$$

Then we encounter the same difficulty caused by $\partial_{xx}\varphi$ in the standard case. We remark that [3] introduced an Ishii's Lemma in this setting to overcome this difficulty.

7 The comparison result: Doubling variable in space

Note that the space \mathcal{X}_6 is not compact. To prove the comparison principle without compactness, we shall use the following variation of the Ekeland-Borwein-Preiss variational principle, see [8]. For this purpose, we first recall the definition of gauge-type function. Let (E, d) be a generic complete metric space.

Definition 7.1. *A function $\Upsilon \in C^0(E^2; [0, \infty))$ is called a gauge-type function on (E, d) if*

$$\Upsilon(x, x) = 0 \text{ for all } x \in E, \text{ and } \lim_{\Upsilon(x_1, x_2) \rightarrow 0} d(x_1, x_2) = 0.$$

Lemma 7.2. *Let $\psi : E \rightarrow \mathbb{R}$ be upper semicontinuous and bounded from above, and Υ a gauge-type function on (E, d) . For any $\varepsilon > 0$ and $x_0 \in E$ satisfying $\psi(x_0) \geq \sup_E \psi - \varepsilon$, there exist $\hat{x} \in E$ and $\{x_i\}_{i \geq 1} \subset E$ such that, denoting $\Psi := \psi - \sum_{i=0}^{\infty} 2^{-i} \Upsilon(\cdot, x_i) \leq \psi$,*

$$\Upsilon(x_i, \hat{x}) \leq \varepsilon 2^{-i}, \text{ for all } i \geq 0, \text{ and } \Psi(\hat{x}) = \text{strict-max}_E \Psi \geq \psi(x_0) \geq \sup_E \psi - \varepsilon.$$

7.1 Smooth gauge-type functions

We now consider the following complete metric space:

$$\Lambda_0 := \{\theta = (t, \xi_{\cdot \wedge t}) : (t, \xi) \in \overline{\mathcal{X}}_{[0, T]}^6\}, \quad d_0(\underline{\theta}, \underline{\theta}') := |t - t'| + \|\xi_{\cdot \wedge t} - \xi'_{\cdot \wedge t'}\|_6. \quad (7.1)$$

The metric d_0 itself is of course a gauge-type function. However, for fixed θ' , the mapping $\theta \mapsto d_0(\underline{\theta}, \underline{\theta}')$ is not smooth in the sense of Definition 3.1. In order to construct smooth test functions later, we need smooth gauge-type functions. For this purpose, we introduce

$$\Upsilon_t(\mathbf{x}) := \frac{(|\mathbf{x}_{\cdot \wedge t}|_{\infty}^6 - |\mathbf{x}_t|^6)^3}{|\mathbf{x}_{\cdot \wedge t}|_{\infty}^{12}} \mathbf{1}_{\{|\mathbf{x}_{\cdot \wedge t}|_{\infty} \neq 0\}} + 3|\mathbf{x}_t|^6, \quad (t, \mathbf{x}) \in [0, T] \times \mathbb{X}; \quad (7.2)$$

$$\Upsilon_0(\underline{\theta}, \underline{\theta}') := \mathbb{E}[\Upsilon_{t \vee t'}(\xi_{\cdot \wedge t} - \xi'_{\cdot \wedge t'})], \text{ and } \bar{\Upsilon}_0(\underline{\theta}, \underline{\theta}') := \Upsilon_0(\underline{\theta}, \underline{\theta}') + |t - t'|^2, \quad \theta, \theta' \in \Lambda_0.$$

We emphasize that the map Υ is defined on deterministic paths, rather than on processes, consequently, their path derivatives should be understood in the sense of Dupire [26]. We also recall from [56] the useful bound on Υ and the triangle-like inequality

$$|\mathbf{x}_{\cdot \wedge t}|_{\infty}^6 \leq \Upsilon_t(\mathbf{x}) \leq 3|\mathbf{x}_{\cdot \wedge t}|_{\infty}^6, \quad \Upsilon_t(\mathbf{x} + \mathbf{x}') \leq 32[\Upsilon_t(\mathbf{x}) + \Upsilon_t(\mathbf{x}')], \quad t \in [0, T], \mathbf{x}, \mathbf{x}' \in \mathbb{X}. \quad (7.3)$$

Moreover, Υ is differentiable in t and twice continuously differentiable in \mathbf{x} with:

$$\partial_t \Upsilon_t(\mathbf{x}) = 0, \quad |\partial_{\mathbf{x}} \Upsilon_t(\mathbf{x})| \leq 18|\mathbf{x}_t|^5, \quad \text{and} \quad |\partial_{\mathbf{x}\mathbf{x}} \Upsilon_t(\mathbf{x})| \leq 306|\mathbf{x}_t|^4. \quad (7.4)$$

Combining this with Example 3.4, we have the following result.

Lemma 7.3. (i) $\bar{\Upsilon}_0$ is a gauge-type function on (Λ_0, d_0) .

(ii) For any $\theta' = (t', \xi') \in \bar{\mathcal{X}}_{[0,T]}^6$, the mapping $\theta \in \bar{\mathcal{X}}_{[t',T]}^6 \mapsto \Upsilon_0(\theta, \theta')$ is in $C^{1,2}(\bar{\mathcal{X}}_{[t',T]}^6)$ such that, for any $\hat{t} \geq t'$, $X \in \mathcal{M}_{[\hat{t},T]}^6$ as in (3.1), and $t \in [\hat{t}, T]$, we have

$$d\Upsilon_0((t, \underline{X}), \underline{\theta}') = \mathbb{E} \left[\partial_{\mathbf{x}} \Upsilon_t(X - \xi'_{\wedge t'}) \cdot dX_t + \frac{1}{2} \partial_{\mathbf{xx}} \Upsilon_t(X - \xi'_{\wedge t'}) : d\langle X \rangle_t \right]. \quad (7.5)$$

Remark 7.4. In the state dependent case $U_t(\underline{\xi}) = U_t(\underline{\xi}_t)$, we may consider a much simpler smooth gauge-type function: $\bar{\Upsilon}_0((t, \underline{\xi}_t), (t', \underline{\xi}'_t)) := |t - t'|^2 + \|\xi_t - \xi'_t\|_2^2$. The related calculations will also be simplified significantly. However, our approach still requires the singular component ϕ in the test functions, so the main arguments for the comparison principle will remain the same. See also Remark 5.3 concerning ϕ in the case without volatility control.

We next extend the space (Λ_0, d_0) by doubling the spatial variable:

$$\begin{aligned} \Lambda_1(\hat{t}) &:= \{ \lambda := (t, \xi_{\wedge t}, \zeta_{\wedge t}) : t \in [\hat{t}, T] \text{ and } \xi, \zeta \in \mathcal{X}_6 \}, \text{ for all } \hat{t} \in [0, T], \\ d_1(\underline{\lambda}, \underline{\lambda}') &:= d_0((t, \underline{\xi}), (t', \underline{\xi}')) + d_0((t, \underline{\zeta}), (t', \underline{\zeta}')), \end{aligned} \quad (7.6)$$

$$\Upsilon_1(\underline{\lambda}, \underline{\lambda}') := \Upsilon_0((t, \underline{\xi}), (t', \underline{\xi}')) + \Upsilon_0((t, \underline{\zeta}), (t', \underline{\zeta}')), \quad \bar{\Upsilon}_1(\underline{\lambda}, \underline{\lambda}') := \Upsilon_1(\underline{\lambda}, \underline{\lambda}') + |t - t'|^2.$$

Similarly, we extend further the space by doubling the temporal and spatial variables:

$$\begin{aligned} \Lambda_2(\hat{t}, \hat{s}) &:= \{ \iota := (\theta, \eta) : \theta = (t, \xi_{\wedge t}) \in \bar{\mathcal{X}}_{[\hat{t},T]}^6, \eta = (s, \zeta_{\wedge s}) \in \bar{\mathcal{X}}_{[\hat{s},T]}^6 \}, \quad \hat{t}, \hat{s} \in [0, T], \\ d_2(\underline{\iota}, \underline{\iota}') &:= d_0(\underline{\theta}, \underline{\theta}') + d_0(\underline{\eta}, \underline{\eta}'), \end{aligned} \quad (7.7)$$

$$\Upsilon_2(\underline{\iota}, \underline{\iota}') := \Upsilon_0(\underline{\theta}, \underline{\theta}') + \Upsilon_0(\underline{\eta}, \underline{\eta}'), \text{ and } \bar{\Upsilon}_2(\underline{\iota}, \underline{\iota}') := \bar{\Upsilon}_0(\underline{\theta}, \underline{\theta}') + \bar{\Upsilon}_0(\underline{\eta}, \underline{\eta}').$$

Clearly $\bar{\Upsilon}_1$ and $\bar{\Upsilon}_2$ are gauge-type functions on $(\Lambda_1(\hat{t}), d_1)$ and $(\Lambda_2(\hat{t}, \hat{s}), d_2)$, respectively.

In this and next section, we prove the comparison principle by using the doubling variable arguments. In $(\Lambda_1(\hat{t}), d_1)$ we double the spatial variables only, while in $(\Lambda_2(\hat{t}, \hat{s}), d_2)$ we double both the spatial and the temporal variables. From now on we consider the setting in Theorem 5.6 and assume all the conditions there hold true. Moreover, we shall assume without loss of generality that U_1 satisfies (2.6) and, denoting:

$$m := \sup_{\theta \in \bar{\mathcal{X}}_{[0,T]}^6} [U_1 - U_2](\theta),$$

we shall assume to contrary that $m > 0$, and work toward a contradiction.

7.2 Doubling variable in space

For any $n \geq 1$ and $\varepsilon > 0$, introduce the function $\psi^{n,\varepsilon}$ defined for $\lambda = (t, \xi, \zeta) \in \Lambda_1(0)$ by:

$$\begin{aligned} \psi^{n,\varepsilon}(\underline{\lambda}) &= \psi_t^{n,\varepsilon}(\underline{\xi}, \underline{\zeta}) := U_1(t, \underline{\xi}) - U_2(t, \underline{\zeta}) - n\mathbb{E}[\Upsilon_t(\xi - \zeta)] - n^{\frac{4}{3}}\mathbb{E}[|\xi_t - \zeta_t|^6] \\ &\quad - \varepsilon \left(1 - \frac{t}{2T}\right) \mathbb{E}[\Upsilon_t(\zeta)]. \end{aligned} \quad (7.8)$$

Proposition 7.5. *Let all the conditions in Theorem 5.6 hold true. Assume U_1 satisfies (2.6), and assume to the contrary that $m > 0$. Then, there exist $0 < \varepsilon_m \leq 1$ such that, for all $0 < \varepsilon \leq \varepsilon_m$, the following hold.*

(i) *There exist $\hat{\lambda}^{n,\varepsilon} = (\hat{t}^{n,\varepsilon}, \hat{\xi}^{n,\varepsilon}, \hat{\zeta}^{n,\varepsilon}) \in \Lambda_1(0)$ and $\Psi^{n,\varepsilon} \in C^0(\Lambda_1(\hat{t}^{n,\varepsilon}))$ s.t. $\Psi^{n,\varepsilon} \leq \psi^{n,\varepsilon}$ and*

$$\Psi^{n,\varepsilon}(\hat{\Delta}^{n,\varepsilon}) = \max_{\lambda \in \Lambda_1(\hat{t}^{n,\varepsilon})} \Psi^{n,\varepsilon}(\lambda) \geq \sup_{\lambda \in \Lambda_1(0)} \psi^{n,\varepsilon}(\lambda) - \frac{1}{n} \geq \frac{m}{2} - \frac{1}{n}. \quad (7.9)$$

(ii) *Moreover, $[\psi^{n,\varepsilon} - \Psi^{n,\varepsilon}](\lambda) = \pi_0^{n,\varepsilon}(t) + \pi_1^{n,\varepsilon}(t, \underline{\xi}) + \pi_2^{n,\varepsilon}(t, \underline{\zeta})$, for some nonnegative functions $\pi_0^{n,\varepsilon} \in C^1([\hat{t}^{n,\varepsilon}, T])$, $\pi_1^{n,\varepsilon}, \pi_2^{n,\varepsilon} \in C^{1,2}(\overline{\mathcal{X}}_{[\hat{t}^{n,\varepsilon}, T]}^6)$, such that: at $\hat{\lambda}^{n,\varepsilon}$,*

$$\begin{aligned} 0 &\leq \pi_j^{n,\varepsilon} \leq Cn^{-1}, \quad j = 0, 1, 2; \quad |\partial_t \pi_0^{n,\varepsilon}| \leq Cn^{-\frac{1}{2}}; \\ \partial_t \pi_j^{n,\varepsilon} &= 0, \quad \|\partial_X \pi_j^{n,\varepsilon}\|_{\frac{6}{5}} \leq Cn^{-\frac{5}{6}}, \quad \|\partial_{\mathbf{x}X} \pi_j^{n,\varepsilon}\|_{\frac{3}{2}} \leq Cn^{-\frac{2}{3}}, \quad j = 1, 2. \end{aligned} \quad (7.10)$$

(iii) *There exists $C_\varepsilon > 0$, which may depend on ε but not on n , such that*

$$\mathbb{E} \left[|\hat{\xi}_{\cdot, \wedge \hat{t}^{n,\varepsilon}}^{n,\varepsilon}|_{\infty}^6 + |\hat{\zeta}_{\cdot, \wedge \hat{t}^{n,\varepsilon}}^{n,\varepsilon}|_{\infty}^6 \right] \leq C_\varepsilon, \quad \text{and} \quad (7.11)$$

$$n \mathbb{E} \left[|\hat{\xi}_{\cdot, \wedge \hat{t}^{n,\varepsilon}}^{n,\varepsilon} - \hat{\zeta}_{\cdot, \wedge \hat{t}^{n,\varepsilon}}^{n,\varepsilon}|_{\infty}^6 \right] + n^{\frac{3}{2}} \mathbb{E} \left[|\hat{\xi}_{\hat{t}^{n,\varepsilon}}^{n,\varepsilon} - \hat{\zeta}_{\hat{t}^{n,\varepsilon}}^{n,\varepsilon}|^6 \right] \leq Cn^{-\frac{1}{30}}, \quad \text{for all } n \geq 1. \quad (7.12)$$

(iv) $\overline{T}_{\varepsilon,m} := \overline{\lim}_{n \rightarrow \infty} \hat{t}^{n,\varepsilon} < T$. *Consequently, there exists $n_{\varepsilon,m}$, which may depend on ε but not on n , such that $\hat{t}^{n,\varepsilon} \leq T_{\varepsilon,m} := \frac{1}{2}(\overline{T}_{\varepsilon,m} + T) < T$, for all $n \geq n_{\varepsilon,m}$.*

Proof. (i) As $m > 0$, there exists $\theta_0 = (t_0, \xi^0) \in \Lambda_0$ s.t. $(U_1 - U_2)(\underline{\theta}_0) \geq \frac{2m}{3}$. Then

$$\sup_{\Lambda_1(0)} \psi^{n,\varepsilon} \geq \psi^{n,\varepsilon}(t_0, \underline{\xi}^0, \underline{\xi}^0) = (U_1 - U_2)(\underline{\theta}_0) - \varepsilon \left(1 - \frac{t_0}{2T}\right) \mathbb{E}[\Upsilon(\theta_0)] \geq \frac{2m}{3} - 3\varepsilon \mathbb{E}[|\xi_{\cdot, \wedge t_0}^0|_{\infty}^6],$$

where the last inequality is due to (7.3). By setting $\varepsilon_m := \frac{m}{1 + 18\mathbb{E}[|\xi_{\cdot, \wedge t_0}^0|_{\infty}^6]}$, we obtain the last inequality in (7.9). Next, by (7.3) we have

$$\mathbb{E}[\Upsilon_t(\zeta) + \Upsilon_t(\xi - \zeta)] \geq \frac{1}{33} \mathbb{E}[\Upsilon_t(\xi) + \Upsilon_t(\zeta)] \geq \frac{1}{33} [\|\xi_{\cdot, \wedge t}\|_6^6 + \|\zeta_{\cdot, \wedge t}\|_6^6].$$

Note that $n \geq 1 \geq \varepsilon$ and $1 - \frac{t}{2T} \geq \frac{1}{2}$, then

$$n \mathbb{E}[\Upsilon_t(\xi - \zeta)] + \varepsilon \left(1 - \frac{t}{2T}\right) \mathbb{E}[\Upsilon_t(\zeta)] \geq \frac{\varepsilon}{2} \mathbb{E}[\Upsilon_t(\zeta) + \Upsilon_t(\xi - \zeta)] \geq \frac{\varepsilon}{66} [\|\xi_{\cdot, \wedge t}\|_6^6 + \|\zeta_{\cdot, \wedge t}\|_6^6].$$

Thus, since U_1, U_2 satisfy the estimates (5.6),

$$\psi^{n,\varepsilon}(\lambda) \leq C[1 + \|\xi_{\cdot, \wedge t}\|_2 + \|\zeta_{\cdot, \wedge t}\|_2] - \frac{\varepsilon}{66} [\|\xi_{\cdot, \wedge t}\|_6^6 + \|\zeta_{\cdot, \wedge t}\|_6^6]. \quad (7.13)$$

This clearly implies $\sup_{\Lambda_1(0)} \psi^{n,\varepsilon} < \infty$, and thus there exists $\lambda_0^{n,\varepsilon} = (t_0^{n,\varepsilon}, \xi_0^{n,\varepsilon}, \zeta_0^{n,\varepsilon}) \in \Lambda_1(0)$ such that $\psi^{n,\varepsilon}(\Delta_0^{n,\varepsilon}) \geq \sup_{\Lambda_1(0)} \psi^{n,\varepsilon} - \frac{1}{n}$. Now apply Lemma 7.2 on $(\Lambda_1(t_0^{n,\varepsilon}), d_1)$, there exist $\hat{\lambda}^{n,\varepsilon} = (\hat{t}^{n,\varepsilon}, \hat{\xi}^{n,\varepsilon}, \hat{\zeta}^{n,\varepsilon}) \in \Lambda_1(t_0^{n,\varepsilon})$ and $\lambda_i^{n,\varepsilon} = (t_i^{n,\varepsilon}, \xi_i^{n,\varepsilon}, \zeta_i^{n,\varepsilon}) \in \Lambda_1(t_0^{n,\varepsilon})$, $i \geq 1$, such that

- $t_i^{n,\varepsilon} \uparrow \hat{t}^{n,\varepsilon}$ as $i \rightarrow \infty$, where the monotonicity of $(t_i^{n,\varepsilon})_i$ is due to [56, Lemma 2.14];
- $\bar{\Upsilon}_1(\hat{\lambda}^{n,\varepsilon}, \lambda_i^{n,\varepsilon}) \leq \frac{1}{n2^i}$, $i \geq 0$, and $\Psi^{n,\varepsilon} := \psi^{n,\varepsilon} - \sum_{i=0}^{\infty} 2^{-i} \bar{\Upsilon}_1(\hat{\lambda}, \lambda_i^{n,\varepsilon}) \leq \psi^{n,\varepsilon}$ satisfies (7.9).

(ii) For notational convenience, in the rest of the proof we omit the superscript (n,ε) , e.g. $\hat{\lambda} = \hat{\lambda}^{n,\varepsilon}$, $\lambda_i = \lambda_i^{n,\varepsilon}$, $\psi = \psi^{n,\varepsilon}$, $\Psi = \Psi^{n,\varepsilon}$. By the definition of Ψ and (7.6), we have $[\psi - \Psi](\underline{\lambda}) = \pi_0(t) + \pi_1(t, \underline{\xi}) + \pi_2(t, \underline{\zeta})$ with nonnegative maps $\pi_0(t) := \sum_{i=0}^{\infty} 2^{-i} (t - t_i)^2$,

$$\pi_1(t, \underline{\xi}) := \sum_{i=0}^{\infty} 2^{-i} \Upsilon_0((t, \underline{\xi}), (t_i, \underline{\xi}_i)), \quad \text{and} \quad \pi_2(t, \underline{\zeta}) := \sum_{i=0}^{\infty} 2^{-i} \Upsilon_0((t, \underline{\zeta}), (t_i, \underline{\zeta}_i)). \quad (7.14)$$

The regularity of $\pi_j^{n,\varepsilon}$ follow from Lemma 7.3, and it only remains to verify (7.10). First,

$$\begin{aligned} \pi_0(t) + \pi_1(t, \underline{\xi}) + \pi_2(t, \underline{\zeta}) &= \sum_{i=0}^{\infty} 2^{-i} \bar{\Upsilon}_1(\hat{\lambda}, \lambda_i) \leq \sum_{i=0}^{\infty} \frac{1}{n2^{2i}} \leq \frac{2}{n}, \\ \text{and } \partial_t \pi_0(\hat{t}) &= \sum_{i=0}^{\infty} 2^{-i} (\hat{t} - t_i) \leq \sum_{i=0}^{\infty} \frac{1}{2^i} \left(\frac{1}{n2^i} \right)^{\frac{1}{2}} \leq \frac{C}{\sqrt{n}}. \end{aligned}$$

Next, by Lemmas 7.3 and the estimates (7.4) we have $\partial_t \pi_1 = 0$, and

$$\|\partial_X \pi_1(\hat{t}, \hat{\underline{\xi}})\|_{\frac{6}{5}} \leq \sum_{i=0}^{\infty} 2^{-i} \|\partial_{\mathbf{x}} \Upsilon_i(\hat{\xi}_{\cdot, \wedge \hat{t}} - (\xi_i)_{\cdot, \wedge t_i})\|_{\frac{6}{5}} \leq C \sum_{i=0}^{\infty} 2^{-i} \|\hat{\xi}_{\hat{t}} - (\xi_i)_{t_i}\|_{\frac{6}{5}}^5$$

Then, it follows from the bounds on Υ in (7.3) that

$$\begin{aligned} \|\partial_X \pi_1(\hat{t}, \hat{\underline{\xi}})\|_{\frac{6}{5}} &\leq C \sum_{i=0}^{\infty} 2^{-i} \|\hat{\xi}_{\hat{t}} - (\xi_i)_{t_i}\|_{\frac{6}{5}}^5 \leq C \sum_{i=0}^{\infty} 2^{-i} \|\Upsilon_i(\hat{\xi}_{\cdot, \wedge \hat{t}} - (\xi_i)_{\cdot, \wedge t_i})\|_1^{\frac{5}{6}} \\ &= C \sum_{i=0}^{\infty} 2^{-i} \left(\Upsilon_0((\hat{t}, \hat{\xi}), (t_i, \xi_i)) \right)^{\frac{5}{6}} \leq C \sum_{i=0}^{\infty} 2^{-i} \left(\frac{1}{n2^i} \right)^{\frac{5}{6}} \leq C n^{-\frac{5}{6}}. \end{aligned}$$

Similarly, as $|\partial_{\mathbf{xx}} \pi_1(\hat{t}, \hat{\underline{\xi}})| \leq \sum_{i=0}^{\infty} 2^{-i} |\partial_{\mathbf{xx}} \Upsilon_i(\hat{\xi}_{\cdot, \wedge \hat{t}} - (\xi_i)_{\cdot, \wedge t_i})|$, we may get $\|\partial_{\mathbf{xx}} \pi_1(\hat{t}, \hat{\underline{\xi}})\|_{\frac{3}{2}} \leq C n^{-\frac{2}{3}}$. The estimates for π_2 can be proved similarly.

(iii) First, by (7.9) we have

$$\psi(\hat{\lambda}) \geq \Psi(\hat{\lambda}) \geq \frac{m}{2} - \frac{1}{n} \geq -1. \quad (7.15)$$

Then, noting that $\|\cdot\|_2 \leq \|\cdot\|_6$, it follows from (7.13) that

$$-1 \leq C [1 + \|\hat{\xi}_{\cdot, \wedge \hat{t}}\|_6 + \|\hat{\zeta}_{\cdot, \wedge \hat{t}}\|_6] - \frac{\varepsilon}{66} [\|\hat{\xi}_{\cdot, \wedge \hat{t}}\|_6^6 + \|\hat{\zeta}_{\cdot, \wedge \hat{t}}\|_6^6].$$

This implies (7.11) with $C_\varepsilon = C\varepsilon^{-\frac{6}{5}}$.

Next, as $\Psi(\hat{\lambda}) \geq \Psi(\hat{t}, \hat{\zeta}, \hat{\xi})$ by the optimality of $\hat{\lambda}$ in (7.9), then by (7.8) and (ii) we have

$$\begin{aligned} 0 &\leq U_1(\hat{t}, \hat{\xi}) - U_1(\hat{t}, \hat{\zeta}) - n\mathbb{E}[\Upsilon_{\hat{t}}(\hat{\xi} - \hat{\zeta})] - n^{\frac{4}{3}}\mathbb{E}[|\hat{\xi}_{\hat{t}} - \hat{\zeta}_{\hat{t}}|^6] \\ &\quad + \sum_{i \geq 0} 2^{-i}\mathbb{E}\left[\Upsilon_{\hat{t}}(\hat{\zeta} - (\xi_i)_{\cdot \wedge \hat{t}_i}) - \Upsilon_{\hat{t}}(\hat{\xi} - (\xi_i)_{\cdot \wedge \hat{t}_i})\right]. \end{aligned} \quad (7.16)$$

Since $\Upsilon \geq 0$, we have

$$\begin{aligned} n\mathbb{E}[\Upsilon_{\hat{t}}(\hat{\xi} - \hat{\zeta})] + n^{\frac{4}{3}}\mathbb{E}[|\hat{\xi}_{\hat{t}} - \hat{\zeta}_{\hat{t}}|^6] &\leq K_1 + K_2, \quad \text{where} \\ K_1 &:= |U_1(\hat{t}, \hat{\xi}) - U_1(\hat{t}, \hat{\zeta})|, \quad K_2 := \sum_{i \geq 0} 2^{-i}\mathbb{E}[\Upsilon_{\hat{t}}(\hat{\zeta}_{\cdot \wedge \hat{t}_i} - (\xi_i)_{\cdot \wedge \hat{t}_i})]. \end{aligned}$$

Since U_1 satisfies (2.6), by (7.11) we have

$$K_1 \leq C\|\hat{\xi}_{\cdot \wedge \hat{t}} - \hat{\zeta}_{\cdot \wedge \hat{t}}\|_6.$$

Moreover, recall from (i) that $\bar{\Upsilon}_1(\hat{\lambda}, \lambda_i) \leq \frac{1}{n2^i}$. Then, by (7.3) we have

$$K_2 \leq C \sum_{i=0}^{\infty} 2^{-i}\mathbb{E}\left[\Upsilon_{\hat{t}}(\hat{\zeta} - \hat{\xi}) + \Upsilon_{\hat{t}}(\hat{\xi} - (\xi_i)_{\cdot \wedge \hat{t}_i})\right] \leq C\|\hat{\xi}_{\cdot \wedge \hat{t}} - \hat{\zeta}_{\cdot \wedge \hat{t}}\|_6^6 + \frac{C}{n}.$$

Recall from (7.3) that $\Upsilon_t(\mathbf{x}) \geq |\mathbf{x}_{\cdot \wedge t}|_{\infty}^6$. Then

$$n\|\hat{\xi}_{\cdot \wedge \hat{t}} - \hat{\zeta}_{\cdot \wedge \hat{t}}\|_6^6 + n^{\frac{4}{3}}\|\hat{\xi}_{\hat{t}} - \hat{\zeta}_{\hat{t}}\|_6^6 \leq C\|\hat{\xi}_{\cdot \wedge \hat{t}} - \hat{\zeta}_{\cdot \wedge \hat{t}}\|_6 + C\|\hat{\xi}_{\cdot \wedge \hat{t}} - \hat{\zeta}_{\cdot \wedge \hat{t}}\|_6^6 + \frac{C}{n}.$$

For $n \geq 3C$ and noting that $Cx \leq \frac{1}{3}nx^6 + Cn^{-\frac{1}{5}}$, this provides

$$\frac{n}{3}\|\hat{\xi}_{\cdot \wedge \hat{t}} - \hat{\zeta}_{\cdot \wedge \hat{t}}\|_6^6 + n^{\frac{4}{3}}\|\hat{\xi}_{\hat{t}} - \hat{\zeta}_{\hat{t}}\|_6^6 \leq Cn^{-\frac{1}{5}}.$$

Note further that $\frac{3}{2} - \frac{4}{3} - \frac{1}{5} = -\frac{1}{30}$. This implies (7.12) immediately.

(iv) First, for $n \geq \frac{6}{m}$, by (7.15) we have $\psi(\hat{\lambda}) \geq \frac{m}{2} - \frac{1}{n} \geq \frac{m}{3}$. Then, by the second inequality in (5.6) and since $U_1(T, \cdot) \leq U_2(T, \cdot)$, we derive from (7.8), (7.11), (7.12) that

$$\begin{aligned} \frac{m}{3} &\leq \psi(\hat{\lambda}) \leq U_1(\hat{t}, \hat{\xi}) - U_2(\hat{t}, \hat{\zeta}) \leq [U_1(\hat{t}, \hat{\xi}_{\cdot \wedge \hat{t}}) - U_2(\hat{t}, \hat{\zeta}_{\cdot \wedge \hat{t}})] - [U_1(T, \hat{\xi}_{\cdot \wedge \hat{t}}) - U_2(T, \hat{\zeta}_{\cdot \wedge \hat{t}})] \\ &\leq 2\rho_{C_\varepsilon}(T - \hat{t}) + C\|\hat{\xi}_{\cdot \wedge \hat{t}} - \hat{\zeta}_{\cdot \wedge \hat{t}}\|_2 \leq 2\rho_{C_\varepsilon}(T - \hat{t}) + C_\varepsilon n^{-\frac{31}{180}}. \end{aligned}$$

The required result follows from taking \liminf in the last inequality. \blacksquare

8 The comparison result: a crucial estimate

To prove the comparison principle, we need another proposition. Recall the notations ε_m , C_ε and $n_{\varepsilon,m}$ defined in Proposition 7.5.

Proposition 8.1. *Let all the conditions in Proposition 7.5 hold true. Assume further that*

$$U_2 \text{ is a viscosity supersolution of } \mathcal{L}U_2(t, \underline{\xi}) \leq -c_1 \text{ for some } c_1 > 0. \quad (8.1)$$

Then there exists a constant $C > 0$, independent of n, ε, m, c_1 , such that

$$c_1 \leq \varepsilon \left(C - \frac{1}{2T} \right) \|\hat{\zeta}_{\cdot, \wedge \hat{t}^{n,\varepsilon}}^{n,\varepsilon}\|_6^6 + C\varepsilon + C_\varepsilon n^{-\frac{1}{45}}, \text{ for all } \varepsilon \leq \varepsilon_m, n \geq n_{\varepsilon,m}. \quad (8.2)$$

We note that $\psi^{n,\varepsilon}$ in (7.8) doubles the spatial variable only. The rather lengthy proof of this proposition requires additional doubling variables in time and space.

8.1 Doubling variable in time

Let $\varepsilon_m, n_{\varepsilon,m}$, $\hat{\lambda} = \hat{\lambda}^{n,\varepsilon} = (\hat{t}^{n,\varepsilon}, \hat{\xi}^{n,\varepsilon}, \hat{\zeta}^{n,\varepsilon})$ be as in Proposition 7.5, and set for $0 < \varepsilon \leq \varepsilon_m$ and $n \geq n_{\varepsilon,m}$:

$$\hat{\mathbf{a}} = \hat{\mathbf{a}}^{n,\varepsilon} := \frac{1}{2} [\hat{\xi}^{n,\varepsilon} + \hat{\zeta}^{n,\varepsilon}], \quad \hat{\boldsymbol{\theta}} = \hat{\boldsymbol{\theta}}^{n,\varepsilon} := (\hat{t}^{n,\varepsilon}, \hat{\xi}^{n,\varepsilon}), \quad \hat{\boldsymbol{\eta}} = \hat{\boldsymbol{\eta}}^{n,\varepsilon} := (\hat{t}^{n,\varepsilon}, \hat{\zeta}^{n,\varepsilon}),$$

where we omit the superscripts n,ε when the contexts are clear. For each $N \geq 1$ and $\iota = (\boldsymbol{\theta}, \boldsymbol{\eta}) \in \Lambda_2(\hat{t})$, with $\boldsymbol{\theta} = (t, \xi)$ and $\boldsymbol{\eta} = (s, \zeta)$, define

$$\begin{aligned} \psi_{n,\varepsilon}^N(\underline{\iota}) &:= U_1(\underline{\boldsymbol{\theta}}) - U_2(\underline{\boldsymbol{\eta}}) - [\varphi_1^{\hat{\boldsymbol{\theta}}}(\underline{\boldsymbol{\theta}}, \hat{\mathbf{a}}) + \varphi_2^{\hat{\boldsymbol{\eta}}}(\underline{\boldsymbol{\eta}}, \hat{\mathbf{a}})] - \pi_0^{n,\varepsilon}(t) - N|s - t|^2 \\ &\quad - \sup_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_-^{\hat{\boldsymbol{\theta}}, \hat{\mathbf{a}}}(\boldsymbol{\theta}, \alpha) - \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_+^{\hat{\boldsymbol{\theta}}, \hat{\mathbf{a}}}(\boldsymbol{\eta}, \alpha), \end{aligned} \quad (8.3)$$

where, recalling the notations in (4.10) and (5.3), the penalties $\varphi_j^{\hat{\boldsymbol{\theta}}}$ and \mathcal{J}_\pm are given by

$$\begin{aligned} \varphi_1^{\hat{\boldsymbol{\theta}}}(\underline{\boldsymbol{\theta}}, \hat{\mathbf{a}}) &:= \bar{\Upsilon}_0(\underline{\boldsymbol{\theta}}, \hat{\boldsymbol{\theta}}) + 2^5 n \Upsilon_0(\underline{\boldsymbol{\theta}}, (\hat{t}, \hat{\mathbf{a}})) + \pi_1^{n,\varepsilon}(\underline{\boldsymbol{\theta}}); \\ \varphi_2^{\hat{\boldsymbol{\eta}}}(\underline{\boldsymbol{\eta}}, \hat{\mathbf{a}}) &:= \bar{\Upsilon}_0(\underline{\boldsymbol{\eta}}, \hat{\boldsymbol{\eta}}) + 2^5 n \Upsilon_0(\underline{\boldsymbol{\eta}}, (\hat{t}, \hat{\mathbf{a}})) + \varepsilon \left(1 - \frac{s}{2T} \right) \mathbb{E}[\Upsilon_s(\zeta)] + \pi_2^{n,\varepsilon}(\underline{\boldsymbol{\eta}}); \\ \mathcal{J}_\pm^{\hat{\boldsymbol{\theta}}, \hat{\mathbf{a}}}(\boldsymbol{\theta}, \alpha) &:= 2^5 n^{\frac{4}{3}} \mathbb{E} [|\hat{\mathcal{I}}_t^\alpha(\xi)|^6] \pm \hat{F}_t^\alpha, \quad \hat{\mathcal{I}}^\alpha := \mathcal{I}^{b^{\hat{\boldsymbol{\theta}}}, \sigma^{\hat{\boldsymbol{\theta}}}, \hat{t}, \hat{\mathbf{a}}, \alpha}, \quad \hat{F}^\alpha := F^{f^{\hat{\boldsymbol{\theta}}}, \hat{t}, \alpha}. \end{aligned} \quad (8.4)$$

We emphasize that the term $\mathcal{J}_+^{\hat{\boldsymbol{\theta}}, \hat{\mathbf{a}}}(\boldsymbol{\eta}, \alpha)$ in (8.3) also relies on $\hat{\boldsymbol{\theta}}$, and not on $\hat{\boldsymbol{\eta}}$.

Proposition 8.2. *Let all the conditions in Proposition 7.5 hold true. Then for any $0 < \varepsilon \leq \varepsilon_m$, $n \geq n_{\varepsilon,m}$, the following hold.*

(i) There exist $\check{t}^N = (\check{\theta}^N, \check{\eta}^N) = ((\check{t}^N, \check{\xi}^N), (\check{s}^N, \check{\zeta}^N)) \in \Lambda_2(\hat{t}, \hat{t})$ (omitting n, ε) and $\Psi_{n, \varepsilon}^N \in C^0(\Lambda_2(\hat{t}, \hat{t}))$ such that $\Psi_{n, \varepsilon}^N \leq \psi_{n, \varepsilon}^N$ and

$$\Psi_{n, \varepsilon}^N(\check{\underline{L}}^N) = \max_{\iota \in \Lambda_2(\hat{t}, \hat{t})} \Psi_{n, \varepsilon}^N(\underline{L}) \geq \left(\sup_{\iota \in \Lambda_2(\hat{t}, \hat{t})} \psi_{n, \varepsilon}^N(\underline{L}) - \frac{1}{N} \right) \vee \Psi^{n, \varepsilon}(\hat{\lambda}). \quad (8.5)$$

(ii) Moreover, $[\psi_{n, \varepsilon}^N - \Psi_{n, \varepsilon}^N](\underline{L}) = \pi_1^{n, \varepsilon, N}(\underline{\theta}) + \pi_2^{n, \varepsilon, N}(\underline{\eta})$, for some nonnegative functions $\pi_1^{n, \varepsilon, N} \in C^{1,2}(\overline{\mathcal{X}}_{[\check{t}^N, T]}^6)$, $\pi_2^{n, \varepsilon, N} \in C^{1,2}(\overline{\mathcal{X}}_{[\check{s}^N, T]}^6)$, such that: for $j = 1, 2$ and at \check{t}^N ,

$$0 \leq \pi_j^{n, \varepsilon, N} \leq \frac{C}{N}; \quad |\partial_t \pi_j^{n, \varepsilon, N}| \leq \frac{C}{\sqrt{N}}; \quad \|\partial_X \pi_j^{n, \varepsilon, N}\|_{\frac{6}{5}} \leq \frac{C}{N^{\frac{5}{6}}}, \quad \|\partial_{\mathbf{x}X} \pi_j^{n, \varepsilon, N}\|_{\frac{3}{2}} \leq \frac{C}{N^{\frac{2}{3}}}. \quad (8.6)$$

(iii) $\sup_{n, N} \mathbb{E} \left[\|\check{\xi}_{\cdot, \wedge \check{t}^N}^N\|^6 + \|\check{\zeta}_{\cdot, \wedge \check{s}^N}^N\|^6 \right] \leq C_\varepsilon$, for some C_ε , which may depend on ε .

Proof. Recall the connection between $\Psi^{n, \varepsilon}$ and $\pi_j^{n, \varepsilon}$ in Proposition 7.5 (ii) and that $\hat{\xi} - \hat{\mathbf{a}} = \hat{\mathbf{a}} - \hat{\zeta} = \frac{1}{2}[\hat{\xi} - \hat{\zeta}]$. Then, by (8.3), (7.8), (4.10), and (7.2) we have

$$\begin{aligned} \psi_{n, \varepsilon}^N(\underline{\hat{\theta}}, \underline{\hat{\eta}}) - \Psi_{n, \varepsilon}^{n, \varepsilon}(\hat{\lambda}) &= \psi_{n, \varepsilon}^N(\underline{\hat{\theta}}, \underline{\hat{\eta}}) - \psi^{n, \varepsilon}(\hat{\lambda}) + \pi_0^{n, \varepsilon}(\hat{t}) + \pi_1^{n, \varepsilon}(\underline{\hat{\theta}}) + \pi_2^{n, \varepsilon}(\underline{\hat{\eta}}) \\ &= n\Upsilon_0(\underline{\hat{\theta}}, \underline{\hat{\eta}}) + n^{\frac{4}{3}} \mathbb{E} \left[|\hat{\xi}_{\hat{t}} - \hat{\zeta}_{\hat{t}}|^6 \right] \\ &\quad - 2^5 n \left[\Upsilon_0(\underline{\hat{\theta}}, (\hat{t}, \hat{\mathbf{a}})) + \Upsilon_0(\underline{\hat{\eta}}, (\hat{t}, \hat{\mathbf{a}})) \right] - 2^5 n^{\frac{4}{3}} \mathbb{E} \left[|\hat{\xi}_{\hat{t}} - \hat{\mathbf{a}}_{\hat{t}}|^6 + |\hat{\zeta}_{\hat{t}} - \hat{\mathbf{a}}_{\hat{t}}|^6 \right] \\ &= n \mathbb{E} \left[\Upsilon_{\hat{t}}(\hat{\xi} - \hat{\eta}) \right] - 2^5 n \mathbb{E} \left[\Upsilon_{\hat{t}} \left(\frac{\hat{\xi} - \hat{\zeta}}{2} \right) + \Upsilon_{\hat{t}} \left(\frac{\hat{\zeta} - \hat{\xi}}{2} \right) \right] = 0, \end{aligned} \quad (8.7)$$

where the last equality is due to the fact that $\Upsilon_t(\frac{\mathbf{x}}{2}) = \Upsilon_t(-\frac{\mathbf{x}}{2}) = \frac{1}{2^6} \Upsilon_t(\mathbf{x})$, which can be verified straightforwardly from (7.2). Then one can prove all the statements following the same arguments as in Proposition 7.5. In particular, we have $\pi_1^{n, \varepsilon, N}(\underline{\theta}) := \sum_{i=0}^{\infty} 2^{-i} \overline{\Upsilon}_0(\underline{\theta}, \check{\theta}_i^N)$ and $\pi_2^{n, \varepsilon, N}(\underline{\eta}) := \sum_{i=0}^{\infty} 2^{-i} \overline{\Upsilon}_0(\underline{\eta}, \check{\eta}_i^N)$, for some appropriate $\check{t}_i^N = (\check{\theta}_i^N, \check{\eta}_i^N)$. Note that the $\pi_j^{n, \varepsilon, N}$ here include the time difference, while the $\pi_j^{n, \varepsilon}$ in (7.14) does not. \blacksquare

The convergence of \check{t}^N is more involved and will be conducted in two alternative cases.

8.2 Proof of Proposition 8.1: Case 1.

In this subsection we prove the proposition in the case that:

$$\text{there is a subsequence, still denoted } (\check{\theta}^N, \check{\eta}^N)_N, \text{ satisfying } \check{s}^N \leq \check{t}^N. \quad (8.8)$$

Surprisingly in this case we do not need to invoke the viscosity subsolution property of U_1 . The viscosity supersolution property of U_2 alone induces the desired estimate.

Lemma 8.3. *In the setting of Proposition 8.2, if $\check{s}^N \leq \check{t}^N$, then*

$$\|\check{\xi}_{\cdot, \wedge \check{t}^N}^N - \hat{\xi}_{\cdot, \wedge \hat{t}}\|_6^6 + \|\check{\zeta}_{\cdot, \wedge \check{s}^N}^N - \hat{\zeta}_{\cdot, \wedge \hat{t}}\|_6^6 + |\check{t}^N - \hat{t}|^2 + |\check{s}^N - \hat{t}|^2 + N |\check{t}^N - \check{s}^N|^2 \leq \rho_{C_\varepsilon} (C_{n, \varepsilon} N^{-\frac{1}{2}}) + C_{n, \varepsilon} N^{-\frac{1}{4}}.$$

where C_ε is defined in Proposition 7.5, and $C_{\varepsilon,n}$ is a constant depending on n, ε only. Consequently, under (8.8), there exists $N_{n,\varepsilon}$, such that

$$\check{s}^N \leq \check{t}^N \leq T'_{\varepsilon,m} := \frac{T_{\varepsilon,m} + T}{2} < T, \quad \text{for all } n \geq n_{\varepsilon,m}, N \geq N_{n,\varepsilon}. \quad (8.9)$$

Proof. Clearly, (8.9) is a direct consequence of the claimed estimate in this lemma, which we now focus on. Fix (n, ε, N) such that $\check{s}^N \leq \check{t}^N$. We omit further the subscripts/superscripts (n, ε) in $\psi^{n,\varepsilon}, \Psi^{n,\varepsilon}, \psi_{n,\varepsilon}^N, \Psi_{n,\varepsilon}^N, \pi_j^{n,\varepsilon}, \pi_j^{n,\varepsilon,N}$. Note that $\check{s}^N \leq \check{t}^N$ implies

$$\Upsilon_{\check{t}^N}(\check{\zeta}_{\wedge \check{s}^N}^N) = \Upsilon_{\check{s}^N}(\check{\zeta}^N), \quad \Upsilon_0(\check{\theta}^N, (\check{t}^N, \check{\zeta}_{\wedge \check{s}^N}^N)) = \Upsilon_0(\check{\theta}^N, \check{\eta}^N), \quad \pi_2(\check{t}^N, \check{\zeta}_{\wedge \check{s}^N}^N) = \pi_2(\check{\eta}^N), \quad (8.10)$$

where we used (7.2) and (7.14). Since $\Psi^N \leq \psi^N$, by (8.5) and (7.9) we have

$$0 \geq \Psi^N(\check{\zeta}^N) - \psi^N(\check{\zeta}^N) \geq \Psi(\hat{\lambda}) - \psi^N(\check{\zeta}^N) \geq \Psi(\check{t}^N, \check{\xi}^N, \check{\zeta}_{\wedge \check{s}^N}^N) - \psi^N(\check{\zeta}^N).$$

Thus, it follows from the definitions of ψ, ψ^N in (7.8) and (8.3), as well as Proposition 7.5 (ii) and (8.10) that

$$\begin{aligned} 0 &\geq \psi(\check{t}^N, \check{\xi}^N, \check{\zeta}_{\wedge \check{s}^N}^N) - [\pi_0(\check{t}^N) + \pi_1(\check{\theta}^N) + \pi_2(\check{\eta}^N)] - \psi^N(\check{\zeta}^N) \\ &= \left[U_2(\check{s}^N, \check{\zeta}^N) - U_2(\check{t}^N, \check{\zeta}_{\wedge \check{s}^N}^N) \right] + \left[\bar{\Upsilon}_0(\check{\theta}^N, \hat{\theta}) + \bar{\Upsilon}_0(\check{\eta}^N, \hat{\eta}) \right] \\ &\quad + \frac{\varepsilon}{2T}(\check{t}^N - \check{s}^N) \mathbb{E}[\Upsilon_{\check{s}^N}(\check{\zeta}^N)] + 2^5 n \left[\Upsilon_0(\check{\theta}^N, (\hat{t}, \hat{\alpha})) + \Upsilon_0(\check{\eta}^N, (\hat{t}, \hat{\alpha})) \right] \\ &\quad - n \Upsilon_0(\check{\theta}^N, \check{\eta}^N) - n^{\frac{4}{3}} \mathbb{E}[|\check{\xi}_{\check{t}^N}^N - \check{\zeta}_{\check{s}^N}^N|^6] + N|\check{s}^N - \check{t}^N|^2 \\ &\quad + \sup_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\check{\theta}^N, \alpha) + \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\check{\eta}^N, \alpha). \end{aligned} \quad (8.11)$$

Recall (4.10). Note that, by (7.2) and (7.3), as well as the fact $32[|x|^6 + |y|^6] \geq |x - y|^6$,

$$\begin{aligned} 2^5 \left[\Upsilon_0(\check{\theta}^N, (\hat{t}, \hat{\alpha})) + \Upsilon_0(\check{\eta}^N, (\hat{t}, \hat{\alpha})) \right] &\geq \Upsilon_0(\check{\theta}^N, \check{\eta}^N), \\ \sup_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\check{\theta}^N, \alpha) + \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\check{\eta}^N, \alpha) &\geq \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left[\mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\check{\theta}^N, \alpha) + \mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\check{\eta}^N, \alpha) \right] \\ &\geq \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left[n^{\frac{4}{3}} \mathbb{E}[|\hat{\mathcal{I}}_{\check{t}^N}^\alpha(\check{\xi}^N) - \hat{\mathcal{I}}_{\check{s}^N}^\alpha(\check{\zeta}^N)|^6] + \hat{F}_{\check{s}^N}^\alpha - \hat{F}_{\check{t}^N}^\alpha \right]. \end{aligned} \quad (8.12)$$

Plug these into (8.11) and note that $\check{t}^N - \check{s}^N \geq 0$, we have

$$\begin{aligned} N|\check{s}^N - \check{t}^N|^2 + \left[\bar{\Upsilon}_0(\check{\theta}^N, \hat{\theta}) + \bar{\Upsilon}_0(\check{\eta}^N, \hat{\eta}) \right] &\leq U_2(\check{t}^N, \check{\zeta}_{\wedge \check{s}^N}^N) - U_2(\check{s}^N, \check{\zeta}^N) \\ &\quad + n^{\frac{4}{3}} \mathbb{E}[|\check{\xi}_{\check{t}^N}^N - \check{\zeta}_{\check{s}^N}^N|^6] - \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left[n^{\frac{4}{3}} \mathbb{E}[|\hat{\mathcal{I}}_{\check{t}^N}^\alpha(\check{\xi}^N) - \hat{\mathcal{I}}_{\check{s}^N}^\alpha(\check{\zeta}^N)|^6] + \hat{F}_{\check{s}^N}^\alpha - \hat{F}_{\check{t}^N}^\alpha \right]. \end{aligned} \quad (8.13)$$

Recall (5.3), (8.4), and denote, for any α ,

$$\Delta_N^\alpha := [\check{\xi}_{\check{t}^N}^N - \check{\zeta}_{\check{s}^N}^N] - [\hat{\mathcal{I}}_{\check{t}^N}^\alpha(\check{\xi}^N) - \hat{\mathcal{I}}_{\check{s}^N}^\alpha(\check{\zeta}^N)] = \int_{\check{s}^N}^{\check{t}^N} b_r^{\hat{\theta}, \alpha} dr + \int_{\check{s}^N}^{\check{t}^N} \sigma_r^{\hat{\theta}, \alpha} dB_r.$$

By (7.11) we have $\mathbb{E}[|\Delta_N^\alpha|^6] \leq C_\varepsilon(\check{t}^N - \check{s}^N)^3$, and then, by Proposition 8.2 (iii),

$$\begin{aligned} & \mathbb{E}\left[|\check{\xi}_{\check{t}^N}^N - \check{\zeta}_{\check{s}^N}^N|^6\right] - \mathbb{E}\left[|\hat{\mathcal{I}}_{\check{t}^N}^\alpha(\check{\xi}^N) - \hat{\mathcal{I}}_{\check{s}^N}^\alpha(\check{\zeta}^N)|^6\right] \\ &= \mathbb{E}\left[|\check{\xi}_{\check{t}^N}^N - \check{\zeta}_{\check{s}^N}^N|^6 - |\check{\xi}_{\check{t}^N}^N - \check{\zeta}_{\check{s}^N}^N - \Delta_N^\alpha|^6\right] \leq C_\varepsilon \|\Delta_N^\alpha\|_6 \leq C_\varepsilon(\check{t}^N - \check{s}^N)^{\frac{1}{2}}, \end{aligned}$$

Moreover, since $-U_2$ satisfies (5.6), it is clear that

$$U_2(\check{t}^N, \check{\underline{\zeta}}_{\check{s}^N}^N) - U_2(\check{s}^N, \check{\underline{\zeta}}^N) \leq \rho_{C_\varepsilon}(\check{t}^N - \check{s}^N); \quad \left|F_{\check{s}^N}^{\hat{\theta}, \alpha} - F_{\check{t}^N}^{\hat{\theta}, \alpha}\right| \leq C_\varepsilon(\check{t}^N - \check{s}^N).$$

Then by (8.13) we have

$$N|\check{s}^N - \check{t}^N|^2 + \left[\bar{\Upsilon}_0(\check{\underline{\theta}}^N, \hat{\underline{\theta}}) + \bar{\Upsilon}_0(\check{\underline{\eta}}^N, \hat{\underline{\eta}})\right] \leq \rho_{C_\varepsilon}(\check{t}^N - \check{s}^N) + C_{n, \varepsilon}(\check{t}^N - \check{s}^N)^{\frac{1}{2}}. \quad (8.14)$$

As $\rho_R(\delta) \leq 1 + C_R\delta$, without loss of generality, the above implies that $0 \leq \check{t}^N - \check{s}^N \leq C_{n, \varepsilon}N^{-\frac{1}{2}}$. The required estimate follows by plugging this into the right side of (8.14). \blacksquare

Proof of Proposition 8.1 under (8.8). Fix $\varepsilon \leq \varepsilon_m$, $n \geq n_{\varepsilon, m}$, $N \geq N_{n, \varepsilon}$ such that $\check{s}^N \leq \check{t}^N$. As before we omit the subscripts/supscripts (n, ε) . Introduce

$$\varphi_2^N(\underline{\eta}) := \varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\underline{\alpha}}) + N|s - \check{t}^N|^2 + \pi_2^N(\underline{\eta}); \quad \phi_2^N(\underline{\eta}) := \inf_{\alpha \in \mathcal{A}_{[\check{t}, T]}} \left[2^5 n^{\frac{4}{3}} \mathbb{E}[|\hat{\mathcal{I}}_s^\alpha(\zeta)|^6] + \hat{F}_t^\alpha\right].$$

Here, in terms of the notation in (4.10) and (4.11), we are setting $(\tilde{t}, \tilde{\xi}) = (t', \xi') = (\hat{t}, \hat{\xi})$, $(\tilde{b}^\alpha, \tilde{\sigma}^\alpha, \tilde{f}^\alpha) = (b^{\hat{\theta}, \alpha}, \sigma^{\hat{\theta}, \alpha}, f^{\hat{\theta}, \alpha})$, and $k = 2^5 n$. Then one can easily check that $\phi_2^N \in C^+(\bar{\mathcal{X}}_{[\check{s}^N, T]}^6)$. By (8.9) we have $\check{s}^N < T$, and by Proposition 7.5 (ii) and Proposition 8.2 (ii) we see that $\varphi_2^N \in C^{1,2}(\bar{\mathcal{X}}_{[\check{t}, T]}^6)$. Then it follows from Proposition 8.2 (i) that $(-\varphi_2^N, -\phi_2^N) \in \mathfrak{F}_6^- U_2(\check{\underline{\eta}}^N)$. Thus, by the viscosity supersolution property of U_2 , and recalling (5.3), we have

$$\begin{aligned} c_1 \leq & \partial_t \varphi_2^N(\check{\underline{\eta}}^N) + \varliminf_{\delta \rightarrow 0} \sup_{\alpha \in \mathcal{A}_{[\check{s}^N, T]}} \frac{1}{\delta} \int_{\check{s}^N}^{\check{s}^N + \delta} \left[\phi_2^N(s, \bar{\mathcal{X}}^{\check{\eta}^N, \alpha}) \right. \\ & \left. - H_s(\check{\underline{\zeta}}_{\check{s}^N}^N, -\partial_X \varphi_2^N(\check{\underline{\eta}}^N), -\partial_{\mathbf{x}X} \varphi_2^N(\check{\underline{\eta}}^N), \underline{\alpha}_s) \right] ds. \end{aligned}$$

Note that $t' = \hat{t} \leq \check{s}^N$. This allows us to apply Proposition 4.3 (i) and obtain

$$\begin{aligned} \int_{\check{s}^N}^{\check{s}^N + \delta} \phi_2^N(s, \bar{\mathcal{X}}^{\check{\eta}^N, \alpha}) ds & \leq \inf_{\tilde{\alpha} \in \mathcal{A}_{[\check{s}^N, T]}} \int_{\check{s}^N}^{\check{s}^N + \delta} \left[Cn^{\frac{4}{3}} I_s^{\check{\eta}^N}(\alpha, \tilde{\alpha}) + f_s^{\hat{\theta}, \tilde{\alpha}} \right] ds \\ & \leq \int_{\check{s}^N}^{\check{s}^N + \delta} \left[Cn^{\frac{4}{3}} I_s^{\check{\eta}^N}(\alpha, \alpha) + f_s^{\hat{\theta}, \alpha} \right] ds, \end{aligned} \quad (8.15)$$

where we denoted

$$I_s^\eta(\alpha, \tilde{\alpha}) := \|b_s^{\eta, \alpha} - b_s^{\hat{\theta}, \tilde{\alpha}}\|_6 \sup_{\alpha' \in \mathcal{A}_{[\hat{t}, T]}} \|\mathcal{I}_s^{\hat{\theta}, \alpha'}(\bar{X}^\eta, \alpha)\|_6^5 + \|\sigma_s^{\eta, \alpha} - \sigma_s^{\hat{\theta}, \tilde{\alpha}}\|_6^2 \sup_{\alpha' \in \mathcal{A}_{[\hat{t}, T]}} \|\mathcal{I}_s^{\hat{\theta}, \alpha'}(\bar{X}^\eta, \alpha)\|_6^4,$$

$$\mathcal{I}_s^{\hat{\theta}, \alpha'}(\bar{X}^{\eta, \alpha}) := \mathcal{I}_s^{b^{\hat{\theta}}, \sigma^{\hat{\theta}}, \hat{t}, \hat{\alpha}, \alpha'}(\bar{X}^{\eta, \alpha}). \quad (8.16)$$

Observe that it is crucial to have $\inf_{\tilde{\alpha}}$ in the right side of the first line of (8.15), which helps to replace $\tilde{\alpha}$ with α in the second line. Then, by (3.5) we have

$$c_1 \leq K_0^N + \liminf_{\delta \rightarrow 0} \sup_{\alpha \in \mathcal{A}_{[\tilde{s}^N, T]}} \frac{1}{\delta} \int_{\tilde{s}^N}^{\tilde{s}^N + \delta} [K_1^N(s) + K_2^N(s) + K_3^N(s) + K_4^N(s)] ds, \quad (8.17)$$

where

$$\begin{aligned} K_0^N &:= \partial_t \varphi_2^N(\underline{\eta}^N), \quad K_1^N(s) := \mathbb{E} \left[b_s^{\tilde{\eta}^N, \alpha} \cdot \partial_X \varphi_2^N(\underline{\eta}^N) \right], \\ K_2^N(s) &:= \frac{1}{2} \mathbb{E} \left[(\sigma \sigma^\top)_s^{\tilde{\eta}^N, \alpha} : \partial_{\mathbf{x}X} \varphi_2^N(\underline{\eta}^N) \right], \\ K_3^N(s) &:= C n^{\frac{4}{3}} I_s^{\tilde{\eta}^N}(\alpha, \alpha), \quad K_4^N(s) := \mathbb{E} \left[f_s^{\hat{\theta}, \alpha} - f_s^{\tilde{\eta}^N, \alpha} \right]. \end{aligned} \quad (8.18)$$

We now estimates the K_i^N separately. We shall send $N \rightarrow \infty$, along the subsequence such that $\tilde{s}^N \leq \tilde{t}^N$, by applying the convergence in Proposition 8.2 (i) and Lemma 8.3 repeatedly. We shall also apply the estimates (7.4), Proposition 7.5 (ii), (iii), Proposition 8.2 (ii), (iii), as well as Assumption 2.1 repeatedly. At below we let $o(1)$ denote generic terms which will vanish when $N \rightarrow \infty$. The convergence rate may depend on n, ε , but is uniform in α, δ, s . Note that, by (8.4), (7.2), and (7.5),

$$\begin{aligned} \partial_t \varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\mathbf{a}}) &= -\frac{\varepsilon}{2T} \mathbb{E}[\Upsilon(\eta)] + 2(s - \hat{t}), \\ \partial_X \varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\mathbf{a}}) &= \partial_{\mathbf{x}} \Upsilon_s(\zeta - \hat{\zeta}_{\cdot, \wedge \hat{t}}) + 2^5 n \partial_{\mathbf{x}} \Upsilon_s(\zeta - \hat{\mathbf{a}}_{\cdot, \wedge \hat{t}}) + \varepsilon \left[1 - \frac{s}{2T} \right] \partial_{\mathbf{x}} \Upsilon_s(\zeta) + \partial_X \pi_2(\underline{\eta}), \\ \partial_{\mathbf{x}X} \varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\mathbf{a}}) &= \partial_{\mathbf{x}X} \Upsilon_s(\zeta - \hat{\zeta}_{\cdot, \wedge \hat{t}}) + 2^5 n \partial_{\mathbf{x}X} \Upsilon_s(\zeta - \hat{\mathbf{a}}_{\cdot, \wedge \hat{t}}) + \varepsilon \left[1 - \frac{s}{2T} \right] \partial_{\mathbf{x}X} \Upsilon_s(\zeta) + \partial_{\mathbf{x}X} \pi_2(\underline{\eta}). \end{aligned} \quad (8.19)$$

First, thanks to the crucial assumption $\tilde{s}^N \leq \tilde{t}^N$,

$$\begin{aligned} K_0^N &= \partial_t \varphi_2^{\hat{\eta}}(\underline{\eta}^N, \hat{\mathbf{a}}) + 2N(\tilde{s}^N - \tilde{t}^N) + \partial_t \pi_2^N(\underline{\eta}^N) \\ &\leq \partial_t \varphi_2^{\hat{\eta}}(\underline{\eta}^N, \hat{\mathbf{a}}) + o(1) = -\frac{\varepsilon}{2T} \mathbb{E}[\Upsilon(\hat{\eta}^N)] + 2(\tilde{s}^N - \hat{t}) + o(1) \\ &= -\frac{\varepsilon}{2T} \mathbb{E}[\Upsilon(\hat{\eta})] + o(1) \leq -\frac{\varepsilon}{2T} \mathbb{E}[|\hat{\zeta}_{\cdot, \wedge \hat{t}}|_\infty^6] + o(1). \end{aligned} \quad (8.20)$$

Next, denote $\|\zeta\|_{2,t} := (\mathbb{E}_t^0[|\zeta_{\cdot, \wedge t}|_\infty^2])^{\frac{1}{2}}$,

$$\begin{aligned} K_1^N(s) &= \mathbb{E} \left[b_s^{\tilde{\eta}^N, \alpha} \cdot [\partial_X \varphi_2^{\hat{\eta}}(\underline{\eta}^N) + \partial_X \pi_2^N(\underline{\eta}^N)] \right] = \mathbb{E} \left[b_s^{\tilde{\eta}^N, \alpha} \cdot \partial_X \varphi_2^{\hat{\eta}}(\underline{\eta}^N) \right] + o(1) \\ &\leq C \mathbb{E} \left[\left(1 + |\check{\zeta}_{\cdot, \wedge \tilde{s}^N}^N|_\infty + \|\check{\zeta}^N\|_{2, \tilde{s}^N} \right) \times \right. \\ &\quad \left. \left(|\partial_{\mathbf{x}} \Upsilon_{\tilde{s}^N}(\check{\zeta}^N - \hat{\zeta}_{\cdot, \wedge \hat{t}})| + n |\partial_{\mathbf{x}} \Upsilon_{\tilde{s}^N}(\check{\zeta}^N - \hat{\mathbf{a}}_{\cdot, \wedge \hat{t}})| + \varepsilon |\partial_{\mathbf{x}} \Upsilon_{\tilde{s}^N}(\check{\zeta}^N)| + |\partial_X \pi_2(\underline{\eta}^N)| \right) \right] + o(1) \end{aligned}$$

$$\begin{aligned}
&= C\mathbb{E}\left[\left(1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty + \|\hat{\zeta}\|_{2,\hat{t}}\right)\left(n|\partial_{\mathbf{x}}\Upsilon_{\hat{t}}(\hat{\zeta} - \hat{\mathbf{a}})| + \varepsilon|\partial_{\mathbf{x}}\Upsilon_{\hat{t}}(\hat{\zeta})| + |\partial_X\pi_2(\hat{\eta})|\right)\right] + o(1) \\
&\leq C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n(\mathbb{E}[|\hat{\zeta}_{\hat{t}} - \hat{\xi}_{\hat{t}}|^6])^{\frac{5}{6}} + C_\varepsilon(\mathbb{E}[|\partial_X\pi_2(\hat{\eta})|^{\frac{6}{5}}])^{\frac{5}{6}} + o(1) \\
&\leq C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n(n^{-\frac{3}{2}-\frac{1}{30}})^{\frac{5}{6}} + C_\varepsilon n^{-\frac{5}{6}} + o(1) \\
&\leq C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n^{-\frac{5}{18}} + o(1). \tag{8.21}
\end{aligned}$$

Similarly,

$$\begin{aligned}
K_2^N(s) &\leq C\mathbb{E}\left[\left(1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^2 + \|\hat{\zeta}\|_{2,\hat{t}}^2\right) \times \right. \\
&\quad \left. \left(n|\partial_{\mathbf{xx}}\Upsilon_{\hat{t}}(\hat{\zeta} - \hat{\mathbf{a}})| + \varepsilon|\partial_{\mathbf{xx}}\Upsilon_{\hat{t}}(\hat{\zeta})| + |\partial_{\mathbf{x}X}\pi_2(\hat{\eta})|\right)\right] + o(1) \\
&\leq C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n(\mathbb{E}[|\hat{\zeta}_{\hat{t}} - \hat{\xi}_{\hat{t}}|^6])^{\frac{2}{3}} + C_\varepsilon(\mathbb{E}[|\partial_X\pi_2(\hat{\eta})|^{\frac{3}{2}}])^{\frac{2}{3}} + o(1) \\
&\leq C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n(n^{-\frac{3}{2}-\frac{1}{30}})^{\frac{2}{3}} + C_\varepsilon n^{-\frac{2}{3}} + o(1) \\
&\leq C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n^{-\frac{1}{45}} + o(1). \tag{8.22}
\end{aligned}$$

To estimate K_3^N , we first recall (4.10) and note that

$$\mathcal{I}_s^{\hat{\theta},\alpha'}(\bar{X}\bar{\eta}^{N,\alpha}) = \check{\zeta}_{\check{s}^N}^N - \hat{\mathbf{a}}_{\hat{t}} + \int_{\check{s}^N}^s b_t^{\bar{\eta}^{N,\alpha}} dt + \int_{\check{s}^N}^s \sigma_t^{\bar{\eta}^{N,\alpha}} dB_t - \int_{\hat{t}}^s b_t^{\hat{\theta},\alpha'} dt - \int_{\hat{t}}^s \sigma_t^{\hat{\theta},\alpha'} dB_t.$$

Then, for $s \in [\check{s}^N, \check{s}^N + \delta]$, we have

$$\begin{aligned}
\mathbb{E}[|\mathcal{I}_s^{\hat{\theta},\alpha'}(\bar{X}\bar{\eta}^{N,\alpha})|^6] &\leq C\mathbb{E}[|\check{\zeta}_{\check{s}^N}^N - \hat{\mathbf{a}}_{\hat{t}}|^6] + C_\varepsilon(s - \hat{t})^3 \\
&= C\mathbb{E}[|\hat{\zeta}_{\hat{t}} - \hat{\xi}_{\hat{t}}|^6] + C_\varepsilon\delta^3 + o(1) \leq C_\varepsilon n^{-\frac{23}{15}} + C_\varepsilon\delta^3 + o(1).
\end{aligned}$$

Moreover, note that

$$\begin{aligned}
&\mathbb{E}\left[|b_s^{\bar{\eta}^{N,\alpha}} - b_s^{\hat{\theta},\alpha'}|^6 + |\sigma_s^{\bar{\eta}^{N,\alpha}} - \sigma_s^{\hat{\theta},\alpha'}|^6 + |f_s^{\bar{\eta}^{N,\alpha}} - f_s^{\hat{\theta},\alpha'}|^6\right] \\
&\leq C\mathbb{E}[|\check{\zeta}_{\check{s}^N}^N - \hat{\xi}_{\cdot,\Lambda\hat{t}}|_\infty^6] = C\mathbb{E}[|\hat{\zeta}_{\cdot,\Lambda\hat{t}} - \hat{\xi}_{\cdot,\Lambda\hat{t}}|_\infty^6] + o(1) \leq C_\varepsilon n^{-\frac{31}{30}} + o(1).
\end{aligned}$$

Plug these into (8.16), we have

$$\begin{aligned}
K_3^N(s) &\leq C_\varepsilon n^{\frac{4}{3}} \left[(n^{-\frac{23}{15}} + \delta^3)^{\frac{5}{6}} (n^{-\frac{31}{30}})^{\frac{1}{6}} + (n^{-\frac{23}{15}} + \delta^3)^{\frac{2}{3}} (n^{-\frac{31}{30}})^{\frac{1}{3}} \right] + o(1) \\
&\leq C_\varepsilon n^{-\frac{1}{30}} + C_{n,\varepsilon}\delta^2 + o(1); \tag{8.23}
\end{aligned}$$

$$K_4^N(s) \leq C_\varepsilon (n^{-\frac{31}{30}})^{\frac{1}{6}} + o(1) = C_\varepsilon n^{-\frac{31}{180}} + o(1).$$

Plug all these estimates (8.20)-(8.23) into (8.17), we obtain

$$\begin{aligned}
c_1 &\leq -\frac{\varepsilon}{2T}\mathbb{E}[|\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + \left[C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n^{-\frac{5}{18}} \right] \\
&\quad + \left[C\varepsilon\mathbb{E}[1 + |\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C_\varepsilon n^{-\frac{1}{45}} \right] + C_\varepsilon n^{-\frac{1}{30}} + C_\varepsilon n^{-\frac{31}{180}} + o(1) \\
&\leq \varepsilon \left[C - \frac{1}{2T} \right] \mathbb{E}[|\hat{\zeta}_{\cdot,\Lambda\hat{t}}|_\infty^6] + C\varepsilon + C_\varepsilon n^{-\frac{1}{45}} + o(1).
\end{aligned}$$

By sending $N \rightarrow \infty$, this implies (8.2) immediately, in the case (8.8). ■

8.3 Proof of Proposition 8.1: Case 2

We now turn to the case that

$$\check{t}^N < \check{s}^N \quad \text{for all large } N. \quad (8.24)$$

In this case, we need to modify (8.3) by adding another penalization. Fix $0 < \varepsilon \leq \varepsilon_m$, $n \geq n_{\varepsilon,m}$, and N large enough such that (8.24) holds. Consider the settings of Propositions 7.5 and 8.2. Again we omit the subscripts/superscripts (n, ε) when the contexts are clear. For each M , recall (8.4) and define:

$$\begin{aligned} \psi^{N,M}(\underline{\mathcal{L}}) &:= U_1(\underline{\theta}) - U_2(\underline{\eta}) - [\varphi_1^{\hat{\theta}}(\underline{\theta}, \hat{\mathbf{a}}) + \varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\mathbf{a}})] - \pi_0^{n,\varepsilon}(t) - N|s - t|^2 \\ &\quad - M|(s - t) - (\check{s}^N - \check{t}^N)|^2 - \inf_{\alpha \in \mathcal{A}_{[\check{t}, T]}} \left(\mathcal{J}_-^{\hat{\theta}, \hat{\mathbf{a}}}(\theta, \alpha) + \mathcal{J}_+^{\hat{\theta}, \hat{\mathbf{a}}}(\eta, \alpha) \right). \end{aligned} \quad (8.25)$$

Proposition 8.4. *In the setting of Proposition 8.2, for any $0 < \varepsilon \leq \varepsilon_m$, $n \geq n_{\varepsilon,m}$, and N large enough such that $\check{t}^N < \check{s}^N$, the following hold.*

(i) *There exist $\mathring{i}^{N,M} = (\mathring{\theta}^{N,M}, \mathring{\eta}^{N,M}) = ((\mathring{t}^{N,M}, \mathring{\xi}^{N,M}), (\mathring{s}^{N,M}, \mathring{\zeta}^{N,M})) \in \Lambda_2(\check{t}^N, \check{s}^N)$ (omitting n, ε again) and $\Psi^{N,M} \in C^0(\Lambda_2(\mathring{t}^{N,M}, \mathring{s}^{N,M}))$ such that $\Psi^{N,M} \leq \psi^{N,M}$ and*

$$\Psi^{N,M}(\underline{\mathcal{L}}^{N,M}) = \max_{\Lambda_2(\mathring{t}^{N,M}, \mathring{s}^{N,M})} \Psi^{N,M} \geq \left(\sup_{\Lambda_2(\mathring{t}^{N,M}, \mathring{s}^{N,M})} \psi^{N,M} - \frac{1}{N} \right) \vee \psi^{N,M}(\underline{\mathcal{L}}^N). \quad (8.26)$$

(ii) *Moreover, $[\psi^{N,M} - \Psi^{N,M}](\underline{\mathcal{L}}) = \pi_1^{N,M}(\underline{\theta}) + \pi_2^{N,M}(\underline{\eta})$, for some nonnegative functions $\pi_1^{N,M} \in C^{1,2}(\overline{\mathcal{X}}_{[\mathring{t}^{N,M}, T]}^6)$, $\pi_2^{N,M} \in C^{1,2}(\overline{\mathcal{X}}_{[\mathring{s}^{N,M}, T]}^6)$, such that: for $j = 1, 2$ and at $\mathring{i}^{N,M}$,*

$$0 \leq \pi_j^{N,M} \leq \frac{C}{N}; \quad |\partial_t \pi_j^{N,M}| \leq \frac{C}{\sqrt{N}}; \quad \|\partial_X \pi_j^{N,M}\|_{\frac{6}{5}} \leq \frac{C}{N^{\frac{5}{6}}}, \quad \|\partial_{\mathbf{x}X} \pi_j^{N,M}\|_{\frac{3}{2}} \leq \frac{C}{N^{\frac{2}{3}}}. \quad (8.27)$$

(iii) $\sup_{n,N,M} \mathbb{E} \left[\|\dot{\xi}_{\cdot, \wedge \mathring{t}^{N,M}}^{N,M}\|^6 + \|\dot{\zeta}_{\cdot, \wedge \mathring{s}^{N,M}}^{N,M}\|^6 \right] \leq C_\varepsilon$, for some C_ε , which may depend on ε .

(iv) $M|(\mathring{s}^{N,M} - \mathring{t}^{N,M}) - (\check{s}^N - \check{t}^N)|^2 \leq C_{n,\varepsilon}$, for some $C_{n,\varepsilon}$ which may depend on n, ε , but not on N, M ; Consequently, there exists $M_N = M_{n,\varepsilon,N}$ large enough such that $\mathring{t}^{N,M_N} < \mathring{s}^{N,M_N}$.

(v) For the M_N in (iv), and abbreviating the notations further that $\mathring{i}^N = \mathring{i}^{N,M_N}$, we have

$$\begin{aligned} \mathbb{E} \left[|\dot{\xi}_{\cdot, \wedge \mathring{t}^N}^N - \hat{\xi}_{\cdot, \wedge \hat{t}}|_\infty^6 + |\dot{\zeta}_{\cdot, \wedge \mathring{s}^N}^N - \hat{\zeta}_{\cdot, \wedge \hat{t}}|_\infty^6 \right] + |\mathring{t}^N - \hat{t}|^2 + |\mathring{s}^N - \hat{t}|^2 \\ + N|\mathring{t}^N - \check{s}^N|^2 + M_N \left| (\mathring{s}^N - \mathring{t}^N) - (\check{s}^N - \check{t}^N) \right|^2 \leq C_{n,\varepsilon} N^{-\frac{1}{3}}, \end{aligned}$$

for some $C_{n,\varepsilon}$, which may depend on n, ε , but not on N . Consequently, there exists $N_{n,\varepsilon}$ such that $\mathring{t}^N < \check{s}^N < T$ for all $N \geq N_{n,\varepsilon}$.

Proof. (i)-(iii) follow similar arguments as in Proposition 7.5. We now fix (n, ε, N, M) and as in the previous proofs, we omit the subscripts/superscripts (n, ε) , but keep N, M .

(iv) First, since $\Psi^{N,M} \leq \psi^{N,M}$ and $\Psi^N \leq \psi^N$, we have

$$\psi^{N,M}(\underline{\dot{\iota}}^{N,M}) \geq \Psi^{N,M}(\underline{\dot{\iota}}^{N,M}) \geq \psi^{N,M}(\underline{\dot{\iota}}^N) \geq \psi^N(\underline{\dot{\iota}}^N) \geq \Psi^N(\underline{\dot{\iota}}^N) \geq \Psi^N(\underline{\dot{\iota}}^{N,M}), \quad (8.28)$$

where the second inequality is due to (8.26), the third one is by comparing (8.3) and (8.25) directly, and the last one is due to (8.5). Then, by (8.3), (8.25), and Proposition 8.2 (ii),

$$\begin{aligned} 0 &\geq \Psi^N(\underline{\dot{\iota}}^{N,M}) - \psi^{N,M}(\underline{\dot{\iota}}^{N,M}) \\ &= \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left(\mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\hat{\theta}^{N,M}, \alpha) + \mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\hat{\eta}^{N,M}, \alpha) \right) + M |(\dot{s}^{N,M} - \dot{t}^{N,M}) - (\dot{s}^N - \dot{t}^N)|^2 \\ &\quad - \left[\sup_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\hat{\theta}^{N,M}, \alpha) + \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\hat{\eta}^{N,M}, \alpha) + \pi_1^N(\hat{\theta}^{N,M}) + \pi_2^N(\hat{\eta}^{N,M}) \right]. \end{aligned}$$

This, together with (iii) and (8.6), implies the estimate in Item (iv) immediately.

(v) Recall again that $\dot{\iota}^N = \dot{\iota}^{N, M_N}$. By (8.28), (8.5), and then (7.9), we have

$$\psi^{N, M_N}(\underline{\dot{\iota}}^N) \geq \psi^N(\underline{\dot{\iota}}^N) \geq \Psi(\hat{\underline{\dot{\iota}}}) \geq \Psi(\dot{s}^N, \underline{\dot{\xi}}_{\wedge \dot{t}^N}^N, \underline{\dot{\zeta}}^N),$$

where $\Psi = \Psi^{n, \varepsilon}$. Since $\dot{t}^N < \dot{s}^N$, similarly to (8.10) we have

$$\Upsilon_{\dot{s}^N}(\underline{\dot{\xi}}_{\wedge \dot{t}^N}^N) = \Upsilon_{\dot{t}^N}(\underline{\dot{\xi}}^N), \quad \Upsilon_0(\underline{\dot{\eta}}^N, (\dot{s}^N, \underline{\dot{\xi}}_{\wedge \dot{t}^N}^N)) = \Upsilon_0(\underline{\dot{\eta}}^N, \underline{\dot{\theta}}^N), \quad \pi_1(\dot{s}^N, \underline{\dot{\xi}}_{\wedge \dot{t}^N}^N) = \pi_1(\underline{\dot{\theta}}^N).$$

Then, by comparing (7.8) and (8.25), and recalling Proposition 7.5 (ii) and (8.4), we have

$$\begin{aligned} 0 &\geq \psi(\dot{s}^N, \underline{\dot{\xi}}_{\wedge \dot{t}^N}^N, \underline{\dot{\zeta}}^N) - \pi_0(\dot{s}^N) - \pi_1(\dot{s}^N, \underline{\dot{\xi}}_{\wedge \dot{t}^N}^N) - \pi_2(\underline{\dot{\eta}}^N) - \psi^{N, M_N}(\underline{\dot{\iota}}^N) \\ &= [U_1(\dot{s}^N, \underline{\dot{\xi}}_{\wedge \dot{t}^N}^N) - U_1(\underline{\dot{\theta}}^N)] + [\bar{\Upsilon}_0(\underline{\dot{\theta}}^N, \underline{\dot{\theta}}) + \bar{\Upsilon}_0(\underline{\dot{\eta}}^N, \underline{\dot{\eta}})] \\ &\quad + 2^5 n [\Upsilon_0(\underline{\dot{\theta}}^N, (\hat{t}, \hat{\alpha})) + \Upsilon_0(\underline{\dot{\eta}}^N, (\hat{t}, \hat{\alpha}))] - n \Upsilon_0(\underline{\dot{\theta}}^N, \underline{\dot{\eta}}^N) - n^{\frac{4}{3}} \mathbb{E}[|\underline{\dot{\xi}}_{\dot{t}^N}^N - \underline{\dot{\zeta}}_{\dot{s}^N}^N|^6] \\ &\quad + \pi_0(\dot{t}^N) - \pi_0(\dot{s}^N) + N |\dot{s}^N - \dot{t}^N|^2 + M_N |(\dot{s}^N - \dot{t}^N) - (\dot{s}^N - \dot{t}^N)|^2 \\ &\quad + \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left(\mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\hat{\theta}^N, \alpha) + \mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\hat{\eta}^N, \alpha) \right) \\ &\geq [U_1(\dot{s}^N, \underline{\dot{\xi}}_{\wedge \dot{t}^N}^N) - U_1(\underline{\dot{\theta}}^N)] + [\bar{\Upsilon}_0(\underline{\dot{\theta}}^N, \underline{\dot{\theta}}) + \bar{\Upsilon}_0(\underline{\dot{\eta}}^N, \underline{\dot{\eta}})] - n^{\frac{4}{3}} \mathbb{E}[|\underline{\dot{\xi}}_{\dot{t}^N}^N - \underline{\dot{\zeta}}_{\dot{s}^N}^N|^6] \\ &\quad + \pi_0(\dot{t}^N) - \pi_0(\dot{s}^N) + N |\dot{s}^N - \dot{t}^N|^2 + M_N |(\dot{s}^N - \dot{t}^N) - (\dot{s}^N - \dot{t}^N)|^2 \\ &\quad + \inf_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left[n^{\frac{4}{3}} \mathbb{E}[|\hat{\mathcal{I}}_{\dot{t}^N}^\alpha(\dot{\xi}^N) - \hat{\mathcal{I}}_{\dot{s}^N}^\alpha(\dot{\zeta}^N)|^6] + [\hat{F}_{\dot{s}^N}^\alpha - \hat{F}_{\dot{t}^N}^\alpha] \right], \end{aligned}$$

where the last inequality is due to estimates similar to (8.12). Then,

$$N |\dot{s}^N - \dot{t}^N|^2 + M_N |(\dot{s}^N - \dot{t}^N) - (\dot{s}^N - \dot{t}^N)|^2 + \bar{\Upsilon}_0(\underline{\dot{\theta}}^N, \underline{\dot{\theta}}) + \bar{\Upsilon}_0(\underline{\dot{\eta}}^N, \underline{\dot{\eta}})$$

$$\begin{aligned}
&\leq |U_1(\dot{s}^N, \underline{\xi}^N) - U_1(\dot{t}^N, \underline{\xi}^N)| + |\pi_0(\dot{t}^N) - \pi_0(\dot{s}^N)| \\
&\quad - \inf_{\alpha \in \mathcal{A}_{[\dot{t}, T]}} \left[n^{\frac{4}{3}} \mathbb{E} [|\hat{\mathcal{I}}_{\dot{t}^N}^\alpha(\dot{\xi}^N) - \hat{\mathcal{I}}_{\dot{s}^N}^\alpha(\dot{\zeta}^N)|^6 - |\dot{\xi}_{\dot{t}^N}^N - \dot{\zeta}_{\dot{s}^N}^N|^6] + [\hat{F}_{\dot{s}^N}^\alpha - \hat{F}_{\dot{t}^N}^\alpha] \right] \\
&\leq C_{n,\varepsilon} |\dot{s}^N - \dot{t}^N|^{\frac{1}{2}},
\end{aligned}$$

where the last inequality is due to (7.10) and similar arguments as for (8.14). In particular, $N|\dot{s}^N - \dot{t}^N|^2 \leq C_{n,\varepsilon} |\dot{s}^N - \dot{t}^N|^{\frac{1}{2}}$, and thus $|\dot{s}^N - \dot{t}^N| \leq C_{n,\varepsilon} N^{-\frac{2}{3}}$. Plugging this into the right side of the last inequality provides the required estimate in Item (v). \blacksquare

Proof of Proposition 8.1 in Case (8.24). Fix $0 < \varepsilon \leq \varepsilon_m$, $n \geq n_{\varepsilon,m}$, $N \geq N_{n,\varepsilon}$, and recall the $M_N = M_{n,\varepsilon,N}$ in Proposition 8.4 (v) and (vi). As before we omit the subscripts/supscripts (n, ε) , and abbreviate $\dot{t}^N := \dot{t}^{N, M_N}$. In particular, by otherwise considering a larger $N_{n,\varepsilon}$, we have $\dot{t}^N < \dot{s}^N < T$. Introduce:

$$\begin{aligned}
\varphi_1^N(\underline{\theta}) &:= \varphi_1^{\hat{\theta}}(\underline{\theta}, \hat{\alpha}) + \pi_0(t) + N|t - \dot{s}^N|^2 + M_N|t - \dot{t}^N - \dot{s}^N + \dot{s}^N|^2 + \pi_1^{N, M_N}(\underline{\theta}); \\
\phi_1^N(\underline{\theta}) &:= \inf_{\alpha \in \mathcal{A}_{[\dot{t}, T]}} \left(\mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\theta, \alpha) + \kappa_1(\alpha) \right), \quad \kappa_1(\alpha) := \mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\hat{\eta}^N, \alpha); \\
\varphi_2^N(\underline{\eta}) &:= \varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\alpha}) + N|s - \dot{t}^N|^2 + M_N|s - \dot{s}^N - \dot{t}^N + \dot{t}^N|^2 + \pi_2^{N, M_N}(\underline{\eta}); \\
\phi_2^N(\underline{\eta}) &:= \inf_{\alpha \in \mathcal{A}_{[\dot{s}, T]}} \left(\mathcal{J}_+^{\hat{\theta}, \hat{\alpha}}(\eta, \alpha) + \kappa_2(\alpha) \right), \quad \kappa_2(\alpha) := \mathcal{J}_-^{\hat{\theta}, \hat{\alpha}}(\hat{\theta}^N, \alpha).
\end{aligned}$$

One can easily see that $(\varphi_1^N, \phi_1^N) \in \mathfrak{F}^+ U_1(\underline{\theta}^N)$ and $(-\varphi_2^N, -\phi_2^N) \in \mathfrak{F}^- U_2(\underline{\eta}^N)$. Then, by the viscosity properties of U_1 and U_2 , we have

$$\begin{aligned}
c_1 &\leq K_0^N + \lim_{\delta \rightarrow 0} \inf_{\alpha \in \mathcal{A}_{[\dot{t}^N, T]}} \frac{1}{\delta} \int_{\dot{t}^N}^{\dot{t}^N + \delta} [K_1^N(\delta, \alpha, t) + K_2^N(\delta, \alpha, t)] dt \\
&\quad + \lim_{\delta \rightarrow 0} \sup_{\alpha \in \mathcal{A}_{[\dot{s}^N, T]}} \frac{1}{\delta} \int_{\dot{s}^N}^{\dot{s}^N + \delta} [K_3^N(\delta, \alpha, s) + K_4^N(\delta, \alpha, s)] ds,
\end{aligned} \tag{8.29}$$

where

$$\begin{aligned}
K_0^N &:= \partial_t \varphi_1^N(\underline{\theta}^N) + \partial_t \varphi_2^N(\underline{\eta}^N); \\
K_1^N(\delta, \alpha, t) &:= \mathbb{E} \left[b_t^{\hat{\theta}^N, \alpha} \cdot \partial_X \varphi_1^N(\underline{\theta}^N) + \frac{1}{2} (\sigma \sigma^\top)_t^{\hat{\theta}^N, \alpha} : \partial_{\mathbf{x}\mathbf{x}} \varphi_1^N(\underline{\theta}^N) \right]; \\
K_2^N(\delta, \alpha, t) &:= \dot{\phi}_1^N(t, \bar{X}^{\hat{\theta}^N, \alpha}) + f_t^{\hat{\theta}^N, \alpha}; \\
K_3^N(\delta, \alpha, s) &:= \mathbb{E} \left[b_s^{\hat{\eta}^N, \alpha} \cdot \partial_X \varphi_2^N(\underline{\eta}^N) + \frac{1}{2} (\sigma \sigma^\top)_s^{\hat{\eta}^N, \alpha} : \partial_{\mathbf{x}\mathbf{x}} \varphi_2^N(\underline{\eta}^N) \right]; \\
K_4^N(\delta, \alpha, s) &:= \dot{\phi}_2^N(s, \bar{X}^{\hat{\eta}^N, \alpha}) - f_s^{\hat{\eta}^N, \alpha}.
\end{aligned} \tag{8.30}$$

We shall estimate K_i^N separately. Similarly to the previous subsection, we shall send $N \rightarrow \infty$, under (8.24), by applying the convergence in Proposition 8.4 (ii), (v) repeatedly.

We shall also apply the estimates (7.4), Proposition 7.5 (ii), (iii), Proposition 8.2 (ii), (iii), Proposition 8.4 (v), as well as Assumption 2.1 repeatedly. At below again we let $o(1)$ denote generic terms which will vanish when $N \rightarrow \infty$. The convergence rate may depend on n, ε , but is uniform in α, δ, t, s . Recall (8.19), and we have similar expressions for the derivatives of $\varphi_1^{\hat{\theta}}(\underline{\theta}, \hat{\mathbf{a}})$.

First, noting the obvious but crucial cancellations, we have

$$\begin{aligned}
K_0^N &= \partial_t \varphi_1^{\hat{\theta}}(\underline{\theta}^N, \hat{\mathbf{a}}) + \partial_t \pi_0(\dot{t}^N) + \partial_t \pi_1^{N, M_N}(\underline{\theta}^N) + \partial_t \varphi_2^{\hat{\eta}}(\underline{\eta}^N, \hat{\mathbf{a}}) + \partial_t \pi_2^{N, M_N}(\underline{\eta}^N) \\
&= -\frac{\varepsilon}{2T} \mathbb{E}[\Upsilon(\dot{\eta}^N)] + 2[(\dot{t}^N - \hat{t}) + (\dot{s}^N - \hat{t})] + \partial_t \pi_0(\dot{t}^N) + o(1) \\
&= -\frac{\varepsilon}{2T} \mathbb{E}[\Upsilon(\hat{\zeta})] + \partial_t \pi_0(\hat{t}) + o(1) \\
&\leq -\frac{\varepsilon}{2T} \mathbb{E}[|\hat{\zeta}_{\cdot, \wedge \hat{t}}|_{\infty}^6] + Cn^{-\frac{1}{2}} + o(1).
\end{aligned} \tag{8.31}$$

Next, similarly to (8.21) and (8.22), we have

$$K_1^N(\delta, \alpha, t) \leq C_\varepsilon n^{-\frac{1}{45}} + o(1); \quad K_3^N(\delta, \alpha, s) \leq C_\varepsilon \mathbb{E}[1 + |\hat{\zeta}_{\cdot, \wedge \hat{t}}|_{\infty}^6] + C_\varepsilon n^{-\frac{1}{45}} + o(1). \tag{8.32}$$

Moreover, note that, in terms of the notations in Proposition 4.3, for ϕ_2^N we have $t' = \dot{t}^N$ and $t = \dot{s}^N$. Since $\dot{t}^N < \dot{s}^N$, we may apply Proposition 4.3 (i) again to estimate K_4^N : for any α and recalling (8.16),

$$\begin{aligned}
\int_{\dot{s}^N}^{\dot{s}^N + \delta} K_4^N(\delta, \alpha, s) ds &\leq \inf_{\tilde{\alpha} \in \mathcal{A}_{[t, T]}} \int_{\dot{s}^N}^{\dot{s}^N + \delta} [Cn^{\frac{4}{3}} I_s^{\dot{\eta}^N}(\alpha, \tilde{\alpha}) + f_s^{\hat{\theta}, \tilde{\alpha}}] ds - \int_{\dot{s}^N}^{\dot{s}^N + \delta} f_s^{\dot{\eta}^N, \alpha} ds \\
&\leq \int_{\dot{s}^N}^{\dot{s}^N + \delta} [Cn^{\frac{4}{3}} I_s^{\dot{\eta}^N}(\alpha, \alpha) + f_s^{\hat{\theta}, \alpha} - f_s^{\dot{\eta}^N, \alpha}] ds.
\end{aligned}$$

Then it follows from the same arguments as in (8.23) that $\frac{1}{\delta} \int_{\dot{s}^N}^{\dot{s}^N + \delta} K_4^N(\delta, \alpha, s) ds \leq C_\varepsilon n^{-\frac{1}{30}} + C_n \delta^2 + o(1)$, and thus

$$\sup_{\alpha \in \mathcal{A}_{[\dot{s}^N, T]}} \frac{1}{\delta} \int_{\dot{s}^N}^{\dot{s}^N + \delta} K_4^N(\delta, \alpha, s) ds \leq C_n n^{-\frac{1}{30}} + C_n \delta^2 + o(1). \tag{8.33}$$

Finally we estimate K_2^N . Recall the notations K_2^N in (8.30), $\bar{X}^{\dot{\theta}^N, \alpha}$ in (5.3), and the setting in (4.10)-(4.11). Note that for ϕ_1^N we have $t' = \dot{s}^N > \dot{t}^N = t$, then we can only apply Proposition 4.3 (ii) here. That is, there exists $\alpha^\delta \in \mathcal{A}_{[\dot{t}^N, T]}$, which may depend on $\underline{X}_{\cdot, \wedge \dot{t}^N}^{\dot{\theta}^N, \alpha} = \underline{\xi}_{\cdot, \wedge \dot{t}^N}^{\dot{\theta}^N}$, but not on $(b_s^{\dot{\theta}^N, \alpha}, \sigma_s^{\dot{\theta}^N, \alpha})_{s \geq \dot{t}^N}$, and thus is independent of α , such that

$$\int_{\dot{t}^N}^{\dot{t}^N + \delta} K_2^N(\delta, \alpha, t) dt \leq \int_{\dot{t}^N}^{\dot{t}^N + \delta} [Cn^{\frac{4}{3}} I_t^{\dot{\theta}^N}(\alpha, \alpha^\delta) - f_t^{\hat{\theta}, \alpha^\delta} + f_t^{\dot{\theta}^N, \alpha}] dt,$$

and thus

$$\inf_{\alpha \in \mathcal{A}_{[\check{t}^N, T]}} \int_{\check{t}^N}^{\check{t}^N + \delta} K_2^N(\delta, \alpha, t) dt \leq \int_{\check{t}^N}^{\check{t}^N + \delta} \left[Cn^{\frac{4}{3}} I_t^{\dot{\theta}^N}(\alpha^\delta, \alpha^\delta) - f_t^{\hat{\theta}, \alpha^\delta} + f_t^{\dot{\theta}^N, \alpha^\delta} \right] dt.$$

Then again by (8.23) we have

$$\inf_{\alpha \in \mathcal{A}_{[\check{t}^N, T]}} \frac{1}{\delta} \int_{\check{t}^N}^{\check{t}^N + \delta} K_2^N(\delta, \alpha, t) dt \leq Cn^{-\frac{1}{30}} + Cn\delta^2 + o(1). \quad (8.34)$$

Plug (8.31), (8.32), (8.33), and (8.34) into (8.29), we obtain

$$c_1 \leq \varepsilon \left[C - \frac{1}{2T} \right] \mathbb{E} [|\hat{\zeta}_{\cdot, \wedge \check{t}}|^6] + C\varepsilon + C_\varepsilon n^{-\frac{1}{45}} + o(1).$$

Send $N \rightarrow \infty$, we obtain (8.2) immediately, in the case (8.24). This, together with the proof in Subsection 8.2, completes the proof of Proposition 8.1. \blacksquare

Remark 8.5. *We remark that the arguments in this subsection do not work when $\check{s}^N < \check{t}^N$, and thus we cannot eliminate Subsection 8.2. Indeed, in this case we can only apply Proposition 4.3 (ii), instead of (i), to estimate $\dot{\phi}_2^N$ and K_4^N . That is, there exists α^δ , independent of α , s.t.*

$$\int_{\check{s}^N}^{\check{s}^N + \delta} K_4^N(\delta, \alpha, s) ds \leq \int_{\check{s}^N}^{\check{s}^N + \delta} \left[Cn^{\frac{4}{3}} I_s^{\dot{\eta}^N}(\alpha, \alpha^\delta) + f_s^{\hat{\theta}, \alpha^\delta} - f_s^{\dot{\eta}^N, \alpha} \right] ds + \delta^2.$$

However, for the term $\int_{\check{s}^N}^{\check{s}^N + \delta} K_4^N(\delta, \alpha, s) ds$, we need to take supremum over α :

$$\sup_{\alpha \in \mathcal{A}_{[\check{s}^N, T]}} \int_{\check{s}^N}^{\check{s}^N + \delta} K_4^N(\delta, \alpha, s) ds \leq \sup_{\alpha \in \mathcal{A}_{[\check{s}^N, T]}} \int_{\check{s}^N}^{\check{s}^N + \delta} \left[Cn^{\frac{4}{3}} I_s^{\dot{\eta}^N}(\alpha, \alpha^\delta) + f_s^{\hat{\theta}, \alpha^\delta} - f_s^{\dot{\eta}^N, \alpha} \right] ds + \delta^2.$$

Recall (8.16), the right side of the above won't be small anymore, and then we won't be able to derive the desired contradiction.

9 The comparison result: Proof of Theorem 5.6

(i) **We first prove the case that U_1 satisfies (2.6).** Fix $\delta_0 \in (0, \frac{1}{2C})$ for the C in (8.2). We proceed in three steps.

Step 1. We first assume (8.1) holds true and $T \leq \delta_0$. Assume by contradiction that $m > 0$. Then, since $C < \frac{1}{2\delta_0} \leq \frac{1}{2T}$, by (8.2) we have

$$c_1 \leq C\varepsilon + C_\varepsilon n^{-\frac{1}{45}}, \quad \text{for } n \geq n_{\varepsilon, m}.$$

By first sending $n \rightarrow \infty$ and then $\varepsilon \rightarrow 0$, we obtain the desired contradiction: $c_1 \leq 0$.

Step 2. We next assume (8.1) holds true but T can be arbitrarily large. Consider a partition $0 = T_0 < \dots < T_k = T$ such that $T_{i+1} - T_i \leq \delta_0$ for all i . By Step 1 we should have $U_1(t, \underline{\xi}) \leq U_2(t, \underline{\xi})$ for all $(t, \underline{\xi}) \in \overline{\mathcal{X}}_{[T_{k-1}, T_k]}^2$. In particular, $U_1(T_{k-1}, \cdot) \leq U_2(T_{k-1}, \cdot)$. Now consider the equation on $[T_{k-2}, T_{k-1}]$, by Step 1 again we obtain $U_1(t, \underline{\xi}) \leq U_2(t, \underline{\xi})$ for all $(t, \underline{\xi}) \in \overline{\mathcal{X}}_{[T_{k-2}, T_{k-1}]}^2$. Repeat the arguments backwardly in time, we prove the result on the whole interval $[0, T]$.

Step 3. In the general case, let $c_1 > 0$ be an arbitrary number. Set $U_2^{c_1} := U_2 + \frac{(1+T)^2}{1+t} c_1$. It is clear that $U_2^{c_1}$ also satisfies (2.6) and, providing U_2 is smooth,

$$\partial_t U_2^{c_1} = \partial_t U_2 - \frac{(1+T)^2}{(1+t)^2} c_1 \leq \partial_t U_2 - c_1, \quad \partial_X U_2^{c_1} = \partial_X U_2, \quad \partial_{\mathbf{x}X} U_2^{c_1} = \partial_{\mathbf{x}X} U_2.$$

Then one can easily see that $U_2^{c_1}$ is a viscosity supersolution of $\mathcal{L}U_2^{c_1} \leq -c_1$. Since $U_1(T, \cdot) \leq U_2(T, \cdot) < U_2^{c_1}(T, \cdot)$, then by Step 2 we have $U_1(t, \cdot) \leq U_2^{c_1}(t, \cdot)$ for all $t \in [0, T]$. Now since $c_1 > 0$ is arbitrary, we see that $U_1(t, \cdot) \leq U_2(t, \cdot)$ for all $t \in [0, T]$.

(ii) We next prove the case that U_2 satisfies (2.6). The proof is almost the same as in (i), with the following modifications. First, (7.8) is modified by changing the last term:

$$\psi^{n,\varepsilon}(\underline{\lambda}) := U_1(t, \underline{\xi}) - U_2(t, \underline{\zeta}) - n\mathbb{E}[\Upsilon_t(\xi - \zeta)] - n^{\frac{4}{3}}\mathbb{E}[|\xi_t - \zeta_t|^6] - \varepsilon\left(1 - \frac{t}{2T}\right)\mathbb{E}[\Upsilon_t(\xi)].$$

This is used to derive the counterpart of (7.16), based on $\Psi(\hat{\underline{\lambda}}) \geq \Psi(\hat{t}, \hat{\underline{\xi}}, \hat{\underline{\zeta}})$:

$$\begin{aligned} 0 \leq & U_2(\hat{t}, \hat{\underline{\xi}}) - U_2(\hat{t}, \hat{\underline{\zeta}}) - n\mathbb{E}[\Upsilon_{\hat{t}}(\hat{\underline{\xi}} - \hat{\underline{\zeta}})] - n^{\frac{4}{3}}\mathbb{E}[|\hat{\xi}_{\hat{t}} - \hat{\zeta}_{\hat{t}}|^6] \\ & + \sum_{i \geq 0} 2^{-i} \mathbb{E}\left[\Upsilon_{\hat{t}}(\hat{\xi} - (\zeta_i)_{\cdot \wedge \hat{t}_i}) - \Upsilon_{\hat{t}}(\hat{\zeta} - (\zeta_i)_{\cdot \wedge \hat{t}_i})\right]. \end{aligned}$$

Then the estimates (2.6) for U_2 will lead to (7.12).

Next, we modify (8.4) by moving the ε -term from $\varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\underline{\mathbf{a}}})$ to $\varphi_1^{\hat{\theta}}(\underline{\theta}, \hat{\underline{\mathbf{a}}})$:

$$\begin{aligned} \varphi_1^{\hat{\theta}}(\underline{\theta}, \hat{\underline{\mathbf{a}}}) &:= \bar{\Upsilon}_0(\underline{\theta}, \hat{\theta}) + 2^5 n \Upsilon_0(\underline{\theta}, (\hat{t}, \hat{\underline{\mathbf{a}}})) + \varepsilon\left(1 - \frac{t}{2T}\right)\mathbb{E}[\Upsilon_t(\xi)] + \pi_1^{n,\varepsilon}(\underline{\theta}); \\ \varphi_2^{\hat{\eta}}(\underline{\eta}, \hat{\underline{\mathbf{a}}}) &:= \bar{\Upsilon}_0(\underline{\eta}, \hat{\eta}) + 2^5 n \Upsilon_0(\underline{\eta}, (\hat{t}, \hat{\underline{\mathbf{a}}})) + \pi_2^{n,\varepsilon}(\underline{\eta}). \end{aligned}$$

This ensures that the calculation in (8.7) remains true. Moreover, in this case the derivatives of $\varphi_2^{\hat{\eta}}$ is simplified slightly. Then (8.2) will become: for all $\varepsilon \leq \varepsilon_m$, and $n \geq n_{\varepsilon,m}$,

$$c_1 \leq C\varepsilon + C_\varepsilon n^{-\frac{1}{45}} + \begin{cases} 0, & \text{under (8.10);} \\ \varepsilon\left(C - \frac{1}{2T}\right)\|\hat{\xi}_{\cdot, \wedge \hat{t}^{n,\varepsilon}}\|_6^6, & \text{under (8.24).} \end{cases}$$

This leads to the desired contradiction as in Case 1. ■

10 Appendix

Proof of Lemma 3.2. We first prove the Lemma in the case that $t \in [\hat{t}, T]$, $\xi \in \mathcal{M}_{[\hat{t}, T]}^p$.

First, let $X \in \mathcal{M}_{[\hat{t}, T]}^p$ be such that $X_{\cdot \wedge t} = \xi_{\cdot \wedge t}$ and $\beta_s = \mathbf{0}, \gamma_s = \mathbf{0}, s \geq t$, in (3.1). Then $X_{\cdot \wedge s} = X_{\cdot \wedge t} = \xi_{\cdot \wedge t}$ for all $s \geq t$, and thus by (3.2) we obtain,

$$\varphi_{t+\delta}(\xi_{\cdot \wedge t}) = \varphi_t(\xi_{\cdot \wedge t}) + \int_t^{t+\delta} \partial_t \varphi_s(\xi_{\cdot \wedge t}) ds, \quad 0 < \delta \leq T - t.$$

Since $\partial_t \varphi$ is continuous, we obtain a representation of $\partial_t \varphi$ which implies its uniqueness:

$$\partial_t \varphi_t(\xi_{\cdot \wedge t}) = \lim_{\delta \rightarrow 0} \frac{\varphi_{t+\delta}(\xi_{\cdot \wedge t}) - \varphi_t(\xi_{\cdot \wedge t})}{\delta}.$$

Next, replace the above X with $\beta_s \equiv \beta_t, s \in [t, T]$ for some arbitrary $\beta_t \in \mathbb{L}^p(\mathcal{F}_t; \mathbb{R}^d)$. By (3.2) we have, for any $0 < \delta \leq T - t$,

$$\varphi_{t+\delta}(\underline{X}) = \varphi_t(\underline{X}) + \int_t^{t+\delta} \partial_t \varphi_s(\underline{X}) ds + \mathbb{E} \left[\int_t^{t+\delta} \partial_X \varphi_s(\underline{X}) ds \cdot \beta_t \right].$$

By the uniqueness of $\partial_t \varphi$ and the continuity of $\partial_t \varphi, \partial_X \varphi$, we see that

$$\mathbb{E}[\partial_X \varphi_t(\xi) \cdot \beta_t] = \lim_{\delta \rightarrow 0} \left[\frac{\varphi_{t+\delta}(\overline{X}) - \varphi_t(\overline{X})}{\delta} - \partial_t \varphi_t(\xi) \right]$$

is unique. Since β_t is arbitrary, we see that $\partial_X \varphi_t(\xi)$ is unique, \mathbb{P} -a.s. Similarly, by considering X with $\beta_s \equiv 0$ but $\gamma_s \equiv \gamma_t, s \in [t, T]$, we obtain the uniqueness of $\mathbb{E}[\partial_{\mathbf{x}X} \varphi_t(\xi) : \gamma_t \gamma_t^\top]$. Since $\partial_{\mathbf{x}X} \varphi$ is symmetric and γ_t is arbitrary, we see that $\partial_{\mathbf{x}X} \varphi_t(\xi)$ is unique, a.s.

Finally, for arbitrary $(t, \xi) \in \overline{\mathcal{X}}_{[\hat{t}, T]}^p$, clearly there exist a sequence $\xi^n \in \mathcal{M}_{[\hat{t}, T]}^p$ such that $\lim_{n \rightarrow \infty} \|\xi^n_{\cdot \wedge t} - \xi_{\cdot \wedge t}\|_p = 0$. By the above arguments $\partial_t \varphi, \partial_X \varphi, \partial_{\mathbf{x}X} \varphi$ are unique at (t, ξ^n) . Then it follows from their continuity that they are unique at (t, ξ) as well. \blacksquare

Proof of Proposition 4.3. For any $\alpha \in \mathcal{A}_{[\hat{t}, T]}$, by the standard Itô formula we have:

$$\begin{aligned} & \left| \mathbb{E} [|\mathcal{I}_{t+\delta}^\alpha(X)|^p - |\mathcal{I}_t^\alpha(X)|^p] \right| \\ &= \left| \mathbb{E} \left[\int_t^{t+\delta} [p |\mathcal{I}_s^\alpha(X)|^{p-2} (\mathcal{I}_s^\alpha(X)) \cdot (\beta_s - \tilde{b}_s^\alpha) + \frac{p}{2} |\mathcal{I}_s^\alpha(X)|^{p-2} |\gamma_s - \tilde{\sigma}_s^\alpha|^2 \right. \right. \\ & \quad \left. \left. + \frac{p(p-2)}{2} |\mathcal{I}_s^\alpha(X)|^{p-4} |\mathcal{I}_s^\alpha(X)| (\gamma_s - \tilde{\sigma}_s^\alpha)^2] ds \right] \right| \\ &\leq C_p \mathbb{E} \left[\int_t^{t+\delta} [|\mathcal{I}_s^\alpha(X)|^{p-1} |\beta_s - \tilde{b}_s^\alpha| + |\mathcal{I}_s^\alpha(X)|^{p-2} |\gamma_s - \tilde{\sigma}_s^\alpha|^2] ds \right]. \end{aligned} \quad (10.1)$$

Then, recalling (4.12) and noting that $\kappa(\alpha)$ does not depend on t ,

$$\left| \phi_{t+\delta}(\underline{X}) - \phi_t(\underline{X}) \right| \leq k \sup_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left| \mathbb{E} [|\mathcal{I}_{t+\delta}^\alpha(X)|^p - |\mathcal{I}_t^\alpha(X)|^p] \right| + \sup_{\alpha \in \mathcal{A}_{[\hat{t}, T]}} \left[\int_t^{t+\delta} |\tilde{f}_s^\alpha| ds \right]$$

$$\begin{aligned}
&\leq C_{p,k} \sup_{\alpha \in \mathcal{A}_{[\tilde{t}, T]}} \mathbb{E} \left[\int_t^{t+\delta} [|\mathcal{I}_s^\alpha(X)|^p + |\beta_s|^p + |\gamma_s|^p + |\tilde{b}_s^\alpha|^p + |\tilde{\sigma}_s^\alpha|^p + 1] ds \right] \\
&\leq C_{p,k} \int_t^{t+\delta} \mathbb{E} \left[|X_s|^p + |\tilde{\xi}_s|^p + |\beta_s|^p + |\gamma_s|^p + \sup_{[\tilde{t}, T] \times A \times \mathcal{A}_T} [|\tilde{b}|^p + |\tilde{\sigma}|^p + 1] \right] ds.
\end{aligned}$$

This implies that $\phi_t(\underline{X})$ is absolutely continuous in t .

We next prove (i). Note that $\mathcal{A}_{[\tilde{t}, T]} = \{\alpha \oplus_t \tilde{\alpha} : \alpha \in \mathcal{A}_{[\tilde{t}, T]}, \tilde{\alpha} \in \mathcal{A}_{[t, T]}\}$, where the concatenation is defined as: $\alpha \oplus_t \tilde{\alpha} := \alpha \mathbf{1}_{[\tilde{t}, t)} + \tilde{\alpha} \mathbf{1}_{[t, T]}$. Recall the $\kappa(\alpha)$ in (4.12) and note that, since $t \geq t'$, we have $\kappa(\alpha \oplus_t \tilde{\alpha}) = \kappa(\alpha)$. Thus

$$\begin{aligned}
\phi_{t+\delta}(\underline{X}) - \phi_t(\underline{X}) &= \inf_{\alpha \in \mathcal{A}_{[\tilde{t}, T]}} \inf_{\tilde{\alpha} \in \mathcal{A}_{[t, T]}} \left[k\mathbb{E}[|\mathcal{I}_{t+\delta}^{\alpha \oplus_t \tilde{\alpha}}(X)|^p] + F_{t+\delta}^{\alpha \oplus_t \tilde{\alpha}} + \kappa(\alpha) \right] \\
&\quad - \inf_{\alpha \in \mathcal{A}_{[\tilde{t}, T]}} \left[k\mathbb{E}[|\mathcal{I}_t^\alpha(X)|^p] + F_t^\alpha + \kappa(\alpha) \right] \\
&\leq \inf_{\tilde{\alpha} \in \mathcal{A}_{[t, T]}} \sup_{\alpha \in \mathcal{A}_{[\tilde{t}, T]}} \left\{ k\mathbb{E}[|\mathcal{I}_{t+\delta}^{\alpha \oplus_t \tilde{\alpha}}(X)|^p - |\mathcal{I}_t^\alpha(X)|^p] + \int_t^{t+\delta} \tilde{f}_s^{\tilde{\alpha}} ds \right\}.
\end{aligned}$$

We remark that the above involves α only on $[\tilde{t}, t)$. By (10.1) we prove (4.13) immediately.

It remains to prove (ii). For any $\delta > 0$, by (4.12) there exists α^δ s.t.

$$\phi_t(\underline{X}) \geq k\mathbb{E}[|\mathcal{I}_t^{\alpha^\delta}(X)|^p] + F_t^{\alpha^\delta} + \kappa(\alpha^\delta) - \delta^2.$$

It is clear that the above property, and hence α^δ , does not involve $(\beta_s, \gamma_s)_{s \geq t}$. Then,

$$\begin{aligned}
&\phi_{t+\delta}(\underline{X}) - \phi_t(\underline{X}) \\
&\leq \left[k\mathbb{E}[|\mathcal{I}_{t+\delta}^{\alpha^\delta}(X)|^p] + F_{t+\delta}^{\alpha^\delta} + \kappa(\alpha^\delta) \right] - \left[k\mathbb{E}[|\mathcal{I}_t^{\alpha^\delta}(X)|^p] + F_t^{\alpha^\delta} + \kappa(\alpha^\delta) \right] + \delta^2 \\
&= k\mathbb{E}[|\mathcal{I}_{t+\delta}^{\alpha^\delta}(X)|^p - |\mathcal{I}_t^{\alpha^\delta}(X)|^p] + \int_t^{t+\delta} \tilde{f}_s^{\alpha^\delta} ds + \delta^2.
\end{aligned}$$

By (10.1) again we can easily prove (4.14). ■

In order to prove Proposition 5.4, we need the following slight generalization of [18, Lemma F.2].

Lemma 10.1. *Let $t \in [0, T]$ and $F : \mathcal{A}_T \rightarrow \mathbb{R}$ be continuous. Then,*

$$\lim_{s \downarrow t} \inf_{\alpha_s \in \mathcal{A}_s} F(\underline{\alpha}_s) = \inf_{\alpha_t \in \mathcal{A}_t} F(\underline{\alpha}_t). \quad (10.2)$$

Proof. Since $\mathcal{A}_t \subset \mathcal{A}_s$ for $s \geq t$, we immediately have $\lim_{s \downarrow t} \inf_{\alpha_s \in \mathcal{A}_s} F(\underline{\alpha}_s) \leq \inf_{\alpha_t \in \mathcal{A}_t} F(\underline{\alpha}_t)$. Assume by contradiction that (10.2) does not hold. Then there exist $\varepsilon > 0$ and $N \in \mathbb{N}$ such that $\mathcal{B}_n \neq \emptyset$ for all $n \geq N$, where denoting $t_n := t + \frac{1}{n}$,

$$\mathcal{B}_n := \left\{ \alpha_{t_n} \in \mathcal{A}_{t_n} : F(\underline{\alpha}_{t_n}) \leq \inf_{\alpha_t \in \mathcal{A}_t} F(\underline{\alpha}_t) - \varepsilon \right\}.$$

Note that $\mathcal{A}_{t_{n+1}} \subset \mathcal{A}_{t_n}$ and thus $\mathcal{B}_{n+1} \subset \mathcal{B}_n$. Moreover, by the continuity of F we see that \mathcal{B}_n is closed. Since \mathcal{A}_T is complete, by Cantor's intersection theorem, there exists $\hat{\alpha} \in \mathcal{A}_T$ such that $\hat{\alpha} \in \bigcap_{n \geq N} \mathcal{B}_n \subset \bigcap_{n \geq N} \mathcal{A}_{t_n}$. Note that $\bigcap_{n \geq N} \mathcal{A}_{t_n}$ is the set of \mathcal{F}_{t+} -measurable A -valued random variables. Since the augmented filtration of \mathbb{F} is right continuous, there exists $\hat{\alpha}' \in \mathcal{A}_t$ such that $\hat{\alpha}' = \hat{\alpha}$, \mathbb{P} -a.s., see e.g. [55, Proposition 1.2.1]. Then $F(\hat{\alpha}') = F(\hat{\alpha}) \leq \inf_{\alpha_t \in \mathcal{A}_t} F(\alpha_t) - \varepsilon$. This is a desired contradiction. \blacksquare

Proof of Proposition 5.4. We shall only prove the equivalence of subsolution properties. The supersolution case follows essentially the same argument.

First, note that $C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^2) \subset C^{1,2}(\overline{\mathcal{X}}_{[0,T]}^6)$. If U is a viscosity subsolution, for any $(t, \xi) \in \overline{\mathcal{X}}_{[0,T]}^6$, it is clear that $(U, \mathbf{0}) \in \mathfrak{F}_6^+ U(t, \xi)$, then it is straightforward to derive from (5.4) the classical subsolution property of U at (t, ξ) . Moreover, by the arguments in Remark 5.3 (ii), we obtain the classical subsolution property at all $(t, \xi) \in \overline{\mathcal{X}}_{[0,T]}^2$.

On the other hand, assume U is a classical subsolution. Fix $(t, \xi) \in \overline{\mathcal{X}}_{[0,T]}^6$ and $(\varphi, \phi) \in \mathfrak{F}_6^+ U(t, \xi)$. For any $\alpha \in \mathcal{A}_{[t,T]}$, recall (5.3) and denote $\bar{X}^\alpha := \bar{X}^{t,\xi,\alpha}$. By the Itô formula (3.2) and (5.1) we have: for $\forall \delta > 0$,

$$\begin{aligned} 0 &\geq \left[[U - (\varphi + \phi)]_{t+\delta}(\bar{X}^\alpha) - [U - (\varphi + \phi)]_t(\xi) \right] \\ &= \int_t^{t+\delta} \mathbb{E} \left[\partial_t(U - \varphi)_s(\bar{X}^\alpha) + \partial_X(U - \varphi)_s(\bar{X}^\alpha) \cdot b_s^{t,\xi,\alpha} \right. \\ &\quad \left. + \frac{1}{2} \partial_{\mathbf{x}X}(U - \varphi)_s(\bar{X}^\alpha) : (\sigma \sigma^\top)_s^{t,\xi,\alpha} - \dot{\phi}_s(\bar{X}^\alpha) \right] ds. \end{aligned}$$

By the smoothness of $U - \varphi$ and the regularity of b, σ, f , including Assumption 3.7, we have

$$\begin{aligned} &\int_t^{t+\delta} \left[\partial_t U_t(\underline{\xi}) + H_t(\underline{\xi}, \partial_X U_t(\underline{\xi}), \partial_{\mathbf{x}X} U_t(\underline{\xi}), \underline{\alpha}_s) \right] ds \\ &\leq \int_t^{t+\delta} \left[\partial_t \varphi_t(\underline{\xi}) + H_s(\underline{\xi}_{\cdot \wedge t}, \partial_X \varphi_t(\underline{\xi}), \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}), \underline{\alpha}_s) + \dot{\phi}_s(\bar{X}^\alpha) \right] ds + o(\delta), \end{aligned}$$

where $o(\delta)$ is uniform in α . Then

$$\begin{aligned} &\inf_{\alpha \in \mathcal{A}_{[t,T]}} \frac{1}{\delta} \int_t^{t+\delta} \left[\partial_t U_t(\underline{\xi}) + H_t(\underline{\xi}, \partial_X U_t(\underline{\xi}), \partial_{\mathbf{x}X} U_t(\underline{\xi}), \underline{\alpha}_s) \right] ds \\ &\leq \inf_{\alpha \in \mathcal{A}_{[t,T]}} \frac{1}{\delta} \int_t^{t+\delta} \left[\partial_t \varphi_t(\underline{\xi}) + H_s(\underline{\xi}_{\cdot \wedge t}, \partial_X \varphi_t(\underline{\xi}), \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}), \underline{\alpha}_s) + \dot{\phi}_s(\bar{X}^\alpha) \right] ds + o(1), \end{aligned}$$

Note that, by Assumption 3.7, we have $H_t(\underline{\xi}, \partial_X U_t(\underline{\xi}), \partial_{\mathbf{x}X} U_t(\underline{\xi}), \cdot)$ is continuous. Send $\delta \rightarrow 0$, by Lemma 10.1 we obtain

$$\mathcal{L}U_t(\underline{\xi}) \leq \partial_t \varphi_t(\underline{\xi}) + \liminf_{\delta \rightarrow 0} \inf_{\alpha \in \mathcal{A}_{[t,T]}} \frac{1}{\delta} \int_t^{t+\delta} \left[H_s(\underline{\xi}_{\cdot \wedge t}, \partial_X \varphi_t(\underline{\xi}), \partial_{\mathbf{x}X} \varphi_t(\underline{\xi}), \underline{\alpha}_s) + \dot{\phi}_s(\bar{X}^\alpha) \right] ds.$$

Then (5.4) follows from the classical subsolution property of U . ■

Proof of Proposition 6.3. (i) We shall prove only $\check{V}_t(\underline{\xi}) \geq \check{V}_t(\underline{\xi}')$. It suffices to prove $\check{J}_t(\underline{\xi}, \underline{\alpha}) \geq \check{V}_t(\underline{\xi}')$ for an arbitrarily fixed $\alpha \in \mathcal{A}_{[t,T]}$. By standard approximation arguments, we may assume without loss of generality that, for some $t = t_0 < \dots < t_n = T$:

$$\alpha_s = \sum_{i=0}^{n-1} \alpha_i(\omega, B_{\cdot \wedge t_i}^1, B_{\cdot \wedge t_i}^0) \mathbf{1}_{[t_i, t_{i+1})}(s), \quad (10.3)$$

where $\alpha_i : \Omega \times \mathbb{X} \rightarrow A$ is \mathcal{F}_0 measurable in ω and uniformly continuous in \mathbf{x} . Moreover, since \mathcal{F}_0 is rich enough, we can generate \mathcal{F}_0 -measurable random variables $\{\mathbf{U}_i\}_{i \geq 0}$ which are i.i.d. with distribution $\text{Uniform}([0, 1])$ and are independent of (ξ, ξ', α) , and hence are independent of $X^{t, \xi, \alpha}$. Denote $X^{\tilde{\alpha}} := X^{t, \xi, \tilde{\alpha}}$, $X'^{\tilde{\alpha}} := X^{t, \xi', \tilde{\alpha}}$ for arbitrary $\tilde{\alpha}$.

First, since $\mathbb{P}_{\xi, \wedge t | \mathcal{F}_t^0} = \mathbb{P}_{\xi', \wedge t | \mathcal{F}_t^0}$, \mathbb{P} -a.s., then $\mathbb{P}_{(\xi, \wedge t_0, B^0)} = \mathbb{P}_{(\xi', \wedge t_0, B^0)}$. By using \mathbf{U}_0 , one can easily construct $\alpha'_{t_0} \in \mathcal{A}_{t_0}$ such that $\mathbb{P}_{(\alpha_{t_0}, \xi, \wedge t_0, B^0)} = \mathbb{P}_{(\alpha'_{t_0}, \xi', \wedge t_0, B^0)}$. This clearly implies that $\mathbb{P}_{(\alpha_{t_0}, X_{\cdot \wedge t_1}^\alpha, B^0)} = \mathbb{P}_{(\alpha'_{t_0}, X'^{\alpha'}_{\cdot \wedge t_1}, B^0)}$. Repeat the arguments by using the \mathbf{U}_i , we may construct $\alpha' = \sum_{i=0}^{n-1} \alpha'_i \mathbf{1}_{[t_i, t_{i+1})} \in \mathcal{A}_{[t, T]}$ such that $\mathbb{P}_{(\alpha, X^\alpha, B^0)} = \mathbb{P}_{(\alpha', X'^{\alpha'}, B^0)}$. Then clearly $\check{J}_t(\underline{\xi}, \underline{\alpha}) = \check{J}_t(\underline{\xi}', \underline{\alpha}') \geq \check{V}_t(\underline{\xi}')$. By the arbitrariness of α we have $\check{V}_t(\underline{\xi}) \geq \check{V}_t(\underline{\xi}')$. Similarly $\check{V}_t(\underline{\xi}') \geq \check{V}_t(\underline{\xi})$, then equality holds and hence (6.4) is well defined.

Moreover, by (2.6) and Proposition 6.2 we obtain part of (6.6) immediately:

$$|\check{V}_t(\mu)| \leq C \left(1 + W_2(\mu, \wedge t, \delta_0)\right), \quad |\check{V}_t(\mu) - \check{V}_t(\mu')| \leq CW_2(\mu, \wedge t, \mu'_{\wedge t'}). \quad (10.4)$$

(ii) We proceed in several steps. For $(t, \xi) \in \overline{\mathcal{X}}_{[0, T]}^2$ and $\alpha \in \mathcal{A}_{[t, T]}$, denote

$$X^\alpha := X^{t, \xi, \alpha}, \quad Y_t^\alpha := \mathbb{E}_{\mathcal{F}_t^0} \left[\check{g}(X^\alpha, \mathbb{P}_{X^\alpha | \mathcal{F}_T^0}) + \int_t^T \check{f}_s(X^\alpha, \alpha_s, \mathbb{P}_{(X^\alpha, \alpha_s) | \mathcal{F}_s^0}) ds \right].$$

Step 1. We first assume $\xi \in \mathcal{X}_2^\perp(\mathbb{F}_t^0)$. Then $\mathbb{P}_{\xi, \wedge t | \mathcal{F}_t^0} = \mathbb{P}_{\xi, \wedge t}$, and thus the claimed result follows directly from (6.4).

Moreover, assume $\alpha \in \mathcal{A}_{[t, T]}$ takes the form (10.3). For each $\mathbf{x}^0 \in C^0([0, T]; \mathbb{R}^{d_0})$ with $\mathbf{x}_0^0 = 0$, where d_0 is the dimension of B^0 , denote

$$\alpha^{\mathbf{x}^0} := \sum_{i=0}^{n-1} \alpha_i(\omega, B_{\cdot \wedge t_i}^1, (\mathbf{x}^0 \otimes_t B^0)_{\cdot \wedge t_i}) \mathbf{1}_{[t_i, t_{i+1})} \in \mathcal{A}_{[t, T]}, \quad \text{and} \quad \check{J}'_t(\underline{\xi}, \underline{\alpha}, \mathbf{x}^0) := \check{V}_t(\underline{\xi}, \underline{\alpha}^{\mathbf{x}^0}),$$

$$\text{where} \quad (\mathbf{x}^0 \otimes_t B^0)_s := \mathbf{x}_s^0 \mathbf{1}_{[0, t)}(s) + [\mathbf{x}_t^0 + B_s^0 - B_t^0] \mathbf{1}_{[t, T]}(s).$$

We note that this is in the spirit of the regular conditional probability distribution, but we avoid it by considering the special form (10.3). Since $\xi \in \mathcal{X}_2^\perp(\mathbb{F}_t^0)$, we can easily show that

$$Y_t^\alpha = J'_t(\underline{\xi}, \underline{\alpha}, B^0), \quad \mathbb{P}\text{-a.s.} \quad (10.5)$$

In particular, this implies that $Y_t^\alpha \geq \check{V}_t(\underline{\xi})$, \mathbb{P} -a.s.

Step 2. We next consider the case $\xi = \sum_{i=1}^n \xi_i \mathbf{1}_{E_i}$, where $\xi_i \in \mathcal{X}_2^\perp(\mathbb{F}_t^0)$ and $\{E_i\}_{1 \leq i \leq n} \subset \mathcal{F}_t^0$ form a partition of Ω . Note that $\mathbb{P}_{\xi|\mathcal{F}_t^0} = \sum_{i=1}^n \mathbb{P}_{\xi_i} \mathbf{1}_{E_i}$, \mathbb{P} -a.s. Then

$$V_t'(\underline{\xi}) := \mathbb{E}[\check{V}_t(\mathbb{P}_{\xi|\mathcal{F}_t^0})] = \sum_{i=1}^n \check{V}_t(\mathbb{P}_{\xi_i}) \mathbb{P}(E_i).$$

We first prove $V_t(\underline{\xi}) \geq V_t'(\underline{\xi})$. For any $\alpha \in \mathcal{A}_{[t,T]}$ taking the form (10.3), by (10.5) we have

$$J_t(\underline{\xi}, \underline{\alpha}) = \mathbb{E}[Y_t^\alpha] = \mathbb{E}\left[\sum_{i=1}^n Y_t^{t,\xi_i,\alpha} \mathbf{1}_{E_i}\right] \geq \mathbb{E}\left[\sum_{i=1}^n \check{V}_t(\mathbb{P}_{\xi_i}) \mathbf{1}_{E_i}\right] = V_t'(\underline{\xi}).$$

Now by standard approximation arguments, we have $J_t(\underline{\xi}, \underline{\alpha}) \geq V_t'(\underline{\xi})$ for all $\alpha \in \mathcal{A}_{[t,T]}$, and thus $V_t(\underline{\xi}) \geq V_t'(\underline{\xi})$.

To see the opposite inequality, fix an arbitrary $\varepsilon > 0$. For each $i = 1, \dots, n$, there exists α^i taking the form (10.3) (with different n there) such that $Y_t^{t,\xi_i,\alpha^i} \leq \check{V}_t(\mathbb{P}_{\xi_i}) + \varepsilon$. Construct $\alpha^\varepsilon := \sum_{i=1}^n \alpha^i \mathbf{1}_{E_i}$. One can easily see that $Y_t^{t,\xi,\alpha^\varepsilon} = \sum_{i=1}^n Y_t^{t,\xi_i,\alpha^i} \mathbf{1}_{E_i}$. Then

$$V_t(\underline{\xi}) \leq \check{J}_t(\underline{\xi}, \alpha^\varepsilon) = \mathbb{E}[Y_t^{t,\xi,\alpha^\varepsilon}] = \mathbb{E}\left[\sum_{i=1}^n Y_t^{t,\xi_i,\alpha^i} \mathbf{1}_{E_i}\right] \leq \mathbb{E}\left[\sum_{i=1}^n [\check{V}_t(\mathbb{P}_{\xi_i}) + \varepsilon] \mathbf{1}_{E_i}\right] = V_t'(\underline{\xi}) + \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, we obtain $V_t(\underline{\xi}) \geq V_t'(\underline{\xi})$, and hence the equality.

Step 3. By the regularity of V in (2.6) and that of \check{V} in (10.4), and by standard approximation arguments, similarly to (10.3) we may assume without loss of generality that ξ takes the following form: for some $0 = t_0 < \dots < t_n = t$,

$$\begin{aligned} \xi_s \equiv \xi'_{t_0}, \quad t \in [t_0, t_1]; \quad \xi_s = \frac{t_{i+1} - t}{t_{i+1} - t_i} \xi'_{t_{i-1}} + \frac{t - t_i}{t_{i+1} - t_i} \xi'_{t_i}, \quad t \in [t_i, t_{i+1}], \quad i = 1, \dots, n-1; \\ \text{where } \xi'_{t_i} = h_i\left(\omega, (B_{t_1}^1, \dots, B_{t_i}^1), (B_{t_1}^0, \dots, B_{t_i}^0)\right), \end{aligned}$$

and $h_i : \Omega \times \mathbb{R}^{di} \rightarrow \mathbb{R}^d$ is \mathcal{F}_0 measurable in ω and uniformly continuous in $y \in \mathbb{R}^{di}$. Fix $\varepsilon > 0$, and let $\{O_j^\varepsilon\}_{1 \leq j \leq n_\varepsilon}$ be a partition of $\{y \in \mathbb{R}^{d_0 n} : |y| \leq \frac{1}{\varepsilon}\}$ such that each O_j^ε has diameter less than ε . Fix an arbitrary $y_j^\varepsilon \in O_j^\varepsilon$ for each j , and denote $O_j^{\varepsilon,i} := \{Proj_i(y) : y \in O_j^\varepsilon\}$, where $Proj_i(y) \in \mathbb{R}^{di}$ denotes the first i components of $y \in \mathbb{R}^{dn}$. Define

$$\begin{aligned} \xi_s^\varepsilon \equiv \xi'_{t_0}, \quad t \in [t_0, t_1]; \quad \xi_s^\varepsilon = \frac{t_{i+1} - t}{t_{i+1} - t_i} \xi'_{t_{i-1}}^\varepsilon + \frac{t - t_i}{t_{i+1} - t_i} \xi'_{t_i}^\varepsilon, \quad t \in [t_i, t_{i+1}], \quad i = 1, \dots, n-1; \\ \text{where } \xi'_{t_i}^\varepsilon = \sum_{j=1}^{n_\varepsilon} \xi_{t_i}^{\varepsilon,i} \mathbf{1}_{\{(B_{t_1}^0, \dots, B_{t_i}^0) \in O_j^{\varepsilon,i}\}}, \quad \xi_{t_i}^{\varepsilon,i} := h_i\left(\omega, (B_{t_1 \wedge t_i}^1, \dots, B_{t_n \wedge t_i}^1), Proj_i(y_j^\varepsilon)\right). \end{aligned}$$

Note that $\xi_{t_i}^{\varepsilon,i}$ is independent of \mathcal{F}_t^0 , and, on $\{(B_{t_1}^0, \dots, B_{t_i}^0) \notin \cup_j O_j^{\varepsilon,i}\}$, $\xi_{t_i}^{\varepsilon,i} = 0$, which is also independent of \mathcal{F}_t^0 . Then one can easily see that ξ^ε takes the form as in Step 2, and

thus it follows from Step 2 that $V_t(\underline{\xi}^\varepsilon) = V_t'(\underline{\xi}^\varepsilon)$. Send $\varepsilon \rightarrow 0$, by the desired regularities we obtain $V_t(\underline{\xi}) = V_t'(\underline{\xi})$.

(iii) follows (ii) and DPP (2.5) for V . Finally, for $0 \leq t < t + \delta \leq T$, $\mu \in \mathcal{P}_2(\mathbb{X})$, and $\xi \in \mathcal{X}_2^\perp(\mathbb{F}^0; t, \mu)$, by (iii) we have, again denoting $X^\alpha := X^{t, \xi, \alpha}$,

$$\check{V}_t(\mu) - \check{V}_{t+\delta}(\mu_{\cdot \wedge t}) = \inf_{\alpha \in \mathcal{A}_{[t, T]}} \mathbb{E} \left[\check{V}_{t+\delta}(\mathbb{P}_{X^\alpha | \mathcal{F}_{t+\delta}^0}) - \check{V}_{t+\delta}(\mathbb{P}_{X_{\cdot \wedge t}^\alpha}) + \int_t^{t+\delta} \check{f}_s(X^\alpha, \alpha_s, \mathbb{P}_{(X^\alpha, \alpha_s) | \mathcal{F}_s^0}) ds \right].$$

Apply (10.4) on $\check{V}_{t+\delta}(\mathbb{P}_{X^\alpha | \mathcal{F}_{t+\delta}^0}) - \check{V}_{t+\delta}(\mathbb{P}_{X_{\cdot \wedge t}^\alpha})$, we can easily obtain

$$|\check{V}_t(\mu) - \check{V}_{t+\delta}(\mu_{\cdot \wedge t})| \leq C(1 + W_2(\mu_{\cdot \wedge t}, \delta_0))\sqrt{\delta}.$$

This, together with (10.4) again, implies (6.6) immediately. \blacksquare

We now turn to the proof of Proposition 6.4. For that purpose we first introduce briefly the path derivatives and specify the space $C_b^{1,2,2}([0, T] \times \mathbb{X} \times \mathcal{P}_2(\mathbb{X}))$, which actually serves as the technical conditions for the proposition. We refer to [14, 54] for more details. We first need to extend U to the càdlàg space. Let \mathbb{D} denote the space of d -dimensional càdlàg paths on $[0, T]$, equipped with the uniform norm, and $\mathcal{P}_2(\mathbb{D})$ the space of square integrable probability measures on \mathbb{D} , equipped with the 2-Wasserstein distance. Given an adapted function $U : [0, T] \times \mathbb{X} \times \mathcal{P}_2(\mathbb{X}) \rightarrow \mathbb{R}$, let $\hat{U} : [0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}) \rightarrow \mathbb{R}$ be an extension. The path derivatives of \hat{U} are functions $\partial_t \hat{U} : [0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}) \rightarrow \mathbb{R}$, $\partial_{\mathbf{x}} \hat{U} : [0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}) \rightarrow \mathbb{R}^d$ and $\partial_\mu \hat{U} : [0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ determined by:

$$\begin{aligned} \partial_t \hat{U}_t(\mathbf{x}, \mu) &:= \lim_{\delta \rightarrow 0} \frac{1}{\delta} \left[\hat{U}_{t+\delta}(\mathbf{x}_{\cdot \wedge t}, \mu_{\cdot \wedge t}) - \hat{U}_t(\mathbf{x}_{\cdot \wedge t}, \mu_{\cdot \wedge t}) \right]; \\ \hat{U}_t(\mathbf{x} + \Delta x \mathbf{1}_{[t, T]}, \mu) - \hat{U}_t(\mathbf{x}, \mu) &= \partial_{\mathbf{x}} \hat{U}_t(\mathbf{x}, \mu) \cdot \Delta x + o(|\Delta x|), \quad \forall \Delta x \in \mathbb{R}^d; \\ \hat{U}_t(\mathbf{x}, \mathbb{P}_{\xi + \eta_t \mathbf{1}_{[t, T]}}) - \hat{U}_t(\mathbf{x}, \mathbb{P}_\xi) &= \mathbb{E} \left[\partial_\mu \hat{U}_t(\mathbf{x}, \mathbb{P}_\xi, \xi_t) \cdot \eta_t \right] + o(\|\eta_t\|_2), \quad \forall \eta_t \in \mathbb{L}^2(\mathcal{F}_t; \mathbb{R}^d). \end{aligned}$$

In particular, it can be proved that the Lions' derivative $\partial_\mu \hat{U}$ takes the above specific structure. We can then define the higher order derivatives in the same manner. In particular, $\partial_{\tilde{x}\mu} \hat{U}_t(\mathbf{x}, \mu, \tilde{x})$ is the standard (finite-dimensional) derivative of $\partial_\mu \hat{U}$ with respect to \tilde{x} , and $\partial_{\mu\mu} \hat{U} : [0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}) \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{S}^d$ involves two additional variables $(\tilde{x}, \bar{x}) \in \mathbb{R}^d \times \mathbb{R}^d$. Let $C_b^{1,2,2}([0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}))$ denote the space of adapted and continuous functions $\hat{U} : [0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}) \times \mathbb{R}^d \rightarrow \mathbb{R}$ such that the derivatives $\partial_t \hat{U}, \partial_{\mathbf{x}} \hat{U}, \partial_\mu \hat{U}, \partial_{\mathbf{x}\mathbf{x}} \hat{U}, \partial_{\mathbf{x}\mu} \hat{U}, \partial_{\tilde{x}, \mu} \hat{U}, \partial_{\mu\mu} \hat{U}$ exist, and are continuous and bounded.

We then define the path derivatives of U by restricting those of \hat{U} to the continuous paths, for example, $\partial_\mu U$ is defined by restricting $\partial_\mu \hat{U}$ on $[0, T] \times \mathbb{X} \times \mathcal{P}_2(\mathbb{X}) \times \mathbb{R}^d$. It can be shown that the path derivatives of U are independent of the choice of the extension \hat{U} and

thus are intrinsic to U . Moreover, we say $U \in C_b^{1,2,2}([0, T] \times \mathbb{X} \times \mathcal{P}_2(\mathbb{X}))$ if there exists an extension $\hat{U} \in C_b^{1,2,2}([0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}))$.

Proof of Proposition 6.4. We shall only prove (6.7) in the integral form on $[0, T]$ for an arbitrary extension $\hat{U} \in C_b^{1,2,2}([0, T] \times \mathbb{D} \times \mathcal{P}_2(\mathbb{D}))$. For notational simplicity we denote $U = \hat{U}$ at below. Set $0 = t_0 < \dots < t_n = T$ with $t_i = \frac{i}{n}T$. For $j = 1, 2$, denote

$$\begin{aligned} X_t^{j,n} &:= \sum_{i=0}^{n-1} X_{t_i}^j \mathbf{1}_{[t_i, t_{i+1})}(t) + X_T^j \mathbf{1}_{\{t_n\}}(t), \quad \mu^n := \mathbb{P} \circ (X^{2,n})^{-1}; \\ X_t^{j,n,i,l} &:= X_{\cdot \wedge t_i}^{j,n} + l X_{t_i, t_{i+1}}^j \mathbf{1}_{[t_{i+1}, T]}, \quad \mu^{n,i,l} := \mathbb{P} \circ (X^{2,n,i,l})^{-1}, \quad l \in [0, 1], \end{aligned}$$

where $X_{t_i, t_{i+1}}^j := X_{t_{i+1}}^j - X_{t_i}^j$. Note that $X_{\cdot \wedge t_{i+1}}^{j,n} = X_{\cdot \wedge t_i}^{j,n} + X_{t_i, t_{i+1}}^j \mathbf{1}_{[t_{i+1}, T]}$. Then,

$$\begin{aligned} U_T(X^{1,n}, \mu^n) - U_0(X^{1,n}, \mu^n) &= \sum_{i=0}^{n-1} \left[U_{t_{i+1}}(X^{1,n}, \mu^n) - U_{t_i}(X^{1,n}, \mu^n) \right] \\ &= \sum_{i=0}^{n-1} \left[[U_{t_{i+1}}(X_{\cdot \wedge t_i}^{1,n}, \mu_{\cdot \wedge t_i}^n) - U_{t_i}(X^{1,n}, \mu^n)] + [U_{t_{i+1}}(X^{1,n}, \mu^n) - U_{t_{i+1}}(X_{\cdot \wedge t_i}^{1,n}, \mu_{\cdot \wedge t_i}^n)] \right] \\ &= \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \partial_t U_t(X_{\cdot \wedge t_i}^{1,n}, \mu_{\cdot \wedge t_i}^n) dt + \sum_{i=0}^{n-1} \int_0^1 [I_1^{n,i}(l) + I_2^{n,i}(l)] dl, \end{aligned}$$

where, denoting by $(\tilde{X}^{2,n,i,l}, \tilde{X}_{t_i, t_{i+1}}^2)$ and $(\bar{X}^{2,n,i,l}, \bar{X}_{t_i, t_{i+1}}^2)$ conditionally independent copies of $(X^{2,n,i,l}, X_{t_i, t_{i+1}}^2)$, conditional on \mathbb{F}^0 ,

$$\begin{aligned} I_1^{n,i}(l) &:= \partial_{\mathbf{x}} U_{t_{i+1}}(X^{1,n,i,l}, \mu^{n,i,l}) \cdot X_{t_i, t_{i+1}}^1, \\ I_2^{n,i}(l) &:= \mathbb{E}_{\mathcal{F}_{t_{i+1}}} [\partial_{\mu} U_{t_{i+1}}(X^{1,n,i,l}, \mu^{n,i,l}, \tilde{X}^{2,n,i,l}) \cdot \tilde{X}_{t_i, t_{i+1}}^2]. \end{aligned}$$

Let $o(\frac{1}{n})$ denote a generic term whose \mathbb{L}^2 -norm vanishes faster than $\frac{1}{n}$ when $n \rightarrow \infty$. By the desired regularities, one can easily check that

$$\begin{aligned} I_1^{n,i}(l) &= \partial_{\mathbf{x}} U_{t_{i+1}}(X_{\cdot \wedge t_i}^{1,n}, \mu_{\cdot \wedge t_i}^n) \cdot X_{t_i, t_{i+1}}^1 \\ &\quad + \left[\partial_{\mathbf{x}} U_{t_{i+1}}(X^{1,n,i,l}, \mu^{n,i,l}) - \partial_{\mathbf{x}} U_{t_{i+1}}(X^{1,n,i,0}, \mu^{n,i,0}) \right] \cdot X_{t_i, t_{i+1}}^1 \\ &= \partial_{\mathbf{x}} U_{t_{i+1}}(X_{\cdot \wedge t_i}^{1,n}, \mu_{\cdot \wedge t_i}^n) \cdot X_{t_i, t_{i+1}}^1 + \int_0^l \left[\partial_{\mathbf{xx}} U_{t_{i+1}}(X^{1,n,i,l'}, \mu^{n,i,l'}) X_{t_i, t_{i+1}}^1 \right. \\ &\quad \left. + \mathbb{E}_{\mathcal{F}_{t_{i+1}}} [\partial_{\mathbf{x}\mu} U_{t_{i+1}}(X^{1,n,i,l'}, \mu^{n,i,l'}, \tilde{X}^{2,n,i,l'}) \tilde{X}_{t_i, t_{i+1}}^2] \right] dl' \cdot X_{t_i, t_{i+1}}^1 \\ &= \int_{t_i}^{t_{i+1}} \partial_{\mathbf{x}} U_t(X^1, \mu) \cdot dX_t^1 + l \partial_{\mathbf{xx}} U_{t_{i+1}}(X_{\cdot \wedge t_i}^{1,n}, \mu_{\cdot \wedge t_i}^n) : (X_{t_i, t_{i+1}}^1)(X_{t_i, t_{i+1}}^1)^\top \\ &\quad + l \mathbb{E}_{\mathcal{F}_{t_{i+1}}} [\partial_{\mathbf{x}\mu} U_{t_{i+1}}(X_{\cdot \wedge t_i}^{1,n}, \mu_{\cdot \wedge t_i}^n, \tilde{X}_{\cdot \wedge t_i}^{2,n}) : (\tilde{X}_{t_i, t_{i+1}}^2)(X_{t_i, t_{i+1}}^1)^\top] + o(\frac{1}{n}) \end{aligned}$$

$$\begin{aligned}
&= \int_{t_i}^{t_{i+1}} \partial_x U_t(X^1, \mu) \cdot dX_t^1 + l \int_{t_i}^{t_{i+1}} \partial_{xx} U_t(X^1, \mu) : \sigma_t^1 (\sigma_t^1)^\top dt \\
&\quad + l \int_{t_i}^{t_{i+1}} \mathbb{E}_{\mathcal{F}_{t_{i+1}}} [\partial_{x\mu} U_t(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^{2,0} (\sigma^{1,0})^\top] dt + o\left(\frac{1}{n}\right); \\
I_2^{n,i}(l) &= \mathbb{E}_{\mathcal{F}_{t_{i+1}}} \left[\partial_\mu U_{t_{i+1}}(X^1, \mu, \tilde{X}^2) \cdot \tilde{X}_{t_i, t_{i+1}}^2 \right. \\
&\quad \left. + [\partial_\mu U_{t_{i+1}}(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^{2,0} (\sigma^{1,0})^\top] \cdot \tilde{X}_{t_i, t_{i+1}}^2 \right] \\
&= \mathbb{E}_{\mathcal{F}_{t_{i+1}}} \left[\partial_\mu U_{t_{i+1}}(X^1, \mu, \tilde{X}^2) \cdot \tilde{X}_{t_i, t_{i+1}}^2 \right] \\
&\quad + \int_0^l \mathbb{E}_{\mathcal{F}_{t_{i+1}}} \left[[\partial_{x\mu} U_{t_{i+1}}(X^1, \mu, \tilde{X}^2) X_{t_i, t_{i+1}}^1 \right. \\
&\quad \quad + \partial_{\mu\mu} U_{t_{i+1}}(X^1, \mu, \tilde{X}^2) \tilde{X}_{t_i, t_{i+1}}^2 \\
&\quad \quad \left. + \partial_{\tilde{x}\mu} U_{t_{i+1}}(X^1, \mu, \tilde{X}^2) \tilde{X}_{t_i, t_{i+1}}^2] \cdot \tilde{X}_{t_i, t_{i+1}}^2 \right] dl' \\
&= \mathbb{E}_{\mathcal{F}_{t_{i+1}}} \left[\int_{t_i}^{t_{i+1}} \partial_\mu U_t(X^1, \mu, \tilde{X}^2) \cdot d\tilde{X}_t^2 \right] \\
&\quad + l \mathbb{E}_{\mathcal{F}_{t_{i+1}}} \left[\int_{t_i}^{t_{i+1}} [\partial_{x\mu} U_t(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^{2,0} (\sigma^{1,0})^\top \right. \\
&\quad \quad \left. + \partial_{\mu\mu} U_t(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^{2,0} (\tilde{\sigma}^{2,0})^\top + \partial_{\tilde{x}\mu} U_t(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^2 (\tilde{\sigma}^2)^\top] dt + o\left(\frac{1}{n}\right). \right.
\end{aligned}$$

Then, noting that $\int_0^1 l dl = \frac{1}{2}$,

$$\begin{aligned}
&U_T(X^1, \mu) - U_0(X^1, \mu) = U_T(X^1, \mu^n) - U_0(X^1, \mu^n) + o(1) \\
&= \int_0^T \left[\partial_t U_t(X^1, \mu) dt + \partial_x U_t(X^1, \mu) \cdot dX_t^1 + \frac{1}{2} \partial_{xx} U_t(X^1, \mu) : \sigma_t^1 (\sigma_t^1)^\top \right] dt \\
&\quad + \int_0^T \mathbb{E}_{\mathcal{F}_t} \left[\partial_{x\mu} U_t(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^{2,0} (\sigma^{1,0})^\top + \frac{1}{2} \partial_{\mu\mu} U_t(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^{2,0} (\tilde{\sigma}^{2,0})^\top \right. \\
&\quad \quad \left. + \frac{1}{2} \partial_{\tilde{x}\mu} U_t(X^1, \mu, \tilde{X}^2) : \tilde{\sigma}_t^2 (\tilde{\sigma}^2)^\top \right] dt + o(1).
\end{aligned}$$

By sending $n \rightarrow \infty$, this is exactly (6.7). ■

References

- [1] Ahuja, S.; Ren, W.; and Yang, T.W. *Forward-backward stochastic differential equations with monotone functionals and mean field games with common noise*, *Stochastic Process. Appl.*, 129 (2019), 3859-3892.
- [2] Aubin, J.-P. and Ekeland, I. *Applied Nonlinear Analysis*, Wiley, 1984.
- [3] Bayraktar, E.; Ekren, I.; and Zhang, X. *Comparison of viscosity solutions for a class of second order PDEs on the Wasserstein space*, preprint, arXiv:2309.05040.

- [4] Bensoussan, A.; Graber, P.J.; and Yam, S.C.P. *Stochastic control on space of random variables*, Preprint, arXiv:1903.12602.
- [5] Bensoussan, A.; Graber, P.J.; and Yam, S.C.P. *Control on Hilbert spaces and application to mean field type control theory*, Preprint, arXiv:2005.10770.
- [6] Bensoussan, A.; Tai, H.M.; and Yam, S.C.P. *Mean Field Type Control Problems, Some Hilbert-space-valued FBSDEs, and Related Equations*, Preprint, arXiv:2305.04019.
- [7] Bertucci, C. *Stochastic optimal transport and Hamilton-Jacobi-Bellman equations on the set of probability measures*, preprint, arXiv:2306.04283.
- [8] Borwein, J. M. and Zhu, Q. J. *Techniques of Variational Analysis*, volume 20 of CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer-Verlag, New York, 2005.
- [9] Burzoni, M.; Ignazio, V.; Reppen, A. M.; and Soner, H. M. *Viscosity solutions for controlled McKean-Vlasov jump-diffusions*, *SIAM J. Control Optim.*, 58 (2020), 1676-1699.
- [10] Caines, P.E.; Huang, M.; and Malhame, R.P. *Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle*, *Commun. Inf. Syst.*, 6 (2006), 221-252.
- [11] Cardaliaguet, P. and Quincampoix, M. *Deterministic differential games under probability knowledge of initial condition*, *Int. Game Theory Rev.*, 10 (2008), 1-16.
- [12] Carmona, R.; Cormier, Q.; and Soner, H.M. *Synchronization in a Kuramoto mean field game*, preprint, arXiv:2210.12912.
- [13] Carmona, R. and Delarue, F. *Probabilistic Theory of Mean Field Games I - Mean Field FBSDEs, Control, and Games*, Springer, 2018.
- [14] Carmona, R. and Delarue, F. *Probabilistic Theory of Mean Field Games II - Mean Field Games with Common Noise and Master Equations*, Springer, 2018.
- [15] Cecchin, A. and Delarue, F. *Weak solutions to the master equation of potential mean field games*, preprint, arXiv:2204.04315.
- [16] Conforti, G.; Kraaij, R.; and Tonon, D. *Hamilton-Jacobi equations for controlled gradient flows: the comparison principle*, preprint, arXiv:2111.13258.

- [17] Cont, R. and Fournié, D.-A. *Functional Itô calculus and stochastic integral representation of martingales*, *Ann. Probab.*, 41 (2013), 109-133.
- [18] Cosso, A.; Gozzi, F.; Kharroubi, I.; Pham, H.; and Rosestolato, M. *Optimal control of path dependent McKean-Vlasov SDEs in infinite dimension*, *Ann. Appl. Probab.*, 33 (2023), 2863-2918.
- [19] Cosso, A.; Gozzi, F.; Kharroubi, I.; Pham, H.; and Rosestolato, M. *Master Bellman equation in the Wasserstein space: Uniqueness of viscosity solutions*, *Trans. Amer. Math. Soc.*, DOI: <https://doi.org/10.1090/tran/8986>, arXiv:2107.10535.
- [20] Cox, A. M.; Kallblad, S.; Larsson, M.; and Svaluto-Ferro, S. *Controlled measure-valued martingales: a viscosity solution approach*, *Ann. Appl. Probab.*, accepted, arXiv:2109.00064.
- [21] Crandall, M. G. and Ishii, H. *The maximum principle for semicontinuous functions*, *Differ. Integral Equ.*, 3 (1990), 1001-1014.
- [22] Crandall, M. G.; Ishii, H.; and Lions, P. L. *User's guide to viscosity solutions of second order partial differential equations*, *Bull. Amer. Math. Soc.*, 27 (1992), 1-67.
- [23] Crandall, M. G. and Lions, P. L. *Viscosity solutions of Hamilton-Jacobi equations in infinite dimensions. V. Unbounded linear terms and B-continuous solutions*, *J. Func. Anal.*, 97 (1991), 417-465.
- [24] Daudin, S. and Seeger, B. *A comparison principle for semilinear Hamilton-Jacobi-Bellman equations in the Wasserstein space*, preprint, arXiv:2308.15174.
- [25] Daudin, S.; Jackson, J.; and Seeger, B. *Wellposedness of Hamilton-Jacobi equations in the Wasserstein space: non-convex Hamiltonians and common noise*, preprint, arXiv:2312.02324.
- [26] Dupire, B. *Functional Itô calculus*, *Quant. Finance*, 19 (2019), 721-729.
- [27] Fadle, A. and Touzi, N. *Itô and Itô-Wentzell chain rule for flows of conditional laws of continuous semimartingales: an easy approach*, Preprint.
- [28] Ekren, I.; Keller, C.; Touzi, N.; and Zhang, J. *On viscosity solutions of path dependent PDEs*, *Ann. Probab.*, 42 (2014), 204-236.

- [29] Ekren, I.; Touzi, N.; and Zhang, J. *Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part I*, *Ann. Probab.*, 44 (2016), 1212-1253.
- [30] Ekren, I.; Touzi, N.; and Zhang, J. *Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part II*, *Ann. Probab.*, 44 (2016), 2507-2553.
- [31] Feng, J. and Katsoulakis, M. *A comparison principle for Hamilton-Jacobi equations related to controlled gradient flows in infinite dimensions*, *Arch. Ration. Mech. An.*, 192 (2009), 275-310.
- [32] Feng, J. and Świąch, A. *Optimal control for a mixed flow of Hamiltonian and gradient type in space of probability measures*, *Trans. Amer. Math. Soc.*, 365 (2013), 3987-4039.
- [33] Fabbri, G.; Gozzi, F.; and Świąch, A. *Stochastic Optimal Control in Infinite Dimension. Dynamic Programming and HJB Equations*, in: *Probability Theory and Stochastic Modelling*, volume 82, Springer, 2017.
- [34] Gangbo, W.; Mayorga, S.; and Swiech, A. *Finite dimensional approximations of Hamilton-Jacobi-Bellman equations in spaces of probability measures*, *SIAM J. Math. Anal.*, 53 (2021), 1320-1356.
- [35] Gangbo, W. and Meszaros, A. R. *Global well-posedness of master equations for deterministic displacement convex potential mean field games*, *Comm. Pure Appl. Math.*, 75 (2022), 2685-2801.
- [36] Gangbo, W.; Nguyen, T.; and Tudorascu, A. *Hamilton-Jacobi equations in the Wasserstein space*, *Methods Appl. Anal.*, 15 (2008), 155-184.
- [37] Gangbo, W. and Tudorascu, A. *On differentiability in the Wasserstein space and well-posedness for Hamilton-Jacobi equations*, *J. Math. Pures Appl.* 125 (2019), 119-174.
- [38] Gomes, D. A. and Voskanyan, V. K., *Extended mean field games*, *Izv. Nats. Akad. Nauk Armenii Mat.*, 48 (2013), 63-76.
- [39] Jimenez, C.; Marigonda, A.; and Quincampoix, M. *Optimal control of multiagent systems in the Wasserstein space*, *Calc. Var. Partial Differential Equations*, 59 (2020), 1-45.
- [40] Jimenez, C.; Marigonda, A.; and Quincampoix, M. *Dynamical systems and Hamilton-Jacobi-Bellman equations on the Wasserstein space and their L^2 representations*, *SIAM J. Math. Anal.*, 55 (2023), 5919-5966.

- [41] Karatzas, I. A. and Shreve, S. E. *Brownian Motion and Stochastic Calculus*, Springer, New York, 1991.
- [42] Lasry, J. and Lions, P.L. *Mean field games*, *Jpn. J. Math.*, 2 (2007), 229-260.
- [43] Li, X. and Yong, J. *Optimal Control Theory for Infinite Dimensional Systems*, Birkhäuser, 1995.
- [44] Lions, P. L. *Viscosity solutions of fully nonlinear second-order equations and optimal stochastic control in infinite dimensions. I. The case of bounded stochastic evolutions*, *Acta Math.*, 161 (1988), 243-278.
- [45] Lions, P. L. *Viscosity solutions of fully nonlinear second-order equations and optimal stochastic control in infinite dimensions. II. Optimal control of Zakai's equation*. in: *Stochastic partial differential equations and applications, II*, eds. G. Da Prato, L. Tubaro, 147-170, Lecture Notes in Mathematics 1390, Springer, 1989.
- [46] Lions, P. L. *Viscosity solutions of fully nonlinear second-order equations and optimal stochastic control in infinite dimensions. III. Uniqueness of viscosity solutions for general second-order equations*, *J. Funct. Anal.*, 86 (1989), 1-18.
- [47] Lions, P.-L. *Cours au Collège de France*, www.college-de-france.fr (2007).
- [48] Mayorga, S. and Świąch, A. *Finite dimensional approximations of Hamilton-Jacobi-Bellman equations for stochastic particle systems with common noise*, *SIAM J. Control Optim.*, 61 (2023), 820-851.
- [49] Ren, Z.; Touzi, N.; and Zhang, J. *Comparison of viscosity solutions of fully nonlinear degenerate parabolic path-dependent PDEs*, *SIAM J. Math. Anal.*, 49 (2017), 4093-4116.
- [50] Soner, H. M. and Yan, Q. *Viscosity solutions for McKean-Vlasov control on a torus*, Preprint, arXiv:2212.11053.
- [51] Soner, H. M. and Yan, Q. *Viscosity Solutions of the Eikonal Equation on the Wasserstein Space*, Preprint, arXiv:2308.04097.
- [52] Talbi, M.; Touzi, N.; and Zhang, J. *Dynamic Programming Equation for the Mean Field Optimal Stopping Problem*, *SIAM J. Control Optim.*, 61 (2023), 2140-2164.
- [53] Talbi, M.; Touzi, N.; and Zhang, J. *Viscosity solutions for obstacle problems on Wasserstein space*, *SIAM J. Control Optim.*, 61 (2023), 1712-1736.

- [54] Wu, C. and Zhang, J. *Viscosity solutions to parabolic master equations and McKean-Vlasov SDEs with closed-loop controls*, *Ann. Appl. Probab.*, 30 (2020), 936-986.
- [55] Zhang, J. *Backward Stochastic Differential Equations-From Linear to Fully Nonlinear Theory*, Springer, New York, 2017.
- [56] Zhou, J. *Viscosity solutions to second order path-dependent Hamilton-Jacobi-Bellman equations and applications*, *Ann. Appl. Probab.*, 33 (2023), 5564-5612.