# ON EFFICIENT INFERENCE OF CAUSAL EFFECTS WITH MULTIPLE MEDIATORS

By Haoyu Wei\*<sup>1,a</sup>, Hengrui Cai\*<sup>2,b\*</sup> Chengchun Shi<sup>3,c</sup> and Rui Song†<sup>4,d†</sup>

<sup>1</sup>Department of Economics, University of California, San Diego, <sup>a</sup>h8wei@ucsd.edu

This paper provides robust estimators and efficient inference of causal effects involving multiple interacting mediators. Most existing works either impose a linear model assumption among the mediators or are restricted to handle conditionally independent mediators given the exposure. To overcome these limitations, we define causal and individual mediation effects in a general setting, and employ a semiparametric framework to develop quadruply robust estimators for these causal effects. We further establish the asymptotic normality of the proposed estimators and prove their local semiparametric efficiencies. The proposed method is empirically validated via simulated and real datasets concerning psychiatric disorders in trauma survivors.

1. Introduction. Causal inference plays a crucial role in various fields, such as epidemiology (Hernán, 2004), medicine (Hernán et al., 2000), education (Card, 1999), and economics (Panizza and Presbitero, 2014). Within this spectrum, Pearl's causal graphical models (Pearl, 2000; Pearl et al., 2009) have recently emerged as a powerful tool for disentangling causal structures among variables (such as confounders, exposure, mediator(s), and outcome). Causal mediation analysis, a core method for examining causal graphical models, aims to reveal the causal mechanisms underlying observed effects from exposure to outcome through the mediator(s), to evaluate the effectiveness of the intervention, and to better understand the roles of mediators (see, for example, Pearl, 2012, 2014).

Existing statistical inferential tools for multiple mediators (see e.g., Robins and Greenland, 1992; Petersen et al., 2006; Imai et al., 2010; VanderWeele, 2015; Chakrabortty et al., 2018; Cai et al., 2020; Shi and Li, 2021) comprise the following three principal steps. Initially, causal structure learning methodologies (see e.g., Spirtes et al., 2000; Chickering, 2002; Nandy et al., 2018; Li et al., 2019; Yuan et al., 2019; Li et al., 2023) are applied to estimate the causal graph, often presented by a directed acyclic graph (DAG), using observational data. In the absence of additional assumptions (Shimizu et al., 2006; Neal, 2020), the graph is only identified up to a Markov equivalence class (MEC), and a completed partially directed acyclic graph (CPDAG) in such a class is often used to represent the graph

MSC2020 subject classifications: Primary 62A09, 62G05, 62G35.

Keywords and phrases: Causal graph, Mediation analysis, Multiply robust estimator, Statistical inference.

<sup>&</sup>lt;sup>2</sup>Department of Statistics, University of California, Irvine, <sup>b</sup>hengrc1@uci.edu

<sup>&</sup>lt;sup>3</sup>Department of Statistics, London School of Economics and Political Science, <sup>c</sup>c.shi7@lse.ac.uk

<sup>&</sup>lt;sup>4</sup>Department of Statistics, North Carolina State University, <sup>d</sup>songray@gmail.com

<sup>\*</sup>Equal contribution.

<sup>&</sup>lt;sup>†</sup>Corresponding author.

structure. The subsequent step is the estimation of the causal effects of mediators based on the DAG or CPDAG obtained from the initial phase. For this task, a variety of estimation techniques have been proposed, including the application of ordinary least squares (OLS) estimators (VanderWeele and Robinson, 2014; Lin and VanderWeele, 2017; Chakrabortty et al., 2018), parametric models (VanderWeele and Vansteelandt, 2014; VanderWeele et al., 2016; Chen et al., 2023), and nonparametric methods (An and VanderWeele, 2022; Brand et al., 2023). The final step is to conduct inferences based on the estimated effects, which often requires finding the exact (asymptotic) distributions of the estimators. As pointed out in Chen et al. (2023), such an inference is often regarded a separate task and has received less attention in recent causal graph literature.

Although most of the existing work on causal mediation inference is limited to scenarios with a single mediator (Tchetgen and Shpitser, 2012; Tchetgen Tchetgen, 2013; Kennedy et al., 2017; Wang and Tchetgen Tchetgen, 2018; Xia and Chan, 2023), there are some studies that employ the three main standard steps to conduct mediation analysis. However, all of them fall short of comprehensive. Theoretical challenges in unknown causal structures have led to current methods for multiple mediators inference being categorized mainly into three types. One approach assumes that multivariate mediators are conditionally independent given the treatment, or a set of transformed, conditionally independent variables, significantly simplifying the analysis (Preacher and Hayes, 2008; Boca et al., 2014; Zhang et al., 2016; Huang and Pan, 2016; Guo et al., 2023; Yuan and Qu, 2023). Another category, which does not impose this condition, relies on linear structural equation models (LSEMs) (Maathuis et al., 2009; Nandy et al., 2017, 2018; Chakrabortty et al., 2018; Zhao et al., 2022; Zhao and Luo, 2022). The last category allows for a general causal structure and correlated mediators, but uses approximations, such as assuming Gaussian conditional distributions under exposure (Daniel et al., 2015; Kim et al., 2019; Tai et al., 2022), or following a Probit/logistic model for odds ratios (VanderWeele and Vansteelandt, 2014; Steen et al., 2017; Park and Kürüm, 2018). However, these approaches present limitations for complex applications where the causal structure may not be correctly specified.

To bridge this significant gap in addressing potential model misspecification, we consider developing a semiparametric framework to infer causal effects, adapting the general causal structure. Extensive research exists on deriving double robust and highly efficient estimates of the total causal effect of exposure when the model is misspecified (Scharfstein et al., 1999; Bickel and Kwon, 2001; Bang and Robins, 2005). Complementary to this, multiple robust estimators have been developed to quantify direct and indirect effects (Goetgeluk et al., 2008; Tchetgen and Shpitser, 2012; Chan et al., 2016; Bhattacharya et al., 2022; Xia and Chan, 2023). A notable benefit of these multiple robust techniques is their integration of dimension reduction strategies with confounding adjustment, such that the estimators are consistent and asymptotically normal, provided that at least one of the strategies is correct (Van Der Laan and Rubin, 2006). These methods also achieve semiparametric efficiency when all included strategies are correct (Van Der Vaart et al., 1996; Bickel and Kwon, 2001; Bang and Robins, 2005). Despite considerable progress in the field, current multiple robust estimators are limited to only a single mediator. Hence, a new semiparametric inference is on demand for inferring causal effects involving multiple interacting mediators under (potentially) unknown causal graphs.

Our Contributions. We conclude our contributions with the following three folds.

- 1. Conceptually, we introduce the causal direct and indirect interventional effects for individual mediators (Definition 3.2 and Equation (3.1)). Our definitions expand upon those existing in various literature, accommodating a more general model setting. Specifically, it is applicable to both linear and non-linear models, thereby extending beyond existing literature such as Nandy et al. (2017); Chakrabortty et al. (2018); Cai et al. (2020). Moreover, our approach allows mediators to take a general value space, making it more flexible than the discrete settings as in Albert and Nelson (2011); Lin and VanderWeele (2017). Importantly, our definitions are consistent with the aforementioned literature when applied to the same settings. We further establish the identifiability results of the proposed definitions based on the estimated CPDAG from the data.
- 2. Methodological-wise, based on the proposed definitions, we firstly introduce the semiparametric framework concerning potential model misspecification under unknown
  graph structure for multiple interacted mediators (Theorem 4.3 and Corollary 4.4). Our
  analytical approach stands out for its novel insights into efficiency and robustness in
  the context of statistical inference of mediators on causal graphs. Specifically, we integrate four different estimating strategies to introduce new quadruply robust estimators
  for the causal effects of mediators. Additionally, we propose two algorithms to calculate
  these estimators together with the confidence intervals provided (Algorithm 1, Algorithm 2, and Proposition 6.1) to handle general noises and to increase computational
  speed, respectively. Under a semi-linear framework (Assumption 5.1), we derive concise parametric expressions for all proposed causal effects, and propose OLS estimators
  that can be computed using standard regressions, allowing for the direct acquisition of
  asymptotically valid confidence intervals simultaneously.
- 3. From a theoretical perspective, we prove the asymptotic properties of both our OLS estimators and quadruply robust estimators under mild conditions. Specifically: (i) Our OLS estimators are asymptotical normal with the analytical form of asymptotic variance provided, even under high-dimensional setting (Theorem 7.3 and Theorem 7.8); (ii) The introduced quadruply robust estimator is consistent to the true as long as at least some of the conditional densities or conditional expectations are correctly specified, even under a potently increasing function class. Moreover, if all the conditional densities and conditional expectations are correctly specified, and if converge at rates that are conservatively permissible by various machine learning approaches, these estimators can assuredly achieve  $n^{1/2}$ -consistency, asymptotic normality, and semiparametric efficiency (Theorem 7.11).

The rest of this paper is organized as follows: Section 2 presents preliminary concepts. Section 3 formally defines the direct and indirect causal effects of mediators. Section 4 outlines the semiparametric efficient scores for these causal effects. Section 5 explores the direct strategy for estimating the causal effects defined in Section 3, along with an OLS estimation procedure for semi-linear structures. Section 6 presents alternative estimation strategies, including the introduction of novel quadruply robust estimators. This section also provides both a general algorithm and, under specific conditions, a faster algorithm

for computing these estimators. Section 7 discusses the asymptotic properties of both the OLS and quadruply robust estimators. Section 8 presents various simulation results that validate the theories proposed for the estimators. In Section 9, an application of the proposed estimators is used to analyze real data collected from trauma survivors. The glossary of notations, all proofs, and additional technical materials are collected in the Appendix.

## 2. Preliminaries.

- 2.1. Graph Terminology. Consider a graph  $\mathcal{G}=(X,E)$  with a set of nodes X and a set of edges E. There is at most one edge between any pair of nodes. If there is an edge between  $X_i$  and  $X_j$ , then  $X_i$  and  $X_j$  are adjacent. The node  $X_i$  is said to be a parent of  $X_j$  if there is a directed edge from  $X_i$  to  $X_j$ . Let the set of all parents of node  $X_j$  in  $\mathcal{G}$  be  $\operatorname{Pa}(X_j) = \operatorname{Pa}_{\mathcal{G}}(X_j) = \operatorname{Pa}_{X_j}(\mathcal{G})$ , and all adjacent nodes of  $X_j$  in  $\mathcal{G}$  by  $\operatorname{adj}(X_j) = \operatorname{adj}_{\mathcal{G}}(X_j)$ . A path from  $X_i$  to  $X_j$  in  $\mathcal{G}$  is a sequence of distinct vertices,  $\pi := \{a_0, a_1, \cdots, a_L\} \subset V$  such that  $a_0 = X_i$ , and  $a_L = X_j$ . A directed path from  $X_i$  to  $X_j$  is a path between  $X_i$  and  $X_j$  where all edges are directed towards  $X_j$ . A directed cycle is formed by the directed path from  $X_i$  to  $X_j$  together with the directed edge  $X_j$  to  $X_i$ . A directed graph that does not contain directed cycles is called a directed acyclic graph (DAG). A directed graph is acyclic if and only if it has a topological ordering.
- 2.2. Causal Graph Structural Assumption. Let A be a **binary** exposure/treatment in  $\{0,1\}$ ,  $M:=(M_1,M_2,\cdots,M_p)^{\top}\in\mathbb{R}^p$  be mediators with dimension p in its support  $\mathcal{M}=\mathcal{M}_1\times\cdots\times\mathcal{M}_p\subseteq\mathbb{R}^p$ , and  $Y\in\mathbb{R}$  be the outcome of interest. Additionally, we also consider that there are t-1 confounders  $C:=(C_1,\ldots,C_{t-1})^{\top}\in\mathbb{R}^{t-1}$  in its support  $C\subseteq\mathbb{R}^{t-1}$ . We would just let t=1 here to represent the absence of confounders, that is  $C=\varnothing$ . Suppose that there exists a DAG  $\mathcal{G}=(X,E)$  that characterizes the causal relationship among  $X=(C^{\top},A,M^{\top},Y)^{\top}$ , where the dimension of X is d=t+p+1. We suppose we observe i.i.d data on  $X=(C^{\top},A,M^{\top},Y)^{\top}$  is collected for n subjects. To characterize our model, we consider the following assumptions.

ASSUMPTION 2.1. The causal graph  $\mathcal{G}$  satisfies Causal Markov Condition, Causal Faithfulness Condition, and Causal Sufficiency (Hasan et al., 2023). The random vector X satisfies the structure assumption: (i) No potential mediator is a direct cause of confounders C; (ii) The outcome Y has no descendant; (iii) The only parents of treatment A are confounders.

In many instances, the accessible data offers an incomplete view of the inherent causal structure. To address this gap, *Causal Markov Condition*, *Causal Faithfulness Condition*, and *Causal Sufficiency* in the above assumption provide a sufficient condition for causal discovery in i.i.d. data contexts (Lee and Honavar, 2020; Assaad et al., 2022; Hasan et al., 2023). The rigorous definitions for them and related details can be found in Section 2.4 in Hasan et al. (2023). Furthermore, our structural assumptions aim at ensuring the identifiability of the causal model, which are similar to Consistency Assumption and Sequential Ignorability Assumption in Tchetgen and Shpitser (2012), and the structure assumptions in Section 2.4 of Chakrabortty et al. (2018).

- 2.3. Markov Equivalence Class. A general causal DAG, G, may not be identifiable from the distribution of X. According to Pearl (2000), a DAG only encodes conditional independence relationships through the concept of d-separation. In general, several DAGs can encode the same conditional independence relationships, and such DAGs form a Markov equivalence class. Two DAGs belong to the same Markov equivalence class if and only if they have the same skeleton and the same v-structures (Kalisch and Bühlmann, 2007). A Markov equivalence class of DAGs can be uniquely represented by a completed partially directed acyclic graph (CPDAG) (Spirtes et al., 2000), which is a graph that can contain both directed and undirected edges. A CPDAG satisfies the following:  $X_i \leftrightarrow X_j$  in the CPDAG if the Markov equivalence class contains a DAG including  $X_i \to X_j$ , as well as another DAG including  $X_i \to X_i$ . CPDAGs can be estimated from observational data using various algorithms, such as the algorithms in Kalisch and Bühlmann (2007), Harris and Drton (2013), and Zhang and Bareinboim (2018). The Markov equivalence class for a fixed CPDAG  $\mathcal{C}$  is denoted by  $MEC(\mathcal{C})$ , which is a set containing all DAGs  $\mathcal{G}$  that have the CPDAG structure C. If we can obtain the true DAG from the data, we can simply treat it as a special case of the "MEC" containing only this DAG, i.e.,  $MEC(\mathcal{G}) = \{\mathcal{G}\}$ . For simplicity, we denote the corresponding causal structure for the mediators M as  $\mathcal{G}_M$ , which can be obtained by deleting nodes C, A, Y, and the corresponding edges from  $\mathcal{G}$ . The CPDAG of mediators is similarly denoted as  $\mathcal{C}_M$ . For simplicity and with a minor stretch of notation, we employ  $\mathcal{G}_M$  and  $\mathcal{C}_M$  to denote the causal DAG and CPDAG of X, respectively, such that their corresponding mediators' causal DAG and CPDAG are represented by  $\mathcal{G}_M$  and  $\mathcal{C}_M$  exactly.
- 3. Definition of Causal Effects. In this section, we will formally give our refined definition of the causal effects of mediators. To begin with, we give the total effect TE, the natural direct effect that is not mediated by mediators DE, and the natural indirect effect that is regulated by mediators IE defined in Pearl et al. (2009).

DEFINITION 3.1 (Pearl et al. (2009)). Natural effects are defined as follows:

$$\begin{split} TE &= \mathbf{E}\big[Y\mid do(A=1)\big] - \mathbf{E}\big[Y\mid do(A=0)\big],\\ DE &= \mathbf{E}\Big[\mathbf{E}\big[Y\mid do(A=1,M=M^{(0)})\big]\Big] - \mathbf{E}\big[Y\mid do(A=0)\big],\\ IE &= \mathbf{E}\Big[\mathbf{E}\big[Y\mid do(A=0,M=M^{(1)})\big]\Big] - \mathbf{E}\big[Y\mid do(A=0)\big]. \end{split}$$

In the above definition,  $do(A=0)=do_{\mathcal{G}}(A=0)$  is a mathematical operator to simulate physical interventions that hold A constant as 0 while keeping the rest of the model unchanged, which corresponds to remove edges into A and replace A by the constant 0 in the original causal graph  $\mathcal{G}$ . Here,  $M^{(0)}$  is the (random) value of M if setting do(A=0), and  $M^{(1)}$  is the (random) value of M if setting do(A=1). One can refer to Pearl et al. (2009) for more details of 'do-operator'. The expectation  $E[\cdot]$  is an abbreviation of  $E_P[\cdot]$  with  $P=P_X$  is the law of X under  $\mathcal{G}$ . Inspired by the above definition, we can give the definition of the causal effects for an individual mediator.

DEFINITION 3.2. Let  $TM_j(\mathcal{G}_M)$  represent total individual mediation effects via an individual mediator  $M_j$  defined as

$$\begin{split} TM_j(\mathcal{G}_M) := & \left\{ \mathbf{E} \big[ Y \mid do(A=1) \ \big] - \mathbf{E} \Big[ \mathbf{E} \big[ Y \mid do_{\mathcal{G}_M}(A=1,M_j) \big] \Big] \right\} \\ & - \left\{ \mathbf{E} \big[ Y \mid do(A=0) \big] - \mathbf{E} \Big[ \mathbf{E} \big[ Y \mid do_{\mathcal{G}_M}(A=0,M_j) \big] \Big] \right\}, \end{split}$$

under any fixed mediators' causal structure  $\mathcal{G}_M$ . Let  $DM_j$  denote the direct interventional effect via an individual mediator  $M_j$  defined as

$$\begin{split} DM_j := \mathbf{E} \Bigg[ \int_{\mathcal{M}} \mathbf{E} \big[ Y \mid C, A &= 1, M = m \big] f_{M_{-j} \mid C, A} (m_{-j} \mid C, A = 0) \\ & \times \big\{ f_{M_j \mid C, A} (m_j \mid C, A = 1) - f_{M_j \mid C, A} (m_j \mid C, A = 0) \big\} \, \mathrm{d}m \Bigg], \end{split}$$

where  $f_{\cdot|\cdot}(\cdot|\cdot)$  are the conditional density (or mass) functions. Then the indirect interventional effect for  $M_j$  under  $\mathcal{G}_M$  is defined as  $IM_j(\mathcal{G}_M) := TM_j(\mathcal{G}_M) - DM_j$ .

The Definition 3.2 serves important meanings when we are concerned with different impacts of mediators. Note that  $TM_j(\mathcal{G}_M)$  in our definition is an extension for the individual mediation effect in Chakrabortty et al. (2018) for LSEMs, denoted as

$$\eta_{j}(\mathcal{G}_{M}) = \frac{\partial}{\partial a} \mathbb{E}\left[Y \mid do(A = a)\right] - \frac{\partial}{\partial a} \mathbb{E}\left[Y \mid do_{\mathcal{G}_{M}}(A = a, M_{j} = m_{j})\right],$$

which can be interpreted as the change in the total causal effect of the exposure A on the response Y when the potential mediator  $M_j$  is removed from the causal graph  $\mathcal G$  through the intervention  $do(M_j=m_j)$ . But under the non-linearity assumption with binary exposure, the above  $\eta_j$  will be a function of j-th mediator  $m_j$  (Remark 2.1 in Chakrabortty et al. (2018)). Therefore, for solving this problem, we take integral with respect to the density  $f_{M_j}(m_j)$  for  $M_j$ , i.e.

$$\begin{split} &\int_{\mathcal{M}_{j}} \left[ \frac{\partial}{\partial a} \mathbf{E} \big[ Y \mid do(A = a) \big] - \frac{\partial}{\partial a} \mathbf{E} \big[ Y \mid do_{\mathcal{G}_{M}}(A = a, M_{j} = m_{j}) \big] \right] f_{M_{j}}(m_{j}) \, \mathrm{d}m_{j} \\ &= \frac{\partial}{\partial a} \mathbf{E} \big[ Y \mid do(A = a) \big] - \mathbf{E} \left[ \frac{\partial}{\partial a} \mathbf{E} \big[ Y \mid do_{\mathcal{G}_{M}}(A = a, M_{j}) \big] \right]. \end{split}$$

Then by  $a \in \{0,1\}$ , we get the expression of  $TM_j(\mathcal{G}_M)$  in Definition 3.2. Our definition of  $DM_j$  is a straightforward extension of Equation (6) in Vansteelandt and Daniel (2017) by replacing summation with integral. The introduction of  $DM_j$  and  $IM_j(\mathcal{G}_M) = TM_j(\mathcal{G}_M) - DM_j$  in the above definition is driven by the need for an orthogonal decomposition of the total causal effect of mediators, a concept crucial for unraveling the intricate relationships among variables in mediation analysis, as argued in Cai et al. (2020). Here  $DM_j$  can be interpreted as the causal effect through a particular mediator from the exposure to the outcome, i.e.,  $\mathrm{E}\big[Y\mid C,A=1,M=m\big]\big\{f_{M_j\mid C,A}(m_j\mid C,A=1)-f_{M_j\mid C,A}(m_j\mid C,A=1)-f_{M_j\mid$ 

C, A=0), that is not regulated by any other mediators, i.e.,  $f_{M_{-j}|C,A}(m_{-j} \mid C, A=0)$ , and thus not regulated by its descendant mediators. Then  $IM_j(\mathcal{G}_M) = TM_j(\mathcal{G}_M) - DM_j$  captures the indirect effect of the particular mediator  $M_j$  on the outcome Y regulated by descendant mediators.

Definition 3.2 is generally defined for any causal graph with binary exposure A. In the context of particular linear causal structures, these definitions transform into concise parametric expressions, providing more intuitive 'do' representations and aligning consistently with existing literature. A comprehensive discussion on this can be found in Section 5.1.

Usually, we do not know the true structure of mediators  $\mathcal{G}_M$  and we can only estimate its corresponding CPDAG  $\mathcal{C}_M$  (Maathuis et al., 2009). If the number of  $\mathrm{MEC}(\mathcal{C}_M)$  is larger than one,  $TM_j$  and  $IM_j$  based on elements  $\mathrm{MEC}(\mathcal{C}_M)$  will be not unique. As a result, we define an identifiable version of  $TM_j$  based on a CPDAG as the average over  $\mathrm{MEC}(\mathcal{C}_{0,M})$ . Specifically,

(3.1) 
$$\overline{TM}_j := \frac{1}{\# \operatorname{MEC}(\mathcal{C}_M)} \sum_{\mathcal{G}_M \in \operatorname{MEC}(\mathcal{C}_M)} TM_j(\mathcal{G}_M).$$

The corresponding identifiable indirect interventional effect of mediator  $M_j$  is  $\overline{IM}_j := \overline{TM}_j - DM_j$  for  $j \in [p]$ .

**4. Semiparametric Efficient Scores.** We start with exploring the definition in Section 3. Define propensity score  $e_{a'}(x_S) := P(A = a' \mid X_S = x_S)$  for  $a' \in \{0,1\}$ , the outcome mean  $\mu(x_S) := \mathrm{E}\big[Y \mid X_S = x_S\big]$ , and the conditional density  $\pi_{x_S}(m_T) := f_{M_T \mid X_S}(m_T \mid X_S = x_S)$  for any subset  $T \subseteq [p]$  and  $S \subseteq [t+p+1]$ . Suppose all these functions belong to  $\ell^2$ -class. Note that in our notation,  $x_S$  and  $m_T$  can represent vectors. At times, we may abbreviate  $\mu\big((x_{S_1}^\top, x_{S_2}^\top, \dots, x_{S_N}^\top)^\top\big)$  as  $\mu(x_{S_1}, x_{S_2}, \dots, x_{S_N})$  and  $\pi_{(x_{S_1}^\top, x_{S_2}^\top, \dots, x_{S_N}^\top)^\top}(m_T)$  as  $\pi_{x_{S_1}, x_{S_2}, \dots, x_{S_N}}(m_T)$  when  $x_S$  is the concatenation of these vectors, i.e.,  $x_S = (x_{S_1}^\top, x_{S_2}^\top, \dots, x_{S_N}^\top)^\top$ . Let  $\mathrm{Pa}_j(\mathcal{G}_M) = \mathrm{Pa}_{\mathcal{G}_M}(M_j) \subseteq \{M_1, \dots, M_{j-1}, M_{j+1}, \dots, M_p\}$  denote the *parent mediators* of  $M_j$  and  $\mathrm{pa}_j(\mathcal{G}_M)$  as its realization. Denote

(4.1) 
$$\kappa(a', C) := E[Y \mid C, A = a'],$$

(4.2) 
$$\zeta_j(a', a, C) := \int_{\mathcal{M}} \mu(C, a^*, m) \pi_{C, a'}(m_j) \pi_{C, a}(m_{-j}) dm,$$

and

$$(4.3) \qquad \varrho_{j}(a',M_{j},C\,;\,\mathcal{G}_{M}) := \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \mu(C,a',\mathrm{pa}_{j},M_{j})\,\pi_{C,a'}(\mathrm{pa}_{j})\,\mathrm{d}\,\mathrm{pa}_{j},$$

for a fixed causal graph  $\mathcal{G}_M$ , where  $\mathcal{M}_{\mathrm{pa}_j}(\mathcal{G}_M)$  is the support of  $\mathrm{Pa}_j(\mathcal{G}_M)$  and  $a^*=0$  is the reference level of the exposure. It is worth noting that all three quantities above are random due to the randomness in C (and  $M_j$ ). Given that the exposure features two levels, 0 and 1, for simplicity, we use the notation  $\langle \cdot \rangle$  to signify the difference evaluated at these two levels. Specifically, let us define

$$\langle g_{\cdot,u}(\cdot,v)\rangle := g_{1,u}(1,v) - g_{0,u}(0,v),$$

for any function  $g_{\cdot,u}(\cdot,v)$ , where u and v are arbitrary parameters. Then we can have the following theorem to characterize the relationship between the interventional effects of mediators as specified in Definition 3.2 and the quantities defined above.

THEOREM 4.1. Suppose Assumption 2.1 holds, then for any  $j \in [p]$ ,

$$DM_j = \mathbb{E}[\langle \zeta_j(\cdot, 0, C) \rangle],$$

and for any fixed  $G_M$ 

$$TM_j(\mathcal{G}_M) = \mathbb{E}[\langle \kappa(\cdot, C) - \varrho_j(\cdot, M_j, C; \mathcal{G}_M) \rangle].$$

Consider  $\mathcal{M}_{nonpar}$  as the full model where the observed data likelihood is not constrained, encompassing all conventional laws  $P_X$  or, equivalently, distribution  $F_X$  of the observed data X. The aforementioned theorem establishes that the causal effects detailed in Section 3 can be represented as regular expectations. Consequently, they function as mappings from  $\mathcal{M}_{nonpar}$  to the real line  $\mathbb{R}$ . We assume  $P_X$  satisfies the positivity assumption given below.

ASSUMPTION 4.2. There exists a  $\varepsilon > 0$  such that for any  $c \in \mathcal{C}$ ,  $a' \in \{0,1\}$ , and  $m \in \mathcal{M}$ ,

$$\varepsilon < e_{a'}(c) < 1 - \varepsilon$$
 and  $\varepsilon < \pi_{c,a',m_i}(m_{-j}) < \infty$ ,

with probability one.

The efficient scores for the functionals DE and IE have been studied in various literature (Tchetgen and Shpitser, 2012; Tchetgen Tchetgen, 2013; Shi et al., 2020). The explicit expressions and detailed analysis can be seen in Theorem 1 of Tchetgen and Shpitser (2012). For finding the efficient scores for the functionals  $DM_i$  and  $IM_i$ , we denote

(4.5) 
$$\tau_{:S}(C, a', M_T) := \int_{\mathcal{M}_S} \mu(C, a', m_S, M_T) \pi_{:S}(m_S) \, \mathrm{d}m_S$$

for any  $a' \in \{0,1\}$  and  $T,S \subseteq [p]$ . Then we can derive the efficient score for  $\kappa(a',C)$ ,  $\zeta_j(a',a,C)$ , and  $\varrho_j(a',M_j,C\,;\mathcal{G}_M)$  on  $\mathscr{M}_{\text{nonpar}}$  in the following theorem.

THEOREM 4.3. Suppose the Assumption 2.1 and 4.2 hold, we have the efficient scores for  $E\kappa(a',C)$ ,  $E\zeta_j(a',0,C)$ , and  $E\varrho_j(a',M_j,C;\mathcal{G}_M)$  as

$$S^{eff,nonpar}\big(\mathrm{E}\kappa(a',C)\big) = \frac{\mathbb{1}(A=a')}{e_{a'}(C)}\Big\{Y - \kappa(a',C)\Big\} + \kappa(a',C) - \mathrm{E}\kappa(a',C),$$

$$S^{eff,nonpar}\left(\mathrm{E}\zeta_{j}(a',0,C)\right) = \frac{\mathbb{1}(A=1)}{e_{1}(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_{j}}(M_{-j})} \left[Y - \mu(C,1,M)\right]$$

$$+ \frac{\mathbb{1}(A=0)}{e_{0}(C)} \left[\tau_{C,a';j}(C,1,M_{-j}) - \zeta_{j}(a',0,C)\right] + \frac{\mathbb{1}(A=a')}{e_{a'}(C)} \left[\tau_{C,0;-j}(C,1,M_{j}) - \zeta_{j}(a',0,C)\right]$$

$$+ \zeta_{j}(a',0,C) - \mathrm{E}\zeta_{j}(a',0,C),$$

and

$$S^{eff,nonpar}\left(\operatorname{E}\varrho_{j}(a',M_{j},C\,;\mathcal{G}_{M})\right) = \frac{\mathbb{1}(A=a')}{e_{a'}(C)}\pi_{C,a'}\left(\operatorname{Pa}_{j}(\mathcal{G}_{M})\right)\left[Y-\mu(C,a',M)\right] \\ + \frac{\mathbb{1}(A=a')}{e_{a'}(C)}\left[\tau_{C\,;j}(C,a',\operatorname{Pa}_{j}(\mathcal{G}_{M}))-\operatorname{E}\left[\varrho_{j}(a',M_{j},C\,;\mathcal{G}_{M})\mid C\right]\right] \\ + \varrho_{j}(a',M_{j},C\,;\mathcal{G}_{M})-\operatorname{E}\left[\varrho_{j}(a',M_{j},C\,;\mathcal{G}_{M})\mid C\right]+\operatorname{E}\left[\varrho_{j}(a',M_{j},C\,;\mathcal{G}_{M})\mid C\right]-\operatorname{E}\varrho_{j}(a',M_{j},C\,;\mathcal{G}_{M})$$
under model. (6) for any  $i\in[n]$  of  $i\in[n]$  and fixed  $C$  is where

under model 
$$\mathcal{M}_{nonpar}$$
 for any  $j \in [p]$ ,  $a' \in \{0,1\}$ , and fixed  $\mathcal{G}_M$ , where

$$E[\varrho_{j}(a', M_{j}, C; \mathcal{G}_{M}) \mid C]$$

$$= \int_{\mathcal{M}_{pa_{j}}(\mathcal{G}_{M}) \cup \mathcal{M}_{j}} \mu(C, a', pa_{j}(\mathcal{G}_{M}), m_{j}) \pi_{C, a'}(pa_{j}(\mathcal{G}_{M})) \pi_{C}(m_{j}) d pa_{j}(\mathcal{G}_{M}) d m_{j}.$$

Here, we let  $\pi_{C,a'}(\operatorname{pa}_i(\mathcal{G}_M)) \equiv 1$  if  $\operatorname{Pa}_i(\mathcal{G}_M) = \emptyset$ .

In Theorem 4.3, we retain the final two terms in  $S^{\text{eff,nonpar}}(\mathbb{E}\varrho_j(a',M_j,C,;,\mathcal{G}_M))$ , because we want to express  $S^{\text{eff,nonpar}}(\mathbb{E}\varrho_j(a',M_j,C;\mathcal{G}_M))$  is composed of four parts. This result can be then combined with Theorem 4.1 to obtain the efficient scores for  $DM_j$  and  $IM_j(\mathcal{G}_M)$ , thus  $TM_j(\mathcal{G}_M) = DM_j + IM_j(\mathcal{G}_M)$ .

COROLLARY 4.4. Suppose the conditions in Theorem 4.3 holds, then we have

$$S^{\textit{eff},nonpar}(TE) = \Big\langle S^{\textit{eff},nonpar}\big(\mathbb{E}\kappa(\boldsymbol{\cdot},C)\big)\Big\rangle,$$

$$S^{\textit{eff},nonpar}(DM_j) = \Big\langle S^{\textit{eff},nonpar}\big(\mathbb{E}\zeta_j(\boldsymbol{\cdot},0,C)\big)\Big\rangle,$$

for any  $j \in [p]$ . Furthermore, for any fixed  $\mathcal{G}_M$ , we have

$$S^{\textit{eff},nonpar}\big(TM_{j}(\mathcal{G}_{M})\big) = S^{\textit{eff},nonpar}(TE) - \left\langle S^{\textit{eff},nonpar}\big(\mathrm{E}\varrho_{j}(\cdot,M_{j},C\,;\,\mathcal{G}_{M})\big)\right\rangle$$
 and  $S^{\textit{eff},nonpar}\big(IM_{j}(\mathcal{G}_{M})\big) = S^{\textit{eff},nonpar}\big(TM_{j}(\mathcal{G}_{M})\big) - S^{\textit{eff},nonpar}(DM_{j})$  for any  $j \in [p]$ .

In Corollary 4.4, the explicit formulas for  $S^{\text{eff,nonpar}}(DM_j)$ ,  $S^{\text{eff,nonpar}}(TM_j(\mathcal{G}_M))$ , and  $S^{\text{eff,nonpar}}(IM_j(\mathcal{G}_M))$  can be directly derived by substituting the relevant expressions from Theorem 4.3. Consequently, the semiparametric efficiency bounds for estimating  $DM_j$ ,  $IM_j(\mathcal{G}_M)$ , and  $TM_j(\mathcal{G}_M)$  within the full nonparametric model  $\mathscr{M}_{\text{nonpar}}$  are respectively  $\mathrm{E}\big[S^{\text{eff,nonpar}}(DM_j)\big]^2$ ,  $\mathrm{E}\big[S^{\text{eff,nonpar}}(IM_j(\mathcal{G}_M))\big]^2$ , and  $\mathrm{E}\big[S^{\text{eff,nonpar}}(TM_j(\mathcal{G}_M))\big]^2$  for any specified  $\mathcal{G}_M$ , all with clearly delineated forms. The asymptotic variances of any regular asymptotic linear estimators in  $\mathscr{M}_{\text{nonpar}}$  must be greater than or equal to these bounds. Given that  $TM_j(\mathcal{G}_M) = DM_j + IM_j(\mathcal{G}_M)$ , we will only focus on  $DM_j$  and  $IM_j(\mathcal{G}_M)$  in the subsequent sections.

**5. Direct Strategy and Ordinary Least Squares (OLS) Estimations.** An important implication of Corollary 4.4 is that all regular and asymptotically linear (RAL) estimators of  $DM_j$  and  $IM_j(\mathcal{G}_M)$  in the model  $\mathcal{M}_{nonpar}$  share the common score  $S^{\text{eff, nonpar}}(DM_j)$  and  $S^{\text{eff, nonpar}}(IM_j(\mathcal{G}_M))$ , respectively. For illustrating this and as a motivation for multiply robust estimation when nonparametric methods are not appropriate, we provide a detailed study of different estimating strategies in this section and the next section.

Theorem 4.1 gives an explicit expression for  $DM_j$  and  $IM_j(\mathcal{G}_M)$ , we can correspondingly give their estimators by (i) replacing the unknown quantities  $\kappa(\cdot,\cdot)$ ,  $\zeta_j(\cdot,\cdot,\cdot)$ ,  $\varrho_j(\cdot,\cdot,\cdot;\mathcal{G}_M)$  with their estimators and then (ii) replacing  $\mathrm{E}[\cdot]$  by  $\mathbb{P}_n[\cdot] = n^{-1} \sum_{i=1}^n [\cdot]_i$  directly. To be specific, in the step (i), we construct the following estimators:

$$\widehat{\kappa}^{\mathcal{M}_0}(a',c) := \widehat{\mu}(c,a'),$$

$$\widehat{\zeta}_j^{\mathcal{M}_0}(a',0,c) := \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \widehat{\pi}_{C,a'}(m_j) \widehat{\pi}_{C,0}(m_{-j}) \,\mathrm{d}m,$$

and

$$\widehat{\varrho}_{j}^{\mathcal{M}_{0}}(a', m_{j}, c; \mathcal{G}_{M}) := \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \widehat{\mu}(C, a', \mathrm{pa}_{j}(\mathcal{G}_{M}), M_{j}) \widehat{\pi}_{C, a'}(\mathrm{pa}_{j}(\mathcal{G}_{M})) \, \mathrm{d}\, \mathrm{pa}_{j}(\mathcal{G}_{M}),$$

which are consistent for  $\kappa(a',C)$ ,  $\zeta_j(a',0,c)$  and  $\varrho_j(a',m_j,c)$  for any  $c \in \mathcal{C}$ ,  $a' \in \{0,1\}$ , and  $m \in \mathcal{M}$ . Note that  $\kappa(a',C)$  can be written as

$$E[Y \mid C, A = a'] = \int_{\mathcal{M}} E[Y \mid C, A = a', M = m] f_{M \mid C, A}(m \mid C, A = a') dm.$$

Therefore, the consistency of  $\widehat{\kappa}^{\mathscr{M}_0}(a',c)$ ,  $\widehat{\zeta}_j^{\mathscr{M}_0}(a',0,c)$ , and  $\widehat{\varrho}_j^{\mathscr{M}_0}(a',m_j,c;\mathcal{G}_M)$  will use the correctly specified information as follows:

•  $\mathcal{M}_0$ : the conditional expectation  $\mathrm{E}[Y \mid C = \cdot, A = \cdot, M = \cdot]$  and the conditional density of the mediator  $f_{M \mid C, A}(m \mid C = \cdot, A = \cdot)$  are correctly specified.

Then in the step (ii), we can construct the estimators

$$\widehat{DM}_{j}^{\mathcal{M}_{0}} = \mathbb{P}_{n} \left[ \left\langle \widehat{\zeta}_{j}^{\mathcal{M}_{0}}(\boldsymbol{\cdot}, 0, C) \right\rangle \right],$$

$$\widehat{TM}_{j}^{\mathcal{M}_{0}}(\mathcal{G}_{M}) = \mathbb{P}_{n} \left[ \left\langle \widehat{\kappa}^{\mathcal{M}_{0}}(\cdot, C) - \widehat{\varrho}_{j}^{\mathcal{M}_{0}}(\cdot, M_{j}, C; \mathcal{G}_{M}) \right\rangle \right]$$

and  $\widehat{IM}_j^{\mathcal{M}_0}(\mathcal{G}_M)=\widehat{TM}_j^{\mathcal{M}_0}(\mathcal{G}_M)-\widehat{DM}_j^{\mathcal{M}_0}$ . Next, the estimators for the identifiable  $\overline{TM}_j$  and  $\overline{IM}_j$  are

$$\widehat{TM}_{j}^{\text{avg},\mathcal{M}_{0}} = \frac{1}{\# \operatorname{MEC}(\widehat{\mathcal{C}}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\widehat{\mathcal{C}}_{M})} \widehat{TM}_{j}^{\mathcal{M}_{0}}(\mathcal{G}_{M}),$$

and  $\widehat{IM}_j^{\text{avg},\mathcal{M}_0} = \widehat{TM}_j^{\text{avg},\mathcal{M}_0} - \widehat{DM}_j^{\mathcal{M}_0}$ , where  $\widehat{\mathcal{C}}_M \in \{0,1\}^{p \times p}$  is the adjacency matrix of the estimated CPDAG for the mediators  $M = (M_1,\dots,M_p)^{\top} \in \mathbb{R}^p$ . The consistent causal

structure  $\widehat{\mathcal{C}}_M$  can be obtained by after we obtain the estimated adjacency matrix  $\widehat{\mathcal{C}}$  for the whole causal graph, and we then extract a subset  $\widehat{\mathcal{C}}_M = [\widehat{\mathcal{C}}_{kk}]_{k \in t+1,\dots,t+p}$  to arrive at the causal structure for the mediators. The estimated adjacency matrix  $\widehat{\mathcal{C}}$  can be achieved through methods such as the PC algorithm (Spirtes et al., 2000), greedy equivalence search (GES) (Chickering, 2002), and adaptively restricted greedy equivalence search (ARGES) (Nandy et al., 2018), among others.

5.1. OLS estimator under semi-linear model. The direct strategy in model  $\mathcal{M}_0$  involves two unknown quantities:  $\mathrm{E}[Y \mid C = \cdot, A = \cdot, M = \cdot]$  and  $f_{M\mid C,A}(m \mid C = \cdot, A = \cdot)$ . Specially, when we have known that the structure follows  $Y \leftarrow h_Y(C,A,M) + \epsilon_Y$  and  $M \leftarrow h_M(C,A,M) + \epsilon_M^{-1}$  with given functions  $h_Y(\cdot)$  and  $h_M(\cdot)$ , and given that the error terms  $\epsilon_Y$  and  $\epsilon_M$  belong to some classes of distributions,  $\mathcal{M}_0$  will be correctly recovered. A commonly used approach for this is assuming Linear Structural Equation Models (LSEMs), i.e.,  $X \leftarrow B^\top X + \epsilon$  with mean-zero and jointly independent error vector  $\epsilon$ , where  $B = (b_{ij})_{1 \le i \le d, 1 \le j \le d}$  be a  $d \times d$  matrix, where  $b_{ij}$  is the weight of the edge  $X_i \to X_j \in E$ , and  $b_{ij} = 0$  otherwise. There are numerous rigorous theoretical findings for LSEMs, as discussed in (Chakrabortty et al., 2018; Cai et al., 2020; Shi and Li, 2022). However, LSEMs are not applicable when dealing with binary exposure, given that the element A in X is constrained to either 0 or 1. In lieu of LSEMs, we propose the following semi-linear structure assumption.

ASSUMPTION 5.1. We assume X is **semi-linear** when it is generated as follows

(5.1) 
$$A \leftarrow h(C, \epsilon_A),$$

$$M \leftarrow B_{MC}^{\top} C + \beta_{MA} A + B_{MM}^{\top} M + \epsilon_M,$$

$$Y \leftarrow \beta_{YC}^{\top} C + \alpha_{YA} A + \beta_{YM}^{\top} M + \epsilon_Y,$$

where  $h: \mathbb{R}^{t-1} \times \mathbb{R} \to \{0,1\}$  is a known link function and  $\epsilon_A, \epsilon_M, \epsilon_Y$  are mean-zero error terms independent with each other as well as C.

Through this paper,  $\alpha$ ,  $\beta$ , and B will always represent a scalar, vector, and matrix, respectively. Define  $\theta_{MA} := \left[ (I_p - B_{MM}^\top)^{-1} \beta_{MA} \right]$  where  $I_p \in \mathbb{R}^{p \times p}$  is the identity matrix, then under the above semi-linear structural assumptions, we have the following propositions for the uniqueness of  $\theta_{MA}$  under MEC, interpretation displays, and neat parametric expressions for the causal effects defined in Section 3.

PROPOSITION 5.2 (Identification). Under Assumption 5.1,  $\theta_{MA}$  is unique in any fixed  $MEC(\mathcal{C}_M)$ . Hence, DE, IE, and  $DM_j$  are also unique in  $MEC(\mathcal{C}_M)$ .

 $<sup>^{1}</sup>$ The symbol  $\leftarrow$  emphasizes that the expressions should be understood as a generating mechanism rather than as a mere equation.

PROPOSITION 5.3 (Interpretation). Under Assumption 2.1 and Assumption 5.1, for any fixed  $\mathcal{G}_M$ , we have

$$DM_{j} = \left\{ \mathbf{E} \left[ Y \mid do(A = 0, M_{j} = m_{j}^{(0)} + 1, M_{-j} = m_{-j}^{(0)}) \right] - \mathbf{E} \left[ Y \mid do(A = 0) \right] \right\} \times \Delta_{j}^{*},$$

and

$$\begin{split} IM_{j}(\mathcal{G}_{M}) &= \left\{ \mathbf{E} \big[ Y \mid do(A=0, M_{j}=m_{j}^{(0)}+1) \big] \\ &- \mathbf{E} \big[ Y \mid do(A=0, M_{j}=m_{j}^{(0)}+1, M_{-j}=m_{-j}^{(0)}) \big] \right\} \times \Delta_{j}^{*}, \end{split}$$

where  $\Delta_j^* = \langle E[M_j \mid do(A = \cdot)] \rangle$ .

PROPOSITION 5.4. *Under Assumption 2.1 and Assumption 5.1, we have (i):* 

$$DE = \alpha_{YA}, \qquad IE = \theta_{MA}^{\top} \beta_{MA}$$

and hence  $TE = \alpha_{YA} + \theta_{MA}^{\top} \beta_{MA}$  for natural effects, and

$$DM_j = \beta_{YM,j}\theta_{MA,j}.$$

(ii) For any fixed  $\mathcal{G}_M$ ,

$$TM_j = \beta_{YM}^{\top} \theta_{MA} - \beta_{YM_{-i}}^{\top} \theta_{M_{-i}A},$$

where  $\theta_{M_{-j}A} := (I - B_{M_{-j}M_{-j}}^{\top})^{-1}\beta_{M_{-j}A}$ , and hence

$$IM_j = \beta_{YM,j}^{\top}(\theta_{MA,-j} - \theta_{M_{-j}A}).$$

(ii') For any fixed  $\mathcal{G}_M$ , we have the alternative expression for  $TM_i$  as

$$TM_j = \theta_{MA,j} \times \mathrm{E}^{reg} [Y \mid M_j \cup \mathrm{Pa}_j(\mathcal{G}_M) \cup A \cup C]_1,$$

hence

$$IM_{j} = \theta_{MA,j} \times \Big( \mathbb{E}^{reg} \big[ Y \mid M_{j} \cup \operatorname{Pa}_{j}(\mathcal{G}_{M}) \cup A \cup C \big]_{1} - \beta_{YM,j} \Big),$$

where  $E^{reg}[X_k \mid X_l \cup X_S]_1$  denotes the true coefficient of  $X_l$  in the linear regression of  $X_k$  on the combined set  $X_l \cup X_S$ .

Proposition 5.2 gives the fact that only  $IM_j$  and  $TM_j$  require specific DAG structure, while other quantities do not require any knowledge of the causal structure under semi-linear assumption. Meanwhile, Proposition 5.3 implies that, under the semi-linear assumption, our definitions for direct/indirect individual mediation effects in Definition 3.2 exactly coincides with the definitions in Cai et al. (2020): the first multiplier is in Proposition 5.3 with the classical meaning of 'natural' in the causal inference literature Pearl (2000). Thus,  $DM_j$  can be interpreted as the causal effect through a particular mediator from the treatment on the outcome that is not regulated by its descendant mediators. Similarly, by the first multiplier in the  $IM_j$ , we know that  $IM_j$  captures the indirect effect of a particular mediator on the outcome regulated by its descendant mediators.

More importantly, Proposition 5.2 can imply a simply OLS estimator for the direct strategy together with Proposition 5.4 as long as the sample size n is larger than the dimension d. Indeed, we can rewrite the part of semi-linear structure (5.1) as follows:

$$(5.2) \quad \begin{cases} M = \underbrace{(I - B_{MM}^{\intercal})^{-1} B_{MC}^{\intercal} C}_{=:\Theta_{MC}} + \underbrace{(I - B_{MM}^{\intercal})^{-1} \beta_{MA}}_{\theta_{MA}} A + \underbrace{(I - B_{MM}^{\intercal})^{-1} \epsilon_{M}}_{=:e_{M}}, \\ Y = (\beta_{YC}^{\intercal}, \alpha_{YA}, \beta_{YM}^{\intercal}) (C^{\intercal}, A, M^{\intercal})^{\intercal} + \epsilon_{Y} \end{cases}.$$

Write  $\widehat{\theta}_{MA}$  as the OLS estimator of unknown parameter  $\theta_{MA}$ , similarly define the other corresponding estimated quantities as follows:

$$\begin{bmatrix} \widehat{\Theta}_{MC}^\top \\ \widehat{\theta}_{MA}^\top \end{bmatrix} \qquad \text{and} \qquad \begin{bmatrix} \widehat{\beta}_{YC} \\ \widehat{\alpha}_{YA} \\ \widehat{\beta}_{YM} \end{bmatrix}.$$

Then we will have OLS estimators for the direct strategy estimators: For DE and IE,  $\widehat{DE}^{\text{OLS}} = \widehat{\alpha}_{YA}$  and  $\widehat{IE}^{\text{OLS}} = \widehat{\beta}_{YM}^{\top}\widehat{\theta}_{MA}$ ; For  $DM_j$  and  $IM_j$ ,  $\widehat{DM}_j^{\text{OLS}} = \widehat{\beta}_{YM,j}\widehat{\theta}_{MA,j}$ ,  $\widehat{IM}_j^{\text{OLS}}(\mathcal{G}_M) = \widehat{\theta}_{MA,j} \big\{ \widehat{\mathbf{E}}^{\text{reg}} \big[ Y \mid M_j \cup \mathrm{Pa}_j(\mathcal{G}_M) \cup A \cup C \big]_1 - \widehat{\beta}_{YM,j} \big\}$ , and

$$\widehat{IM}_{j}^{\text{avg, OLS}} = \frac{1}{\# \operatorname{MEC}(\widehat{\mathcal{C}}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\widehat{\mathcal{C}}_{M})} \widehat{IM}_{j}^{\text{OLS}}(\mathcal{G}_{M}),$$

where  $\widehat{\mathrm{E}}^{\mathrm{reg}}[X_k \mid X_l \cup X_S]_1$  is the estimated coefficient for  $X_l$  obtained from the linear regression of  $X_k$  on  $X_l \cup X_S$ , as determined from the data. Thus, when the semi-linear structure is determined, we can simplify direct strategy estimators to OLS estimators. All the these OLS estimators can be easily obtained by just applying simple regressions with nice properties, we will discuss their asymptotic properties in Section 7.

## 6. Multiple Robust Estimators.

- 6.1. Several Alternative Strategies. For a fixed  $j \in [p]$ , beyond the direct strategy above, there are alternative identification formulas for  $E\kappa(a',C)$ ,  $E\zeta_j(a',0,C)$ , and  $E\varrho_j(a',M_j,C;\mathcal{G}_M)$ . Based on these formulations, we can derive the corresponding estimators. We will discuss them one by one in the subsequent sections.
- 6.1.1. Alternative Strategy 1. The first one is using propensity score to construct the inverse probability weighting estimator. Note that we have<sup>2</sup>

$$E\left[\frac{\mathbb{1}(A=a')}{e_{a'}(C)}Y\right] = E\kappa(a',C),$$

(6.1) 
$$\operatorname{E}\left[\frac{\mathbb{1}(A=1)}{e_1(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_j}(M_{-j})} Y\right] = \operatorname{E}\zeta_j(a',0,C),$$

<sup>&</sup>lt;sup>2</sup>The calculation details are shown in C.3.

and

(6.2) 
$$\operatorname{E}\left[\frac{\mathbb{1}(A=a')}{e_{a'}(C)}\pi_{C,a'}\left(\operatorname{Pa}_{j}(\mathcal{G}_{M})\right)Y\right] = \operatorname{E}\varrho_{j}(a',M_{j},C\,;\,\mathcal{G}_{M}).$$

Thus, corresponding estimators take the form

$$\widehat{\kappa}^{\mathcal{M}_1}(a',C) = \frac{\mathbb{1}(A=a')}{\widehat{e}_{a'}(C)}Y,$$

$$\widehat{\zeta}_{j}^{\mathcal{M}_{1}}(a',0,C) = \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \frac{\widehat{\pi}_{C,a'}(M_{-j})}{\widehat{\pi}_{C,1,M_{j}}(M_{-j})} Y,$$

and

$$\widehat{\varrho}_{j}^{\mathcal{M}_{1}}(a', M_{j}, C; \mathcal{G}_{M}) = \frac{\mathbb{1}(A = a')}{\widehat{e}_{a'}(C)} \widehat{\pi}_{C, a'} (\operatorname{Pa}_{j}(\mathcal{G}_{M})) Y,$$

respectively. Here, the propensity scores  $e_{a'}(\cdot)$  and conditional densities  $\pi.(\cdot)$  appearing in  $\widehat{\kappa}^{\mathscr{M}_1}(a',C)$ ,  $\widehat{\zeta}_j^{\mathscr{M}_1}(a',0,C)$ , and  $\widehat{\varrho}_j^{\mathscr{M}_1}(a',M_j,C\,;\mathcal{G}_M)$  should be correctly estimated in the collection of quantities  $\mathscr{M}_{j,1}$  such that

•  $\mathcal{M}_{j,1}$ : The propensity scores  $P(A = \cdot \mid C = \cdot)$ , and the conditional density of the mediator  $f_{M_{-j}\mid C,A}(\cdot \mid C = \cdot, A = \cdot)$  and  $f_{M_{-j}\mid C,A,M_j}(\cdot \mid C = \cdot, A = 1, M_j = \cdot)$  are correctly specified for any  $\mathcal{G}_M \in \mathrm{MEC}(\mathcal{C}_M)$ .

Define

(6.3)

$$\overline{\mathbb{P}_n}\widehat{\varrho}_{j;1-0}^{\mathcal{M}_1}(M_j,C;\mathcal{C}_M) = \frac{1}{\#\operatorname{MEC}(\mathcal{C}_M)} \sum_{\mathcal{G}_M \in \operatorname{MEC}(\mathcal{C}_M)} \mathbb{P}_n\Big[ \big\langle \widehat{\varrho}_j^{\mathcal{M}_1}(\cdot,M_j,C;\mathcal{G}_M) \big\rangle \Big],$$

then, we can construct the estimators under  $\mathcal{M}_{j,1}$  is  $\widehat{DM}_j^{\mathcal{M}_1} = \mathbb{P}_n\left[\left\langle \widehat{\zeta}_j^{\mathcal{M}_1}(\cdot,0,C)\right\rangle\right]$ ,

$$\widehat{TM}_{j}^{\text{avg}, \mathcal{M}_{1}} = \frac{1}{\# \operatorname{MEC}(\widehat{\mathcal{C}}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\widehat{\mathcal{C}}_{M})} \widehat{TM}_{j}^{\mathcal{M}_{1}}(\mathcal{G}_{M})$$

$$= \mathbb{P}_{n} [\langle \widehat{\kappa}^{\mathcal{M}_{1}}(\cdot, C) \rangle] - \overline{\mathbb{P}_{n}} \widehat{\varrho}_{j; 1-0}^{\mathcal{M}_{1}}(M_{j}, C; \widehat{\mathcal{C}}_{M}),$$

and  $\widehat{IM}_j^{\mathrm{avg},\mathscr{M}_1} = \widehat{TM}_j^{\mathrm{avg},\mathscr{M}_1} - \widehat{DM}_j^{\mathscr{M}_1}$ , where  $\widehat{\mathcal{C}}_M$  is the estimated adjacency matrix consistent with the true  $\mathcal{C}_M$ . Here, the superscript  $\mathscr{M}_1$  associated with these estimators signifies that their consistency relies on the correct specification of information in  $\mathscr{M}_{j,1}$ . For clarity and where there is no risk of confusion, we will also employ  $\mathscr{M}_1$  to represent the estimation methodology behind these estimators. In the subsequent two subsections, the notations  $\mathscr{M}_2$  and  $\mathscr{M}_3$  bear analogous meanings.

6.1.2. Alternative Strategy 2. Similarly, we can verify that

(6.4) 
$$E\left[\frac{\mathbb{1}(A=0)}{e_0(C)} \int_{\mathcal{M}_j} \mu(C, 1, m_j, M_{-j}) \pi_{a', C}(m_j) \, \mathrm{d}m_j\right] = E\zeta_j(a', 0, C)$$

and

(6.5) 
$$\mathrm{E}\left[\frac{\mathbb{1}(A=a')}{e_{a'}(C)}\int_{\mathcal{M}_{j}}\mu(C,a',\mathrm{Pa}_{j}(\mathcal{G}_{M}),m_{j})\pi_{C}(m_{j})\,\mathrm{d}m_{j}\right]=\mathrm{E}\varrho_{j}(a',M_{j},C\,;\,\mathcal{G}_{M}).$$

Thus, corresponding estimators take the forms

$$\widehat{\zeta}_{j}^{\mathcal{M}_{2}}(a',0,C) = \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \int_{\mathcal{M}_{i}} \widehat{\mu}(C,1,m_{j},M_{-j}) \widehat{\pi}_{C,a'}(m_{j}) \, \mathrm{d}m_{j},$$

and

$$\widehat{\varrho}_{j}^{\mathcal{M}_{2}}(a', M_{j}, C; \mathcal{G}_{M}) = \frac{\mathbb{1}(A = a')}{\widehat{e}_{a'}(C)} \int_{\mathcal{M}_{i}} \widehat{\mu}(C, a', \operatorname{Pa}_{j}(\mathcal{G}_{M}), m_{j}) \widehat{\pi}_{C}(m_{j}) \, dm_{j}$$

with estimators  $\widehat{e}_{\cdot}(\cdot)$ ,  $\widehat{\mu}(\cdot)$ , and  $\widehat{\pi}_{\cdot}(\cdot)$  appear in  $\widehat{\kappa}^{\mathcal{M}_1}(a',C)$ ,  $\widehat{\zeta}_j^{\mathcal{M}_2}(a',0,C)$ , and  $\widehat{\varrho}_j^{\mathcal{M}_2}(a',M_j,C)$ ;  $\mathcal{G}_M$ ). They use the information in  $\mathcal{M}_{j,2}$  such that

•  $\mathcal{M}_{j,2}$ : The propensity scores  $\mathrm{P}(A=\cdot\mid C=\cdot)$ , conditional density of j-th mediator  $f_{M_j\mid C,A}(\cdot\mid C=\cdot,A=\cdot)$ , and the conditional expectations  $\mathrm{E}[Y\mid C=\cdot,A=\cdot,M=\cdot]$  and  $\mathrm{E}[Y\mid C=\cdot,A=\cdot,\mathrm{Pa}_j(\mathcal{G}_M)=\cdot,M_j=\cdot]$  are correctly specified for any  $\mathcal{G}_M\in\mathrm{MEC}(\mathcal{C}_M)$ .

Here we notice the fact that  $\pi_c(m_j) = \pi_{c,0}(m_j) + \pi_{c,1}(m_j)$ , and thus,  $\widehat{\kappa}^{\mathcal{M}_1}(a',C)$  will also be consistent in  $\mathcal{M}_{j,2}$ . Then  $\widehat{DM}_j^{\mathcal{M}_2} = \mathbb{P}_n\big[\big\langle \widehat{\zeta}_j^{\mathcal{M}_2}(\cdot,0,C) \big\rangle \big]$ ,

$$\widehat{TM}_{j}^{\text{avg}, \mathcal{M}_{2}} = \mathbb{P}_{n} \left[ \left\langle \widehat{\kappa}^{\mathcal{M}_{1}}(\cdot, C) \right\rangle \right] - \overline{\mathbb{P}_{n}} \widehat{\varrho}_{j; 1-0}^{\mathcal{M}_{2}}(M_{j}, C; \widehat{\mathcal{C}}_{M}),$$

and  $\widehat{IM}_j^{\text{avg},\mathcal{M}_2} = \widehat{TM}_j^{\text{avg},\mathcal{M}_2} - \widehat{DM}_j^{\mathcal{M}_2}$  are consistent provided that the estimated adjacency matrix  $\widehat{\mathcal{C}}_M$  is consistent to  $\mathcal{C}_M$ .  $\overline{\mathbb{P}_n}\widehat{\varrho}_{j;1-0}^{\mathcal{M}_2}(M_j,C\,;\widehat{\mathcal{C}}_M)$  is similarly defined by substituting  $\mathcal{M}_1$  with  $\mathcal{M}_2$  as detailed in (6.3).

6.1.3. *Alternative Strategy 3*. The last strategy is based on the third representation of the functional as follows:

(6.6) 
$$E\left[\frac{\mathbb{1}(A=a')}{e_{a'}(C)}\int_{\mathcal{M}_{-j}}\mu(C,1,M_j,m_{-j})\pi_{C,0}(m_{-j})\,\mathrm{d}m_{-j}\right] = E\zeta_j(a',0,C),$$

and

(6.7) 
$$\mathbb{E}\Big[\mathbb{E}\big[\varrho_j(a',M_j,C;\mathcal{G}_M)\mid C\big]\Big] = \mathbb{E}\varrho_j(a',M_j,C;\mathcal{G}_M).$$

Similarly, we can consider the estimators

$$\widehat{\zeta}_{j}^{\mathcal{M}_{3}}(a',0,C) = \frac{\mathbb{1}(A=a')}{\widehat{e}_{a'}(C)} \int_{\mathcal{M}_{-j}} \widehat{\mu}(C,1,M_{j},m_{-j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m_{-j},$$

and

$$\widehat{\varrho}_{j}^{\mathcal{M}_{3}}(a', M_{j}, C; \mathcal{G}_{M}) = \widehat{\mathbf{E}} \left[ \varrho_{j}(a', M_{j}, C; \mathcal{G}_{M}) \mid C \right]$$

$$= \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M}) \cup \mathcal{M}_{j}} \widehat{\mu} \left( C, a', \mathrm{pa}_{j}(\mathcal{G}_{M}), m_{j} \right) \widehat{\pi}_{C, a'} \left( \mathrm{pa}_{j}(\mathcal{G}_{M}) \right) \widehat{\pi}_{C}(m_{j}) \, \mathrm{d} \, \mathrm{pa}_{j}(\mathcal{G}_{M}) \, \mathrm{d} m_{j}.$$

Thus, our estimators under the third identification formulas can be written as  $\widehat{DM}_j^{\mathcal{M}_3} = \mathbb{P}_n \big[ \widehat{\zeta}_j^{\mathcal{M}_3}(1,0,C) - \widehat{\zeta}_j^{\mathcal{M}_3}(0,0,C) \big],$ 

$$\widehat{TM}_{j}^{\text{avg},\mathcal{M}_{3}} = \mathbb{P}_{n} \left[ \left\langle \widehat{\kappa}^{\mathcal{M}_{1}}(\boldsymbol{\cdot},C) \right\rangle \right] - \overline{\mathbb{P}_{n}} \widehat{\varrho}_{j;1-0}^{\mathcal{M}_{3}}(M_{j},C;\widehat{\mathcal{C}}_{M}),$$

and  $\widehat{IM}_j^{\mathrm{avg},\mathcal{M}_3} = \widehat{TM}_j^{\mathrm{avg},\mathcal{M}_3} - \widehat{DM}_j^{\mathcal{M}_3}$ , where the estimated adjacency matrix  $\widehat{\mathcal{C}}_M$  is consistent to  $\mathcal{C}_M$ , and  $\overline{\mathbb{P}_n}\widehat{\mathcal{C}}_{j;1-0}^{\mathcal{M}_3}(M_j,C\,;\widehat{\mathcal{C}}_M)$  is by replacing  $\mathcal{M}_1$  with  $\mathcal{M}_3$  in (6.3). The estimators  $\widehat{e}.(\cdot)$ ,  $\widehat{\mu}(\cdot)$ , and  $\widehat{\pi}.(\cdot)$  use the following information:

•  $\mathcal{M}_{j,3}$ : The propensity scores  $\mathrm{P}(A=\cdot\mid C=\cdot)$ , the conditional densities  $f_{M_{-j}\mid C,A}(\cdot\mid C=\cdot,A=\cdot)$  and  $f_{M_{j}\mid C}(\cdot\mid C=\cdot)$ , and the conditional expectation  $\mathrm{E}[Y\mid C=\cdot,A=\cdot,M=\cdot]$  are correctly specified for any  $\mathcal{G}_{M}\in\mathrm{MEC}(\mathcal{C}_{M})$ .

Here we note the fact that  $\widehat{\kappa}^{\mathcal{M}_1}(a',C)$  will be consistent in  $\mathcal{M}_{i,3}$  again.

6.2. Quadruply Robust Estimator. Denote  $\mathcal{M}_{j,\mathrm{union}} := \mathcal{M}_0 \cup \mathcal{M}_{j,1} \cup \mathcal{M}_{j,2} \cup \mathcal{M}_{j,3}$ , then  $\bigcup_{j=1}^p \mathcal{M}_{j,\mathrm{union}} \subsetneq \mathcal{M}_{\mathrm{nonpar}}$ , and  $\widehat{DM}_j^{\mathcal{M}_\ell}$  and  $\widehat{IM}_j^{\mathcal{M}_\ell}(\mathcal{G}_M)$  are all mapping the estimated distribution  $\widehat{F}_X$  to the true  $DM_j$  and  $IM_j(\mathcal{G}_M)$  defined in Definition 3.2 for  $\ell=0,1,2,3$  and any fixed  $\mathcal{G}_M$ , since all these representations agree on the nonparametric model  $\mathcal{M}_{\mathrm{nonpar}}$ . Therefore, we may conclude that both direct strategy and alternative strategies are in fact asymptotically efficient in  $\mathcal{M}_{\mathrm{nonpar}}$  with common scores  $S^{\mathrm{eff, nonpar}}(DM_j)$  and  $S^{\mathrm{eff, nonpar}}(IM_j(\mathcal{G}_M))$ . Furthermore, from this observation, one further concludes that (asymptotic) inferences obtained using one of the four representations are identical to inferences using either of the other three representations for a fixed  $\mathcal{G}_M$ . However, to achieve this, each strategy need exactly correctly specified for the conditional expectation and conditional density, i.e., correctly specified for the corresponding collections  $\mathcal{M}_{j,\ell}$  in Section 5, 6.1.1, 6.1.2, and 6.1.3, where we denote  $\mathcal{M}_{j,0} \equiv \mathcal{M}_0$  for each  $j \in [p]$ . In general,  $\widehat{DM}_j^{\mathcal{M}_\ell}$ ,  $\widehat{IM}_j^{\mathcal{M}_\ell}(\mathcal{G}_M)$  fail to be consistent outside of the corresponding submodel  $\mathcal{M}_{j,\ell}$  for each  $\ell \in \{0,1,2,3\}$ .

Note that the alternative strategy 1 in Section 6.1.1 in  $\mathcal{M}_{j,1}$  induces Inverse Probability Weighted (IPW) estimator. A commonly-used method is combining the direct strategy estimator in Section 5 correctly specified with the model  $\mathcal{M}_0$  and IPW estimator in 6.1.1 with the model  $\mathcal{M}_0$ , and getting the double robust estimator. But the double robust estimators only combine two estimation strategies,  $\mathcal{M}_0$  and  $\mathcal{M}_{j,1}$ , and ignore use other two alternative strategies. Hence, the double robust estimator may be inconsistent outside of  $\mathcal{M}_0 \cup \mathcal{M}_{j,1}$ . To overcome this problem, we propose an approach that produces a quadruply robust estimator by *combining the above all four strategies* as follows:

• 
$$\widehat{DM}_{j}^{QR}$$
 solves

$$\mathbb{P}_n \widehat{S}^{\text{eff, nonpar}}(\widehat{DM}_i^{\text{QR}}) = 0;$$

• For a fixed DAG  $\mathcal{G}_M$ ,  $\widehat{IM}_j^{\mathrm{QR}}(\mathcal{G}_M)$  solves

$$\mathbb{P}_n \widehat{S}^{\text{eff, nonpar}}(\widehat{IM}_j^{QR}(\mathcal{G}_M)) = 0,$$

where  $\widehat{S}^{\text{eff, nonpar}}(\cdot)$  is equal to  $S^{\text{eff, nonpar}}(\cdot)$  evaluated at the given consistent estimators  $\widehat{e}_{a'}(\cdot)$ ,  $\widehat{\pi}.(\cdot)$ , and  $\widehat{\mu}(\cdot)$  for all propensity scores, the conditional densities, and the conditional expectations appearing in  $S^{\text{eff, nonpar}}(\cdot)$ . Denote

$$\widehat{\tau}_{\cdot,S}(C,a',M_T) := \int_{\mathcal{M}_{-T}} \widehat{\mu}(C,a',m_{-T},M_T)\widehat{\pi}_{\cdot}(m_{-T}) dm_{-T},$$

as the corresponding estimator for  $\tau_{\cdot;S}(C,a',M_T)$  defined in (4.5), then we have the following explicit expressions for the quadruply estimators as

$$\widehat{DM}_{j}^{QR} := \widehat{DM}_{j}^{\mathcal{M}_{0}} + \mathbb{P}_{n} \left[ \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \frac{\langle \widehat{\pi}_{C,\cdot}(M_{-j}) \rangle}{\widehat{\pi}_{C,1,M_{j}}(M_{-j})} \left[ Y - \widehat{\mu}(C,1,M) \right] \right]$$

$$+ \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \left\langle \widehat{\tau}_{C,\cdot;j}(C,1,M_{-j}) - \widehat{\zeta}_{j}^{\mathcal{M}_{0}}(\cdot,0,C) \right\rangle$$

$$+ \left\langle \frac{\mathbb{1}(A=\cdot)}{\widehat{e}_{\cdot}(C)} \right\rangle \widehat{\tau}_{C,0;-j}(C,1,M_{j}) - \left\langle \frac{\mathbb{1}(A=\cdot)}{\widehat{e}_{\cdot}(C)} \widehat{\zeta}_{j}^{\mathcal{M}_{0}}(\cdot,0,C) \right\rangle \right],$$

$$\widehat{TM}_{j}^{QR}(\mathcal{G}_{M}) := \widehat{TM}_{j}^{\mathcal{M}_{0}} + \mathbb{P}_{n} \left[ \left\langle \frac{\mathbb{1}(A=\cdot)}{\widehat{e}_{\cdot}(C)} \left( Y - \widehat{\mu}(C,\cdot) \right) \right\rangle \right]$$

$$(6.9) \quad - \mathbb{P}_{n} \left[ \left\langle \frac{\mathbb{1}(A=\cdot)}{\widehat{e}_{\cdot}(C)} \widehat{\pi}_{C,\cdot} \left( \operatorname{Pa}_{j}(\mathcal{G}_{M}) \right) \left[ Y - \widehat{\mu}(C,\cdot,\operatorname{Pa}_{j}(\mathcal{G}_{M}),M_{j}) \right] \right\rangle$$

$$+ \left\langle \frac{\mathbb{1}(A=\cdot)}{\widehat{e}_{\cdot}(C)} \left( \widehat{\tau}_{C;j} \left( C,\cdot,\operatorname{Pa}_{j}(\mathcal{G}_{M}) \right) - \widehat{\operatorname{E}} \left[ \varrho_{j}(\cdot,M_{j},C;\mathcal{G}_{M}) \mid C \right] \right) \right\rangle \right],$$

and  $\widehat{IM}_j^{\mathrm{QR}}(\mathcal{G}_M) = \widehat{TM}_j^{\mathrm{QR}}(\mathcal{G}_M) - \widehat{DM}_j^{\mathrm{QR}}$ . Then the quadruply estimator for indirect interventional mediation effect with a consistent estimated  $\widehat{\mathcal{C}}_M$  is defined as  $\widehat{IM}_j^{\mathrm{avg,QR}} = \frac{1}{\#\mathrm{MEC}(\widehat{\mathcal{C}}_M)} \sum_{\mathcal{G}_M \in \mathrm{MEC}(\widehat{\mathcal{C}}_M)} \widehat{IM}_j^{\mathrm{QR}}(\mathcal{G}_M)$ . Compared to double robust estimators, our novel quadruply robust estimators can tolerate a higher degree of misspecification outside of  $\mathcal{M}_0 \cup \mathcal{M}_{j,1}$  and still achieve consistency. We will see this in Section 7.2.

Subject to some mild regularity conditions, delineated in Section 7.2, our quadruply estimators are asymptotic normal and efficient. Thus, based on the semiparametric efficient

scores, we get the score-based variance estimators for  $\widehat{DM}_j^{\rm QR}$  and  $\widehat{IM}_j^{\rm avg,\,QR}$  as

$$\widehat{\mathrm{var}}(\widehat{DM}_j^{\mathrm{QR}}) := \frac{1}{n^2} \sum_{i=1}^n \left[ \widehat{S}^{\mathrm{eff,\,nonpar}}(\widehat{DM}_j^{\mathrm{QR}}) - \widehat{DM}_j^{\mathrm{QR}} \right]^2$$

and

$$\begin{split} \widehat{\operatorname{var}}(\widehat{IM}_{j}^{\operatorname{avg,QR}}) \\ &:= \frac{1}{n^{2}} \sum_{i=1}^{n} \left[ \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} \left( \widehat{S}^{\operatorname{eff, nonpar}} \left( \widehat{IM}_{j}^{\operatorname{QR}}(\mathcal{G}_{M}) \right) - \widehat{IM}_{j}^{\operatorname{QR}}(\mathcal{G}_{M}) \right) \right]^{2} \end{split}$$

correspondingly. However, in practical scenarios, confidence intervals (CIs) derived using the Wald-type method, especially when grounded on score-based variance estimators, tend to be more narrow (Boos and Stefanski, 2013). This can potentially result in anticonservatism. To achieve more concise statistical inference for our quadruply estimators, we consider utilizing the variances derived from the symmetric t-bootstrap approach (Hall, 1988) here. A pseudocode summarizing the proposed algorithm for these quadruply estimators and their bootstrap CIs is given in Algorithm 1. The  $\log n$  truncations in Algorithm 1 aims to achieve the numerical stability, which is a technique widely recognized in statistical literature (Heckman, 1976; Sun et al., 2020; Chinot et al., 2020).

6.3. Practical fast implement. The formulas for the quadruply robust estimators, as shown in equations (6.8) and (6.9), require several numerical integrals for each  $i \in [n]$ , which may be computationally demanding. To address this challenge, we purpose Algorithm 1 in the above section, in which we employ the Monte Carlo method to evaluate these integrals. However, when the data partly satisfy the semi-linear structure and both  $\epsilon_M$  and  $\epsilon_Y$  adhere to a mean-zero Gaussian distribution, explicit expressions for these numerical integrals can be derived, facilitating faster computation. Indeed, if we assume the linear structure in  $M \leftarrow C \oplus A \oplus M$  and denote the density (or mass) function of  $\epsilon_M = (\epsilon_{M,1}, \dots, \epsilon_{M,p})^{\top}$  as  $f(x) = f(x_1, \dots, x_p)$ , then the conditional density of M given C and A = 1 is  $f(x - \Theta_{MC}C - \theta_{MA})$  from (5.2). This allows us to compute (6.10)

$$\int_{\mathcal{M}_{j}} \mu(C, 1, m_{j}, M_{-j}) \pi_{C,1}(m_{j}) dm_{j}$$

$$= \int_{\mathcal{M}_{j}} \left[ \beta_{YC}C + \alpha_{YA} + \beta_{YM,j} m_{j} + \beta_{YM,-j}^{\mathsf{T}} M_{-j} \right] f\left(m_{j} - \left[\Theta_{MC}C\right]_{j} - \theta_{MA,j}\right) dm_{j}$$

$$= \beta_{YC}C + \alpha_{YA} + \beta_{YM,j} \left\{ \left[\widehat{\Theta}_{MC}C\right]_{j} + \theta_{MA,j} \right\} + \beta_{YM,-j}^{\mathsf{T}} M_{-j}.$$

Similarly, we can derive explicit expressions for some other integrals in equations (6.8) and (6.9) as long as the linear structure in  $M \leftarrow C \oplus A \oplus M$  holds. One step more, when  $\epsilon_M$  is a mean-zero Gaussian distribution, any integral in (6.8) and (6.9) will have an explicit expression. This leads to a more efficient implementation of (6.8) and (6.9). The following Algorithm 2 and Proposition 6.1 elaborates on this.

## **Algorithm 1** General algorithm for quadruply robust estimations

**INPUT:** The data  $\{X_i\}_{i=1}^n$ , the treatment index t, and the Monte Carlo sample size N.

- 1: Apply any suitable structure learning algorithm to learn the CPDAG of X and obtain the corresponding estimated adjacency matrix  $\mathcal{C}$ , then set the estimated adjacency matrix of mediators as  $\mathcal{C}_M =$  $\left[\widehat{\mathcal{C}}_{kk}\right]_{k\in(t+1):(t+p)}$ .
- 2: Using any proper estimating procedure to estimate the propensity score  $e_{a'}(c) = P(A = a' \mid C = c)$  with
- 3: Using any proper estimation method to estimate the conditional densities  $\pi_{\bullet}(\cdot)$  and conditional expectations
- 4: For estimating the causal effect of j-th mediator, set  $\widehat{DM}_j^{\mathrm{QR}} = \widehat{TM}_j^{\mathrm{avg},\mathrm{QR}} = 0$ .
- Sample  $M_{\widehat{\pi}_{C_i,1}}^{(1)},\dots,M_{\widehat{\pi}_{C_i,1}}^{(N)} \overset{\text{i.i.d.}}{\sim} \widehat{\pi}_{C,1}(m) := \widehat{\pi}_{C,1}(m_j)\widehat{\pi}_{C,1,m_j}(m_{-j})$ , and similarly sample  $M_{\boldsymbol{\cdot},\widehat{\pi}_{\cdot_i}}^{(1)},\dots,M_{\boldsymbol{\cdot},\widehat{\pi}_{\cdot_i}}^{(N)}$  from any other conditional densities displaying in (6.8).  $\widehat{DM}_j^{\text{QR}} \longleftarrow \widehat{DM}_j^{\text{QR}} + n^{-1}\widehat{QR}_{DM_j}^{\text{MC}}(X_i)\mathbb{1}\left(|\widehat{QR}_{DM_j}^{\text{MC}}(X_i)| \leq \log n\right), \text{ where } \widehat{QR}_{DM_j}^{\text{MC}}(X_i) \text{ is defined in (B.1).}$
- for  $\mathcal{G}_M \in \mathrm{MEC}(\widehat{\mathcal{C}}_M)$  do
- Obtain  $M_j$ 's parent mediators  $\operatorname{Pa}_j(\mathcal{G}_M) = \left[\mathcal{G}_M^\top \circ 1_p 1_p^\top\right]_{i}$ .
- Sample  $M^{(1)}_{\boldsymbol{\cdot},\widehat{\pi}_{\cdot_i}},\dots,M^{(N)}_{\boldsymbol{\cdot},\widehat{\pi}_{\cdot_i}}$  from all conditional densities displaying in (6.9) and in  $\widehat{TM}_j^{\mathcal{M}_0}$ .  $\widehat{TM}_j^{\mathrm{QR}}(\mathcal{G}_M) \longleftarrow \widehat{TM}_j^{\mathrm{QR}}(\mathcal{G}_M) + n^{-1}\widehat{QR}_{TM_j}^{\mathrm{MC}}(X_i;\mathcal{G}_M)\mathbb{1}\big(|\widehat{QR}_{TM_j}^{\mathrm{MC}}(X_i)| \leq \log n\big), \text{ where } 1$ 11:  $TM_j$   $(g_M) \leftarrow TM_j$   $(g_M) + R$   $\mathcal{Q}_{M,j} + R$
- 13: Symmetric *t*-bootstrap (Hall, 1988) is applied to construct confidence intervals for  $\sqrt{n}(\widehat{DM}_{i}^{QR} DM_{i})$ and  $\sqrt{n}(\widehat{IM}_j^{\mathrm{avg,\,QR}}-\overline{IM}_j).$

# Algorithm 2 Fast implement algorithm for quadruply robust estimations

**INPUT:** The data  $\{X_i\}_{i=1}^n$  and the treatment index t.

- 1: Apply any suitable structure learning algorithm (such as GES or PC) to learn the CPDAG of X and obtain  $\widehat{\mathcal{C}}$ , then obtain the CPDAG of mediators  $\left[\widehat{\mathcal{C}}_{kk}\right]_{k\in(t+1):(t+p)}$
- 2: Using any proper estimating procedure to estimate the propensity score  $e_1(c) = P(A = 1 \mid C = c)$  with  $\widehat{e}_1(c)$  and  $\widehat{e}_0(c) = 1 - \widehat{e}_1(c)$ .
- 3: Regress M on  $(C^{\top}, A)^{\top}$  obtain OLS estimator  $(\widehat{\Theta}_{MC}, \widehat{\theta}_{MA})^{\top}$  and the estimated covariance of error
- 4: Regress Y on  $(C^{\top}, A, M^{\top})^{\top}$  and  $(C^{\top}, A, M^{\top})^{\top}$ , obtain OLS estimators  $(\widehat{\beta}_{YC}^{\top}, \widehat{\alpha}_{YC}, \widehat{\beta}_{YM}^{\top})^{\top}$  and  $(\widehat{\eta}_{YA}^{\dagger}, \widehat{\gamma}_{YC}^{\dagger\top})^{\top}$  correspondingly. 5: **for**  $j \in [p]$  **do**
- Set  $\widehat{DM}_j^{\mathrm{QR}} = \widehat{\beta}_{YM,j} \widehat{\theta}_{MA,j}$  and  $\widehat{TM}_j^{\mathrm{avg},\,\mathrm{QR}} = 0$ . for  $i \in [n]$  do
- $\widehat{DM}_{j}^{\mathrm{QR}} \longleftarrow \widehat{DM}_{j}^{\mathrm{QR}} + n^{-1} \widehat{QR}_{DM_{j}}^{\mathrm{fast}}(X_{i}), \text{ where } \widehat{QR}_{DM_{j}}^{\mathrm{fast}}(X_{i}) \text{ is defined in (B.3)}.$
- for  $\widetilde{\mathcal{G}}_M\in\mathrm{MEC}(\widehat{\mathcal{C}}_M)$  do 9:
- Obtain  $M_j$ 's parent mediators  $\operatorname{Pa}_j(\mathcal{G}_M) = \left[\mathcal{G}_M^\top \circ 1_p 1_p^\top\right]_{j}$ . 10:
- OLS estimator
- $\widehat{TM}_{j}^{\mathrm{QR}}(\mathcal{G}_{M}) \longleftarrow n^{-1}\left(\widehat{\eta}_{YM_{j}} + \widehat{QR}_{TM_{j}}^{\mathrm{fast}}(X_{i};\mathcal{G}_{M})\right), \text{ where } \widehat{QR}_{TM_{j}}^{\mathrm{fast}}(X_{i},\mathcal{G}_{M}) \text{ is defined}$ 12:
- **return**  $\widehat{DM}_{j}^{\mathrm{QR}}$  and  $\widehat{IM}_{j}^{\mathrm{avg,QR}} = \frac{1}{\# \mathrm{MEC}(\widehat{\mathcal{C}}_{M})} \sum_{\mathcal{G}_{M} \in \mathrm{MEC}(\widehat{\mathcal{C}}_{M})} \widehat{TM}_{j}^{\mathrm{QR}}(\mathcal{G}_{M}) \widehat{DM}_{j}^{\mathrm{QR}}$  with their symmetric t-bootstrap CIs.

PROPOSITION 6.1. Assume that at least one linear structure in Assumption 5.1 holds, and that  $\epsilon_M$  and  $\epsilon_Y$  are both mean-zero Gaussian distributed, Algorithm 2 produces valid quadruply robust estimators  $\{\widehat{DM}_j^{QR}, \widehat{IM}_j^{avg, QR}\}_{j=1}^p$  as defined in Section 6.

In practical scenarios where the sample size n is sufficiently large, it becomes reasonable to treat the sample means  $\overline{\epsilon}_M := \frac{1}{n} \sum_{i=1}^n \epsilon_{M,i}$  and  $\overline{\epsilon}_Y := \frac{1}{n} \sum_{i=1}^n \epsilon_{Y,i}$  as if they follow mean-zero Gaussian distributions. This permits the utilization of Proposition 6.1, particularly when empirical evidence can support the linear structural relationships for  $M \leftarrow C \oplus A \oplus M$  or  $Y \leftarrow C \oplus A \oplus M$ .

- **7. Asymptotic Behavior.** In this section, we first give the asymptotic properties of the OLS estimators when the model satisfies Assumption 5.1. Then we will establish the asymptotic normality of quadruply robust estimators, allowing the model misspecification.
- 7.1. Asymptotic Properties of OLS estimators. In section 5.1, we highlighted that given the causal structure is appropriately specified as semi-linear according to Assumption 5.1, one can employ OLS estimators by just applying two simple regressions. As we allow the number of mediators p can grow with sample size n, some assumptions are required. The following assumptions come from Portnoy (1984) and Portnoy (1985). They control the behavior of minimum eigenvalue will hold in probability if the observations are a sample from an appropriate distribution in  $\mathbb{R}^p$ . Denote the error vector as  $\epsilon = (\epsilon_A^{\mathsf{T}}, \epsilon_M^{\mathsf{T}}, \epsilon_Y^{\mathsf{T}})^{\mathsf{T}}$ .

ASSUMPTION 7.1. (Assumptions for Error Distributions)  $\epsilon$  is marginal sub-Gaussian with finite Orlicz norm (Definition (6.18) in Wainwright (2019)).

ASSUMPTION 7.2. (Restricted Eigenvalue Condition)  $\lim_{n\to\infty} \operatorname{var} (Y \mid C \cup A \cup M) > 0$  and  $\lim_{n\to\infty} \operatorname{Evar}(M \mid C \cup A) \succ 0$ .

We use the bold symbol  $\mathbf{X}$  to represent the data matrix of any i.i.d. random observations  $\{X_i\}_{i=1}^n$ . i.e.  $\mathbf{X} = [X_1, \dots, X_n]^{\top}$ . Denote the transformation  $\widehat{\Gamma}_{\cdot, \cdot} : \mathbb{R}^{n \times p_1} \times \mathbb{R}^{n \times p_2} \to \mathbb{R}^{p_1 \times n}$  of two data matrix with sample size n as

$$\widehat{\Gamma}_{X,Z} := \left[ \mathbf{X}^{\top} (I_n - P_{\mathbf{Z}}) \mathbf{X} \right]^{-1} \mathbf{X}^{\top} (I_n - P_{\mathbf{Z}}),$$

where  $P_{\mathbf{Z}} = \mathbf{Z}(\mathbf{Z}^{\top}\mathbf{Z})^{-1}\mathbf{Z}^{\top} \in \mathbb{R}^{n \times n}$  is the projection matrix of  $\mathbf{Z}$ . This transformation streamlines our representation of the asymptotics for our OLS estimators.

THEOREM 7.3. Suppose the model satisfies Assumptions 2.1, 5.1, 7.1, and 7.2. Let  $\{\widehat{e}_{M,i}\}_{i=1}^n$  and  $\{\widehat{\epsilon}_{Y,i}\}_{i=1}^n$  be the residuals from the OLS estimator in regression (5.2). Then for any  $\alpha \in (0,1)$ , we have

$$\lim_{n\to\infty} \mathbf{P}\Bigg( \left| \sqrt{n} (\widehat{DE}^{OLS} - DE) \right| \le \Phi^{-1} (1 - \alpha/2) \sqrt{\widehat{\Gamma}_{A,(M,C)} \widehat{\Gamma}_{A,(M,C)}^{\top} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2}} \Bigg) = 1 - \alpha$$

Furthermore, denote  $\widehat{\Sigma}_{\beta_{YM}} = \sum_{i=1}^n \widehat{\epsilon}_{Y,i}^2 \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{M,(C,A)}^{\top}$  and  $\widehat{\Sigma}_{\theta_{MA}} = \sum_{i=1}^n \widehat{e}_{M,i} \widehat{e}_{M,i}^{\top} \widehat{\Gamma}_{A,C} \widehat{\Gamma}_{A,C}^{\top}$ , then

$$\lim_{n\to\infty} \mathbf{P}\Bigg( \left| \sqrt{n} (\widehat{IE}^{OLS} - IE) \right| \le \Phi^{-1} (1 - \alpha/2) \sqrt{\widehat{\beta}_{YM}^{\top} \widehat{\Sigma}_{\theta_{MA}} \widehat{\beta}_{YM} + \widehat{\theta}_{MA}^{\top} \widehat{\Sigma}_{\beta_{YM}} \widehat{\theta}_{MA}} \Bigg) \ge 1 - \alpha,$$

and

$$\lim_{n\to\infty} \mathbf{P}\Bigg( \Big| \sqrt{n} (\widehat{DM}_j^{OLS} - DM_j) \Big| \le \Phi^{-1} (1 - \alpha/2) \sqrt{\widehat{\beta}_{YM,j}^2 \widehat{\Sigma}_{\theta_{MA},jj} + \widehat{\theta}_{MA,j}^2 \widehat{\Sigma}_{\beta_{YM},jj}} \Bigg) \ge 1 - \alpha$$

for any  $j \in [p]$ .

The above theorem ensures that under mild conditions we can construct valid confidence intervals for  $\widehat{DE}^{OLS}$ ,  $\widehat{IE}^{OLS}$ , and  $\widehat{DM_j}^{OLS}$  when n is large enough. It is worthy to note that the probabilities for  $\widehat{IE}^{OLS}$  and  $\widehat{DM_j}^{OLS}$  is  $\geq$  instead of =. This distinction arises from the dual nature of the limiting distributions for these two OLS estimators: one is the standard normal, the other is not. However, as argued in Chakrabortty et al. (2018), the non-standard asymptotic distributions here are more conservative than  $\mathcal{N}(0,1)$ . Thus, we obtain  $\geq$  instead of =. The details can be found in the proof. Notably, these asymptotic confidence intervals can be derived concurrently with the regression estimators and residuals. When applying the regression to procure these estimators, no additional steps are needed to obtain these confidence intervals.

these confidence intervals. For the estimators  $\widehat{IM}_j^{\text{OLS}}$ , additional assumptions are needed due to their reliance on the unknown DAG structure. This necessitates consistent CPDAG estimation, as well as more strong sparsity assumptions and restricted eigenvalue conditions, which are common in high-dimensional settings (Portnoy, 1985; Van de Geer et al., 2014; Zhang and Zhang, 2014; Chakrabortty et al., 2018).

ASSUMPTION 7.4. (Structure learning consistency) Consistency of learning structure:  $P(\widehat{C}_M \neq C_M) \longrightarrow 0$ .

ASSUMPTION 7.5. The sparsity of maximum degree in  $C_M$ ,  $\max_{j \in [p]} q_j = \max_{j \in [p]} |\operatorname{adj}(M_j)| = O(n^{1-b_1})$  for some  $0 < b_1 \le 1$ .

ASSUMPTION 7.6.  $\lim_{n\to\infty} \max_{j\in[p]} n^{-1/2} \left\{ q_j + \log\left(L_{\mathrm{distinct},j}\right) \right\} = 0$ , where  $L_{\mathrm{distinct},j}$  is the number of distinct elements of the set  $\left\{ X_{S_{j1}}, \dots, X_{S_{jL_j}} \right\} = \left\{ (M_j, \mathrm{Pa}_j(\mathcal{G}_M), A, C)^\top : \mathcal{G}_M \in \mathrm{MEC}(\mathcal{C}_M) \right\}$ .

Assumption 7.7.  $\lim_{n\to\infty} \min_{j\in[p]} \operatorname{var}(Y\mid\operatorname{adj}(M_j)\cup C\cup M_j)>0$  and  $\lim_{n\to\infty} \min_{j\in[p]} \operatorname{E}\operatorname{var}(M_j\mid\operatorname{adj}(M_j)\cup C)>0$ .

THEOREM 7.8. Suppose Assumption 2.1, 5.1, 7.1, 7.2 and Assumption 7.4, 7.5, 7.6, 7.7 hold, then

$$\lim_{n \to \infty} P\left(\sqrt{n} | \widehat{IM}_j^{OLS} - IM_j| \ge \widehat{\sigma}_{\overline{IM}_j} \Phi^{-1}(1 - \alpha/2)\right) \ge 1 - \alpha$$

for any  $\alpha \in (0,1)$ . The explicit formula for  $\widehat{\sigma}_{\overline{IM}_i}^2$  can be found in (C.6) in Appendix C.

We now therefore obtain a valid asymptotic confidence interval for  $\overline{IM}_j$  for any  $j \in [p]$  alongside the regression from Theorem 7.8.

7.2. Asymptotic Properties of Quadruply Robust Estimators. The quadruply robust estimators aim to obtain the robust estimators even when the model is misspeficied. The double robust estimators, which combines the direct and IPW strategies, possess commendable properties and have been the subject of extensive research as evidenced in literature such as (Laan and Robins, 2003; Tsiatis, 2006; Kang and Schafer, 2007). In this section, we will show that the proposed novel quadruply robust estimators exhibit more favorable asymptotic properties.

To present the results, we assume that the propensity score  $e_{a'}(\cdot) \in \mathcal{E}$  with some function classes  $\mathcal{E}$ . Similarly, for each  $j \in [p]$ , we assume any conditional density employed in (6.8) and (6.9) is

$$f_{M_T|X_S}(m_T \mid x_S) \in \mathcal{F}_{j,T|S},$$

and any conditional mean used in (6.8) and (6.9) adheres to

$$E[Y \mid x_S] \in \mathcal{U}_{j,S}$$

with some specific function classes  $\mathcal{F}_{j,T|S}$  and  $\mathcal{U}_{j,S}$ . We propose the following assumptions concerning these function classes and the convergence rates of the estimators within these classes.

ASSUMPTION 7.9. For any fixed  $j \in [p]$ , any subset  $S \subseteq [t+p+1]$  and  $T \subseteq [p]$  used in (6.8) and (6.9), the function classes  $\mathcal{E}$ ,  $\mathcal{F}_{j,T|S}$ , and  $\mathcal{U}_{j,S}$  are bounded and belong to VC type classes (Definition 2.1 in Chernozhukov et al. (2014)) with VC indices upper bounded by  $v_j = O(n^{\vartheta_j})$  for some  $\vartheta_j$  such that  $\vartheta_j \in [0, 1/2)$ .

ASSUMPTION 7.10. For any fixed  $j \in [p]$ , any subset  $S \subseteq [t+p+1]$  and  $T \subseteq [p]$  used in (6.8) and (6.9), the estimators  $\widehat{\pi}_{x_S}(m_T)$  and  $\widehat{\mu}(x_s)$  converge with  $\ell_2$ -norm to their true values at the rates of  $n^{-\vartheta_{j,\pi}^*}$  and  $n^{-\vartheta_{j,\mu}^*}$ , and the propensity score estimator  $\widehat{e}_{a'}(\cdot)$  with  $a' \in \{0,1\}$  converge with  $\ell_2$ -norm to their true values at the rates of  $n^{-\vartheta_{j,e}^*}$ . Here the positive numbers  $\vartheta_{j,e}^*, \vartheta_{j,\pi}^*, \vartheta_{j,\mu}^*$  satisfies: (i)  $\min\{\vartheta_{j,e}^*, \vartheta_{j,\pi}^*, \vartheta_{j,\mu}^*\} > \vartheta_j/2$ ; (ii)  $v_1^* + v_2^* > 1/2$  for any  $\{v_1^*, v_2^*\} \subsetneq \{\vartheta_{j,e}^*, \vartheta_{j,\pi}^*, \vartheta_{j,\mu}^*\}$ .

Assumption 7.9 is reasonably moderate, as the function classes are user-defined. VC-type classes encompass a broad spectrum of functional categories, including but not limited to classic parametric model, neural networks and regression trees. The VC index governs the complexity of the model, typically escalating with an increase in the number of parameters within the model. We permit the VC index to diverge alongside the sample size, which serves to minimize the estimator's bias arising from model misspecification. On the other hand, an important feature of Assumption 7.10 is that the required estimators' convergence rates can only be nonparametric (slower than  $n^{-1/2}$ ) and no metric entropy

condition (Donsker class for instance) is needed. In particular,  $\vartheta_{j,e}^*$ ,  $\vartheta_{j,\pi}^*$ ,  $\vartheta_{j,\mu}^* > 1/4$  will perfectly admit Assumption 7.10. Therefore, the estimators can be computed via standard nonparametric estimation (Fan and Yao, 2003) and supervised learning algorithms (including random forests and deep learning, Wager and Athey, 2018; Schmidt-Hieber, 2020). The reason both assumptions regarding the sizes of the function classes and the rate of convergence for  $\widehat{DM}_j^{\mathrm{QR}}$  and  $\widehat{IM}_j^{\mathrm{avg},\,\mathrm{QR}}$  are identical, which are different from conditions in Theorem 7.3 and Theorem 7.8, stems from the uniform convergence characteristics of our estimations for any  $\mathcal{G}_M$  in  $\mathrm{MEC}(\mathcal{C}_M)$ .

THEOREM 7.11. Let the conditions in Theorem 4.3 and Assumption 7.9 hold. Suppose the estimators  $\widehat{e}_{a'}(x_s)$ ,  $\widehat{\mu}(x_s)$ , and  $\widehat{\pi}_{x_s}(m_T)$  in either  $\mathcal{M}_0$ ,  $\mathcal{M}_{j,1}$ ,  $\mathcal{M}_{j,2}$ , or  $\mathcal{M}_{j,3}$  converges in  $\ell_2$ -norm to their true values for each  $j \in [p]$ . Then

- $\widehat{DM}_{j}^{QR}$  is the consistent estimator of  $DM_{j}$  under the model  $\mathcal{M}_{j,union}$  for any  $j \in [p]$ . Furthermore, if Assumption 7.10 holds, then  $\sqrt{n} (\widehat{DM}_{j}^{QR} DM_{j})$  is asymptotic normally distributed under model  $\mathcal{M}_{nonpar}$  with asymptotic variance  $\mathbb{E} \left\{ \left[ S^{eff, nonpar}(DM_{j}) \right]^{2} \right\}$
- mally distributed under model  $\mathcal{M}_{nonpar}$  with asymptotic variance  $\mathrm{E}\Big\{\big[S^{eff,\ nonpar}(DM_j)\big]^2\Big\}$ .

   If Assumption 7.4 also hold,  $\widehat{IM}_j^{avg,QR}$  is the consistent estimator of  $\overline{IM}_j$  under the model  $\mathcal{M}_{j,union}$  for any  $j\in[p]$ . Furthermore, if Assumption 7.10 holds, then  $\sqrt{n}\big(\widehat{IM}_j^{avg,QR}-\overline{IM}_j\big)$  is asymptotically normally distributed under model  $\mathcal{M}_{nonpar}$  with asymptotic variance

$$\mathbb{E}\Bigg\{\Bigg[\frac{1}{\#\operatorname{MEC}(\mathcal{C}_{M})}\sum_{\mathcal{G}_{M}\in\operatorname{MEC}(\mathcal{C}_{M})}S^{\textit{eff, nonpar}}(IM_{j}(\mathcal{G}_{M}))\Bigg]^{2}\Bigg\}.$$

An important result of Theorem 7.11 is that: for any  $j \in [p]$ , the quadruply robust estimators  $\widehat{DM}_j^{\mathrm{QR}}$  and  $\widehat{IM}_j^{\mathrm{avg,QR}}$  are semiparametric locally efficient in the sense that they are regular and asymptotically linear under model  $\mathcal{M}_{j,\mathrm{union}}$ , and achieve the semiparametric efficiency bound for  $DM_j$  and  $\overline{IM}_j$  under model at the intersection submodel  $\mathcal{M}_0 \cap \mathcal{M}_{j,1} \cap \mathcal{M}_{j,2} \cap \mathcal{M}_{j,3}$ . Hence, when all models are correct,  $\widehat{DM}_j^{\mathrm{QR}}$  and  $\widehat{IM}_j^{\mathrm{avg,QR}}$  are semiparametric efficient in the model  $\mathcal{M}_{\mathrm{nonpar}}$  at the intersection submodel  $\mathcal{M}_0 \cap \mathcal{M}_{j,1} \cap \mathcal{M}_{j,2} \cap \mathcal{M}_{j,3}$  by part iv in Bickel and Kwon (2001) for any  $j \in [p]$ .

**8. Simulation Studies.** In this section, we assess the finite-sample performance of our proposed quadruply robust estimators across two simulation scenarios. The first scenario seeks to illustrate the robustness characteristics of our estimator in comparison to other estimation strategies, particularly when certain model specifications to a specific mediator are not met. In the second simulation study, we demonstrate that our method can also be superior to any other estimation strategies in estimating the both direct and indirect interventional effects across all mediators, on average, within commonly adopted model configurations.

- 8.1. Simulation for a single mediator. We consider the finite-sample performance of the proposed quadruply robust estimators in comparison to the estimators under direct strategy, and the alternative strategies in Section 5, 6.1.1, 6.1.2, and 6.1.3 for a single mediator. We describe the detailed setting as follows: we set t=p=3, and fix the pre-specified j randomly sampled from  $U\{[p]\}$ . Then we design the following four data generating processes (DGPs), here  $\Phi(\cdot)$  and  $\operatorname{logit}(\cdot)$  are the standard normal distribution function and the inverse of the standard logistic function, and  $B_{j:}$  and  $B_{-j:}$  represent the j-th row of the matrix B and the matrix B with j-th row removed.
- All correct:  $C \leftarrow \mathcal{N}(0, I_{t-1}), A \leftarrow \mathbb{1}\left\{U[0, 1] \leq \Phi(\beta_{AC}^{\top}C)\right\}, M \leftarrow B_{MC}^{\top}C + \beta_{MA}A + B_{MM}^{\top}M + \mathcal{N}(0, I_p), \text{ and } Y \leftarrow \beta_{YC}^{\top}C + \alpha_{YA}A + \beta_{YM}^{\top}M + \mathcal{N}(0, 1);$
- $\mathcal{M}_0$  is correct: the exposure A comes from  $A \leftarrow \mathbb{1}(\operatorname{logit}(U[0,1]) \leq \beta_{AC}^{\top}C)$  instead;
- $\mathcal{M}_{j,1}$  is correct: the outcome Y comes from  $Y \leftarrow (\beta_{YC}^\top C + \alpha_{YA}A + \beta_{YM}^\top M)^{2/3} + \mathcal{N}(0,1)$  instead;
- $\mathcal{M}_{j,2}$  is correct: the mediators have the alternative structure  $M_j \leftarrow \Theta_{MC,j:}C + \theta_{MA,j}A + \left[ (I B_{MM}^\top)^{-1}\mathcal{N}(0,I_p) \right]_j$  and  $M_k \leftarrow (\Theta_{MA,k:}C + \theta_{MA,k}A)^{2/3} + \left[ (I B_{MM}^\top)^{-1}\mathcal{N}(0,I_p) \right]_k$  for  $k \neq j$ ;
- $\mathcal{M}_{j,3}$  is correct: the mediators have the alternative structure  $M_j \leftarrow \Theta_{MC,j}$ :  $C + \frac{1}{2}\theta_{MA,j} + \left[ (I B_{MM}^\top)^{-1} \mathcal{N}(0,I_p) \right]_j$  and  $M_{-j} \leftarrow \Theta_{MC,-j}$ :  $C + \theta_{MA,-j}A + \left[ (I B_{MM}^\top)^{-1} \mathcal{N}(0,I_p) \right]_{-j}$ .

Here the true adjacency matrix of mediators is generated from the Erdős-Rényi (ER) model with an expected degree as  $\lfloor p/2 \rfloor$ , and the non-zero entries in  $B_{MM} \in \mathbb{R}^{p \times p}$  and all the elements in  $\alpha_{YA}, \beta_{MA}, \beta_{YC}, \beta_{YM}, B_{MC}$  are independently sampled from U(-1,1). In each estimation method, we consistently treat  $\mathcal{M}_0$  as the underlying true model by default. We generate n=1000 simulation samples, each comprising N=100 independent observations, and the result for estimating the direct and indirect interventional effect of the prespecified mediator  $M_j$  is shown in Table 1. Here, we use the PC algorithm (Harris and Drton, 2013) to estimate the adjacency matrix of CPDAGs.

As illustrated in Table 1, the simulation results align with the theoretical predictions made in previous sections. Specifically, when the entire distribution  $F_X(\cdot)$  is correctly specified, all estimators display consistency. However, in the presence of at least one misspecified component, only the quadruply robust estimator retains consistency. In contrast, one among the other estimators,  $\mathcal{M}_\ell$  for  $\ell=0,1,2,3$ , becomes inconsistent. Although we present only the continuous scenario in this part, our simulations under discrete C or M settings yielded similar outcomes. Importantly, under this simulation scenario, the estimator  $\widehat{TM}_j^{\mathcal{M}_0} = \widehat{DM}_j^{\mathcal{M}_0} + \widehat{IM}_j^{\mathcal{M}_0}$  corresponds precisely to the estimator utilized for the individual mediation effect  $\eta_j$  proposed in Chakrabortty et al. (2018). Thus, our quadruply robust estimators outperform the estimator defined in Chakrabortty et al. (2018).

		all correct	$\mathcal{M}_0$ is correct	$\mathcal{M}_{j,1}$ is correct	$\mathcal{M}_{j,2}$ is correct	$\mathcal{M}_{j,3}$ is correct
direct	$\mathcal{M}_0$	0.006 (0.009)	0.004 (0.004)	0.187 (0.069)	0.032 (0.038)	0.014 (0.001)
indirect		0.033 (0.028)	0.041 (0.033)	1.930 (0.027)	0.028 (0.035)	0.001 (0.002)
direct	$\mathcal{M}_1$	0.227 (0.074)	0.064 (0.069)	0.234 (0.485)	0.899 (0.087)	0.383 (0.087)
indirect		0.683 (0.344)	0.756 (0.395)	0.418 (0.511)	2.214 (0.131)	1.675 (0.140)
direct	$-M_2$	0.007 (0.009)	0.004 (0.004)	0.187 (0.068)	0.032 (0.038)	0.014 (0.001)
indirect		0.853 (0.081)	0.914 (0.104)	2.040 (0.134)	0.237 (0.102)	1.292 (0.095)
direct	- M3	0.038 (0.039)	0.045 (0.057)	0.229 (0.179)	0.033 (0.042)	0.050 (0.062)
indirect		0.176 (0.128)	0.068 (0.082)	0.127 (0.156)	1.030 (0.079)	0.069 (0.086)
direct	- QR	0.031 (0.043)	0.008 (0.012)	0.145 (0.212)	0.054 (0.065)	0.025 (0.031)
indirect		0.124 (0.118)	0.087 (0.109)	0.281 (0.571)	0.053 (0.064)	0.024 (0.030)

Table 1 The average Bias (Standard Error) under simulation with sample size n=1000 under N=100 replications.

- 8.2. Simulation for all mediators. Next, we consider the average performance of our quadruply robust estimators compared with other estimations under a fair model misspecification scenario in both continuous case (Section 1.4 in Kang and Schafer (2007)) and discrete case (Section 4.1 in Xia and Chan (2023)). The DGPs are defined as follows:
- Continuous  $M: Z \leftarrow \mathcal{N}(0, I_{t-1}), A \leftarrow \mathbb{1}\{U[0, 1] \leq \Phi(\beta_{AC}^{\top}Z)\},\$

$$M \leftarrow B_{MC}^{\top} Z + \beta_{MA} A + B_{MM}^{\top} M + \mathcal{N} \left( 0, \begin{bmatrix} \sigma_1^2 \\ \ddots \\ \sigma_p^2 \end{bmatrix} \right),$$

and 
$$Y \leftarrow \beta_{YC}^{\top} Z + (\alpha_{YA} A + \beta_{YM}^{\top} M)^{2/3} + \mathcal{N}(0, 1);$$

• Discrete  $M: C \leftarrow \mathcal{N}(0, I_{t-1}), A \leftarrow \mathbb{1}\{U[0, 1] \leq \Phi(\beta_{AC}^{\top}C)\},\$ 

$$M_j \leftarrow \mathbb{1}\left\{ \text{logit}(U[0,1]) \leq \Theta_{MC,j:}C + \theta_{MA,j}A + \left[ (I - B_{MM}^\top)^{-1} \mathcal{N} \left( 0, \begin{bmatrix} \sigma_1^2 \\ & \ddots \\ & \sigma_p^2 \end{bmatrix} \right) \right]_j \right\},$$

$$\text{ and } Y \leftarrow \beta_{YC}^\top C + \alpha_{YA} A + \beta_{YM}^\top M + \beta_{YC}^\top AC + \beta_{YM}^\top AM + \mathcal{N}(0,1).$$

Here  $\sigma_1^2,\ldots,\sigma_p^2$  are independently drawn from the uniform distribution in [0.5,1], whereas the other setting is the same as previous. In the continuous setting, instead of observing the  $Z_i$ 's, we observe  $C_i$  as the transformations of  $Z_i$ . We will always leave out the interaction and  $x^{2/3}$  when fitting each model, and we also assume the link functions are all Probit. For computations in the continuous M setting, we implement Algorithm 2. While in the discrete M context, we employ Algorithm 1, setting the Monte Carlo sample size to L=100.

As illustrated in Figure 1, aside from the quadruply estimators (QR), other methods fail to yield consistent results. Furthermore, in most cases, our quadruply estimators exhibit a lower standard error compared to other methods. Thus, this also shows the robustness of our estimators.




FIG 1. Simulation results for both continuous (top row) and discrete (bottom row) scenarios, showcasing the estimated average causal direct mediation effect (left column) and indirect mediation effect (right column) over N=100 replications. The dots represent estimated averages, error bars detail the standard error derived from the replications, and the red line represents the true average value, calculated via numeric integrals according to Definition 3.2.

**9. Empirical Study.** In this section, we illustrate our estimator in a real world application from AURORA study to explore the causal association of psychiatric disorders among trauma survivors, which is also studied in Watson et al. (2023). In the study, our primary response of interest is the post-traumatic stress disorder (PTSD), which was assessed three months post-trauma Y. The focal event, in this case, is the pre-trauma insomnia A that trauma survivors often experience: A = 1 represents survivor does have insomnia




FIG 2. Estimated DAG of mediators

and A=0 represents does not. The 4-dimensional potential mediator M including Peritraumatic PT (PTSD), stress, acute distress (ASD), and depression, gauged two weeks subsequent to the traumatic incident, are included in our analysis. This study also accounts for various confounders C is a 9-dimensional vector such as age, gender, race, education level, pre-trauma physical and mental health, perceived stress level, neuroticism, and childhood trauma. The same as Watson et al. (2023), before employing our methodology, categorical variables underwent one-hot encoding, numerical variables were centralized, and any missing data was excluded. The total number of observations is n=1494 with t=10 and p=4. The estimated DAG of the mediators by PC algorithm (Harris and Drton, 2013) is shown in Figure 2. Results from the quadruply robust estimators, as obtained using Algorithm 1 with Monte Carlo sample size L=100 and a bootstrap number of B=500, along with other estimation methods employing a  $\log n$  truncation and the same Monte Carlo sample size and bootstrap number, are presented in Table 2.

Table 2
Estimated direct and indirect interventional effects of mediators using Algorithm 1. Values in bold denote statistical significance at the 95% confidence level.

		$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathscr{M}_2$	$\mathcal{M}_3$	QR
distress	direct	0.012 (0.022)	0.564 (17.827)	0.004 (0.677)	0.555 (144.040)	0.080 (0.055)
	indirect	0.004 (0.029)	-12.970 (146.383)	-13.672 (1150.637)	-14.021 (246.881)	0.101 (0.089)
ASD	direct	0.134 (0.112)	-3.284 (33.808)	0.043 (6.680)	0.557 (162.481)	0.166 (0.140)
	indirect	0.810 (0.633)	-8.846 (149.805)	-14.406 (2454.382)	-13.265 (264.681)	0.781 (0.278)
PTSD	direct	0.604 (0.358)	0.120 (22.549)	0.194 (96.854)	0.711 (164.921)	0.737 (0.278)
	indirect	0.004 (0.164)	-12.742 (153.783)	-14.883 (2367.793)	-13.630 (313.450)	0.012 (0.093)
depression	direct	0.104 (0.087)	-3.410 (56.095)	0.034 (5.875)	1.108 (142.886)	0.134 (0.105)
	indirect	0.120 (0.280)	-8.307 (148.382)	-13.731 (1522.495)	-14.517 (247.827)	0.198 (0.108)

As demonstrated in Table 2, for each mediator under consideration, a substantial discrepancy is observed between the estimates of  $\mathcal{M}_{\ell_1}$  and those of  $\mathcal{M}_{\ell_2}$  for  $\ell_2 \neq \ell_1$  when employ-

ing any of the four estimation methods  $\mathcal{M}_\ell$  for  $\ell=0,1,2,3$ . Moreover, none of these estimation methods manage to identify significant direct or indirect interventional effects for any of the mediators. This highlights the pressing need for robust estimation approaches in this dataset. Notably, with the quadruply robust estimation, we discern that both the indirect interventional effect of acute distress and the direct interventional effect of peritraumatic PT are significant at the 95% confidence level, while other effects remain non-significant. These results also indicate that preventive intervention of 3-month PTSD after trauma exposure that focuses on reducing acute distress and peritraumatic PT is more likely to be effective for trauma survivors.

10. Discussion. The main contribution of this article is the introduction of direct and indirect interventional effects of mediators, alongside their semiparametric bounds and quadruply robust estimators. Our method accommodates continuous, categorical, and multivariate pre-treatments, mediators, and outcomes. Moreover, extending our methodology and theory to polytomous exposures is straightforward. However, extending to continuous exposure, even under LSEMs, is non-trivial in theoretical sense. A potential method is suggested in Cai et al. (2021) to replace the indicator function  $\mathbb{1}(A=a)$  with some kernel function K((A-a)/h) under bandwidth h, but as discussed in Díaz and van der Laan (2013); Kennedy et al. (2017, 2023), pathwise differentiability will fail in this case, necessitating alternative estimation procedures and techniques. On the other hand, note that our framework is dimensional-free, as long as the conditional densities and expectations meet the mild convergence rate, our quadruply robust estimations will always achieve semiparametric efficiency. However, in high-dimensional cases, non-parametric estimation mentioned in this article may not achieve the rate, necessitating additional assumptions like symmetry and shape constraints, as discussed in Deng et al. (2021); Xu and Samworth (2021); Rodríguez-Casal and Saavedra-Nieves (2022). Introducing these assumptions still validates the semiparametric framework in our article under the full nonparametric model M<sub>nonpar</sub>, but our quadruply estimators may not be the most efficient under these added conditions.

## **REFERENCES**

- Albert, J. M. and S. Nelson (2011). Generalized causal mediation analysis. *Biometrics* 67(3), 1028–1038.
- An, W. and T. J. VanderWeele (2022). Opening the blackbox of treatment interference: Tracing treatment diffusion through network analysis. *Sociological Methods & Research* 51(1), 141–164.
- Assaad, C. K., E. Devijver, and E. Gaussier (2022). Survey and evaluation of causal discovery methods for time series. *Journal of Artificial Intelligence Research* 73, 767–819.
- Bang, H. and J. M. Robins (2005). Doubly robust estimation in missing data and causal inference models. *Biometrics* 61(4), 962–973.
- Bhattacharya, R., R. Nabi, and I. Shpitser (2022). Semiparametric inference for causal effects in graphical models with hidden variables. *Journal of Machine Learning Research* 23, 1–76.
- Bickel, P. J. and J. Kwon (2001). Inference for semiparametric models: some questions and an answer. *Statistica Sinica*, 863–886.
- Boca, S. M., R. Sinha, A. J. Cross, S. C. Moore, and J. N. Sampson (2014). Testing multiple biological mediators simultaneously. *Bioinformatics* 30(2), 214–220.
- Boos, D. D. and L. A. Stefanski (2013). Essential statistical inference: theory and methods, Volume 591. Springer.

- Brand, J. E., X. Zhou, and Y. Xie (2023). Recent developments in causal inference and machine learning. Annual Review of Sociology 49.
- Cai, H., C. Shi, R. Song, and W. Lu (2021). Deep jump learning for off-policy evaluation in continuous treatment settings. *Advances in Neural Information Processing Systems* 34, 15285–15300.
- Cai, H., R. Song, and W. Lu (2020). Anoce: Analysis of causal effects with multiple mediators via constrained structural learning. In *International Conference on Learning Representations*.
- Card, D. (1999). The causal effect of education on earnings. In *Handbook of labor economics*, Volume 3, pp. 1801–1863. Elsevier.
- Chakrabortty, A., P. Nandy, and H. Li (2018). Inference for individual mediation effects and interventional effects in sparse high-dimensional causal graphical models. *arXiv preprint arXiv:1809.10652*.
- Chan, K. C. G., S. C. P. Yam, and Z. Zhang (2016). Globally efficient non-parametric inference of average treatment effects by empirical balancing calibration weighting. *Journal of the Royal Statistical Society*. Series B, Statistical methodology 78(3), 673.
- Chen, L., C. Li, X. Shen, and W. Pan (2023). Discovery and inference of a causal network with hidden confounding. *Journal of the American Statistical Association* (just-accepted), 1–27.
- Chernozhukov, V., D. Chetverikov, and K. Kato (2014). Gaussian approximation of suprema of empirical processes. The Annals of Statistics, 1564–1597.
- Chickering, D. M. (2002). Optimal structure identification with greedy search. *Journal of machine learning research* 3(Nov), 507–554.
- Chinot, G., G. Lecué, and M. Lerasle (2020). Robust statistical learning with lipschitz and convex loss functions. *Probability Theory and related fields* 176(3-4), 897–940.
- Daniel, R. M., B. L. De Stavola, S. N. Cousens, and S. Vansteelandt (2015). Causal mediation analysis with multiple mediators. *Biometrics* 71(1), 1–14.
- Deng, H., Q. Han, and C.-H. Zhang (2021). Confidence intervals for multiple isotonic regression and other monotone models. *The Annals of Statistics* 49(4), 2021–2052.
- Díaz, I. and M. J. van der Laan (2013). Targeted data adaptive estimation of the causal dose–response curve. *Journal of Causal Inference 1*(2), 171–192.
- Fan, J. and Q. Yao (2003). *Nonlinear time series: nonparametric and parametric methods*, Volume 20. Springer. Goetgeluk, S., S. Vansteelandt, and E. Goetghebeur (2008). Estimation of controlled direct effects. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 70(5), 1049–1066.
- Guo, X., R. Li, J. Liu, and M. Zeng (2023). Statistical inference for linear mediation models with high-dimensional mediators and application to studying stock reaction to covid-19 pandemic. *Journal of Econometrics* 235(1), 166–179.
- Hall, P. (1988). On symmetric bootstrap confidence intervals. *Journal of the Royal Statistical Society Series B:* Statistical Methodology 50(1), 35–45.
- Harris, N. and M. Drton (2013). Pc algorithm for nonparanormal graphical models. *Journal of Machine Learning Research* 14(11).
- Hasan, U., E. Hossain, and M. O. Gani (2023). A survey on causal discovery methods for i.i.d. and time series data. *Transactions on Machine Learning Research*. Survey Certification.
- Heckman, J. J. (1976). The common structure of statistical models of truncation, sample selection and limited dependent variables and a simple estimator for such models. In *Annals of economic and social measurement, volume 5, number 4*, pp. 475–492. NBER.
- Hernán, M. A. (2004). A definition of causal effect for epidemiological research. *Journal of Epidemiology & Community Health* 58(4), 265–271.
- Hernán, M. Á., B. Brumback, and J. M. Robins (2000). Marginal structural models to estimate the causal effect of zidovudine on the survival of hiv-positive men. *Epidemiology*, 561–570.
- Huang, Y.-T. and W.-C. Pan (2016). Hypothesis test of mediation effect in causal mediation model with highdimensional continuous mediators. *Biometrics* 72(2), 402–413.
- Imai, K., L. Keele, and D. Tingley (2010). A general approach to causal mediation analysis. *Psychological methods* 15(4), 309.
- Kalisch, M. and P. Bühlmann (2007). Estimating high-dimensional directed acyclic graphs with the pcalgorithm. *Journal of Machine Learning Research* 8(Mar), 613–636.

- Kang, J. D. and J. L. Schafer (2007). Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statistical Science* 22(4), 523–539.
- Kennedy, E., S. Balakrishnan, and L. Wasserman (2023). Semiparametric counterfactual density estimation. Biometrika, asad017.
- Kennedy, E. H., Z. Ma, M. D. McHugh, and D. S. Small (2017). Non-parametric methods for doubly robust estimation of continuous treatment effects. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 79(4), 1229–1245.
- Kim, C., M. J. Daniels, J. W. Hogan, C. Choirat, and C. M. Zigler (2019). Bayesian methods for multiple mediators: Relating principal stratification and causal mediation in the analysis of power plant emission controls. *The annals of applied statistics* 13(3), 1927.
- Kuroki, M. and M. Miyakawa (1999). Identifiability criteria for causal effects of joint interventions. *Journal of the Japan Statistical Society* 29(2), 105–117.
- Laan, M. J. and J. M. Robins (2003). Unified methods for censored longitudinal data and causality. Springer. Lee, S. and V. Honavar (2020). Towards robust relational causal discovery. In Uncertainty in Artificial Intelligence, pp. 345–355. PMLR.
- Li, C., X. Shen, and W. Pan (2019). Likelihood ratio tests for a large directed acyclic graph. *Journal of the American Statistical Association*.
- Li, C., X. Shen, and W. Pan (2023). Inference for a large directed acyclic graph with unspecified interventions. *Journal of Machine Learning Research* 24(73), 1–48.
- Lin, S.-H. and T. VanderWeele (2017). Interventional approach for path-specific effects. *Journal of Causal Inference* 5(1).
- Maathuis, M. H., M. Kalisch, P. Bühlmann, et al. (2009). Estimating high-dimensional intervention effects from observational data. *The Annals of Statistics* 37(6A), 3133–3164.
- Nandy, P., A. Hauser, and M. H. Maathuis (2018). High-dimensional consistency in score-based and hybrid structure learning. *The Annals of Statistics* 46(6A), 3151–3183.
- Nandy, P., M. H. Maathuis, T. S. Richardson, et al. (2017). Estimating the effect of joint interventions from observational data in sparse high-dimensional settings. *The Annals of Statistics* 45(2), 647–674.
- Neal, B. (2020). Introduction to causal inference from a machine learning perspective. *Course Lecture Notes* (*draft*).
- Panizza, U. and A. F. Presbitero (2014). Public debt and economic growth: is there a causal effect? *Journal of Macroeconomics* 41, 21–41.
- Park, S. and E. Kürüm (2018). Causal mediation analysis with multiple mediators in the presence of treatment noncompliance. Statistics in medicine 37(11), 1810–1829.
- Pearl, J. (2000). Causality: models, reasoning and inference, Volume 29. Springer.
- Pearl, J. (2012). The causal mediation formula—a guide to the assessment of pathways and mechanisms. *Prevention science 13*, 426–436.
- Pearl, J. (2014). Interpretation and identification of causal mediation. Psychological methods 19(4), 459.
- Pearl, J. et al. (2009). Causal inference in statistics: An overview. Statistics surveys 3, 96-146.
- Petersen, M. L., S. E. Sinisi, and M. J. van der Laan (2006). Estimation of direct causal effects. *Epidemiology*, 276–284.
- Portnoy, S. (1984). Asymptotic behavior of m-estimators of p regression parameters when  $p^2/n$  is large. i. consistency. *The Annals of Statistics*, 1298–1309.
- Portnoy, S. (1985). Asymptotic behavior of m estimators of p regression parameters when  $p^2/n$  is large; ii. normal approximation. The Annals of Statistics 13(4), 1403–1417.
- Preacher, K. J. and A. F. Hayes (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior research methods* 40(3), 879–891.
- Robins, J. M. and S. Greenland (1992). Identifiability and exchangeability for direct and indirect effects. *Epidemiology* 3(2), 143–155.
- Rodríguez-Casal, A. and P. Saavedra-Nieves (2022). A data-adaptive method for estimating density level sets under shape conditions. *The Annals of Statistics* 50(3), 1653–1668.
- Scharfstein, D. O., A. Rotnitzky, and J. M. Robins (1999). Adjusting for nonignorable drop-out using semi-parametric nonresponse models. *Journal of the American Statistical Association 94*(448), 1096–1120.

- Schmidt-Hieber, J. (2020). Nonparametric regression using deep neural networks with relu activation function. The Annals of Statistics 48(4), 1875–1897.
- Shi, C. and L. Li (2021). Testing mediation effects using logic of boolean matrices. *Journal of the American Statistical Association*, 1–14.
- Shi, C. and L. Li (2022). Testing mediation effects using logic of boolean matrices. *Journal of the American Statistical Association* 117(540), 2014–2027.
- Shi, X., W. Miao, J. C. Nelson, and E. J. Tchetgen Tchetgen (2020). Multiply robust causal inference with double-negative control adjustment for categorical unmeasured confounding. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 82(2), 521–540.
- Shimizu, S., P. O. Hoyer, A. Hyvärinen, and A. Kerminen (2006). A linear non-gaussian acyclic model for causal discovery. *Journal of Machine Learning Research* 7(Oct), 2003–2030.
- Spirtes, P., C. Glymour, R. Scheines, S. Kauffman, V. Aimale, and F. Wimberly (2000). Constructing bayesian network models of gene expression networks from microarray data.
- Steen, J., T. Loeys, B. Moerkerke, and S. Vansteelandt (2017). Flexible mediation analysis with multiple mediators. American journal of epidemiology 186(2), 184–193.
- Sun, Q., W.-X. Zhou, and J. Fan (2020). Adaptive huber regression. Journal of the American Statistical Association 115(529), 254–265.
- Tai, A.-S., P.-H. Lin, Y.-T. Huang, and S.-H. Lin (2022). Path-specific effects in the presence of a survival outcome and causally ordered multiple mediators with application to genomic data. Statistical Methods in Medical Research 31(10), 1916–1933.
- Tchetgen, E. J. T. and I. Shpitser (2012). Semiparametric theory for causal mediation analysis: efficiency bounds, multiple robustness, and sensitivity analysis. *Annals of statistics* 40(3), 1816.
- Tchetgen Tchetgen, E. J. (2013). Inverse odds ratio-weighted estimation for causal mediation analysis. *Statistics in medicine* 32(26), 4567–4580.
- Tsiatis, A. A. (2006). Semiparametric theory and missing data.
- Van de Geer, S., P. Bühlmann, Y. Ritov, and R. Dezeure (2014). On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics* 42(3), 1166–1202.
- Van Der Laan, M. J. and D. Rubin (2006). Targeted maximum likelihood learning. *The international journal of biostatistics* 2(1).
- Van Der Vaart, A. W., J. A. Wellner, A. W. van der Vaart, and J. A. Wellner (1996). Weak convergence. Springer. VanderWeele, T. (2015). Explanation in causal inference: methods for mediation and interaction. Oxford University Press.
- VanderWeele, T. and S. Vansteelandt (2014). Mediation analysis with multiple mediators. Epidemiologic methods 2(1), 95–115.
- VanderWeele, T. J., J. W. Jackson, and S. Li (2016). Causal inference and longitudinal data: a case study of religion and mental health. *Social psychiatry and psychiatric epidemiology* 51, 1457–1466.
- VanderWeele, T. J. and W. R. Robinson (2014). On causal interpretation of race in regressions adjusting for confounding and mediating variables. *Epidemiology (Cambridge, Mass.)* 25(4), 473.
- Vansteelandt, S. and R. M. Daniel (2017). Interventional effects for mediation analysis with multiple mediators. *Epidemiology (Cambridge, Mass.)* 28(2), 258.
- Wager, S. and S. Athey (2018). Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association* 113(523), 1228–1242.
- Wainwright, M. J. (2019). High-dimensional statistics: A non-asymptotic viewpoint, Volume 48. Cambridge university press.
- Wang, L. and E. Tchetgen Tchetgen (2018). Bounded, efficient and multiply robust estimation of average treatment effects using instrumental variables. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 80(3), 531–550.
- Watson, R. A., H. Cai, X. An, S. McLean, and R. Song (2023). On heterogeneous treatment effects in heterogeneous causal graphs. *arXiv preprint arXiv:2301.12383*.
- Xia, F. and K. C. G. Chan (2023). Identification, semiparametric efficiency, and quadruply robust estimation in mediation analysis with treatment-induced confounding. *Journal of the American Statistical Associa*tion 118(542), 1272–1281.

- Xu, M. and R. J. Samworth (2021). High-dimensional nonparametric density estimation via symmetry and shape constraints. *The Annals of Statistics* 49(2), 650–672.
- Yu, Y., J. Chen, T. Gao, and M. Yu (2019). Dag-gnn: Dag structure learning with graph neural networks. arXiv preprint arXiv:1904.10098.
- Yuan, Y. and A. Qu (2023). De-confounding causal inference using latent multiple-mediator pathways. arXiv preprint arXiv:2302.05513.
- Yuan, Y., X. Shen, W. Pan, and Z. Wang (2019). Constrained likelihood for reconstructing a directed acyclic gaussian graph. *Biometrika* 106(1), 109–125.
- Zhang, C.-H. and S. S. Zhang (2014). Confidence intervals for low dimensional parameters in high dimensional linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 76(1), 217–242.
- Zhang, H., Y. Zheng, Z. Zhang, T. Gao, B. Joyce, G. Yoon, W. Zhang, J. Schwartz, A. Just, E. Colicino, et al. (2016). Estimating and testing high-dimensional mediation effects in epigenetic studies. *Bioinformatics* 32(20), 3150–3154.
- Zhang, J. and E. Bareinboim (2018). Non-parametric path analysis in structural causal models. In *Proceedings* of the 34th Conference on Uncertainty in Artificial Intelligence.
- Zhao, Y., L. Li, and A. D. N. Initiative (2022). Multimodal data integration via mediation analysis with high-dimensional exposures and mediators. *Human Brain Mapping* 43(8), 2519–2533.
- Zhao, Y. and X. Luo (2022). Pathway lasso: pathway estimation and selection with high-dimensional mediators. *Statistics and its interface 15*(1), 39.

### APPENDIX A: GLOSSARY OF TERMS AND NOTATIONS

Symbol	Definition	Symbol	Definition
X	Observed variables	$\mathcal{G}$	Causal DAG (or corresponding adjacency matrix) of X
C	Confounders	С	Causal CPDAG (or corresponding adjacency matrix) of X
A	0-1 exposure	$\mathrm{MEC}(\mathcal{C})$	Adjacency matrix in the Markov equivalence class of $C$
M	Mediators	$\mathcal{G}_{M}$	Causal DAG (or corresponding adjacency matrix) of $M$
Y	Univariate outcome	$\mathcal{C}_{M}$	$ \begin{array}{cccc} \text{Causal} & \text{CPDAG} & \text{(or} \\ \text{corresponding} & \text{ad-} \\ \text{jacency} & \text{matrix)} & \text{of} \\ M \end{array} $
$   \operatorname{Pa}_{j}(\mathcal{G}_{M}), \operatorname{pa}_{j}(\mathcal{G}_{M}) $	Parents of $M_j$ in mediators $\{M_1, \ldots, M_p\}$ , its realization	$e_{a'}(x_S)$	$P(A = a' \mid X_S = x_S)$

$\mu(oldsymbol{\cdot})$	Conditional expectation of Y given	$\pi.(m_T)$	Conditional density of $M_T$ given $\cdot$ evaluated at $m_T$
$S^{ ext{eff, nonpar}}(oldsymbol{\cdot})$	Efficient score for • on full model	$\mathbb{P}_n[oldsymbol{\cdot}]$	$\frac{1}{n}\sum_{i=1}^{n}[\cdot]_{i}$
$\kappa(a',C)$	Formula (4.1)	$\zeta_j(a',a,C)$	Formula (4.2)
$arrho_j(a',M_j,C;\mathcal{G}_M)$	Formula (4.3)	$\langle g_{\cdot,u}(\cdot,v)\rangle$	Difference in $g_{\cdot,u}(\cdot,v)$ evaluated at different exposure defined in Equation (4.4)
$ au_{:S}(C,a',M_T)$	Formula (4.5)		

### APPENDIX B: EXPLICIT FORMULAS FOR ALGORITHM 1 AND 2

We first give the formulas in Algorithm 1, the formulas in Algorithm 2 will be shown in the proof of Proposition 6.1 later in this section. For calculating the formulas in Algorithm 1, by Monte Carlo approximation, we have

$$\begin{split} \widehat{\zeta}_{j}^{\mathcal{M}_{0}}(a',0,C) &= \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \widehat{\pi}_{C,a'}(m_{j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m \\ &= \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \frac{\widehat{\pi}_{C,a'}(m_{j}) \widehat{\pi}_{C,0}(m_{-j})}{\widehat{\pi}_{C,1}(m)} \widehat{\pi}_{C,1}(m) \, \mathrm{d}m \\ &\approx \frac{1}{N} \sum_{\ell=1}^{N} \widehat{\mu}(C,1,M_{\widehat{\pi}_{C,0}}^{(\ell)}) \frac{\widehat{\pi}_{C,a'}(M_{\widehat{\pi}_{C,0},j}^{(\ell)}) \widehat{\pi}_{C,0}(M_{\widehat{\pi}_{C,0},-j}^{(\ell)})}{\widehat{\pi}_{C,1}(M_{\widehat{\pi}_{C,1}}^{(\ell)})}. \end{split}$$

where the sampling density  $\widehat{\pi}_{C,1}(m)$  is calculated by  $\widehat{\pi}_{C,1}(m) = \widehat{\pi}_{C,1}(m_j)\widehat{\pi}_{C,1,m_j}(m_{-j})$  and does not need to estimate additionally. Similarly, we have

$$\langle \widehat{\tau}_{C,\cdot;j}(C,1,M_{-j}) \rangle \approx \frac{1}{N} \sum_{\ell=1}^{N} \left[ \widehat{\mu}(C,1,M_{j,\widehat{\pi}_{C,1}}^{(\ell)},M_{-j}) - \widehat{\mu}(C,1,M_{j,\widehat{\pi}_{C,0}}^{(\ell)},M_{-j}) \right]$$

and

$$\widehat{\tau}_{C,0;-j}(C,1,M_j) \approx \frac{1}{N} \sum_{\ell=1}^{N} \widehat{\mu}(C,1,M_j,M_{-j,\widehat{\pi}_{C,0}}^{(\ell)}).$$

Note that we can rewrite  $\widehat{DM}_j^{\mathrm{QR}} = \frac{1}{n} \sum_{i=1}^n \widehat{QR}_{DM_i}(X_i)$  with

$$\begin{split} \widehat{QR}_{DM_j}(X) &:= \frac{\mathbbm{1}(A=1)}{\widehat{e}_1(C)} \frac{\left\langle \widehat{\pi}_{C,\cdot}(M_{-j}) \right\rangle}{\widehat{\pi}_{C,1,M_j}(M_{-j})} \big[ Y - \widehat{\mu}(C,1,M) \big] + \frac{\mathbbm{1}(A=0)}{\widehat{e}_0(C)} \left\langle \widehat{\tau}_{C,\cdot;j}(C,1,M_{-j}) \right\rangle \\ &+ \left\langle \frac{\mathbbm{1}(A=\cdot)}{\widehat{e}_0(C)} \right\rangle \widehat{\tau}_{C,0;\,-j}(C,1,M_j) + \left[ 1 - \frac{\mathbbm{1}(A=0)}{\widehat{e}_0(C)} - \frac{\mathbbm{1}(A=1)}{\widehat{e}_1(C)} \right] \widehat{\zeta}_j^{\mathcal{M}_0}(1,0,C) \\ &+ \left[ 2 \frac{\mathbbm{1}(A=0)}{\widehat{e}_0(C)} - 1 \right] \widehat{\zeta}_j^{\mathcal{M}_0}(0,0,C). \end{split}$$

Therefore, we have the approximation as  $\widehat{DM}_j^{QR} \approx \frac{1}{n} \sum_{i=1}^n \widehat{QR}_{DM_j}^{MC}(X_i)$  with the following explicit formula (B.1)

$$\begin{split} \widehat{QR}_{DM_{j}}^{\text{MC}}(X) &:= \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \bigg[ \frac{\widehat{\pi}_{C,1}(M_{-j}) - \widehat{\pi}_{C,0}(M_{-j})}{\widehat{\pi}_{C,1,M_{j}}(M_{-j})} \bigg] \big\{ Y - \widehat{\mu}(C,1,M) \big\} \\ &+ \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \frac{1}{N} \sum_{\ell=1}^{N} \bigg[ \widehat{\mu}(C,1,M_{j,\widehat{\pi}_{C,1}}^{(\ell)},M_{-j}) - \widehat{\mu}(C,1,M_{j,\widehat{\pi}_{C,0}}^{(\ell)},M_{-j}) \bigg] \\ &+ \bigg[ \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} - \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \bigg] \frac{1}{N} \sum_{\ell=1}^{N} \widehat{\mu}(C,1,M_{j},M_{-j,\widehat{\pi}_{C,0}}^{(\ell)}) \\ &+ \bigg[ 1 - \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} - \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \bigg] \frac{1}{N} \sum_{\ell=1}^{N} \widehat{\mu}(C,1,M_{\widehat{\pi}_{C,0}}^{(\ell)}) \frac{\widehat{\pi}_{C,1}(M_{\widehat{\pi}_{C,0},j}^{(\ell)}) \widehat{\pi}_{C,0}(M_{\widehat{\pi}_{C,0},-j}^{(\ell)})}{\widehat{\pi}_{C,1}(M_{\widehat{\pi}_{C,0},-j}^{(\ell)})} \\ &+ \bigg[ 2 \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} - 1 \bigg] \frac{1}{N} \sum_{\ell=1}^{N} \widehat{\mu}(C,1,M_{\widehat{\pi}_{C,0}}^{(\ell)}) \frac{\widehat{\pi}_{C,0}(M_{\widehat{\pi}_{C,0},j}^{(\ell)}) \widehat{\pi}_{C,0}(M_{\widehat{\pi}_{C,0},-j}^{(\ell)})}{\widehat{\pi}_{C,1}(M_{\widehat{\pi}_{C,1}}^{(\ell)})}. \end{split}$$

Similarly, note that the double integral can be rewritten as

$$\begin{split} \mathrm{E}\big[\varrho_{j}(a',M_{j},C\,;\,\mathcal{G}_{M})\,|\,C\big] &= \int \mu\big(a',m_{j},\mathrm{pa}_{j}(\mathcal{G}_{M}),C\big)\pi_{C,a'}\big(\,\mathrm{pa}_{j}(\mathcal{G}_{M})\big)\pi_{C}(m_{j})\,\mathrm{d}(m_{j},\mathrm{pa}_{j}(\mathcal{G}_{M})) \\ &\approx \frac{1}{N^{2}}\sum_{\ell_{1}=1}^{N}\sum_{\ell_{2}=1}^{N}\mu\big(a',M_{j,\pi_{C}}^{(\ell_{1})},\mathrm{Pa}_{j,\pi_{C,a'},\mathcal{G}_{M}}^{(\ell_{2})},C\big). \end{split}$$

Thus, given the fact  $\widehat{TM}_j^{QR}(\mathcal{G}_M) = \frac{1}{n} \sum_{i=1}^n \widehat{QR}_{TM_i}(X_i;\mathcal{G}_M)$  with

$$\begin{split} \widehat{QR}_{TM_{j}}(X\,;\mathcal{G}_{M}) := & \left\langle \frac{\mathbb{1}(A=\cdot)}{\widehat{e}.(C)} \big[ Y - \widehat{\mu}(C,\cdot) \big] - \frac{\mathbb{1}(A=\cdot)}{\widehat{e}.(C)} \widehat{\pi}_{C,\cdot} \big( \operatorname{Pa}_{j}(\mathcal{G}_{M}) \big) \Big( Y - \widehat{\mu}(C,\cdot, \operatorname{Pa}_{j}(\mathcal{G}_{M}), M_{j}) \Big) \right\rangle \\ & - \left\langle \frac{\mathbb{1}(A=\cdot)}{\widehat{e}.(C)} \Big\{ \widehat{\tau}_{C\,;\,j} \big( C, 1, \operatorname{Pa}_{j}(\mathcal{G}_{M}) \big) - \operatorname{E} \big[ \widehat{\varrho}_{j}(\cdot, M_{j}, C\,;\,\mathcal{G}_{M}) \mid C \big] \right\} \right\rangle \\ & + \left\langle \widehat{\mu}(C,\cdot) - \widehat{\tau}_{C,\cdot\,;\,\operatorname{pa}_{j}(\mathcal{G}_{M})}(C,\cdot, M_{j}) \right\rangle, \end{split}$$

we have the approximation  $\widehat{DM}_j^{\rm QR} \approx \frac{1}{n} \sum_{i=1}^n \widehat{QR}_{TM_j}^{\rm MC}(X_i)$ , where (B.2)

$$\begin{split} \widehat{QR}_{TM_{j}}^{\text{MC}}(X;\mathcal{G}_{M}) := & \left[ \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \Big\{ Y - \widehat{\mu}(C,1) \Big\} - \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \Big\{ Y - \widehat{\mu}(C,0) \Big] \Big\} \right] \\ & - \left[ \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \widehat{\pi}_{C,1} \Big( \operatorname{Pa}_{j}(\mathcal{G}_{M}) \Big) \Big\{ Y - \widehat{\mu}(C,1,\operatorname{Pa}_{j}(\mathcal{G}_{M}),M_{j}) \Big\} \\ & - \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \widehat{\pi}_{C,0} \Big( \operatorname{Pa}_{j}(\mathcal{G}_{M}) \Big) \Big\{ Y - \widehat{\mu}(C,0,\operatorname{Pa}_{j}(\mathcal{G}_{M}),M_{j}) \Big\} \\ & + \frac{1}{N^{2}} \sum_{\ell_{1}=1}^{N} \sum_{\ell_{2}=1}^{N} \Big[ \widehat{\mu}(C,1,M_{j,\widehat{\pi}_{C}}^{(\ell_{1})},\operatorname{Pa}_{j}(\mathcal{G}_{M})) - \widehat{\mu}(C,1,M_{j,\widehat{\pi}_{C}}^{(\ell_{1})},\operatorname{Pa}_{j,\widehat{\pi}_{C,1,\mathcal{G}_{M}}^{(\ell_{2})} \Big) \Big] \\ & - \frac{1}{N^{2}} \sum_{\ell_{1}=1}^{N} \sum_{\ell_{2}=1}^{N} \Big[ \widehat{\mu}(C,0,M_{j,\widehat{\pi}_{C}}^{(\ell_{1})},\operatorname{Pa}_{j}(\mathcal{G}_{M})) - \widehat{\mu}(C,0,M_{j,\widehat{\pi}_{C}}^{(\ell_{1})},\operatorname{Pa}_{j,\widehat{\pi}_{C,0,\mathcal{G}_{M}}^{(\ell_{2})},C) \Big] \Big] \\ & + \widehat{\mu}(C,1) - \widehat{\mu}(C,0) - \Big[ \frac{1}{N} \sum_{\ell=1}^{N} \Big[ \widehat{\mu}(C,1,\operatorname{Pa}_{j,\widehat{\pi}_{C,1,\mathcal{G}_{M}}^{(\ell)},M_{j}) - \widehat{\mu}(C,0,\operatorname{Pa}_{j,\widehat{\pi}_{C,0,\mathcal{G}_{M}}^{(\ell)},M_{j}) \Big]. \end{split}$$

## **Proof of Proposition 6.1:**

PROOF. Denote  $\phi(\cdot; \mu, \Sigma)$  as the density of distribution  $\mathcal{N}(\mu, \Sigma)$ . Similar to (6.10), we have

$$\begin{split} &\int_{\mathcal{M}_{j}}\mu(C,1,m_{j},M_{-j})\pi_{C,1}(m_{j})\,\mathrm{d}m_{j}\\ &=\int_{\mathcal{M}_{j}}\left[\beta_{YC}C+\alpha_{YA}+\beta_{YM,j}m_{j}+\beta_{YM,-j}^{\intercal}M_{-j}\right]\phi\Big(m_{j}\,;\,\left[\Theta_{MC}C\right]_{j}+\theta_{MA,j},\,\left[\mathrm{var}(e_{M})\right]_{jj}\Big)\,\mathrm{d}m_{j}\\ &=\beta_{YC}C+\alpha_{YA}+\beta_{YM,j}\Big\{\big[\widehat{\Theta}_{MC}C\big]_{j}\Big\}+\beta_{YM,-j}^{\intercal}M_{-j}, \end{split}$$

thus

$$\begin{split} \left\langle \tau_{C,\cdot\,;j}(C,1,M_{-j}) \right\rangle &= \left[ \beta_{YC}C + \alpha_{YA} + \beta_{YM,j} \Big\{ \left[ \Theta_{MC}C \right]_j + \theta_{MA,j} \Big\} + \beta_{YM,-j}^\top M_{-j} \right] \\ &- \left[ \beta_{YC}C + \alpha_{YA} + \beta_{YM,j} \left[ \Theta_{MC}C \right]_j + \beta_{YM,-j}^\top M_{-j} \right] = \beta_{YM,j} \theta_{MA,j}. \end{split}$$

Similarly, we have

$$\begin{split} &\int_{\mathcal{M}_{-j}} \mu(C, 1, M_j, m_{-j}) \pi_{C,0}(m_{-j}) \, \mathrm{d} m_{-j} \\ &= \int_{\mathcal{M}_j} \left[ \beta_{YC} C + \alpha_{YA} + \beta_{YM,j} M_j + \beta_{YM,-j}^\top m_{-j} \right] \phi \Big( m_{-j} \, ; \, \left[ \Theta_{MC} C \right]_{-j}, \, \left[ \operatorname{var}(e_M) \right]_{-j,-j} \Big) \, \mathrm{d} m_{-j} \\ &= \beta_{YC} C + \alpha_{YA} + \beta_{YM,j} M_j + \beta_{YM,-j}^\top \left[ \Theta_{MC} C \right]_{-j} \end{split}$$

and

$$\begin{split} &\zeta_{j}(a',0,C) \\ &= \int_{\mathcal{M}} \mu(C,1,m) \pi_{C,a'}(m_{j}) \pi_{C,0}(m_{-j}) \, \mathrm{d}m \\ &= \int_{\mathcal{M}} \left[ \beta_{YC}C + \alpha_{YA} + \beta_{YM,j} m_{j} + \beta_{YM,-j}^{\top} m_{-j} \right] \phi \left( m_{j} \, ; \, \left[ \Theta_{MC}C \right]_{j} + \theta_{MA,j} a', \, \left[ \operatorname{var}(e_{M}) \right]_{j,j} \right) \\ &\qquad \qquad \phi \left( m_{-j} \, ; \, \left[ \Theta_{MC}C \right]_{-j}, \, \left[ \operatorname{var}(e_{M}) \right]_{-j,-j} \right) \, \mathrm{d}m_{j} \, \mathrm{d}m_{-j} \\ &= \beta_{YC}C + \alpha_{YA} + \beta_{YM,j} \left\{ \left[ \Theta_{MC}C \right]_{j} + \theta_{MA,j} a' \right\} + \beta_{YM,-j}^{\top} \left[ \Theta_{MC}C \right]_{-j}. \end{split}$$

Therefore, under Assumption 5.1 and Gaussian assumption,

$$\widehat{DM}_{j}^{\mathrm{QR}} = \mathbb{P}_{n} \widehat{QR}_{DM_{j}}^{\mathrm{fast}}(X) + \widehat{DM}_{j}^{\mathrm{OLS}},$$

where the (estimated) quadruply adjustment terms  $\widehat{QR}_{DM_j}(X)$  is defined as (B.3)

$$\begin{split} \widehat{QR}_{DM_{j}}^{\text{fast}}(X) &= \widehat{QR}_{DM_{j}}^{\text{fast}}(C, A, M, Y) \\ &= \frac{\mathbb{1}(A = 1)}{\widehat{e}_{1}(C)} \left[ \frac{\phi\left(M_{-j}; \left[\widehat{\Theta}_{MC}C\right]_{-j} + \widehat{\theta}_{MA, -j}, \left[\widehat{\text{var}}(e_{M})\right]_{-j, -j}\right) - \phi\left(M_{-j}; \left[\widehat{\Theta}_{MC}C\right]_{-j}, \left[\widehat{\text{var}}(e_{M})\right]_{-j, -j}\right)}{\phi\left(M_{-j} \mid M_{j}; \left[\widehat{\Theta}_{MC}C + \widehat{\theta}_{MA}\right]_{-j \mid j}, \left[\widehat{\text{var}}(e_{M})\right]_{-j \mid j}\right)} \right] \\ &\times \left\{ Y - \left[\widehat{\beta}_{YC}C + \widehat{\alpha}_{YA} + \widehat{\beta}_{YM}^{\top}M\right] \right\} \\ &+ \left[ \frac{\mathbb{1}(A = 1)}{\widehat{e}_{1}(C)} - \frac{\mathbb{1}(A = 0)}{\widehat{e}_{0}(C)} \right] \left[\widehat{\beta}_{YC}C + \widehat{\alpha}_{YA} + \widehat{\beta}_{YM,j}M_{j} + \widehat{\beta}_{YM, -j}^{\top}[\widehat{\Theta}_{MC}C]_{-j}\right] \\ &- \left[ \frac{\mathbb{1}(A = 1)}{\widehat{e}_{1}(C)} \left[\widehat{\beta}_{YC}^{\top}C + \widehat{\alpha}_{YA} + \widehat{\beta}_{YM,j} \left\{ \left[\widehat{\Theta}_{MC}C\right]_{j} + \widehat{\theta}_{MA,j} \right\} + \widehat{\beta}_{YM, -j}^{\top}[\widehat{\Theta}_{MC}C]_{-j} \right] \right] \\ &- \frac{\mathbb{1}(A = 0)}{\widehat{e}_{2}(C)} \left[\widehat{\beta}_{YC}^{\top}C + \widehat{\alpha}_{YA} + \widehat{\beta}_{YM,j} \left[\widehat{\Theta}_{MC}C\right]_{j} + \widehat{\beta}_{YM, -j}^{\top}[\widehat{\Theta}_{MC}C]_{-j} \right] \right]. \end{split}$$

Similarly, we can show that

$$\begin{split} \pi_{C}(M_{j}) &= e_{0}(C)\pi_{C,0}(M_{j}) + e_{1}(C)\pi_{C,1}(M_{j}) \\ &= e_{0}(C)\phi\Big(m_{j}\,;\, \left[\Theta_{MC}C\right]_{j}, \left[\operatorname{var}(e_{M})\right]_{j,j}\Big) + e_{1}(C)\phi\Big(m_{j}\,;\, \left[\Theta_{MC}C\right]_{j} + \theta_{MA,j}, \left[\operatorname{var}(e_{M})\right]_{j,j}\Big), \\ \pi_{C,a'}\big(\operatorname{Pa}_{j}(\mathcal{G}_{M})\big) &= \phi\Big(\operatorname{Pa}_{j}(\mathcal{G}_{M})\,;\, \left[\Theta_{MC}C\right]_{k:k\in\operatorname{pa}_{j}(\mathcal{G}_{M})} + \theta_{MA,k\in\operatorname{pa}_{j}(\mathcal{G}_{M})}a', \left[\operatorname{var}(e_{M})\right]_{kk:k\in\operatorname{pa}_{j}(\mathcal{G}_{M})}\Big), \\ \tau_{C\,;\,j}\big(C,a',\operatorname{Pa}_{j}(\mathcal{G}_{M})\big) &= \int_{\mathcal{M}_{j}}\Big\{\eta_{YM_{j}}m_{j} + \gamma_{Y\operatorname{Pa}_{j}(\mathcal{G}_{M})}^{\top}\operatorname{Pa}_{j}(\mathcal{G}_{M}) + \eta_{YA}a' + \gamma_{YC}^{\top}C\Big\} \\ \Big\{e_{0}(C)\phi\Big(m_{j}\,;\, \left[\Theta_{MC}C\right]_{j}, \left[\operatorname{var}(e_{M})\right]_{j,j}\Big) + e_{1}(C)\phi\Big(m_{j}\,;\, \left[\Theta_{MC}C\right]_{j} + \theta_{MA,j}, \left[\operatorname{var}(e_{M})\right]_{j,j}\Big)\Big\}\,\mathrm{d}m_{j} \\ &= \gamma_{Y\operatorname{Pa}_{j}(\mathcal{G}_{M})}^{\top}\operatorname{Pa}_{j}(\mathcal{G}_{M}) + \eta_{YA}a' + \gamma_{YC}^{\top}C + \eta_{YM_{j}}\Big[e_{0}(C)\left[\Theta_{MC}C\right]_{j} + e_{1}(C)\left(\left[\Theta_{MC}C\right]_{j} + \theta_{MA,j}\right)\Big], \end{split}$$

and

$$\begin{split} & \operatorname{E}\left[\varrho_{j}(a',M_{j},C\,;\mathcal{G}_{M})\,|\,C\right] \\ &= \int_{\mathcal{M}_{\operatorname{pa}_{j}}(\mathcal{G}_{M})} \operatorname{dpa}_{j}(\mathcal{G}_{M}) \int_{\mathcal{M}_{j}} \mu\big(C,a',\operatorname{pa}_{j}(\mathcal{G}_{M}),m_{j}\big) \pi_{C,a'}\big(\operatorname{pa}_{j}(\mathcal{G}_{M})\big) \pi_{C}(m_{j}) \operatorname{d}m_{j} \\ &= \int_{\mathcal{M}_{\operatorname{pa}_{j}}(\mathcal{G}_{M})} \operatorname{dpa}_{j}(\mathcal{G}_{M}) \int_{\mathcal{M}_{j}} \left\{ \eta_{YM_{j}} m_{j} + \gamma_{Y}^{\top} \operatorname{Pa}_{j}(\mathcal{G}_{M}) \operatorname{pa}_{j} + \eta_{YA} a' + \gamma_{YC}^{\top} C \right\} \\ & \times \phi\Big(\operatorname{pa}_{j}\,;\, \left[\Theta_{MC}C\right]_{k:k\in\operatorname{pa}_{j}(\mathcal{G}_{M})} + \theta_{MA,k\in\operatorname{pa}_{j}(\mathcal{G}_{M})} a',\, \left[\operatorname{var}(e_{M})\right]_{kk:k\in\operatorname{pa}_{j}(\mathcal{G}_{M})} \big) \\ & \times \Big\{e_{0}(C)\phi\Big(m_{j}\,;\, \left[\Theta_{MC}C\right]_{j},\, \left[\operatorname{var}(e_{M})\right]_{j,j}\Big) + e_{1}(C)\phi\Big(m_{j}\,;\, \left[\Theta_{MC}C\right]_{j} + \theta_{MA,j},\, \left[\operatorname{var}(e_{M})\right]_{j,j}\Big) \Big\} \operatorname{d}m_{j} \\ &= \eta_{YA}a' + \gamma_{YC}^{\top}C + \gamma_{Y}^{\top} \operatorname{Pa}_{j}(\mathcal{G}_{M}) \Big\{ \left[\Theta_{MC}C\right]_{k:k\in\operatorname{pa}_{j}(\mathcal{G}_{M})} + \theta_{MA,k\in\operatorname{pa}_{j}(\mathcal{G}_{M})} a' \Big\} \\ &\quad + \eta_{YM_{j}} \Big\{e_{0}(C) \left[\Theta_{MC}C\right]_{j} + e_{1}(C) \left[\Theta_{MC}C\right]_{j} + e_{1}(C) \theta_{MA,j} \Big\}. \end{split}$$

Note that  $\mu(C, a') = \gamma_{YA}^{\dagger} a' + \eta_{YC}^{\dagger \top} C$ . Hence, we conclude

$$\widehat{TM}_{j}^{QR}(\mathcal{G}_{M}) = \mathbb{P}_{n}\widehat{QR}_{TM_{j}}^{fast}(X;\mathcal{G}_{M}) + \widehat{TM}_{j}^{OLS},$$

with the estimated adjustment term  $\widehat{QR}_{TM_j}^{\mathrm{fast}}(X;\mathcal{G}_M)$  defined as (B.4)

$$\begin{split} \widehat{QR}_{TM_{j}}^{\text{fast}}(X;\mathcal{G}_{M}) &= \left\lfloor \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \Big\{ Y - \widehat{\gamma}_{YA}^{\dagger} - \widehat{\eta}_{YC}^{\dagger\top}C \Big\} - \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \Big\{ Y - \widehat{\eta}_{YC}^{\dagger\top}C \Big\} \right\} \\ &- \left\lfloor \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \phi \Big( \operatorname{Pa}_{j}(\mathcal{G}_{M}); \left[ \widehat{\Theta}_{MC}C \right]_{k:k \in \operatorname{pa}_{j}(\mathcal{G}_{M})} + \widehat{\theta}_{MA,k \in \operatorname{pa}_{j}(\mathcal{G}_{M})}, \left[ \widehat{\operatorname{var}}(e_{M}) \right]_{kk:k \in \operatorname{pa}_{j}(\mathcal{G}_{M})} \right) \\ &\times \left\{ Y - \widehat{\eta}_{YM_{j}} M_{j} - \widehat{\gamma}_{Y}^{\top} \operatorname{Pa}_{j}(\mathcal{G}_{M}) \operatorname{Pa}_{j}(\mathcal{G}_{M}) - \widehat{\eta}_{YA}A - \widehat{\gamma}_{YC}^{\top}C \right\} \\ &- \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \phi \Big( \operatorname{Pa}_{j}(\mathcal{G}_{M}); \left[ \widehat{\Theta}_{MC}C \right]_{k:k \in \operatorname{pa}_{j}(\mathcal{G}_{M})}, \left[ \widehat{\operatorname{var}}(e_{M}) \right]_{kk:k \in \operatorname{pa}_{j}(\mathcal{G}_{M})} \right) \\ &\times \left\{ Y - \widehat{\eta}_{YM_{j}} M_{j} - \widehat{\gamma}_{Y}^{\top} \operatorname{Pa}_{j}(\mathcal{G}_{M}) \operatorname{Pa}_{j}(\mathcal{G}_{M}) - \widehat{\gamma}_{YC}^{\top}C \right\} \\ &+ \frac{\mathbb{1}(A=1)}{\widehat{e}_{1}(C)} \Big\{ \widehat{\gamma}_{Y}^{\top} \operatorname{Pa}_{j}(\mathcal{G}_{M}) \Big[ \operatorname{Pa}_{j}(\mathcal{G}_{M}) - \left[ \widehat{\Theta}_{MC}C \right]_{k:k \in \operatorname{pa}_{j}(\mathcal{G}_{M})} - \widehat{\theta}_{MA,k:k \in \operatorname{pa}_{j}(\mathcal{G}_{M})} \Big] \right\} \\ &- \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \Big\{ \widehat{\gamma}_{Y}^{\top} \operatorname{Pa}_{j}(\mathcal{G}_{M}) \Big[ \operatorname{Pa}_{j}(\mathcal{G}_{M}) - \left[ \widehat{\Theta}_{MC}C \right]_{k:k \in \operatorname{pa}_{j}(\mathcal{G}_{M})} \Big] \right\} \Big] \end{split}$$

where  $\phi(\cdot \mid X_{S_1}; \mu_{S_2 \mid S_1}, \Sigma_{S_2 \mid S_1})$  is the density for the conditional distribution  $X_{S_2} \mid X_{S_1}$  where  $(X_{S_1}^\top, X_{S_2}^\top)^\top \sim \mathcal{N}(\mu, \Sigma)$ . Finally, by  $\widehat{IM}_j^{QR}(\mathcal{G}_M) = \widehat{TM}_j^{QR}(\mathcal{G}_M) - \widehat{DM}_j^{QR}$ , we conclude the proposition when both linear structures hold in Assumption 5.1. Finally, it is noteworthy, as seen in Theorem 7.11, that even if only one linear assumption holds in Assumption 5.1, our estimator remains consistent.

## APPENDIX C: TECHNIQUE PROOFS

In the proof, to avoid any misunderstanding, we abbreviate the conditional density  $f_{Z_2|Z_1}(z_2 \mid z_1)$  as  $f(z_2 \mid z_1)$ , and omit the interval of the integral, assuming the interval is the support of the integrand variable.

#### C.1. The Proofs of Section 4.

#### **Proof of Theorem 4.1:**

PROOF. Fixing a DAG  $\mathcal{G}_M$ , we drop the DAG index in these quantities for simplicity. Note that the joint intervention density can be written as

$$\begin{split} f\big(y\mid do(A=a,\,M_j=m_j)\big) &= \int \frac{f(y,c,a,m_j,\mathrm{pa}_j)}{f(a\mid c)f(m_j\mid c,a,\mathrm{pa}_j)}\,\mathrm{d}c\,\mathrm{d}\,\mathrm{pa}_j \\ &= \int \frac{f(y,c,a,m_j,\mathrm{pa}_j)}{f(m_j,c,a,\mathrm{pa}_j)}\,\frac{f(c,a,\mathrm{pa}_j)}{f(a,c)}f(c)\,\mathrm{d}c\,\mathrm{d}\,\mathrm{pa}_j \\ &= \int f(y\mid c,a,m_j,\mathrm{pa}_j)f(\mathrm{pa}_j\mid c,a)f(c)\,\mathrm{d}c\,\mathrm{d}\,\mathrm{pa}_j, \end{split}$$

where the first equation is applying Pa(A) = C, and  $Pa(M_j) = (C, A, Pa_j)$  with Theorem 6 in Kuroki and Miyakawa (1999). Thus,

$$E[Y \mid do(A = a, M_j = m_j), C] = \int E[Y \mid A = a, M_j = m_j, Pa_j = pa_j, C] f(pa_j \mid a, C) dpa_j$$
$$= \varrho_j(a, m_j, C),$$

then

$$E[Y \mid do(A = 1, M_i), C = c] - E[Y \mid do(A = 0, M_i), C = c] = \varrho_i(1, M_i, c) - \varrho_i(0, M_i, c).$$

Denote

$$\begin{split} TM_{j}(c) := & \Big\{ \mathbf{E} \big[ Y \mid A = 1, C = c \big] - \mathbf{E} \big[ Y \mid A = 0, C = c \big] \Big\} \\ & - \Big\{ \mathbf{E} \big[ Y \mid do(A = 1, M_{j}), C = c \big] - \mathbf{E} \big[ Y \mid do(A = 0, M_{j}), C = c \big] \Big\}. \end{split}$$

Since  $TM_j = ETM_j(C)$ , we have

$$TM_{j} = E[TM_{j}(C)]$$

$$= \left\{ E[Y \mid A = 1, C] - E[Y \mid A = 0, C] \right\} - \left\{ \varrho_{j}(1, M_{j}, C) - \varrho_{j}(0, M_{j}, C) \right\}.$$

The formula for  $DM_i$  is just by the definition.

#### **Proof of Theorem 4.3:**

PROOF. Let  $F_{X;t}$  denote a one dimensional regular parametric submodel of  $\mathcal{M}_{\text{nonpar}}$ , with  $F_{X;0} = F_X$ , and let  $E_t$  be the expectation with respect to  $F_{X;t}$ . Denote U the score of  $F_{X;t}$  at t=0 and  $\nabla_{t=0}$  denoting differentiation with respect to t at t=0. For a fixed DAG  $\mathcal{G}_M$ , we drop  $\mathcal{G}_M$  in  $\varrho_j(a', M_j, C; \mathcal{G}_M)$  as  $\varrho_j(a', M_j, C)$ . Note that we can write

$$\begin{split} & & \quad \mathbf{E}\varrho_j(a',M_j,C) \\ & = \int f(m_j,c) \mathbf{d}(m_j,c) \int \mu(C,a',\mathbf{pa}_j,M_j) \, \pi_{C,a'}(\mathbf{pa}_j) \, \mathrm{d}\, \mathbf{pa}_j \\ & = \int \mathbf{E}\big[Y \mid A=a',M_j=m_j,\mathbf{Pa}_j=\mathbf{pa}_j,C=c\big] f(\mathbf{pa}_j \mid a',c) f(m_j \mid c) f(c) \, \mathrm{d}\, \mathbf{pa}_j \, \mathrm{d}m_j \, \mathrm{d}c. \end{split}$$

Denote  $d(c, pa_j, m_j) = d\mu$ , then we can get

$$\begin{split} \frac{\partial}{\partial t} & \mathbf{E}_{t} \varrho_{j}(a', M_{j}, C) \Big|_{t=0} \\ &= \int \nabla_{t=0} \mathbf{E}_{t} \big[ Y \mid A = a', M_{j} = m_{j}, \mathbf{P} \mathbf{a}_{j} = \mathbf{p} \mathbf{a}_{j}, C = c \big] f(\mathbf{p} \mathbf{a}_{j} \mid a', c) f(m_{j} \mid c) f(c) \mathrm{d}\mu \, \mathrm{d}F_{X, t} \\ &+ \int \mathbf{E} \big[ Y \mid A = a', M_{j} = m_{j}, \mathbf{P} \mathbf{a}_{j} = \mathbf{p} \mathbf{a}_{j}, C = c \big] \nabla_{t=0} f_{t}(\mathbf{p} \mathbf{a}_{j} \mid a', c) f(m_{j} \mid c) f(c) \, \mathrm{d}\mu \, \mathrm{d}F_{X, t} \\ &+ \int \mathbf{E} \big[ Y \mid A = a', M_{j} = m_{j}, \mathbf{P} \mathbf{a}_{j} = \mathbf{p} \mathbf{a}_{j}, C = c \big] f(\mathbf{p} \mathbf{a}_{j} \mid a', c) \nabla_{t=0} f_{t}(m_{j} \mid c) f(c) \, \mathrm{d}\mu \, \mathrm{d}F_{X, t} \\ &+ \int \mathbf{E} \big[ Y \mid A = a', M_{j} = m_{j}, \mathbf{P} \mathbf{a}_{j} = \mathbf{p} \mathbf{a}_{j}, C = c \big] f(\mathbf{p} \mathbf{a}_{j} \mid a', c) f(m_{j} \mid c) \nabla_{t=0} f_{t}(c) \, \mathrm{d}\mu \, \mathrm{d}F_{X, t} \\ &=: \sum_{k=1}^{4} A_{j,k}. \end{split}$$

Here  $A_{j,k}$ , k = 1, 2, 3, 4 can be calculated straightforwardly by the method in Tchetgen and Shpitser (2012). First,

$$\begin{split} A_{j,1} &= \int \nabla_t \mathbf{E}_t \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] f(\mathbf{p} \mathbf{a}_j \, | \, a', c) f(m_j \, | \, c) f(c) \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \mathrm{d}(m_j, c) \, \mathrm{d} F_{X,t} \\ &= \mathbf{E} \Bigg[ U \int \frac{\mathbbm{1}}{f(c) f(a' \, | \, c) f(\mathbf{p} \mathbf{a}_j \, | \, a', c) f(m_j \, | \, \mathbf{p} \mathbf{a}_j, a', c)} \\ &\qquad \times \Big\{ Y - \mathbf{E} \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] \Big\} f(\mathbf{p} \mathbf{a}_j \, | \, a', c) f(m_j, c) \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \mathrm{d}(m_j, c) \Bigg] \\ &= \mathbf{E} \Bigg[ U \int \frac{\mathbbm{1}}{a'} \frac{(a' \, = A, m_j \, = M_j, \mathbf{p} \mathbf{a}_j \, = \mathbf{P} \mathbf{a}_j, c \, = C) f(m_j \, | \, c)}{f(a' \, | \, c) f(m_j \, | \, \mathbf{p} \mathbf{a}_j, a', c)} \\ &\qquad \times \Big\{ Y - \mathbf{E} \big[ Y \, | \, A = a', M_j \, = m_j, \mathbf{P} \mathbf{a}_j \, = \mathbf{p} \mathbf{a}_j, C \, = c \big] \Big\} \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \, \mathrm{d}(m_j, c) \Bigg] \\ &= \mathbf{E} \Bigg[ U \frac{\mathbbm{1}}{a'} \frac{(A \, = \, a')}{c_{a'}(C)} \mathbbm{1}_{C,a'}(\mathbf{P} \mathbf{a}_j) \Big\{ Y - \mathbf{E} \big[ Y \, | \, A \, = \, a', M_j, \mathbf{P} \mathbf{a}_j, C \big] \Big\} \Bigg]. \end{split}$$

Similarly, one can easily obtain that

$$\begin{split} A_{j,2} &= \int \mathbf{E} \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] \nabla_t f_t(\mathbf{p} \mathbf{a}_j \, | \, a', c) f(m_j, c) \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \, \mathrm{d}(m_j, c) \, \mathrm{d} F_{X,t} \\ &= \mathbf{E} \left[ U \int \mathbf{E} \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] \\ &\qquad \times \frac{\mathbb{1}(a' = A, c = C)}{f(a' \, | \, c) f(c)} \left\{ \mathbb{1}(\mathbf{p} \mathbf{a}_j = \mathbf{P} \mathbf{a}_j) - f(\mathbf{p} \mathbf{a}_j \, | \, a', c) \right\} f(m_j \, | \, c) f(c) \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \mathrm{d} m_j \, \mathrm{d} c \right] \\ &= \mathbf{E} \left[ U \frac{\mathbb{1}(A = a')}{e_{a'}(C)} \left\{ \int \mathbf{E} \big[ Y \, | \, C, A = a', \mathbf{P} \mathbf{a}_j, M_j = m_j \big] f(m_j \, | \, C) \, dm_j - \mathbf{E} \big[ \varrho_j(a', M_j, C) \, | \, C \big] \right\} \right], \\ A_{j,3} &= \int \mathbf{E} \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] f(\mathbf{p} \mathbf{a}_j \, | \, a', c) \nabla_t f_t(m_j \, | \, c) f(c) \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \mathrm{d} (m_j, c) \, \mathrm{d} F_{X,t} \\ &= \mathbf{E} \left[ U \int \mathbf{E} \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] f(\mathbf{p} \mathbf{a}_j \, | \, a', c) \\ &\qquad \times \frac{\mathbb{1}(c = C)}{f(c)} \left\{ \mathbb{1}(m_j = M_j) - f(m_j \, | \, c) \right\} f(c) \, \mathrm{d} m_j \, \mathrm{d} \mathbf{p} \mathbf{a}_j \, \mathrm{d} c \right] \\ &= \mathbf{E} \left[ U \left\{ \varrho_j(a', M_j, C) - \mathbf{E} \big[ \varrho_j(a', M_j, C) \, | \, C \big] \right\} \right], \\ \mathbf{and} \\ A_{j,4} &= \int \mathbf{E} \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] f(\mathbf{p} \mathbf{a}_j \, | \, a', c) f(m_j \, | \, c) \nabla_t f_t(c) \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \mathrm{d}(m_j, c) \, \mathrm{d} F_{X,t} \right. \\ &= \mathbf{E} \left[ U \int \mathbf{E} \big[ Y \, | \, A = a', M_j = m_j, \mathbf{P} \mathbf{a}_j = \mathbf{p} \mathbf{a}_j, C = c \big] \\ &\qquad \qquad \times f(\mathbf{p} \mathbf{a}_j \, | \, a', c) f(m_j \, | \, c) \left\{ \mathbb{1}(c = C) - f(c) \right\} \, \mathrm{d} \, \mathbf{p} \mathbf{a}_j \, \mathrm{d}(m_j, c) \right] \\ &= \mathbf{E} \left[ U \left\{ \mathbf{E} \big[ \varrho_j(a', M_j, C) \, | \, C \big] - \mathbf{E} \varrho_j(a', M_j, C) \right\} \right]. \end{aligned}$$

Similarly, we can decompose

$$\begin{split} &\frac{\partial}{\partial t} \mathbf{E}_{t} \zeta_{j}(a',0,C) \bigg|_{t=0} \\ &= \int \nabla_{t=0} \mathbf{E}_{t} \big[ Y \mid A = 1, M = m, C = c \big] f(m_{-j} \mid A = 0, C = c) f(m_{j} \mid A = 1, C = c) f(c) \, \mathrm{d}\mu \, \mathrm{d}F_{X,t} \\ &+ \int \mathbf{E}_{t} \big[ Y \mid A = 1, M = m, C = c \big] \nabla_{t=0} f(m_{-j} \mid A = 0, C = c) f(m_{j} \mid A = 1, C = c) f(c) \, \mathrm{d}\mu \, \mathrm{d}F_{X,t} \\ &+ \int \mathbf{E}_{t} \big[ Y \mid A = 1, M = m, C = c \big] f(m_{-j} \mid A = 0, C = c) \nabla_{t=0} f(m_{j} \mid A = 1, C = c) f(c) \, \mathrm{d}\mu \, \mathrm{d}F_{X,t} \\ &+ \int \mathbf{E}_{t} \big[ Y \mid A = 1, M = m, C = c \big] f(m_{-j} \mid A = 0, C = c) f(m_{j} \mid A = 1, C = c) \nabla_{t=0} f(c) \, \mathrm{d}\mu \, \mathrm{d}F_{X,t} \\ &+ \sum_{k=1}^{4} B_{j,k}, \end{split}$$

$$=: \sum_{k=1}^{4} B_{j,k},$$

and similarly verify that

$$B_{j,1} = \mathbf{E} \left[ U \frac{\mathbb{1}(A=1)}{f(A=1 \mid C)} \frac{f(M_j \mid A=0, C) f(M_{-j} \mid A=a', C)}{f(M_j \mid A=1, C, M_{-j}) f(M_{-j} \mid A=1, C)} \left\{ Y - \mathbf{E} \left[ Y \mid A=1, M, C \right] \right\} \right],$$

$$B_{j,2} = \mathrm{E} \Bigg[ U \frac{\mathbbm{1}(A=0)}{f(A=0 \mid C)} \Bigg\{ \int \mathrm{E} \big[ Y \mid A=1, M_{-j}, C, M_j = m_j \big] f(m_j \mid A=a', C=c) \, \mathrm{d} m_j - \zeta_j(a', 0, C) \Bigg\} \Bigg],$$

$$B_{j,3} = \mathbb{E}\left[U\frac{\mathbb{1}(A=a')}{f(A=a'\mid C)} \left\{ \int \mathbb{E}\left[Y\mid A=1, M_j, C, M_{-j}=m_{-j}\right] f(m_{-j}\mid A=0, C) \, \mathrm{d}m_{-j} - \zeta_j(a', 0, C) \right\} \right],$$

and

$$B_{j,4} = \mathbb{E}\left[U\left(\zeta_j(a',0,C) - \mathbb{E}\zeta_j(a',0,C)\right)\right].$$

Finally, the efficient score for  $E\kappa(a',C)$  equal to

$$\frac{\mathbb{1}(A=a')}{e_{a'}(C)} \left\{ Y - \kappa(a',C) \right\} + \kappa(a',C) - \operatorname{E}\kappa(a',C)$$

has been well studied in various literature, see Section 2.2 in Tchetgen and Shpitser (2012) for example. Thus, we conclude the results in the theorem.  $\Box$ 

### **Proof of Corollary 4.4:**

PROOF. The results are directly derived due to the linear property of efficient scores.

### C.2. The Proofs of Section 5.

Let  $E_{\xi}[\cdot]$  represent the expected value computed with respect to the random variable  $\xi$ , while treating other variables as constants. Employing graphical techniques, we can demonstrate the lemma as follows.

LEMMA C.1. Suppose the model satisfies Assumption 5.1, then natural effects defined in Definition 3.1 satisfy

$$TE = E_C[E\{Y \mid do(A=1), C\} - E\{Y \mid do(A=0), C\}],$$

$$DE = E_C[E\{Y \mid do(A = 1, M = m^{(0)}), C\} - E\{Y \mid do(A = 0), C\}],$$

and

$$IE = E_C[E\{Y \mid do(A = 0, M = m^{(1)}), C\} - E\{Y \mid do(A = 0), C\}].$$

### **Proof of Lemma C.1:**

PROOF. Introduce  $A^{\dagger}$  has the same law as A except that  $A^{\dagger} \perp \!\!\! \perp C$ , then by the definition of do operator,

$$\begin{split} \mathbf{E}_{C} \big[ \mathbf{E} \{ Y \mid do(A = a), C \} \big] &= \mathbf{E}_{C} \mathbf{E} \{ Y \mid A^{\dagger} = a, C \} \\ &= \mathbf{E} \{ Y \mid A^{\dagger} = a \} = \mathbf{E} \{ Y \mid do(A = a) \}. \end{split}$$

for any  $a \in \{0,1\}$  This furthermore implies

$$\begin{split} TE &= \mathbb{E}\{Y \mid do(A=1)\} - \mathbb{E}\{Y \mid do(A=0)\} \\ &= \mathbb{E}_{C} \big[ \mathbb{E}\{Y \mid do(A=1), C\} \big] - \mathbb{E}_{C} \big[ \mathbb{E}\{Y \mid do(A=0), C\} \big]. \end{split}$$

This proves the result of TE. We can similarly prove the result for DE and IE.

Lemma C.1 shows the definition of natural effects defined in Pearl (2000) can actually be written as the average on confounders. We require some DAG lemmas in order to study the formula for indirect effects of mediators. To the best of our knowledge, these lemmas are also novel, and they can be helpful resources for pertinent research.

LEMMA C.2. Suppose B is the weight matrix of a DAG on  $\{X_1, ..., X_q\}$ . Define  $\Delta_{-i} := \left[ (I_q - B^\top)^{-1} u \right]_{-i} - (I_{q-1} - B^\top_{-i,-i})^{-1} u_{-i}$ , then

(1) (Path Representation)

$$\Delta_{-i} = u_i \left[ \frac{\partial}{\partial x_i} \mathbb{E}[X_j \mid do_B(X_i = x_i)] \right]_{j \neq i} + \left[ \sum_{k \neq i} u_k \frac{\partial}{\partial x_k} \mathbb{E}[X_j \mid do_{B_{-i,-i}}(X_k = x_k)] \right]_{j \neq i},$$

where  $do_A$  is the do operator on the DAG weight matrix A.

(2) (Matrix Expression)

$$\Delta_{-i} = u_i \left[ (I_q - B^\top)^{-1} \right]_{-i,i} + \left[ (I_q - B^\top)^{-1} \right]_{-i,i} \left[ (I_q - B^\top)^{-1} \right]_{i,-i} u_{-i}.$$

Before proving this lemma, we need the following result for digraph (do not require acyclic). We denote |w(x)| as the product of the weights of the edges for any path x.

LEMMA C.3 (Yu et al. (2019)). If A is an adjacency matrix of a weighted digraph on m vertices, then

$$\left[A^{k}\right]_{ij} = \sum_{x \in \pi_{ij}^{+}(k)} |w(x)| - \sum_{y \in \pi_{ij}^{-}(k)} |w(y)|,$$

where  $\pi_{ij}^+(k)$  is the set of path with positive direction from i to j and  $\pi_{ij}^-(k)$  is the set of path with negative direction from i to j with length k.

#### **Proof of Lemma C.2:**

PROOF. Denote j as the major index. And for any nonempty set  $S = \{l_1, \dots, l_{|S|}\}$ , we denote

$$\left[u_{j}\right]_{S} = \left[u_{j}\right]_{j \in S} = \left(u_{l_{1}}, \dots, u_{l_{|S|}}\right)^{\top}$$

as a vector with dimension |S|. Specially, define  $[u_j] := [u_j]_{j \in [q]}$ . By lemma C.3, for any  $u \in \mathbb{R}^q$ , we have

$$(I - B^{\top})^{-1}u = u + \sum_{m=1}^{\infty} \left[B^{\top}\right]^{m}u$$

$$= [u_{j}] + \sum_{m=1}^{\infty} \left[\sum_{k=1}^{q} \left[(B^{m})^{\top}\right]_{jk} u_{k}\right]$$

$$= [u_{j}] + \sum_{m=1}^{\infty} \left[\sum_{k=1}^{q} \left[B^{m}\right]_{kj} u_{k}\right]$$

$$\stackrel{\text{By applying Lemma C.3}}{=} [u_{j}] + \sum_{m=1}^{\infty} \left[\sum_{k=1}^{q} u_{k} \sum_{\Pi \in \pi_{kj}^{(m)}} \left|w_{B}(\Pi)\right|\right]$$

$$= [u_{j}] + \sum_{k=1}^{q} u_{k} \left[\sum_{m=1}^{\infty} \sum_{\Pi \in \pi_{kj}} \left|w_{B}(\Pi)\right|\right]$$

$$= \left[u_{j} + \sum_{k=1}^{q} u_{k} \sum_{\exists \Pi \in \pi_{kj}} \left|w_{B}(\Pi)\right|\right],$$

where we drop the superscript in the path  $\pi_{kj}^+$  by the fact that there is no negative path in DAG. Similarly, one can prove

$$(I - B_{-i,-i}^{\top})^{-1} u_{-i} = \left[ u_j + \sum_{k=1}^p u_k \sum_{\exists \Pi \in \pi_{kj} \text{ and } i \notin \Pi} \left| w_B(\Pi) \right| \right]_{j \neq i}.$$

Therefore,

$$\begin{split} &\Delta_{-i} = \left[ (I - B^\top)^{-1} u \right]_{-i} - (I - B^\top_{-i,-i})^{-1} u_{-i} \\ &= \left[ u_j + \sum_{k=1}^q u_k \sum_{\exists \Pi \in \pi_{kj}} \left| w_B(\Pi) \right| \right]_{j \neq i} - \left[ u_j + \sum_{k=1}^p u_k \sum_{\exists \Pi \in \pi_{kj} \text{ and } i \notin \Pi} \left| w_B(\Pi) \right| \right]_{j \neq i} \\ &= \left[ \sum_{k=1}^p u_k \sum_{\exists \Pi \in \pi_{kj} \text{ and } i \in \Pi} \left| w_B(\Pi) \right| \right]_{j \neq i} \\ &= \left[ u_i \sum_{\exists \Pi \in \pi_{ij}} \left| w_B(\Pi) \right| + \sum_{k \neq i}^q u_k \sum_{\exists \Pi \in \pi_{kj} \text{ and } i \notin \Pi} \left| w_B(\Pi) \right| \right]_{j \neq i} \\ &= \left[ u_i \sum_{\exists \Pi \in \pi_{ij}} \left| w_B(\Pi) \right| \right]_{j \neq i} + \left[ \sum_{k \neq i}^q u_k \sum_{\exists \Pi \in \pi_{kj} \text{ and } i \notin \Pi} \left| w_B(\Pi) \right| \right]_{j \neq i} \\ &= : \Delta_{-i,1} + \Delta_{-i,2}. \end{split}$$

By the definition of do operator and Proposition 3.1 in the Supplementary of Nandy et al. (2017), one have

$$\Delta_{-i,1} = \left[ u_i \sum_{\exists \Pi \in \pi_{ij}} \left| w_B(\Pi) \right| \right]_{j \neq i}$$

$$= u_i \left[ \sum_{\exists \Pi \in \pi_{ij}} \left| w_B(\Pi) \right| \right]_{j \neq i}$$

$$= u_i \left[ \frac{\partial}{\partial x_i} E[X_j \mid do_B(X_i = x_i)] \right]_{j \neq i}.$$

Similarly,

$$\begin{split} \Delta_{-i,2} &= \left[ \sum_{k \neq i}^q u_k \sum_{\exists \, \Pi \in \pi_{kj} \text{ and } i \notin \Pi} \left| w_B(\Pi) \right| \right]_{j \neq i} \\ &= \left[ \sum_{k \neq i}^q u_k \sum_{\exists \, \Pi \in \pi_{kj}(B_{-i,-i})} \left| w_{B_{-i,-i}}(\Pi) \right| \right]_{j \neq i} \\ &= \left[ \sum_{k \neq i} u_k \frac{\partial}{\partial x_k} \mathrm{E}[X_j \mid do_{B_{-i,-i}}(X_k = x_k)] \right]_{j \neq i}, \end{split}$$

This proves the path representation. For the matrix representation, we use Woodbury matrix identity

$$\left(A_{-j,-j}^{-1}\right)^{-1} = \left[A^{-1}\right]_{-j,-j} - \left[A^{-1}\right]_{-j,j} \left(\left[A^{-1}\right]_{jj}\right)^{-1} \left[A^{-1}\right]_{j-j}.$$

Thus, take  $A = (I_q - B^\top)$ , we have

$$(A_{-i,-i}^{-1})^{-1} = (I_{q-1} - B_{-i,-i})^{-1}$$

$$= [(I_q - B^\top)^{-1}]_{-i,-i} - [(I_q - B^\top)^{-1}]_{-i,i} ([(I_q - B^\top)^{-1}]_{i,i})^{-1} [(I_q - B^\top)^{-1}]_{i,-i}$$

$$= [(I_q - B^\top)^{-1}]_{-i,-i} - [(I_q - B^\top)^{-1}]_{-i,i} [(I_q - B^\top)^{-1}]_{i,-i}.$$

where we use the fact that  $[A^{-1}]_{i,i} \equiv 1$  for  $i \in [p]$  as  $B^{\top}$  is acyclic. Then,

$$(I_{q-1} - B_{-i,-i})^{-1} u_{-i}$$

$$= [(I_q - B^\top)^{-1}]_{-i,-i} u_{-i} - [(I_q - B^\top)^{-1}]_{-i,i} [(I_q - B^\top)^{-1}]_{i,-i} u_{-i}.$$

On the other hand, note that, for any  $j \neq i$ ,  $(Au)_j = \sum_{k=1}^q A_{jk}u_k$ . Thus,

$$[Au]_{-i} = [Au]_{j \neq i} = \left[ A_{ji}u_i + \sum_{k \neq i} A_{jk}u_k \right]_{j \neq i}$$
$$= u_i[A]_{-i,i} + A_{-i,-i}u_{-i}.$$

Note that

$$[(I_q - B^{\top})^{-1}u]_{-i}$$

$$= u_i [(I_q - B^{\top})^{-1}]_{-i,i} + [(I_q - B^{\top})^{-1}]_{-i,-i}u_{-i}.$$

Combine the two results, we obtain that

$$\Delta_{-i} = \left[ (I_q - B^\top)^{-1} u \right]_{-i} - (I_{q-1} - B_{-i,-i}^\top)^{-1} u_{-i}$$

$$= \left( u_i \left[ (I_q - B^\top)^{-1} \right]_{-i,i} + \left[ (I_q - B^\top)^{-1} \right]_{-i,-i} u_{-i} \right)$$

$$- \left( \left[ (I_q - B^\top)^{-1} \right]_{-i,-i} u_{-i} - \left[ (I_q - B^\top)^{-1} \right]_{-i,i} \left[ (I_q - B^\top)^{-1} \right]_{i,-i} u_{-i} \right)$$

$$= u_i \left[ (I_q - B^\top)^{-1} \right]_{-i,i} + \left[ (I_q - B^\top)^{-1} \right]_{-i,i} \left[ (I_q - B^\top)^{-1} \right]_{i,-i} u_{-i}.$$

This implies the matrix expression.

An important result from Lemma C.2 is that it guarantees the following two identities. See the following corollary.

COROLLARY C.4. Suppose the conditions are the same as Lemma C.3, then we have

(C.1) 
$$\left[ \mathbb{E}[X_j \mid X_i \cup \text{Pa}_B(X_i)]_1 \mathbb{1}(j \notin \text{Pa}_B(X_i)) \right]_{j \neq i} = \left[ (I - B^\top)^{-1} \right]_{-i,i},$$

and

(C.2) 
$$\left[ \mathbb{E}[X_i \mid X_j \cup \text{Pa}_B(X_j)]_1 \mathbb{1}(i \notin \text{Pa}_B(X_j)) \right]_{j \neq i} = \left[ (I - B^\top)^{-1} \right]_{i,-i}.$$

## **Proof of Corollary C.4:**

PROOF. Compare the two expressions in Lemma C.3, take arbitrary  $u \in \mathbb{R}^q$ , we have

$$\left[ (I - B^{\top})^{-1} \right]_{-i,i} = \left[ \frac{\partial}{\partial x_i} \mathbb{E}[X_j \mid do_B(X_i = x_i)] \right]_{j \neq i}.$$

On the other hand, the Pearl's back-door adjustment implies

$$\frac{\partial}{\partial x_i} \mathbb{E}[X_j \mid do_B(X_i = x_i)] = \mathbb{E}[X_j \mid X_i \cup \mathrm{Pa}_B(X_i)]_1 \mathbb{1}(j \notin \mathrm{Pa}_B(X_i)).$$

The two results give (C.1). Similarly, we can get (C.2).

## **Proof of Proposition 5.2:**

PROOF. Consider the M-estimator defined as

(C.3) 
$$\left[ \Theta^* \quad \theta^* \right] \in \arg \min_{[\Theta \ \theta] \in \mathbb{R}^{p \times t}} \operatorname{tr} \mathbf{E} \left[ M - \Theta C - \theta A \right] \left[ M - \Theta C - \theta A \right]^{\top}$$

Since the distribution of X is non-degenerate,

$$\operatorname{tr} \operatorname{E} [M - \Theta C - \theta A] [M - \Theta C - \theta A]^{\top}$$

$$= \operatorname{E} \operatorname{tr} [M - \Theta C - \theta A] [M - \Theta C - \theta A]^{\top}$$

$$= \operatorname{E} [M - \Theta C - \theta A]^{\top} [M - \Theta C - \theta A]$$

$$= \operatorname{E} [M - \Theta_{MC} C - \theta_{MA} A + (\Theta_{MC} - \Theta) C + (\theta_{MA} - \theta) A]^{\top}$$

$$[M - \Theta_{MC} C - \theta_{MA} A + (\Theta_{MC} - \Theta) C + (\theta_{MA} - \theta) A]$$

$$= \operatorname{E} [e_{M} + (\Theta_{MC} - \Theta) C + (\theta_{MA} - \theta) A]^{\top} [e_{M} + (\Theta_{MC} - \Theta) C + (\theta_{MA} - \theta) A]$$

$$= \operatorname{E} e_{M}^{\top} e_{M} + \operatorname{E} [(\Theta_{MC} - \Theta) C + (\theta_{MA} - \theta) A]^{\top} [(\Theta_{MC} - \Theta) C + (\theta_{MA} - \theta) A].$$

Therefore,  $\begin{bmatrix} \Theta_{MC} & \theta_{MA} \end{bmatrix}$  is a solution of (C.3). It is enough to show that the solution  $\begin{bmatrix} \Theta^* & \theta^* \end{bmatrix}$  is unique over MEC. Indeed, use the fact in Assumption 5.1 that the components of  $\epsilon$  are mutually independent,

$$\begin{split} \mathbf{E}\big[M - \Theta C - \theta A\big] \big[M - \Theta C - \theta A\big]^\top &= \mathbf{E} e_M e_M^\top \\ &= (I - B_{MM}^\top) \operatorname{cov}(\epsilon) (I - B_{MM}) \\ &= \left(\frac{1}{\sigma_{\ell_1}^2} [I - B_{MM}]_{\ell_1}^\top [I - B_{MM}]_{:\ell_2}\right)_{1 \le \ell_1, \ell_2 \le p}. \end{split}$$

When  $\ell_1 = \ell_2 = \ell$ ,  $\frac{1}{\sigma_{\ell_1}^2} [I - B_{MM}]_{\ell_1:1}^{\top} [I - B_{MM}]_{:\ell_2} = \frac{1}{\sigma_{\ell}^2} \sum_{k=1}^p \left( B_{MM} \right)_{k\ell}^2$ . Thus, we know that

$$\operatorname{tr} \mathbf{E} [M - \Theta C - \theta A] [M - \Theta C - \theta A]^{\top} = \sum_{1 \le i, k \le n} \frac{1}{\sigma_j^2} (B_{MM})_{kj}^2.$$

Since any DAG in the same MEC shares the same skeleton, we know that the above optimization problem (C.3) is unique over MEC. The unique solution comes from the theory of M-estimator, see Van de Geer et al. (2014).

## **Proof of Proposition 5.4:**

PROOF. The basic idea is implying Lemma C.1. Denote the conditional version of the natural direct effect as

$$DE(c) := E\{Y \mid do(A = 1, M = m^{(0)}), C = c\} - E\{Y \mid do(A = 0), C = c\}.$$

Note that under Assumption 5.1, we can write

$$A \leftarrow h(C, \epsilon_A),$$

$$M \leftarrow \left(I_{t-1} - B_{MM}^{\top}\right)^{-1} B_{MC}^{\top} C + \left(I_{t-1} - B_{MM}^{\top}\right)^{-1} \beta_{MA} A + \left(I_{t-1} - B_{MM}^{\top}\right)^{-1} \epsilon_{M}.$$

Thus, we have

$$E\{Y \mid do(A = 1, M = m^{(0)}), C = c\} = E\{Y \mid A = 1, M = m^{(0)}, C = c\}$$
$$= \beta_{YC}^{\top} c + \alpha_{YA} + \beta_{YM}^{\top} m^{(0)},$$

and

$$E\{Y \mid do(A=0), C=c\} = E\{Y \mid A=0, C=c\} = \beta_{YC}^{\top}C + \beta_{YM}^{\top}m^{(0)}.$$

These equations together with the first display in Lemma C.1 indicate

$$DE = \mathcal{E}_C DE(C)$$

$$= \mathcal{E}_C \left[ \left( \beta_{YC}^\top C + \alpha_{YA} + \beta_{YM}^\top m^{(0)} \right) - \left( \beta_{YC}^\top C + \beta_{YM}^\top m^{(0)} \right) \right]$$

$$= \mathcal{E}_C \alpha_{YA} = \alpha_{YA}.$$

And similarly,

$$\begin{split} IE &= \mathbf{E}_{C}IE(C) \\ &= \mathbf{E}_{C} \left[ \mathbf{E}\{Y \mid A = 0, M = m^{(1)}, C\} - \mathbf{E}\{Y \mid A = 0, C\} \right] \\ &= \mathbf{E}_{C} \left[ \beta_{YC}^{\top} C + \beta_{YM}^{\top} m^{(1)} - \left( \beta_{YC}^{\top} C + \beta_{YM}^{\top} m^{(0)} \right) \right] \\ &= \mathbf{E}_{C} \left[ \beta_{YM}^{\top} \left( m^{(1)} - m^{(0)} \right) \right] \\ &= \mathbf{E}_{C} \left[ \beta_{YM}^{\top} \left( m^{(1)} - m^{(0)} \right) \right] \\ &\stackrel{\text{by } C \perp \perp \in_{M}}{=} \mathbf{E}_{C} \left[ \left( I_{t-1} - B_{MM}^{\top} \right)^{-1} B_{MC}^{\top} C + \left( I_{t-1} - B_{MM}^{\top} \right)^{-1} \beta_{MA} \cdot 1 \right. \\ &\left. - \left( \left( I_{t-1} - B_{MM}^{\top} \right)^{-1} B_{MC}^{\top} C + \left( I_{t-1} - B_{MM}^{\top} \right)^{-1} \beta_{MA} \cdot 0 \right) \right] \\ &= \mathbf{E}_{C} \left[ \beta_{YM}^{\top} \left( I - B_{MM}^{\top} \right)^{-1} \beta_{MA} \right] = \beta_{YM}^{\top} \left( I - B_{MM}^{\top} \right)^{-1} \beta_{MA}. \end{split}$$

This completes part (i).

For proving part (ii), we first note that Theorem F.4 in Cai et al. (2020) implies  $DM_j$  defined in Definition 3.2 is equivalent to Definition 3.2 in Cai et al. (2020), which derives the expression  $DM_j$  directly from Part (i). On the other hand, recall we have  $TM_j = TM_j(c) = IE - IE_{\mathcal{G}_{(-j)}}$ . Hence,  $TM_j$  also has the exactly same as Definition 3.3 in Cai et al. (2020) by Corollary F.1 in Cai et al. (2020). Therefore, we denote the total effect's condition version as

$$TE(c) = E\{Y \mid do(A=1), C=c\} - E\{Y \mid do(A=0), C=c\}.$$

Then, by Theorem E.2 in Watson et al. (2023), that is  $TM_j(c)$  can be interpreted as the effect of treatment A on the outcome Y that is mediated by the mediator  $M_j$ , or inversely as the change in total treatment effect caused by  $M_j$  being removed from the causal graph, i.e.

$$TM_j(c) = TE(c) - TE_{\mathcal{G}_{(-j)}}(c) = IE(c) - IE_{\mathcal{G}_{(-j)}}(c) = IE - IE_{\mathcal{G}_{(-j)}}$$

Similar to Lemma C.1, one can show  $TM_j = E_C TM_j(C)$ . Thus,

$$\begin{split} \dot{T}M_{j} &= \mathbf{E}_{C}TM_{j}(C) \\ &- \mathbf{cov}(\mathbf{E}\{Y \mid do(M_{j} = m_{j} + 1), C\} - \mathbf{E}\{Y \mid do(M_{j} = m_{j}), C\}, \Delta_{j}(C)) \\ &= \mathbf{E}_{C}TM_{j}(C) = IE - IE_{\mathcal{G}_{(-j)}}. \end{split}$$

Therefore.

$$IM_{j} = TM_{j} - DM_{j} = IE - IE_{\mathcal{G}_{(-j)}} - DM_{j}$$

$$= \beta_{YM}^{\top} (I - B_{MM}^{\top})^{-1} \beta_{MA} - \beta_{YM_{-j}}^{\top} (I - B_{M_{-j}M_{-j}}^{\top})^{-1} \beta_{M_{-j}A} - DM_{j}$$

$$= \beta_{YM,-j}^{\top} \theta_{MA,-j} - \beta_{YM_{-j}}^{\top} \theta_{M_{-j}A} = \beta_{YM,-j}^{\top} (\theta_{MA,-j} - \theta_{M_{-j}A}).$$

Finally, for proving (ii'), note that Y is still linear about C, A, and M, given C = c, the proof argument in proof of Theorem 3.1 and Theorem 3.2 of Nandy et al. (2017) still holds. Thus, fixed C = c, Proposition 2.1 in Chakrabortty et al. (2018) guarantees, that is

$$\eta_{j}(c) = \mathbf{E} \big[ M_{j} \mid A \cup \{C = c\} \big]_{1} \times \mathbf{E} \big[ Y \mid M_{j}, \operatorname{Pa}(M_{j}), A, \{C = c\} \big]_{1}$$

$$\stackrel{\text{by (5.2)}}{=} \theta_{MA, j} \times \mathbf{E} \big[ Y \mid M_{j}, \operatorname{Pa}(M_{j}), A, \{C = c\} \big]_{1}$$

for any fixed C = c, which combined with the expression of  $DM_j$  gives the result for the alternative expression for  $IM_j$ . Hence, we complete our proof.

#### C.3. The Proofs of Section 6.

#### **Proof of Alternative Strategies:**

PROOF. Just by the definition, we can prove the equivalence for alternative strategy 1 as follows:

$$\begin{split} & \operatorname{E}\left[\frac{1(A=a')}{f(A=a'\mid c)}Y\right] = \int \frac{1(a=a')}{f(a=a'\mid c)}yf(c,a,y)\mathrm{d}(a,c) \\ & = \int \frac{1}{f(a=a'\mid c)}yf(y\mid a',c)f(a=a'\mid c)f(c)\,\mathrm{d}(a,c) \\ & = \int yf(y\mid a',c)f(c)\,\mathrm{d}(a,c) \\ & = \operatorname{E}\kappa(a',C), \end{split} \\ & \operatorname{E}\left[\frac{1(A=0)}{f(A=0\mid C)}\left\{\int \operatorname{E}[Y\mid A=1,M_{-j},C,M_j=m_j]f(m_j\mid A=a',C=c)\,\mathrm{d}m_j\right\}\right] \\ & = \int \frac{1(a=0)}{f(A=a\mid C)}f(c,a,m_{-j})\,\mathrm{d}c\,\mathrm{d}a\,\mathrm{d}m_{-j} \\ & \times \left\{\int \operatorname{E}[Y\mid A=1,M_{-j}=m_{-j},C=c,M_j=m_j]f(m_j\mid A=a',C=c)\,\mathrm{d}m_j\right\} \\ & = \int \frac{1}{f(A=0\mid C)}f(c)f(A=0\mid c)f(m_{-j}\mid C=c,A=0)\,\mathrm{d}c\,\mathrm{d}m_{-j} \\ & \times \left\{\int \operatorname{E}[Y\mid A=1,M_{-j}=m_{-j},C=c,M_j=m_j]f(m_j\mid A=a',C=c)\,\mathrm{d}m_j\right\} \\ & = \int \operatorname{E}[Y\mid A=1,M=m,C]f(m_j\mid C=c,A=a')f(m_{-j}\mid C=c,A=0)f(c)\,\mathrm{d}m\,\mathrm{d}c \\ & = \operatorname{E}[\zeta_j(a',0,C)] \\ \operatorname{and} & \operatorname{E}\left[\frac{1(A=a')}{f(A=a'\mid c)}\left\{\int \operatorname{E}[Y\mid C,A=a',\operatorname{Pa}_j(\mathcal{G}_M),M_j=m_j]f(m_j\mid C)\,\mathrm{d}m_j\right\}\right] \\ & = \int \frac{1(A=a')}{f(A=a'\mid c)}f(c,a,\operatorname{pa}_j)\,\mathrm{d}c\,\mathrm{d}a\,\mathrm{d}\operatorname{pa}_j \\ & \times \left\{\int \operatorname{E}[Y\mid C=c,A=a',\operatorname{Pa}_j(\mathcal{G}_M)=\operatorname{pa}_j,M_j=m_j]f(m_j\mid C=c)\,\mathrm{d}m_j\right\} \\ & = \int \operatorname{E}[Y\mid C=c,A=a',\operatorname{Pa}_j(\mathcal{G}_M)=\operatorname{pa}_j,M_j=m_j]f(m_j\mid C=c)\,\mathrm{d}m_j \\ & = \int \operatorname{E}[Y\mid C=c,A=a',\operatorname{Pa}_j(\mathcal{G}_M)=\operatorname{pa}_j,M_j=m_j]f(p_a\mid A=a',c)f(m_j,c)\,\mathrm{d}\operatorname{pa}_j\,\mathrm{d}(m_j,c) \\ & = \int \operatorname{e}_j(a',m_j,c;\mathcal{G}_M)\,\mathrm{d}(m_j,c)=\operatorname{E}[\varrho_j(a',M_j,C;\mathcal{G}_M)]. \end{split}$$

Thus, all identities for alternative strategy 1 hold. Similarly, for Strategy 2 and Strategy 3, we have

$$\begin{split} & \mathbb{E}\left[\frac{1(A=a')C)}{f(A=a')C)} \left\{ \int \mathbb{E}[Y \mid A=1, M_j, C, M_{-j}=m_{-j}] f(m_{-j} \mid A=0, C) \, \mathrm{d}m_{-j} \right\} \right] \\ & = \int \frac{1(A=a')}{f(A=a')C)} f(c, a, m_j) \, \mathrm{d}c \, \mathrm{d}a \, \mathrm{d}m_j \\ & \qquad \times \left\{ \int \mathbb{E}[Y \mid A=1, M_j=m_j, C=c, M_{-j}=m_{-j}] f(m_{-j} \mid A=0, C=c) \, \mathrm{d}m_{-j} \right\} \\ & = \int \mathbb{E}[Y \mid A=1, C=c, M=m] f(m_{-j} \mid A=0, c) f(m_j \mid A=a', c) f(c) \, \mathrm{d}m_{-j} \mathrm{d}c \, \mathrm{d}m_j \\ & = \mathbb{E}[\zeta_j(a', 0, C)], \\ & \mathbb{E}\left[\frac{1(A=1)}{f(A=1 \mid C)} \frac{f(M_{-j} \mid A=a', C)}{f(M_{-j} \mid A=1, C, M_j)} Y\right] \\ & = \int \frac{1(a=1)}{f(a=1 \mid c)} \frac{f(m_j \mid a=1, c) f(m_{-j} \mid A=a', c)}{f(m_j \mid a=1, c) f(m_{-j} \mid A=a', c)} \\ & \qquad \times y f(y \mid a=1, c, m) f(m \mid a=1, c) f(a=1 \mid c) f(c) \, \mathrm{d}\mu \\ & = \int y f(y \mid a=1, c, m) f(m_j \mid a=1, c) f(m_{-j} \mid A=a', c) f(c) \, \mathrm{d}\mu \\ & = \int \mathbb{E}[Y \mid A=1, M=m, C=c] f(m_j \mid a=1, c) f(m_{-j} \mid A=a', c) f(c) \, \mathrm{d}\mu = \mathbb{E}\left[\zeta_j(a', 0, C)\right], \\ & \text{and} \\ & \mathbb{E}\left[\frac{1(A=a')}{f(A=a' \mid c)} f\left(\operatorname{Pa}_j(\mathcal{G}_M) \mid C, A=a'\right) Y\right] \\ & = \int \frac{1}{f(a=a' \mid c)} f\left(\operatorname{Pa}_j(\mathcal{G}_M) \mid C, a=a'\right) y f(y \mid a=a', c, m_j, \operatorname{pa}_j) f(m_j \mid a=a', c) f(a=a' \mid c) f(c) \, \mathrm{d}\mu \\ & = \int y f(y \mid c, a=a', m_j, \operatorname{pa}_j) f\left(\operatorname{pa}_j \mid c, a=a'\right) y f(m_j, c) \, \mathrm{d}\mu_j \, \mathrm{d}(m_j, c) \\ & = \mathbb{E}\left[\varrho_j(a', M_j, C; \mathcal{G}_M)\right], \end{split}$$

which complete the calculations.

**C.4. Proof of Section 7.1.** Denote  $a = \text{plim } a_n \text{ if } a_n \xrightarrow{P} a$ .

LEMMA C.5. Let X are mean-zero random vector and Y are univariate random variable. Suppose  $E(\epsilon \mid X) = 0$  and  $var(\epsilon \mid X) = \sigma^2$  is free of X in the regression

$$Y = \beta^{\top} X + \epsilon = \beta_1^{\top} X_1 + \beta_2^{\top} X_2 + \epsilon.$$

Suppose  $X \in \mathbb{R}^p$ , and  $\widehat{\beta}$  is the OLS estimator with sample size p < n,

$$\sqrt{n}(\widehat{\beta}_1 - \beta_1) = \left[ \operatorname{plim} \widehat{\Gamma}_{X_1, X_2} \widehat{\Gamma}_{X_1, X_2}^{\top} \right]^{1/2} \frac{1}{\sqrt{n}} \sum_{i=1}^n \varepsilon_i + o_p(1) \rightsquigarrow \mathcal{N}\left(0, \operatorname{plim} \sum_{i=1}^n \epsilon_i^2 \left[ \widehat{\Gamma}_{X_1, X_2} \widehat{\Gamma}_{X_1, X_2}^{\top} \right] \right),$$

where  $\{\widehat{\epsilon}_i\}_{i=1}^n$  are the residuals and  $\{\varepsilon_i\}_{i=1}^n$  are the i.i.d.  $\mathcal{N}(0, \sigma^2 I)$  error vector independent with X.

PROOF. We first note, we have

(C.5) 
$$\widehat{\beta}_1 - \beta_1 = \left[ \mathbf{X}_1^{\top} (I_n - P_{\mathbf{X}_2}) \mathbf{X}_1 \right]^{-1} \mathbf{X}_1^{\top} (I_n - P_{\mathbf{X}_2}) \boldsymbol{\varepsilon} = \widehat{\Gamma}_{X_1, X_2} \boldsymbol{\varepsilon}.$$

$$\left[n\widehat{\Gamma}_{X_1,X_2}\widehat{\Gamma}_{X_1,X_2}^{\top}\right]^{-1} = \frac{1}{n}\mathbf{X}_1^{\top}\mathbf{X}_1 - \left[\frac{1}{n}\mathbf{X}_1^{\top}\mathbf{X}_2\right] \left[\frac{1}{n}\mathbf{X}_2^{\top}\mathbf{X}_2\right]^{-1} \left[\frac{1}{n}\mathbf{X}_2^{\top}\mathbf{X}_1\right],$$

we have

$$\begin{split} & \left[ n \widehat{\Gamma}_{X_1, X_2} \widehat{\Gamma}_{X_1, X_2}^{\top} \right]^{-1} - \left[ \widehat{\text{var}}(X_1) - \widehat{\text{cov}}(X_1, X_2) \widehat{\text{var}}^{-1}(X_2) \widehat{\text{cov}}(X_2, X_1) \right] \\ &= \left( \frac{1}{n} \mathbf{X}_1^{\top} \mathbf{X}_1 - \widehat{\text{var}}(X_1) \right) \\ &- \left( \left[ \frac{1}{n} \mathbf{X}_1^{\top} \mathbf{X}_2 \right] \left[ \frac{1}{n} \mathbf{X}_2^{\top} \mathbf{X}_2 \right]^{-1} \left[ \frac{1}{n} \mathbf{X}_2^{\top} \mathbf{X}_1 \right] - \widehat{\text{cov}}(X_1, X_2) \widehat{\text{var}}^{-1}(X_2) \widehat{\text{cov}}(X_2, X_1) \right) \\ &= 0, \end{split}$$

which gives

$$n\widehat{\Gamma}_{X_1,X_2}\widehat{\Gamma}_{X_1,X_2}^{\top} = \left[\widehat{\operatorname{var}}(X_1) - \widehat{\operatorname{cov}}(X_1,X_2)\widehat{\operatorname{var}}^{-1}(X_2)\widehat{\operatorname{cov}}(X_2,X_1)\right]^{-1}.$$

On the other hand, from (C.5), we get that

$$\widehat{\beta}_1 - \beta_1 = \widehat{\Gamma}_{X_1, X_2} \varepsilon.$$

Since  $\sqrt{n}(\widehat{\beta} - \beta)$  is asymptotic normal and  $\mathrm{E}(\epsilon^2 \mid X) = \sigma^2$ , its asymptotic variance is

$$\left( \operatorname{E} \begin{bmatrix} X_{1}X_{1}^{\top} X_{1}X_{2}^{\top} \\ X_{2}X_{1}^{\top} X_{2}X_{2}^{\top} \end{bmatrix} \right)^{-1} \operatorname{E} \left( \epsilon_{Y}^{2} \begin{bmatrix} X_{1}X_{1}^{\top} X_{1}X_{2}^{\top} \\ X_{2}X_{1}^{\top} X_{2}X_{2}^{\top} \end{bmatrix} \right) \left( \operatorname{E} \begin{bmatrix} X_{1}X_{1}^{\top} X_{1}X_{2}^{\top} \\ X_{2}X_{1}^{\top} X_{2}X_{2}^{\top} \end{bmatrix} \right)^{-1} \\
= \sigma^{2} \left( \begin{bmatrix} \operatorname{var}(X_{1}) & \operatorname{cov}(X_{1}, X_{2}) \\ \operatorname{cov}(X_{2}, X_{1}) & \operatorname{var}(X_{2}) \end{bmatrix} \right)^{-1} \\
= \sigma^{2} \begin{bmatrix} \left( \operatorname{var}(X_{1}) - \operatorname{cov}(X_{1}, X_{2}) \operatorname{var}^{-1}(X_{2}) \operatorname{cov}(X_{2}, X_{1}) \right)^{-1} * \\ * \end{bmatrix}.$$

Combining the above result, we have

$$\begin{split} \sqrt{n} \left( \widehat{\beta}_1 - \beta_1 \right) &\leadsto \mathcal{N} \left( 0, \sigma^2 \left[ \operatorname{var}(X_1) - \operatorname{cov}(X_1, X_2) \operatorname{var}^{-1}(X_2) \operatorname{cov}(X_2, X_1) \right]^{-1} \right) \\ &= \mathcal{N} \left( 0, \sigma^2 \operatorname{plim} n \widehat{\Gamma}_{X_1, X_2} \widehat{\Gamma}_{X_1, X_2}^\top \right) \\ &= \mathcal{N} \left( 0, \operatorname{plim} \frac{\sum_{i=1}^n \widehat{\epsilon}_i^2}{n} \operatorname{plim} n \widehat{\Gamma}_{X_1, X_2} \widehat{\Gamma}_{X_1, X_2}^\top \right), \end{split}$$

by the fact that  $\frac{1}{n}\sum_{i=1}^{n}\hat{\epsilon}_{i}^{2}\overset{\mathrm{P}}{\longrightarrow}\mathrm{E}\epsilon^{2}=\sigma^{2}.$  The asymptotic linear expression comes from the above display directly.

### **Proof of Theorem 7.3:**

PROOF. From the regression (5.2), we know that

$$\begin{pmatrix} \widehat{\beta}_{YC} \\ \widehat{\alpha}_{YA} \\ \widehat{\beta}_{YM} \end{pmatrix} - \begin{pmatrix} \beta_{YC} \\ \alpha_{YA} \\ \beta_{YM} \end{pmatrix} = \begin{pmatrix} \begin{bmatrix} \mathbf{C}^{\top} \\ \mathbf{A}^{\top} \\ \mathbf{M}^{\top} \end{bmatrix} \begin{bmatrix} \mathbf{C} \mathbf{A} \mathbf{M} \end{bmatrix} \end{pmatrix}^{-1} \begin{bmatrix} \mathbf{C}^{\top} \\ \mathbf{A}^{\top} \\ \mathbf{M}^{\top} \end{bmatrix} \boldsymbol{\epsilon}_{Y}.$$

Thus, by (C.5), we have

$$\widehat{\alpha}_{YA} - \alpha_{YA} = \widehat{\Gamma}_{A,(M,C)} \epsilon_Y$$

and

$$\widehat{\beta}_{YM} - \beta_{YM} = \widehat{\Gamma}_{M,(C,A)} \epsilon_{Y}.$$

where  $var(\epsilon_Y \mid C, A, M) = \sigma_Y^2$  is a constant free of C, A, M. Similarly,

$$\widehat{\boldsymbol{\theta}}_{MA}^{\top} - \boldsymbol{\theta}_{MA}^{\top} = \widehat{\boldsymbol{\Gamma}}_{A,C} \mathbf{e}_{M}$$

it is worthy to note that

$$\operatorname{var}(e_M \mid C, A) = \operatorname{var}\left[ (I - B_{MM}^{\top}) \epsilon_M \mid (I - B_{CC}^{\top}) \epsilon_C, \beta_{AC}^{\top} (I - B_{CC}^{\top}) \epsilon_C + \epsilon_A \right]$$
$$= \operatorname{var}((I - B_{MM}^{\top}) \epsilon_M) = \sigma_{\epsilon_M}^2 (I - B_{MM}^{\top}) (I - B_{MM})$$

is also a constant free of C and A by  $\epsilon_M \perp \!\!\! \perp (C, \epsilon_A)$ . Therefore, by Lemma C.5, we have

$$\sqrt{n} (\widehat{\alpha}_{YA} - \alpha_{YA}) \rightsquigarrow \mathcal{N} \left( 0, \text{plim} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \left[ \widehat{\Gamma}_{A,(M,C)} \widehat{\Gamma}_{A,(M,C)}^{\top} \right] \right),$$

$$\sqrt{n}(\widehat{\beta}_{YM} - \beta_{YM}) \rightsquigarrow \mathcal{N}\left(0, \text{plim}\sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \left[\widehat{\Gamma}_{M,(C,A)}\widehat{\Gamma}_{M,(C,A)}^{\top}\right]\right),$$

and

$$\begin{split} \sqrt{n} \begin{pmatrix} \widehat{\alpha}_{YA} - \alpha_{YA} \\ \widehat{\beta}_{YM} - \beta_{YM} \end{pmatrix} &= \begin{pmatrix} \widehat{\Gamma}_{A,(M,C)} \\ \widehat{\Gamma}_{M,(C,A)} \end{pmatrix} \epsilon_{Y} \\ & \rightsquigarrow N \begin{pmatrix} 0, \begin{bmatrix} * & \text{plim} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \left[ \widehat{\Gamma}_{A,(M,C)} \widehat{\Gamma}_{M,(C,A)}^{\top} \right] \\ \text{plim} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \left[ \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{A,(M,C)}^{\top} \right] \\ * \end{pmatrix}. \end{split}$$

Besides,

$$\sqrt{n}(\widehat{\theta}_{MA} - \theta_{MA}) \rightsquigarrow \mathcal{N}\left(0, \operatorname{plim} \widehat{\mathbf{e}}_{M}^{\top} \widehat{\mathbf{e}}_{M} \Gamma_{A,C} \Gamma_{A,C}^{\top}\right).$$

Hence, we have

$$\begin{split} \sqrt{n} \left( \frac{\widehat{DE}^{OLS} - DE}{\widehat{IE}^{OLS} - IE} \right) &= \begin{pmatrix} 1 \\ \theta_{MA}^{\top} & \beta_{YM}^{\top} \end{pmatrix} \begin{pmatrix} \widehat{\alpha}_{YA} - \alpha_{YA} \\ \widehat{\beta}_{YM} - \beta_{YM} \\ \widehat{\theta}_{MA} - \widehat{\theta}_{MA} \end{pmatrix} \\ & \sim N \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \begin{bmatrix} \text{plim} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \left[ \widehat{\Gamma}_{A,(M,C)} \widehat{\Gamma}_{A,(M,C)}^{\top} \right] & \text{plim} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \left[ \widehat{\Gamma}_{A,(M,C)} \widehat{\Gamma}_{M,(C,A)}^{\top} \right] \theta_{MA} \\ \text{plim} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \theta_{MA}^{\top} \left[ \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{A,(M,C)} \right] \theta_{MA}^{\top} \left[ \text{plim} \sum_{i=1}^{n} \widehat{\epsilon}_{Y,i}^{2} \left[ \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{M,(C,A)}^{\top} \right] \right] \theta_{MA} \\ + \beta_{YM}^{\top} \left[ \text{plim} \widehat{\mathbf{e}}_{M}^{\top} \widehat{\mathbf{e}}_{M} \Gamma_{A,C} \Gamma_{A,C}^{\top} \right] \beta_{YM} \end{bmatrix} \end{split}$$

by  $\epsilon_Y \perp \!\!\! \perp e_M$ , which gives the result for  $\widehat{DE}^{OLS}$  as well as the result for  $\widehat{IE}^{OLS}$  when  $\theta_{MA}$  and  $\beta_{YM}$  are not both equal to zero. When  $\theta_{MA} = \beta_{YM} = 0$ , we write

$$\frac{\sqrt{n}(\widehat{IE}^{\text{OLS}} - IE)}{\sqrt{\widehat{\beta}_{YM}^{\top}\widehat{\Sigma}_{\theta_{MA}}\widehat{\beta}_{YM} + \widehat{\theta}_{MA}^{\top}\widehat{\Sigma}_{\beta_{YM}}\widehat{\theta}_{MA}}} = \frac{\sqrt{n}\widehat{\theta}_{MA}^{\top}\sqrt{n}\beta_{YM}}{\sqrt{\sqrt{n}\widehat{\beta}_{YM}^{\top}\widehat{\Sigma}_{\theta_{MA}}\sqrt{n}\widehat{\beta}_{YM} + \sqrt{n}\widehat{\theta}_{MA}^{\top}\widehat{\Sigma}_{\beta_{YM}}\sqrt{n}\widehat{\theta}_{MA}}}$$

which leads to the result by the continuous mapping theorem. Similar method applies to the proof of  $\widehat{DM}_j^{OLS}$ . Therefore, we have Then

$$\frac{\sqrt{n}(\widehat{DE}^{OLS} - DE)}{\sqrt{\widehat{\Gamma}_{A,(M,C)}\widehat{\Gamma}_{A,(M,C)}^{\top}\sum_{i=1}^{n}\widehat{\epsilon}_{Y,i}^{2}} \rightsquigarrow \mathcal{N}(0,1).$$

and

$$\frac{\sqrt{n}(\widehat{IE}^{\text{OLS}} - IE)}{\sqrt{\widehat{\beta}_{YM}^{\top} \widehat{\Sigma}_{\theta_{MA}} \widehat{\beta}_{YM} + \widehat{\theta}_{MA}^{\top} \widehat{\Sigma}_{\beta_{YM}} \widehat{\theta}_{MA}}} \rightsquigarrow \begin{cases} \frac{Z_{\theta}^{\top} Z_{\beta}}{\sqrt{Z_{\theta}^{\top} \Sigma_{\beta} Z_{\theta} + Z_{\beta}^{\top} \Sigma_{\theta} Z_{\beta}}} &, \text{if } \theta_{MA} = \beta_{YM} = 0, \\ \mathcal{N}(0, 1), & \text{otherwise.} \end{cases}$$

where

$$\begin{pmatrix} Z_{\theta} \\ Z_{\beta} \end{pmatrix} \sim \mathcal{N} \left( 0, \begin{pmatrix} \Sigma_{\theta_{MA}} \\ \Sigma_{\beta_{YM}} \end{pmatrix} \right), \qquad \Sigma_{\theta_{MA}} = \operatorname{plim} \widehat{\Sigma}_{\theta_{MA}}, \qquad \Sigma_{\beta_{YM}} = \operatorname{plim} \widehat{\Sigma}_{\beta_{YM}}.$$

Note that the  $\frac{Z_{\theta}^{\top}Z_{\beta}}{\sqrt{Z_{\theta}^{\top}\Sigma_{\beta}Z_{\theta}+Z_{\beta}^{\top}\Sigma_{\theta}Z_{\beta}}}$  is much more concentrated around zero compared to the standard normal distribution Chakrabortty et al. (2018), we get the " $\geq$ " instead of "=" for the confidence interval of  $\widehat{IE}^{OLS}$ . Similar result can be applied to  $\widehat{DM}_{j}^{OLS}$ .

### **Proof of Theorem 7.8:**

PROOF. Let  $\mu = EX$  is the mean of X. Denote

$$\xi_j(\mathcal{G}_M) := \mathrm{E}^{\mathrm{reg}} \big[ Y \mid M_j \cup \mathrm{Pa}_j(\mathcal{G}_M) \cup A \cup C \big]_1$$

and corresponding quantities

$$\overline{\xi}_{j}(\mathcal{C}_{M}) := \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} \operatorname{E}^{\operatorname{reg}} \left[ Y \mid M_{j} \cup \operatorname{Pa}_{j}(\mathcal{G}_{M}) \cup A \cup C \right]_{1}$$

and

$$\widetilde{\xi}_{j}(\mathcal{C}_{M}) := \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} \widehat{\xi}(\mathcal{G}_{M}), \qquad \widehat{\xi}(\mathcal{G}_{M}) := \widehat{\operatorname{E}}^{\operatorname{reg}} \big[ Y \mid M_{j} \cup \operatorname{Pa}_{j}(\mathcal{G}_{M}) \cup A \cup C \big]_{1}$$

From (ii') in Proposition 5.4, we have

$$\begin{split} \widehat{IM}_{j}^{OLS} - \overline{IM}_{j} \\ = &\widehat{\theta}_{MA,j} \Big( \widetilde{\xi}_{j}(\widehat{\mathcal{C}}_{M}) - \widehat{\beta}_{YM,j} \Big) - \theta_{MA,j} \Big( \overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j} \Big) \\ = &\widehat{\theta}_{MA,j} \Big( \big( \widetilde{\xi}_{j}(\widehat{\mathcal{C}}_{M}) - \overline{\xi}_{j}(\mathcal{C}_{M}) \big) - \big( \widehat{\beta}_{YM,j} - \beta_{YM,j} \big) \Big) + \big( \widehat{\theta}_{MA,j} - \theta_{MA,j} \big) \Big( \overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j} \Big) \\ = &\widehat{\theta}_{MA,j} \Big( \widetilde{\xi}_{j}(\widehat{\mathcal{C}}_{M}) - \widetilde{\xi}_{j}(\mathcal{C}_{M}) \Big) + \widehat{\theta}_{MA,j} \Big( \widetilde{\xi}_{j}(\mathcal{C}_{M}) - \overline{\xi}_{j}(\mathcal{C}_{M}) \Big) \\ - &\widehat{\theta}_{MA,j} \Big( \widehat{\beta}_{YM,j} - \beta_{YM,j} \Big) + \Big( \widehat{\theta}_{MA,j} - \theta_{MA,j} \Big) \Big( \overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j} \Big) \\ = : &D_{j,1} + D_{j,2} - D_{j,3} + D_{j,4} \end{split}$$

$$=:D_{j,1}+D_{j,2}-D_{j,3}+D_{j,4}$$

for  $j \in [p]$ . What we want to do next is finding the asymptotic expression of  $\widehat{IM}_j - \overline{IM}_j$ . First, for any non-negative sequence  $\{a_n\}$ ,

$$P\left(a_{n} \max_{j \in [p]} \left| D_{j,1} \right| > \varepsilon\right) \leq P\left(\exists j \in [p] : \widetilde{\xi}_{j}(\widehat{C}_{M}) - \widetilde{\xi}_{j}(C_{M}) \neq 0\right)$$

$$\leq P\left(\widehat{C}_{M} \neq C_{M}\right) \xrightarrow{\text{by Assumption (7.4)}} 0.$$

Thus, we have  $\max_{j \in [p]} D_{j,1} = o_p(a_n^{-1})$ . We first consider the case at least one  $\theta_{MA,j}$  or  $\overline{\xi}_j(\mathcal{C}_M) - \beta_{YM,j}$  is not equal to zero. From the proof of Theorem 7.3 and Lemma C.5, we know that

$$D_{j,2} = \theta_{MA,j} \left( \widetilde{\xi}_j(\mathcal{C}_M) - \overline{\xi}_j(\mathcal{C}_M) \right) + o_p \left( \frac{1}{n} \right),$$

$$\begin{split} D_{j,3} &= \theta_{MA,j} \left[ \widehat{\Gamma}_{M,(C,A)} \boldsymbol{\epsilon}_Y \right]_j + o_p \left( \frac{1}{n} \right) \\ &= \theta_{MA,j} \left[ \text{plim } \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{M,(C,A)}^\top \right]_{jj}^{1/2} \frac{1}{n} \sum_{i=1}^n \boldsymbol{\epsilon}_Y + o_p \left( \frac{1}{n} \right), \end{split}$$

and

$$\begin{split} D_{j,4} &= \left(\widehat{\theta}_{MA,j}^{\top} - \theta_{MA,j}^{\top}\right) \left(\overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j}\right) \\ &= \left[\widehat{\Gamma}_{A,C} \mathbf{e}_{M}\right]_{:j} \left(\overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j}\right) + o_{p} \left(\frac{1}{n}\right) \\ &= \left(\overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j}\right) \left[\widehat{\Gamma}_{A,C} \epsilon_{M} (I_{p} - B_{MM})^{-1}\right]_{:j} + o_{p} \left(\frac{1}{n}\right) \\ &= \left(\overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j}\right) \left[\operatorname{plim} \widehat{\Gamma}_{A,C} \widehat{\Gamma}_{A,C}^{\top}\right]^{1/2} \\ &= \left[\left(I_{p} - B_{MM}^{\top}\right)^{-1} \Sigma_{M} (I_{p} - B_{MM})^{-1}\right]_{jj}^{1/2} \frac{1}{n} \sum_{i=1}^{n} \varepsilon_{M,j,i} + o_{p} \left(\frac{1}{n}\right). \end{split}$$

uniformly in  $j \in [p]$ , where  $\varepsilon_M := \Sigma_M^{-1/2} \epsilon_M$ . We denote  $X_{S_{M,j,1}}, \dots, X_{S_{M,j,L_{\text{distinct},j}}}$  be distinct parent sets of  $M_j$  with  $m_{j,1}, \dots, m_{j,L_{\text{distinct},j}}$  times. Let

$$\begin{split} \mathcal{S}_{j} := \left\{ S_{jr} = (j+t,1,\dots,t,X_{S_{M,j,r}}) : r \in \{1,\dots,L_{\text{distinct},j}\}, \, t \notin X_{S_{M,j,r}} \right\} \\ & \cup \left\{ S_{jr} = (j+t,1,\dots,t-1,X_{S_{M,j,r}}) : r \in \{1,\dots,L_{\text{distinct},j}\}, \, t \in X_{S_{M,j,r}} \right\} \epsilon_{Y} \end{split}$$

with  $a_{S_{jr}} = (m_{jr}/L_j, 0, \dots, 0)^{\top} \in \mathbb{R}$  and  $L_j = \sum_{r=1}^{L_{\text{distinct},j}} m_{jr}$ . Since

$$\sum_{S_{jr} \in \mathcal{S}_j} \|a_{S_{jr}}\|_2 = \sum_{r=1}^{L_{\text{distinct},j}} \frac{m_{jr}}{L_j} = 1$$

for any  $j \in [p]$ , we can apply REMARK 5.2 in Chakrabortty et al. (2018), we have

$$\widetilde{\xi}_j(\mathcal{C}_M) - \overline{\xi}_j(\mathcal{C}_M) = \frac{1}{n} \sum_{i=1}^n W_i^{(j)}(\mathcal{C}_M) + o_p\left(\frac{1}{\sqrt{n}}\right)$$

where

$$W^{(j)}(\mathcal{C}_M)$$

$$\begin{split} &= \frac{1}{L_{j}} \sum_{\ell=1}^{L_{j}} e_{1,|S_{j\ell}|}^{\top} \left( \Sigma_{S_{j\ell}S_{j\ell}} \right)^{-1} \left( X_{S_{j\ell}} - \mu_{S_{j\ell}} \right) \left( Y - EY - \Sigma_{pS_{j\ell}} \left( \Sigma_{S_{j\ell}S_{j\ell}} \right)^{-1} \left( X_{S_{j\ell}} - \mu_{S_{j\ell}} \right) \right) \\ &= \frac{1}{L_{j}} \sum_{\ell=1}^{L_{j}} \frac{m_{r}}{L_{j} \sigma_{t+j|(1,\dots,t,S_{M,j,\ell})}^{2}} \left( (M_{j,i} - EM_{j}) - \beta_{t+j|(1,\dots,t,S_{M,j,\ell})}^{\top} \left( X_{(1,\dots,t,S_{M,\ell,j}),i} - EX_{(1,\dots,t,S_{M,\ell,j})} \right) \right) \\ &\times \left( Y - EY - \Sigma_{pS_{j\ell}} \left( \Sigma_{S_{j\ell}S_{j\ell}} \right)^{-1} \left( X_{S_{j\ell}} - \mu_{S_{j\ell}} \right) \right) \end{split}$$

where  $e_{i,k}$  the i-th column of the  $k \times k$  identity matrix and  $\beta_{t+j|(1,\dots,t,S_{M,j,\ell})}^{\top} := (\Sigma)_{j(1,\dots,t,S_{M,j,\ell})}$   $(\Sigma_{(1,\dots,t,S_{M,j,\ell})(1,\dots,t,S_{M,j,\ell})})^{-1}$ . Therefore, we can obtain

$$\sqrt{n} \left( \widehat{IM}_j^{OLS} - \overline{IM}_j \right)$$

$$= \frac{1}{\sqrt{n}} \sum_{i=1}^{n} \theta_{MA,j} \left\{ W_i^{(j)}(\mathcal{C}_M) + \left[ \operatorname{plim} \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{M,(C,A)}^{\top} \right]_{jj}^{1/2} \epsilon_{Y,i} \right\}$$

$$+ \left(\overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j}\right) \left[\operatorname{plim}\widehat{\Gamma}_{A,C}\widehat{\Gamma}_{A,C}^{\top}\right]^{1/2} \left[ (I_{p} - B_{MM}^{\top})^{-1} \Sigma_{M} (I_{p} - B_{MM})^{-1} \right]_{jj}^{1/2} \frac{1}{\sqrt{n}} \sum_{i=1}^{n} \varepsilon_{M,j,i} + o_{p}(1) \rightsquigarrow \mathcal{N}\left(0, \sigma_{\overline{IM}}^{2}\right)$$

where

$$\begin{split} &\sigma_{\overline{IM}_{j}}^{2} := \mathbf{E} \bigg[ \theta_{MA,j} \Big\{ W^{(j)}(\mathcal{C}_{M}) + \left[ \operatorname{plim} \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{M,(C,A)}^{\top} \right]_{jj}^{1/2} \epsilon_{Y} \Big\} \\ &+ \left( \overline{\xi}_{j}(\mathcal{C}_{M}) - \beta_{YM,j} \right) \left[ \operatorname{plim} \widehat{\Gamma}_{A,C} \widehat{\Gamma}_{A,C}^{\top} \right]^{1/2} \left[ (I_{p} - B_{MM}^{\top})^{-1} \Sigma_{M} (I_{p} - B_{MM})^{-1} \right]_{jj}^{1/2} \varepsilon_{M,j} \bigg]^{2}, \end{split}$$

which implies  $\sigma_{\overline{IM}_j}^2$  can be estimated by

$$\widehat{\sigma}_{\overline{IM}_{j}}^{2} = \frac{1}{n} \sum_{i=1}^{n} \left[ \widehat{\theta}_{MA,j} \left\{ \widehat{W}_{i}^{(j)}(\widehat{\mathcal{C}}_{M}) + \left[ \widehat{\Gamma}_{M,(C,A)} \widehat{\Gamma}_{M,(C,A)}^{\top} \right]_{jj}^{1/2} \widehat{\epsilon}_{Y,i} \right\}$$

$$+ \left( \widetilde{\xi}_{j}(\widehat{\mathcal{C}}_{M}) - \widehat{\beta}_{YM,j} \right) \left[ \widehat{\Gamma}_{A,C} \widehat{\Gamma}_{A,C}^{\top} \right]^{1/2} \left[ (I_{p} - \widehat{B}_{MM}^{\top})^{-1} \widehat{\Sigma}_{M} (I_{p} - \widehat{B}_{MM})^{-1} \right]_{jj}^{1/2} \widehat{\epsilon}_{M,j,i} \right]^{2}$$

where  $\widehat{B}_{MM}$  can be estimated incidentally by a version of PC algorithm (Harris and Drton, 2013). Finally, when  $\theta_{MA,j} = \overline{\xi}_j(\mathcal{C}_M) - \beta_{YM,j} = 0$ , we will have  $\lim_{n \to \infty} \mathrm{P} \big( \sqrt{n} \big| \widehat{IM}_j^{OLS} - IM_j \big| \ge \widehat{\sigma}_{\overline{IM}_j} \Phi^{-1} (1 - \alpha/2) \big) > 1 - \alpha$  by using the similar argument in the proof of Theorem 7.3.

## **Proof of Theorem 7.11:**

PROOF. We will prove the consistency for  $\widehat{DM}_j^{QR}$  and semiparametric efficiency for  $\widehat{TM}_j^{avg,QR}$ . The proof of other parts can be similarly derived. We first prove the consistency.

## Consistency of $\widehat{DM}_{j}^{\mathrm{QR}}$ :

Note that 
$$\widehat{DM}_j^{\mathrm{QR}} - DM_j = \mathbb{P}_n \langle \widehat{S}^{\mathrm{eff, nonpar}} \big( \mathrm{E} \zeta_j(\cdot, 0, C) \big) \rangle$$
, so it is sufficient to prove  $\mathbb{P}_n \widehat{S}^{\mathrm{eff, nonpar}} \big( \mathrm{E} \zeta_j(a', 0, C) \big) \xrightarrow{\mathrm{P}} 0$  for any  $a' \in \{0, 1\}$ . Denote  $\pi_{x_S}^{(T)}(m_T) = \pi_{x_S}(m_T)$ ,

Then we write

$$\psi_{j}^{(0)}(\mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)}) = \int_{\mathcal{M}} \mu(C, 1, m) \pi_{C,1}(m_{j}) \pi_{C,0}(m_{-j}) \, dm - E\zeta_{j}(a', 0, C)$$

$$\psi_{j}^{(1)}(e_{0}, \mu, \pi_{C,a'}^{(j)}) = \frac{\mathbb{1}(A = 0)}{e_{0}(C)} \left\{ \int_{\mathcal{M}_{j}} \mu(C, 1, m_{j}, M_{-j}) \pi_{C,a'}(m_{j}) \, dm_{j} - \zeta_{j}(a', 0, C) \right\}$$

$$\psi_{j}^{(2)}(e_{a'}, \mu, \pi_{C,0}^{(-j)}) = \frac{\mathbb{1}(A = a')}{e_{a'}(C)} \left\{ \int_{\mathcal{M}_{-j}} \mu(C, 1, M_{j}, m_{-j}) \pi_{C,0}(m_{-j}) \, dm_{-j} - \zeta_{j}(a', 0, C) \right\}$$

$$\psi_{j}^{(3)}(e_{1}, \pi_{C,a'}^{(-j)}, \pi_{C,1,M_{j}}^{(-j)}, \mu) = \frac{\mathbb{1}(A = 1)}{e_{1}(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_{j}}(M_{-j})} \left\{ Y - \mu(C, 1, M) \right\}$$

then  $\mathbb{P}_n S^{\mathrm{eff,\,nonpar}} \big( \mathrm{E} \zeta_j(a',0,C) \big) = \sum_{\ell=0}^3 \mathbb{P}_n \psi_j^{(\ell)}(\cdot)$ . We break the proof into four parts, **Part**  $\ell$  gives the consistency under different model  $\mathcal{M}_{j,\,\ell}$ .

**Part 0.** When model  $\mathcal{M}_0$  is correctly specified, i.e.,  $\widehat{\mu}$ ,  $\widehat{\pi}_{C,a'}^{(j)}$ , and  $\widehat{\pi}_{C,0}^{(-j)}$  are consistency. We have

$$\mathbb{P}_n \psi_j^{(0)} \left( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \right) = \mathbb{P}_n \left[ \psi_j^{(0)} \left( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \right) - \psi_j^{(0)} \left( \mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)} \right) \right] + \mathbb{P}_n \psi_j^{(0)} \left( \mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)} \right),$$

with  $\mathbb{P}_n \psi_j^{(0)} \left(\mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)}\right) \stackrel{\mathrm{P}}{\longrightarrow} P \psi_j^{(0)} \left(\mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)}\right) = 0$ . For another part, we will use empirical processes technique. Indeed, we have

$$\begin{split} & \left| \mathbb{P}_{n} \left[ \psi_{j}^{(0)} \left( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \right) - \psi_{j}^{(0)} \left( \mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)} \right) \right] \right| \\ \leq & \left| \mathbb{P}_{n} \int_{\mathcal{M}} \left\{ \widehat{\mu}(C, 1, m) - \mu(C, 1, m) \right\} \pi_{C,a}(m_{j}) \pi_{C,0}(m_{-j}) \, \mathrm{d}m \right| \\ & + \left| \mathbb{P}_{n} \int_{\mathcal{M}} \mu(C, 1, m) \left\{ \widehat{\pi}_{C,a}(m_{j}) - \pi_{C,a}(m_{j}) \right\} \pi_{C,0}(m_{-j}) \, \mathrm{d}m \right| \\ & + \left| \mathbb{P}_{n} \int_{\mathcal{M}} \mu(C, 1, m) \pi_{C,a}(m_{j}) \left\{ \widehat{\pi}_{C,0}(m_{-j}) - \pi_{C,0}(m_{-j}) \right\} \, \mathrm{d}m \right|. \end{split}$$

For sufficient small constant  $\varepsilon > 0$ , we define a set of functions  $\mathcal{U}(\varepsilon)$  that contains conditional expectation  $\mu_{\varepsilon}$  such that

$$E \int_{\mathcal{M}} \left\{ \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right\}^{2} dm \leq \varepsilon^{2}.$$

then we consider

$$\begin{split} & \operatorname{E} \sup_{\mu_{\varepsilon} \in \mathcal{U}(\varepsilon)} \left| \mathbb{P}_{n} \int_{\mathcal{M}} \left\{ \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right\} \pi_{C, a}(m_{j}) \pi_{C, 0}(m_{-j}) \, \mathrm{d}m \right| \\ \leq & \operatorname{E} \sup_{\mu_{\varepsilon} \in \mathcal{U}(\varepsilon)} \left[ \mathbb{P}_{n} \int_{\mathcal{M}} \left| \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right| \pi_{C, a}(m_{j}) \pi_{C, 0}(m_{-j}) \, \mathrm{d}m \right] \\ \leq & \mathbb{P}_{n} \sup_{\mu_{\varepsilon} \in \mathcal{U}(\varepsilon)} \operatorname{E} \int_{\mathcal{M}} \left| \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right| \pi_{C, a}(m_{j}) \pi_{C, 0}(m_{-j}) \, \mathrm{d}m \\ & + \operatorname{E} \sup_{\mu_{\varepsilon} \in \mathcal{U}(\varepsilon)} \mathbb{P}_{n} \left[ \int_{\mathcal{M}} \left| \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right| \pi_{C, a}(m_{j}) \pi_{C, 0}(m_{-j}) \, \mathrm{d}m \right] \\ & - \operatorname{E} \int_{\mathcal{M}} \left| \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right| \pi_{C, a}(m_{j}) \pi_{C, 0}(m_{-j}) \, \mathrm{d}m \right] \end{split}$$

Cauchy Inequality and  $\ell^2$  assumption <  $\varepsilon^2$ 

$$+ \operatorname{E} \sup_{\mu_{\varepsilon} \in \mathcal{U}(\varepsilon)} \mathbb{P}_{n} \left[ \int_{\mathcal{M}} \left| \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right| \pi_{C, a}(m_{j}) \pi_{C, 0}(m_{-j}) \, \mathrm{d}m \right. \\ \left. - \operatorname{E} \int_{\mathcal{M}} \left| \mu_{\varepsilon}(C, 1, m) - \mu(C, 1, m) \right| \pi_{C, a}(m_{j}) \pi_{C, 0}(m_{-j}) \, \mathrm{d}m \right]$$

Corollary 5.1 in Chernozhukov et al. (2014)  $\lesssim^2 + n^{-1/2} \sqrt{n^\vartheta} \varepsilon \big(\log(\varepsilon) + \log n\big)$ 

Let  $\varepsilon \to 0$  with Assumption 7.9 0.

Therefore, we must have  $\mathbb{P}_n \int_{\mathcal{M}} \{\widehat{\mu}(C,1,m) - \mu(C,1,m)\} \pi_{C,a}(m_j) \pi_{C,0}(m_{-j}) dm = o_p(1)$ . Similarly, we have

$$\mathbb{P}_{n} \int_{\mathcal{M}} \mu(C, 1, m) \{ \widehat{\pi}_{C, a}(m_{j}) - \pi_{C, a}(m_{j}) \} \pi_{C, 0}(m_{-j}) \, dm$$

$$= \mathbb{P}_{n} \int_{\mathcal{M}} \mu(C, 1, m) \pi_{C, a}(m_{j}) \{ \widehat{\pi}_{C, 0}(m_{-j}) - \pi_{C, 0}(m_{-j}) \} \, dm = o_{p}(1),$$

which implies

$$\mathbb{P}_n \Big[ \psi_j^{(0)} \big( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \big) - \psi_j^{(0)} \big( \mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)} \big) \Big] = o_p(1)$$

This yields  $\mathbb{P}_n \psi_j^{(0)} (\widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)}) = o_p(1)$ . Note that the expressions of  $\psi_j^{(3)}(\cdot)$  and Assumption 4.2 ensure

$$\left| \mathbb{P}_n \psi_j^{(3)} \left( \widehat{e}_1, \widehat{\pi}_{C, a'}^{(-j)}, \widehat{\pi}_{C, 1, M_j}^{(-j)}, \widehat{\mu} \right) \right| \lesssim \left| \frac{1}{n} \sum_{i=1}^n \left\{ Y_i - \widehat{\mu}(C_i, 1, M_i) \right\} \right| = o_p(1),$$

by  $\widehat{\mu}(\cdot)$  is correctly estimated, which gives  $\mathbb{P}_n \psi_j^{(3)} \left( \widehat{e}_1, \widehat{\pi}_{C,0}^{(j)}, \widehat{\pi}_{C,a'}^{(-j)}, \widehat{\pi}_{C,1,M_{-j}}^{(j)}, \widehat{\pi}_{C,1}^{(-j)}, \widehat{\mu} \right) = o_p(1)$ . Similarly,

$$\begin{split} & \left| \mathbb{P}_{n} \psi_{j}^{(1)} \left( \widehat{e}_{0}, \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)} \right) \right| \\ \lesssim & \mathbb{P}_{n} \left[ \int_{\mathcal{M}_{j}} \widehat{\mu}(C, 1, m_{j}, M_{-j}) \widehat{\pi}_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \zeta_{j}(a', 0, C) \right] \\ \leq & \left| \mathbb{P}_{n} \left[ \int_{\mathcal{M}_{j}} \mu(C, 1, m_{j}, M_{-j}) \pi_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \zeta_{j}(a', 0, C) \right] \right| \\ & + \left| \mathbb{P}_{n} \int_{\mathcal{M}_{j}} \left\{ \widehat{\mu}(C, 1, m_{j}, M_{-j}) \widehat{\pi}_{C,a'}(m_{j}) - \mu(C, 1, m_{j}, M_{-j}) \pi_{C,a'}(m_{j}) \right\} \, \mathrm{d}m_{j} \right|. \end{split}$$

The first term above is  $o_p(1)$  obviously by the definition of  $\zeta_j(a',0,C)$ . By using the same steps for proving  $\mathbb{P}_n\left[\psi_j^{(0)}\left(\widehat{\mu},\widehat{\pi}_{C,a'}^{(j)},\widehat{\pi}_{C,0}^{(-j)}\right)-\psi_j^{(0)}\left(\mu,\pi_{C,a'}^{(j)},\pi_{C,0}^{(-j)}\right)\right]=o_p(1)$  above, we can prove the second term is also  $o_p(1)$  by both  $\widehat{\mu}$  and  $\widehat{\pi}_{C,a'}(m_j)$  are correctly estimated. This yields  $\mathbb{P}_n\psi_j^{(1)}\left(\widehat{e}_0,\widehat{\mu},\widehat{\pi}_{C,a'}^{(j)}\right)=o_p(1)$ , and similarly gives  $\mathbb{P}_n\psi_j^{(2)}\left(\widehat{e}_{a'},\widehat{\mu},\widehat{\pi}_{C,0}^{(-j)}\right)=o_p(1)$ . The proof for **Part 0** is thus completed.

Part 1. When the model  $\mathcal{M}_{j,1}$  is correctly specified, we have consistency estimator  $\widehat{e}_{a'}$ ,  $\widehat{\mu}$ , and  $\widehat{\pi}_{C,a'}^{(j)}$ . We first handle  $\psi_j^{(0)}(\cdot) + \psi_j^{(1)}(\cdot)$ . We rewrite

$$\begin{split} \psi_{j}^{(0)} \left( \mu, \pi_{C,a'}^{(j)}, \pi_{C,0}^{(-j)} \right) + \psi_{j}^{(1)} \left( e_{0}, \mu, \pi_{C,a'}^{(j)} \right) \\ = & \zeta_{j}(a', 0, C) - \mathrm{E}\zeta_{j}(a', 0, C) + \frac{\mathbb{1}(A = 0)}{e_{0}(C)} \left\{ \int_{\mathcal{M}_{j}} \mu(C, 1, m_{j}, M_{-j}) \pi_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \zeta_{j}(a', 0, C) \right\} \\ = & \frac{\zeta_{j}(a', 0, C)}{e_{0}(C)} \left[ \mathbb{1}(A = 0) - e_{0}(C) \right] \\ & + \frac{\mathbb{1}(A = 0)}{e_{0}(C)} \left\{ \int_{\mathcal{M}_{j}} \mu(C, 1, m_{j}, M_{-j}) \pi_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \mathrm{E}\zeta_{j}(a', 0, C). \end{split}$$

Therefore,

$$\begin{split} & \left| \mathbb{P}_{n} \left[ \psi_{j}^{(0)} \left( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \right) + \psi_{j}^{(1)} \left( \widehat{e}_{0}, \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)} \right) \right] \right| \\ & \lesssim \left| \mathbb{P}_{n} \left[ \frac{\int_{\mathcal{M}} \widehat{\mu}(C, 1, m) \widehat{\pi}_{C,a}(m_{j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m}{\widehat{e}_{0}(C)} \left\{ \mathbb{1}(A = 0) - \widehat{e}_{0}(C) \right\} \right] \right| \\ & + \left| \mathbb{P}_{n} \left[ \frac{\mathbb{1}(A = 0)}{\widehat{e}_{0}(C)} \int_{\mathcal{M}_{j}} \widehat{\mu}(C, 1, m_{j}, M_{-j}) \widehat{\pi}_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \mathrm{E}\zeta_{j}(a', 0, C) \right] \right|. \end{split}$$

By the bounded assumption in Assumption 7.9, we have

$$\left| \mathbb{P}_n \left[ \frac{\int_{\mathcal{M}} \widehat{\mu}(C, 1, m) \widehat{\pi}_{C, a}(m_j) \widehat{\pi}_{C, 0}(m_{-j}) \, \mathrm{d}m}{\widehat{e}_0(C)} \left\{ \mathbb{1}(A = 0) - \widehat{e}_0(C) \right\} \right] \right|$$

$$\lesssim \left| \mathbb{P}_n \left\{ \mathbb{1}(A = 0) - \widehat{e}_0(C) \right\} \right| = o_p(1)$$

by  $\widehat{e}_0(\cdot)$  convergences to  $e_0(\cdot)$  in  $\ell^2$  norm and  $\mathrm{E}\big\{\mathbb{1}(A=0)-e_0(C)\big\}=0$ . On the other hand, we have

$$\begin{split} \mathbb{P}_{n} \bigg[ \frac{\mathbb{1}(A=0)}{\widehat{e}_{0}(C)} \bigg\{ \int_{\mathcal{M}_{j}} \widehat{\mu}(C,1,m_{j},M_{-j}) \widehat{\pi}_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \mathrm{E}\zeta_{j}(a',0,C) \bigg] \\ & = \sum_{j=0}^{n} \mathbb{1}(A=0) \sum_{C,a'} \mathbb{P}_{n} \bigg[ \frac{\mathbb{1}(A=0)}{e_{0}(C)} \int_{\mathcal{M}_{j}} \mu(C,1,m_{j},M_{-j}) \pi_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \mathrm{E}\zeta_{j}(a',0,C) \bigg] + o_{p}(1) \\ & = \mathrm{E} \bigg[ \frac{\mathbb{1}(A=0)}{e_{0}(C)} \int_{\mathcal{M}_{j}} \mu(C,1,m_{j},M_{-j}) \pi_{C,a'}(m_{j}) \, \mathrm{d}m_{j} - \mathrm{E}\zeta_{j}(a',0,C) \bigg] + o_{p}(1) \\ & = \sum_{j=0}^{n} o_{p}(1), \end{split}$$

which gives  $\mathbb{P}_n \big[ \psi_j^{(0)} \big( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \big) + \psi_j^{(1)} \big( \widehat{e}_0, \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)} \big) \big] = o_p(1)$ . It remains to show  $\mathbb{P}_n \big[ \psi_j^{(2)} \big( \widehat{e}_{a'}, \widehat{\mu}, \widehat{\pi}_{C,0}^{(-j)} \big) + \psi_j^{(3)} \big( \widehat{e}_1, \widehat{\pi}_{C,a'}^{(-j)}, \widehat{\pi}_{C,1,M_j}^{(-j)}, \widehat{\mu} \big) \big] = o_p(1)$ . Indeed, the consistency of  $\widehat{\mu}$  ensures  $\mathbb{P}_n \big[ \psi_j^{(3)} \big( \widehat{e}_1, \widehat{\pi}_{C,a'}^{(-j)}, \widehat{\pi}_{C,1,M_j}^{(-j)}, \widehat{\mu} \big) \big] = o_p(1)$  as stated in **Part 0**, and

$$\begin{split} \left| \mathbb{P}_n \left[ \psi_j^{(2)} \left( \widehat{e}_{a'}, \widehat{\mu}, \widehat{\pi}_{C,0}^{(-j)} \right) \right] \right| \\ &= \left| \mathbb{P}_n \left[ \frac{\mathbb{1}(A = a')}{\widehat{e}_{a'}(C)} \left\{ \int_{\mathcal{M}_{-j}} \widehat{\mu}(C, 1, M_j, m_{-j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m_{-j} \right. \right. \\ &\left. \left. \left. \left. - \int_{\mathcal{M}} \widehat{\mu}(C, 1, m) \widehat{\pi}_{C,1}(m_j) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m \right\} \right] \right| \\ &\leq \mathbb{P}_n \left[ \frac{\mathbb{1}(A = a')}{\widehat{e}_{a'}(C)} \int_{\mathcal{M}_{-j}} \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m_{-j} \right| \int_{\mathcal{M}_j} \widehat{\mu}(C, 1, m_j, m_{-j}) \left[ \pi_{C,1}(m_j) - \widehat{\pi}_{C,1}(m_j) \right] \, \mathrm{d}m_j \right| \end{split}$$
by  $\ell^2$  assumption and Cauchy

by 
$$\ell^2$$
 assumption and Cauchy 
$$\lesssim \mathbb{P}_n \bigg| \int_{\mathcal{M}_j} \big[ \pi_{C,1}(m_j) - \widehat{\pi}_{C,1}(m_j) \big]^2 \, \mathrm{d}m_j \bigg|$$

$$= \mathrm{E} \bigg| \int_{\mathcal{M}_j} \big[ \pi_{C,1}(m_j) - \widehat{\pi}_{C,1}(m_j) \big]^2 \, \mathrm{d}m_j \bigg| + o_p(1) = o_p(1),$$

which completes the proof required for Part 1.

**Part 2.** Exactly the same as **Part 1**, one can prove when model  $\mathcal{M}_{j,2}$  is correctly specified, we have

$$\begin{split} \left| \mathbb{P}_{n} \left[ \psi_{j}^{(0)} \left( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \right) + \psi_{j}^{(2)} \left( \widehat{e}_{a'}, \widehat{\mu}, \widehat{\pi}_{C,0}^{(-j)} \right) \right] \right| \\ = \left| \mathbb{P}_{n} \left[ \int_{\mathcal{M}} \widehat{\mu}(C, 1, m) \widehat{\pi}_{C,1}(m_{j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m - \mathrm{E}\zeta_{j}(a', 0, C) \right. \\ \left. + \frac{\mathbb{1}(A = a')}{\widehat{e}_{a'}(C)} \left\{ \int_{\mathcal{M}_{-j}} \widehat{\mu}(C, 1, M_{j}, m_{-j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m_{-j} - \zeta_{j}(a', 0, C) \right\} \right] \right| \\ \leq \left| \mathbb{P}_{n} \frac{\int_{\mathcal{M}} \widehat{\mu}(C, 1, m) \widehat{\pi}_{C,a}(m_{j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m}{e_{a'}(C)} \left[ \mathbb{1}(A = a') - \widehat{e}_{a'}(C) \right] \right| \\ \left. + \left| \mathbb{P}_{n} \left[ \frac{\mathbb{1}(A = a')}{\widehat{e}_{a'}(C)} \int_{\mathcal{M}_{-j}} \widehat{\mu}(C, 1, M_{j}, m_{-j}) \widehat{\pi}_{C,0}(m_{-j}) \, \mathrm{d}m_{-j} - \mathrm{E}\zeta_{j}(a', 0, C) \right] \right| \\ \lesssim \mathrm{E} \left[ \mathbb{1}(A = a') - \widehat{e}_{a'}(C) \right]^{2} \\ \left. + \mathrm{E} \left[ \frac{\mathbb{1}(A = a')}{e_{a'}(C)} \int_{\mathcal{M}_{-j}} \mu(C, 1, M_{j}, m_{-j}) \pi_{C,0}(m_{-j}) \, \mathrm{d}m_{-j} - \mathrm{E}\zeta_{j}(a', 0, C) \right] + o_{p}(1) \right. \\ \overset{\mathrm{by}}{=} \underbrace{\left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right.} \right. \\ \left. + \left. \left[ 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \\ \left. \begin{array}{c} 0 \\ 0 \end{array} \right]$$

Similarly, we can prove

$$\begin{split} & \left| \mathbb{P}_{n} \psi_{j}^{(1)}(\widehat{e}_{0}, \widehat{\mu}, \widehat{\pi}_{C, a'}^{(j)}) \right| \\ \leq & \left| \mathbb{P}_{n} \left[ \frac{\mathbb{1}(A = 0)}{e_{0}(C)} \left\{ \int_{\mathcal{M}_{j}} \widehat{\pi}_{C, a'}(m_{j}) \, \mathrm{d}m_{j} \int_{\mathcal{M}_{-j}} \widehat{\mu}(C, 1, m_{j}, m_{-j}) \left[ \pi_{C, 0}(m_{-j}) - \widehat{\pi}_{C, 0}(m_{-j}) \right] \, \mathrm{d}m_{-j} \right] \right| \\ \lesssim & \mathbb{P}_{n} \left| \int_{\mathcal{M}_{-j}} \left[ \pi_{C, 0}(m_{-j}) - \widehat{\pi}_{C, 0}(m_{-j}) \right]^{2} \, \mathrm{d}m_{-j} \right| = o_{p}(1) \end{split}$$

and  $\mathbb{P}_n \big[ \psi_j^{(3)} \big( \widehat{e}_1, \widehat{\pi}_{C,a'}^{(-j)}, \widehat{\pi}_{C,1,M_j}^{(-j)}, \widehat{\mu} \big) \big] = o_p(1)$ , which completes the proof of this part.

**Part 3.** When the model  $\mathcal{M}_{j,3}$  is correctly specified, we have  $\widehat{e}_1$ ,  $\widehat{\pi}_{C,a'}^{(-j)}$ , and  $\widehat{\pi}_{C,1,M_j}^{(-j)}$  are consistent, as well as  $\widehat{\pi}_{C,a',\operatorname{Pa}_i(\mathcal{G}_M)}^{(j)}$  thus  $\widehat{\pi}_{C,a'}^{(j)}$ , but not  $\widehat{\mu}$ . In this case, we first rewrite

$$\begin{split} \psi_{j}^{(0)}\left(\mu,\pi_{C,a'}^{(j)},\pi_{C,0}^{(-j)}\right) + \psi_{j}^{(3)}\left(e_{1},\pi_{C,a'}^{(-j)},\pi_{C,1,M_{j}}^{(-j)},\mu\right) \\ = & \left[\frac{\mathbb{1}(A=1)}{e_{1}(C)}\frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_{j}}(M_{-j})}Y - \mathrm{E}\zeta_{j}(a',0,C)\right] \\ & + \int_{\mathcal{M}}\mu(C,1,m)\pi_{C,1}(m_{j})\pi_{C,a'}(m_{-j})\,\mathrm{d}m - \frac{\mathbb{1}(A=1)}{e_{1}(C)}\frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_{j}}(M_{-j})}\mu(C,1,M). \end{split}$$

By using the same empirical processes technique in Part 0, we can prove

$$\begin{split} & \left| \mathbb{P}_n \left[ \psi_j^{(0)} \left( \widehat{\mu}, \widehat{\pi}_{C,a'}^{(j)}, \widehat{\pi}_{C,0}^{(-j)} \right) + \psi_j^{(3)} \left( \widehat{e}_1, \widehat{\pi}_{C,a'}^{(-j)}, \widehat{\pi}_{C,1,M_j}^{(-j)}, \widehat{\mu} \right) \right] \right| \\ & \leq \left| \mathbb{P}_n \left[ \frac{\mathbb{I}(A=1)}{\widehat{e}_1(C)} \frac{\widehat{\pi}_{C,a'}(M_{-j})}{\widehat{\pi}_{C,1,M_j}(M_{-j})} Y - \mathbb{E}\zeta_j(a',0,C) \right] \right| \\ & + \left| \mathbb{P}_n \left[ \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \widehat{\pi}_{C,1}(m_j) \widehat{\pi}_{C,a'}(m_{-j}) \, \mathrm{d}m - \frac{\mathbb{I}(A=1)}{\widehat{e}_1(C)} \frac{\widehat{\pi}_{C,a'}(M_{-j})}{\widehat{\pi}_{C,1,M_j}(M_{-j})} \widehat{\mu}(C,1,M) \right] \right| \\ & = \left| \mathbb{E} \left[ \frac{\mathbb{I}(A=1)}{e_1(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_j}(M_{-j})} Y - \mathbb{E}\zeta_j(a',0,C) \right] + o_p(1) \right| \\ & + \left| \mathbb{E} \left[ \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \pi_{C,1}(m_j) \pi_{C,a'}(m_{-j}) \, \mathrm{d}m - \frac{\mathbb{I}(A=1)}{e_1(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_j}(M_{-j})} \widehat{\mu}(C,1,M) \right] + o_p(1) \right| \\ & = \mathbb{E} \left[ \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \pi_{C,1}(m_j) \pi_{C,a'}(m_{-j}) \, \mathrm{d}m - \mathbb{E} \left[ \frac{\mathbb{I}(A=1)}{e_1(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_j}(M_{-j})} \widehat{\mu}(C,1,M) \right] \right| + o_p(1) \\ & = \left| \mathbb{E} \left[ \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \pi_{C,1}(m_j) \pi_{C,a'}(m_{-j}) \, \mathrm{d}m - \mathbb{E} \left[ \frac{\mathbb{I}(A=1)}{e_1(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_j}(M_{-j})} \widehat{\mu}(C,1,M) \right] C \right] \right] \right| + o_p(1) \\ & = 0 + o_p(1) = o_p(1), \end{split}$$

where we use the fact that

$$\begin{split} & \operatorname{E}\left[\frac{\mathbbm{1}(A=1)}{e_{1}(C)} \frac{\pi_{C,a'}(M_{-j})}{\pi_{C,1,M_{j}}(M_{-j})} \widehat{\mu}(C,1,M) \mid C\right] \\ & = \int_{\mathcal{M}} \frac{1}{f(a=1\mid C)} \frac{f(m_{-j}\mid C, a=a')}{f(m_{-j}\mid C, a=1, m_{j})} \widehat{\mu}(C,1,m) \mathbbm{1}(a=1) f(a,m\mid C) \, \mathrm{d}m \\ & = \int_{\mathcal{M}} \frac{1}{f(a=1\mid C)} \frac{f(m_{-j}\mid C, a=a')}{f(m_{-j}\mid C, a=1, m_{j})} \widehat{\mu}(C,1,m) \\ & \qquad \times f(a=1\mid C) f(m_{j}\mid C, a=1) f(m_{-j}\mid C, a=1, m_{j}) \, \mathrm{d}m \\ & = \int_{\mathcal{M}} \widehat{\mu}(C,1,m) f(m_{j}\mid C, a=1) f(m_{-j}\mid C, a=a') \, \mathrm{d}m = \int_{\mathcal{M}} \widehat{\mu}(C,1,m) \pi_{C,1}(m_{j}) \pi_{C,a'}(m_{-j}) \, \mathrm{d}m. \end{split}$$

We can also prove that

$$\left| \mathbb{P}_n \psi_j^{(1)}(\widehat{e}_0, \widehat{\mu}, \widehat{\pi}_{C, a'}^{(j)}) \right| \lesssim \mathbb{P}_n \left| \int_{\mathcal{M}_{-j}} \left[ \pi_{C, 0}(m_{-j}) - \widehat{\pi}_{C, 0}(m_{-j}) \right]^2 dm_{-j} \right| = o_p(1)$$

and

$$\begin{split} \left| \mathbb{P}_n \left[ \psi_j^{(2)} \left( \widehat{e}_{a'}, \widehat{\mu}, \widehat{\pi}_{C,0}^{(-j)} \right) \right] \right| &\lesssim \mathbb{P}_n \left| \int_{\mathcal{M}_j} \left[ \pi_{C,1}(m_j) - \widehat{\pi}_{C,1}(m_j) \right]^2 dm_j \right| \\ &= \mathbb{P}_n \left| \int_{\mathcal{M}_j} \int_{\mathcal{M}_{\mathrm{pa}_j}(\mathcal{G}_M)} \left[ \pi_{C,1,\mathrm{pa}_j(\mathcal{G}_M)}(m_j) - \widehat{\pi}_{C,1,\mathrm{pa}_j(\mathcal{G}_M)}(m_j) \right]^2 dm_j \right| \\ &= o_p(1). \end{split}$$

Thus, we finish proof of **Part 3**. The proof of consistency of  $\widehat{DM}_{j}^{QR}$  is hence completed.

# Semiparametric Efficiency of $\widehat{TM}_{i}^{\text{avg, QR}}$ :

Next, we want to prove the semiparametric efficiency for  $\widehat{TM}_j^{\text{avg, QR}}$ . Here we have all estimators in model  $\mathcal{M}_{j,\ell}, \ell=0,1,2,3$  are correct. We use  $\widehat{P}$  to denote these correct estimated functions or estimated distribution, which is a consistent estimator of the true law of X, P. First note that, for any fixed  $\mathcal{G}_M \in \mathrm{MEC}(\widehat{\mathcal{C}}_M)$  and  $j \in [p]$ ,  $\widehat{TM}_i^{\mathrm{QR}}(\mathcal{G}_M)$  can be written as the following one-step estimators

$$\begin{split} \widehat{TM}_{j}^{\mathrm{QR}}(\mathcal{G}_{M}) \\ =& \mathbb{P}_{n} \left[ \frac{\mathbb{I}(A=1)}{\widehat{e}_{1}(C)} \Big\{ Y - \widehat{\mu}(C,1) \Big\} - \frac{\mathbb{I}(A=0)}{\widehat{e}_{0}(C)} \Big\{ Y - \widehat{\mu}(C,0) \Big] \Big\} \right] \\ -& \mathbb{P}_{n} \left[ \frac{\mathbb{I}(A=1)}{\widehat{e}_{1}(C)} \widehat{\pi}_{C,1} \Big( \mathrm{Pa}_{j}(\mathcal{G}_{M}) \Big) \Big\{ Y - \widehat{\mu}(C,1,\mathrm{Pa}_{j}(\mathcal{G}_{M}),M_{j}) \Big\} \\ & - \frac{\mathbb{I}(A=0)}{\widehat{e}_{0}(C)} \widehat{\pi}_{C,0} \Big( \mathrm{Pa}_{j}(\mathcal{G}_{M}) \Big) \Big\{ Y - \widehat{\mu}(C,0,\mathrm{Pa}_{j}(\mathcal{G}_{M}),M_{j}) \Big\} \\ +& \int_{\mathcal{M}_{j}} \left( \frac{\mathbb{I}(A=1)}{\widehat{e}_{1}(C)} \widehat{\mu}(C,1,\mathrm{Pa}_{j}(\mathcal{G}_{M}),m_{j}) - \frac{\mathbb{I}(A=0)}{\widehat{e}_{0}(C)} \widehat{\mu}(C,0,\mathrm{Pa}_{j}(\mathcal{G}_{M}),m_{j}) \Big) \widehat{\pi}_{C}(m_{j}) \, \mathrm{d}m_{j} \\ -& \left( \frac{\mathbb{I}(A=1)}{\widehat{e}_{1}(C)} \widehat{\mathbb{E}} \Big[ \varrho_{j}(1,M_{j},C\,;\mathcal{G}_{M}) \mid C \Big] - \frac{\mathbb{I}(A=0)}{\widehat{e}_{0}(C)} \widehat{\mathbb{E}} \Big[ \varrho_{j}(0,M_{j},C\,;\mathcal{G}_{M}) \mid C \Big] \right) \Big] + \widehat{TM}_{j}^{\mathcal{M}_{0}} \\ =&: \Big[ \mathbb{P}_{n} \varphi_{j,1}^{\mathrm{QR}}(X,\widehat{P}\,;\mathcal{G}_{M}) + \psi_{j,1}(\widehat{P}\,;\mathcal{G}_{M}) \Big] - \Big[ \mathbb{P}_{n} \varphi_{j,0}^{\mathrm{QR}}(X,\widehat{P}\,;\mathcal{G}_{M}) + \psi_{j,0}(\widehat{P}\,;\mathcal{G}_{M}) \Big] \\ \text{where } \psi_{j,a'} \text{ with } a' = 0,1 \text{ is defined as} \\ \psi_{j,a'} : F_{X} \times \mathcal{G}_{M} \mapsto \mathbb{E} \Big[ \Big\{ \mu(a',C) - \int_{\mathbb{M}_{n}(\mathcal{G}_{N})} \mu(C,a',\mathrm{pa}_{j},M_{j}) \, \pi_{C,a'}(\mathrm{pa}_{j}) \, \mathrm{d}\, \mathrm{pa}_{j} \Big\} \Big]. \end{split}$$

$$\psi_{j,a'}: F_X \times \mathcal{G}_M \mapsto \mathbf{E}\left[\left\{\mu(a',C) - \int_{\mathcal{M}_{\mathrm{pa}_j}(\mathcal{G}_M)} \mu(C,a',\mathrm{pa}_j,M_j) \,\pi_{C,a'}(\mathrm{pa}_j) \,\mathrm{d}\,\mathrm{pa}_j\right\}\right].$$

and  $\varphi_{j,a'}$  is the mapping from the observation and the underlying distribution of X to the efficient score for  $\psi_{j,a'}$ , from Theorem 4.3, we know that

$$\begin{split} & \varphi_{j,a'}^{QR}(X,Q;\mathcal{G}_{M}) \\ := & \frac{\mathbb{1}(A=a')}{e_{a'}^{Q}(C)} \Big\{ Y - \kappa^{Q}(a',C) \Big\} + \kappa^{Q}(a',C) - \left[ \frac{\mathbb{1}(A=a')}{e_{a'}^{Q}(C)} \pi_{C,a'}^{Q} \Big( \operatorname{Pa}_{j}(\mathcal{G}_{M}) \Big) \Big\{ Y - \mu^{Q}(C,a',M) \Big\} \right] \\ & - \left[ \frac{\mathbb{1}(A=a')}{e_{a'}^{Q}(C)} \left\{ \int_{\mathcal{M}_{j}} \mu^{Q}(C,a',\operatorname{Pa}_{j}(\mathcal{G}_{M}),m_{j}) \pi_{C}^{Q}(m_{j}) \, \mathrm{d}m_{j} - \operatorname{E}_{P} \Big[ \varrho_{j}^{Q}(a',M_{j},C;\mathcal{G}_{M}) \mid C \Big] \right\} \\ & + \varrho_{j}^{Q}(a',M_{j},C;\mathcal{G}_{M}) \right] - \psi_{j,a'}(Q;\mathcal{G}_{M}) \end{split}$$

$$= \varphi_{j,a'}^{(1)}(X,Q;\mathcal{G}_M) - \varphi_{j,a'}^{(2)}(X,Q;\mathcal{G}_M) - \psi_{j,a'}(Q;\mathcal{G}_M)$$

for arbitrary distribution Q for X, where  $e^Q_{a'}(c) := Q(A=a'\mid c)$  and similar definition for  $k^Q(a',c),\,\pi^Q_S(m_T),\,\mu^Q(\cdot),$  and  $\rho^Q_j(a',m_j,c\,;\mathcal{G}_M),$  where

$$\varphi_{j,a'}^{(1)}(X,Q;\mathcal{G}_{M}) := \frac{\mathbb{1}(A=a')}{e_{a'}^{Q}(C)} \Big\{ Y - \kappa^{Q}(a',C) \Big\} + \kappa^{Q}(a',C) - \left[ \frac{\mathbb{1}(A=a')}{e_{a'}^{Q}(C)} \pi_{C,a'}^{Q} \Big( \operatorname{Pa}_{j}(\mathcal{G}_{M}) \Big) \Big\{ Y - \mu^{Q}(C,a',M) \Big\} \right] \\
+ \varrho_{j}^{Q}(a',M_{j},C;\mathcal{G}_{M}),$$

and

$$\varphi_{j,a'}^{(2)}(X,Q;\mathcal{G}_{M}) := \frac{\mathbb{1}(A=a')}{e^{Q}(C)} \left\{ \int_{M} \mu^{Q}(C,a',\operatorname{Pa}_{j}(\mathcal{G}_{M}),m_{j}) \pi_{C}^{Q}(m_{j}) \, \mathrm{d}m_{j} - \operatorname{E}_{P} \left[ \varrho_{j}^{Q}(a',M_{j},C;\mathcal{G}_{M}) \mid C \right] \right\}$$

We can see that  $\psi_{j,1} - \psi_{j,0}$  the functional mapping of the underlying distribution of X to  $TM_j(\mathcal{G}_M)$  for a fixed DAG  $\mathcal{G}_M$ . Therefore, we can decompose

$$\begin{split} &\widehat{TM}_{j}^{\mathrm{QR}}(\mathcal{G}_{M}) - TM_{j}(\mathcal{G}_{M}) \\ = & \Big\langle \left( \mathbb{P}_{n} - P \right) \phi_{j,\cdot}^{\mathrm{QR}}(X, P \, ; \, \mathcal{G}_{M}) + \left( \mathbb{P}_{n} - P \right) \left[ \phi_{j,\cdot}^{\mathrm{QR}}(X, \widehat{P} \, ; \, \mathcal{G}_{M}) - \phi_{j,\cdot}^{\mathrm{QR}}(X, P \, ; \, \mathcal{G}_{M}) \right] + R_{j,\cdot}^{\mathrm{QR}}(\widehat{P}, P \, ; \, \mathcal{G}_{M}) \Big\rangle \end{split}$$
 where

$$R_{j,a'}^{QR}(\widehat{P},P;\mathcal{G}_M) = \psi_{j,a'}(\widehat{P};\mathcal{G}_M) - \psi_{j,a'}(P;\mathcal{G}_M) + \int \varphi_{j,a'}^{QR}(x,\widehat{P};\mathcal{G}_M) dP(x).$$

Given Assumption 7.4, by using the argument in Proof of Theorem 7.8, we can prove (C.7)

$$\begin{split} \widehat{TM}_{j}^{\text{avg, QR}} - \overline{TM}_{j} \\ = & \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} (\mathbb{P}_{n} - P) \left\{ \varphi_{j,1}^{\operatorname{QR}}(X, P; \mathcal{G}_{M}) - \varphi_{j,0}^{\operatorname{QR}}(X, P; \mathcal{G}_{M}) \right\} \\ + & \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} (\mathbb{P}_{n} - P) \left[ \left\{ \varphi_{j,1}^{\operatorname{QR}}(X, \widehat{P}; \mathcal{G}_{M}) - \varphi_{j,1}^{\operatorname{QR}}(X, P; \mathcal{G}_{M}) \right\} \\ & - \left\{ \varphi_{j,0}^{\operatorname{QR}}(X, \widehat{P}; \mathcal{G}_{M}) - \varphi_{j,0}^{\operatorname{QR}}(X, P; \mathcal{G}_{M}) \right\} \right] \\ + & \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} \left\{ R_{j,1}^{\operatorname{QR}}(\widehat{P}, P; \mathcal{G}_{M}) - R_{j,0}^{\operatorname{QR}}(\widehat{P}, P; \mathcal{G}_{M}) \right\} + o_{p}(\ell_{n}^{-1}) \end{split}$$

with any  $\ell_n$  such that  $\lim_{n\to\infty}\ell_n=\infty$ . The (vector-)function classes for  $\varphi_{j,a'}^{\operatorname{QR}}(X,P;\mathcal{G}_M)$  with  $a'\in\{0,1\}$  is  $q=(q_k)_{k\in\mathcal{K}}\in\mathcal{Q}:=\bigotimes_{\{T,S\}\text{ used in }(6.8)\text{ and }(6.9)\text{ for a fixed }j\in[p]}\mathcal{E}\otimes\mathcal{F}_{T\mid S}\otimes\mathcal{U}_S$ , which is also VC-class with index the same index  $\vartheta_j\in[0,1/2)$  by Lemma 2.6.18 in Van Der Vaart et al. (1996). On the other hand, note that the consistency in Assumption 7.10 ensures there exists sufficient large n such that all denominators in  $\varphi_{j,a'}^{\operatorname{QR}}(X,\widehat{P};\mathcal{G}_M)$  will be larger than  $\varepsilon/2$ , then by using  $|a_1a_2-b_1b_2|\leq |b_1||a_2-b_2|+|b_2||a_1-b_1|+|a_1-a_2||b_1-b_2|$ 

for any real number  $a_1, a_2, b_1, b_2$ , we can show that (C.8)

$$\begin{split} &\left| (\mathbb{P}_{n} - P) \left\{ \varphi_{j,a'}^{QR}(X, \widehat{P}; \mathcal{G}_{M}) - \varphi_{j,a'}^{QR}(X, P; \mathcal{G}_{M}) \right\} \right| \\ & \text{by $\ell^{2}$ assumption} \\ & \lesssim \left| (\mathbb{P}_{n} - P) \left\{ \widehat{q}_{k}(X; \mathcal{G}_{M}) - q_{k}(X; \mathcal{G}_{M}) \right\} \right| \\ & \text{comes from $\varphi_{j,a'}^{(1)}(X, \widehat{P}; \mathcal{G}_{M}) - \varphi_{j,a'}^{(1)}(X, P; \mathcal{G}_{M})$} \\ & + \left| (\mathbb{P}_{n} - P) \int_{\mathcal{M}_{\mathrm{ps}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\pi}_{C,a'} \left( \mathrm{pa}_{j}(\mathcal{G}_{M}) \right) - \pi_{C,a'} \left( \mathrm{pa}_{j}(\mathcal{G}_{M}) \right) \right] \mathrm{dpa}_{j}(\mathcal{G}_{M}) \\ & + \left| (\mathbb{P}_{n} - P) \int_{\mathcal{M}_{\mathrm{ps}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\mu}(C, a, \mathrm{pa}_{j}, M_{j}) - \mu(C, a, \mathrm{pa}_{j}, M_{j}) \right] \mathrm{dpa}_{j} \right| \\ & \text{comes from $\varphi_{j,a'}^{(2)}(X, \widehat{P}; \mathcal{G}_{M}) - \varphi_{j,a'}^{(2)}(X, P; \mathcal{G}_{M})$} \\ & + \left| (\mathbb{P}_{n} - P) \int_{\mathcal{M}_{\mathrm{ps}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\pi}_{C,a}(\mathrm{pa}_{j}) - \pi_{C,a}(\mathrm{pa}_{j}) \right] \mathrm{dpa}_{j} \right| \\ & \text{comes from $\psi_{j,a'}(\widehat{P}; \mathcal{G}_{M}) - \psi_{j,a'}(P; \mathcal{G}_{M})$} \\ & + \left| (\mathbb{P}_{n} - P) \left[ \widehat{\mu}(a', C) - \mu(a', C) \right] \right| \\ & \text{comes from $\psi_{j,a'}(\widehat{P}; \mathcal{G}_{M}) - \psi_{j,a'}(P; \mathcal{G}_{M})$} \\ & + \left| (\mathbb{P}_{n} - P) \left[ \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\mu}(C, a', \mathrm{pa}_{j}, M_{j}) - \mu(C, a', \mathrm{pa}_{j}, M_{j}) \right] \mathrm{dpa}_{j} \right] \right| \\ & \text{comes from $\psi_{j,a'}(\widehat{P}; \mathcal{G}_{M}) - \psi_{j,a'}(P; \mathcal{G}_{M})$} \\ & + \left| (\mathbb{P}_{n} - P) \left[ \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\pi}_{C,a'}(\mathrm{pa}_{j}) - \pi_{C,a'}(\mathrm{pa}_{j}) \right] \mathrm{dpa}_{j} \right] \right| . \end{aligned}$$

For any  $k \in \mathcal{K}$ ,  $\widehat{q}_k$  converge with  $\ell^2$ -norm to  $q_k$  at a rate of  $n^{-\vartheta_{q_k}^*}$  in class  $\mathcal{Q}$  with  $\vartheta_{q_k}^* \in \{\vartheta_{j,e}^*, \vartheta_{j,\pi}^*, \vartheta_{j,\mu}^*\}$ . Then by Corollary 5.1 in Chernozhukov et al. (2014), we have

$$\mathbb{E}\left| (\mathbb{P}_n - P) \left\{ \widehat{q}_k(X; \mathcal{G}_M) - q_k(X; \mathcal{G}_M) \right\} \right| \\
\approx n^{-1/2} \sqrt{n^{\vartheta_j}} n^{-\vartheta_{q_k}^*} \left( \log n^{-\vartheta_{q_k}^*} + \log n \right) \\
\approx O(n^{-1/2 + \vartheta_j/2 - \vartheta_{q_k}^*} \log n) = o(n^{-1/2})$$

by  $\vartheta_j/2 - \min\{\vartheta_{j,e}^*, \vartheta_{j,\pi}^*, \vartheta_{j,\mu}^*\} < 0$  in Assumption 7.10, which implies  $(\mathbb{P}_n - P)\{\widehat{q}_k(X; \mathcal{G}_M) - q_k(X; \mathcal{G}_M)\} = o_p(n^{-1/2})$  for any  $\mathcal{G}_M \in \mathrm{MEC}(\mathcal{C}_M)$ . Thus,

$$\begin{split} & (\mathbb{P}_n - P) \left[ \widehat{\mu}(a', C) - \mu(a', C) \right] = o_p(n^{-1/2}), \\ & (\mathbb{P}_n - P) \left[ \int_{\mathcal{M}_{\mathrm{pa}_j}(\mathcal{G}_M)} \left[ \widehat{\mu}(C, a', \mathrm{pa}_j, M_j) - \mu(C, a', \mathrm{pa}_j, M_j) \right] \mathrm{d} \, \mathrm{pa}_j \right] = o_p(n^{-1/2}), \\ & (\mathbb{P}_n - P) \left[ \int_{\mathcal{M}_{\mathrm{pa}_j}(\mathcal{G}_M)} \left[ \widehat{\pi}_{C, a'}(\mathrm{pa}_j) - \pi_{C, a'}(\mathrm{pa}_j) \right] \mathrm{d} \, \mathrm{pa}_j = o_p(n^{-1/2}). \end{split}$$

Similarly, for any  $\mathcal{G}_M \in \mathrm{MEC}(\mathcal{C}_M)$  we can prove

$$\begin{aligned}
& \mathbf{E} \left[ (\mathbb{P}_{n} - P) \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\pi}_{C,a'} \left( \mathrm{pa}_{j}(\mathcal{G}_{M}) \right) - \pi_{C,a'} \left( \mathrm{pa}_{j}(\mathcal{G}_{M}) \right) \right] \mathrm{d} \, \mathrm{pa}_{j}(\mathcal{G}_{M}) \right] = o(n^{-1/2}), \\
& \mathbf{E} \left[ (\mathbb{P}_{n} - P) \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\mu}(C, a, \mathrm{pa}_{j}, M_{j}) - \mu(C, a, \mathrm{pa}_{j}, M_{j}) \right] \mathrm{d} \, \mathrm{pa}_{j} \right] = o(n^{-1/2}), \\
& \mathbf{E} \left[ (\mathbb{P}_{n} - P) \int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \left[ \widehat{\pi}_{C,a}(\mathrm{pa}_{j}) - \pi_{C,a}(\mathrm{pa}_{j}) \right] \mathrm{d} \, \mathrm{pa}_{j} \right] = o(n^{-1/2}).
\end{aligned}$$

Therefore, we conclude

(C.9)
$$\frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} (\mathbb{P}_{n} - P) \Big[ \{ \varphi_{j,1}^{\operatorname{QR}}(X, \widehat{P}; \mathcal{G}_{M}) - \varphi_{j,1}^{\operatorname{QR}}(X, P; \mathcal{G}_{M}) \} \Big] = o_{p} (n^{-1/2}),$$

$$\frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} (\mathbb{P}_{n} - P) \{ \varphi_{j,0}^{\operatorname{QR}}(X, \widehat{P}; \mathcal{G}_{M}) - \varphi_{j,0}^{\operatorname{QR}}(X, P; \mathcal{G}_{M}) \} = o_{p} (n^{-1/2}).$$

It remains to deal with the reminder  $R_{j,a'}^{QR}(\widehat{P},P)$  for  $a' \in \{0,1\}$ . Note that for any distribution Q, we rewrite

$$\begin{split} R_{j,a'}^{\mathsf{RR}}(Q,P;\mathcal{G}_{M}) &= \psi_{j,a'}(P;\mathcal{G}_{M}) + \int \varphi_{j,a'}^{\mathsf{RR}}(x,Q;\mathcal{G}_{M}) \, \mathrm{d}P(x) \\ &= -\psi_{j,a'}(P;\mathcal{G}_{M}) + \int \left\{ \frac{1(a=a')}{e_{a'}^{Q}(c)} \left\{ y - \kappa^{Q}(a',c) \right\} + \kappa^{Q}(a',c) \right. \\ &- \left[ \frac{1(a=a')}{e_{a'}^{Q}(c)} \pi_{C,a'}^{Q}(\operatorname{Pa}_{j}(\mathcal{G}_{M})) \left\{ y - \mu^{Q}(c,a',m) \right\} \right. \\ &+ \frac{1(a=a')}{e_{a'}^{Q}(c)} \left( \int_{\mathcal{M}_{j}} \mu^{Q}(c,a',\operatorname{pa}_{j}(\mathcal{G}_{M}),m_{j}) \pi_{c}^{Q}(m_{j}) \, \mathrm{d}m_{j} - \operatorname{E}_{P} \left[ \varrho_{j}^{Q}(a',M_{j},c;\mathcal{G}_{M}) \mid c \right] \right) \\ &+ \varrho_{j}^{Q}(a',m_{j},c;\mathcal{G}_{M}) \right] \right\} \, \mathrm{d}P(x) \\ &= - \int \left[ \kappa(a',c) - \varrho_{j}(a',m_{j},c;\mathcal{G}_{M}) \right] \, \mathrm{d}P(x) + \int \left\{ \frac{1(a=a')}{e_{a'}^{Q}(c)} \left\{ y - \kappa^{Q}(a',c) \right\} + \kappa^{Q}(a',c) \right\} + \kappa^{Q}(a',c) \right. \\ &- \left[ \frac{1(a=a')}{e_{a'}^{Q}(c)} \frac{\pi_{c}^{Q}(m_{j})}{\pi_{c,a',\operatorname{pa}_{j}}^{Q}(\mathcal{G}_{M})(m_{j})} \left\{ y - \mu^{Q}(c,a',m) \right\} \right. \\ &+ \frac{1(a=a')}{e_{a'}^{Q}(c)} \left( \int_{\mathcal{M}_{j}} \mu^{Q}(c,a',\operatorname{pa}_{j}(\mathcal{G}_{M}),m_{j}) \pi_{c}^{Q}(m_{j}) \, \mathrm{d}m_{j} - \operatorname{E}_{P} \left[ \varrho_{j}^{Q}(a',M_{j},c;\mathcal{G}_{M}) \mid c \right] \right) \\ &+ \varrho_{j}^{Q}(a',m_{j},c;\mathcal{G}_{M}) \right] \right\} \, \mathrm{d}P(x) \\ \text{by rearranging} \int \left\{ \frac{e_{a'}(c)}{e_{a'}^{Q}(c)} \left\{ \kappa(a',c) - \kappa^{Q}(a',c) \right\} + \left\{ \kappa^{Q}(a',c) - \kappa(a',c) \right\} \right. \\ &- \left[ \frac{e_{a'}(c)}{e_{a'}^{Q}(c)} \frac{\pi_{c}^{Q}(m_{j})}{\pi_{c,a',\operatorname{pa}_{j}}^{Q}(g_{M})(m_{j})} \left\{ \mu(c,a',m) - \mu^{Q}(c,a',m) \right\} \right. \\ &+ \frac{e_{a'}(c)}{e_{a'}^{Q}(c)} \left( \int_{\mathcal{M}_{j}} \mu^{Q}(c,a',\operatorname{pa}_{j}(\mathcal{G}_{M}),m_{j}) \pi_{c}^{Q}(m_{j}) \, \mathrm{d}m_{j} - \operatorname{E}_{P} \left[ \varrho_{j}^{Q}(a',M_{j},c;\mathcal{G}_{M}) \mid c \right] \right) \\ &+ \varrho_{j}^{Q}(a',m_{j},c;\mathcal{G}_{M}) - \varrho_{j}(a',\operatorname{pa}_{j}(\mathcal{G}_{M}),m_{j}) \pi_{c}^{Q}(m_{j}) \, \mathrm{d}m_{j} - \operatorname{E}_{P} \left[ \varrho_{j}^{Q}(a',M_{j},c;\mathcal{G}_{M}) \mid c \right] \right) \\ &+ \varrho_{j}^{Q}(a',m_{j},c;\mathcal{G}_{M}) - \varrho_{j}(a',m_{j},c;\mathcal{G}_{M}) \right] \right\} \, \mathrm{d}P(x) \\ =: R_{j,n'}^{(1)}(Q,P) + R_{j,n'}^{(2)}(Q,P;\mathcal{G}_{M}) - R_{j,n'}^{(3)}(Q,P;\mathcal{G}_{M}) + R_{j,n'}^{(4)}(Q,P;\mathcal{G}_{M}). \end{split}$$

In the equation, we use the identities

$$\begin{split} &\int \mu(c,a',m) \Big\{ \pi_{c,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{c,a'}^Q \big( \operatorname{pa}_j(\mathcal{G}_M) \big) \Big\} \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &+ \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \left( \int_{\mathcal{M}_j} \mu^Q(c,a',\operatorname{pa}_j(\mathcal{G}_M),m_j) \pi_c^Q(m_j) \operatorname{d} m_j - \operatorname{E}_P \big[ \varrho_j^Q(a',M_j,c;\mathcal{G}_M) \mid c \big] \right) \operatorname{d} P(x) \\ &+ \operatorname{expanding the expectation} \int \operatorname{d} P(x) \int \mu(c,a',m) \Big\{ \pi_{c,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{c,a'}^Q \big( \operatorname{pa}_j(\mathcal{G}_M) \big) \Big\} \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &- \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \operatorname{d} P(x) \int_{\mathcal{M}_j} \mu^Q(c,a',m) \pi_c^Q(m_j) \operatorname{d} m_j \\ &- \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big\{ \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q \big( \operatorname{pa}_j(\mathcal{G}_M) \big) \Big\} \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &- \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \operatorname{d} P(x) \int_{\mathcal{M}_j} \mu(c,a',m) \pi_c^Q(m_j) \operatorname{d} m_j \\ &- \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \operatorname{d} P(x) \int_{\mathcal{M}_j} \mu(c,a',m) \pi_c^Q(m_j) \operatorname{d} m_j \\ &- \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big\{ \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q \big( \operatorname{pa}_j(\mathcal{G}_M) \big) \Big\} \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &+ \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \operatorname{d} P(x) \int_{\mathcal{M}_j} \Big\{ \mu(c,a',m) - \mu^Q(c,a',m) \Big\} \pi_c^Q(m_j) \operatorname{d} m_j \\ &- \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big\{ \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q \big( \operatorname{pa}_j(\mathcal{G}_M) \big) \Big\} \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &+ \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \pi_c^Q(m_j) \operatorname{d} P(x) \int_{\mathcal{M}_j} \mu(c,a',m) \pi_c^Q(m_j) \operatorname{d} m_j \\ &- \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big\{ \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q \big( \operatorname{pa}_j(\mathcal{G}_M) \big) \Big\} \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &+ \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \pi_c^Q(m_j) \operatorname{d} P(x) \\ &- \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big\{ \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q \big( \operatorname{pa}_j(\mathcal{G}_M) \big) \Big\} \Big( \mu(c,a',m) - \mu^Q(c,a',m) \Big) \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &+ \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \pi_c^Q(m_j) \operatorname{d} P(x) \\ &- \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big( \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q(\operatorname{pa}_j(\mathcal{G}_M) \big) \Big) \Big( \mu(c,a',m) - \mu^Q(c,a',m) \Big) \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &+ \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big( \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q(\operatorname{pa}_j(\mathcal{G}_M) \big) \Big) \Big( \mu(c,a',m) - \mu^Q(c,a',m) \Big) \operatorname{d} \operatorname{pa}_j(\mathcal{G}_M) \\ &+ \int_{\mathcal{M}_{\operatorname{pa}_j}(\mathcal{G}_M)} \Big( \pi_{C,a'} \big( \operatorname{pa}_j(\mathcal{G}_M) \big) - \pi_{C,a'}^Q(\operatorname{pa}_j(\mathcal{G}_M) \big) \Big) \Big( \mu(c,a',m)$$

and then the components  $R_{j,a'}^{(\ell)}(\cdot)$  with  $\ell=1,2,3,4$  can be rewritten as following

$$\begin{split} R_{j,a'}^{(1)}(Q,P) &= \int \left[ \frac{e_{a'}(c)}{e_{a'}^Q(c)} \Big\{ \kappa(a',c) - \kappa^Q(a',c) \Big\} + \Big\{ \kappa^Q(a',c) - \kappa(a',c) \Big\} \right] \mathrm{d}P(x) \\ &= \int \left( \frac{1}{e_{a'}^Q(c)} - \frac{1}{e_{a'}(c)} \right) \Big( \kappa(a',c) - \kappa^Q(a',c) \Big) e_{a'}(c) \, \mathrm{d}P(x), \end{split}$$

$$\begin{split} &R_{j,a'}^{(2)}(Q,P;\mathcal{G}_M)\\ &=\int \mathrm{d}P(x) \bigg[\frac{e_{a'}(c)}{e_{a'}^Q(c)}-1\bigg] \Big\{\mu(c,a',m)-\mu^Q(c,a',m)\Big\} \pi_{c,a'}^Q\big(\operatorname{pa}_j(\mathcal{G}_M)\big) \operatorname{d}\operatorname{pa}_j(\mathcal{G}_M)\\ &=\int \bigg(\frac{1}{e_{a'}^Q(c)}-\frac{1}{e_{a'}(c)}\bigg) \Big(\mu(c,a',m)-\mu^Q(c,a',m)\Big) e_{a'}(c) \pi_{c,a'}^Q\big(\operatorname{pa}_j(\mathcal{G}_M)\big) \operatorname{d}\operatorname{pa}_j(\mathcal{G}_M) \operatorname{d}P(x), \end{split}$$

and

$$\begin{split} R_{j,a'}^{(3)}(Q,P;\mathcal{G}_{M}) &= \int e_{a'}^{Q}(c) \, \mathrm{d}P(x) \int_{\mathcal{M}_{j}} \mu(c,a',m) \pi_{c}^{Q}(m_{j}) \, \mathrm{d}m_{j} \\ &\int_{\mathcal{M}_{\mathrm{pa}_{j}}(\mathcal{G}_{M})} \left( \frac{1}{e_{a'}^{Q}(c)} - \frac{1}{e_{a'}(c)} \right) \left( \pi_{C,a'} \left( \mathrm{pa}_{j}(\mathcal{G}_{M}) \right) - \pi_{C,a'}^{Q} \left( \mathrm{pa}_{j}(\mathcal{G}_{M}) \right) \right) \mathrm{d} \, \mathrm{pa}_{j}(\mathcal{G}_{M}), \end{split}$$

$$R_{j,a'}^{(4)}(Q, P; \mathcal{G}_M) = \int \frac{e_{a'}(c)}{e_{a'}^Q(c)} \pi_c^Q(m_j) \, \mathrm{d}P(x)$$

$$\int_{\mathcal{M}_{\mathrm{pa}_j}(\mathcal{G}_M)} \left( \pi_{C,a'} \left( \mathrm{pa}_j(\mathcal{G}_M) \right) - \pi_{C,a'}^Q \left( \mathrm{pa}_j(\mathcal{G}_M) \right) \right) \left( \mu(c,a',m) - \mu^Q(c,a',m) \right) \, \mathrm{d}\, \mathrm{pa}_j(\mathcal{G}_M).$$

Therefore, by Assumption 7.10, we have for sufficient large n such that

$$\left| \widehat{f}(x_{S_2} \mid x_{S_1}) - f(x_{S_2} \mid x_{S_1}) \right| \le f(x_{S_2} \mid x_{S_1}) \quad \text{and} \quad \left| \widehat{\mathbb{E}} \left[ x_{S_2} \mid x_{S_1} \right] - \mathbb{E} \left[ x_{S_2} \mid x_{S_1} \right] \right| \le \mathbb{E} \left[ x_{S_2} \mid x_{S_1} \right]$$

almost surely. Then components  $R_{j,a'}^{(\ell)}(\cdot)$  with  $\ell=1,2,3,4$  can be furthermore bounded by

$$\begin{split} \left| R_{j,a'}^{(1)}(\widehat{P},P) \right| &= \left| \int \left( \frac{1}{\widehat{e}_{a'}(c)} - \frac{1}{e_{a'}(c)} \right) \left( \kappa(a',c) - \widehat{\kappa}(a',c) \right) e_{a'}(c) \, \mathrm{d}P(x) \right| \\ & \qquad \qquad \leq \frac{1}{\varepsilon} \int \left| e_{a'}(c) - \widehat{e}_{a'}(c) \right| \left| \kappa(a',c) - \widehat{\kappa}(a',c) \right| \, \mathrm{d}P(x) \\ & \qquad \qquad \otimes n^{-\vartheta_{j,e}^* + \vartheta_{j,\mu}^*} = o_p(n^{-1/2}), \end{split}$$

and similarly

$$\begin{split} \left| R_{j,a'}^{(2)}(\widehat{P},P\,;\,\mathcal{G}_M) \right| \\ &= \left| \int \left( \frac{1}{\widehat{e}_{a'}(c)} - \frac{1}{e_{a'}(c)} \right) \left( \mu(c,a',m) - \widehat{\mu}(c,a',m) \right) e_{a'}(c) \pi_{c,a'}^Q \left( \operatorname{pa}_j(\mathcal{G}_M) \right) \operatorname{dpa}_j(\mathcal{G}_M) \operatorname{d}P(x) \right| \\ & \lesssim \int \left| e_{a'}(c) - \widehat{e}_{a'}(c) \right| \left| \mu(c,a',m) - \widehat{\mu}(c,a',m) \right| \operatorname{d}P(x) \end{split}$$

by Assumption 7.10 
$$\stackrel{\text{7.10}}{\asymp} n^{-\vartheta_{j,e}^* + \vartheta_{j,\mu}^*} = o_p(n^{-1/2}),$$

$$\left|R_{j,a'}^{(3)}(\widehat{P},P;\mathcal{G}_M)\right|$$

by 
$$\ell^2$$
 assumption and Assumption 4.2 
$$\lesssim \int \left| e_{a'}(c) - \widehat{e}_{a'}(c) \right| \left| \pi_{C,a'} \left( \operatorname{pa}_j(\mathcal{G}_M) \right) - \widehat{\pi}_{C,a'} \left( \operatorname{pa}_j(\mathcal{G}_M) \right) \right| dP(x)$$
by Assumption 7.10 
$$n^{-\vartheta_{j,e}^* + \vartheta_{j,\pi}^*} = o_n(n^{-1/2}),$$

and

$$\left|R_{a'}^{(4)}(\widehat{P},P;\mathcal{G}_M)\right|$$

by 
$$\ell^2$$
 assumption and Assumption 4.2 
$$\lesssim \int \left| \pi_{C,a'} \left( \operatorname{pa}_j(\mathcal{G}_M) \right) - \widehat{\pi}_{C,a'} \left( \operatorname{pa}_j(\mathcal{G}_M) \right) \right| \left| \mu(c,a',m) - \widehat{\mu}(c,a',m) \right| dP(x)$$
by Assumption 7.10 
$$\approx n^{-\vartheta_{j,\pi}^* + \vartheta_{j,\mu}^*} = o_n(n^{-1/2}).$$

Therefore, for the reminder, we have

(C.10) 
$$\frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} R_{j,1}^{\operatorname{QR}}(\widehat{P}, P; \mathcal{G}_{M})$$

$$\approx \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} R_{j,0}^{\operatorname{QR}}(\widehat{P}, P; \mathcal{G}_{M})$$

$$= o_{p}(n^{-1/2})$$

for our quadruply robust estimators. Plug (C.9) and (C.10) into (C.7), we obtain

$$\begin{split} &\sqrt{n} \big( \widehat{TM}_{j}^{\text{avg, QR}} - \overline{TM}_{j} \big) \\ &= \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} \left[ \mathbb{G}_{n} \big\{ \varphi_{j,1}^{\operatorname{QR}}(X, P \, ; \, \mathcal{G}_{M}) - \varphi_{j,0}^{\operatorname{QR}}(X, P \, ; \, \mathcal{G}_{M}) \big\} + o_{p}(1) \right] \\ & \overset{\text{by } o_{p}(1) \text{ is uniform on } \operatorname{MEC}(\mathcal{C}_{M})}{=} \mathbb{G}_{n} \left\{ \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{M})} \big\{ \varphi_{j,1}^{\operatorname{QR}}(X, P \, ; \, \mathcal{G}_{M}) - \varphi_{j,0}^{\operatorname{QR}}(X, P \, ; \, \mathcal{G}_{M}) \big\} \right\} \\ & + o_{p}(1) \\ & \overset{}{\sim} \mathcal{N} \left( 0, \operatorname{E} \left[ \frac{1}{\# \operatorname{MEC}(\mathcal{C}_{0,M})} \sum_{\mathcal{G}_{M} \in \operatorname{MEC}(\mathcal{C}_{0,M})} S^{\operatorname{eff, nonpar}} \big( TM_{j}(\mathcal{G}_{M}) \big) \right]^{2} \right) \end{split}$$

which completes the proof of the semiparametric efficiency for  $\widehat{TM}_j^{\rm avg,\,QR}.$   $\hfill\Box$