

LS-CATEGORY AND TOPOLOGICAL COMPLEXITY OF REAL TORUS MANIFOLDS AND DOLD MANIFOLDS OF REAL TORUS TYPE

KOUSHIK BRAHMA, NAVNATH DAUNDKAR, AND SOUMEN SARKAR

ABSTRACT. The real torus manifolds are a generalization of small covers and generalized real Bott manifolds. We compute the LS-category of these manifolds under some constraints and obtain sharp bounds on their topological complexities. We obtain a simplified description of their cohomology ring and discuss a relation on the cup-product of its generators. We obtain the sharp bounds on their zero-divisors-cup-lengths. We improve the dimensional upper bound on their topological complexity. We also show that under certain hypotheses, the topological complexity of real torus manifolds of dimension n is either $2n$ or $2n + 1$. We compute the \mathbb{Z}_2 -equivariant LS-category of any small cover when the \mathbb{Z}_2 -fixed points are path connected. We then compute the LS-category of Dold manifolds of real torus type and obtain sharp bounds on their topological complexity. In the end, we obtain sharp bounds on the symmetric topological complexity of a class of these manifolds.

1. INTRODUCTION

Farber [16] introduced the concept of topological complexity to study the robot motion planning problem through the topological lens. For a topological space X , the topological complexity $\text{TC}(X)$ is a numerical homotopy invariant. Let X be a path-connected space and PX be the space of all paths in X equipped with a compact open topology. Let $\gamma: [0, 1] \rightarrow X$ be any path in X . Then there is a fibration $\pi: PX \rightarrow X \times X$ defined by $\pi(\gamma) = (\gamma(0), \gamma(1))$. The *topological complexity* of X is the smallest k for which $X \times X$ admit an open cover U_1, \dots, U_k , such that there exist continuous sections of π on U_i for $1 \leq i \leq k$. In general, determining the exact value of $\text{TC}(X)$ is a hard problem even for the spaces having nice CW-complex structures like Dold manifolds. Since Farber first introduced this invariant in 2003, numerous methods have been created to determine bounds on $\text{TC}(X)$, and the precise value of this invariant has been calculated for several spaces, see for example [3], [13], [14], [16], [18], [20], [22], [23]. This invariant is closely related to the well-known invariant, the *Lusternik-Schnirelmann category* (in short, *LS-category*) [26]. For a path connected space X , the LS-category of X is denoted $\text{cat}(X)$. The $\text{cat}(X)$ is the smallest integer r such that X can be covered by r open subsets V_1, \dots, V_r with inclusion $V_i \hookrightarrow X$ is null-homotopic for $1 \leq i \leq r$. In particular, it was shown in [7] that,

$$\text{cat}(X) \leq \text{TC}(X) \leq 2\text{cat}(X) - 1.$$

The cohomological methods have been used to obtain some lower bounds on $\text{TC}(X)$ and $\text{cat}(X)$. We describe them here briefly. Let R be a commutative ring with unity and X be a path-connected topological space with its cohomology ring $H^*(X; R)$. The cup-length of X over R is the maximal integer ℓ such that there exists $x_i \in H^*(X; R)$ for $i = 1, \dots, \ell$ satisfying $\prod_{i=1}^{\ell} x_i \neq 0$. We denote this number by $\text{cl}_R(X)$. It was proved in [7, Proposition

2020 *Mathematics Subject Classification.* 55M30, 57S12.

Key words and phrases. real torus manifold, generalized Bott manifold, small cover, LS-Category, Topological Complexity, Dold manifold of real torus type.

1.5] that the number gives a lower bound for the $\text{cat}(X)$. Let

$$\cup: H^*(X; R) \otimes H^*(X; R) \longrightarrow H^*(X; R)$$

be the map induced by the cup product. Then the zero-divisors-cup-length of X with respect to the coefficient ring R is defined as the maximal integer k such that there exist cohomology classes $u_i \in H^*(X; R) \otimes H^*(X; R)$ satisfying $\cup(u_i) = 0$ for all $1 \leq i \leq k$ and $\prod_{i=1}^k u_i \neq 0$. This integer is denoted by $\text{zcl}_R(X)$. It was proved in [16, Theorem 7] that $\text{zcl}_R(X)$ gives a lower bound for the $\text{TC}(X)$. That is, we have the following;

$$\text{cat}(X) \geq \text{cl}_R(X) + 1 \quad \text{and} \quad \text{TC}(X) \geq \text{zcl}_R(X) + 1. \quad (1)$$

A well-studied manifolds are generalized Bott manifolds. However, little is known about their real counterpart. Briefly, A *generalized real Bott tower* of height m is a sequence

$$B_m \xrightarrow{\pi_m} B_{m-1} \xrightarrow{\pi_{m-1}} \dots \xrightarrow{\pi_2} B_1 \xrightarrow{\pi_1} B_0 = \{\text{pt}\} \quad (2)$$

of manifolds $B_j = \mathbb{P}(\underline{\mathbb{R}} \oplus E_j^{(1)} \oplus \dots \oplus E_j^{(n_j)})$, where $\underline{\mathbb{R}}$ is the trivial line bundle over B_{j-1} , $E_j^{(i)}$ is a real line bundle over B_{j-1} for $i = 1, \dots, n_j$, and $j = 1, \dots, m$. Here $\mathbb{P}(\cdot)$ denotes the projectivization. The space B_j is called a j -th stage generalized real Bott manifold. Note that the fibre of the map π_j is $\mathbb{R}P^{n_j}$ for $j = 1, \dots, m$. In particular, $B_1 = \mathbb{R}P^{n_1}$. Also, when $n_j = 1$ for every $j \in \{1, \dots, m\}$, then B_j is called an m th-stage real Bott manifold. Interestingly, generalized real Bott manifolds are small covers. We note that the small covers are central objects in Toric Topology introduced by Davis and Januszkiewicz [11]. They are smooth manifolds with effective locally standard real torus actions and have rich combinatorial properties. The concept of small covers has been extended in [25] where the author called them 2-torus manifolds. If the real torus action on a 2-torus manifold is locally standard and the orbit space is a nice manifold with corners, we may call the 2-torus manifold a ‘real torus manifold’. Following [25], one can get a combinatorial pair (P, λ) from a real torus manifold where P is the orbit space which is a nice manifold with corners and λ is called a \mathbb{Z}_2 -characteristic function on P .

The other class of manifolds we are interested in are called *generalized projective product spaces*. These manifolds were recently introduced by S. Sarkar and P. Zvengrowski in [29], extending the concept of (generalized) Dold manifolds [12, 27] and projective product spaces [9]. Let N be a manifold equipped with a free \mathbb{Z}_2 -action and M a \mathbb{Z}_2 -manifold. Then the diagonal \mathbb{Z}_2 -action on the product $M \times N$ is free. Then the orbit space $D(M, N) := (M \times N)/\mathbb{Z}_2$ is a manifold. In particular, if M is a real torus manifold, we call $D(M, N)$ a *Dold manifold of real torus type*. Several properties of these manifolds remain to explore for many M and N . The goal of this article is to compute the LS-category and topological complexity of real torus manifolds (which contain generalized real Bott manifolds and small covers) and Dold manifolds of real torus type. We explain the details of this in the organization.

In Section 2, we recall a few basic topological properties of real torus manifolds. we prove that the LS-category of a real torus manifold M is $\dim(M) + 1$ if the orbit space contains a boundary of a simple polytope, see Theorem 2.5. If M is an n -dimensional real torus manifold such that a facet of the orbit space is an $(n - 1)$ -simplex, then the lower bound on the topological complexity of M can be given by the zero-divisors-cup-length of $\mathbb{R}P^{n-1}$, see Theorem 2.7. In addition, if the \mathbb{Z}_2 -characteristic function λ associated with M satisfies certain conditions, then the lower bound on the topological complexity of M can be given by the zero-divisors-cup-length of $\mathbb{R}P^n$, see Theorem 2.7. Then, we show that a certain product of a generating set of $H^*(M(P, \lambda); \mathbb{Z}_2)$ is non-zero if P is a finite product of simplices, see Lemma 2.12. As a result, we use cohomological methods to obtain

a sharp lower bound on the topological complexity of generalized real Bott manifolds, see Theorem 2.13 and Corollary 2.14. Then we prove that if the product of simplices contains k -many odd dimensional simplices, then the dimensional upper bound on the topological complexity of generalized real Bott manifolds can be improved by k , see Theorem 2.15.

In Section 3, we compute the \mathbb{Z}_2 -equivariant LS-category of small covers when the fixed point sets are path connected, see Theorem 3.2. For the other kinds of \mathbb{Z}_2 -actions, when fixed point sets are not connected, we show that the \mathbb{Z}_2 -equivariant category of small covers could be much greater than the dimension plus one, see Theorem 3.3. In general, it remains lesser than the number of vertices of the corresponding polytope. Example 3.4 observes that, in certain cases, it can be exactly the number of vertices of the polytope. Then we compute the LS-category of Dold manifolds of real torus type and obtain sharp bounds on their topological complexity, see Theorem 3.7 and Theorem 3.8. In many specific cases, the topological complexity of these manifolds is either $a - 1$, a , or $a + 1$ for some $a \in \mathbb{N}$, see Corollary 3.9.

In Section 4, first we recall the definition and a few properties of symmetric topological complexities. Then, we obtain sharp bounds on the symmetric topological complexity of a class of real torus manifolds and Dold manifolds of real torus type (see Theorem 4.4, Theorem 4.5 and Theorem 4.6). Moreover, in Theorem 4.5, we compute the exact value of symmetric topological complexity of many real torus manifolds.

2. LS-CATEGORY AND TOPOLOGICAL COMPLEXITY OF REAL TORUS MANIFOLDS

In this section, we recall the concepts of real torus manifolds and generalized real Bott manifolds. We give a nice presentation of the cohomology ring of generalized real Bott manifolds with \mathbb{Z}_2 -coefficients. Then, we compute some lower and upper bounds for the topological complexity of these manifolds. We begin by stating the concept of real torus manifolds.

We recall the definition of a nice manifold with corners following [10] and [2]. The codimension function c associates a point $x = (x_1, \dots, x_n) \in (\mathbb{R}_{\geq 0})^n$ to the number of x_i which are zero. An n -dimensional manifold with corners is a Hausdorff, second-countable topological space together with a maximal atlas of local charts onto open subsets of $(\mathbb{R}_{\geq 0})^n$ such that the overlapping maps are homeomorphisms that preserve codimension function. That means the codimension function is a well-defined map from a manifold with corners P to non-negative integers. A connected component of $c^{-1}(m)$ is called an open face of P . The closure of an open face is called a face. We can define the dimension of a face by subtracting its codimension from n . In an n -dimensional nice manifold with corners, a face of dimension zero and $n - 1$ are called a vertex and a facet. The vertex set and the facet set of P are denoted by $V(P)$ and $\mathcal{F}(P)$, respectively.

A manifold with corners P is called nice if, for every $p \in P$ with $c(p) = 2$, the number of facets of P which contains p is also 2. Therefore, a codimension- k face of a nice manifold with corners P is a connected component of the intersection of the unique collection of k many facets of P .

An easy example of a manifold with corners is a convex polytope (a convex hull of finitely many points in \mathbb{R}^n for some $n \in \mathbb{Z}_{\geq 0}$). If a convex polytope is nice as in the above sense, then it is called a simple polytope i.e., at each vertex exactly n many facets intersect.

Consider \mathbb{R}^n with standard \mathbb{Z}_2 -action induced by the reflections on coordinate hyperplanes. A smooth action of \mathbb{Z}_2^n on an n -dimensional smooth manifold M is called *locally standard*, if $x \in M$ then it has a \mathbb{Z}_2^n -invariant open neighbourhood U_x and a diffeomorphism

$\phi: U_x \rightarrow V$, where V is \mathbb{Z}_2^n -invariant open set in \mathbb{R}^n , and an isomorphism $\delta_x: \mathbb{Z}_2^n \rightarrow \mathbb{Z}_2^n$ such that $\phi(t \cdot x) = \delta_x(t) \cdot \phi(x)$ for all $t \in \mathbb{Z}_2^n$ and $x \in U_x$.

Definition 2.1. A closed, connected, and smooth n -dimensional manifold M with locally standard action of \mathbb{Z}_2^n is called a real torus manifold over a nice manifold with corners P of dimension n , if the following conditions are satisfied:

- (1) The boundary ∂P is non-empty.
- (2) There is a projection map $\mathfrak{q}: M \rightarrow P$ constant on orbits which maps every ℓ -dimensional orbit to a point in the interior of an ℓ -dimensional face of P .

If P is a simple polytope, then the real torus manifold M is called a small cover. Let P be a nice manifold with corners, and $\mathcal{F}(P) := \{F_1, \dots, F_r\}$.

Definition 2.2. A function $\lambda: \mathcal{F}(P) \rightarrow \mathbb{Z}_2^n$ is called a characteristic function if the submodule of \mathbb{Z}_2^n generated by $\{\lambda(F_{i_1}), \dots, \lambda(F_{i_\ell})\}$ is an ℓ -dimensional direct summand of \mathbb{Z}_2^n whenever $F_{i_1} \cap \dots \cap F_{i_\ell} \neq \emptyset$. The vector $\lambda_i := \lambda(F_i)$ is called the characteristic vector associated with the facet $F_i \in \mathcal{F}(P)$, and the pair (P, λ) is called a characteristic pair.

We discuss the construction of a real torus manifold from a characteristic pair (P, λ) . For each point $p \in P$, let $F(p)$ be the unique face of P , which contains p in its relative interior. Then $F(p)$ is a connected component of $F_{i_1} \cap \dots \cap F_{i_k}$ for some unique facets F_{i_1}, \dots, F_{i_k} of P . Let $G_{F(p)}$ be the subgroup of \mathbb{Z}_2^n generated by $\{\lambda(F_{i_1}), \dots, \lambda(F_{i_k})\}$. We define an equivalence relation on $P \times \mathbb{Z}_2^n$ as follows:

$$(p, g) \sim (q, h) \Leftrightarrow p = q, g^{-1}h \in G_{F(p)}. \quad (3)$$

The identification space

$$M^n(P, \lambda) := (P \times \mathbb{Z}_2^n) / \sim$$

has an n -dimensional manifold structure with a natural locally standard \mathbb{Z}_2^n -action induced by the group operation on the second factor of $P \times \mathbb{Z}_2^n$. Thus $M^n(P, \lambda)$ is a real torus manifold over P (see [11] for details). The projection onto the first factor gives the orbit map

$$\mathfrak{q}: M^n(P, \lambda) \rightarrow P \text{ defined by } [p, g]_{\sim} \mapsto p,$$

where $[p, g]_{\sim}$ is the equivalence class of (p, g) .

Example 2.3. Let $Q = \{(x_1, \dots, x_n, x) \in \mathbb{R}^{n+1} \mid \sum_1^n x_i^2 + x^2 = 1, x_i \geq 0 \text{ for } i = 1, \dots, n\}$ for $n \geq 2$. Then Q is a nice manifold with corners homeomorphic to a suspension on $(n-1)$ -simplex. Note that Q is not a simple polytope. The facet set of Q is $\mathcal{F}(Q) = \{F_1, \dots, F_n\}$, where

$$F_i := \{(x_1, \dots, x_n, x) \in Q \mid x_i = 0 \text{ for } i = 1, \dots, n\}.$$

Define $\lambda: \mathcal{F}(Q) \rightarrow \mathbb{Z}_2^n$ by $\lambda(F_i) = (1, \dots, 1, -1, 1, \dots, 1)$, where -1 is at the i -th component. Then $M^n(Q, \lambda)$ is a real torus manifold homeomorphic to S^n , which is not a small cover.

Let M be a real torus manifold over a nice manifold with corners P of dimension n , and $\mathfrak{q}: M \rightarrow P$ be the orbit map. Let $\mathcal{F}(P) = \{F_1, \dots, F_r\}$. Then the isotropy subgroup of $\mathfrak{q}^{-1}(F_i)$ is a subgroup of \mathbb{Z}_2^n and isomorphic to \mathbb{Z}_2 . This isotropy subgroup is uniquely determined by $\lambda_i \in \mathbb{Z}_2^n \setminus \{(1, \dots, 1)\}$ for $i = 1, \dots, m$. Thus, we have an assignment $\lambda: \mathcal{F}(P) \rightarrow \mathbb{Z}_2^n$ satisfying Definition 2.2. Therefore, by [25, Lemma 3.1], we have the following.

Proposition 2.4. There is an equivariant homomorphism from $M^n(P, \lambda)$ onto M .

Now, we will compute the LS-category of real torus manifolds over the manifolds with corners and sharp bounds on their topological complexity.

Theorem 2.5. *Let M be an n -dimensional real torus manifold over P . If a connected component of ∂P is a boundary of an n -dimensional simple polytope Q , then $\text{cat}(M) = n + 1$.*

Proof. Let v be a vertex in $\partial Q \subseteq \partial P$. So there exist unique facets F_1, \dots, F_n of P such that $v = F_1 \cap \dots \cap F_n$ and $F_{i_1} \cap \dots \cap F_{i_k}$ is an $(n - k)$ -dimensional simple polytope for any $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$. Let $q: M \rightarrow P$ be the orbit map and x_i the Poincaré dual of $q^{-1}(F_i)$. It follows from the given hypotheses on P , that the intersection $F_1 \cap \dots \cap F_n$ is transversal. Thus, the product $x_1 \cdots x_n$ of cohomology classes is nonzero in $H^*(M; \mathbb{Z}_2)$. Therefore, the result follows from the cup-length lower bound (1) and the dimensional upper bound as given in [7, Theorem 1.7]. \square

Theorem 2.6. *Let M be an n -dimensional real torus manifold over P with $n > 1$. Suppose Δ^{n-1} is a facet of P . Then,*

$$n \leq \text{cat}(M) \leq n + 1, \text{ and } n \leq \text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{n-1}) + 1 \leq \text{TC}(M) \leq 2n + 1. \quad (4)$$

In particular, if $n = 2^s + 1$, then

$$2^{s+1} \leq \text{TC}(M) \leq 2^{s+1} + 3.$$

Proof. The characteristic submanifold $M_{\Delta^{n-1}}$ corresponding to the facet Δ^{n-1} is equivariantly homeomorphic to $\mathbb{R}P^{n-1}$. Then the natural inclusion $\iota: M_{\Delta^{n-1}} \hookrightarrow M(P, \lambda)$ induces a surjective homomorphism

$$\iota^*: H^*(M; \mathbb{Z}_2) \rightarrow H^*(\mathbb{R}P^{n-1}; \mathbb{Z}_2).$$

Precisely, if F_i be the i -th facet of P and $x_i \in H^1(M; \mathbb{Z}_2)$ corresponds to F_i , then $\iota^*(x_i) \in H^1(\mathbb{R}P^{n-1}; \mathbb{Z}_2)$ corresponds to $F_i \cap \Delta^{n-1}$. Note that $F_i \cap \Delta^{n-1}$ is empty then $\iota^*(x_i) = 0$. Since ι^* is surjective, we have

$$\text{cl}_{\mathbb{Z}_2}(\mathbb{R}P^{n-1}) \leq \text{cl}_{\mathbb{Z}_2}(M), \text{ and } \text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{n-1}) \leq \text{zcl}_{\mathbb{Z}_2}(M).$$

Thus, the left inequality of (4) follows. The right inequality of (4) follows from [16, Theorem 4].

If $n = 2^s + 1$, then $\text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{n-1}) \geq 2^{s+1} - 1$, see [18]. So, the particular claim follows. \square

Theorem 2.7. *Let M be an n -dimensional real torus manifold over P with $n > 1$ and Δ^{n-1} a facet of P . Suppose F_1, \dots, F_n be the facets of P such that $\Delta^{n-1} \cap F_i \neq \emptyset$ and $\det[\lambda(F_1) \cdots \lambda(F_n)] = 1$. Then $\text{cat}(M) = n + 1$, and*

$$\text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^n) + 1 \leq \text{TC}(M) \leq 2n + 1. \quad (5)$$

In particular, if $n = 2^s$, then $2^{s+1} \leq \text{TC}(M) \leq 2^{s+1} + 1$.

Proof. We can have $M = M' \# \mathbb{R}P^n$ following the discussion in Subsection 1.11 in [11]. Then $H^*(\mathbb{R}P^n; \mathbb{Z}_2)$ is a subring of $H^*(M; \mathbb{Z}_2)$. Then $n + 1 = \text{cl}_{\mathbb{Z}_2}(\mathbb{R}P^n) \leq \text{cl}_{\mathbb{Z}_2}(M) \leq n + 1$, and the left inequality of (5) follows from the observation $\text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^n) \leq \text{zcl}_{\mathbb{Z}_2}(M^n(P, \lambda))$. The right inequality of (5) follows from [16, Theorem 4].

If $n = 2^s$, then $\text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^n) \geq 2^{s+1} - 1$, see [18]. So, the particular claim follows. \square

In the remaining, we focus on computing LS-category and topological complexity of generalized real Bott manifolds. We recall the results, which show that these manifolds are small covers over the product of simplices. Eminently, we obtain sharp bounds on their topological complexity.

Proposition 2.8. [24, Corollary 4.6] *The j -th stage generalized real Bott manifold B_j of the tower (2) is a small cover over $\prod_{i=1}^j \Delta^{n_i}$ where Δ^{n_i} is the n_i -simplex.*

The converse statement of this result also holds by the following proposition.

Proposition 2.9. [15, Proposition 2.7] *Every small cover over a product of simplices is a generalized real Bott manifold.*

Let $P = \prod_{j=1}^m \Delta^{n_j}$ be a product of m simplices. Let $\mathcal{F}(\Delta^{n_j}) := \{F_1^{\Delta_j}, \dots, F_{n_j+1}^{\Delta_j}\}$ be the facets of Δ^{n_j} for $j = 1, \dots, m$. Therefore, the facet set of P is

$$\mathcal{F}(P) := \{F_{r_j}^j \mid 1 \leq r_j \leq n_j + 1, j = 1, \dots, m\}, \quad (6)$$

where $F_{r_j}^j := \Delta^{n_1} \times \dots \times \Delta^{n_{j-1}} \times F_{r_j}^{\Delta_j} \times \Delta^{n_{j+1}} \times \dots \times \Delta^{n_m}$. Let $\mathcal{N}_j := \sum_{s=1}^j n_s$, for $j = 1, \dots, m$. Thus $\mathcal{N}_1 = n_1$ and $\mathcal{N}_m = n$. Let us define $\mathcal{N}_0 := 0$. Note that $|\mathcal{F}(P)| = n_1 + 1 + n_2 + 1 + \dots + n_m + 1 = n + m$. Define a \mathbb{Z}_2 -characteristic function on P by the following

$$\lambda: \mathcal{F}(P) \rightarrow \mathbb{Z}_2^n, \quad (7)$$

$$[\lambda(F_1^1), \dots, \lambda(F_{n_1+1}^1), \lambda(F_1^2), \dots, \lambda(F_{n_2+1}^2), \dots, \lambda(F_1^m), \dots, \lambda(F_{n_m+1}^m)]$$

is given by

$$[e_1, \dots, e_{\mathcal{N}_1}, \boldsymbol{\alpha}_1, e_{\mathcal{N}_1+1}, \dots, e_{\mathcal{N}_2}, \boldsymbol{\alpha}_2, \dots, e_{\mathcal{N}_{m-1}+1}, \dots, e_{\mathcal{N}_m}, \boldsymbol{\alpha}_m].$$

The function λ determines the following $m \times m$ vector matrix A .

$$A := \begin{pmatrix} \alpha_1^1 & \alpha_2^1 & \dots & \alpha_m^1 \\ \alpha_1^2 & \alpha_2^2 & \dots & \alpha_m^2 \\ \vdots & \vdots & \dots & \vdots \\ \alpha_1^m & \alpha_2^m & \dots & \alpha_m^m \end{pmatrix}_{m \times m} = (\boldsymbol{\alpha}_1 \quad \boldsymbol{\alpha}_2 \quad \dots \quad \boldsymbol{\alpha}_m)_{1 \times m}.$$

where $\boldsymbol{\alpha}_\ell^j = (\alpha_{\ell_1}^j, \alpha_{\ell_2}^j, \dots, \alpha_{\ell_{n_j}}^j)^t \in \mathbb{Z}_2^{n_j}$.

Next, we recall the cohomology ring of the generalized real Bott manifolds from [11]. We denote the facet $F_{r_j}^j$ by $F_{\mathcal{N}_{j-1}+r_j+j-1}$. Consider an indeterminate x_i corresponding to the facet F_i for each $i = 1, 2, \dots, n + m$.

Proposition 2.10 ([11]). *The cohomology ring of the generalized real Bott manifold is given by*

$$H^*(M^n(P, \lambda); \mathbb{Z}_2) \cong \frac{\mathbb{Z}_2[x_1, \dots, x_{n+m}]}{\tilde{I} + \tilde{J}}. \quad (8)$$

The ideal \tilde{I} is given by

$$\tilde{I} = \langle \{x_{\mathcal{N}_{j-1}+1} x_{\mathcal{N}_{j-1}+2} \dots x_{\mathcal{N}_j} x_{n+j} \mid j = 1, \dots, m\} \rangle, \quad (9)$$

and the ideal \tilde{J} is generated by the coordinates of an n -tuple

$$\Lambda_{\tilde{J}} = (\lambda(F_1)^t \quad \lambda(F_2)^t \quad \dots \quad \lambda(F_{n+m})^t)_{(n \times (n+m))} \cdot (x_1 \quad x_2 \quad \dots \quad x_{n+m})_{(n+m) \times 1}^t. \quad (10)$$

The i -th coordinate of $\Lambda_{\tilde{J}}$ is given by

$$x_i + \alpha_{1k_j}^j x_{n+1} + \alpha_{2k_j}^j x_{n+2} + \dots + \alpha_{mk_j}^j x_{n+m},$$

where $i = \mathcal{N}_{j-1} + k_j$; $k_j = 1, \dots, n_j$ and $j = 1, \dots, m$. Thus any x_i can be written as a \mathbb{Z}_2 -linear combination of x_{n+1}, \dots, x_{n+m} for $i = 1, \dots, n$. For simplicity, we denote the indeterminate x_{n+j} by y_j for $j = 1, \dots, m$. Thus,

$$x_i = \sum_{\ell=1}^m \alpha_{\ell k_j}^j y_\ell \text{ where } i = \mathcal{N}_{j-1} + k_j, k_j = 1, \dots, n_j \text{ and } j = 1, \dots, m, \quad (11)$$

in $H^*(M^n(P, \lambda); \mathbb{Z}_2)$. Therefore, using the generators of \tilde{I} we can express The cohomology ring of the generalized Bott tower $H^*(M^n(P, \lambda); \mathbb{Z}_2)$ as a quotient of the polynomial ring $\mathbb{Z}[y_1, y_2, \dots, y_n]$. Moreover, the generators of the ideal \tilde{I} in (9) can be described in terms of y_j 's.

On the other hand, using the arguments in [5, Proposition 5.1], the matrix A is conjugate to a unipotent lower triangular vector matrix of the following form:

$$\tilde{A} := \begin{pmatrix} \mathbf{1} & \mathbf{0} & \cdots & \mathbf{0} \\ \beta_1^2 & \mathbf{1} & \cdots & \mathbf{0} \\ \vdots & \vdots & \cdots & \vdots \\ \beta_1^m & \beta_2^m & \cdots & \mathbf{1} \end{pmatrix}_{m \times m}, \quad (12)$$

where $\beta_j^k = (\beta_{j_1}^k, \beta_{j_2}^k, \dots, \beta_{j_{n_k}}^k)^t \in \mathbb{Z}_2^{n_k}$ and $\mathbf{1} = (1, \dots, 1)^t \in \mathbb{Z}_2^{n_k}$ for $k = 1, \dots, m$. The matrix \tilde{A} is called the Bott matrix.

Thus, the ideal \tilde{J} is generated by the coordinates of the following matrix.

$$\begin{pmatrix} 1 & 0 & \cdots & 0 & 1 & 0 & \cdots & 0 & \cdots & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 1 & 0 & \cdots & 0 & \cdots & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 1 & 0 & \cdots & 0 & \cdots & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & \beta_{1n_1}^2 & 1 & \cdots & 1 & \cdots & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & \beta_{1n_2}^2 & 0 & \cdots & 1 & \cdots & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & \beta_{1n_m}^m & 0 & \cdots & \beta_{2n_m}^m & \cdots & \cdots & 0 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & \beta_{1n_m}^m & 0 & \cdots & \beta_{2n_m}^m & \cdots & \cdots & 1 & 1 \end{pmatrix}_{n \times (n+m)} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{N_1} \\ y_1 \\ x_{N_1+1} \\ \vdots \\ y_2 \\ \vdots \\ \vdots \\ \vdots \\ x_{N_m} \\ y_m \end{pmatrix}_{(n+m) \times 1}.$$

From the above matrix multiplication, the first n_1 elements are $x_1 + y_1 = 0, x_2 + y_1 = 0, \dots, x_{N_1} + y_1 = 0$. Therefore, we get $x_1 = x_2 = \cdots = x_{N_1} = y_1$. For $j \geq 2$,

$$x_{N_{j-1}+k_j} = \beta_{1k_j}^j y_1 + \beta_{2k_j}^j y_2 + \cdots + \beta_{(j-1)k_j}^j y_{j-1} + y_j$$

for $1 \leq k_j \leq n_j$.

Therefore, we have the following presentation of the cohomology ring with \mathbb{Z}_2 coefficients.

Proposition 2.11 ([4]). *The cohomology ring of the generalized Bott tower $M^n(P, \lambda)$ with \mathbb{Z}_2 coefficients is given by*

$$H^*(M^n(P, \lambda); \mathbb{Z}_2) \cong \mathbb{Z}_2[y_1, y_2, \dots, y_m]/I,$$

where

$$I = \langle \Gamma_1, \Gamma_2, \dots, \Gamma_m \rangle \text{ and } \Gamma_j = \prod_{k_j=1}^{n_j} \left(y_j + \sum_{\ell=1}^{j-1} \beta_{\ell k_j}^j y_\ell \right) y_j \text{ for } j = 1, 2, \dots, m.$$

Lemma 2.12. *Let $M^n(P, \lambda)$ be a small cover over the product of simplices $\prod_{j=1}^m \Delta^{n_j}$ with the characteristic function λ . Then, for $j \in \{1, \dots, m\}$, $y_j^{n_j} \neq 0$ in the cohomology ring $H^*(M^n(P, \lambda); \mathbb{Z}_2)$. Moreover, $y_1^{n_1} y_2^{n_2} \cdots y_m^{n_m} \neq 0$.*

Proof. Let $M^n(P, \lambda)$ be the small cover over the product of simplices $P = \prod_{j=1}^m \Delta^{n_j}$ and \tilde{A} be the Bott matrix as in (12) corresponding to the characteristic function λ on P . We recall the cohomology ring of a small cover over a product of simplices from Proposition 2.11.

Now we look at the generators Γ_j 's of the ideal I in the cohomology ring in Proposition 2.11. Note that,

$$\Gamma_1 = x_1 x_2 \cdots x_{n_1} y_1 = y_1^{n_1+1}, \text{ and}$$

$$\begin{aligned} \Gamma_j &= x_{\mathcal{N}_{j-1}+1} x_{\mathcal{N}_{j-1}+2} \cdots x_{\mathcal{N}_j} y_j \\ &= (y_j + \beta_{11}^j y_1 + \beta_{21}^j y_2 + \cdots + \beta_{(j-1)1}^j y_{j-1}) (y_j + \beta_{12}^j y_1 + \beta_{22}^j y_2 + \\ &\quad \cdots + \beta_{(j-1)2}^j y_{j-1}) \cdots (y_j + \beta_{1n_j}^j y_1 + \beta_{2n_j}^j y_2 + \cdots + \beta_{(j-1)n_j}^j y_{j-1}) y_j, \end{aligned}$$

for $j = 2, \dots, m$. Now the least power of y_j in Γ_j is $n_j + 1$. Our claim is that $y_j^{n_j} \neq 0$. If not, let $y_j^{n_j} = 0$. Then $y_j^{n_j} \in I$. But the least power of y_j , which appears as a term in a polynomial in the ideal I is $y_j^{n_j+1}$. This is a contradiction. Hence $y_j^{n_j} \notin I$, i.e., $y_j^{n_j} \neq 0$ in $H^*(M^n(P, \lambda); \mathbb{Z}_2)$ for $j = 1, 2, \dots, m$.

Using the Poincare duality, we have $x_1 \cdots x_{\mathcal{N}_1} x_{\mathcal{N}_1+1} \cdots x_{\mathcal{N}_2} \neq 0$. Now $x_{\mathcal{N}_1+1} = \dots = x_{\mathcal{N}_2}$ and all of them are either y_2 or, $y_1 + y_2$. If all of them are equal to y_2 , then $y_1^{n_1} y_2^{n_2} \neq 0$. If all of them are equal to $y_1 + y_2$ then

$$x_1 \cdots x_{\mathcal{N}_1} x_{\mathcal{N}_1+1} \cdots x_{\mathcal{N}_2} = y_1^{n_1} (y_1 + y_2)^{n_2} = y_1^{n_1} (y_2^{n_2} + y_1 \cdot f(y_1, y_2)) \neq 0.$$

We have $y_1^{n_1+1} = 0$. Therefore, $y_1^{n_1} y_2^{n_2} \neq 0$ holds for both the cases.

A similar computation as above implies that $y_1^{n_1} y_2^{n_2+1} = 0$, since $x_1 \cdots x_{\mathcal{N}_1} \Gamma_2 = 0$. Again using Poincare duality $x_1 \cdots x_{\mathcal{N}_1} x_{\mathcal{N}_1+1} \cdots x_{\mathcal{N}_2} \cdots x_{\mathcal{N}_1} x_{\mathcal{N}_2+1} \cdots x_{\mathcal{N}_3} \neq 0$ and we have two other condition $y_1^{n_1+1} = 0$ and $y_1^{n_1} y_2^{n_2+1} = 0$. Using this

$$y_1^{n_1} (y_1 + y_2)^{n_2} (y_1 + y_2 + y_3)^{n_3} = y_1^{n_1} y_2^{n_2} y_3^{n_3}.$$

Then we have $y_1^{n_1} y_2^{n_2} y_3^{n_3} \neq 0$ and $y_1^{n_1} y_2^{n_2} y_3^{n_3+1} = 0$.

Now inductively, we have $y_1^{n_1+1} = 0$, $y_1^{n_1} y_2^{n_2+1} = 0, \dots, y_1^{n_1} y_2^{n_2} \cdots y_{m-1}^{n_{m-1}+1} = 0$. Moreover, using the Poincare duality we get,

$$x_1 \cdots x_{\mathcal{N}_1} x_{\mathcal{N}_1+1} \cdots x_{\mathcal{N}_2} x_{\mathcal{N}_2+1} \cdots x_{\mathcal{N}_{m-1}+1} \cdots x_{\mathcal{N}_m} \neq 0.$$

This implies $y_1^{n_1} y_2^{n_2} \cdots y_m^{n_m} \neq 0$. □

We now use Lemma 2.12 to get the sharp lower bound on the topological complexity of generalized real Bott manifolds.

Theorem 2.13. *Let $M^n(P, \lambda)$ be a small cover over $P = \prod_{j=1}^m \Delta^{n_j}$. If $n_j \leq 2^{r_j} - 1 < 2n_j$, then*

$$\text{TC}(M^n(P, \lambda)) \geq (2^{r_1} + \cdots + 2^{r_m}) - (m - 1).$$

Proof. Let $d_j = 2^{r_j} - 1$ such that $n_j \leq d_j < 2n_j$ for $j = 1, \dots, m$. Let $\mathfrak{a}_j := 1 \otimes y_j - y_j \otimes 1$. Then \mathfrak{a}_j is in the ideal of the zero-divisors of $H^*(M^n(P, \lambda); \mathbb{Z}_2)$ because \mathfrak{a}_j belongs to the kernel of the homomorphism

$$\cup: H^*(M^n(P, \lambda); \mathbb{Z}_2) \otimes H^*(M^n(P, \lambda); \mathbb{Z}_2) \rightarrow H^*(M^n(P, \lambda); \mathbb{Z}_2). \quad (13)$$

Now,

$$\begin{aligned}\mathfrak{a}_j^{d_j} &= (1 \otimes y_j - y_j \otimes 1)^{d_j} \\ &= \sum_{s=0}^{d_j} (-1)^{d_j-s} \binom{d_j}{s} (1 \otimes y_j)^s (y_j \otimes 1)^{d_j-s} \\ &= \sum_{s=0}^{d_j} (-1)^{d_j-s} \binom{d_j}{s} (y_j^{d_j-s} \otimes y_j^s).\end{aligned}$$

Note that the binomial coefficients $\binom{2^{r_j}-1}{i}$ are odd for all $0 \leq i \leq d_j$. The binomial expansion of $\mathfrak{a}_j^{d_j}$ contains the term $(y_j^{d_j-n_j} \otimes y_j^{n_j})$ which is non-zero and there is no other same term in the expression of $\mathfrak{a}_j^{d_j}$. So $\mathfrak{a}_j^{d_j}$ is nonzero. Therefore $\mathfrak{a}_1^{d_1} \cdots \mathfrak{a}_m^{d_m}$ is non-zero in the kernel of the map in (13). Hence zero-divisors-cup-length of $H^*(M^n(P, \lambda); \mathbb{Z}_2)$ is greater than or equal to $d_1 + \cdots + d_m$. Therefore, we have $\text{TC}(M^n(P, \lambda)) \geq d_1 + \cdots + d_m + 1$, i.e.,

$$\text{TC}(M^n(P, \lambda)) \geq 2^{r_1} + \cdots + 2^{r_m} - (m - 1).$$

This proves the claim. \square

Theorem 2.13 gives a better lower bound of the LS-category of a generalized real Bott manifold if each n_j is a power of 2.

Corollary 2.14. *Let $M^n(P, \lambda)$ be a small cover over $P = \prod_{j=1}^m \Delta^{n_j}$. If $n_j = 2^{r_j-1}$ for $j = 1, 2, \dots, m$ then*

$$\text{TC}(M^n(P, \lambda)) \geq 2n - m + 1.$$

Proof. If $n_j = 2^{r_j-1}$ then r_j satisfies $n_j \leq 2^{r_j} - 1 < 2n_j$. Using Theorem 2.13, we get

$$\text{TC}(M^n(P, \lambda)) \geq 2^{r_1} + \cdots + 2^{r_m} - m + 1.$$

Since $n = n_1 + \cdots + n_m$, we have $\text{TC}(M^n(P, \lambda)) \geq 2n - m + 1$. \square

We now compute the sharp upper bound on $\text{TC}(M^n(P, \lambda))$.

Theorem 2.15. *Let $M^n(P, \lambda)$ be a small cover over $P = \prod_{j=1}^m \Delta^{n_j}$. If $n_j > 1$ is odd for each $j \in \{i_1, \dots, i_k\} \subseteq \{1, \dots, m\}$ then*

$$\text{TC}(M^n(P, \lambda)) \leq 2n - k + 1.$$

Proof. The real moment angle manifold corresponding to the polytope $P = \prod_{j=1}^m \Delta^{n_j}$ is $\prod_{j=1}^m S^{n_j}$. Let $m_j = \frac{n_j+1}{2}$. Then, for each $j \in \{i_1, \dots, i_k\} \subset \{1, \dots, m\}$, there is a free S^1 -action on

$$S^{n_j} = \{(z_1, \dots, z_{m_j}) \in \mathbb{C}^{m_j} \mid \sum_{j=1}^{m_j} |z_j|^2 = 1\}$$

defined by $\eta(z_1, \dots, z_{m_j}) \mapsto (\eta z_1, \dots, \eta z_{m_j})$. Thus, there exists a free $(S^1)^k$ -action on $\prod_{j=1}^m S^{n_j}$. Now $M^n(P, \lambda)$ can be realized as the orbit space of the moment angle manifold $\prod_{j=1}^m S^{n_j}$ by a free $(\mathbb{Z}_2)^m$ -action, see [15, Remark 2.3]. Since both of these actions are a multiplication by a complex number, they commute and induce a free action of $(S^1 / \langle e^{\pi\sqrt{-1}} \rangle)^k$ -action on $M^n(P, \lambda)$ if $n_j > 1$ is odd for each $j \in \{i_1, \dots, i_k\} \subset \{1, \dots, m\}$. Then, the result follows from [21, Corollary 5.3], since $S^1 / \langle e^{\pi\sqrt{-1}} \rangle \cong S^1$. \square

Corollary 2.16. *Let $M^n(P, \lambda)$ be a small cover over $P = \prod_{j=1}^m \Delta^{n_j}$. If $1 < n_j \leq 2^{r_j} - 1 < 2n_j$ and n_j is odd for each $j \in \{i_1, \dots, i_k\} \subset \{1, \dots, m\}$ then*

$$2^{r_1} + \cdots + 2^{r_m} - m + 1 \leq \text{TC}(M^n(P, \lambda)) \leq 2n - k + 1.$$

Proof. This follows from Theorem 2.13 and 2.15. \square

Corollary 2.17. *Let $M^n(P, \lambda)$ be a small cover over $P = \prod_{j=1}^m \Delta^{n_j}$. If $n_j = 2^{r_j} + 1$ for all $j \in \{1, 2, \dots, m\}$ and $r_j \geq 1$ then*

$$2n - 3m + 1 \leq \text{TC}(M^n(P, \lambda)) \leq 2n - m + 1.$$

Example 2.18. (1) $47 \leq \text{TC}(M^n(P, \lambda)) \leq 49$ if $P = \Delta^{16} \times \Delta^8$.
(2) $47 \leq \text{TC}(M^n(P, \lambda)) \leq 50$ if $P = \Delta^{16} \times \Delta^9$ or $\Delta^{17} \times \Delta^8$.
(3) $47 \leq \text{TC}(M^n(P, \lambda)) \leq 51$ if $P = \Delta^{17} \times \Delta^9$.

3. LS-CATEGORY AND TOPOLOGICAL COMPLEXITY OF DOLD MANIFOLDS OF REAL TORUS TYPE

In this section, we recall Dold manifolds of Bott type from [29] and study some of its basic properties. Then, we compute some lower and upper bounds for the topological complexity of these manifolds.

Recall the construction of small covers from Section 2. We now describe the involutions on small cover $M^n(P, \lambda)$ generated by \mathbb{Z}_2 -subgroups of \mathbb{Z}_2^n . Note that any \mathbb{Z}_2 -subgroup G of \mathbb{Z}_2^n is given by $G = \langle (a_1, \dots, a_n) \rangle$ where $a_i \in \{1, -1\}$ for $1 \leq i \leq n$. Then define $\tau: M^n(P, \lambda) \rightarrow M^n(P, \lambda)$ by

$$\tau([x, (t_1, \dots, t_n)]) = [x, (a_1 t_1, \dots, a_n t_n)] \quad (14)$$

where $x \in P$, $(t_1, \dots, t_n) \in \mathbb{Z}_2^n$, and $[x, (t_1, \dots, t_n)]$ is the equivalence class of $(x, (t_1, \dots, t_n))$ under \sim in (3). One can observe that the i -th coordinate subgroup $1 \times \dots \times 1 \times \mathbb{Z}_2 \times 1 \times \dots \times 1$ of \mathbb{Z}_2^n determine a nontrivial involution on $M^n(P, \lambda)$, say i -th involution. We observe that, if $(t_1, \dots, t_n, p) \sim (s_1, \dots, s_n, p)$, then $(a_1 t_1, \dots, a_n t_n, p) \sim (a_1 s_1, \dots, a_n s_n, p)$ as $a_i^{-1} t_i^{-1} a_i s_i = t_i^{-1} s_i$. This shows that the relation \sim is preserved under any involution generated by the \mathbb{Z}_2 -subgroup of \mathbb{Z}_2^n . The following example gives involution on some $M(P, \lambda)$ outside of these types.

Example 3.1. *Let $M(P, \lambda)$ be an n -dimensional real torus manifold such that there is an involution ζ' on P satisfying $\lambda(\zeta'(F)) = \lambda(F)$ for any facet F of P . Then, ζ' induces an involution ζ on $M(P, \lambda)$ which commutes with the \mathbb{Z}_2^n -action on $M(P, \lambda)$. Let $\mathfrak{q}: M(P, \lambda) \rightarrow P$ be the orbit map. Suppose that the set $\{\lambda(F) \mid F \in \mathcal{F}(P) \text{ and } F \cap P^{(\zeta')} \neq \emptyset\}$ generates \mathbb{Z}_2^n . Then one can show from the equivalence relation \sim in (3) that $\mathfrak{q}^{-1}(P^{(\zeta')})$ is path-connected.*

Recall the height function f on a simple polytope P that was introduced in the proof of Theorem 3.1 of [11]. One can orient the edges of polytope so that the f increases along them. This orientation makes the 1-skeleton of P into a directed graph.

Theorem 3.2. *Let M be an n -dimensional small cover such that $M^{(\tau)}$ is path-connected for some involution τ on M and the $\langle \tau \rangle$ -action commutes with the \mathbb{Z}_2^n -action on M . Then, there exist $(n+1)$ many $\langle \tau \rangle$ -invariant categorical sets of X and*

$$\text{cat}_{\langle \tau \rangle}(M) = n + 1. \quad (15)$$

Proof. We have $M \cong M^n(P, \lambda)$ for a \mathbb{Z}_2 -characteristic pair (P, λ) by Proposition 2.4. We also have $n + 1 = \text{cat}(M^n(P, \lambda))$ by . Therefore, using [1, Section 2], one can get $n + 1 = \text{cat}(M^n(P, \lambda)) \leq \text{cat}_{\langle \tau \rangle}(M^n(P, \lambda))$, since the fixed point set M^τ is a path-connected space.

Now we describe the τ -invariant categorical cover of $M^n(P, \lambda)$ containing $n + 1$ many sets. For any vertex v of P , let $U_v := P \setminus \cup_{v \notin F} F$ be the open subset of P . Let $V(P)$ be the set of vertices of P and

$$V_j = \{v \in V(P) : f(v) = j\},$$

where f is the height function of P . The DJ-construction of M gives us $q^{-1}(U_v) = \mathbb{Z}_2^n \times U_v / \sim$. Let $Y_j = \cup_{v \in V_j} q^{-1}(U_v)$ for $j = 0, \dots, n$. Since the actions of $\langle \tau \rangle$ and \mathbb{Z}_2^n commute, the space $q^{-1}(V_j)$ is τ -invariant. Observe that,

$$q^{-1}(U_v) = \frac{\mathbb{Z}_2^n \times U_v}{\sim} \simeq \frac{\mathbb{Z}_2^n \times \{v\}}{(g, v) \sim (h, v)},$$

for any $g, h \in \mathbb{Z}_2^n$ as v is a fixed point of an action of \mathbb{Z}_2^n on X . Observe that $P = \cup_{v \in V(P)} U_v$ and $U_v \cap U_w = \emptyset$ if $f(v) = j \neq f(w)$. Therefore, for each $0 \leq j \leq n$, one can see that

$$Y_j \simeq \frac{\mathbb{Z}_2^n \times V_j}{\sim} \simeq \{[(1, \dots, 1, v)] : v \in V_j\} \subset M^n(P, \lambda)^\tau. \quad (16)$$

Since $M^n(P, \lambda)^\tau$ is path connected, $q^{-1}(Y_j)$ is τ -invariantly contractible in Y . This proves the first part of the claim.

Now, it is evident that the homotopy equivalence in (16) is τ -equivariant for $j = 0, \dots, n$. Therefore, the cover $\{Y_j : 0 \leq j \leq n\}$ is a τ -equivariant categorical cover of X . This gives $\text{cat}_{\langle \tau \rangle}(X) \leq n + 1$. \square

We note that $n + 1 \leq \text{cat}_{\langle \tau \rangle}(M) \leq \#M^{\mathbb{Z}_2^n}$ for any involution determined by a \mathbb{Z}_2 -subgroup of \mathbb{Z}_2^n . Consider a small cover $M = M^n(P, \lambda)$. Observe that the column vector of λ determines a \mathbb{Z}_2 -action of $M^n(P, \lambda)$. With this, we have the following result.

Theorem 3.3. *Let $M := M^n(P, \lambda)$ be a small cover. Then*

$$\text{cat}(M_{F_{i_1}} \cup \dots \cup M_{F_{i_k}} \cup M^{\mathbb{Z}_2^n}) \leq \text{cat}_{\mathbb{Z}_2}(M) \leq \#M^{\mathbb{Z}_2^n},$$

where $M_{F_{i_j}}$ is the characteristic submanifold corresponding to the facet F_{i_j} for $1 \leq j \leq k$ such that $T_{F_{i_j}} = T_{F_{i_l}}$ for $1 \leq j \neq l \leq k$.

Proof. Observe that the fixed point set $M^{\mathbb{Z}_2} = M_{F_{i_1}} \cup \dots \cup M_{F_{i_1}} \cup M^{\mathbb{Z}_2^n}$. Then, the left inequality follows from [2, Corollary 2.9]. Let $q: M \rightarrow P$ be an orbit map. Then one can observe that $\{q^{-1}(U_v) \mid v \in V(P)\}$ forms a \mathbb{Z}_2 -categorical cover of M , where

$$q^{-1}(U_v) = \frac{\mathbb{Z}_2^n \times U_v}{\sim} \simeq \frac{\mathbb{Z}_2^n \times \{v\}}{(g, v) \sim (h, v)}.$$

This gives the right inequality. \square

Example 3.4. (1) Let $P = I^2$ and consider that facets are labelled by $\{F_1, F_2, F_3, F_4\}$. Define the characteristic function as follows $\lambda(F_1) = (1, 0) = \lambda(F_3)$, $\lambda(F_2) = (0, 1)$ and $\lambda(F_4) = (1, 1)$. Note that $M^n(P, \lambda) = M$ is the Klein bottle. Let \mathbb{Z}_2 -action on M be given by the subgroup $\mathbb{Z}_2 \times \{1\}$. Then one can observe that $M^{\mathbb{Z}_2} = M_{F_1} \sqcup M_{F_3} \cong S^1 \sqcup S^1$. Therefore, from Theorem 3.3 we get $\text{cat}_{\mathbb{Z}_2}(M) = 4$.

(2) One can generalize the above example. Let $P = P_{2m}$ be the $2m$ -gon. We label facets of P_{2m} by $\{F_1, F_2, \dots, F_{2m-1}, F_{2m}\}$. Then note that we can define a characteristic function λ such that $\lambda(F_{2i-1}) = (1, 0)$ for $1 \leq i \leq m$. Then the \mathbb{Z}_2 -action on $M = M^n(P, \lambda)$ determined by the \mathbb{Z}_2 -subgroup $\mathbb{Z}_2 \times \{1\}$ has fixed point $M^{\mathbb{Z}_2} = \cup_{i=1}^m M_{F_{2i-1}}$. Note that $M^{\mathbb{Z}_2}$ contains all vertices of P_{2m} and it is homeomorphic to disjoint union of m -many circles as $M_{F_i} \cong S^1$ for each $1 \leq i \leq m$. Therefore, Theorem 3.3 gives us

$$2m = \text{cat}(\sqcup_m S^1) \leq \text{cat}_{\mathbb{Z}_2}(M) \leq 2m.$$

Therefore, $\text{cat}_{\mathbb{Z}_2}(M) = 2m$.

The following result is a consequence of [6, Corollary 5.8] and Theorem 3.2.

Corollary 3.5. *Let M be an n -dimensional small cover such that $M^{\langle \tau \rangle}$ is path-connected for some involution τ on M and the $\langle \tau \rangle$ -action commutes with the \mathbb{Z}_2^n -action on M . Then,*

$$\mathrm{TC}_{\langle \tau \rangle}(M) \leq 2n + 1.$$

Next, we discuss the main objectives of this section. The projective product spaces were introduced by Davis in [9] as follows:

$$P(p_1, \dots, p_r) = \frac{S^{p_1} \times \dots \times S^{p_r}}{(x_1, \dots, x_r) \sim (-x_1, \dots, -x_r)}.$$

Davis computed the mod-2 cohomology algebra of this space. Recall that an n -dimensional real torus manifold admits \mathbb{Z}_2^n -action such that the orbit space of this action is a nice manifold with corners. Then one can note that any \mathbb{Z}_2 -subgroup of \mathbb{Z}_2^n induces an involution on M . Now, we consider the following identification spaces. Define

$$D(M; p_1, \dots, p_r) := \frac{M \times S^{p_1} \times \dots \times S^{p_r}}{(y, x_1, \dots, x_r) \sim (\tau(y), -x_1, \dots, -x_r)},$$

where $\tau: M \rightarrow M$ be an involution. Then $D(M; p_1, \dots, p_r)$ is a manifold of dimension $n + p_1 + \dots + p_r$. We call this manifold a Dold manifold of real torus type. Note that we have a fibre bundle

$$M \hookrightarrow D(M, p_1, \dots, p_r) \xrightarrow{p} P(p_1, \dots, p_r),$$

where $P(p_1, \dots, p_r)$ is the projective product space with $p_1 \leq \dots \leq p_r$.

The following result describes the mod-2 cohomology ring of $D(M; p_1, \dots, p_r)$. Let P be a retractable nice manifold with corners in the sense of [28, Definition 3.2], and $M^n(P, \lambda)$ a real torus manifold. Then $M^n(P, \lambda)$ has a \mathbb{Z}_2^n -equivariant cell-structure. Thus, the associated \mathbb{Z}_2^n acts trivially on $H^*(M^n(P, \lambda); \mathbb{Z}_2)$. We note that any simple polytope and the polytope in Example 2.3 are retractable nice manifolds with corners.

Proposition 3.6 ([29, Proposition 4.6]). *Let $p_1 \leq \dots \leq p_r$ and $M^n(P; \lambda)$ be a real torus manifold over a retractable P . Then the cohomology ring $H^*(D(M, p_1, \dots, p_r); \mathbb{Z}_2)$ is isomorphic as a graded \mathbb{Z}_2 -algebra to*

$$H^*(D(M, p_1, \dots, p_r); \mathbb{Z}_2) \cong \mathbb{Z}_2[\alpha] / (\alpha^{p_1+1}) \otimes \Lambda[\alpha_2, \dots, \alpha_r] \otimes H^*(M; \mathbb{Z}_2),$$

where $|\alpha| = p_1$, $|\alpha_i| = p_i$ for $i > 1$, and Λ denotes the mod-2-exterior algebra.

We now compute the LS-category of $D(M; p_1, \dots, p_r)$.

Theorem 3.7. *Let M be a real torus manifold over a retractable P and $p_1 \leq \dots \leq p_r$. Then*

$$\mathrm{cat}(D(M; p_1, \dots, p_r)) \geq \mathrm{cl}_{\mathbb{Z}_2}(M) + p_1 + r. \quad (17)$$

Moreover, if M is a small cover of dimension n with $M^{\langle \tau \rangle}$ is path connected, then

$$\mathrm{cat}(D(M; p_1, \dots, p_r)) = n + p_1 + r.$$

Proof. It follows from [9, Theorem 2.1] that $\mathrm{cl}_{\mathbb{Z}_2}(P(p_1, \dots, p_r)) = p_1 + r - 1$. Therefore, by Proposition 3.6, we get that $\mathrm{cl}_{\mathbb{Z}_2}(D(M; p_1, \dots, p_r)) = \mathrm{cl}_{\mathbb{Z}_2}(M) + \mathrm{cl}_{\mathbb{Z}_2}(P(p_1, \dots, p_r))$. Thus, the inequality in (17) follows from the cup-length lower bound on category.

Now if M is small cover, then it follows from [8, Proposition 2.5] and [19, Theorem 1.2]

$$\mathrm{cat}(D(M; p_1, \dots, p_r)) \leq q + \mathrm{cat}(P(p_1, \dots, p_r)) - 1 \leq q + p_1 + r - 1,$$

where q is the smallest integer such that M is covered by q -many τ -invariant categorical sets. From Theorem 3.2 it follows that $q = n + 1$. This proves the claim. \square

Next, we obtain some bounds for $\text{TC}(D(M; p_1, \dots, p_r))$.

Theorem 3.8. *Let $p_1 \leq \dots \leq p_r$ and M be a real torus manifold over a retractable P such that $M^{\langle \tau \rangle}$ is path-connected for some involution τ on M and the $\langle \tau \rangle$ -action commutes with the \mathbb{Z}_2^n -action on M . Then*

$$\text{zcl}_{\mathbb{Z}_2}(M) + \text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{p_1}) + r \leq \text{TC}(D(M; p_1, \dots, p_r)) \leq 2(\text{cat}_{\langle \tau \rangle}(M) + p_1 + r) - 1. \quad (18)$$

Moreover, $\text{TC}(D(M; p_1, \dots, p_r)) \leq 2(n + p_1 + r) + 1$ if M is an n -dimensional small cover.

Proof. Note that, using Proposition 3.6 we have

$$\text{zcl}_{\mathbb{Z}_2}(D(M, p_1, \dots, p_r)) = \text{zcl}_{\mathbb{Z}_2}(M) + \text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{p_1}) + r - 1.$$

Therefore, we get the left inequality of (18). The right inequality of (18) then follows from [8, Theorem 2.5]. The remaining follows from Theorem 3.2. \square

Corollary 3.9. *Let M be a real torus manifold satisfying the hypotheses in Theorem 2.7. Then,*

$$\text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^n) + \text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{p_1}) + 1 \leq \text{TC}(D(M; p_1)) \leq 2(n + p_1) + 1. \quad (19)$$

In particular, if $n = 2^s$ and $p_1 = 2^t$, then $2^{s+1} + 2^{t+1} - 1 \leq \text{TC}(D(M; p_1)) \leq 2^{s+1} + 2^{t+1} + 1$.

Example 3.10. *Consider S^n as a real toric manifold as in Example 2.3 for $n \geq 2$. Then $\text{cat}_{\langle \tau \rangle}(S^n) = 2$. Also, $\text{zcl}_{\mathbb{Z}_2}(S^{2k+1}) = 1$ and $\text{zcl}_{\mathbb{Z}_2}(S^{2k}) = 2$ for any $k \in \mathbb{N}$. Thus,*

$$\text{zcl}_{\mathbb{Z}_2}(S^n) + \text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{p_1}) + 1 \leq \text{TC}(D(M; p_1)) \leq 2p_1 + 5.$$

If p_1 is the power of 2, then $2p_1 + r + \text{zcl}_{\mathbb{Z}_2}(S^n) \leq \text{TC}(D(M; p_1, \dots, p_r)) \leq 2p_1 + 2r + 3$.

4. SYMMETRIC TOPOLOGICAL COMPLEXITY OF SMALL COVERS

In [17], Farber and Grant studied the symmetric analogue of the motion planning problem and introduced the notion of symmetric topological complexity. In this section, we recall the basic terminologies related to symmetric topological complexity and state some results that we are going to use further. We compute the symmetric topological complexity of some classes of small covers.

Let Y be a path-connected space and consider the corresponding ordered configuration space of 2 points

$$F(Y; 2) = \{(x, y) \in Y \times Y \mid x \neq y\}.$$

Observe that the free path space fibration $\pi: PY \rightarrow Y \times Y$ restricts to a fibration

$$\pi': P'Y \rightarrow F(Y; 2), \quad (20)$$

where $P'Y$ is a subspace of PY consisting paths with distinct endpoints. That is $P'Y = \{\gamma: I \rightarrow Y \mid \gamma(0) \neq \gamma(1)\}$. Note that $\pi': P'Y \rightarrow F(Y; 2)$ is a \mathbb{Z}_2 -equivariant map, where the group \mathbb{Z}_2 acts freely on $F(Y; 2)$ via switching the coordinates, and on $P'Y$ by sending a path γ to its reverse path $\bar{\gamma}$. Therefore, we have the following fibration

$$\pi'': P'Y/\mathbb{Z}_2 \rightarrow B(Y; 2), \quad (21)$$

where $B(Y; 2)$ is the orbit space $F(Y; 2)/\mathbb{Z}_2$, also called an unordered configuration space of 2 points on Y . This map is also a fibration.

Definition 4.1. *The symmetric topological complexity of Y , $\text{TC}^S(Y)$, is defined by*

$$\text{TC}^S(Y) := 1 + \text{secat}(\pi'').$$

Note that the space $B(Y; 2)$ is empty if and only if Y is a single point. It was mentioned in [17] that, in this case, $\text{TC}^S(Y) = 1$ and if Y contains more than one point, then $\text{TC}^S(Y) \geq 2$. Moreover, it follows from [17, Example 7] that the symmetric topological complexity of a contractible space with more than two points is 2.

We now describe a lower bound on the symmetric topological complexity given in [17]. Let N_Y be the sub-ring of $H^*(Y) \otimes H^*(Y)$ spanned by the norm elements (i.e., the elements of the form $x \otimes y + y \otimes x$ with $x \neq y$). The following result follows from Corollary 9, Proposition 10, and Theorem 17 in [17].

Proposition 4.2. *Let Y be a closed smooth manifold. Then*

$$\max\{\text{TC}(Y), \text{cl}(N_Y) + 2\} \leq \text{TC}^S(Y) \leq 2\dim(Y) + 1.$$

Remark 4.3. *The element $1 \otimes y_j + y_j \otimes 1$ is same as $1 \otimes y_j - y_j \otimes 1$ in $N_{M^n(P, \lambda)}$. Thus the zero-divisors-cup-length of $M^n(P, \lambda)$ is the same as the cup-length of $N_{M^n(P, \lambda)}$.*

Theorem 4.4. *Let $M^n(P, \lambda)$ be a small cover over $P = \prod_{j=1}^m \Delta^{n_j}$. If $n_j \leq 2^{r_j} - 1 < 2n_j$, then*

$$\text{TC}^S(M^n(P, \lambda)) \geq (2^{r_1} + \cdots + 2^{r_m}) - m + 2.$$

In particular, if $n_j = 2^{r_j-1}$ for $j = 1, 2, \dots, m$, then

$$\text{TC}^S(M^n(P, \lambda)) \geq 2n - m + 2.$$

Proof. This follows from Theorem 2.13, Proposition 4.2, and Remark 4.3. \square

Theorem 4.5. *Let $M^n(P, \lambda)$ be a real torus manifold satisfying the hypotheses in Theorem 2.7. Then,*

$$\text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^n) + 2 \leq \text{TC}^S(M^n(P, \lambda)) \leq 2n + 1.$$

In particular, if $n = 2^s$, then

$$\text{TC}^S(M^n(P, \lambda)) = 2n + 1.$$

Proof. This follows from Theorem 2.7, Proposition 4.2 and Remark 4.3. \square

Theorem 4.6. *Let $M^n(P, \lambda)$ be a real torus manifold satisfying the hypotheses in Theorem 2.7. Then,*

$$\text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^n) + \text{zcl}_{\mathbb{Z}_2}(\mathbb{R}P^{p_1}) + 2 \leq \text{TC}^S(D(M(P, \lambda); p_1)) \leq 2(n + p_1) + 1.$$

In particular, if $n = 2^s$ and $p_1 = 2^t$, then $2^{s+1} + 2^{t+1} \leq \text{TC}^S(M^n(P, \lambda)) \leq 2^{s+1} + 2^{t+1} + 1$.

Proof. This follows from Theorem 2.7, Proposition 4.2 and Remark 4.3. \square

Acknowledgement. The authors would like to thank Bikramaditya Naskar and Subhankar Sau for some helpful discussion. Koushik Brahma thanks the Indian Statistical Institute Kolkata and Chennai Mathematical Institute for financial support. Navnath Daundkar thanks NBHM for the support through the grant 0204/10/(16)/2023/R&D-II/2789.

REFERENCES

- [1] A. Angel and H. Colman. Equivariant topological complexities. *Topological complexity and related topics, Contemp. Math*, 702:1–15, 2018.
- [2] M. Bayeh and S. Sarkar. Some aspects of equivariant LS-category. *Topology Appl.*, 196:133–154, 2015.
- [3] I. Bernstein. On the Lusternik–Schnirelmann category of Grassmannians. *Math. Proc. Cambridge Philos. Soc.*, 79(1):129–134, 1976.
- [4] K. Brahma, B. Naskar, S. Sarkar, and S. Sau. Various topological complexities of small covers and real bott manifolds. *arXiv preprint arXiv:2306.00916*, 2023.

- [5] S. Choi, M. Masuda, and D. Y. Suh. Quasitoric manifolds over a product of simplices. *Osaka J. Math.*, 47(1):109–129, 2010.
- [6] H. Colman and M. Grant. Equivariant topological complexity. *Algebr. Geom. Topol.*, 12(4):2299–2316, 2012.
- [7] O. Cornea, G. Lupton, J. Oprea, and D. Tanré. *Lusternik–Schnirelmann category*, volume 103 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [8] N. Daundkar and S. Sarkar. LS-category and topological complexity of several families of fibre bundles. *Accepted for publication in homology homotopy and applications, arXiv:2302.00468*, 2023.
- [9] D. M. Davis. Projective product spaces. *J. Topol.*, 3(2):265–279, 2010.
- [10] M. W. Davis. Groups generated by reflections and aspherical manifolds not covered by Euclidean space. *Ann. of Math. (2)*, 117(2):293–324, 1983.
- [11] M. W. Davis and T. Januszkiewicz. Convex polytopes, Coxeter orbifolds and torus actions. *Duke Math. J.*, 62(2):417–451, 1991.
- [12] A. Dold. Erzeugende der Thomschen Algebra \mathfrak{R} . *Math. Z.*, 65:25–35, 1956.
- [13] A. Dranishnikov. On topological complexity of non-orientable surfaces. *Topology Appl.*, 232:61–69, 2017.
- [14] A. Dranishnikov. On topological complexity of hyperbolic groups. *Proc. Amer. Math. Soc.*, 148(10):4547–4556, 2020.
- [15] R. Dsouza and V. Uma. Results on the topology of generalized real Bott manifolds. *Osaka J. Math.*, 56(3):441–458, 2019.
- [16] M. Farber. Topological complexity of motion planning. *Discrete Comput. Geom.*, 29(2):211–221, 2003.
- [17] M. Farber and M. Grant. Symmetric motion planning. *Contemporary Mathematics*, 438:85–104, 2007.
- [18] M. Farber, S. Tabachnikov, and S. Yuzvinsky. Topological robotics: motion planning in projective spaces. *Int. Math. Res. Not.*, (34):1853–1870, 2003.
- [19] S. Fişekci and L. Vandembroucq. On the LS-category and topological complexity of projective product spaces. *J. Homotopy Relat. Struct.*, 16(4):769–780, 2021.
- [20] J. González, M. Grant, E. Torres-Giese, and M. Xicoténcatl. Topological complexity of motion planning in projective product spaces. *Algebr. Geom. Topol.*, 13(2):1027–1047, 2013.
- [21] M. Grant. Topological complexity, fibrations and symmetry. *Topology Appl.*, 159(1):88–97, 2012.
- [22] M. Grant, G. Lupton, and J. Oprea. Spaces of topological complexity one. *Homology Homotopy Appl.*, 15(2):73–81, 2013.
- [23] M. Grant and S. Mescher. Topological complexity of symplectic manifolds. *Math. Z.*, 295(1-2):667–679, 2020.
- [24] S. Kuroki and Z. Lü. Projective bundles over small covers and the bundle triviality problem. *Forum Math.*, 28(4):761–781, 2016.
- [25] Z. Lü and M. Masuda. Equivariant classification of 2-torus manifolds. *Colloq. Math.*, 115(2):171–188, 2009.
- [26] L. Lyusternik and L. Schnirelmann. Topological methods in variational problems and their application to the differential geometry of surfaces. *Uspehi Matem. Nauk (N.S.)*, 2(1(17)):166–217, 1947.
- [27] A. Nath and P. Sankaran. On generalized Dold manifolds. *Osaka J. Math.*, 56(1):75–90, 2019.
- [28] S. Sarkar. Equivariant K -theory of divisive torus orbifolds. *Rocky Mountain J. Math.*, 50(1):255–266, 2020.
- [29] S. Sarkar and P. Zvengrowski. On generalized projective product spaces and Dold manifolds. *Homology Homotopy Appl.*, 24(2):265–289, 2022.

DEPARTMENT OF MATHEMATICS, CHENNAI MATHEMATICAL INSTITUTE, SIRUSERI 603103, INDIA.
 Email address: koushikbrahma95@gmail.com

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH PUNE, INDIA.
 Email address: navnath.daundkar@acads.iiserpune.ac.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY MADRAS, CHENNAI 600036, INDIA.
 Email address: soumen@iitm.ac.in