

Robust Genomic Prediction and Heritability Estimation using Density Power Divergence

Upama Paul Chowdhury, Susmita Das and Abhik Ghosh
Indian Statistical Institute, Kolkata, India.

January 17, 2024

Abstract

This manuscript delves into the intersection of genomics and phenotypic prediction, focusing on the statistical innovation required to navigate the complexities introduced by noisy covariates and confounders. The primary emphasis is on the development of advanced robust statistical models tailored for genomic prediction from single nucleotide polymorphism (SNP) data collected from genome-wide association studies (GWAS) in plant and animal breeding and multi-field trials. The manuscript explores the limitations of traditional marker-assisted recurrent selection, highlighting the significance of incorporating all estimated effects of marker loci into the statistical framework and aiming to reduce the high dimensionality of GWAS data while preserving critical information. This paper introduces a new robust statistical framework for genomic prediction, employing one-stage and two-stage linear mixed model analyses along with utilizing the popular robust minimum density power divergence estimator (MDPDE) to estimate genetic effects on phenotypic traits. The study illustrates the superior performance of the proposed MDPDE-based genomic prediction and associated heritability estimation procedures over existing competitors through extensive empirical experiments on artificial datasets and application to a real-life maize breeding dataset. The results showcase the robustness and accuracy of the proposed MDPDE-based approaches, especially in the presence of data contamination, emphasizing their potential applications in improving breeding programs and advancing genomic prediction of phenotyping traits.

Keywords: Genome-wide Association Study (GWAS), Genomic Prediction, Field Trials, Robust Estimation, Minimum Density Power Divergence Estimator, Linear Mixed Models.

1 Introduction

In the expansive landscape of statistical innovation, the intersection of genomics and phenotypic prediction constitutes a pivotal frontier. The ascent of high-throughput genotyping technologies, epitomized by the widespread adoption of single nucleotide polymorphism (SNP) data, has propelled a statistical revolution geared toward unraveling the genetic basis of complex traits. This manuscript unfolds as an answer to the escalating demand for advanced statistical models, meticulously tailored to exploit the richness of genomic data while navigating the intricacies introduced by covariates and confounders. In the realm of statistical genomics, genome-wide association studies (GWAS) serve as the vanguard and are a vital scientific method for identifying genetic variations associated with specific traits, diseases, or conditions in both animal and plant breeding (Uffelmann et al., 2021 [28]; Lourenço et al., 2020 [19]). This process is meticulously governed by an arsenal of statistical models and methods, each calibrated to distill meaningful insights from the vast genomic dataset. Unlike traditional marker-assisted recurrent selection, GWAS transcends the constraints of significance, incorporating all estimated effects of marker loci as well as genomic breeding values of individuals into its statistical framework to predict marker effects, irrespective of their individual association with the phenotype.

The integration of genomics into phenotypic prediction assumes a pivotal role in deciphering the intricate statistical code governing trait variations. In the dynamic interplay of genomics and statistics, the significance of statistical models that transcend traditional limitations becomes paramount. Advanced statistical techniques enhance the power and precision of genetic association analyses, identifying subtle genetic effects often missed by traditional methods when dealing with numerous genetic markers. Many of these techniques aim to reduce high-dimensionality of GWAS data while preserving critical information, aiding in the identification of key genetic variants associated with traits (Giraud, 2021 [13];

Wainwright, 2019 [29]; Buhlmann et al., 2014 [5]). Some models, however, do take into account the high-dimensional nature of the marker matrix and attempt to fit the entire genome using methods such as penalized-likelihood or Bayesian shrinkage estimation. However, several issues have been identified when employing these models for effect estimation including computational intensity and modeling complexity. Shen et al. (2013) [26] proposed that utilizing heteroscedastic marker variances in ridge regression offers an alternative to Bayesian genome-wide prediction models. To illustrate the superior effectiveness of the heteroscedastic effect model, Hofheinz and Frisch (2014) [15] employed two methods within a one-stage model analysis where a single model incorporates all random effects (genetic effects, block effects, replicate effects, etc.) following classical literature. Alternatively, to avoid the computational complexity of this one-stage method, many authors proposed a second two-stage mixed model framework, where the random components are partitioned into two distinct stages; firstly genetic effects are considered as fixed effects while block and replicate effects are considered as random effects, and subsequently, in the second stage, genetic effects are regarded as random variables (Piepho et al., 2012 [21]; Estaghirou et al., 2013 [6]). These types of data analyses have been commonly applied in plant and animal breeding techniques and multi-factor field trials in previous literature (Westhues et al., 2021 [30]; Tanaka, 2020 [27]; Resende, 2016 [24]; Bernal-Vasquez et al., 2016 [4]; Estaghirou et al., 2015 [7]; Falke et al., 2014 [8]; Shen et al., 2013 [26]; Piepho, 2009 [22]; Möhring and Piepho, 2009 [20]; Barbosa et al., 2005[1]).

This manuscript embarks on a statistical journey, navigating the challenges posed by noisy covariates and confounders to refine genomic prediction even under possible data contamination, aligning with the evolving demands of precision statistical analyses. We present a statistically robust framework for genomic prediction, building upon the foundation laid by GWAS. The overarching objectives revolve around the development and validation of a so-

phisticated robust statistical model adept at seamlessly integrating SNP data with a myriad of covariates. Through a statistically rigorous journey, we aim to showcase the model’s superiority in capturing the nuanced interplay between genetic markers, environmental factors, and covariate effects. The primary goal is to provide a powerful tool that not only enhances the precision and reliability of genomic predictions but also stands as a testament to the evolving landscape of statistical methodologies in genomics. Two distinct model based approaches are employed for this purpose, namely one-stage and two-stage linear mixed model analysis; in both the approaches we additionally propose to suitably utilize the popular robust minimum density power divergence estimator (MDPDE) to estimate genetic effects on various phenotypic traits. The MDPDE is a robust statistical procedure employed to address the complexities of the linear mixed model setup, emphasizing robustness and accuracy in obtaining reliable estimates of genetic effects (Lee and Jo, 2023 [18]; Saraceno et al., 2020 [25]; Goegebeur et al., 2014 [14]). Density power divergence (DPD) is pivotal to our analysis, quantifying the dissimilarity between probability density functions. It is a fundamental component of the robust estimation process, particularly suited for addressing contaminated data and model misspecification (Ghosh and Basu, 2013 [10]; Piepho et al., 2012b [21]). Notably, the MDPDE has gained widespread popularity due to its high robustness, computational simplicity, and its capacity to offer a direct interpretation akin to the maximum likelihood estimator (MLE). In our study, the robust statistical methods are deliberately designed to withstand influential factors, including the presence of outliers in various directions, deviations from normality, and issues stemming from model misspecification. To underscore the robustness of these techniques, we subject them to examination using Monte-Carlo simulations involving both uncontaminated and contaminated data. Furthermore, our investigation extends to practical application as we evaluate these methods by simulating data in two distinct setups and assessing their performance on a real dataset

related to maize (*Zea mays*) genetics taken from Hofheinz and Frisch (2014) [15].

Additionally, in the present study, we propose a robust procedure for estimating heritability, as a by-product of genomic estimation, that leads to more accurate result than traditional heritability estimators especially when the data is contaminated with outliers. Heritability is unequivocally entangled to bioethics. To extent that twin heritability estimates played a central role in debates of science from the 70s through the 90s, the advent of SNP-heritability warrants a renewed assessment. It quantifies the proportion of phenotypic variation in a population that can be attributed to genetic factors (Zhu and Zhou, 2020 [32]), yielding an estimate of the extent to which genetic differences contribute to the observed variability in a particular trait within a population. We assess the effectiveness of the proposed robust estimator of heritability through Monte Carlo simulation studies and by applying them to an actual maize dataset mentioned above.

2 Methodology

2.1 Mathematical Model for Genomic Prediction

Suppose that our data consist of N individual phenotype observations in r replicate and p biallelic single-nucleotide polymorphism (SNP) markers where genotype effects are distributed in B blocks and L confounder or controllers. Let \mathbf{y}_k denotes the vector of n_k phenotype data for the k -th replicate for each $k = 1, \dots, r$, \mathbf{X}_g denotes the matrix of p SNP markers along the genome for N individuals (\mathbf{X}_g has p columns for the SNP observations, usually coded as 0, 1, -1 for homo zygote aa, the hetero-zygote Aa and the other homo-zygote AA, respectively) \mathbf{X}_b denotes the design matrix for random block effect and \mathbf{Z} denotes the $N \times L$ matrix of confounders. We denote the k -th partition of rows of \mathbf{Z} , \mathbf{X}_g and \mathbf{X}_b associated with each \mathbf{y}_k by \mathbf{Z}_k , \mathbf{X}_{gk} and \mathbf{X}_{bk} , respectively, for all $k = 1, \dots, r$. Following

preceding works, e.g., Tanaka, 2020 [27]; Lourenço et al., 2020 [19]; Estaghvirou et al., 2014 [6]; Piepho et al., 2012 [23], let us assume the a linear mixed model (LMM) for SNP data in the k -th replicate as given by

$$\mathbf{y}_k = \mathbf{Z}_k\boldsymbol{\gamma} + \mathbf{X}_{gk}\mathbf{u}_g + \mathbf{X}_{bk}\mathbf{u}_b + \boldsymbol{\epsilon}_k, \quad k = 1, \dots, r, \quad (1)$$

where the p -vector of random SNP effect $\mathbf{u}_g \sim N(\mathbf{0}, \boldsymbol{\Sigma}_g)$, the random block effects $\mathbf{u}_b \sim N(\mathbf{0}, \sigma_b^2 \mathbb{I}_B)$, $\boldsymbol{\gamma}$ denotes the l -vector of the confounder effects and the random error $\boldsymbol{\epsilon}_k \sim N(\mathbf{0}, \sigma_e^2 \mathbb{I}_{n_k})$ for $k = 1, \dots, r$. Note that $N = \sum_{k=1}^r n_k$ which is the total number of observation and individual replication is assumed to be independent of each other. Put $\mathbf{y} = (\mathbf{y}'_1, \mathbf{y}'_2, \dots, \mathbf{y}'_r)'$ to denote the vector of all phenotype data. Then, we can rewrite the LMM in (1) as

$$\mathbf{y} = \mathbf{Z}\boldsymbol{\gamma} + \begin{bmatrix} \mathbf{X}_g & \mathbf{X}_b \end{bmatrix} \begin{bmatrix} \mathbf{u}_g \\ \mathbf{u}_b \end{bmatrix} + \boldsymbol{\epsilon} = \mathbf{Z}\boldsymbol{\gamma} + \mathbf{X}\mathbf{u} + \boldsymbol{\epsilon}, \quad (2)$$

where $\boldsymbol{\epsilon} = (\boldsymbol{\epsilon}'_1, \dots, \boldsymbol{\epsilon}'_r)'$, $\mathbf{X} = \begin{bmatrix} \mathbf{X}_g & \mathbf{X}_b \end{bmatrix}$ and $\mathbf{u} = \begin{bmatrix} \mathbf{u}_g \\ \mathbf{u}_b \end{bmatrix}$. Based on the observed data, our aim is to first estimate the fixed effect $\boldsymbol{\gamma}$ and the random effect \mathbf{u} and then use them for genomic prediction and heritability estimation.

2.2 Estimating Marker Effects and Breeding Values

For ordinary Ridge regression the fixed effect $\boldsymbol{\gamma}$ and the random effect \mathbf{u} can be estimated jointly via Henderson's mixed model equation (Henderson 1953) given by

$$\begin{bmatrix} \mathbf{Z}^T \mathbf{Z} & \mathbf{Z}^T \mathbf{X} \\ \mathbf{X}^T \mathbf{Z} & \mathbf{X}^T \mathbf{X} + \Lambda \end{bmatrix} \begin{bmatrix} \boldsymbol{\gamma} \\ \mathbf{u} \end{bmatrix} = \begin{bmatrix} \mathbf{Z}^T \mathbf{y} \\ \mathbf{X}^T \mathbf{y} \end{bmatrix}, \quad (3)$$

where $\mathbf{\Lambda} = \text{Diag}(\lambda_1, \lambda_2, \dots, \lambda_p)$, with λ_j being the shrinkage parameter of j -th SNP for $j = 1, \dots, p$, which is indeed taken as a function of σ_e^2 and $\mathbf{\Sigma}_g$.

For estimating the marker-effects, the traditional approach is to minimize the likelihood based loss function associated with the model (2) which can be simplified to have the form

$$L(\mathbf{u}, \boldsymbol{\gamma}) = (\mathbf{y} - \mathbf{X}\mathbf{u} - \mathbf{Z}\boldsymbol{\gamma})^T(\mathbf{y} - \mathbf{X}\mathbf{u} - \mathbf{Z}\boldsymbol{\gamma}) + \mathbf{u}^T \mathbf{\Lambda} \mathbf{u}.$$

This loss function can be minimized analytically to get an estimator of the random-effect u (including both SNP-marker effects and block effects) as given by

$$\hat{\mathbf{u}} = (\mathbf{X}^T \mathbf{M}_{\mathbf{Z}} \mathbf{X} + \mathbf{\Lambda})^{-1} \mathbf{X}^T \mathbf{M}_{\mathbf{Z}} \mathbf{y}, \quad (4)$$

where $\mathbf{M}_{\mathbf{Z}} = \mathbf{I} - \mathbf{Z}(\mathbf{Z}^T)^{-1}\mathbf{Z}^T$ is the projection matrix removing the effect of confounding or controller variables, $\mathbf{\Lambda}$ is a diagonal matrix of shrinkage parameters λ_j , $j = 1, \dots, p$. These shrinkage parameters λ_j , being a function of the variance parameters, can be estimated from different types of variance estimators and subsequently we can get the estimated $\hat{\mathbf{u}}$ from (4) using estimated values of λ_j s in $\mathbf{\Lambda}$. Then, for genomic prediction, the estimate of breeding value will be given by [26]

$$\hat{\mathbf{g}} = \mathbf{X}_g \cdot \hat{\mathbf{u}}_g, \quad (5)$$

and the estimated phenotype effects can be obtained as

$$\hat{\mathbf{y}} = \mathbf{Z} \cdot \hat{\boldsymbol{\gamma}} + \mathbf{X} \cdot \hat{\mathbf{u}}. \quad (6)$$

We would like to emphasize that there exist several approaches for obtaining the estimates of random effects and error variances leading to different estimates of the shrinkage parameters in the literature of genomic prediction. For ordinary ridge regression, we assume the

simplest form $\Sigma_g = \sigma_g^2 \mathbb{I}_p$ and then term Λ can be replaced by $\lambda \mathbb{I}$ with the constant shrinkage parameter being computed as $\hat{\lambda} = \frac{\hat{\sigma}_e^2}{\hat{\sigma}_g^2}$. However, while using the likelihood based estimation approaches, it has been shown that the heteroscedastic values of λ_j works significantly better where we assume $\Sigma_g = \text{Diag}(\sigma_{g1}^2, \dots, \sigma_{gp}^2)$ and estimate different shrinkage parameters, corresponding to different SNPs, as $\hat{\lambda}_j = \frac{\hat{\sigma}_e^2}{\hat{\sigma}_{gj}^2}$ for every $j = 1, \dots, p$. The most common one is to use the restricted maximum likelihood estimator under the assumed LMM which is often referred to as the ridge regression (RR) estimator. Hofheinz and Frisch (2014) [15] proposed two methods, to be referred to as RMLV and RMLA, that are shown to have significantly improved performances over the classical RR estimator for genomic prediction; they both estimate the random effect $\hat{\mathbf{u}}$ using (4) with λ_j replaced by its estimators $\hat{\lambda}_j$ for all j under the model assumption $\Sigma_g = \text{Diag}(\sigma_{g1}^2, \dots, \sigma_{gp}^2)$, but these $\hat{\lambda}_j$ are obtained in two ways as described below:

- RMLA: First, a moment estimator of marker-specific variance component is obtained from random single-factor analysis of variance (ANOVA) for j -th marker as given by

$$\hat{\sigma}_{gj}^{2*} = \frac{MQM_j - MQE_j}{\frac{1}{2} \cdot (N - \sum_i \frac{n_i^2}{N})}, \quad (7)$$

where MQM_j denotes the mean square due to j -th SNP marker, MQE_j denotes the error for the j -th SNP marker, and n_i is the numbers of individuals in the i -th SNP-marker factors for $i = 1, 2, 3$ corresponding to three SNP-marker factors 0, 1, -1, respectively. Then, the shrinkage parameters are estimated from the resulting variance estimates as given by

$$\hat{\lambda}_j = \frac{\sigma_e^2}{\sigma_u^2} \cdot \frac{\sum_j \hat{\sigma}_{gj}^{2*}}{\hat{\sigma}_{gj}^{2*}}, \quad j = 1, \dots, p. \quad (8)$$

Here, σ_u^2 total genetic variance estimated from the model (2)

- RMLV: This method is a modification of the restricted maximum likelihood procedure, producing heteroscedastic SNP-variance through appropriate expectation-maximization algorithm. In this approach, we first obtain the estimates of the error variance and the marker variance for the j -th SNP marker (for each $j = 1, \dots, p$) as

$$\hat{\sigma}_e^2 = (\mathbf{y}'\mathbf{y} - \hat{\boldsymbol{\gamma}}'\mathbf{Z}'\mathbf{y} - \hat{\mathbf{u}}'X'\mathbf{y})/(N - 1), \quad \text{and} \quad \hat{\sigma}_{gj}^2 = (\hat{\mathbf{u}}_j'\hat{\mathbf{u}}_j - \sigma_e^2 \cdot \text{tr}(\mathbf{C}_{jj}))/j,$$

where $\text{tr}(\mathbf{C}_{jj})$ is the trace of inverse of the left most (coefficient) matrix in (3). Then the shrinkage parameters are estimated as $\hat{\lambda}_j = \frac{\hat{\sigma}_e^2}{\hat{\sigma}_{gj}^2}$ for each $j = 1, \dots, p$.

These two approaches, RMLV and RMLA, are considered as robust by the authors in [15] since the RMLV is robustified by taking different marker-variance and the RMLA is robustified with both different marker-variance and shrinkage parameters for each marker effects. However, these estimators are seen to be highly unstable against data contamination or noises in the observed datasets, which is the primary focus of the present work. Our proposed estimation approaches based on the minimum DPD estimation, that yield robust results in the presence of data contamination, are described in the following subsection.

2.3 The proposed Robust Estimation Approaches

2.3.1 The Minimum DPD Estimation for General set-up

The minimum DPD estimator (MDPDE) was initially developed by Basu et al. (1998)[2] for independent and identically distributed data, say $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ from a true distribution G , to be modelled by a parametric family of distributions indexed by $\boldsymbol{\theta} \in \Theta$. This approach relies on the minimization of a divergence measure, namely the DPD, between the true density g and the model density $f_{\boldsymbol{\theta}}$ to obtain a robust and highly efficient estimate of the unknown parameter vector $\boldsymbol{\theta}$. This particular divergence is defined in terms of a tuning parameter

$\alpha \geq 0$ as follows

$$d_\alpha(g, f_\theta) = \begin{cases} \int f_\theta^{\alpha+1} - (1 + (1/\alpha)) \int f_\theta^\alpha g + \frac{1}{\alpha} \int g^{1+\alpha}, & \alpha > 0 \\ \int g \ln\left(\frac{g}{f_\theta}\right), & \alpha = 0. \end{cases}$$

In practice, based on the observed data, we replace the unknown density g suitably by the help of the empirical distribution function estimate which leads to the following simplified objective function for obtaining the MDPDE as (see, e.g., Basu et al., 1998 [2])

$$\int f_\theta^{(1+\alpha)} - \left(1 + \frac{1}{\alpha}\right) \frac{1}{n} \sum_{i=1}^n f_\theta^\alpha(\mathbf{y}_i).$$

This leads to robust parameter estimates for $\alpha > 0$ with the tuning parameter α controlling the trade-off between the (asymptotic) efficiency under pure data and the robustness under contaminated data. It can be seen that the MDPDE at $\alpha = 0$ coincides (in a limiting sense) to the most efficient by highly non-robust maximum likelihood estimator (MLE) as the DPD measure at $\alpha = 0$ is indeed the Kullback-Leibler divergence. As $\alpha > 0$ increases, the corresponding MDPDE achieves greater robustness with a little loss in its pure data (asymptotic) efficiency; see Basu et al. (2011) [3] for more details about the MDPDE along with its applications under standard model set-ups.

The theory of the MDPDE has been extended for the independent but non-homogeneous data by Ghosh and Basu (2013) [10] and subsequently found many applications including regression and mixed models. Whenever our observed data $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ are independent but non-homogeneous with each $\mathbf{y}_i \sim g_i$ for all $i = 1, \dots, n$ and g_i is being modeled by the parametric density $f_i(\cdot, \theta)$, the MDPDE of the common unknown parameter θ can be obtained by minimizing the average divergence between the data points and the respective model densities; this becomes equivalent to the minimization of a simpler objective function

given by (Ghosh and Basu, 2013 [10])

$$H_n(\boldsymbol{\theta}) = \frac{1}{n} \sum_{i=1}^n \left[\int f_i(\mathbf{y}, \boldsymbol{\theta})^{(1+\alpha)} dy - \left(1 + \frac{1}{\alpha}\right) f_i(\mathbf{y}_i, \boldsymbol{\theta})^\alpha \right] = \frac{1}{n} \sum_{i=1}^n H_i(\mathbf{y}_i, \boldsymbol{\theta}) \quad (9)$$

Once again the case $\alpha = 0$ coincides with the MLE and we get improved robust estimators at larger values of $\alpha > 0$ with only a little loss in efficiency; see, e.g., Ghosh and Basu (2013, 2015, 2016) [10, 11, 12] and Ghosh (2019) [9] among many others, for further details and applications. In particular, Saraceno et al. (2020) [25] used this general theory of the MDPDE to develop robust estimators under the general set-up of linear mixed-models which we would be using in our context to get robust estimators of marker effects and breeding values and eventually a robust genomic prediction procedure.

2.3.2 The MDPDE based One-stage Approach for Genomic Prediction

For robust genomic prediction in the one-stage approach, we consider the mathematical model described in (1) with $\boldsymbol{\Sigma}_g = \sigma_g^2 \mathbb{I}_p$ and propose to estimate the associated parameters robustly using the MDPDE instead of the likelihood based approach described in Section 2.2. From (1), we get that $\mathbf{y}_k \sim N(\mathbf{Z}_k \boldsymbol{\gamma}, \mathbf{V}_k)$ independently for each $k = 1, \dots, r$, where

$$\mathbf{V}_k = \sigma_e^2 \mathbb{I}_{n_k} + (\mathbf{X}_g \mathbf{X}_g^T \sigma_g^2 + \mathbf{X}_b \mathbf{X}_b^T \sigma_b^2).$$

This set-up clearly belongs to the independent non-homogeneous set-up described in the preceding subsection and hence the robust MDPDE of the model parameters $\boldsymbol{\theta} = (\boldsymbol{\gamma}', \sigma_g^2, \sigma_b^2, \sigma_e^2)'$ can be obtained by minimizing the objective function given in (9) which, for the present case, further simplifies to

$$H_n(\boldsymbol{\theta}) = \frac{1}{r} \sum_{k=1}^r \frac{1}{(2\pi)^\alpha |\mathbf{V}_k|^{\alpha/2}} \left[\frac{1}{(\alpha + 1)} - \left(1 + \frac{1}{\alpha}\right) e^{\frac{\alpha}{2} (\mathbf{y}_k - \mathbf{Z}_k \boldsymbol{\gamma})^T \mathbf{V}_k^{-1} (\mathbf{y}_k - \mathbf{Z}_k \boldsymbol{\gamma})} \right]. \quad (10)$$

In our implementation, we have minimized the above objective function numerically in \mathbf{R} to obtain the MDPDE of model parameter in our numerical exploration presented in this paper. One can alternatively compute these MDPDEs by solving the associated estimating equations obtained as $\frac{\partial H_n(\boldsymbol{\theta})}{\partial \boldsymbol{\gamma}} = \mathbf{0}$ and $\frac{\partial H_n(\boldsymbol{\theta})}{\partial \boldsymbol{\delta}} = 0$ with $\boldsymbol{\delta} = (\sigma_g^2, \sigma_b^2, \sigma_e^2)$. We refer to Saraceno et al. (2020) [25] for more details including the asymptotic and robustness properties of the resulting parameter estimates.

Once we have obtained the robust MDPDE of the variance parameters as $\widehat{\sigma}_g^2$ and $\widehat{\sigma}_e^2$, we can use them to get an estimate of $\boldsymbol{\Lambda}$ as in the case of ordinary ridge regression, by assuming $\boldsymbol{\Lambda} = \lambda \mathbb{I}$ and estimating λ by $\widehat{\lambda} = \frac{\widehat{\sigma}_e^2}{\widehat{\sigma}_g^2}$. Then the random effects can be robustly estimated from (4) with $\boldsymbol{\Lambda} = \widehat{\lambda} \mathbb{I}$ and subsequently be used for robust estimation of the breeding values and prediction of phenotype effects following (5) and (6), respectively.

2.3.3 The MDPDE based Two-stage Approach for Genomic Prediction

For robust genomic prediction in the two-stage approach, we start with the mathematical model described in (2) with $\boldsymbol{\Sigma}_g = \sigma_g^2 \mathbb{I}_p$ and split it in two sequential models following Estaghevrou et al. (2013) [7] and Piepho et al. (2012) [23]. In the first stage, we consider a simpler model for the observed phenotype response as given by

$$\mathbf{y} = \boldsymbol{\eta} \boldsymbol{\mu} + \mathbf{X}_b \mathbf{u}_b + \boldsymbol{\epsilon}. \quad (11)$$

As described in Piepho et al. (2012) [23], $\boldsymbol{\eta}$ can be taken in such way that $\boldsymbol{\eta} \cdot \mathbf{Z}_1 = \mathbf{Z}$ and $\boldsymbol{\eta} \cdot \mathbf{X}_1 = \mathbf{X}_g$ so that it may be thought of as the design matrix in this simpler model for the genotypic means $\boldsymbol{\mu} = \mathbf{Z}_1 \boldsymbol{\gamma} + \mathbf{X}_1 \mathbf{u}_g$. In this stage, the marker effects as well as the genotypic mean are considered as fixed parameters and only the block effects are treated as random effects in the model (11). The adjusted mean ($\boldsymbol{\mu}$) can then be estimated easily from the first stage model (11) by classical likelihood based or suitable robust estimators. Following

Lourenco et al. (2020) [19], we would use the robust M-estimators with Huber’s weight function, as proposed and studied in Koller (2013) , in this first stage modeling to estimate $\boldsymbol{\mu}$ and associated variance components σ_b^2 and σ_e^2 , which are computed using the function `rlmer` from the R-package `robustlmm` (Koller, 2016 [17]) in our numerical experiments.

Once we get the estimated adjusted mean as $\hat{\boldsymbol{\mu}}$, it is then used as response in the second stage model given by

$$\hat{\boldsymbol{\mu}} = \mathbf{Z}_1\boldsymbol{\gamma} + \mathbf{X}_1\mathbf{u}_g + \tilde{\boldsymbol{\epsilon}}. \quad (12)$$

Assuming normality of these new error terms, we can now estimate $\boldsymbol{\gamma}$, σ_g^2 and the new error variance (say $\tilde{\sigma}_e^2$) by a suitable method. We propose to use the robust MDPDE for this purpose which can be obtained by solving the associated objective function as described previously. Once we get the robust estimates of all the parameters, we can obtain the robust estimates of the breeding values and prediction of phenotype effects as before following Eqns. (5) and (6), respectively.

2.4 Measuring Heritability

Mathematically, one can explain heritability to describe the difference of variation in a given trait that can be attributed to genetic variation. To measure how well the differences in genes account for differences in several kind of phenotypic traits, SNP-based heritability can be measured by the ratio of genetic variance (σ_g^2) to the total phenotypic variance as well as effect of environment (e.g., block effect); it lies between 0 to 1 (Wray et al., 2008 [31]). An estimate of the heritability of a trait is specific for populations as well as environment and changes with circumstances. Once we get the robust MDPDE of the variance parameters as $\hat{\sigma}_g^2$, $\hat{\sigma}_b^2$ and $\hat{\sigma}_e^2$, we can estimate the heritability following Wray et al. (2008 [31]) as

$$\hat{H}_p = \frac{\hat{\sigma}_g^2}{\hat{\sigma}_g^2 + \hat{\sigma}_b^2 + \hat{\sigma}_e^2}. \quad (13)$$

Note that, a high value 0.5 of \widehat{H}_p implies that on average half of the differences among phenotypes depend on genetic effect whereas a value close to 0 indicates that almost all of the variability in a trait among population are not under the genetic control and due to some other factors such as environment. We will empirically illustrate that use of the proposed MDPDE based variance estimators leads to a much stable and accurate heritability estimator than the existing measures.

3 Experiments

We will illustrate the superior performance of the proposed MDPDE based one and two-stage genomic prediction and heritability estimation over its recent competitors through extensive empirical experiments involving artificial datasets from different known model setups as well as a real-life breeding dataset. These simulation models and datasets are described in the following subsections, along with the performance metrics and the competing methods considered in all our experiments.

3.1 Experimental Setup: Artificial Datasets

We consider the simulation model setups resembling real-life situations similar to the one used in Lourenco et al. (2020) [19]. The underlying true data-generating model is taken as

$$y_{ijk} = \phi + g_i + b_{jk} + e_{ijk}, \quad (14)$$

where y_{ijk} is the phenotypic trait value for i -th genotype in the j -th block within k -th replicate, ϕ is the overall mean, $g_i = \sum_{l=1}^p z_{il}u_{gl}$ be the i -th genotypic value with z_{il} being the SNP-marker values and u_{gl} being the random effects generated from $N(0, \sigma_g^2)$ independently for all $j = 1, \dots, p$, b_{jk} is random block effect of j -th block of k -th replicate, computed based

on random effects associated with B blocks generated independently from $N(0, \sigma_b^2)$ along with the block design matrix, and e_{ijk} is random error generated from $N(0, \sigma_e^2)$ independently for all i, j, k . Each artificial dataset is generated with $k = 2$ replicates, $B = 39$ blocks and $p = 11646$ SNP markers, where 4 blocks contain 17 observations each and remaining 35 blocks contain 18 observations each leading to a total of $N = 715$ observations. We also take $\phi = 0.05$ and $\sigma_r^2 = 0$ but the other parameter values are taken differently for two setups as described below.

- First setup for one-stage approach: Following Lourenco et al. (2020) [19], we take $\sigma_b^2 = 6.27$, $\sigma_e^2 = 53.8715$ and $\sigma_g^2 = 0.005892$. Also the SNP marker values are generated as $z_{il} \sim \text{Bin}(1, 0.5)$ independently for all i, l .
- Second setup for two-stage approach: We take $\sigma_b^2 = 6.3148$, $\sigma_e^2 = 20.5$, $\sigma_s^2 = 0.5892$, and $z_{il} \sim \text{Bin}(2, 0.5) - 1$ independently for all i, l . Note that the minus 1 transform the binomial random variables so that each SNP marker takes value within $\{-1, 0, 1\}$.

Additionally, to illustrate the claimed robustness properties, we also generate artificial contaminated datasets from each simulated pure datasets (as described above) based on the following two contamination schemes:

- Random contamination: For each dataset, 5% phenotype observations are randomly chosen and replaced by its original value plus 5 times of the standard deviation of the residual error.
- Block contamination: For each dataset, 5 blocks are chosen randomly and the phenotype observations under those blocks (for all replications) are replaced by its original value plus 8 times of the standard deviation of the residual error.

3.2 Competitive Methods

For one-stage approach, we consider the methods described as RMLA and RMLV in Section 2.2 since they are shown to outperform other ridge regression based method by Hofheinz and Frisch (2014) [15]. However, since our primary focus is on robust inference against data contamination, we also compare our MDPDE based one-stage proposal with robust variants of RMLA and RMLV methods, which we would refer to as Rob-RMLA and Rob-RMLV; they are computed as in RMLA and RMLV, respectively, but using the Huber’s M-estimation approach (Koller, 2013 [16]) to estimate the parameters of the underlying linear mixed models instead of the likelihood based approach. These robust M-estimators are considered as classical benchmark against our proposed MDPDE based approach and are computed using the function `rlmer` from the R-package `robustlmm` (Koller, 2016 [17]).

For two-stage approach, we consider two existing robust competitors based on the work of Lourenço et al. (2020) [19] which we refer to as Rob1 and Rob2. Both these methods robustify only the first stage estimation by using the M-estimation approach as described in Section 2.3.3; but Rob1 uses the classical likelihood based estimation in the second stage modeling while Rob2 uses the robust M-estimation approach also in the second stage modelling. Recall that our proposal involves using the novel MDPDE based approach in the second stage modeling, while keeping the the first stage the same as Rob1 and Rob2.

3.3 Performance Metrics

In order to compare our proposed robust procedures with other existing benchmark methodologies of genomic predictions, we illustrate how close the estimated SNP marker effects ($\hat{\mathbf{u}}_g$) are in comparison to their original values (\mathbf{u}_g). So, for each artificial datasets, we compute the following two measures to illustrate the accuracy of genomic predictions:

- Pearsonian correlation coefficient $\hat{\rho} = \frac{Cov(\mathbf{u}_g, \hat{\mathbf{u}}_g)}{\sqrt{Var(\mathbf{u}_g)Var(\hat{\mathbf{u}}_g)}}$. Note that, the higher the value

of $\hat{\rho}$ higher is the predictive accuracy. This measure has been traditionally used in the related literature on genomic prediction (e.g., Lourenco et al., 2020 [19]).

- Median absolute deviation (MAD): a robust accuracy measure obtained as the median of the absolute difference between the predicted and the original marker effects. A low value of MAD close to zero indicates good performance. We use this measure to get a robust indication of deviations in the predicted trait values.

Note that the values of original \mathbf{u}_g are known for each artificial dataset through our simulations. We compute one value of $\hat{\rho}$ and MAD for each such dataset and report their average over 100 simulation replications in Tables 1–2.

Further, for each artificial dataset we get one final estimated value of the heritability measures \hat{H}_p . So, its performances under different existing and proposed approaches are examined by computing its Mean-square deviations (MSD) against its true value (known for each simulation setups) over 100 simulation replications (Table 3).

3.4 Real Data Analysis

We consider a real life breeding dataset generated by the International Maize and Wheat Improvement Center (CIMMYT) that contains 300 tropical maize lines in 10 blocks. Genotyping was conducted using 1135 SNP markers and the target phenotypic trait was grain yield assessed under severe drought stress and well-watered conditions. These two conditions are treated as two replicates in our analyses.

Within the 300 tropical maize lines, 16 lines have missing values in some SNP markers. To avoid computational challenges due to missing values, these datapoints are excluded from our analysis presented in this paper. Thus all methods of genomic prediction, both existing and proposed, and associated heritability estimation are applied to the remaining

284 datapoints to predict their grain yields from 1135 SNP markers suitably adjusted from the block and the replicate effects.

Since the underlying true values of the marker effects are unknown for a real dataset, in order to examine the performance of genomic predictions, we have computed the correlation coefficient $\hat{\rho}$ and the MAD between the original and the predicted values of the target phenotypic trait, which is grain yield in our dataset. The resulting values of $\hat{\rho}$ and MAD, along with the estimated heritability measure \hat{H}_p , are reported in Table 4 for different existing and proposed approaches.

4 Results and Discussions

4.1 Performance of the Proposed One-stage Method

Prediction accuracy measures, namely $\hat{\rho}$ and MAD, obtained for different one-stage procedures are presented in Table 1 for artificial datasets generated under the first simulation setup; the results for pure data as well as two contaminated scenarios are presented. Examining these results, we observe that the predictive accuracy of the proposed MDPDE based approach improves ($\hat{\rho}$ increases and MAD decrease) as α increases for both pure and contaminated situations; we get the best performances at $\alpha = 1$ in all three cases.

In the context of pure data, the proposed MDPDE based prediction method clearly outperforms other existing methods, namely both RMLA and RMLV and their robust counterparts. Although the performances of RMLV and Rob-RMLV are similar and better than RMLA and Rob-RMLA, they are yet significantly worse compared to our proposed MDPDE based approach which can achieve a high correlation of 0.8258 and a small MAD of 4.0430 at $\alpha = 1$.

Under contamination, the performances of classical method of RMLA and RMLV dete-

Table 1: Prediction accuracy measures obtained from different methods under the one-stage approach with artificial datasets (first setup).

Methods	Pure Data		Random Contamination		Block Contamination	
	$\hat{\rho}$	MAD	$\hat{\rho}$	MAD	$\hat{\rho}$	MAD
RMLA	0.5216	26.0458	0.3754	46.452	0.2635	59.9679
RMLV	0.7182	5.0496	0.6008	3.962	0.3678	54.744
Rob-RMLA	0.5183	8.4920	0.4139	4.904	0.2898	83.6232
Rob-RMLV	0.7308	5.0485	0.6008	3.862	0.3678	53.0470
Proposed MDPDE based approach						
$\alpha = 0.1$	0.8253	4.1311	0.7291	4.633	0.7180	13.2800
$\alpha = 0.3$	0.8255	4.1310	0.7292	4.632	0.7292	7.5980
$\alpha = 0.5$	0.8255	4.1301	0.7327	4.645	0.7341	6.9385
$\alpha = 0.7$	0.8257	4.0880	0.7395	4.604	0.7346	6.7637
$\alpha = 1$	0.8258	4.0430	0.7399	4.570	0.7352	6.4935

riorate drastically while those of their robust counterparts (Rob-RMLA and Rob-RMLV) remain somewhat stable in case of random contamination. In fact, Rob-MLVA achieves the smallest MAD value of 3.8624 among all methods under random contamination, making it a superior choice in terms of predictive accuracy measured by MAD. Nevertheless, the proposed MDPDE with $\alpha = 1$ produces a competitive MAD value of 4.570 and a significantly higher correlation value (0.7399 compared to 0.6 of Rob-RMLV) and hence remains the best method in terms of correlation measure of accuracy.

For block contaminated data, however, Rob-RMLV produces quite low correlation and high MAD values and the proposed MDPDE based procedure clearly emerges as the superior choice among all methods in terms of both predictive accuracy measures with improved performances as α increases. Comparing all set-ups and accuracy measures, the MDPDE based procedure remains most stable illustrating their significantly improved robustness properties over the existing procedures.

4.2 Performance of the Proposed Two-stage Method

The two-stage model analyses are conducted for the second experimental setup, and Table 2 presents the results of the predictive accuracy measures $\hat{\rho}$ and the MAD for both pure and contaminated data. Based on the tabulated values, it is evident that the MDPDE method provides the best estimates among all the second stage methods and the effect of α is almost negligible on their performances although $\alpha = 1$ gives marginally best results. Further, the results obtained by two-stage procedures are also significantly better than the one-stage results as intuitively expected; our proposed MDPDE based two-stage procedure can achieve an impressively high correlation of 0.9998, indicating strong correlation between the predicted and the original marker effects. Our method also leads to marker effect estimates having smaller deviations (MAD) from original values as compared with the existing Rob1 and Rob2 procedures. Further, the results from Table 2 also indicate that, when MDPDE is used in the second stage of the two-stage modeling, there is a significant reduction in MAD values and an increase in estimated correlation coefficients as compared with the existing for both block and response contaminated datasets. These observations suggest that the proposed MDPDE-based method exhibits even greater robustness compared to the only existing robust procedures.

Table 2: Prediction accuracy measures obtained from different methods under the two-stage approach with artificial datasets (second setup).

Methods	Pure Data		Random Contamination		Block Contamination	
	$\hat{\rho}$	MAD	$\hat{\rho}$	MAD	$\hat{\rho}$	MAD
Rob1	0.9623	3.76	0.8304	5.792	0.9231	3.8604
Rob2	0.9622	3.76	0.8405	5.789	0.9601	3.8130
Proposed MDPDE based approach						
$\alpha = 0.1$	0.9983	2.41	0.9914	2.679	0.9983	2.4074
$\alpha = 0.3$	0.9983	2.26	0.9915	2.678	0.9983	2.4069
$\alpha = 0.5$	0.9983	2.25	0.9923	2.678	0.9984	2.4066
$\alpha = 0.7$	0.9984	2.24	0.9924	2.675	0.9984	2.4064
$\alpha = 1$	0.9998	2.23	0.9925	2.679	0.9989	2.3062

4.3 One-stage vs. Two-stage Modeling

In this paper, we have examined two distinct types of model analysis, namely one-stage and two-stage modeling approaches. Based on the empirical analyses, it becomes evident that the two-stage approach yields superior predictive outcomes compared to the one-stage approach as the incorporation of different random effects in two different stages enhances the robustness of marker-effect estimates.

However, it is to be noted that we have employed two different simulation setups for illustrating the accuracy of one-stage and two-stage modeling approaches. This is because, as we learned from extensive empirical explorations, performances of the proposed MDPDE based robust two-stage approach relies heavily on both error variance (σ_e^2) and (random) SNP-marker-effect variance (σ_g^2). To illustrate this dependence, in Figure 1, we have plotted the correlation measure ($\hat{\rho}$) between the estimated and the original SNP marker-effects under the second experimental setup with different values of σ_e^2 and σ_g^2 for both pure and contaminated cases. It can be clearly seen from these figures (and many similar numerical explorations not presented here for brevity) that the accuracy of the proposed method in-




Figure 1: Plots of the accuracy measure $\hat{\rho}$ obtained by the MDPDE-based two-stage genomic prediction (with $\alpha = 1$) over different SNP marker-effect variance (σ_g^2) in the second experimental setup for two different σ_e^2 .

creases as σ_g^2 increases and σ_e^2 increases for both pure and contaminated datasets, which is equivalent to say that the genotypic signal-to-noise ratio of the underlying model increases. Therefore, for datasets with a high genotypic effect variance compared to the noise variance, genomic prediction using the proposed MDPDE based two-stage modeling approach will be more accurate and robust. In other cases with lower SNP effect variance or higher noise variances, the use of robust one-stage procedure is expected to produce better results.

4.4 Accuracy and Robustness of the Heritability Estimator

In Table 3, we present the mean square deviation (MSD) of the heritability estimate (\hat{H}_p), obtained by different existing and proposed methods in one-stage and two-stage approaches based on artificial pure and contaminated datasets in respective simulation set-ups. For one-stage analysis with the first experimental setup, the MSD of \hat{H}_p is minimized by the MDPDE method with $\alpha = 0.3$ which is deemed to be the best choice for pure dataset as well as for noisy dataset with random contamination. But for the artificial dataset with incorporated block contamination, the desired minimum heritability achieved by the MDPDE method with $\alpha = 1$ (which is 5.3731×10^{-10}) in one-stage analysis. The proposed MDPDE-based two-stage analysis, however, produces significantly lower MSDs, indicating more accurate heritability estimation, for all $\alpha \geq 1$ in cases of both pure and contaminated datasets.

4.5 Comparative Performances for the Maize Dataset

We have presented the correlation measure $\hat{\rho}$ between the predicted and the true responses and their MAD obtained by different one-stage and two-stage methods in Table 4 along with the estimated values of the heritability measure (\hat{H}_p). In case of this real data, we can see that the proposed MDPDE based one-stage modeling approach yields the best fit in terms of both correlation and MAD; the prediction performance further improves as α increases

Table 3: The MSD (in 10^{-10}) of heritability estimators obtained from different one-stage and two-stage procedures applied on pure and contaminated (cont.) experimental datasets.

Methods	Pure Data	Random Cont.	Block Cont.
One-stage Procedures			
RMLA	3.5235	5.378	8.3625
RMLV	98.1462	69.482	78.3400
ROB-RMLA	20.4525	9.7652	90.3215
ROB-RMLV	84.0401	59.3341	58.330
Proposed MDPDE based approach			
$\alpha = 0.1$	1.7173	8.0626	7.3721
$\alpha = 0.3$	1.9353	2.5649	6.5677
$\alpha = 0.5$	2.9861	2.5407	8.4062
$\alpha = 0.7$	3.5678	5.1520	9.4702
$\alpha = 1$	4.3594	5.7077	5.3731
Two-stage Procedures			
Rob1	19.7689	24.9496	25.8832
Rob2	60.2620	24.9373	25.0535
Proposed MDPDE based approach			
$\alpha = 0$	2.9425	2.6456	2.9885
$\alpha = 0.1$	0.6341	0.2944	0.6324
$\alpha = 0.3$	0.4843	0.2982	0.7317
$\alpha = 0.5$	0.6657	0.3182	0.8079
$\alpha = 0.7$	0.9088	0.3677	0.8006
$\alpha = 1$	0.5741	0.4768	1.0044

with $\alpha = 1$ giving the highest correlation of 0.828 and the lowest MAD value 0.47, indicating minimal deviation between observed and predicted values.

It's worth noting that the two-stage approach have not produced better predictions for this dataset; this is because of the low marker effect variance compared to error variance as evident from extremely low estimated values of heritability given in Table 4. As per our observations presented in Section 4.3, one-stage modeling performs well when dealing with data exhibiting low variance in random SNP effects, which is the case in this real dataset. Consequently, one-stage model analysis with MDPDE proved to deliver desirable prediction accuracy measures.

Table 4: Prediction accuracy and heritability measures obtained by different one-stage and two-stage procedures applied on Maize dataset.

One-stage Procedures				Two-stage Procedures			
Method	$\hat{\rho}$	MAD	\hat{H}_p	Method	$\hat{\rho}$	MAD	\hat{H}_p
RMLA	0.561	0.595	0.003770	Rob1	0.726	2.43	0.002310
RMLV	0.047	4.985	0.015600	Rob2	0.726	2.42	0.002050
ROB-RMLA	0.573	0.594	0.003760	Proposed MDPDE based approach			
ROB-RMLV	0.578	0.453	0.016400	$\alpha = 0.1$	0.294	0.75	0.000196
Proposed MDPDE based approach				$\alpha = 0.3$	0.282	1.88	0.000234
$\alpha = 0.1$	0.682	1.117	0.000666	$\alpha = 0.5$	0.307	2.30	0.000266
$\alpha = 0.3$	0.684	1.116	0.001420	$\alpha = 0.7$	0.311	2.26	0.000360
$\alpha = 0.5$	0.705	1.132	0.001550	$\alpha = 1$	0.314	2.14	0.000384
$\alpha = 0.7$	0.710	1.132	0.001505				
$\alpha = 1$	0.828	0.407	0.000706				

Further, as the heritability estimate in both one-stage and two-stage model analyses tends toward zero, it also suggests that the variability of this phenotypic trait within the tested maize population is mostly due to environmental factors and not under genetic control.

5 Conclusions

In this paper, we have devised robust estimation techniques for genomic prediction based on an underlying linear mixed model. In pursuit of more robust estimation, our proposal also employs the popular MDPDE of the parameter of mixed models in two distinct modeling structure with one and two-stage approaches. We have compared our proposed methods to existing approaches using estimated marker effects in simulation experiments and predicted trait values in a real dataset. Remarkably, for both pure and contaminated data, the two-stage modeling structure consistently outperformed other approaches in terms of prediction accuracy whenever the random SNP-marker effects have greater variance compared to the random noise in the model. Our simulation study across various experimental setups demonstrates the sensitivity of results to the parameter α in the MDPDE based pro-

positional with increasing value of α enhancing the accuracy and robustness of parameter and marker-effect estimation, ultimately yielding the more desirable predictive accuracy measure. Therefore, our comprehensive analysis establishes that, with a high correlation value and minimal deviation, the proposed MDPDE-based approaches with $\alpha = 1$ proves to be the optimal choice for both one-stage and two-stage model structures offering more accurate and reliable genomic prediction and heritability estimation compared to existing approaches.

The promising results and insights derived from this research pave the way for exciting future prospects in the field of linear mixed models and genotypic-phenotypic effect estimation. Future studies can delve deeper into refining and improving the robustness of the proposed estimation methods. This could involve exploring and devising new techniques that adapt dynamically to data quality and contamination levels which would help to frame valuable contributions to genomics, genetics, and associated fields, ultimately advancing our comprehension of the intricate relationship between genetic factors and traits.

We also hope to prepare an R-package for the proposed MDPDE-based procedures for genomic prediction and heritability estimation for dissemination among scientists and practitioners. They would then be able to apply our proposals easily in new experimental datasets to get more accurate and stable results, even in the presence of any possible noises and/or contamination, eventually leading to new insights and innovations.

Acknowledgments:

This research is supported by a SRG research grant no. SRG/2020/000072 from the Science and Engineering Research Board (SERB), Government of India, India.

References

- [1] BARBOSA, M. H. P., DE RESENDE, M. D. V., BRESSIANI, J. A., DA SILVEIRA,

- L. C. I., PETERNELLI, L. A., ET AL. Selection of sugarcane families and parents by reml/blup. *Crop Breeding and Applied Technology* 5, 4 (2005), 443.
- [2] BASU, A., HARRIS, I. R., HJORT, N. L., AND JONES, M. Robust and efficient estimation by minimising a density power divergence. *Biometrika* 85, 3 (1998), 549–559.
- [3] BASU, A., SHIOYA, H., AND PARK, C. *Statistical inference: the minimum distance approach*. CRC press, 2011.
- [4] BERNAL-VASQUEZ, A.-M., UTZ, H.-F., AND PIEPHO, H.-P. Outlier detection methods for generalized lattices: a case study on the transition from anova to reml. *Theoretical and Applied Genetics* 129 (2016), 787–804.
- [5] BÜHLMANN, P., KALISCH, M., AND MEIER, L. High-dimensional statistics with a view toward applications in biology. *Annual Review of Statistics and Its Application* 1 (2014), 255–278.
- [6] ESTAGHVIROU, S. B. O., OGUTU, J. O., AND PIEPHO, H.-P. Influence of outliers on accuracy estimation in genomic prediction in plant breeding. *G3: Genes, Genomes, Genetics* 4, 12 (2014), 2317–2328.
- [7] ESTAGHVIROU, S. B. O., OGUTU, J. O., AND PIEPHO, H.-P. How genetic variance and number of genotypes and markers influence estimates of genomic prediction accuracy in plant breeding. *Crop Science* 55, 5 (2015), 1911–1924.
- [8] FALKE, K. C., MAHONE, G. S., BAUER, E., HASENEYER, G., MIEDANER, T., BREUER, F., AND FRISCH, M. Genome-wide prediction methods for detecting genetic effects of donor chromosome segments in introgression populations. *BMC genomics* 15 (2014), 1–10.

- [9] GHOSH, A. Robust inference under the beta regression model with application to health care studies. *Statistical methods in medical research* 28, 3 (2019), 871–888.
- [10] GHOSH, A., AND BASU, A. Robust estimation for independent non-homogeneous observations using density power divergence with applications to linear regression.
- [11] GHOSH, A., AND BASU, A. Robust estimation for non-homogeneous data and the selection of the optimal tuning parameter: the density power divergence approach. *Journal of Applied Statistics* 42, 9 (2015), 2056–2072.
- [12] GHOSH, A., AND BASU, A. Robust estimation in generalized linear models: the density power divergence approach. *Test* 25 (2016), 269–290.
- [13] GIRAUD, C. *Introduction to high-dimensional statistics*. CRC Press, 2021.
- [14] GOEGEBEUR, Y., GUILLOU, A., AND VERSTER, A. Robust and asymptotically unbiased estimation of extreme quantiles for heavy tailed distributions. *Statistics & Probability Letters* 87 (2014), 108–114.
- [15] HOFHEINZ, N., AND FRISCH, M. Heteroscedastic ridge regression approaches for genome-wide prediction with a focus on computational efficiency and accurate effect estimation. *G3: Genes, Genomes, Genetics* 4, 3 (2014), 539–546.
- [16] KOLLER, M. *Robust estimation of linear mixed models*. PhD thesis, ETH Zurich, 2013.
- [17] KOLLER, M. robustlmm: an r package for robust estimation of linear mixed-effects models. *Journal of statistical software* 75 (2016), 1–24.
- [18] LEE, S., AND JO, M. Robust estimation for bivariate integer-valued autoregressive models based on minimum density power divergence. *Journal of Statistical Computation and Simulation* (2023), 1–29.

- [19] LOURENÇO, V. M., OGUTU, J. O., AND PIEPHO, H.-P. Robust estimation of heritability and predictive accuracy in plant breeding: evaluation using simulation and empirical data. *BMC genomics* 21, 1 (2020), 1–18.
- [20] MÖHRING, J., AND PIEPHO, H.-P. Comparison of weighting in two-stage analysis of plant breeding trials. *Crop Science* 49, 6 (2009), 1977–1988.
- [21] PIEPHO, H., OGUTU, J., SCHULZ-STREECK, T., ESTAGHVIROU, B., GORDILLO, A., AND TECHNOW, F. Efficient computation of ridge-regression best linear unbiased prediction in genomic selection in plant breeding. *Crop Science* 52, 3 (2012), 1093–1104.
- [22] PIEPHO, H.-P. Ridge regression and extensions for genomewide selection in maize. *Crop Science* 49, 4 (2009), 1165–1176.
- [23] PIEPHO, H.-P., MÖHRING, J., SCHULZ-STREECK, T., AND OGUTU, J. O. A stage-wise approach for the analysis of multi-environment trials. *Biometrical Journal* 54, 6 (2012), 844–860.
- [24] RESENDE, M. D. V. D. Software selegen-reml/blup: a useful tool for plant breeding. *Crop Breeding and Applied Biotechnology* 16 (2016), 330–339.
- [25] SARACENO, G., GHOSH, A., BASU, A., AND AGOSTINELLI, C. Robust estimation under linear mixed models: The minimum density power divergence approach. *arXiv preprint arXiv:2010.05593* (2020).
- [26] SHEN, X., ALAM, M., FIKSE, F., AND RÖNNEGÅRD, L. A novel generalized ridge regression method for quantitative genetics. *Genetics* 193, 4 (2013), 1255–1268.
- [27] TANAKA, E. Simple outlier detection for a multi-environmental field trial. *Biometrics* 76, 4 (2020), 1374–1382.

- [28] UFFELMANN, E., HUANG, Q. Q., MUNUNG, N. S., DE VRIES, J., OKADA, Y., MARTIN, A. R., MARTIN, H. C., LAPPALAINEN, T., AND POSTHUMA, D. Genome-wide association studies. *Nature Reviews Methods Primers* 1, 1 (2021), 59.
- [29] WAINWRIGHT, M. J. *High-dimensional statistics: A non-asymptotic viewpoint*, vol. 48. Cambridge university press, 2019.
- [30] WESTHUES, C. C., MAHONE, G. S., DA SILVA, S., THORWARTH, P., SCHMIDT, M., RICHTER, J.-C., SIMIANER, H., AND BEISSINGER, T. M. Prediction of maize phenotypic traits with genomic and environmental predictors using gradient boosting frameworks. *Frontiers in plant science* 12 (2021), 699589.
- [31] WRAY, N., AND VISSCHER, P. Estimating trait heritability. *Nature education* 1, 1 (2008), 29.
- [32] ZHU, H., AND ZHOU, X. Statistical methods for snp heritability estimation and partition: A review. *Computational and Structural Biotechnology Journal* 18 (2020), 1557–1568.